

# Hypersaline tidal flats as important “Blue Carbon” systems: A case study from three ecosystems

5 Dylan R. Brown<sup>1</sup>, Humberto Marotta<sup>2,3,4</sup>, Roberta B. Peixoto<sup>2,3</sup>, Alex Enrich-Prast<sup>2,5,6</sup>, Glenda C. Barroso<sup>3</sup>, Mario L. G. Soares<sup>7</sup>, Wilson Machado<sup>3</sup>, Alexander Pérez<sup>3,8</sup>, Joseph M. Smoak<sup>9</sup>, Luciana M. Sanders<sup>10</sup>, Stephen Conrad<sup>1</sup>, James Z. Sippo<sup>1,10,11</sup>, Isaac R. Santos<sup>1,12</sup>, Damien T. Maher<sup>1,10,11</sup>, Christian J. Sanders<sup>1,13\*</sup>

10 <sup>1</sup>National Marine Science Centre, School of Environment, Science and Engineering, Southern Cross University, PO Box 4321, Coffs Harbour, NSW 2450, Australia

<sup>2</sup>Ecosystems and Global Change Laboratory (LEMG-UFF) / International Laboratory of Global Change (LINCGlobal), Biomass and Water Management Research Center (NAB), Universidade Federal Fluminense, Av. Edmundo March, s/n°, Niterói, RJ, 24210-310, Brazil

15 <sup>3</sup>Graduate Program in Geosciences (Environmental Geochemistry), Department of Geochemistry, Universidade Federal Fluminense, Niterói, RJ, 24020-141, Brazil

<sup>4</sup>Physical Geography Laboratory (LAGEF-UFF), Department of Geography, Graduate Program in Geography, Universidade Federal Fluminense (UFF), Av. Gal. Milton Tavares de Souza, s/n°, Niterói, RJ, 24210-346, Brazil.

<sup>5</sup>Department of Thematic Studies–Environmental Change, Linköping University, 581 83, Linköping, Sweden.

<sup>6</sup>Department of Botany, Universidade Federal do Rio de Janeiro, 21941-902, Rio de Janeiro, Brazil.

20 <sup>7</sup>Laboratory For Mangrove Studies (NEMA-UERJ) / International Laboratory of Global Change (LINCGlobal), Department of Biological Oceanography, Faculty of Oceanography, Universidade do Estado do Rio de Janeiro (UERJ), Rua São Francisco Xavier. 524, sala 4019-E, Rio de Janeiro, RJ, 20550-900, Brazil

25 <sup>8</sup>Universidad Peruana Cayetano Heredia, Centro de investigación para el desarrollo integral y sostenible (CIDIS), Facultad de Ciencias y Filosofía, Laboratorios de investigación y desarrollo (LID), Laboratorio de Biogeociencias, Av. Honorio Delgado 430, Urb. Ingeniería, Lima 31 – Perú.

<sup>9</sup>Environmental Science, University of South Florida, St. Petersburg, Florida 33701, USA

<sup>10</sup>Southern Cross Geoscience, Southern Cross University, PO Box 157, Lismore NSW 2480, Australia

<sup>11</sup>School of Environment, Science and Engineering, Southern Cross University, PO Box 157, Lismore, NSW, 2480, Australia.

<sup>12</sup>Department of Marine Sciences, University of Gothenburg, Gothenburg, Sweden

30 <sup>13</sup>State Key Laboratory of Estuarine and Coastal Research and Institute of Eco-Chongming, East China Normal University, Shanghai 201100, P. R. China

\*Corresponding author: Christian Sanders ([christian.sander@scu.edu.au](mailto:christian.sander@scu.edu.au))

35 **Abstract.** Hypersaline tidal flats (HTFs) are coastal ecosystems with freshwater deficits often occurring in arid or semi-arid regions near mangrove supratidal zones with no major fluvial contributions. Here, we estimate that organic carbon (OC), total nitrogen (TN) and total phosphorus (TP) are being buried at rates averaging 21 ( $\pm$  6), 1.7 ( $\pm$  0.3), and 1.4 ( $\pm$  0.3) g m<sup>-2</sup> y<sup>-1</sup>, respectively, during the previous century in three contrasting HTFs systems, one in Brazil (eutrophic) and two in Australia (oligotrophic). Although these rates are lower than those from nearby mangrove, saltmarsh and seagrass systems, the  
40 importance of HTFs as sinks for OC, TN and TP may be significant given their extensive coverage. Despite the measured short-term variability between net air-saltpan CO<sub>2</sub> influx and emission estimates found during the dry and wet season in the

Brazilian HTF, the only site with seasonal CO<sub>2</sub> fluxes measurements, the OC sedimentary profiles over several decades suggests efficient OC burial at all sites. Indeed, the stable isotopes of OC and TN ( $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$ ) along with C:N ratios show that microphytobenthos are the major source of the buried OC in these HTFs. Our findings highlight a previously unquantified carbon as well as nutrient sink and suggest that coastal HTF ecosystems could be included in the emerging blue carbon framework.

## 1 Introduction

Hypersaline tidal flats (HTFs), supratidal flats, salt pans, sabkhas, and salt flats are all terms used to define the shallow coastal ecosystems on the upper fringe of fluvio-marine plains in estuaries showing freshwater deficits (Ridd and Stieglitz, 2002; Albuquerque et al., 2014). These environments are generally located in an intermediary position between mangrove forests or saltmarshes and the terrestrial environment which are common in many tropical arid, and to a lesser extent non arid, intertidal zones. These systems occur in many regions around the world including northern Australia, Africa, Spain, the Gulf of Mexico and throughout Brazil where they are referred to as *apicum* ecosystems (Ridd and Stieglitz, 2002; Albuquerque et al., 2013; Albuquerque et al., 2014; Soares et al., 2017). In arid and semi-arid estuaries (Ridd and Stieglitz, 2002) or humid tropical supratidal zones with less fluvial contribution (Soares et al., 2017), HTF ecosystems cover an area that exceeds mangrove forests and occupy a substantial proportion of tropical intertidal zones. HTFs occupy the area just below the highest astronomical tides and are thus only flooded for short periods of the year (Ridd and Stieglitz, 2002; Bento et al., 2017). Evaporation, the flat topography, and pronounced hydraulic deficit results in hypersaline conditions with salinity as high as five times that of seawater (Ridd and Stieglitz, 2002; Shen et al., 2018).

Despite the extreme conditions and the apparent absence of vegetation, microphytobenthos are commonly found on the surface of HTFs (usually in the form of microbial mats dominated by cyanobacteria from the Oscillatoriales order including *Microcoleus* spp., *Leptolyngbya* spp. and *Lyngbya* sp.) (Adame et al., 2012; Masuda and Enrich-Prast, 2016). These microphytobenthos are well adapted to the extreme conditions (Paerl et al., 2000) and are considered to be the main primary producers in HTFs. Similar to traditional vegetated blue carbon systems (Ouyang and Lee, 2014; Sanders et al., 2016a; Macreadie et al., 2019; Serrano et al., 2019), these microphytobenthos ~~and~~ are capable of high rates of carbon (C) and nitrogen (N) fixation from the atmosphere, particularly after periods of flooding and/or rainfall (Chairi et al., 2010; Adame et al., 2012; Burford et al., 2016). Their ability to sequester and potentially store carbon and nutrients in their soils for long periods of time (centuries to millennia) makes them noteworthy contenders to be included in blue carbon framework (Lovelock and Duarte, 2019). Upon inundation, the fixed C, N and other nutrients such as phosphorus (P) may be leached from the microbial mats and transported to adjacent coastal areas, where nutrient subsidies can enhance the overall productivity of the receiving ecosystems (Adame et al., 2012; Burford et al., 2016).

Given the few studies on HTFs, there is limited understanding on the role that these ecosystems play in the coastal zone, and whether they are currently under threat from global change (Halpern et al., 2008; Martinez-Porchas and Martinez-

Cordova, 2012). To date, there has been large scale destruction and degradation of these systems on a global scale as a result of anthropogenic pressures on coastal areas including infilling for urban and agricultural/aquaculture development (Halpern et al., 2008). Although there has been the implementation of various laws in some parts of the world to prevent the loss of coastal vegetated systems, these legislations rarely extend to protect HTFs that are viewed as being ecological deserts with no obvious vegetation (Albuquerque et al., 2013). Furthermore, the landward encroachment of mangrove forests as a response to rising sea levels, coupled to barriers preventing landward migration of HTF (i.e. the “coastal squeeze”), may also contribute to the loss of these ecosystems (Alongi, 2008; Saintilan et al., 2014; Kelleway et al., 2017).

Given the substantial areal extent of these HTFs and the fact that they remain relatively undisturbed in many regions around the world, HTFs may have unrecognized ecological values (Burford et al., 2016). However, information on OC, nitrogen and phosphorus burial and sediment CO<sub>2</sub> fluxes from these ecosystems remains scarce (e.g. Bento et al., 2017; Schile et al., 2017). Determining if HTFs are a source or sink of carbon is critical to understanding their importance and value in regards to climate change and coastal carbon sequestration (Lovelock and Duarte, 2019). Here, we quantify carbon and nutrient burial and atmospheric CO<sub>2</sub> fluxes in HTFs in Australia and Brazil. We hypothesize that microphytobenthos in HTFs sequester CO<sub>2</sub> from the atmosphere and a portion of this organic matter (and associated nitrogen and phosphorus) is buried, similar to the traditional vegetated blue carbon systems.

## 2 Methods

### 2.1 Study site

This study was conducted in three tropical HTFs in Australia and Brazil (Fig. 1). In Australia, the HTF study sites were located near Karumba, Queensland (17°25'12"S, 140°51'36"E) and Curtis Island, Gladstone, Queensland (Site 1: 23°45'41"S, 151°16'34"E; Site 2: 23°45'18"S, 151°16'49"E), and in Brazil the study site was located in Guaratiba, Rio de Janeiro (23°00'29"S, 43°36'31"W).

In Australia, the Karumba HTF is located adjacent to the oligotrophic mouth of the Norman River estuary in the southeastern coast of the Gulf of Carpentaria. The study site consists of a large continuous HTF (16.9 km<sup>2</sup>) in the high upper intertidal zone. The southeastern Gulf of Carpentaria has a diurnal tidal cycle (typical range <0.1 – 4.5 m) and a 23-year average annual rainfall of 833 mm (based on monthly averages from 1938 to 2010) with most falling in the summer monsoon period (790 mm from December to March) (Bureau of Meteorology, 2019). The 19-year average maximum monthly temperatures (from 1993 to 2019) vary from 27 °C in the dry winter months to 33 °C in the wet summer months (Bureau of Meteorology 2019). A narrow strip of mangrove forest followed by extensive tidal mudflats fringe the HTFs on the seaward side.

The Gladstone Harbour experiences similar tidal and climatic conditions to Karumba with semidiurnal tides (typical range 0.1 – 4.7 m) and a 25-year average annual rainfall of 846 mm (based on monthly averages from 1994 to 2019); with most falling also in the summer monsoon period (537 mm from December to March) (Bureau of Meteorology, 2019). The 26-

year average maximum monthly temperatures (from 1993 to 2019) vary from 23 °C in the winter months to 31 °C in the summer months (Bureau of Meteorology, 2019). The sheltered strait between Curtis Island and the mainland of Australia is largely occupied by mangrove forests and large continuous expanses of HTFs. The Gladstone site contained two HTF study areas; Site 1 (2.84 km<sup>2</sup>) was situated in the higher tidal area and is inundated less frequently and for shorter periods of time  
110 than Site 2 (0.95 km<sup>2</sup>).

In Brazil, the tropical HTF was located in the Guaratiba State Biological Reserve, ~40 km south of Rio de Janeiro City which forms part of the Sepetiba Bay estuary system. This conservation area is surrounded by the urban expansion area of Rio de Janeiro City, and Sepetiba Bay receives discharges of nutrients and organic matter from its watershed dominated by agriculture, pasture and urban uses (Rezende et al., 2010). The HTF covers an area of approximately 7.4 km<sup>2</sup> equivalent to  
115 almost 36% of the fringing mangrove forest (Estrada et al., 2013; Soares et al., 2017) (Table 1). There is little variation in topography and the tidal range is 0.1 – 2.0 m (Masuda and Enrich-Prast, 2016; Bento et al., 2017). The 32-year monthly average rainfall and temperature vary from 36 mm and 21 °C in dry winter months and 138 mm and 27 °C in rainy summer months, reaching annual averages of accumulated rainfall of 1058 mm (Estevam, 2019) (Table 1).

## 2.2 Sediment core sampling and analysis

120 Sediment cores (one core per site for a total of four cores) were collected from the middle of HTFs by using either a 50 cm long, 5 cm diameter Russian Peat Auger (Karumba core) or by inserting a PVC tube (8.7 cm diameter) into the substratum using manual percussion (Gladstone and Guaratiba cores). Only cores with no observed compaction were retained for further analysis. The sediment cores were sectioned at 1 cm intervals (with the exception of the Karumba core which was sectioned at 2 cm intervals). Dry bulk density (DBD, g cm<sup>-3</sup>) was determined as the dry sediment weight (g) divided by the initial volume  
125 (cm<sup>3</sup>) (Ravichandran et al., 1995). From the original dry section, a non-homogenized portion was rewetted and treated with 30% hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) to remove organic matter without altering grain size. A solution of sodium hexametaphosphate was used as a deflocculating agent to separate aggregates prior to grain size analysis. Grain size analyses were conducted using a CILAS 1090L diffraction laser unit or wet sieving following the methods used by Conrad et al., (2019). Total phosphorous (TP) was measured after acid digestion (H<sub>2</sub>O/HF/HClO<sub>4</sub>/HNO<sub>3</sub>, 2:2:1:1) using a Perkin Elmer ELAN DRCe ICPMS.

130 Organic carbon (OC) and total nitrogen (TN) stable isotope ratios of mangrove leaves, microphytobenthos, and HTF sediments were measured to identify the sources of organic matter (OM) contributing the sediment column at each site. Fresh green leaves from mangrove trees (n = 3 for each dominant species: *Rhizophora mangle*, *Avicenna shaueriana*, and *Laguncularia racemose*) were collected at 1-2m above the soil, and washed with deionized water soon after sampling in the Brazilian HTF. Samples were then lyophilized, crushed, sieved, and ~6-8 mg encapsulated in tin capsules to determine the  
135 OC, TN and their isotopic composition ( $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$ ). Microphytobenthos samples, in the form of dense algal mats, were collected from the surface of HTF sediments, scrapped and thoroughly washed with deionized water to avoid sediment contamination. A total of 6 microphytobenthos samples were collected and analyzed (3 from Brazil, 2 from Karumba, and 1 from Gladstone). A homogenized portion was acidified to remove carbonate material, washed in deionized water, dried (60

140 °C) and then ground to powder for OC and  $\delta^{13}\text{C}$  analyses using a Leco Flash Elemental Analyzer coupled to a Thermo Fisher  
Delta V IRMS (isotope ratio mass spectrometer). A non-acidified homogenized portion was also analyzed for TN and  $\delta^{15}\text{N}$ .  
Analytical precision: C = 0.1%, N = 0.1%,  $\delta^{13}\text{C}$  = 0.1‰ and  $\delta^{15}\text{N}$  = 0.15 ‰. We assess whether HTFs accumulate carbon and  
then to compare HTF with well-established, nearby mangrove systems.

145 Radionuclides from the uranium-238 ( $^{238}\text{U}$ ) decay series were measured in a high-purity germanium (HPGe) gamma  
detectors, a planar for the Gladstone and Guaratiba and a well detector for the Karumba samples. ~~Radionuclides from the  
uranium-238 ( $^{238}\text{U}$ ) decay series were measured in a high-purity germanium (HPGe) planar or well gamma detectors.~~ Identical  
geometry was used for all samples and sample dry weights were between 20 and 30 g. Sealed and packed samples were set  
aside for at least 21 days to allow for radon-222 ( $^{222}\text{Rn}$ ) ingrowth and to establish secular equilibrium between radium-226  
( $^{226}\text{Ra}$ ) and its granddaughter lead-214 ( $^{214}\text{Pb}$ ). Lead-210 ( $^{210}\text{Pb}$ ) activity was determined by the direct measurement of the 46.5  
KeV gamma peak.  $^{226}\text{Ra}$  activity was determined via the  $^{214}\text{Pb}$  daughter at 351.9 KeV.  $^{210}\text{Pb}$  and  $^{226}\text{Ra}$  activities were calculated  
150 by multiplying the counts per minute by a correction factor that includes the gamma-ray intensity and detector efficiency  
determined from NIST Rocky Flat soils reference material. ~~were calculated by multiplying the counts per minute by a correction~~  
~~factor that includes the gamma ray intensity and detector efficiency determined from standard calibrations.~~ Excess  $^{210}\text{Pb}$  was  
used to determine ages of sediment intervals using the Constant Initial Concentration (CIC) model (Appleby and Oldfield,  
1992). Mass accumulation rates were multiplied by the percent OC, N and TP to calculate burial rates.

### 155 2.3 Air-sediment gas flux measurements

CO<sub>2</sub> fluxes at the air-sediment interface were measured in July 2009 & 2010 and February 2015 (Guaratiba, Brazil), August  
2016 & 2018 (Karumba, Australia), and June 2018 (Gladstone, Australia), encompassing the annual variation of emissions  
between dry and rainy seasons in the HTF in Brazil and non-monsoon months in Australia. In all sampling sites, we used  
sediment chambers connected in a closed system with an infrared or cavity ring-down analyzer as reported in Lovelock (2008).  
160 The sediment chambers were composed of transparent plexiglass (light chamber) or an opaque material such as PVC or covered  
by layers of aluminium foil (dark chamber) for measurements of light and dark air-sediment CO<sub>2</sub> fluxes respectively (Leopold  
et al., 2015). Before each measurement, the chambers were gently pushed into the sediment (~2 cm) to form a gas tight seal.  
Each short-term incubation lasted 5-15 min to achieve a linear change in CO<sub>2</sub> concentration within the chambers, and was  
associated with a maximum increased temperature ~ 2°C in relation to external conditions, indicating no bias due to warming  
165 and subsequent changes in the inner pressure and biological activity. ~~Each short term incubation lasted 5-15 min to achieve a~~  
~~linear change in CO<sub>2</sub> concentration within the chambers.~~ Gas concentrations were measured using either a Los Gatos Research  
(LGR) Ultra-Portable Greenhouse Gas Analyzer (UGGA) or Picarro G4301 GasScouter recorded at 1-second intervals in the  
Australian sites, and a PPSystems EGM-4 or a Vaisala GMT222 at 1-minute intervals in the Brazilian sites. Equipment were  
previously calibrated with CO<sub>2</sub> standards of 400 and 1000 ppm in the laboratory.

170 CO<sub>2</sub> fluxes were measured at dark and light conditions in Brazil (n = 51 and 94, respectively) and Australian HTFs  
(n = 46 and 32, respectively). The air-sediment CO<sub>2</sub> fluxes were calculated from the maximum linear change in CO<sub>2</sub>

concentration over the duration of the measurement using the following formula (Rosentreter et al., 2017 and references therein):

$$F = (s(V/RT_{\text{air}}))A \quad (1)$$

175 where  $s$  is the regression slope for each chamber incubation deployments ( $\text{ppm sec}^{-1}$  or  $\text{ppm min}^{-1}$ , converted to  $\text{ppm h}^{-1}$ ),  $V$  is the chamber volume ( $\text{m}^3$ ),  $R$  is the universal gas constant,  $T_{\text{air}}$  is the air temperature inside the chamber ( $K$ ),  $A$  is the surface area of sediment inside the chamber ( $\text{m}^2$ ). Negative values represent net sediment  $\text{CO}_2$  uptake while those positive represent net  $\text{CO}_2$  emission from sediments to the atmosphere. We assume that pressure in the chamber is 1 atm. To determine the net ecosystem exchange (NEE), we integrate diurnal and night fluxes from light and dark chambers for each sampling day, 180 respectively. To test the normality of  $\text{CO}_2$  emissions data we performed a Komogorov-Smirnov test. For non-normally distributed data, a Mann Whitney test (significance level;  $p < 0.05$ ) was undertaken to compare light and dark fluxes at the combined Brazil and Australian samples, and also to compare wet and dry season Brazil fluxes.

### 3 Results

#### 3.1 Sediment accretion rates (SAR)

185 All four sediment profiles showed a net down-core decrease in excess  $^{210}\text{Pb}$  activity reaching background levels at the bottom of each sediment core (Fig. 2), enabling the use of the CIC  $^{210}\text{Pb}$  dating methodology. All cores were dated back to between 50 and 110 years with constant sediment accretion rates estimated at  $0.11 \pm 0.05$  (1903),  $0.18 \pm 0.06$  (1955),  $0.21 \pm 0.05$  (1931), and  $0.23 \pm 0.05 \text{ cm yr}^{-1}$  (1964) for Guaratiba, Karumba, Gladstone site 1, and Gladstone site 2 HTF sediment cores, respectively.

#### 190 3.2 Carbon, nitrogen and phosphorus burial rate estimates

Most of the parameters remained relatively constant throughout the sediment profiles, with no clear vertical trends in grain size, OC, TN or TP (Fig. 3). Sand content was generally  $< 20\%$  and OC, TN and TP contents ranged from 0.09 to 1.40, 0.01 to 0.16, and 0.02 to 0.12%, respectively across all sites and depth intervals (Fig. 3). By multiplying the average sedimentation rate, DBD, and OC content in these cores, we obtained carbon burial rates of  $17.8 (\pm 0.8)$ ,  $31.7 (\pm 4.3)$ ,  $11.3 (\pm 2.1)$ , and  $25.2$  195  $(\pm 2.9) \text{ g m}^{-2} \text{ y}^{-1}$  in the Guaratiba, Karumba, Gladstone site 1, and Gladstone site 2 cores respectively for the past ~50 years (Table 2). Average TN burial rates were  $2.3 (\pm 0.2)$ ,  $2.8 (\pm 0.3)$ ,  $0.8 (\pm 0.1)$ , and  $1.2 (\pm 0.1) \text{ g m}^{-2} \text{ y}^{-1}$  and average TP burial rates were  $2.0 (\pm 0.1)$ ,  $1.3 (\pm 0.1)$ ,  $1.4 (\pm 0.0)$ , and  $1.4 (\pm 0.3) \text{ g m}^{-2} \text{ y}^{-1}$  in the Guaratiba, Karumba, Gladstone site 1, and Gladstone site 2 cores, respectively (Table 2).

#### 3.3 Organic matter source

200 To assess the source of organic matter (OM), sediment, HTF microphytobenthos, and nearby mangrove end member samples were analysed for  $\delta^{13}\text{C}$  stable isotopes and cross-plotted against molar C:N ratios (Fig. 4). Microphytobenthos samples showed a small spread in  $\delta^{13}\text{C}$  and molar C:N ratios ranging from  $-13.4$  to  $-19.0 \text{ ‰}$  and 7.9 to 14.8 respectively (Fig. 4). Similarly, values of  $\delta^{13}\text{C}$  and molar C:N ratios showed little down-core variation in both the Guaratiba ( $-17.7$  to  $-18.4 \text{ ‰}$  and 7.6 to 9.7

respectively) and Karumba sediment cores (-15.5 to -20.5 ‰ and 10.5 to 14.6 respectively). In contrast, both the Gladstone  
205 sediment cores showed a considerable range in the  $\delta^{13}\text{C}$  and molar C:N values (-20.1 to -24.2 ‰ and 13.6 to 21.8 at site 1; -  
16.6 to -24.4 ‰ and 10.7 to 34.8 at site 2). Higher  $\delta^{15}\text{N}$  and lower C:N ratio values were noted in the Guaratiba HTF compared  
to other sites (Fig. 4).

### 3.4 CO<sub>2</sub> fluxes at the air-sediment interface

Median ( $\pm$  SE) hourly CO<sub>2</sub> fluxes measured at the air-sediment interface varied with HTF location and type of measurement  
210 (Light vs Dark) (Fig. 5). Median light CO<sub>2</sub> values were -2.1 ( $\pm$  4.1), 38.2 ( $\pm$  0.0), 13.7 ( $\pm$  1.9), and 29.3 ( $\pm$  2.1) mg C m<sup>-2</sup> h<sup>-1</sup>  
for the Brazilian (Guaratiba) and Australian (Karumba, Gladstone site 1, and Gladstone site 2) HTFs, respectively (Fig. 5).  
Median CO<sub>2</sub> fluxes in the dark chambers were significantly higher than those estimated in the light chambers (Mann Whitney  
test;  $p < 0.05$ ), *i.e.* 2.1 ( $\pm$  1.0), 39.6 ( $\pm$  9.2), 45.7 ( $\pm$  4.5), and 34.6 ( $\pm$  3.1) mg C m<sup>-2</sup> h<sup>-1</sup> for Guaratiba, Karumba, Gladstone site  
1, and Gladstone site 2, respectively (Fig. 5). In Brazil, significantly higher CO<sub>2</sub> uptake rates (median  $\pm$  SE) were recorded in  
215 the light chambers during the dry season compared to the wet season ( $-3.0 \pm 1.3$  and  $48.9 \pm 7.2$  mg C m<sup>-2</sup> h<sup>-1</sup>, respectively;  
Mann Whitney;  $p < 0.05$ ).

## 4 Discussion

### 4.1 C, N and P burial in HTFs versus vegetated blue carbon ecosystems

Considerable differences in OC burial rates between the two Gladstone sites were observed in this study. The likely difference  
220 between sites is due to the tidal area of each site, *i.e.* upper vs lower tidal areas are expected to accumulate carbon at different  
rates (Sanders et al., 2014). By averaging the sediment burial rates on a centennial scale (*i.e.* entire core) of the four sediment  
cores across all the study sites, we estimate that HTF ecosystems accumulate OC, TN and TP at rates of 21 ( $\pm$  6), 1.7 ( $\pm$  0.3),  
and 1.4 ( $\pm$  0.3) g m<sup>-2</sup> y<sup>-1</sup>, respectively. These centennial scale averages reduce short term variations allowing comparisons with  
saltmarsh, mangrove forests, and seagrass beds which have been studied extensively using similar methodologies and  
225 timeframes (McLeod et al., 2011). The average OC accumulation rates in HTF systems were ~12, ~8 and ~7 fold lower than  
the global averages reported for saltmarsh ( $245 \pm 26$  g m<sup>-2</sup> y<sup>-1</sup>; Ouyang and Lee (2014)), mangrove forests ( $163 \pm 40$  g m<sup>-2</sup> y<sup>-1</sup>;  
Breithaupt et al. (2012)), and seagrasses ( $138 \pm 38$  g m<sup>-2</sup> y<sup>-1</sup>; McLeod et al. (2011)), respectively. These lower burial rates may  
be related to the lower organic matter supply (including no contribution from below ground productivity) and/or lower  
sediment accretion rates than the traditional blue carbon systems. Furthermore, the reduced structural complexity and ability  
230 of the microalgae to trap sediments, the lower primary production rates, the lack of underground root protection, and the fact  
that microalgae organic material is more labile can explain the lower burial and sediment accretion rates of HTFs than  
traditional, vegetated blue carbon systems.

Hypersaline tidal flats can be a significant source of nutrient export to adjacent ecosystems which may potentially  
fuel primary productivity in nutrient-limited receiving marine ecosystems (Lovelock et al., 2010; Burford et al., 2016). Here,  
235 we find that these HTF ecosystems are also sites for the long-term storage of nitrogen and phosphorus (Table 2). The high TP  
burial rates compared to TN observed are likely due to the lack of anthropogenic nitrogen inputs observed in other systems.

Although the average TN accumulation rates reported here ( $1.7 \pm 0.3 \text{ g m}^{-2} \text{ y}^{-1}$ ) were also relatively low when compared to mangrove sediments ( $12.5 \pm 1.9 \text{ g m}^{-2} \text{ y}^{-1}$ ; Breithaupt et al. (2014)), the average TP accumulation rates in both Australian pristine ( $1.4 \pm 0.3 \text{ g m}^{-2} \text{ y}^{-1}$ ) and Brazilian eutrophic HTFs ( $2.0 \pm 0.1 \text{ g m}^{-2} \text{ y}^{-1}$ ) were higher than conserved mangrove sites with little anthropogenic nutrient discharges ( $0.5 \pm 0.2 \text{ g m}^{-2} \text{ y}^{-1}$ ; Breithaupt et al. (2014)). However, the HTF TP accumulation rates were not as high as those found in anthropogenically disturbed mangrove sites such as the heavily urbanized Jiulongjiang Estuary, China with TP accumulation rates reaching  $48.1 \text{ g m}^{-2} \text{ y}^{-1}$  (Alongi et al., 2005). Anthropogenic activities such as urbanization and major industrial developments drive degradation and increased primary production in mangrove forests (Sanders et al., 2014). Nutrients such as iron and phosphorus may be limiting to mangrove growth (Alongi, 2010; Reef et al., 2010), and those forests receiving high nutrient loads from highly concentrated anthropogenic nutrient discharges accumulate OC, TN, and TP at rates much higher than those from the undisturbed mangrove (Sanders et al., 2014). Nevertheless, the nitrogen and phosphorus burial in HTFs as shown here over long periods of time may play an important role in nutrient sequestration from other coastal anthropogenic activities, e.g. shrimp farming activities (Ashton, 2008; Marchand et al., 2011).

By upscaling the average OC, TN and TP accumulation results for the past century in this study to the regional areas of HTFs, we can provide a first-order estimate of the amount of OC, TN and TP being stored annually in these HTFs. Ridd and Stieglitz (2002) identify the areal extent of both HTFs and mangrove forests for five estuaries in Queensland, Australia with the HTFs identified as having a ~10-fold higher areal extent ( $279 \text{ km}^2$ ) than mangrove forests ( $29 \text{ km}^2$ ) over the five estuaries. In these estuaries alone, HTFs would contribute to the annual accumulation of approximately  $5.76 \pm 1.57$ ,  $0.46 \pm 0.09$  and  $0.40 \pm 0.08 \text{ Gg y}^{-1}$  of OC, TN and TP respectively which is similar to the contribution of mangrove forests ( $4.73 \pm 1.16$ ,  $0.36 \pm 0.06$  and  $0.26 \pm 0.03 \text{ Gg y}^{-1}$  for OC, TN and TP respectively) when based on global average accumulation rates (Breithaupt et al., 2012; Breithaupt et al., 2014). In contrast to Australia, the mangrove forests of Guaratiba ( $20.9 \text{ km}^2$ ) have been identified to have a ~3 fold higher area than local HTFs ( $7.4 \text{ km}^2$ ) (Soares et al., 2017); resulting in annual OC, TN and TP accumulation in HTFs ( $0.15 \pm 0.04$ ,  $0.01 \pm 0.00$  and  $0.01 \pm 0.00 \text{ Gg y}^{-1}$  respectively) equivalent to 4-6% of those estimated for mangrove forests ( $3.41 \pm 0.84$ ,  $0.26 \pm 0.04$  and  $0.19 \pm 0.02 \text{ Gg y}^{-1}$  for OC, TN and TP respectively) when based on global average accumulation rates (Breithaupt et al., 2012; Breithaupt et al., 2014).

Our estimates suggest HTFs are capable of long-term storage of OC, TN and TP and given their large areal extent, have the potential to store as much OC, TN and TP as traditional coastal blue carbon systems in arid regions such as Queensland, Australia. To improve these estimates, there is clearly a need for determining carbon and nutrient accumulation rates from additional coastal HTFs and assess their areal cover in Australia, Brazil and elsewhere. To improve these estimates, there is clearly a need for determining carbon and nutrient accumulation rates from additional coastal HTFs and assess their areal cover in Australia, Brazil and elsewhere. Furthermore, microphytobenthos also exist in the arid or semi-arid areas near saltmarshes, as well as in the lower intertidal flats inundated daily, which are often areas greater than vegetated areas and may contribute to blue carbon burial.

#### 4.2 Organic matter source



270 Microphytobenthos associated with coastal HTF ecosystems were an important source of OM accumulation in each of the sediment profiles. Microscopic examinations in previous studies have identified the cyanobacteria *Oscillatoria* spp., *Lyngbya* spp., *Microcoleus* spp., and *Phormidium* spp. as the dominant microphytobenthos in HTF ecosystems (Adame et al., 2012; Burford et al., 2016; Masuda and Enrich-Prast, 2016; Bento et al., 2017). These microphytobenthos are likely to be the important species contributing to the accumulation of OM, particularly in Guaratiba and Karumba where the  $\delta^{13}\text{C}$  and C:N ratio values were consistently similar to that of the HTF microphytobenthos end member values (Fig. 4). Therefore, we suggest that microphytobenthos are the dominate source of OM accumulating in the sedimentary substrates during the past century.

In contrast to the Guaratiba and Karumba profiles, the considerable spread in  $\delta^{13}\text{C}$  and molar C:N ratio values along the Gladstone sedimentary profiles suggests the OM accumulation inputs are from a combination of microphytobenthos and mangrove material (Fig. 4). These results are not surprising given the vast areal extent of mangrove systems in the Gladstone Harbour and their close proximity to the HTFs. Effective N consumption in coastal wetland sediments (Wadnerkar et al., 2019) may increase overall sedimentary C:N ratios. Sedimentary N and the relatively higher  $\delta^{15}\text{N}$  values observed in the Guaratiba HTF sediments (Fig. 3) may be indicative of eutrophication (Sanders et al., 2014). Indeed, wastewater inputs typically have elevated  $\delta^{15}\text{N}$  values due to elevated nitrogen cycling including denitrification (Costanzo et al., 2005). Anthropogenic wastewater inputs high in N and P loads are also of growing concern across the globe, particularly in HTF areas near shrimp farming (Ashton, 2008; Marchand et al., 2011). While there are no shrimp farms near our study sites, the release of high N and P loads may drive eutrophication of adjacent coastal areas (Ashton, 2008; Marchand et al., 2011) and modify carbon burial rates (Sanders et al., 2014). In addition to the increase of the N and P release, shrimp farms would drive a reduction of the HTFs area that may remove N and P.

### 4.3 CO<sub>2</sub> fluxes at the air-sediment interface

290 The great variability of air-saltpan CO<sub>2</sub> fluxes here suggests a highly dynamic and productive metabolism along the HTFs. The oligotrophic Gladstone's sites were net sources of CO<sub>2</sub> to the atmosphere in the dry season ( $0.72 \pm 0.01 \text{ g C m}^{-2} \text{ d}^{-1}$ ), while the eutrophic Guaratiba's HTF experienced net CO<sub>2</sub> uptake and source in the dry and rainy season ( $-0.03 \pm 0.01$  and  $0.71 \pm 0.22 \text{ g C m}^{-2} \text{ d}^{-1}$ , respectively). These estimates of net seasonal fluxes of CO<sub>2</sub> contribute to reduce the scarcity of studies quantifying this gas exchange at the air-sediment interface in HTFs (Table 3). The net CO<sub>2</sub> source observed during rainy compared to the net influx during dry seasons in Brazil (Mann-Whitney,  $p < 0.05$ ) may be attributed to higher temperature and cloud cover over sampling days in the rainy summer than the dry winter. Previous evidence indicates that the light attenuation by clouds may reduce microphytobenthos photosynthetic activity (Blackford, 2002; Barnett et al., 2020), while warmer sampling conditions on average  $\pm$  SE,  $26.7 \pm 0.02$  and  $21.5 \pm 0.02^\circ\text{C}$  during the wet and dry season, respectively, may stimulate heterotrophy in tidal flat systems (Laviale et al., 2015; Lin et al., 2020). The CO<sub>2</sub> source to the atmosphere found during the rainy summer still contrasted with previous evidence in the same Brazilian HTF on enhanced CO<sub>2</sub> sink after rain events in winter (Bento et al., 2017), suggesting that factors other than the occurrence of precipitation (*e.g.*, rainfall duration and intensity) may cause the dynamic short-term changes on microphytobenthic production. In addition, higher values of air-saltpan CO<sub>2</sub> influx during similar sunnier periods in the Brazilian than Australian HTFs may be attributed to more eutrophic

conditions, which could stimulate microphytobenthic production in saltpan sediments (Xie et al., 2019). These findings highlight the high temporal variability and the need for future seasonal sampling due to the short-term shifts in air-saltpan CO<sub>2</sub> exchange, specifically considering the potential net atmospheric CO<sub>2</sub> sink in HTFs as indicated by the autochthonous OM ~~is~~ found in the sedimentary profiles. As such, gaining a clearer understanding of the drivers of net primary production in HTFs during changing climatic and anthropogenic conditions is critical to determine their global relevance as atmospheric carbon sinks.

#### 4.4 Can HTFs be considered “Blue Carbon” systems?

While much of the research on blue carbon systems continues to focus on mangrove forests, tidal marshes, and seagrass meadows, there are suggestions of considering other ecosystems in the blue carbon framework (Raven, 2018; Trevathan-Tackett et al., 2015; Lovelock and Duarte, 2019). Tidally influenced freshwater forests, marine macroalgae and kelp beds, and HTFs for instance, are all ecosystems where blue carbon stocks and sequestration rates may be conceptually equivalent to conventional blue carbon systems (Raven, 2018; Krause-Jensen et al., 2018; Krauss et al., 2018; Lovelock and Duarte, 2019).

Lovelock and Duarte (2019) discuss several key assessment criteria for the inclusion of an ecosystem in the blue carbon framework. First, an ecosystem needs to be capable of long term-storage of CO<sub>2</sub> resulting in significant greenhouse gas (GHG) removal from the atmosphere. The results from this study indicate that HTF ecosystems are capable of long-term-storage of fixed CO<sub>2</sub> at rates averaging  $21 \pm 6 \text{ g C m}^{-2} \text{ y}^{-1}$ . Given that HTFs are extensively distributed in coastal areas showing freshwater deficit such as in northern Australia and Brazil, the scale of CO<sub>2</sub> removal can be significant and comparable to traditional blue carbon systems in some key arid regions. While this study demonstrates carbon burial in three HTF systems, accurate estimates on the magnitude of this carbon sink on national or global scales will require further studies and improved areal estimates.

The second consideration for inclusion into the blue carbon framework is that management of an ecosystem is possible. Management should maintain or enhance carbon and nitrogen stocks and thereby reduce GHG emissions (Lovelock and Duarte, 2019). Over the past few decades, HTFs have experienced large scale destruction and degradation on a global scale as a result of anthropogenic pressures such as urban and agricultural/aquaculture development (Ashton, 2008; Halpern et al., 2008; Martinez-Porchas and Martinez-Cordova, 2012) which may ultimately lead to large scale release of CO<sub>2</sub> to the atmosphere. Local, national and/or international management actions, therefore, have the potential to reduce and possibly revert these losses and destruction, thereby maintaining or even enhancing C sequestration similar to adjacent mangroves and saltmarshes. These management practices include regulating urban development or the construction of shrimp farming to prevent HTF ecosystem decline (Halpern et al., 2008; Martinez-Porchas and Martinez-Cordova, 2012). Moreover, current frameworks and management strategies in place for coastal vegetated ecosystems have the potential to incorporate HTFs given their close association. Therefore, we suggest that HTF ecosystems can be classified as blue carbon systems and should be included in global management and mitigation policies and are likely to be important contributors on regional scales.

## 5 Conclusions

The investigated HTF ecosystems have accumulated significant amounts of OC, TN and TP during the previous century. Although these accumulation rates are lower than other vegetated blue carbon systems per unit area, a substantial amount of carbon and nutrients are sequestered in HTFs considering their extensive global areal extent and should not be overlooked.

340 Stable isotope analysis along with the molar C:N ratios indicate that the microphytobenthos associated with these HTFs are an important source of the organic material accumulated along the sediment columns of these systems. To improve the robustness of our observations, there is a need for determining carbon and nutrient accumulation rates and CO<sub>2</sub> fluxes from additional coastal HTFs and to determine a more precise areal estimate of HTFs in Australia, Brazil and other parts of the world. However, our initial data implies that these coastal HTF ecosystems fit the definition of blue carbon systems and could be included in

345 global and regional management and mitigation policies.

## Acknowledgments

Field and laboratory investigations were funded by the Australian Research Council (DE160100443, DP180101285 & LE140100083). The data used in this research are available in the Tables. HM and RBP were funded by Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES - Código 001). HM was awarded by CNPq Research Productivity and

350 FAPERJ Young Scientist of Rio de Janeiro State fellowships. AP is supported by the “Fondo Nacional de Desarrollo Científico Tecnológico y de Innovación Tecnológica” (FONDECYT - PERU) through the Magnet program (Grant n° 007-2017-FONDECYT) and the “Incorporación de Investigadores” program (Grant n° E038-2019-02-FONDECYT-BM).

## Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be

355 construed as a potential conflict of interest.

## Author Contributions

DRB, HRM, and CJS designed and obtained funding for this work. DRB, CJS, HRM, RBP, DTM and LSM contributed to acquisition of data, and contributed to the analysis and interpretation of data. All of the authors made contributions to the drafting of the article and revisions critically for important intellectual content. All authors gave the final approval of the

360 version to be published.

## References

- Adame, M. F., Reef, R., Grinham, A., Holmes, G., and Lovelock, C. E.: Nutrient exchange of extensive cyanobacterial mats in an arid subtropical wetland, *Mar. Freshw. Res.*, 63, 457-467, 2012.
- Albuquerque, A., Ferreira, T., Nóbrega, G., Romero, R., Júnior, V. S., Meireles, A., and Otero, X.: Soil genesis on hypersaline tidal flats (apicum ecosystem) in a tropical semi-arid estuary (Ceará, Brazil), *Soil Res.*, 52, 140-154, 2014.
- 365

- Albuquerque, A. G. B. M., Ferreira, T. O., Cabral, R. L., Nóbrega, G. N., Romero, R. E., Meireles, A. J. d. A., and Otero, X. L.: Hypersaline tidal flats (apicum ecosystems): the weak link in the tropical wetlands chain, *Environ. Rev.*, 22, 99-109, 2013.
- 370 Alongi, D., Pfitzner, J., Trott, L., Tirendi, F., Dixon, P., and Klumpp, D.: Rapid sediment accumulation and microbial mineralization in forests of the mangrove *Kandelia candel* in the Jiulongjiang Estuary, China, *Estuar. Coast. Shelf Sci.*, 63, 605-618, 2005.
- Alongi, D. M.: Mangrove forests: resilience, protection from tsunamis, and responses to global climate change, *Estuar. Coast. Shelf Sci.*, 76, 1-13, 2008.
- Alongi, D. M.: Dissolved iron supply limits early growth of estuarine mangroves, *Ecology*, 91, 3229-3241, 2010.
- 375 Appleby, P. G., and Oldfield, F.: Application of lead-210 to sedimentation studies, in: Uranium Series Disequilibrium: Application to Earth, Marine and Environmental Science, edited by: Ivanovich, M., and Harmon, S., Oxford Science Publications, 731-783, 1992.
- Ashton, E. C.: The impact of shrimp farming on mangrove ecosystems, *CAB Rev: Persp. Agr. Vet. Sci. Nut. Nat. Res.*, 3, 2008.
- 380 Barnett, A., Méléder, V., Dupuy, C., and Lavaud, J.: The vertical migratory rhythm of intertidal microphytobenthos in sediment depends on the light photoperiod, intensity, and spectrum: Evidence for a positive effect of blue wavelengths, *Front. Mar. Sci.*, 7, 2020.
- Bento, L., Masuda, L. S. M., Peixoto, R. B., and Enrich-Prast, A.: Regulation in the metabolism and community structure of a tropical salt flat after rainfall, *J. Coast. Res.*, 33, 304-308, 2017.
- 385 Blackford, J. C.: The influence of microbenthos on the Northern Aridic ecosystem: A modelling study, *Estuar. Coast. Shelf Sci.*, 55, 2002.
- Breithaupt, J. L., Smoak, J. M., Smith, T. J., and Sanders, C. J.: Temporal variability of carbon and nutrient burial, sediment accretion, and mass accumulation over the past century in a carbonate platform mangrove forest of the Florida Everglades, *J. Geophys. Res. Biogeosci.*, 119, 2014.
- 390 Breithaupt, J. L., Smoak, J. M., Smith, T. J., Sanders, C. J., and Hoare, A.: Organic carbon burial rates in mangrove sediments: Strengthening the global budget, *Global Biogeochem. Cy.*, 26, 2012.
- Bureau of Meteorology: Climate statistics for Australian locations, available at [http://www.bom.gov.au/climate/averages/tables/cw\\_029028.shtml](http://www.bom.gov.au/climate/averages/tables/cw_029028.shtml), last accesses: 12 December 2019.
- Burford, M., Valdez, D., Curwen, G., Faggotter, S., Ward, D., and Brien, K. O.: Inundation of saline supratidal mudflats provides an important source of carbon and nutrients in an aquatic system, *Mar. Ecol. Prog. Ser.*, 545, 21-33, 2016.
- 395 Chairi, R., Derenne, S., Abdeljaoued, S., and Largeau, C.: Sediment cores representative of contrasting environments in salt flats of the Moknine continental sabkha (Eastern Tunisia): sedimentology, bulk features of organic matter, alkane sources and alteration, *Org. Geochem.*, 41, 637-652, 2010.

- 400 Conesa, H., María-Cervantes, A., Álvarez-Rogel, J., and González-Alcaraz, M.: Influence of soil properties on trace element availability and plant accumulation in a Mediterranean salt marsh polluted by mining wastes: implications for phytomanagement, *Sci. Total Environ.*, 409, 4470-4479, 2011.
- Conrad, S. R., Santos, I. R., White, S., and Sanders, C. J.: Nutrient and trace metal fluxes into estuarine sediments linked to historical and expanding agricultural activity (Hearnes Lake, Australia), *Estuar. Coasts*, 42, 944-957, 2019.
- 405 Costanzo, S. D., Udy, J., Longstaff, B., and Jones, A.: Using nitrogen stable isotope ratios ( $\delta^{15}\text{N}$ ) of macroalgae to determine the effectiveness of sewage upgrades: changes in the extent of sewage plumes over four years in Moreton Bay, Australia, *Mar. Pollut. Bull.*, 51, 212-217, 2005.
- Estevam, R. M. E.: Os manguezais de Guaratiba Frente às mudanças climáticas globais: análise da influência da variabilidade climática sobre a dinâmica de comunidades pioneiras, Universidade do Estado do Rio de Janeiro, Programa de Pós-Graduação em Meio Ambiente, 99 pp., 2019.
- 410 Estrada, G. C. D., Soares, M. L. G., de Oliveira Chaves, F., and Cavalcanti, V. F.: Analysis of the structural variability of mangrove forests through the physiographic types approach, *Aquat. Bot.*, 111, 135-143, 2013.
- Fourqurean, J. W., Duarte, C. M., Kennedy, H., Marbà, N., Holmer, M., Mateo, M. A., Apostolaki, E. T., Kendrick, G. A., Krause-Jensen, D., and McGlathery, K. J.: Seagrass ecosystems as a globally significant carbon stock, *Nat. Geosci.*, 5, 505-509, 2012.
- 415 Halpern, B. S., Walbridge, S., Selkoe, K. A., Kappel, C. V., Micheli, F., D'agrosa, C., Bruno, J. F., Casey, K. S., Ebert, C., and Fox, H. E.: A global map of human impact on marine ecosystems, *Sci.*, 319, 948-952, 2008.
- Kelleway, J. J., Cavanaugh, K., Rogers, K., Feller, I. C., Ens, E., Doughty, C., and Saintilan, N.: Review of the ecosystem service implications of mangrove encroachment into salt marshes, *Global Change Biol.*, 23, 3967-3983, 2017.
- Krause-Jensen, D., Lavery, P., Serrano, O., Marbà, N., Masque, P., and Duarte, C. M.: Sequestration of macroalgal carbon: 420 the elephant in the Blue Carbon room, *Biol. Lett.*, 14, 2018.
- Krauss, K. W., Noe, G. B., Duberstein, J. A., Conner, W. H., Stagg, C. L., Cormier, N., Jones, M. C., Bernhardt, C. E., Graeme Lockaby, B., and From, A. S.: The role of the upper tidal estuary in wetland blue carbon storage and flux, *Global Biogeochem. Cy.*, 32, 817-839, 2018.
- Laviale, M., Barnett, A., Ezequiel, J., Lepetit, B., Frankenbach, S., Méléder, V., Serôdio, J., and Lavaud, J.: Response of 425 intertidal benthic microalgal biofilms to a coupled light-temperature stress: evidence for latitudinal adaptation along the Atlantic coast of Southern Europe, *Environ. Microbiol.*, 17, 3662-3677, 2015.
- Leopold, A., Marchand, C., Deborde, J., and Allenbach, M.: Temporal variability of  $\text{CO}_2$  fluxes at the sediment-air interface in mangroves (New Caledonia), *Sci. Total Environ.*, 502, 617-626, 2015.
- Leopold, A., Marchand, C., Deborde, J., Chaduteau, C., and Allenbach, M.: Influence of mangrove zonation on  $\text{CO}_2$  fluxes at 430 the sediment-air interface (New Caledonia), *Geoderma*, 202-203, 62-70, 2013.
- Lin, W. J., Wu, J., and Lin, H. J.: Contribution of unvegetated tidal flats to coastal carbon flux, *Global Change Biol.*, 26, 3443-3454, 2020.

- Lovelock, C. E.: Soil respiration and belowground carbon allocation in mangrove forests, *Ecosystems*, 11, 342-354, 2008.
- Lovelock, C. E., and Duarte, C. M.: Dimensions of Blue Carbon and emerging perspectives, *Biol. Lett.*, 15, 2019.
- 435 Lovelock, C. E., Grinham, A., Adame, M. F., and Penrose, H. M.: Elemental composition and productivity of cyanobacterial mats in an arid zone estuary in north Western Australia, *Wetl. Ecol. Manag.*, 18, 37-47, 2010.
- Macreadie, P. I., Anton, A., Raven, J. A., Beaumont, N., Connolly, R. M., Friess, D. A., Kelleway, J. J., Kennedy, H., Kuwae, T., and Lavery, P. S.: The future of Blue Carbon science, *Nat. Commun.*, 10, 1-13, 2019.
- MapBiomass: Land cover data, available at <http://mapbiomas.org/map#coverage>, last access: 12 August 2019.
- 440 Marchand, C., Lallier-Vergès, E., and Allenbach, M.: Redox conditions and heavy metals distribution in mangrove forests receiving effluents from shrimp farms (Teremba Bay, New Caledonia), *J. Soils Sediments.*, 11, 529-541, 2011.
- Martinez-Porchas, M., and Martinez-Cordova, L. R.: World aquaculture: Environmental impacts and troubleshooting alternatives, *Sci. World J.*, 9, 2012.
- Masuda, L., and Enrich-Prast, A.: Benthic microalgae community response to flooding in a tropical salt flat, *Brazilian J. Biol.*,
- 445 76, 577-582, 2016.
- McLeod, E., Chmura, G. L., Bouillon, S., Salm, R., Björk, M., Duarte, C. M., Lovelock, C. E., Schlesinger, W. H., and Silliman, B. R.: A blueprint for blue carbon: Toward an improved understanding of the role of vegetated coastal habitats in sequestering CO<sub>2</sub>, *Front. Ecol. Environ.*, 9, 552-560, 2011.
- Ouyang, X., and Lee, S. Y.: Updated estimates of carbon accumulation rates in coastal marsh sediments, *Biogeosciences*, 11,
- 450 5057-5071, 2014.
- Paerl, H. W., Pinckney, J. L., and Steppe, T. F.: Cyanobacterial–bacterial mat consortia: Examining the functional unit of microbial survival and growth in extreme environments, *Environ. Microbiol.*, 2, 11-26, 2000.
- R Development Core Team: R: A language and environment for statistical computing, R Foundation for Statistical Computing, Vienna, Austria, ISBN 3-900051-07-0, URL <http://www.R-project.org/>, 2011.
- 455 Radabaugh, K. R., Moyer, R. P., Chappel, A. R., Powell, C. E., Bociu, I., Clark, B. C., and Smoak, J. M.: Coastal blue carbon assessment of mangroves, salt marshes, and salt barrens in Tampa Bay, Florida, USA, *Estuar. Coasts.*, 41, 1496-1510, 2018.
- Raven, J.: Blue carbon: past, present and future, with emphasis on macroalgae, *Biol. Lett.*, 14, 2018.
- Ravichandran, M., Baskaran, M., Santschi, P. H., and Bianchi, T. S.: Geochronology of sediments in the Sabine-Neches
- 460 estuary, Texas, USA, *Chem. Geol.*, 125, 291-306, 1995.
- Reef, R., Feller, I. C., and Lovelock, C. E.: Nutrition of mangroves, *Tree physiol.*, 30, 1148-1160, 2010.
- Rezende, C., Pfeiffer, W., Martinelli, L., Tsamakis, E., Hedges, J., and Keil, R.: Lignin phenols used to infer organic matter sources to Sepetiba Bay–RJ, Brasil, *Estuar. Coast. Shelf Sci.*, 87, 479-486, 2010.
- Ridd, P., and Stieglitz, T.: Dry season salinity changes in arid estuaries fringed by mangroves and saltflats, *Estuar. Coast. Shelf*
- 465 *Sci.*, 54, 1039-1049, 2002.

- Rosentreter, J., Maher, D. T., Ho, D., Call, M., Barr, J., and Eyre, B. E.: Spatial and temporal variability of CO<sub>2</sub> and CH<sub>4</sub> gas transfer velocities and quantification of the CH<sub>4</sub> microbubble flux in mangrove dominated estuaries, *Limnol. Oceanogr.*, 62, 561-578, 2017.
- 470 Saintilan, N., Wilson, N. C., Rogers, K., Rajkaran, A., and Krauss, K. W.: Mangrove expansion and salt marsh decline at mangrove poleward limits, *Global Change Biol.*, 20, 147-157, 2014.
- Sanders, C. J., Maher, D. T., Tait, D. R., Williams, D., Holloway, C., Sippo, J. Z., and Santos, I. R.: Are global mangrove carbon stocks driven by rainfall? *J. Geophys. Res. Biogeosci.*, 121, 2600-2609, 2016a.
- Sanders, C. J., Santos, I. R., Maher, D. T., Breithaupt, J. L., Smoak, J. M., Ketterer, M. E., Call, M., Sanders, L., and Eyre, B. D. Examining <sup>239+240</sup>Pu, <sup>210</sup>Pb and historical events to determine carbon, nitrogen and phosphorus burial in mangrove  
475 sediments of Moreton Bay, Australia, *J. Environ. Radioact.*, 151, 623-629, 2016b.
- Sanders, C. J., Eyre, B. E., Santos, I. R., Machado, W., Luiz-Silva, W., Smoak, J. M., Breithaupt, J. L., Ketterer, M. E., Sanders, L., Marotta, H., and Silva-Filho, E.: Elevated rates of organic carbon, nitrogen, and phosphorus accumulation in a highly impacted mangrove wetland, *Geophys. Res. Lett.*, 41, 2475-2480, 2014.
- Schile, L. M., Kauffman, J. B., Crooks, S., Fourqurean, J. W., Glavan, J., and Megonigal, J. P.: Limits on carbon sequestration  
480 in arid blue carbon ecosystems, *Ecol. Appl.*, 27, 859-874, 2017.
- Serrano, O., Lovelock, C. E., Atwood, T. B., et al.: Australian vegetated coastal ecosystems as global hotspots for climate change mitigation. *Nat. Commun.*, 10, 1-10, 2019.
- Shen, C., Zhang, C., Xin, P., Kong, J., and Li, L.: Salt dynamics in coastal marshes: Formation of hypersaline zones, *Water Resour. Res.*, 54, 3259-3276, 2018
- 485 Soares, M. L. G., Chaves, F. D. O., Estrada, G. C. D., and Fernandez, V.: Mangrove forests associated with salt flats: A case study from southeast Brazil, *Braz. J. Oceanogr.*, 65, 102-115, 2017.
- Spalding M., Kainuma, M., and Collins, L. *World Atlas of Mangroves (version 2.0)*. A collaborative project of ITTO, ISME, FAO, UNEP-WCMC, UNESCO-MAB, UNU-INWEH and TNC, London (UK): Earthscan, London, 319 pp. URL: <http://www.routledge.com/books/details/9781844076574>, 2010.
- 490 Trevathan-Tackett, S. M., Kelleway, J., Macreadie, P. I., Beardall, J., Ralph, P., and Bellgrove, A.: Comparison of marine macrophytes for their contributions to blue carbon sequestration, *Ecology*, 96, 3043-3057, 2015.
- Wadnerkar, P. D., Santos, I. R., Looman, I., Sanders, C. J., White, S., Tucker, J. P., and Holloway, C. Significant nitrate attenuation in a mangrove-fringed estuary during a flood-chase experiment, *Environ. Pollut.*, 253, 1000-1008, 2019.
- 495 Xie, Y., Wang, L., Liu, X., Li, X., Wang, Y., and Huang, B.: Contrasting responses of intertidal microphytobenthos and phytoplankton biomass and taxonomic composition to the nutrient loads in the Jiulong River Estuary, *Phycological Res.*, 67, 152-163, 2019.

**Table 1. Characterization of study sites. Rainfall (mm) and Temperature (°C) data are based on annual averages derived from monthly measurements (n = number of years data) and HTF: Mangrove Area indicates the ratio of hypersaline tidal flat area to mangrove area.**

	Hypersaline Tidal Flat Study Sites			
	Guaratiba	Karumba	Gladstone Site 1 High Tidal Area	Gladstone Site 2 Low Tidal Area
Location	23°00'29"S, 43°36'31"W	17°25'12"S, 140°51'36"E	23°45'41"S, 151°16'34"E	23°45'18"S, 151°16'49"E
Rainfall (mm)	1058 (n=32)	883 (n=23)	846 (n=25)	846 (n=25)
Temperature (°C)	21 – 27 (n=32)	27 – 33 (n=19)	23 – 31 (n=26)	23 – 31 (n=26)
Tidal Range (m)	0.1 – 2.0	0.1 – 4.5	0.1 – 4.7	0.1 – 4.7
HTF Area (Km <sup>2</sup> )	7.4	16.9	2.84	0.95
HTF: Mangrove Area	1:3	10:1	1:1	1:1

**Table 2. Mean (± standard error) ~50 year organic carbon (OC), total nitrogen (TN) and total phosphorus (TP) burial rates in the four hypersaline tidal flat sediment cores. Means are based on one core per site**

Study site	OC (g m <sup>-2</sup> y <sup>-1</sup> )	TN (g m <sup>-2</sup> y <sup>-1</sup> )	TP (g m <sup>-2</sup> y <sup>-1</sup> )
Guaratiba	17.8 ± 0.8	2.3 ± 0.2	2.0 ± 0.1
Karumba	31.7 ± 4.3	2.8 ± 0.3	1.3 ± 0.1
Gladstone site 1	11.3 ± 2.1	0.8 ± 0.1	1.4 ± 0.0
Gladstone site 2	25.2 ± 2.9	1.2 ± 0.1	1.4 ± 0.3



510 **Table 3. Mean sediment organic carbon content (%) and CO<sub>2</sub> fluxes (mg C m<sup>-2</sup> h<sup>-1</sup>) at the air-sediment interface in hypersaline tidal flat sediments reported in the literature. Values are means ± standard error unless otherwise stated.**

Location	Air-sediment CO <sub>2</sub> fluxes (light) (mg C m <sup>-2</sup> h <sup>-1</sup> )	Air-sediment CO <sub>2</sub> fluxes (dark) (mg C m <sup>-2</sup> h <sup>-1</sup> )	Sediment organic carbon content (%)	Reference
New Caledonia	20.4 ± 3.2	20.0 ± 3.3	1.6 ± 0.2 <sup>d</sup>	Leopold et al. (2013)
Guaratiba, Brazil				
Dry season	-6.7 ± 1.3	2.3 ± 0.3	0.6 ± 0.0	This study
Wet season	44.2 ± 7.2	20.9 ± 3.7	0.6 ± 0.0	This study
Guaratiba, Brazil	-5.3 ± 3.2 <sup>b</sup>	4.8 ± 3.6 <sup>b</sup>	NA	Bento et al. (2017)
Gladstone, Australia	24.3 ± 2.0	36.5 ± 2.7	0.5 ± 0.1	This study
Karumba, Australia	38.2 ± 0.0	44.8 ± 9.2	0.8 ± 0.1	This study
Karumba, Australia	14.5 ± 12.2 <sup>a,b</sup>	0.9 ± 0.5 <sup>a,b</sup>	NA	Burford et al. (2016)
Exmouth Gulf, Australia	64.8 ± 7.4 <sup>a</sup>	40.2 ± 3.7 <sup>a</sup>	0.7	Lovelock et al. (2010)
Arabian Gulf, United Arab Emirates	38.0 ± 15.1 <sup>b,c</sup>		1.6 ± 0.7 <sup>b</sup>	Schile et al. (2017)
Murcia, Spain	NA	NA	5.2 ± 0.5	Conesa et al. (2011)
Tunisia, Africa	NA	NA	0.6 ± 0.1	Chairi et al. (2010)
Ceará, Brazil	NA	NA	0.5 ± 0.3	Albuquerque et al. (2014)
Ceará, Brazil	NA	NA	0.7 ± 0.8	Albuquerque et al. (2013)
Bahia, Brazil	NA	NA	0.7 ± 0.1 <sup>c</sup>	Albuquerque et al. (2013)
Teremba Bay, New Caledonia	NA	NA	5.9 ± 1.2 <sup>b</sup>	Marchand et al. (2011)
Tempa Bay, Florida	NA	NA	0.7 ± 0.5	Radabaugh et al. (2018)

<sup>a</sup> Fluxes calculated from measurements of oxygen (O<sub>2</sub>) assuming a molar CO<sub>2</sub>:O<sub>2</sub> ratio of 1:1


<sup>b</sup> Values from figures were estimated using WebPlotDigitizer (<https://automeris.io/WebPlotDigitizer/>)

<sup>c</sup> Organic carbon value = Organic matter/1.724


515 <sup>d</sup> Organic carbon value = 95% of total carbon value

° Study did not clarify if CO<sub>2</sub> flux was measured in a light or dark chamber


520




525 **Figure 1.** Study sites (red arrows); Guaratiba, Brazil (left), Karumba, Australia (center), and Gladstone, Australia (right). Stars show the location of literature data summarized in Table 3. Purple stars are areas with only sediment carbon content data, green stars are areas with only gas flux data, and yellow stars are areas with both sediment carbon content and gas flux data from hypersaline tidal flat (HTF) studies (satellite images were taken from ©Google Earth).




530 **Figure 2.**  $^{226}\text{Ra}$  (x) and  $^{210}\text{Pb}$  (circles) depth profiles of the four hypersaline tidal flat sediment cores in this work. Error bars indicate counting uncertainties.



535 **Figure 3.** Vertical distribution of sand (>63  $\mu\text{m}$ ), organic carbon (OC), total nitrogen (TN) and total phosphorous (TP) contents (%) as well as  $\delta^{13}\text{C}$ ,  $\delta^{15}\text{N}$  and molar C:N ratios of the four hypersaline tidal flat sediment cores.



**Figure 4. Distribution of  $\delta^{13}\text{C}$  vs C:N molar ratio in the four hypersaline tidal flat sediment cores. Endmember values were taken from HTF surface microphytobenthos and nearby mangrove vegetation.**



540

Figure 5. Median air-sediment CO<sub>2</sub> fluxes (mg C m<sup>-2</sup> h<sup>-1</sup>) from hypersaline tidal flat sediments of Guaritiba, Brazil (Dry season represented yellow triangles, n = 44 to dark chambers and n = 53 to light chambers, and wet season represented by blue triangles, n = 6 to dark chambers and n = 41 to light chambers) and Australia (Gladstone site represented by light grey circles, n = 33 to dark chambers and n = 31 to light chambers, and Karumba site represented by dark grey circles, n = 13 to dark chambers and n = 1 to light chambers). Negative values represent net CO<sub>2</sub> influx to sediments while positive represents net CO<sub>2</sub> emission to the atmosphere. Average values represented by crosses and error bars denotes whiskers minimum to maximum. Different letters indicate significant differences (Mann Whitney; p<0.05).

545