

An improved process-oriented hydro-biogeochemical model for simulating dynamic fluxes of methane and nitrous oxide in alpine ecosystems with seasonally frozen soils

Wei Zhang¹, Zhisheng Yao¹, Siqi Li¹, Xunhua Zheng^{1, 2}, Han Zhang^{1, 3}, Lei Ma^{1, 4}, Kai Wang¹, Rui Wang¹, Chunyan Liu¹, Shenghui Han¹, Jia Deng⁵, Yong Li⁶

¹State Key Laboratory of Atmospheric Boundary Layer Physics and Atmospheric Chemistry, Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing 100029, P. R. China

²College of Earth and Planetary Science, University of Chinese Academy of Sciences, Beijing 100049, P. R. China

³School of Geographic and Environmental Sciences, Tianjin Normal University, Tianjin 300387, P. R. China

⁴Institute of Meteorology and Climate Research, Atmospheric Environmental Research (IMK-IFU), Karlsruhe Institute of Technology, Kreuzeckbahnstrasse 19, 82467 Garmisch-Partenkirchen, Germany

⁵Complex Systems Research Center, Institute for the Study of Earth, Oceans and Space, University of New Hampshire, 39 College Road, Durham, NH 03824, USA

⁶Key Laboratory of Agro-ecological Processes in Subtropical Region, Institute of Subtropical Agriculture, Chinese Academy of Sciences, Hunan 410125, P. R. China

Correspondence to: Xunhua Zheng (xunhua.zheng@post.iap.ac.cn)

Abstract. The hydro-biogeochemical model Catchment Nutrient Management Model - DeNitrification-DeComposition (CNMM-DNDC) was established to simultaneously quantify ecosystem productivity and losses of nitrogen and carbon at the site or catchment scale. As a process-oriented model, this model is expected to be universally applied to different climate zones, soils, land uses and field management practices. This study is one of many efforts to fulfil such an expectation, which was performed to improve the CNMM-DNDC by incorporating a physical-based soil thermal module to simulate the soil thermal regime in the presence of freeze-thaw cycles. The modified model was validated with simultaneous field observations in three typical alpine ecosystems (wetlands, meadows and forests) within a catchment located in the seasonally frozen region of the eastern Tibetan Plateau, including soil profile temperature, topsoil moisture and fluxes of methane (CH₄) and nitrous oxide (N₂O). The validation showed that the modified CNMM-DNDC was able to simulate the observed seasonal dynamics and magnitudes of above variables in the three typical alpine ecosystems, with index of agreement values of 0.91–1.00, 0.49–0.83, 0.57–0.88 and 0.26–0.47, respectively. Consistent with the emissions determined from the field observations, the simulated aggregate emissions of CH₄ and N₂O were highest for the wetland among three alpine ecosystems, which were dominated by the CH₄ emissions. This study indicates the possibility for utilizing the process-oriented model CNMM-DNDC to predict hydro-biogeochemical processes, as well as related gas emissions, in seasonally frozen regions. As the original CNMM-DNDC was previously validated in some unfrozen regions, the modified CNMM-DNDC could be potentially applied to estimate the emissions of CH₄ and N₂O from various ecosystems under different climate zones at the site or catchment scale.

35 1 Introduction

36 During the last 50 years, the extraordinary changes in the nitrogen and carbon cycles have occurred globally, which are
37 essential components of ecosystems (e.g., Galloway *et al.*, 2008; Canfield *et al.*, 2010). Climate changes due to warming and
38 human anthropogenic activities derived from food production have significantly altered the cycling of nitrogen and carbon
39 and led to increased reactive nitrogen availability and carbon losses, which result in a series of environmental problems at
40 the catchment, regional and even global scales (e.g., Galloway *et al.*, 2004; Galloway *et al.*, 2008; Ju *et al.*, 2009). Excessive
41 reactive nitrogen in soils can be lost in the forms of nitrogen gases, such as nitrous oxide (N_2O), nitric oxide (NO) and
42 ammonia (NH_3), and nitrogen pollution, such as nitrate (NO_3^-) and ammonium (NH_4^+), in water through leaching or surface
43 runoff (e.g., Seitzinger, 2008; Collins *et al.*, 2016). In the face of increased air temperatures and intensive land use changes,
44 especially in cold regions, the soil organic carbon stored since the Last Glacial Maximum has been lost to the atmosphere via
45 methane (CH_4) and carbon dioxide (CO_2) (e.g., Piao *et al.*, 2009; Fenner and Freeman, 2011; Schuur *et al.*, 2015). These
46 nitrogen and carbon losses contribute to potential global warming (CO_2 , CH_4 and N_2O), air pollution (NO and NH_3) and
47 surface/groundwater pollution (NO_3^- and NH_4^+). Therefore, sustainable ecosystems urgently need to be established that not
48 only focus on net primary productivity but also are friendly to the environment with the minimal hazards, including
49 greenhouse gases, air pollution and water pollutants (e.g., Cui *et al.*, 2018; Zhang *et al.*, 2019).

50 The cycling of nitrogen and carbon is closely related to soil water processes (e.g., Breuer *et al.*, 2010; Vereecken *et al.*,
51 2016; Zhang *et al.*, 2018b). Thus, interactions among soil waters and the cycling of nitrogen and carbon govern biological
52 productivity and environmental outcomes (e.g., Zhu *et al.*, 2018). The interactions consist of the redox potential for different
53 transformation processes influenced by the spatiotemporal variation in soil water content and the lateral transport of water
54 and dissolved nitrogen or carbon controlled by surface and subsurface flow (e.g., McClain *et al.*, 2003; Castellano *et al.*,
55 2013; Bechmann, 2014). For example, the variation in soil water content can create hot spots or moments of nitrogen and
56 carbon losses by influencing plant nitrogen uptake, redox potential, and the transport of dissolved nitrogen and carbon (e.g.,
57 Zhu *et al.*, 2012; Keiluweit *et al.*, 2017). Therefore, a complete understanding of biogeochemical processes will inevitably
58 involve interactions among soil water and the cycling of nitrogen and carbon (e.g., Breuer *et al.*, 2010; Vereecken *et al.*,
59 2016; Zhu *et al.*, 2018).

60 Biogeochemical models, such as DNDC, LandscapeDNDC, WNMM, MOMOS, CENTURY and DayCent, are
61 effective tools for simulating the cycling of nitrogen and carbon and quantifying the effects of climate change and
62 anthropogenic activities on ecosystems (e.g., Foereid *et al.*, 2007; Haas *et al.*, 2012; Li, 2007; Li *et al.*, 2007; Pansu *et al.*,
63 2010; Cheng *et al.*, 2014; Pansu *et al.*, 2014). In recent years, some new conceptual approaches are applied in the
64 biogeochemical models, such as centering on the functional role of the soil microbial biomass (Pansu *et al.*, 2010; Pansu *et al.*,
65 2014) and detailing the lateral transport of water and nutrients (Haas *et al.*, 2012; Zhang *et al.*, 2018b). Generally,
66 comprehensive hydrological processes, especially for the lateral transport of water and nutrients, are simplified or ignored in
67 most models due to specific questions that must be addressed (e.g., Li, 2007; Li *et al.*, 2007; Chen *et al.*, 2008; Deng *et al.*,

2014). For the land surface or hydrological models at large scales, they are designed with explicit mechanisms of hydrology and generally focus on vertical and lateral nutrient transport, such as nitrate loads into rivers (e.g., Liu *et al.*, 2019). However, the simulations of nitrogen and carbon processes are usually based on empirical functions even without predicting gas loss. Due to the various purposes of different models, coupling soil hydrological models with biogeochemical models can be an effective strategy for integrating soil water and cycling of nitrogen and carbon to improve model performance. Thus, the coupled model with improved performance can be applied to simultaneously predict productivity and potential negative environmental effects (e.g., Chen *et al.*, 2008; Zhu *et al.*, 2018).

In recent years, efforts have been implemented to couple models, such as SWAT-N, LandscapeDNDC-CMF, APSIM, SWAT-DayCent, and CNMM-DNDC (e.g., Pohlert *et al.*, 2007; Haas *et al.*, 2012; Holzworth *et al.*, 2014; Wu *et al.*, 2016; Zhang *et al.*, 2016; Zhang *et al.*, 2018b). The models derived from SWAT were all based on semi-distributed hydrological models using hydrologic response units and did not perform better in estimating non-point source pollution (e.g., Pohlert *et al.*, 2007; Bosch *et al.*, 2011 ;Wu *et al.*, 2016). A coupler was used to couple two models for LandscapeDNDC-CMF, which realized the simulation of horizontal movement of water and nutrients (e.g., Haas *et al.*, 2012; Klatt *et al.*, 2017; Schroeck *et al.*, 2019). Compared with other models, the Catchment Nutrient Management Model - DeNitrification-DeComposition (CNMM-DNDC), which was established by incorporating the core biogeochemical processes of DNDC into the hydrological framework of the CNMM, was validated at a catchment with complex landscapes in the subtropical region and showed good performance for simultaneously simulating various variables, including ecosystem productivity, hydrological nitrogen losses and nitrate discharge in streams, and emissions of gaseous carbon and nitrogenous gases (Zhang *et al.*, 2018b). Therefore, the CNMM-DNDC has the capacity to simulate the various variables closely related to both productivity and environmental hazards.

However, as a process-oriented hydro-biogeochemical model designed to be applicable to different climate zones, soils, land uses and field management practices, CNMM-DNDC testing is still lacking due to limited observations for model validation. In this study, the model was applied to a catchment in a seasonally frozen region located on the eastern Tibetan Plateau with the land use types of alpine wetlands, meadows and forests to test its ability to simulate hydro-biogeochemical processes. However, scientific descriptions of soil thermal dynamics due to freeze-thaw cycles are still lacking for the CNMM-DNDC. This gap may hinder model application in seasonally frozen regions, which account for 56% of the exposed land surface of the Northern Hemisphere (Jiang *et al.*, 2020). In addition, the soil freeze-thaw cycles that occur in these mid-high latitude regions exert important influences on soil thermal dynamics, as well as on related hydrological processes, thus increasing the availability of substrates and stimulating the processes of CH₄ and N₂O production and emissions in soils (e.g., Song *et al.*, 2019). Therefore, we hypothesize that adding the missing scientific processes of soil thermal dynamics into the internal model program codes can improve the performance of the CNMM-DNDC in simulating the soil thermal dynamics, hydrological processes and CH₄ and N₂O fluxes in seasonally frozen regions. Filling this gap is especially necessary to broaden model applicability.

101 To test the above hypothesis, the catchment simulation in the Rierlangshan was conducted using a unique experimental
102 dataset, which was obtained by Zhang *et al.* (2018a, 2019) and Yao *et al.* (2019) for the catchment that involved three typical
103 alpine ecosystems, wetlands, meadows and forests, on the eastern Tibetan Plateau. The aims of this study were to (i) attempt
104 to address the gap in the CNMM-DNDC by improving the scientific processes of soil thermal dynamics for seasonally
105 frozen regions and (ii) compare the performances of the original and modified models in simulating the soil profile
106 temperature, topsoil moisture and CH₄ and N₂O fluxes in three typical alpine ecosystems in the Rierlangshan catchment with
107 field observations. Therefore, the validated model with modifications provides a mechanism for not only interpreting
108 observations but also predicting the CH₄ and N₂O fluxes in alpine ecosystems.

109 2 Materials and methods

110 2.1 Model description

111 2.1.1 Overview of the CNMM-DNDC model

112 The CNMM-DNDC is a process-oriented model developed for simulating hydro-biogeochemical interactions at the
113 catchment or site scale, and this model is designed following the basic theories of physics, chemistry, and biogeochemistry
114 and has the capacity to simulate the complex transport and transformation of water, nitrogen and carbon in terrestrial
115 ecosystems under both aerobic and anaerobic conditions. The model can be applied to simultaneously quantify ecosystem
116 productivity, net emissions of nitrogen and carbon gases and hydrological nitrogen losses through soil leaching and
117 discharge in streams from an entire catchment or individual landscape unit (Zhang *et al.*, 2018b). The model was established
118 to address the bottleneck issue associated with most biogeochemical models, i.e., the inability to simulate the lateral flows of
119 water and nutrients, by incorporating the core biogeochemical processes of DNDC (including the processes of
120 decomposition, nitrification, denitrification and fermentation) into the hydrological framework of the CNMM, which is fully
121 distributed. For the new generation of biogeochemical models, the microbial ecology was integrated into the biogeochemical
122 models, which represents direct microbial control over decomposition, such as MOMOS (Pansu *et al.*, 2010; Treseder *et al.*,
123 2011; Todd-Brown *et al.*, 2012; Pansu *et al.*, 2014). The biogeochemical processes simulated by the DNDC were generally
124 based on first-order kinetics for decomposition and Michaelis-Menten kinetics of two substrates for nitrification and
125 denitrification, which only the parameterized growth and death of nitrifiers and denitrifiers were considered (Li, 2000).
126 However, due to the global application and validation of DNDC (e.g., Chen *et al.*, 2008; Giltrap *et al.*, 2010; Cui *et al.*, 2014,
127 Zhang *et al.*, 2015), the biogeochemical processes of DNDC were selected in the CNMM-DNDC despite some deficiencies
128 in simulating microbial biomass.

129 The simulated soil depth (including bedrock) is user-defined. The temporal and spatial resolutions are also user-
130 defined according to the driving data of climate (generally in 3 hours) and digital elevation model (DEM). The soil moisture
131 was calculated based on the mass balance of precipitation, irrigation, evapotranspiration, vertical flow, lateral flow and water

132 from a rising water table. The total water that can be infiltrated during each time step was determined by a defined maximum
133 infiltration rate. Darcy's law was applied for predicting the vertical water flow in the soil profile. A cell-by-cell approach
134 using a kinematic approximation was applied to route the saturated overland and subsurface flow based on DEM. The stream
135 flow was estimated using a cascade of linear channel reservoirs (Wigmosta *et al.*, 1994). For plant growth, gross primary
136 production was simulated using Farquhar *et al.* (1980) for C_3 and Collatz *et al.* (1992) for C_4 , with annual primary
137 productivity calculated as the residue of gross primary production and autotrophic respiration. The processes related to the
138 production of N_2O include nitrification and denitrification, which occur simultaneously at aerobic and anaerobic microsites,
139 respectively. The concept of an "anaerobic balloon" was adopted to determine the microsites and allocate substrates for
140 nitrification and denitrification. The sizes of the aerobic (nitrification) and anaerobic (denitrification) microsites were
141 determined by the soil redox potential (Eh) using the Nernst equation (Li, 2007). The "hole in the pipe" concept was applied
142 to calculate N_2O production during nitrification, which is influenced by the soil moisture, temperature and pH (Li, 2016).
143 The production of N_2O during denitrification was predicted with Michaelis-Menten kinetics and Pirt functions following the
144 reaction chain of denitrification. The predicted CH_4 flux was influenced by CH_4 production, oxidation and transportation
145 derived from the module of fermentation in the DNDC (Li, 2007). Methane production and oxidation occurred
146 simultaneously and were determined by the sizes of the aerobic (production) and anaerobic (oxidation) microsites, which
147 were defined by an Eh calculator in terms of an "anaerobic balloon" (" CH_4 balloon") (Li, 2007). The predicted CH_4
148 production was calculated from the carbon substrates resulting from decomposed soil organic carbon (SOC) and plant root
149 biomass with the effects of soil temperature (Li, 2000, 2016). For more details, please see Li. (2000, 2007) and Zhang *et al.*
150 (2018b).

151 2.1.2 Modifications of the CNMM-DNDC model

152 In the CNMM-DNDC, the soil temperature was predicted by solving the one-dimensional heat conduction equation
153 with the implicit method of Crank-Nicholson. However, despite the simple parameterization used for the calculation of soil
154 heat capacity and thermal conductivity, the variations of soil temperature induced by the freeze-thaw cycles were also not
155 considered (Table S1 of the online supplementary materials), which inevitably hindered its application in seasonally frozen
156 regions. In this study, the CNMM-DNDC was modified by replacing the above soil thermal module by a physical based
157 module of Northern Ecosystem Soil Temperature (Zhang *et al.*, 2003; Deng *et al.*, 2014), which can explicitly describe the
158 energy exchange within the soil, the active layer dynamics and the soil thermal regime in the presence of freeze-thaw cycles.
159 These modifications are indispensable for accurately simulating freeze-thaw cycles in seasonally frozen regions, which are
160 crucial for characterizing the active layer and soil thermal dynamics, soil hydrology and nitrogen or carbon cycling in these
161 regions. Therefore, the CNMM-DNDC with and without the above modifications are hereafter referred to as the original and
162 modified model, respectively.

163 The modified thermal dynamics of the soil are calculated by the one-dimensional heat conduction equation (Eq. 1).
164 The equation is solved numerically by converting to an explicit form (Eqs. 2–4), which is more efficient for considering the

165 freeze-thaw cycles (Zhang *et al.*, 2003). In the above equations, C ($\text{J m}^{-3} \text{ }^{\circ}\text{C}^{-1}$), k ($\text{W m}^{-1} \text{ }^{\circ}\text{C}^{-1}$), T ($^{\circ}\text{C}$) and G (W m^{-2}) denote
166 the soil heat capacity, thermal conductivity, soil temperature and heat fluxes between layers, respectively. Both Z and D are
167 the thicknesses of the soil layer (m), Δt is the time step of the calculation, and l denotes the soil layer l . S is the internal heat
168 exchange due to freezing or thawing (W m^{-3}) when the soil temperature is around $0 \text{ }^{\circ}\text{C}$. The soil temperature changes
169 affected by freezing or thawing are determined on the basis of energy conservation, which indicate that the latent heat
170 released during freezing equalled the amount of heat required for the increased soil temperature and vice versa. The dynamic
171 soil heat capacity (C_l , $\text{J m}^{-3} \text{ }^{\circ}\text{C}^{-1}$) is the weighted average of the heat capacity for five constituents, including organic matter
172 ($C_{l, \text{OM}}$), minerals ($C_{l, \text{Min}}$), water ($C_{l, \text{Water}}$), ice ($C_{l, \text{Ice}}$) and air ($C_{l, \text{Air}}$) (Eq. 5). The values of heat capacity for organic matter,
173 minerals, water, ice and air were 2.5×10^6 , 2.0×10^6 , 4.2×10^6 , 2.1×10^6 and $1.2 \times 10^3 \text{ J m}^{-3} \text{ }^{\circ}\text{C}^{-1}$, respectively (Huang, 2000). The
174 weight is the relative volumetric fraction of each constituent ($\theta_{l, \text{OM}}$, $\theta_{l, \text{Min}}$, $\theta_{l, \text{Water}}$, $\theta_{l, \text{Ice}}$, $\theta_{l, \text{Air}}$) in the soil. The dynamic
175 thermal conductivity (k_l , $\text{W m}^{-1} \text{ }^{\circ}\text{C}^{-1}$) is calculated using the thermal conductivities of above five constituents (Eq. 6–13),
176 with values of 0.25 ($k_{l, \text{OM}}$), 2.9 ($k_{l, \text{Min}}$), 0.57 ($k_{l, \text{Water}}$), 2.2 ($k_{l, \text{Ice}}$) and 0.025 ($k_{l, \text{Air}}$) $\text{W m}^{-1} \text{ }^{\circ}\text{C}^{-1}$ for organic matter, minerals,
177 water, ice and air, respectively (Johansen, 1975). ST_l denotes the soil temperature of layer l ($^{\circ}\text{C}$). The upper and lower
178 boundary conditions of the thermal dynamics were determined by the surface energy balance and the defined geothermal
179 heat flux at a soil depth of 35 m.

$$C \frac{\partial T}{\partial t} = \frac{\partial}{\partial Z} \left(k \frac{\partial T}{\partial Z} \right) + S \quad (1)$$

$$C_l \frac{\Delta T_l}{\Delta t} = \frac{G_{l-1, l} - G_{l, l+1}}{D_l} + S_l \quad (2)$$

$$G_{l-1, l} = \frac{(0.5k_l + 0.5k_{l-1})(T_{l-1} - T_l)}{0.5D_{l-1} + 0.5D_l} \quad (3)$$

$$G_{l, l+1} = \frac{(0.5k_l + 0.5k_{l+1})(T_l - T_{l+1})}{0.5D_l + 0.5D_{l+1}} \quad (4)$$

$$C_l = C_{l, \text{OM}}\theta_{l, \text{OM}} + C_{l, \text{Min}}\theta_{l, \text{Min}} + C_{l, \text{Water}}\theta_{l, \text{Water}} + C_{l, \text{Ice}}\theta_{l, \text{Ice}} + C_{l, \text{Air}}\theta_{l, \text{Air}} \quad (5)$$

$$k_l = \frac{\theta_{l, \text{Water}}k_{l, \text{Water}} + F_{l, \text{Air}}\theta_{l, \text{Air}}k_{l, \text{Air_adj}} + F_{l, \text{OM+Min}}(\theta_{l, \text{OM}} + \theta_{l, \text{Min}})k_{l, \text{OM+Min}} + F_{l, \text{Ice}}\theta_{l, \text{Ice}}k_{l, \text{Ice}}}{\theta_{l, \text{Water}} + F_{l, \text{Air}}\theta_{l, \text{Air}} + F_{l, \text{OM+Min}}(\theta_{l, \text{OM}} + \theta_{l, \text{Min}}) + F_{l, \text{Ice}}\theta_{l, \text{Ice}}} \quad (6)$$

$$k_{l, \text{Air_adj}} = \begin{cases} k_{l, \text{Air}} + 0.0238e^{0.0536ST_l} & (\theta_{l, \text{Water}} > 0.09) \\ 0.418 \times (0.0615 + 1.96\theta_{l, \text{Water}}) & (\theta_{l, \text{Water}} \geq 0.09) \end{cases} \quad (7)$$

$$g_a = \begin{cases} 0.333 - \frac{0.298\theta_{l, \text{Air}}}{1 - \theta_{l, \text{OM}} - \theta_{l, \text{Min}}} & (\theta_{l, \text{Water}} > 0.09) \\ 0.013 + 0.944\theta_{l, \text{Water}} & (\theta_{l, \text{Water}} \geq 0.09) \end{cases} \quad (8)$$

$$g_c = 1.0 - 2.0g_a \quad (9)$$

$$k_{l, \text{OM+Min}} = k_{l, \text{OM}} \frac{\theta_{l, \text{OM}}}{\theta_{l, \text{OM}} + \theta_{l, \text{Min}}} + k_{l, \text{Min}} \frac{\theta_{l, \text{Min}}}{\theta_{l, \text{OM}} + \theta_{l, \text{Min}}} \quad (10)$$

$$F_{l, \text{Air}} = 0.333 \left(\frac{2.0}{1.0 + g_a \left(\frac{k_{l, \text{Air adj}}}{k_{l, \text{Water}}} - 1.0 \right)} + \frac{1.0}{1.0 + g_c \left(\frac{k_{l, \text{Air adj}}}{k_{l, \text{Water}}} - 1.0 \right)} \right) \quad (11)$$

$$F_{l, \text{OM+Min}} = 0.333 \left(\frac{2.0}{1.0 + 0.125 \left(\frac{k_{l, \text{OM+Min}}}{k_{l, \text{Water}}} - 1.0 \right)} + \frac{1.0}{1.0 + 0.75 \left(\frac{k_{l, \text{OM+Min}}}{k_{l, \text{Water}}} - 1.0 \right)} \right) \quad (12)$$

$$F_{l, \text{Ice}} = 0.333 \left(\frac{2.0}{1.0 + 0.125 \left(\frac{k_{l, \text{Ice}}}{k_{l, \text{Water}}} - 1.0 \right)} + \frac{1.0}{1.0 + 0.75 \left(\frac{k_{l, \text{Ice}}}{k_{l, \text{Water}}} - 1.0 \right)} \right) \quad (13)$$

Compared to the original thermal module, the internal heat exchange due to freezing or thawing (S) was included with improved algorithm for thermal conductivity (k). In addition, the one-dimensional heat conduction equation (Eq. 1) was solved by converting it to an explicit form in the modified model (Eqs. 2–4), while was solved with the implicit method in the original models (Table S1). The modified CNMM-DNDC was able to simulate the thermal dynamics in seasonally frozen regions as well as their impacts on biogeochemical processes, such as the emissions of nitrogen and carbon gases.

2.2 Catchment and field descriptions

The study area is the Rierlangshan catchment (34°02'N, 102°43'E) on the eastern Tibetan Plateau with an area of 189 ha (Yao *et al.*, 2019). This catchment is located in the source region of the Pai-Lung River, which is a sub-branch of the upper Yangtze River (Zhang *et al.*, 2018a; 2019). This region is subject to a cold humid continental monsoon climate, and it had an annual mean air temperature of 1.6 ± 0.7 °C and average annual precipitation of 649 ± 94 mm in 1980–2012 as observed at the Zoige Meteorological Station (~80 km south of the catchment) (Ma *et al.*, 2018). The catchment consists of alpine wetlands, meadows and forests (Yao *et al.*, 2019). The alpine wetlands in the catchment are part of the Zoige wetland and are degraded due to anthropogenic drainage and climate warming (Dong *et al.*, 2010; Li *et al.*, 2014). Degraded alpine wetlands are commonly distributed throughout the Zoige wetland, and nearly 83% of the permanently inundated wetlands have been converted into “wet grassland” (Xiang *et al.*, 2009; Li *et al.*, 2014).

CH₄ and N₂O fluxes were manually measured once or twice per week using the gas chromatograph-based static opaque chamber method (Zhang *et al.*, 2018a) at three sites in alpine wetlands (34°02'6.53"N, 102°43'29.66"E, 3304 m a.s.l.), meadows (34°02'01"N, 102°43'28"E, 3326 m a.s.l.) and forests (34°01'47.13"N, 102°44'0.87"E, 3415 m a.s.l.) in the Rierlangshan catchment from 2013 to 2015 (Zhang *et al.*, 2018a; Yao *et al.*, 2019; Zhang *et al.*, 2019) (Fig. S1). Each chamber was wrapped with a layer of styrofoam and aluminium foil to mitigate temperature increases inside the enclosures due to the heating of solar radiation. The alpine wetland site is located at a slope base with a slope of 2°. The wetland has suffered from anthropogenic drainage and climate warming, and thus degraded to be seasonally inundated. The alpine meadow site neighbours the alpine wetland site, which is located on the north-facing slope with gradient of 11°. In addition, soil temperatures at different depths and topsoil moisture in the alpine wetlands, meadows and forests were observed daily

205 and twice per week, respectively. The details regarding the available field observations of the CH₄ and N₂O fluxes and the
206 relevant auxiliary variables are described in Table S2.

207 2.3 Model simulation

208 The modified CNMM-DNDC was applied in the Rierlangshan catchment with the three alpine ecosystems: wetlands,
209 meadows and forests. The dataset required for the catchment simulation included (1) a digital elevation model (DEM) with a
210 resolution of 30 × 30 m² from the geospatial data cloud (Fig. S1; <http://www.gscloud.cn/>); (2) a map of alpine ecosystems,
211 including wetlands, meadows and forests; (3) a climate dataset of 3-hour weather data (air temperature, precipitation, wind
212 speed, solar radiation, longwave radiation, and humidity), which were obtained from the meteorological station in the target
213 catchment for the years with field observations (2013.11–2015.10) and were adapted from the daily data at the Zoige
214 Meteorological Station (provided by the National Meteorological Information Center: <http://data.cma.cn/>; last access: 10th
215 June, 2020) for other years; (4) a soil properties dataset of the observed clay fraction, organic matter content, total nitrogen,
216 pH and bulk density of the three alpine ecosystems in 1 m soil profile (Ma *et al.*, 2018; Zhang *et al.*, 2018a; Yao *et al.*, 2019;
217 Zhang *et al.*, 2019; Table S3); and (5) a management practices dataset including grazing time and intensity for the alpine
218 wetlands and meadows (Table S3). In addition, other required soil inputs of field capacity, wilting point and saturated
219 hydrological conductivity were calculated by pedo-transfer functions (Li *et al.*, 2019; Table S4). The simulated soil depth
220 was defined as 35 m due to the lower boundary conditions of the thermal dynamics, which was set as the geothermal heat
221 flux at a soil depth of 35 m. The simulated soil profile (0–35 m depth) was divided into 23 layers, including the soil (0–1.5
222 m) and bedrock (1.5–35 m). The layer thicknesses of the soil (0–1.5 m) were 1, 5 10 and 50 cm for the depth of 0–10, 10–20,
223 20–100 and 100–150 cm, respectively. The layer thicknesses of the bedrock (1.5–35 m) were 3.5 and 31m for the depth of
224 1.5–4.0 and 4.0–35 m, respectively. The geothermal heat flux in the catchment was estimated at 0.053 W m⁻² (Pollack and
225 Chapman, 1977). For the target catchment, the soil water dynamics of the alpine ecosystems were determined by the
226 precipitation, evapotranspiration, infiltration, penetration and lateral flow. Using the database, a catchment simulation of
227 hydro-biogeochemical processes was performed with spatial and temporal resolutions of 30×30 m² and 3 hours, respectively,
228 by the modified CNMM-DNDC from 2012 to 2015, which could reflect the influences of hydrological processes on soil
229 water dynamics. Thus, the soil water dynamics of the seasonally inundated wetlands were determined by the hydrological
230 processes without any artificial disturbances in the catchment simulation.

231 2.4 Statistics and analysis

232 The statistical criteria applied for evaluating the model performance in this study included (i) the index of agreement
233 (IA), (ii) the Nash–Sutcliffe efficiency (NSE), and (iii) the determination coefficient (R^2) and slope of the zero-intercept
234 univariate linear regression (ZIR) of the observations against the simulations (e.g., Nash and Sutcliffe, 1970; Willmott and
235 Matsuura, 2005; Moriasi *et al.*, 2007; Congreves *et al.*, 2016; Jiang, 2010; Dubache *et al.*, 2019). A value of IA (0–1) closer
236 to 1 showed a better simulation. An NSE value (ranging from minus infinity to 1) closer to 1 was better. Better model

performance was indicated by a slope and an R^2 value that were both closer to 1 in a significant ZIR. For more details on these criteria, refer to the Eqs. S1–4 in Table S5. In addition, the SPSS Statistics Client 19.0 (SPSS Inc., Chicago, USA) and Origin 8.0 (OriginLab, Northampton, MA, USA) software packages were applied for the statistical analysis and graphical comparison.

3 Results

3.1 Model validation

3.1.1 Soil temperature and moisture

The profile soil temperatures were observed for alpine wetland and meadow, but only topsoil temperature was observed for the alpine forest, which could be used for model validation. The simulated soil temperatures of the three typical alpine ecosystems were significantly improved by including the scientific processes of soil thermal dynamics suitable for seasonally frozen regions (Figs. 1 and S2). The simulated seasonal dynamics and magnitudes were consistent with those from the field observations for various soil depths, with IA, NSE, and ZIR slopes and R^2 values of 0.91–1.00, 0.68–0.99, 0.83–1.09 and 0.73–1.00 for the three alpine ecosystems, respectively (Table 1). For the observed alpine wetlands and meadows, the simulation showed that the freezing of soil started in early November and continued to the end of April in the next year. The frozen depth reached a maximum in the middle of February. However, the simulated maximum frozen depths for the observed alpine meadows (0.69–0.74 m) were approximately double those for the alpine wetlands (0.30–0.39 m) (Fig. S3).

For the soil moisture, only topsoil moisture was observed in the three alpine ecosystems, which could be applied for model validation. The simulated topsoil moisture dynamics were comparable to those from the field observations, with IA and NSE values of 0.49–0.83 and -0.80–0.32 for the three alpine ecosystems, respectively (Fig. 2 and Table 1). In comparison to the other alpine ecosystems, the alpine wetlands had higher soil moisture, which ranged from 0.41 to 0.98 and from 0.38 to 0.93 for the observations and simulations in the water-filled pore space (WFPS), respectively. The soil moisture values of the alpine meadows and forests were highly variable and depended on the variation trend in precipitation for both observations and simulations. However, an underestimation of soil moisture in the winter period occurred for both alpine meadows and forests due to a possible overestimation of evapotranspiration. The performances of the modified model in simulating the soil profile temperature and topsoil moisture indicate that the modified CNMM-DNDC can generally predict the soil thermal and topsoil moisture dynamics in the three alpine ecosystems, which is crucial for correctly simulating soil hydrology, plant growth and biogeochemical processes.

265 3.1.2 Methane fluxes

266 The daily observed CH₄ emissions from the alpine wetlands were highly variable and showed a clear seasonal cycle,
267 with intensive CH₄ emissions from May to November and weak emissions in other periods (Fig. 3a). The observed alpine
268 meadows and forests functioned exclusively as sinks of atmospheric CH₄ with higher rates of uptake during the growing
269 season and lower uptake rates in the dormant season (Figs. 3b–c). The original model significantly overestimated CH₄
270 emissions from the alpine wetlands. The modified CNMM-DNDC accurately identified the functions of the sources or sinks
271 in the three alpine ecosystems and generally captured the magnitude and seasonal characteristics of the daily CH₄ fluxes,
272 with an IA of 0.57–0.88 for the three alpine ecosystems (Figs. 3a–c and Table 1). However, the CH₄ uptake rates during the
273 dormant season were obviously underestimated by the modified model at both sites, especially at the alpine forest site, which
274 was responsible for the underestimation of cumulative CH₄ uptake. The observed cumulative CH₄ emissions ranged from -
275 2.60 to 33.5 kg C ha⁻¹ yr⁻¹ and the modelled values ranged from -1.90 to 31.0 kg C ha⁻¹ yr⁻¹ (Fig. 4a). For the catchment
276 simulation, the simulated annual CH₄ emissions ranged from -2.35 to 73.0 kg C ha⁻¹ yr⁻¹ from November 2013 to November
277 2014 (Fig. S4a). These results indicate that the modified CNMM-DNDC well simulated the CH₄ fluxes of the three typical
278 alpine ecosystems.

279 3.1.3 Nitrous oxide fluxes

280 The daily observed N₂O emissions from the alpine wetlands were higher than those from the alpine meadows but
281 lower than those from the alpine forests (Figs. 3d–f). Similar seasonal patterns of N₂O fluxes were observed for the three
282 alpine ecosystems with intensive emissions in the growing season. The N₂O emission peak during the dormant season was
283 observed in the alpine meadows, which was the major contributor to annual emissions. The modified CNMM-DNDC
284 generally captured the seasonal dynamics of daily N₂O fluxes with an IA of 0.26–0.47 for the three alpine ecosystems (Figs.
285 3d–f and Table 1), but the N₂O emissions from the alpine wetlands were significantly overestimated by the original model.
286 For the modified model, the simulated N₂O emissions from the alpine wetlands and forests showed obvious seasonal patterns
287 with higher emissions during the growing season, but no abrupt emission peak was captured at the end of the growing season
288 for the alpine wetlands. In addition, compared with the original model, the modified model captured the peak emissions that
289 occurred during the freeze-thaw period from the alpine meadows due to the death of microbes, but the dynamics of the peak
290 emissions were not well simulated. The observed cumulative N₂O emissions ranged from 0.14 to 0.58 kg N ha⁻¹ yr⁻¹ and the
291 modelled values ranged from 0.12 to 0.32 kg N ha⁻¹ yr⁻¹ (Fig. 4b). For the catchment simulation, the simulated annual N₂O
292 emissions ranged from 0.01 to 0.74 kg N ha⁻¹ yr⁻¹ from November 2013 to November 2014 (Fig. S4b). These results indicate
293 that the modified CNMM-DNDC has the potential to estimate N₂O emissions in seasonally frozen regions.

294 3.2 Annual aggregate emissions of CH₄ and N₂O

295 Annual aggregate emissions of CH₄ and N₂O in carbon dioxide (CO₂) equivalents were calculated for the three alpine
296 ecosystems from November 2013 to November 2014 for alpine wetlands and meadow and from April 2014 to April 2015 for
297 alpine forests, and the global warming potentials were 34 for CH₄ and 298 for N₂O on a 100-year time horizon (IPCC, 2013).
298 The simulated aggregate emissions by the modified model were 1.5, 0.015, and 0.061 Mg CO₂eq ha⁻¹ yr⁻¹ for the observed
299 alpine wetlands, meadows and forests, respectively, which were consistent with those from the field observations (1.6, 0.014,
300 and 0.15 Mg CO₂eq ha⁻¹ yr⁻¹ for the alpine wetlands, meadows and forests, respectively) (Fig. 4c). However, the original
301 model significantly overestimated the aggregate emissions due to the high predicted CH₄ and N₂O emissions. In comparison,
302 the observed seasonally inundated wetlands functioned as the sources of aggregate emissions of CH₄ and N₂O, but the
303 aggregate emissions from adjacent wet alpine meadows were much lower.

304 4 Discussions

305 4.1 Model performance in simulating thermal dynamics

306 The soil freeze-thaw cycles in seasonally frozen regions determine the soil profile temperature and hydrological
307 processes, which are key factors that regulate the cycling of nitrogen and carbon (e.g., Zhang *et al.*, 2015; Hugelius *et al.*,
308 2020). Therefore, improving the scientific processes of soil thermal dynamics in the presence of active layer dynamics is
309 essential for applying the CNMM-DNDC to simulate the biogeochemical processes in seasonally frozen regions, which are
310 sensitive and vulnerable to climate change and human activities (Hatano, 2019; Hugelius *et al.*, 2020; Jiang *et al.*, 2020). The
311 original model adopted a relatively simple module to calculate thermal transportation within the soil profile and did not
312 consider the effects of freeze-thaw cycles on soil temperature and moisture. The newly incorporated module was based on
313 explicit energy conservation and exchange in the soil profile and successfully captured the variations in soil temperature and
314 topsoil moisture for the three alpine ecosystems during the freeze-thaw period. The simulated lower soil frozen depth for the
315 observed alpine wetland was primarily attributed to the higher soil profile moisture level, as the thermal conductivity and
316 heat capacity for water-filled pores were higher than those for air-filled pores. In order to quantify the impacts of climate
317 change on the cycling of carbon and water on the regional and global scales, several large scale ecosystem models or
318 macroscale hydrological models, such as Terrestrial Ecosystem Model, Lund-Potsdam-Jena dynamic global vegetation
319 model and Variable Infiltration Capacity model, have been enhanced to simulate the soil thermal dynamics at northern high
320 latitude (e.g., Wania *et al.*, 2009; Zhuang *et al.*, 2001; Cuo *et al.*, 2015; Jiang *et al.*, 2020). In addition, the soil thermal
321 modules were also improved in some biogeochemical models, such as DNDC and Mobile-DNDC, to evaluate the influences
322 of climate warming on the biogeochemical processes in high latitude regions (e.g., Zhang *et al.*, 2003; de Bruijn *et al.*, 2009;
323 Wolf *et al.*, 2011; Zhang *et al.*, 2012; Deng *et al.*, 2014). Compared with the simulated soil profile temperatures by above
324 models at different scales, the simulations in this study by the modified CNMM-DNDC were equally well, especially for

deeper soil layers (e.g., Wania *et al.*, 2009). For the validated topsoil moisture in this study, the modified model generally captured the variation trends, which were comparable with the performances of other models (e.g., de Bruijn *et al.*, 2009; Wolf *et al.*, 2011; Cuo *et al.*, 2015). However, compared with the studies focused on simulating soil moisture (e.g., Ford *et al.*, 2014), further improvements are still required to improve the model performance in simulating the soil moisture. These results indicate the efficiency of the incorporated module in simulating soil thermal and topsoil moisture dynamics in seasonally frozen regions.

4.2 Model performance in simulating CH₄ fluxes

Compared with the annually inundated wetlands, the seasonally inundated wetlands had relatively low observed and simulated CH₄ emissions due to the significant influences of the water table level on CH₄ emissions (e.g., Hatano, 2019; Zhang *et al.*, 2019). The CH₄ emissions simulated by the CNMM-DNDC were determined by the processes of production, oxidation and transpiration. The unsaturated soil with moisture levels ranging from 0.41 to 0.98 WFPS resulted in a small CH₄ balloon and thus reduced CH₄ production. At the same time, relatively dry conditions caused the upper soil layer to act as an efficient oxidative methanotrophic barrier for the diffusion of CH₄ from the subsoil and thus decreased CH₄ emissions (Kandel *et al.*, 2018; Tan *et al.*, 2020). In addition, the highly fluctuating CH₄ emissions simulated by the modified model were also attributed to the high dependency of CH₄ production on soil moisture, which controlled the size of the CH₄ balloon. Theoretically, the CH₄ emissions simulated by the original model should not be higher than those simulated by the modified model due to the lower predicted soil moisture level. The overestimated CH₄ emissions simulated by the original model were mainly attributed to the overestimated soil temperature due to their influences on mineralized substrates for CH₄ production, as well as the processes of CH₄ production. This result implies that global warming may trigger intensive CH₄ emissions from degraded wetlands, which could partly serve as a trade-off for the decreased CH₄ emissions due to the lower water table level in degraded wetlands (e.g., Gong *et al.*, 2020). For the studies focused on simulating CH₄ emissions from wetlands by the large-scale ecosystem models, the model validation with field observation is difficult due to coarse spatial resolution (e.g., Zhuang *et al.*, 2004). For the biogeochemical model, such as DNDC, the dynamics of CH₄ emissions from wetland and peatland in the northern permafrost regions were well simulated (Zhang *et al.*, 2012; Deng *et al.*, 2014), which showed consistent seasonal variations and magnitudes as those in this study. Both observations and simulations showed that the CH₄ uptake in alpine forests was higher than that in alpine meadows, which was mainly attributed to the high SOC content of the alpine forests in the simulation. Methane uptake by upland soils is a biological process governed by the availability of CH₄ and oxygen as well as the activity and quantity of methanotrophic bacteria in soils (e.g., Liu *et al.*, 2007; Zhang *et al.*, 2014). In the model, the simulated CH₄ uptake was positively related to the SOC content, which is closely related to the population size of methanotrophic bacteria. Thus, the SOC content primarily contributed to the differences in CH₄ uptake from alpine meadows and forests, as the values for forests were more than twice of those for meadows (Table S3). As the simulated dynamic characteristics of CH₄ uptake were primarily regulated by soil temperature and moisture, the inhibitory effects of low soil temperature (< 0.0 °C) on CH₄ uptake rates resulted in obvious underestimations in the dormant season for both

358 alpine meadows and forests. Therefore, an improved parameterization for simulating CH₄ uptake under low soil temperatures
359 is required for the model to better capture the dynamics of CH₄ uptake in the dormant season.

360 4.3 Model performance in simulating N₂O fluxes

361 In comparison, the N₂O emissions from the alpine wetlands and forests were higher than those from the alpine
362 meadows for both the observations and simulations due to the high SOC content and nitrogen availability. Natural wetlands
363 are large carbon reserves and play a crucial role in mitigating global warming (e.g., Deng *et al.*, 2014; Kang *et al.*, 2020; Tan
364 *et al.*, 2020). The intentional drainage of annually inundated wetlands alters not only the water regime but also nutrient
365 availability (e.g., Hoffmann *et al.*, 2016). The simulated relatively low soil moisture for the alpine wetlands stimulated the
366 decomposition of SOC and nitrogen (or peat oxidation) under aerobic conditions, thus improving nitrogen mineralization for
367 nitrification and denitrification and enhancing N₂O emissions (e.g., Tan *et al.*, 2020; Zhang *et al.*, 2020). The intensive N₂O
368 emissions simulated by the original model resulted from the overestimated soil temperature for the alpine wetlands. Firstly,
369 as the presence of ice could impede the water movement, the water lateral flows were promoted by the original model due to
370 the neglecting of freeze-thaw cycles. These further resulted in the lower simulated soil moisture as compared with the
371 modified model (Fig. S5), which provided favorable oxygen conditions for N₂O production. Meanwhile, the simulated high
372 soil moisture by the modified model provided feasible anaerobic conditions for thoroughly denitrification. Secondly, higher
373 simulated soil temperature by the original model also facilitated the mineralization, which provided more available mineral
374 nitrogen. Field studies showed that high SOC concentrations could stimulate the processes of mineralization and nitrification
375 in the forests (e.g., Li *et al.*, 2005; Yao *et al.*, 2019). The model input of soil organic matter measured in the observed alpine
376 forests was more than twice that in the observed alpine meadows (Table S3). Thus, the high SOC content at the alpine forest
377 site provided more available nitrogen through mineralization and thus stimulated the nitrification processes in the simulation.
378 Furthermore, the seasonal grazing that occurred in the alpine meadows resulted in constant loss of available nitrogen and
379 thus hindered the N₂O emissions from the biological processes in the simulation. Field observations showed that the soil
380 freeze-thaw cycles occurred in seasonally frozen regions not only increased the availability of nitrogen and carbon substrates
381 by disrupting of soil aggregates but also affected the structure, population and activity of the microbes, and thus influencing
382 the emissions of N₂O (e.g., Song *et al.*, 2019). de Bruijn *et al.* (2009) have explored the combined mechanisms for
383 simulating freeze-thaw related N₂O emissions, which were the promoted anaerobiosis and denitrification due to reduced gas
384 diffusion derived from soil frost and snow cover, and the stimulated microbial growth due to easy decomposable organic
385 carbon and nitrogen derived from the dead microbes during freeze-thaw cycles. Wolf *et al.* (2011) introduced an impedance
386 factor to parameterize the reduced water flow between layers in the presence of ice, which could captured the freeze-thaw
387 related N₂O emissions for ungrazed steppe. In the CNMM-DNDC, threshold values of soil temperature were set to trigger
388 the death of microbes during the freezing period and stimulate the production of NO, N₂O and N₂ using substrates derived
389 from the dead microbes during the thawing period, which was similar to one of the mechanisms explored by de Bruijn *et al.*
390 (2009). However, compared with the simulated freeze-thaw related N₂O emissions by other studies, the simulated dynamics

391 of peak emissions due to freeze-thaw cycles in this study were inconsistent with those from the field observations. Thus,
392 improvements are required to incorporate some other effective mechanisms to better capture the dynamic characteristics. The
393 peak emissions during the freeze-thaw period were not captured by the original model due to the significantly overestimated
394 soil temperature. The low evaluation statistics for the daily fluxes, especially for the alpine forests, were also attributed to the
395 underestimation of background emissions, which resulted from both measurement errors due to low fluxes around detection
396 limits ($\pm 0.41 \text{ g N ha}^{-1} \text{ d}^{-1}$) and model deficiencies in the simulation of tight nitrogen cycling in natural ecosystems.

397 Compared with the empirical model, one key advantage of the process-oriented models is that the models are
398 independent of the local parameterization (Zhang *et al.*, 2015). In this study, default internal parameter combinations of
399 biogeochemical processes were used for the original and modified models, which have been applied in the catchment
400 simulation in the subtropical region (Zhang *et al.*, 2018b), due to the limited field observations (only one year) for both
401 calibration and validation. The biogeochemical processes were predicted by the first-order and Michaelis-Menten kinetics in
402 the CNMM-DNDC based on some defined parameters of flow fractionation. For instance, there are 17 parameters related
403 with N_2O emission in the module of denitrification (Table S6), which would inevitably increase the uncertainty of simulation.
404 Houska *et al.* (2017) found that hydro-biogeochemical models can be right for the wrong reasons, such as matching
405 greenhouse gas emissions while failing to simulate soil moisture, which emphasized the importance for simultaneous
406 validations of multi-variables. Thus, simultaneous validations of CH_4 and N_2O fluxes, as well as soil environment variables,
407 were necessary for comprehensive evaluation of the model performance. In addition, the microbial ecology was recently
408 recommended to be integrated into the biogeochemical model using a smaller number of well-defined kinetic parameters,
409 such as MOMOS (Pansu *et al.*, 2010; Treseder *et al.*, 2011). Therefore, direct control of microbial on biogeochemical
410 processes, such as the stoichiometry of decomposer, is required to be included in the CNMM-DNDC in near future. The
411 model performances of simulating various variables for three typical alpine ecosystems in the Rierlangshan catchment imply
412 that the modified CNMM-DNDC can be applied to predict the thermal dynamics and fluxes of CH_4 and N_2O from alpine
413 ecosystems in seasonally frozen regions.

414 5 Conclusions

415 To apply the process-oriented hydro-biogeochemical model Catchment Nutrient Management Model - DeNitrification-
416 DeComposition (CNMM-DNDC) in seasonally frozen regions, an improved module of soil thermal dynamics for describing
417 the soil thermal regime in the presence of freeze-thaw cycles was incorporated in this study. Using the unique experimental
418 dataset obtained in the Rierlangshan catchment with the typical alpine wetland, meadow and forest ecosystems, the modified
419 model was evaluated for simulating soil thermal dynamics (soil profile temperature), topsoil moisture and fluxes of methane
420 (CH_4) and nitrous oxide (N_2O) in seasonally frozen regions of the Tibetan Plateau. The modified CNMM-DNDC could
421 generally capture the seasonal dynamics and magnitudes of profile soil temperature, topsoil moisture and fluxes of CH_4 and
422 N_2O in seasonally frozen regions. Both the observed and simulated CH_4 and N_2O fluxes from three alpine ecosystems

423 indicate that the aggregate emissions of CH₄ and N₂O were highest for the wetland among three alpine ecosystems. The
424 intensive aggregate emissions of CH₄ and N₂O were regulated by the high soil moisture, which was primarily determined by
425 the CH₄ emissions. This study implies that a hydro-biogeochemical model, such as the modified CNMM-DNDC, are able to
426 predict soil thermal dynamics, topsoil moisture and fluxes of CH₄ and N₂O in seasonally frozen regions with an improved
427 physical-based soil thermal module.

428 **Data availability**

429 The model, input and output databases can be obtained from the first author and all the observed data sets used in this study
430 can be available from the co-authors.

431 **Author contribution**

432 Zheng, X. and Zhang, W. contributed to developing the idea and enhancing the science of this study. Zhang, W. improved
433 the scientific processes of the model, implemented the model simulations and prepared the manuscript with contributions
434 from all co-authors. Li, S. improved the model structure for standard input. Yao, Z., Zhang, H., Ma, L., Wang, K., Wang, R.
435 and Liu, C. designed and carried out the field experiments. Han, S. collected and established the input database for modelling.
436 Deng, J and Li, Y contributed to the modification of the model and the improvement of the manuscript.

437 **Competing interests**

438 The authors declare that they have no conflict of interest.

439 **Acknowledgement**

440 This study was jointly supported by the National Key R&D Program of China (2016YFA0602303), the Chinese Academy of
441 Sciences (ZDBS-LY-DQC007), the National Key Scientific and Technological Infrastructure project “Earth System Science
442 Numerical Simulator Facility” (EarthLab) and the National Natural Science Foundation of China (41603075, 41861134029).

443 **References**

444 Bechmann, M., 2014. Long-term monitoring of nitrogen in surface and subsurface runoff from small agricultural dominated
445 catchments in Norway. *Agric. Ecosyst. Environ.* 198, 13–24.
446 Bosch, N., Allan, J., Dolan, D., Han, H., Richards, R., 2011. Application of the Soil and Water Assessment Tool for six
447 watersheds of Lake Erie: Model parameterization and calibration. *J. Great Lakes Res.* 37, 263–271.
448 Breuer, L., VachÉ, K., Julich, S., Frede, H., 2010. Current concepts in nitrogen dynamics for mesoscale catchments. *Hydrol.*
449 *Sci. J.-J. Sci. Hydrol.* 53, 1059–1074.

Canfield, D., Glazer, A., Falkowski, P., 2010. The evolution and future of Earth's nitrogen cycle. *Science* 330, 192–196.

Castellano, M., Lewis, D., Kaye, J., 2013. Response of soil nitrogen retention to the interactive effects of soil texture, hydrology, and organic matter. *J. Geophys. Res.-Biogeosci.* 118, 280–290.

Chen, D., Li, Y., Grace, P., Mosier, A., 2008. N₂O emissions from agricultural lands: a synthesis of simulation approaches. *Plant Soil* 309, 169–189.

Cheng, K., Ogle, S., Parton, W., Pan, G., 2014. Simulating greenhouse gas mitigation potentials for Chinese Croplands using the DAYCENT ecosystem model. *Glob. Change Biol.* 20, 948–962.

Collatz, G., Ribas-Carbo, M., Berry, J., 1992. Coupled photosynthesis-stomatal conductance model for leaves of C₄ plants. *Aust. J. Plant Physiol.*, 19, 519–538.

Collins, A., Zhang, Y., Winter, M., Inman, A., Jones, J., Johnes, P., Cleasby, W., Vrain, E., Lovett, A., Noble, L., 2016. Tackling agricultural diffuse pollution: What might uptake of farmer-preferred measures deliver for emissions to water and air? *Sci. Total Environ.* 547, 269–281.

Congreves, K., Grant, B., Dutta, B., Smith, W., Chantigny, M., Rochette, Desjardins, R., 2016. Prediction ammonia volatilization after field application of swine slurry: DNDC model development, *Agric. Ecosyst. Environ.* 219, 179–189.

Cui, Z., Zhang, H., Chen, X., Zhang, C., Ma, W., Huang, C., Zhang, W., Mi, G., Miao, Y., Li, X., Gao, Q., Yang, J., Wang, Z., Ye, Y., Guo, S., Lu, J., Huang, J., Lv, S., Sun, Y., Liu, Y., Peng, X., Ren, J., Li, S., Deng, X., Shi, X., Zhang, Q., Yang, Z., Tang, L., Wei, C., Jia, L., Zhang, J., He, M., Tong, Y., Tang, Q., Zhong, X., Liu, Z., Cao, N., Kou, C., Ying, H., Yin, Y., Jiao, X., Zhang, Q., Fan, M., Jiang, R., Zhang, F., Dou, Z., 2018. Pursuing sustainable productivity with millions of smallholder farmers. *Nature* 555, 363–366.

Cui, F., Zheng, X., Liu, C., Wang, K., Zhou, Z., Deng, J., 2014. Assessing biogeochemical effects and best management practice for a wheat-maize cropping system using the DNDC model. *Biogeosciences* 11, 91–107.

Cuo, L., Zhang, Y., Bohn, T., Zhao, L., Li, J., Liu, Q., Zhou, B., 2015. Frozen soil degradation and its effects on surface hydrology in the northern Tibetan Plateau. *J. Geophys. Res. Atmos.* 120, 8276–8298.

de Bruijn, A.M.G., Butterbach-Bahl, K., Blagodatsky, S., Grote, R., 2009. Model evaluation of different mechanisms driving freeze–thaw N₂O emissions. *Agric. Ecosyst. Environ.* 133, 196–207.

Deng, J., Li, C., Frolking, S., Zhang, Y., Bäckstrand, K., Crill, P., 2014. Assessing effects of permafrost thaw on C fluxes based on multiyear modeling across a permafrost thaw gradient at Stordalen, Sweden. *Biogeosciences* 11, 4753–4770.

Dong, Z., Hu, G., Yan, C., Wang, W., Lu, J., 2010. Aeolian desertification and its causes in the Zoige Plateau of China's Qinghai–Tibetan Plateau. *Environ. Earth Sci.* 59, 1731–1740.

Dubache, G., Li, S., Zheng, X., Zhang, W., Deng, J., 2019. Modeling ammonia volatilization following urea application to winter cereal fields in the United Kingdom by improving a biogeochemical model, *Sci. Total Environ.* 660, 1403–1418.

Farquhar, G., Caemmerer, S., Berry, J., 1980. A biochemical model of photosynthetic CO₂ assimilation in leaves of C₃ species. *Planta*, 149, 78–90.

Fenner, N., Freeman, C., 2011. Drought-induced carbon loss in peatlands. *Nat. Geosci.* 4, 895–900.

Foereid, B., Barthram, G., Marriott, C., 2007. The CENTURY model failed to simulate soil organic matter development in an acidic grassland. *Nutr. Cycl. Agroecosyst.* 78, 143–153.

Ford, T., Harris, E., Quiring, S., 2014. Estimating root zone soil moisture using near-surface observations from SMOS. *Hydrol. Earth Syst. Sci.* 18, 139–154.

Galloway, J., Dentener, F., Capone, D., Boyer, E., Howarth, R., Seitzinger, S., Asner, G., Cleveland, C., Green, P., Holland, E., Karl, D., Michaels, A., Porter, J., Townsend, A., Vorosmarty, C., 2004. Nitrogen Cycles: past, present, and future. *Biogeochemistry* 70, 153–226.

Galloway, J., Townsend, A., Erisman, J., Bekunda, M., Cai, Z., Freney, J., Martinelli, L., Seitzinger, S., Sutton, M., 2008. Transformation of the nitrogen cycle: recent trends, questions, and potential solutions. *Science* 320, 889–892.

Giltrap, D.L., Li, C., Sagggar, S., 2010. DNDC: A process-based model of greenhouse gas fluxes from agricultural soils. *Agric. Ecosyst. Environ.* 136, 292–300.

Gong, Y., Wu, J., Vogt, J., Ma, W., 2020. Greenhouse gas emissions from peatlands under manipulated warming, nitrogen addition, and vegetation composition change: a review and data synthesis. *Environ. Rev.* 28, 428–437.

Haas, E., Klatt, S., Fröhlich, A., Kraft, P., Werner, C., Kiese, R., Grote, R., Breuer, L., Butterbach-Bahl, K., 2012. LandscapeDNDC: a process model for simulation of biosphere–atmosphere–hydrosphere exchange processes at site and regional scale. *Landsc. Ecol.* 28, 615–636.

500 Hatano, R., 2019. Impact of land use change on greenhouse gases emissions in peatland: a review. *Int. Agrophys.* 33, 167–
501 173.

502 Holzworth, D., Huth, N., deVoil, P., Zurcher, E., Herrmann, N., McLean, G., Chenu, K., van Oosterom, E., Snow, V.,
503 Murphy, C., Moore, A., Brown, H., Whish, J., Verrall, S., Fainges, J., Bell, L., Peake, A., Poulton, P., Hochman, Z.,
504 Thorburn, P., Gaydon, D., Dalgliesh, N., Rodriguez, D., Cox, H., Chapman, S., Doherty, A., Teixeira, E., Sharp, J.,
505 Cichota, R., Vogeler, I., Li, F., Wang, E., Hammer, G., Robertson, M., Dimes, J., Whitbread, A., Hunt, J., van Rees, H.,
506 McClelland, T., Carberry, P., Hargreaves, J., MacLeod, N., McDonald, C., Harsdorf, J., Wedgwood, S., Keating, B.,
507 2014. APSIM – Evolution towards a new generation of agricultural systems simulation. *Environ. Modell. Softw.* 62,
508 327–350.

509 Houska, T., Kraus, D., Kiese, R., Breuer, L., 2017. Constraining a complex biogeochemical model for CO₂ and N₂O
510 emission simulations from various land uses by model–data fusion. *Biogeosciences* 14, 3487–3508.

511 Huang, C., 2000. *Soil Science*. China Agriculture Press, Beijing. Pp. 125. (In Chinese)

512 Hugelius, G., Loisel, J., Chadburn, S., Jackson, R., Jones, M., MacDonald, G., Marushchak, M., Olefeldt, D., Maara, P.,
513 Siewert, M., Treat, C., Turetsky, M., Voigt, C., Yu, Z., 2020. Large stocks of peatland carbon and nitrogen are
514 vulnerable to permafrost thaw. *Proc. Natl. Acad. Sci. U. S. A.* doi: 10.1073/pnas.1916387117

515 Intergovernmental Panel on Climate Change (IPCC): Climate Change 2013: The Physical Science Basis, Contribution of
516 Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change (eds. Stocker TF,
517 Qin D, Plattner G-K, et al.), Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2013.

518 Jiang, H., Yi, Y., Zhang, W., Yang, K., Chen, D., 2020. Sensitivity of soil freeze/thaw dynamics to environmental conditions
519 at different spatial scales in the central Tibetan Plateau. *Sci. Total Environ.* 734, 139261. doi:
520 org/10.1016/j.scitotenv.2020.139261

521 Jiang, Z., 2010. Analysis on the establishment conditions of the square sum decomposition formular of regression model, *J.*
522 *Industr. Techn. Econ.* 29(4), 116–119 (in Chinese).

523 Johansen, O., 1975. *Thermal conductivity of soils*. Ph.D. thesis, Univ. of Trondheim, Trondheim, Norway.

524 Ju, X., Xing, G., Chen, X., Zhang, S., Zhang, L., Liu, X., Cui, Z., Yin, B., Christie, P., Zhu, Z., Zhang, F., 2009. Reducing
525 environmental risk by improving N management in intensive Chinese agricultural systems. *Proc. Natl. Acad. Sci. U.*
526 *S. A.* 106, 3041–3046.

527 Kandel, T., Lærke, P., Elsgaard, L., 2018. Annual emissions of CO₂, CH₄ and N₂O from a temperate peat bog: comparison of
528 an undrained and four drained sites under permanent grass and arable crop rotations with cereals and potato. *Agric. For.*
529 *Meteorol.* 256–257, 470–481.

530 Kang, X., Li, Y., Wang, J., Yan, L., Zhang, X., Wu, H., Yan, Z., Zhang, K., Hao, Y., 2020. Precipitation and temperature
531 regulate the carbon allocation process in alpine wetlands: quantitative simulation. *J. Soils Sediments* 20, 3300–3315.

532 Keiluweit, M., Wanzek, T., Kleber, M., Nico, P., Fendorf, S., 2017. Anaerobic microsites have an unaccounted role in soil
533 carbon stabilization. *Nat. Commun.* 8. doi: 10.1038/s41467-017-01406-6

534 Klatt, S., Kraus, D., Kraft, P., Breuer, L., Wlotzka, M., Heuveline, V., Haas, E., Kiese, R., Butterbach-Bahl, K., 2017.
535 Exploring impacts of vegetated buffer strips on nitrogen cycling using a spatially explicit hydro-biogeochemical
536 modeling approach. *Environ. Modell. Softw.* 90, 55–67.

537 Li, B., Yu, Z., Liang, Z., Song, K., Li, H., Wang, Y., Zhang, W., Acharya, K., 2014. Effects of climate variations and human
538 activities on runoff in the Zoige alpine wetland in the eastern edge of the Tibetan Plateau. *J. Hydrol. Eng.* 19, 1026–
539 1035.

540 Li, C., 2000. Modeling trace gas emissions from agricultural ecosystems. *Nutr. Cycl. Agroecosyst.* 58, 259–276.

541 Li, C., 2007. Quantifying greenhouse gas emissions from soils: scientific basis and modeling approach. *Soil Sci. Plant Nutr.*
542 53, 344–352.

543 Li, C., 2016. *Biogeochemistry: Scientific Fundamentals and Modelling Approach*. Tsinghua University Press, Beijing. Pp.
544 530. (In Chinese)

545 Li, C., Frolking, S., Butterbach-Bahl, K., 2005. Carbon sequestration in arable soils is likely to increase nitrous oxide
546 emissions, offsetting reductions in climate radiative forcing. *Clim. Change* 72, 321–338.

547 Li, S., Zheng, X., Zhang, W., Han, S., Deng, J., Wang, K., Wang, R., Yao, Z., Liu, C., 2019. Modeling ammonia
548 volatilization following the application of synthetic fertilizers to cultivated uplands with calcareous soils using an
549 improved DNDC biogeochemistry model. *Sci. Total Environ.* 660, 931–946.

Li, Y., White, R., Chen, D., Zhang, J., Li, B., Zhang, Y., Huang, Y., Edis, R., 2007. A spatially referenced water and nitrogen management model (WNMM) for (irrigated) intensive cropping systems in the North China Plain. *Ecol. Model.* 203, 395–423.

Liu, C., Holst, J., Brüggemann, N., Butterbach-Bahl, K., Yao, Z., Yue, J., Han, S., Han, X., Krümmelbein, J., Horn, R., Zheng, X., 2007. Winter-grazing reduces methane uptake by soils of a typical semi-arid steppe in Inner Mongolia, China. *Atmos. Environ.* 41, 5948–5958.

Liu, S., Xie, Z., Zeng, Y., Liu, B., Li, R., Wang, Y., Wang, L., Qin, P., Jia, B., Xie, J., 2019. Effects of anthropogenic nitrogen discharge on dissolved inorganic nitrogen transport in global rivers. *Glob. Change Biol.* 25, 1493–1513.

Ma, L., Yao, Z., Zheng, X., Zhang, H., Wang, K., Zhu, B., Wang, R., Zhang, W., Liu, C., 2018. Increasing grassland degradation stimulates the non-growing season CO₂ emissions from an alpine meadow on the Qinghai-Tibetan Plateau. *Environ. Sci. Pollut. Res.* 25, 26576–26591.

McClain, M., Boyer, E., Dent, C., Gergel, S., Grimm, N., Groffman, P., Hart, S., Harvey, J., Johnston, C., Mayorga, E., 2003. Biogeochemical hot spots and hot moments at the interface of terrestrial and aquatic ecosystems. *Ecosystems* 6, 301–312.

Moriasi, D., Arnold, J., Van Liew, M., Bingner, R., Harmel, R., Veith, T., 2007. Model evaluation guidelines for systematic quantification of accuracy in watershed simulation, *T. Am. Soc. Agr. Biol. Eng.* 50, 885–900.

Nash, J., Sutcliffe, J., 1970. River flow forecasting through conceptual models: part I- a discussion of principles, *J. Hydrol.* 10, 282–290.

Pansu, M., Sarmiento, L., Rujano, M., Ablan, M., Acevedo, D., Bottner, P., 2010. Modelling Organic transformations by Micro-Organisms of Soils in six contrasting ecosystems: validation of the MOMOS model. *Glob. Biogeochem. Cycl.* 24, GB1008. doi: org/10.1029/2009GB003527

Pansu, M., Machado, D., Bottner, P., Sarmiento, L., 2014. Modelling microbial exchanges between forms of soil nitrogen in contrasting ecosystems. *Biogeosciences* 11, 915–927.

Piao, S., Fang, J., Ciais, P., Peylin, P., Huang, Y., Sitch, S., Wang, T., 2009. The carbon balance of terrestrial ecosystems in China. *Nature* 458, 1009–1013.

Pohler, T., Huisman, J., Breuer, L., Frede, H., 2007. Integration of a detailed biogeochemical model into SWAT for improved nitrogen predictions—Model development, sensitivity, and GLUE analysis. *Ecol. Model.* 203, 215–228.

Pollack, H., Chapman, D., 1977. On the regional variation of heat flow, geotherms, and lithospheric thickness. *Tectonophysics* 38, 279–296.

Schroock, A., Gaube, V., Haas, E., Winiwarter, W., 2019. Estimating nitrogen flows of agricultural soils at a landscape level - A modelling study of the Upper Enns Valley, a long-term socio-ecological research region in Austria. *Sci. Total Environ.* 665, 275–289.

Schuur, E., McGuire, A., Schadel, C., Grosse, G., Harden, J., Hayes, D., Hugelius, G., Koven, C., Kuhry, P., Lawrence, D., Natali, S., Olefeldt, D., Romanovsky, V., Schaefer, K., Turetsky, M., Treat, C., Vonk, J., 2015. Climate change and the permafrost carbon feedback. *Nature* 520, 171–179.

Seitzinger, S., 2008. Nitrogen cycle - Out of reach. *Nature* 452, 162–163.

Song, L., Yao, Y., Lin, L., Gao, W., Cai, T., Liang, H., Gao, D., 2019. The potential source of nitrous oxide in the pristine riparian marsh during freeze-thaw cycles, case study in Northeast China. *Ecol. Eng.* 134, 18–25.

Tan, L., Ge, Z., Zhou, X., Li, S., Li, X., Tang, J., 2020. Conversion of coastal wetlands, riparian wetlands, and peatlands increases greenhouse gas emissions: a global meta-analysis. *Glob. Change Biol.* 26, 1638–1653.

Todd-Brown, K., Hopkins, F., Kivlin, S., Jennifer, M., Talbot, J., Allison, S., 2012. A framework for representing microbial decomposition in coupled climate models. *Biogeochemistry* 109, 19–33.

Treseder, K., Balser, T., Bradford, M., Brodie, E., Dubinsky, E., Eviner, V., Hofmockel, K., Lennon, J., Levine, U., MacGregor, B., Pett-Ridge, J., Waldrop, M., 2011. Integrating microbial ecology into ecosystem models: challenges and priorities. *Biogeochemistry* 109, 7–18.

Vereecken, H., Schnepf, A., Hopmans, J., Javaux, M., Or, D., Roose, T., Vanderborght, J., Young, M., Amelung, W., Aitkenhead, M., Allison, S., Assouline, S., Baveye, P., Berli, M., Brüggemann, N., Finke, P., Flury, M., Gaiser, T., Govers, G., Ghezzehei, T., Hallett, P., Hendricks Franssen, H., Heppell, J., Horn, R., Huisman, J., Jacques, D., Jonard, F., Kollet, S., Lafolie, F., Lamorski, K., Leitner, D., McBratney, A., Minasny, B., Montzka, C., Nowak, W., Pachepsky, Y., Padarian, J., Romano, N., Roth, K., Rothfuss, Y., Rowe, E., Schwen, A., Šimůnek, J., Tiktak, A., Van Dam, J., van

der Zee, S., Vogel, H., Vrugt, J., Wöhling, T., Young, I., 2016. Modeling Soil Processes: Review, Key Challenges, and New Perspectives. *Vadose Zone J.* 15, doi:10.2136/vzj2015.2009.0131.

Wania, R., Ross, I., Prentice, I., 2009. Integrating peatlands and permafrost into a dynamic global vegetation model: 1. Evaluation and sensitivity of physical land surface processes. *Glob. Biogeochem. Cycl.* 23, GB3014, doi:10.1029/2008GB003412.

Wigmosta, M., Vail, L., Lettenmaier, D., 1994. A distributed hydrology-vegetation model for complex terrain. *Water Resour. Res.*, 30, 1665–1679.

Willmott, C., Matsuura, K., 2005. Advantages of the mean absolute error (MAE) over the root mean square error (RMSE) in assessing average model performance. *Clim. Res.*, 30, 79–82.

Wolf, B., Kiese, R., Chen, W., Grote, R., Zheng, X., Butterbach-Bahl, K., 2011. Modelling N₂O emissions from steppe in Inner Mongolia, China, with consideration of spring thaw and grazing intensity. *Plant Soil* 350, 297–310.

Wu, Y., Liu, S., Qiu, L., Sun, Y., 2016. SWAT-DayCent coupler: An integration tool for simultaneous hydro-biogeochemical modeling using SWAT and DayCent. *Environ. Modell. Softw.* 86, 81–90.

Xiang, S., Guo, R., Wu, N., Sun, S., 2009. Current status and future prospects of Zoige Marsh in Eastern Qinghai-Tibet Plateau. *Ecol. Eng.* 35, 553–562.

Yao, Z., Ma, L., Zhang, H., Zheng, X., Wang, K., Zhu, B., Wang, R., Wang, Y., Zhang, W., Liu, C., Butterbach-Bahl, K., 2019. Characteristics of annual greenhouse gas flux and NO release from alpine meadow and forest on the eastern Tibetan Plateau. *Agric. For. Meteorol.* 272–273, 166–175.

Zhang, H., Yao, Z., Ma, L., Zheng, X., Wang, R., Wang, K., Liu, C., Zhang, W., Zhu, B., Tang, X., Hu, Z., Han, S., 2019. Annual methane emissions from degraded alpine wetlands in the eastern Tibetan Plateau. *Sci. Total Environ.* 657, 1323–1333.

Zhang, H., Yao, Z., Wang, K., Zheng, X., Ma, L., Wang, R., Liu, C., Zhang, W., Zhu, B., Tang, X., Hu, Z., Han, S., 2018a. Annual N₂O emissions from conventionally grazed typically alpine grass meadows in the eastern Qinghai-Tibetan Plateau. *Sci. Total Environ.* 625, 885–899.

Zhang, W., Li, Y., Zhu, B., Zheng, X., Liu, C., Tang, J., Su, F., Zhang, C., Ju, X., Deng, J., 2018b. A process-oriented hydro-biogeochemical model enabling simulation of gaseous carbon and nitrogen emissions and hydrologic nitrogen losses from a subtropical catchment. *Sci. Total Environ.* 616–617, 305–317.

Zhang, W., Liu, C., Zheng, X., Fu, Y., Hu, X., Cao, G., Butterbach-Bahl, K., 2014. The increasing distribution area of zokor mounds weaken greenhouse gas uptakes by alpine meadows in the Qinghai-Tibetan Plateau. *Soil Biol. Biochem.* 71, 105–112.

Zhang, W., Liu, C., Zheng, X., Wang, K., Cui, F., Wang, R., Li, S., Yao, Z., Zhu, J., 2019. Using a modified DNDC biogeochemical model to optimize field management of a multi-crop (cotton, wheat, and maize) system: a site-scale case study in northern China. *Biogeosciences* 16, 2905–2922.

Zhang, W., Liu, C., Zheng, X., Zhou, Z., Cui, F., Zhu, B., Haas, E., Klatt, S., Butterbach-Bahl, K., Kiese, R., 2015. Comparison of the DNDC, LandscapeDNDC and IAP-N-GAS models for simulating nitrous oxide and nitric oxide emissions from the winter wheat-summer maize rotation system. *Agric. Syst.* 140, 1–10.

Zhang, W., Wang, J., Hu, Z., Li, Y., Yan, Z., Zhang, X., Wu, G., Yan, L., Zhang, K., Kang, X., 2020. The primary drivers of greenhouse gas emissions along the water table gradient in the Zoige alpine peatland. *Water Air Soil Pollut.* 231: 224. doi: org/10.1007/s11270-020-04605-y

Zhang, Y., Chen, W., Cihlar, J., 2003. A process-based model for quantifying the impact of climate change on permafrost thermal regimes. *J. Geophys. Res.-Atmos.* 108, 4695.

Zhang, Y., Sachs, T., Li, C., Boike, J., 2012. Upscaling methane fluxes from closed chambers to eddy covariance based on a permafrost biogeochemistry integrated model. *Glob. Change Biol.* 18, 1428–1440.

Zhang, Y., Shao, Q., Ye, A., Xing, H., Xia, J., 2016. Integrated water system simulation by considering hydrological and biogeochemical processes: model development, with parameter sensitivity and autocalibration. *Hydrol. Earth Syst. Sci.* 20, 529–553.

Zhu, Q., Castellano, M., Yang, G., 2018. Coupling soil water processes and the nitrogen cycle across spatial scales: potentials, bottlenecks and solutions. *Earth-Sci. Rev.* 187, 248–258.

Zhu, Q., Schmidt, J.P., Bryant, R., 2012. Hot moments and hot spots of nutrient losses from a mixed land use watershed. *J. Hydrol. Eng.* 414, 393–404.

650 Zhuang, Q., Romanovsk, V., McGuire, A., 2001. Incorporation of a permafrost model into a large-scale ecosystem model:
651 Evaluation of temporal and spatial scaling issues in simulating soil thermal dynamics. *J. Geophys. Res.* 106, 33649–
652 33670.

653 Zhuang, Q., Melillo, J., Kicklighter, D., Prinn, R., McGuire, A., Steudler, P., Felzer, B., Hu, S., 2004. Methane fluxes
654 between terrestrial ecosystems and the atmosphere at northern high latitudes during the past century: a retrospective
655 analysis with a process-based biogeochemistry model. *Glob. Biogeochem. Cycl.* 18, GB3010,
656 doi:10.1029/2004GB002239.

657

658 Table 1 Statistics of the validated variables by the modified CNMM-DNDC for three typical alpine ecosystems.

659

Item	Ecosystem	n^a	DR ^b	IA		NSE		ZIR-slope		ZIR- R^{2c}		ZIR-P	
				O ^c	M ^d	O	M	O	M	O	M	O	M
Soil temperature													
5 cm	Meadow	500	Daily	0.90	0.96	0.82	0.95	0.84	1.09	0.89	0.96	< 0.001	< 0.001
	Forest	48	Daily	0.85	0.91	0.37	0.68	0.64	0.83	0.68	0.73	< 0.001	< 0.001
10 cm	Wetland	366	Daily	0.90	0.98	0.57	0.92	0.72	1.07	0.81	0.93	< 0.001	< 0.001
	Meadow	500	Daily	0.93	0.99	0.71	0.95	0.80	1.08	0.85	0.96	< 0.001	< 0.001
20 cm	Wetland	366	Daily	0.82	0.99	0.18	0.96	0.64	1.05	0.66	0.97	< 0.001	< 0.001
	Meadow	500	Daily	0.87	0.99	0.48	0.97	0.74	1.06	0.76	0.98	< 0.001	< 0.001
50 cm	Wetland	366	Daily	0.66	0.99	-1.01	0.97	0.51	1.05	0.43	0.97	< 0.001	< 0.001
	Meadow	401	Daily	0.70	1.00	-0.48	0.99	0.58	1.06	0.53	1.00	< 0.001	< 0.001
70 cm	Wetland	366	Daily	0.58	0.98	-2.23	0.93	0.47	1.05	0.38	0.93	< 0.001	< 0.001
	Meadow	401	Daily	0.64	1.00	-1.19	0.99	0.54	1.03	0.49	1.00	< 0.001	< 0.001
90 cm	Wetland	366	Daily	0.52	0.98	-4.07	0.90	0.44	1.03	0.36	0.90	< 0.001	< 0.001
Soil moisture	Wetland	74	Daily	0.63	0.83	-1.65	0.20	1.31	1.13	–	0.60	–	< 0.001
	Meadow	128	Daily	0.78	0.78	0.28	0.32	0.96	0.93	0.30	0.41	< 0.001	< 0.001
	Forest	40	Daily	0.48	0.49	-1.04	-0.80	1.21	1.19	–	–	–	–
Daily CH ₄ flux	Wetland	180	Daily	0.37	0.74	-11.1	-0.73	0.46	0.87	–	–	–	–
	Meadow	168	Daily	0.87	0.88	0.42	0.38	1.09	0.94	0.44	0.39	< 0.001	< 0.001
	Forest	49	Daily	0.59	0.57	-2.79	-3.39	0.92	0.79	–	–	–	–
Daily N ₂ O flux	Wetland	180	Daily	0.01	0.26	-323	-0.07	0.01	0.59	–	–	–	–
	Meadow	168	Daily	0.23	0.44	-0.16	-1.76	0.99	0.35	–	–	–	–
	Forest	58	Daily	0.47	0.47	-1.85	-1.64	0.44	0.47	–	–	–	–

660 ^a n indicates the number of the observations. ^b DR denotes the time resolution of the observed data. ^c O indicates the simulations by the original
661 model. ^d M indicates the simulations by the modified model. ^e “–” indicated no value due to the sum of regression square are larger than the sum
662 of the total square for the regression). IA, NSE, ZIR-slope, ZIR- R^2 and ZIR-P indicate the index of agreement, Nash–Sutcliffe efficiency,
663 determination coefficient and slope of the zero-intercept univariate linear regression (ZIR) of the observations against the simulations, as well as
664 the significance level of the ZIR.




Figure: 1 Observed and simulated daily profile soil temperature from the alpine wetlands and meadows by the original and modified models. The legends in panel a apply for all panels.




Figure: 2 Observed and simulated daily topsoil (0–6 cm) moisture in the water-filled pore space (WFPS) from the alpine wetlands, meadows and forests by the original and modified models. The legends in panel a apply for all panels.





Figure: 3 Observed and simulated daily methane (CH_4) and nitrous oxide (N_2O) fluxes from the alpine wetlands, meadows and forests by the original and modified models. The vertical bar for each observation indicates the standard error of six spatial replicates. The legends in panel a apply for all panels.



675

676 **Figure: 4 Observed and simulated annual emissions of methane (CH₄), nitrous oxide (N₂O) and aggregate emissions of both from**
 677 **the alpine wetlands (Wetland), meadows (Meadow) and forests (Forest). The legends in panel a apply for all panels.**