

Validation of a coupled $\delta^2\text{H}_{n\text{-alkane}}\text{-}\delta^{18}\text{O}_{\text{sugar}}$ paleohygrometer approach based on a climate chamber experiment

Johannes Hepp^{a,b,1,*}, Christoph Mayr^{c,d}, Kazimierz Rozanski^e, Imke Kathrin Schäfer^f, Mario Tuthorn^{g,2},
Bruno Glaser^b, Dieter Juchelka^g, Willibald Stichler^h, Roland Zech^{i,j,3}, Michael Zech^{b,j,4,#}

^aChair of Geomorphology and BayCEER, University of Bayreuth, Universitätsstrasse 30, D-95440 Bayreuth, Germany

^bInstitute of Agricultural and Nutritional Sciences, Soil Biogeochemistry, Martin Luther University Halle-Wittenberg, Von-Seckendorff-Platz 3, D-06120 Halle (Saale), Germany

^cInstitute of Geography, Friedrich-Alexander-University Erlangen-Nürnberg, Wetterkreuz 15, D-91058 Erlangen, Germany

^dGeoBio-Center & Earth and Environmental Sciences, Ludwig-Maximilian University Munich, Richard-Wagner-Str. 10, D-80333 München, Germany

^eFaculty of Physics and Applied Computer Science, AGH University of Science and Technology, Al. Mickiewicza 30, PL-30-059 Kraków, Poland

^fInstitute of Geography and Oeschger Centre for Climate Research, University of Bern, Hallerstrasse 12, CH-3012 Bern, Switzerland

^gThermo Fisher Scientific, Hanna-Kunath-Str. 11, D-28199 Bremen, Germany

^hHelmholtz Zentrum München, German Research Center for Environmental Health, Ingolstädter Landstrasse 1, D-85764 Neuherberg, Germany

ⁱInstitute of Geography, Chair of Physical Geography, Friedrich-Schiller University of Jena, Löbdergraben 32, D-07743 Jena, Germany

^jInstitute of Geography, Heisenberg Chair of Physical Geography with focus on paleoenvironmental research, Technische Universität Dresden, Helmholtzstrasse 10, D-01062 Dresden, Germany

*corresponding author: johannes-hepp@gmx.de

¹Present address: Chair of Geomorphology and BayCEER, University of Bayreuth, Universitätsstrasse 30, D-95440 Bayreuth, Germany

²Present address: Thermo Fisher Scientific, Hanna-Kunath-Str. 11, D-28199 Bremen, Germany

³Present address: Institute of Geography, Chair of Physical Geography, Friedrich-Schiller University of Jena, Löbdergraben 32, D-07743 Jena, Germany

⁴Present address: Institute of Geography, Heisenberg Chair of Physical Geography with focus on paleoenvironmental research, Technische Universität Dresden, Helmholtzstrasse 10, D-01062 Dresden, Germany

Gelöscht: Agronomy

Gelöscht: -

Gelöscht: -

Keywords

hydrogen stable isotopes, oxygen stable isotopes, hemicellulose sugars, leaf waxes, leaf water enrichment, deuterium-excess, relative humidity

Abstract

The hydrogen isotope composition of leaf wax-derived biomarkers, e.g. long chain n -alkanes ($\delta^2\text{H}_{n\text{-alkane}}$), is widely applied in paleoclimatology. However, a direct reconstruction of the isotope composition of source water based on $\delta^2\text{H}_{n\text{-alkane}}$ alone is challenging due to the enrichment of heavy-isotopes during evaporation. The coupling of $\delta^2\text{H}_{n\text{-alkane}}$ with $\delta^{18}\text{O}$ of hemicellulose-derived sugars ($\delta^{18}\text{O}_{\text{sugar}}$) has the potential to disentangle this limitation and additionally to allow relative humidity reconstructions. Here, we present $\delta^2\text{H}_{n\text{-alkane}}$ as well as $\delta^{18}\text{O}_{\text{sugar}}$ results obtained from leaves of *Eucalyptus globulus*, *Vicia faba*, and *Brassica oleracea*, which grew under controlled conditions. We addressed the following questions (i) do $\delta^2\text{H}_{n\text{-alkane}}$ and $\delta^{18}\text{O}_{\text{sugar}}$ values allow reconstructions of leaf water isotope composition, (ii) how accurately does the reconstructed leaf-water-isotope composition enable relative humidity (RH) reconstruction, and (iii) does the coupling of $\delta^2\text{H}_{n\text{-alkane}}$ and $\delta^{18}\text{O}_{\text{sugar}}$ enable a robust source water calculation? For all investigated species, the n -alkane $n\text{-C}_{29}$ was most abundant and therefore used for compound-specific $\delta^2\text{H}$ measurements. For *Vicia faba*, additionally the $\delta^2\text{H}$ values of $n\text{-C}_{31}$ could be evaluated robustly. With regard to hemicellulose-derived monosaccharides, arabinose and xylose were most abundant and their $\delta^{18}\text{O}$ values were therefore used to calculate weighted mean leaf $\delta^{18}\text{O}_{\text{sugar}}$ values. Both $\delta^2\text{H}_{n\text{-alkane}}$ and $\delta^{18}\text{O}_{\text{sugar}}$ yielded significant correlations with $\delta^2\text{H}_{\text{leaf-water}}$ and $\delta^{18}\text{O}_{\text{leaf-water}}$, respectively ($r^2 = 0.45$ and 0.85 , respectively; $p < 0.001$, $n = 24$). Mean fractionation factors between biomarkers and leaf water were found to be -156‰ (ranging from -133 to -192‰) for $\epsilon_{n\text{-alkane/leaf-water}}$ and $+27.3\text{‰}$ (ranging from $+23.0$ to 32.3‰) for $\epsilon_{\text{sugar/leaf-water}}$, respectively. Modelled RH_{air} values from a Craig-Gordon model using measured T_{air} , $\delta^2\text{H}_{\text{leaf-water}}$ and $\delta^{18}\text{O}_{\text{leaf-water}}$ as input correlate highly significantly with modeled RH_{air} values ($R^2 = 0.84$, $p < 0.001$, $\text{RMSE} = 6\%$). When coupling $\delta^2\text{H}_{n\text{-alkane}}$ and $\delta^{18}\text{O}_{\text{sugar}}$ values, the correlation of modeled RH_{air} values with measured RH_{air} values is weaker but still highly significant with $R^2 = 0.54$ ($p < 0.001$, $\text{RMSE} = 10\%$). Finally, the reconstructed source water isotope composition ($\delta^2\text{H}_s$ and $\delta^{18}\text{O}_s$) as calculated from our coupled approach matches the source water in the climate chamber experiment ($\delta^2\text{H}_{\text{tank-water}}$ and $\delta^{18}\text{O}_{\text{tank-water}}$). This highlights the great potential of the coupled $\delta^2\text{H}_{n\text{-alkane}}\text{-}\delta^{18}\text{O}_{\text{sugar}}$ paleohygrometer approach for paleoclimate and relative humidity reconstructions.

Gelöscht: isotopic

Gelöscht: ology research

Gelöscht: isotopic

Gelöscht: can be

Gelöscht: alteration of the soil water isotopic signal by leaf-water

Gelöscht: enrichment

Gelöscht: effect

1 Introduction

Leaf wax-derived biomarkers such as long chain n -alkanes and their stable hydrogen isotope composition ($\delta^2\text{H}_{n\text{-alkane}}$) are widely applied in paleoclimatology. Sedimentary $\delta^2\text{H}_{n\text{-alkane}}$ values correlate with $\delta^2\text{H}$ of precipitation (Huang et al., 2004; Mügler et al., 2008; Sachse et al., 2004; Sauer et al., 2001), confirming the high potential of $\delta^2\text{H}_{n\text{-alkane}}$ to establish $\delta^2\text{H}$ records of past precipitation (Hou et al., 2008; Rao et al., 2009; Sachse et al., 2012). However, this interpretation is challenged by heavy isotope enrichment during evapotranspiration (e.g. Zech et al., 2015). Apart from studies of sedimentary cellulose (Heyng et al., 2014; Wissel et al., 2008), the oxygen stable isotope composition of sugar biomarkers ($\delta^{18}\text{O}_{\text{sugar}}$) emerged as complementary paleoclimate proxy during the last decade (Hepp et al., 2015, 2017, Zech et al., 2013a, 2014a; Zech and Glaser, 2009). The interpretation of the $\delta^{18}\text{O}_{\text{sugar}}$ values is comparable to those of $\delta^2\text{H}_{n\text{-alkane}}$. When sugars originate primarily from leaf biomass of higher terrestrial plants, they reflect the plant source water (which is often directly linked to the local precipitation) modified by evapotranspirative enrichment of the leaf water (Tuthorn et al., 2014; Zech et al., 2014a). The coupling of $\delta^2\text{H}_{n\text{-alkane}}$ with $\delta^{18}\text{O}_{\text{sugar}}$ values allows quantifying the leaf-water isotope enrichment and relative air humidity (Zech et al., 2013a). This approach was validated by Tuthorn et al. (2015) by applying it to topsoil samples along a climate transect in Argentina. Accordingly, the biomarker-derived relative air humidity values correlate significantly with actual relative air humidity, highlighting the potential of the $\delta^2\text{H}_{n\text{-alkane}}-\delta^{18}\text{O}_{\text{sugar}}$ paleohygrometer approach. The coupled approach is based on the observation that the isotope signature of precipitation ($\delta^2\text{H}_{\text{precipitation}}$ and $\delta^{18}\text{O}_{\text{precipitation}}$) typically plots near to the global meteoric water line (GMWL) in a two-dimensional $\delta^2\text{H}-\delta^{18}\text{O}$ diagram. The GMWL is characterized by the equation $\delta^2\text{H}_{\text{precipitation}} = 8 \cdot \delta^{18}\text{O}_{\text{precipitation}} + 10$ (Craig, 1961; Dansgaard, 1964). In many cases, the local precipitation is directly linked to the source water of plants, which is indeed soil water and shallow groundwater. The isotope composition of xylem water readily reflects these sources (e.g. Dawson, 1993). However, leaf-derived biomarkers reflect the leaf water isotope composition, which is, unlike xylem water, prone to evapotranspiration (e.g. Barbour and Farquhar, 2000; Helliker and Ehleringer, 2002; Cernusak et al., 2003; Barbour et al., 2004; Cernusak et al., 2005; Feakins and Sessions, 2010; Kahmen et al., 2011; Sachse et al., 2012; Kahmen, et al., 2013; Tipple et al., 2013; Lehmann et al., 2017; Liu et al., 2017). During daytime, the leaf water is typically enriched in the heavy isotope compared to the source water because of evapotranspiration through stomata. Thereby, lighter water isotopes evaporate preferentially, which leads to gradual enrichment of heavier isotopes compared to precipitation. The degree of enrichment by evapotranspiration is mainly controlled by the relative air humidity (RH_{air}) in the direct surrounding of the plant leaves (e.g. Cernusak et al., 2016). Although the biomarkers reflect the isotope composition of leaf water, there is still a modification of its isotope signature by fractionation during biosynthesis, leading to an offset between leaf water and biomarker isotope compositions. In case the biosynthetic fractionation is known and constant, RH_{air} can be calculated from coupling $\delta^2\text{H}_{n\text{-alkane}}$ with $\delta^{18}\text{O}_{\text{sugar}}$ values.

First applications of this approach for paleoclimatic reconstruction (Hepp et al., 2017, 2019; Zech et al., 2013a) and climate transect validation studies (Hepp et al., 2020; Lemma et al., 2021; Strobel et al., 2020; Tuthorn et al., 2015) revealed promising results. Furthermore, the stability of the biomarker isotope signals during degradation was studied (Zech et al., 2011, 2012). In brief, n -alkanes and sugars can be extracted compound-specifically from plants, soils and a wide range of different sediments, retaining the isotope signal of intact plant tissues.

Gelöscht: (Hepp et al., 2015, 2017, Zech et al., 2013a, 2014a)

Gelöscht: (Craig, 1961)

Gelöscht: isotopes

Gelöscht: Dansgaard, 1964)

Gelöscht:

Gelöscht: Indeed, there are already some f

Gelöscht: paleo

Gelöscht: as well as first

Gelöscht: (Hepp et al., 2020; Tuthorn et al., 2015)

Gelöscht: published

Gelöscht: amongst others by our working group

Gelöscht: investigated

Gelöscht: . This

Gelöscht: without the need means nofor

Gelöscht: of fossilized leaves are necessary

The overall aim of this study is to evaluate the $\delta^2\text{H}_{n\text{-alkane}}\text{-}\delta^{18}\text{O}_{\text{sugar}}$ paleohygrometer approach by applying it to plant leaf material from three different plants grown in a climate chamber experiment under controlled conditions. More specifically, we addressed the following questions:

- (i) which n -alkanes and monosaccharides can be used to obtain $\delta^2\text{H}_{n\text{-alkane}}$ and $\delta^{18}\text{O}_{\text{sugar}}$ results of the leaves grown in our climate chamber experiment, respectively,
- (ii) how precisely do $\delta^2\text{H}_{n\text{-alkane}}$ and $\delta^{18}\text{O}_{\text{sugar}}$ values allow reconstructing $\delta^2\text{H}_{\text{leaf-water}}$ and $\delta^{18}\text{O}_{\text{leaf-water}}$, respectively,
- (iii) how accurately does the leaf water isotope composition reflect RH_{air} ,
- (iv) does the coupling of $\delta^2\text{H}_{n\text{-alkane}}$ and $\delta^{18}\text{O}_{\text{sugar}}$ enable a RH_{air} reconstruction, and
- (v) how robust are source water calculations?

2 Material and Methods

2.1 Climate chamber experiment

A phytotron experiment was conducted at the Helmholtz Zentrum München in Neuherberg during winter 2000/2001 (Mayr, 2002). Three different dicotyledon plant species (*Eucalyptus globulus*, *Vicia faba* var. *minor* and *Brassica oleracea* var. *medullosa*) were grown in eight chambers for 56 days under seven distinct climatic conditions (same conditions in chambers 4 and 8). All three species belong to C_3 plants and were primarily chosen because they are different in terms of morphology and physiology. Two of the species are herbaceous (*Vicia faba* var. *minor* and *Brassica oleracea* var. *medullosa*), while *Eucalyptus globulus* is a lignifying tree. Nevertheless, all three species form a stem. The different habitus of the three plants allows to check for inter-species differences in isotope fractionation (Mayr, 2002). Similar isotope patterns in the three plants could allow generalizing our results for a wide range of C_3 plants, independent of their habitus. An additional, equally important criterion for the choice of the plants was the resilience of the taxa to the experimental climatic conditions and sufficient growth during the experiments. Air temperature (T_{air}) were set to 14, 18, 24 and 30°C and RH_{air} to around 20, 30, 50, and 70% during the daytime of constant climate conditions (between 11 a.m. and 4 p.m.) (Fig. 1A). During the rest of the day, diurnal variations typical for natural conditions were aimed for (for more details see Mayr, 2002).




Fig. 1: A: Climate chamber conditions. Error bars represent analytical standard deviation of the respective measurements. For more details see section 2.2 and Mayr (2002).

Gelöscht: The

Formatiert: Tiefgestellt

Gelöscht: very

Gelöscht: is

Gelöscht: best

Gelöscht: species-dependencies

Formatiert: Tiefgestellt

Gelöscht:

Gelöscht: between

Gelöscht: typical natural

Gelöscht: in

[2] verschoben

Gelöscht: (leaf temperature and relative humidity in green and air temperature and relative humidity in red)

Gelöscht: (see

Gelöscht: ,

Feldfunktion geändert

The chambers had a high fresh air supply rate ($750 \text{ m}^3 \text{ h}^{-1}$), avoiding therefore an evaporative isotope enrichment in the chambers. Furthermore, uniform irrigation conditions were guaranteed via an automatic irrigation system, which was controlled by tensiometers installed in 9 cm substrate depth. The tank water used for irrigation was sampled periodically (intervals of one to three days) over the whole experiment and revealed only minor variability in its isotope composition ($\delta^{18}\text{O}_{\text{tank-water}} = -10.7 \pm 0.3\text{‰}$ standard deviation (σ); $\delta^2\text{H}_{\text{tank-water}} = -7 \pm 1\text{‰}$ σ). Once a week, soil water (via ceramic cups in 13 cm soil depth) and atmospheric water vapor (via dry ice condensation traps) was sampled ($\delta^2\text{H}_{\text{soil-water}}$, $\delta^{18}\text{O}_{\text{soil-water}}$ and $\delta^2\text{H}_{\text{atmospheric-water-vapor}}$, $\delta^{18}\text{O}_{\text{atmospheric-water-vapor}}$). Additionally, leaf temperature (T_{leaf}) was calculated from gas exchange measurements, at least once a week (Mayr, 2002).

In order to analyze stable hydrogen and oxygen isotope composition of leaf ($\delta^2\text{H}_{\text{leaf-water}}$, $\delta^{18}\text{O}_{\text{leaf-water}}$) and stem water, the plants were harvested at the end of the experiment. The vacuum distillation method was used for the extraction of the plant water. It should be noted that stem water is a mixture between phloem and xylem water. Only the latter reflects the isotope composition of soil water. For simplification, stem water is referred to as xylem water in the following ($\delta^2\text{H}_{\text{xylem-water}}$, $\delta^{18}\text{O}_{\text{xylem-water}}$).

For more details about the experiment, [the \(plant\) water isotope results, and gas exchange measurements](#) we refer to the original publication (Mayr, 2002).

2.2 Leaf biomarker extraction and compound-specific stable isotope analysis

A total of 24 leaf samples were prepared according to Schäfer et al. (2016) for compound specific $\delta^2\text{H}$ measurements of *n*-alkanes, at the Institute of Geography, Group of Biogeochemistry and Paleoclimate, University of Bern. Microwave extraction with 15 ml dichloromethane (DCM)/methanol (MeOH) 9:1 (v:v) at 100°C for 1 h was conducted. The resulting total lipid extracts were purified and separated using aminopropyl-silica-gel (Supelco, $45 \mu\text{m}$) pipette columns. The hydrocarbon fractions (containing *n*-alkanes) were eluted with *n*-hexane and cleaned via silver nitrate-coated silica gel pipettes (Supelco, 60-200 mesh) and zeolite (Geokleen Ltd.) columns. The $\delta^2\text{H}$ measurements of the dominant *n*-alkanes (*n*- C_{29} and *n*- C_{31}) were performed on a GC- ^2H -pyrolysis-IRMS system, consisting of an Agilent 7890A gas chromatograph (GC) coupled to an IsoPrime 100 isotope-ratio-mass spectrometer (IRMS) via a GC5 pyrolysis/combustion interface operating in pyrolysis mode with a Cr (ChromeHD) reactor at 1000°C . The compound-specific $\delta^2\text{H}$ values were calibrated against a standard alkane mix (*n*- C_{27} , *n*- C_{29} , *n*- C_{33}) with known isotope composition (A. Schimmelmann, University of Indiana), measured twice every six sample injections. Standard deviation of the triplicate measurements was typically $\leq 5\text{‰}$. The H^{3+} factor stayed constant during the course of the measurements.

Additionally, the leaf samples were dried and finely ground in preparation for $\delta^{18}\text{O}$ analysis of hemicellulose-derived sugars (modified from Zech and Glaser, 2009) at the Institute of Agricultural and Nutritional Sciences, Soil Biogeochemistry, Martin Luther University Halle-Wittenberg. The hemicellulose sugars were hydrolytically extracted for 4 h at 105°C using 4M trifluoroacetic acid (Amelung et al., 1996) and purified via XAD-7 and Dowex 50WX8 columns. Prior to the methylboronic-acid (MBA) derivatization (4 mg of MBA in $400 \mu\text{l}$ dry pyridine for 1 h at 60°C), the cleaned sugars were frozen and freeze-dried overnight (Gross and Glaser, 2004; Knapp, 1979). Compound-specific $\delta^{18}\text{O}$ measurements were performed on a Trace GC 2000 coupled to a Delta V Advantage IRMS via an ^{18}O -pyrolysis reactor (GC IsoLink) and a ConFlo IV interface (all devices from Thermo Fisher Scientific, Bremen, Germany). The sample batches were measured along with embedded co-derivatized standard batches, which contained arabinose, fucose, xylose, and rhamnose in different concentrations of known $\delta^{18}\text{O}$ value. The $\delta^{18}\text{O}$ values of the standard sugars were determined via temperature

Gelöscht: /

Formatiert: Hochgestellt

Gelöscht: was avoided

Gelöscht: s

Gelöscht: were

Gelöscht: derived

Gelöscht: isotopic

Gelöscht: isotopic

Gelöscht: (Knapp, 1979)

230 conversion/elemental analysis-IRMS coupling at the Institute of Plant Sciences, ETH Zurich, Switzerland (Zech and Glaser, 2009). This procedure allows corrections for possible amount dependencies (Zech and Glaser, 2009) and ensures the “Principle of Identical Treatment” (Werner and Brand, 2001). Standard deviations for the triplicate measurements were 0.9‰ and 2.2‰ (average over all investigated samples) for arabinose and xylose, respectively. We focus on arabinose and xylose in this
 235 study because they were (i) the dominant peaks in all chromatograms, and (ii) previously found to strongly predominate over fucose (and rhamnose) in terrestrial plants and soils (Hepp et al., 2016).

Gelöscht: ,

All δ values are expressed in ‰ as isotope ratios ($R = {}^{18}\text{O}/{}^{16}\text{O}$ or ${}^2\text{H}/{}^1\text{H}$) relative to the Vienna Standard Mean Ocean Water (VSMOW) standard in the common delta notation ($\delta = (R_{\text{sample}} - R_{\text{standard}})/R_{\text{standard}}$; e.g. Coplen, 2011).

Gelöscht: Figure 1B summarizes isotope data obtained for the 24 analysed leaf samples where a

2.3 Framework for coupling $\delta^2\text{H}_{n\text{-alkane}}$ with $\delta^{18}\text{O}_{\text{sugar}}$ results

2.3.1 Deuterium-excess of leaf water and relative humidity

245 The coupled approach is based on the observation that isotope composition of global precipitation plots typically close to the GMWL ($\delta^2\text{H}_{\text{precipitation}} = 8 \cdot \delta^{18}\text{O}_{\text{precipitation}} + 10\text{‰}$; Craig, 1961; Fig. 4). The soil water and shallow groundwater, which acts as source water for plants, can often directly be related to the local precipitation. However, especially during daytime, leaf water is typically enriched in heavy isotopes compared to the precipitation due to evapotranspiration through the stomata, therefore plotting to the right of the GMWL (Fig. 4; e.g. Allison et al., 1985; Bariac et al., 1994; Walker and Brunel, 1990). During natural climatic conditions, the leaf water reservoir at the evaporative sites is frequently assumed to be in isotope steady-state (Allison et al., 1985; Bariac et al., 1994; Gat et al., 2007; Walker and Brunel, 1990), meaning that isotope composition of transpired water vapour is equal to the isotope composition of the source water utilized by plants during the evapotranspiration process. The Craig-Gordon model (e.g. Flanagan et al., 1991; Roden and Ehleringer, 1999) approximates the isotope
 255 processes in leaf water in δ terms (e.g. Barbour et al., 2004):

$$\delta_e \approx \delta_s + \epsilon^* + \epsilon_k + (\delta_a - \delta_s - \epsilon_k) \frac{e_a}{e_i}, \quad (\text{Equation 1})$$

Gelöscht: 2

Gelöscht: 2

Gelöscht: stable

Gelöscht: isotopic

Gelöscht: isotopic

Gelöscht: isotopic

where δ_e , δ_s and δ_a are the hydrogen and oxygen isotope compositions of leaf water at the evaporative sites, source water and atmospheric water vapor, respectively. The equilibrium enrichment (ϵ^*) is expressed as $(1 - 1/\alpha_{L/V}) \cdot 10^3$, where $\alpha_{L/V}$ is the equilibrium fractionation between liquid and vapor in ‰. The kinetic fractionation parameter (ϵ_k) describes the water vapor diffusion from intracellular air space through the stomata and the boundary layer into to the atmosphere, and e_a/e_i is the ratio of the
 260 atmospheric to intracellular vapor pressure.

Gelöscht: isotopic

In a $\delta^2\text{H}$ - $\delta^{18}\text{O}$ diagram, the isotope composition of the leaf water as well as the source water can be described as deuterium-excess (d) values by using the equation of Dansgaard (1964), with $d = \delta^2\text{H} - 8 \cdot \delta^{18}\text{O}$. This allows rewriting Eq. 1, in which hydrogen and oxygen isotopes have to be handled in separate equations, in one equation:

$$d_e \approx d_s + (\epsilon_2^* - 8 \cdot \epsilon_{18}^*) + (C_k^2 - 8 \cdot C_k^{18}) + [d_a - d_s - (C_k^2 - 8 \cdot C_k^{18})] \cdot \frac{e_a}{e_i}, \quad (\text{Equation 2})$$

where d_e , d_s and d_a are the deuterium-excess values of leaf water at evaporative sites, source water and atmospheric water vapor, respectively. The kinetic fractionation (ϵ_k) is typically related to stomata and boundary layer resistances to water flux (Farquhar et al., 1989). We used the kinetic enrichment factor (C_k) instead of ϵ_k to be close to paleo studies where direct measurements of plant physiology is not possible. The kinetic enrichment factor is derived from a more generalized form of the Craig-Gordon model for describing the kinetic isotope enrichment for ${}^2\text{H}$ and ${}^{18}\text{O}$ (C_k^2 and C_k^{18} , respectively)

(Craig and Gordon, 1965; Gat and Bowser, 1991). If the plant source water and the local atmospheric water vapor are in isotope equilibrium, the term $\delta_a - \delta_s$ in Eq. 1 can be approximated by $-\epsilon^*$. Thus, Eq. 2 can be reduced to:

$$d_e \approx d_s + (\epsilon_2^* - 8 \cdot \epsilon_{18}^* + C_k^2 - 8 \cdot C_k^{18}) \cdot \left(1 - \frac{e_a}{e_i}\right). \quad (\text{Equation 3})$$

285 The actual atmospheric vapor pressure (e_a) and the leaf vapor pressure (e_i) in kPa can be derived from Eqs. 4 and 5 by using T_{air} (Buck, 1981):

$$e_a = 0.61121 \cdot e^{[17.502 \cdot T_{\text{air}} / (T_{\text{air}} + 240.97)]} \cdot RH_{\text{air}} \quad (\text{Equation 4})$$

$$e_i = 0.61121 \cdot e^{[17.502 \cdot T_{\text{air}} / (T_{\text{air}} + 240.97)]}. \quad (\text{Equation 5})$$

When e_i is calculated as in Eq. 5, e_a/e_i represents RH_{air} (ranging between 0 and 1, representing 0 to 100% relative humidity). We are aware, that the Craig-Gordon model would require T_{leaf} values for calculating e_i values. However, the RH reconstruction methodological framework presented is attempted to paleo studies for which the T_{leaf} parameter is probably rather difficult to achieve.

However, as can be seen in Fig. 1, leaf temperature is very close to air temperature.

With rearranging Eq. 3, an equation is given to derive relative humidity values (Eq. 6):

$$RH_{\text{air}} \approx 1 - \frac{d_e - d_s}{(\epsilon_2^* - 8 \cdot \epsilon_{18}^* + C_k^2 - 8 \cdot C_k^{18})}. \quad (\text{Equation 6})$$

Equilibrium fractionation parameters (ϵ_2^* and ϵ_{18}^*) can be calculated from empirical equations of Horita and Wesolowski (1994) by using the climate chamber T_{air} values. The kinetic fractionation parameters (C_k^2 and C_k^{18}) for ^2H and ^{18}O , respectively, are set to 25.1 and 28.5‰ according to Merlivat (1978), who reported maximum values during the molecular diffusion process of water through a stagnant boundary layer. When using supplementary data of Cernusak et al., (2016), ϵ_k values of broadleaf trees and shrubs over broad climatic conditions can be calculated which are well in the range with the used C_k^2 and C_k^{18} values (23.9 ± 0.9 and $26.7\text{‰} \pm 1.0$ for ϵ_k^2 and ϵ_k^{18} , respectively).

If $\delta^2\text{H}_{\text{leaf-water}}$ and $\delta^{18}\text{O}_{\text{leaf-water}}$ can be reconstructed from the measured δ values of n -alkane and sugar biomarkers, this framework provides a powerful tool to establish relative humidity records from sedimentary archives (Hepp et al., 2017; Zech et al., 2013a).

305 To reconstruct the isotope composition of leaf water it is assumed that fractionation factors of -160‰ for ^2H of alkanes $n\text{-C}_{29}$ and $n\text{-C}_{31}$ (ϵ_{bio}^2 ; Sachse et al., 2012; Sessions et al., 1999), and $+27\text{‰}$ for ^{18}O of the hemicellulose-derived sugars arabinose and xylose ($\epsilon_{\text{bio}}^{18}$; Cernusak et al., 2003; Schmidt et al., 2001; Sternberg et al., 1986; Yakir and DeNiro, 1990) can be applied:

$$\text{alkane-based } \delta^2\text{H}_{\text{leaf-water}} = (\delta^2\text{H}_{n\text{-alkane}} - \epsilon_{\text{bio}}^2) / (1 + \epsilon_{\text{bio}}^2/1000) \quad (\text{Equation 7})$$

$$\text{sugar-based } \delta^{18}\text{O}_{\text{leaf-water}} = (\delta^{18}\text{O}_{\text{sugar}} - \epsilon_{\text{bio}}^{18}) / (1 + \epsilon_{\text{bio}}^{18}/1000). \quad (\text{Equation 8})$$

310 2.3.2 Isotope composition of plant source water

In a $\delta^2\text{H}$ - $\delta^{18}\text{O}$ diagram, the hydrogen and oxygen isotope composition of the plant source water ($\delta^2\text{H}_s$ and $\delta^{18}\text{O}_s$, respectively) can be reconstructed via the slope of the individual leaf water evapotranspiration lines (LEL's; Craig and Gordon, 1965; Gat and Bowser, 1991). The LEL slope (S_{LEL}) can be derived from Eq. 9:

Gelöscht: than the

Gelöscht: can

Gelöscht: derived

$$S_{LEL} \approx \frac{\epsilon_2^* + C_k^2 \cdot \left(1 - \frac{e_a}{e_i}\right)}{\epsilon_{18}^* + C_k^{18} \cdot \left(1 - \frac{e_a}{e_i}\right)} \approx \frac{\epsilon_2^* + C_k^2}{\epsilon_{18}^* + C_k^{18}}, \quad (\text{Equation 9})$$

where all **variables** are defined as in section 2.3.1. The $\delta^2\text{H}_s$ and $\delta^{18}\text{O}_s$ values can then be calculated for each leaf water data point via the intersect between the individual LEL's with the GMWL. The $\delta^2\text{H}_s$ and $\delta^{18}\text{O}_s$ model results can then be compared to the measured $\delta^2\text{H}_{\text{tank-water}}$ and $\delta^{18}\text{O}_{\text{tank-water}}$ values.

2.4 Modeling and isotope fractionation calculations

The d_e values are modeled using Eq. 3 and measured RH_{air} as input, which can be compared to the deuterium-excess via the measured $\delta^2\text{H}_{\text{leaf-water}}$ and $\delta^{18}\text{O}_{\text{leaf-water}}$ values. From the latter, also the RH_{air} can be derived from Eq. 6 and compared to the measured ones. In a next step, reconstructed (biomarker-based) deuterium-excess_{leaf-water} was used as input for Eq. 6 and compared to the measured RH_{air} values. **This model** represent a simplified approach because $\delta_a - \delta_s$ are approximated by $-\epsilon^*$ (see section 2.3). In all equations where δ_s and d_s are needed as input the measured $\delta^2\text{H}_{\text{tank-water}}$ and $\delta^{18}\text{O}_{\text{tank-water}}$ were used for calculations. **The equilibrium and kinetic fractionation factors** were set as described in section 2.3.

Model quality was overall assessed by calculating the coefficient of determination $[R^2 = 1 - \sum(\text{modeled} - \text{measured})^2 / \sum(\text{measured} - \text{measured mean})^2]$ and the root mean square error $\left[\text{RMSE} = \sqrt{\left(\frac{1}{n} \cdot \sum(\text{modeled} - \text{measured})^2\right)}\right]$. The R^2 is not equal to the r^2 , which provides here the fraction of variance explained by a linear regression between a dependent (y) and an explanatory variable $[r^2 = 1 - \sum(y - \text{fitted } y)^2 / \sum(y - \text{mean } y)^2]$ (R Core Team, 2015).

The fractionation between the measured leaf biomarkers and leaf water can be described by the following equations (e.g. Coplen, 2011):

$$\epsilon_{n\text{-alkane/leaf-water}} = (\delta^2\text{H}_{n\text{-alkane}} - \delta^2\text{H}_{\text{leaf-water}}) / (1 + \delta^2\text{H}_{\text{leaf-water}}/1000) \quad (\text{Equation 10})$$

$$\epsilon_{\text{sugar/leaf-water}} = (\delta^{18}\text{O}_{\text{sugar}} + \delta^{18}\text{O}_{\text{leaf-water}}) / (1 + \delta^{18}\text{O}_{\text{leaf-water}}/1000). \quad (\text{Equation 11})$$

In order to provide a 1σ range bracketing the modeled results and calculations they were additionally run with values generated by subtracting/adding the individual σ to the average.

All calculations and statistical analysis were realized in R (version 3.2.2; R Core Team, 2015).

Gelöscht: parameters

Gelöscht: calculated

Gelöscht: All

Gelöscht: s

Gelöscht: All

Gelöscht: other input parameters

3 Results and Discussion

3.1 Compound-specific isotope results

3.1.1 Leaf wax-derived *n*-alkanes

The investigated leaf material shows a dominance of the *n*-C₂₉ alkane homologue. Such a dominance of *n*-C₂₉ in *Brassica oleracea* and *Eucalyptus globulus* was also reported by Ali et al. (2005) and Herbin and Robins (1968). *Vicia faba* leaf samples additionally revealed a high abundance of *n*-C₃₁. This agrees with results from Maffei (1996) and enables a robust determination of compound-specific $\delta^2\text{H}$ values for C₂₉ and C₃₁, respectively. The $\delta^2\text{H}_{n\text{-alkane}}$ values of *Vicia faba* are therefore calculated as weighted mean. Figure 2 illustrates the $\delta^2\text{H}_{n\text{-alkane}}$ results along with isotope data for leaf, xylem and soil water (the latter were originally published in Mayr 2002). The $\delta^2\text{H}_{n\text{-alkane}}$ values ranged from -213 to -144‰ across all three plant species. As revealed by overlapping notches in the respective box plots, there is no statistically significant difference of the median between the three species (Fig. A1A; McGill et al., 1978). Figures A1A and 2, moreover shows that $\delta^2\text{H}_{n\text{-alkane}}$ values range largest for *Eucalyptus globulus*. However, the low number of samples per plant species prohibits a robust interpretation.




Fig. 2: $\delta^2\text{H}$ values of plant water (leaf water, xylem water and soil water) isotope compositions (in green, orange and brown, respectively) and the isotope composition of the investigated leaf wax *n*-alkanes *n*-C₂₉ and *n*-C₃₁ as dots and open squares, respectively.

3.1.2 Hemicellulose-derived sugars

The investigated leaf samples yielded substantially higher amounts of arabinose and xylose compared to fucose and rhamnose. This is in agreement with sugar patterns reported for higher plants (D'Souza et al., 2005; Hepp et al., 2016; Jia et al., 2008; Prietzel et al., 2013; Zech et al., 2012, 2014a) and hampers a robust data evaluation of fucose and rhamnose. Therefore, the $\delta^{18}\text{O}$ values were investigated for the pentoses arabinose and xylose and range from 30 to 47‰ and 30 to 50‰ (Fig. 3), respectively. Additionally, the isotope data for leaf, xylem and soil water are shown in the figure (originally published in Mayr 2002). No considerable difference in the $\delta^{18}\text{O}$ values of arabinose and xylose can be seen in the $\delta^{18}\text{O}$ data ($r^2 = 0.7$, $p < 0.001$, $n = 24$). This is in line with findings of Zech and

Gelöscht: leaf ...eaf wax-derived *n*-alkanes and hemicellulose-derived sugars¶


Gelöscht: *n*-alkanes... Such The A

Formatiert: Schriftart: Kursiv


Formatiert: Schriftart: Kursiv

Gelöscht: *n*-alkanes... This agrees with results from Maffei (1996) and enables a robust determination of compound-specific $\delta^2\text{H}$ values for C₂₉ and C₃₁, respectively....The $\delta^2\text{H}_{n\text{-alkane}}$ values of *Vicia faba* are therefore calculated as weighted mean. Figure 1B ... illustrates the $\delta^2\text{H}_{n\text{-alkane}}$ results along with isotopic ...isotope data for leaf, xylem and soil

A) climate chamber conditions



B) water and biomarker $\delta^2\text{H}$ and $\delta^{18}\text{O}$ values



Gelöscht:

[2] nach oben: Fig. 1: A: Climate chamber conditions (leaf temperature and relative humidity in green and air temperature and relative humidity in red). Error bars represent analytical standard deviation of the respective measurements (see section 2.2 and Mayr, 2002).

Gelöscht: B: P...lant water (leaf water, xylem water and soil water) isotope compositions (in green, orange and brown, respectively) and the isotope composition of the investigated leaf biomarkers (...eaf wax *n*-alkanes *n*-C₂₉ and *n*-C₃₁ as dots and open squares diamonds and triangles

Gelöscht: of the...investigated for the pentoses arabinose and xylose and range from 30 to 47‰ and 30 to 50‰ (Fig. 3), respectively. , ...dditionally, and are shown along with...he isotopic ...isotope data for leaf, xylem and soil water are shown in the figure (the latter were ...riginally published in Mayr 2002)1B... No considerable difference in the $\delta^{18}\text{O}$

Glaser (2009), Zech et al. (2012), Zech et al. (2013b) and Zech et al. (2014b) but contradicting with slightly more positive $\delta^{18}\text{O}_{\text{arabinose}}$ values compared to $\delta^{18}\text{O}_{\text{xylose}}$ values reported by Zech et al. (2013a) and Tuthorn et al. (2014). Zech et al. (2013a) and Tuthorn et al. (2014) argue that a biosynthetic fractionation could be the reason for that difference. This is based on the fact that arabinose is biosynthesized via an epimerase from xylose (Altermatt and Neish, 1956; Burget et al., 2003; Harper and Bar-Peled, 2002). However, the $\delta^{18}\text{O}$ values of arabinose and xylose can here combined as weighted mean (as $\delta^{18}\text{O}_{\text{sugar}}$ values) for further data interpretation. Overall, the $\delta^{18}\text{O}_{\text{sugar}}$ values are not significantly different between the three investigated plant species (Fig. A1B).




Fig. 3: $\delta^{18}\text{O}$ values of plant water (leaf water, xylem water and soil water) isotope compositions (in green, orange and brown, respectively) and the isotope composition of the investigated hemicellulose-derived sugars: arabinose and xylose as dots and open squares, respectively.

3.1.3 $\delta^2\text{H}$ - $\delta^{18}\text{O}$ diagram

A comparison of compound-specific isotope results of leaf hemicellulose-derived sugars and leaf wax-derived *n*-alkanes with leaf, xylem, soil and tank water (compare Figs. 2, 3 and Fig. 4) reveals that soil and xylem water plot close to the tank water. Only a slight heavy-isotope enrichment in the soil and xylem water was observed (c.f. the offset between soil and xylem water compared to the tank water). A larger evaporation effect of this water pools was inhibited by the high fresh air supply to the climate chambers (see section 2.1). Thus, the leaf water shows a clear heavy-isotope enrichment due to evapotranspiration. This enrichment strongly differs between the climate chambers, depending mainly on T and RH conditions. The biomarker results furthermore follow the leaf water with a certain offset, which is ϵ_{bic}

Gelöscht: Overall, the two sugars display very similar results (Fig. 1B; $r^2 = 0.7$, $p < 0.001$, $n = 24$).

Gelöscht: argue,

Gelöscht: ,

Gelöscht: The

Gelöscht: therefore

Gelöscht: be

Gelöscht: a

Gelöscht: The

Gelöscht: and circles

Gelöscht:) for the three plants *Eucalyptus globulus*, *Vicia faba* and *Brassica oleracea* grown in the climate chambers.

Gelöscht: ¶

Formatiert: Schriftart: Fett

Gelöscht: 4

Gelöscht: 1B

Gelöscht: and

Gelöscht: 2

Gelöscht: in Fig. 2

Gelöscht: (

Gelöscht:)





Fig. 4: $\delta^2\text{H}$ - $\delta^{18}\text{O}$ diagram illustrating the isotope composition of the biomarkers: $\delta^2\text{H}$ values of the leaf wax n -alkanes (C_{29} for *Eucalyptus globulus* and *Brassica oleracea* and weighted mean of C_{29} and C_{31} for *Vicia faba*) and $\delta^{18}\text{O}$ values of the hemicellulose-derived sugars arabinose and xylose (red crosses = *Brassica oleracea*, orange crosses = *Eucalyptus globulus*, blue crosses = *Vicia faba*). The measured isotope compositions of leaf water (green squares), xylem water (orange squares), soil water (brown squares), atmospheric water vapor (red squares) and the tank water used for irrigation (blue triangle), which plot very close to the global meteoric water line (GMWL). The deuterium-excess with respect to the GMWL is marked as d in ‰.



Gelöscht:

Gelöscht: 2

Gelöscht:

Gelöscht: black

3.2 Do n -alkane and sugar biomarkers reflect the isotope composition of leaf water?

3.2.1 $\delta^2\text{H}_{n\text{-alkane}}$ vs. $\delta^2\text{H}_{\text{leaf-water}}$

The $\delta^2\text{H}_{n\text{-alkane}}$ dataset reveals a significant correlation with $\delta^2\text{H}_{\text{leaf-water}}$ ($r^2 = 0.45$) using all plant species with $p < 0.001$ (Fig. 5A). A slope of 1.1 and an intercept of -152‰ furthermore characterize the linear relationship. Since it is well known that measured leaf water is not always equal to the specific water pool in which the n -alkanes were biosynthesized (e.g. Tipple et al., 2015), this could explain the observed rather low r^2 value. The correlation between the $\delta^2\text{H}_{n\text{-alkane}}$ and $\delta^2\text{H}_{\text{leaf-water}}$ presented here is still well in range with the literature. Feakins and Sessions (2010) presented n -alkane (C_{29} and C_{31}) and leaf water $\delta^2\text{H}$ data from typical plant species (excluding grasses) along a southern California aridity gradient, revealing that only $\delta^2\text{H}$ of $n\text{-C}_{29}$ is significantly correlated with leaf water ($r^2 = 0.24$, $p < 0.1$, $n = 16$; based on the associated supplementary data). Another field dataset from the temperate forest at Brown's Lake Bog, Ohio, USA revealed significant correlations between $\delta^2\text{H}$ of $n\text{-C}_{29}$ or $n\text{-C}_{31}$ and leaf water of *Prunus serotina*, *Acer saccharinum*, *Quercus rubra*, *Quercus alba*, and *Ulmus americana* ($r^2 = 0.49$, $p < 0.001$, $n = 38$; $r^2 = 0.59$, $p < 0.001$, $n = 29$; as derived from the supplement material of Freimuth et al., 2017). Data from a controlled climate chamber experiment using two tree species show a highly significant relationship between $\delta^2\text{H}$ of leaf wax n -alkanes and leaf water (with C_{31} of *Betula*

Gelöscht: ¶

Gelöscht: significant

Gelöscht: from

Gelöscht:

Formatiert: Tiefgestellt

Gelöscht:

Formatiert: Tiefgestellt

occidentalis and C₂₉ of *Populus fremontii*; $r^2 = 0.96$, $p < 0.001$, $n = 24$; derived from supplementary data of Tipple et al., 2015).





Fig. 5: (A) Scatterplot depicting the relationship between $\delta^2\text{H}_{n\text{-alkane}}$ and $\delta^2\text{H}_{\text{leaf-water}}$. Error bars of the δ values represent standard deviation of repeated measurements (see section 2.2 and Mayr, 2002). (B) Boxplot comprising the plant-specific fractionation $\epsilon_{n\text{-alkane/leaf-water}}$ according Eq. 10, showing median (thick black line), interquartile range (IQR) with upper (75%) and lower (25%) quartiles, lower and upper whiskers, which are restricted to $1.5 \cdot \text{IQR}$. Outside the $1.5 \cdot \text{IQR}$ space, the data points are marked with a dot. The notches are extend to $\pm 1.58 \cdot \text{IQR} / \sqrt{n}$, by convention and give a 95% confidence interval for the difference of two medians (McGill et al., 1978). *Brassica oleracea*, *Eucalyptus globulus* and *Vicia faba* samples are shown in red, orange and blue, respectively.

It seems that each plant type shows a different $\delta^2\text{H}_{n\text{-alkane}}$ to $\delta^2\text{H}_{\text{leaf-water}}$ relation, with the highest slope for *Vicia faba* and the lowest slope for *Brassica oleracea* (Fig. 5A). However, we argue that the number of replicates for each plant species is simply too low to interpret this finding robustly. In order to explore possible species-specific effects on the fractionation between the biomarkers and the leaf water in more detail, boxplots of the individual plant species of $\epsilon_{n\text{-alkane/leaf-water}}$ values are shown in Fig. 5B. Median $\epsilon_{n\text{-alkane/leaf-water}}$ values are -155‰ for *Brassica oleracea*, -164‰ for *Eucalyptus globulus* and -149‰ for *Vicia faba*, with an overall mean value of -156‰ (ranging from -133 to -192‰). The boxplots of $\epsilon_{n\text{-alkane/leaf-water}}$ reveal that the median of the three investigated plant species can be statistically not distinguished, due to overlapping notches. It should be noted that due to the low sample number from each species, the 95% confidence interval is larger than the interquartile range in some cases. However, it seems that at least small species-specific differences cannot be ruled out. Our $\epsilon_{n\text{-alkane/leaf-water}}$ values resemble well the data from a laboratory study (Kahmen et al., 2011), reporting a median value of -162‰ for $n\text{-C}_{25}$, $n\text{-C}_{27}$ and $n\text{-C}_{29}$ of *Populus trichocarpa*. Furthermore, they are well comparable to climate chamber data of *Betula occidentalis* ($n\text{-C}_{31}$) and *Populus fremontii* ($n\text{-C}_{29}$) from Tipple et al. (2015), reporting a median $\epsilon_{n\text{-alkane/leaf-water}}$ value of -155‰. In addition, field experiments reveal similar median values of -151‰ (for $n\text{-C}_{29}$) and -142‰ (for $n\text{-C}_{31}$) from typical plant species (excluding grasses) from southern California (Feakins and Sessions, 2010) and -144‰ (for $n\text{-C}_{29}$, of the species *Prunus serotina*, *Acer saccharinum*, *Quercus rubra*, *Quercus alba* and *Ulmus americana*) from the temperate forest at Brown's Lake Bog, Ohio, USA. The large range in $\epsilon_{\text{xylem-water/leaf-water}}$ values from our study (-192 to -133‰) is also obvious in the respective laboratory and field studies (-198 to -115‰,



Gelöscht:
Gelöscht: 3
Gelöscht: s
Gelöscht: s
Gelöscht: the compound-specific biomarker isotope composition and the respective leaf water values (A):
Gelöscht: vs.
Gelöscht: ;
Gelöscht: B: $\delta^{18}\text{O}_{\text{sugar}}$ vs. $\delta^{18}\text{O}_{\text{leaf-water}}$. *Brassica oleracea*, *Eucalyptus globulus* and *Vicia faba* samples are shown in purple, orange and black, respectively.
Gelöscht: .

Gelöscht: 4
Gelöscht: (Fig. 4A)

derived from n -C₂₉ and n -C₃₁ data from Feakins and Sessions, 2010; Kahmen et al., 2011a; Tipple et al., 2015; Freimuth et al., 2017).

The observed large range in $\epsilon_{n\text{-alkane/leaf-water}}$ and the rather low r^2 of the relationship between $\delta^2\text{H}_{n\text{-alkane}}$ and $\delta^2\text{H}_{\text{leaf-water}}$ could point to a more specific water pool being used rather than bulk leaf water during biosynthesis (Sachse et al., 2012; Schmidt et al., 2003). Furthermore, NADPH is acting also as hydrogen source during n -alkane biosynthesis, its $\delta^2\text{H}$ is clearly more negative than the biosynthetic water pool (Schmidt et al., 2003), further contributing to a weakening of the $\delta^2\text{H}_{n\text{-alkane}}$ to $\delta^2\text{H}_{\text{leaf-water}}$ correlation and enlarging the range of $\epsilon_{n\text{-alkane/leaf-water}}$. In more detail, alkane synthesis takes place by modifying/expanding fatty acids in the cytosol, while fatty acids are synthesized in the chloroplasts (Schmidt et al., 2003). Thus, the cytosol and chloroplast waters are two hydrogen sources. However hydrogen can additionally be added to the alkanes and fatty acids by NADPH which originates from different sources (photosynthesis and pentose phosphate cycle, Schmidt et al., 2003). It is therefore challenging to measure directly the water pool, from which the alkanes are biosynthesized (Tipple et al., 2015). Moreover, biosynthetic and metabolic pathways in general (Kahmen et al., 2013; Sessions et al., 1999; Zhang et al., 2009), the carbon and energy metabolism of plants more specifically (Cormier et al., 2018) and the number of carbon atoms of the n -alkane chains (Zhou et al., 2010) may have an influence on the fractionation. Our $\epsilon_{n\text{-alkane/leaf-water}}$ values correlate with T_{air} (Fig. A2A), whereas the correlation with RH_{air} (Fig. A2B) is not significant. This could point to a relationship between $\epsilon_{\text{xylem-water/leaf-water}}$ and plant physiological processes (affecting various plants differently). In summary, the fractionation between leaf water and n -alkanes is strongly influenced by the metabolic pathway of the n -alkane biosynthesis including direct hydrogen transfers, exchange reactions and NADPH as hydrogen source.

3.2.2 $\delta^{18}\text{O}_{\text{sugar}}$ vs. $\delta^{18}\text{O}_{\text{leaf-water}}$

A highly significant correlation is observed for the correlation between $\delta^{18}\text{O}_{\text{sugar}}$ and $\delta^{18}\text{O}_{\text{leaf-water}}$ ($r^2 = 0.84$, $p < 0.001$; Fig. 6A). The regression reveals a slope of 0.74 and an intercept of 30.7‰. The observed slope of the $\delta^{18}\text{O}_{\text{sugar}}$ vs. $\delta^{18}\text{O}_{\text{leaf-water}}$ relationship could serve as indicator for a leaf water (heavy-isotope enrichment) signal transfer damping of approximately 26%.

The theory behind the signal damping is adopted from the cellulose research (e.g. Barbour and Farquhar, 2000). Barbour and Farquhar (2000) related the extent of the signal damping to the proportion of unenriched source water, which contribute to the local synthesis water pool and to the proportion of exchangeable oxygen during cellulose synthesis. Our damping of 26% is within the range with values reported for cellulose synthesis in *Gossypium hirsutum* leaves (between 35 and 38%; Barbour and Farquhar, 2000), for *Eucalyptus globulus* leaf samples (38%; Cernusak et al., 2005) and for five C₃ and C₄ grasses (25%; Helliker and Ehleringer, 2002). Recently, Cheesman and Cernusak (2017) provided damping for leaf cellulose synthesis based on plant data grown under same conditions at Jerusalem Botanical Gardens published by Wang et al. (1998), ranging between 4 and 100% with a mean of 49%, revealing large variations among and between ecological groups (namely conifers, deciduous, evergreen and shrubs). A large range of signal damping associated with leaf cellulose was also reported by Song et al. (2014) for *Ricinus communis* grown under controlled conditions. A common disadvantage of the above-mentioned studies is the absence of direct measurements of the proportion of depleted source water contribution to the local synthesis water (as noticed by Liu et al., 2017), which largely contribute to the extent of the signal damping (Barbour and Farquhar, 2000). However, when transferring cellulose results to pentoses, such as hemicellulose-derived arabinose and xylose, it should be noted that they are biosynthesized via decarboxylation of the carbon at position six (C6) from

Gelöscht: The observed large range in $\epsilon_{n\text{-alkane/leaf-water}}$ and the rather low r^2 of the relationship between $\delta^2\text{H}_{n\text{-alkane}}$ and $\delta^2\text{H}_{\text{leaf-water}}$

Gelöscht: ¶

Gelöscht:

Gelöscht: Since it is well known that measured leaf water is not always equal to the specific water pool in which the n -alkanes are biosynthesized (e.g. Tipple et al., 2015), the correlation reveals a rather low r^2 (Fig. 3A). Furthermore, NADPH is acting also as hydrogen source during n -alkane biosynthesis, which is clearly more negative than the biosynthetic water pool (Schmidt et al., 2003), further contributing to a weakening of the $\delta^2\text{H}_{n\text{-alkane}}$ to $\delta^2\text{H}_{\text{leaf-water}}$ correlation. The correlation between the deuterium contents of leaf wax n -alkanes and leaf water presented here is still well in range with the literature. Feakins and Sessions (2010) presented n -alkane (C₂₉ and C₃₁) and leaf water $\delta^2\text{H}$ data from typical plant species (excluding grasses) along a southern California aridity gradient, revealing that only $\delta^2\text{H}$ of n -C₂₉ is significantly correlated with leaf water ($r^2 = 0.24$, $p < 0.1$, $n = 16$; based on the associated supplementary data). Another field dataset from the temperate forest at Brown's Lake Bog, Ohio, USA revealed significant correlations between $\delta^2\text{H}$ of n -C₂₉ and n -C₃₁ with leaf water of the species *Prunus serotina*, *Acer saccharinum*, *Quercus rubra*, *Quercus alba*, and *Ulmus americana* ($r^2 = 0.49$, $p < 0.001$, $n = 38$; $r^2 = 0.59$, $p < 0.001$, $n = 29$; as derived from the supplement material of Freimuth et al., 2017). Data from a controlled climate chamber experiment using two tree species show a highly significant relationship between leaf wax n -alkanes $\delta^2\text{H}$ and leaf water (with C₃₁ of *Betula occidentalis* and C₂₉ of *Populus fremontii*; $r^2 = 0.96$, $p < 0.001$, $n = 24$; derived from supplementary data of Tipple et al., 2015). It is conformed that leaf wax n -alkanes of dicotyledonous plants largely incorporate the leaf water isotope signal, while in monocotyledonous plants (e.g. grasses) the n -alkanes are more strongly affected by the source water due to the leaf growth at the intercalary meristem (Kahmen et al., 2013). The boxplots of $\epsilon_{n\text{-alkane/leaf-water}}$ reveal that the median of the three investigated plant species can be statistically not distinguished, due to overlapping notches (Fig. 4A). It should be noted that due to the low sample number from each species, the 95% confidence interval is larger than the interquartile range in some cases. However, it seems that at least small species-specific differences cannot be ruled out. Our $\epsilon_{n\text{-alkane/leaf-water}}$ values resemble well the data from a laboratory study (Kahmen et al., 2011), reporting a median

Gelöscht: in

Gelöscht: ¶

Gelöscht: 3B

Gelöscht: to

Gelöscht: (Fig. 3B)

Gelöscht: The here observed

Gelöscht: well in

Gelöscht: factors

Gelöscht: factors

Gelöscht: factor

glucose (Altermatt and Neish, 1956; Burget et al., 2003; Harper and Bar-Peled, 2002). Waterhouse et al. (2013) showed that the oxygen atoms at C6 position in glucose moieties, used for heterotrophic cellulose synthesis, are strongly affected by the exchange with local water (up to 80%). Based on these findings, it can be suggested that the influence of the non-enriched source water during the synthesis of leaf hemicelluloses is rather small.




Fig. 6: (A) Scatterplot depicting the relationship between $\delta^{18}\text{O}_{\text{sugar}}$ vs. $\delta^{18}\text{O}_{\text{leaf-water}}$. Error bars of the δ values represent standard deviation of repeated measurements (see section 2.2 and Mayr, 2002). (B) Boxplot comprising the plant-specific fractionation $\epsilon_{\text{sugar/leaf-water}}$ according to Eq. 11, showing median (thick black line), interquartile range (IQR) with upper (75%) and lower (25%) quartiles, lower and upper whiskers, which are restricted to $1.5 \cdot \text{IQR}$. Outside the $1.5 \cdot \text{IQR}$ space, the data points are marked with a dot. The notches are extend to $\pm 1.58 \cdot \text{IQR} / \sqrt{n}$, by convention and give a 95% confidence interval for the difference of two medians (McGill et al., 1978). *Brassica oleracea*, *Eucalyptus globulus* and *Vicia faba* samples are shown in red, orange and blue, respectively.

The oxygen isotope composition of leaf sugars (e.g. sucrose and cellulose) are strongly influenced by O exchange processes, which cause more positive $\delta^{18}\text{O}$ values of sugars compared to leaf water (Barbour et al., 2004; Cernusak et al., 2003). Therefore, median $\epsilon_{\text{sugar/leaf-water}}$ values of +27.0‰ for *Brassica oleracea*, +26.6‰ for *Eucalyptus globulus*, +26.8‰ for *Vicia faba* are observed (Fig. 6B). The overall $\epsilon_{\text{sugar/leaf-water}}$ average value of the three investigated species is +27.3‰ (ranging from +23.0 to +32.3‰). No systematic difference between the individual species is evident. Moreover, the $\epsilon_{\text{sugar/leaf-water}}$ values do not correlate significantly with T_{air} , but significantly with RH_{air} (Fig. A2C and D). A temperature dependence of the $\epsilon_{\text{sugar/leaf-water}}$ is not supported by this experiment, in contrast to results from Sternberg and Ellsworth (2011), where a temperature effect on oxygen fractionation during heterotrophic cellulose biosynthesis is observed. The here observed fractionation between hemicellulose-derived sugars and leaf water, with regard to $\epsilon_{\text{sugar/leaf-water}}$ values, is well in range with values reported for sucrose (exported from photosynthesizing leaves) and leaf water, which was shown to be +27‰ (Cernusak et al., 2003). Also the cellulose biosynthesis is associated with an heavy-isotope enrichment of around +27‰ compared to the synthesis water as shown in growth experiments (Sternberg et al., 1986; Yakir and DeNiro, 1990). The relatively uniform fractionation is explained via the isotope exchange between the carbonyl oxygens of the organic molecules and the surrounding water (cf. Schmidt et al., 2001). This equilibrium fractionation effect was indeed described earlier by the reversible hydration reaction of acetone in water by Sternberg and DeNiro (1983) to be +28, +28

Gelöscht: s

Gelöscht: between the biomarkers and the leaf water (A: $\epsilon_{\text{n-alkane/leaf-water}}$ according Eq. 10; B:

Gelöscht:).

Gelöscht: *Brassica oleracea*, *Eucalyptus globulus* and *Vicia faba* samples are shown in purple, orange and black, respectively. Boxplots

Gelöscht: ic

Formatiert: Hochgestellt

Gelöscht: Median

Gelöscht: shown

Gelöscht: in

Gelöscht: 4B

Gelöscht: In both plots, n

Gelöscht: seems to be observable

Gelöscht: ¶

T

and +26‰ at 15, 25 and 35°C, respectively. However, the observed range of approximately 9‰ (Fig. 6B) could indicate that partially more than the oxygen equilibrium fractionation between organic molecules and medium water have to be considered. Presumably, isotope as well as sucrose synthesis gradients within the leaf have to be taken into account when interpreting leaf sugar oxygen isotope compositions and their correlation to leaf water (Lehmann et al., 2017). Lehmann et al. (2017) reported on a fractionation between sucrose and leaf water of +33.1‰. Based on this they proposed a conceptual scheme how such gradients can lead to discrepancies between the isotope composition of the bulk leaf water and the synthesis water, while the latter is incorporated into the carbohydrates, and thus fractionation determination based on bulk leaf water can exceed the common average of +27‰. Also Mayr et al. (2015) found a fractionation between aquatic cellulose $\delta^{18}\text{O}$ and lake water larger than this value of around +29‰.

3.2.3 Inter-species variability

The differences observed in $\epsilon_{n\text{-alkane/leaf-water}}$ (Fig. 5B) and $\epsilon_{\text{sugar/leaf-water}}$ (Fig. 6B) between the taxa might be explained by different leaf sizes and geometries. Please note that these differences are statistically not significant. We do not know whether this is caused by the rather low sample number of our study ($n = 24$) or not. Still, the yielded mean biosynthetic fractionation factors over all species are -156‰ and +27‰ for $\delta^2\text{H}_{n\text{-alkanes}}$ and $\delta^{18}\text{O}_{\text{sugars}}$, respectively. This is well in agreement with data from the literature (usually -160‰ and +27‰, respectively).

When reviewing Fig. 2 and 3, we observe differences between experiment 4 and 8 for *Eucalyptus globulus*, which is mainly related to different sampling daytime of leaves between the two experiments. Because of the diurnal course of light, relative humidity and temperature which was simulated in each chamber, steady-state conditions prevailed only during 11 am and 4 pm (see section 2.1). Due to time restrictions, the leaf samples of *Eucalyptus globulus* from experiments 1 to 4 were sampled after that daytime explaining the deviations between experiment 4 and 8 in leaf water isotope composition only. This, however, does not influence isotope values of tissue samples which represent the integrated signal over the entire growing period and does not explain the observed differences in $\delta^2\text{H}_{n\text{-alkane}}$. It is also obvious that the $\delta^2\text{H}_{n\text{-alkane}}$ values show this difference for all plant types whereas $\delta^{18}\text{O}_{\text{sugar}}$, $\delta^{18}\text{O}_{\text{leaf-water}}$ do not. The differences in $\delta^2\text{H}_{n\text{-alkane}}$ are most likely explainable via the fractionation occurring during biosynthesis of *n*-alkanes, which is depending not only on leaf water isotope composition but also on plant physiological factors (e.g. water pressure deficit between air and leaf, transpiration rate, assimilation rate; cf. discussion in 3.2.1).

3.2.4 Implications for paleoclimatic reconstructions

The damping of the leaf water oxygen isotope signal caused by the exchange between sugars and water is an important issue, especially in stem, trunk and root tissues. While these tissues hardly produce *n*-alkanes, (hemi-)celluloses and sugars extracted from such tissues do not show the full leaf water heavy-isotope-enrichment because of a partial oxygen exchange with non-enriched stem water (e.g. Zech et al., 2014a). This has to be kept in mind for paleoclimatic reconstructions, especially for the sugars of grasses, which do not record the full leaf water enrichment (e.g. Helliker and Ehleringer, 2002b). In the case of grasses, this signal damping affects *n*-alkanes, too. However, such uncertainties can be included in the RH reconstruction via assumptions and sensitivity analysis of the used model, thus allowing quantifying uncertainties of reconstructed RH records as shown by Hepp et al. (2019).

Gelöscht: 4

Gelöscht: isotopic

Gelöscht: isotopic

Gelöscht: isotopic

Gelöscht: As a

Gelöscht: In experiments 1 to 4,

Formatiert: Schriftart: Kursiv

Gelöscht: due to time restrictions

Gelöscht: became

Gelöscht: show this as well as $\delta^2\text{H}_{\text{leaf-water}}$ for *Vicia faba* and *Brassica oleracea*.

Gelöscht: (as stated in the first reply)

Gelöscht: ic

Gelöscht: application

Gelöscht: O-

Gelöscht: causing the fractionation

Gelöscht: and thus also the signal damping

Gelöscht: signal

Gelöscht: applications

Gelöscht: herbaceous plants like

Gelöscht: (

Gelöscht: .,

Gelöscht: ¶

¶

3.3 Fractionation factors between biomarkers and leaf water

Gelöscht: In order to explore possible species-specific effects on the fractionation between the biomarkers and the leaf water, boxplots of the individual plant species of $\epsilon_{n\text{-alkane/leaf-water}}$ and $\epsilon_{\text{sugar/leaf-water}}$ values are shown in Fig. 4. Median $\epsilon_{n\text{-alkane/leaf-water}}$ values are -155‰ for *Brassica oleracea*, -164‰ for *Eucalyptus globulus* and -149‰ for *Vicia faba* (Fig. 4A), with an overall mean value of -156‰ (ranging from -133 to -192‰). Median $\epsilon_{\text{sugar/leaf-water}}$ values of +27.0‰ for *Brassica oleracea*, +26.6‰ for *Eucalyptus globulus*, +26.8‰ for *Vicia faba* are shown in Fig. 4B. The overall $\epsilon_{\text{sugar/leaf-water}}$ average value of the three investigated species is +27.3‰ (ranging from +23.0 to +32.3‰). In both plots, no systematic difference between the individual species seems to be observable. ¶

Gelöscht: ¶

3.3 Strong control of relative humidity over deuterium-excess of leaf water

3.3.1 Calculating relative humidity based on measured leaf water

The correlations between leaf water-based and measured RH_{air} and modeled d_e and measured deuterium-excess_{leaf-water} are illustrated in Fig. 7A and B, respectively. Since the climate chambers had a high fresh air supply (see section 2.1) the variability of the leaf water deuterium-excess is predominantly the result of leaf water heavy-isotope enrichment and thus strongly driven by relative humidity.




Fig. 7: Scatterplots illustrating the correlation between (A) modeled (leaf water-based) and measured air relative humidity (RH_{air}) and modeled (T_{air} -based) vs. measured leaf water deuterium-excess. Black lines indicate the 1:1 relationship. R^2 and RMSE are calculated as described in section 2.4, while the RMSE values have the dimensions of the respective variables. Error bars for the measured RH values represent analytical standard deviations (see Mayr, 2002). See section 2.4 for the uncertainties of the calculated and modeled results.

Evidence for the strong control of relative humidity on deuterium-excess of leaf water comes from multivariate regression analysis between the measured deuterium-excess_{leaf-water} values versus RH_{air} and T_{air} . The results reveal that the deuterium-excess_{leaf-water} significantly correlates with RH_{air} of the climate chambers ($p < 0.001$), with an r^2 of 0.92. The strong control of RH on the deuterium-excess of leaf water is furthermore supported by the significant correlations between calculated versus measured RH_{air} values (Fig. 7A). This is in line with the strong correlation between modeled d_e based on T_{air} and measured deuterium-excess_{leaf-water} values (Fig. 7B).

The modeled d_e values show a high agreement with measured deuterium-excess of leaf water without being too positive. This could be expected from the literature, because the output of the Craig-Gordon-based leaf water enrichment model (e.g. Allison et al., 1985; Barbour et al., 2004; Cernusak et al., 2016; section 2.3) reflects leaf water at the evaporative sites, which should be more positive than the measured bulk leaf water. Especially under low relative humidity conditions, the discrepancy between Craig-Gordon model results and the measured values is shown to be more pronounced, associated with higher transpiration fluxes and higher isotope heterogeneity within the leaf water due to a non-uniform closure of the stomata (Flanagan et al., 1991; Santrucek et al., 2007). An overestimation of the Craig-Gordon models can hardly be observed here (Fig. 7B). However, based on the accepted leaf water enrichment theory (e.g. Cernusak et al., 2016), higher transpiration rates (e.g. under low

Gelöscht: 4


Gelöscht: 5A

Gelöscht: for

Gelöscht: the

Gelöscht:

Gelöscht: driven



Gelöscht:

Gelöscht: 5

Gelöscht: ,

Gelöscht: (T_{air} -based)

Gelöscht: 5A

Gelöscht: 5B

Gelöscht: Overall, t

Gelöscht: ,

Gelöscht: which

Gelöscht: can be

Gelöscht: . This is

Gelöscht: however,

Gelöscht: 5B

humidity conditions) should still lead to a larger discrepancy between Craig-Gordon modelled and measured leaf water, because the back diffusion of enriched leaf water from the evaporative sites should get lower the higher the transpiration flux is.

It should be noted, that there is the possibility to build up a more detailed model based on Eq. 2. With this, a model is given to derive d_e values with the usage of d_a and d_s , which can be compared to the measured deuterium-excess_{leaf-water} values. However, when modeling d_e without the simplification $\delta_a - \delta_s \approx -\epsilon^*$ the R^2 results to 0.86 and RMSE equals 13.07‰ compared to the presented 0.88 and 12.31‰. Furthermore, in Eq. 5 T_{air} can be replaced by T_{leaf} . With this, Eq. 2 results to d_e values based on leaf temperature. This would take into account that the Craig-Gordon model requires the temperature of the evaporating surface rather than the air temperature for e_i . However, with this model extension the R^2 and the RMSE results to 0.55 and 23.54%, respectively. By rearranging Eq. 2, RH values can be modeled which can be compared to RH_{air} as well as RH_{leaf} values (e_a/e_i multiplied by 100 with T_{leaf}). The respective model characteristics are again lower for the RH_{leaf} case ($R^2 = 0.27$ and $RMSE = 11.84\%$) than for the RH_{air} comparison ($R^2 = 0.81$ and $RMSE = 6.56\%$). Still Eq. 6 provides better results, as presented in this paragraph ($R^2 = 0.84$ and $RMSE = 6.04\%$). This discussion is in line the differences between T_{leaf} vs. T_{air} and RH_{leaf} vs. RH_{air} conditions in the climate chambers. They reveal the same trends and magnitude, but T_{leaf} is consequently higher than T_{air} along with higher RH_{leaf} values compared to RH_{air} (Fig. 1; Mayr, 2002). Summarized we therefore argue, that the model presented in Eq. 6 (including the simplifications of $\delta_a - \delta_s \approx -\epsilon^*$ and using T_{air} in Eq. 5) is able to reconstruct RH_{air} values based on $\delta^2H_{leaf-water}$ and $\delta^{18}O_{leaf-water}$ values.

3.3.2 Calculating relative humidity based on reconstructed leaf water

In order to test the proposed paleohygrometer approach, the alkane and sugar-based (reconstructed) leaf-water values were used to calculate RH_{air} . The measured RH_{air} is well reflected by the biomarker-based air relative air humidity values ($R^2 = 0.54$; Fig. 8). Overall, a lower coefficient of determination of the biomarker-based model results compared to the leaf water-based reconstructions (compare black with grey data points in Fig. 8) is observed. This can be attributed to the uncertainties in reconstructed leaf water using $\delta^2H_{n-alkane}$ and $\delta^{18}O_{sugar}$ datasets as discussed in sections 3.2.1 and 3.2.2. Briefly, the limitations regarding δ^2H arose from the rather weak relationship between the δ^2H of the n -alkanes and the leaf water, probably linked with the large range in $\epsilon^2_{n-alkane/leaf-water}$ (Fig. 5B). The applied equation for reconstructing $\delta^2H_{leaf-water}$ by using $\delta^2H_{n-alkane}$ using a constant biosynthetic fractionation of -160‰ (Eq. 10) was considered to be suitable (Sachse et al., 2012; Sessions et al., 1999), but contributes also some uncertainty to the final relative humidity reconstruction. With regard to $\delta^{18}O$, the relatively large variations in $\epsilon_{sugar/leaf-water}$ of 9‰ have to be considered (Fig. 6B), because in the $\delta^{18}O_{leaf-water}$ reconstructions a fixed value of +27‰ is used (Eq. 11). Such a uniform biosynthetic fractionation is just an approximation of real probably more variable values. The fact that the measured bulk leaf water does not capture the isotope value of the water in which the biomarkers are biosynthesized, could thus partly explain the weaker relationship between measured and calculated RH_{air} when biomarkers are used to derive leaf water. Especially the underestimation of the biomarker-based RH_{air} values under the 68% relative humidity conditions, as well as the large range in reconstructed RH_{air} values for the 48, 49, 50% RH_{air} chambers can be attributed to the leaf water reconstruction uncertainties. It should be mentioned that using Eqs. 7 and 8 to calculate leaf water isotope composition based on the biomarkers via a biosynthetic fractionation values implies that the fractionation process in principle can be treated as single process with a unique source. While this approximation can be questioned (see discussion in section 3.2), the overall correlation between

Gelöscht: with

Gelöscht: for the e_i parameter

Formatiert: Tiefgestellt

Gelöscht: A

[1] verschoben

Gelöscht: Finally

Gelöscht:

Gelöscht: captured

Gelöscht: 7

Gelöscht: 7

Gelöscht: reconstructed

Gelöscht: 3.2

Gelöscht: T

Formatiert: Schriftart: Nicht Kursiv, Nicht Hochgestellt/
Tiefgestellt

Gelöscht: deuterium

Gelöscht: the fractionation between n -alkanes and leaf water (

Gelöscht:)

Gelöscht: to

Gelöscht: ed

Gelöscht: and

Gelöscht: was

Gelöscht: introduces

Gelöscht: for

Gelöscht: oxygen

Gelöscht: 4B

Gelöscht: which may not always be fulfilled, as shown in the literature (e.g. Sternberg and Ellsworth, 2011; Lehmann et al., 2017)

Gelöscht:

Gelöscht: full

Gelöscht: variability of the water, which

Gelöscht: lead to a certain difference to the water important for the biosynthesis of the n -alkanes and sugars

Gelöscht: .

biomarker-based and measured RH_{air} highlights the potential of the approach (Hepp et al., 2017; Tuthorn et al., 2015; Zech et al., 2013a), also for future paleoclimate reconstructions.





Fig. 8: Scatterplot depicting the relationship between modeled (biomarker-based) and measured relative air humidity (RH_{air}). The black line indicates the 1:1 relationship. R^2 and RMSE was calculated as described in section 2.4, while the RMSE values have the dimensions of the respective variables. Error bars for the measured values represent analytical standard deviations (see Mayr, 2002). For uncertainty calculation of the results, see section 2.4. In addition, the leaf water-based relative air humidity results are shown in light grey for comparison.

3.4 Coupling $\delta^2H_{n\text{-alkane}}$ and $\delta^{18}O_{sugar}$ for $\delta^2H_{source\text{-water}}$ and $\delta^{18}O_{source\text{-water}}$ calculation

The second advantage of the proposed coupled $\delta^2H_{n\text{-alkane}}\text{-}\delta^{18}O_{sugar}$ approach is a more robust reconstruction of the isotope composition of the source water, which can often be directly linked to the local precipitation signal (Hepp et al., 2015, 2017; Tuthorn et al., 2015; Zech et al., 2013a). Therefore we calculated the source water isotope compositions via the slopes of the LEL's and the GMWL. In order to show the difference to approaches using either $\delta^2H_{n\text{-alkane}}$ or $\delta^{18}O_{sugar}$ for source water reconstruction, figure 9 depicts (i) the leaf water isotope composition reconstructed from $\delta^2H_{n\text{-alkane}}$ and $\delta^{18}O_{sugar}$ results using the biosynthetic fractionation factors, (ii) the measured $\delta^2H_{leaf\text{-water}}$ and $\delta^{18}O_{leaf\text{-water}}$ values, (iii) the source water isotope composition based on reconstructed leaf water, (iv) the source water isotope composition based on measured leaf water along with the (v) tank water isotope composition.

For δ^2H , neither the range nor the median of the $\delta^2H_{leaf\text{-water}}$ are well captured by the alkane-based leaf water values. However, the overlapping notches do not support a statistical difference in the median values (Fig. 9A). The median is on average 13‰ more positive than the measured $\delta^2H_{tank\text{-water}}$. A higher agreement between measured and modeled values is observed from leaf water-based δ^2H_s compared to $\delta^2H_{tank\text{-water}}$. The average offset is reduced to 2‰ and the range is reduced by approximately 70‰, compared to the biomarker-based reconstruction.

Gelöscht: -applications



Gelöscht:

Gelöscht: 7

Gelöscht: (modeled)

Gelöscht: air

Formatiert: Tiefgestellt

Gelöscht: modeled properties

Gelöscht: air

Gelöscht: ¶

Gelöscht: 5

Gelöscht: – potential and limitations

Gelöscht: One of the

Gelöscht: s

Gelöscht: Source

Gelöscht: were calculated

Gelöscht: The ff

Gelöscht: shows

Gelöscht: clearly that the n -alkane and sugar biomarkers reflect leaf water rather than

Gelöscht: used for irrigation, when the biomarker results are corrected for the individual biosynthetic fractionation

Gelöscht: 6

Gelöscht: s are still

Gelöscht: ¶




Fig. 9: Boxplots showing the measured leaf water in comparison to the biomarker-based leaf water, source water calculated with biomarker-based leaf water values and source water based on measured leaf water (A: $\delta^2\text{H}_{\text{leaf-water}}$, B: $\delta^{18}\text{O}_{\text{leaf-water}}$). Tank water is given as a black solid line in both plots. Source water isotope compositions were calculated via the slopes of the LEL's (either with biomarker-based or measured leaf water values) and the GMWL. Boxplots show median (thick black line), interquartile range (IQR) with upper (75%) and lower (25%) quartiles, lower and upper whiskers, which are restricted to $1.5 \cdot \text{IQR}$. Outside the $1.5 \cdot \text{IQR}$ space, the data points are marked with a dot. The notches extend to $\pm 1.58 \cdot \text{IQR} / \sqrt{n}$, by convention and give a 95% confidence interval for the difference of two medians (McGill et al., 1978).

For $\delta^{18}\text{O}$, the sugar-based leaf water values are in agreement with the measured ones with regard to the median values, as supported by the largely overlapping notches (Fig. 9B). The range of the reconstructed leaf water is about 6‰ lower than for the measured $\delta^{18}\text{O}_{\text{leaf-water}}$ dataset. All reconstructed $\delta^{18}\text{O}_s$ values, regardless whether they are biomarker- or leaf water-based, are comparable to the measured $\delta^{18}\text{O}_{\text{tank-water}}$. While the biomarker-based datasets depict an average offset of 2‰, the leaf water-based values only differ by 0.3‰ from the tank water $\delta^{18}\text{O}$ values, referring to the median.

The overall larger range of modeled $\delta^2\text{H}_s$ and $\delta^{18}\text{O}_s$ compared to measured $\delta^2\text{H}_{\text{tank-water}}$ and $\delta^{18}\text{O}_{\text{tank-water}}$ can be related to uncertainties in S_{LEL} modeling (see equations in section 2.3.2). Bariac et al. (1994) found no agreement between the intersects of modeled LEL's with the GMWL and the plant source water. Allison et al. (1985) explained such results with changing environmental conditions, leading to various LEL's with a locus line not necessarily passing the $\delta^2\text{H}_s$ and $\delta^{18}\text{O}_s$ data point, in a system that approaches rapidly new steady-state conditions. However, when a larger isotope range of source water is used in the studies, leaf water seems to be a good tracer for source water changes when a Craig-Gordon model is applied (Benettin et al., 2021; Bush et al., 2017).

Gelöscht:

Gelöscht: 6

Gelöscht: tank water,

Gelöscht: marked

Gelöscht: with

Gelöscht: 6

Gelöscht: in the order of

Gelöscht: smaller

Gelöscht: mentioned that they

Gelöscht: ic

Feldfunktion geändert

[1] nach oben: Finally, the alkane and sugar-based leaf water values were used to reconstruct RH_{air} . The measured RH_{air} is well captured by the biomarker-based air relative humidity values ($R^2 = 0.54$; Fig. 7). Overall, a lower coefficient of determination of the biomarker-based model results compared to the leaf water-based reconstructions (compare black with grey data points in Fig. 7) is observed. This can be attributed to the uncertainties in leaf water reconstructed using $\delta^2\text{H}_{n\text{-alkane}}$ and $\delta^{18}\text{O}_{\text{sugar}}$ datasets as discussed in section 3.2. The limitations regarding deuterium arose from the rather weak relationship between the $\delta^2\text{H}$ of the n -alkanes and the leaf water, probably linked with the large range in the fractionation between n -alkanes and leaf water ($\epsilon^2_{n\text{-alkane/leaf-water}}$). The applied equation to reconstructed $\delta^2\text{H}_{\text{leaf-water}}$ by using $\delta^2\text{H}_{n\text{-alkane}}$ and a constant biosynthetic fractionation of -160‰ (Eq. 10) was considered to be suitable (Sachse et al., 2012; Sessions et al., 1999), but introduces also some uncertainty for the final relative humidity reconstruction. With regard to oxygen, the relatively large variations in $\epsilon_{\text{sugar/leaf-water}}$ of 9‰ have to be considered (Fig. 4B), because in the $\delta^{18}\text{O}_{\text{leaf-water}}$ reconstructions a fixed value of +27‰ is used (Eq. 11). Such a uniform biosynthetic fractionation is an approximation which may not always be fulfilled, as shown in the literature (e.g. Sternberg and Ellsworth, 2011; Lehmann et al., 2017).

4 Conclusions

The climate chamber results and discussion suggest that leaf wax-derived n -alkane and hemicellulose-derived sugar biomarkers are valuable $\delta^2\text{H}_{\text{leaf-water}}$ and $\delta^{18}\text{O}_{\text{leaf-water}}$ recorders, respectively. The coupling of $\delta^2\text{H}_{n\text{-alkane}}$ and $\delta^{18}\text{O}_{\text{sugar}}$ results allows moreover a more robust RH_{air} reconstruction of the chambers in which the plants were grown, by using a simplified Craig-Gordon equation, compared to single isotopes. With regard to the research questions, we conclude the following:

- (i) $n\text{-C}_{29}$ predominated and occurred at abundances suitable for compound-specific $\delta^2\text{H}$ measurements in the leaf samples from all investigated species (*Eucalyptus globulus*, *Vicia faba* var. *minor* and *Brassica oleracea* var. *medullosa*). For *Vicia faba*, additionally $n\text{-C}_{31}$ could be evaluated robustly. $\delta^{18}\text{O}_{\text{sugar}}$ values could be obtained for the hemicellulose-derived monosaccharides arabinose and xylose.
- (ii) Both the $\delta^2\text{H}_{n\text{-alkane}}$ and $\delta^{18}\text{O}_{\text{sugar}}$ values yielded highly significant correlations with $\delta^2\text{H}_{\text{leaf-water}}$ and $\delta^{18}\text{O}_{\text{leaf-water}}$ ($r^2 = 0.45$ and 0.85 , respectively; $p < 0.001$, $n = 24$). Mean fractionation factors between biomarkers and leaf water were found to be -156‰ (ranging from -133 to -192‰) for $\epsilon_{n\text{-alkane/leaf-water}}$ and $+27.3\text{‰}$ (ranging from $+23.0$ to $+32.3\text{‰}$) for $\epsilon_{\text{sugar/leaf-water}}$.
- (iii) RH_{air} can be derived robustly by using the measured leaf water isotope composition ($\delta^2\text{H}_{\text{leaf-water}}$ and $\delta^{18}\text{O}_{\text{leaf-water}}$) and a rearranged Craig-Gordon model, ($R^2 = 0.84$; $p < 0.001$; $\text{RMSE} = 6\%$).
- (iv) Biomarker-based and measured RH_{air} correlation with R^2 of 0.54 ($p < 0.001$) and RMSE of 10% highlights the great potential of the coupled $\delta^2\text{H}_{n\text{-alkane}}$ - $\delta^{18}\text{O}_{\text{sugar}}$ paleohygrometer approach for reliable relative humidity reconstructions. Uncertainties with regard to relative humidity reconstructions via biomarker-based leaf water isotope composition arose from leaf water reconstructions and model uncertainties, as shown in the conclusions ii) and iii).
- (v) The coupled $\delta^2\text{H}$ - $\delta^{18}\text{O}$ approach enables a better back calculation of the plant source water compared to single isotopes. Reconstructed source water isotope composition ($\delta^2\text{H}_s$, $\delta^{18}\text{O}_s$) is in range with the measured tank water ($\delta^2\text{H}_{\text{tank-water}}$, $\delta^{18}\text{O}_{\text{tank-water}}$). However, modeled $\delta^2\text{H}_s$ and $\delta^{18}\text{O}_s$ showed a clearly larger range compared to $\delta^2\text{H}_{\text{tank-water}}$ and $\delta^{18}\text{O}_{\text{tank-water}}$. The uncertainties for source water determination are thus considerably higher compared to the relative humidity reconstructions.

Acknowledgements

We would like to thank M. Bliedtner and J. Zech (both University of Bern) for help during lipid biomarker and $\delta^2\text{H}_{n\text{-alkane}}$ analysis. We thank M. Benesch (Martin-Luther-University Halle-Wittenberg) and M. Schaarschmidt (University of Bayreuth) for laboratory assistance during sugar biomarker and $\delta^{18}\text{O}_{\text{sugar}}$ analysis. The research was partly funded by the Swiss National Science Foundation (PP00P2 150590). We also acknowledge N. Orlowski (University of Freiburg), M. M. Lehmann (Swiss Federal Institute WSL, Birmensdorf) and L. Wüthrich (University of Bern) for helpful discussions. We are very grateful for the constructive discussion on an earlier version of this manuscript as preprint at Biogeosciences Discussions (<https://doi.org/10.5194/bg-2019-427>). Involvement of K. Rozanski was supported by Polish Ministry of Science and Higher Education, project no. 16.16.220.842 B02. J. Hepp greatly acknowledges the support given by the German Federal Environmental Foundation. The experiment carried out by C. Mayr was gratefully supported by the HGF-project "Natural climate variations from 10,000 years to the present" (project no. 01SF9813). The experiments were possible

Formatiert: Deutsch (Deutschland)

Formatiert: Deutsch (Deutschland)

Formatiert: Deutsch (Deutschland)

Gelöscht: as follows

Gelöscht: Alkanes with the chain-length

Gelöscht: Reconstructed source water isotope composition ($\delta^2\text{H}_s$, $\delta^{18}\text{O}_s$) is in range with the measured tank water ($\delta^2\text{H}_{\text{tank-water}}$, $\delta^{18}\text{O}_{\text{tank-water}}$). However, modeled $\delta^2\text{H}_s$ and $\delta^{18}\text{O}_s$ show a clearly larger range compared to $\delta^2\text{H}_{\text{tank-water}}$ and $\delta^{18}\text{O}_{\text{tank-water}}$. The uncertainties for source water determination are thus considerably higher compared to the relative humidity reconstructions. Still, t

Gelöscht: Uncertainties with regard to relative humidity reconstructions via biomarker-based leaf water isotope composition arose from leaf water reconstructions and model uncertainties, as shown in the conclusions ii) and iii). Overall, the biomarker-based and measured RH_{air} correlation with R^2 of 0.54 ($p < 0.001$) and RMSE of 10% highlights the great potential of the coupled $\delta^2\text{H}_{n\text{-alkane}}$ - $\delta^{18}\text{O}_{\text{sugar}}$ paleohygrometer approach for reliable relative humidity reconstructions.

due to the assistance of J.B. Winkler, H. Lowag, D. Strube, A. Kruse, D. Arthofer, H. Seidlitz, D. Schneider, H. D. Payer, and other members of the Helmholtz Zentrum München.

Author contributions

J. Hepp, M. Zech and C. Mayr wrote the paper; C. Mayr was responsible for the climate chamber experiment together with W. Stichler and provided the leaf samples and the data; M. Zech and R. Zech were responsible for compound-specific isotope analysis on the biomarkers; J. Hepp, M. Tuthorn and I. K. Schäfer did laboratory work and data evaluation of the biomarker compound-specific isotope analysis; B. Glaser, D. Juchelka, K. Rozanski and all co-authors contributed to the discussion and commented on the manuscript.

Appendix




Fig. A1: Boxplots comprising the plant-specific $\delta^2\text{H}_{n\text{-alkane}}$ (A) and $\delta^{18}\text{O}_{\text{sugar}}$ values (B). *Brassica oleracea*, *Eucalyptus globulus* and *Vicia faba* samples are shown in red, orange and blue, respectively. Boxplots show median (thick black line), interquartile range (IQR) with upper (75%) and lower (25%) quartiles, lower and upper whiskers, which are restricted to 1.5 · IQR. Outside the 1.5 · IQR space, the data points are marked with a dot. The notches extend to $\pm 1.58 \cdot \text{IQR} / \sqrt{n}$, by convention and give a 95% confidence interval for the difference of two medians (McGill et al., 1978).

Gelöscht: and

Gelöscht: ¶

Gelöscht:

Gelöscht: purple

Gelöscht: black




Fig. A2: Scatterplots of the fractionation between the biomarkers and leaf water vs. air temperature, air relative humidity (A and B: $\epsilon_{n\text{-alkane/leaf-water}}$ according Eq. 10; C and D $\epsilon_{\text{sugar/leaf-water}}$ according Eq. 11). *Brassica oleracea*, *Eucalyptus globulus* and *Vicia faba* samples are shown in red, orange and blue, respectively. Error bars for the measured values represent analytical standard deviations of repeated measurements (see section 2.2 and Mayr, 2002). For uncertainty calculation of the ϵ values, see section 2.4.

Gelöscht:

- Gelöscht: purple
- Gelöscht: black

References

- Ali, H. A. M., Mayes, R. W., Hector, B. L., Verma, A. K. and Ørskov, E. R.: The possible use of *n*-alkanes, long-chain fatty alcohols and long-chain fatty acids as markers in studies of the botanical composition of the diet of free-ranging herbivores, *The Journal of Agricultural Science*, 143(1), 85–95, doi:10.1017/S0021859605004958, 2005.
- 1305 Allison, G. B., Gat, J. R. and Leaney, F. W. J.: The relationship between deuterium and oxygen-18 delta values in leaf water, *Chemical Geology*, 58, 145–156, 1985.
- Altermatt, H. A. and Neish, A. C.: The biosynthesis of cell wall carbohydrates: III. Further studies on formation of cellulose and xylan from labeled monosaccharides in wheat plants, *Canadian Journal of Biochemistry and Physiology*, 34(3), 405–413, doi:10.1139/o56-042, 1956.
- 1310 Amelung, W., Cheshire, M. V. and Guggenberger, G.: Determination of neutral and acidic sugars in soil by capillary gas-liquid chromatography after trifluoroacetic acid hydrolysis, *Soil Biology and Biochemistry*, 28(12), 1631–1639, 1996.
- Barbour, M. M. and Farquhar, G. D.: Relative humidity-and ABA-induced variation in carbon and oxygen isotope ratios of cotton leaves, *Plant, Cell & Environment*, 23(5), 473–485, 2000.
- 1315 Barbour, M. M., Roden, J. S., Farquhar, G. D. and Ehleringer, J. R.: Expressing leaf water and cellulose oxygen isotope ratios as enrichment above source water reveals evidence of a Péclet effect, *Oecologia*, 138(3), 426–435, doi:10.1007/s00442-003-1449-3, 2004.
- Bariac, T., Gonzalez-Dunia, J., Katerji, N., Béthenod, O., Bertolini, J. M. and Mariotti, A.: Spatial variation of the isotopic composition of water (^{18}O , ^2H) in the soil-plant-atmosphere system, 2. Assessment under field conditions, *Chemical Geology*, 115, 317–333, 1994.
- 1320 Benettin, P., Nehemy, M. F., Cernusak, L. A., Kahmen, A. and Jeffrey, J.: On the use of leaf water to determine plant water source: A proof of concept, *Scientific Briefing to Hydrological Processes*, 35(3), e14073, doi:10.1002/hyp.14073, 2021.
- 1325 Buck, A. L.: New Equations for Computing Vapor Pressure and Enhancement Factor, *Journal of Applied Meteorology*, 20, 1527–1532, 1981.
- Burget, E. G., Verma, R., Mølhøj, M. and Reiter, W.-D.: The Biosynthesis of L-Arabinose in Plants: Molecular Cloning and Characterization of a Golgi-Localized UDP-D-Xylose 4-Epimerase Encoded by the MUR4 Gene of Arabidopsis, *Plant Cell*, 15(February), 523–531, doi:10.1105/tpc.008425.response, 2003.
- 1330 Bush, R. T., Berke, M. A. and Jacobson, A. D.: Plant water dD and $\delta^{18}\text{O}$ of tundra species from West Greenland, Arctic, Antarctic, and Alpine Research, 49(3), 341–358, 2017.
- Cernusak, L. A., Wong, S. C. and Farquhar, G. D.: Oxygen isotope composition of phloem sap in relation to leaf water in *Ricinus communis*, *Functional Plant Biology*, 30(10), 1059–1070, 2003.
- 1335 Cernusak, L. A., Farquhar, G. D. and Pate, J. S.: Environmental and physiological controls over oxygen and carbon isotope composition of Tasmanian blue gum, *Eucalyptus globulus*, *Tree Physiology*, 25(2), 129–146, doi:10.1093/treephys/25.2.129, 2005.
- Cernusak, L. A., Barbour, M. M., Arndt, S. K., Cheesman, A. W., English, N. B., Feild, T. S., Helliker, B. R., Holloway-Phillips, M. M., Holtum, J. A. M., Kahmen, A., Mcinerney, F. A., Munksgaard, N. C., Simonin, K. A., Song, X., Stuart-Williams, H., West, J. B. and Farquhar, G. D.: Stable isotopes in leaf water of terrestrial plants, *Plant Cell and Environment*, 39(5), 1087–1102, doi:10.1111/pce.12703, 2016.
- 1340 Cheesman, A. W. and Cernusak, L. A.: Infidelity in the outback: Climate signal recorded in $\Delta^{18}\text{O}$ of leaf but not branch cellulose of eucalypts across an Australian aridity gradient, *Tree Physiology*, 37(5), 554–564, doi:10.1093/treephys/tpw121, 2017.
- 1345 Coplen, T. B.: Guidelines and recommended terms for expression of stable-isotope-ratio and gas-ratio measurement results, *Rapid Communications in Mass Spectrometry*, 25(17), 2538–2560, doi:10.1002/rcm.5129, 2011.
- 1350 Cormier, M.-A., Werner, R. A., Sauer, P. E., Gröcke, D. R., M.C., L., Wieloch, T., Schleucher, J. and Kahmen, A.: ^2H fractionations during the biosynthesis of carbohydrates and lipids imprint a metabolic signal on the $\delta^2\text{H}$ values of plant organic compounds, *New Phytologist*, 218(2), 479–491, doi:10.1111/nph.15016, 2018.

- Craig, H.: Isotopic Variations in Meteoric Waters, *Science*, 133, 1702–1703, 1961.
- 1355 Craig, H. and Gordon, L. I.: Deuterium and oxygen-18 variations in the ocean and the marine atmosphere, in *Proceedings of a Conference on Stable Isotopes in Oceanographic Studies and Palaeotemperatures*, edited by E. Tongiorgi, pp. 9–130, Lischi and Figli, Pisa., 1965.
- D'Souza, F., Garg, A. and Bhosle, N. B.: Seasonal variation in the chemical composition and carbohydrate signature compounds of biofilm, *Aquatic Microbial Ecology*, 41(2), 199–207, doi:10.3354/ame041199, 2005.
- 1360 Dansgaard, W.: Stable isotopes in precipitation, *Tellus*, 16(4), 436–468, doi:10.1111/j.2153-3490.1964.tb00181.x, 1964.
- Dawson, T. E.: Hydraulic lift and water use by plants: implications for water balance, performance and plant-plant interactions, *Oecologia*, 95(4), 565–574, 1993.
- 1365 Farquhar, G. D., Hubick, K. T., Condon, A. G. and Richards, R. A.: Carbon Isotope Fractionation and Plant Water-Use Efficiency, in *Stable Isotopes in Ecological Research. Ecological Studies (Analysis and Synthesis)*, vol. 68, edited by P. W. Rundel, J. R. Ehleringer, and K. A. Nagy, pp. 21–40, Springer-Verlag, New York., 1989.
- Feakins, S. J. and Sessions, A. L.: Controls on the D/H ratios of plant leaf waxes in an arid ecosystem, *Geochimica et Cosmochimica Acta*, 74(7), 2128–2141, doi:http://dx.doi.org/10.1016/j.gca.2010.01.016, 2010.
- 1370 Flanagan, L. B., Comstock, J. P. and Ehleringer, J. R.: Comparison of Modeled and Observed Environmental Influences on the Stable Oxygen and Hydrogen Isotope Composition of Leaf Water in *Phaseolus vulgaris* L., *Plant Physiology*, (96), 588–596, 1991.
- Freimuth, E. J., Diefendorf, A. F. and Lowell, T. V.: Hydrogen isotopes of *n*-alkanes and *n*-alkanoic acids as tracers of precipitation in a temperate forest and implications for paleorecords, *Geochimica et Cosmochimica Acta*, 206, 166–183, doi:10.1016/j.gca.2017.02.027, 2017.
- 1375 Gat, J. R. and Bowser, C. J.: The heavy isotope enrichment of water in coupled evaporative systems, in *Stable Isotope Geochemistry: A Tribute to Samuel Epstein*, vol. 3, edited by H. P. Taylor, J. R. O'Neil, and I. R. Kaplan, pp. 159–168, The Geochemical Society, Lancaster., 1991.
- 1380 Gat, J. R., Yakir, D., Goodfriend, G., Fritz, P., Trumborn, P., Lipp, J., Gev, I., Adar, E. and Waisel, Y.: Stable isotope composition of water in desert plants, *Plant and Soil*, 298(1–2), 31–45, doi:10.1007/s11104-007-9321-6, 2007.
- Gross, S. and Glaser, B.: Minimization of carbon addition during derivatization of monosaccharides for compound-specific $\delta^{13}\text{C}$ analysis in environmental research, *Rapid Communications in Mass Spectrometry*, 18, 2753–2764, doi:10.1002/rcm.1684, 2004.
- 1385 Harper, A. D. and Bar-Peled, M.: Biosynthesis of UDP-Xylose. Cloning and Characterization of a Novel Arabidopsis Gene Family, UXS, Encoding Soluble and Putative Membrane-Bound UDP-Glucuronic Acid Decarboxylase Isoforms, *Gene*, 130(December), 2188–2198, doi:10.1104/pp.009654.2188, 2002.
- 1390 Helliker, B. R. and Ehleringer, J. R.: Differential ^{18}O enrichment of leaf cellulose in C3 versus C4 grasses, *Functional Plant Biology*, 29, 435–442, 2002a.
- Helliker, B. R. and Ehleringer, J. R.: Grass blades as tree rings: environmentally induced changes in the oxygen isotope ratio of cellulose along the length of grass blades, *New Phytologist*, 155, 417–424, 2002b.
- 1395 Hepp, J., Tuthorn, M., Zech, R., Mügler, I., Schlütz, F., Zech, W. and Zech, M.: Reconstructing lake evaporation history and the isotopic composition of precipitation by a coupled $\delta^{18}\text{O}$ – $\delta^2\text{H}$ biomarker approach, *Journal of Hydrology*, 529, 622–631, 2015.
- Hepp, J., Rabus, M., Anhäuser, T., Bromm, T., Laforsch, C., Sirocko, F., Glaser, B. and Zech, M.: A sugar biomarker proxy for assessing terrestrial versus aquatic sedimentary input, *Organic Geochemistry*, 98, 98–104, doi:10.1016/j.orggeochem.2016.05.012, 2016.
- 1400 Hepp, J., Zech, R., Rozanski, K., Tuthorn, M., Glaser, B., Greule, M., Keppler, F., Huang, Y., Zech, W. and Zech, M.: Late Quaternary relative humidity changes from Mt. Kilimanjaro, based on a coupled ^2H – ^{18}O biomarker paleohygrometer approach, *Quaternary International*, 438, 116–130, doi:10.1016/j.quaint.2017.03.059, 2017.
- 1405 Hepp, J., Wüthrich, L., Bromm, T., Bliedtner, M., Schäfer, I. K., Glaser, B., Rozanski, K., Sirocko, F., Zech,

- R. and Zech, M.: How dry was the Younger Dryas? Evidence from a coupled $\delta^2\text{H}$ – $\delta^{18}\text{O}$ biomarker paleohygrometer applied to the Gemündener Maar sediments, Western Eifel, Germany, *Climate of the Past*, 15, 713–733, doi:10.5194/cp-15-713-2019, 2019.
- 1410 Hepp, J., Schäfer, I. K., Lanny, V., Franke, J., Bliedtner, M. and Rozanski, K.: Evaluation of bacterial glycerol dialkyl glycerol tetraether and ^2H – ^{18}O biomarker proxies along a central European topsoil transect, *Biogeosciences*, 17, 741–756, doi:10.5194/bg-17-741-2020, 2020.
- Herbin, G. A. and Robins, P. A.: Studies on plant cuticular waxes-II. Alkanes from members of the genus *Agave* (Agavaceae), the genera *Kalanchoe*, *Echeveria*, *Crassula* and *Sedum* (Crassulaceae) and the genus *Eucalyptus* (Myrtaceae) with an examination of Hutchinson, *Phytochemistry*, 7(1951), 257–268, 1968.
- 1415 Heyng, A., Mayr, C., Lücke, A., Wissel, H. and Striewski, B.: Late Holocene hydrologic changes in northern New Zealand inferred from stable isotope values of aquatic cellulose in sediments from Lake Pupuke, *Journal of Paleolimnology*, 51(4), 485–497, doi:10.1007/s10933-014-9769-3, 2014.
- 1420 Horita, J. and Wesolowski, D. J.: Liquid-vapor fractionation of oxygen and hydrogen isotopes of water from the freezing to the critical temperature, *Geochimica et Cosmochimica Acta*, 58(16), 3425–3437, doi:http://dx.doi.org/10.1016/0016-7037(94)90096-5, 1994.
- Hou, J., D’Andrea, W. J. and Huang, Y.: Can sedimentary leaf waxes record D/H ratios of continental precipitation? Field, model, and experimental assessments, *Geochimica et Cosmochimica Acta*, 72, 3503–3517, doi:10.1016/j.gca.2008.04.030, 2008.
- 1425 Huang, Y., Shuman, B., Wang, Y. and Ili, T. W.: Hydrogen isotope ratios of individual lipids in lake sediments as novel tracers of climatic and environmental change: a surface sediment test, *Journal of Paleolimnology*, 31, 363–375, 2004.
- 1430 Jia, G., Dungait, J. A. J., Bingham, E. M., Valiranta, M., Korhola, A. and Evershed, R. P.: Neutral monosaccharides as biomarker proxies for bog-forming plants for application to palaeovegetation reconstruction in ombrotrophic peat deposits, *Organic Geochemistry*, 39(12), 1790–1799, doi:10.1016/j.orggeochem.2008.07.002, 2008.
- Kahmen, A., Dawson, T. E., Vieth, A. and Sachse, D.: Leaf wax *n*-alkane δD values are determined early in the ontogeny of *Populus trichocarpa* leaves when grown under controlled environmental conditions, *Plant, Cell and Environment*, 34(10), 1639–1651, doi:10.1111/j.1365-3040.2011.02360.x, 2011.
- 1435 Kahmen, A., Schefuß, E. and Sachse, D.: Leaf water deuterium enrichment shapes leaf wax *n*-alkane δD values of angiosperm plants I: Experimental evidence and mechanistic insights, *Geochimica et Cosmochimica Acta*, 111, 39–49, 2013.
- 1440 Knapp, D. R.: *Handbook of Analytical Derivatization Reactions*, John Wiley & Sons, New York, Chichester, Brisbane, Toronto, Singapore., 1979.
- Lehmann, M. M., Gamarra, B., Kahmen, A., Siegwolf, R. T. W. and Saurer, M.: Oxygen isotope fractionations across individual leaf carbohydrates in grass and tree species, *Plant Cell and Environment*, 40(8), 1658–1670, doi:10.1111/pce.12974, 2017.
- 1445 Lemma, B., Bittner, L., Glaser, B., Kebede, S., Nemomissa, S., Zech, W. and Zech, M.: $\delta^2\text{H}_{n\text{-alkanes}}$ and $\delta^{18}\text{O}_{\text{sugar}}$ biomarker proxies from leaves and topsoils of the Bale Mountains, Ethiopia, and implications for paleoclimate reconstructions, *Biogeochemistry*, 153(2), 135–153, doi:10.1007/s10533-021-00773-z, 2021.
- 1450 Liu, H. T., Schäufele, R., Gong, X. Y. and Schnyder, H.: The $\delta^{18}\text{O}$ and $\delta^2\text{H}$ of water in the leaf growth-and-differentiation zone of grasses is close to source water in both humid and dry atmospheres, *New Phytologist*, 214(4), 1423–1431, doi:10.1111/nph.14549, 2017.
- Maffei, M.: Chemotaxonomic significance of leaf wax *n*-alkanes in the umbelliferae, cruciferae and leguminosae (subf. *Papilionoideae*), *Biochemical Systematics and Ecology*, 24(6), 531–545, doi:10.1016/0305-1978(96)00037-3, 1996.
- 1455 Mayr, C.: Möglichkeiten der Klimarekonstruktion im Holozän mit $\delta^{13}\text{C}$ - und $\delta^2\text{H}$ -Werten von Baum-Jahrringen auf der Basis von Klimakammerversuchen und Zentstudien, PhD thesis, Ludwig-Maximilians-Universität München. GSF-Bericht 14/02, 152 pp., 2002.
- Mayr, C., Laprida, C., Lücke, A., Martín, R. S., Massaferró, J., Ramón-Mercau, J. and Wissel, H.: Oxygen

Formatiert: Deutsch (Deutschland)

Formatiert: Deutsch (Deutschland)

Formatiert: Deutsch (Deutschland)

isotope ratios of chironomids, aquatic macrophytes and ostracods for lake-water isotopic reconstructions - Results of a calibration study in Patagonia, *Journal of Hydrology*, 529(P2), 600–607, doi:10.1016/j.jhydrol.2014.11.001, 2015.

McGill, R., Tukey, J. W. and Larsen, W. A.: Variations of Box Plots, *The American Statistician*, 32(1), 12–16, 1978.

Merlivat, L.: Molecular diffusivities of H₂¹⁶O, HD¹⁶O, and H₂¹⁸O in gases, *The Journal of Chemical Physics*, 69(6), 2864–2871, doi:http://dx.doi.org/10.1063/1.436884, 1978.

Mügler, I., Sachse, D., Werner, M., Xu, B., Wu, G., Yao, T. and Gleixner, G.: Effect of lake evaporation on δ D values of lacustrine *n*-alkanes: A comparison of Nam Co (Tibetan Plateau) and Holzmaar (Germany), *Organic Geochemistry*, 39(6), 711–729, 2008.

Prietz, J., Dechamps, N. and Spielvogel, S.: Analysis of non-cellulosic polysaccharides helps to reveal the history of thick organic surface layers on calcareous Alpine soils, *Plant and Soil*, 365(1–2), 93–114, doi:10.1007/s11104-012-1340-2, 2013.

R Core Team: R: A Language and Environment for Statistical Computing, [online] Available from: <https://www.r-project.org/>, 2015.

Rao, Z., Zhu, Z., Jia, G., Henderson, A. C. G., Xue, Q. and Wang, S.: Compound specific δ D values of long chain *n*-alkanes derived from terrestrial higher plants are indicative of the δ D of meteoric waters: Evidence from surface soils in eastern China, *Organic Geochemistry*, 40(8), 922–930, doi:http://dx.doi.org/10.1016/j.orggeochem.2009.04.011, 2009.

Roden, J. S. and Ehleringer, J. R.: Observations of Hydrogen and Oxygen Isotopes in Leaf Water Confirm the Craig-Gordon Model under Wide-Ranging Environmental Conditions, *Plant Physiology*, 120(August), 1165–1173, 1999.

Sachse, D., Radke, J. and Gleixner, G.: Hydrogen isotope ratios of recent lacustrine sedimentary *n*-alkanes record modern climate variability, *Geochimica et Cosmochimica Acta*, 68(23), 4877–4889, doi:http://dx.doi.org/10.1016/j.gca.2004.06.004, 2004.

Sachse, D., Billault, I., Bowen, G. J., Chikaraishi, Y., Dawson, T. E., Feakins, S. J., Freeman, K. H., Magill, C. R., McInerney, F. A., van der Meer, M. T. J., Polissar, P., Robins, R. J., Sachs, J. P., Schmidt, H.-L., Sessions, A. L., White, J. W. C. and West, J. B.: Molecular Paleohydrology: Interpreting the Hydrogen-Isotopic Composition of Lipid Biomarkers from Photosynthesizing Organisms, *Annual Reviews*, 40, 221–249, doi:10.1146/annurev-earth-042711-105535, 2012.

Santrucek, J., Kveton, J., Setlik, J. and Bulickova, L.: Spatial Variation of Deuterium Enrichment in Bulk Water of Snowgum Leaves, *Plant Physiology*, 143(1), 88–97, doi:10.1104/pp.106.089284, 2007.

Sauer, P. E., Eglinton, T. I., Hayes, J. M., Schimmelmann, A. and Sessions, A. L.: Compound-specific D/H ratios of lipid biomarkers from sediments as a proxy for environmental and climatic conditions, *Geochimica et Cosmochimica Acta*, 65(2), 213–222, doi:http://dx.doi.org/10.1016/S0016-7037(00)00520-2, 2001.

Schäfer, I. K., Lanny, V., Franke, J., Eglinton, T. I., Zech, M., Vysloužilová, B. and Zech, R.: Leaf waxes in litter and topsoils along a European transect, *SOIL*, 2, 551–564, doi:10.5194/soil-2-551-2016, 2016.

Schmidt, H.-L., Werner, R. A. and Roßmann, A.: ¹⁸O Pattern and biosynthesis of natural plant products, *Phytochemistry*, 58(1), 9–32, doi:http://dx.doi.org/10.1016/S0031-9422(01)00017-6, 2001.

Schmidt, H.-L., Werner, R. A. and Eisenreich, W.: Systematics of ²H patterns in natural compounds and its importance for the elucidation of biosynthetic pathways, *Phytochemistry Reviews*, 2(1–2), 61–85, doi:10.1023/B:PHYT.0000004185.92648.ae, 2003.

Sessions, A. L., Burgoyne, T. W., Schimmelmann, A. and Hayes, J. M.: Fractionation of hydrogen isotopes in lipid biosynthesis, *Organic Geochemistry*, 30, 1193–1200, 1999.

Song, X., Farquhar, G. D., Gessler, A. and Barbour, M. M.: Turnover time of the non-structural carbohydrate pool influences δ 18O of leaf cellulose, *Plant Cell and Environment*, 37(11), 2500–2507, doi:10.1111/pce.12309, 2014.

Sternberg, L. and Ellsworth, P. F. V.: Divergent Biochemical Fractionation, Not Convergent Temperature, Explains Cellulose Oxygen Isotope Enrichment across Latitudes, *PLoS ONE*, 6(11), e28040, doi:10.1371/journal.pone.0028040, 2011.

- Sternberg, L. da S. L. O. and DeNiro, M. J. D.: Biogeochemical implications of the isotopic equilibrium fractionation factor between the oxygen atoms of acetone and water, *Geochimica et Cosmochimica Acta*, 47(12), 2271–2274, doi:10.1016/0016-7037(83)90049-2, 1983.
- 1515 Sternberg, L. S. L., DeNiro, M. J. and Savidge, R. A.: Oxygen Isotope Exchange between Metabolites and Water during Biochemical Reactions Leading to Cellulose Synthesis, *Plant Physiology*, 82, 423–427, 1986.
- Strobel, P., Haberzettl, T., Bliedtner, M., Struck, J., Glaser, B., Zech, M. and Zech, R.: The potential of $\delta^2\text{H}_{n\text{-alkanes}}$ and $\delta^{18}\text{O}_{\text{sugar}}$ for paleoclimate reconstruction – A regional calibration study for South Africa, *Science of the Total Environment*, 716, 137045, doi:https://doi.org/10.1016/j.scitotenv.2020.137045, 2020.
- 1520 Tipple, B. J., Berke, M. A., Doman, C. E., Khachaturyan, S. and Ehleringer, J. R.: Leaf-wax *n*-alkanes record the plant-water environment at leaf flush, *Proceedings of the National Academy of Sciences*, 110(7), 2659–2664, doi:10.1073/pnas.1213875110, 2013.
- 1525 Tipple, B. J., Berke, M. A., Hambach, B., Roden, J. S. and Ehleringer, J. R.: Predicting leaf wax *n*-alkane $^2\text{H}/^1\text{H}$ ratios: Controlled water source and humidity experiments with hydroponically grown trees confirm predictions of Craig-Gordon model, *Plant, Cell and Environment*, 38(6), 1035–1047, doi:10.1111/pce.12457, 2015.
- Tuthorn, M., Zech, M., Ruppenthal, M., Oelmann, Y., Kahmen, A., del Valle, H. F., Wilcke, W. and Glaser, B.: Oxygen isotope ratios ($^{18}\text{O}/^{16}\text{O}$) of hemicellulose-derived sugar biomarkers in plants, soils and sediments as paleoclimate proxy II: Insight from a climate transect study, *Geochimica et Cosmochimica Acta*, 126, 624–634, doi:http://dx.doi.org/10.1016/j.gca.2013.11.002, 2014.
- 1530 Tuthorn, M., Zech, R., Ruppenthal, M., Oelmann, Y., Kahmen, A., del Valle, H. F., Eglinton, T., Rozanski, K. and Zech, M.: Coupling $\delta^2\text{H}$ and $\delta^{18}\text{O}$ biomarker results yields information on relative humidity and isotopic composition of precipitation - a climate transect validation study, *Biogeosciences*, 12, 3913–3924, doi:10.5194/bg-12-3913-2015, 2015.
- 1535 Walker, C. D. and Brunel, J.-P.: Examining Evapotranspiration in a Semi-Arid Region using Stable Isotopes of Hydrogen and Oxygen, *Journal of Hydrology*, 118, 55–75, 1990.
- Wang, X.-F., Yakir, D. and Avisha, M.: Non-climatic variations in the oxygen isotopic composition of plants, *Global Change Biology*, 4, 835–849, 1998.
- 1540 Waterhouse, J. S., Cheng, S., Juchelka, D., Loader, N. J., McCarroll, D., Switsur, V. R. and Gautam, L.: Position-specific measurement of oxygen isotope ratios in cellulose: Isotopic exchange during heterotrophic cellulose synthesis, *Geochimica et Cosmochimica Acta*, 112(0), 178–191, doi:http://dx.doi.org/10.1016/j.gca.2013.02.021, 2013.
- 1545 Werner, R. A. and Brand, W. A.: Referencing strategies and techniques in stable isotope ratio analysis, *Rapid Communications in Mass Spectrometry*, 15(7), 501–519, doi:10.1002/rcm.258, 2001.
- Wissel, H., Mayr, C. and Lücke, A.: A new approach for the isolation of cellulose from aquatic plant tissue and freshwater sediments for stable isotope analysis, *Organic Geochemistry*, 39(11), 1545–1561, doi:http://dx.doi.org/10.1016/j.orggeochem.2008.07.014, 2008.
- 1550 Yakir, D. and DeNiro, M. J.: Oxygen and Hydrogen Isotope Fractionation during Cellulose Metabolism in *Lemna gibba* L., *Plant Ecology*, 93, 325–332, 1990.
- Zech, M. and Glaser, B.: Compound-specific $\delta^{18}\text{O}$ analyses of neutral sugars in soils using gas chromatography-pyrolysis-isotope ratio mass spectrometry: problems, possible solutions and a first application, *Rapid Communications in Mass Spectrometry*, 23, 3522–3532, doi:10.1002/rcm, 2009.
- 1555 Zech, M., Pedentchouk, N., Buggle, B., Leiber, K., Kalbitz, K., Markovic, S. B. and Glaser, B.: Effect of leaf litter degradation and seasonality on D/H isotope ratios of *n*-alkane biomarkers, *Geochimica et Cosmochimica Acta*, 75(17), 4917–4928, doi:http://dx.doi.org/10.1016/j.gca.2011.06.006, 2011.
- 1560 Zech, M., Werner, R. A., Juchelka, D., Kalbitz, K., Buggle, B. and Glaser, B.: Absence of oxygen isotope fractionation/exchange of (hemi-) cellulose derived sugars during litter decomposition, *Organic Geochemistry*, 42(12), 1470–1475, doi:http://dx.doi.org/10.1016/j.orggeochem.2011.06.006, 2012.
- Zech, M., Tuthorn, M., Detsch, F., Rozanski, K., Zech, R., Zöller, L., Zech, W. and Glaser, B.: A 220 ka

1565 terrestrial $\delta^{18}\text{O}$ and deuterium excess biomarker record from an eolian permafrost paleosol
sequence, NE-Siberia, Chemical Geology, 360–361, 220–230,
doi:http://dx.doi.org/10.1016/j.chemgeo.2013.10.023, 2013a.

Zech, M., Tuthorn, M., Glaser, B., Amelung, W., Huwe, B., Zech, W., Zöller, L. and Löffler, J.: Natural
abundance of $\delta^{18}\text{O}$ of sugar biomarkers in topsoils along a climate transect over the Central
1570 Scandinavian Mountains, Norway, Journal of Plant Nutrition and Soil Science, 176(1), 12–15,
doi:10.1002/jpln.201200365, 2013b.

Zech, M., Mayr, C., Tuthorn, M., Leiber-Sauheitl, K. and Glaser, B.: Oxygen isotope ratios ($^{18}\text{O}/^{16}\text{O}$) of
hemicellulose-derived sugar biomarkers in plants, soils and sediments as paleoclimate proxy I:
Insight from a climate chamber experiment, Geochimica et Cosmochimica Acta, 126(0), 614–
1575 623, doi:http://dx.doi.org/10.1016/j.gca.2013.10.048, 2014a.

Zech, M., Mayr, C., Tuthorn, M., Leiber-Sauheitl, K. and Glaser, B.: Reply to the comment of Sternberg
on “Zech et al. (2014) Oxygen isotope ratios ($^{18}\text{O}/^{16}\text{O}$) of hemicellulose-derived sugar
biomarkers in plants, soils and sediments as paleoclimate proxy I: Insight from a climate
chamber experiment. GCA, Geochimica et Cosmochimica Acta, 141(0), 680–682,
1580 doi:10.1016/j.gca.2014.04.051, 2014b.

Zech, M., Zech, R., Rozanski, K., Gleixner, G. and Zech, W.: Do *n*-alkane biomarkers in soils/sediments
reflect the $\delta^2\text{H}$ isotopic composition of precipitation? A case study from Mt. Kilimanjaro and
implications for paleoaltimetry and paleoclimate research, Isotopes in Environmental and
Health Studies, 51(4), 508–524, doi:10.1080/10256016.2015.1058790, 2015.

1585 Zhang, X., Gillespie, A. L. and Sessions, A. L.: Large D/H variations in bacterial lipids reflect central
metabolic pathways, PNAS, 106(31), 12580–12586, 2009.

Zhou, Y., Grice, K., Stuart-Williams, H., Farquhar, G. D., Hocart, C. H., Lu, H. and Liu, W.: Biosynthetic
origin of the saw-toothed profile in $\delta^{13}\text{C}$ and $\delta^2\text{H}$ of *n*-alkanes and systematic isotopic
differences between *n*-, *iso*- and *anteiso*-alkanes in leaf waxes of land plants, Phytochemistry,
1590 71(4), 388–403, doi:10.1016/j.phytochem.2009.11.009, 2010.