# Competing effects of nitrogen deposition and ozone exposure on Northern hemispheric terrestrial carbon uptake and storage, 1850-2099

Martina Franz<sup>1,2</sup> and Sönke Zaehle<sup>1,3</sup>

<sup>1</sup>Biogeochemical Signals Department, Max Planck Institute for Biogeochemistry, Jena, Germany
 <sup>2</sup>International Max Planck Research School (IMPRS) for Global Biogeochemical Cycles, Jena, Germany
 <sup>3</sup>Michael Stifel Center Jena for Data-driven and Simulation Science, Jena, Germany

Correspondence: Martina Franz (mfranz@bgc-jena.mpg.de)

#### Abstract.

Tropospheric ozone  $(O_3)$  and nitrogen deposition affect vegetation growth and thus thereby the ability of the land biosphere to take up and store carbon. However, the magnitude of this effect these effects on the contemporary and future terrestrial carbon balance is insufficiently understood. Here, we apply an extended version of the O-CN terrestrial biosphere model that

- 5 simulates the atmosphere to canopy transport of O<sub>3</sub>, its surface and stomatal uptake, the O<sub>3</sub>-induced leaf injury, as well as the ozone-induced leaf injurycoupled terrestrial carbon and nitrogen cycles. We use this model to simulate past and future impacts of air pollution (ozone and nitrogen deposition) against a background of concurrent changes in climate and carbon dioxide concentrations (CO<sub>2</sub>) for two contrasting representative concentration pathways (RCP) scenarios (RCP2.6 and RCP8.5). The simulations show that O<sub>3</sub>-related damage considerably reduced Northern hemispheric gross primary production (GPP)
- and long-term carbon storage between 1850 and the 2010s. The ozone simulated  $O_3$  effect on GPP in the Northern hemisphere peaks at peaked towards the end of the 20<sup>th</sup> century with reductions of 4 %, causing a reduction in the Northern hemispheric carbon sink of 0.4 PgCyr<sup>-1</sup>. During the 21<sup>st</sup> century, ozone-induced  $O_3$ -induced reductions in GPP and carbon storage is are projected to decline through a combination of direct air pollution control methods that reduce tropospheric near surface  $O_3$ and the indirect effects of rising atmospheric CO<sub>2</sub>, which reduces stomatal uptake of ozone- $O_3$  concurrent with increases of
- 15 leaf-level water-use efficiency.

However, in hotspot regions such as East Asia, the model simulations suggest a sustained decrease of GPP by more than 8% during % throughout the  $21^{st}$  century. Regionally, ozone exposure reduces  $O_3$  exposure reduces projected carbon storage at the end of the  $21^{st}$  century by up to 15 % in parts of Europe, the US and East Asia. These estimates are lower compared to previous studies, which partially results from the explicit representation of non-stomatal ozone destruction, which considerably reduces cimulated ozone uptake by lower and incurred injury.

20 reduces simulated ozone uptake by leaves and incurred injury.

Our simulations suggest that ozone damage largely offsets the growth stimulating effect induced by nitrogen deposition in the Northern hemisphere until the 2050s. Thus, accounting stimulating effect of nitrogen deposition on regional GPP and carbon storage is lower in magnitude compared to the detrimental effect of  $O_3$  during most of the simulation period for both RCPs. In the second half of the 21<sup>st</sup> century, the detrimental effect of  $O_3$  on GPP is outweighed by nitrogen deposition, but the 25 <u>effect of nitrogen deposition on land carbon storage remains lower than the effect of  $O_3$ . Accounting for the stimulating effects of nitrogen deposition but omitting the detrimental effect of  $O_3$  <u>might may</u> lead to an over estimation of <u>projected</u> carbon uptake and storage.</u>

### 1 Introduction

Productivity and carbon storage in many Northern hemispheric terrestrial ecosystems are affected by the limited availability of

- 30 nitrogen (Vitousek and Howarth, 1991; LeBauer and Treseder, 2008; Zaehle, 2013)(N; Vitousek and Howarth, 1991; LeBauer and Treseder, 2008; Zaehle, 2013)(N; Vitousek and Howarth, 1991; LeBauer and Treseder, As a side-effect of air pollution, increased deposition of reactive nitrogen from e.g. anthropogenic fossil fuel burning and increased soil emissions associated with fertiliser use (Galloway et al., 2004) have the potential to fertilise these N-limited ecosystems and thereby enhance productivity and carbon storage (Norby, 1998; Zaehle et al., 2011; Thomas et al., 2010). However, oxidised forms of reactive nitrogen in the atmosphere (collectively referred to as NO<sub>y</sub>) are also a precursor of tro-
- 35 pospheric ozone (O<sub>3</sub>). Ozone (O<sub>3</sub>) is a toxic air pollutant that enters plants primarily though the leaves' stomata where it can induce cellular damage (Fiscus et al., 2005; Tausz et al., 2007; McAinsh et al., 2002). Commonly observed effects are visible injury (Langebartels et al., 1991; Wohlgemuth et al., 2002), reductions in photosynthetic capacity (Tjoelker et al., 1995; Wittig et al., 2007) - and growth or yield (Grantz et al., 2006; Hayes et al., 2007; Feng and Kobayashi, 2009; Wittig et al., 2009; Leisner and Ainsworth, 2012). Ozone induced plant damage can reduce the terrestrial carbon uptake and storage and through this
- 40 cause an increase in atmospheric  $CO_2$  concentrations and an intensification of climate change (Sitch et al., 2007; Ainsworth et al., 2012).

Ozone concentrations over the mid- and high-latitudes of Eurasia and North America Ozone mixing ratios in Europe have approximately doubled between the pre-industrial period (around 1860) and the year 2000 (Akimoto, 2003; Cooper et al., 2014; Marenco et a during the 20<sup>th</sup> century (Cooper et al., 2014). The anthropogenic increase in NO<sub>y</sub> emissions primarily from combustion

- sources has been identified as the major cause for the increasing near-surface ozone concentrations between 1970–1995  $O_3$ concentrations between 1970–1995 in the mid-latitudes of the Northern Hemisphere (Fusco and Logan, 2003). Ozone levels are projected to decline until the end of the  $21^{st}$  century due to assumed stringent air pollution policies, but future climate conditions with increasing temperatures as well as reduced cloudiness and precipitation will tend to increase ozone  $O_3$  formation with increasing daily ozone  $O_3$  peaks and average concentrations in summer (Meleux et al., 2007; van Vuuren et al., 2011).
- 50 The application of the RCP scenarios (Moss et al., 2010; van Vuuren et al., 2011) in 14 global chemistry transport models results in the projection of declining annual global mean surface O<sub>3</sub> concentrations in most regions of the globe except South Asia where increases are simulated (Wild et al., 2012). Projections of nitrogen deposition in the 21<sup>st</sup> century suggest little change across all scenarios of the Representative concentration pathways (RCP), despite notable regional differences (Lamarque et al., 2013). Only under the most optimistic scenario RCP2.6 a A small decline in deposition rates is proposed only under

# 55 the scenario RCP2.6.

Simulations with nitrogen-enabled terrestrial biosphere models suggest that N deposition may be responsible for 10 to 50 % of the global residual land carbon uptake (Zaehle et al., 2011; Quéré et al., 2018). Several models including the ozone

 $O_3$  effect on carbon cycle suggest that simulated present-day and future ozone  $O_3$  exposure can reduce regional and global scale productivity (Felzer et al., 2005; Sitch et al., 2007; Franz et al., 2017; Lombardozzi et al., 2015; Oliver et al., 2018).

For instance, modelling studies by Sitch et al. (2007) and Oliver et al. (2018) suggest a reduction in O<sub>3</sub> induced damage of global gross primary production (GPP) by 4-15-4-15 % and an associated reduction of land carbon storage by 3-10 %. 3-10 %. Where Sitch et al. (2007) simulated global O<sub>3</sub> impacts between 1901–2100 and Oliver et al. (2018) focused on a European scale damage between 1901–2050.

Here, we assess the combined effect of ozone- $O_3$  and nitrogen deposition on the Northern hemispheric terrestrial biosphere against the background of simulated changes due to increasing atmospheric  $CO_2$  and climate change. Elevated levels of atmospheric  $CO_2$  stimulate leaf photosynthesis and reduce stomatal conductance (Medlyn et al., 2001; Ainsworth and Long, 2005), and therefore can increase plant growth and plant nitrogen limitation (Oren et al., 2001; Norby et al., 2009; Zaehle et al., 2014). However, reductions in stomatal conductance reduce the leaf-level uptake of air pollutants like  $O_3$  and thereby have the potential to restrict ozone- $O_3$  induced damage to plants (Paoletti and Grulke, 2005; Barnes and Pfirrmann, 1992; Isebrands et al., 2001; Talhelm et al., 2014; Zak et al., 2011; Noormets et al., 2010).

We analyse the response of the Northern hemispheric carbon cycle to changes in climate, atmospheric  $CO_2$  and  $O_3$  as well as N deposition for the historical period (1850-2005) and two future scenarios (2006-2099), the most optimistic and most pessimistic RCP scenario(2006-2099), a high and a low climate-change mitigation scenario, RCP2.6 and RCP8.5 respectively). In a factorial analysis, we investigate the impact of the single drivers ( $O_3$ ,  $CO_2$  and N deposition), as well as

their interaction (specifically the interaction between O<sub>3</sub> and CO<sub>2</sub>, and O<sub>3</sub> and N deposition) on plant growth and terrestrial carbon storage. We employ a significantly enhanced version of the O-CN terrestrial biosphere model (Zaehle and Friend, 2010), which explicitly accounts for the O<sub>3</sub> transport and deposition from the free atmosphere into the stomatestroposphere into the stomatestroposphere into the stomatestroposphere into the stomatestroposphere into the stomate O<sub>3</sub> removal by other processes (such as soil and leaf surface uptake) (Franz et al., 2017). This model has been evaluated against biomass damage relationships observed in a range of fumigation/filtration experiments with European tree species (Büker et al., 2015; Franz et al., 2018).

#### 2 Methods

Simulations are conducted with the O-CN terrestrial biosphere model (Zaehle and Friend, 2010; Franz et al., 2017), version  $tun_{VC}$  where ozone O<sub>3</sub> damage is calculated based on injury functions to the maximum carboxylation capacity of the leaf  $V_{cmax}$  (Franz et al., 2018). The  $tun_{VC}$  injury functions were calibrated to reproduce observed biomass damage relationships

85 of experiments with a range of European tree species in fumigation/filtration experiments (Franz et al., 2018). Contrary to Franz et al. (2018), (Franz et al., 2018; Büker et al., 2015).

The O-CN model includes an  $O_3$  deposition scheme that explicitly accounts for the  $O_3$  transport and deposition from the free troposphere into the stomata (Franz et al., 2017). Here, we use the ozone deposition scheme described in Franz et al. (2017) is applied in the simulations here (referred to as D-model version in Franz et al. (2017)) in Franz et al. (2017), contrary to

90 Franz et al. (2018) where the  $O_3$  deposition scheme was turned off.

#### 2.1 The O-CN model

O-CN (Zaehle and Friend, 2010) is a further development of the biogeochemistry in the land-surface-scheme ORCHIDEE (Krinner et al., 2005), and simulates the coupled terrestrial carbon (C), nitrogen (N) and water cycles for twelve plant functional types. The model accounts for the effects of nitrogen availability on growth, root:shoot allocation, litter and soil organic matter

- 95 decay, and represents a comprehensive nitrogen cycle including process-oriented formulations for nitrogen leaching and gas losses, and its ability to reproduce N fertilisation experiments has been evaluated by (Meyerholt and Zaehle, 2015). O-CN compares well to a range of regional to global terrestrial biosphere benchmarks (Quéré et al., 2018). O-CN is driven by climate data, N deposition, atmospheric composition including the deposition, atmospheric CO<sub>2</sub> and O<sub>3</sub> burden, concentrations and land use information(land cover, land cover change, and fertiliser application).
- O-CN simulates a multi-layer canopy with up to 20 layers (each with a thickness of up to 0.5 leaf area index) where net photosynthesis is calculated for shaded and sun-lit leaves with consideration of the light profiles of diffuse and direct radiation (Kull and Kruijt, 1998; Friend, 2001; Zaehle and Friend, 2010). Stomatal conductance to water,  $CO_2$  and  $O_3$  is calculated coupled to net photosynthesis following a Ball-&-Berry-type formulation (see Sect. 2.2). Leaf nitrogen concentration and leaf area determine the photosynthetic capacity, which are both affected by ecosystem available N. The maximum carboxylation
- 105 capacity ( $V_{cmax}$ ) and electron transport capacity ( $J_{max}$ ) of the leaf increase with an increased leaf nitrogen concentration, leading to an increase in the maximum net photosynthesis and stomatal conductance per unit leaf area (Zaehle and Friend, 2010). The highest leaf N content is simulated at the top of the canopy and exponentially decreases with increasing canopy depth (Friend, 2001; Niinemets et al., 2015). Following this, the net photosynthesis, stomatal conductance and O<sub>3</sub> uptake are generally highest in the top of the canopy and lowest in the bottom of the canopy. Changes in stomatal conductance affect 110 transpiration rates and estimates of O<sub>3</sub> uptake and ozone O<sub>3</sub> damage.

#### 2.2 Ozone injury calculation in O-CN

Leaf-level ozone  $O_3$  uptake is determined by stomatal conductance and atmospheric  $O_3$  concentrations, as described in Franz et al. (2017). In contrast to Franz et al. (2017), the stomatal conductance  $g_{st}$  is calculated based on the Ball and Berry formulation (Ball et al., 1987) as

115 
$$g_{st,l} = g_0 + g_1 \times \frac{A_{n,l} \times RH \times f(height_l)}{C_a} \frac{A_{n,l} \times RH \times f(height_l)}{[CO_2]}$$
(1)

where net photosynthesis  $(A_{n,l})$  RH is the atmospheric relative humidity,  $f(height_l)$  the water-transport limitation with canopy height,  $[CO_2]$  the atmospheric CO<sub>2</sub> concentration,  $A_{n,l}$  the net photosynthesis,  $g_0$  the residual conductance when  $A_n$ approaches zero, and  $g_1$  the stomatal-slope parameter as in Krinner et al. (2005). The index l indicates that  $g_{st}$  and  $A_n$  are calculated separately for each canopy layer.  $A_{n,l}$  is calculated as described in Zaehle and Friend (2010) as a function of the

120 leaf-internal partial pressure of CO<sub>2</sub>, absorbed photosynthetic photon flux density on shaded and sunlit leaves, leaf temperature, the nitrogen-specific rates of maximum light harvesting, electron transport ( $J_{max}$ ) and carboxylation rates ( $V_{cmax}$ ). RH is the atmospheric relative humidity,  $f(height_l)$  the water-transport limitation with canopy height,  $C_a$  the atmospheric concentration,  $g_0$  the residual conductance when  $A_n$  approaches zero, and  $g_1$  the stomatal-slope parameter as in Krinner et al. (2005). The index l indicates that  $g_{st}$  is calculated separately for each canopy layer.

125 The stomatal conductance to  $\frac{O_3}{O_3} = O_3 g_{st,l}^{O_3}$  is calculated as

140

145

$$g_{st,l}^{O_3} = \frac{g_{st,l}}{1.51} \tag{2}$$

where the factor 1.51 accounts for the different diffusivity of  $O_3$  from water vapour (Massman, 1998).

For each canopy layer, the O<sub>3</sub> stomatal flux  $(f_{st,l}, \text{nmol m}^{-2} \text{ s}^{-1})$  is calculated from the canopy  $O_3$  concentration  $(\chi_{can}^{O_3}[O_3]_{can}^{can})$ , and  $g_{st,l}$  is calculated as

130 
$$f_{st,l} = (\underbrace{\chi^{O_3}_{atm}}_{atm}[O_3]^{can} - \underbrace{\chi^{O_3}_{i}}_{i}[O_3]^{i})g^{O_3}_{st,l}.$$
(3)

where the leaf-internal  $O_3$  concentration  $(\chi_i^{O_3}[O_3]^i)$  is assumed to be zero (Laisk et al., 1989).

The ozone  $O_3$  deposition module calculates  $\chi_{can}^{O_3} [Q_3]_{can}^{can}$  from the  $O_3$  concentration in 45m height  $(\chi_{atm}^{O_3})$  as provided by the chemical transport models as input for terrestrial biosphere models like O-CN (Franz et al., 2017).  $\chi_{can}^{O_3}$ , is calculated based m height  $([O_3]^{atm})$ , which represents the lowest level  $O_3$  concentrations in the forcing data (Franz et al., 2017). Based on the constant flux assumption  $[O_3]^{can}$  in units of nmol m<sup>-3</sup> is calculated as

$$\underline{\chi_{can}^{O_3}}[\mathcal{O}_3]_{\sim\sim}^{can} = \underline{\chi_{atm}^{O_3}}[\mathcal{O}_3]_{\sim\sim}^{atm} \left(1 - \frac{R_a}{R_a + R_b + R_c}\right) \tag{4}$$

with  $R_a$  the aerodynamic resistance,  $R_b$  the canopy-scale quasi-laminar layer resistance and  $R_c$  the compound surface resistance to O<sub>3</sub> deposition.  $R_c$  is calculated as the sum of the canopy scale stomatal and the non-stomatal resistance to O<sub>3</sub> uptake (Franz et al., 2017). The non-stomatal resistance is defined by the O<sub>3</sub> destruction on the leaf surface, within-canopy resistance to O<sub>3</sub> transport, and ground surface resistance (Franz et al., 2017).

Part of the  $O_3$  taken up into the leaves is assumed to get detoxified and to cause no damage to the plant. Ozone fluxes exceeding the detoxification threshold of X ( $f_{st,l,X}$ ,) are accumulated over time to give the cumulative Without the application of the  $O_3$  uptake above a flux deposition module, the  $O_3$  uptake inside the leaves would be calculated based on the near surface  $O_3$  concentrations from the forcing data without accounting for the turbulent transport between the lower troposphere and the leaves, as well as the deposition and destruction of ozone on other surfaces.

The accumulation of O<sub>3</sub> fluxes into the leaves above a threshold of X nmol m<sup>-2</sup> s<sup>-1</sup> ( $CUOX_l$ ) with  $f_{st,l,1}$ , nmol m<sup>-2</sup> s<sup>-1</sup>) with

$$f_{\underline{st,l,Xst,l}}(X) = \underline{MAX} \underbrace{\max}_{0}(0, f_{st,l} - X).$$
(5)

gives the  $CUOX_L$ . Summing  $CUOX_l$  over all canopy layers gives the canopy value CUOX (Franz et al., 2017). In this study, a flux threshold of 1 nmol m<sup>-2</sup> s<sup>-1</sup>, i.e. CUO1, is applied to account for the plants ability to detoxify part of the taken up O<sub>3</sub> (Franz et al., 2018; LRTAP-Convention, 2017; Büker et al., 2015). The cumulative uptake of O<sub>3</sub> without detoxification, i.e. a threshold of zero, is represented by CUO0.

Emerging leaves are assumed undamaged and accumulate CUOX during the growing season. The  $CUOX_l$  is reduced by the fraction of newly developed leaves per time step and canopy layer. Deciduous PFTs shed all CUOX at the end of the

155 growing season and grow uninjured leaves the next spring. Evergreen PFTs shed proportionate amounts of *CUOX* during the entire year when new leaves or needles are grown or old foliage is replaced.

The ozone  $O_3$  injury fraction  $(d_l^{O_3})$ , is calculated as a linear function of  $CUO1_l$ 

$$d_l^{O_3} = 1 - b \times CUO1_l \tag{6}$$

where the slope of the injury function (b) is set to 0.075 for broadleaf species and 0.025 for needleleaf species (Franz et al., 2018).  $d_l^{O_3}$  is calculated separately for each canopy layer l according to the specific accumulated ozone-O<sub>3</sub> uptake of the respective canopy layer ( $CUO1_l$ ), and takes values between 0 and 1. Within-canopy gradients in stomatal conductance and photosynthetic capacity cause variations of  $CUO1_l$  and hence  $d_l^{O_3}$  between canopy layers.

The effect of  $\frac{1}{2}$  or  $O_3$  injury on plant carbon uptake is calculated by

$$V_{cmax,l}^{O_3} = V_{cmax,l}(1 - d_l^{O_3}).$$
<sup>(7)</sup>

165 with the maximum carboxylation capacity of the leaf in the respective canopy layer  $(V_{cmax,l})$ , which is used in the calculation of  $A_{n,l}A_{n,l}$ . J<sub>max,l</sub> is reduced in proportion to  $V_{cmax,l}$  such that the ratio between both keeps-is maintained.

Ozone induced reductions in  $A_{n,l}A_{n,l}$  cause a decline in  $g_{st,l}$  as both are tightly coupled. Lower values of  $g_{st,l}$  diminish the O<sub>3</sub> uptake into the plant  $(f_{st,l})$  and slow the increase in  $CUO1_l$  and hence ozone O<sub>3</sub> induced injury.

## 2.3 Model forcing

- 170 The model is driven by climate model output of the Institute Pierre Simon Laplace (IPSL) general circulation model IPSL-CM5A-LR (Dufresne et al., 2013), bias-corrected according to the Inter-Sectoral Impact Model Intercomparison Project (Hempel et al., 2013). Downward nitrogen deposition velocity and near surface ozone. The applied meteorological forcing for near-surface conditions comprises daily data of specific humidity, incoming long wave radiation, incoming short wave radiation, cloudiness, wind speed, maximum temperature, minimum temperature and total precipitation, which are disaggregated
- to the 30 min time step of the model using a statistical weather generator (Krinner et al., 2005). Reduced and oxidised monthly mean nitrogen deposition in wet and dry form and monthly mean near surface  $O_3$  concentrations are provided by CAM, the community atmosphere model (Lamarque et al., 2010; Cionni et al., 2011). Land cover, soil, and N fertiliser application are used as in Zaehle et al. (2011) and kept at 2000 values throughout the simulation. Data on atmospheric  $CO_2$  concentrations are obtained from Meinshausen et al. (2011). Through all simulations present day land-use information are applied for the year
- 180 2000 (Hurtt et al., 2011).

Figure 1 provides and overview over the scenarios applied in this study. Note that there are important regional patterns behind the changes in N deposition and tropospheric ozonenear surface  $O_3$ , which are shown in the Appendix Supplement Fig. S1 and Fig. S2.



Figure 1. Time series of the terrestrial Northern hemispheric ( $\geq 30^{\circ}$ N) - mean a) tropospheric ozone near surface O<sub>3</sub> concentration, b) summed nitrogen deposition, c) air temperature at 2 m height, and d) atmospheric CO<sub>2</sub> concentration according to the RCP2.6 and RCP8.5 pollution emission scenario. For visual clarity, the effect of the seasonal cycles of tropospheric ozone near surface O<sub>3</sub> and N deposition are smoothed by a moving average of 12 months. See SI for spatial patterns of N deposition and tropospheric ozone near surface O<sub>3</sub> (Appendix Supplement Fig. S1 and Fig. S2).

#### 2.4 Modelling protocol

185 The model is run at a spatial resolution of  $1^{\circ} \times 1^{\circ}$  and operates on a half hourly time step. As the injury functions developed by Franz et al. (2018) are based on manipulation exposure experiments with boreal and temperate European tree species, the simulation scope domain is restricted to the temperate and boreal region of the Northern Hemisphere  $\geq 30^{\circ}$ N.

To achieve an equilibrium in terms of the terrestrial vegetation and soil carbon and nitrogen pools, O-CN is run for 1291 years (including 10 iterations of 1000 years soil biogeochemistry and 100 years vegetation+soil biogeochemistry) by using the

190

forcing data of the year 1850 data. Prior to year 1901 climate years are randomly iterated from the period of 1901 to 1930, as 1901 is the first year of the climate data set.

From this equilibrium, five factorial simulation runs are simulated where key drivers of plant growth and carbon sequestration  $(CO_2, climate, nitrogen deposition, O_3)$  are simulated either as fixed to the reference year, or transient (i.e. progressively changing within the simulation period) (see Tab. 1). These simulations run from the year 1850 to 2099. The period up to the

**Table 1.** Forcing setting of the factorial runs. Climate forcing for the years prior to 1901 is always drawn from the same random sequence of years between 1901 and 1930.

Factorial run	$\mathrm{CO}_2$	Climate	Nitrogen deposition	$O_3$	
<b>S</b> 1	<del>1850-2099</del> _1850_2099	<del>1901-1930</del> _1901-1930	1850	1850	
S2	<del>1850-2099_1850_2099</del>	<del>1901-1930_1901-1930</del>	1850	<del>1850-2099_1850_2099</del>	
<b>S</b> 3	<del>1850-2099_1850_2099</del>	<del>1901-2099_1901_2099</del>	1850	<del>1850-2099_</del> 1850_2099	
S4	<del>1850-2099_1850_2099</del>	<del>1901-2099_1901_2099</del>	<del>1850-2099_1850_2099</del>	1850	
S5	<del>1850-2099_1850_2099</del>	<del>1901-2099</del> - <u>1901</u> -2099	<del>1850-2099_1850_2099</del>	<del>1850-2099_1850_2099</del>	

195 year 2005 is simulated identical for both <u>RCPsRCP scenarios</u>. From 2006 until 2099 simulations are run using the forcing according to either the RCP2.6 or the RCP8.5 forcing scenario (Moss et al., 2010; van Vuuren et al., 2011).

To investigate the impact of the ozone  $O_3$  deposition scheme on the simulation results, the factorial runs are repeated with a model version where the ozone  $O_3$  deposition scheme is turned off (see ATM model version in (Franz et al., 2017). In simulations where the ozone  $O_3$  deposition module is turned – off the canopy ozone  $O_3$  concentration equals the  $O_3$  concentration at 45 m above the surface which is the hight of the lowest level of the atmospheric chemistry transport model (CTM) that deliver

200

45 m above the surface which is the <u>hight of the</u> lowest level of the <u>atmospheric chemistry transport model (CTM) that deliver</u> the forcing for our runs hereforcing data.

# 2.5 Factorial analysis

The impact of a single forcing driver on the simulation results can be approximated by subtracting the simulation results of suitable combination of factorial runs from one another (see Tab.2). In the following, the term 'forcing driver' is used to refer
to the input variables of the conducted simulations and 'single driver' refers to the approximated impact of a single forcing driver on the simulation results. The described approach is an approximation of the impact of the single drivers and assumes that the drivers effect effects on the analysed output variables is are additive. The assumption of additive effects is a necessary simplification to restrict the number of simulations and computation time (Zaehle et al., 2010). For O<sub>3</sub>, a main driver of interest, two different approaches to calculate the single driver were realised. In one approach, the O<sub>3</sub> impact is calculated from the
two factorial runs with only one/ two transient drivers (S1 and S2), and a second time from the factorial runs where all and all but one driver (S5 and S4 respectively) are simulated transient. The comparison of these two approaches to calculate the single drivers of interacting forcing drivers on the estimate of the O<sub>3</sub> single driver. The relative changes between two simulation runs SX and SY are calculated as (SX – SY)/SY.

# 3 Results

215 The simulations show a strong increase in gross primary production (GPP) in the Northern Hemisphere ( $\geq 30^{\circ}$ N) between the year 1850 and 2099 with all forcings considered in this study (S5; Fig. 2). There is a notable difference between the scenarios,

**Table 2.** Calculation of the single driver effects (CO<sub>2</sub>, climate, nitrogen deposition, O<sub>3</sub>) from the conducted simulations.  $S1_{ref}$  refers to the mean of the years 1850 to 1859 of the S1 simulation. The relative changes between simulation SX and SY reported in Section 3 are ealculated as (SX - SY)/SY See Tab. 1 for info on the forcing setting of the factorial runs S1–S5.

Attributed single driver	Simulations		
$\mathrm{CO}_2$	$S1 - S1_{ref}$		
O <sub>3</sub> approach 1	S2-S1		
$O_3$ approach 2	S5-S4		
Climate	S3-S2		
Nitrogen deposition	S5-S3		

with smaller simulated changes for the RCP 2.6 scenario, which level of at an increase by about a third. GPP continues to increase In simulations based on the RCP8.5 scenario, GPP increases throughout the  $21^{st}$  centuryunder the RCP8.5 scenario, roughly doubling by the year 2099 relative to 1850 values by the year 2099. Reflecting the changes in productivity, . In the RCP

220 2.6 scenario, the simulated increase in GPP levels off in the 2040s at approximately a third of the simulated increase at the end of the  $21^{st}$  century in the RCP8.5 scenario. The changes in terrestrial carbon storage also increases reflect these changes in GPP (see Fig. 2d).

Concurrent with the strong increase in GPP is an increased foliar uptake of  $\frac{1}{0}$ , which parallels the increase in tropospheric near surface and canopy-air  $O_3$  concentrations (Fig.3). Simulated changes in the cumulative  $O_3$  uptake without a flux threshold (CUO0, see Methods) strongly follow changes in the O3 concentrations during the entire simulation

- period. Accounting for the ability of leaves to detoxify part of the  $\frac{1}{2}$  of the  $\frac{1}{2}$  by  $O_3$  taken up, the cumulative canopy  $O_3$  uptake above a flux threshold of  $1 \text{ nmol m}^{-2} \text{ s}^{-1}$  (CUO1) does not remain at relative-relatively constant values during the  $21^{st}$  century under RCP8.5. Instead, it reaches a maximum at the end of the 20<sup>th</sup> century and steadily declines afterwards. This difference between CUO0 and CUO1 for simulations based on RCP8.5 implies that the frequency at which the detoxification threshold is exceed gradually declines during the  $21^{st}$  century. This results from a decline in peak uptake rates  $(F_{st})$  in the  $21^{st}$  century 230
- (Fig.3b) despite fairly constant tropospheric near surface  $O_3$  concentrations (Fig.3a). The decline in peak uptake rates is the consequence of the reduced ratio of stomatal conductance to net photosynthesis under high atmospheric  $CO_2$ .

#### 3.1 Factorial Analysis

225

We next decompose the simulations into the effects of the different model drivers with a special focus on the effects of  $\Theta_3$ O<sub>3</sub> and nitrogen deposition. In all five factorial runs the simulated GPP increases strongly between 1850 and 2099 and ap-235 proximately doubles for the run S5 based on RCP8.5 (see Fig. 2a). The primary cause for this simulated increase is the  $CO_2$ fertilisation effect induced by increasing atmospheric  $CO_2$  concentrations (see Fig. 4c and Tab. 3). Climate change is the second most import-important factor for the simulated increase, whereas the positive effect of N deposition is less pronounced. Ozone injury causes a modest decrease in productivity, which manifests strongest during the 1990s. During the 20<sup>th</sup> century



**Figure 2.** Simulated Northern hemispheric gross primary production (GPP), vegetation carbon (vegetation-C), soil carbon, and net land C flux of the factorial run S5 (all variables are simulated transient, see Tab. 1). Displayed are the period of 1850-2099 1850-2099 for both the RCP2.6 and RCP8.5. See also Tab. 3.

240 the decline gradually reverses. The land carbon sink strongly responds to elevated levels of  $CO_2$  (see Fig. 4f), whereas climate change induces a varying impact on the land carbon sink. During the second half of the  $21^{st}$  century, the effect of climate mainly causes a reduction in the simulated land carbon sink.

#### 3.2 Magnitude of nitrogen deposition impact

Nitrogen deposition has a positive effect on the simulated carbon uptake, storage and ozone  $O_3$  uptake and accumulation in plants, but the magnitude varies between the different scenarios. N deposition increases simulated increased summed regional GPP by about 2.1 % (0.7 PgC yr<sup>-1</sup>) in the present at the beginning of the 21<sup>st</sup> century compared to pre-industrial values (see Fig. 5, Appendix Supplement Fig. S3, and Tab. ??.4). At the end of the 21<sup>st</sup> century, GPP is increased by approximately 2.5 % under both RCPs.

Carbon stored in vegetation (vegetation-C) is increased by nitrogen deposition by about 2 % (2.1 PgC) in the present at the beginning of the 21<sup>st</sup> century and by approximately 3 % (4.4 -6.1 and 6.1 PgC yr<sup>-1</sup> for RCP2.6 and RCP8.5 respectively) at the end of the 21<sup>st</sup> century. This positive effect of N deposition keeps growing in China East Asia until the of the 21<sup>st</sup> century (see Tab. ??). However, 4), whereas in Europe and the USAtemperate North America, the GPP and vegetation-C enhancement by nitrogen deposition reduces at the end of the 21<sup>st</sup> century compared to values simulated during the middle of



**Figure 3.** Simulated canopy  $O_3$  concentration, ozone  $O_3$  uptake ( $F_{st}$ ), cumulative  $O_3$  uptake without a flux threshold (CUO0) and cumulative  $O_3$  uptake above a flux threshold of 1 nmol m<sup>-2</sup> s<sup>-1</sup> (CUO1) of the factorial run S5 (all forcing variables are simulated transient). Displayed are monthly values (solid lines) and smothed values (broken line) where the effect of the seasonal cycle is smoothed by the application of a moving average of 12 month.Brown lines: RCP2.6, red lines: RCP8.5

the peaks around the mid  $21^{st}$  century and declines thereafter. The soil-C is less affected by nitrogen deposition and maximal increases of  $\approx 1 \%$  (6 PgC) compared to pre-industrial values are simulated at the end of the  $21^{st}$  century. Nitrogen deposition stimulates enhances the simulated land carbon sink (land C flux) the strongest most in the period between 1950 and 2050 by 5-25% (-0.02-..., -0.15 PgC yr<sup>-1</sup>) compared to pre-industrial values.

Nitrogen deposition steadily increases plant nitrogen uptake until the middle of the 21<sup>st</sup> century, with maximum simulated increases of about 40 TgNH<sub>4</sub>yr<sup>-1</sup> or 9% (see Fig. 6 and Appendix Supplement Fig. S4). Concurrently, N deposition increases
260 N<sub>2</sub>O emissions by 0.7 TgN<sub>2</sub>Oyr<sup>-1</sup> (20%) and NH<sub>4</sub> leaching of 10 TgNH<sub>4</sub>yr<sup>-1</sup> (100%) compared to pre-industrial values are simulated. NO<sub>3</sub> leaching increases until the end of the 20<sup>th</sup> century by 6 TgNO<sub>3</sub>yr<sup>-1</sup> (80%) compared to pre-industrial values and declines again thereafter afterwards.

#### 3.3 Magnitude of ozone O<sub>3</sub> deposition impact

#### 3.3.1 Ozone uptake

Projections of  $\frac{1}{02000}O_3$  uptake and damage are primarily controlled by the scenarios of tropospheric near surface  $O_3$  (Fig. 4a,b). However, foliar ozone uptake is reduced by increasing atmospheric concentrations because of its effect on the relationship



**Figure 4.** Single drivers obtained by subtracting factorial runs for selected output variables. Displayed are the results for simulated regional mean ozone-O<sub>3</sub> uptake ( $F_{st}$ ), regional mean cumulative canopy O<sub>3</sub> uptake above a flux threshold of 1 nmol m<sup>-2</sup> s<sup>-1</sup> (CUO1), regional summed GPP, regional summed stocks of total carbon biomass (vegetation-C), soil organic matter carbon (soil-C), and summed land carbon flux (land C flux) for simulations based on RCP2.6 and RCP8.5. The effect of the seasonal cycle is smoothed by the application of a moving average of 12 months(a,b).

between stomatal conductance and net photosynthesis. The effect of can also be seen in reduction of CUO1 during the  $21^{st}$  century shown in Fig. 3. Lower stomatal conductances reduce  $F_{st}$  and CUO1, even if the concentrations slightly increase in simulations based on RCP8.5.

Nitrogen deposition slightly increases  $F_{st}$  by maximal 0.008 and 0.01 for RCP2.6 and RCP8.5, respectively (see Fig. 5a). This relates to an increase in CUO1 of about 1 %. Climate change increases stomatal ozone uptake about 0.04 during the  $21^{st}$  century and considerably stronger compared to -deposition.

The simulated mean ozone  $O_3$  uptake  $(F_{st})$  due to changes in near surface  $O_3$  concentration increases by approximately 0.27 nmol m<sup>-2</sup> s<sup>-1</sup> (70 %) between the pre-industrial period and the year 2000 (see Fig. 7a). Under the RCP8.5 scenario,  $F_{st}$ 



**Figure 5.** Nitrogen deposition induced absolute change in regional mean ozone- $O_3$  uptake ( $F_{st}$ ), mean cumulative  $O_3$  uptake above a flux threshold of 1 nmol m<sup>-2</sup> s<sup>-1</sup> (CUO1), and change in % -change-in summed GPP, summed carbon biomass (vegetation-C), summed carbon soil organic matter (soil-C), and summed land carbon flux (land C flux) compared to pre-industrial values in the simulation region. The nitrogen deposition induced change is calculated from the simulation runs S3 and S5 (see Tab. 2). Solid lines indicate results from simulations based on RCP8.5, dotted lines results from simulations based on RCP2.6. The effect of the seasonal cycle is smoothed by the application of a moving average of 12 months (a,b).

- 275 increases until the end of the  $21^{st}$  century, reaching more than a 90% increase compared pre-industrial times. Conversely, in simulations based on RCP2.6,  $F_{st}$  declines strongly and by the end of the  $21^{st}$  century comparable values to simulations based on pre-industrial O<sub>3</sub> concentrations are reached. The mean CUO1 increases by approximately 2.3 mmol O<sub>3</sub> m<sup>-2</sup> until the year 2000 and strongly declines until the end of the  $21^{st}$  century (see Fig. 7b). In simulations based on RCP2.6 the CUO1 values reach comparable values to simulations based on pre-industrial O<sub>3</sub> concentrations by 2099.
- However, foliar O<sub>3</sub> uptake is reduced by increasing atmospheric CO<sub>2</sub> concentrations because of its effect on the relationship between stomatal conductance and net photosynthesis. The effect of CO<sub>2</sub> can also be seen in reduction of CUO1 during the  $21^{st}$  century shown in Fig. 3. Lower stomatal conductances reduce  $F_{st}$  and CUO1, even if the O<sub>3</sub> concentrations slightly increase in simulations based on RCP8.5. Nitrogen deposition slightly increases  $F_{st}$  by at most 0.008 nmol m<sup>-2</sup> s<sup>-1</sup> and



**Figure 6.** Nitrogen deposition induced change in % -change-in regional summed total N uptake by region,  $N_2O$  emission,  $NH_4$  leaching and  $N_2O$ - $NO_3$  leaching compared to pre-industrial values in the simulation region. The nitrogen deposition induced change in % -change is calculated from the simulation runs S3 and S5 (see Tab. 2). Solid lines indicate results from simulations based on RCP8.5, dotted lines results from simulations based on RCP8.6.

0.01 nmol m<sup>-2</sup> s<sup>-1</sup> for RCP2.6 and RCP8.5, respectively (see Fig. 5a). This relates to an increase in CUO1 of about 1 %.
 285 Climate change increases stomatal O<sub>3</sub> uptake about 0.04 nmol m<sup>-2</sup> s<sup>-1</sup> during the 21<sup>st</sup> century and considerably stronger compared to N-deposition.

The two different approaches to assess the contribution of  $O_3$  to the simulated trends in the carbon cycle based on analysing alternative combinations of model drivers (see Tab. 2) yield similar but not identical results (see Fig. 7). Typically, the differences between the two approaches do not exceed 1 % except for CUO1, where larger relative changes occur for small absolute changes (see Fig. 7b).

#### 3.3.2 Ozone damage

290

295

In the period of 1970-1990, i.e. the time of the peak increase on tropospheric ozone near surface  $O_3$  concentrations, the simulated detrimental effects of  $O_3$  on photosynthesis GPP nearly completely counteract the positive effect of rising  $CO_2$  concentrations (see Fig. 4c). The negative impact of  $O_3$  on GPP shows a maximum approximately in the 1990s at approximately -1.5 PgC yr<sup>-1</sup> (4 %) compared to pre-industrial values (see Fig. 7c, Appendix Supplement Fig. S5 and Tab. 4).

In the subsequent decades, the simulated ozone  $O_3$  induced reduction in GPP declines to 1 % by the end of the 21<sup>st</sup> century for RCP8.5 and to close to zero for RCP2.6.

During the period 2000 to 2030, increasing dominates the change in GPP. After that time, GPP stagnates at 2030 levels due

Due to the stabilisation of atmospheric  $CO_2$  in RCP 2.6 the RCP2.6 scenario, the increase in GPP levels off at 2040s levels,

300

but continues to rise under RCP 8.5 with increasing  $CO_2$ . The growth-stimulating effect of N-deposition is smaller than the negative impact induced by  $O_3$  during the  $20^{th}$  century (see Fig. 4c). This pattern is reversed during the course of the  $21^{st}$  century (see Section 3.4).

The  $O_3$  effect on GPP propagates to vegetation and thus considerably affects the simulated above- and below-ground biomass total carbon biomass in vegetation (vegetation-C), and to a limited extend extend also soil-C storage. In the simulations with transient  $O_3$  (S2,S3,S5), the regionally integrated vegetation-C vegetation-C ceases to grow in the 1950s for 30-50 vears

- transient  $O_3$  (S2,S3,S5), the regionally integrated vegetation-C vegetation-C ceases to grow in the 1950s for 30-50, 30-50 years (see in Fig. 2b), causing a loss of carbon storage compared to the simulations without increasing  $O_3$  of about 8 Pg C (8 %). Despite the declining effect of  $O_3$  on GPP, vegetation-C vegetation-C remains reduced for much of the first half of the  $21^{st}$ century and only recovers very slowly thereafter afterwards (see Fig. 4c and d). The strongest ozone- $O_3$  induced reduction in mean simulated vegetation-C in the simulation area occurs in the period of 2000-2020-2020 at approximately 10 % (see
- 310 Fig. 7d and Appendix Supplement Fig. S5). The ozone O<sub>3</sub> effect on vegetation-C declines to 5 % by 2099 for RCP8.5 and 4 % for RCP2.6 (see Tab. 4). The soil-C is less strongly impacted by O<sub>3</sub> with simulated maximal reductions of less than 2 % (10 Pgc).

 $O_3$  impact on land C storage (land C flux) peaks at the end of the  $20^{th}$  century. In the 1990s, the effect of  $O_3$  on the land sink is about 0.4 Pg C yr<sup>-1</sup> or approx. 20 %. During the  $21^{st}$  century, the ozone  $O_3$  effect on the land C flux is reversed and

315 eventually becomes positive. In the 2090s, the simulated land C flux is increased by about 4 to 7% for RCP8.5 and 16 % for RCP2.6. This seemingly counter-intuitive effect is the result of lower ozone-induced O<sub>3</sub>-induced net primary production, which reduces the formation of soil carbon. The resulting lower stock in soil carbon in simulations accounting for ozone-O<sub>3</sub> damage results in lower increases in heterotrophic respiration due to climate change during the  $21^{st}$  century, which causes the reversal of the O<sub>3</sub> effect on the land C sink.

# 320 3.3.3 Regional patterns

The simulated cumulative  $O_3$  uptake above a flux threshold of 1 nmol m<sup>-2</sup> s<sup>-1</sup> (CUO1) shows a strong geographic variation. Highest values of CUO1 are simulated during the 1990s in the eastern and north-eastern US, large parts of Europe centralCentral Europe, and eastern Asia (see Appendix Supplement Fig. S6a). Evergreen trees accumulate  $O_3$  damage over several years, because of the longer life time of their leaves compared to deciduous trees. This can result in high values of

325 <u>CUO1, even if  $O_3$  concentrations are moderate</u>. At the end of the  $21^{st}$  century simulated CUO1 values reach comparable values to pre-industrial times in large parts of the simulation region and slightly lower values in large parts of the US and Eurasia in simulations based on RCP2.6.

The highest ozone  $O_3$  induced absolute reductions in GPP occur in Europe, Eastern US and Eastern Asia where the respective increase in CUO1 is highest. <u>Peak During the 1990s</u>, peak reductions of about <u>150-220 gCm<sup>2</sup>yr<sup>-1</sup></u> (<u>8-11 8-11</u> %)

are simulated in the eastern US, southern Europe and eastern Asia during the decade of 1990 (see Fig. 8). Simulated ozone induced damage to GPP declines  $O_3$  induced reductions in GPP decline in the decades of 2040 and 2090 for both RCPs ,

Table 3. Absolute and relative change in GPP, total carbon biomass in vegetation (vegetation-C), soil organic matter carbon (soil-C) and land carbon flux (land C flux) ir
deposition (Ndep), and O <sub>3</sub> concentrations. The differences in GPP, vegetation-C, soil-C and Land C flux are presented for simulations of the past years of 1850 to 2005
2009, and for the entire simulation period 1850 to 2099. The reported change refers to the change between the last and the first year of the respective time periods. The giver
the O <sub>3</sub> impact.

RCP and time span	$CO_2$	$CO_2$	Clima
GPP	$[PgC yr^{-1}]$	[%]	[PgCyr
Past 1850:2005	3.4	11.5	3.1
RCP2.6 1850:2099	6.3	21.5	5.2
RCP8.5 1850:2099	16.3	55.6	12.9
RCP2.6 2006:2099 2:47.3 37.8 0.3 0.4 1.21.5 3.64.2 RCP8.5 2006:2099 12.4 37.4 12.9 28.4 0.8 0.5 0.9 2.83.2 vegetation-C	[PgC]	[%]	[PgC
Past 1850:2005	18.3	23	4.9
RCP2.6 1850:2099	42.7	53.7	20.2
RCP8.5 1850:2099	83.8	105.4	41.9
RCP2.6 2006:2099 24.6 25.2 14.9 11.2 2.3 1 4.45 5.56.1 RCP8.5 2006:2099 65 66.1 37.5 22.1 4 0.8 0.10.6 44.9 soil-C	[PgC]	[%]	[PgC
Past 1850:2005	23.2	4.4	-2.2
RCP2.6 1850:2099	64.8	12.2	-8.4
RCP8.5 1850:2099	100.5	18.9	-37.6
RCP2.6 2006:2099 41.77.5 -6.3 -1 3.4 0.6 2.62.8 0.50.6 RCP8.5 2006:2099 76.9 13.8 -35.5 -5.6 3.6 0.10.6 0.3 0.2 Land C flux	$[\rm PgCyr^{-1}]$	[%]	[PgCyr
Past 1850:2005	-0.5	72.9	-0.4
RCP2.6 1850:2099	0.3	-39.1	-1
RCP8.5 1850:2099	-1.3	201.9	-1.1
RCP2.6 2006:2099 1.2 -76 -1.3 205.7 0.1 -5.1 -0.30.4 29.445.5RCP8.5 2006:2099 -0.5 30.8 -1.9 116 0 -12.8 -0.20.3 15.232.6			



**Figure 7.** Ozone induced absolute change in regional mean ozone-O<sub>3</sub> uptake ( $F_{st}$ ), mean cumulative O<sub>3</sub> uptake above a flux threshold of 1 nmol m<sup>-2</sup> s<sup>-1</sup> (CUO1) and change in % -change in summed GPP, summed total carbon biomass in vegetation (vegetation-C), summed carbon soil organic matter (soil-C), and summed land carbon flux (land C flux) compared to pre-industrial values in the simulation region. Different colors indicate different approaches to calculate the ozone-O<sub>3</sub> induced change from the factorial runs. Orange lines represent approach 1: (S2-S1)/S1, blue lines approach 2:(S5-S4)/S4. Solid lines indicate results from simulations based on RCP8.5, dotted lines results from simulations based on RCP2.6. The effect of the seasonal cycle is smoothed by the application of a moving average of 12 months.

but considerable ozone and in many regions, while considerable  $O_3$  induced reductions in GPP are simulated until the end of the 21<sup>st</sup> century in eastern Asia. In the decade of 2040-2040s under the RCP8.5 scenario relative reductions in GPP of 4-8-4-8 % are simulated in southern Europe, parts of the eastern and western US in simulations based on RCP8.5. Peak USA, while peak relative decreases of 8-11-8 – 11 % are simulated in eastern Asia. At Towards the end of the 21<sup>st</sup> century ozone-O<sub>3</sub> induced reductions in GPP continue to decline, but reductions of above 8 % are still simulated in small parts of eastern Asia. Slight increases in GPP are simulated in a large fraction of the Eastern US and small scattered areas in Asia.

335

Simulations based on RCP2.6 indicate for the end of the 21<sup>st</sup> century close to no ozone O<sub>3</sub> induced damage compared to pre-industrial values over large parts of the simulation scopedomain. Small absolute reductions are simulated in parts of Europe
 and small absolute increases are simulated in the Eastern US induced by lower CUO1 values compared to pre-industrial values

17

(see Appendix-Supplement Fig. S6). Increased atmospheric  $CO_2$  concentrations compared to pre-industrial values reduce the stomatal conductance, restrict ozone  $O_3$  uptake and enable the increase in GPP values.

As expected,  $O_3$  impacts on vegetation-C and soil-C peak later compared to GPP (see Fig. 7c-e and Tab. 4). For both scenarios, the strongest ozone  $O_3$  induced absolute reductions in vegetation-C of 1400–1600 gCm<sup>2</sup> occur in the

345 decade of 2040 in the eastern US, southern Europe and eastern Asia (see Fig. 10). For both RCPs the ozone  $O_3$  induced vegetation-C reductions exceed 20% in parts of Europe, eastern and western US and eastern Asia in the middle of the 21<sup>st</sup> century. By the end of the 21<sup>st</sup> century the ozone  $O_3$  induced vegetation-C reduction attenuates in these hotspots for both RCPs, though stronger in simulations based on RCP2.6.

# Ozone impacts on vegetation-and soil-peak later compared to GPP (see Fig. 7c-e and Tab. 4).

**Table 4.** Mean percent change in GPP, total carbon biomass in vegetation (vegetation-C), and land carbon flux (land C flux) induced by ozone  $O_3$  during the decades of 1990 (1990–1999), 2040 (2040–20492040–2049) and 2090 (2090–20992090–2099) compared to pre-industrial values for the Northern Hemisphere north of 30°N (NH30), Europe, USA and China. The given range indicates the estimates according to both approaches to calculate the ozone  $O_3$  impact. The spread in the effect sizes due to inter-annual variability, derived from error propagation of the yearly estimates is displayed in Supplement Tab. S1.

Region	Pollutant	1990	2040 RCP8.5	2040 RCP2.6	2090 RCP8.5	2090 RCP2.6
GPP						
NH30	$O_3$	-3.84.3	-2.02.8	-1.32.0	-0.71.0	00.2
NH30	Ndep	1.8	2.7	2.3	2.5	2.4
Europe	$O_3$	-4.54.9	-2.22.6	-1.41.9	-0.8	-0.20.3
Europe	Ndep	2.7	3.8	<u>2.9</u>	<u>2.9</u>	2.5
USA	$O_3$	-4.75.0	-2.12.7	-1.72.1	-0.81.1	0.31.0
USA	Ndep	1.4	1.1	0.9	0.6	0.9
China	$O_3$	-9.210.1	-7.810.8	-7.18.6	-1.62.8	-3.85.7
China	Ndep	2.9	5.5	<u>6.1</u>	<u>6.4</u>	7
vegetation-Vegetation-C						
NH30	$O_3$	-8.58.9	-8.48.8	-7.47.6	-5.15.4	-3.83.9
NH30	Ndep	1.8	3.3	3	<u>3.2</u>	3.2
Europe	$O_3$	-10.811.5	-9.910.6	-8.89.3	-6.16.4	-4.9
Europe	Ndep	3.2	4.7	4.2	3.6	<u>4</u>
USA	$O_3$	-11.912.5	-10.811.7	-9.710.2	-6.56.8	-4.14.3
USA	Ndep	1.6	1.8	1.7	1.5	1.3
China	$O_3$	-15.115.9	-26.329.3	-22.924.4	-15.818.5	-16.216.4
China	Ndep	1.6	2.8	$\overset{4}{\sim}$	<u>3.9</u>	<u>6.2</u>
Land CLand C flux						
NH30	$O_3$	-20.721.2	02.2	6.27.4	3.56.9	15.716.2
NH30	Ndep	9.7	5	<u>4.9</u>	1.6	3
Europe	$O_3$	-23.725.6	0.41.7	99.9	4.615.7	1517.1
Europe	Ndep	15.6	5.9	<u>4.4</u>	- <u>6.7</u>	-0.2
USA	$O_3$	-18.420.4	0.61.1	8.310.1	2.96.4	16.419.9
USA	Ndep	4.3	0.7	-0.7	-1.9	0.9
China	$O_3$	-58.862.9	-7.312.8	-1.11.7	11.324.8	24.930.7
China	Ndep	24.6	9.9 19	13.8	14.1	15.4

#### 3.4 Comparative impact of N deposition and O<sub>3</sub>

The magnitude of ozone the  $O_3$  induced damage on GPP exceeded the growth stimulating effect induced by nitrogen deposition over large parts of the 20th century and until the beginning of the  $21^{st}$  century (see Fig. 4c). Contrary to the tropospheric In contrast to the near surface  $O_3$  concentrations, the regional mean nitrogen deposition does not decline during the  $21^{st}$  century but slightly increases in RCP8.5 and RCP2.6. The growth stimulating effect on GPP induced by nitrogen deposition becomes

but slightly increases in RCP8.5 and RCP2.6. The growth stimulating effect on GPP induced by nitrogen deposition becomes higher in magnitude during the  $21^{st}$  century compared to the detrimental effect of ozone O<sub>3</sub> (see Fig. 4c, <u>Supplement Fig. S8</u> and Tabs. 4 and ??4).

The growth stimulating effect of nitrogen deposition on vegetation-C remains lower in magnitude compared to the detrimental effects of ozone for both pollution  $O_3$  for both emission scenarios throughout the entire simulation period (see Fig. 4d

and Tab. 3). However, in simulations based on RCP2.6 the ozone  $O_3$  induced reduction on vegetation-C is at the end of the  $21^{st}$  century only slightly higher in magnitude compared to the growth stimulating effect induced by nitrogen deposition (see Tabs. 4and ??).

The extend extent of simulated impact of ozone O<sub>3</sub> and nitrogen deposition on the terrestrial carbon uptake (GPP) and storage (vegetation-C) differs strongly within the simulated region. Nitrogen deposition stimulates GPP compared to simulations run with pre-industrial deposition values mainly in Europe and Eastern Asia. Simulated increases of GPP in these regions constitute about 80-140 gC m<sup>2</sup> yr<sup>-1</sup> for simulations run based on RCP8.5 (see left column in Fig. 8). In relative terms, peak increases of 10-16-10-16 % are found in parts of eastern, central and northern Asia and small parts of Europe (see Appendix Supplement Fig. S7). Simulated increases in GPP are higher, and hotspot areas more extended, in the decade of 2090 compared to the 2040 decade for both RCPs (see Supplement Fig. S8). Simulations based on RCP2.6 exhibit similar patterns compared to simulations based on RCP8.5, but show a less strong increase in GPP induced by to nitrogen deposition.

Nitrogen deposition induces peak increases in vegetation-C of  $\frac{500-600}{500-600}$  gC m<sup>2</sup> compared to pre-industrial values in parts of Europe and eastern Asia (see Fig. 10). Highest relative increases in vegetation-C of  $\frac{14-17-14-17}{14-17}$ % are simulated in the decades of 2040 and 2090 in regions of southern and northern Asia, where absolute changes are mostly small.

Mean percent change in GPP, vegetation-, and land carbon flux (land flux) induced by nitrogen deposition during the decades

of 1990 (1990-1999), 2040 (2040-2049) and 2090 (2090-2099) compared to pre-industrial values for the Northern Hemisphere north of 30°N (), Europe, USA and China. Region 1990 2040 RCP8.5 2040 RCP2.6 2090 RCP8.5 2090 RCP2.6 GPP 1.8 2.7 2.3 2.5 2.4 Europe 2.7 3.8 2.9 2.9 2.5 USA 1.4 1.1 0.9 0.6 0.9 China 2.9 5.5 6.1 6.4 7-vegetation-1.8 3.3 3 3.2 3.2 Europe 3.2 4.7 4.2 3.6 4 USA 1.6 1.8 1.7 1.5 1.3 China 1.6 2.8 4 3.9 6.2-

Land Cflux 9.7 5 4.9 1.6 3Europe 15.6 5.9 4.4 - 6.7 - 0.2 USA 4.3 0.7 - 0.7 - 1.9 0.9 China 24.6 9.9 13.8 14.1 15.4

# 380 **3.5** Impact of the ozone-O<sub>3</sub> deposition scheme

Simulations run with a model version where the ozone deposition scheme is turned offresult in considerably higher estimates of  $F_{st}$  and CUO1, leading to higher damage estimates (see Fig. 9). In these simulations In the simulations presented in the previous sections, O-CN was applied with its O<sub>3</sub> deposition scheme turned on. To test the impact of the application of the O<sub>3</sub> deposition scheme on the estimated ozone damage, we reran the simulations with the O<sub>3</sub> deposition scheme turned off. In
simulations where the O<sub>3</sub> deposition scheme is turned off the O<sub>3</sub> is assumed to enter leaves directly without accounting for the turbulent transport in between the lower troposphere and the leaves, as well as the deposition and destruction of ozone O<sub>3</sub> on other surfaces. Turning off the O<sub>3</sub> deposition scheme result in considerably higher estimates of F<sub>st</sub> and CUO1, resulting in higher damage estimates (see Fig. 9). In simulations where the ozone O<sub>3</sub> deposition scheme is turned off. Part and CUO1, resulting O<sub>3</sub>-induced reductions in GPP and vegetation-C are approximately twice as high compared to simulations where the ozone O<sub>3</sub> deposition scheme is turned on. Peak reductions of GPP amount 3 PgC yr<sup>-1</sup> (≈8%) compared to approximately 1.5 PgC yr<sup>-1</sup> (≈4%) in simulations where the deposition scheme is turned off the O<sub>3</sub> constitute 25 PgC (≈11%) in simulations where the deposition scheme is turned off and 10 PgC (≈5%) in simulations where the deposition scheme is turned off and 10 PgC (≈5%) in simulations where the deposition scheme is turned off the ozone O<sub>3</sub> the ozone O<sub>3</sub> deposition scheme is turned off the ozone O<sub>3</sub> the ozone O<sub>3</sub> deposition scheme is turned off the ozone O<sub>3</sub> the ozone O<sub>3</sub> deposition scheme is turned off the ozone O<sub>3</sub> deposition scheme is turned driven by the RCP8.5 scenario. As for GPP and vegetation-C, the omission of the ozone O<sub>3</sub> deposition scheme causes roughly a doubling of the simulated damage to carbon

 $\ensuremath{$  soil organic matter (soil-C) and summed land carbon flux (land C flux).

# 4 Discussion

The simulations of the Northern Hemisphere biosphere from 1850-2099-1850-2099 according to the representative concentration pathway scenarios RCP8.5 and RCP2.6 2.6 and 8.5 indicate that air pollution (ozone O<sub>3</sub> and nitrogen deposition) may have considerably affected carbon uptake and plant growth in the past and has the potential to will continue to have a considerable impact during the  $21^{st}$  century. We simulate an ozone-

400 imp

In our simulations, nitrogen deposition stimulates the simulated land carbon sink of the Northern Hemisphere  $\geq 30^{\circ}$ N the strongest in the period between 1950 and 2050 by 5–25 % (0.02... 0.15 PgC yr<sup>-1</sup>) compared to pre-industrial values.

These values are broadly consistent with a meta-analysis by Schulte-Uebbing and de Vries (2018), who estimated that nitrogen stimulates the global land carbon sink in above ground and below ground woody biomass by 0.112-0.243 PgCyr<sup>-1</sup>.

- 405 Global carbon storage in forests was estimated to increase about 0.27  $PgCyr^{-1}$  induced by N deposition for the period 1997—2013 in simulations based on RCP4.5 (Wang et al., 2017). Thomas et al. (2010) found that above-ground biomass increment increased by 40 % compared to pre-industrial conditions in the northeastern and north-central USA during the 1980s and 1990s, from which they estimate that N deposition stimulates global tree carbon storage by 0.3 PgCyr<sup>-1</sup>.
- Our simulations indicate an  $O_3$  induced reduction in the land C flux of  $0.4 \text{ PgCyr}^{-1}$  in the 1990s, decade of 1990. During the  $21^{st}$  century the ozone  $O_3$  effect on the land C flux is reversed and becomes positive. This is caused by lower increases of soil respiration due to climate change as a result of ozone-induced  $O_3$ -induced declines of net primary production and thus litterfall. This highlights the importance of investigating interactive processes on longer time scales together to get a better understanding of their net effect on the land carbon sink.

The stimulating effect of nitrogen deposition on regional mean GPP and biomass is lower in magnitude compared to the detri-415 mental effect of O<sub>3</sub> during most of the simulation period for both RCPs (results for RCP2.6 not shown). Both effects approximately even out in their impact on the mean regional GPP by 2030–2050. By the end of the 21<sup>st</sup> century nitrogen deposition stronger increases GPP than O<sub>3</sub> impacts decline it. However, regions that experience strong ozone O<sub>3</sub> induced negative effects do not always coincide with regions that benefit from the stimulating effect of nitrogen deposition (see Supplement Fig. S8).

- 420
- During the  $21^{st}$  century the cumulative  $O_3$  uptake above a flux threshold of  $1 \text{ nmol } \text{m}^{-2} \text{ s}^{-1}$  (CUO1), on which the damage calculations base, declines due to the impact of the  $CO_2$  fertilisation effect on stomatal conductance and ozone- $O_3$  uptake. This result is in agreement with Oliver et al. (2018), who found in Europe-wide simulations that elevated future  $CO_2$  levels and reductions in  $O_3$  concentrations result in reduced  $O_3$  induced damage values by 2050. Induced by the simulated decline in CUO1 the mean regional reduction in GPP deceases decreases in the decade of 2050 to approximately 2 % in simulations based on RCP8.5 and 1-1.5 % in simulations based on RCP2.6. By the end of the 21st century damage induced by elevated 425 levels of O<sub>3</sub> decreases to approximately 1 % in simulations based on RCP8.5 and close to zero for RCP2.6. Simulations with the JULES model estimate a 8–15 % reduction in global GPP between 1901–2100 (Sitch et al., 2007). A more recent version of the JULES model suggest a 4–9 % reduction in European GPP due to  $\frac{1}{02000}$  between 1901 and 2050 (Oliver et al., 2018). Both estimates are higher compared to the simulation results here (see Tab. 4).

Our estimates of the impact of  $\frac{1}{2}$  on the land C sink is smaller than that by Oliver et al. (2018), who simulated

- 430 an ozone  $O_3$  induced reduction of the land C sink by -0.7-... -1.3 PgC in the decade of 1970. The simulated detrimental ozone-O<sub>3</sub> effect declines in the following decades to -0.3-... -0.5 PgC in the period of 2002-2011. A possible reason for the higher estimates by Sitch et al. (2007) and Oliver et al. (2018) is the absence of an ozone  $O_3$  deposition scheme in JULES, what might have caused higher surface  $\frac{0}{2000}$  concentrations and hence increased  $\frac{0}{2000}$ , uptake and incurred damage. The tropospheric near surface  $O_3$  concentrations used in the simulations here to force the model are provided by CTMs which
- 435 report  $O_3$  concentrations in a height of approximately 45 m above the surface. The ozone- $O_3$  deposition scheme included into O-CN uses the  $O_3$  concentration of the free atmosphere troposphere to calculate the  $O_3$  concentration at canopy level.

If the  $O_3$  concentration provided by the CTMs from the forcing is used as if being at canopy level the O-CN model simulates a higher ozone-O<sub>3</sub> uptake. Following this twice as high damage values to GPP and vegetation-C are calculated compared to simulations where the deposition scheme is applied to calculate the canopy level  $O_3$  concentration. This highlights the importance of using canopy level O3 concentrations to calculate ozone-O3 uptake and damage to prevent a considerable 440 overestimation of  $\frac{1}{2}$  over  $O_3$  induced damage.

#### 4.1 Air pollution impacts on GPP and total carbon biomass

The average ozone-O<sub>3</sub> effect on GPP in the Northern Hemisphere ( $\geq 30^{\circ}$ N) increases until the decade of 19901990s, when GPP is reduced by approximately 4 % compared to the pre-industrial period. Regional hotspots in southern Europe, eastern Asia and the eastern US exhibit ozone-O3 induced reductions of 8–11 % for the decade of 1990. In a meta-analyses by Wittig 445 et al. (2009) net photosynthesis damage of trees grown in ambient  $O_3$  concentrations vs. charcoal filtered air is estimated to amount 11 % and 19 % for trees grown in elevated  $O_3$  concentrations vs. charcoal filtered air. Lombardozzi et al. (2013) estimates damage to net photosynthesis of temperate deciduous trees to amount 12 % and 16 % for temperate evergreen trees. A reduction of 28 % in net photosynthesis is estimated for woody plants grown in elevated O<sub>3</sub> compared to a control by Li

et al. (2017). Simulated  $\frac{1}{2}$  of  $O_3$  damage values in hotspot areas are close to the lower damage estimates suggested by Wittig 450

et al. (2009) and Lombardozzi et al. (2013), while the regional means including many areas with low  $O_3$  exposure, results in lower average ozone  $O_3$  damage than estimated by these meta-analyses.

Several process based models estimated ozone-process-based models estimated  $O_3$  induced damage to NPP/GPP GPP or net primary production (NPP) on global or regional scale: a mean global ozone- $O_3$  induced reduction in NPP of 0.8–2.9 %

- 455 from 1989 to 1993 is estimated by the Terrestrial Ecosystem Model (Felzer et al., 2005). Simulations with the Community Land Model suggest a 10.8 % reduction of global mean GPP for present day O<sub>3</sub> concentrations (Lombardozzi et al., 2015). A mean reduction in NPP of 4.5 % in China between 1961-2000-1961-2000 is estimated by a process-based Dynamic Land Ecosystem Model (Ren et al., 2007). The simulation of ozone-O<sub>3</sub> damage to China's forests suggest a 0.2–1.6 % decrease in NPP from the 1960s to 2000-05-2005 (Ren et al., 2011). Simulations using the Terrestrial Ecosystem Model estimate a
- 460 mean reduction in NPP of 2.6–6.8 % in the United States for the period of the late 1980s to early 1990s (Felzer et al., 2004). The YIBS model simulates a 4–8 % damage to GPP due to  $O_3$  in the eastern US and 8–17 % damage in hot spots for the decade of 1998–2007 (Yue and Unger, 2014). In the Euro-Mediterranean region a reduction in GPP of 22 % is estimated for the year 2002 by the ORCHIDEE model (Anav et al., 2011). The mean GPP of the years 2001–2010 in Europe is simulated to be reduced by 7.6 % compared to not accounting for ozone  $O_3$  damage by the O-CN model (Franz et al., 2017). Here, on a
- regional mean basis ozone  $O_3$  induced reductions of about 4 % are simulated at the end of the 20<sup>th</sup> century and beginning of the 21<sup>st</sup> century compared to pre-industrial values. At the end of the 21<sup>st</sup> century close to zero ozone  $O_3$  induced reductions in GPP are simulated by O-CN here. An exception are hot spots like Eastern Asia where peak decreases of more than 8 % are simulated for both RCPs at the end of 21<sup>st</sup> century. Our damage estimates here are lower compared to at least most of the previous estimates suggested by biosphere models.
- 470 The ozone induced simulated mean In our simulations, the  $O_3$  induced regional reduction in total above- and below-ground carbon biomass carbon biomass in vegetation (vegetation-C) reaches peak values of 8–10% at the end of the 20<sup>th</sup> and first half of the 21<sup>st</sup> century. Damage values of 20–23% are simulated in damage hotspots in southern Europe, eastern Asia and the eastern and western US for the decade of 1990 (see Appendix Supplement Fig. S9). A meta-analyses conducted with trees suggests a 7% reduction in total biomass for trees grown in ambient air compared to charcoal filtered air and a 17% reduction
- for trees grown in elevated  $O_3$  concentrations compared to charcoal filtered air (Wittig et al., 2009). In a meta-analyses by Li et al. (2017) a 14 % reduction in total biomass is calculated for trees grown in elevated  $O_3$  concentrations (mean of 116 ppb) compared to controls grown in a mean  $O_3$  concentration of 21 ppb. The simulated regional mean estimate of ozone  $O_3$  induced damage to vegetation-C is higher compared to the estimate of trees grown in ambient vs. charcoal filtered air by Wittig et al. (2009) and lower compared to trees grown in elevated  $O_3$  vs. charcoal filtered air or a mean of 21 ppb  $O_3$  (Wittig et al., 2009;
- 480 Li et al., 2017). Simulated damage values in the hotspots are higher compared to the estimates by the meta-analyses. Our simulated declines in ozone- $O_3$  induced damage to GPP and vegetation-C during the 21<sup>st</sup> century generally agree with simulated reductions in potential threat to vegetation by Klingberg et al. (2014).

Klingberg et al. (2014) report that by 2050 the ozone O<sub>3</sub> exposure index AOT40 (Accumulated exposure Over a Threshold of 40 ppb O<sub>3</sub>) is projected to decrease over wide areas of Europe below critical levels defined by the EU directive 2008/50/EC
and the LRTAP convention in simulations of a chemical transport model (CTM) the chemistry transport model MATCH driven

by the RCP4.5 emission scenario. The Their simulations suggest that the more physiological based ozone  $O_3$  damage index POD1 (Phytotoxic Ozone Dose above a threshold of  $1 \mod m^{-2} s^{-1}$ ) is projected to decline declines as well, however to a lesser extend extent compared to the AOT40 index and not below critical levels defined for forest trees (Klingberg et al., 2014).

An ensemble of six global atmospheric chemistry transport models project improvements of the AOT40 index in the Northern

490

Hemisphere by 2099 under the RCP2.6 and RCP4.5, while critical levels continue to be exceeded over many areas (Sicard et al., 2017). By 2099 the potential impact of O<sub>3</sub> on photosynthesis and carbon assimilation is projected to decline by 61 % under the RCP2.6 scenario, by 47<sup>%</sup> under RCP4.5 and increase by 70<sup>%</sup> under the RCP8.5 scenario compared to the early 2000s (Sicard et al., 2017).

### 4.2 Interactive effects of O<sub>3</sub> and CO<sub>2</sub>

- Elevated levels of CO<sub>2</sub> (eCO<sub>2</sub>) have the potential to induce stomatal closure (Paoletti and Grulke, 2005) what might limit O<sub>3</sub> uptake and damage. Contradictory evidence exists showing that either eCO<sub>2</sub> ameliorated the negative effects of O<sub>3</sub> on plants (Barnes and Pfirrmann, 1992; Broadmeadow and Jackson, 2000; Isebrands et al., 2001; Riikonen et al., 2004) or that there was little interaction between both gases and the stimulating effect of eCO<sub>2</sub> on NPP persisted (Talhelm et al., 2014; Zak et al., 2011). Results from the Aspen FACE indicate that stomatal conductance and ozone-O<sub>3</sub> uptake were not reduced by eCO<sub>2</sub> in their experiment (Uddling et al., 2010), and that ozone-O<sub>3</sub> fumigation completely offset the growth enhancement observed in
  - the  $eCO_2$  treatment for <u>ozone</u>  $O_3$  sensitive and tolerant clones (Karnosky et al., 2003).

Several studies find species specific positive or negative impacts of  $+eCO_2$  and elevated levels of  $O_3$  (eO<sub>3</sub>) on photosynthesis (Noormets et al., 2001), growth (Isebrands et al., 2001) and biomass (King et al., 2005). An amplification of the negative effects of  $O_3$  under  $eCO_2$  on leaf chlorophyll content, nitrogen content and electron transport capacity ( $J_{max}$ ) was observed in ozone

- $O_3$  sensitive and tolerant aspen clones (Noormets et al., 2010). A possible reason for the amplification of ozone- $O_3$  induced negative effects under  $eCO_2$  is a possible down regulation or suppression of antioxidant production under  $eCO_2$  and hence increased injury (Wustman et al., 2001; Karnosky et al., 2003). All in all, a clear picture of the joint effects of  $eCO_2 + eO_3$  on plants or plant groups is still lacking.
- Terrestrial biosphere models often assume a tight coupling between net photosynthesis and stomatal conductance what which
  induces stomatal closure in case of simulated eCO<sub>2</sub> and restricts O<sub>3</sub> uptake and damage (Felzer et al., 2004, 2005; Sitch et al., 2007; Oliver et al., 2018; Yue and Unger, 2014). For example Sitch et al. (2007) simulated a 6–9 % reduction in O<sub>3</sub> induced damage to GPP due to elevated levels of CO<sub>2</sub> and a 5–10 % reduction in the ozone-related effect on land carbon storage between the years 1901 and 2100. 2100 in a scenario with strongly rising atmospheric CO<sub>2</sub>. Oliver et al. (2018) simulated a 1–2 % decrease in O<sub>3</sub> induced damage to GPP and land carbon storage caused by elevated levels of CO<sub>2</sub> between 1901 and 2050. The largest simulated impact of ozone O<sub>3</sub> on the land carbon sink occurred during the 20<sup>th</sup> century when the atmospheric tropospheric O<sub>3</sub> concentration rose quickly (Oliver et al., 2018). During the 21th century simulated O<sub>3</sub> concentrations changed less and the simulated elevated levels of CO<sub>2</sub> restricted O<sub>3</sub> O<sub>3</sub> uptake and induced damage (Oliver et al., 2018). This agrees well with our findings here that ozone O<sub>3</sub> induced damage increases from pre-industrial times until the end of the 20<sup>th</sup> century

(GPP) or beginning of the  $21^{st}$  century (vegetation-C) and afterwards decreases again (see Fig. 7).

However, the very simplistic simulation of reduced ozone O<sub>3</sub> uptake and incurred damage induced by eCO<sub>2</sub> does not mirror all the effects observed in field experiments (Wustman et al., 2001; Karnosky et al., 2003; Noormets et al., 2010). Similar to other terrestrial biosphere models, O-CN does not account for observed effects like an exacerbation of ozone O<sub>3</sub> induced damage due to eCO<sub>2</sub> (Wustman et al., 2001; Karnosky et al., 2003) or unaltered rates of stomatal conductance and O<sub>3</sub> uptake under eCO<sub>2</sub> (Uddling et al., 2010). Following this the presented low values of simulated future ozone O<sub>3</sub> damage represent a possible future scenario under the assumption that the large majority of plants react to the the combined exposure to elevated levels of CO<sub>2</sub> and O<sub>3</sub> by a reduced stomatal uptake of O<sub>3</sub> and reduced incurred damage.

#### 4.3 Limitations of comparisons between publications

When interpreting the comparison of the results here our results and previously published simulation results one has to keep in mind that the different modelling approaches usually set-ups and approaches differ in several aspects that might considerably impact considerably affect the damage estimate. Models often apply different injury functions which relate ozone O<sub>3</sub> uptake to plant damage (Lombardozzi et al., 2012, 2015; Franz et al., 2017; Oliver et al., 2018). However, injury functions have the potential to induce considerable over- or underestimation of simulated biomass damage compared to measured damage values (Franz et al., 2018). Simulations often differ in the simulated time period time period covered, e.g. Sitch et al. (2007) (1901-21001901-2100), Lombardozzi et al. (2015) 25 years with an average O<sub>3</sub> concentration of the years 2002-20092002-2009, Franz et al. (2017) (1961-2011)961-2011), and Oliver et al. (2018) (1901-20501901-2050). They differ in also differ in the forcing considered e.g. the representation of changing CO<sub>2</sub> concentrations, nitrogen deposition and land-

- cover/ land-use change. For instance, Sitch et al. (2007) simulate changing  $CO_2$  concentrations, Lombardozzi et al. (2015) do include neitherwhile Lombardozzi et al. (2015) does not, Franz et al. (2017) account for changing  $CO_2$  concentrations, nitrogen deposition but use static land-cover (kept fixed at 2005 levels), and Oliver et al. (2018) simulate changing  $CO_2$  con-
- centrations and a partly fixed land-cover, but no effect of N deposition. Furthermore damage estimates are calculated based on different references reference periods and conditions. Damage might be given as the difference between a simulation accounting for O<sub>3</sub> damage compared to a reference simulation not accounting for ozone O<sub>3</sub> damage (Lombardozzi et al., 2015; Franz et al., 2017). Another approach is to report the damage simulated between a specific time period. Sitch et al. (2007) calculate ozone O<sub>3</sub> induced damage between 1901-2100-1901-2100 and Oliver et al. (2018) between 1901-2001 and 2001-20501901-2001
  and 2001-2050.
  - Different modelling studies apply differing pollution emission scenarios, e.g. IPCC SRES (Sitch et al., 2007) and the RCP scenarios used here, what which might impact simulated ozone  $O_3$  uptake and incurred damage. The application of the IPCC SRES scenarios(, which assume a large increase in  $O_3$  precursor emissions) results in a simulated , implies an increase in annual global mean surface  $O_3$  concentrations by 4-6-4-6 ppb (Wild et al., 2012). Contrary to this, the application of the
- 550 RCP scenarios (Moss et al., 2010; van Vuuren et al., 2011) in 14 global chemistry transport models results in the projection of declining annual global mean surface O<sub>3</sub> concentrations of as much as 2 ppb by 2050 in most regions of the globe except South Asia where increases are simulated (Wild et al., 2012). Lower projected ozone-induced damage in our study compared to Sitch et al. (2007) is therefore also a consequence of the assumed scenario.

Turnock et al. (2020) found that the CMIP6 models overestimate observed surface O<sub>3</sub> concentrations by up to 16 ppb across
 most regions of the globe. This will likely lead to a general overestimation of simulated O<sub>3</sub> damage by terrestrial biosphere models. However, the ozone deposition scheme included into O-CN has the potential to ameliorate this observed discrepancy. The calculation of canopy level O<sub>3</sub> concentrations from the lowest level O<sub>3</sub> concentrations of the forcing data are lower and thus probably closer to the observations.

A further important difference between the published results is the time resolution of the ozone  $O_3$  forcing applied in the simulations. Some studies used hourly ozone  $O_3$  forcing (e.g. Lombardozzi et al. (2015), Franz et al. (2017), and Oliver et al. (2018)) and others are forced by monthly diurnal mean values (e.g. Sitch et al. (2007) and the simulations here). As the formation of ozone  $O_3$  shows a pronounced diurnal cycle (Sanz et al., 2007), the use of monthly mean ozone  $O_3$  concentrations probably impacts the simulated estimates of ozone  $O_3$  uptake. However, to which extend extent the omission of a diurnal cycle impacts ozone  $O_3$  uptake, accumulation and damage estimates is yet uncertain.

#### 565 4.4 Limits to the parameterisation of ozone O<sub>3</sub> damage in O-CN

570

Plants can activate defence mechanism and physiological pathways to produce protective compounds like ascorbate and polyamines which can detoxify at least part of the ozone taken up (Kangasjärvi et al., 1994; Kronfuß et al., 1998; Tausz et al., 2007). In the simulations conducted here we account for detoxification by introducing a flux threshold but do not account for the cost to produce protective compounds like antioxidants due to the lack of suitable data. This could potentially introduce a bias bias towards underestimating damage to GPP if the leaf-injury parameterisations are based on leaf-level data.

Ozone sensitivity is known to differ between plant groups, plant species and between genotypes (Wittig et al., 2007; Lombardozzi et al., 2013; Li et al., 2017; Hayes et al., 2007; Karnosky et al., 2003). The assumed injury function is a key aspect of the simulation of ozone- $O_3$  damage and has a large impact on the <u>extend extent</u> of the estimated damage (Franz et al., 2018). However, the scarcity of suitable data restricts the possibility to parameterise injury functions for all simulated PFTs (e.g. 12

- 575 PFTs in O-CN), let alone a variation of the ozone-sensitivity  $O_3$ -sensitivity within PFTs. Furthermore it restricts the evaluation of ozone-submodels  $O_3$ -submodels and the included injury functions. The injury functions used for the simulations here are tuned to reproduce observed biomass damage from filtration/fumigation experiments of broadleaved and needle-leaved tree species (Franz et al., 2018). The simulations are restricted to the Northern Hemisphere  $\geq 30^{\circ}N$  to limit the domain of simulation to temperate/boreal forests and thus similar species as used for the tuning of the injury functions. Due to the lack of
- 580 suitable damage functions for grass species we here applied the damage functions developed to match damage to trees. This induces a bias in the damage estimates and will likely results in an underestimation of simulated damage for example for the crop plant functional types.

The biomass damage experiments used to parameterise the injury function are-in O-CN were conducted with young trees grown in monocultures. The common attempt to estimate responses of adult trees grown under natural conditions by the extrap-

olation of results from short-term experiments with young trees is subject to several issues, e.g. due to the differing environmental conditions and changing ozone- $O_3$  sensitivities with increasing tree size or age (Schaub et al., 2005; Cailleret et al., 2018) (Schaub et al., 2005; Cailleret et al., 2018; Franz et al., 2018). It is yet uncertain if the simulation of injury to photosynthesis based on experiments with young trees can be transferred to adult trees to obtain realistic biomass damage estimates.

Differing ozone O3 sensitivities can induce changes in community composition (Barbo et al., 1998; Kubiske et al., 2007; Zak

- 590 et al., 2011) as well as the interactive effects of changed  $CO_2$  and  $O_3$  concentrations (Karnosky et al., 2003). The responses of plants grown under interspecific competition, e.g. in forests, may not be transferred from results of filtration/fumigation experiments (with elevated  $CO_2$  and/or  $O_3$ ) of plants grown in monoculture (Kozovits et al., 2005). For instance, Zak et al. (2011) found that initial declines in forest productivity induced by elevated levels of  $O_3$  were compensated for by the growth of ozone  $O_3$  tolerant individuals resulting in an equivalent NPP between ambient and elevated levels of  $O_3$ . Simulations by an
- individual-based forest model indicate that  $O_3$  damage might not reduce the carbon sequestration capacity in forests might not be reduced by ozone damage if at the ecosystem level of forests if the reduced carbon fixation of ozone-sensitive  $O_3$ -sensitive species is compensated for by an by increased carbon fixation of less ozone-sensitive species  $O_3$ -sensitive species at the ecosystem level (Wang et al., 2016). The simulation of community dynamics is limited in O-CN, as it does not account for species, and therefore acclimation processes at the ecosystem level are not accounted for. The effect of interspecific competition
- 600 on  $\frac{1}{2}$  on  $\frac{1}{2}$

The application of present day land-use information fixed to the year 2000 in our simulations may affect simulated trends of GPP, canopy conductance and biomass production in regions where land cover and/or land-use have historically changed or are projected to change during the scenario period. This can lead to a discrepancy in the simulated effect of nitrogen deposition

and  $O_3$  damage. For example  $O_3$  damage differs between plant functional types and a shift to highly productive crops would results in an increase in damage.

Holding the N fertiliser application at the year 2000 levels in our simulations here imposes a bias on the simulated GPP, biomass production and  $O_3$  damage in regions where fertiliser application changed. Regions where fertiliser application decreased would show a reduction in growth stimulation along with a reduction in  $O_3$  damage. Regions exposed to increases in fertiliser application would exhibit a stimulation in growth along with an increase in  $O_3$  damage.

The simulations conducted here are run offline and following this atmosphere and biosphere do not feedback on one another. Forcing variables like  $O_3$  concentrations and nitrogen deposition are provided by a different model than the climate. This imposes an inconsistency between the biosphere, climate and the abundance of the air pollutants whose formation depends on climate variables. This contributes to unavoidable inconsistencies between the atmospheric forcing and the land fluxes when

615 making offline simulations compared to a simulation with a fully coupled Earth System Model. However, these limitations, do not invalidate the simulated sensitivity of the land carbon cycle simulation to the forcing applied.

# 5 Conclusion

610

 $O_3$  damage considerably reduced simulated carbon uptake (GPP) and storage (vegetation-C) in the simulation area where the maximal impact occurs at the end of the 20<sup>th</sup> century and beginning of the 21<sup>st</sup> century respectively. The detrimental ozone- $O_3$ 

- 620 impact declines during the  $21^{st}$  century and reaches mean regional reductions of 0-1% for GPP and 4-5% for vegetation-C by the end of the  $21^{st}$  century compared to pre-industrial values. However, in hotspots decreases in GPP of more than 8%(eastern Asia) and decreases in vegetation-C of more than 15% (parts of Europe, eastern and western US and eastern Asia) are simulated at the end of the  $21^{st}$  century. Nitrogen deposition increases GPP less than  $O_3$  impacts decrease it for most of the simulated period. The increasing effect of nitrogen deposition on vegetation-C is lower compared to the decreasing effect of
- $O_3$  for the entire simulation period. Accounting for the stimulating effects of nitrogen deposition but omitting the detrimental effect of  $O_3$  can lead to an over estimation of carbon uptake and storage.

*Acknowledgements.* The research leading to this publication was supported by the EU Framework programme through grant no. 282910 (ECLAIRE) and the Max Planck Society for the Advancement of Science e.V. through the ENIGMA project. This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement no. 647204; QUINCY).

630



**Figure 8.** Absolute change in GPP compared to pre-industrial values induced by nitrogen deposition (left column) and ozone-O<sub>3</sub> calculated according to approach 2 (right column). The induced change in GPP is displayed for the decades 1990 (mean of the years 1990–1999), 2040 (mean of the years 2040–20492040–2049) and 2090 (mean of the years 2090–20992090–2099). For the decades 2040 and 2090 results from simulations based on RCP8.5 and RCP2.6 are displayed. See Tab. 2 for details on the calculation of the single drivers.



**Figure 9.** Ozone impacts on the regional mean ozone  $O_3$  uptake  $(F_{st})$ , mean cumulative  $O_3$  uptake above a flux threshold of 1 nmol m<sup>-2</sup> s<sup>-1</sup> (CUO1), summed GPP, summed total carbon biomass in vegetation (vegetation-C), summed carbon soil organic matter (soil-C), and summed land carbon flux (land C flux) compared to pre-industrial values in the simulation region. The displayed ozone  $O_3$  impact is calculated based on approach 2. Orange lines: Results based on a model version where the ozone  $O_3$  deposition scheme is turned on. Magenta lines: Results based on a model version where the ozone  $O_3$  deposition scheme is from simulations based on RCP8.5, dotted lines results from simulations based on RCP2.6. The effect of the seasonal cycle is smoothed by the application of a moving average of 12 months (a,b).



**Figure 10.** Absolute change in the total carbon biomass in vegetation (vegetation-C) compared to pre-industrial values induced by nitrogen deposition (left column) and ozone  $O_3$  calculated according to approach 2 (right column). The induced change in the total carbon biomass is displayed for the decades 1990 (mean of the years 1990-1999), 2040 (mean of the years 2040-20492040-2049) and 2090 (mean of the years 2090-20992090-2099). For the decades 2040 and 2090 results from simulations based on RCP8.5 and RCP2.6 are displayed. See Tab. 2 for details on the calculation of the single drivers.

#### References

- Ainsworth, E. A. and Long, S. P.: What have we learned from 15 years of free-air CO<sub>2</sub> enrichment (FACE)? A meta-analytic review of the responses of photosynthesis, canopy properties and plant production to rising CO<sub>2</sub>, New Phytologist, 165, 351–372, https://doi.org/10.1111/j.1469-8137.2004.01224.x, https://nph.onlinelibrary.wiley.com/doi/abs/10.1111/j.1469-8137.2004.01224.x, 2005.
- 635 Ainsworth, E. A., Yendrek, C. R., Sitch, S., Collins, W. J., and Emberson, L. D.: The Effects of Tropospheric Ozone on Net Primary Productivity and Implications for Climate Change, Annual Review of Plant Biology, 63, 637–661, 2012. Akimoto, H.: Global Air Ouality and Pollution, Science, 302, 1716–1719, 2003.

Anav, A., Menut, L., Khvorostyanov, D., and Viovy, N.: Impact of tropospheric ozone on the Euro-Mediterranean vegetation, Global Change Biology, 17, 2342–2359, 2011.

- 640 Ball, J., Woodrow, I., and Berry, J.: A Model Predicting Stomatal Conductance and its Contribution to the Control of Photosynthesis under Different Environmental Conditions, in: Prog. Photosynthesis Res. Proc. Int. Congress 7th, Providence. 10-15 Aug 1986, edited by Biggins, J., pp. 221–224, Progress in Photosynthesis Research, 1987.
  - Barbo, D., Chappelka, A., Somers, G., Miller-Goodman, M., and Stolte, K.: Diversity of an early successional plant community as influenced by ozone, The New Phytologist, 138, 653–662, 1998.
- 645 Barnes, J. and Pfirrmann, T.: The influence of *CO*<sub>2</sub> and *O*<sub>3</sub>, singly and in combination, on gas exchange, growth and nutrient status of radish (*Raphanus sativus* L.), New Phytologist, 121, 403–412, 1992.
  - Broadmeadow, M. S. and Jackson, S.: Growth responses of *Quercus petraea*, *Fraxinus excelsior* and *Pinus sylvestris* to elevated carbon dioxide, ozone and water supply, The New Phytologist, 146, 437–451, 2000.
  - Büker, P., Feng, Z., Uddling, J., Briolat, A., Alonso, R., Braun, S., Elvira, S., Gerosa, G., Karlsson, P., Le Thiec, D., Marzuoli, R., Mills, G.,
- 650 Oksanen, E., Wieser, G., , Wilkinson, M., and Emberson, L.: New flux based dose-response relationships for ozone for European forest tree species, Environmental Pollution, 206, 163–174, https://doi.org/10.1016/j.envpol.2015.06.033, http://tinyurl.sfx.mpg.de/ug3q, 2015. Cailleret, M., Ferretti, M., Gessler, A., Rigling, A., and Schaub, M.: Ozone effects on European forest growth—Towards an integrative

approach, Journal of Ecology, 106, 1377-1389, 2018.

Cionni, I., Eyring, V., Lamarque, J.-F., Randel, W., Stevenson, D., Wu, F., Bodeker, G., Shepherd, T., Shindell, D., and Waugh, D.: Ozone

- database in support of CMIP5 simulations: results and corresponding radiative forcing, Atmospheric Chemistry and Physics, 11, 11267– 11292, 2011.
- Cooper, O. R., Parrish, D., Ziemke, J., Balashov, N., Cupeiro, M., Galbally, I., Gilge, S., Horowitz, L., Jensen, N., Lamarque, J.-F., Naik, V., Oltmans, S., Schwab, J., Shindell, D., Thompson, A., Thouret, V., Wang, Y., and Zbinden, R.: Global distribution and trends of tropospheric ozone: An observation-based review, Elementa: Science of the Anthropocene, 2, 000029, https://doi.org/10.12952/journal.elementa.000029, 2014.
- Dufresne, J.-L., Foujols, M.-A., Denvil, S., Caubel, A., Marti, O., Aumont, O., Balkanski, Y., Bekki, S., Bellenger, H., Benshila, R., Bony, S., Bopp, L., Braconnot, P., Brockmann, P., Cadule, P., Cheruy, F., Codron, F., Cozic, A., Cugnet, D., de Noblet, N., Duvel, J.-P., Ethé, C., Fairhead, L., Fichefet, T., Flavoni, S., Friedlingstein, P., Grandpeix, J.-Y., Guez, L., Guilyardi, E., Hauglustaine, D., Hourdin, F., Idelkadi, A., Ghattas, J., Joussaume, S., Kageyama, M., Krinner, G., Labetoulle, S., Lahellec, A., Lefebvre, M.-P., Lefevre, F., Levy, C., Li, Z. X.,
- 665 Lloyd, J., Lott, F., Madec, G., Mancip, M., Marchand, M., Masson, S., Meurdesoif, Y., Mignot, J., Musat, I., Parouty, S., Polcher, J., Rio, C., Schulz, M., Swingedouw, D., Szopa, S., Talandier, C., Terray, P., Viovy, N., and Vuichard, N.: Climate change projections using the IPSL-CM5 Earth System Model: from CMIP3 to CMIP5, Climate Dynamics, 40, 2123–2165, 2013.

Felzer, B., Kicklighter, D., Melillo, J., Wang, C., Zhuang, O., and Prinn, R.: Effects of ozone on net primary production and carbon sequestration in the conterminous United States using a biogeochemistry model, Tellus B, 56, 230-248, 2004.

- 670 Felzer, B., Reilly, J., Melillo, J., Kicklighter, D., Sarofim, M., Wang, C., Prinn, R., and Zhuang, Q.: Future Effects of Ozone on Carbon Sequestration and Climate Change Policy Using a Global Biogeochemical Model, Climatic Change, 73, 345–373, 2005.
  - Feng, Z. and Kobayashi, K.: Assessing the impacts of current and future concentrations of surface ozone on crop yield with meta-analysis, Atmospheric Environment, 43, 1510–1519, 2009.
  - Fiscus, E., Booker, F., and Burkey, K.: Crop responses to ozone: uptake, modes of action, carbon assimilation and partitioning, Plant, Cell &

675 Environment, 28, 997-1011, 2005.

> Franz, M., Simpson, D., Arneth, A., and Zaehle, S.: Development and evaluation of an ozone deposition scheme for coupling to a terrestrial biosphere model, Biogeosciences, 14, 45–71, https://doi.org/10.5194/bg-14-45-2017, http://www.biogeosciences.net/14/45/2017/, 2017.

> Franz, M., Alonso, R., Arneth, A., Büker, P., Elvira, S., Gerosa, G., Emberson, L., Feng, Z., Le Thiec, D., Marzuoli, R., Oksanen, E., Uddling, J., Wilkinson, M., and Zaehle, S.: Evaluation of simulated ozone effects in forest ecosystems against biomass damage estimates from

680 fumigation experiments, Biogeosciences, 15, 6941-6957, https://doi.org/10.5194/bg-15-6941-2018, https://www.biogeosciences.net/15/ 6941/2018/, 2018.

Friend, A.: Modelling canopy CO<sub>2</sub> fluxes: are 'big-leaf'simplifications justified?, Global Ecology and Biogeography, 10, 603–619, 2001.

Fusco, A. and Logan, J.: Analysis of 1970–1995 trends in tropospheric ozone at Northern Hemisphere midlatitudes with the GEOS-CHEM model, J. Geophys. Res. 108, 1988-1997, 2003.

- Galloway, J. N., Dentener, F. J., Capone, D. G., Boyer, E. W., Howarth, R. W., Seitzinger, S. P., Asner, G. P., Cleveland, C. C., Green, 685 P., Holland, E. A., Karl, D., Michaels, A., Porter, J., Townsend, A., and Vöosmarty, C.: Nitrogen Cycles: Past, Present, and Future, Biogeochemistry, 70, 153-226, 2004.
  - Grantz, D., Gunn, S., and VU, H.: O<sub>3</sub> impacts on plant development: a meta-analysis of root/shoot allocation and growth, Plant, Cell & Environment, 29, 1193-1209, 2006.
- 690 Hayes, F., Jones, M., Mills, G., and Ashmore, M.: Meta-analysis of the relative sensitivity of semi-natural vegetation species to ozone, Environmental Pollution, 146, 754-762, 2007.

Hempel, S., Frieler, K., Warszawski, L., Schewe, J., and Piontek, F.: A trend-preserving bias correction-the ISI-MIP approach, Earth System Dynamics, 4, 219-236, 2013.

Hurtt, G. C., Chini, L. P., Frolking, S., Betts, R. A., Feddema, J., Fischer, G., Fisk, J. P., Hibbard, K., Houghton, R. A., Janetos, A., Jones,

695 C. D., Kindermann, G., Kinoshita, T., Klein Goldewijk, K., Riahi, K., Shevliakova, E., Smith, S., Stehfest, E., Thomson, A., Thornton, P., van Vuuren, D. P., and Wang, Y. P.: Harmonization of land-use scenarios for the period 1500-2100: 600 years of global gridded annual land-use transitions, wood harvest, and resulting secondary lands, Climatic Change, 109, 117, https://doi.org/10.1007/s10584-011-0153-2, https://doi.org/10.1007/s10584-011-0153-2, 2011.

Isebrands, J., McDonald, E., Kruger, E., Hendrey, G., Percy, K., Pregitzer, K., Sober, J., and Karnosky, D.: Growth responses of Populus 700 tremuloides clones to interacting elevated carbon dioxide and tropospheric ozone, Environmental Pollution, 115, 359–371, 2001.

- Kangasjärvi, J., Talvinen, J., Utriainen, M., and Karjalainen, R.: Plant defence systems induced by ozone, Plant, Cell & Environment, 17. 783-794, 1994.
  - Karnosky, D. F., Zak, D. R., Pregitzer, K. S., Awmack, C. S., Bockheim, J. G., Dickson, R. E., Hendrey, G. R., Host, G. E., King, J. S., Kopper, B. J., Kruger, E. L., Kubiske, M. E., Lindroth, R. L., Mattson, W. J., Mcdonald, E. P., Noormets, A., Oksanen, E., Parsons, W. F. J., Percy,
- 705 K. E., Podila, G. K., Riemenschneider, D. E., Sharma, P., Thakur, R., Sôber, A., Sôber, J., Jones, W. S., Anttonen, S., Vapaavuori, E.,

Mankovska, B., Heilman, W., and Isebrands, J. G.: Tropospheric  $O_3$  moderates responses of temperate hardwood forests to elevated  $CO_2$ : a synthesis of molecular to ecosystem results from the Aspen FACE project, Functional Ecology, 17, 289–304, 2003.

- King, J., Kubiske, M., Pregitzer, K., Hendrey, G., McDonald, E., Giardina, C., Quinn, V., and Karnosky, D.: Tropospheric O<sub>3</sub> compromises net primary production in young stands of trembling aspen, paper birch and sugar maple in response to elevated atmospheric CO<sub>2</sub>, New Phytologist, 168, 623–636, 2005.
- Klingberg, J., Engardt, M., Karlsson, P., Langner, J., and Pleijel, H.: Declining risk of ozone impacts on vegetation in Europe 1990–2050 due to reduced precursor emissions in a changed climate, Biogeosciences Discussions, 11, 625–655, 2014.
- Kozovits, A. R., Matyssek, R., Blaschke, H., Göttlein, A., and Grams, T. E.: Competition increasingly dominates the responsiveness of juvenile beech and spruce to elevated  $CO_2$  and/or  $O_3$  concentrations throughout two subsequent growing seasons, Global Change Biology,
- 715 11, 1387–1401, 2005.

710

- Krinner, G., Viovy, N., de Noblet-Ducoudré, N., Ogée, J., Polcher, J., Friedlingstein, P., Ciais, P., Sitch, S., and Prentice, I.: A dynamic global vegetation model for studies of the coupled atmosphere-biosphere system, Global Biogeochemical Cycles, 19, GB1015, https://doi.org/10.1029/2003GB002199, 2005.
- Kronfuß, G., Polle, A., Tausz, M., Havranek, W., and Wieser, G.: Effects of ozone and mild drought stress on gas exchange, antioxidants
- 720 and chloroplast pigments in current-year needles of young Norway spruce [*Picea abies* (L.) Karst.], Trees-Structure and Function, 12, 482–489, 1998.
  - Kubiske, M., Quinn, V., Marquardt, P., and Karnosky, D.: Effects of Elevated Atmospheric CO<sub>2</sub> and/or O<sub>3</sub> on Intra-and Interspecific Competitive Ability of Aspen, Plant Biology, 9, 342–355, 2007.
- Kull, O. and Kruijt, B.: Leaf photosynthetic light response: a mechanistic model for scaling photosynthesis to leaves and canopies, Functional Ecology, 12, 767–777, 1998.
  - Laisk, A., Kull, O., and Moldau, H.: Ozone Concentration in Leaf Intercellular Air Spaces is Close to Zero, Plant Physiology, 90, 1163–1167, 1989.
  - Lamarque, J. F., Bond, T. C., Eyring, V., Granier, C., Heil, A., Klimont, Z., Lee, D., Liousse, C., Mieville, A., Owen, B., Schultz, M. G., Shindell, D., Smith, S. J., Stehfest, E., Van Aardenne, J., Cooper, O. R., Kainuma, M., Mahowald, N., McConnell, J. R., Naik, V., Riahi,
- K., and van Vuuren, D. P.: Historical (1850-2000) gridded anthropogenic and biomass burning emissions of reactive gases and aerosols: methodology and application, Atmos. Chem. Physics, 10, 7017–7039, https://doi.org/10.5194/acp-10-7017-2010, 2010.
  - Lamarque, J.-F., Dentener, F., McConnell, J., Ro, C.-U., Shaw, M., Vet, R., Bergmann, D., Cameron-Smith, P., Dalsoren, S., Doherty, R., Faluvegi, G., Ghan, S. J., Josse, B., Lee, Y. H., MacKenzie, I. A., Plummer, D., Shindell, D. T., Skeie, R. B., Stevenson, D. S., Strode, S., Zeng, G., Curran, M., Dahl-Jensen, D., Das, S., Fritzsche, D., and Nolan, M.: Multi-model mean nitrogen and sulfur deposition from the
- 735 Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP): evaluation of historical and projected future changes, Atmospheric Chemistry and Physics, 13, 7997–8018, https://doi.org/10.5194/acp-13-7997-2013, https://acp.copernicus.org/articles/13/ 7997/2013/, 2013.
  - Langebartels, C., Kerner, K., Leonardi, S., Schraudner, M., Trost, M., Heller, W., and Sandermann Jr, H.: Biochemical Plant Responses to Ozone: I. Differential Induction of Polyamine and Ethylene Biosynthesis in Tobacco, Plant Physiology, 95, 882, 1991.
- 740 LeBauer, D. and Treseder, K.: Nitrogen limitation of net primary productivity in terrestrial ecosystems is globally distributed, Ecology, 89, 371–379, 2008.
  - Leisner, C. and Ainsworth, E.: Quantifying the effects of ozone on plant reproductive growth and development, Global Change Biology, 18, 606–616, 2012.

Li, P., Feng, Z., Catalayud, V., Yuan, X., Xu, Y., and Paoletti, E.: A meta-analysis on growth, physiological and biochemical responses of

- woody species to ground-level ozone highlights the role of plant functional types, Plant, Cell & Environment, 40, 2369–2380, 2017.
  - Lombardozzi, D., Levis, S., Bonan, G., and Sparks, J.: Predicting photosynthesis and transpiration responses to ozone: decoupling modeled photosynthesis and stomatal conductance, Biogeosciences, 9, 3113–3130, 2012.
  - Lombardozzi, D., Sparks, J. P., and Bonan, G.: Integrating O<sub>3</sub> influences on terrestrial processes: photosynthetic and stomatal response data available for regional and global modeling, Biogeosciences, 10, 6815–6831, https://doi.org/10.5194/bgd-10-6973-2013, http://www.
- 750 biogeosciences-discuss.net/10/6973/2013/, 2013.
  - Lombardozzi, D., Levis, S., Bonan, G., Hess, P., and Sparks, J.: The Influence of Chronic Ozone Exposure on Global Carbon and Water Cycles, Journal of Climate, 28, 292–305, 2015.
  - LRTAP-Convention: Manual on Methodologies and Criteria for Modelling and Mapping Critical Loads and Levels; and Air Pollution Effects, Risks and Trends, https://icpvegetation.ceh.ac.uk/, https://icpvegetation.ceh.ac.uk/publications/documents/FinalnewChapter3v4Oct2017\_

755 000.pdf, 2017.

Marenco, A., Gouget, H., Nédélec, P., Pagés, J., and Karcher, F.: Evidence of a long-term increase in tropospheric ozone from Pic du Midi data series: Consequences: Positive radiative forcing, Journal of Geophysical Research: Atmospheres (1984–2012), 99, 16617–16632, 1994.

Massman, W.: A review of the molecular diffusivities of  $H_2O$ ,  $CO_2$ ,  $CH_4$ , CO,  $O_3$ ,  $SO_2$ ,  $NH_3$ ,  $N_2O$ , NO, and  $NO_2$  in air,  $O_2$  and  $N_2$  near

- 760 STP, Atmospheric Environment, 32, 1111–1127, https://doi.org/10.1016/S1352-2310(97)00391-9, http://tinyurl.sfx.mpg.de/ucen, 1998. McAinsh, M., Evans, N., Montgomery, L., and North, K.: Calcium signalling in stomatal responses to pollutants, New Phytologist, 153, 441–447, 2002.
  - Medlyn, B., Barton, C., Broadmeadow, M., Ceulemans, R., De Angelis, P., Forstreuter, M., Freeman, M., Jackson, S., Kellomäki, S., Laitat, E., et al.: Stomatal conductance of forest species after long-term exposure to elevated CO<sub>2</sub> concentration: A synthesis, New Phytologist, 149, 247–264, 2001.
- 765 14
  - Meinshausen, M., Smith, S. J., Calvin, K., Daniel, J. S., Kainuma, M., Lamarque, J.-F., Matsumoto, K., Montzka, S., Raper, S., Riahi, K., Thomson, A., Velders, G. J. M., and van Vuuren, D. P.: The RCP greenhouse gas concentrations and their extensions from 1765 to 2300, Climatic Change, 109, 213, 2011.

Meleux, F., Solmon, F., and Giorgi, F.: Increase in summer European ozone amounts due to climate change, Atmospheric Environment, 41,

770 7577–7587, 2007.

- Meyerholt, J. and Zaehle, S.: The role of stoichiometric flexibility in modelling forest ecosystem responses to nitrogen fertilization, New Phytologist, 208, 1042–1055, https://doi.org/10.1111/nph.13547, https://nph.onlinelibrary.wiley.com/doi/abs/10.1111/nph.13547, 2015.
- Moss, R. H., Edmonds, J. A., Hibbard, K. A., Manning, M. R., Rose, S. K., Van Vuuren, D. P., Carter, T. R., Emori, S., Kainuma, M., Kram, T., Meehl, G. A., Mitchell, J. F. B., Nakicenovic, N., Riahi, K., Smith, S. J., Stouffer, R. J., Thomson, A. M., Weyant, J. P., and Wilbanks,
- T. J.: The next generation of scenarios for climate change research and assessment, Nature, 463, 747, 2010.
   Niinemets, Ü., Keenan, T. F., and Hallik, L.: A worldwide analysis of within-canopy variations in leaf structural, chemical and physiological traits across plant functional types, New Phytologist, 205, 973–993, 2015.
- Noormets, A., Sober, A., Pell, E., Dickson, R., Podila, G., Sober, J., Isebrands, J., and Karnosky, D.: Stomatal and non-stomatal limitation to photosynthesis in two trembling aspen (*Populus tremuloides* Michx.) clones exposed to elevated CO<sub>2</sub> and/or O<sub>3</sub>, Plant, Cell & Environment, 24, 327–336, 2001.

Noormets, A., Kull, O., Sôber, A., Kubiske, M., and Karnosky, D.: Elevated CO<sub>2</sub> response of photosynthesis depends on ozone concentration in aspen, Environmental Pollution, 158, 992-999, 2010.

Norby, R.: Nitrogen deposition: a component of global change analyses, New Phytologist, 139, 189–200, 1998.

Norby, R., Warren, J., Iversen, C., Garten, C., Medlyn, B., and McMurtrie, R.: CO<sub>2</sub> Enhancement of Forest Productivity Constrained by 785 Limited Nitrogen Availability, Nature Precedings, https://doi.org/10.1038/npre.2009.3747.1, 2009.

- Oliver, R. J., Mercado, L. M., Sitch, S., Simpson, D., Medlyn, B. E., Lin, Y.-S., and Folberth, G. A.: Large but decreasing effect of ozone on the European carbon sink, Biogeosciences, 15, 4245–4269, 2018.
  - Oren, R., Ellsworth, D. S., Johnsen, K. H., Phillips, N., Ewers, B. E., Maier, C., Schäfer, K. V., McCarthy, H., Hendrey, G., McNulty, S. G., and Katul, G. G.: Soil fertility limits carbon sequestration by forest ecosystems in a CO2-enriched atmosphere, Nature, pp. 469-472, https://doi.org/10.1038/35078064, 2001.
- 790
  - Paoletti, E. and Grulke, N.: Does living in elevated CO<sub>2</sub> ameliorate tree response to ozone? A review on stomatal responses, Environmental Pollution, 137, 483-493, 2005.
  - Quéré, C. L., Andrew, R. M., Friedlingstein, P., Sitch, S., Hauck, J., Pongratz, J., Pickers, P. A., Korsbakken, J. I., Peters, G. P., Canadell, J. G., Arneth, A., Arora, V. K., Barbero, L., Bastos, A., Bopp, L., Chevallier, F., Chini, L. P., Ciais, P., Doney, S. C., Gkritzalis, T., Goll,
- 795 D. S., Harris, I., Haverd, V., Hoffman, F. M., Hoppema, M., Houghton, R. A., Hurtt, G., Ilvina, T., Jain, A. K., Johannessen, T., Jones, C. D., Kato, E., Keeling, R. F., Goldewijk, K. K., Landschützer, P., Lefèvre, N., Lienert, S., Liu, Z., Lombardozzi, D., Metzl, N., Munro, D. R., Nabel, J. E. M. S., ichiro Nakaoka, S., Neill, C., Olsen, A., Ono, T., Patra, P., Peregon, A., Peters, W., Peylin, P., Pfeil, B., Pierrot, D., Poulter, B., Rehder, G., Resplandy, L., Robertson, E., Rocher, M., Rödenbeck, C., Schuster, U., Schwinger, J., Séférian, R., Skjelvan, I., Steinhoff, T., Sutton, A., Tans, P. P., Tian, H., Tilbrook, B., Tubiello, F. N., van der Laan-Luijkx, I. T., van der Werf, G. R., Viovy,
- 800 N., Walker, A. P., Wiltshire, A. J., Wright, R., Zaehle, S., and Zheng, B.: Global Carbon Budget 2018, Earth System Science Data, 10, 2141-2194, https://doi.org/10.5194/essd-10-2141-2018, https://ueaeprints.uea.ac.uk/id/eprint/69293/, 2018.
  - Ren, W., Tian, H., Liu, M., Zhang, C., Chen, G., Pan, S., Felzer, B., and Xu, X.: Effects of tropospheric ozone pollution on net primary productivity and carbon storage in terrestrial ecosystems of China, Journal of Geophysical Research: Atmospheres, 112, 2007.
- Ren, W., Tian, H., Tao, B., Chappelka, A., Sun, G., Lu, C., Liu, M., Chen, G., and Xu, X.: Impacts of tropospheric ozone and climate change 805 on net primary productivity and net carbon exchange of China's forest ecosystems, Global Ecology and Biogeography, 20, 391-406, 2011.
  - Riikonen, J., Lindsberg, M.-M., Holopainen, T., Oksanen, E., Lappi, J., Peltonen, P., and Vapaavuori, E.: Silver birch and climate change: variable growth and carbon allocation responses to elevated concentrations of carbon dioxide and ozone. Tree Physiology, 24, 1227–1237, 2004.
    - Sanz, M., Calatayud, V., and Sánchez-Peña, G.: Measures of ozone concentrations using passive sampling in forests of South Western Europe,

810 Environmental Pollution, 145, 620-628, 2007.

- Schaub, M., Skelly, J., Zhang, J., Ferdinand, J., Savage, J., Stevenson, R., Davis, D., and Steiner, K.: Physiological and foliar symptom response in the crowns of Prunus serotina, Fraxinus americana and Acer rubrum canopy trees to ambient ozone under forest conditions, Environmental Pollution, 133, 553-567, 2005.
- Schulte-Uebbing, L. and de Vries, W.: Global-scale impacts of nitrogen deposition on tree carbon sequestration in tropical, temperate,
- 815 and boreal forests: A meta-analysis, Global Change Biology, 24, e416-e431, https://doi.org/https://doi.org/10.1111/gcb.13862, https:// onlinelibrary.wiley.com/doi/abs/10.1111/gcb.13862, 2018.
  - Sicard, P., Anav, A., Marco, A. D., and Paoletti, E.: Projected global ground-level ozone impacts on vegetation under different emission and climate scenarios, Atmospheric Chemistry and Physics, 17, 12177-12196, 2017.

Sitch, S., Cox, P., Collins, W., and Huntingford, C.: Indirect radiative forcing of climate change through ozone effects on the land-carbon

820 sink, Nature, 448, 791–794, 2007.

850

- Staehelin, J., Thudium, J., Buehler, R., Volz-Thomas, A., and Graber, W.: Trends in surface ozone concentrations at Arosa (Switzerland), Atmospheric Environment, 28, 75–87, 1994.
- Talhelm, A. F., Pregitzer, K. S., Kubiske, M. E., Zak, D. R., Campany, C. E., Burton, A. J., Dickson, R. E., Hendrey, G. R., Isebrands, J. G., Lewin, K. F., Nagy, J., and Karnosky, D. F.: Elevated carbon dioxide and ozone alter productivity and ecosystem carbon content
- 825 in northern temperate forests, Global Change Biology, 20, 2492–2504, https://doi.org/10.1111/gcb.12564, https://onlinelibrary.wiley.com/ doi/abs/10.1111/gcb.12564, 2014.
  - Tausz, M., Grulke, N., and Wieser, G.: Defense and avoidance of ozone under global change, Environmental Pollution, 147, 525–531, 2007.
  - Thomas, R. Q., Canham, C. D., Weathers, K. C., and Goodale, C. L.: Increased tree carbon storage in response to nitrogen deposition in the US, Nature Geoscience, 3, 13–17, 2010.
- 830 Tjoelker, M., Volin, J., Oleksyn, J., and Reich, P.: Interaction of ozone pollution and light effects on photosynthesis in a forest canopy experiment, Plant, Cell & Environment, 18, 895–905, 1995.
  - Turnock, S. T., Allen, R. J., Andrews, M., Bauer, S. E., Deushi, M., Emmons, L., Good, P., Horowitz, L., John, J. G., Michou, M., Nabat, P., Naik, V., Neubauer, D., O'Connor, F. M., Olivié, D., Oshima, N., Schulz, M., Sellar, A., Shim, S., Takemura, T., Tilmes, S., Tsigaridis, K., Wu, T., and Zhang, J.: Historical and future changes in air pollutants from CMIP6 models, Atmospheric Chemistry and Physics, 20,
- 835 14 547–14 579, https://doi.org/10.5194/acp-20-14547-2020, https://acp.copernicus.org/articles/20/14547/2020/, 2020.
  - Uddling, J., Hogg, A., Teclaw, R., Carroll, M., and Ellsworth, D.: Stomatal uptake of O<sub>3</sub> in aspen and aspen-birch forests under free-air CO<sub>2</sub> and O<sub>3</sub> enrichment, Environmental Pollution, 158, 2023–2031, 2010.
- van Vuuren, D. P., Edmonds, J., Kainuma, M., Riahi, K., Thomson, A., Hibbard, K., Hurtt, G. C., Kram, T., Krey, V., Lamarque, J.-F., Masui, T., Meinshausen, M., Nakicenovic, N., Smith, S. J., and Rose, S. K.: The representative concentration pathways: an overview, Climatic Change, 109, 5, https://doi.org/10.1007/s10584-011-0148-z, https://doi.org/10.1007/s10584-011-0148-z, 2011.
  - Vitousek, P. M. and Howarth, R. W.: Nitrogen limitation on land and in the sea: how can it occur?, Biogeochemistry, 13, 87–115, 1991.
  - Wang, B., Shugart, H. H., Shuman, J. K., and Lerdau, M. T.: Forests and ozone: productivity, carbon storage, and feedbacks, Scientific Reports, 6, 22133, 2016.

- 845 L., Vuichard, N., Zhou, F., Li, B., Piao, S., Peng, S., Huang, Y., and Tao, S.: Global forest carbon uptake due to nitrogen and phosphorus deposition from 1850 to 2100, Global Change Biology, 23, 4854–4872, https://doi.org/https://doi.org/10.1111/gcb.13766, https://onlinelibrary.wiley.com/doi/abs/10.1111/gcb.13766, 2017.
  - Wild, O., Fiore, A. M., Shindell, D., Doherty, R., Collins, W., Dentener, F., Schultz, M., Gong, S., MacKenzie, I., Zeng, G., Hess, P., Duncan, B. N., Bergmann, D. J., Szopa, S., Jonson, J. E., Keating, T. J., and Zuber, A.: Modelling future changes in surface ozone: a parameterized approach, Atmospheric Chemistry and Physics, 12, 2037–2054, 2012.
  - Wittig, V., Ainsworth, E., and Long, S.: To what extent do current and projected increases in surface ozone affect photosynthesis and stomatal conductance of trees? A meta-analytic review of the last 3 decades of experiments, Plant, Cell & Environment, 30, 1150–1162, 2007.
  - Wittig, V., Ainsworth, E., Naidu, S., Karnosky, D., and Long, S.: Quantifying the impact of current and future tropospheric ozone on tree biomass, growth, physiology and biochemistry: a quantitative meta-analysis, Global Change Biology, 15, 396–424, 2009.

Wang, R., Goll, D., Balkanski, Y., Hauglustaine, D., Boucher, O., Ciais, P., Janssens, I., Penuelas, J., Guenet, B., Sardans, J., Bopp,

- 855 Wohlgemuth, H., Mittelstrass, K., Kschieschan, S., Bender, J., Weigel, H., Overmyer, K., Kangasjärvi, J., Sandermann, H., and Langebartels, C.: Activation of an oxidative burst is a general feature of sensitive plants exposed to the air pollutant ozone, Plant, Cell & Environment, 25, 717–726, 2002.
- Wustman, B., Oksanen, E., Karnosky, D., Noormets, A., Isebrands, J., Pregitzer, K., Hendrey, G., Sober, J., and Podila, G.: Effects of elevated CO<sub>2</sub> and O<sub>3</sub> on aspen clones varying in O<sub>3</sub> sensitivity: can CO<sub>2</sub> ameliorate the harmful effects of O<sub>3</sub>?, Environmental Pollution, 115, 473–481, 2001.
  - Yue, X. and Unger, N.: Ozone vegetation damage effects on gross primary productivity in the United States, Atmospheric Chemistry and Physics, 14, 9137–9153, 2014.
  - Zaehle, S.: Terrestrial nitrogen-carbon cycle interactions at the global scale, Philosophical Transactions of the Royal Society B: Biological Sciences, 368, 20130125, https://doi.org/10.1098/rstb.2013.0125, https://royalsocietypublishing.org/doi/abs/10.1098/rstb.2013.0125,

865 2013.

- Zaehle, S. and Friend, A.: Carbon and nitrogen cycle dynamics in the O-CN land surface model: 1. Model description, site-scale evaluation, and sensitivity to parameter estimates, Global Biogeochemical Cycles, 24, GB1005, http://dx.doi.org/10.1029/2009GB003521, 2010.
- Zaehle, S., Friend, A., Friedlingstein, P., Dentener, F., Peylin, P., and Schulz, M.: Carbon and nitrogen cycle dynamics in the O-CN land surface model: 2. Role of the nitrogen cycle in the historical terrestrial carbon balance, Global Biogeochemical Cycles, 24, GB1006,

870 2010.

- Zaehle, S., Ciais, P., Friend, A. D., and Prieur, V.: Carbon benefits of anthropogenic reactive nitrogen offset by nitrous oxide emissions, Nature Geoscience, 4, 601–605, 2011.
- Zaehle, S., Medlyn, B. E., De Kauwe, M. G., Walker, A. P., Dietze, M. C., Hickler, T., Luo, Y., Wang, Y.-P., El-Masri, B., Thornton, P., Jain, A., Wang, S., Warlind, D., Weng, E., Parton, W., Iversen, C. M., Gallet-Budynek, A., McCarthy, H., Finzi, A., Hanson, P. J., Prentice, I. C.,
- 875 Oren, R., and Norby, R. J.: Evaluation of 11 terrestrial carbon–nitrogen cycle models against observations from two temperate Free-Air CO2 Enrichment studies, New Phytologist, 202, 803–822, https://doi.org/10.1111/nph.12697, https://nph.onlinelibrary.wiley.com/doi/abs/ 10.1111/nph.12697, 2014.
  - Zak, D. R., Pregitzer, K. S., Kubiske, M. E., and Burton, A. J.: Forest productivity under elevated *CO*<sub>2</sub> and *O*<sub>3</sub>: positive feedbacks to soil N cycling sustain decade-long net primary productivity enhancement by *CO*<sub>2</sub>, Ecology Letters, 14, 1220–1226, 2011.

880 Appendix A



**Figure S1.** Mean simulated change in nitrogen deposition rates for the temperate and boreal Northern Hemisphere ( $\geq 30^{\circ}$ N) in the decades of the years of 1990, 2040 and 2090 compared to the decade of the year 1850, each according to the RCP2.6 and RCP8.5 pollution emission scenario.



**Figure S2.** Mean simulated change in canopy level  $O_3$  concentration for the temperate and boreal Northern Hemisphere ( $\geq 30^\circ N$ ) in the decades of the years of 1850, 1990, 2040 and 2090 compared to the decade of the year 1850, each according to the RCP2.6 and RCP8.5 pollution emission scenario.



**Figure S3.** Nitrogen deposition induced absolute change in regional mean ozone  $O_3$  uptake ( $F_{st}$ ), mean cumulative  $O_3$  uptake above a flux threshold of 1 nmol m<sup>-2</sup> s<sup>-1</sup> (CUO1), summed GPP, summed total carbon biomass in vegetation (vegetation-C), summed carbon soil organic matter (soil-C), and summed land carbon flux (land C flux) compared to pre-industrial values in the simulation region. The nitrogen deposition induced change is calculated from the simulation runs S3 and S5 (see Tab. 2). Solid lines indicate results from simulations based on RCP8.5, dotted lines results from simulations based on RCP2.6. The effect of the seasonal cycle is smoothed by the application of a moving average of 12 months (a,b).



**Figure S4.** Nitrogen deposition induced absolute change in regional summed total N uptake by region,  $N_2O$  emission,  $NH_4$  leaching and  $N_2O$  leaching compared to pre-industrial values in the simulation region. The nitrogen deposition induced absolute change is calculated from the simulation runs S3 and S5 (see Tab. 2). Solid lines indicate results from simulations based on RCP8.5, dotted lines results from simulations based on RCP2.6.



**Figure S5.** Ozone induced absolute change in regional mean ozone  $O_3$  uptake ( $F_{st}$ ), mean cumulative  $O_3$  uptake above a flux threshold of 1 nmol m<sup>-2</sup> s<sup>-1</sup> (CUO1), summed GPP, summed total carbon biomass in vegetation (vegetation-C) and summed carbon soil organic matter (soil-C) compared to pre-industrial values in the simulation region. Different colors indicate different approaches to calculate the ozone  $O_3$  induced change from the factorial runs. Orange lines represent approach 1: (S2-S1)/S1, blue lines approach 2:(S5-S4)/S4. Solid lines indicate results from simulations based on RCP8.5, dotted lines results from simulations based on RCP2.6. The effect of the seasonal cycle is smoothed by the application of a moving average of 12 months.



**Figure S6.** Absolute change in CUO1 compared to pre-industrial values induced by  $\frac{\text{ozone}O_3}{\text{ozone}O_3}$ , calculated according to approach 2. Displayed are the decade 1990 (mean of the years 1990-1999), 2040 (mean of the years 2040-2049) and of 2090 (mean of the years 2090-2099). For the decades 2040 and 2090 results from simulations based on RCP8.5 and RCP2.6 are displayed. See Tab. 2 for details on the calculation of the <u>ozone-O\_3</u> impact.



**Figure S7.** Relative change in GPP compared to pre-industrial values induced by nitrogen deposition (left column) and  $\frac{1}{02000}$  O<sub>3</sub> calculated according to approach 2 (right column). The induced change in GPP is displayed for the decades 1990 (mean of the years 1990-1999), 2040 (mean of the years 2040-2049) and 2090 (mean of the years 2090-2099). For the decades 2040 and 2090 results from simulations based on RCP8.5 and RCP2.6 are displayed. See Tab. 2 for details on the calculation of the single drivers.



**Figure S8.** Combined absolute impact of nitrogen deposition and  $O_3$  calculated according to approach 2 on GPP compared to pre-industrial values. The induced change in GPP is displayed for the decades 1990 (mean of the years 1990-1999), 2040 (mean of the years 2040-2049) and 2090 (mean of the years 2090-2099). For the decades 2040 and 2090 results from simulations based on RCP8.5 and RCP2.6 are displayed. See Tab. 2 for details on the calculation of the single drivers.



**Figure S9.** Relative change in the <u>total carbon biomass in vegetation</u> (vegetation-C) compared to pre-industrial values induced by nitrogen deposition (left column) and <del>ozone</del>- $O_3$  calculated according to approach 2 (right column). The induced change in vegetation-C is displayed for the decades 1990 (mean of the years 1990-1999), 2040 (mean of the years 2040-2049) and 2090 (mean of the years 2090-2099). For the decades 2040 and 2090 results from simulations based on RCP8.5 and RCP2.6 are displayed. See Tab. 2 for details on the calculation of the single drivers.

**Table S1.** The spread in the mean percent change displayed in Tab. 4 due to inter-annual variability, derived from error propagation of the yearly estimates in GPP, total carbon biomass in vegetation (vegetation-C), and land carbon flux (land C flux) induced by  $O_3$  during the decades of 1990 (1990–1999), 2040 (2040–2049) and 2090 (2090–2099). For  $O_3$  two values are displayed which refer to the two approaches to calculate the  $O_3$  impact.

Region	Pollutant	1990	2040 RCP8.5	2040 RCP2.6	2090 RCP8.5	2090 RCP2.6
GPP						
NH30	$O_3$	±0.17;±0.14	±0.11;±0.19	±0.09;±0.22	±0.10;±0.10	±0.10;±0.08
NH30	Ndep	±0.13	±0.18	±0.11	±0.20	±0.12
Europe	$O_3$	±0.25;±0.33	$\pm 0.14;\pm 0.19$	±0.14;±0.24	±0.11;±0.13	$\pm 0.04; \pm 0.08$
Europe	Ndep	$\pm 0.26$	$\pm 0.30$	$\pm 0.21$	$\pm 0.21$	±0.12
USA	$O_3$	±0.41;±0.39	$\pm 0.19; \pm 0.26$	±0.17;±0.30	$\pm 0.05;\pm 0.23$	±0.20;±0.17
USA	Ndep	±0.23~	±0.19	±0.23	$\pm 0.30$	±0.13
China	$O_3$	$\pm 0.95; \pm 0.76$	$\pm 0.67; \pm 1.31$	±0.45;±1.02	±0.41;±0.40	$\pm 0.43; \pm 0.44$
China	Ndep	±0.23	±0.53	±0.23	±0.40	±0.59
Vegetation-C						
NH30	$O_3$	$\pm 0.38; \pm 0.40$	±0.14;±0.27	±0.15;±0.17	±0.17;±0.12	±0.19;±0.18
NH30	Ndep	±0.13	±0.15	±0.09	$\pm 0.08$	±0.04
Europe	$O_3$	$\pm 0.34;\pm 0.42$	±0.13;±0.33	±0.19;±0.21	±0.22;±0.15	$\pm 0.18; \pm 0.24$
Europe	Ndep	±0.15	±0.20	±0.11	$\pm 0.09$	$\pm 0.05$
USA	$O_3$	<u>±0.62;±0.50</u>	±0.26;±0.27	±0.27;±0.29	±0.16;±0.24	$\pm 0.28; \pm 0.25$
USA	Ndep	$\pm 0.10$	±0.07	$\pm 0.03$	$\pm 06$	$\pm 0.04$
China	$O_3$	±1.63;±1.86	$\pm 0.43;\pm 0.60$	$\pm 0.23;\pm 0.32$	$\pm 0.45;\pm 0.44$	$\pm 0.44;\pm 0.48$
China	Ndep	±0.13	±0.22	$\pm 0.19$	$\pm 0.24$	±0.21
Land C flux						
NH30	$O_3$	<u>±4.28;±4.48</u>	<u>±0.78;±0.98</u>	±2.45;±2.07	±0.88;±2.15	±3.87;±7.86
NH30	Ndep	±2.20	±1.42	±1.46	±2.04	±1.51
Europe	$O_3$	$\pm 12.48;\pm 14.84$	±1.19;±1.31	±3.57;±4.70	±1.45;±12.10	±8.66;±11.30
Europe	Ndep	±13.34	±2.06	$\pm 1.69$	±5.06	±2.10
USA	$O_3$	±8.60;±12.22	±1.17;±3.20	±2.13;±6.93	$\pm 0.72; \pm 6.25$	±5.30;±12.86
USA	Ndep	$\pm 4.50$	±3.85	$\pm 3.33$	$\pm 4.95$	±2.95
China	$O_3$	±13.23;±25.46	<u>±6.35;±13.3</u>	$\pm 6.85; \pm 11.77$	$\pm 6.64; \pm 13.77$	$\pm 16.81; \pm 14.57$
China	Ndep	±15.81	±7.00	±4.89	±6.46	±8.29