

Arctic Ocean acidification over the 21st century co-driven by anthropogenic carbon increases and freshening in the CMIP6 model ensemble

Jens Terhaar^{1,2}, Olivier Torres³, Timothée Bourgeois⁴, and Lester Kwiatkowski⁵

¹Climate and Environmental Physics, Physics Institute, University of Bern, Bern, Switzerland

²Oeschger Center for Climate Change Research, University of Bern, Bern, Switzerland

³LMD/IPSL, Ecole Normale Supérieure/PSL Université, CNRS, Ecole Polytechnique, Sorbonne Université, Paris, France

⁴NORCE Norwegian Research Centre and Bjerknes Centre for Climate Research, Bergen, Norway

⁵LOCEAN/IPSL, Sorbonne Université, CNRS, IRD, MNHN, Paris, France

Correspondence: Jens Terhaar (jens.terhaar@climate.unibe.ch)

Abstract. The uptake of anthropogenic carbon (C_{ant}) by the ocean leads to ocean acidification, causing the reduction of pH and the calcium carbonate saturation states of aragonite (Ω_{arag}) and calcite (Ω_{calc}). The Arctic Ocean is particularly vulnerable to ocean acidification due to its naturally low pH and saturation states and due to ongoing freshening and the concurrent reduction in alkalinity in this region. Here, we analyse ocean acidification in the Arctic Ocean over the 21st century across 14 Earth System Models (ESMs) from the latest Coupled Model Intercomparison Project Phase 6 (CMIP6). Compared to the previous model generation (CMIP5), [models generally better simulate maximum sea surface densities in the Arctic Ocean and consequently the transport of \$C_{\text{ant}}\$ into the Arctic Ocean interior, with simulated historical increases in \$C_{\text{ant}}\$ in improved agreement with observational products. Moreover, in CMIP6 the inter-model uncertainty of projected end-of-century changes over the 21st century in Arctic Ocean \$\Omega_{\text{arag/calc}}\$ averaged over the upper 1000 m is reduced by 44-64%. The strong reduction in projection uncertainties of \$\Omega_{\text{arag/calc}}\$ can be attributed to compensation between \$C_{\text{ant}}\$ uptake and alkalinity reduction in the latest models. Specifically, ESMs with a large increase in Arctic Ocean \$C_{\text{ant}}\$ over the 21st century tend to simulate a relatively weak concurrent freshening and alkalinity reduction, while ESMs with a small increase in \$C_{\text{ant}}\$ simulate a relatively strong freshening and concurrent alkalinity reduction. Although both mechanisms contribute to Arctic Ocean acidification over the 21st century, the increase in \$C_{\text{ant}}\$ remains the dominant driver. Even under the low-emissions shared socioeconomic pathway SSP1-2.6, basin-wide averaged \$\Omega_{\text{arag}}\$ undersaturation \[in the upper 1000 m\]\(#\) occurs before the end of the century. While under the high-emissions pathway SSP5-8.5, the Arctic Ocean mesopelagic is projected to even become undersaturated with respect to calcite. An emergent constraint, identified in CMIP5, which relates present-day maximum sea surface densities in the Arctic Ocean to the projected end-of-century Arctic Ocean \$C_{\text{ant}}\$ inventory, is found to generally hold in CMIP6. However, a coincident constraint on Arctic declines in \$\Omega_{\text{arag/calc}}\$ is not apparent in the new generation of models. This is due to both the reduction in \$\Omega_{\text{arag/calc}}\$ projection uncertainty and the weaker direct relationship between projected changes in Arctic Ocean \$C_{\text{ant}}\$ and \$\Omega_{\text{arag/calc}}\$. ~~In CMIP6, models generally better simulate maximum sea surface densities in the Arctic Ocean and consequently the transport of \$C_{\text{ant}}\$ into the Arctic Ocean interior, with simulated historical increases in \$C_{\text{ant}}\$ in improved agreement with observational products.~~](#)

25 1 Introduction

Human activities such as the burning of fossil fuels, cement production, and land use change have released large amounts of carbon into the atmosphere that cause global warming. The ocean mitigates global warming by taking up around one quarter of this anthropogenic carbon (C_{ant}) (Friedlingstein et al., 2019). However, the increase of carbon in the ocean causes ocean acidification, a process that decreases pH, carbonate ion (CO_3^{2-}) concentrations, and in consequence the calcium carbonate (CaCO₃) saturation states, of calcite and aragonite minerals (Haugan and Drange, 1996; Orr et al., 2005). The Arctic Ocean is particularly vulnerable to ocean acidification due to its naturally high dissolved inorganic carbon concentrations, low carbonate ion concentrations and its thus naturally low saturation states (Orr et al., 2005; Fabry et al., 2009; Gattuso and Hansson, 2011; Riebesell et al., 2013; AMAP, 2018).

In contrast to most of the global ocean, Arctic Ocean acidification is not solely caused by increasing C_{ant} concentrations (Anderson et al., 2010; Ulfsbo et al., 2018; Terhaar et al., 2020a, b) but also by freshening (Koenigk et al., 2013; Nummelin et al., 2016; Shu et al., 2018; Brown et al., 2020; Woosley and Millero, 2020). Freshwater from rivers, precipitation, and sea ice typically has much lower total alkalinity (A_T) and total dissolved inorganic carbon (C_T) concentrations than the ocean and therefore, in the absence of indirect impacts on other fluxes, dilutes both marine A_T and C_T (Xue and Cai, 2020). As freshwater A_T and C_T concentrations are generally similar, freshwater fluxes into the ocean typically act to reduce the difference between A_T and C_T , ~~which is approximately equal to the CO_3^{2-} concentration (Waldbusser and Salisbury, 2014; Wanninkhof et al., 2015; Xue and Cai, 2020).~~ Freshening thus reduces ~~decreasing~~ marine CO_3^{2-} concentrations and ~~enhances ocean acidification (Bates et al., 2009; Bates and Mathis, 2009).~~ ocean pH (Bates et al., 2009; Bates and Mathis, 2009; Yamamoto-Kawai et al., 2011; Waldbusser and Salisbury, 2014; Wanninkhof et al., 2015). In the Arctic Ocean, projected freshening over the 21st century is larger than in most other ocean regions due to ongoing sea ice melt, positive precipitation minus evaporation, and large river runoff (Rawlins et al., 2010; Rudels, 2015; Shu et al., 2018).

Due to freshening and increasing C_{ant} concentrations, the Arctic Ocean is projected to be the first large-scale ocean region to become undersaturated with respect to the metastable CaCO₃ polymorph aragonite ($\Omega_{\text{arag}} < 1$) (Steinacher et al., 2009). Under the RCP8.5 high-emissions scenario, Arctic Ocean mesopelagic waters may even become undersaturated with respect to the more stable CaCO₃ polymorph calcite ($\Omega_{\text{calc}} < 1$) before 2100 (Terhaar et al., 2020a). Aragonite and calcite undersaturation is likely to affect the growth, reproduction, and survival of calcifying organisms, such as sea butterflies (Comeau et al., 2010) and foraminifera (Davis et al., 2017), and could have ramifications for the wider Arctic ecosystem (Armstrong et al., 2005; Karnovsky et al., 2008) including some of its most iconic predators, such as gray whales ~~or walruses (Bates et al., 2009)~~ and walruses (Jay et al., 2011; AMAP, 2018).

Projections of Arctic Ocean ~~ocean~~ acidification over the 21st century had considerable subsurface uncertainties in the simulations conducted as part of the Coupled Model Intercomparison Project Phase 5 (CMIP5) (Steiner et al., 2013; Terhaar et al., 2020a), with projected end-of-century basin-wide Ω_{arag} in mesopelagic waters ranging from 0.61 to 1.05. This large uncertainty has been attributed to multiple factors including variable inflow of Atlantic waters and their subsequent subduction, difficulties

resolving the narrow passages between the Arctic Ocean and its surrounding basins and differences in brine rejection during sea ice formation, which is critical to the formation of dense Arctic waters (Terhaar et al., 2019b).

1.1 Emergent constraints on Arctic Ocean carbon uptake and acidification

60 Emergent constraints are a suite of statistical techniques that relate observable trends or sensitivities across multi-model ensembles to differences in model projections in order to reduce future uncertainties (Allen and Ingram, 2002; Hall and Qu, 2006; Hall et al., 2019). Such (Allen and Ingram, 2002; Hall et al., 2019). An elegant early application that demonstrates many of the principals of emergent constraints is that of Hall and Qu (2006). In their study of models that contributed to the Intergovernmental Panel on Climate Change's Fourth Assessment Report (AR4), Hall and Qu found a strong positive correlation between the magnitude of a model's snow albedo feedback on present-day seasonal timescales and under future climate change. They concluded that relevant model biases were consistent across these contrasting timescales and therefore observations of the seasonal snow albedo feedback could be used to constrain the ensemble range of the projected snow albedo feedback under climate change. Since the publication of Hall and Qu (2006), emergent constraint approaches have been applied extensively within the earth sciences to constrain projections, amongst others, projections of climate sensitivity (Caldwell et al., 2018), Arctic sea ice extent (Hall and Qu, 2006) (Boé et al., 2009), precipitation extremes (O'Gorman, 2012; DeAngelis et al., 2016), carbon cycle feedbacks (Cox et al., 2013; Wenzel et al., 2014), and marine primary production (Kwiatkowski et al., 2017).

Recently Terhaar et al. (2020a) showed that an emergent constraint could be applied to CMIP5 projections of the Arctic Ocean C_{ant} inventory and coincident acidification over the twenty-first century. As the C_{ant} increase in the Arctic Ocean is mainly driven by the inflow of C_{ant} -rich waters from the Atlantic and their subsequent subduction in the Barents Sea (Midttun, 1985; Rudels et al., 1994, 2000; Jeansson et al., 2011; Smedsrud et al., 2013), the capability of each model to form dense surface waters in the Barents Sea was shown to strongly influence the future Arctic Ocean C_{ant} inventory. By constraining simulated surface water densities with observations, uncertainties related to the end-of-century Arctic Ocean C_{ant} inventory in 2100 were reduced by around one third and the best estimate under RCP8.5 was increased by 20% to 9.0 ± 1.6 Pg C (Terhaar et al., 2020a). Along with the projected C_{ant} inventory, uncertainties in the projected associated basin-wide Arctic Ocean acidification could also be reduced in CMIP5. It should be noted however, that in CMIP5, projected freshening and reductions in alkalinity were of minor importance for Arctic Ocean acidification over the 21st century. Moreover, the models have been shown to underestimate historical freshwater fluxes (1992–2012) in the Arctic Ocean by around 50% (Shu et al., 2018), which suggests they might also have underestimated freshwater fluxes over the 21st century.

Given that emergent constraints in many cases conflict with one another (Caldwell et al., 2018; Brient, 2020) and can even be derived from data mined pseudocorrelations (Caldwell et al., 2014), it is critical to test published constraints, and the mechanisms that underpin them, across ESM generations (Eyring et al., 2019; Hall et al., 2019). The CMIP6 simulations provide such an opportunity (Schlund et al., 2020).

1.2 From CMIP5 to CMIP6 models and simulations

During the transition from CMIP5 to CMIP6, ESMs have generally improved the simulation of ocean dynamics and marine biogeochemistry (Séférián et al., 2020). Across most ESMs, the horizontal and/or vertical resolution of ocean models has increased, which potentially has large effects on the representation of Arctic Ocean circulation, sea ice dynamics (Docquier et al., 2019), and the carbon cycle (Terhaar et al., 2019b). Ocean biogeochemical model components in CMIP6 also tend to have a more complex representation of the carbon and nutrient cycles than in CMIP5. In particular, the treatment of organic matter carbon cycling has generally evolved, with ~~sedimentation now explicitly~~ remineralization of particles in sediments now simulated in 10 out of 14 ESMs. These developments will likely have a large effect on simulating the Arctic Ocean biogeochemistry given that 50% of the Arctic Ocean is made up of shelf seas (Jakobsson, 2002), where sedimentation and sediment remineralization are crucial components of the carbon and nutrient cycle (Brüchert et al., 2018; Grotheer et al., 2020). Furthermore, the external carbon and nutrient sources from glaciers, atmospheric deposition, and rivers are represented in more models in CMIP6 (Table 1) (Séférián et al., 2020). Riverine inputs in particular have been shown to be of importance for present-day Arctic Ocean acidification (Anderson et al., 2010; Tank et al., 2012) and its future changes (Terhaar et al., 2019a).

In this study, we extend recent CMIP6 ocean biogeochemical assessments (e.g. Séférián et al. (2020); Kwiatkowski et al. (2020)) and previous attempts to constrain projected Arctic Ocean C_{ant} uptake by,

1. Assessing projections of the Arctic Ocean C_{ant} inventory over the 21st century in CMIP6 simulations,
2. Exploring the role of C_{ant} inventory increases and freshening in driving concurrent basin-wide ocean acidification in the Arctic basin,
3. Reevaluating previous emergent constraints on the Arctic Ocean C_{ant} inventory and associated acidification using the CMIP6 model ensemble and multiple future emissions scenarios.

2 Methods

2.1 Arctic Ocean

The Arctic Ocean was defined as the water north of the Fram Strait, the Barents Sea Opening, the Bering Strait and the Baffin Bay following Bates and Mathis (2009). This is consistent with the previously published emergent constraint on projected Arctic Ocean C_{ant} and acidification (Terhaar et al., 2020a).

2.2 Earth System Models

An ensemble of 14 ESMs from the Coupled Model Intercomparison Project Phase 6 (CMIP6) (Table 1) was used with one ensemble member per model. All models follow the the biogeochemical protocols outlined in Orr et al. (2017). Riverine input

of C_T and A_T is included in 6 ESMs (Table 1). The absence of C_T and A_T in riverine freshwater input causes an overly strong reduction of CO_3^{2-} concentrations and thus low biased saturation states in coastal regions, but is of minor importance on the pan-Arctic scale (Terhaar et al., 2019a). The spin-up length for each model varied between 500 years (IPSL-CM6A-LR) and 12000 years (MPI-ESM1-2-LR) (Séférian et al., 2020).

For each model, monthly 3D-fields of dissolved inorganic carbon, total alkalinity, dissolved inorganic phosphorus and silicon, temperature and salinity were used. All 3D-fields were regridded to the regular $1^\circ \times 1^\circ$ grid with 33 depth levels used in the GLObal Ocean Data Analysis Project Version 2 (GLODAPv2) observational product (Lauvset et al., 2016) to add simulated changes of these variables over the 21st century to observations of the present day mean state (see below).

C_{ant} was defined as the difference between annual means of dissolved inorganic carbon in the historical (1850–2014) simulations merged with the respective Shared Socioeconomic Pathway (SSP1-2.6, SSP2-4.5, SSP3-7.0, and SSP5-8.5) (2015–2100) (Riahi et al., 2017), and the concurrent pre-industrial control simulations of each model. Output from 13 models was available for SSP1-2.6, output from 12 models for SSP2-4.5 and SSP3-7.0, and output from 14 models for SSP5-8.5 (Table 1).

Changes in ~~total alkalinity~~ (A_T) over the 21st century were calculated by subtracting changes in the pre-industrial control simulations from changes in the respective SSP. To quantify the effect of freshening on changes in A_T , the A_T anomalies for each model were further decomposed into changes resulting from freshening and from the combined effect of other biogeochemical processes by calculating the temporal evolution of salinity corrected alkalinity with a reference salinity of 35 following Lovenduski et al. (2007). A zero alkalinity end-member was assumed for freshwater. This assumption is correct for models with no alkalinity in freshwater but an overestimation for models with finite alkalinity concentrations in freshwater inputs (Table 1). Unfortunately, information on alkalinity concentrations in freshwater is not available for all models. Moreover, with the available model output, we cannot quantify the individual contributions of land ice melt, sea ice melt, precipitation minus evaporation, and riverine input to freshwater changes. Thus, for simplicity a zero alkalinity end-member was assumed for all models.

Ocean carbon chemistry variables in ESMs commonly exhibit mean state biases (Orr et al., 2005; Steiner et al., 2013). Therefore, observations of C_T , A_T , dissolved inorganic phosphorus and silicon, temperature and salinity from GLODAPv2 (Lauvset et al., 2016), which is normalized to the year 2002, were used to calculate present day $\Omega_{arag/calc}$, pH and pCO_2 using the *mocsy2.0* routine (Orr and Epitalon, 2015) and the equilibrium constants recommended for best practices (Dickson et al., 2007). Future $\Omega_{arag/calc}$, pH and pCO_2 were calculated for each model as the sum of the simulated changes in C_T , A_T , dissolved inorganic phosphorus and silicon, temperature and salinity from 2002 onwards and the observed quantities in 2002.

The present-day maximum sea surface density in the Arctic Ocean was calculated from monthly climatologies over 1986–2005, constructed from the respective salinity and temperature outputs of each model. Maximum sea surface density was calculated, as in Terhaar et al. (2020a), as the mean density of the densest 5% of Arctic surface waters (95th percentile waters) over all 12 months of the year.

2.3 Simulations

150 The simulations performed within CMIP5 were not forced with the same atmospheric CO₂ concentrations as the simulations performed under CMIP6. In CMIP5, historically observed atmospheric CO₂ concentrations were used from 1850 to 2005 (Meinshausen et al., 2011). From 2006 onwards, the CO₂ concentrations follow the different RCPs. In CMIP6, the historical period was extended until 2014 and thereafter CO₂ concentrations follow the different SSPs (Meinshausen et al., 2019).

The different land and energy use assumptions in the SSPs (Riahi et al., 2017) compared to the RCPs (O'Neill et al., 2016) lead to higher atmospheric CO₂ trajectories over the 21st century for the Tier 1 SSPs (Meinshausen et al., 2019) compared to their RCP counterparts (O'Neill et al., 2016) (Meinshausen et al., 2011), which results in globally greater surface ocean acidification in CMIP6 compared to CMIP5 (Kwiatkowski et al., 2020). ~~Furthermore, historical~~ Historical atmospheric CO₂ concentrations were also refined with additional data available since CMIP5 (Meinshausen et al., 2017). This refinement did not change the average atmospheric CO₂ concentration from 1850 to 2005 ($\Delta p\text{CO}_2^{\text{tm}} = 0 \pm 1$ ppm), but changed annual CO₂ concentration for single years by up to ± 2 ppm. Furthermore, global CO₂ concentrations were additionally provided as monthly latitudinally resolved concentrations with model groups free to choose the forcing files they use (Meinshausen et al., 2017, 2019)

~

2.4 C_{ant} scaling

The different atmospheric CO₂ trajectories over the 21st century between CMIP5 and CMIP6 complicates a comparison of simulated Arctic Ocean C_{ant} inventories between model generations. To nevertheless compare the simulated Arctic Ocean C_{ant}, we used the commonly applied scaling approach that assumes that the change in marine C_{ant} is proportional to the atmospheric C_{ant} concentration (Gloor et al., 2003; Mikaloff Fletcher et al., 2006; Gerber et al., 2009; Gruber et al., 2009). Under this assumption, the C_{ant} inventory in 2100 for each scenario (RCP8.5, SSP1-2.6, SSP2-4.5, SSP3-7.0) is linearly rescaled to that of SSP5-8.5 by multiplying the simulated Arctic Ocean C_{ant} inventory under the respective scenario by the ratio of the mean atmospheric C_{ant} concentration from 1850 to 2100 in SSP5-8.5 and the respective scenario.

This approximation is likely imprecise when very different scenarios are compared, such as SSP1-2.6 and SSP5-8.5, as the effects of circulation changes, sea ice melt and warming are not considered. However, when comparing scenarios with the same radiative forcing, such as SSP5-8.5 and RCP8.5, it permits a first order comparison.

2.5 Observational constraints

175 As for the ESMs, the maximum sea surface density was calculated based on a monthly sea surface density climatology on a regular 1° × 1° grid, which was constructed from observed monthly salinity and temperature climatologies in the World Ocean Atlas 2018 (Locarnini et al., 2018; Zweng et al., 2018).

The density uncertainty was calculated from the temperature and salinity uncertainties that were reported by the World Ocean Atlas following standard propagation of uncertainty. The total uncertainty is a combination of (1) the standard deviations for sea surface density derived from published standard deviations of sea surface temperature and salinity for each grid cell and

each month in the World Ocean Atlas 2018, and (2) the standard deviation from the weighted mean of the 95th percentile density waters (Terhaar et al., 2020a).

2.6 ~~Probability~~ Emergent constraints and probability density functions of C_{ant}

~~The probability density functions (PDFs) To calculate the emergent constraint, first an ordinary least squares regression was~~
185 ~~calculated between the simulated present-day maximum sea surface density and the Arctic Ocean C_{ant} inventory in 2100 for~~
~~each ESM of the CMIP6 model ensemble. The uncertainty range was estimated using the 1σ prediction interval. In a second~~
~~step the probability density function (PDF) from the observations was convoluted with the PDF from the linear regression,~~
~~assuming a Gaussian distribution in both cases. The convolution of both PDFs is the constrained projection of the Arctic Ocean~~
 ~~C_{ant} inventory following previous studies (Cox et al., 2013; Wenzel et al., 2014; Kwiatkowski et al., 2017). The PDFs for un-~~
190 ~~constrained projections of the C_{ant} inventory were calculated using equal weights for each model and assuming a Gaussian~~
~~distribution.~~

~~The PDFs for the constrained projections of the C_{ant} inventory were calculated as the convolution of the observational PDF~~
~~and the PDF that results from the emergent relationship between present-day maximum sea surface density and projected C_{ant}~~
~~following previous studies (Cox et al., 2013; Wenzel et al., 2014; Kwiatkowski et al., 2017).~~

195 Extending the analysis of Terhaar et al. (2020a), PDFs for the constrained projections of the C_{ant} inventory were calculated
not only for the year 2100 but for each year from 2002 to 2100 and not only for the highest emission scenario (RCP8.5) but for
the four SSPs (SSP1-2.6, SSP2-4.5, SSP3-7.0, and SSP5-8.5).

3 Results

3.1 Arctic Ocean C_{ant} inventory

200 3.1.1 Multi-model mean

Over the historical period from 1850 to 2005, the CMIP6 ESMs simulate an Arctic Ocean C_{ant} increase of 2.1 ± 0.6 Pg C (inter-
model standard deviation; Fig. 1a). As such, the CMIP6 ESMs simulate an Arctic Ocean C_{ant} inventory in 2005 that is ~~61%~~
~~higher than the inventory that was simulated by the previous model generation in CMIP5 (1.3 ± 0.7 Pg C; Terhaar et al. (2020a)~~
~~).~~ ~~This is still~~ 36% below the data-based estimate for the period from 1765 to 2005 of 3.3 ± 0.3 Pg C (Terhaar et al., 2020b).

205 Over the 21st century the Arctic Ocean C_{ant} inventory increases depending on the SSP. Following the low-emission pathway
SSP1-2.6 leads to a projected Arctic Ocean C_{ant} inventory in 2100 of 6.2 ± 1.3 Pg C. With increasing atmospheric C_{ant}
concentrations from SSP1-2.6 to SSP5-8.5, the projected Arctic Ocean C_{ant} inventory in 2100 also increases, resulting in 8.0
 ± 1.7 Pg C for SSP2-4.5, 9.6 ± 1.6 Pg C for SSP3-7.0, and 10.3 ± 2.2 Pg C for SSP5-8.5. The relatively large uncertainties
across the model ensemble (17-21%) result in an overlap of the simulated C_{ant} inventories in 2100 for SSP2-4.5, SSP3-7.0,
210 and SSP5-8.5 within \pm one standard deviation.




Figure 1. Arctic Ocean C_{ant} inventory over the 21st century. **a)** Time-series of multi-model mean Arctic Ocean C_{ant} inventory over the 21st century for SSP1-2.6 (blue), SSP2-4.5 (green), SSP3-7.0 (orange), and SSP5-8.5 (red) with the respective standard deviation across the model ensemble (shading) ($n=12-14$). The bars on the right side of panels indicate the inter-model standard deviation in year 2100. **b)** Time-series of Arctic Ocean C_{ant} inventory for each scenario after the emergent constraint is applied.

3.1.2 Constrained results

As was shown for RCP8.5 in the CMIP5 model ensemble (Terhaar et al., 2020a), a linear relationship between maximum sea surface density and the Arctic Ocean C_{ant} inventory in 2100 is found across the CMIP6 model ensemble for all four of the SSPs ($r^2=0.63-0.72$; Fig. 2a,c,e,g). By deriving a similar relationship for the projected C_{ant} inventory in all years from 2000 to 2100 and combining this with observations of present-day sea surface density, the uncertainty of the projected Arctic Ocean C_{ant} inventory can be reduced throughout the 21st century (Fig. 1b). The emergent relationship is significant over all years from 2015 to 2100 ($p<0.05$) and the r^2 increases from 0.39 in 2014 to 0.63-0.72 in 2100 depending on the scenario. In the year 2100, this results in Arctic Ocean C_{ant} inventory estimates of 6.3 ± 0.9 Pg C (SSP1-2.6), 8.2 ± 1.2 Pg C (SSP2-4.5), 9.8 ± 1.1 Pg C (SSP3-7.0), and 10.7 ± 1.4 Pg C (SSP5-8.5) (Fig. 2b,d,f,h). As such, the emergent constraint is shown to slightly increase the CMIP6 multi-model mean projected C_{ant} inventory for each SSP (+2 to +4 %) and substantially reduce associated uncertainties (-29 to -31 %) resulting in greater separation of the SSPs (Fig. 1).

3.2 Ocean acidification

3.2.1 Multi-model mean

Over the 21st century, ocean acidification leads to a reduction in Ω_{arag} and Ω_{calc} (Fig. 3). Until around 2040, the reduction of both saturation states averaged over the first-upper 1000 m is independent of the SSP. From 2040 onwards, the rate of the reduction depends on the pathway and by 2100 basin-wide Ω_{arag} averaged over the first-upper 1000 m reaches 1.00 ± 0.06 (SSP1-2.6), 0.82 ± 0.05 (SSP2-4.5), 0.66 ± 0.04 (SSP3-7.0), and 0.59 ± 0.05 (SSP5-8.5), while Ω_{calc} reaches 1.56 ± 0.09 (SSP1-2.6), 1.29 ± 0.08 (SSP2-4.5), 1.06 ± 0.06 (SSP3-7.0), and 0.92 ± 0.05 (SSP5-8.5).

The first-upper 1000 m of the Arctic Ocean are projected to be on average undersaturated with respect to aragonite by 2100 under all of the SSPs with Ω_{arag} in SSP1-2.6 just below 1 (0.995). Vertically, the entire water column will be undersaturated with respect to aragonite in 2100 for all pathways but SSP1-2.6 (Fig 4). Under this low-emissions pathway, Arctic Ocean




Figure 2. Emergent constraint on Arctic Ocean C_{ant} inventory in 2100. Maximum Arctic Ocean sea surface density (95th percentile) and Arctic Ocean C_{ant} inventory in 2100 for 12-14 ESMs for **a)** SSP1-2.6 (blue), **c)** SSP2-4.5 (green), **e)** SSP3-7.0 (orange), and **g)** SSP5-8.5 (red). The ordinary least squares regressions (dashed lines) and the \pm one σ prediction intervals (shaded area) are shown for each SSP. Probability density functions before and after applying the emergent constraint for **b)** SSP1-2.6, **d)** SSP2-4.5, **f)** SSP3-7.0, and **h)** SSP5-8.5.

waters above 500 m are projected to remain supersaturated with respect to aragonite, while those below 500 are projected to be undersaturated. For the more stable calcium carbonate polymorph calcite, basin-wide Arctic Ocean undersaturation is only




Figure 3. Basin-wide Arctic Ocean calcium carbonate saturation states over the 21st century. Time series of multi-model mean basin-wide saturation states of aragonite from 2000 to 2014 (black) and from 2015 to 2100 for SSP1-2.6 (blue), SSP2-4.5 (green), SSP3-7.0 (orange), and SSP5-8.5 (red) averaged **a)** from 0 to 1000 m, **c)** from 0 to 50 m, **e)** from 50 to 500 m, and **g)** from 500 to 1000 m with \pm one standard deviation ($n=12-14$) shown as shaded area. The differences when changing from historical simulations to the SSPs result from the different number of ensemble members that are available for each simulation. **b), d), f), h)** the same time series but for calcite.

projected under SSP5-8.5. Under this high-emissions pathway, basin-wide undersaturation is projected for the water masses
 235 between 50 and 800 m.

Alongside declines in calcium carbonate saturation states, pH and $p\text{CO}_2$ are also projected to change over the 21st century. Basin-wide averaged pH in the first upper 1000 m decrease from 8.06 in 2002 to 7.88 ± 0.02 (SSP1-2.6), 7.79 ± 0.03 (SSP2-4.5), 7.68 ± 0.03 (SSP3-7.0), and 7.61 ± 0.03 (SSP5-8.5), while basin-wide averaged $p\text{CO}_2$ in the first upper 1000 m increases to $739 \pm 38 \mu\text{atm}$ (SSP1-2.6), $918 \pm 75 \mu\text{atm}$ (SSP2-4.5), $1209 \pm 92 \mu\text{atm}$ (SSP3-7.0), and $1428 \pm 124 \mu\text{atm}$ (SSP5-8.5). The
 240 lowest pH values and highest $p\text{CO}_2$ values are projected to occur between 100 and 1000 m from the surface (Fig. 4).




Figure 4. Vertical profile of basin-wide Arctic Ocean calcium carbonate saturation states, pH and $p\text{CO}_2$ in 2100. Multi-model mean vertical profiles of **a)** aragonite and **b)** calcite saturation states, **c)** pH, and **d)** $p\text{CO}_2$ in 2100 for SSP1-2.6 (blue), SSP2-4.5 (green), SSP3-7.0 (orange), and SSP5-8.5 (red) averaged **a)** with \pm one standard deviation ($n=12-14$) shown as shaded area. Observed profiles in 2002 from GLODAPv2 are shown in black (Lauvset et al., 2016)

3.2.2 Emergent constraints on acidification

The emergent relationship between observed maximum sea surface density and end-of-century Arctic Ocean acidification that was previously identified in CMIP5 models (Terhaar et al., 2020a) does not exist anymore in the new CMIP6 model generation (Fig. 5). Despite the emergent constraint approach still functioning for projections of the Arctic Ocean C_{ant} inventory in 245 CMIP6, this no longer translates into an emergent constraint for Arctic Ocean $\Omega_{\text{arag/calc}}$. However, if all changes in ocean biogeochemistry with the exception of alkalinity, which is kept constant at observed levels from 2002 (Lauvset et al., 2016), are considered when estimating changes in $\Omega_{\text{arag/calc}}$, emergent relationships are still found between maximum sea surface density and Ω_{arag} across all SSPs ($r^2 = 0.54-0.73$). When decreases in alkalinity (A_T) are also taken into account, $\Omega_{\text{arag/calc}}$ decrease further and the emergent relationship disappears ($r^2 = 0.01-0.12$).

250 3.3 Comparison between CMIP5 and CMIP6

Compared to ESMs from CMIP5, the new generation of ESMs (CMIP6) has improved in simulating the maximum Arctic Ocean sea surface density. Specifically, negative density biases have been reduced and the inter-model range in maximum sea surface density has substantially decreased from 3.6 kg m^{-3} in CMIP5 (Terhaar et al., 2020a) to 0.9 kg m^{-3} in CMIP6 (Fig. 2). As a result, the inter-model range of the C_{ant} inventory in the CMIP6 model ensemble is also reduced (Fig. 2). Moreover, 255 without the negative maximum sea surface density bias, the simulated multi-model mean Arctic Ocean C_{ant} inventory in 2005 is 61% higher than the inventory that was simulated by the previous model generation ($1.3 \pm 0.7 \text{ Pg C}$) (Terhaar et al., 2020a).




Figure 5. The CMIP6 ensemble relationship between end-of-century Ω_{arag} and the C_{ant} inventory in the Arctic Ocean. Basin-wide Ω_{arag} averaged over the first upper 1000 m with (orange) and without (blue) taking into account concurrent changes in alkalinity (A_T) against the Arctic Ocean C_{ant} inventory in year 2100 for **a)** SSP1-2.6, **b)** SSP2-4.5, **c)** SSP3-7.0, and **d)** SSP5-8.5.

At the end of the 21st century, the unconstrained simulated Arctic Ocean C_{ant} inventory under SSP5-8.5 is 37% larger and the uncertainty is 19% smaller than the unconstrained simulated C_{ant} inventory in 2100 under RCP8.5 (7.5 ± 2.9 Pg C) (Terhaar et al., 2020a). After applying the constraint, the Arctic Ocean C_{ant} inventory in 2100 under SSP5-8.5 (10.7 ± 1.4 Pg C) is 19% larger than the constrained Arctic Ocean C_{ant} inventory under RCP8.5 (9.0 ± 1.6 Pg C). This difference is of the same order of magnitude as the difference in prescribed atmospheric CO_2 concentration over the 21st century, which is higher in SSP5-8.5 (CMIP6) than RCP8.5 (CMIP5) (Meinshausen et al., 2011, 2019) and therefore results in greater surface ocean acidification for approximately the same radiative forcing (Kwiatkowski et al., 2020).

To compare the emergent constraint across scenarios with different atmospheric CO_2 concentrations, the simulated Arctic Ocean C_{ant} inventory in 2100 for each scenario was rescaled to SSP5-8.5 using the mean atmospheric C_{ant} concentration from 1850 to 2100 as a linear scaling factor (Fig. 6). The relationship remains robust ($r^2=0.63\text{--}0.74$) for all five analysed scenarios. The slope of the emergent relationship is however substantially steeper in CMIP6 ($9.4\text{--}12.6$ Pg C kg^{-1} m^3) than in CMIP5 (3.3 Pg C kg^{-1} m^3). However, the slope in CMIP5 increases to 8.9 Pg C kg^{-1} m^3 if the three CMIP5 models with particularly low maximum sea surface densities (<27.5 kg m^{-3}) are excluded (dotted line in Fig. 6). The resulting constrained estimate for the rescaled Arctic Ocean C_{ant} inventory decreases from the low-emission scenario to the high-emission scenario from 12.3 to 10.7 Pg C. When comparing the two high-emission scenarios, the rescaled Arctic Ocean C_{ant} inventories are 10.7 Pg C (SSP5-8.5) and 10.4 Pg C (RCP8.5). The latter remains unchanged if the three CMIP5 models with particularly low maximum sea surface densities (<27.5 kg m^{-3}) are excluded.




Figure 6. Arctic Ocean C_{ant} inventory in 2100, rescaled to SSP5-8.5 atmospheric C_{ant} concentrations (see Methods), against maximum Arctic Ocean sea surface density (95th percentile) for SSP1-2.6 (blue; $r^2=0.70$), SSP2-4.5 (green; $r^2=0.63$), SSP3-7.0 (orange; $r^2=0.70$), SSP5-8.5 (red; $r^2=0.72$), and RCP8.5 (magenta; $r^2=0.74$) **a**) over the entire density range and **b**) over the density range from 27.5 to 28.3 kg m^{-3} . The simulated Arctic Ocean C_{ant} inventory in 2100 for each scenario (Fig. 2) was rescaled to SSP5-8.5 using the mean atmospheric C_{ant} concentration from 1850 to 2100 as a scaling factor. Linear fits use all available models for each scenario (dashed lines). For RCP8.5 an additional fit is shown excluding the three models with density anomalies below 27.5 kg m^{-3} **b**) (dotted line; $r^2=0.79$).

4 Discussion

275 4.1 Arctic Ocean C_{ant} inventory

Across the CMIP6 model ensemble, the Arctic Ocean C_{ant} storage over the 21st century is highly related to maximum sea surface densities (Fig. 2), which predominately occur in the Barents Sea (Midttun, 1985; Smedsrud et al., 2013; Terhaar et al., 2020a). The inter-model range in maximum sea surface density in the Barents Sea is mainly explained by differences in sea surface ~~salinities~~ salinities ($r^2=0.93$), which are influenced by brine rejection (Midttun, 1985; Årthun et al., 2011) and the strength of inflowing, saltier Atlantic waters through the Barents Sea Opening (Fig. 7). Compared to CMIP5 models, the reduced negative bias of simulated maximum sea surface densities (Fig. 6) indicates model improvement in simulating the circulation of Atlantic and Arctic surface waters. Despite the reduced inter-model range, the robust relationship between maximum sea surface density and C_{ant} across model generations (Fig. 6) supports evidence that inflowing Atlantic waters through the Barents Sea Opening and their transformation into deep and intermediate waters via brine rejection are the dominant process governing Arctic Ocean C_{ant} increases (Midttun, 1985; Rudels et al., 1994, 2000; Jeansson et al., 2011; Smedsrud et al., 2013). The application of observational constraints to this emergent multi-model relationship in order to constrain the projected Arctic Ocean C_{ant} inventory (Fig. 2) and focus model development therefore remains promising.

However, the slope of the linear relationship between maximum sea surface density and the Arctic Ocean C_{ant} inventory over the 21st century in the CMIP5 model ensemble ($3.3 \pm 0.6 \text{ Pg C kg}^{-1} \text{ m}^3$, scaled to SSP5-8.5 atmospheric C_{ant} concentrations) is three to four times less than that in the CMIP6 model ensemble ($12.6 \pm 2.6 \text{ Pg C kg}^{-1} \text{ m}^3$ for SSP1-2.6, $12.1 \pm 2.8 \text{ Pg C kg}^{-1} \text{ m}^3$ for SSP2-4.5, $9.4 \pm 1.9 \text{ Pg C kg}^{-1} \text{ m}^3$ for SSP3-7.0, and $11.3 \pm 2.0 \text{ Pg C kg}^{-1} \text{ m}^3$ for SSP5-8.5) (Fig. 6). The reduced slope in the CMIP5 ensemble is mainly caused by three models with maximum surface density anomalies well below




Figure 7. Annual mean sea surface salinity in the Barents Sea and its neighbouring waters from **a)** the World Ocean Atlas 2018, **b)** the CMIP6 model with the highest (UKESM1-0-LL) and **c)** lowest (CESM2-WACCM) maximum sea surface density and salinity. The 0°C isotherm indicates approximately the extent of Atlantic waters (black solid line).

27.5 kg m⁻³. Excluding these three models, the remaining CMIP5 models follow a slope of 8.9 ± 2.2 Pg C kg⁻¹ m³, in broad agreement with the CMIP6 model ensemble (Fig. 6). This suggests that the linear emergent relationship does not hold below a certain value of maximum sea surface density below which the impact on deep-water formation and subsequent C_{ant} storage in the Arctic Ocean is limited. However, as the two linear relationships happen to cross the observed maximum sea surface density in nearly the same location (Fig. 6), the constrained C_{ant} inventory for the CMIP5 model ensemble remains almost entirely unchanged when the three low-density models are excluded.

Even without the low-density bias in the Barents Sea, the constrained Arctic Ocean C_{ant} inventory in 2005 in CMIP6 remains 36% below the data-based estimate (Terhaar et al., 2020b). This underestimation is partly due to the different definition of C_{ant} in data-based estimates and ESMs. While the historical simulations in CMIP5 and CMIP6 typically start in 1850, data-based estimates account for all C_{ant} since 1765. This leads to an underestimation of the global ocean C_{ant} inventory by ESMs of around 30% (Bronsele et al., 2017) and of around 20% in the Arctic Ocean (Terhaar et al., 2019b). Even if we increased the constrained Arctic Ocean C_{ant} inventory in 2005 by 20%, an underestimation of around 16% remains compared to the data-based estimate. This underestimation of the data-based estimate suggests that all ESMs are missing additional pathways of C_{ant} entry into the Arctic Ocean, other than the principal pathway via the Barents Sea. Indeed, small-scale density flows along continental slopes can be observed in different regions of the Arctic Ocean (Rudels et al., 1994; Jones et al., 1995) but cannot be simulated by the coarse resolution of most ESMs. As such, the constrained estimates of the Arctic Ocean C_{ant} inventory presented here are likely still a lower boundary.

Recent observation of dilution of A_T and C_T in surface waters of the Amerasian basins caused by freshening have led to the hypothesis that continuous freshening might turn the Arctic Ocean from a sink of C_{ant} into a source over the 21st century (Woosley and Millero, 2020). However, observations in the Eurasian basins, which receive more saline Atlantic water and less freshwater input, still show increases of C_{ant} concentrations (Ulfso et al., 2018) over a depth of 1500 m in the last 20 years. The CMIP5 and CMIP6 model ensembles both simulate continuous accumulation of C_{ant} in the Arctic Ocean under all SSPs (Fig. 1), suggesting that the subduction of C_{ant}-rich Atlantic waters in the Barents Sea remains larger than any loss of C_{ant} in surface waters over the 21st century. Nevertheless, the reduction of the storage rate of C_{ant} under SSP5-8.5 (Fig. 1) in combination with constantly increasing atmospheric CO₂ concentrations (Riahi et al., 2017) indicates that dilution may reduce the capacity of the Arctic Ocean to store further C_{ant} as suggested by Woosley and Millero (2020).

4.2 Arctic Ocean acidification

320 4.2.1 $\Omega_{\text{arag/calc}}$, pH, and $p\text{CO}_2$ in 2100

Even under the most optimistic scenario assessed (SSP1-2.6), the Arctic Ocean will become on average undersaturated with respect to aragonite with possible consequences for calcifying organisms (Comeau et al., 2010) and the food chain (Armstrong et al., 2005; Karnovsky et al., 2008). In this scenario only water masses above 500 m remain supersaturated this century. These findings are in good agreement with idealized estimates of $\Omega_{\text{arag/calc}}$ that project the future C_{ant} inventory based on the
325 Transient Time Distribution method and observed CFC-12 concentrations (Anderson et al., 2010; Terhaar et al., 2020b).

In addition to widespread Arctic Ocean undersaturation with respect to aragonite, water masses between 50 and 800 m are even projected to become undersaturated with respect to calcite by the end of the century under SSP5-8.5. Calcite undersaturation in Arctic Ocean subsurface waters will likely further enhance the pressure on the Arctic Ocean ecosystem as calcite forming organisms, such as foraminifera (Davis et al., 2017) and coccolithophores (Kottmeier et al., 2016), experience potential
330 impacts on growth and survival.

Furthermore, the projected increases in $p\text{CO}_2$ (Fig. 4c, d) alongside projected increases in its seasonal amplitude (McNeil and Sasse, 2016; Kwiatkowski and Orr, 2018) is likely to lead to hypercapnic conditions that might directly affect the growth and survival of Arctic fish (Frommel et al., 2012; Schmidt et al., 2017; Kunz et al., 2018) under high emissions scenarios.

4.2.2 Driving processes of acidification

335 The CMIP6 model ensemble shows astonishingly good agreement with respect to projections of Arctic Ocean acidification over the 21st century. Compared to the CMIP5 model ensemble, the uncertainties in projected Ω_{arag} averaged over the [first upper](#) 1000 m in 2100 have been reduced from 0.13 to 0.04–0.06 and those for Ω_{calc} from 0.21 to 0.05–0.09.

The main reason for the reduced uncertainty appears to be that the ESMs that have a lower maximum sea surface density and that thus take up less C_{ant} in the Arctic Ocean over the 21st century (Fig. 2) are the same models that simulate a stronger
340 reduction in A_{T} (Fig. 8). To explain this negative correlation between C_{ant} uptake and reductions in A_{T} , we propose the following mechanism: Over the 21st century, melting of sea ice, land ice, and increased river runoff are projected to freshen the Arctic Ocean (Koenigk et al., 2013; Nummelin et al., 2016; Shu et al., 2018) and to reduce A_{T} (Fig. 8) (Woosley and Millero, 2020). In the CMIP6 ensemble, this freshening and reduction of A_{T} tends to be stronger in models with lower sea surface salinities and densities, i.e. the models that simulate less inflow of saline Atlantic waters such as CESM2-WACCM (Fig. 7),
345 have lower Barents Sea surface salinity and density, weak deep-water formation, and therefore less C_{ant} storage (Fig. 2). In contrast, models with a larger inflow of saline Atlantic water have stronger deep-water formation, greater C_{ant} storage, and a smaller freshening and reduction in A_{T} . This compensation then results in a similar simulated reduction of $\Omega_{\text{arag/calc}}$ over the 21st century across all models in the CMIP6 ensemble, either through a reduction in A_{T} or an increase in C_{T} . Other possible drivers, such as changes in temperature and salinity, are likely of minor importance given the relatively good correlation
350 between the projected C_{ant} inventory and basin-wide reduction of $\Omega_{\text{arag/calc}}$ over the 21st century when A_{T} is not taken into account ($r^2=0.54\text{--}0.73$).




Figure 8. CMIP6 basin-wide changes in alkalinity (A_T) averaged over the first-upper 1000 m from 2002 to 2100 caused by fresh water dilution (blue) and changes in biogeochemistry (green) against the Arctic Ocean C_{ant} inventory for **a)** SSP1-2.6, **b)** SSP2-4.5, **c)** SSP3-7.0, and **d)** SSP5-8.5. A_T anomalies were decomposed by calculating the temporal evolution of salinity corrected alkalinity following Lovenduski et al. (2007).

In the CMIP5 model ensemble, this compensation effect did not exist and projected saturation states and pH were mainly driven by the projected C_{ant} storage in each model (Terhaar et al., 2020a) with changes in A_T , temperature, and salinity of minor importance. The relatively small changes in A_T in CMIP5 compared to CMIP6 might have been caused by an underestimation of Arctic Ocean freshening over the 21st century given that the CMIP5 models also underestimated historical Arctic Ocean freshening by around 50% in the Arctic Ocean (Shu et al., 2018). In the CMIP6 model ensemble this negative bias with respect to freshwater fluxes and subsequent dilution of A_T appears to have been reduced. This is probably a consequence of better representation of riverine fluxes as well as increased model resolution and hence improved circulation (Séférian et al., 2020).

To estimate the likely impacts of C_T increases and A_T decreases to Arctic Ocean acidification over the 21st century, one can assess their relative contributions to acidification at constrained estimates of end-of-century C_{ant} (dotted line in Fig 5). In SSP5-8.5 for example, increasing C_T along with changes in temperature and salinity reduce Ω_{arag} from 1.38 to 0.75, while decreasing A_T further reduces this to 0.59. Thus, the future storage of C_{ant} in the Arctic Ocean still remains the main driver of Arctic Ocean acidification, with acidification further enhanced by decreases in A_T .

5 Conclusions

Earth System Models tend to have improved their performance in the Arctic Ocean from CMIP5 to CMIP6. The negative bias with respect to present-day maximum sea surface density is substantially reduced in CMIP6 and the historical C_{ant} inventory is closer to the data-based estimate (Terhaar et al., 2020b). Furthermore, the inter-model range of projected $\Omega_{arag/calc}$ and the

C_{ant} inventory have been reduced. Uncertainties in the future C_{ant} inventory under four shared socioeconomic pathways can be further reduced by applying the previously identified emergent constraint between present-day maximum Arctic Ocean sea surface density and the future C_{ant} inventory (Terhaar et al., 2020a).

Hall et al. (2019) define “confirmed” emergent constraints as fulfilling the following criteria: 1) a plausible mechanism, 2) verification of that mechanism, and 3) out-of-sample testing. The mechanism underpinning the relationship between maximum sea surface densities and the projected C_{ant} inventory in the Arctic Ocean is consistent with observations and has been verified in an ocean-only model at different resolutions (Terhaar et al., 2020a). The testing of this emergent constraint in the CMIP6 ensemble can be considered out-of-sampling, although the extent of model independence between generations remains questionable (Knutti et al., 2013; Sanderson et al., 2015). Despite the overall emergent constraint on projected Arctic Ocean C_{ant} being similar between CMIP6 and CMIP5, the change in the multi-model emergent relationship indicates that further validation is required. Specifically, it appears that the linear relationship between maximum sea surface densities and the projected C_{ant} inventory breaks down in models with extremely low-biased sea surface densities, highlighting a potential limitation to this emergent constraint that requires further assessment.

While the mechanisms underlying emergent constraints on the future Arctic Ocean C_{ant} inventory appear to be consistent between CMIP5 and CMIP6, this is not the case for the previously identified emergent constraint on ocean acidification variables ($\Omega_{\text{arag/calc}}$, pH, and $p\text{CO}_2$). In the CMIP6 model ensemble, projected model uncertainties in $\Omega_{\text{arag/calc}}$, pH, and $p\text{CO}_2$ are dramatically reduced [compared to the uncertainties](#) in CMIP5 and are not only driven by the increase in C_{ant} , but also by a reduction in A_T due to pan-Arctic freshening. A weak inflow of saline Atlantic waters in ESMs results in lighter surface waters and less C_{ant} storage over the 21st century but also in a stronger reduction of alkalinity caused by a stronger freshening. A strong Atlantic inflow however, appears to limit freshening and reductions in alkalinity but results in greater C_{ant} storage. Although both C_{ant} storage and alkalinity reductions contribute considerably to Arctic Ocean acidification, our results suggest that C_{ant} remains the dominant process. In the CMIP5 model ensemble, the influence of freshening on emergent constraints on acidification variables might have been of limited importance because a) freshening was largely underestimated (Shu et al., 2018) and b) differences in simulating the inflow of Atlantic waters through the Barents Sea Opening were considerably larger across the model ensemble (Terhaar et al., 2020a). As such, the absence of an emergent constraint on projected Arctic acidification in CMIP6 could be viewed as a consequence of successful model development. Indeed, there is clearly no need for such constraints when models are in broad agreement.

Independent of the driving mechanism, the projected Arctic Ocean acidification over the 21st century has possibly grave consequences for the wider Arctic Ocean ecosystem (Gattuso and Hansson, 2011; Riebesell et al., 2013; AMAP, 2018). Under all socio-economic pathways, the upper 1000 m of the Arctic Ocean will be on average undersaturated with respect to aragonite. As such, keystone species like the sea butterfly (aragonitic pteropod *Limacina helicina*) may lose most to all of their suitable habitat in the Arctic Ocean (Comeau et al., 2010). However, under the low-emissions pathway SSP1-2.6 aragonite undersaturation is projected to be avoided in the upper 500 m of the water column, providing a potential refuge. Under the high-emissions pathway SSP5-8.5, water masses between 50 and 800 m are additionally projected to become undersaturated with respect to calcite. While isolated regions of the surface Arctic Ocean, strongly influenced by riverine inputs, are already seasonally un-

undersaturated with respect to calcite (Bates and Mathis, 2009), subsurface Arctic waters would be the first to exhibit annually averaged and basin-wide calcite undersaturation over a depth of several hundred meters. Vertically migrating organisms that form calcite shells and skeletons, such as coccolithophores (Kottmeier et al., 2016) and foraminifera (Davis et al., 2017), may lose their natural refugia ~~area~~ to which they migrate during the day (Berge et al., 2015). These hostile conditions for calcifying organisms will likely cause their decline with unknown consequences for the wider Arctic food web (Armstrong et al., 2005; Karnovsky et al., 2008) and its iconic species (~~Bates et al., 2009~~)([Jay et al., 2011](#); [AMAP, 2018](#)).

Data availability. The Earth system model output used in this study is available via the Earth System Grid Federation (<https://esgf-node.ipsl.upmc.fr/projects/esgf-ipsl/>). Observations from the World Ocean Atlas 2013 (<https://www.nodc.noaa.gov/OC5/woa18/>) and GLODAPv2 (https://www.nodc.noaa.gov/ocads/oceans/GLODAPv2_2019/) are available via the National Oceanic and Atmospheric Administration.

Author contributions. The study was led by JT, who made the figures and wrote the initial manuscript. TB and LK provided help for the analyses and interpretation of the results. OT processed the CMIP6 model data (download, regriding). All authors contributed to the final manuscript.

415 *Competing interests.* No competing interests are present.

Acknowledgements. This project has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 821003 (4C), No 641816 (CRESCENDO), and No 820989 (COMFORT). The work reflects only the authors' view; the European Commission and their executive agency are not responsible for any use that may be made of the information the work contains. This project has also received funding from the Swiss National Science Foundation (grant no. PP00P2_170687), the Agence Nationale de la Recherche grant ANR-18-ERC2-0001-01 (CONVINCE) and the Research Council of Norway under grant number 275268 (COLUMBIA). We acknowledge the World Climate Research Programme's Working Group on Coupled Modelling, which is responsible for CMIP. For CMIP the US Department of Energy's Program for Climate Model Diagnosis and Intercomparison provided coordinating support and led the development of software infrastructure in partnership with the Global Organisation for Earth System Science Portals. We also thank the IPSL modelling group for the software infrastructure, which facilitated CMIP6 analysis.

425 References

- Allen, M. R. and Ingram, W. J.: Constraints on future changes in climate and the hydrologic cycle, *Nature*, 419, 228–232, <https://doi.org/https://doi.org/10.1038/nature01092>, 2002.
- AMAP: AMAP Assessment 2018: Arctic Ocean Acidification, Tech. rep., Arctic Monitoring and Assessment Programme (AMAP), Tromsø, Norway, 2018.
- 430 Anderson, L., Tanhua, T., Björk, G., Hjalmarsson, S., Jones, E., Jutterström, S., Rudels, B., Swift, J., and Wählström, I.: Arctic ocean shelf–basin interaction: An active continental shelf CO₂ pump and its impact on the degree of calcium carbonate solubility, *Deep-Sea Res. Pt. I*, 57, 869 – 879, <https://doi.org/https://doi.org/10.1016/j.dsr.2010.03.012>, 2010.
- Armstrong, J. L., Boldt, J. L., Cross, A. D., Moss, J. H., Davis, N. D., Myers, K. W., Walker, R. V., Beauchamp, D. A., and Haldorson, L. J.: Distribution, size, and interannual, seasonal and diel food habits of northern Gulf of Alaska juvenile pink salmon, *Oncorhynchus gorbuscha*, *Deep-Sea Res. Pt. II*, 52, 247–265, <https://doi.org/https://doi.org/10.1016/j.dsr2.2004.09.019>, 2005.
- 435 Bates, N. R. and Mathis, J. T.: The Arctic Ocean marine carbon cycle: evaluation of air-sea CO₂ exchanges, ocean acidification impacts and potential feedbacks, *Biogeosciences*, 6, 2433–2459, <https://doi.org/https://doi.org/10.5194/bg-6-2433-2009>, 2009.
- Bates, N. R., Mathis, J. T., and Cooper, L. W.: Ocean acidification and biologically induced seasonality of carbonate mineral saturation states in the western Arctic Ocean, *J. Geophys. Res.*, 114, <https://doi.org/https://doi.org/10.1029/2008JC004862>, 2009.
- 440 Berge, J., Renaud, P. E., Darnis, G., Cottier, F., Last, K., Gabrielsen, T. M., Johnsen, G., Seuthe, L., Weslawski, J. M., Leu, E., Moline, M., Nahrang, J., Søreide, J. E., Varpe, Ø., Lønne, O. J., Daase, M., and Falk-Petersen, S.: In the dark: A review of ecosystem processes during the Arctic polar night, *Prog. Oceanog.*, 139, 258 – 271, <https://doi.org/https://doi.org/10.1016/j.pocean.2015.08.005>, overarching perspectives of contemporary and future ecosystems in the Arctic Ocean, 2015.
- Boé, J., Hall, A., and Qu, X.: September sea-ice cover in the Arctic Ocean projected to vanish by 2100, *Nat. Geosci.*, 2, 341–343, <https://doi.org/https://doi.org/10.1038/ngeo467>, 2009.
- 445 Boucher, O., Denvil, S., Caubel, A., and Foujols, M. A.: IPSL IPSL-CM6A-LR model output prepared for CMIP6 CMIP, Tech. rep., Version 20180803. Earth System Grid Federation, <https://doi.org/https://doi.org/10.22033/ESGF/CMIP6.1534>, 2018a.
- Boucher, O., Denvil, S., Caubel, A., and Foujols, M. A.: IPSL IPSL-CM6A-LR model output prepared for CMIP6 ScenarioMIP, Tech. rep., Version20190119. Earth System Grid Federation, <https://doi.org/https://doi.org/10.22033/ESGF/CMIP6.1532>, 2018b.
- 450 Boucher, O., Servonnat, J., Albright, A. L., Aumont, O., Balkanski, Y., Bastrikov, V., Bekki, S., Bonnet, R., Bony, S., Bopp, L., Braconnot, P., Brockmann, P., Cadule, P., Caubel, A., Cheruy, F., Codron, F., Cozic, A., Cugnet, D., D’Andrea, F., Davini, P., de Lavergne, C., Denvil, S., Deshayes, J., Devilliers, M., Ducharne, A., Dufresne, J.-L., Dupont, E., Éthé, C., Fairhead, L., Falletti, L., Flavoni, S., Foujols, M.-A., Gardoll, S., Gastineau, G., Ghattas, J., Grandpeix, J.-Y., Guenet, B., Guez, Lionel, E., Guilyardi, E., Guimberteau, M., Hauglustaine, D., Hourdin, F., Idelkadi, A., Joussaume, S., Kageyama, M., Khodri, M., Krinner, G., Lebas, N., Levavasseur, G., Lévy, C., Li, L., Lott, F., Lurton, T., Luysaert, S., Madec, G., Madeleine, J.-B., Maignan, F., Marchand, M., Marti, O., Mellul, L., Meurdesoif, Y., Mignot, J., Musat, I., Otlé, C., Peylin, P., Planton, Y., Polcher, J., Rio, C., Rochetin, N., Rousset, C., Sepulchre, P., Sima, A., Swingedouw, D., Thiéblemont, R., Traore, A. K., Vancoppenolle, M., Vial, J., Vialard, J., Viovy, N., and Vuichard, N.: Presentation and Evaluation of the IPSL-CM6A-LR Climate Model, *J. Adv. Model. Eearth Sy.*, 12, e2019MS002 010, <https://doi.org/https://doi.org/10.1029/2019MS002010>, 2020.
- 460 Brient, F.: Reducing uncertainties in climate projections with emergent constraints: Concepts, Examples and Prospects, *Adv. Atm. Sci.*, 37, 1–15, <https://doi.org/https://doi.org/10.1002/2016GL069465>, 2020.

- Bronselaer, B., Winton, M., Russell, J., Sabine, C. L., and Khatiwala, S.: Agreement of CMIP5 simulated and observed ocean anthropogenic CO₂ uptake, *Geophys. Res. Lett.*, 44, 12,298–12,305, <https://doi.org/https://doi.org/10.1002/2017GL074435>, 2017.
- 465 Brown, K. A., Holding, J. M., and Carmack, E. C.: Understanding Regional and Seasonal Variability Is Key to Gaining a Pan-Arctic Perspective on Arctic Ocean Freshening, *Front. Mar. Sci.*, 7, 606, <https://doi.org/https://doi.org/10.3389/fmars.2020.00606>, 2020.
- Brüchert, V., Bröder, L., Sawicka, J. E., Tesi, T., Joye, S. P., Sun, X., Semiletov, I. P., and Samarkin, V. A.: Carbon mineralization in Laptev and East Siberian sea shelf and slope sediment, *Biogeosciences*, 15, 471–490, <https://doi.org/https://doi.org/10.5194/bg-15-471-2018>, 2018.
- 470 Caldwell, P. M., Bretherton, C. S., Zelinka, M. D., Klein, S. A., Santer, B. D., and Sanderson, B. M.: Statistical significance of climate sensitivity predictors obtained by data mining, *Geophys. Res. Lett.*, 41, 1803–1808, <https://doi.org/https://doi.org/10.1002/2014GL059205>, 2014.
- Caldwell, P. M., Zelinka, M. D., and Klein, S. A.: Evaluating Emergent Constraints on Equilibrium Climate Sensitivity, *J. Clim.*, 31, 3921–3942, <https://doi.org/https://doi.org/10.1175/JCLI-D-17-0631.1>, 2018.
- Christian, J. R., Holdsworth, A., Steiner, N., Lee, W. G., Swart, N. C., Denman, K. L., Hayashida, H., and Riche, O. G. J.: The Canadian 475 Ocean Ecosystem, *Geosci. Model Dev.*, in preparation, 2020.
- Comeau, S., Jeffree, R., Teyssié, J.-L., and Gattuso, J.-P.: Response of the Arctic Pteropod *Limacina helicina* to Projected Future Environmental Conditions, *Plos One*, 5, 1–7, <https://doi.org/10.1371/journal.pone.0011362>, 2010.
- Cox, P. M., Pearson, D., Booth, B. B., Friedlingstein, P., Huntingford, C., Jones, C. D., and Luke, C. M.: Sensitivity of tropical carbon to climate change constrained by carbon dioxide variability, *Nature*, 494, 341–344, <https://doi.org/https://doi.org/10.1038/nature11882>, 480 2013.
- Danabasoglu, G.: NCAR CESM2 model output prepared for CMIP6 CMIP, Tech. rep., Version 20190308. Earth System Grid Federation, <https://doi.org/https://doi.org/10.22033/ESGF/CMIP6.2185>, 2019a.
- Danabasoglu, G.: NCAR CESM2 model output prepared for CMIP6 ScenarioMIP, Tech. rep., Version 20190308. Earth System Grid Federation, <https://doi.org/https://doi.org/10.22033/ESGF/CMIP6.2201>, 2019b.
- 485 Danabasoglu, G.: NCAR CESM2-WACCM model output prepared for CMIP6 CMIP, Tech. rep., Version 20190808. Earth System Grid Federation, <https://doi.org/https://doi.org/10.22033/ESGF/CMIP6.10024>, 2019c.
- Danabasoglu, G.: NCAR CESM2-WACCM model output prepared for CMIP6 ScenarioMIP, Tech. rep., Version 20190808. Earth System Grid Federation, <https://doi.org/https://doi.org/10.22033/ESGF/CMIP6.10026>, 2019d.
- 490 Danabasoglu, G., Lamarque, J.-F., Bacmeister, J., Bailey, D. A., DuVivier, A. K., Edwards, J., Emmons, L. K., Fasullo, J., Garcia, R., Gettelman, A., Hannay, C., Holland, M. M., Large, W. G., Lauritzen, P. H., Lawrence, D. M., Lenaerts, J. T. M., Lindsay, K., Lipscomb, W. H., Mills, M. J., Neale, R., Oleson, K. W., Otto-Bliesner, B., Phillips, A. S., Sacks, W., Tilmes, S., van Kampenhout, L., Vertenstein, M., Bertini, A., Dennis, J., Deser, C., Fischer, C., Fox-Kemper, B., Kay, J. E., Kinnison, D., Kushner, P. J., Larson, V. E., Long, M. C., Mickelson, S., Moore, J. K., Nienhouse, E., Polvani, L., Rasch, P. J., and Strand, W. G.: The Community Earth System Model Version 2 (CESM2), *J. Adv. Model. Earth Sy.*, 12, e2019MS001916, <https://doi.org/10.1029/2019MS001916>, 2020.
- 495 Davis, C. V., Rivest, E. B., Hill, T. M., Gaylord, B., Russell, A. D., and Sanford, E.: Ocean acidification compromises a planktic calcifier with implications for global carbon cycling, *Sci. Rep.*, 7, 1–8, <https://doi.org/https://doi.org/10.1038/s41598-017-01530-9>, 2017.
- DeAngelis, A. M., Qu, X., and Hall, A.: Importance of vegetation processes for model spread in the fast precipitation response to CO₂ forcing, *Geophys. Res. Lett.*, 43, 12 550–12 559, <https://doi.org/https://doi.org/10.1002/2016GL071392>, 2016.

- Dickson, A. G., Sabine, C. L., and Christian, J. R.: Guide to best practices for ocean CO₂ measurements, Tech. rep., PICES Special Publication 3, 191 pp., 2007.
- 500 Docquier, D., Grist, J. P., Roberts, M. J., Roberts, C. D., Semmler, T., Ponsoni, L., Massonnet, F., Sidorenko, D., Sein, D. V., Iovino, D., et al.: Impact of model resolution on Arctic sea ice and North Atlantic Ocean heat transport, *Clim. Dynam.*, 53, 4989–5017, <https://doi.org/https://doi.org/10.1007/s00382-019-04840-y>, 2019.
- Dunne, J. P., Bociu, I., Bronselaer, B., Guo, H., John, J. G., Krasting, J. P., Stock, C. A., Winton, M., and Zadeh, N.: Simple Global Ocean Biogeochemistry with Light, Iron, Nutrients and Gas version 2 (BLINGv2): Model description and simulation characteristics in GFDL's CM4.0, *J. Adv. Model. Earth Sy.*, p. e2019MS002008, <https://doi.org/https://doi.org/10.1029/2019MS002008>, 2020a.
- 505 Dunne, J. P., Horowitz, L. W., Adcroft, A. J., Ginoux, P., Held, I. M., John, J. G., Krasting, J. P., Malyshev, S., Naik, V., Paulot, F., Shevliakova, E., Stock, C. A., Zadeh, N., Balaji, V., Blanton, C., Dunne, K. A., Dupuis, C., Durachta, J., Dussin, R., Gauthier, P. P. G., Griffies, S. M., Guo, H., Hallberg, R. W., Harrison, M., He, J., Hurlin, W., McHugh, C., Menzel, R., Milly, P. C. D., Nikonov, S., Paynter, D. J., Ploshay, J., Radhakrishnan, A., Rand, K., Reichl, B. G., Robinson, T., Schwarzkopf, D. M., Sentman, L. T., Underwood, S., Vahlenkamp, H., Winton, M., Wittenberg, A. T., Wyman, B., Zeng, Y., and Zhao, M.: The GFDL Earth System Model version 4.1 (GFDL-ESM 4.1): Overall coupled model description and simulation characteristics, *J. Adv. Model. Earth Sy.*, p. e2019MS002015, <https://doi.org/https://doi.org/10.1029/2019MS002015>, 2020b.
- 510 Eyring, V., Cox, P. M., Flato, G. M., Gleckler, P. J., Abramowitz, G., Caldwell, P., Collins, W. D., Gier, B. K., Hall, A. D., Hoffman, F. M., et al.: Taking climate model evaluation to the next level, *Nat. Clim. Change*, 9, 102–110, <https://doi.org/https://doi.org/10.1038/s41558-018-0355-y>, 2019.
- Fabry, V. J., McClintock, J. B., Mathus, J. T., and Grebmeier, J. M.: Ocean Acidification at High Latitudes: The Bellwether, *Oceanography*, 22, 160–171, <https://doi.org/https://doi.org/10.5670/oceanog.2009.105>, 2009.
- Friedlingstein, P., Jones, M. W., O'Sullivan, M., Andrew, R. M., Hauck, J., Peters, G. P., Peters, W., Pongratz, J., Sitch, S., Le Quéré, C., Bakker, D. C. E., Canadell, J. G., Ciais, P., Jackson, R. B., Anthoni, P., Barbero, L., Bastos, A., Bastrikov, V., Becker, M., Bopp, L., Buitenhuis, E., Chandra, N., Chevallier, F., Chini, L. P., Currie, K. I., Feely, R. A., Gehlen, M., Gilfillan, D., Gkritzalis, T., Goll, D. S., Gruber, N., Gutekunst, S., Harris, I., Haverd, V., Houghton, R. A., Hurtt, G., Ilyina, T., Jain, A. K., Joetzjer, E., Kaplan, J. O., Kato, E., Klein Goldewijk, K., Korsbakken, J. I., Landschützer, P., Lauvset, S. K., Lefèvre, N., Lenton, A., Lienert, S., Lombardozzi, D., Marland, G., McGuire, P. C., Melton, J. R., Metzl, N., Munro, D. R., Nabel, J. E. M. S., Nakaoka, S.-I., Neill, C., Omar, A. M., Ono, T., Peregón, A., Pierrot, D., Poulter, B., Rehder, G., Resplandy, L., Robertson, E., Rödenbeck, C., Séférian, R., Schwinger, J., Smith, N., Tans, P. P., Tian, H., Tilbrook, B., Tubiello, F. N., van der Werf, G. R., Wiltshire, A. J., and Zaehle, S.: Global Carbon Budget 2019, *Earth Sys. Sci. Data*, 11, 1783–1838, <https://doi.org/10.5194/essd-11-1783-2019>, 2019.
- 525 Frommel, A. Y., Maneja, R., Lowe, D., Malzahn, A. M., Geffen, A. J., Folkvord, A., Piatkowski, U., Reusch, T. B., and Clemmesen, C.: Severe tissue damage in Atlantic cod larvae under increasing ocean acidification, *Nat. Clim. Change*, 2, 42–46, <https://doi.org/https://doi.org/10.1038/nclimate1324>, 2012.
- Gattuso, J.-P. and Hansson, L.: Ocean acidification, Oxford University Press, Oxford, 2011.
- Gerber, M., Joos, F., Vázquez-Rodríguez, M., Touratier, F., and Goyet, C.: Regional air-sea fluxes of anthropogenic carbon inferred with an Ensemble Kalman Filter, *Global Biogeochem. Cy.*, 23, <https://doi.org/https://doi.org/10.1029/2008GB003247>, 2009.
- Gloor, M., Gruber, N., Sarmiento, J., Sabine, C. L., Feely, R. A., and Rödenbeck, C.: A first estimate of present and preindustrial air-sea CO₂ flux patterns based on ocean interior carbon measurements and models, *Geophys. Res. Lett.*, 30, 10–1–10–4, <https://doi.org/https://doi.org/10.1029/2002GL015594>, 2003.
- 535

- Good, P., Sellar, A., Tang, Y., Rumbold, S., Ellis, R., Kelley, D., Kuhlbrodt, T., and Walton, J.: MOHC UKESM1.0-LL model output prepared for CMIP6 ScenarioMIP, Tech. rep., Version 20190726. Earth System Grid Federation, <https://doi.org/https://doi.org/10.22033/ESGF/CMIP6.1567>, 2019.
- 540 Grotheer, H., Meyer, V., Riedel, T., Pfalz, G., Mathieu, L., Hefter, J., Gentz, T., Lantuit, H., Mollenhauer, G., and Fritz, M.: Burial and Origin of Permafrost-Derived Carbon in the Nearshore Zone of the Southern Canadian Beaufort Sea, *Geophys. Res. Lett.*, 47, e2019GL085897, <https://doi.org/https://doi.org/10.1029/2019GL085897>, 2020.
- Gruber, N., Gloor, M., Mikaloff Fletcher, S. E., Doney, S. C., Dutkiewicz, S., Follows, M. J., Gerber, M., Jacobson, A. R., Joos, F., Lindsay, K., Menemenlis, D., Mouchet, A., Müller, S. A., Sarmiento, J. L., and Takahashi, T.: Oceanic sources, sinks, and transport of atmospheric
- 545 CO₂, *Global Biogeochem. Cy.*, 23, <https://doi.org/https://doi.org/10.1029/2008GB003349>, 2009.
- Guo, H., John, J. G., Blanton, C., McHugh, C., Nikonov, S., Radhakrishnan, A., Rand, K., Zadeh, N. T., Balaji, V., Durachta, J., Dupuis, C., Menzel, R., Robinson, T., Underwood, S., Vahlenkamp, H., Bushuk, M., Dunne, K. A., Dussin, R., Gauthier, P. P., Ginoux, P., Griffies, S. M., Hallberg, R., Harrison, M., Hurlin, W., Malyshev, S., Naik, V., Paulot, F., Paynter, D. J., Ploshay, J., Reichl, B. G., Schwarzkopf, D. M., Seman, C. J., Shao, A., Silvers, L., Wyman, B., Yan, X., Zeng, Y., Adcroft, A., Dunne, J. P., Held, I. M., Krasting, J. P., Horowitz,
- 550 L. W., Milly, P., Shevliakova, E., Winton, M., Zhao, M., and Zhang, R.: NOAA-GFDL GFDL-CM4 model output prepared for CMIP6 CMIP, Tech. rep., Version 20180701. Earth System Grid Federation, <https://doi.org/https://doi.org/10.22033/ESGF/CMIP6.1402>, 2018a.
- Guo, H., John, J. G., Blanton, C., McHugh, C., Nikonov, S., Radhakrishnan, A., Zadeh, N. T., Balaji, V., Durachta, J., Dupuis, C., Menzel, R., Robinson, T., Underwood, S., Vahlenkamp, H., Dunne, K. A., Gauthier, P. P., Ginoux, P., Griffies, S. M., Hallberg, R., Harrison, M., Hurlin, W., Lin, P., Malyshev, S., Naik, V., Paulot, F., Paynter, D. J., Ploshay, J., Schwarzkopf, D. M., Seman, C. J., Shao, A., Silvers, L.,
- 555 Wyman, B., Yan, X., Zeng, Y., Adcroft, A., Dunne, J. P., Held, I. M., Krasting, J. P., Horowitz, L. W., Milly, C., Shevliakova, E., Winton, M., Zhao, M., and Zhang, R.: NOAA-GFDL GFDL-CM4 model output prepared for CMIP6 ScenarioMIP, Tech. rep., Version 20180701. Earth System Grid Federation, <https://doi.org/https://doi.org/10.22033/ESGF/CMIP6.9242>, 2018b.
- Hajima, T., Abe, M., Arakawa, O., Suzuki, T., Komuro, Y., Ogura, T., Ogochi, K., Watanabe, M., Yamamoto, A., Tatebe, H., Noguchi, M. A., Ohgaito, R., Ito, A., Yamazaki, D., Ito, A., Takata, K., Watanabe, S., Kawamiya, M., and Tachiiri, K.:
- 560 MIROC MIROC-ES2L model output prepared for CMIP6 CMIP, Tech. rep., Version 20190823. Earth System Grid Federation, <https://doi.org/https://doi.org/10.22033/ESGF/CMIP6.902>, 2019.
- Hajima, T., Watanabe, M., Yamamoto, A., Tatebe, H., Noguchi, M. A., Abe, M., Ohgaito, R., Ito, A., Yamazaki, D., Okajima, H., Ito, A., Takata, K., Ogochi, K., Watanabe, S., and Kawamiya, M.: Development of the MIROC-ES2L Earth system model and the evaluation of biogeochemical processes and feedbacks, *Geosci. Model Dev.*, 13, 2197–2244, <https://doi.org/https://doi.org/10.5194/gmd-13-2197-2020>,
- 565 2020.
- Hall, A. and Qu, X.: Using the current seasonal cycle to constrain snow albedo feedback in future climate change, *Geophys. Res. Lett.*, 33, <https://doi.org/https://doi.org/10.1029/2005GL025127>, 2006.
- Hall, A., Cox, P., Huntingford, C., and Klein, S.: Progressing emergent constraints on future climate change, *Nat. Clim. Change*, 9, 269–278, <https://doi.org/https://doi.org/10.1038/s41558-019-0436-6>, 2019.
- 570 Haugan, P. M. and Drange, H.: Effects of CO₂ on the ocean environment, *Energ. Convers. Manag.*, 37, 1019–1022, [https://doi.org/https://doi.org/10.1016/0196-8904\(95\)00292-8](https://doi.org/https://doi.org/10.1016/0196-8904(95)00292-8), proceedings of the International Energy Agency Greenhouse Gases: Mitigation Options Conference, 1996.
- Held, I. M., Guo, H., Adcroft, A., Dunne, J. P., Horowitz, L. W., Krasting, J., Shevliakova, E., Winton, M., Zhao, M., Bushuk, M., Wittenberg, A. T., Wyman, B., Xiang, B., Zhang, R., Anderson, W., Balaji, V., Donner, L., Dunne, K., Durachta, J., Gauthier, P. P. G., Ginoux, P., Golaz,

- 575 J.-C., Griffies, S. M., Hallberg, R., Harris, L., Harrison, M., Hurlin, W., John, J., Lin, P., Lin, S.-J., Malyshev, S., Menzel, R., Milly, P. C. D., Ming, Y., Naik, V., Paynter, D., Paulot, F., Rammawamy, V., Reichl, B., Robinson, T., Rosati, A., Seman, C., Silvers, L. G., Underwood, S., and Zadeh, N.: Structure and Performance of GFDL's CM4.0 Climate Model, *J. Adv. Model. Earth Sy.*, 11, 3691–3727, <https://doi.org/https://doi.org/10.1029/2019MS001829>, 2019.
- Jakobsson, M.: Hypsometry and volume of the Arctic Ocean and its constituent seas, *Geochem. Geophys. Geosy.*, 3, 1–18, <https://doi.org/https://doi.org/10.1029/2001GC000302>, 2002.
- 580 Jay, C. V., Marcot, B. G., and Douglas, D. C.: Projected status of the Pacific walrus (*Odobenus rosmarus divergens*) in the twenty-first century, *Polar Biol.*, 34, 1065–1084, 2011.
- Jeansson, E., Olsen, A., Eldevik, T., Skjelvan, I., Omar, A. M., Lauvset, S. K., Nilsen, J. E. O., Bellerby, R. G. J., Johannessen, T., and Falck, E.: The Nordic Seas carbon budget: Sources, sinks, and uncertainties, *Global Biogeochem. Cy.*, 25, <https://doi.org/https://doi.org/10.1029/2010GB003961>, 2011.
- 585 John, J. G., Blanton, C., McHugh, C., Radhakrishnan, A., Rand, K., Vahlenkamp, H., Wilson, C., Zadeh, N. T., Gauthier, P. P., Dunne, J. P., Dussin, R., Horowitz, L. W., Lin, P., Malyshev, S., Naik, V., Ploshay, J., Silvers, L., Stock, C., Winton, M., and Zeng, Y.: NOAA-GFDL GFDL-ESM4 model output prepared for CMIP6 ScenarioMIP, Tech. rep., Version 20180701. Earth System Grid Federation, <https://doi.org/https://doi.org/10.22033/ESGF/CMIP6.1414>, 2018.
- 590 Jones, E., Rudels, B., and Anderson, L.: Deep waters of the Arctic Ocean: origins and circulation, *Deep-Sea Res. Pt. I*, 42, 737–760, [https://doi.org/https://doi.org/10.1016/0967-0637\(95\)00013-V](https://doi.org/https://doi.org/10.1016/0967-0637(95)00013-V), 1995.
- Jungclaus, J., Bittner, M., Wieners, K.-H., Wachsmann, F., Schupfner, M., Legutke, S., Giorgetta, M., Reick, C., Gayler, V., Haak, H., de Vrese, P., Raddatz, T., Esch, M., Mauritsen, T., von Storch, J.-S., Behrens, J., Brovkin, V., Claussen, M., Crueger, T., Fast, I., Fiedler, S., Hagemann, S., Hohenegger, C., Jahns, T., Kloster, S., Kinne, S., Lasslop, G., Kornblueh, L., Marotzke, J., Matei, D., Meraner, K., 595 Mikolajewicz, U., Modali, K., Müller, W., Nabel, J., Notz, D., Peters, K., Pincus, R., Pohlmann, H., Pongratz, J., Rast, S., Schmidt, H., Schnur, R., Schulzweida, U., Six, K., Stevens, B., Voigt, A., and Roeckner, E.: MPI-M MPIESM1.2-HR model output prepared for CMIP6 CMIP, Tech. rep., Version 20190710. Earth System Grid Federation, <https://doi.org/https://doi.org/10.22033/ESGF/CMIP6.741>, 2019.
- Karnovsky, N. J., Hobson, K. A., Iverson, S., and Hunt Jr, G. L.: Seasonal changes in diets of seabirds in the North Water Polynya: a multiple-indicator approach, *Mar. Ecol. Prog. Ser.*, 357, 291–299, <https://doi.org/https://doi.org/10.3354/meps07295>, 2008.
- 600 Knutti, R., Masson, D., and Gettelman, A.: Climate model genealogy: Generation CMIP5 and how we got there, *Geophys. Res. Lett.*, 40, 1194–1199, <https://doi.org/https://doi.org/10.1002/grl.50256>, 2013.
- Koenigk, T., Brodeau, L., Graverson, R. G., Karlsson, J., Svensson, G., Tjernström, M., Willén, U., and Wyser, K.: Arctic climate change in 21st century CMIP5 simulations with EC-Earth, *Clim. Dyn.*, 40, 2719–2743, <https://doi.org/https://doi.org/10.1007/s00382-012-1505-y>, 2013.
- 605 Kottmeier, D. M., Rokitta, S. D., and Rost, B.: H⁺-driven increase in CO₂ uptake and decrease in uptake explain coccolithophores' acclimation responses to ocean acidification, *Limnol. Oceanogr.*, 61, 2045–2057, <https://doi.org/https://doi.org/10.1002/lno.10352>, 2016.
- Krasting, J. P., John, J. G., Blanton, C., McHugh, C., Nikonov, S., Radhakrishnan, A., Rand, K., Zadeh, N. T., Balaji, V., Durachta, J., Dupuis, C., Menzel, R., Robinson, T., Underwood, S., Vahlenkamp, H., Dunne, K. A., Gauthier, P. P., Ginoux, P., Griffies, S. M., Hallberg, R., Harrison, M., Hurlin, W., Malyshev, S., Naik, V., Paulot, F., Paynter, D. J., Ploshay, J., Schwarzkopf, D. M., Seman, C. J., Silvers, L., 610 Wyman, B., Zeng, Y., Adcroft, A., Dunne, J. P., Dussin, R., Guo, H., He, J., Held, I. M., Horowitz, L. W., Lin, P., Milly, P., Shevliakova, E., Stock, C., Winton, M., Xie, Y., and Zhao, M.: NOAA-GFDL GFDL-ESM4 model output prepared for CMIP6 CMIP, Tech. rep., Version 20190726. Earth System Grid Federation, <https://doi.org/https://doi.org/10.22033/ESGF/CMIP6.1407>, 2018.

- 615 Kunz, K. L., Claireaux, G., Pörtner, H.-O., Knust, R., and Mark, F. C.: Aerobic capacities and swimming performance of polar cod (*Boreogadus saida*) under ocean acidification and warming conditions, *J. Exp. Biol.*, 221, <https://doi.org/https://doi.org/10.1242/jeb.184473>, 2018.
- Kwiatkowski, L. and Orr, J. C.: Diverging seasonal extremes for ocean acidification during the twenty-first century, *Nat. Clim. Change*, 8, 141–145, <https://doi.org/https://doi.org/10.1038/s41558-017-0054-0>, 2018.
- 620 Kwiatkowski, L., Bopp, L., Aumont, O., Ciais, P., Cox, P. M., Laufkötter, C., Li, Y., and Séférian, R.: Emergent constraints on projections of declining primary production in the tropical oceans, *Nat. Clim. Change*, 7, 355–358, <https://doi.org/https://doi.org/10.1038/nclimate3265>, 2017.
- Kwiatkowski, L., Torres, O., Bopp, L., Aumont, O., Chamberlain, M., Christian, J. R., Dunne, J. P., Gehlen, M., Ilyina, T., John, J. G., Lenton, A., Li, H., Lovenduski, N. S., Orr, J. C., Palmieri, J., Santana-Falcón, Y., Schwinger, J., Séférian, R., Stock, C. A., Tagliabue, A., Takano, Y., Tjiputra, J., Toyama, K., Tsujino, H., Watanabe, M., Yamamoto, A., Yool, A., and Ziehn, T.: Twenty-first century ocean warming, acidification, deoxygenation, and upper-ocean nutrient and primary production decline from CMIP6 model projections, *Biogeosciences*, 625 17, 3439–3470, <https://doi.org/https://doi.org/10.5194/bg-17-3439-2020>, 2020.
- Lauvset, S. K., Key, R. M., and Perez, F. F.: A new global interior ocean mapped climatology: the $1^\circ \times 1^\circ$ GLODAP version 2, *Earth Syst. Sci. Data*, 8, 325, <https://doi.org/https://doi.org/10.5194/essd-8-325-2016>, 2016.
- 630 Locarnini, R. A., Mishonov, A. V., Baranova, O. K., Boyer, T. P., Zweng, M. M., Garcia, H. E., Reagan, J. R., Seidov, D., Weathers, K., Paver, C. R., and Smolyar, I.: World Ocean Atlas 2018, Volume 1: Temperature, Tech. rep., A. Mishonov Technical Ed.; NOAA Atlas NESDIS 81, 2018.
- Lovenduski, N. S., Gruber, N., Doney, S. C., and Lima, I. D.: Enhanced CO₂ outgassing in the Southern Ocean from a positive phase of the Southern Annular Mode, *Gl. Biogeochem. Cy.*, 21, <https://doi.org/https://doi.org/10.1029/2006GB002900>, 2007.
- 635 Mauritsen, T., Bader, J., Becker, T., Behrens, J., Bittner, M., Brokopf, R., Brovkin, V., Claussen, M., Crueger, T., Esch, M., Fast, I., Fiedler, S., Fläschner, D., Gayler, V., Giorgetta, M., Goll, D. S., Haak, H., Hagemann, S., Hedemann, C., Hohenegger, C., Ilyina, T., Jahns, T., Jimenéz-de-la Cuesta, D., Jungclaus, J., Kleinen, T., Kloster, S., Kracher, D., Kinne, S., Kleberg, D., Lasslop, G., Kornbluh, L., Marotzke, J., Matei, D., Meraner, K., Mikolajewicz, U., Modali, K., Möbis, B., Müller, W. A., Nabel, J. E. M. S., Nam, C. C. W., Notz, D., Nyawira, S.-S., Paulsen, H., Peters, K., Pincus, R., Pohlmann, H., Pongratz, J., Popp, M., Raddatz, T. J., Rast, S., Redler, R., Reick, C. H., Rohrschneider, T., Schemann, V., Schmidt, H., Schnur, R., Schulzweida, U., Six, K. D., Stein, L., Stemmler, I., Stevens, B., von Storch, J.-S., Tian, F., Voigt, A., Vrese, P., Wieners, K.-H., Wilkenskjaeld, S., Winkler, A., and Roeckner, E.: Developments in the 640 MPI-M Earth System Model version 1.2 (MPI-ESM1.2) and Its Response to Increasing CO₂, *J. Adv. Model. Earth Sy.*, 11, 998–1038, <https://doi.org/https://doi.org/10.1029/2018MS001400>, 2019.
- McNeil, B. I. and Sasse, T. P.: Future ocean hypercapnia driven by anthropogenic amplification of the natural CO₂ cycle, *Nature*, 529, 383–386, <https://doi.org/https://doi.org/10.1038/nature16156>, 2016.
- 645 Meinshausen, M., Smith, S. J., Calvin, K., Daniel, J. S., Kainuma, M., Lamarque, J.-F., Matsumoto, K., Montzka, S., Raper, S., Riahi, K., et al.: The RCP greenhouse gas concentrations and their extensions from 1765 to 2300, *Climatic change*, 109, 213, <https://doi.org/https://doi.org/10.1007/s10584-011-0156-z>, 2011.
- Meinshausen, M., Vogel, E., Nauels, A., Lorbacher, K., Meinshausen, N., Etheridge, D. M., Fraser, P. J., Montzka, S. A., Rayner, P. J., Trudinger, C. M., Krummel, P. B., Beyerle, U., Canadell, J. G., Daniel, J. S., Enting, I. G., Law, R. M., Lunder, C. R., O’Doherty, S., Prinn, R. G., Reimann, S., Rubino, M., Velders, G. J. M., Vollmer, M. K., Wang, R. H. J., and Weiss, R.: Historical greenhouse gas 650 concentrations for climate modelling (CMIP6), *Geosci. Model Dev.*, 10, 2057–2116, <https://doi.org/10.5194/gmd-10-2057-2017>, 2017.

- Meinshausen, M., Nicholls, Z., Lewis, J., Gidden, M. J., Vogel, E., Freund, M., Beyerle, U., Gessner, C., Nauels, A., Bauer, N., Canadell, J. G., Daniel, J. S., John, A., Krummel, P., Luderer, G., Meinshausen, N., Montzka, S. A., Rayner, P., Reimann, S., Smith, S. J., van den Berg, M., Velders, G. J. M., Vollmer, M., and Wang, H. J.: The SSP greenhouse gas concentrations and their extensions to 2500, *Geosci. Model Dev. Discussions*, 2019, 1–77, <https://doi.org/https://doi.org/10.5194/gmd-2019-222>, 2019.
- 655 Midttun, L.: Formation of dense bottom water in the Barents Sea, *Deep-Sea Res. Pt. I*, 32, 1233–1241, [https://doi.org/https://doi.org/10.1016/0198-0149\(85\)90006-8](https://doi.org/https://doi.org/10.1016/0198-0149(85)90006-8), 1985.
- Mikaloff Fletcher, S. E., Gruber, N., Jacobson, A. R., Doney, S. C., Dutkiewicz, S., Gerber, M., Follows, M., Joos, F., Lindsay, K., Mendenlis, D., Mouchet, A., Müller, S. A., and Sarmiento, J. L.: Inverse estimates of anthropogenic CO₂ uptake, transport, and storage by the ocean, *Global Biogeochem. Cy.*, 20, <https://doi.org/https://doi.org/10.1029/2005GB002530>, 2006.
- 660 Müller, W. A., Jungclaus, J. H., Mauritzen, T., Baehr, J., Bittner, M., Budich, R., Bunzel, F., Esch, M., Ghosh, R., Haak, H., Ilyina, T., Kleine, T., Kornbluh, L., Li, H., Modali, K., Notz, D., Pohlmann, H., Roeckner, E., Stemmler, I., Tian, F., and Marotzke, J.: A Higher-resolution Version of the Max Planck Institute Earth System Model (MPI-ESM1.2-HR), *J. Adv. Model. Earth Sy.*, 10, 1383–1413, <https://doi.org/https://doi.org/10.1029/2017MS001217>, 2018.
- Nummelin, A., Ilicak, M., Li, C., and Smedsrud, L. H.: Consequences of future increased Arctic runoff on Arctic Ocean stratification, circulation, and sea ice cover, *J. Geophys. Res. Oceans*, 121, 617–637, <https://doi.org/https://doi.org/10.1002/2015JC011156>, 2016.
- 665 O’Neill, B. C., Tebaldi, C., van Vuuren, D. P., Eyring, V., Friedlingstein, P., Hurtt, G., Knutti, R., Kriegler, E., Lamarque, J.-F., Lowe, J., Meehl, G. A., Moss, R., Riahi, K., and Sanderson, B. M.: The Scenario Model Intercomparison Project (ScenarioMIP) for CMIP6, *Geosci. Model Dev.*, 9, 3461–3482, <https://doi.org/https://doi.org/10.5194/gmd-9-3461-2016>, 2016.
- Orr, J. C. and Epitalon, J.-M.: Improved routines to model the ocean carbonate system: mocsy 2.0, *Geosci. Model Dev.*, 8, 485–499, <https://doi.org/https://doi.org/10.5194/gmd-8-485-2015>, <https://www.geosci-model-dev.net/8/485/2015/>, 2015.
- 670 Orr, J. C., Fabry, V. J., Aumont, O., Bopp, L., Doney, S. C., Feely, R. A., Gnanadesikan, A., Gruber, N., Ishida, A., Joos, F., et al.: Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms, *Nature*, 437, 681–686, <https://doi.org/https://doi.org/10.1038/nature04095>, 2005.
- Orr, J. C., Najjar, R. G., Aumont, O., Bopp, L., Bullister, J. L., Danabasoglu, G., Doney, S. C., Dunne, J. P., Dutay, J.-C., Graven, H., Griffies, S. M., John, J. G., Joos, F., Levin, I., Lindsay, K., Matear, R. J., McKinley, G. A., Mouchet, A., Oschlies, A., Romanou, A., Schlitzer, R., Tagliabue, A., Tanhua, T., and Yool, A.: Biogeochemical protocols and diagnostics for the CMIP6 Ocean Model Intercomparison Project (OMIP), *Geosci. Model Dev.*, 10, 2169–2199, <https://doi.org/10.5194/gmd-10-2169-2017>, 2017.
- O’Gorman, P. A.: Sensitivity of tropical precipitation extremes to climate change, *Nat. Geosci.*, 5, 697–700, <https://doi.org/https://doi.org/10.1038/ngeo1568>, 2012.
- 680 Rawlins, M. A., Steele, M., Holland, M. M., Adam, J. C., Cherry, J. E., Francis, J. A., Groisman, P. Y., Hinzman, L. D., Huntington, T. G., Kane, D. L., Kimball, J. S., Kwok, R., Lammers, R. B., Lee, C. M., Lettenmaier, D. P., McDonald, K. C., Podest, E., Pundsack, J. W., Rudels, B., Serreze, M. C., Shiklomanov, A., Skagseth, O., Troy, T. J., Vörösmarty, C. J., Wensnahan, M., Wood, E. F., Woodgate, R., Yang, D., Zhang, K., and Zhang, T.: Analysis of the Arctic System for Freshwater Cycle Intensification: Observations and Expectations, *J. Clim.*, 23, 5715–5737, <https://doi.org/https://doi.org/10.1175/2010JCLI3421.1>, 2010.
- 685 Riahi, K., van Vuuren, D. P., Kriegler, E., Edmonds, J., O’Neill, B. C., Fujimori, S., Bauer, N., Calvin, K., Dellink, R., Fricko, O., Lutz, W., Popp, A., Cuaresma, J. C., KC, S., Leimbach, M., Jiang, L., Kram, T., Rao, S., Emmerling, J., Ebi, K., Hasegawa, T., Havlik, P., Humpenöder, F., Da Silva, L. A., Smith, S., Stehfest, E., Bosetti, V., Eom, J., Gernaat, D., Masui, T., Rogelj, J., Strefler, J., Drouet, L., Krey, V., Luderer, G., Harmsen, M., Takahashi, K., Baumstark, L., Doelman, J. C., Kainuma, M., Klimont, Z., Marangoni, G., Lotze-Campen, H.,

- Obersteiner, M., Tabeau, A., and Tavoni, M.: The Shared Socioeconomic Pathways and their energy, land use, and greenhouse gas emissions implications: An overview, *Global Environ. Chang.*, 42, 153–168, <https://doi.org/https://doi.org/10.1016/j.gloenvcha.2016.05.009>, 2017.
- 690 Riebesell, U., Gattuso, J.-P., Thingstad, T. F., and Middelburg, J. J.: Arctic ocean acidification: pelagic ecosystem and biogeochemical responses during a mesocosm study, *Biogeosciences*, 10, 5619–5626, <https://doi.org/https://doi.org/10.5194/bg-10-5619-2013>, 2013.
- Rudels, B.: Arctic Ocean circulation, processes and water masses: A description of observations and ideas with focus on the period prior to the International Polar Year 2007–2009, *Prog. Oceanog.*, 132, 22–67, <https://doi.org/https://doi.org/10.1016/j.pocean.2013.11.006>, 2015.
- 695 Rudels, B., Jones, E. P., Anderson, L. G., and Kattner, G.: On the Intermediate Depth Waters of the Arctic Ocean, pp. 33–46, American Geophysical Union (AGU), <https://doi.org/https://doi.org/10.1029/GM085p0033>, 1994.
- Rudels, B., Muench, R. D., Gunn, J., Schauer, U., and Friedrich, H. J.: Evolution of the Arctic Ocean boundary current north of the Siberian shelves, *J. Mar. Sys.*, 25, 77–99, [https://doi.org/https://doi.org/10.1016/S0924-7963\(00\)00009-9](https://doi.org/https://doi.org/10.1016/S0924-7963(00)00009-9), 2000.
- 700 Sanderson, B. M., Knutti, R., and Caldwell, P.: A Representative Democracy to Reduce Interdependency in a Multimodel Ensemble, *J. Clim.*, 28, 5171–5194, <https://doi.org/https://doi.org/10.1175/JCLI-D-14-00362.1>, 2015.
- Schlund, M., Lauer, A., Gentine, P., Sherwood, S. C., and Eyring, V.: Emergent constraints on Equilibrium Climate Sensitivity in CMIP5: do they hold for CMIP6?, *Earth Syst. Dynam. Discussions*, 2020, 1–40, <https://doi.org/https://doi.org/10.5194/esd-2020-49>, 2020.
- Schmidt, M., Windisch, H. S., Ludwichowski, K.-U., Seegert, S. L. L., Pörtner, H.-O., Storch, D., and Bock, C.: Differences in neurochemical profiles of two gadid species under ocean warming and acidification, *Front. Zool.*, 14, 1–13, <https://doi.org/https://doi.org/10.1186/s12983-017-0238-5>, 2017.
- 705 Schupfner, M., Wieners, K.-H., Wachsmann, F., Steger, C., Bittner, M., Jungclaus, J., Früh, B., Pankatz, K., Giorgetta, M., Reick, C., Legutke, S., Esch, M., Gayler, V., Haak, H., de Vrese, P., Raddatz, T., Mauritsen, T., von Storch, J.-S., Behrens, J., Brovkin, V., Claussen, M., Crueger, T., Fast, I., Fiedler, S., Hagemann, S., Hohenegger, C., Jahns, T., Kloster, S., Kinne, S., Lasslop, G., Kornblueh, L., Marotzke, J., Matei, D., Meraner, K., Mikolajewicz, U., Modali, K., Müller, W., Nabel, J., Notz, D., Peters, K., Pincus, R., Pohlmann, H., Pongratz, J., Rast, S., Schmidt, H., Schnur, R., Schulzweida, U., Six, K., Stevens, B., Voigt, A., and Roeckner, E.: DKRZ MPI-ESM1.2-HR model output prepared for CMIP6 ScenarioMIP, <https://doi.org/https://doi.org/10.22033/ESGF/CMIP6.2450>, 2019.
- Seferian, R.: CNRM-CERFACS CNRM-ESM2-1 model output prepared for CMIP6 CMIP, Tech. rep., Version 20181206. Earth System Grid Federation, <https://doi.org/https://doi.org/10.22033/ESGF/CMIP6.1391>, 2018a.
- 715 Seferian, R.: CNRM-CERFACS CNRM-ESM2-1 model output prepared for CMIP6 ScenarioMIP, Tech. rep., Version 20191021. Earth System Grid Federation, <https://doi.org/https://doi.org/10.22033/ESGF/CMIP6.1395>, 2018b.
- Séférian, R., Berthet, S., Yool, A., Palmieri, J., Bopp, L., Tagliabue, A., Kwiatkowski, L., Aumont, O., Christian, J., Dunne, J., et al.: Tracking improvement in simulated marine biogeochemistry between CMIP5 and CMIP6, *Curr. Clim. Change Rep.*, 6, 95–119, <https://doi.org/https://doi.org/10.1007/s40641-020-00160-0>, 2020.
- 720 Sellar, A. A., Jones, C. G., Mulcahy, J. P., Tang, Y., Yool, A., Wiltshire, A., O’Connor, F. M., Stringer, M., Hill, R., Palmieri, J., Woodward, S., de Mora, L., Kuhlbrodt, T., Rumbold, S. T., Kelley, D. I., Ellis, R., Johnson, C. E., Walton, J., Abraham, N. L., Andrews, M. B., Andrews, T., Archibald, A. T., Berthou, S., Burke, E., Blockley, E., Carslaw, K., Dalvi, M., Edwards, J., Folberth, G. A., Gedney, N., Griffiths, P. T., Harper, A. B., Hendry, M. A., Hewitt, A. J., Johnson, B., Jones, A., Jones, C. D., Keeble, J., Liddicoat, S., Morgenstern, O., Parker, R. J., Predoi, V., Robertson, E., Siahann, A., Smith, R. S., Swaminathan, R., Woodhouse, M. T., Zeng, G., and Zerroukat, M.: UKESM1: Description and Evaluation of the U.K. Earth System Model, *J. Adv. Model. Earth Syst.*, 11, 4513–4558, <https://doi.org/https://doi.org/10.1029/2019MS001739>, 2019.
- 725

- Shu, Q., Qiao, F., Song, Z., Zhao, J., and Li, X.: Projected Freshening of the Arctic Ocean in the 21st Century, *J. Geophys. Res. Oceans*, 123, 9232–9244, <https://doi.org/https://doi.org/10.1029/2018JC014036>, 2018.
- 730 Smedsrud, L. H., Esau, I., Ingvaldsen, R. B., Eldevik, T., Haugan, P. M., Li, C., Lien, V. S., Olsen, A., Omar, A. M., Otterå, O. H., Risebrobakken, B., Sandø, A. B., Semenov, V. A., and Sorokina, S. A.: The role of the Barents Sea in the Arctic climate system, *Rev. Geophys.*, 51, 415–449, <https://doi.org/https://doi.org/10.1002/rog.20017>, 2013.
- Steinacher, M., Joos, F., Frolicher, T. L., Plattner, G.-K., and Doney, S. C.: Imminent ocean acidification in the Arctic projected with the NCAR global coupled carbon cycle-climate model, *Biogeosciences*, 6, 515–533, <https://doi.org/https://doi.org/10.5194/bg-6-515-2009>, 2009.
- 735 Steiner, N. S., Christian, J. R., Six, K. D., Yamamoto, A., and Yamamoto-Kawai, M.: Future ocean acidification in the Canada Basin and surrounding Arctic Ocean from CMIP5 earth system models, *J. Geophys. Res.*, 119, 332–347, <https://doi.org/https://doi.org/10.1002/2013JC009069>, 2013.
- Stock, C. A., Dunne, J. P., Fan, S., Ginoux, P., John, J., Krasting, J. P., Laufkötter, C., Paulot, F., and Zadeh, N.: Ocean Biogeochemistry in GFDL’s Earth System Model 4.1 and its Response to Increasing Atmospheric CO₂, *J. Adv. Model. Earth Sy.*, n/a, e2019MS002043, <https://doi.org/https://doi.org/10.1029/2019MS002043>, 2020.
- 740 Swart, N. C., Cole, J. N., Kharin, V. V., Lazare, M., Scinocca, J. F., Gillett, N. P., Anstey, J., Arora, V., Christian, J. R., Jiao, Y., Lee, W. G., Majaess, F., Saenko, O. A., Seiler, C., Seinen, C., Shao, A., Solheim, L., von Salzen, K., Yang, D., Winter, B., and Sigmond, M.: CCCma CanESM5 model output prepared for CMIP6 CMIP, Tech. rep., Version v20190429. Earth System Grid Federation, <https://doi.org/https://doi.org/10.22033/ESGF/CMIP6.1303>, <https://doi.org/10.22033/ESGF/CMIP6.3673>, 2019a.
- 745 Swart, N. C., Cole, J. N., Kharin, V. V., Lazare, M., Scinocca, J. F., Gillett, N. P., Anstey, J., Arora, V., Christian, J. R., Jiao, Y., Lee, W. G., Majaess, F., Saenko, O. A., Seiler, C., Seinen, C., Shao, A., Solheim, L., von Salzen, K., Yang, D., Winter, B., and Sigmond, M.: CCCma CanESM5 model output prepared for CMIP6 ScenarioMIP, Tech. rep., Version v20190429. Earth System Grid Federation, <https://doi.org/https://doi.org/10.22033/ESGF/CMIP6.1317>, 2019b.
- Swart, N. C., Cole, J. N., Kharin, V. V., Lazare, M., Scinocca, J. F., Gillett, N. P., Anstey, J., Arora, V., Christian, J. R., Jiao, Y., Lee, W. G., Majaess, F., Saenko, O. A., Seiler, C., Seinen, C., Shao, A., Solheim, L., von Salzen, K., Yang, D., Winter, B., and Sigmond, M.: CCCma CanESM5-CanOE model output prepared for CMIP6 CMIP, Tech. rep., Version v20190429. Earth System Grid Federation, <https://doi.org/https://doi.org/10.22033/ESGF/CMIP6.10205>, 2019c.
- 750 Swart, N. C., Cole, J. N., Kharin, V. V., Lazare, M., Scinocca, J. F., Gillett, N. P., Anstey, J., Arora, V., Christian, J. R., Jiao, Y., Lee, W. G., Majaess, F., Saenko, O. A., Seiler, C., Seinen, C., Shao, A., Solheim, L., von Salzen, K., Yang, D., Winter, B., and Sigmond, M.: CCCma CanESM5-CanOE model output prepared for CMIP6 ScenarioMIP, Tech. rep., Version v20190429. Earth System Grid Federation, <https://doi.org/https://doi.org/10.22033/ESGF/CMIP6.10207>, 2019d.
- 755 Swart, N. C., Cole, J. N. S., Kharin, V. V., Lazare, M., Scinocca, J. F., Gillett, N. P., Anstey, J., Arora, V., Christian, J. R., Hanna, S., Jiao, Y., Lee, W. G., Majaess, F., Saenko, O. A., Seiler, C., Seinen, C., Shao, A., Sigmond, M., Solheim, L., von Salzen, K., Yang, D., and Winter, B.: The Canadian Earth System Model version 5 (CanESM5.0.3), *Geosci. Model Dev.*, 12, 4823–4873, <https://doi.org/https://doi.org/10.5194/gmd-12-4823-2019>, 2019e.
- 760 Séférian, R., Nabat, P., Michou, M., Saint-Martin, D., Voldoire, A., Colin, J., Decharme, B., Delire, C., Berthet, S., Chevallier, M., Sénési, S., Franchisteguy, L., Vial, J., Mallet, M., Joetzjer, E., Geoffroy, O., Guérémy, J.-F., Moine, M.-P., Msadek, R., Ribes, A., Rocher, M., Roehrig, R., Salas-y Mélia, D., Sanchez, E., Terray, L., Valcke, S., Waldman, R., Aumont, O., Bopp, L., Deshayes, J., Éthé, C., and Madec,

- G.: Evaluation of CNRM Earth System Model, CNRM-ESM2-1: Role of Earth System Processes in Present-Day and Future Climate, *J. Adv. Model. Earth Syst.*, 11, 4182–4227, <https://doi.org/https://doi.org/10.1029/2019MS001791>, 2019.
- 765 Tachiiri, K., Abe, M., Hajima, T., Arakawa, O., Suzuki, T., Komuro, Y., Ogochi, K., Watanabe, M., Yamamoto, A., Tatebe, H., Noguchi, M. A., Ohgaito, R., Ito, A., Yamazaki, D., Ito, A., Takata, K., Watanabe, S., and Kawamiya, M.: MIROC MIROC-ES2L model output prepared for CMIP6 ScenarioMIP, Tech. rep., Version 20190823. Earth System Grid Federation, <https://doi.org/https://doi.org/10.22033/ESGF/CMIP6.936>, 2019.
- 770 Tang, Y., Rumbold, S., Ellis, R., Kelley, D., Mulcahy, J., Sellar, A., Walton, J., and Jones, C.: MOHC UKESM1.0-LL model output prepared for CMIP6 CMIP, Tech. rep., Version 20190627. Earth System Grid Federation, <https://doi.org/https://doi.org/10.22033/ESGF/CMIP6.1569>, 2019.
- Tank, S. E., Raymond, P. A., Striegl, R. G., McClelland, J. W., Holmes, R. M., Fiske, G. J., and Peterson, B. J.: A land-to-ocean perspective on the magnitude, source and implication of DIC flux from major Arctic rivers to the Arctic Ocean, *Global Biogeochem. Cy.*, 26, <https://doi.org/https://doi.org/10.1029/2011GB004192>, 2012.
- 775 Terhaar, J., Orr, J. C., Ethé, C., Regnier, P., and Bopp, L.: Simulated Arctic Ocean Response to Doubling of Riverine Carbon and Nutrient Delivery, *Global Biogeochem. Cy.*, 33, 1048–1070, <https://doi.org/https://doi.org/10.1029/2019GB006200>, 2019a.
- Terhaar, J., Orr, J. C., Gehlen, M., Ethé, C., and Bopp, L.: Model constraints on the anthropogenic carbon budget of the Arctic Ocean, *Biogeosciences*, 16, 2343–2367, <https://doi.org/10.5194/bg-16-2343-2019>, 2019b.
- 780 Terhaar, J., Kwiatkowski, L., and Bopp, L.: Emergent constraint on Arctic Ocean acidification in the twenty-first century, *Nature*, 582, 379–383, <https://doi.org/https://doi.org/10.1038/s41586-020-2360-3>, 2020a.
- Terhaar, J., Tanhua, T., Stöven, T., Orr, J. C., and Bopp, L.: Evaluation of Data-Based Estimates of Anthropogenic Carbon in the Arctic Ocean, *J. Geophys. Res. Oceans*, 125, e2020JC016 124, <https://doi.org/https://doi.org/10.1029/2020JC016124>, 2020b.
- Ulfso, A., Jones, E. M., Casacuberta, N., Korhonen, M., Rabe, B., Karcher, M., and van Heuven, S. M.: Rapid Changes in Anthropogenic Carbon Storage and Ocean Acidification in the Intermediate Layers of the Eurasian Arctic Ocean: 1996–2015, *Global Biogeochem. Cy.*, 32, 1254–1275, <https://doi.org/https://doi.org/10.1029/2017GB005738>, 2018.
- 785 Waldbusser, G. G. and Salisbury, J. E.: Ocean Acidification in the Coastal Zone from an Organism’s Perspective: Multiple System Parameters, Frequency Domains, and Habitats, *Annu. Rev. Mar. Sci.*, 6, 221–247, <https://doi.org/https://doi.org/10.1146/annurev-marine-121211-172238>, 2014.
- 790 Wanninkhof, R., Barbero, L., Byrne, R., Cai, W.-J., Huang, W.-J., Zhang, J.-Z., Baringer, M., and Langdon, C.: Ocean acidification along the Gulf Coast and East Coast of the USA, *Cont. Shelf Res.*, 98, 54–71, <https://doi.org/https://doi.org/10.1016/j.csr.2015.02.008>, 2015.
- Wenzel, S., Cox, P. M., Eyring, V., and Friedlingstein, P.: Emergent constraints on climate-carbon cycle feedbacks in the CMIP5 Earth system models, *J. Geophys. Res. Biogeosciences*, 119, 794–807, <https://doi.org/https://doi.org/10.1002/2013JG002591>, 2014.
- 795 Wieners, K.-H., Giorgetta, M., Jungclaus, J., Reick, C., Esch, M., Bittner, M., Gayler, V., Haak, H., de Vrese, P., Raddatz, T., Mauritsen, T., von Storch, J.-S., Behrens, J., Brovkin, V., Claussen, M., Crueger, T., Fast, I., Fiedler, S., Hagemann, S., Hohenegger, C., Jahns, T., Kloster, S., Kinne, S., Lasslop, G., Kornbluh, L., Marotzke, J., Matei, D., Meraner, K., Mikolajewicz, U., Modali, K., Müller, W., Nabel, J., Notz, D., Peters, K., Pincus, R., Pohlmann, H., Pongratz, J., Rast, S., Schmidt, H., Schnur, R., Schulzweida, U., Six, K., Stevens, B., Voigt, A., and Roeckner, E.: MPI-M MPIESM1.2-LR model output prepared for CMIP6 ScenarioMIP, Tech. rep., Version 20190710. Earth System Grid Federation, <https://doi.org/https://doi.org/10.22033/ESGF/CMIP6.793>, 2019a.
- 800 Wieners, K.-H., Giorgetta, M., Jungclaus, J., Reick, C., Esch, M., Bittner, M., Legutke, S., Schupfner, M., Wachsmann, F., Gayler, V., Haak, H., de Vrese, P., Raddatz, T., Mauritsen, T., von Storch, J.-S., Behrens, J., Brovkin, V., Claussen, M., Crueger, T., Fast, I., Fiedler,

- S., Hagemann, S., Hohenegger, C., Jahns, T., Kloster, S., Kinne, S., Lasslop, G., Kornbluh, L., Marotzke, J., Matei, D., Meraner, K., Mikolajewicz, U., Modali, K., Müller, W., Nabel, J., Notz, D., Peters, K., Pincus, R., Pohlmann, H., Pongratz, J., Rast, S., Schmidt, H., Schnur, R., Schulzweida, U., Six, K., Stevens, B., Voigt, A., and Roeckner, E.: MPI-M MPIESM1.2-LR model output prepared for CMIP6
805 CMIP, Tech. rep., Version 20190710. Earth System Grid Federation, <https://doi.org/https://doi.org/10.22033/ESGF/CMIP6.742>, 2019b.
- Woosley, R. J. and Millero, F. J.: Freshening of the western Arctic negates anthropogenic carbon uptake potential, *Limn. Oceanog.*, n/a, <https://doi.org/https://doi.org/10.1002/lno.11421>, 2020.
- Xue, L. and Cai, W.-J.: Total alkalinity minus dissolved inorganic carbon as a proxy for deciphering ocean acidification mechanisms, *Mar. Chem.*, 222, 103–791, <https://doi.org/https://doi.org/10.1016/j.marchem.2020.103791>, 2020.
- 810 Yamamoto-Kawai, M., McLaughlin, F. A., and Carmack, E. C.: Effects of ocean acidification, warming and melting of sea ice on aragonite saturation of the Canada Basin surface water, *Geophys. Res. Lett.*, 38, <https://doi.org/https://doi.org/10.1029/2010GL045501>, 2011.
- Yukimoto, S., Kawai, H., Koshiro, T., Oshima, N., Yoshida, K., Urakawa, S., Tsujino, H., Deushi, M., Tanaka, T., Hosaka, M., Yabu, S., Yoshimura, H., Shindo, E., Mizuta, R., Obata, A., Adachi, Y., and Ishii, M.: The Meteorological Research Institute Earth System Model Version 2.0, MRI-ESM2.0: Description and Basic Evaluation of the Physical Component, *J. Meteorol. Soc. Jpn. Ser. II*, 97, 931–965,
815 <https://doi.org/https://doi.org/10.2151/jmsj.2019-051>, 2019a.
- Yukimoto, S., Koshiro, T., Kawai, H., Oshima, N., Yoshida, K., Urakawa, S., Tsujino, H., Deushi, M., Tanaka, T., Hosaka, M., Yoshimura, H., Shindo, E., Mizuta, R., Ishii, M., Obata, A., and Adachi, Y.: MRI MRI-ESM2.0 model output prepared for CMIP6 CMIP, Tech. rep., Version 20191205. Earth System Grid Federation, <https://doi.org/https://doi.org/10.22033/ESGF/CMIP6.621>, 2019b.
- Yukimoto, S., Koshiro, T., Kawai, H., Oshima, N., Yoshida, K., Urakawa, S., Tsujino, H., Deushi, M., Tanaka, T., Hosaka, M., Yoshimura,
820 H., Shindo, E., Mizuta, R., Ishii, M., Obata, A., and Adachi, Y.: MRI MRI-ESM2.0 model output prepared for CMIP6 ScenarioMIP, Tech. rep., Version 20200120. Earth System Grid Federation, <https://doi.org/https://doi.org/10.22033/ESGF/CMIP6.638>, 2019c.
- Ziehn, T., Chamberlain, M., Lenton, A., Law, R., Bodman, R., Dix, M., Wang, Y., Dobrohotoff, P., Srbinovsky, J., Stevens, L., Vohralik, P., Mackallah, C., Sullivan, A., O'Farrell, S., and Druken, K.: CSIRO ACCESS-ESM1.5 model output prepared for CMIP6 CMIP, Tech. rep., Version 20191112. Earth System Grid Federation, <https://doi.org/https://doi.org/10.22033/ESGF/CMIP6.2288>, 2019a.
- 825 Ziehn, T., Chamberlain, M., Lenton, A., Law, R., Bodman, R., Dix, M., Wang, Y., Dobrohotoff, P., Srbinovsky, J., Stevens, L., Vohralik, P., Mackallah, C., Sullivan, A., O'Farrell, S., and Druken, K.: CSIRO ACCESS-ESM1.5 model output prepared for CMIP6 ScenarioMIP, Tech. rep., Version 20191115. Earth System Grid Federation, <https://doi.org/https://doi.org/10.22033/ESGF/CMIP6.2291>, 2019b.
- Ziehn, T., Chamberlain, M., Law, R., Lenton, A., Bodman, R., Dix, M., Stevens, L., Wang, Y., and Srbinovsky, J.: The Australian Earth System Model: ACCESS-ESM1. 5., *Journal of Southern Hemisphere Earth Systems Science (JSHESS)*,
830 <https://doi.org/https://doi.org/10.1071/ES19035>, 2020.
- Zweng, M. M., Reagan, J. R., Seidov, D., Boyer, T. P., Locarnini, R. A., Garcia, H. E., Mishonov, A. V., Baranova, O. K., Weathers, K., Paver, C. R., and Smolyar, I.: World Ocean Atlas 2018, Volume 2: Salinity, Tech. rep., A. Mishonov Technical Ed.; NOAA Atlas NESDIS 81, 2018.
- Årthun, M., Ingvaldsen, R., Smedsrud, L., and Schrum, C.: Dense water formation and circulation in the Barents Sea, *Deep-Sea Res. Pt. I*,
835 58, 801–817, <https://doi.org/https://doi.org/10.1016/j.dsr.2011.06.001>, 2011.

Table 1. The CMIP6 ESMs used in this study with their ocean-sea ice and marine-biogeochemical (MBG) model components.

Model	Ocean-sea ice	MBG	Riverine BGC fluxes	Data DOI
ACCESS-ESM1.5 (Ziehn et al., 2020)	MOM5, CICE4	WOMBAT	none	Ziehn et al. (2019a, b)
CanESM5 (Swart et al., 2019e)	NEMO 3.4.1-LIM2	CMOC	none	Swart et al. (2019a, b)
CanESM5-CanOE (Swart et al., 2019e; Christian et al., 2020)	NEMO 3.4.1-LIM2	CanOE	none	Swart et al. (2019c, d)
CESM2 (Danabasoglu et al., 2020)	POP2-CICE5	MARBL-BEC	C, N, P, Fe, Si, Al	Danabasoglu (2019a, b)
CESM2-WACCM (Danabasoglu et al., 2020)	POP2-CICE5	MARBL-BEC	C, N, P, Fe, Si, Al	Danabasoglu (2019c, d)
CNRM-ESM2-1 (S��ferian et al., 2019)	NEMOv3.6-GELATOv6	PISCESv2-gas	C, P, Ar	Seferian (2018a, b)
GFDL-CM4 ^a (Held et al., 2019; Dunne et al., 2020a)	MOM6, SIS2	BLINGv2	C, N, P, Fe, Ar	Guo et al. (2018a, b)
GFDL-ESM4 (Dunne et al., 2020b; Stock et al., 2020)	MOM6, SIS2	COBALTv2	C, P, N, Ar	Krasting et al. (2018); John et al. (2018)
IPSL-CM6A-LR (Boucher et al., 2020)	NEMOv3.6-LIM3	PISCESv2	C, N, P, Fe, Si, Ar	Boucher et al. (2018a, b)
MIROC-ES2L (Hajima et al., 2020)	COCO	OECO2	N, P	Hajima et al. (2019); Tachiiri et al. (2019)
MPI-ESM1.2-HR (M��ller et al., 2018; Mauritsen et al., 2019)	MPIOM	HAMOCC6	none	Schupfner et al. (2019); Jungclauss et al. (2019)
MPI-ESM1.2-LR (Mauritsen et al., 2019)	MPIOM	HAMOCC6	none	Wieners et al. (2019b, a)
MRI-ESM2 ^b (Yukimoto et al., 2019a)	MRICOM4	NPZD	none	Yukimoto et al. (2019b, c)
UKESM1-0-LL (Sellar et al., 2019)	NEMO v3.6, CICE	MEDUSA-2	none	Tang et al. (2019); Good et al. (2019)

^a Only SSP2-4.5 and SSP5-8.5^b Only SSP5-8.5