



Assessing MODIS Vegetation Continuous Fields tree cover product (collection 6): performance and applicability in tropical forests and savannas.

- 5 Rahayu Adzhar^{1,2*}, Douglas I Kelley^{1*}, Ning Dong³, Mireia Torello Raventos⁴, Elmar Veenendaal⁵, Ted R Feldpausch⁶, Oliver L Philips⁷, Simon Lewis⁷, Bonaventure Sonké⁸, Herman Taedoum⁹, Beatriz Schwantes Marimon¹⁰, Tomas Domingues¹¹, Luzmila Arroyo¹², Gloria Djagbletey¹³, Gustavo Saiz¹⁴, France Gerard¹
- 10 ¹ U.K. Centre for Ecology & Hydrology, Wallingford, Oxfordshire, U.K.
² Department of Life Sciences, Imperial College London, Berkshire, U.K.
³ Department of Biological Sciences, Macquarie University, North Ryde, Australia
⁴ School of Earth and Environmental Science, James Cook University, Cairns, Australia
⁵ Plant Ecology and Nature Conservation, Wageningen University, Wageningen, The Netherlands
15 ⁶ College of Life and Environmental Sciences, University of Exeter, Exeter, U.K.
⁷ School of Geography, University of Leeds, U.K.
⁸ Plant Systematics and Ecology Laboratory, Department of Biology, Higher Teachers' Training College, University of Yaoundé, Yaoundé, Cameroon
⁹ Consultative Group on International Agricultural Research | CGIAR · Bioversity International, Yaoundé,
20 Cameroon
¹⁰ State University of Mato Grosso, Mato Grosso, Brazil
¹¹ Departamento de Biología (Ribeirão Preto), University of São Paulo, São Paulo, Brazil
¹² Universidad Autónoma Gabriel René Moreno, Santa Cruz, Bolivia
¹³ Forest and Climate Change Division, Forestry Research Institute of Ghana, Ghana
25 ¹⁴ Facultad de Ciencias, Universidad Católica de la Santísima Concepción, Concepción, Chile

Correspondence to: rahayu.adzhar@gmail.com; doukel@ceh.ac.uk

Abstract. The Moderate Resolution Imaging Spectroradiometer vegetation continuous fields (MODIS VCF)
30 Earth observation product is widely used to estimate forest cover changes, parameterise vegetation and Earth System models, and as a reference for validation or calibration where field data is limited. However, although limited independent validations of MODIS VCF have shown that MODIS VCF's accuracy decreases when estimating tree cover in sparsely-vegetated areas such as tropical savannas, no study has yet assessed the impact this may have on the VCF-based tree cover distributions used by many in their research. Using tropical forest
35 and savanna inventory data collected by the TROPical Biomes in Transition (TROBIT) project, we produce a series of corrections that take into account (i) the spatial disparity between the in-situ plot size and the MODIS VCF pixel, and (ii) the trees' spatial distribution within in-situ plots. We then applied our corrections to areas identified as forest or savanna in the International Geosphere-Biosphere Programme (IGBP) land cover mapping product. All IGBP classes identified as 'savanna' show substantial increases in cover after correction, indicating
40 that the most recent version of MODIS VCF consistently underestimates woody cover in tropical savannas. We estimate that MODIS VCF could be underestimating tropical tree cover by between 9 to 15 %. Models that use MODIS VCF as their benchmark could therefore be underestimating the carbon uptake in forest-savanna areas and misrepresenting forest-savanna dynamics. While more detailed in-situ field data is necessary to produce more accurate and reliable corrections, we recommend caution when using MODIS VCF in tropical savannas.

45

1 Introduction

Tree cover values derived from Earth observation (E.O.) data form a fundamental part of ecological research. They are used to estimate forest cover change, biomass, and carbon stocks (Bastin et al., 2019; Giriraj et al.,
50 2017; Saatchi et al., 2011; Song et al., 2014); help identify key areas for conservation efforts (Miles et al.,



2006); and are used as a basis for climatic and vegetation modelling and model evaluation (Brovkin et al., 2013; Burton et al., 2019; Kelley et al., 2013). All this research, in turn, plays a vital role in informing local, regional, and global environmental policies (Harris et al., 2012). As such, an E.O. product's accuracy is important to consider, as any errors in the initial tree cover estimate can be further compounded in downstream work.

55

Only a handful of E.O. products provide global maps of percentage tree cover or forest and shrub cover distributions (Bartholomé and Belward, 2005; Bicheron et al., 2008), and fewer still provide information stretching over at least a decade (Friedl et al., 2002; Hansen et al., 2003). Of these, one of the products most widely used in ecological modelling is the Moderate Resolution Imaging Spectroradiometer Vegetation Continuous Fields (MODIS VCF) product (DiMiceli, 2017). MODIS VCF is a yearly product that provides percent tree cover globally at a spatial resolution of 250 m and is available for the years 2000 through to 2020. Its quantitative measure of woody cover is recorded annually and is described as a percentage of ground cover, making it particularly suited for use in evaluating dynamic global models (Lasslop et al., 2018; Rabin et al., 2017), as a proxy for in-situ data that is harder to collect (Kelley et al., 2019), and to help define parameters for calculating global tree restoration potential (Bastin et al., 2019). Collection 6 is the most recent iteration of the product.

As the VCF product has progressed from Collection 1 to its current Collection 6, several validations using in-situ field data or higher-resolution remotely sensed data as a reference measurement have been carried out. These have been few and limited to sites within a biome (Montesano et al., 2009a), a region (Hansen et al., 2005; White et al., 2005), or within a country (Gao et al., 2014; Sexton et al., 2013). The MODIS VCF product evaluated was the most recent collection available at the time (i.e. Hansen et al., 2005 and White et al., 2005 for Collection 3; Montesano et al., 2009a for Collection 4; and Gao et al., 2015 and Sexton et al., 2013 for Collection 5). To our knowledge, no such independent validation experiment has yet been conducted on Collection 6, which produces tree cover estimates in the same manner as Collection 5 but with improvements made to the upstream inputs to enhance its accuracy (DiMiceli, 2017).

The validations found that MODIS VCF may be less suitable for estimating tree cover in sparsely-vegetated areas. Huang & Siegert (2006) noted that MODIS VCF classified large areas of land as 'bare' where their land cover classification system identified it as sparsely-vegetated. Montesano et al. (2009) found that MODIS VCF data (Collection 4) overestimated cover in areas of low tree cover in taiga-tundra transition zones. Sexton et al. (2013) found that the Collection 5 product overestimated cover in areas of low cover (below 20 %) and underestimated in areas of higher tree cover, while Gao et al. (2015) found that MODIS VCF can only partially discriminate between tropical forest and non-forest, struggling in areas that have greater heterogeneity. What is clear from the history of these validation and comparison experiments is that MODIS VCF has accuracy issues in areas with low woody vegetation cover, which has implications when its tree cover estimates are treated as accurately representative of real-world conditions. Failure to accurately account for the product's difficulty in estimating low woody covers can, therefore, lead to miscalibrated models and estimations that do not reflect real-world conditions. This, in turn, has knock-on effects on environmental policy-making, conservation efforts, and future ecological research, especially in areas with vegetation cover types that are most prone to error.

Tropical savannas have woody covers that fall within the range particularly affected by the reported MODIS VCF errors. A large proportion of these savannas can be found in tropical developing countries (Boval and Dixon, 2012), and are predicted to be home to half of the world's population by 2050 (State of the Tropics, 2020). Tropical savannas are therefore highly vulnerable to anthropogenic change. In the face of a growing population, land fragmentation, and changing climate, a savanna's ability to maintain robust ecosystem functions is directly linked to the amount of woody cover present (Sankaran et al., 2006). As a result, the ability to accurately monitor the state, dynamics, and woody cover trends of tropical savannas is a vital part of understanding how and why savannas are changing in the tropics (Harris et al., 2012; Miles et al., 2006), while also improving modelled climate projections and vegetation dynamics for this complex biome.

100



105 In this study, we validate the accuracy of MODIS VCF Collection 6 in tropical savannas and forests by comparing the tree cover percentage of the product to corresponding field data. We then characterise the observed bias in woody covers across both savanna and forest ecosystems and apply our corrections across the tropics to highlight the regions most likely to be affected by these inaccuracies in the MODIS VCF product.

2 Methods

110 We used the MODIS VCF Collection 6 product (spatial resolution of 250 m, DiMiceli, 2017) with tree cover values averaged across the years 2006 through to 2009 to reflect the range of the field data collection period. The in-situ field data was sourced from the ‘TROpical Biomes InTransition’ project (TROBIT) (www.geog.leeds.ac.uk/TROBIT, Torello-Raventos et al., 2013) which include the corner locations and the Canopy Area Index (CAI) values for 24 forest and 24 savanna sites distributed across Australia, Brazil, Bolivia, Cameroon, and Ghana (Fig. A1 and Table A1, Fig. 2 in Torello-Raventos et al., 2013). The TROBIT field
115 campaigns were carried out over a 3-year period, from 2006 to 2009, and the field sites used in this study are 100 m x 100 m in size (except 3 sites from Ghana that are 50 m x 100 m). All the sites fall within the tropics, that is, within 23.5 degrees north and south of the equator.

120 The classification of the TROBIT sites as either ‘forest’ or ‘savanna’ is based on the parameters described in Torello-Raventos et al. (2013) and Veenendaal et al. (2015). A ‘savanna’ is a natural land cover that is not a forest, bare ground, or a body of water, with ‘forest’ being defined as woody vegetation with a mean height of or exceeding 6 m for trees with a diameter at breast height (dbh) exceeding 10 cm, and a tree cover of or exceeding ~60 %. There is some ambiguity in how ‘savannas’ and ‘grasslands’ are defined, with some modelling-based
125 research treating the two biomes as different (Whitley et al., 2017) while studies based on plant functional traits group them together (Solofondranohatra et al., 2018; White et al., 2000). As there is some concern that MODIS VCF will struggle to pick up woody cover in areas with really sparse vegetation, in this paper we have decided to treat ‘grasslands’ as part of the savanna domain.

130 CAI is defined as the sum of the projected areas of individual tree canopies divided by the ground area. In the TROBIT project, the upper-stratum, mid-stratum, and subordinate-stratum crown areas are factored into the CAI values and were recorded using methods outlined in Torello-Raventos et al. (2013). Membership to the strata is determined by the tree’s dbh (upper-stratum: $dbh > 10$ cm, mid-stratum: $2.5 \text{ cm} < dbh < 10$ cm, and subordinate-stratum: $dbh < 2.5$ cm). However, CAI values do not account for within-site tree canopy distribution patterns and the overlap between individual tree canopies. We account for this by converting each CAI value into a
135 probability distribution function incorporating the following two extreme scenarios: “enforced overlap”, where the location probability of individual canopies increases linearly from 0 to 1 across a site; and “unenforced overlap”, where individual canopies follow a uniform random distribution pattern and canopy overlap is not purposefully introduced (Fig. A2). We repeated this 1000 times per CAI measurement to determine the probability distribution of expected CAI for each site.

140 Unlike CAI, which is the fraction of ground covered by tree crowns, the percent tree cover value from MODIS VCF is defined as “the portion of the skylight orthogonal to the surface which is intercepted by trees” (Hansen et al. 2002). To make the MODIS VCF tree cover value comparable to the tree cover derived from the CAI of a TROBIT site, we divided the MODIS VCF values by 0.8 as suggested by Hansen et al. (2002). Next, to account
145 for the difference in size between the MODIS VCF pixel (250 m x 250 m) and the smaller field plot size (100 m x 100 m), we calculated the possible percent tree cover an area the size of a TROBIT field site could have, given the MODIS VCF percent tree cover for a MODIS-sized pixel. This was done for two extreme scenarios: “enforced clumping,” where all the tree cover for the given MODIS VCF value is forcibly ‘clumped’ on one side of the pixel, or “unenforced clumping,” where ‘clumping’ is not enforced and tree cover is distributed
150 randomly within the pixel (Fig. A3). Basically, the clumping scenarios introduce possible variations in percent cover due to the area and location mismatch between a TROBIT field site and a MODIS pixel. A probability distribution was generated for each MODIS VCF pixel by calculating percent tree cover values for 1000 samples (100 m x 100 m) randomly placed within the 250 m x 250 m MODIS VCF pixel.



155 We thereby compared MODIS VCF and TROBIT under four different scenarios: 1) unenforced overlap and
clumping; 2) enforce overlap and unenforced clumping; 3) unenforced overlap and enforced clumping; 4)
enforced overlap and clumping. Comparisons were conducted by fitting the following function:

$$\text{logit}(VCF) = C_0 + \Delta \times \log(C^{\tau_1}/(1 - C^{\tau_2})) \quad (\text{Equation 1})$$

160 Where $C_0, \Delta, \tau_1, \tau_2$ are optimised parameters and VCF and C are the MODIS VCF pixel and TROBIT site
probability distributions, respectively. This is similar to a standard linear regression of logit transformed data,
accounting for maximum and minimum bounds of 0-100 % tree cover, with τ_1, τ_2 allowing for a non-symmetric
transformation of tree cover. To account for the probability density of each point, we inferred the parameters in
165 Equation 1 using a Total Least Squares Bayesian Inference technique using a Metropolis-Hastings Markov
Chain Monte Carlo step. Priors were uninformed but physically bounded (i.e. $\Delta, \tau_1, \tau_2 > 0$) to assume an
increasing relationship between MODIS VCF and C . Our logit transformed data in Equation 1, along with τ_1, τ_2
correcting for symmetry, allowed us to assume normally distributed model errors, and so our conditional
probability of observations for a given parameter combination was described by a normal distribution (Gelman
170 et al., 2013). Each combination was run over 10 chains, with 1000 warm-up iterations and 10,000 sampling
iterations. Optimisation was performed using the rstan2.19.2 (Stan Development Team, 2019) package in R3.5.2
(R Core Team, 2018).

We evaluated the impact of the MODIS VCF biases inferred from this correction across the tropics by inverting
175 our calculation of MODIS VCF bias (Fig. A4) as follows: first, the inverse (i.e. solving for C) of equation 1 was
applied to MODIS VCF values after conversion to a 100 m x 100 m pixel size grid (matching the field site area);
then this corrected value was translated back to the original 250 m x 250 m VCF pixel size. As the inverse of
Equation 1 has no analytical solution, we found the rounded percent value of C that minimises the absolute
180 difference between the left- and right-hand side of the equation. For computational feasibility, we constructed
maps of the tropics with corrected MODIS VCF values (Fig. 2) by sampling 5 iterations that were randomly
sampled from each of our 10 optimisation chains (50 in total) and masking out pixels with cover types not
considered as ‘forest’ or ‘savanna’.

We used the 500 m MODIS Land Cover Type (MCD12Q1) product to identify the areas of ‘forest’ and
185 ‘savanna’ across the tropics in the MODIS VCF product. MCD12Q1 is widely used by the global land surface
modelling community (e.g. Sellar et al., 2019; Wiltshire et al., 2020) and describes land cover in terms of 17
global land cover classes as per the International Geosphere-Biosphere Programme (IGBP, Table 3 in Sulla-
Menashe and Friedl, 2018). The product is based on the same spectroradiometer (MODIS) and temporal
resolution as the VCF product. Referring to the definition of ‘savanna’ of Veenendaal et al. (2015), the
190 following land cover classes were chosen to represent ‘savanna’: Closed Shrubland, Open Shrubland, Woody
Savanna, Savanna, and Grassland, while ‘forest’ encompasses: Evergreen Needleleaf Forests, Evergreen
Broadleaf Forests, Deciduous Needleleaf Forests, Deciduous Broadleaf Forests, and Mixed Forests.


For a more detailed land-cover-specific evaluation, we resampled the corrected 250 m MODIS VCF pixels to a
195 500 m grid and combined it with the MCD12Q1 product to construct land-cover-specific MODIS VCF tree
cover frequency distributions (Fig. A5). Our tree cover correction by cover type (Fig. 3) for the four
clumping/overlap regression combinations was then calculated by multiplying each cover type MODIS VCF
frequency distribution (Fig. A5) with curves representing the median, 5 %, and 95 % confidence lines of the
correction equation ensembles.

200



3 Results

3.1 Comparing MODIS VCF to tree cover from TROBIT field sites



205

Figure 1. MODIS VCF percent tree cover versus percent tree cover from TROBIT field data, taking into account uncertainties associated with tree cover spatial distributions within a MODIS pixel and/or field site. The 4 combinations are: (1) no overlap and no clumping, where tree canopies are randomly distributed within both pixel and site; (2) no overlap and maximum clumping, where tree canopies are clustered in one area of the pixel, and randomly distributed throughout the field site; (3) with overlap and no clumping, where tree canopies are randomly distributed within the pixel, but overlap substantially within the field site; and (4) with overlap and maximum clumping, where tree canopies are clustered to one side within a pixel, and overlap substantially within the site. The bolded dashed line in black shows the 1:1 relationship. The solid lines represent the median of the respective regressions (green for forest; orange for savanna; black for forest and savanna combined), and the thin lines represent the 5 and 95 % confidence interval of their respective regression lines. The vertical error bars represent uncertainty introduced by clumping; the horizontal error bars represent the uncertainty introduced by overlap.

210

215

MODIS VCF underestimates tree cover within the 23 % to 72 % range across all four combinations of enforced-unenforced overlap and clumping (black line, Fig. 1). Below 12 %, MODIS VCF tree cover values do not significantly disagree with TROBIT field data, and may instead be overestimating tree cover (50 % confidence, dashed line, Fig. 1). A similar pattern is seen when tree cover exceeds ~80 %: MODIS VCF does not differ significantly from TROBIT when there is no enforced overlap (i.e. when tree canopies are spaced randomly within the site - Fig. A2 left), but may overestimate tree cover when overlap is enforced (i.e. trees are clustering towards one side increasing the degree of canopy overlap - Fig. A2 right).

220

There is a clear difference in how accurately MODIS VCF estimates tree cover in forested areas (in green, Figure 1) as opposed to areas identified as savannas (in orange, Fig. 1). In savanna sites, MODIS VCF significantly underestimates tree cover regardless of the amount of overlap and clumping. Significant underestimation (at 95 % confidence) occurs when *in-situ* tree cover exceeds 18 % (without enforced clumping) or 9-10 % (with enforced clumping). In forest sites, MODIS VCF generally estimates tree cover well. Divergence does occur at high covers, depending on the enforcement of overlap or clumping. MODIS VCF


225



230 underestimates tree cover where tree cover exceeds 79 % (at the 95 % confidence interval) when neither overlap
nor clumping is enforced, and overestimates where tree cover exceeds 58 % (at 5 % confidence interval) when
both overlap and clumping are enforced.

3.2 Global estimates of change in tropical tree cover

235 We assessed the impact of VCFs underestimation of intermediate tree covers across the tropics (IGBP forest and
savanna land cover classes), using a correction based on the combined forest and savanna (black curve, Fig. 1)
instead of using the savanna-only sites for a savanna-specific correction (orange curve, Fig. 1). This is
because there were few TROBIT sites representing savanna with MODIS VCF tree cover values exceeding
40 %, and global land cover maps disagree on the distribution of savannas within the forest-savanna ecotone
(Herold et al., 2008).




240 Figure 2: Distribution of tree cover across the tropics according to original MODIS VCF values (top left), the change in tree
cover post-correction for all four scenarios (bottom four maps), and the change in tree cover that was statistically significant
(95 % interval) in the same direction (positive or negative correction) across all four scenarios (top right). Black dots on the
scenario maps indicate areas where the post-correction values have a 95 % certainty of being positive or negative
245 corrections.

The distribution of tree cover change post-correction is similar across the four scenarios (Fig. 2), and the regions
where all four scenarios agree on the direction of change (positive and negative) are substantial. However, there
are some differences caused by the uncertainty introduced by different extents of overlap and clumping. While
we see a significant increase in tree cover across all clumping-overlap combinations in many regions of tropical
savannas and grasslands (Pennington et al., 2018), such as in the forest-savanna mosaics that surround
250 Congolian rainforests, we do not see the same pattern in the Cerrado of Brazil. This is likely because the African
forest-savanna regions fall within the range of MODIS VCF values that consistently undergo a positive
correction (~30-50 %, see Fig. A4), while the Cerrado of Brazil does not.

255 We also see significant tree cover decrease in the Sahel post-correction in most or all of the scenarios, which
runs counter to the results of Brandt et al. (2020) that found that tree cover was underestimated in the region.
This disparity may be explained by our lack of field sites in more arid regions. As these corrections were based
on a limited number of sites in a limited number of regions, it is important to note that the maps shown in Figure
2 are not definitive. Instead, it should be used to identify areas where MODIS VCF estimates may be more or
less reliable.



260 **3.3 Change in tree cover within different vegetation classes in tropical ecosystems**



265

Figure 3. Percent change in tree cover after the application of the relevant correction (clockwise: no enforced clumping or overlap (black); enforced clumping and no enforced overlap (blue); no enforced clumping and enforced overlap (red); enforced clumping and overlap (pink) in the ‘forest’ supercategory and the 5 savanna classes. Palest tone indicates positive change, mid-tone indicates negative change, and the darkest tone indicates net change. Error bars denote the 5-95 % confidence interval; if the error bar extends past the x-axis, the post-correction change is not considered significant.

270

When looking at our correction in more detail, we see that MODIS VCF significantly underestimates tree cover in all the IGBP land cover classes that we considered, regardless of overlap or clumping (95 % confidence interval)(Fig. 3). The most substantial and significant underestimation is in the classes ‘woody savannas’ and ‘savannas’. The underestimation is the largest in woody savannas, except when clumping and overlap are enforced (in purple, Fig. 3). This is because the peak in the tree cover distribution for savannas aligns with



where the correction for maximum overlap and clumping is the largest (i.e. at about 20 % tree cover, see Fig. A5), while the peak in cover distribution for woody savannas (30-70 %, Fig. A5) aligns with the cover range that undergoes the greatest correction (Fig. 4, Fig. A5) in the other clumping and overlap scenarios.

275 Open shrubland only shows a small underestimation of tree cover, despite having a tree cover range that closely resembles the range where VCF most underestimates tree cover (i.e. 20-60 % cover). The discrepancy may be because IGBP class definitions may not entirely correspond with the tree cover percentage identified by MODIS VCF, or there is some issue with how apt the IGBP description of shrublands is in terms of tree cover. For example, IGBP distinguishes ‘shrublands’ from ‘savannas’ based on a woody perennial height threshold of < 2
280 m height, while MODIS VCF only observes canopies of trees exceeding 5 m in height. MODIS VCF tree cover in open shrubland is therefore likely to be lower than the IGBP tree cover (see Fig. A5), resulting in corrections that are more conservative.

We found significant increases in tree cover for ‘forests’ in every correction scenario, though net change is only significant (95 % confidence) when overlap is unenforced. This can be explained by the presence of both
285 negative and positive corrections in the higher ranges of tree cover when overlap is enforced. Similarly, the net change is insignificant across all clumping and overlap scenarios for the IGBP classes matching the lower ranges of tree cover (grassland, close shrubland and open shrubland).

4 Discussion

290 While MODIS VCF is a powerful and accessible tool to map tree cover, our field data-based corrections indicate that the latest MODIS VCF collection 6 is missing a lot of woody cover even when uncertainty introduced by site canopy overlap and clumping within the MODIS VCF pixel are accounted for. Our maps (Fig. 2) highlight that this potential underestimation of woody cover is mainly occurring in tropical savannas. Moreover, the highest underestimation in the savanna classes occurs when there is no enforced overlap (i.e. when there is a uniform random distribution of trees) which is the most likely scenario for the TROBIT savanna plots as evidenced by work done by Veenendaal et al. (2015) to test the plots for complete spatial randomness. Only minor indications of overlap were found. Woody savannas, as an example, may have their tree cover
295 underestimated by up to 27 % (95 % confidence) when neither clumping nor overlap is enforced (in black, Fig. 3). If our results are representative of the tropics, then overall, MODIS VCF may be underestimating tropical tree cover by between 5-22 % for unenforced clumping and overlap or 0-22 % for when either clumping or
300 overlap are enforced (5-95 % confidence).

An overestimation at the lower end of the cover (< 20 %) (Hansen et al., 2002; Sexton et al., 2013) and underestimation in the middle range of cover have been identified in validations of previous MODIS VCF collections (Gross et al., 2018; Yang and Crews, 2019). MODIS VCF only maps trees that are 5 m or taller (Hansen et al. 2003), while the TROBIT CAI includes all trees. This could explain the observed underestimation
305 in the lower tree cover ranges for the savanna sites. However, the discrepancies we found between the percent tree cover ranges of the IGBP class descriptions (i.e. MCD12Q1 product) and the corresponding MODIS VCF suggests that the 5 m height threshold may not always apply. Unlike in MODIS VCF, an IGBP ‘tree’ has a minimum height of 2 m, and ‘shrubs’ are shorter. Therefore, MODIS VCF recording tree cover in the open and closed shrubland classes of MCD12Q1 (Fig. A5) indicates one of three things: these shrublands contain trees tall enough to be measured by MODIS VCF (> 5 m), that MODIS VCF is able to spot shorter trees (< 5 m), or some combination of both scenarios. More importantly, for the MCD12Q1 savanna class, MODIS VCF yields a percent tree cover range that matches closely with the savanna class definition (between 10 % and 30 %), even though the MODIS VCF and IGBP tree thresholds are different. This would suggest that in areas classed as ‘savanna’, trees grow taller than 5 m, or, similar to the shrubland classes, MODIS VCF is picking up trees with
315 heights below 5 m. Because of how our field reference CAI is derived, we were not able to apply a 5 m threshold to the reference data. As a result, we cannot conclusively link the 5 m threshold to our observed underestimation. We expect that the TROBIT CAI percent tree cover value will include tree cover that is not identifiable by MODIS VCF, and that our observed underestimation is not as extensive as shown. More work needs to be done to evaluate how effective MODIS VCF is at distinguishing trees above the 5 m threshold.

320 Our results suggest that the biases found in the previous collections may have persisted in collection 6, despite reported improvement in accuracy (DiMiceli et al., 2017). This indicates that the biases introduced by binning the training data (Gerard et al. 2017) and using a CART (Classification and Regression Tree) model (Hanan et al., 2013) are inherent and still present within this version of MODIS VCF. Models calibrated using MODIS VCF (Brandt et al., 2017; Burton et al., 2019; Kelley et al., 2019, 2020) risk inheriting these biases and should
325 therefore be validated using other sources of data. We suggest that while MODIS VCF gives a good overview of



330 tree cover on a global scale, it should be re-calibrated before it is used as a reference or training data. Special care should be taken in savannas, a biome that has long been noted as being challenging for E.O. products to characterise, as solitary trees in the landscape tend to be missed by global tree cover products (Jung et al., 2006, Brandt et al., 2020). The poor performance of MODIS VCF in savannas in particular (Gaughan et al., 2013; Gross et al., 2018; Kumar et al., 2019) emphasises the importance of continuous independent validation and re-calibration of the product. The ecosystem functions of savannas can vary drastically with just a slight difference in tree cover (Gaughan et al., 2013) and even slight errors may create issues in how we interpret the state and dynamics of the biome, which in turn affects how the land is managed.

335 Work on forest restoration potential would also be impacted. Bastin et al. (2019), for example, used MODIS VCF to estimate tree cover in agricultural land. As this tree cover is likely to have been underestimated substantially, the derived available land space for replanting may be less than projected, with the restoration potential overestimated. However, our results also indicate an underestimated tree cover in woodier savannas and forests. Accounting for this, the restoration potential could actually be greater than anticipated, because the carrying capacity of a unit of land could be greater than previously thought. The MODIS VCF correction could also result in a more uniform cover distribution across regions, producing a more gradual transition between 340 low-cover savannas and high-cover forests. This could have implications for work that, for example, uses MODIS VCF to study forest-savanna dynamics and bi-stability (Lasslop et al., 2018; Wuyts et al., 2017; Xu et al., 2016).


345 To ensure the appropriate use of the product, we suggest that where field data is available, the height threshold of MODIS VCF should be evaluated, and the product itself should be calibrated for use in the target region. Alternatively, comparing MODIS VCF to other land cover maps or higher-resolution remotely sensed data is recommended (Gross et al., 2018; Lary and Lait, 2006), though without a large-scale effort to re-calibrate MODIS VCF, the question of how appropriate MODIS VCF is for use in both forests and savannas in the tropics will remain. By highlighting the extent to which MODIS VCF struggles to estimate tree cover in tropical 350 forests and savannas, we hope to inform the future use of this product to improve its usability.

5 Conclusion

355 We found that MODIS VCF significantly underestimates tree cover in tropical forests and savannas, even when within-field site and field site-pixel variation are accounted for during validation. As MODIS VCF is a product that is commonly used in a wide variety of ecological research including vegetation modelling, estimating restoration potential, and identifying forest-savanna bimodality, we stress that more independent work on validating and re-calibrating is required before its tree cover estimates can be relied upon in the tropics.



Appendix



360 **Figure A1.** Location of sampling sites in Africa, Australia, and South America from the TROBIT Project (based on Fig. 2, Torello-Raventos et al., 2013). Of the 63 field sites, only the 48 sites with available GPS coordinates were selected (<https://www.forestplots.net>).

No.	Site Name	Country	Latitude	Longitude	MODIS VCF Percent Tree Cover (%)	Canopy Area Index	Cover Type
1	ALC-01	Brazil	-2.53	-54.91	12.50	0.32	Savanna
2	ALF-01	Brazil	-9.60	-55.94	77.00	2.31	Forest
3	ALF-02	Brazil	-9.58	-55.92	76.00	2.65	Forest
4	ASU-01	Ghana	7.14	-2.45	41.33	2.54	Forest
5	BBI-01	Burkina Faso	12.73	-1.17	1.33	0.52	Savanna
6	BBI-02	Burkina Faso	12.73	-1.16	1.50	0.99	Savanna
7	BDA-01	Burkina Faso	10.94	-3.15	6.17	0.30	Savanna
8	BDA-02	Burkina Faso	10.94	-3.15	4.50	0.18	Savanna
9	BFI-01	Ghana	7.71	-1.69	15.00	1.22	Savanna
10	BFI-02	Ghana	7.71	-1.69	12.83	1.08	Savanna
11	BFI-03	Ghana	7.71	-1.70	25.83	2.54	Forest
12	CTC-01	Australia	-16.10	145.45	72.67	2.35	Forest
13	DCR-01	Australia	-17.02	145.58	21.67	1.67	Savanna
14	DCR-02	Australia	-17.03	145.60	65.67	0.71	Forest
15	EKP-01	Australia	-18.07	145.99	43.50	0.74	Forest
16	FLO-01	Brazil	-12.81	-51.85	65.67	2.40	Forest
17	FMS-01	Australia	-18.09	144.84	7.67	0.32	Forest
18	FMS-02	Australia	-18.11	144.82	44.17	1.21	Forest
19	HOM-01	Mali	15.34	-1.47	0.50	0.05	Savanna
20	HOM-02	Mali	15.33	-1.55	0.83	0.16	Savanna
21	IBG-01	Brazil	-15.95	-47.87	20.83	0.22	Savanna
22	IBG-02	Brazil	-15.95	-47.87	20.00	0.02	Savanna



23	IBG-03	Brazil	-15.93	-47.87	20.50	0.12	Savanna
24	IBG-04	Brazil	-15.94	-47.86	27.17	0.77	Savanna
25	KBL-01	Australia	-17.77	145.54	75.00	1.69	Forest
26	KBL-02	Australia	-17.85	145.53	61.17	0.81	Forest
27	KBL-03	Australia	-17.69	145.53	79.50	3.00	Forest
28	KCR-01	Australia	-17.11	145.60	78.83	2.44	Forest
29	LFB-03	Bolivia	-14.60	-60.85	28.17	0.39	Savanna
30	MDJ-01	Cameroon	6.17	12.83	42.00	3.24	Forest
31	MDJ-02	Cameroon	6.16	12.82	18.67	0.44	Savanna
32	MDJ-03	Cameroon	5.98	12.87	64.67	2.97	Forest
33	MDJ-04	Cameroon	6.00	12.87	15.00	0.37	Savanna
34	MDJ-05	Cameroon	5.98	12.87	70.33	2.85	Forest
35	MDJ-06	Cameroon	6.00	12.89	20.50	0.68	Savanna
36	MDJ-07	Cameroon	6.01	12.89	57.33	1.75	Forest
37	MDJ-08	Cameroon	6.21	12.75	15.00	0.48	Savanna
38	MLE-01	Ghana	9.30	-1.86	10.00	0.34	Savanna
39	NXV-02	Brazil	-14.70	-52.35	20.83	1.82	Forest
40	RSC-01	Australia	-20.16	146.54	28.00	1.15	Forest
41	SMT-01	Brazil	-12.82	-51.77	36.67	1.55	Savanna
42	SMT-02	Brazil	-12.82	-51.77	41.50	1.44	Forest
43	SMT-03	Brazil	-12.83	-51.77	19.33	0.53	Savanna
44	TUC-01	Bolivia	-18.52	-60.81	50.33	1.29	Forest
45	TUC-02	Bolivia	-18.53	-60.63	21.67	0.81	Savanna
46	TUC-03	Bolivia	-18.19	-60.86	10.83	0.37	Savanna
47	VCR-01	Brazil	-14.83	-52.16	69.50	2.81	Forest
48	VCR-02	Brazil	-14.83	-52.17	69.67	2.74	Forest

365 **Table A1.** Site names, locations, Canopy Area Index values, MODIS VCF percent tree cover values, and cover type of the 48 TROBIT Project sites used in this study. Details regarding the vegetation structure for each site can be found in Fig. 1 of Torello-Raventos et al., 2013.






Figure A2. Visual representation of the effects of enforcing overlap within a (100 m x 100 m) TROBIT site with a given Canopy Area Index (CAI). Left: Overlap is not enforced, and individual crowns follow a uniform random distribution. Right: Overlap is enforced by linearly increasing the probability of a canopy being located more on one side of the site (i.e. here the right side of the site) than the other. This results in tree canopies ‘overlapping’ to a greater extent, which affects how accurately CAI represents actual canopy cover.



380 **Figure A3.** Visual representation of the effects of unenforced and enforced clumping in a 250 m x 250 m MODIS VCF pixel with 50 % tree cover. Clumping all the cover to one side of the pixel (bottom) affects the average canopy cover value of a 100 m x 100 m-sized average TROBIT site, illustrated here as the black boxes.



385

Figure A4. The correction curves developed for MODIS VCF based on 4 pixel-site mismatch scenarios (no clumping and no overlap; enforced clumping no overlap; no clumping enforced overlap; and enforced clumping and enforced overlap). The dashed line signifies the 'ideal' 1:1 relationship wherein corrected MODIS VCF is unchanged from the original MODIS VCF values. The shaded regions represent 5 to 95 % confidence intervals for the respective corrected MODIS VCF values.




Figure A5. Frequency distributions of percent tree cover value as estimated by MODIS VCF across the ‘forest’ supercategory and the following IGBP classes that by our definition count as part of the ‘savanna’ domain: Closed Shrublands, Open Shrublands, Woody Savannas, Savannas, and Grasslands. Specific class definitions as per the User Guide for the MODIS Land Cover Product (Sulla-Menashe and Friedl, 2018).

390



Code/Data Availability.

395 The code and data used to support the findings of this study are archived at
https://github.com/douglask3/VCF_vs_sites revision number [fdda3ff](#)

Author Contribution.

400 R.A., D.K., F.G. designed the TROBIT, MODIS VCF pixel-site comparison technique. R.A., F.G. collated
TROBIT and corresponding MODIS VCF values. D.K. and N.D. performed regression analysis and constructed
global maps. R.A. wrote the first draft of the paper with input from D.K. and F.G. D.K. plotted the figures.
M.T.R., E.V., T.R.F., O.L.P., S.L., B.S., H.T., B.S.M., T.D., L.A., G.D., and J.K. carried out the extensive
TROBIT field campaigns. M.T.R. was the main person responsible for field data quality checking and
digitising. R.A., D.K., F.G. and N.D. contributed to the final manuscript.

Competing Interests.

405 The authors declare that they have no conflict of interest.

Acknowledgements.

The contribution by D.K. was supported by the U.K. Natural Environment Research Council through The U.K.
Earth System Modelling Project (UKESM, grant no. NE/N017951/1), F.G. was supported by the SUNRISE
project (grant no. NE/R000131/1). N.D. was supported by the Australia Research Council (DP170103410).

410 Maps in Fig. 2 and A1 were constructed using raster2.6-7 (Hijmans, 2017) and mapproj1.2-5 (Brownrigg et al.,
2017) in R 3.2.0 (R Core Team, 2015). Coastlines were obtained from mapsv3.1.0 (Becker et al., 2016).

Special thanks to Prof. Jon Lloyd for supervising R.A. at the start of the project, to Jeanette Kemp for her work
in the Australian field sites, and to Azemiyah Abdul Rahim for her support during the production of this
manuscript.

References.

- 415 Archer, C., Penny, A., Templeman, S., McKenzie, M., Hunt, E., Toral, D., Diakhite, M., Nhlapo, T., Mawoko,
D., Vergnani, L., Chamdimba, C., Diop, H., Kalanzi, B., Touitha, Y., Jackson, A., Mchugh, J., Chang, O.,
Mohamad, A., Hunter, E., and Lopez, C.: State of the Tropics 2020 Report, 2020.
- 420 Bartholomé, E. and Belward, A. S.: GLC2000: a new approach to global land cover mapping from Earth
observation data, *International Journal of Remote Sensing*, 26(9), 1959–1977,
doi:10.1080/01431160412331291297, 2005.
- 425 Bastin, J.F., Finegold, Y., Garcia, C., Mollicone, D., Rezende, M., Routh, D., Zohner, C. M. and Crowther, T.
W.: The global tree restoration potential, *Science*, 365(6448), 76–79, doi:10.1126/science.aax0848, 2019.
- Becker, R. A., Minka, T. P., Wilks, A. R., Brownrigg, R. and Deckmyn, A.: maps: Draw Geographical Maps.
[online] Available from: <http://CRAN.R-project.org/package=maps> (Accessed 1 July 2016), 2016.
- 430 Bicheron, P., Defourny, P., Brockmann, C., Schouten, L., Vancutsem, C., Huc, M., Bontemps, S., Leroy, M.,
Frédéric, A., Herold, M., Ranera, F. and Arino, O.: GLOBCOVER: products description and validation report,
2008.
- 435 Boval, M. and Dixon, R. M.: The importance of grasslands for animal production and other functions: a review
on management and methodological progress in the tropics, *Animal*, 6(5), 748–762,
doi:10.1017/S1751731112000304, 2012.
- 440 Brandt, M., Tucker, C., Kariryaa, A., Rasmussen, K., Abel, C., Small, J., Chave, J., Rasmussen, L., Hiernaux,
P., Diouf, A., Kergoat, L., Mertz, O., Igel, C., Gieseke, F., Schöning, J., Li, S., Melocik, K., Meyer, J., Sinno, S.,
and Fensholt, R.: An unexpectedly large count of trees in the West African Sahara and Sahel, *Nature*, 1–5.



10.1038/s41586-020-2824-5, 2020.

- 445 Brandt, M., Rasmussen, K., Penuelas, J., Tian, F., Schurgers, G., Verger, A., Mertz, O., Palmer, J. and Fensholt, R.: Human population growth offsets climate-driven increase in woody vegetation in sub-Saharan Africa, *Nature Ecology & Evolution*, 1, 0081, doi:10.1038/s41559-017-0081, 2017.
- 450 Brovkin, V., Boysen, L., Raddatz, T., Gayler, V., Loew, A. and Claussen, M.: Evaluation of vegetation cover and land-surface albedo in MPI-ESM CMIP5 simulations, *Journal of Advances in Modeling Earth Systems*, 5(1), 48–57, doi:10.1029/2012MS000169, 2013.
- Brownrigg, R., McIlroy, D., Minka, T. P. and Bivand, R.: mapproj: Map Projections. [online] Available from: <http://CRAN.R-project.org/package=mapproj> (Accessed 15 March, 2018), 2017.
- 455 Burton, C., Betts, R., Cardoso, M., Feldpausch, T. R., Harper, A., Jones, C. D., Kelley, D. I., Robertson, E. and Wiltshire, A.: Representation of fire, land-use change and vegetation dynamics in the Joint U.K. Land Environment Simulator vn4.9 (JULES), *Geoscientific Model Development*, 12(1), 179–193, doi:https://doi.org/10.5194/gmd-12-179-2019, 2019.
- 460 DiMiceli, M. C.: MOD44B MODIS/Terra Vegetation Continuous Fields Yearly L3 Global 250m SIN Grid V006, doi:10.5067/MODIS/MOD44B.006, 2017.
- 465 DiMiceli, C., Carroll, M., Sohlberg, R. A., Huang, C., Hansen, M. C. and Townshend, J. R. G.: Annual global automated MODIS vegetation continuous fields (MOD44B) at 250m spatial resolution for data years beginning day 65, 2000-2014, collection 5 percent tree cover, version 6, University of Maryland, 2017.
- 470 Friedl, M. A., McIver, D. K., Hodges, J. C. F., Zhang, X. Y., Muchoney, D., Strahler, A. H., Woodcock, C. E., Gopal, S., Schneider, A., Cooper, A., Baccini, A., Gao, F. and Schaaf, C.: Global land cover mapping from MODIS: algorithms and early results, *Remote Sensing of Environment*, 83(1), 287–302, doi:10.1016/S0034-4257(02)00078-0, 2002.
- 475 Gao, Y., Mas, J. F., Paneque-Gálvez, J., Skutsch, M., Ghilardi, A., Pacheco, J. A. N. and Paniagua, I.: Validation of MODIS vegetation continuous fields in two areas in Mexico, in *2014 Third International Workshop on Earth Observation and Remote Sensing Applications (EORSA)*, pp. 14–18., 2014.
- Gao, Y., Ghilardi, A., Paneque-Gálvez, J., Skutsch, M. M. and Mas, J.: Validation of MODIS Vegetation Continuous Fields for monitoring deforestation and forest degradation: two cases in Mexico, *Geocarto International*, doi:10.1080/10106049.2015.1110205, 2015.
- 480 Gaughan, A., Holdo, R. and Anderson, T.: Using short-term MODIS time-series to quantify tree cover in a highly heterogeneous African savanna, *International Journal of Remote Sensing*, 1–18, doi:10.1080/01431161.2013.810352, 2013.
- 485 Gelman, A., Carlin, J. B., Stern, H. S., Dunson, D. B., Vehtari, A. and Rubin, D. B.: *Bayesian Data Analysis*, Third Edition, CRC Press., 2013.
- 490 Gerard, F., Hooftman, D., Langevelde, F., Veenendaal, E., White, S. and Lloyd, J.: MODIS VCF should not be used to detect discontinuities in tree cover due to binning bias. A comment on Hanan et al. (2014) and Staver and Hansen (2015), *Global Ecology and Biogeography*, 26, doi:10.1111/geb.12592, 2017.
- Giriraj, A., Babar, S. and Murthy, M.: Evaluating MODIS-vegetation continuous field products to assess tree cover change and forest fragmentation in India – A multi-scale satellite remote sensing approach, *The Egyptian Journal of Remote Sensing and Space Science*, 20, doi:10.1016/j.ejrs.2017.05.004, 2017.
- 495 Gross, D., Achard, F., Dubois, G., Brink, A. and Prins, H. H. T.: Uncertainties in tree cover maps of Sub-Saharan Africa and their implications for measuring progress towards CBD Aichi Targets, *Remote Sensing in Ecology and Conservation*, 4(2), 94–112, doi:10.1002/rse2.52, 2018.
- 500 Hanan, N., Tredennick, A., Prihodko, L., Bucini, G., and Dohn, J.: Analysis of stable states in global savannas: Is the CART pulling the horse?, *Global Ecology and Biogeography*, doi: 23. 10.1111/geb.12122, 2013.



- Hansen, M. C., Townshend, J., DeFries, R., and Mark Carroll: Estimation of tree cover using MODIS data at global, continental and regional/local scales, *International Journal of Remote Sensing*, 26:19, 4359–4380, doi: 10.1080/01431160500113435, 2005.
- 505 Hansen, M.C., DeFries, R. S., Townshend, J.R.G, Carroll, M., Dimiceli, C., and Sohlberg, R.: Global Percent Tree Cover at a Spatial Resolution of 500 Meters: First Results of the MODIS Vegetation Continuous Fields Algorithm, *Earth Interactions*, 7(10), 1–15, 2003.
- 510 Hansen, M.C., Defries, R., Townshend, J., Marufu, L., and Sohlberg, R.: Development of a MODIS tree cover validation data set for Western Province, Zambia, *Remote Sensing of Environment*, 83(1), doi: 10.1016/S0034-4257(02)00080-9, 2002.
- 515 Harris, N. L., Brown, S., Hagen, S. C., Saatchi, S. S., Petrova, S., Salas, W., Hansen, M. C., Potapov, P. V. and Lotsch, A.: Baseline map of carbon emissions from deforestation in tropical regions, *Science*, 336(6088), 1573–1576, doi:10.1126/science.1217962, 2012.
- 520 Herold, M., Mayaux, P., Woodcock, C., Baccini, A. and Schmullius, C.: Some challenges in global land cover mapping: An assessment of agreement and accuracy in existing 1 km datasets, *Remote Sensing of Environment*, 112, 2538–2556, doi:10.1016/j.rse.2007.11.013, 2008.
- Hijmans, R. J.: raster: Geographic Data Analysis and Modeling. [online] Available from: <http://CRAN.R-project.org/package=raster> (Accessed 15 March 2018), 2017.
- 525 Huang, S. and Siegert, F.: Land cover classification optimised to detect areas at risk of desertification in North China based on SPOT VEGETATION imagery, *Journal of Arid Environments*, 67(2), 308–327, doi:10.1016/j.jaridenv.2006.02.016, 2006.
- 530 Jung, M., Henkel, K., Herold, M. and Churkina, G.: Exploiting synergies of global land cover products for carbon cycle modeling, *Remote Sensing of Environment*, 101(4), 534–553, doi:10.1016/j.rse.2006.01.020, 2006.
- 535 Kelley, D. I., Prentice, I. C., Harrison, S. P., Wang, H., Simard, M., Fisher, J. B. and Willis, K. O.: A comprehensive benchmarking system for evaluating global vegetation models, *Biogeosciences*, 10(5), 3313–3340, doi:<https://doi.org/10.5194/bg-10-3313-2013>, 2013.
- Kelley, D. I., Bistinas, I., Whitley, R., Burton, C., Marthews, T. R. and Dong, N.: How contemporary bioclimatic and human controls change global fire regimes, *Nature Climate Change*, 9(9), 690–696, doi:10.1038/s41558-019-0540-7, 2019.
- 540 Kelley, D. I., Burton, C., Huntingford, C., Brown, M. A. J., Whitley, R. and Dong, N.: Technical note: Low meteorological influence found in 2019 Amazonia fires, *Biogeosciences Discussions*, 1–17, doi:<https://doi.org/10.5194/bg-2020-123>, 2020.
- 545 Kumar, S. S., Hanan, N. P., Prihodko, L., Anchang, J., Ross, C. W., Ji, W. and Lind, B. M.: Alternative Vegetation States in Tropical Forests and Savannas: The Search for Consistent Signals in Diverse Remote Sensing Data, *Remote Sensing*, 11(7), 815, doi:10.3390/rs11070815, 2019.
- 550 Lary, D. and Lait, L.: Using probability distribution functions for satellite validation, *Geoscience and Remote Sensing, IEEE Transactions on*, 44, 1359–1366, doi:10.1109/TGRS.2005.860662, 2006.
- 555 Lasslop, G., Moeller, T., D’Onofrio, D., Hantson, S. and Kloster, S.: Tropical climate–vegetation–fire relationships: multivariate evaluation of the land surface model JSBACH, *Biogeosciences*, 15(19), 5969–5989, doi:<https://doi.org/10.5194/bg-15-5969-2018>, 2018.
- Miles, L., Newton, A. C., DeFries, R. S., Ravilious, C., May, I., Blyth, S., Kapos, V. and Gordon, J. E.: A global overview of the conservation status of tropical dry forests, *Journal of Biogeography*, 33(3), 491–505, doi:10.1111/j.1365-2699.2005.01424.x, 2006.
- 560 Montesano, P., Nelson, R., Sun, G., Margolis, H., Kerber, A. and Ranson, K. J.: MODIS tree cover validation



- for the circumpolar taiga–tundra transition zone, *Remote Sensing of Environment - REMOTE SENS ENVIRON*, 113, 2130–2141, doi:10.1016/j.rse.2009.05.021, 2009.
- 565 Pennington, R. T., Lehmann, C. E. R. and Rowland, L. M.: Tropical savannas and dry forests, *Current Biology*, 28(9), R541–R545, doi:10.1016/j.cub.2018.03.014, 2018.
- R Core Team: R: A language and environment for statistical computing, R Foundation for Statistical Computing, Vienna, Austria. [online] Available from: <https://www.R-project.org/>, 2018.
- 570 Rabin, S. S., Melton, J. R., Lasslop, G., Bachelet, D., Forrest, M., Hantson, S., Kaplan, J. O., Li, F., Mangeon, S., Ward, D. S., Yue, C., Arora, V. K., Hickler, T., Kloster, S., Knorr, W., Nieradzic, L., Spessa, A., Folberth, G. A., Sheehan, T., Voulgarakis, A., Kelley, D. I., Prentice, I. C., Sitch, S., Harrison, S. and Arneth, A.: The Fire Modeling Intercomparison Project (FireMIP), phase 1: experimental and analytical protocols with detailed model descriptions, *Geoscientific Model Development*, 10(3), 1175–1197, doi:https://doi.org/10.5194/gmd-10-1175-2017, 2017.
- 575 Saatchi, S., Harris, N., Brown, S., Lefsky, M., Mitchard, E., Salas, W., Zutta, B., Buermann, W., Lewis, S., Hagen, S., Petrova, S., White, L., Silman, M. and Morel, A.: Benchmark map of forest carbon stocks in tropical regions across three continents, *Proceedings of the National Academy of Sciences of the United States of America*, 108, 9899–904, doi:10.1073/pnas.1019576108, 2011.
- 580 Sankaran, M., Hanan, N., Scholes, R., Ratnam, J., Augustine, D., Cade, B., Gignoux, J., Higgins, S., Roux, X., Ludwig, F., Ardö, J., Banyikwa, F., Bronn, A., Bucini, G., Caylor, K., Coughenour, M., Diouf, A., Ekaya, W., Feral, C. and Zambatis, N.: Determinants of woody cover in African Savannas, *Nature*, 438, 846–9, doi:10.1038/nature04070, 2006.
- 585 Sellar, A. A., Jones, C. G., Mulcahy, J. P., Tang, Y., Yool, A., Wiltshire, A., O’Connor, F. M., Stringer, M., Hill, R., Palmieri, J., Woodward, S., Mora, L. de, Kuhlbrodt, T., Rumbold, S. T., Kelley, D. I., Ellis, R., Johnson, C. E., Walton, J., Abraham, N. L., Andrews, M. B., Andrews, T., Archibald, A. T., Berthou, S., Burke, E., Blockley, E., Carslaw, K., Dalvi, M., Edwards, J., Folberth, G. A., Gedney, N., Griffiths, P. T., Harper, A. B., Hendry, M. A., Hewitt, A. J., Johnson, B., Jones, A., Jones, C. D., Keeble, J., Liddicoat, S., Morgenstern, O., Parker, R. J., Predoi, V., Robertson, E., Siahann, A., Smith, R. S., Swaminathan, R., Woodhouse, M. T., Zeng, G. and Zerroukat, M.: UKESM1: Description and Evaluation of the U.K. Earth System Model, *Journal of Advances in Modeling Earth Systems*, 11(12), 4513–4558, doi:10.1029/2019MS001739, 2019.
- 595 Sexton, J. O., Song, X.-P., Feng, M., Noojipady, P., Anand, A., Huang, C., Kim, D.-H., Collins, K. M., Channan, S., DiMichele, C. and Townshend, J. R.: Global, 30-m resolution continuous fields of tree cover: Landsat-based rescaling of MODIS vegetation continuous fields with lidar-based estimates of error, *International Journal of Digital Earth*, 6(5), 427–448, doi:10.1080/17538947.2013.786146, 2013.
- 600 Solofondranohatra, C. L., Vorontsova, M. S., Hackel, J., Besnard, G., Cable, S., Williams, J., Jeannoda, V. and Lehmann, C. E. R.: Grass Functional Traits Differentiate Forest and Savanna in the Madagascar Central Highlands, *Front. Ecol. Evol.*, 6, doi:10.3389/fevo.2018.00184, 2018.
- 605 Song, X. P., Huang, C., Feng, M., Sexton, J. O., Channan, S. and Townshend, J. R.: Integrating global land cover products for improved forest cover characterisation: an application in North America, *International Journal of Digital Earth*, 7(9), 709–724, doi:10.1080/17538947.2013.856959, 2014.
- 610 Stan Development Team: RStan: The R Interface to Stan. [online] Available from: <http://mc-stan.org/>, 2019.
- Sulla-Menashe, D. and Friedl, M. A.: User Guide to Collection 6 MODIS Land Cover (MCD12Q1 and MCD12C1) Product. Available from: https://lpdaac.usgs.gov/sites/default/files/public/product_documentation/mcd12_user_guide_v6.pdf, 2018.
- 615 Torello-Raventos, M., Feldpausch, T., Veenendaal, E., Schrod, F., Saiz, G., Domingues, T., Djangbletey, G., Ford, A., Kemp, J., Marimon, B., Marimon-Junior, B. H., Lenza, E., A Ratter, J., Maracahipes, L., Sasaki, D., Sonké, B., Zapfack, L., Taedoumg, H., Daniel, V. and Lloyd, J.: On the delineation of tropical vegetation types with an emphasis on forest/savanna transitions, *Plant Ecology & Diversity*, 6(1), 101–137, doi:10.1080/17550874.2012.762812, 2013.
- 620



- 625 Veenendaal, E. M., Torello-Raventos, M., Feldpausch, T. R., Domingues, T. F., Gerard, F., Schrod, F., Saiz, G., Quesada, C. A., Djagbletey, G., Ford, A., Kemp, J., Marimon, B. S., Marimon-Junior, B. H., Lenza, E., Ratter, J. A., Maracahipes, L., Sasaki, D., Sonké, B., Zapfack, L., Villarreal, D., Schwarz, M., Yoko Ishida, F., Gilpin, M., Nardoto, G. B., Affum-Baffoe, K., Arroyo, L., Bloomfield, K., Ceca, G., Compaore, H., Davies, K., Diallo, A., Fyllas, N. M., Gignoux, J., Hien, F., Johnson, M., Mougou, E., Hiernaux, P., Killeen, T., Metcalfe, D., Miranda, H. S., Steininger, M., Sykora, K., Bird, M. I., Grace, J., Lewis, S., Phillips, O. L. and Lloyd, J.: Structural, physiognomic and above-ground biomass variation in savanna–forest transition zones on three continents – how different are co-occurring savanna and forest formations?, *Biogeosciences*, 12(10), 2927–2951, doi:10.5194/bg-12-2927-2015, 2015.
- 630 White, M., Shaw, J. and Ramsey, R.: Accuracy assessment of the vegetation continuous field tree cover product using 3954 ground plots in the south-western USA, *International Journal of Remote Sensing*, 26, 2699–2704, doi:10.1080/01431160500080626, 2005.
- 635 White, R. P., Murray, S. and Rohweder, M.: *Pilot Analysis of Global Ecosystems: Grassland Ecosystems*, World Resources Institute., 2000.
- 640 Whitley, R., Beringer, J., Hutley, L. B., Abramowitz, G., De Kauwe, M. G., Evans, B., Haverd, V., Li, L., Moore, C., Ryu, Y., Scheiter, S., Schymanski, S. J., Smith, B., Wang, Y.-P., Williams, M. and Yu, Q.: Challenges and opportunities in land surface modelling of savanna ecosystems, *Biogeosciences*, 14(20), 4711–4732, doi:https://doi.org/10.5194/bg-14-4711-2017, 2017.
- 645 Wiltshire, A. J., Burke, E. J., Chadburn, S. E., Jones, C. D., Cox, P. M., Davies-Barnard, T., Friedlingstein, P., Harper, A. B., Liddicoat, S., Sitch, S. A. and Zaehle, S.: JULES-CN: a coupled terrestrial Carbon-Nitrogen Scheme (JULES vn5.1), *Geoscientific Model Development Discussions*, 1–40, doi:https://doi.org/10.5194/gmd-2020-205, 2020.
- 650 Wuyts, B., Champneys, A. R. and House, J. I.: Amazonian forest-savanna bistability and human impact, *Nature Communications*, 8(1), 15519, doi:10.1038/ncomms15519, 2017.
- Xu, C., Hantson, S., Holmgren, M., H. van Nes, E., Staal, A. and Scheffer, M.: Remotely sensed canopy height reveals three pantropical ecosystem states, *Ecology*, 97, 2518–2521, doi:10.1002/ecy.1470, 2016.
- 655 Yang, X. and Crews, K.: Applicability analysis of MODIS tree cover product in Texas savanna, *International Journal of Applied Earth Observation and Geoinformation*, 81, 186–194, doi:10.1016/j.jag.2019.05.003, 2019.