

1 **Reviews and syntheses: Heterotrophic fixation of inorganic carbon –**
2 **significant but invisible flux in environmental carbon cycling**

3

4 Alexander Braun¹, Marina Spona-Friedl¹, Maria Avramov¹, Martin Elsner^{1,2}, Federico Baltar³,
5 Thomas Reinthaler³, Gerhard J. Herndl^{3,4} & Christian Griebler^{1,3*}

6

7 ¹ Helmholtz Zentrum München, Institute of Groundwater Ecology, Ingolstaedter Landstrasse 1, D-85764
8 Neuherberg, Germany

9 ² Technical University of Munich, Department of Analytical Chemistry and Water Chemistry, Munich, Germany

10 ³ University of Vienna, Department of Functional and Evolutionary Ecology, Althanstrasse 14, 1090 Vienna,
11 Austria

12 ⁴ Department of Marine Microbiology and Biogeochemistry, Royal Netherlands Institute for Sea Research,
13 Utrecht University, PO Box 59, 1790 AB Den Burg, The Netherlands

14 * Author for correspondence: christian.griebler@univie.ac.at

15

16 **Abstract**

17 Heterotrophic CO₂ fixation is a significant, yet underappreciated CO₂ flux in environmental
18 carbon cycling. In contrast to photosynthesis and chemolithoautotrophy – the main
19 recognized autotrophic CO₂ fixation pathways - the importance of heterotrophic CO₂
20 fixation remains enigmatic. All heterotrophs – from microorganisms to humans – take up
21 CO₂ and incorporate it into their biomass. Depending on the availability and quality of
22 growth substrates, and drivers such as the CO₂ partial pressure, heterotrophic CO₂ fixation
23 contributes at least 1-5% and in the case of methanotrophs up to 50% of the carbon
24 biomass. Assuming a standing stock of global heterotrophic biomass of 47-85 Pg C, we
25 roughly estimate that up to 5 Pg C might be derived from heterotrophic CO₂ fixation and up
26 to 12 Pg C yr⁻¹ originating from heterotrophic CO₂ fixation are funneled into the global
27 annual heterotrophic production of 34-245 Pg C yr⁻¹. These first estimates on the
28 importance of heterotrophic fixation of inorganic carbon indicate that this pathway should
29 be incorporated in present and future carbon cycling budgets.

30

31 **Key words:** CO₂ fixation, heterotrophs, anaplerosis, carbon cycling

32

33 **1. Introduction**

34 Fixation of CO₂ is a fundamental biosynthetic process in nature (Beer et al. 2010, Berg et al.
35 2007) providing the main source of metabolic energy on Earth (Giovannoni and Stingl 2005).
36 At the same time, it acts as a sink for atmospheric CO₂, the most important greenhouse gas,
37 which is responsible for more than 60% of the 'enhanced greenhouse effect' resulting in
38 global warming (Beer et al. 2010, Berg 2011, Houghton 2007, Le Quéré et al. 2016).

39 While photosynthesis and chemosynthesis are the most important processes of carbon
40 fixation, non-autotrophic carbon fixation, i.e., the carbon fixation mediated by
41 heterotrophic organisms might also be relevant albeit uncommonly quantified. While
42 heterotrophs are, per definition, organisms that respire organic compounds to gain energy
43 and build up biomass, CO₂ fixation plays also an essential role in heterotrophic carbon
44 metabolism. The diversity of carboxylating enzymes in nature reaches far beyond
45 autotrophy and virtually all heterotrophs harbor numerous enzymes fixing dissolved
46 inorganic carbon. Even though the first carboxylase in heterotrophs was discovered already
47 more than 80 years ago (Wood and Werkman 1936), the role of heterotrophs in carbon
48 cycling has so far largely focused on the oxidation of organic substrates using oxygen or
49 alternative electron acceptors (e.g. nitrate, ferric iron, sulfate) and the production of CO₂.
50 Similar to the CO₂ fixation by autotrophs, "heterotrophic CO₂ fixation" might, however,
51 constitute a significant carbon flux in specific habitats. The relevance of this process has
52 hardly been quantified due to the lack of reliable estimates of heterotrophic CO₂ fixation for
53 most organisms and habitats, and the presumption that CO₂ fixation in natural
54 environments is restricted to autotrophic organisms.


55 To fill this gap, we review the current knowledge on (i) the role of heterotrophic CO₂ fixation
56 for cellular metabolism, (ii) respiration and non-autotrophic CO₂ fixation, (iii) CO₂ fixation in
57 habitats dominated by heterotrophs, and provide (iv) quantitative estimates of
58 heterotrophic CO₂ fixation in different environments.

59

60 **2. Role of heterotrophic CO₂ fixation for cellular metabolism**

61 The non-autotrophic uptake of inorganic carbon has been reported for a wide range of
62 organisms from prokaryotes and fungi to vertebrates (Woods & Werkman 1938, Kleiber et
63 al. 1952, Cochrane 1958, Hartman et al. 1972, Perez & Matin 1982, Schinner et al. 1982,
64 Parkinson et al. 1990, Roslev et al. 2004, Hesselsoe et al. 2005, Feisthauer et al. 2008,
65 Spona-Friedl et al. 2020) and plants (Melzer and O'leary 1987). Currently, more than twenty
66 carboxylases are known forming an integral part of the central and peripheral metabolic
67 pathways of heterotrophic metabolism (Fig. 1), e.g., in gluconeogenesis, the synthesis of
68 fatty acids, amino acids, vitamins and nucleotides, the assimilation of leucine, and in
69 anaplerosis (Evans and Slotin 1940, Krebs 1941, Wood and Werkman 1941, Werkman and

70 Wood 1942, Kornberg and Krebs 1957, Wood and Stjernholm 1962, Kornberg 1965, Scrutton
 71 1971, Hartman et al. 1973, Dijkhuizen and Harder 1985, Parkinson et al. 1991, Attwood
 72 1995, Han et al 2000, Sauer and Eikmanns 2005, Erb et al. 2009, Schink 2009, Erb 2011, Bar-
 73 Even et al. 2012). Carboxylation in heterotrophs not just compensates for the dependence
 74 on organic matter, rather CO₂ fulfills the role of a “co-substrate” providing an effective and
 75 simple way to extend an existing organic carbon substrate by a single C1 unit as part of the
 76 secondary production (Erb 2011).



77


78 **Figure 1:** Selected heterotrophic CO₂ fixation reactions and pathways. PEP: phosphoenolpyruvate, DAPA: 7,8-
 79 diaminononanoate, AIR: 1-(5'-phosphoribosyl)-5-aminoimidazole, CAIR: 1-(5-phospho-D-ribose)-5-amino-4-
 80 imidazolecarboxylate, CoA: Coenzyme-A.

81 The most important CO₂ fixation pathway in all organisms is anaplerosis. Anaplerosis
 82 replenishes intermediates in the tricarboxylic acid (TCA) cycle, which have been released for
 83 biosynthesis. TCA metabolites are used as building blocks for macromolecular compounds,
 84 e.g. almost half of all amino acids in prokaryotes are directly synthesized from oxaloacetate
 85 and α-ketoglutarate (Fuchs 1999). For this purpose, heterotrophs use the enzymes pyruvate
 86 carboxylase present in a large variety of organisms, including prokaryotes, archaea, yeasts,

87 fungi and higher organisms (e.g. mammals), and phosphoenol pyruvate (PEP) carboxylase,
88 widely distributed in bacteria (Attwood 1995; Jitrapakdee and Wallace 1999; Sauer and
89 Eikmanns 2005; Jitrapakdee et al. 2008) (Fig. 1). The replenishment of metabolites
90 continuously withdrawn from the TCA cycle via the anaplerotic reaction of PEP carboxylase
91 entails an assimilation of CO₂ corresponding to 25% of the initial substrate's carbon content.
92 In a systematic stable isotope labelling experiments with *Bacillus subtilis*, a gram-positive
93 heterotrophic bacterium widespread in the environment, the interdependency of pathways
94 and rates of CO₂-fixation on the concurrent utilization of organic substrate(s) was explored
95 (Spona-Friedl et al. 2020). Over the course of the experiments *B. subtilis* assimilated 6% and
96 5% of carbon biomass from the external H¹³CO₃ pool when growing on glucose and lactate,
97 respectively (Spona-Friedl et al. 2020). Growth on malate, an intermediate of the TCA cycle,
98 expected to serve directly to refill the oxaloacetate pool of the TCA cycle, still revealed a
99 contribution to biomass production from inorganic carbon of 3% (Spona-Friedl et al. 2020).
100 PEP carboxylase was still actively transforming pyruvate to oxaloacetate. Heterotrophic CO₂-
101 fixation continued to a lower extent even in the absence of cell growth during the stationary
102 phase (Spona-Friedl et al. 2020), indicating that anaplerotic reactions are important in low-
103 productivity habitats (see below).

104 Overall, heterotrophic CO₂ fixation via anaplerosis in microorganisms contributes around 1
105 to 8% to the carbon biomass (Romanenko 1964, Perez and Martin 1982, Doronina and
106 Trotsenko 1984, Miltner et al. 2004, Roslev et al. 2004, Hesselsoe et al. 2005, Sandruckova
107 et al. 2005, Feisthauer et al. 2008, Akyniede et al. 2020, Spona-Friedl et al. 2020). Under
108 particular environmental conditions even higher contributions were reported (Perez and
109 Martin 1982). The advantage that CO₂ is readily available to the cell either as atmospheric
110 gas or, more commonly, in its hydrated form HCO₃⁻, obviously outcompetes the
111 disadvantage that carboxylation is generally an endergonic reaction (Faber et al. 2015). This
112 thermodynamic obstacle may be less important when carboxylation supports the
113 assimilation of organic substrates more reduced than the organism's biomass, resulting in
114 carbon-limited but excess-energy conditions (Heijnen and Roels, 1981, Ensign et al. 1998,
115 von Stockar et al. 2006, Battley 2013). In this case, in addition to anaplerosis further
116 carboxylation reactions are induced (Fig. 1) to add oxidized C (from CO₂) to the reduced
117 organic substrate for adjusting the degree of reduction to that of the biomass (Fig. 2). For
118 example, the assimilation of leucine and propionate into biomass entails carboxylation of
119 the initial C-6 and C-3 carbon bodies, respectively and thus, triggers an assimilation of
120 dissolved inorganic carbon (DIC) that corresponds to 17% and 33% of the initial substrate's
121 carbon content, respectively (Erb 2011). In aerobic methane oxidation, the full oxidation
122 potential of one molecule of CO₂ is needed to adjust the high degree of reduction of
123 methane to that of biomass during its assimilation. Consequently, methanotrophs derive up
124 to 50% of their carbon biomass from CO₂ (Strong, et al. 2015, Battley 2013).

125



127

128 **Figure 2:** Anaplerotic CO₂ fixation contributes 1-8% of carbon biomass (indicated by the grey band) in
 129 heterotrophic cells. Dependent to the organism and in relation to the uptake of the individual organic
 130 compounds and their entry into the TCA cycle and central metabolic pathways the relative amount of
 131 inorganic carbon assimilated varies, as highlighted by the red arrows. See examples for malic and lactic acid.
 132 With organic carbon sources more reduced than the organism’s biomass (dashed line) further carboxylation
 133 reactions are induced (indicated by black line), increasing the overall carbon contribution from CO₂ beyond
 134 anaplerosis (grey band). In methanotrophs, 50% of the cell’s carbon may originate from CO₂ fixation. For
 135 further explanations, see text.

136

137 Besides the degree of reduction of organic carbon sources, the partial pressure of CO₂ plays
 138 a role. Carboxylases may catalyze carboxylation as well as decarboxylation of organic
 139 compounds and the equilibrium of the reaction depends on the concentrations of all
 140 compounds involved. An increase in the CO₂ concentration may move the equilibrium of the
 141 reaction toward the product of the carboxylation, and thus leading to an increase in CO₂
 142 fixation.

143 In a physiological context, the amount of inorganic carbon fixed by heterotrophs, either
 144 from an endogenous or exogenous source, may be dependent on the metabolic state of the
 145 organisms and the specific environmental conditions. In their early work, Romanenko et al.
 146 (1972) suggested that the rate of heterotrophic anaplerotic fixation of DIC is strictly
 147 proportional to the heterotrophic bacterial carbon production. Since then, a number of

148 factors have been identified potentially influencing the relative contribution of anaplerotic
149 and other non-autotrophic CO₂ fixation reactions on biomass production. In laboratory
150 experiments with the bacterial strain *Thiobacillus novellus*, for example, a higher amount of
151 CO₂ was fixed under nutrient limited conditions (Perez and Matin 1982). Moreover,
152 mixotrophic bacterial strains fixed more DIC compared to those grown autotrophically
153 (Perez and Matin 1982). Fungi fixed relatively more CO₂ at lower organic carbon (glucose
154 and maltose) concentrations (Schinner et al. 1982). The degree of heterotrophic CO₂ fixation
155 highly depended on the availability of easy degradable organic carbon sources (Schinner et
156 al. 1982).

157 Studies on the possible relationship between heterotrophic DIC fixation and the activity of
158 prokaryotic cells revealed contradicting results. While Roslev et al. (2004) mentioned
159 actively growing cells fix more DIC than resting cells, Merlin et al. (2003) report enhanced
160 uptake of DIC by heterotrophic bacteria during slow growth and starvation. A relationship
161 between DIC and heterotrophic bacterial production has been reported frequently as
162 exemplified below.

163

164 **2. Respiration and non-autotropic CO₂ fixation**

165 The production of CO₂ via respiration and the parallel fixation of CO₂ in heterotrophs take
166 place simultaneously. The heterotrophic fixation of CO₂ is thus generally considered a back-
167 reaction, i.e., part of the originally produced CO₂ from respiration is re-assimilated.
168 Following this line of arguments, the more reduced an organic substrate is the less CO₂ is
169 released (Fig. 2). Heterotrophic fixation of DIC does not necessarily lead to a net carbon
170 biomass production, however, if microbes oxidize geogenic methane, this would result in a
171 net carbon biomass production. Experimentally it is difficult to differentiate respiratory CO₂
172 flux from concurrent anaplerotic CO₂ fixation. As a consequence, there are numerous
173 experiments and field studies determining dark CO₂ fixation, but only a few studies
174 quantified the assimilation of DIC by non-autotrophs.

175 Respiration in aquatic systems is frequently determined via the consumption of dissolved
176 oxygen (Robinson and Williams 2005) potentially underestimating the carbon use efficiency
177 of heterotrophs. Depending on the substrate, the respiration quotient ($\Delta\text{CO}_2/-\Delta\text{O}_2$) varies
178 between 0.7 – 1.3 (Robinson 2019) leading to an error between 20 and 40% with regard to
179 CO₂ production from respiration. Moreover, the respiration quotient also varies because
180 other oxygen consuming processes are potentially taking place simultaneously (e.g.
181 nitrification) (Robinson 2019). For instance, it is 138 O₂ for 106 CO₂ for ideal Redfield type
182 organic matter, and 150 O₂ for 106 CO₂ for more realistic marine organic matter (Fraga et al.
183 1998; Paulmier et al. 2009). Calculations based on a study on temperate forest soils
184 revealed a reduction of overall CO₂ emissions due to dark CO₂ fixation by mainly

185 heterotrophic microbes (Akinyede et al. 2020). Collectively, with respect to C cycling,
186 heterotrophic CO₂ fixation and the carbon flux from the inorganic pool into heterotrophic
187 biomass can be regarded as a process more important than hitherto assumed.

188

189 **3. CO₂ fixation in habitats dominated by heterotrophs**

190 In contrast to sunlit habitats, where photoautotrophs make up a significant portion of the
191 total biomass and photosynthesis is of major importance in carbon cycling, heterotrophs
192 and chemolithoautotrophs represent the prevailing biota in the “dark habitats”, i.e., soils,
193 subsurface environments and the deep sea. These dark environments, **although**
194 **characterized by disproportionately lower biological activity**, exceed their photic counterparts
195 in both, volume and biomass. In the oceans, the deep sea (below 200 m) exceeds the sunlit
196 surface layer by a factor of 18 in volume and, remarkably, by a factor of two in biomass
197 (Aristegui et al. 2009). Therefore, the so-called “dark CO₂ fixation” does not only occur in
198 specific 'hot spots' on the seafloor (hydrothermal vents, cold seeps and mud volcanoes), or
199 in anoxic waters, but **also** throughout the entire oxygenated 'dark' water column (Reinthal
200 et al., 2010, Yakimov et al., 2014). In limnic environments, the dark groundwater ecosystems
201 outnumber surface waters 100-fold in terms of water volume (Danielopol et al. 2003), and
202 similarly, also soils are with the exception of their surface exclusively dark habitats.

203 Yet, heterotrophic CO₂ fixation does not occur only in the dark environments since
204 heterotrophs are also found in the photic zone. This is particularly relevant in the ocean
205 because the photic zone is where the highest biomass concentrations are found. Recently, it
206 has been estimated that the inclusion of dark CO₂ fixation (integrated over the euphotic
207 layer, 0-150 m depth) would increase oceanic primary production estimates by 2.5–22 %
208 (Baltar et al., 2019). A similar situation might be assumed for surface inland waters,
209 however, global estimations are missing so far.

210 Dark DIC fixation has been reported for all types of ecosystems, including marine habitats
211 (Wuchter et al. 2003, Middelburg 2011, DeLorenzo et al. 2012, Molari et al. 2013, Baltar and
212 Herndl 2019, Lengger et al. 2019, Smith et al. 2019, Vasquez-Cardenas et al. 2020), brackish
213 and freshwater systems (Bräuer et al. 2013, Santoro et al. 2013, Noguera et al. 2015,
214 Signori et al. 2017, Vick-Majors and Priscu 2019, Zhao et al. 2020), cave waters and
215 groundwater ecosystems (Pedersen & Ekendahl 1992a, 1992b; Kotelnikova & Pedersen
216 1998, Kellermann et al. 2012, Lazar et al. 2017), and soil habitats (Ehleringer et al. 2000,
217 Miltner et al. 2004, 2005, Šantrůčková et al. 2005, 2018, Akinyede et al. 2020 and references
218 therein). In the absence of solar radiation, particularly in the dark ocean, CO₂ fixation rates
219 of up to ~125 mg C m⁻³ d⁻¹ have been measured, amounting to 30% (on a per volume basis)
220 of the phototrophic CO₂ fixation in ocean surface waters (Zopfi et al. 2001, Detmer et al.
221 1993, Casamayor et al. 2001, Baltar et al. 2010). In a eutrophic lagoon, dark DIC fixation

222 accounted for 31% of total DIC fixation in the water column (Lliros et al. 2011). Recently it
223 was shown that the ratio between dark/light CO₂ fixation in oceanic surface waters which is
224 usually around 0.1 increases with depth reaching a ratio of 1 at 120-160 m depth (Baltar et
225 al., 2019). In the past, however, dark DIC fixation has frequently been attributed to the
226 activity of chemoautotrophs only. **Only a few studies so far provided strong quantitative**
227 **evidence for heterotrophic CO₂ fixation in aquatic and terrestrial ecosystems** (Tab. 1).

228 As indicated, part of the dark CO₂ fixation in oceans has been attributed to
229 chemolithoautotrophic archaea (Wuchter et al. 2003, Ingalls et al. 2006) obtaining the
230 energy required for the endergonic carboxylation through the oxidation of reduced
231 inorganic compounds, such as ammonia or hydrogen sulfide (Swan et al. 2011; Zhang et al.
232 2020). A total annual chemolithoautotrophic CO₂ fixation rate of 0.77Pg C was calculated for
233 the oceans (Middelburg 2011). The observed fluxes of the reduced inorganic compounds
234 available as energy sources, however, seem largely insufficient to explain the relatively high
235 dark CO₂ fixation rates (Overbeck 1979, Tuttle and Jannasch 1979, Baltar et al. 2010,
236 Reinthaler et al. 2010, Herndl and Reinthaler 2013). In some cases, the supply rates of the
237 reduced inorganic compounds used as an energy source explain less than 40% of the
238 observed dark CO₂ fixation rates (Zopfi et al. 2001). Recently, chemoautotrophic nitrification
239 was estimated to explain <13% of the dark CO₂ fixation (integrated over the euphotic zone)
240 with the rest coming from either heterotrophic DIC fixation or other chemoautotrophic
241 processes (Baltar and Herndl 2019).

242 The potential energy sources for the unexplained proportion of the dark CO₂ fixation remain
243 enigmatic. Possible explanations could be either an underestimation of the supply rates of
244 reduced inorganic compounds or the uptake of CO₂ by heterotrophic organisms (Zopfi et al.
245 2001, Baltar et al. 2019). In the surface ocean in particular, DIC incorporation via anaplerotic
246 reactions might play an important role in compensating metabolic imbalances in marine
247 bacteria under oligotrophic conditions, contributing > 30 % of the carbon incorporated into
248 biomass (González et al. 2008; Palovaara et al., 2014). Evidence for the latter comes from
249 experiments with Arctic seawater, which exhibited high DIC fixation rates (0.5–2.5 µg C L⁻¹ d⁻¹)
250 correlating with heterotrophic bacterial production (Alonso-Sáez et al. 2010). Using
251 different molecular tools, DIC uptake was attributed mainly to heterotrophic *Gamma*- and
252 *Betaproteobacteria* rather than to typical chemoautotrophs, thus showing that
253 chemolithoautotrophs were not the main drivers of CO₂ fixation in this habitat (Alonso-
254 Sáez et al. 2010). Further evidence comes from the genome of *Polaribacter* sp. MED152, a
255 representative of Bacteroidetes, which typically comprise about 10–20% of the prokaryotic
256 abundance in seawater (González et al. 2008). A unique combination of membrane
257 transporters and carboxylases in these organisms indicates the importance of anaplerosis
258 besides other DIC fixation pathways (González et al. 2008). If the heterotrophic metabolism
259 of bacteria is suddenly intensified (e.g., after an input of organic matter), dark DIC fixation
260 rates and the expression of transcripts associated with key anaplerotic enzymes increase

261 proportionally (Baltar et al., 2016). As mentioned above, contradicting results were obtained
262 on the relationship between heterotrophic CO₂ fixation and the availability of organic
263 matter. A few studies suggest a relative increase in dark DIC fixation in oligotrophic habitats
264 harboring slow-growing or starving bacterial populations (Perez and Matin 1982, Schinner et
265 al. 1982, Merlin et al. 2003, Alonso-Sáez et al. 2010, Santoro et al. 2013). Considering the
266 slow community-wide specific growth rates of heterotrophic bacteria in oligotrophic and/or
267 cold waters, such as the marine aphotic zone, the Arctic Ocean, deep sea sediments,
268 groundwater systems and the terrestrial subsurface, alpine limnic systems and deep-lake
269 sediments, enhanced anaplerotic DIC uptake can be expected. However, there is also
270 evidence for the stimulation of dark DIC fixation in response to organic matter enrichment
271 in different types of soils (Miltner et al. 2005, Šantrůčková et al. 2018). Hence, these
272 contradictory findings require further, more systematic research.

273 Other environmental factors that may influence dark DIC fixation include the concentrations
274 of CO₂ and bicarbonate as inorganic carbon sources. An increase in the CO₂ concentration
275 may shift the equilibrium of the carboxylation-decarboxylation reactions increasing CO₂
276 fixation. Elevated partial pressure of CO₂ might stimulate dark DIC fixation. In temperate
277 forest soils, rates of dark microbial CO₂ fixation were positively correlated with the CO₂
278 concentration (Spohn et al. 2019). Similarly, with increasing CO₂ concentrations, higher dark
279 DIC fixation was observed in wetland soils affected by subcrustal CO₂ degassing (Beuling et
280 al. 2015). Here, besides known chemoautotrophs, CO₂ fixation via anaplerotic reactions was
281 shown for putatively heterotrophs, i.e., subdivision 1 Acidobacteriaceae, lacking enzymatic
282 pathways for autotrophic CO₂ fixation (Beuling et al. 2015). In experiments with two marine
283 heterotrophic bacterial isolates, elevation of CO₂ concentration provoked an increase in CO₂
284 fixation along with a decrease in respiration (Teiro et al. 2012). Thus, we may assume that a
285 rise in CO₂ concentrations and CO₂-induced geochemical changes will alter carbon turnover
286 in affected ecosystems with dark DIC fixation and anaplerotic reactions becoming more
287 important.

288

289 **4. Quantitative estimates of heterotrophic CO₂ fixation in different environments**

290 *Quantification of heterotrophic DIC fixation*

291 It is difficult to properly quantify heterotrophic fixation of inorganic carbon in the
292 environment. Not surprisingly, quantitative data almost exclusively originate from
293 laboratory experiments using cultures and tissues in combination with carbon isotopic
294 labeling (e.g. Spona-Friedl et al. 2020). Field studies generally report on dark carbon fixation,
295 including the activity of chemoautotrophs and heterotrophs. So far, evidence for the
296 significant contribution of heterotrophic CO₂ fixation, as highlighted for selected studies in
297 Tab. 1, is based on additional measures complementing the quantification of dark carbon

298 fixation, i.e. molecular microbial community analysis occasionally including the quantitative
299 assessment of functional genes involved in carbon fixation and characterization of
300 environmental conditions. In fact, significant dark carbon fixation (i) in the obvious absence
301 of chemoautotrophs and related genes or (ii) in well oxygenated environments that lack
302 potential electron donors (e.g. H₂, H₂S, NH₄), led authors to conclude on the quantitative
303 importance of heterotrophic carbon fixation (e.g. Miltner et al. 2005; Alonso-Saéz et al.
304 2010; Yakimov et al. 2014; Šantrůčková et al. 2018; Akinyede et al. 2020). Individual studies
305 succeeded to follow the label of inorganic carbon in environmental samples into amino
306 acids of microorganisms (Spohn et al. 2019; Spona-Friedl et al. 2020). In future, the
307 combined application of metabolomics and isotope tracing may help further developing this
308 field of research.

309

310 *Heterotrophic CO₂ fixation in different habitats*

311 Measurements of dark DIC fixation with a strong evidence of a significant contribution of
312 heterotrophic assimilation of DIC are scarce. In Table 1, we provide a compilation of studies
313 conducted in soils, marine and limnic ecosystems. Where possible, we compared dark DIC
314 fixation rates with heterotrophic activity. In marine and limnic systems, heterotrophic
315 carbon production as a widely applied activity measurement was used. In soils, we
316 compared dark DIC fixation rates with respiration, i.e., CO₂ production. Dark DIC fixation
317 rates in different marine systems range between 0.1 and 206 µg C L⁻¹ d⁻¹ with highest values
318 found in a eutrophic lagoon and lowest values in the deep waters of the Mediterranean Sea
319 (Tab. 1). Data from limnic systems originate from lake sediments with dark DIC fixation rates
320 between 0.12 and 48 mg C m⁻² d⁻¹ (Tab. 1). Projecting these numbers to only the top 10 cm
321 of sediment in the different lakes (which is a gross simplification), values of 1.2-480 µg C L⁻¹
322 sediment d⁻¹ are obtained. When compared to rates of bacterial carbon production, dark DIC
323 fixation rates in these habitats accounted for a considerable fraction of total carbon
324 assimilation, occasionally even exceeding it (Tab. 1). In soils, the dark DIC fixation rates
325 which were attributed mainly to the activity of heterotrophs amounted to 0.04-39% of the
326 overall respiration rate (Tab. 1). Dark DIC fixation rates range from 36 ng C to 23.6 µg C g⁻¹ d⁻¹
327 ranging over three orders of magnitude (Tab. 1). The contribution of heterotrophically
328 fixed DIC to biomass carbon of microbes ranged from 0.2-1.1% in temperate forest soil
329 (Akinyede et al. 2020), 0.2-4.6% in temperate forest and field soils (Santruckova et al. 2005),
330 to 7% in arable soil (Miltner et al. 2004). Santruckova et al. (2005) estimated the overall
331 heterotrophic CO₂ fixation to be even higher, i.e., 1.9-11.3% taking into account that the
332 labile fraction of the biodegradable organic carbon resulted from metabolites released by
333 spilling reactions of microorganisms due to a limitation in inorganic nutrients or due to the
334 presence of highly reduced energy-rich carbon sources (e.g. Tempest et al. 1992). A

335 contribution of heterotrophic CO₂ fixation to biomass carbon of 6.5±2.8% was found in
 336 drinking water biofilms and activated sludge (Roslev et al. 2004).

337

338 **Tab. 1.:** Dissolved inorganic carbon (DIC) assimilation rates from a range of aquatic (marine and limnic) and soil
 339 environments. Dark carbon fixation (DCF) is shown as fraction of either bacterial heterotrophic production (BP)
 340 or respiration. Original data were converted to similar units whenever possible to allow comparison.

Aquatic ecosystems	Depth [m]	DIC fixation [$\mu\text{g C L}^{-1} \text{d}^{-1}$]	BP [$\mu\text{g C}^{-1} \text{d}^{-1}$]	DCF/BP [%]	Source	Remarks
Arctic	Seawater cultures	0.5-2.3	0.4-2.5	100%	Alonso-Saéz et al. 2010	Only potential for DCF
Mediterranean Sea	4900	0.096 ± 0.02	0.048	200%	Yakimov et al. 2014	Only anaplerotic
Tropical South China Sea	200-1500	0.72-1.68	0.48- 4.8	40-105%	Zhou et al. 2017	Probably a large fraction anaplerotic
Tropical Estuary	1-18	4.8-14.4	55.2-1142	1.3-9%	Signori et al. 2018	Probably mostly anaplerotic
Eutrophic lagoon	1-5	206			Lliros et al. 2011	Probably mostly anaplerotic
Boreal lakes sediments	1-3	13.2-48 mg C m ⁻² d ⁻¹	BP 96-216 mg C m ⁻² d ⁻¹	8.4-37.4%	Santoro et al. 2013	Probably a large fraction anaplerotic
Tropical lakes sediments	1-3	0.12-20.4 mg C m ⁻² d ⁻¹	BP 14.4- 583 mg C m ⁻² d ⁻¹	0.4-80.4%	Santoro et al. 2013	Probably a large fraction anaplerotic
Deep granitic groundwater biofilms	812-1240	0. 2-2 $\mu\text{g C m}^{-2} \text{d}^{-1}$	n.d.	n.d.	Ekendahl and Pedersen 1994	Probably a large fraction anaplerotic
Terrestrial ecosystems		DIC fixation [$\mu\text{g C g}^{-1} \text{d}^{-1}$]	R [$\mu\text{g CO}_2\text{-C g}^{-1} \text{d}^{-1}$]	DCF/R [%]		
Temperate forest soil	0-0.7	0.036-0.32	0.95-19.1	1.2-3.9%	Spohn et al. 2019	¹³ C label mainly in AA, indicating anaplerosis
	0-1	0.06-0.86	n.d.	n.d.	Akinyede et al. 2020	Dominance of heterotrophs
Temperate agricultural soil	0-0.3	0.26	.63	2.7%	Miltner et al. 2004	Probably a large fraction anaplerotic
	0-0.3	0.19	9.82	1-5%	Miltner et al. 2005	DCF mainly driven by aerobic heterotrophs
Range of temperate forest & field soils	0.05-0.15	1.82-23.6*	0.65-9.16	3-39%	Šantrůčková et al. 2005	Probably a large fraction anaplerotic
	0-0.15	0.035-0.4	n.d.	n.d.	Nel and Cramer 2019	Probably mostly anaplerotic
Arctic tundra soils		0.04-0.08	0.79-10.7	0.04-16%	Šantrůčková et al. 2018	Anaplerotic enzymes comprised the majority of carboxylase genes.

341

342

343

344

*Values taken from Table 2 in Akinyede et al. 2020
 n.d. not determined

345

346

347 *Carbon biomass stock originating from heterotrophic CO₂ fixation*

348 While it is difficult to derive global estimations from the few studies that measured
349 heterotrophic CO₂ fixation rates in marine, limnic and terrestrial ecosystems, we may use a
350 conservative approach assuming that at least 1-5% of carbon biomass of all heterotrophs
351 originates from anaplerotic DIC fixation. Earth's total living biomass is estimated to amount
352 to about 499 – 738 Pg C, of which approx. 451 – 653 Pg C is photoautotrophic biomass (Bar-
353 On et al. 2018). Heterotrophic biomass thus contributes 47 – 85 Pg C (Table SI-1). The,
354 uncertainties of the estimates of heterotrophic biomass of the terrestrial subsurface,
355 however, are high (Whitman et al. 1998, McMahon and Parnell 2014, Bar-On et al. 2018).
356 Nevertheless, following this line of evidence anaplerotic CO₂ fixation contributes between
357 0.5 – 5 Pg C to the living biomass.

358

359 *Carbon flux related to heterotrophic CO₂ fixation*

360 In terms of annual global heterotrophic production rates, oceans and the terrestrial
361 subsurface (including soils) are the main habitats of heterotrophic CO₂ fixation (Cole et al.
362 2002; Magnabosco et al. 2018) (Table SI-2). Recently, Akinyede et al. (2020) estimated a
363 global dark CO₂ fixation rate of all temperate forest soils of 0.26 ± 0.07 Pg C yr⁻¹. We
364 calculated a global heterotrophic C production of 34 – 245 Pg C yr⁻¹, which would translate
365 into 0.34 – 12.3 Pg of DIC bound by heterotrophic CO₂ fixation each year. Interestingly,
366 these numbers are consistent with the recently calculated contribution of CO₂ fixation for
367 the integrated epipelagic ocean of ca. 1.2– 11 Pg C yr⁻¹ (Baltar and Herndl 2019). This is a
368 significant carbon flux amounting to 0.3-14% of the global net amount of carbon produced
369 annually by photoautotrophs (90 – 110 Pg C yr⁻¹; Ciais et al. 2013).

370 Our estimates are subject to a high uncertainty, which, on the one hand, results from the
371 dependency of the extent of heterotrophic CO₂ fixation on the organic carbon oxidized and,
372 on the other hand, on the predominant environmental conditions. Moreover, data on
373 terrestrial and marine subsurface environments, although large in dimension, are scarce.
374 For these environments, no detailed information on the abundance, growth (yield) and
375 metabolic activity of microbial communities is available, particularly with increasing depth.
376 Most of the deeper subsurface environments, even when harboring considerable living
377 biomass, do not participate in the global carbon cycle on a short and medium time scales
378 (years to decades), but rather in centennial to geological time scales. Nevertheless, in order
379 to provide a first estimate and to be able to roughly evaluate the relevance of heterotrophic
380 CO₂ fixation for all habitats of high uncertainty (e.g. the continental subsurface) we adopted
381 a conservative approach (see also Tables SI-1 and SI-2).

382

383 **5. Conclusions**

384 Current models of carbon cycling and carbon sequestration do not account for
385 heterotrophic CO₂ fixation (Gruber et al. 2004, Le Quéré et al. 2009). Despite the
386 uncertainties in the data on heterotrophic biomass and production rates for some habitats
387 (e.g. the terrestrial subsurface), the numbers presented here represent the first attempt to
388 quantify the global contribution and relevance of heterotrophic CO₂ fixation to carbon
389 cycling. Our results indicate that heterotrophs significantly contribute to global CO₂ fixation
390 – especially (although not restricted to) in habitats experiencing elevated CO₂
391 concentrations and/or lacking a sufficient supply of degradable organic carbon. In specific
392 environments, this may explain the mismatch between autotrophic C input, consumption,
393 and sequestration that has been observed in marine systems (Baltar et al. 2009, Burd et al.
394 2010, Reinthaler et al. 2010, Morán et al. 2007, Hoppe et al. 2002, Tait and Schiel 2013).
395 Particularly in aphotic habitats (which outnumber the photic habitats in both size and
396 volume) such as the dark ocean, seafloor sediments, soils, as well as the sediments and
397 rocks of the terrestrial subsurface (Miltner et al. 2004, Miltner et al. 2005, Yakimov et al.
398 2014, Wegener et al. 2012), carbon cycling needs to be re-evaluated taking into account
399 anaplerotic CO₂ fixation and other inorganic carbon uptake pathways in heterotrophs. In
400 seafloor sediments, wetlands and marshes, as well as in other habitats where methane
401 oxidation is a key process, a large fraction (10-50%) of heterotrophic biomass potentially
402 originates from heterotrophic DIC fixation. Recently, a time-series study showed a tendency
403 towards higher ratios of dark to light DIC fixation in the top half of the euphotic layer (0– 65
404 m) in the years 2012-2019 than in the preceding years (data started in 1989), which was
405 linked to oceanographic changes (i.e., a deepening of the mixed zone) (Baltar et al., 2019).
406 Moreover, the metabolic theory of ecology posits that heterotrophic metabolism increases
407 more than gross primary production in the ocean in response to warming (see Baltar et al.,
408 2019 and reference therein), which might also make heterotrophic DIC fixation relatively
409 more important in a warmer ocean. In the light of global warming leading to an extensive
410 thawing of permafrost soils and providing new habitats for methanotrophs, these processes
411 are expected to become more important in the future. Hence, the potential contribution of
412 heterotrophic CO₂ fixation under climate change conditions clearly deserves further
413 investigations.

414

415 **Author contributions**

416 A.B., M.E. and C.G. conceived the idea for the manuscript. A.B., G.J.H. and C.G. wrote the
417 manuscript. M.S.F., M.E., M.A. F.B. and T.R. substantially commented on and edited the
418 manuscript. M.A., M.S.F. and C.G. did the literature search on available global carbon data.
419 C.G. and M.A. performed the estimation of heterotrophic CO₂ fixation on a global scale.

420

421 **Acknowledgments**

422 We acknowledge B.B. Jørgensen for commenting on an earlier draft of the manuscript. We
423 thank R. Thauer and W. Eisenreich for fruitful discussions on heterotrophic CO₂ fixation.
424 Financial support was provided by the Wittgenstein Prize (Austrian Science Fund, project
425 number Z194-B17), by the European Research Council under the European Community's
426 Seventh Framework Program (FP7/2007-2013) / ERC grant agreement No. 268595 (MEDEA
427 project) and the Austrian Science Fund (P 28781-B21) to G.J.H. Financial support was further
428 provided by the Helmholtz Center Munich to A.B., M.E., M.S.F. and C.G.

429

430 **References:**

- 431 Akyniede, R., Taubert, M., Schruppf, M., Trumbore, S. & Küsel, K. Rates of dark CO₂ fixation are
432 driven by microbial biomass in a temperate forest soil. *Soil Biol. Biochem.* 150, 107950, 2020.
- 433 Alonso-Sáez, L., Galand, P. E., Casamayor, E. O., Pedrós-Alió, C., and Bertilsson, S.: High bicarbonate
434 assimilation in the dark by Arctic bacteria, *ISME J.*, 4, 1581–1590, 2010.
- 435 Arístegui, J., Gasol, J. M., Duarte, C. M., and Herndl, G. J. Microbial oceanography of the dark ocean's
436 pelagic realm. *Limnol. Oceanogr.*, 54, 1501-1529, 2009.
- 437 Attwood, P. V. The structure and the mechanism of action of pyruvate-carboxylase. *Int. J. Biochem.*
438 *Cell B* 27, 231–249, 1995.
- 439 Baltar, F., and Herndl, G. J. Ideas and perspectives: Is dark carbon fixation relevant for oceanic
440 primary production estimates? *Biogeosci.*, 16, 3793-3799, 2019.
- 441 Baltar, F., Bayer, B., Bednarsek, N., Deppeler, S., Escribano, R., Gonzalez, C. E., ... , and Robinson, C.
442 Towards integrating evolution, metabolism, and climate change studies of marine ecosystems.
443 *Trends Ecol. Evol.*, 34, 1022-1033, 2019.
- 444 Baltar, F., Arístegui, J., Sintes, E., Gasol, J. M., Reinthaler, T., and Herndl, G. J. Significance of non-
445 sinking particulate organic carbon and dark CO₂ fixation to heterotrophic carbon demand in the
446 mesopelagic northeast Atlantic. *Geophys. Res. Lett.*, 37, 1-6, 2010.
- 447 Baltar, F., Lundin, D., Palovaara, J., Lekunberri, I., Reinthaler, T., Herndl, G. J., and Pinhassi, J.
448 Prokaryotic responses to ammonium and organic carbon reveal alternative CO₂ fixation pathways
449 and importance of alkaline phosphatase in the mesopelagic North Atlantic. *Front. Microbiol.*, 7,
450 1670, 2016.
- 451 Bar-Even, A., Noor, E., and Milo, R. A survey of carbon fixation pathways through a quantitative lens.
452 *J. Exp. Bot.* 63, 2325–2342, 2012.
- 453 Bar-On, Y. M., Phillips, R., and Milo, R. The biomass distribution on Earth. *PNAS*, 115, 6506-6511,
454 2018.
- 455 Battley, E. H. A theoretical study of the thermodynamics of microbial growth using *Saccharomyces*
456 *cerevisiae* and a different free energy equation. *Quart. Rev. Biol.*, 88, 69-96, 2013.
- 457 Beer, C., Reichstein, M., Tomelleri, E., Ciais, P., Jung, M., Carvalhais, N., Rödenbeck, C., Arain, M. A.,
458 Baldocchi, D., Bonan, G. B., Bondeau, A., Cescatti, A., Lasslop, G., Lindroth, A., Lomas, M., Luysaert,

459 S., Margolis, H., Oleson, K. W., Roupsard, O., Veenendaal, E., Viovy, N., Williams, C., Woodward, F. I.,
460 and Papale, D. Terrestrial gross carbon dioxide uptake: global distribution and covariation with
461 climate. *Science*, 329, 834-838, 2010.

462 Berg, I. A. Ecological aspects of the distribution of different autotrophic CO₂ fixation pathways. *Appl.*
463 *Environ. Microbiol.*, 77, 1925-1936, 2011.

464 Berg, I. A., Kockelkorn, D., Buckel, W., and Fuchs, G. A 3-hydroxypropionate/4-hydroxybutyrate
465 autotrophic carbon dioxide assimilation pathway in Archaea. *Science*, 318, 1782-1786, 2007.

466 Beulig, F., Heuer, V.B., Akob, D.M., Viehweger, B., Elvert, M., Herrmann, M., Hinrichs, K.-U., and
467 Küsel, K. Carbon flow from volcanic CO₂ into soil microbial communities of a wetland mofette. *ISME*
468 *J.* 9, 746–759, 2015.

469 Bräuer, S.L., Kranzler, K., Goodson, N., Murphy, D., Simon, H.M., Baptista, A.M., and Tebo, B.M. Dark
470 carbon fixation in the Columbia River's estuarine turbidity maxima: molecular characterization of
471 red-type *cbbI* genes and measurement of DIC uptake rates in response to added electron donors.
472 *Estuarine, Coast. Shelf Sci.* 36, 1073-1083, 2013.

473 Burd, A. B., Hansell, D. A., Steinberg, D. K., Anderson, T. R., Arístegui, J., Baltar, F., Beupre, S. R.,
474 Buesseler, K. O., De- Hairs, F., Jackson, G. A., Kadko, D. C., Koppelman, R., Lampitt, R. S., Nagata, T.,
475 Reinthaler, T., Robinson, C., Robison, B. H., Tamburini, C., and Tanaka, T.: Assessing the apparent
476 imbalance between geochemical and biochemical indicators of meso- and bathypelagic biological
477 activity: What the @ \$?! Is wrong with present calculations of carbon budgets?, *Deep-Sea Res. Pt. II*,
478 57, 1557–1571, 2010.

479 Casamayor, E. O., García-Cantizano, J., Mas, J., and Pedrós-Alió, C. Primary production in estuarine
480 oxic/anoxic interfaces: contribution of microbial dark CO₂ fixation in the Ebro River Salt Wedge
481 Estuary. *Mar. Ecol. Prog. Ser.*, 215, 49-56, 2001.

482 Ciais, P., Sabine, G. Bala, L. Bopp, V. Brovkin, J. Canadell, A. Chhabra, R. DeFries, J. Galloway, M.
483 Heimann, C. Jones, C. Le Quéré, R. B. Myneni, S. Piao, and P. Thornton. Carbon and other
484 biogeochemical cycles. In *Climate change 2013: The physical science basis. Contribution of working*
485 *group I to the fifth assessment report of the Intergovernmental Panel on Climate Change*, eds. T. F.
486 Stocker, D. Qin, G.-K. Plattner, M. Tignor, S. K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex & P. M.
487 Midgley, 465-570. Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press,
488 2013.

489 Cochrane, V.W. *Physiology of fungi*. John Wiley, New York, 1958.

490 Cole, J. J., Findlay, S. E. G., and Pace, M. L. Bacterial production in fresh and saltwater ecosystems : a
491 cross-system overview. *Mar. Ecol. Prog. Ser.*, 43, 1-10, 1988.

492 Danielopol, D. L., Griebler, C., Gunatilaka, A., and Notenboom, J. Present state and future prospects
493 for groundwater ecosystems. *Environ. Conserv.* 30, 104-130, 2003.

494 DeLorenzo, S., Bräuer, S. L., Edgmont, C. A., Herfort, L., Tebo, B.M., and Zuber, P. Ubiquitous
495 dissolved inorganic carbon assimilation by marine bacteria in the Pacific Northwest Coastal Ocean as
496 determined by stable isotope probing. *PLoS ONE* 7, e46695, 2012.

497 Detmer, A. E., Giesenhausen, H. C., Trenkel, V. M., Auf dem Venne, H., and Jochem, F. J. Phototrophic
498 and heterotrophic pico- and nanoplankton in anoxic depths of the central Baltic Sea. *Mar. Ecol.*
499 *Progr. Ser.*, 99, 197-203, 1993.

500 Dijkhuizen, L., and Harder, W. Current views on the regulation of autotrophic carbon dioxide fixation
501 via the Calvin cycle in bacteria. *Antonie van Leeuwenhoek*, 50, 473-87, 1984.

502 Doronina, N. V., and Trotsenko, Y. A. The levels of carbon dioxide assimilation in bacteria with
503 different pathways of 1-carbon metabolism. *Mikrobiologiya*, 53, 885-889, 1984.

504 Ensign, S. A., Small, F. J., Allen, J. R., Sluis, M. K. New roles for CO₂ in the metabolism of aliphatic
505 epoxides and ketones. *Arch. Microbiol.* 169, 179–187, 1998.

506 Erb, T. J., Brecht, V., Fuchs, G., Muller, M., Alber, B. E. Carboxylation mechanism and stereochemistry
507 of crotonyl-CoA carboxylase/reductase, a carboxylating enoyl-thioester reductase. *PNAS* 106, 8871–
508 8876, 2009.

509 Erb, T. J. Carboxylases in natural and synthetic microbial pathways. *Appl. Environ. Microbiol.*, 77,
510 8466-8477, 2011.

511 Evans, E. A., Jr., and Slotin, L. The utilization of carbon dioxide in the synthesis of α -ketoglutaric acid.
512 *J. Biol. Chem.*, 136, 301, 1940.

513 Faber, K., Fessner, W. D., and Turner, N. J. Science of synthesis: biocatalysis in organic synthesis Vol.
514 2. 672. Thieme Chemistry, 2015.

515 Feisthauer, S., Wick, L. Y., Kastner, M., Kaschabek, S. R., Schlomann, M., Richnow, H. H., Differences
516 of heterotrophic ¹³CO₂ assimilation by *Pseudomonas knackmussii* strain B13 and *Rhodococcus opacus*
517 1CP and potential impact on biomarker stable isotope probing. *Environ. Microbiol.* 10, 1641–1651,
518 2008.

519 Fraga, F., Rios, A., Perez, F., and Figueras, F. Theoretical limits of oxygen:carbon and oxygen:nitrogen
520 ratios during photosynthesis and mineralisation of organic matter in the sea. *Mar. Chem.*, 62, 161–
521 168, 1998.

522 Fuchs, G. Biosynthesis of building blocks. In *Biology of the prokaryotes*, eds. Lengeler, J. W., Drews,
523 G., and Schlegel, H. G., 110-160, Stuttgart, New York: Thieme, 1999.

524 González, J. M., Fernández-Gómez, B., Fernández-Guerra, A., Gómez-Consarnau, L., Sánchez, O., Coll-
525 Lladó, M., del Campo, J., Escudero, L., Rodríguez-Martínez, R., Alonso-Sáez, L., Latasa, M., Paulsen, I.,
526 Nedashkovskaya, O., Lekumberri, I., Pinhassi, J., and Pedrós-Alió, C.: Genome analysis of the
527 proteorhodopsin-containing marine bacterium *Polaribacter* sp. MED152 (Flavobacteria), *P. Natl.*
528 *Acad. Sci. USA*, 105, 8724– 8729, 2008.

529 Gruber, N., Friedlingstein, P., Field, C., Valentini, R., Heimann, M., Richey, J. E., Romero-Lankao, P.,
530 Schulze, E. D. & Chen, C.-T. A. The vulnerability of the carbon cycle in the 21st century: an assessment
531 of carbon-climate-human interactions. In: *The global carbon cycle: integrating humans, climate, and*
532 *the natural world*, eds. Field, C. B., and Raupach, M. R., 45-76. Washington D.C., London: Island
533 Press, 2004.

534 Han, L., Yang, K., Kulowski, K., Wendt-Plienkowski, E., Hutchinson, C. R., and Vining, L. C. An acyl-
535 coenzyme A carboxylase encoding gene associated with jadomycin biosynthesis in *Streptomyces*
536 *venezuelae* ISP5230. *Microbiol. UK* 146, 903–910, 2000.

537 Hartman, R. E., and Keen, N. T. Enzymes catalysing anaplerotic carbon dioxide fixation in *Verticillium*
538 *albo-atrum*. *Phytopathol.* 63, 947-953, 1973.

539 Hartman, R. E., Keen, N. T., and Long, M. Carbon dioxide fixation by *Verticillium albo-atrum*. *J. Gen.*
540 *Microbiol.* 73, 29-34, 1972.

541 Heijnen, J. J., and Roels, J. A. A macroscopic model describing yield and maintenance relationship in
542 aerobic fermentation processes. *Biotechnol. Bioeng.* 23, 739–763, 1981.

543 Herndl, G. J., and Reinthaler, T. Microbial control of the dark end of the biological pump. *Nat. Geosc.*,
544 6, 718-724, 2013.

545 Hesselsoe, M., Nielsen, J. L., Roslev, P., and Nielsen, P. H. Isotope labeling and microautoradiography
546 of active heterotrophic bacteria on the basis of assimilation of ¹⁴CO₂. *Appl. Environ. Microbiol.*, 71,
547 646-655, 2005.

548 Hoppe, H. G., Gocke, K., Koppe, R., and Begler, C. Bacterial growth and primary production along a
549 north-south transect of the Atlantic Ocean. *Nature*, 416, 168-171, 2002.

550 Houghton, R. A. Balancing the global carbon budget. *Ann. Rev. Earth Planet. Sci.*, 35, 313-347, 2007.

551 Ingalls, A. E., Shah, S. R., Hansman, R. L., Aluwihare, L. I., Santos, G. M., Druffel, E. R., and Pearson, A.
552 Quantifying archaeal community autotrophy in the mesopelagic ocean using natural radiocarbon.
553 *PNAS*, 103, 6442-6447, 2006.

554 Jitrapakdee, S., and Wallace, J.C. Structure, function and regulation of pyruvate carboxylase.
555 *Biochem. J.* 340, 1–16, 1999.

556 Jitrapakdee, S., St. Maurice, M., Rayment, I., Cleland, W.W., Wallace, J.C., and Attwood, P.V.
557 Structure, mechanism and regulation of pyruvate carboxylase. *Biochem. J.* 413, 369–387, 2008.

558 Kellermann, C., Selesi, D., Lee, N., Hügler, M., Esperschütz, J., Hartmann, A. & Griebler, C. Microbial
559 CO₂ fixation potential in a tar-oil-contaminated porous aquifer. *FEMS Microbiol. Ecol.* 81, 172-187,
560 2012.

561 Kleiber, M., Smith, A. H., and Black, A. L. Carbonate as precursor of milk constituents in the intact
562 dairy cow. *J. Biol. Chem.*, 195, 707-714, 1952.

563 Kornberg, H. L., and Krebs, E. H. Synthesis of cell constituents from C₂-units by a modified
564 tricarboxylic acid cycle. *Nature* 179, 988–991, 1957.

565 Kornberg, H.L. Anaplerotic sequences in microbial metabolism. *Angew. Chem. internat. Edit.* 4, 558-
566 565, 1965.

567 Kotelnikova, S., and Pedersen, K. Distribution and activity of methanogens and homoacetogens in
568 deep granitic aquifers at Äspö Hard Rock Laboratory, Sweden. *FEMS Microbiol. Ecol.*, 26, 121-134,
569 1998.

570 Krebs, H.A. Carbon dioxide assimilation in heterotrophic organisms. *Nature* 147, 560-563, 1941.

571 Lazar, C.S., Stoll, W., Lehmann, R., Herrmann, M., Schwab, V.F., Akob, D.M., Nawaz, A., Wubet, T.,
572 Buscot, F., Totsche, K.-U., and Küsel, K. Archaeal diversity and CO₂ fixers in carbonate-/siliciclastic-
573 rock groundwater ecosystems. *Archaea* 2136287, 1-13, 2017.

574 Le Quéré, C., M. R. Raupach, J. G. Canadell, G. Marland, L. Bopp, P. Ciais, T. J. Conway, S. C. Doney, R.
575 A. Feely, P. Foster, P. Friedlingstein, K. Gurney, R. A. Houghton, J. I. House, C. Huntingford, P. E. Levy,
576 M. R. Lomas, J. Majkut, N. Metzl, J. P. Ometto, G. P. Peters, I. C. Prentice, J. T. Randerson, S. W.
577 Running, J. L. Sarmiento, U. Schuster, S. Sitch, T. Takahashi, N. Viovy, G. R. van der Werf & F. I.
578 Woodward. Trends in the sources and sinks of carbon dioxide. *Nat. Geosci.*, 2, 831-836, 2009.

579 Le Quéré, C., R. M. Andrew, J. G. Canadell, S. Sitch, J. I. Korsbakken, G. P. Peters, A. C. Manning, T. A.
580 Boden, P. P. Tans, R. A. Houghton, R. F. Keeling, S. Alin, O. D. Andrews, P. Anthoni, L. Barbero, L.
581 Bopp, F. Chevallier, L. P. Chini, P. Ciais, K. Currie, C. Delire, S. C. Doney, P. Friedlingstein, T. Gkritzalis,
582 I. Harris, J. Hauck, V. Haverd, M. Hoppema, K. Klein Goldewijk, A. K. Jain, E. Kato, A. Körtzinger, P.
583 Landschützer, N. Lefèvre, A. Lenton, S. Lienert, D. Lombardozzi, J. R. Melton, N. Metzl, F. Millero, P.
584 M. S. Monteiro, D. R. Munro, J. E. M. S. Nabel, S. I. Nakaoka, K. O'Brien, A. Olsen, A. M. Omar, T. Ono,
585 D. Pierrot, B. Poulter, C. Rödenbeck, J. Salisbury, U. Schuster, J. Schwinger, R. Séférian, I. Skjelvan, B.
586 D. Stocker, A. J. Sutton, T. Takahashi, H. Tian, B. Tilbrook, I. T. van der Laan-Luijkx, G. R. van der Werf,
587 N. Viovy, A. P. Walker, A. J. Wiltshire & S. Zaehle. Global Carbon Budget 2016. *Earth System Science*
588 *Data*, 8, 605-649, 2016.

589 Lengger, S.K., Rush, D., Mayser, J.P., Blewett, J., Schwartz-Narbonne, R., Talbot, H.B., Middelburg,
590 J.J., Jetten, M.S.M., Schouten, S., Sinninghe Damsté, J.S., and Pancost, R.D. Dark carbon fixation in
591 the Arabian Sea oxygen minimum zone contributes to sedimentary organic carbon (SOM). *Global*
592 *Biogeochem. Cycl.* 33, 1715-1732, 2019.

593 Lliros, M., Alonso-Saéz, L., Gich, F., Plasencia, A., Auguet, O., Casamayor, E.O., and Borrego, C.M.
594 Active bacteria and archaea cells fixing bicarbonate in the dark along the water column of a stratified
595 eutrophic lagoon. *FEMS Microbiol. Ecol.* 77, 370–384, 2011.

596 Magnabosco, C., Lin, L. H., Dong, H., Bomberg, M., Ghiorse, W., Stan-Lotter, H., Pedersen, K., Kieft, T.
597 L., van Heerden, E., and Onstott, T. C. The biomass and biodiversity of the continental subsurface.
598 *Nature Geoscience*, 11, 707-717, 2018.

599 McMahon, S., and Parnell J. Weighing the deep continental biosphere. *FEMS Microbiol. Ecol.*, 87,
600 113-120, 2014.

601 Melzer, E., and O'Leary M. H. Anapleurotic CO₂ fixation by phosphoenolpyruvate carboxylase in C3
602 plants. *Plant Physiol.*, 84, 58-60, 1987.

603 Merlin, C., Masters, M., McAteer, S., and Coulson, A. Why is carbonic anhydrase essential to
604 *Escherichia coli*? *J. Bacteriol.* 185, 6415–6424, 2003.

605 Middelburg, J. J. Chemoautotrophy in the ocean. *Geophy. Res. Lett.*, 38, 1-4, 2011.

606 Miltner, A., Kopinke, F.-D., Kindler, R., Selesi, D., Hartmann, A., and Kästner, M. Non-phototrophic
607 CO₂ fixation by soil microorganisms. *Plant Soil*, 269, 193-203, 2005.

608 Miltner, A., Richnow H.-H., Kopinke F.-D., and Kästner, M. Assimilation of CO₂ by soil microorganisms
609 and transformation into soil organic matter. *Org. Geochem.*, 35, 1015-1024, 2004.

610 Molari, M., Manini, E., and Dell'Anno, A. Dark inorganic carbon fixation sustains the functioning of
611 benthic deep-sea ecosystems. *Glob. Biogeochem. Cycl.* 27, 212-221, 2013.

612 Morán, X. A. G., Pérez, V. & Fernández, E. Mismatch between community respiration and the
613 contribution of heterotrophic bacteria in the NE Atlantic open ocean: What causes high respiration
614 in oligotrophic waters? *J. Mar. Res.*, 65, 545-560, 2007.

615 Nel, J.A., and Cramer, M.D. Soil microbial anapleurotic CO₂ fixation in temperate soils. *Geoderma* 335,
616 170-178, 2019.

617 Noguera, I., Picazo, A., Lliros, M., Camacho, A., and Borrego, C.M. Diversity of freshwater
618 Epsilonproteobacteria and dark inorganic carbon fixation in the sulphidic redoxcline of a meromictic
619 karstic lake. *FEMS Microbiol. Ecol.* 91, fiv086, 2015.

620 Overbeck, J. Dark CO₂ uptake - biochemical background and its relevance to in situ bacterial
621 production. *Arch. Hydrobiol. Beiheft*, 12, 38-47, 1979.

622 Palovaara, J., Akram, N., Baltar, F., Bunse, C., Forsberg, J., Pedrós- Alió, C., González, J. M., and
623 Pinhassi, J.: Stimulation of growth by proteorhodopsin phototrophy involves regulation of central
624 metabolic pathways in marine planktonic bacteria, *P. Natl. Acad. Sci. USA*, 111, E3650–E3658, 2014.

625 Parkinson, S. M., Jones, R., Meharg, A. A., Wainwright, M., and Killham, K. The quantity and fate of
626 carbon assimilated from ¹⁴CO₂ by *Fusarium oxysporum* grown under oligotrophic and near
627 oligotrophic conditions. *Mycol. Res.* 95, 1345–1349, 1991

628 Parkinson, S. M., Killham, K., and Wainwright, M. Assimilation of ¹⁴CO₂ by *Fusarium oxysporum*
629 grown under oligotrophic conditions. *Mycol. Res.* 94, 959–964, 1990.

630 Paulmier, A., Kriest, I. & Oschlies, A. Stoichiometries of remineralisation and denitrification in global
631 biogeochemical ocean models. *Biogeoosci.* 6, 923–935, 2009.

- 632 Pedersen, K., and Ekendahl, S. Assimilation of CO₂ and introduced organic compounds by bacterial
633 communities in groundwater from southeastern Sweden deep crystalline bedrock. *Microb. Ecol.* 23,
634 1-14, 1992.
- 635 Pedersen, K., and Ekendahl, S. Incorporation of CO₂ and introduced organic compounds by bacterial
636 populations in groundwater from deep crystalline bedrock of Stripa mine. *J. Gen. Microbiol.* 138,
637 369-376, 1992.
- 638 Perez, R.C., and Matin, A. Carbon dioxide assimilation by *Thiobacillus novellus* under nutrient-limited
639 mixotrophic conditions. *J. Bacteriol.* 150, 46-51, 1982.
- 640 Reinthaler, T., Van Aken, H. M., and Herndl, G. J. Major contribution of autotrophy to microbial
641 carbon cycling in the deep North Atlantic, Åôs interior, *Deep-Sea Res. Pt. II*, 57, 1572– 1580, 2010.
- 642 Robinson, C., and Williams, P.J. Respiration and its measurement in surface marine waters. In:
643 Respiration in aquatic ecosystems (eds. P. A. del Giorgio and P. J. Williams) Oxford: Oxford University
644 Press, 2005.
- 645 Robinson, C. Microbial respiration, the engine of ocean deoxygenation. *Front. Mar. Sci.*, 5, 533, 2019.
- 646 Romanenko, V. I. Heterotrophic CO₂ assimilation by water bacterial flora. *Mikrobiologiya*, 33, 679-
647 683, 1964.
- 648 Romanenko, V. I., Overbeck, J., and Sorokin, Y. I. Estimation of production of heterotrophic bacteria
649 using ¹⁴C. In: Sorokin, Y. I., Kadota, H. (eds.) *Techniques for the assessment of microbial production*
650 *and decomposition in fresh waters.* IBP Handbook No. 23, Blackwell, Oxford, pp. 82-85, 1972.
- 651 Roslev, P., Larsen, M. B., Jørgensen, D. & Hesselsoe, M. Use of heterotrophic CO₂ assimilation as a
652 measure of metabolic activity in planktonic and sessile bacteria. *J. Microbiol. Meth.*, 59, 381-393,
653 2004.
- 654 Santoro, A.L., Bastviken, D., Gudasz, C., Tranvik, L., Enrich-Prast, A. Dark carbon fixation: an
655 important process in lake sediments. *PLoS ONE* 8: e65813, 2013.
- 656 Šantrůčková, H., Bird, M. I., Elhottova, D., Novak, J., Pícek, T., Simek, M., and Tykva, R. Heterotrophic
657 fixation of CO₂ in soil. *Microb. Ecol.* 49, 218–225, 2005.
- 658 Šantrůčková, H., Kotas, P., Bárta, J., Urich, T., Čapek P., Palmtag J., Eloy Alves, R. J., Biasi, C., Diáková,
659 K., Gentsch, N., Gittel, A., Guggenberger, G., Hugelius, G., Lashchinsky, N., Martikainen, P. J.,
660 Mikutta, R., Schleper, C., Schneckner, J., Schwab, C., Shibistova, O., Wild, B., and Richter, A.
661 Significance of dark CO₂ fixation in arctic soils. *Soil Biol. Biochem.* 119, 11–21, 2018.
- 662 Sauer, U., and Eikmanns B. J. The PEP–pyruvate–oxaloacetate node as the switch point for carbon
663 flux distribution in bacteria. *FEMS Microbiol. Rev.*, 29, 765-794, 2005.
- 664 Schink, B. An alternative to the glyoxylate shunt. *Mol. Microbiol.* 73, 975–977, 2009.
- 665 Schinner, F., Concin, R., & Binder, H. Heterotrophic CO₂ -fixation by fungi in dependence on the
666 concentration of the carbon source. *Phyton* 22, 81-85, 1982.
- 667 Scrutton, M. C. Assay of enzymes of CO₂ metabolism. *Methods in Microbiology* Vol 6, Part A, 479-
668 541, 1971.
- 669 Signori, C. N., Valentin, J. L., Pollery, R. C. G., and Enrich-Prast, A. Temporal variability of dark carbon
670 fixation and bacterial production and their relation with environmental factors in a tropical estuarine
671 system, *Estuaries and Coasts*, 41, 1089–1101, 2018.
- 672 Smith, A. R., Kieft, B., Mueller, R., Fisk, M. R., Mason, O. U., Popa, R., and Colwell, F. S. Carbon
673 fixation and energy metabolisms of a subseafloor olivine biofilm. *ISME J.*, 13, 1737-1749, 2019.

674 Spohn, M., Müller, K., Höschen, C., Mueller, C.W., and Marhan, S. Dark microbial CO₂ fixation in
675 temperate forest soils increases with CO₂ concentrations. *Global Change Biology* 26, 1926-1935,
676 2019.

677 Spona-Friedl, M., Braun, A., Huber, C., Eisenreich, W., Griebler, C., Kappler, A., and Elsner M.
678 Substrate-dependent CO₂-fixation in heterotrophic bacteria revealed by stable isotope labelling.
679 *FEMS Microbiol. Ecol.*, 96, fiae080, 2020.

680 Strong, P. J., Xie, S., and Clarke, W. P. Methane as a resource: can the methanotrophs add value?
681 *Environ. Sci. Technol.*, 49, 4001-4018, 2015.

682 Swan, B. K., Martinez-Garcia M., Preston C. M., Sczyrba A., Woyke T., Lamy D., et al. Potential for
683 chemolithoautotrophy among ubiquitous bacteria lineages in the dark ocean. *Science* 333, 1296-
684 1300, 2011.

685 Tait, L. W., and Schiel D. R. Impacts of temperature on primary productivity and respiration in
686 naturally structured macroalgal assemblages. *PLoS ONE*, 8, e74413, 2013.

687 Teiro, E., Fernández, A., Álvarez-Salgado, X. A., García-Martín, E. E., Serret, P., and Sobrino, C.
688 Response of two marine bacterial isolates to high CO₂ concentration *Mar. Ecol. Prog. Ser.*, 453, 27–36,
689 2012.

690 Tempest, D. W., and Neijssel, O. M. Physiological and energetic aspects of bacterial metabolite
691 overproduction. *FEMS Microbiol. Lett.* 100, 169–176, 1992.

692 Tuttle, J. H., and Jannasch H. W. Microbial dark assimilation of CO₂ in the Cariaco Trench. *Limnol.*
693 *Oceanogr.*, 24, 746-753, 1979.

694 Vasquez-Cardenas, D., Meysman, F. J. R., & Boschker, H. T. S. A cross-system comparison of dark
695 carbon fixation in coastal sediments. *Glob. Biogeochem.Cycl.* 34, 1-14, 2020.

696 Vick-Majors, T. J., and Priscu, J. C. Inorganic carbon fixation in ice-covered lakes of the McMurdo Dry
697 Valleys. *Antarctic Sci.* 1-10, 2019.

698 von Stockar, U., Maskow, T., Liu, J., Marison, I. W., and Patiño, R. Thermodynamics of microbial
699 growth and metabolism: An analysis of the current situation. *J. Biotechnol.*, 121, 517-533, 2006.

700 Wegener, G., Bausch, M., Holler, T., Thang, N. M., Mollar, X. P., Kellermann, M. Y., Hinrichs, K. U.,
701 and Boetius, A. Assessing sub-seafloor microbial activity by combined stable isotope probing with
702 deuterated water and ¹³C-bicarbonate. *Environ. Microbiol.*, 14, 1517-1527, 2012.

703 Werkman, C.H., and Wood, H.G. Heterotrophic assimilation of carbon dioxide. In: *Advances in*
704 *Enzymology and Related Areas of Molecular Biology* (Nord, F.F. and Werkman, C.H., eds.), 2, 135-
705 182, 1942.

706 Whitman, W. B., Coleman, D. C., and Wiebe, W. J. Prokaryotes: The unseen majority. *PNAS*, 95,
707 6578-6583, 1998.

708 Wood, H. G., and Werkman, C. H. The utilisation of CO₂ in the dissimilation of glycerol by the
709 propionic acid bacteria. *Biochem. J.*, 30, 48-53, 1936.

710 Wood, H.G., and Werkman, C.H. The utilization of CO₂ by the propionic acid bacteria. *Biochem. J.*, 32,
711 1262–1271, 1938.

712 Wood, H.G., and Werkman, C.H. The position of carbon dioxide-carbon in succinic acid synthesized
713 by heterotrophic bacteria. *Jour. Biol. Chem.*, 139, 377–381, 1941.

714 Wood, H. G., and Stjernholm, R. L. Assimilation of carbon dioxide by heterotrophic organisms. In
715 *Gunsalus, IC, Stanier, RY (Eds.) The Bacteria: A Treatise on Structure and Function*, vol 3. Biosynthesis
716 Academic Press, New York, 41–117, 1962.

- 717 Wuchter, C., Schouten, S., Boschker, H. T. S., and Sinninghe Damsté, J. S. Bicarbonate uptake by
718 marine Crenarchaeota. *FEMS Microbiol. Lett.*, 219, 203-207, 2003.
- 719 Yakimov, M. M., La Cono, V., Smedile, F., Crisafi, F., Arcadi, E., Leonardi, M., Decembrini, F.,
720 Catalfamo, M., Bargiela, R., Ferrer, M., Golyshin, P. N., and Giuliano, L. Heterotrophic bicarbonate
721 assimilation is the main process of de novo organic carbon synthesis in hadal zone of the Hellenic
722 Trench, the deepest part of Mediterranean Sea. *Environ. Microbiol. Rep.*, 6, 709–722, 2014.
- 723 Zhang, Y., Qin, W., Hou, L., Zakem, E.J., Wan, X., Zhao, Z., Liu, L., Hunt, K.A., Jiao, N., Kao, S.-J., Tang,
724 K., Xie, X., Shen, J., Li, Y., Chen, M., Dai, X., Liu, C., Deng, W., Dai, M., Ingalls, A.E., Stahl, D.A., and
725 Herndl, G.J. Nitrifier adaptation to low energy flux controls inventory of reduced nitrogen in the dark
726 ocean. *PNAS* 117, 4823-4830, 2020.
- 727 Zhao, Y., Liu, P., Rui, J., Cheng, L., Wang, Q., Liu, X., and Yuan, Q. Dark carbon fixation and
728 chemolithotrophic microbial community in surface sediments of the cascade reservoirs, Southwest
729 China. *Sci.Tot. Environ.* 698, 134316, 2020.
- 730 Zhou, W., Liao, J., Guo, Y., Yuan, X., Huang, H., Yuan, T., and Liu, S.: High dark carbon fixation in the
731 tropical South China Sea, *Cont. Shelf Res.*, 146, 82–88, 2017.
- 732 Zopfi, J., Ferdelman, T. G., Jørgensen, B. B., Teske, A., and Thamdrup, B. Influence of water column
733 dynamics on sulfide oxidation and other major biogeochemical process in the chemocline of
734 Mariager Fjord (Denmark). *Mar. Chem.*, 74, 29-51, 2001.