

# Management induced changes of soil organic carbon on global croplands

Kristine Karstens<sup>1,3</sup>, Benjamin Leon Bodirsky<sup>1</sup>, Jan Philipp Dietrich<sup>1</sup>, Marta Dondini<sup>2</sup>, Jens Heinke<sup>1</sup>, Matthias Kuhnert<sup>2</sup>, Christoph Müller<sup>1</sup>, Susanne Rolinski<sup>1</sup>, Pete Smith<sup>2</sup>, Isabelle Weindl<sup>1</sup>, Hermann Lotze-Campen<sup>1,3</sup>, and Alexander Popp<sup>1</sup>

<sup>1</sup>Potsdam Institute for Climate Impact Research (PIK), Member of the Leibniz Association, P.O. Box 60 12 03, 14412 Potsdam, Germany

<sup>2</sup>Institute of Biological & Environmental Sciences, University of Aberdeen, Aberdeen, UK

<sup>3</sup>Humboldt-Universität zu Berlin, Department of Agricultural Economics, Unter den Linden 6, 10099 Berlin, Germany

**Correspondence:** Kristine Karstens (kristine.karstens@pik-potsdam.de)

**Abstract.** Soil organic carbon (SOC) is one of the largest terrestrial carbon (C) stocks on Earth. The first meter of the Earth's soils profile stores three times as much carbon C as the vegetation and twice the amount of C in the atmosphere. SOC has been depleted by anthropogenic land-cover change and agricultural management. However, the latter has so far not been well represented in global carbon C stock assessments. While SOC models often simulate detailed biochemical processes that lead to the accumulation and decay of SOC, the management decisions driving these biophysical processes are still little investigated at the global scale. Here we develop a spatially explicit data set for agricultural management on cropland, considering crop production levels, residue returning rates, manure application, and the adoption of irrigation and tillage practices. We combine it with a reduced-complexity model based on the IPCC Tier 2 steady-state soil model method to create a half-degree resolution data set of SOC stocks and SOC stock changes for the first 30 cm of mineral soils. We estimate that due to arable farming, soils have lost around 26 GtC 34.6 GtC relative to a counterfactual hypothetical natural state in 1975. Yet, within the period 1975–2010 this SOC debt has been decreasing again by a net quantity of 4 Gt SOC, which can be mainly traced back to an increased input of C in crop residues due to higher crop productivity continued to expand by 5 GtC (0.14 GtC yr<sup>-1</sup>) to around 39.6 GtC. However, accounting for historical management led to 2.1 GtC less (0.06 GtC yr<sup>-1</sup>) emissions than under the assumption of constant management. We also find that SOC is very sensitive to management decisions such as residue returning indicating the necessity to incorporate better management data in soil model simulations management decisions have influenced the historical SOC trajectory most strongly by residue returning, indicating that increasing SOC sequestration by biomass retention may be a promising negative emissions technique. The reduced-complexity SOC model may allow to simulate management-induced SOC sequestration also within computationally demanding integrated (land-use) assessment modeling.

## 20 1 Introduction

Soil ~~Organic Carbon~~ organic carbon (SOC), the amount of organic carbon stored in the Earth's soil, ~~is~~ constitutes the largest terrestrial organic carbon pool. It exceeds the carbon in the atmospheric and vegetation pools multiple times (Batjes, 1996). Even small changes in processes affecting SOC lead therefore to substantial shifts in the terrestrial carbon cycle and influence the amount of CO<sub>2</sub> in the atmosphere (~~Friedlingstein et al., 2019; Minasny et al., 2017~~)(Friedlingstein et al., 2020; Minasny et al., 2017)

25 . The specific amount of carbon stored in soils globally is quantified with estimates ranging from 1500 to 2400 GtC for the first meter of the soil profile (Batjes, 1996; Sanderman et al., 2017).

Natural properties like climatic, biophysical, and landscape characteristics clearly play the most important roles to determine SOC variations over space and time. Recent studies have focused on the evaluation of total SOC stocks of the world as well as on the spatial disaggregation of soil properties such as SOC content (Batjes, 2016; Hengl et al., 2017; FAO, 2018).

30 However, these studies often do not include human interventions, like land cover change and agricultural management, in their analysis. Compared to climatic and geological ~~drivers, they~~ driving forces, human interventions alter terrestrial carbon pools over much shorter time scales and are currently one of the most dominant drivers of SOC changes on managed land (Hansis et al., 2015; Bastos et al., 2021).

The anthropogenic impact can be measured by the SOC debt (also referred to as SOC component of land-use change emissions, see Pongratz et al. (2014)), which is the amount of organic carbon soils have lost under cultivation compared to a ~~potential natural vegetated~~ hypothetical potential natural vegetation state. Sanderman et al. (2017) identified the anthropogenic SOC debt for the first meter of the soil profile due to land cover change at around 116 GtC (37 GtC for the first 30 cm), compared to previous estimates of 60–130 GtC for the first meter (Lal, 2001).

Global assessments of the carbon cycle via dynamic global vegetation models (DGVMs) ~~and~~ , Earth System Models (ESMs) or bookkeeping models (BKMs) have analyzed SOC losses as part of a comprehensive evaluation of the global carbon budget and land-use change (LUC) emissions (~~Friedlingstein et al., 2019~~)(Friedlingstein et al., 2020). While providing estimates of the magnitude of SOC losses due to ~~land cover~~ land-cover change, most models lack a detailed ~~consideration~~ representation of agricultural management. Earlier ~~DGVM and ESM based~~ DGVM- and ESM-based assessments only considered changes in land cover, but ignored the removal of biomass at harvest (Strassmann et al., 2008; Betts et al., 2015). BKMs are designed to estimate LUC related ~~emissions but~~ emissions but often ignore changes in SOC due to climate change, CO<sub>2</sub> fertilization , ~~and agricultural management~~ (Friedlingstein et al., 2019; Houghton et al., 2012; Hansis et al., 2015) and N deposition. Whereas BKM have largely improved in estimating additional emissions from wood harvest and shifting cultivation, state of the art models do not consider impacts of varying agricultural management (Friedlingstein et al., 2020; Houghton et al., 2012; Hansis et al., 20

45 .

50 Managed agricultural systems were introduced in greater detail to DGVMs and ESMs to improve the assessment of the terrestrial carbon balance (e.g. Bondeau et al., 2007; Lindeskog et al., 2013). Pugh et al. (2015) explicitly consider agricultural management in the form of tillage, irrigation and biomass extraction at harvest, but worked with stylized scenarios rather than with ~~historie~~ historical management data. They also showed the importance of accounting for the land-use history, as many

carbon emissions from agricultural soils are caused by ~~historie~~-historical LUC and the slow decline of SOC under cropland  
55 before ~~it reaches~~-a new equilibrium is reached.

In global-scale carbon cycle assessments, management systems are typically represented as spatially explicit patterns that are static in time (e.g. growing seasons (Portmann et al., 2010), multiple cropping systems (Waha et al., 2020), irrigation systems (Jägermeyr et al., 2015)) or as stylized scenarios (~~e.g. Pugh et al., 2015; Lutz et al. (2019))~~)(e.g. Pugh et al., 2015; Lutz et al., 2019)  
60 . Herzfeld et al. (2021) account for historical changes in fertilizer and manure inputs, residue removal rates and tillage systems and report SOC losses from cropland expansion over the period from 1700–2018 of 215 GtC. Within their stylized future management scenarios they find that none of the management aspects considered (residue management, no-tillage) can create a net carbon sink on current cropland areas under future climate change.

More data sets on spatially explicit agricultural management time series with global coverage ~~become~~-are becoming available (e.g. on tillage systems, see (~~Porwollik et al., 2018~~), (~~Prestele et al., 2018~~))~~and model~~ (Porwollik et al., 2019; Prestele et al., 2018)  
65 ) and modeling approaches are increasingly being developed to project the dynamics of management systems into the future (e.g. (~~Iizumi et al., 2019~~), (~~Minoli et al., 2019~~))(Iizumi et al., 2019; Minoli et al., 2019)), but have — to our knowledge — not yet found their way into comprehensive assessments of the terrestrial carbon cycle in DGVMs and BKM.

Field-scale models (~~Del Grosso et al., 2001; Coleman et al. (1997); Smith et al. (2010); Taghizadeh-Toosi et al. (2014))~~ (Del Grosso et al.  
70 are able to better account for ~~historie~~-historical agricultural management if detailed information on crop yield levels, fertilizer inputs and various other on-farm measures is available for the studied sites. However, due to the lack of comprehensive global management data as input to these models, scaling up to the global domain remains a complex challenge (Morais et al., 2019).

~~The objective of our study is to provide the first global, spatially explicit SOC and SOC debt map that considers spatially explicit and time-variant historic agricultural management. To achieve this objective we create and provide~~ Managed soils have been increasingly studied not only for their carbon emitting behavior, but also because of their capacity to re-store carbon (soil carbon sequestration (SCS) techniques). However, assessing SCS dynamically considering the interdependency with environmental, social and economic sustainability targets has been difficult so far, as integrated assessment models (IAMs) (Popp et al., 2016; Rogelj et al., 2018; Forster et al., 2018) have not integrated soil management into their mitigation pathways. More detailed process-based models are typically computationally too demanding to be integrated into optimization-based IAMs. Better accounting for soil carbon management in IAMs thus requires a light-weight model suitable for iterative modeling  
75 with detailed options to represent agricultural soil management.  
80

The objectives of our study are (1) to develop a reduced-complexity SOC model able to account for SCS in IAM frameworks; (2) to create a comprehensive data set of the global gridded management time series~~data~~, including crop production levels, residue ~~returning~~-input rates, manure ~~application~~amendments, and the adoption of irrigation and tillage practices.~~We simulate SOC stocks, dynamics and the SOC debt for 1975–2010. Using a scenario analysis, we~~; and (3) to provide global as well  
85 as spatially explicit SOC and SOC debt estimates that consider spatially explicit and time-variant agricultural management. We decompose the contribution of different management activities through a scenario analysis, identifying the most impacting management decisions for SOC development. Moreover, we compare our model performance against other SOC stock and SOC emission estimates, to evaluate the suitability of this reduced-complexity approach for integration into IAM modeling.

## 2 Methods

90 In Sect. 2.1 we introduce the basic concept of SOC dynamics as applied in this study and explained in more detail [within in](#) the refinement of the IPCC guidelines vol. 4 ([Calvo-Buendia et al., 2019](#)) [Chapter 5 on “Cropland” \(Ogle et al., 2019\)](#). We additionally describe how we configured and extended the [steady-state method \(for model code see \(Karstens and Dietrich, 2020\) Tier 2 modeling approach \(for model code see Karstens and Dietrich, 2020\)\)](#). In Sect. 2.2 we shortly refer to the concept of stock change factors as outlined in the Tier 1 approach of the IPCC guidelines (Eggleston et al., 2006; Calvo Buendia et al., 95 2019). Section 2.3 provides a detailed description of the global, gridded management data used to drive the model, including crop production levels, residue input rates, manure amendments, and the adoption of irrigation and tillage practices (for model code see Bodirsky et al., 2020a). In Sect. 2.4 we define the [management](#) scenarios used to [complement our historic model results](#) [analyze the role of different management aspects in historical cropland SOC dynamics](#).

### 2.1 SOC stocks and stock changes following the Tier 2 modeling approach

100 Following the Tier 2 [steady-state modeling](#) approach of the refinement of the IPCC guidelines vol. 4 ([Calvo-Buendia et al., 2019](#)) [Chapter 5 on “Cropland” \(Ogle et al., 2019\)](#); referred to as [steady-state method](#) [Tier 2 modeling approach](#) in the following), we estimate [soil organic carbon \(SOC\)](#) stocks and their change over time for cropland at half-degree resolution from 1975 to 2010. We restrict our analysis to the first 0-30 cm of the soil profile. Moreover, we assume the current *SOC* state converges towards a steady state, which itself depends on biophysical, climatic and agronomic conditions. Therefore, we take the following 105 three steps for each year of our simulation period: (1) We calculate annual land-use type-specific steady states and decay rates for *SOC* stocks (Sect. 2.1.1); (2) we account for land conversion by transferring *SOC* from and to natural vegetation (Sect. 2.1.2), (3) we estimate *SOC* stocks and changes based on the stocks of the previous time step, the steady state stocks and the decay rate (Sect. 2.1.3). To initialize the first year of our simulation period we use a spin-up period of 74 years (Sect. 2.1.4).

#### 2.1.1 Steady-state SOC stocks and decay rates

110 In a simple first order kinetic approach the steady-state soil organic carbon stocks  $SOC^{eq}$  are given by

$$SOC_{i,t,sub,lu}^{eq} = \frac{C_{i,t,sub,lu}^{in}}{k_{i,t,sub,lu}} \quad (1)$$

with  $C^{in}$  being the carbon inputs to the soil,  $k$  denotes the soil organic carbon decay rate. This equation is valid for all grid cells  $i$  and all years  $t$ . We use the [steady-state method](#) [Tier 2 modeling approach](#) for our calculations, which assumes three soil carbon sub-pools *sub* (active, slow and passive) and interactions between them, following the approach in the Century model 115 (Parton et al., 1987). Annual carbon inflow to each sub-pool and annual decay rates of each sub-pool are land-use type *lu* specific. We distinguish two land-use types: cropland and uncropped land under potential natural vegetation as representative for all other land-use types including forestry and pastures (referred to as natural vegetation in the following).

Carbon inputs for cropland are below- and above-ground [crop](#) residues left or returned to the field (see Sect. 2.3.2) and manure inputs (see Sect. 2.3.3); for natural vegetation, [litterfall](#) including fine root turnover (Schaphoff et al., 2018b) is the

120 only source of carbon inflow to the soil. Following the IPCC guidelines (~~Calvo-Buendia et al., 2019~~)(Ogle et al., 2019), carbon inputs are disaggregated into metabolic and structural components depending on their lignin and nitrogen content. For each component the sum of all carbon input sources is allocated to the respective *SOC* sub-pools via transfer coefficients. This implies that both the amount of carbon and its structural composition determine the effective inflow into the different pools.

125 Whereas residue and manure default lignin and nitrogen fractions are given by the IPCC guidelines (Ogle et al., 2019), we use plant-functional type and plant-organ specific parameterization for the natural litterfall. Global distribution of plant-functional types is given by (Schaphoff et al., 2018b) as well as separation of litter into leaf, fine root and wood litter compartments excluding litter biomass burnt in wild fires. Leaf litter parameters are given by Brovkin et al. (2012), fine root to leaf litter lignin ratio by Guo et al. (2021), lignin content of wood litter by Rahman et al. (2013) and nitrogen content scaling factors for leaf to fine roots and leaf to wood litter by von Bloh et al. (2018). Data sources for all considered carbon inputs as well as for 130 lignin and nitrogen ~~content~~ parameterization are listed in Table 1.

**Table 1.** ~~Type~~ type and data sources for carbon inputs and parameterization to different land-use types

land-use types	source of carbon inputs	data source	nitrogen and lignin content
cropland	above-ground residues, below-ground residues, manure	FAOSTAT (2016), Schaphoff et al. (2018b), Weindl et al. (2017)	LG:C generic values according to Table 5.5B, 5.5C from IPCC (Ogle et al., 2019), crop-specific N:C from Bodirsky et al. (2012)
<del>Natural</del> <u>natural</u> vege- tation	annual litterfall	Schaphoff et al. (2018b)	<del>IPCC (Calvo-Buendia et al., 2019) and CENTURY ((NREL, 2000))</del> <u>leaf N and LG concentration from Brovkin et al. (2012), root to leaf litter LG ratio Guo et al. (2021), lignin content of wood litter Rahman et al. (2013) and nitrogen scaling factors for leaf to root and wood litter from von Bloh et al. (2018)</u>

The sub-pool specific decay rates  $k_{sub}$  are influenced by climatic conditions, biophysical and biochemical soil properties as well as management factors that all vary over space ( ~~$i$~~ ) and time (~~and time~~  $t$ ). Following the ~~steady-state method~~ (~~Calvo-Buendia et al., 2019~~) Tier 2 modeling approach (Ogle et al., 2019), we consider temperature ( ~~$temp$~~ ), ~~water~~ (~~water~~  $wat$ ), ~~sand-fraction~~ (~~sand-fraction~~  $sf$ ) and tillage (~~and tillage~~  $till$ ) effects to account for spatial and temporal variation 135 of decay rates. Thus,  $k_{sub}$  rates are given by

$$\begin{aligned}
k_{i,t,active,lu} &= k_{active} \cdot temp_{i,t} \cdot wat_{i,t,lu} \cdot till_{i,t,lu} \cdot sf_i \\
k_{i,t,slow,lu} &= k_{slow} \cdot temp_{i,t} \cdot wat_{i,t,lu} \cdot till_{i,t,lu} \cdot \\
k_{i,t,passive,lu} &= k_{passive} \cdot temp_{i,t} \cdot wat_{i,t,lu}
\end{aligned} \tag{2}$$

For natural vegetation, we assume rainfed and non-tilled conditions, whereas for cropland, we distinguish the effect of different tillage (see Sect. 2.3.5) and irrigation (see Sect. 2.3.4) practices on decay rates. We ~~calculated area-weighted~~ calculate area-weighted means for *till* and *wat* on cropland for each grid cell, using area shares for the different tillage and irrigation ~~practices~~ practices. Data sources as well as used parameters for the different decay drivers for all land-use types are listed in Table 2; equations are displayed by ~~equation~~ equations 5.0B–5.0F in ~~Calvo-Buendia-Ogle et al. (2019)~~ (2019).

**Table 2.** ~~Type~~ type and data sources for carbon inputs to different land-use types

land-use types	type of decay driver	parameter use to represent driver	data source
all	<del>Soil</del> <u>soil</u> quality	<del>Sand</del> <u>sand</u> fraction of the first 0-30 cm	Hengl et al. (2017)
	<del>Mireobial</del> <u>mircobial</u> activity	air temperature	Harris et al. (2020)
	<del>Water-restriction</del> <u>soil moisture</u>	precipitation & potential evapotranspiration	Harris et al. (2020)
cropland	<del>Water-restriction*</del> <u>soil moisture*</u>	irrigation	Sect. 2.3.4
(additionally)	<del>Soil</del> <u>soil</u> disturbance	tillage	Sect. <del>2.3.4</del> <u>2.3.5</u>

### 2.1.2 SOC transfer between land-use types

We calculate *SOC* stocks based on the area shares of land-use types (*lu*) within our grid cells (*i*). If land is converted from one land-use type  $lu = \{crop, natveg\}$  into the other  $!lu = \{natveg, crop\}$ , a respective share of the *SOC* is reallocated within our budget. We do not distinguish between newly converted and existing cropland, but can work with the average carbon content as the relative decay of SOC is proportional to the SOC stock (see 1). We account for land conversion at the beginning of each time step *t* by calculating a preliminary stock  $SOC_{t^*}$  via


$$SOC_{i,t^*,sub,lu} = SOC_{i,t-1,sub,lu} - \frac{SOC_{i,t-1,sub,lu}}{A_{i,t-1,lu}} \cdot AR_{i,t,lu} + \frac{SOC_{i,t-1,sub,!lu}}{A_{i,t-1,!lu}} \cdot AE_{i,t,lu} \quad (3)$$

with  $A_{lu}$  being the land-use type specific areas,  $AR_{lu}$  ~~and~~  $AE_{!lu}$  ~~the area reduction resp. and~~  $AE_{lu}$  the area expansion of the two land-use types. Data sources and methodology on land-use states and changes are described in Sect. 2.3.1.

### 2.1.3 Total SOC stocks and stock changes

*SOC* ~~converge~~ converges towards the calculated steady-state stock  $SOC^{eq}$  for each grid cell *i*, each annual time step *t*, each land-use type *lu* and each sub-pool *sub* like

$$SOC_{i,t,sub,lu} = SOC_{i,t^*,sub,lu} + (SOC_{i,t,sub,lu}^{eq} - SOC_{i,t^*,sub,lu}) \cdot k_{i,t,sub,lu} \cdot 1a. \quad (4)$$



**Figure 1.** Scheme of land-use transition representation. Given an initial land-use pattern (as in this example 2 ha land under natural vegetation and 1 ha of cropland), there are separate *SOC* stocks for natural vegetation and cropland. While in this example we assume *SOC* under natural vegetation to be in steady state, the cropland *SOC* stock approaches its steady state without having reached it yet (a). Upon cropland expansion (in this example half of the natural vegetation is cleared to be used as cropland), *SOC* stocks on cropland increase due to a transfer of land from natural vegetation (b). Explicitly representing newly converted cropland and existing cropland to account for *SOC* dynamics (c) leads to the same weighted mean value as averaging *SOC* stocks (d), due to the linearity of Eq. 4 and cropland-age independent decay rates (see Eq. 2).

155 Note that the decay rates have to be multiplied by one year ( $1a$ ) to form a dimensionless factor. Reformulating this equation, we obtain a mass balance equation as follows

$$SOC_{i,t,sub,lu} = SOC_{i,t^*,sub,lu} - \underbrace{SOC_{i,t^*,sub,lu} \cdot k_{i,t,sub,lu} \cdot 1a}_{\text{outflow}} + \overbrace{SOC_{i,t,sub,lu}^{eq} \cdot k_{i,t,sub,lu} \cdot 1a}^{\text{input (using equation (1))}}. \quad (5)$$

The global  $SOC$  stock for each time step  $t$  can then be calculated via

$$SOC_t = \sum_i \underbrace{SOC_{i,t} - \text{total } SOC \text{ stock within cell}}_{SOC_{i,t} - \text{total } SOC \text{ stock within cell}} \sum_{lu} \underbrace{\sum_{sub} SOC_{i,t,sub,lu}}_{SOC_{i,t} - \text{total } SOC \text{ stock within cell}} \quad (6)$$

160 According to the IPCC guidelines  $SOC$  changes can be expressed as the difference of two consecutive years (see Eq. 5.0A in Calvo-Bueno et al., 2019). This, however, will also include naturally occurring changes due to climatic variation over time. For our study, we defined the absolute and relative  $SOC$  changes in relation to a potential natural state  $SOC^{pnv}$  under the same climatic conditions in grid cell  $i$  at time  $t$ , that is based on the natural vegetation  $SOC$  calculations as defined above without accounting for land conversion from cropland at any time. The absolute changes  $\Delta SOC$  and relative changes  $F^{SCF}$  are thus given by

$$\Delta SOC_{i,t} = SOC_{i,t} - SOC_{i,t}^{pnv} \quad \text{and} \quad F_{i,t}^{SCF} = \frac{SOC_{i,t}}{SOC_{i,t}^{pnv}}. \quad (7)$$

Note that the absolute changes  $\Delta SOC$  can be also interpreted as the SOC debt (Sanderman et al., 2017) due to human cropping activities; whereas relative changes  $F^{SCF}$  can be considered stock change factors as defined within the IPCC guidelines of 2006 (Eggleston et al., 2006). Moreover,  $\Delta SOC$  is equivalent to the negated cumulative SOC component of human land-use change emissions (Pugh et al., 2015).

#### 2.1.4 Initialization of SOC pools

175 ~~To initialize all SOC~~ The initialization of  $SOC$  pools is very important and has to include the proper accounting for the land-use history, as many  $CO_2$  emissions from agricultural soils are caused by historical land-use change (LUC) and the slow decline of  $SOC$  under crop cultivation, before it reaches a new equilibrium. We initialize our  $SOC$  sub-pools ~~we assume that cropland and natural vegetation are in a~~ using a three-step approach, since input data availability is limited for climate and litter estimates (starting only in 1901) as well as for agricultural management data (starting only in 1965):

180 Firstly, in order to account for the impacts of legacy fluxes from land-use type-specific steady-state for the initialization year 1901. While land-use, climate and litter input information are available from changes long before the time horizon of interest, we consider land-use change from 1510 onwards. In 1510, we assume all  $SOC$  pools to be in natural steady-state, implying that all land-use change prior to that time occurs in 1510. We assume that by 1901, all cropland converted in 1510 has reached its new steady state, so that it is not necessary to explicitly account for even older land conversion. Model inputs for 1901–1930 for climate and natural vegetation litterfall are repeated for 1510–1900 to mimic constant climate conditions for this first initializing period. Similarly, agricultural management data ~~on residue and manure inputs~~ are held constant at the level of 1965 until 1965. After a spin-up period of 64 years (1901–1965) with This approach follows others studies looking on effects of land-use change and management (e.g. Schaphoff et al., 2018a; Herzfeld et al., 2021).



Secondly, with the availability of transient climate data after 1901, we account not only for land-use change, but also for historical climate change and consequently natural litter inputs to the soil from 1901 to 1965 still considering constant agricultural input data, which are not available prior to 1965.

190 Thirdly, we run the model for ~~additional~~ 10 years ~~with historic input data~~ from 1965 to 1975 with historical dynamic data on agricultural management and start analyzing results from 1975 onwards. ~~Irrigation areas are part of the land-use data set and therefore dynamic from onward.~~ This is in line with the IPCC guidelines vol. 4 method suggestion to have a 5-20 year spin-up period (Ogle et al., 2019).

195 With transient climate considered after 1901 ~~on, whereas data~~, decay rates  $k_{sub}$  become dynamic in time. As the decay rates are also affected by irrigation and tillage (see Sect. 2.1.1), we also account for transient changes in irrigated areas after 1901. Data on no-tillage ~~area is~~ practices are only available after ~~1974~~ 1974 and we assume conventional tillage on all cropland prior to 1975.

## 2.2 SOC stocks and stock changes following Tier 1

Additionally to the ~~steady-state method (Calvo Buendia et al., 2019)~~ Tier 2 modeling approach (Ogle et al., 2019) and the detailed analysis of management data coming with it, ~~SOC~~ SOC changes can be estimated using the IPCC Tier 1 approach of IPCC guidelines (Eggleston et al., 2006; Calvo Buendia et al., 2019). Here, stocks are calculated via stock change factors ( $F^{SCF}$ ) given by the IPCC for the topsoil (0-30 cm) and based on observational data. Estimates of  $F^{SCF}$  are differentiated by ~~different~~ crops, management and input systems (here summarized under  $m$ ) reflecting different dynamics under changed in- and outflows without explicitly tracking these flows. Moreover, estimates of  $F^{SCF}$  vary for different climatic zones ( $c$ ) specified by the IPCC (see Fig. A1). The actual SOC stocks are thus calculated based on a given reference stock  $SOC^{ref}$  by

$$205 \quad SOC_{i,t} = \sum_{c,m} T_{c,i} \cdot SOC_{i,t}^{ref} \cdot F_{c,m}^{SCF} \quad (8)$$

with  $T_{c,i}$  being the translation matrix for grid cells  $i$  into corresponding climate zones  $c$ . For this analysis, we use the default  $F^{SCF}$  from the Tier 1 method of Eggleston et al. (2006) and Calvo Buendia et al. ~~(Calvo Buendia et al., 2019)~~ (2019) as a comparison and consistency check for our more detailed Tier 2 steady-state approach.

## 2.3 Agricultural management data at 0.5 degree resolution

210 We compile country-specific FAO production and cropland statistics (FAOSTAT, 2016) to a harmonized and consistent data set. The data is prepared in 5-year time steps from 1965 to 2010, which restricts our analysis to the time span from 1975 to 2010 (after a spin-up phase from ~~1901-1974~~ 1510–1974). For all the following data, if not declared differently, we interpolate values linearly between the time steps and keep them constant before 1965.

### 2.3.1 Land use and land-use change

Land-use patterns are based on the Land-Use Harmonization 2 (Hurt et al., 2020) data set ([short LUH2](#)), which we sum up from quarter-degree to half-degree resolution. We disaggregate the physical area ([given as total land area in million ha](#)) of the five different cropland subcategories (c3ann: [C3 annual crops](#), c3per: [C3 perennial crops](#), c4ann: [C4 annual crops](#), c4per: [C4 perennial crops](#), c3nfx: [C3 nitrogen-fixing crops](#)) of LUH2 into our 17 crop groups ([see FAO2LUH2MAG\\_croptypes.csv in Bodirsky et al., 2020a](#)) ([see Table “FAO2LUH2MAG\\_croptypes.csv” in Karstens, 2020a](#)), applying the relative shares for each grid cell based on the country- and year-specific area harvested shares of FAOSTAT data (FAOSTAT, 2016). By calculating country-specific multi-cropping factors [MCF](#) using FAOSTAT data, we are able to compute crop-group specific area harvested on grid cell level. Land-use transitions are calculated as net area differences of the land-use data at half-degree resolution, considering no split up into crop-group specific areas but only total cropland and natural vegetation areas.

### 2.3.2 Crop and crop residues production

Crop production patterns are compiled crop group specific using half-degree yield data from LPJmL (Schaphoff et al., 2018b) as well as half-degree cropland patterns (see Sect. 2.3.1). We calibrate cellular yields with a country-level calibration factor for each crop group to meet [historie-historical](#) FAOSTAT production (FAOSTAT, 2016). [By using physical cropland areas in combination with harvested areas, we account for multiple cropping systems as well as for fallow land.](#) [Note](#)

Crop residue production and management is based on a revised methodology of Bodirsky et al. (2012) and key aspects are explained [here given its central role for as they play a central role in](#) soil carbon modeling. Starting from [harvested-crop production \(CP\)](#) estimates and their respective harvested crop area ([crop yield estimates  \$Y\$  and respective physical crop area  \$CA\$](#) ), we estimate [total](#) above-ground ([AGR](#)) and below-ground ([BGR](#)) [residual-biomass-residue biomass \(in tonnes\)](#) using crop group ([cg](#)) [specific harvest index values \(HI\) and root:shoot ratios \(RS\)](#) [specific ratios for above-ground residues to harvested biomass  \$r\_{cg}^{ag,prod}\$  in  \$\(tDM\ ha^{-1}\)\(tDM\ ha^{-1}\)^{-1}\$  above-ground residues to harvested area  \$r\_{cg}^{ag,area}\$  in  \$tDM\ ha^{-1}\$  and below-ground residues to above-ground biomass  \$r\_{cg}^{bg}\$  in  \$tDM\ tDM^{-1}\$](#)  as follows

$$\begin{aligned} AGR_{i,t,cg} &= CA_{i,t,cg} \cdot (Y_{i,t,cg} \cdot r_{cg}^{ag,prod} + MCF_{i,t} \cdot r_{cg}^{ag,area}) \quad \text{and} \\ BGR_{i,t,cg} &= (CA_{i,t,cg} \cdot Y_{i,t,cg} + AGR_{i,t,cg}) \cdot r_{cg}^{bg} \end{aligned} \quad (9)$$

Following the IPCC guidelines, we split the [harvest index into a production \( \$HI^{prod}\$ \) and an area dependent \( \$HI^{area}\$ \) fraction](#) (Eggleston et al., 2006) [above-ground residue calculations into a yield-dependent slope \( \$r\_{cg}^{ag,prod}\$ \) and a positive intercept \( \$r\_{cg}^{ag,area}\$ \) fraction](#) (Hergoualc’h, Kristell et al., 2019). Residues biomass therefore increases under-proportionally with rising yields, reflecting a shifting harvest index of higher-yielding breeds. Deviating from Bodirsky et al. (2012) we use harvested instead of physical crop area ([denoted in Eq. \(9\) by MCF described in Sect. 2.3.1](#)) to account for increased residue biomass due to multiple cropping ([multiple harvests with each lower yields](#)) and decreased residue amounts due to fallow land. We assume that all  $BGR$  are left in the soil, whereas  $AGR$  can be burned or harvested for other purposes such as feeding animals (Weindl et al., 2017), fuel or for material use.

245 A country-specific fixed share of the *AGR* is assumed to be burned on field depending on the per-capita income of the country. Following Smil (1999b) we assume a burn share of ~~25%~~ 25% for low-income countries according to World Bank definitions ( ~~$< 1000 \frac{USD}{yr\ cap}$~~   $< 10000 \frac{USD}{yr\ cap}$   $< 10000 \text{ USD yr}^{-1} \text{ cap}^{-1}$ ), 15% for high-income ( ~~$> 10000 \frac{USD}{yr\ cap}$~~   $> 10000 \frac{USD}{yr\ cap}$   $> 10000 \text{ USD yr}^{-1} \text{ cap}^{-1}$ ) and linearly interpolate shares for all middle-income countries depending on their per-capita income for the periods before 1995. After 1995 we estimate a linear decline of burn shares to 10% for low-income countries and 0% for high-income countries till 2025  
 250 to account for recent increases in air pollution regulation. The estimated trends show good agreement with fire-satellite-image derived estimates by the Global Fire Database (van der Werf et al., 2017). Depending on the crop group, 80–90% of the carbon in the crop residues burned in the fields is lost within the combustion process (Eggleston et al., 2006).

From our 17 crop groups, we compile four residue groups (straw, high- and low-lignin residues, residues without dual use), of which the first three are taken away from the field for other purposes (see mappingCrop2Residue.csv in Bodirsky et al. (2020a)).  
 255 Residue feed demand for five different livestock groups is based on country-specific feed baskets (see Weindl et al., 2017), that differentiate between the residue groups and take available *AGR* biomass as well as livestock productivity into account. We estimate a material-use share for the straw-residue group of 5% and a fuel-share of 10% for all used residue groups in low-income countries. For high-income countries, no withdrawal for material or fuel use is assumed, and use shares of middle-income countries are linearly interpolated based on per-capita income, following the same rationale as for the share  
 260 of burnt residues described above. The remaining *AGR* as well as all *BGR* are expected to be left on the field. We limit high residue return rates to at most  ~~$10 \text{ tC ha}^{-1}$~~   $10 \text{ tC ha}^{-1}$  in order to correct for outliers.

To transform dry matter estimates into carbon and nitrogen, we compiled crop-group and plant-part specific carbon ~~resp.~~ and nitrogen to dry matter (~~e/dm, n/dm~~) ratios (see Table A1).

### 2.3.3 Livestock distribution and manure excretion

265 Manure especially from ruminants is often excreted at pastures and rangelands, but due to the intensification of livestock systems ~~at the present day~~ a lot of the manure has to be stored and can be applied on ~~croplands~~ cropland. We assume that manure ~~application happens at its excretion place~~ is applied in close proximity to its excretion, so that the ~~livestock distribution~~ distribution of livestock is the driving factor ~~of for~~ the spatial pattern of ~~manuring~~ manure application.

To disaggregate country level FAOSTAT livestock production data to half-degree resolution, we use the following rule-based  
 270 assumptions, drawing from the approach of Robinson et al. (2014) and applying feed basket assumptions based on a revised methodology from Weindl et al. (2017). We differentiate between ruminant and monogastric systems, as well as extensive and intensive systems. Due to ~~great~~ high feed demand of ruminants, we assume that ruminant livestock is located where the production of feed occurs to minimize transport of feed. We distinguish between grazed pasture, which is converted into livestock products in extensive systems, and primary-crop feed stuff, which we consider to be consumed in intensive systems.  
 275 For poultry, egg and monogastric meat production we use the per-capita income of the country to distinguish between intensive and extensive production systems. For low-income countries, we assume only extensive production systems. We locate them according to the share of built-up areas based on the assumption that these animals are held in subsistence or small-holder farming systems with a high labor-per-animal ratio. Intensive production associated with high-income countries, is distributed

within a country using the share of primary-crop production, assuming that feed availability is the most determining factor for  
280 livestock location. For middle-income countries we split the livestock production into extensive and intensive systems based  
on the per-capita income.

Manure production and management is based on a revised methodology of Bodirsky et al. (2012) and is presented here due  
to its central role in soil carbon modeling. Based on the gridded livestock distribution we calculate spatially explicit excretion  
by estimating the nitrogen balance of the livestock systems on the basis of comprehensive livestock feed baskets (Weindl et al.,  
285 2017), assuming that all nitrogen in protein feed intake, minus the nitrogen in the slaughter mass, is excreted. Carbon in excreted  
manure is estimated by applying fixed C:N ratios, which range from 10 for poultry up to 19 for beef cattle (for full detail see  
Calvo Buendia et al. (2019)(2019). Depending on the feed system we assume manure to be handled in four different ways: All  
manure originated from pasture feed intake is excreted directly on pastures and rangelands (pasture grazing), deducting manure  
collected as fuel. Whereas for low-income countries, we adopt a share of 25% of crop residues in feed intake directly consumed  
290 and excreted on crop fields (stubble grazing), we do not consider any stubble grazing in high-income countries; middle-income  
countries see linearly interpolated shares depending on their per-capita income. For all other feed items, we assume the manure  
to be stored in animal waste management systems associated with livestock housing. To estimate the carbon actually returned  
to the soil, we account for carbon losses during storage, where return shares depend on different animal waste management and  
grazing systems. Whereas we assume no losses for pasture and stubble grazing, we consider that the manure collected as fuel  
295 is not returned to the fields. For manure stored in different animal waste management systems we compiled carbon loss rates  
(see calcClossConfinement.R in Bodirsky et al. (2020a) for more details) depending on the different systems and the associated  
nitrogen loss rates as specified in Bodirsky et al. (2012). We limit high application shares at  $10\text{tC ha}^{-1}$  to  $10\text{tC ha}^{-1}$  to correct  
for outliers, that can occur due to inconsistencies between FAO production and 0.5 degree land-use data.

#### 2.3.4 Irrigation

300 The LUH2v2 (Hurt et al., 2020) data set provides irrigated fractions for ~~their~~its cropland subcategories. We sum up irrigation  
area shares for all crop groups within a grid cell, and calculate the water effect coefficient *wat* on decay rates using these  
shares to compute the weighted mean between rainfed and irrigated *wat* factors. As a result *wat* is the same for all crop  
groups within a grid cell. Furthermore, we suppose the irrigation effect to be present for all 12 months of a year ~~in a grid cell~~  
~~including irrigated areas~~, since we do not have consistent crop group specific growing periods available. This will lead to an  
305 overestimation of the irrigation effect. We expect, however, water limitations to be a minor problem during the off-season in  
temperature limited cropping regions, causing our assumption to not dramatically overestimate the moisture effects. In tropical,  
water-limited cropping areas, irrigated growing periods might even span over the whole year.

#### 2.3.5 Tillage

In order to derive a spatial distribution of the three different tillage types specified by the IPCC — full tillage, reduced tillage  
310 and no tillage —, we assume that all natural land and pastures are not tilled, whereas annual crops are under full and perennials  
under reduced tillage per default. Furthermore, we assume no tillage in cropland cells specified as no tillage cell based on the

~~historie-historical~~ global gridded tillage data set from Porwollik et al. (2018)(2019). This data set is extended to the period of 1975–2010 by combining country-level data on areas under conservation agriculture from FAO (2016) and half-degree resolution physical crop areas from Hurtt et al. (2020), applying the methodology of Prowollik et al. (2018) (2019) to identify potential no-tillage grid cells.

~~To highlight the impact of changing management effects and to assess the sensitivity of the model towards different initialization and parameter choices, we perform a set of scenario runs. In the following sections we outline name and idea of these scenarios (for technical implementation see Karstens and Dietrich, 2020).~~

## 2.4 Management scenario definitions

To single out the impact of tillage practices, residue and manure inputs, we defined scenario with constant values for these three drivers: In the *constTillage* scenario the adoption of no-tillage practices are neglected (adoption starts in 1974 according to the available data set). The *constResidues* and the *constManure* scenario assume constant input rates from residues resp. manure (in  $\text{tha}^{-1}$ ) at the level of 1975 ~~onwards~~onward. Within the *constResidue* scenario at different effects overlay each other: yields and with them residue biomass increase due to productivity gains; rates of residue left or returned to fields are raising; and shifts of cropping pattern change the amount of residue biomass due to crop-group specific harvest index values. The *constManagement* combines all three scenarios *constTillage*, *constResidues* and *constManure*.

~~As outlined in Sect. 2.1.4 we assume landuse-type specific steady-state SOC stocks for the start year 1901 followed by a long spin-up period of 74 years (*Initial-lu*). As some SOC compartments decay over very long timescales, the initialisation setting might strongly effect the overall outcome of SOC stocks and changes. Thus we conduct counterfactual scenario *Initial-natveg*, initializing the start year with the steady-state SOC under potential natural vegetation for all land-use types. We additionally combined the *Initial-natveg* scenario with the *constManagement* scenario.~~

~~Due to missing information about the litterfall composition, we assume woody potential natural vegetation for the lignin-to-carbon (short LG:C) and nitrogen-to-carbon (short N:C) ratios for all land under natural vegetation. This is however a strong simplification. Therefore, we test the sensitivity of the model for other natural vegetation parameterizations. Perennial grasses are given by the IPCC guidelines (Calvo Buendia et al., 2019), but for woody biomass we used the CENTURY configuration file (NREL, 2000) for trees as parameterization were not given. Whereas LG:C ratios ranges between 0.09 — 0.35 with an average of 0.24 (default value) for different tree compartments and different tree types, N:C ratios range between 0.001 — 0.05 with an average of 0.015 (default value). We decided to average over all tree compartments and tree types equally for the default values, since we have no data available on tree type distribution and tree compartment composition. We test however a range of different plant types as specified in Table ??.~~

~~Scenarios for sensitivity analysis on litter parameterization Scenario name plant type N:C LG:C *LitterPNV-Mean* mix over all tree types 0.015 0.25 *LitterPNV-Grass* perennial grasses 0.029 0.11 *LitterPNV-AASS* Arctic-alpine subshrub 0.022 0.22 *LitterPNV-BLEAF* Temperate broadleaf evergreen 0.013 0.27 *LitterPNV-BRCF* Boreal coniferous forest 0.013 0.17 *LitterPNV-MTCF* Maritime coniferous forest 0.0094 0.22 *LitterPNV-TREF* Tropical evergreen forest 0.016 0.30~~

Detailed results for the spatially explicit global SOC budget including intermediate results on input data as well as SOC stock results for all scenario runs can be found in Karstens (2020a). In the following, the most important results (~~see Karstens, 2020b) for post-processing script~~) (see Karstens, 2020b, for post-processing script) are summarized.


### 3.1 SOC distribution and depletion

350 ~~In~~ The global SOC debt has increased by about 14% in the period between 1975 and 2010 from 34.6 to 39.6 GtC (Fig. 2(a)~~we provide the first~~). This corresponds to an average loss rate of  $0.14 \text{ GtC yr}^{-1}$  in comparison to a hypothetical potential natural vegetation (PNV) state. Considering our estimate of the global SOC stock of around 705 GtC in the upper 30 cm in 1975, global SOC decreased by 0.2 per 1000 per year for the period between 1975–2010. The speed of this SOC loss has decreased towards the end of the modeling period.

355 In Fig. 2(b) we provide a world map of SOC stocks ~~estimates~~ for the first ~~30 cm on croplands considering historic management data at the global scale~~ 30 cm on cropland considering historical management data for the year 2010. Values ~~ranging between well over  $100 \text{ t ha}^{-1}$  range between over  $100 \text{ t ha}^{-1}$  in northern temperate croplands~~ ~~cropland~~ to less than  ~~$5 \text{ t ha}^{-1}$   $5 \text{ t ha}^{-1}$  for arid and semiarid croplands~~ ~~cropland~~. Our spatially explicit results show hotspots of SOC losses and gains compared to SOC under ~~potential natural vegetation~~ PNV in two complementary ways: 1. Absolute SOC changes  $\Delta \text{SOC}$  (see Fig. 2(bc))  
360 indicate areas with high importance for the global ~~SOC losses. They might SOC loss. They can~~ be driven by large relative changes (e.g. in Central Africa) or by a high natural stock, from which even small relative deviations could lead to substantial absolute losses (e.g. North-East Asia). ~~In contrast, large net gains of SOC occur primarily in developed countries of the Global North according to our results.~~ 2. Relative SOC changes measured as stock changes factors  $F^{\text{SCF}}$  (see Fig. 2(ed)) are a helpful metric to analyze the impact of human cropping activities. They indicate areas with large differences in carbon inflows or  
365 SOC decay compared to natural vegetation, that may hold potential to be overcome ~~due to by~~ improved agricultural practices. Large parts of tropical ~~croplands~~ ~~cropland~~ seem to suffer from ~~low stock changes factors, meaning high relative SOC losses and maybe strongly reduced relative stocks,~~ indicating SOC degradation. Conversely, ~~not only temperate croplands of Central Europe, Japan and western areas of the USA have high stock change factors, but irrigated croplands~~ ~~irrigated cropland~~ at the border to dry, unsuitable areas worldwide ~~as well~~ shows a strong relative increase in SOC stocks.

370 ~~The global SOC debt has decreased by about 15% in the period between 1975 and 2010 to 22 GtC (Fig. 2(d)). This corresponds to a sequestration rate of  $0.11 \text{ GtC yr}^{-1}$ . Considering our estimate of the global SOC stock of around 660 GtC in 1975, global SOC increased by 0.2 per 1000 per year for the period between 1975–2010.~~

The spatial distribution of the total  $\Delta \text{SOC}$  summed over all land-use types (Fig. 3(a)) and its change from 1975 to 2010 (Fig. 3(b)) reveals areas of SOC debt decline and increase. Regions with large cropland expansion (e.g. Brazil, Southeast Asia,  
375 Canada) continue to lose SOC, whereas regions with cropland reduction (and thus SOC restoration) or with accumulating cropland SOC can be found e.g. in highly productive areas of Europe and Central USA.




**Figure 2.** Global SOC stocks and SOC stock changes on cropland for the first 30 cm of the soil profile considering historical management data. Panel (a) ~~:-Distribution-~~ shows global  $\Delta SOC$  between historical land use and potential natural vegetation (PNV). The distribution of total global SOC stocks for the first 30 cm on cropland ~~:-Carbon stocks are large for the year 2010 is depicted in high-yielding areas-~~ panel (b) ~~)+(e)-~~. Absolute (bc) and relative (ed) SOC stocks changes for the year 2010 are compared to a potential natural state identify different hotspots of SOC dynamics. Whereas absolute losses  $\Delta SOC$  are often highest in temperate dry regions, relative losses  $F^{SCF}$  are often larger in tropical moist areas. (d):  $\Delta SOC$  between SOC under historic land use and potential natural vegetation is decreasing over time, meaning net SOC gains on global croplands over the the period 1975–2010.

### 3.2 Carbon flows in the agricultural system

C is sequestered from the atmosphere via plant growth and allocated to ~~three~~ different plant parts (~~harvest~~, which we aggregate to three pools (harvested organ, above- and below-ground residues). Whereas harvested organs as well as above ground-residues are taken (partially) from the field to be used for other purposes, below-ground residues (~~785 MtC~~ 729 MtC in 2010)






**Figure 3.** Global total  $\Delta SOC$  and  $\Delta SOC$  change for the first 30 cm of the soil profile. Panel (a) shows global  $\Delta SOC$  as the difference between  $SOC$  under historical land use and  $SOC$  under potential natural vegetation (PNV) in the year 2010 summed over all land-use types. Computing the difference between the  $\Delta SOC$  estimate for 2010 and for 1975 (b) depicts areas of soil depletion (SOC debt increase, red) and net-sequestration (SOC debt decline, blue).

are directly returned to the field. We ~~split up usage for crop biomass~~ divide crop biomass usage into feed usage and aggregate all other usage types (e.g. ~~like~~ food, bioenergy and material) into a human demand category. Livestock feed demand for crop organ harvest and above-ground residues of ~~1141 MtC is almost 1136 MtC is roughly~~ equal to the human demand of ~~1144 MtC~~ 1129 MtC. Whereas large parts of feed intake are ~~reeyeled~~ returned to the soils via manure (C input from manure at ~~384 MtC~~ 384 MtC), we assume the carbon demanded from humans (ending up as e.g. compost, night soil and sewage) is not ~~reeyeled~~ returned to soils. Besides manure C and below-ground residues, above-ground residues form the largest C input to the soil with ~~1200 MtC~~ 1350 MtC returned to the ~~field in fields in the year~~ 2010. However, around 60% of this organic C decomposes before ~~its it is~~ integrated into soils ~~at the the litter-soil barrier~~. Due to the different ~~C composition~~, proportional composition






**Figure 4.** Global carbon flows within the agricultural system for the year 2010 (in MtC). Carbon is first photosynthesised by crop plants and then used depending on the plant part by humans for feed of livestock feed and various other usages subsumed under human demand. After accounting for losses within the agricultural system, there are three major C inputs are applied to croplands cropland SOC: manure, above- and below-ground residues. Large parts of C, however, get are mineralized on the field before entering the soil. Additionally, C is transferred to and from the global agricultural soils soil stock via land-use change to between cropland and from natural vegetation. Finally, SOC gets is mineralized and flows back into to the atmosphere.

of organic C, proportionally more C enters the slow pool from manure compared to than from crop residue. According to our

390 model results, land-use change dynamics led to a C transfer from ~~cropland to natural vegetation of 58 MtC~~ natural vegetation  
to cropland of 257 MtC in 2010. ~~4764 MtC sequestered~~ The agricultural system receives 4585 MtC assimilated by crop plants  
~~face 3897 MtC released within the agricultural system and releases 3554 MtC mostly through respiration.~~ Accounting for SOC  
transfer, ~~SOC increase under and decomposition, the net SOC decrease of global~~ cropland is around 809 MtC. 33 MtC for the  
year 2010.

395 **3.3 Agricultural management effects on SOC debt**



**Figure 5.** (a) Global  $\Delta SOC$  in GtC GtC for different management scenarios. The stylized scenarios ~~deviate deviate~~ from ~~historie~~ historical agricultural management patterns (histManagement) by holding effects of carbon inflows from residues (constResidues) ~~or~~ manure (constManure) constant at the 1975 level, or neglecting adoption of no-tillage practices over time (constTillage). ~~ConstManagement~~ The scenario constManagement combines all three modifications. Note that  $\Delta SOC$  is defined as the difference of SOC under land-use compared to a hypothetical natural vegetation state. Figure Panel (b) shows the carbon inflows from crop residue and manure, underlining the strong impact of residues for SOC stock and SOC stock changes.

We analyze the relative impact of ~~individual management aspects~~ different management practices by comparing the actual ~~historie~~ historical management scenario with counterfactual scenarios, where individual management ~~aspects~~ practices (residues in constResidues, manure in constManure, tillage practices in constTillage, all three in constManagement) are kept static at the 1975 values (Figure 5(a)). ~~Without changes in management regimes, the global ΔSOC on cropland would be~~

400 ~~increasing at a rate of  $0.1 \text{ GtC yr}^{-1}$ . As shown by the `constResidue` scenario difference between the `constResidue` scenario and the other counterfactuals, changes in residue return rates dominate the management effects. Without the `historie-increase in-residue-returning` historical increase in C inputs from residues to agricultural soils, the global  $\Delta \text{SOC}$  would still increase decrease to  $41.7 \text{ GtC}$  at a rate of  $0.06 \text{ GtC yr}^{-1}$   $0.20 \text{ GtC yr}^{-1}$  — a 35% increase compared to  $0.14 \text{ GtC yr}^{-1}$  for the `histManagement` estimates. Both the `constManure` and `constTillage` scenarios show only small deviations from the `historie-values (sequestration rate of  $0.09 \text{ GtC yr}^{-1}$  for both)` historical agricultural management values with  $0.15 \text{ GtC yr}^{-1}$ . The effect of no-tillage has been particularly strong since the 1990s. The strong impact of almost doubling C inputs from crop residue biomass only becomes discernible from 2000 onwards. The large contribution of residues relative to manure also becomes visible when considering the annual C inputs of residues and manure to soils over a period of 35 years on agricultural SOC stocks is shown in (Fig. 5(b)).~~

Our sensitivity analysis shows that the management impact is robust to the initialization of SOC stocks at the beginning of the spin-up phase (Fig. ??). The `Initial-natveg` scenario initializes the start year with the steady-state SOC under potential natural vegetation for all land-use types, compared to the default assumption (`Initial-lu`), which assumes landuse-type specific steady-state SOC stocks in 1901. The  $\Delta \text{SOC}$  estimation almost halves (from  $\sim 26 \text{ GtC}$  to  $\sim 14 \text{ GtC}$ ), as does the decrease in  $\Delta \text{SOC}$  for the period of 1975–2010 (from  $\sim 4 \text{ GtC}$  to  $\sim 2.5 \text{ GtC}$ ) for the `Initial-natveg` scenario. Global SOC stocks range between 645–700 GtC for different parameterization of the natural litterfall, whereas  $\Delta \text{SOC}$  vary 19–24 GtC for the year 2010 (see Fig. ??). The increase of  $\Delta \text{SOC}$  during the period of 1975–2010 is independent of the parameterization choice at  $\sim 4 \text{ GtC}$ .

### 3.4 Model evaluation

To evaluate our model results against reference data in five steps: (1) we compare our stock change factors (see Sect. 2.2) to IPCC default assumptions (Lasco et al., 2006; Ogle et al., 2019); (2) we compare our global (and climate-zone specific) total SOC stocks to other literature estimates; (3) we compare our results to point measurements. To evaluate the representation of our natural SOC stocks (4) we correlated LPJmL4 SOC stocks for PNV with our natural state SOC results on grid level; and (5) we do a similar correlation analysis for our modeled actual SOC stocks in comparison to the results of SoilGrids 2.0 (Poggio et al., 2021), which accounts for actual land use too.

#### 3.4.1 Stock change factors compared to IPCC assumptions

To evaluate our modeled SOC stocks and stock changes under agricultural management, we compare our results to the default IPCC stock change factors  $F^{\text{SCF}}$  of 2006 (Lasco et al., 2006) and their refinements in 2019 (Ogle et al., 2019). Both estimates are based on measurement data for cropland (see Table 3). To allow for comparison, we aggregate our stock change factors weighted by grid-level  $F^{\text{SCF}}$  cropland area to derive average factors for the four IPCC climate zones (Fig. A1).


Stock change factors for temperate climate zones of this study are lower than the default values of the IPCC. For the tropical regions the IPCC factors changed notably from the guidelines in 2006 (Lasco et al., 2006) to the update in 2019 (Ogle et al., 2019). Our results are in good agreement with the 2006 IPCC factors. Modeled  $F^{\text{SCF}}$  have increased or stayed constant for all climate zones over time (1975–2010).

**Table 3.**  $F^{SCF}$  in comparison to IPCC Tier 1 default factors from the guidelines in 2006 (Lasco et al., 2006) and the update in 2019 (Ogle et al., 2019).

	Source	Input	Year	tropical moist	tropical dry	temperate dry	temperate moist
1	IPCC2006	low	invariant	0.44	0.55–0.61	0.74	0.66
2	IPCC2006	medium	invariant	0.48	0.58–0.64	0.80	0.69
3	IPCC2006	high	invariant	0.53	0.60–0.67	0.83	0.77
4	IPCC2019	low	invariant	0.76	0.87	0.70–0.71	0.66–0.67
5	IPCC2019	medium	invariant	0.83	0.92	0.76–0.77	0.69–0.70
6	IPCC2019	high	invariant	0.92	0.96	0.79–0.80	0.77–0.78
7	This Study	hist	1975	0.48	0.59	0.64	0.59
8	This Study	hist	2010	0.5	0.64	0.64	0.59

### 3.4.2 Global SOC stocks comparison

We compare our global SOC stocks with a wide range of global SOC stock estimates for the first 30 cm of the soil profile, using data from WISE (Batjes, 2016), SoilGrids (Hengl et al., 2017), GSOC (FAO, 2018), LPJmL4 (Schaphoff et al., 2018a), SoilGrids 2.0 (Poggio et al., 2021), and SOCDebtPaper (Sanderman et al., 2017) in Fig. 6.




**Figure 6.** Modeled as well as observation-based estimates for global SOC stock in GtC for the first 30 cm of soil aggregated over all land area. The comparison against observation-based data (SoilGrids, SoilGrids 2.0, GSOC and WISE) is supplemented by modeled data from LPJmL4 (Schaphoff et al., 2018a) and estimates from (Sanderman et al., 2017). We show values of this study for the year 2010 accounting for the historical land-use dynamics as well as for an hypothetical PNV.

The global estimates of the total SOC stock of the upper 30 cm from this study are in the middle of the wide range of other modeled or observation-based estimates. SoilGrids (Hengl et al., 2017) especially stands out with its high estimate, whereas

SoilGrids 2.0 (Poggio et al., 2021) marks the lower end. Regional results (Fig. A2) show that our estimates are well within the range of other estimates for most regions, but at the lower end for tropical moist and tropical wet areas

440 **3.4.3 Point-based evaluation**

We correlate our SOC results for natural vegetation and cropland in 2010 with literature values from point measurements (for data base see appendix of (Sanderman et al., 2017)).




**Figure 7.** Correlation between modeled and measured *SOC* stocks. Given the wide span between minimum and maximum measured *SOC* stocks within in a given cell, we correlated median values with our modeled results. Both cropland and areas with natural vegetation tend to be lower in our results than in the point measurements.

**3.4.4 Natural SOC stock comparison with LPJmL4**

Estimates of SOC stocks under natural vegetation influence our modeled results for cropland, which has been converted from natural vegetation at some point in time. We therefore also compare our modeled results for SOC under natural vegetation (derived using litterfall of LPJmL4) against estimates of SOC by LPJmL4 for a PNV simulation. Both models are driven by the same climate conditions and the same natural litterfall and just differ in the representation of SOC and litter dynamics. With

445

our focus on cropland SOC dynamics, we compare only cells with more than 1000 ha of cropland (capturing 99.9% of global cropland area).




**Figure 8.** Correlation between modeled SOC stocks of LPJmL4 and this study for an hypothetical potential natural state (PNV) for the year 2010. The grey lines indicate the 1:1 line.

Spatial correlations of PNV SOC stock values are high (global  $R^2 = 0.81$ ), especially for dry climate zones (Fig. 8). For temperate and tropical moist areas estimates of this study tend to be a bit lower compared to LPJmL4 results.


### 3.4.5 Actual SOC stock comparison with SoilGrids 2.0

SoilGrids 2.0 (Poggio et al., 2021) is a digital soil mapping approach that uses over 240 000 soil profile observations to produce high resolution soil maps including SOC stocks and estimates of their uncertainties. To evaluate the performance of our model at the global scale, we correlate SoilGrids 2.0 SOC stock values, which were aggregated to 0.5 degree resolution, to our

estimates for the year 2010 in Fig. 9. To focus our comparison on cropland areas, we mask out grid cells with less than 1000 ha of cropland. Spatial correlation is moderate for tropical climate zones, whereas it is low for temperate moist areas. In tropical dry and temperate dry areas, we simulate also very low SOC values (below  $10\text{tC ha}^{-1}$ ), which is not found in SoilGrids 2.0 whereas our modeled SOC stocks can be substantially higher in temperate moist areas than reported by SoilGrids 2.0. Additionally, we use the uncertainty estimates from SoilGrids 2.0 in Fig. 10 to identify areas, where our modeled SOC stocks that are below the 5th or above the 95th percentile of the SoilGrids 2.0 data. For the vast majority of grid cells our model results are between the 5th and 95th percentile of SoilGrids 2.0 estimates. We underestimate SOC stocks especially in dry areas (e.g. close to the Sahara). Overestimated stocks are often situated in mountainous regions.



**Figure 9.** Correlation between modeled SOC stocks of this study and projected values from SoilGrids 2.0.



**Figure 10.** [Global map on SOC results compared to uncertainty estimates from SoilGrids 2.0.](#)

#### 4 Discussion

465 This study shows that spatially explicit and time-variant historic agricultural management data considerably alter estimates of the states and trends of SOC compared to the often-used constant management assumptions. This result remains robust to variations of central model parameters and variations in the initialization of SOC stocks. While land cover change has depleted SOC stocks and increased the SOC debt, our analysis points out that the high increases in agricultural productivity may have even led to a net reduction of the SOC debt since 1975.

470 The evaluation of SOC stock gains and losses is complex and has several dimensions as climatic and anthropogenic effects overlap. If defining the SOC debt of 1975 as the baseline, and measuring land-use emissions on croplands as the difference between a potential natural state and the state under human interventions (see Pugh et al., 2015), global croplands have acted as a carbon sink since 1975 according to our study. However, annual C sequestration rates of 0.2 per 1000 are well below the promoted 4 per 1000 (Minasny et al., 2017), indicating that productivity gains on historic levels alone are not enough to  
 475 meet ambitious climate targets. According to Sanderman et al. (2017), the SOC debt since the beginning of human cropping activities has been at around 37 GtC for the first 30 cm of the soil with half of it attributed to SOC depletion on grasslands. Our estimate of 22 GtC in 2010 for cropland debt is higher as Sanderman et al. (2017) estimations. However, there are large uncertainties in modeling SOC.



We have (1) developed a reduced-complexity model and (2) compiled a spatially explicit time series data set of agricultural management data in order to (3) analyze the role of agricultural management in historical cropland SOC dynamics. Our study shows that information on agricultural management alters estimates of the SOC debt and slows down loss of SOC compared to the often used constant management assumptions.

It is important to evaluate the validity of our results as modeling management effects on SOC dynamics at the global scale, and Sanderman et al. (2017) pointed out that their results might be conservatively low compared to experimental results. This suggests that our results are within a plausible range, comes with large uncertainties. The model includes a large number of parameters, and for most of these the uncertainty distributions have not been quantified so far. Moreover, we think that beyond parameter uncertainty, the structural uncertainty from the model design is high. The management data itself is prone to uncertainties as well, as most of it is only indirectly calculated from reported data.

Furthermore, Sanderman et al. (2017) modeled historic trends based on agricultural land expansion without considering SOC variations due to time-variant agricultural management. Pugh et al. (2015) considered management effects like tillage and residue returning in a static way, but neither changes over time nor alignment to observed historic data like yield levels or no-tillage areas were taken into account. Their study moreover concludes that crop productivity gains (increasing yield levels by 18% in their simulations) do not lead to a substantial decline in SOC debt (less than 1% change). Historic productivity increases were, however, notably larger. Despite large spatial heterogeneity aggregate yield increase rates are often estimated to be well above 50% (Pellegrini and Fernández, 2018; Ray et al., 2012; Rudel et al., 2009). In the following, we give a qualitative assessment of the uncertainties and limitations of this study as well as discuss our three study objectives and results against existing literature.

Our study for the first time uses a dynamic management dataset as driver for SOC dynamics. We show that the moderate global cropland expansion of around 11% between 1974 and 2010 and the resulting depletion of SOC stocks in converted cropland has been outweighed by improvements in agricultural productivity and practices. This is in contrast to Pugh et al. (2015) findings of only small effects due to improved practices.

#### 4.1 Comprehensive historical agricultural management data set

Looking on historic cropland intensification rates of above 50% (Rudel et al., 2009), a large fraction of increased C input from residue biomass is attributed to productivity improvements. Our spatially explicit time series data set of agricultural management is based on country-specific FAO production and cropland statistics (FAOSTAT, 2016) as well as 0.5 degree land-use data from LUH2 (Hurt et al., 2020). Starting from these two sources, we derive a harmonized and consistent data set for the major C flows within the agricultural system (4) using a mass balance approach from the IPCC guidelines Vol. 4 (Eggleston et al., 2006; Calvo Buendia et al., 2019) and other auxiliary data sets (e.g. Porwollik et al., 2019).

For some of the aspects covered in our data set, for example livestock distribution (Robinson et al., 2014) or manure production and application (Zhang et al., 2017), well-compiled data sets in high resolution exist that capture real world conditions much better than our estimates. However, they often come with the caveats of either being static in time, demanding large sets of auxiliary data or being inconsistent with each other.

Modeling management effects at the global scale comes with parametric and structural uncertainties. High SOC gains in the first 30 cm of the soil compared to natural vegetation in e. g. the temperate croplands of Central Europe and the United Kingdom indicate suspiciously high estimates of SOC inputs. For most of the parameters used in our management estimates no uncertainty estimates exist. This is why, in our view, most of the uncertainty with respect to the impacts of SOC management is included in the management data itself, and especially in the residue and manure production and application numbers, as these are only indirectly derived from crop and livestock production, feed and area data (FAOSTAT, 2016; Weindl et al., 2017). The uncertainty of recycling shares adds on top of the uncertain total numbers of manure and residue biomass. Previous modeling studies of SOC carbon on cropland often only used stylized scenarios of management practices (Pugh et al., 2015; Lutz et al., 2019), rather than trying to estimate real management.

While our data set, by including crop residues and manure, likely the largest carbon inputs to soils, it does not account for a list of minor carbon inputs from cover crops, agroforestry, green manure, weed biomass as well as application of human excreta, sewage sludge, processing wastes, forestry residues or biochar. Including these sources would correct our estimates upwards and bring our estimates closer to the IPCC stock change factors (see Sect. 3.4.1). Unfortunately, data on the quantity of these inputs is very scarce and does not exist with global coverage.

SOC inputs from above-ground residues had the strongest management effect on SOC debt dynamics on cropland (see Fig. 5). As pointed out by Keel et al. (2017) and Smith et al. (2020), carbon input calculations are highly sensitive to the choice of allometric functions determining below- and above-ground residue estimates from harvested quantities (see A1 for coefficients used in this study). Keel et al. (2017) question whether below-ground residues might increase with a fixed root:shoot ratio rather than being independent of productivity gains. Moreover, the study pointed out that plant breeding shifts allometries, which might not be reflected in outdated data sources. While our study considers a dynamic harvest index with rising yields for several crops, we may still overestimate residue biomass, in particular for below-ground biomass. However, looking on the evaluation of management effects (see Sect. 3.4.1), there is no general indication of overestimating.

## 4.2 Reduced complexity SOC model

Our reduced-complexity SOC model is based on a Tier 2 modeling approach. This reduces the computational and data demand of the model in comparison to DGVMs, while still allowing for the explicit representation of agricultural management practices. Along with the effects of various C inputs, the impacts of water supply from rainfall and irrigation as well as tillage systems can also be accounted for in the computation of SOC decay rates. As such, the model can reflect the spatial and temporal heterogeneity in both management and biophysical conditions.

The substantial impact of changing management practices through time is indicated by the development of our estimated stock change factors at least compared to IPCC default assumptions. Moreover, it is also likely that we are still missing carbon inputs to the soil from e. g. cover and inter-cropping practices (see Table 3) as well as by the time trend of the SOC debt (see Fig. 2(a)). Residue management has changed over the last decades, especially with the phasing out of residue burning practices in several regions and increased general productivity, showing a clear impact on SOC dynamics — underlining the importance to account for these effects in soil carbon modeling.

The Tier 2 approach (Ogle et al., 2019) used here is explicitly designed for agricultural soils, whereas we apply it also to soils under PNV. This is necessary in order to represent SOC losses under land-use change in a dynamic way, as this is an important driver of SOC dynamics. The comparison of simulated PNV data with LPJmL4 shows the model's substantial capability in reproducing PNV SOC stocks (Fig. 8).

Another uncertainty is connected with the initialization of SOC stocks in 1901. Using litterfall estimates from LPJmL4, we have been able to estimate the total SOC stocks of the world, which is assumed to be in steady state considering the land-use pattern of 1901 and agricultural management data of 1965. As shown, dominated by SOC under natural vegetation. However, as the world's SOC stock is highly uncertain, which is seen in the wide range of global SOC stock estimates for the first 30 cm of the soil profile (Batjes, 2016; Hengl et al., 2017; FAO, 2018; Schaphoff et al., 2018a; Poggio et al., 2021; Sanderman et al., 2017) in Fig. ?? the SOC debt estimate almost halves (from -26 GtC to -14 GtC), and the SOC debt reduction is strongly reduced (from -4 GtC to -2.5 GtC), if considering initialization SOC stocks under undisturbed natural vegetation. Pugh 6, the only conclusion we can draw from this is that our result is within a plausible range. To avoid a strong impact of natural land representation and its uncertainties on our results, we focus on SOC changes on cropland. Pristine natural vegetated areas (like permafrost and rain forests) without human land management drop out in our calculation of SOC debt and stock change factors. Natural SOC estimates only influence results when natural land is converted to cropland.

Comparing the geographic SOC patterns to Soil Grids 2.0 (Poggio et al., 2021) (see Fig. 10), we find that our model estimates values of SOC stocks greater than the estimated confidence intervals in Soil Grid 2.0 for some mountainous regions across the globe. This could indicate that we are not capable in capturing specific processes that would reduce the vegetation's productivity (such as erosion on steep slopes or shallow soils (Borrelli et al., 2017)). A large swathe of eastern North America was heavily affected by the dust bowl event, with wind erosion removing large parts of the topsoils, a process not considered in our model. Similarly, we likely overestimate SOC stocks for the loess soils in northern China and the Altiplano in Latin America; in both cases erosion is a likely reason. In contrast, we estimate lower SOC stocks at the edges of the Sahara, where uncertain local water availability and artificial irrigation may dominate spatial SOC patterns.

In our model, erosion should however only affect the spatial pattern but not the aggregate SOC pool. As pointed out by Doetterl et al. (2015) pointed out the importance of accounting for the land-use history, as many CO<sub>2</sub> emissions from agricultural soils are caused by historic land-use change (LUC) and the slow decline of SOC under cropland before it reaches a new equilibrium. Our results of the Initial-natveg scenario show that the qualitative finding of a reduction of SOC debt through improved agricultural management is robust to changing the initialisation of soil organic carbon, even though the level of SOC debt is sensitive to the initialization setting (2016), the final fate of leached or eroded carbon is uncertain and might even offset LUC emissions (Wang et al., 2017). Whereas for soil quality analysis SOC displacement might play an important role, in this budget approach focused especially on the SOC debt, displaced but not emitted SOC can be treated as SOC that remains on the cropland. Erosion and degradation impacts on yields and therefore on soil C inputs are captured by our method as we base them on FAO statistics of actual production. Yet the distribution of production below the country level - which we allocate proportional to LPJmL production potentials that do not reflect erosion feedback to yields - will overestimate yields and therefore biomass inputs to eroded soils.

In comparison with default stock change factors of the IPCC guidelines, our model estimates a stronger decline of SOC stocks (Table 3) for almost all regions. Tropical soils might suffer from low C input rates due to large yield gaps (Global Yield Gap and Water Productivity Atlas. Available URL: [www.yieldgap.org](http://www.yieldgap.org) (accessed on: 03/01/2022)) and high shares of residue removal and burning in lower-income countries (Smil, 1999a; Williams et al., 1997; Jain et al., 2014). Yet, even when comparing our estimates to the low-input stock change factors of the IPCC, our SOC loss is roughly twice as large as the revised 2019 IPCC default values, while it shows very good agreement with the older default values from IPCC (2006). Don et al. (2011) estimated SOC losses for tropical soils of around 25% on average corresponding to a stock change factor of 0.75, but also reported a wide range of measured SOC changes from -80% to +58%. Fujisaki et al. (2015) however found much lower loss rates of around 9%, attributing the difference to the different time period length since the conversion to cropland. As our results do not specifically account for cropland age and most of the cropland is older than 20 years (as assumed for the default IPCC Tier 1 stock change factors) our stock change factors have to be lower by definition following the steady-state assumption that cropland will continue to approach a new equilibrium. For the same reason, our estimates for temperate regions might be lower than both IPCC (2006) and IPCC (2019) default values. With the production-increasing impact of irrigation and fertilization on carbon-poor dryland soils, SOC under cropland can also be higher than under PNV with stock change factors above 1 (see Fig. 2(d)), but these areas are much smaller than where the stock change factors are well below unity.

Generally, ~~the limit limiting the analysis~~ to the first 30 cm of the soil profile follows the IPCC guidelines (Eggleston et al., 2006; Calvo Buendia et al., 2019) and assumes that most of the SOC ~~dynamic happens~~ dynamics happen in the topsoil. In this regard several aspects are strongly simplified within our approach. Firstly, distribution of carbon inputs into different soil layers are neglected and all carbon inputs are allocated to the topsoil. This particularly overestimates SOC stocks in the first 30 cm of soil below deeper rooting vegetation, which is certainly the case for most of the woody natural vegetated areas. ~~Secondly~~ Second, changes to the subsoil due to tillage are neglected. As Powlson et al. (2014) have shown, the subsoil can be ~~a game changer~~ make a large difference in evaluating total SOC losses or gains for no-tillage systems. No-tillage effects may seem larger than ~~the actually are, if only focusing on the topsoil. SOC transfer~~ they actually are if only topsoil is considered. SOC transfers to deeper soil layers under tillage, might enhance subsoil SOC compared to no-till practices. ~~Thirdly~~ Finally, organic soils (like peat- and wetlands) and drained cropland areas are not explicitly considered and emissions from these cropland areas are thus likely substantially underestimated.

~~This study excludes not only peatland degradation, it also does not account for carbon displacement via leaching and erosion. However, as pointed out by Doetterl et al. (2016), the final fate of leached or eroded carbon is uncertain and might even offset LUC emissions (Wang et al., 2017). Whereas for soil quality analysis SOC displacement might play an important role, in this budget approach focusing especially on the SOC debt, displaced but not emitted SOC can be treated as SOC that remains on the cropland. Moreover, the exclusive focus on croplands ignores LUC emission on other land-use types such as pastures, rangelands and forestry. Human interventions have led to large changes in SOC stocks there as well (Sanderman et al., 2017; Friedlingstein et al., 2019). This study does not intend to be a comprehensive LUC emission analysis and acknowledges that land-use changes comes with large overall emissions.~~

To validate our modeled SOC stocks and stock changes under management, we compare our results to default IPCC stock changes factors of 2006 (Eggleston et al., 2006) and their refinements in 2019 (Calvo Buendia et al., 2019). Both estimates are based on measurement data for croplands (see Table 3). To allow for comparison, we aggregate our stock change factors to the four IPCC climate zones (Fig. A1).

### 4.3 SOC debt and SOC drivers

$F^{SCF}$  in comparison to IPCC Tier 1 default factors: Stock change factors are in good agreement with the default values of the IPCC in general. For the tropical regions the assumptions changed notably from the guidelines in 2006 (Eggleston et al., 2006) to the update in 2019 (Calvo Buendia et al., 2019), leaving our results in very good agreement with the old default assumptions. Default assumption are given under the assumption of medium input systems, which, considering the yield gap in mainly developing regions in the tropics, might be an overestimation and decrease  $F^{SCF}$  by additional 5-8 percent. Also modelled  $F^{SOC}$  have increased for all climates over time. Source-Input-Year tropical moist tropical dry temperate dry temperate moist

	1-IPCC2006 low invariant	0.44	0.55	0.61	0.74	0.66	2-IPCC2006 medium invariant	0.48	0.58	0.64	0.80	0.69	3-IPCC2006 high invariant	0.53	0.60	0.67	0.83	0.77
The analysis of SOC stock gains and losses is complex and has several dimensions as climatic and anthropogenic effects overlap. There is broad consensus that land conversion to cropland has caused substantial C emissions over the historical period (e.g. Friedlingstein et al., 2020). There is uncertainty with respect to the overall size of these emissions from different methods and reference points and with respect to the contribution of cropland and agricultural management to these emissions. In order to mitigate greenhouse gas emissions, it is essential to stop the decline of SOC stocks or even transform cropland management to sequester atmospheric C in cropland soils (Minasny et al., 2017). Defining the SOC debt of 1975 as the baseline, and measuring land-use emissions on cropland as the difference between a potential natural state and the state under human interventions (see Pugh et al., 2015), we find that global cropland has acted as a emissions source since 1975. Annual C loss rates of 0.2 per 1000 C still have the opposite trend as the promoted																		
	4-IPCC2019 low invariant	0.76	0.87	0.70	0.71	0.66	0.67	5-IPCC2019 medium invariant	0.83	0.92	0.76	0.77	0.69	0.70	6-IPCC2019 high invariant	0.92	0.96	0.79
	7-This Study hist 1990	0.57	0.61	0.78	0.76	8-This Study hist 2010	0.64	0.68	0.83	0.83								

Whereas our estimates are lower in the two tropical climate zones, temperate zone default factors are higher than Calvo Buendia et al. (2019). The tropical factors differ substantially between the IPCC guidelines in 2006 (Eggleston et al., 2006) and their refinement in 2019 (Calvo Buendia et al., 2019), with our estimates within this range. Taking the simplified assumption, that tropical soils might suffer from insufficient C input rates (low) due to yield gaps, whereas temperate soils in developed regions might be highly managed and fertilized (high), the difference vanishes partially. Our results still show especially in temperate dry regions — the smallest region area-wise — small deviations from natural SOC stocks. Considering the impact of irrigation and fertilization on carbon-poor dryland soils, even factors above 1 (see Fig. 2(c)) may be expected.

With regard to the time trend, our study per 1000 C sequestration rate target (Minasny et al., 2017). Dedicated efforts to increase cropland SOC are thus necessary, as management improvements at historical rates are not enough to counteract ongoing SOC degradation on cropland. Yet our study also shows the substantial impact of changing management factors on the development of stock change factors as also indicated by the time trend of the SOC debt SOC debt (Fig. 5).

650 The world's SOC stock and its changes are highly uncertain, which is seen in the wide range of global SOC stock estimates. According to Sanderman et al. (2017), the SOC debt since the beginning of human cropping activities has been at around 37 GtC for the first 30 cm of the soil profile (Batjes, 2016; Hengl et al., 2017; FAO, 2018; Schaphoff et al., 2018a) in Table 6).

Modeled as well as data-based estimation for global SOC stock in GtC for the first 30 cm of soil aggregated over all land area: Note that SoilGrids, GSOC and WISE do not consider land-use as well as changes over time and rely on soil profile data

655 gather over a long period of time. This makes it hard to pinpoint a specific year for these SOC estimations. In this context they will be compared to modeled data from LPJmL4 for potential natural vegetation and this study for the year 2010.

The global estimates of the total SOC stock from this study are on the lower end compared to other modeled results or data driven estimates. SoilGrids (Hengl et al., 2017) especially stands out for their high estimation, whereas all other sources (including our study) are comparably similar. Looking at regional results in Fig. A2, our estimates turn out to be in good

660 agreement for most of the world, with the largest deviations for boreal moist and tropical moist areas. To avoid that this bias influences our results, which originates from uncertainties in the representation of natural land, we focus on SOC changes on cropland. Pristine natural vegetated areas (like permafrost and rain forests) without human land management thus drop out in our calculation of SOC debt and stock change factors.

Additionally, our estimates for total SOC stocks of the world (as well as our SOC initialization) are dominated by the

665 representation of natural vegetation, which are only estimated in a basic manner. For example, we do not differentiate the parameterization of nitrogen and lignin content of litterfall for woody and grass plant types. This renders carbon inputs and decay dynamics for natural litterfall rather uncertain. The absolute values of SOC stocks and debt from land-use change have to be interpreted with caution. As our default litter parameterization accounts for woody plant types, larger uncertainty in natural land SOC dynamics may arise especially in less forested areas.

670 We conducted a sensitivity analysis (Fig. ??) based on various plant parameterizations from the Century model (see Sect. ??). This shows that the general trend of decreasing SOC debt of ~4 GtC within the period of 1975–2010 is not altered under various estimates for natural SOC stocks.

with half of it attributed to SOC depletion on grasslands. Our estimate of 39.6 GtC in 2010 for cropland debt is thus twice as high as their estimate. However, there are large uncertainties in modeling SOC at the global scale, and Sanderman et al. (2017)

675 pointed out that their results might be conservatively low compared to experimental results.

Our global SOC model is able to estimate spatially explicit SOC stocks, SOC debts and stock change factor considering agricultural management. It is — to our knowledge — the first study that quantifies the impact of

Furthermore, Sanderman et al. (2017) modeled historical trends based on agricultural land expansion without considering SOC variations due to time-variant and spatially explicit historic agricultural management on global SOC stocks

680 considered management effects like tillage and incorporation of residues in stylized and static scenarios only, so that they could not account for historical management effects on SOC dynamics. Their study moreover concludes that yield gains (by 18% in their simulations) do not lead to a substantial decline in SOC debt (less than 1% change). Historical yield increases, however, are often estimated to be well above 50% (Pellegrini and Fernández, 2018; Ray et al., 2012; Rudel et al., 2009). While we find

substantially larger effects of productivity gains than the 1% reported by Pugh et al. (2015), this is not sufficient to compensate SOC losses from moderate global cropland expansion of around 11% between 1974 and 2010.

The effects of agricultural productivity on cropland SOC dynamics, including historical yield trends and associated increases in residue inputs, can be directly accounted for in our modeling approach. In contrast, process-based studies (Pugh et al., 2015; Herzfeld et al. 2021) often lack data on relevant management aspects that drive production increases. Herzfeld et al. (2021) also consider historical management trends for fertilizer and manure inputs as well as on residue removal rates and tillage systems, but cannot reproduce the substantial increase in agricultural productivity over the last decades. Still, they find that compared to no-tillage systems, residue management has much larger potential to affect the strength of their projected future global cropland SOC decline. This is consistent with our finding that increasing SOC inputs from above-ground residues had the strongest effect on the slowing-down of the SOC debt increase (Fig. 5).

Our results clearly demonstrate that agricultural management needs to be explicitly considered in global carbon assessments and models. That also implies that we need better monitoring of agricultural practices to create this data, but also better accessibility of existing data. Our data set and the MADRaT package (Dietrich et al., 2020) constitute a starting point for building comprehensive data sets on agricultural management aspects. Elliott et al. (2018) show that yield trends in the USA can be reproduced by models, but require information on inputs that are not available at the global scale, such as annual data on sowing dates, planting densities, and genetic traits such as kernel number and radiation use efficiency. As such, it will remain challenging for process-based DGVMs to capture the trend of agricultural productivity on cropland SOC dynamics.

Our study again highlights emphasizes again that the expansion of croplands cropland is still a major source of CO<sub>2</sub> emissions — not only by through the removal of vegetation, but also by a slow depletion of C stocks in soils. Our estimates indicate a SOC debt of 22–39.6 GtC in 2010, and every additional deforestation deforested hectare adds to this debt.

However, our results also indicate that the changes in cropland management have led to increased SOC stocks in global cropland soils as a continuous trend since 1975. Even more, this trend of improved management on existing croplands more than overcompensates the depletion of SOC stocks on newly converted soils. The finding of this recovery of cropland SOC stocks challenges the assumption that cropland soils are a CO<sub>2</sub> source. Only under the assumption that cropland management is static over time, as typically assumed in other studies on cropland SOC stocks, we can reproduce their finding that cropland soils are a source of CO<sub>2</sub>. The estimated increase in cropland SOC is therefore caused by changing agricultural management, with the largest contribution stemming from increased carbon inputs to soils from crop residues.

Continuing the historical development, a further closure of the yield gaps may be beneficial to SOC stocks. Forest protection schemes for climate change mitigation would not only reduce SOC losses on newly converted land, they would likely also require further productivity improvements in existing croplands to meet crop demand (Popp et al., 2014). Avoided deforestation and other environmental regulation leads to intensification on existing cropland (Humpenöder et al., 2018) and our results show that such intensification could lead to increased cropland SOC, if residues are returned to the soil — and thereby usually further reduce the SOC debt on croplands.



While further crop productivity gains lead to co-benefits for SOC stocks, they are likely not enough to help meeting ambitious climate mitigation targets. Our findings on sequestration rates are still more than an order of magnitude lower than promoted by the 4 per 1000 initiative amplifying the C sequestration potential of avoided deforestation.

720 ~~Yet, there is~~ There is also ample potential for further improved SOC management. As shown in Fig. 4, approximately one fifth of total annual C sequestration by crops is lost through soils (0.8 GtC per year). ~~Large losses in fact~~ However, even larger losses occur at the end of the food supply chain (1.2 GtC year), ~~and at the litter soil barrier at the soil surface~~ (1.4 GtC), during residue burning (0.3 Gt C) and with manure management (0.2). Improved management could include, firstly, a circular flow from the food supply chain back to soils. Waste composting or excreta recycling could represent a major additional ~~N~~ C input to cropland soils (Brenzinger et al., 2018). Secondly, soil carbon sequestration techniques (Smith, 2016), deep ploughing (Alcántara et al., 2016) or the transformation of ~~e~~ C inputs to more recalcitrant biochar (Woelf et al., 2010) may transfer larger parts of the biomass at the litter soil barrier into permanent soil pools. Thirdly, reducing the share of residue burning and improved manure recycling could further increase C inputs. Finally, other carbon-accumulating practices, such as the cultivation of cover crops (~~Poeplau and Don, 2015~~) (Poeplau and Don, 2015; Porwollik et al., 2022) and agroforestry (Lorenz and Lal, 2014) could increase total C sequestration on ~~croplands~~ cropland.

725

730



## 5 Conclusions

We have compiled a spatially explicit and time-variant data set on agricultural management aspects relevant for cropland SOC dynamics. We have also developed a reduced-complexity SOC model that is able to be applied in optimization-based IAM frameworks, for which detailed process-based models are computationally too expensive. Making use of these data and model, we are able to estimate spatially explicit SOC stocks, SOC debts, and stock change factors considering agricultural management. It is — to our knowledge — the first study that analyzes the role of time-variant and spatially explicit historical agricultural management for global SOC dynamics.


Our results demonstrate that historical changes in agricultural management have shaped the SOC debt on cropland. It is thus necessary to explicitly consider agricultural management in a dynamic manner in global carbon assessments and models, especially when exploring climate mitigation pathways with so-called land-based solutions (e.g. Popp et al., 2016). That also implies that we need better monitoring of agricultural practices to create this data, but also better accessibility of existing data. Our open-source model (Karstens and Dietrich, 2020), published data-set (Karstens, 2020a) and the flexible data processing with the MADRaT package (Dietrich et al., 2020) constitute a starting point for building comprehensive data sets on agricultural management aspects.

With the reduced-complexity SOC model we are able to account for agricultural management effects on cropland SOC dynamics within optimization-based IAM frameworks. Reduced input data requirements such as accounting for changes in productivity rather than reproducing the processes that lead to such changes in productivity (Elliott et al., 2018) will help to explore the role of agricultural management in sustainable development pathway analyses (Sörgel et al., 2021). However, we clearly see that increases in agricultural productivity are not sufficient to create positive net SOC sequestration in cropland soils. More management options that explicitly target the sequestration of C in cropland soils need to be considered. Our open-source model can be expanded to account for additional management options for carbon farming, such as cover crops, agroforestry, or biochar applications.


*Code and data availability.* We compile calculations as open-source R packages available at [github.com/pik-piam/mrcommons](https://github.com/pik-piam/mrcommons) (Bodirsky et al., 2020a) for the management related functions, [github.com/pik-piam/mrsoil](https://github.com/pik-piam/mrsoil) (Karstens and Dietrich, 2020) for soil dynamic related functions and [github.com/pik-piam/mrvalidation](https://github.com/pik-piam/mrvalidation) (Bodirsky et al., 2020b) for validation data. All libraries are based on the MADRaT package at [github.com/pik-piam/madratt](https://github.com/pik-piam/madratt) (Dietrich et al., 2020), a framework which aims to improve reproducibility and transparency in data processing. Model results including C input data are accessible under <https://doi.org/10.5281/zenodo.4320663> (Karstens, 2020a). Software code for paper and result preparation can be found under [www.github.com/k4rst3ns/historicalsocmanagement](https://www.github.com/k4rst3ns/historicalsocmanagement).

**Table A1.** Parameterization of harvested organs and their corresponding residues parts as well as allometric coefficients: This table is mainly based on Bodirsky et al. (2012) together with simple carbon to dry matter assumptions. Allometric coefficients are used as described in Eggleston et al. (2006) with  $HI^{area}$  being slope<sub>(T)</sub>,  $HI^{area}$  intercept<sub>(T)</sub> and RS  $R_{BG-BIO}$ .

Crop code	Crop Type	Harvested Organs			Above-ground Residues			Below-ground Residues			Allometric coefficients	
		nr/dm	wm/dm	c/dm	nr/dm	wm/dm	c/dm	nr/dm	c/dm	$HI^{area}$	$HI^{prod}$	RS
tece	Temperate cereals	0.0217	1.14	0.42	0.0074	1.11	0.42	0.0098	0.38	0.58	1.36	0.24
maiz	Maize	0.016	1.14	0.42	0.0088	1.18	0.42	0.007	0.38	0.61	1.03	0.22
trce	Tropical cereals	0.0163	1.14	0.42	0.007	1.18	0.42	0.006	0.38	0.79	1.06	0.22
rice_pro	Rice	0.0128	1.15	0.42	0.007	1.11	0.42	0.009	0.38	2.46	0.95	0.16
soybean	Soybean	0.0629	1.13	0.42	0.008	1.11	0.42	0.008	0.38	1.35	0.93	0.19
rapeseed	Other oil crops (incl rapeseed)	0.0334	1.08	0.42	0.0081	1.11	0.42	0.0081	0.38	0	1.86	0.22
groundnut	Groundnuts	0.0299	1.06	0.42	0.0224	1.11	0.42	0.008	0.38	1.54	1.07	0.19
sunflower	Sunflower	0.0216	1.08	0.42	0.008	1.11	0.42	0.008	0.38	0	1.86	0.22
oilpalm	Oilpalms	0.0027	1.01	0.49	0.0052	1.11	0.48	0.0053	0.47	0	1.86	0.24
puls_pro	Pulses	0.0421	1.1	0.42	0.0105	1.16	0.42	0.008	0.38	0.79	0.89	0.19
potato	Potatoes	0.0144	4.55	0.42	0.0133	6.67	0.42	0.014	0.38	1.06	0.1	0.2
cassav_sp	Tropical roots	0.0053	2.95	0.42	0.0101	6.67	0.42	0.014	0.38	0	0.85	0.2
sugr_cane	Sugar beet	0.0024	3.7	0.42	0.008	3.82	0.42	0.008	0.38	0	0.67	0.07
sugr_beet	Sugar beet	0.0056	4.17	0.42	0.0176	5	0.42	0.014	0.38	0	0.54	0.2
others	Fruits, Vegetables, Nuts	0.0267	5.49	0.42	0.0081	1.88	0.42	0.007	0.38	0	0.39	0.22
foddr	Forage	0.0201	4.29	0.42	0.0192	4.1	0.42	0.0141	0.38	0	0.28	0.45
cottn_pro	Cotton seed	0.0365	1.09	0.42	0.0093	1.18	0.42	0.007	0.38	0	1.48	0.13
		nr/dm – nitrogen to dry matter ratio			$HI^{area}$ – harvest index per area							
		wm/dm – wet matter to dry matter ratio			$HI^{prod}$ – harvest index per production							
		c/dm – carbon to dry matter ratio			RS – root:shoot ratio							



**Figure A1.** Climate zone map adapted from IPCC: The climate zone classification is based on the classification scheme of the IPCC guidelines (Eggleston et al., 2006) and has been reimplemented by Carre et al. (2010), which is the source of this data. Note that the reduced set, used for the comparison of stock change factors is included in the color code with temperate moist in light blue, temperate dry in dark violet, tropical moist in red and tropical dry in orange.



**Figure A2.** Modelled as well as data based estimation for climate zone specific SOC stock in GtC for the first 30 cm of soil aggregated over all land area: SoilGrids, GSOC and WISE do not consider changes over time and rely on soil profile data gather over a long period of time, which makes it hard to pinpoint a specific year to these SOC estimations. In this context they will be compared to modelled data (LPJmL4, this study) for the year 2010. PNV denotes the potential natural vegetation state without considering human cropping activities, calculated as reference stock within our model. We use the climate zone specification of the IPCC (Eggleston et al., 2006).

~~Global  $\Delta$ SOC for different litter parameterization choices: Whereas perennial grasses (scenario name *LitterPNV-PerennialGrasses*) are given by the IPCC guidelines (Calvo Buendia et al., 2019), we used CENTURY configuration file for woody biomass parameterization (see tree.100 in NREL 2000). *LitterPNV-CenturyAverage* forms the baseline parameterization as used within the study and is calculated as the average over all tree compartments and tree types equally. Litter parameterization has the ability to change global SOC stocks and SOC debt, but is robust concerning trends and sequestration rates.~~

*Author contributions.* KK, BLB and AP designed the study and the model idea. KK wrote the code build on work of BLB, IW. JPD revised and improved the model code. CM, JH and SR provided the LPJmL simulation data. KK wrote the paper with important contributions of BLB and CM. MK, JS, SR and IW provided extensive feedback to outline of the study. All authors discussed the results and commented on the manuscript.

*Competing interests.* The authors declare no competing interests.

*Acknowledgements.* Thanks to Vera Porwollik for contributing the time resolved tillage data set based on her previous work. Additional thanks to the rticles contributors for providing a R Markdown template. The authors thank for the data provided by FAOSTAT and LUH2v2. The work of KK was funded by the DFG Priority Program “Climate Engineering: Risks, Challenges, Opportunities?” (SPP 1689) and specifically the CEMICS2 project (grant no. ED78/3-2). The research leading to these results has received funding for BLB from the European Union’s Horizon 2020 research and innovation program under grant agreement no. 776479 (COACCH) and no. 821010 (CASCADES). The work of SR, JS and IW was also supported by CLIMASTEPPPE (01DJ8012), EXIMO (01LP1903D) and FOCUS (031B0787B) funded by the German Federal Ministry of Education and Research (BMBF). The input of PS, MK and MD contributes to the Soils-R-GGREAT project (NE/P019455/1) and CIRCASA (EU H2020; grant agreement no. 774378).

- Alcántara, V., Don, A., Well, R., and Nieder, R.: Deep Ploughing Increases Agricultural Soil Organic Matter Stocks, *Global Change Biology*, 22, 2939–2956, <https://doi.org/10.1111/gcb.13289>, 2016.
- Bastos, A., Hartung, K., Nützel, T. B., Nabel, J. E. M. S., Houghton, R. A., and Pongratz, J.: Comparison of Uncertainties in Land-Use Change Fluxes from Bookkeeping Model Parameterisation, *Earth System Dynamics*, 12, 745–762, [https://doi.org/10.5194/esd-12-745-](https://doi.org/10.5194/esd-12-745-2021)  
780 2021, 2021.
- Batjes, N.: Total Carbon and Nitrogen in the Soils of the World, *European Journal of Soil Science*, 47, 151–163, <https://doi.org/10.1111/j.1365-2389.1996.tb01386.x>, 1996.
- Batjes, N. H.: Harmonized Soil Property Values for Broad-Scale Modelling (WISE30sec) with Estimates of Global Soil Carbon Stocks, *Geoderma*, 269, 61–68, <https://doi.org/10.1016/j.geoderma.2016.01.034>, 2016.
- 785 Betts, R. A., Golding, N., Gonzalez, P., Gornall, J., Kahana, R., Kay, G., Mitchell, L., and Wiltshire, A.: Climate and Land Use Change Impacts on Global Terrestrial Ecosystems and River Flows in the HadGEM2-ES Earth System Model Using the Representative Concentration Pathways, *Biogeosciences*, 12, 1317–1338, <https://doi.org/10.5194/bg-12-1317-2015>, 2015.
- Bodirsky, B. L., Popp, A., Weindl, I., Dietrich, J. P., Rolinski, S., Scheffele, L., Schmitz, C., and Lotze-Campen, H.: N&lt;Sub&gt;2&lt;/Sub&gt;O Emissions from the Global Agricultural Nitrogen Cycle – Current State and Future Scenarios, *Biogeo-*  
790 sciences, 9, 4169–4197, <https://doi.org/10.5194/bg-9-4169-2012>, 2012.
- Bodirsky, B. L., Karstens, K., Baumstark, L., Weindl, I., Wang, X., Mishra, A., Wirth, S., Stevanovic, M., Steinmetz, N., Kreidenweis, U., Rodrigues, R., Popov, R., Humpenoeder, F., Giannousakis, A., Levesque, A., Klein, D., Araujo, E., Beier, F., Oeser, J., Pehl, M., Leip, D., Molina Bacca, E., Martinelli, E., Schreyer, F., and Dietrich, J. P.: Mrcommons: MadRat Commons Input Data Library, <https://doi.org/10.5281/zenodo.3822009>, 2020a.
- 795 Bodirsky, B. L., Wirth, S., Karstens, K., Humpenoeder, F., Stevanovic, M., Mishra, A., Biewald, A., Weindl, I., Chen, D., Molina Bacca, E., Kreidenweis, U., W. Yalaw, A., Humpenoeder, F., Wang, X., and Dietrich, J. P.: Mrvalidation: Madrat Data Preparation for Validation Purposes, <https://doi.org/10.5281/zenodo.4317826>, 2020b.
- Bondeau, A., Smith, P. C., Zaehle, S., Schaphoff, S., Lucht, W., Cramer, W., Gerten, D., Lotze-Campen, H., Müller, C., Reichstein, M., and Smith, B.: Modelling the Role of Agriculture for the 20th Century Global Terrestrial Carbon Balance, *Global Change Biology*, 13,  
800 679–706, <https://doi.org/10.1111/j.1365-2486.2006.01305.x>, 2007.
- Borrelli, P., Robinson, D. A., Fleischer, L. R., Lugato, E., Ballabio, C., Alewell, C., Meusburger, K., Modugno, S., Schütt, B., Ferro, V., Bagarello, V., Oost, K. V., Montanarella, L., and Panagos, P.: An assessment of the global impact of 21st century land use change on soil erosion, *Nature Communications*, 8, <https://doi.org/10.1038/s41467-017-02142-7>, 2017.
- Brenzinger, K., Drost, S. M., Korthals, G., and Bodelier, P. L. E.: Organic Residue Amendments to Modulate Greenhouse Gas Emissions From Agricultural Soils, *Frontiers in Microbiology*, 9, <https://doi.org/10.3389/fmicb.2018.03035>, 2018.
- 805 Brovkin, V., van Bodegom, P. M., Kleinen, T., Wirth, C., Cornwell, W. K., Cornelissen, J. H. C., and Kattge, J.: Plant-Driven Variation in Decomposition Rates Improves Projections of Global Litter Stock Distribution, *Biogeosciences*, 9, 565–576, <https://doi.org/10.5194/bg-9-565-2012>, 2012.
- Calvo Buendia, E., Tanabe, K., Kranjc, A., Baasansuren, J., Fukuda, M., Ngarize, S., A.Osako, Pyrozhenko, Y., Shermanau, P., and (eds),  
810 S. F., eds.: IPCC 2019, 2019 Refinement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories,, Published: IPCC, Switzerland, 2019.

- Carre, F., Hiederer, R., Blujdea, V., and Koeble, R.: Background Guide for the Calculation of Land Carbon Stocks in the Biofuels Sustainability Scheme : Drawing on the 2006 IPCC Guidelines for National Greenhouse Gas Inventories., Reference reports, Publications Office of the European Union, <https://doi.org/10.2788/34463>, 2010.
- 815 Coleman, K., Jenkinson, D. S., Crocker, G. J., Grace, P. R., Klir, J., Körschens, M., Poulton, P. R., and Richter, D. D.: Simulating Trends in Soil Organic Carbon in Long-Term Experiments Using RothC-26.3, *Geoderma*, 81, 29–44, [https://doi.org/10.1016/S0016-7061\(97\)00079-7](https://doi.org/10.1016/S0016-7061(97)00079-7), 1997.
- Del Grosso, S., Parton, W., Mosier, A., Hartman, M., Brenner, J., Ojima, D., and Schimel, D.: Simulated Interaction of Carbon Dynamics and Nitrogen Trace Gas Fluxes Using the DAYCENT Model, pp. 303–332, <https://doi.org/10.1201/9781420032635.ch8>, 2001.
- 820 Dietrich, J. P., Baumstark, L., Wirth, S., Giannousakis, A., Rodrigues, R., Bodirsky, B. L., Kreidenweis, U., and Klein, D.: Madrat: May All Data Be Reproducible and Transparent (MADRaT), <https://doi.org/10.5281/zenodo.1115490>, 2020.
- Doetterl, S., Berhe, A. A., Nadeu, E., Wang, Z., Sommer, M., and Fiener, P.: Erosion, Deposition and Soil Carbon: A Review of Process-Level Controls, Experimental Tools and Models to Address C Cycling in Dynamic Landscapes, *Earth-Science Reviews*, 154, 102–122, <https://doi.org/10.1016/j.earscirev.2015.12.005>, 2016.
- 825 Don, A., Schumacher, J., and Freibauer, A.: Impact of Tropical Land-Use Change on Soil Organic Carbon Stocks – a Meta-Analysis, *Global Change Biology*, 17, 1658–1670, <https://doi.org/10.1111/j.1365-2486.2010.02336.x>, 2011.
- Eggleston, H., Buendia, L., Miwa, K., Ngara, T., and (eds), K. T., eds.: IPCC 2006, 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Prepared by the National Greenhouse Gas Inventories Programme, Published: IGES, Japan., 2006.
- Elliott, J., Glotter, M., Ruane, A. C., Boote, K. J., Hatfield, J. L., Jones, J. W., Rosenzweig, C., Smith, L. A., and Foster, I.: Characterizing agricultural impacts of recent large-scale US droughts and changing technology and management, *Agricultural Systems*, 159, 275–281, <https://doi.org/https://doi.org/10.1016/j.agsy.2017.07.012>, <https://www.sciencedirect.com/science/article/pii/S0308521X16306643>, 2018.
- FAO: AQUASTAT Main Database, Food and Agriculture Organization of the United Nations (FAO), Rome, 2016.
- FAO: Global Soil Organic Carbon Map (GSOCmap) : Technical Report, FAO, Rome, Italy, 2018.
- FAOSTAT: FAOSTAT Database, The Food and Agriculture Organization of the United Nations (FAO), Rome, 2016.
- 835 Forster, P., Huppmann, D., Kriegler, E., Mundaca, L., Smith, C., Rogelj, J., and Séférian, R.: Mitigation Pathways Compatible with 1.5°C in the Context of Sustainable Development Supplementary Material. In: Global Warming of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change, sustainable development, and efforts to eradicate poverty [Masson-Delmotte, V., P. Zhai, H.-O. Pörtner, D. Roberts, J. Skea, P.R. Shukla, A. Pirani, W. Moufouma-Okia, C. Péan, R. Pidcock, S. Connors, J.B.R. Matthews,
- 840 Y. Chen, X. Zhou, M.I. Gomis, E. Lonnoy, T. Maycock, M. Tignor, and T. Waterfield (eds.)]. Available from <https://www.ipcc.ch/sr15>, 2018.
- Friedlingstein, P., Jones, M. W., O’Sullivan, M., Andrew, R. M., Hauck, J., Peters, G. P., Peters, W., Pongratz, J., Sitch, S., Le Quéré, C., Bakker, D. C. E., Canadell, J. G., Ciais, P., Jackson, R. B., Anthoni, P., Barbero, L., Bastos, A., Bastrikov, V., Becker, M., Bopp, L., Buitenhuis, E., Chandra, N., Chevallier, F., Chini, L. P., Currie, K. I., Feely, R. A., Gehlen, M., Gilfillan, D., Gkritzalis, T., Goll, D. S., Gruber, N., Gutekunst, S., Harris, I., Haverd, V., Houghton, R. A., Hurtt, G., Ilyina, T., Jain, A. K., Joetzjer, E., Kaplan, J. O., Kato, E., Klein Goldewijk, K., Korsbakken, J. I., Landschützer, P., Lauvset, S. K., Lefèvre, N., Lenton, A., Lienert, S., Lombardozi, D., Marland, G., McGuire, P. C., Melton, J. R., Metzl, N., Munro, D. R., Nabel, J. E. M. S., Nakaoka, S.-I., Neill, C., Omar, A. M., Ono, T., Peregon, A., Pierrot, D., Poulter, B., Rehder, G., Resplandy, L., Robertson, E., Rödenbeck, C., Séférian, R., Schwinger, J., Smith, N., Tans, P. P.,

- Tian, H., Tilbrook, B., Tubiello, F. N., van der Werf, G. R., Wiltshire, A. J., and Zaehle, S.: Global Carbon Budget 2019, *Earth System Science Data*, 11, 1783–1838, <https://doi.org/10.5194/essd-11-1783-2019>, 2019.
- Friedlingstein, P., O’Sullivan, M., Jones, M. W., Andrew, R. M., Hauck, J., Olsen, A., Peters, G. P., Peters, W., Pongratz, J., Sitch, S., Le Quéré, C., Canadell, J. G., Ciais, P., Jackson, R. B., Alin, S., Aragão, L. E. O. C., Arneeth, A., Arora, V., Bates, N. R., Becker, M., Benoit-Cattin, A., Bittig, H. C., Bopp, L., Bultan, S., Chandra, N., Chevallier, F., Chini, L. P., Evans, W., Florentie, L., Forster, P. M., Gasser, T., Gehlen, M., Gilfillan, D., Gkritzalis, T., Gregor, L., Gruber, N., Harris, I., Hartung, K., Haverd, V., Houghton, R. A., Ilyina, T., Jain, A. K., Joetzjer, E., Kadono, K., Kato, E., Kitidis, V., Korsbakken, J. I., Landschützer, P., Lefèvre, N., Lenton, A., Lienert, S., Liu, Z., Lombardozzi, D., Marland, G., Metzl, N., Munro, D. R., Nabel, J. E. M. S., Nakaoka, S.-I., Niwa, Y., O’Brien, K., Ono, T., Palmer, P. I., Pierrot, D., Poulter, B., Resplandy, L., Robertson, E., Rödenbeck, C., Schwinger, J., Séférian, R., Skjelvan, I., Smith, A. J. P., Sutton, A. J., Tanhua, T., Tans, P. P., Tian, H., Tilbrook, B., van der Werf, G., Vuichard, N., Walker, A. P., Wanninkhof, R., Watson, A. J., Willis, D., Wiltshire, A. J., Yuan, W., Yue, X., and Zaehle, S.: Global Carbon Budget 2020, *Earth System Science Data*, 12, 3269–3340, <https://doi.org/10.5194/essd-12-3269-2020>, <https://essd.copernicus.org/articles/12/3269/2020/>, 2020.
- Fujisaki, K., Perrin, A.-S., Desjardins, T., Bernoux, M., Balbino, L. C., and Brossard, M.: From Forest to Cropland and Pasture Systems: A Critical Review of Soil Organic Carbon Stocks Changes in Amazonia, *Global Change Biology*, 21, 2773–2786, <https://doi.org/10.1111/gcb.12906>, 2015.
- Guo, L., Deng, M., Yang, S., Liu, W., Wang, X., Wang, J., and Liu, L.: The coordination between leaf and fine root litter decomposition and the difference in their controlling factors, *Global Ecology and Biogeography*, 30, 2286–2296, <https://doi.org/https://doi.org/10.1111/geb.13384>, <https://onlinelibrary.wiley.com/doi/abs/10.1111/geb.13384>, 2021.
- Hansis, E., Davis, S. J., and Pongratz, J.: Relevance of Methodological Choices for Accounting of Land Use Change Carbon Fluxes, *Global Biogeochemical Cycles*, 29, 1230–1246, <https://doi.org/10.1002/2014GB004997>, 2015.
- Harris, I., Osborn, T. J., Jones, P., and Lister, D.: Version 4 of the CRU TS Monthly High-Resolution Gridded Multivariate Climate Dataset, *Scientific Data*, 7, 109, <https://doi.org/10.1038/s41597-020-0453-3>, 2020.
- Hengl, T., de Jesus, J. M., Heuvelink, G. B. M., Gonzalez, M. R., Kilibarda, M., Blagotic, A., Shangguan, W., Wright, M. N., Geng, X., Bauer-Marschallinger, B., Guevara, M. A., Vargas, R., MacMillan, R. A., Batjes, N. H., Leenaars, J. G. B., Ribeiro, E., Wheeler, I., Mantel, S., and Kempen, B.: SoilGrids250m: Global Gridded Soil Information Based on Machine Learning, *Plos One*, 12, e0169748, <https://doi.org/10.1371/journal.pone.0169748>, 2017.
- Hergoualc’h, Kristell, Akiyama, Hiroko, Bernoux, Martial, Chirinda, Ngonidzashe, Del Prado, Agustin, Kasimir, Åsa, MacDonald, Douglas, Ogle, Stephen M., Regina, Kristiina, van der Weerden, TonyHergoualc’h, Kristell, Akiyama, Hiroko, Bernoux, Martial, Chirinda, Ngonidzashe, Del Prado, Agustin, Kasimir, Åsa, MacDonald, Douglas, Ogle, Stephen M., Regina, Kristiina, and van der Weerden, Tony: N<sub>2</sub>O Emissions from Managed Soils, and CO<sub>2</sub> Emissions from Lime and Urea Application, in: IPCC 2019, 2019 Refinement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories., Published: IPCC, Switzerland., 2019.
- Herzfeld, T., Heinke, J., Rolinski, S., and Müller, C.: Soil organic carbon dynamics from agricultural management practices under climate change, *Earth System Dynamics*, 12, 1037–1055, <https://doi.org/10.5194/esd-12-1037-2021>, <https://esd.copernicus.org/articles/12/1037/2021/>, 2021.
- Houghton, R. A., House, J. I., Pongratz, J., van der Werf, G. R., DeFries, R. S., Hansen, M. C., Le Quéré, C., and Ramankutty, N.: Carbon Emissions from Land Use and Land-Cover Change, *Biogeosciences*, 9, 5125–5142, <https://doi.org/10.5194/bg-9-5125-2012>, 2012.



- 885 Humpenöder, F., Popp, A., Bodirsky, B. L., Weindl, I., Biewald, A., Lotze-Campen, H., Dietrich, J. P., Klein, D., Kreidenweis, U., Müller, C., Rolinski, S., and Stevanovic, M.: Large-Scale Bioenergy Production: How to Resolve Sustainability Trade-Offs?, *Environmental Research Letters*, 13, 024011, <https://doi.org/10.1088/1748-9326/aa9e3b>, 2018.
- Hurt, G. C., Chini, L., Sahajpal, R., Frohling, S., Bodirsky, B. L., Calvin, K., Doelman, J. C., Fisk, J., Fujimori, S., Goldewijk, K. K., Hasegawa, T., Havlik, P., Heinemann, A., Humpenöder, F., Jungclaus, J., Kaplan, J., Kennedy, J., Kristzin, T., Lawrence, D., Lawrence, P., 890 Ma, L., Mertz, O., Pongratz, J., Popp, A., Poulter, B., Riahi, K., Shevliakova, E., Stehfest, E., Thornton, P., Tubiello, F. N., van Vuuren, D. P., and Zhang, X.: Harmonization of Global Land-Use Change and Management for the Period 850–2100 (LUH2) for CMIP6, *Geoscientific Model Development Discussions*, pp. 1–65, <https://doi.org/10.5194/gmd-2019-360>, 2020.
- Iizumi, T., Kim, W., and Nishimori, M.: Modeling the Global Sowing and Harvesting Windows of Major Crops Around the Year 2000, *Journal of Advances in Modeling Earth Systems*, 11, 99–112, <https://doi.org/10.1029/2018MS001477>, 2019.
- 895 Jägermeyr, J., Gerten, D., Heinke, J., Schaphoff, S., Kummer, M., and Lucht, W.: Water Savings Potentials of Irrigation Systems: Global Simulation of Processes and Linkages, *Hydrology and Earth System Sciences*, 19, 3073–3091, <https://doi.org/10.5194/hess-19-3073-2015>, 2015.
- Jain, N., Bhatia, A., and Pathak, H.: Emission of Air Pollutants from Crop Residue Burning in India, *Aerosol and Air Quality Research*, 14, 422–430, <https://doi.org/10.4209/aaqr.2013.01.0031>, 2014.
- 900 Karstens, K.: Model output data of the paper: "Management induced changes of soil organic carbon on global croplands", <https://doi.org/10.5281/zenodo.4320663>, <https://doi.org/10.5281/zenodo.4320663>, 2020a.
- Karstens, K.: Result and paper preparation scripts of the Paper: Management induced changes of soil organic carbon on global croplands, <https://github.com/k4rst3ns/historicalsocbudget>, 2020b.
- Karstens, K. and Dietrich, J. P.: mrsoil: MadRat Soil Organic Carbon Budget Library, <https://doi.org/10.5281/zenodo.4317933>, <https://github.com/pik-piam/mrsoil>, r package version 1.1.0, 2020.
- 905 Keel, S. G., Leifeld, J., Mayer, J., Taghizadeh-Toosi, A., and Olesen, J. E.: Large Uncertainty in Soil Carbon Modelling Related to Method of Calculation of Plant Carbon Input in Agricultural Systems, *European Journal of Soil Science*, 68, 953–963, <https://doi.org/10.1111/ejss.12454>, 2017.
- Lal, R.: World Cropland Soils as a Source or Sink for Atmospheric Carbon, vol. 71, pp. 145–191, Academic Press, 910 [https://doi.org/10.1016/S0065-2113\(01\)71014-0](https://doi.org/10.1016/S0065-2113(01)71014-0), 2001.
- Lasco, R. D., Ogle, Stephen, Raison, John, Verchot, Louis, Wassmann, Reiner, Yagi, Kazuyuki, Bhattacharya, Sumana, Brenner, John S., Daka, Julius Partson, González, Sergio P., Krug, Thelma, Li, Yue, Martino, Daniel L., McConkey, Brian G., Smith, Pete, Tyler, Stanley C., and Zhakata, Washington: Cropland, in: IPCC 2006, 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Prepared by the National Greenhouse Gas Inventories Programme, Published: IGES, Japan., 2006.
- 915 Lindeskog, M., Arneth, A., Bondeau, A., Waha, K., Seaquist, J., Olin, S., and Smith, B.: Implications of Accounting for Land Use in Simulations of Ecosystem Carbon Cycling in Africa, *Earth System Dynamics*, 4, 385–407, <https://doi.org/10.5194/esd-4-385-2013>, 2013.
- Lorenz, K. and Lal, R.: Soil Organic Carbon Sequestration in Agroforestry Systems. A Review, *Agronomy for Sustainable Development*, 34, 443–454, <https://doi.org/10.1007/s13593-014-0212-y>, 2014.
- Lutz, F., Herzfeld, T., Heinke, J., Rolinski, S., Schaphoff, S., von Bloh, W., Stoorvogel, J. J., and Müller, C.: Simulating the Effect of 920 Tillage Practices with the Global Ecosystem Model LPJmL (Version 5.0-Tillage), *Geoscientific Model Development*, 12, 2419–2440, <https://doi.org/10.5194/gmd-12-2419-2019>, 2019.

- Minasny, B., Malone, B. P., McBratney, A. B., Angers, D. A., Arrouays, D., Chambers, A., Chaplot, V., Chen, Z.-S., Cheng, K., Das, B. S., Field, D. J., Gimona, A., Hedley, C. B., Hong, S. Y., Mandal, B., Marchant, B. P., Martin, M., McConkey, B. G., Mulder, V. L., O'Rourke, S., Richer-de-Forges, A. C., Odeh, I., Padarian, J., Paustian, K., Pan, G., Poggio, L., Savin, I., Stolbovoy, V., Stockmann, U., Sulaeman, Y., Tsui, C.-C., Vågen, T.-G., van Wesemael, B., and Winowiecki, L.: Soil Carbon 4 per Mille, *Geoderma*, 292, 59–86, <https://doi.org/10.1016/j.geoderma.2017.01.002>, 2017.
- Minoli, S., Egli, D. B., Rolinski, S., and Müller, C.: Modelling Cropping Periods of Grain Crops at the Global Scale, *Global and Planetary Change*, 174, 35–46, <https://doi.org/10.1016/j.gloplacha.2018.12.013>, 2019.
- Morais, T. G., Teixeira, R. F. M., and Domingos, T.: Detailed Global Modelling of Soil Organic Carbon in Cropland, Grassland and Forest Soils, *PLOS ONE*, 14, e0222604, <https://doi.org/10.1371/journal.pone.0222604>, 2019.
- NREL: CENTURY model 4.0, <https://www2.nrel.colostate.edu/projects/century/obtain2.htm>, (accessed December 12, 2020), 2000.
- Ogle, S. M., Wakelin, Stephen J., Buendia, Leandro, McConkey, Brian, Baldock, Jeffrey, Akiyama, Hiroko, Kishimoto-Mo, Ayaka W., Chirinda, Ngonidzashe, Bernoux, Martial, Bhattacharya, Sumana, Chuersuan, Nares, Goheer, Muhammad Arif Rashid, Hergoualc'h, Kristell, Ishizuka, Shigehiro, Lasco, Rodel D., Pan, Xuebiao, Pathak, Himanshu, Regina, Kristiina, Sato, Atsushi, Vazquez-Amabile, Gabriel, Wang, Changke, and Zheng, Xunhua: Cropland, in: IPCC 2019, 2019 Refinement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories., Published: IPCC, Switzerland., 2019.
- Parton, W., Schimel, D., Cole, C., and Ojima, D.: Analysis of Factors Controlling Soil Organic-Matter Levels in Great-Plains Grasslands, *Soil Science Society of America Journal*, 51, 1173–1179, <https://doi.org/10.2136/sssaj1987.03615995005100050015x>, 1987.
- Pellegrini, P. and Fernández, R. J. a. a.: Crop Intensification, Land Use, and on-Farm Energy-Use Efficiency during the Worldwide Spread of the Green Revolution, *Proceedings of the National Academy of Sciences*, 115, 2335–2340, <https://doi.org/10.1073/pnas.1717072115>, 2018.
- Poeplau, C. and Don, A.: Carbon Sequestration in Agricultural Soils via Cultivation of Cover Crops – A Meta-Analysis, *Agriculture, Ecosystems & Environment*, 200, 33–41, <https://doi.org/10.1016/j.agee.2014.10.024>, 2015.
- Poggio, L., de Sousa, L. M., Batjes, N. H., Heuvelink, G. B. M., Kempen, B., Ribeiro, E., and Rossiter, D.: SoilGrids 2.0: Producing Soil Information for the Globe with Quantified Spatial Uncertainty, *SOIL*, 7, 217–240, <https://doi.org/10.5194/soil-7-217-2021>, 2021.
- Pongratz, J., Reick, C. H., Houghton, R. A., and House, J. I.: Terminology as a Key Uncertainty in Net Land Use and Land Cover Change Carbon Flux Estimates, *Earth System Dynamics*, 5, 177–195, <https://doi.org/10.5194/esd-5-177-2014>, 2014.
- Popp, A., Humpenöder, F., Weindl, I., Bodirsky, B. L., Bonsch, M., Lotze-Campen, H., Müller, C., Biewald, A., Rolinski, S., Stevanovic, M., and Dietrich, J. P.: Land-Use Protection for Climate Change Mitigation, *Nature Climate Change*, 4, 1095–1098, <https://doi.org/10.1038/nclimate2444>, 2014.
- Popp, A., Calvin, K., Fujimori, S., Havlik, P., Humpenöder, F., Stehfest, E., Bodirsky, B. L., Dietrich, J. P., Doelmann, J. C., Gusti, M., Hasegawa, T., Kyle, P., Obersteiner, M., Tabeau, A., Takahashi, K., Valin, H., Waldhoff, S., Weindl, I., Wise, M., Kriegler, E., Lotze-Campen, H., Fricko, O., Riahi, K., and van Vuuren, D. P.: Land-Use Futures in the Shared Socio-Economic Pathways, *Global Environmental Change*, 42, 331–345, <https://doi.org/10.1016/j.gloenvcha.2016.10.002>, 2016.
- Portmann, F. T., Siebert, S., and Döll, P.: MIRCA2000—Global Monthly Irrigated and Rainfed Crop Areas around the Year 2000: A New High-Resolution Data Set for Agricultural and Hydrological Modeling, *Global Biogeochemical Cycles*, 24, <https://doi.org/10.1029/2008GB003435>, 2010.
- Porwollik, V., Rolinski, S., Heinke, J., and Müller, C.: Generating a Global Gridded Tillage Dataset, *Earth System Science Data Discussions*, pp. 1–28, <https://doi.org/10.5194/essd-2018-152>, 2018.

- 960 Porwollik, V., Rolinski, S., Heinke, J., and Müller, C.: Generating a Rule-Based Global Gridded Tillage Dataset, *Earth System Science Data*, 11, 823–843, <https://doi.org/10.5194/essd-11-823-2019>, 2019.
- Porwollik, V., Rolinski, S., Heinke, J., von Bloh, W., Schaphoff, S., and Müller, C.: The role of cover crops for cropland soil carbon, nitrogen leaching, and agricultural yields – a global simulation study with LPJmL (V. 5.0-tillage-cc), *Biogeosciences*, 19, 957–977, <https://doi.org/10.5194/bg-19-957-2022>, <https://bg.copernicus.org/articles/19/957/2022/>, 2022.
- 965 Powlson, D. S., Stirling, C. M., Jat, M. L., Gerard, B. G., Palm, C. A., Sanchez, P. A., and Cassman, K. G.: Limited Potential of No-till Agriculture for Climate Change Mitigation, *Nature Climate Change*, 4, 678–683, <https://doi.org/10.1038/NCLIMATE2292>, 2014.
- Prestele, R., Hirsch, A. L., Davin, E. L., Seneviratne, S. I., and Verburg, P. H.: A Spatially Explicit Representation of Conservation Agriculture for Application in Global Change Studies, *Global Change Biology*, 24, 4038–4053, <https://doi.org/10.1111/gcb.14307>, 2018.
- Pugh, T. A. M., Arneth, A., Olin, S., Ahlström, A., Bayer, A. D., Goldewijk, K. K., Lindeskog, M., and Schurgers, G.: Simulated Carbon  
970 Emissions from Land-Use Change Are Substantially Enhanced by Accounting for Agricultural Management, *Environmental Research Letters*, 10, 124 008, <https://doi.org/10.1088/1748-9326/10/12/124008>, 2015.
- Rahman, M. M., Tsukamoto, J., Rahman, M. M., Yoneyama, A., and Mostafa, K. M.: Lignin and its effects on litter decomposition in forest ecosystems, *Chemistry and Ecology*, 29, 540–553, <https://doi.org/10.1080/02757540.2013.790380>, <https://doi.org/10.1080/02757540.2013.790380>, 2013.
- 975 Ray, D. K., Ramankutty, N., Mueller, N. D., West, P. C., and Foley, J. A.: Recent Patterns of Crop Yield Growth and Stagnation, *Nature Communications*, 3, 1293, <https://doi.org/10.1038/ncomms2296>, 2012.
- Robinson, T. P., Wint, G. R. W., Conchedda, G., Boeckel, T. P. V., Ercoli, V., Palamara, E., Cinardi, G., D’Aietti, L., Hay, S. I., and Gilbert, M.: Mapping the Global Distribution of Livestock, *PLOS ONE*, 9, e96 084, <https://doi.org/10.1371/journal.pone.0096084>, 2014.
- Rogelj, J., Popp, A., Calvin, K. V., Luderer, G., Emmerling, J., Gernaat, D., Fujimori, S., Strefler, J., Hasegawa, T., Marangoni, G., Krey,  
980 V., Krieglner, E., Riahi, K., van Vuuren, D. P., Doelman, J., Drouet, L., Edmonds, J., Fricko, O., Harmsen, M., Havlík, P., Humpenöder, F., Stehfest, E., and Tavoni, M.: Scenarios towards Limiting Global Mean Temperature Increase below 1.5 °C, *Nature Climate Change*, 8, 325–332, <https://doi.org/10.1038/s41558-018-0091-3>, 2018.
- Rudel, T. K., Schneider, L., Uriarte, M., Turner, B. L., DeFries, R., Lawrence, D., Geoghegan, J., Hecht, S., Ickowitz, A., Lambin, E. F., Birkenholtz, T., Baptista, S., and Grau, R.: Agricultural Intensification and Changes in Cultivated Areas, 1970–2005, *Proceedings of the  
985 National Academy of Sciences*, 106, 20 675–20 680, <https://doi.org/10.1073/pnas.0812540106>, 2009.
- Sanderman, J., Hengl, T., and Fiske, G. J.: Soil Carbon Debt of 12,000 Years of Human Land Use, *Proceedings of the National Academy of Sciences*, 114, 9575–9580, <https://doi.org/10.1073/pnas.1706103114>, 2017.
- Schaphoff, S., Forkel, M., Müller, C., Knauer, J., von Bloh, W., Gerten, D., Jägermeyr, J., Lucht, W., Rammig, A., Thonicke, K., and Waha, K.: LPJmL4 – a Dynamic Global Vegetation Model with Managed Land – Part 2: Model Evaluation, *Geoscientific Model Development*,  
990 11, 1377–1403, <https://doi.org/10.5194/gmd-11-1377-2018>, 2018a.
- Schaphoff, S., von Bloh, W., Rammig, A., Thonicke, K., Biemans, H., Forkel, M., Gerten, D., Heinke, J., Jägermeyr, J., Knauer, J., Langerwisch, F., Lucht, W., Müller, C., Rolinski, S., and Waha, K.: LPJmL4 – a Dynamic Global Vegetation Model with Managed Land – Part 1: Model Description, *Geoscientific Model Development*, 11, 1343–1375, <https://doi.org/10.5194/gmd-11-1343-2018>, 2018b.
- Smil, V.: Crop Residues: Agriculture’s Largest Harvest: Crop Residues Incorporate More than Half of the World’s Agricultural Phytomass,  
995 *BioScience*, 49, 299–308, <https://doi.org/10.2307/1313613>, 1999a.
- Smil, V.: Nitrogen in Crop Production: An Account of Global Flows, *Global Biogeochemical Cycles*, 13, 647–662, <https://doi.org/10.1029/1999GB900015>, 1999b.

- Smith, J., Gottschalk, P., Bellarby, J., Chapman, S., Lilly, A., Towers, W., Bell, J., Coleman, K., Nayak, D., Richards, M., Hillier, J., Flynn, H., Wattenbach, M., Aitkenhead, M., Yeluripati, J., Farmer, J., Milne, R., Thomson, A., Evans, C., Whitmore, A., Falloon, P., and Smith, P.: Estimating Changes in Scottish Soil Carbon Stocks Using ECOSSE. I. Model Description and Uncertainties, *Climate Research*, 45, 179–192, <https://doi.org/10.3354/cr00899>, 2010.
- Smith, P.: Soil Carbon Sequestration and Biochar as Negative Emission Technologies, *Global Change Biology*, 22, 1315–1324, <https://doi.org/10.1111/gcb.13178>, 2016.
- Smith, P., Soussana, J.-F., Angers, D., Schipper, L., Chenu, C., Rasse, D. P., Batjes, N. H., van Egmond, F., McNeill, S., Kuhnert, M., Arias-Navarro, C., Olesen, J. E., Chirinda, N., Fornara, D., Wollenberg, E., Alvaro-Fuentes, J., Sanz-Cobena, A., and Klumpp, K.: How to Measure, Report and Verify Soil Carbon Change to Realize the Potential of Soil Carbon Sequestration for Atmospheric Greenhouse Gas Removal, *Global Change Biology*, 26, 219–241, <https://doi.org/10.1111/gcb.14815>, 2020.
- Sörgel, B., Kriegler, E., Weindl, I., Rauner, S., Dirnaichner, A., Ruhe, C., Hofmann, M., Bauer, N., Bertram, C., Bodirsky, B. L., et al.: A sustainable development pathway for climate action within the UN 2030 Agenda, *Nature Climate Change*, 11, 656–664, 2021.
- Strassmann, K. M., Joos, F., and Fischer, G.: Simulating Effects of Land Use Changes on Carbon Fluxes: Past Contributions to Atmospheric CO<sub>2</sub> Increases and Future Commitments Due to Losses of Terrestrial Sink Capacity, *Tellus B: Chemical and Physical Meteorology*, 60, 583–603, <https://doi.org/10.1111/j.1600-0889.2008.00340.x>, 2008.
- Taghizadeh-Toosi, A., Christensen, B. T., Hutchings, N. J., Vejlin, J., Kätterer, T., Glendining, M., and Olesen, J. E.: C-TOOL: A Simple Model for Simulating Whole-Profile Carbon Storage in Temperate Agricultural Soils, *Ecological Modelling*, 292, 11–25, <https://doi.org/10.1016/j.ecolmodel.2014.08.016>, 2014.
- van der Werf, G. R., Randerson, J. T., Giglio, L., van Leeuwen, T. T., Chen, Y., Rogers, B. M., Mu, M., van Marle, M. J. E., Morton, D. C., Collatz, G. J., Yokelson, R. J., and Kasibhatla, P. S.: Global Fire Emissions Estimates during 1997–2016, *Earth System Science Data*, 9, 697–720, <https://doi.org/10.5194/essd-9-697-2017>, 2017.
- von Bloh, W., Schaphoff, S., Müller, C., Rolinski, S., Waha, K., and Zaehle, S.: Implementing the nitrogen cycle into the dynamic global vegetation, hydrology, and crop growth model LPJmL (version 5.0), *Geoscientific Model Development*, 11, 2789–2812, <https://doi.org/10.5194/gmd-11-2789-2018>, <https://gmd.copernicus.org/articles/11/2789/2018/>, 2018.
- Waha, K., Dietrich, J. P., Portmann, F. T., Siebert, S., Thornton, P. K., Bondeau, A., and Herrero, M.: Multiple Cropping Systems of the World and the Potential for Increasing Cropping Intensity, *Global Environmental Change*, 64, 102 131, <https://doi.org/10.1016/j.gloenvcha.2020.102131>, 2020.
- Wang, Z., Hoffmann, T., Six, J., Kaplan, J., Govers, G., Doetterl, S., and Oost, K.: Human-Induced Erosion Has Offset One-Third of Carbon Emissions from Land Cover Change, *Nature Climate Change*, 7, <https://doi.org/10.1038/nclimate3263>, 2017.
- Weindl, I., Popp, A., Bodirsky, B. L., Rolinski, S., Lotze-Campen, H., Biewald, A., Humpenöder, F., Dietrich, J. P., and Stevanović, M.: Livestock and Human Use of Land: Productivity Trends and Dietary Choices as Drivers of Future Land and Carbon Dynamics, *Global and Planetary Change*, 159, 1–10, <https://doi.org/10.1016/j.gloplacha.2017.10.002>, 2017.
- Williams, T. O., Fernández-Rivera, S., and Kelley, T. G.: The Influence of Socioeconomic Factors on the Availability and Utilization of Crop Residues as Animal Feeds, CAB International in association with the ICRISAT and ILRI, Oxon, 1997.
- Woolf, D., Amonette, J. E., Street-Perrott, F. A., Lehmann, J., and Joseph, S.: Sustainable Biochar to Mitigate Global Climate Change, *Nature Communications*, 1, 56, <https://doi.org/10.1038/ncomms1053>, 2010.

Zhang, B., Tian, H., Lu, C., Dangal, S. R. S., Yang, J., and Pan, S.: Global Manure Nitrogen Production and Application in Crop-  
land during 1860–2014: A 5 Arcmin Gridded Global Dataset for Earth System Modeling, *Earth System Science Data*, 9, 667–678,  
1035 <https://doi.org/10.5194/essd-9-667-2017>, 2017.