



Interactive comment on “A reactive nitrogen budget of the Bohai Sea based on an isotope mass balance model” by Shichao Tian et al.

Shichao Tian et al.

shichao.tian@uni-hamburg.de

Received and published: 15 April 2021

Referee's comments: Overall the topic is of interest and fits well to the scope of the journal Biogeosciences. Moreover, a budget using the LOICZ approach was constructed using measurements but mainly data from other studies. The budget balances sources and sinks of nitrate and is supported by stable isotope data of nitrate. As a central conclusion the overwhelming role of nitrification as the major source of nitrate is presented. Authors' reply: We thank the referee for his/her ideas on improving the manuscript and appreciate the careful review. In our opinion, the reviewer may not have appreciated the power of combined mass- and isotope budgets, which add a completely new dimension to standard mass-only budgets due to the process-specific isotope fractionation

C1

in the nitrogen cycle. This and an apparently poor choice of title on our side may have led to misunderstanding, which we hope to remedy in a revision. We address this point and others raised by the referee below.

Referee's comments: The most significant error is the definition of equations (1) and (2). Sources and sinks are listed and supposed to be balanced. However, the source terms list nitrification and the loss terms list sedimentation. Both of these do not fulfill the criteria of a source or sink, respectively. Nitrification is neither a source nor a sink for nitrate but simply a microbial process that converts ammonium via nitrite to nitrate. Nitrification does not generate new DIN for a system simply because the substrate of the nitrification process is ammonium and comes from internal turnover processes of organic matter. The LOICZ report (no 5 LOICZ BIOGEOCHEMICAL Modelling Guidelines, 1996) states “The important point to note with this reaction (nitrification) is that carbon and phosphorus are not directly involved in the net reaction. Again this makes the point that the relationship between NO₃ and NH₄ may be considered an “internal cycle” which need not be dealt with directly.” Authors' reply: We are sorry that our approach is apparently open to misunderstanding, and indeed will change the title of our manuscript to “A nitrate budget of the Bohai Sea based on an isotope mass balance model” in order to make our focus on reactive nitrogen clearer. Because anthropogenic impacts on biogeochemical cycles of marginal seas is always seen in amplified inputs of reactive N, we focus on this cycle. The budget in the manuscript thus is basically a budget of nitrate ion in the water mass, neither organic particles nor the sediment are included as active compartments. Instead, we expand the mass-based budget with an isotope-based budget to employ the added possibilities of dual nitrate isotopes to quantify nitrate sources and sinks. That is why nitrification and sedimentation are considered as sources and sinks, because they affect the isotope budget. The “LOICZ approach” that the reviewer refers to links water- and salt-balance to construct carbon, nitrogen and phosphorus budget models and is an established methodology to standardize mass flux estimates of these biogeochemically important elements in coastal systems on regional to local scales (D.C. Gordon et al., 1996; Smith et al., 2005). The

C2

underlying box model approach diagnoses water, salt and CNP-fluxes, for example to decide if systems are autotrophic (production exceeds respiration) or heterotrophic (respiration exceeds production) on the basis of deviations from stoichiometric Redfield ratios. We do not aim to decide if Bohai Sea nutrient cycles indicate net autotrophy or net heterotrophy, and thus specifically do not aim to include the carbon balance associated with the LOICZ approach. We establish a more generic box model approach specifically of the reactive nitrogen pools to provide mass flux estimates of inputs and outputs to the nitrate pool. Our approach thus goes beyond a mass flux estimate (as done in previous LOICZ-type budgets for Bohai Sea, e.g., Zhang et al. (2004)) by constraining some of the branches of reactive nitrogen in this coastal sea based on the tell-tale changes in dual isotope composition of nitrate, and the dual isotopic properties of sources and sinks. This approach thus differs fundamentally from the solely mass-balance approach that have been done previously and that have large uncertainties specifically in the internal nitrogen turnover. In our opinion the approach taken in our study is significantly more specific and is diagnostic of several important pathways of reactive nitrogen. Nitrification is indeed an "internal" source, as the referee says: "nitrification does not generate new DIN for a system simply because the substrate of the nitrification process is ammonium and comes from internal turnover processes of organic matter", the initial source of newly nitrified nitrate is ammonium or organic matter which releases ammonium. However, nitrification affects the stable isotopic ratios and therefore, has to be included into our budget. For our study, N in organic matter is not involved in our box model because it is simply not present as nitrate ions.

Interactive comment on Biogeosciences Discuss., <https://doi.org/10.5194/bg-2020-471>, 2020.