

Long distance particle transport to the central Ionian Sea

Léo Berline¹, Andrea Michelangelo Doglioli¹, Anne Petrenko¹, Stéphanie Barrillon¹, Boris Espinasse², Frederic A.C. Le Moigne¹, François Simon-Bot¹, Melilotus Thyssen¹, François Carlotti¹

5

¹Aix Marseille Université, Université de Toulon, CNRS, IRD, Mediterranean Institute of Oceanography (MIO), 13288, Marseille, France

²Department of Arctic and Marine Biology, UiT The Arctic University of Norway, Tromsø, Norway

10 *Correspondence to:* Leo Berline (leo.berline@mio.osupytheas.fr)

Abstract

15 ~~Together with T-S properties, particle abundance *in situ* measurements are useful to discriminate water masses and derive circulation.~~In the upper layers of the Ionian Sea, young Mediterranean Atlantic Waters (MAW) flowing eastward from the Sicily channel meet old MAW.~~In the upper layers of the Ionian Sea, the fresher Atlantic Waters (AW) recently crossing the Sicily channel meet the resident and saltier AW which have cyclonically circulated in the Eastern basin and come back modified after evaporation and eventually cooling.~~ In May 2017, during the PEACETIME cruise, fluorescence and particle content abundance sampled at high resolution revealed unexpected heterogeneity in the central Ionian. Surface salinity measurements, together with altimetry-derived and hull-mounted ADCP currents, describe a zonal pathway of AW entering the Ionian Sea, consistent with the so-called cyclonic mode in the North Ionian Gyre. The ION-Tr transect, located ~19-20 20°E- ~36°N turned out to be at the crossroad of three water masses, mostly coming from the west, north and from an isolated anticyclonic eddy northeast of ION-Tr. Using Lagrangian numerical simulations, we suggest that the contrast in particle loads along ION-Tr originates from particles transported from these three different water masses. Waters from the west, identified as young AW carried by a strong southwestward jet, were intermediate moderate in particle load, probably originating from the Sicily channel. Water mass originating from the north was carrying abundant particles, probably originating from northern Ionian, or further from the south Adriatic. Waters from the eddy, depleted in particles and Chl-a may originate from south of Peloponnese, where the Pelops eddy forms.

25
30 The central Ionian Sea hence appears as a mosaic area, where waters of contrasted biological history meet. This contrast is particularly clear in spring, when blooming and non-blooming areas co-occur. ~~Particle abundance *in situ* measurements are useful to discriminate water masses and derive circulation, together with T-S properties.~~

Interpreting the complex dynamics of physical-biogeochemical coupling from discrete measurements made at isolated stations at sea is a **big** challenge. The combination of multiparametric *in situ* measurements at high resolution with remote sensing and Lagrangian modeling appears as one proper way to address this challenge.

1 Introduction

35

The study of particle abundance, size structure and dynamics is a key to understand the marine ecosystem, from primary and secondary production, export and remineralization all the way through food-web dynamics (McDonnell et al 2015, Giering et al 2020). In the semi-enclosed Mediterranean Sea, the main particles sources are the biological production in the open ocean and over continental shelves, rivers discharge, atmospheric deposition as well as sediment resuspension. From West to East and North to South, the Mediterranean Sea harbours areas of very distinct productivity, going from eutrophic (e.g. in the Alboran Sea) to ultra oligotrophic (e.g. in the Levantine basin, Moutin et al 2017). This heterogeneity is further developed by the complex general and mesoscale circulation that favor exchanges between these contrasted areas, leading to a fine scale spatial distribution of biogeochemical properties and particles (Karageorgis et al 2008, Durrieu de Madron et al, 1992, Rousselet et al 2019).

40

45

Located in the center of the Mediterranean Sea, the Ionian Sea is a cross road where three different surface water masses meet: i) recent Atlantic water (AW) flowing from the Sicily channel ii) older AW flowing from the East after its cyclonic circulation in the Levantine basin and iii) fresher (Po influenced) water coming from the Adriatic Sea (Malanotte-Rizzoli et al 1997). Circulation in the North Ionian is strongly variable, as is the path of AW spreading. The most stable branch of the Atlantic Ionian Stream entering the Ionian Sea follows a rather direct pathway to the SouthEast, connecting the Sicily Channel with the Cretan Passage (Mid Ionian Jet, MIJ, Menna et al 2019). South of the MIJ, circulation pattern is stable and anticyclonic, known also as the Syrte Gyre (Pinardi et al, 2015). North of the MIJ, two alternate circulation states have been proposed, the anticyclonic and cyclonic mode of the so called North Ionian Gyre (NIG, Gačić et al 2010). In addition, semi-permanent mesoscale gyres are observed in the western part (Maltese Channel Crest, Medina Gyre, Messina Rise Vortex) as well as in the eastern part (Pelops Gyre), some of them triggered by wind. Although the AW circulation has been documented according to seasons and NIG modes (Malanotte-Rizzoli et al 1997, Gačić et al 2011, Menna et al 2019), the fine-scale pathways of AW crossing the Ionian Sea have only seldom been sampled. The mode of the NIG has a strong influence on the dispersal of water masses and properties in the Ionian basin, also impacting its productivity (Lavigne et al 2018).

50

55

60

The Ionian Sea is generally considered oligotrophic (Boldrin et al 2002), with a north-south gradient of Chl-a (d'Ortenzio et Ribera d'Alcala 2009). However, it is not homogeneous, as distinct phytoplankton communities associated to the main water masses have been described (Casotti et al 2003). Three main phytoplankton communities were associated with water coming

from Adriatic in the north west, water from the Eastern Mediterranean in the north east and AW from the Sicily channel to the south. Zooplankton community was also contrasted between northwestern and eastern Ionian (Mazzocchi et al 2003).

65 Particle distribution in the Ionian Sea is less described. The particle concentration in the water column can be measured with several instruments. Discrete sampling is carried out using bottles or pumps followed by filtration, allowing bulk mass measurements (McDonnell et al 2015). Continuous sampling can be carried out using optical measurements that give either bulk measurements (transmissometer, backscatter (Briggs et al 2013)) or size-resolved measurements: Laser In Situ Scattering and Transmissometry- LISST, (Karageorgis et al 2012), Underwater Vision Profiler – UVP (Picheral et al 2010), Laser Optical Particle Counter – LOPC, (Herman, 2004) for the most common instruments, see Giering et al (2020) for a
70 comprehensive review. In the Ionian Sea, few data are available on the horizontal distribution of particles. These data have a coarse horizontal resolution and are mostly from transmissometry, water filtration (Rabitti et al 1994, Boldrin et al 2002, Karageorgis et al 2008, 2012) while few data are from optical devices (Karageorgis et al 2012, Ramondenc et al 2016). Here we used the LOPC, providing size resolved abundances over the size range [100-2500] μm . This instrument was mounted on a free-fall MVP fish, allowing high resolution horizontal and vertical sampling of particles and water mass properties along
75 the ship track, to investigate their joint distribution in the Central Ionian Sea.

The objective of this paper is to document particle distribution in the central Ionian Sea, based on a high-resolution multi-parametric transect. In particular we analyse particle distribution with regard to water mass properties and transport, to propose scenarios of histories of these waters masses and discuss implications for the functioning of the Ionian Sea ecosystem.

80 2 Material and Methods

The PEACETIME cruise (<http://peacetime-project.org/>, Guieu et al 2020) was conducted during late spring conditions from May 10th to June 11th, 2017 on board the R/V *Pourquoi Pas ?*. The overall aim of PEACETIME was to investigate the role of atmospheric dust deposition on biogeochemical fluxes and pelagic ecosystem functioning. Along the 4300 km transect covering western and eastern Mediterranean basin, 10 short stations (with an average duration of 8 hours) and 3 long stations
85 (duration of 4-5 days) were performed. The ship entered the Ionian Sea on May 22nd and the last station (ST8) in the region was occupied on May 30th as detailed in table 1. SAV station was planned to deploy a float, which explains its distance with other stations.

At stations, Rosette casts were carried out (0-500 and 0-bottom). The "classical" Rosette was composed of a CTD (pressure, temperature and salinityconductivity of seawater, dissolved oxygen concentration, photosynthetically active radiation (PAR),
90 beamlight transmission (at 650 nm) (Karageorgis et al 2008), chlorophyll a fluorescence. Fluorescence was calibrated to report Chl-a in mg.m^{-3} . This instrumental package was also composed of a sampling system : 24 12-L Niskin bottles could be fired at specified levels during upcasts. Water from the classical Rosette was used to quantify oxygen, nutrients, Dissolved

Organic Carbon, Particulate Organic Carbon and Nitrogen and pigments using HPLC (see Guieu et al 2020; and Marañón et al 2020 for detail on HPLC).

95 2.1 High resolution vertical sampling

Between stations, transects with a Moving Vessel Profiler (MVP) were carried out. In the Ionian Sea two transects were achieved, ION-Tr and a very short ST8-Tr (figure 1). MVP200 ODIM Brooke Ocean performed vertical profiles from surface to ~300 m, at a spatial resolution of ~1 nm, with the nearly vertical free falls of the Multi Sensor Free Fall Fish type I equipped with a AML microCTD, a WetLabs ECO fluorimeter, and a ODIM Brooke Ocean Laser Optical Plankton Counter (LOPC). The T and S of the microCTD were calibrated with the rosette CTD. The LOPC recorded particle size and abundance distributions: the instrument records the cross-sectional area of each particle passing through its laser beam (Herman et al., 2004; Herman & Harvey, 2006) for a size range between 100 μ m and a few centimeters. If a particle is recorded by at least 3 diode elements, it will be recorded as a multi-element plankton (MEP), in contrast to single element plankton (SEP). LOPC also provides information about the MEPs allowing to compute an attenuation index (AI), which is essentially a normalized opacity index (an AI of 1 means a fully opaque object) although the MEP shape also has an impact (Espinasse et al 2018). The living fraction of particles can be estimated, using a calibration based on comparison with plankton net tows (Espinasse et al., 2018), as this fraction is linked with the percent of MEP with respect to the total particle count (%MEP). The sampling volume was estimated using the LOPC sampling tunnel surface multiplied by the depth increment estimated with the pressure sensor. Given the fish free-fall velocity of ~4 m/s^{-1} and the LOPC acquisition frequency (2 Hz), we binned the LOPC data over 5-m vertical bins. As the surface docking depth of the MVP fish was not constant, the first bin (0–7.5 m) was discarded from the analysis. The LOPC abundances were redistributed into log-scale size bins for easier analysis. Total abundance (e.g., abundance over the full LOPC size range) and parameters of the size distribution were computed (slope and intercept of the Normalized Biomass Size Spectra (NBSS), e.g. Espinasse et al., (2018). We thus have access to vertical profiles of total abundance of particles (particles per cubic meter) at almost each 115 kilometer during the MVP transects.

2.2 Underway measurements

The FSGThermoSalinoGraph, equipped with a SeaBird SBE21, was connected to a continuous surface water flow-through system that pumped seawater at 2-m depth. Output data were sea surface temperature and Salinity every minute all along the cruise route. Horizontal velocity was measured with a hull mounted ADCP.: RDI Ocean Surveyor 150kHz. Data treatment used CASCADE software version 7.1 (Le Bot et al 2011) and LATEXtools toolbox (Petrenko et al., 2017, Doglioli et al., 2013, <https://people.mio.osupytheas.fr/~doglioli/latextools.htm>).

2.3 Automated cytometry

Surface phytoplankton community was analysed using an automated flow cytometer (Cytosense benchtop flow cytometer from CytoBuoy b.v) installed on a dedicated continuous sampling system set up to pump surface water (at 5m depth). The Cytosense automated flow cytometer was equipped with a 120 mW, 488 nm laser beam. The volume analyzed was controlled by a calibrated peristaltic pump and each particle passed in front of the laser beam at a speed of 2 m.s⁻¹. The particle resolved size range varied from <1 µm up to 800 µm in width and several hundreds of µm in length for chain forming cells. The trigger to record a signal was based on pigment fluorescence. Stability of the optical unit was controlled before and after the cruise using fluorescing 2 µm Polyscience beads, and a set of silica beads of 1.0 , 2.0 ,3.0, 5.0, 7.0 µm. The size estimates of cells were retrieved from the silica beads FWS signature, and enabled to separate pico-nano-microphytoplankton size classes. The instrument allowed for the quantification of pico-, and nanophytoplankton populations, up to microphytoplankton when abundant enough in the 5 mL analyzed. Seven groups were resolved. One group of picoeukaryotes (mean size 2.81±0.19 µm), three groups of nanoeukaryotes (Nanoeukaryotes1, 2 and 3, mean sizes of 4.76±0.13, 7.16± 0.35 and 10.64 ± 2.18 µm respectively), Synechococcus like (1.04± 0.12 µm), Chryptophytes like (6.00 ±0.93 µm) and a coccolithophore like group (5.37± 0.14 µm). Each phytoplankton group was manually clustered thanks to the Cytoclus® (<http://www.cytobuoy.com/products/software/>) dedicated software.

2.4 Satellite data

Several remote sensing datasets were exploited using the SPASSO (Software Package for an Adaptive Satellite-based Sampling for Ocean campaigns, <https://spasso.mio.osupytheas.fr/>; last access: 2 Oct. 2020). Altimetry data was obtained from the AVISO Mediterranean regional product. Delayed-time L34 maps of SST Sat (Mediterranean Sea – Ultra High Resolution L34 SST Reprocessed; Nardelli et al., 2013; Pisano et al., 2016) and Chl Sat a (Mediterranean Sea Reprocessed Surface Chlorophyll Concentration from Multi Satellite observations) were retrieved for the period of the cruise, on regular grids of 0.014 × 0.014 ° resolution for SST Sat and of 1 × 1 km resolution for Chl Sat a from CMEMS - Copernicus Marine Environment Monitoring Service (<http://marine.copernicus.eu/>). SPASSO provided several Lagrangian diagnostics of the altimetry-derived current field, such as the Finite scale Lyapunov Exponent (FSLE) computed using the algorithm of d'Ovidio et al. (2004), in order to help in positioning the sampling transect and stations taking into account the ocean dynamics. This approach was successfully adopted during previous cruises such as LATEX (Petrenko et al, 2017), KEOPS2 (d'Ovidio et al., 2015), OUTPACE (Moutin et al, 2017).

2.5 Lagrangian numerical simulations

In order to estimate the origins of the sampled surface water along transect ION-Tr, the ARIANE package (Blanke and Raynaud, 1997) was used to predict surface trajectories of water particles, using surface geostrophic velocities derived from altimetry and wind from WRF model. The daily gridded surface geostrophic velocity fields from Copernicus Monitoring

Environment Marine Service (CMEMS) were used, with a resolution of $1/8^\circ$, using the near real time data, validated by an accurate comparison with both the hull-mounted ADCP-data and the [Surface Velocity Program \(SVP\)](#) drifters deployed during the cruise (Barrillon et al 2019). ~~The numerical particle positions were initialized in a polygon around the ship's track on May 29 (longitude:18-20.1, latitude: 35.4-35.9, $\pm 0.2^\circ$ in latitude). This polygon takes into account possible offset between in situ velocity and satellite derived geostrophic velocity.~~ Given the daily time series of the surface velocity field, the ARIANE package computes numerical particle trajectories through a backward or forward integration time. The so-called "qualitative" mode is chosen: each particle trajectory can be followed step by step through the integration time. Only surface trajectories were considered. ~~The simulated particle positions were initialized in a polygon around the ship's track on May 29th (longitude from 18° to 20.1° E, latitude from 35.4° to 35.9° N with a 0.2° widening in latitude). This polygon surrounds the actual track, to take into account a potential offset between *in situ* velocity and satellite derived geostrophic velocity.~~ 10000 particles (sufficient for a comprehensive sampling of the polygon area) randomly chosen inside the launch area were advected backward for one month. This duration was chosen as a trade-off between particle source identification, and error accumulation along the trajectories ~~and residence time in the Ionian Sea (39 days according to Celentano et al 2020).~~ The particle trajectories and final positions were analyzed to determine the main source areas.

The cruise data are available on LEFE-CYBER database (<http://www.obs-vlfr.fr/proof/php/PEACETIME/peacetime.php>).

3 Results

170 3.1 Oceanographic context

~~Considering the surface circulation over the northern Ionian Sea (figure 1), averages of satellite Chl-a and surface temperature for the region 16° to 22° E, 34° to 40° N were computed for the period before our sampling.~~ Time series of satellite Chl-a showed that surface phytoplankton concentration was decaying since mid-March (figure 2). ~~Maximum Chl-a concentration was 0.24 mg.m^{-3} in January.~~ Since the end of April, the Ionian Sea was steadily warming while Chl-a concentration remained low (0.09 mg.m^{-3}) indicating the absence of significant surface production events. On average over April-May, maximum Chl-a concentrations were observed in the area South of the Strait of Messina and in the Gulf of Taranto (not shown).

At the end of May in the Ionian basin, satellite imagery (figures 3A and 3B) shows a slight large scale meridional gradient of SST and Chl-a, with warmer and low-Chl water to the south and colder and medium Chl-water to the north. In addition, a strong variability was observed at the mesoscale with cold and Chl-rich filaments south of the Strait of Messina and south of Sicily, extending toward the center of the basin. Several cyclonic (~~located at 38° N 17.5° E and 34.5° N 21.3° E~~) and anti-cyclonic (~~located at 35.7° N 19° E, 34.7° N 17.5° E~~) structures were visible in the center of the basin.

185 The along route salinity showed a general increase toward the east and the north. On the northern part of the route (SAV-
ST7-ION), there was a transition from medium salinity to the north to low (<38.2) then high (>38.8) salinity at the
easternmost end of the route close to long duration station ION. On the southern part of the route (ION to ST8), there was an
alternation of low and high salinity water, sampled during ION-Tr transect. Then west of 18°E, salinity was stable lower than
190 38.2 up to the Sicily channel. Therefore SAV and ST7 were in medium salinity waters, ION (ST8) was in high (low) salinity
waters, while ION-Tr crossed waters with contrasted salinities.

The two cytometric groups showing contrasted distribution and ~~sufficient~~ abundant enough for the counts to be accurate are shown (figures 3C and 3D). In the Ionian Sea, nanoeukaryotes¹ and coccolithophore like showed a distribution
consistent with salinity, with higher values in saltier water except southeast of Sicily, where lower salinities were associated
195 to large abundance. The pattern was also mostly consistent with satellite Chl-a, with high abundances southeast of Sicily and
south of the Strait of Messina. However, along ION-Tr, abundance was variable while satellite Chl-a was fairly constant.

Altimetry-derived surface geostrophic current showed a general circulation with an anticyclonic flow across the
southwestern part of the basin (figure 4A). South of Sicily, water entered the Ionian Sea around 36°N along a southeastward
200 path, progressed eastward along 35.5°N, then veered southwestward close to 19.5°E. East of 18°E, the currents show a
pattern consistent with the intense (~0,5 m/s) currents measured by the hull-mounted ADCP along the ship route,
depicting an anti-cyclonic meander, more intense on its eastern flank. There is a small spatial offset between altimetry and
ADCP (~0.4° longitude, figure 4B), and current magnitude estimated from AVISO was about half the value of ADCP
derived current magnitude. The southwestward branch of this meander was located approximately in the center of the ION-
205 Tr transect, while the ION station was located on the eastern border of the meander. Altimetry-derived FSLE field (figure 3),
as well as the time series of ADT (annex A) showed that this anticyclonic meander east of 18°E was stable during May. The
southwestward jet sampled during ION-Tr coincided with a slightly cooler and higher Chl-a filament than surrounding
waters, as shown on figures 3A and 3B.

3.2 Distribution of water masses and biological properties along transect ION-Tr

210 Based on the salinity depth distribution (figure 5), ION-Tr can be divided into three parts. From West to East, a western part
(noted W) with a fresher surface layer extending 0-30 m covering a deeper layer with salinity >39, except the fresher top
layer 0-30m, a center part (noted C) with a fresher low salinity vein (<38.7) extending from surface down to 70 m deep, and
an eastern part (noted E) with quasi-homogeneous salinity on the vertical, slightly lower than in W (39). Hereafter we define
three pseudo-stations along ION-Tr, named ION-W, ION-C, ION-E. Note that ION long duration station was located very
215 close to the eastern part of ION-Tr (figure 1), thus ION-E and ION can be approximated as a single station.

This salinity distribution coincided with pattern of ADCP currents magnitude, with ION-W and ION-E being in quiescent water (current $< 0.2 \text{ m.s}^{-1/6}$), while ION-C correspond to a strong ($> 0.3 \text{ m.s}^{-1/6}$) southwestward jet extending from the surface to 300_m depth. The total abundance of particles, as well as Chl-a fluorescence clearly contrasted between ION-W and ION-E. In ION-W, particle abundance was low ($< 5 \cdot 10^4 \text{ \#.m}^{-3}$), essentially concentrated within the 20-30_m layer. Fluorescence was rather homogeneous with a broad maxima around 110_m. In ION-E part, abundance was much higher ($> 2 \cdot 10^4 \text{ \#.m}^{-3}$) with a deep maximum at 100_m depth (values around $1 \cdot 10^5 \text{ \#.m}^{-3}$) close to the depth of the Deep Chlorophyll Maximum (DCM) and also high particle abundance in the upper 20-90_m layer ($\sim 3 \cdot 10^4 \text{ \#.m}^{-3}$). ION-C was distinct from ION-E and ION-W in terms of vertical distribution and abundance of particle as well as fluorescence, with the low salinity vein coinciding with higher particle abundance and the maximum fluorescence being located ca 70_m, at the interface between low salinity waters and underlying waters.

The T-S distribution of ION-Tr (figure 6) mirrored the distributions of stations ST8 and ION. Water at ION-C is similar to ST8 for potential densities greater than 1028 kg.m^{-3} (ie deeper than 25m). ION-E is similar to ION with more scatter. ION-W does not match the other stations. Northern stations ST7 and SAV had similar water masses properties, distinct from ST8, ION and the ION-Tr profiles.

At the surface, cytometric groups also showed contrasting abundances along ION-Tr (figures 3C and 3D). Horizontal sampling resolution did not allow to distinguish ION-C region. From west to east with respect to ION-C, nanoeukaryotes increased from low values (as mean \pm SD, $487 \pm 52 \text{ cells.cm}^{-3}$, $n=27$, $18-19.5^\circ\text{E}$) to higher values ($659 \pm 50 \text{ cells.cm}^{-3}$, $n=117$, $> 19^\circ\text{E}$). Coccolithophore like group also increased ($176 \pm 18 \text{ cells.cm}^{-3}$ against $192.47 \pm 13 \text{ cells.cm}^{-3}$). Conversely, picoeukaryotes abundances decreased eastward ($653 \pm 94 \text{ cells.cm}^{-3}$ vs $462.8 \pm 76 \text{ cells.cm}^{-3}$, not shown). The other groups did not evidence high changes in abundances.

3.3 Comparison of vertical distribution across stations

CTD and bottle Carouse measurements performed at the SAV, ST7, ST8 stations together with the one from the ION transect are shown in figure 7. These vertical profiles allow a finer analysis of the contrast between stations and among parts of ION-Tr. Consistent with figure 6, Salinity increased from ST8 to ION-E (figure 7, panels A, F). If we take apart the upper 0-25_m depth corresponding to the mixed layer (ML), couples of stations were similar: ION-C and ST8, ION-E and ION-W, ST7 and SAV. A narrow (< 10 _m thick), lower salinity layer was present at 30_m on every profile except ION-E. The surface pycnocline was located at 10 to 25_m, followed by a steep thermocline down to 60-80_m. Surface nitrate (figure 7, panel C) was depleted everywhere. Nitracline depths increased from 60_m at SAV up to 90_m at ION, together with nitrate concentrations below the euphotic zone.

DCM was always present (figure 7, panels D, H), but its intensity was highest at ST8 and lowest at ION-W and ION-C, where fluorescence was more diluted. The depth of DCM corresponded to the top of nitracline and to the bottom of the euphotic zone (~90_m at ION).

250 Profiles of particle abundance were strongly variable among stations (figure 7, panel E, I). The depth-integrated abundance was highest at SAV and lowest at ION-W. Some profiles had several maxima. At ST8 and ION-E, the main peak was just above the DCM depth, and a secondary peak was present above DCM at ~20_m, also present at ION-W. SAV profile was different with a large peak at 50_m, above DCM, and a surface peak at 10_m.

255 Considering particle parameters (table 2), the NBSS slopes were similar for stations SAV, ION-W and ION-E (0.85), higher for ST8 (0.93) and intermediate for ION-C (0.89) indicating more abundant smaller particles at ST8. On a depth average, using the calibration given in Espinasse et al (2018), %MEP indicated that at ION-C and ION-W, particles counted were ~60% zooplankton, while at ION-E, ST8 and SAV particles were mainly detritus (~75-90%) ie non-living particles such as aggregates produced after intense primary production. AI was similar across depth layers but higher at ST8 and ION-E (0.11-0.10 against 0.05 and 0.08), while MEP was always more abundant in the 50-150_m layer, where the particle maxima
260 were observed.

3.4 Lagrangian Origin of particles

The focus here is on the particles produced before sampling and transported to the region of sampling. Figure 8 show the full trajectories of particles originating along the ION transect, splitted in four panels according to the main path followed by particles and therefore their origin. Each color identifies a distinct path. Panel E show the particle positions one month before our sampling, colored according to the origin. Note that as the altimetry-derived velocity was shifted ~0.4° in longitude west with respect to *in situ* velocity from ADCP (figure 4B), the initial positions of numerical particles were extended westward to 18°E. These simulations showed that the water parcels observed along ION-Tr originate from four main regions (figure 8), so called source regions (South origin So, West origin Wo, North origin No, Eddy origin Eo) with Wo, No and Eo being dominant.- Along ION-Tr, the proportion of each origin region showed a rather clear pattern, with some overlap in the center
265 (figure 8F). Referring to the three parts identified along ION-Tr, ION-W was in waters mainly originating from Eo. ION-C was mainly originating from Wo but also from Eo and to a lesser extent from So. ION-E was only originating from No. Finally, west of 18.6°E, beyond the limit of the actual transect, So was the main contributor.

4 Discussion

4.1 Surface circulation and distribution of water masses in the Ionian Sea

275 Surface circulation, together with the along track surface salinity distribution, describe one pathway of MAW penetrating into the Ionian Sea (figures 3 and 4). Here, this pathway is approximately zonal along latitude 36°N from the Sicily Strait up

to 19°E, then veering southwestward. This pathway is supported by the similarity of water masses properties sampled at ST8 and ION-C (figures 6 and 7) and illustrated by backtracked trajectories (figure 8, Western origin), indicating a *ca* one month travel between 15°E and 19.5°E. It is also supported by the absence of transport barriers east of ST8 shown by FSLE (figure 3). North of 36°N, the similarity of ST7 and SAV water masses properties (figure 6) also supports a single origin of water masses in the Northern Ionian.

This distribution of MAW in the Ionian Sea is consistent with a cyclonic mode of the so called North Ionian Gyre (NIG, Gačić et al 2010), that characterizes the circulation in the Ionian Sea north of *ca* 35.5°N. In the cyclonic mode of the MAW follows a rather direct East-SouthEast path, while saline water from the Levantine basin enters the Northern Ionian Sea from the East, and current flows southwestward along Calabria (Menna et al 2019). As observed by altimetry, the NIG was cyclonic since 2011 (Menna et al 2019), and reversed to anticyclonic in the second half of 2017 (von Schuckmann et al., 2019), after our sampling.

It is interesting to note that the southwestward jet sampled along ION-Tr is a persistent feature of the Ionian Sea circulation, generally located at *ca* 20°E, 36°N and directed southward (see figure 2 in Menna et al 2019, Celentano et al 2020, Berline et al 2014). It is present during cyclonic and anticyclonic modes of the NIG. It pertains to the northern branch of the Mid Ionian Jet (MIJ).

Cyclonic circulation of the NIG implies a downwelling of nutricline at the northern border of the NIG, and an upwelling of nutricline in the central NIG (Civitaresse et al 2010, Lavigne et al, 2018). Our observations of nutricline depth (isoline 1 μM at ~ 55 m for SAV and ~ 95 m for ST8) are consistent with the shallow nutriclines during cyclonic mode shown by Lavigne et al, (2018). Cyclonic mode also favors higher Chl-a in the center NIG, and a late winter bloom (Lavigne et al, 2018). However the Chl-a evolution over 2017 only shows a weak bloom in the Northern Ionian (figure 2). This may result from a weak winter buoyancy loss, independent of the NIG circulation, as pointed out by Lavigne et al (2018). The last increase of Chl-a occurred in the last week of April 2017, then Chl-a decreased and reached its minimum value around the end of May, during our sampling.

4.2 Origin of water masses sampled along ION-Tr

In the center of ION-Tr, a strong current is associated to a low salinity vein (down to 38.4, figure 6). The center current dynamically separates the two parts of the transect, and this separation is stable as supported by the FSLE ridge (figure 3). While the density stratification is similar along the transect, particle and Chl-a distributions show a striking difference from E to W. At ION-E, there is 2.8 times higher particle abundance, and 2.4 times higher biovolume compared to ION-W (table 2), essentially due to particles with and equivalent spherical diameter (ESD) smaller than 500 μm . At ION-E particles are filling the pycnocline from 1028 to 1028.7 $\text{kg}\cdot\text{m}^{-3}$ (potential density, figure 5d), with a maximum associated to a narrow DCM at

310 | 100_m, while at ION-W particles are only present in the density layer 1028-1028.2 kg.m^{-3} , and the DCM is broad and weak, not associated to a particle peak.

The distinct Chl-a and particle distributions along ION-Tr suggest that waters sampled at ION-E and ION-W have distinct histories of biological production. One hypothesis is that these water masses originate from distinct locations. To explore this hypothesis, backtracking of particle trajectories was carried out and led to three main origins of water sampled along ION-Tr
315 (Wo, No and Eo, as So is only marginally contributing to the transect), that correspond to different part of the transect (figure 8). When two origins overlap, we only consider the dominant origin. Water found at ION-C would be transported from the West (Wo), ie the Sicily channel, while water found at ION-E (ION-W) would be transported mostly from the Northern Ionian (No), and from an eddy located North (Eo) respectively. Referring back to the general circulation in the Ionian, the ION-Tr appears to be at the crossroad between waters from the North Ionian, waters from the MIJ coming from the West and
320 waters trapped into an anticyclonic eddy.

4.3 Biological history of water masses sampled along ION-Tr

LOPC detects particles of ESD >100 μm (median ESD was 130-145 μm along ION-Tr).Typically, such large aggregates form as a result of intense primary production (Briggs et al., 2020; Martin et al., 2011). In addition, the presence of transparent exopolymer particles (so called TEP) represents an important upper ocean process by which dissolved organic matter is converted into particles. Besides, these compounds act as glue enhancing small particles aggregation and leading the formation of marine snow aggregates (Engel et al., 2004) of the size we observe here, typically aggregates formed as a result of intense primary production (Briggs et al., 2020; Martin et al., 2011). Once produced, aggregates can fragment or coagulate, be grazed, be degraded by bacteria attached to them and sink out of the surface layer (Le Moigne 2019, Stemmann et al 2004). Therefore the particle abundance in the euphotic layer generally peaks shortly after a primary
325 production event, then decreases with a speed depending on several factors.
330

Although the age of particles cannot be determined from their size distribution, we can make some hypothesis. At the time of sampling, the whole Ionian Sea was stratified, mixed layer was shallow (~25_m) and nitrate was fully depleted over the euphotic zone except at SAV, indicating stable, post-bloom conditions. As surface Chl-a continuously decreased since the
335 end of April (figure 2), this suggests that the last surface primary production event around ION station occurred about a month before our sampling. This is longer than the time lag observed between primary production and subsequent export in large mesocosms experiments (Stange et al., 2017). Therefore, freshly formed particles are not likely.

The sinking speed of particles depends on their mineral composition (Le Moigne 2013), shape (Laurenceau et al., 2019) and
340 on the phytoplankton community (Riley et al 2012, Guidi et al 2009). Consistent with our cytometric results, the pigment composition available at ST7, ST8 and ION show a community dominated by nanophytoplankton (>60% of total Chl-a) in

the upper 80_m, while diatoms were abundant at DCM depth (41-51% of total Chl-a, see Marañón et al 2020). The dominance of nanophytoplankton and coccolithophores were also found by Boldrin et al (2002) in the Northern Ionian Sea (38,5°N). Thus, in the upper 80_m small cells (typically <5_μm in diameter) do not favor the production of large aggregates and direct export (Richardson and Jackson 2007), conversely to DCM.

Second, aggregates are slowed down by density gradients, because of the equilibration time of their interstitial water (MacIntyre et al 1995, Prairie et al 2015). Therefore pronounced stratification can locally increase their abundance, as reported in numerous settings (Espinasse et al 2014, Marcolin et al 2013, Ohman et al 2012). This explains that all profiles have a particle peak at the main density gradient, just below the mixed layer depth (figure 7). However, this mechanism does not explain deeper peaks observed at 60_m and 90_m.

Finally, zooplankton can also contribute to the particle production through fecal pellet; and transformation (sloppy feeding), but we are lacking zooplankton abundance data for ION-C and ION-W.

We now review the potential processes that led to the observed particles and Chl-a distributions along ION-Tr, from west to east.

4.3.1 Western part

At ION-W, below the weak particle peak (figure 7, $2 \cdot 10^4 \# \cdot m^{-3}$) associated with the density gradient at the bottom of the mixed layer, abundances steeply decrease and reach minimum values at 80_m ($1 \cdot 10^4 \# \cdot m^{-3}$). Such a low particle abundance in the top 100_m was not observed anywhere else across all PEACETIME LOPC profiles (~800 casts, Guieu et al, 2020). The Chl-a concentration is also lower and very patchy compared to ION-E. As nitracline depth was shown to be highly correlated with the depth of the $1028,9 \text{ kg} \cdot \text{m}^{-3}$ isopycnal in the Ionian (Lavigne et al 2018), the steep deepening of this isopycnal from ION-E to ION-W suggests a much deeper nutricline at ION-W than at ION-E (figure 5). Together, these observations emphasize the very oligotrophic character of station ION-W. Considering the backward trajectories (figure 8), this low abundance may result from two origins (i) waters trapped into an anticyclonic eddy (Eo), or (ii) waters from the southern Ionian (So), which is more oligotrophic and where phytoplankton concentration peak earlier in the year (D’Ortenzio and Ribera d’Alcala 2009). For hypothesis (i), analysis of the evolution of altimetry (annex A) revealed that the eddy stayed coherent during the first six months of 2017 while slowly moving from the eastern Ionian. This eddy may have trapped waters from the more oligotrophic eastern Ionian (Casotti et al 2003), and the vertical velocities in the center of the anticyclonic eddy may have accelerated the export of particles, as observed in the eastern Mediterranean (Waite et al 2016). For hypothesis (ii), it was not supported by particles profiles obtained with Underwater Vision Profiler (UVP) in the

southern Ionian in July 2008 (Picheral et al 2010) (ecotaxa.obs-vlfr.fr, Kiko et al 2020), as they did not show such low abundances, therefore not supporting this hypothesis.

375 4.3.2 Center part

At ION-C, the backtracked trajectories indicate two overlapping origins (Eo and Wo mainly). However the low salinity vein extending down to 70_m is a clear footprint of AW observed at ST8. Given the distribution of surface salinity from TSG (figure 3A and 3B) this gives more support to Wo. ST8 and ION-C also share higher NBSS slopes (0.93 and 0.89). As with ION-E, significant particles abundance above DCM suggest that particles were produced upstream, near ST8 or further west, possibly in the enriched area south of Sicily, then advected to central Ionian.

4.3.3 Eastern part (ION station)

At ION-E DCM, microphytoplankton was dominant (Marañón et al 2020) and associated to a particle peak. This high ratio of particle to phytoplankton was also noted using other optical instruments (Barbieux et al 2021). The DCM was associated to the bottom of the euphotic zone (~90_m at ION), but not to a density gradient. Therefore the particle peak associated with DCM probably result from aggregates produced locally, from diatoms decay and fecal pellet production by zooplankton. Association of DCM with a particle peak was not observed in similar settings (see station A in Espinasse et al 2018), which suggests that the plankton community at ION-E DCM enhances aggregate production, perhaps through mucous production, as mucous was reported in sediment traps in the Northern Ionian (Boldrin et al 2002).

Above DCM, significant particle abundance ($2-3 \cdot 10^4 \text{ \#} \cdot \text{m}^{-3}$) was observed together with phytoplankton dominated by nanophytoplankton (Marañón et al 2020). This significant particle abundance above DCM is also observed at SAV and ST8, with much higher peak concentration at SAV than at ST8 ($1.5 \cdot 10^5 \text{ \#} \cdot \text{m}^{-3}$ and $3-4 \cdot 10^4 \text{ \#} \cdot \text{m}^{-3}$ resp). When particles are suspended or sink slowly with respect to the horizontal velocity, particle abundance act as tracer for horizontal advection (e.g. Karageorgis et al 2012, Chronis et al 2000). We argue that at stations SAV, ST8 as well as at ION-E, the abundance of particles above DCM is a remainder of past surface production event that has not been exported out of the surface layer yet. In the Northern Ionian, a POM maximum above DCM (0-50_m) was also reported by Rabitti et al (1994), and attributed to relict bloom. As above DCM waters are depleted in nutrients, above-DCM particles at ION-E could be transported from an area where phytoplankton production was more intense and shallower, such as SAV according to the backward trajectories and the vertical profiles. This continuity is supported by the similar slope of the particle size spectra at SAV and ION-E (0.86 and 0.85, table 2). It is also supported by the higher surface abundance of cytometric counts of nanoeukaryotes and coccolithophores at SAV, along Calabria and at ION-E with respect to the other parts of the route (figures 3C, 3D). During transport from SAV to ION-E, particle abundance above DCM may have decreased because of the processes mentioned earlier.

405 | In the upper 0-90_m layer, nutrients were depleted and nanophytoplankton was dominant. Nanoeukaryotes abundances were stable during the five days of occupation of ION station, according to cytometric counts (not shown). This implies an active nutrient recycling, probably linked to the particle associated microbial community, also supported by the high bacterial production in the layer 0-100_m reported by Marañón et al (2020) at ION.

410 | ION-E particle characteristics are closest to the ‘continental shelf’ habitat type as defined by Espinasse et al (2014), in the Gulf of Lion in early May. The integrated particle abundance is about half ($18.5 \cdot 10^3 \# / m^2$ vs $38 \cdot 10^3 \# / m^2$), the average AI is lower (0.1 vs 0.2), the %MEP is higher (6.1 vs 1.8), and the NBSS slope is similar but steeper (0.85 vs 0.79). Zooplankton abundance was slightly higher ($300 \cdot 10^3 \text{ ind. m}^{-2}$ vs $206 \cdot 10^3 \text{ ind. m}^{-2}$, see Feliu et al 2020, but this value may be biased high due to the combination of two mesh sizes (100 and 200_μm) instead of one (120_μm in Espinasse et al 2018). Thus ION-E was characterized by porous aggregates, with a rather high contribution of large aggregates (%MEP).

415 | Taking into account Chl-a, nanophytoplankton abundances, significant particle load above DCM and high bacterial production, ION station was not strongly oligotrophic, possibly under the influence of particles possibly transported from North Ionian.

45 Conclusions

420 | High resolution sampling of water masses, including fluorescence and particle contents revealed unexpected heterogeneity in the central Ionian Sea. Surface salinity measurements, together with altimetry and ADCP currents, describe a zonal pathway of AW entering into the Ionian Sea, consistent with the cyclonic mode of the North Ionian Gyre. The ION-Tr appears precisely located at the crossroad of three water masses, coming from the West, North and from an isolated anticyclonic eddy. Water mass originating from the north carried abundant particles, probably originating from North Ionian, or further from the Adriatic. Waters from the eddy were depleted in particles, and probably nutrients, and may originate from the Pelops eddy area, south of Peloponnese. Waters from the West, carried by a strong jet, were intermediate in particle load, probably originating from the Sicily channel, carried by AW.

430 | A particle properties signature on water masses can be drawn (figure 7 and table 2). Comparing the three water masses observed at ST7-SAV, ST8 and ION, they correspond to three distinct particle signatures in terms of vertical profile, abundance, %MEP and AI, especially for the top 5-50 m layer which has the most contrasted particle loads. Comparing water masses at ION-W, ION-C and ION-E, only two distinct water masses emerge (ION-C and the others). In this case, particle signatures allow to distinguish ION-E and ION-W, with distinct vertical profiles, %MEP and AI. Thus particle properties are complementary to water masses properties as they bring information on their biological history along their pathways.

435 Even away from the coasts, the central Ionian Sea appears as a mosaic area, where waters of contrasted biological history
meet. ION station is influenced by particles potentially transported from the North Ionian Sea. This contrast is probably
amplified in spring, when blooming and non-blooming areas co-occur. Long distance particle transport appears as a
significant contribution to particulate matter load, together with atmospheric input.

The small scale heterogeneity of particle abundance revealed here emphasizes the spatial decoupling between particle
440 production and particle distribution. Such decoupling added to the time lag between production and export of particles
(Stange et al 2017) may have large impact on assessing the efficiency of carbon export from the surface ocean (Henson et al
2011). This also implies that neutrally buoyant particles can sustain production away, further horizontally and to deeper
depths. Interpreting the complex dynamics of physical-biogeochemical coupling from discrete measurements made at
isolated stations at sea is a challenge. The combination of multiparametric in-situ measurements at high resolution with
445 remote sensing and Lagrangian modeling appears as one proper way to address this challenge.

Author contribution

LB : Conceptualization, Writing – original draft preparation, Formal analysis, FSB : Software, Formal analysis, AD, AP, SB,
BE, FLM, FSB, FC, MT: Formal analysis, Writing – review & editing.

Acknowledgments

450 This study is a contribution to the PEACETIME project (<http://peacetime-project.org>), a joint initiative of the MERMEX and
ChArMEx components supported by CNRS-INSU, IFREMER, CEA, and Météo-France as part of the programme
MISTRALS coordinated by INSU. PEACETIME was endorsed as a process study by GEOTRACES (PEACETIME cruise
<https://doi.org/10.17600/17000300>). Jean-Luc Fuda is thanked for his help on CTD calibration.
455 We thank Dominique Lefèvre, Olivier Grosso and Gérald Grégori (MIO) for the set up and handling (DL) of the automated
and continuous sea water surface sampling system. We also thank DL and Thibault Wagener for regular checks on the
Cytosense flow cytometer during the cruise. GG contributed to the installation and uninstallation of the cytometer.

References

460 Barrillon S., Rousselet L., Petrenko A.A., Doglioli A., *Lagrangian study of the water masses dynamics at the stations of the*
PEACETIME cruise in the Mediterranean Sea. 7th LAPCOD Meeting. Venice, Italy. 17-21 June 2019.

- 465 [Barbieux, M., Uitz, J., Mignot, A., Roesler, C., Claustre, H., Gentili, B., Taillandier, V., D'Ortenzio, F., Loisel, H., Poteau, A., Leymarie, E., Penkerch, C., Schmechtig, C., and Bricaud, A.: Biological production in two contrasted regions of the Mediterranean Sea during the oligotrophic period: An estimate based on the diel cycle of optical properties measured by BGC-Argo profiling floats, *Biogeosciences Discuss.* <https://doi.org/10.5194/bg-2021-123>, in review, 2021.](#)
- Berline L, Rammou A-M, Doglioli A, Molcard A, Petrenko A A *Connectivity-Based Eco-Regionalization Method of the Mediterranean Sea*. PLoS ONE 9(11): e111978. doi:10.1371/journal.pone.0111978 2014
- 470 Blanke, B., & Raynaud, S. Kinematics of the Pacific equatorial undercurrent: An Eulerian and Lagrangian approach from GCM results. *Journal of Physical Oceanography*, 27(6), 1038-1053. 1997.
- Boldrin, A., Miserocchi, S., Rabitti, S., Turchetto, M. M., Balboni, V., & Socal, G. Particulate matter in the southern Adriatic and Ionian Sea: characterisation and downward fluxes. *Journal of Marine Systems*, 33, 389-410. 2002.
- [Briggs, N. T., Slade, W. H., Boss, E., and Perry, M. J. \(2013\). Method for estimating mean particle size from high-frequency fluctuations in beam attenuation or scattering measurements. *Appl. Opt.* 52:6710. doi: 10.1364/AO.52.006710](#)
- 475 Briggs, N., Dall'Olmo, G., Claustre, H., Major role of particle fragmentation in regulating biological sequestration of CO₂ by the oceans. *Science* (80-.). <https://doi.org/10.1126/science.aay1790> 2020
- Casotti, R., Landolfi, A., Brunet, C., D'Ortenzio, F., Mangoni, O., d'Alcalà, M. R., & Denis, M. Composition and dynamics of the phytoplankton of the Ionian Sea (eastern Mediterranean). *Journal of Geophysical Research: Oceans*, 108(C9). 2003
- 480 Celentano, P., Falco, P., & Zambianchi, E. Surface connection between the Ionian Sea and different areas of the Mediterranean derived from drifter data. *Deep Sea Research Part I: Oceanographic Research Papers*, 166, 103431. 2020
- Chronis, G., Lykousis, V., Georgopoulos, D., Zervakis, V., Stavrakakis, S., & Poulos, S. . Suspended particulate matter and nepheloid layers over the southern margin of the Cretan Sea (NE Mediterranean): seasonal distribution and dynamics. *Progress in oceanography*, 46(2-4), 163-185. [https://doi.org/10.1016/S0079-6611\(00\)00017-3](https://doi.org/10.1016/S0079-6611(00)00017-3) 2000
- 485 Civitarese, G., Gačić, M., Lipizer, M., and Eusebi Borzelli, G. L.: On the impact of the Bimodal Oscillating System (BiOS) on the biogeochemistry and biology of the Adriatic and Ionian Seas (Eastern Mediterranean), *Biogeosciences*, 7, 3987–3997, <https://doi.org/10.5194/bg-7-3987-2010>, 2010.
- Feliú, G., Pagano, M., Hidalgo, P., & Carlotti, F. Structure and function of epipelagic mesozooplankton and their response to dust deposition events during the spring PEACETIME cruise in the Mediterranean Sea. *Biogeosciences*, 17(21), 5417-5441. 2020
- 490 Doglioli, A.M., Nencioli, F., Petrenko, A. A. , Fuda, J.-L., Rougier, G., Grima, N., *A software package and hardware tools for in situ experiments in a Lagrangian reference frame*. *J. Atmos. Ocean. Tech.*, [doi: 10.1175/JTECH-D-12-00183.1](https://doi.org/10.1175/JTECH-D-12-00183.1). 2013

- 495 d'Ortenzio, F., & d'Alcalà, M. R. On the trophic regimes of the Mediterranean Sea: a satellite analysis. *Biogeosciences*, 6(2).
2009
- d'Ovidio, F., Della Penna, A., Trull, T. W., Nencioli, F., Pujol, M.-I., Rio, M.-H., Park, Y.-H., Cotté, C., Zhou, M., and
Blain, S.: The biogeochemical structuring role of horizontal stirring: Lagrangian perspectives on iron delivery
downstream of the Kerguelen Plateau, *Biogeosciences*, 12, 5567–5581, <https://doi.org/10.5194/bg-12-5567-2015> 2015.
- 500 d'Ovidio, F., Fernández, V., Hernández-García, E., and López, C.: Mixing structures in the Mediterranean Sea from finite-
size Lyapunov exponents, *Geophys. Res. Lett.*, 31, L17203, doi:10.1029/2004GL020328, 2004.
- da Rocha Marcolin, C., Schultes, S., Jackson, G. A., & Lopes, R. M. . Plankton and seston size spectra estimated by the
LOPC and ZooScan in the Abrolhos Bank ecosystem (SE Atlantic). *Continental Shelf Research*, 70, 74-87. 2013
- [Engel, A. \(2004\). Distribution of transparent exopolymer particles \(TEP\) in the northeast Atlantic Ocean and their potential
505 significance for aggregation processes. *Deep Sea Research Part I: Oceanographic Research Papers*, 51\(1\), 83-92.](#)
- Espinasse, B., Carlotti, F., Zhou, M., & Devenon, J. L. . Defining zooplankton habitats in the Gulf of Lion (NW
Mediterranean Sea) using size structure and environmental conditions. *Marine Ecology Progress Series*, 506, 31-46.
2014
- Espinasse, B., Basedow, S., Schultes, S., Zhou, M., Berline, L., & Carlotti, F.. Conditions for assessing zooplankton
510 abundance with LOPC in coastal waters. *Progress in Oceanography*, 163, 260-270. 2018
- Gačić, M., Borzelli, G. E., Civitarese, G., Cardin, V., & Yari, S. .-Can internal processes sustain reversals of the ocean upper
circulation? The Ionian Sea example. *Geophysical research letters*, 37(9). 2010
- Gačić, M., G. Civitarese, G. L. Eusebi Borzelli, V. Kovačević, P.-M. Poulain, A. Theocharis, M. Menna, A. Catucci, and N.
Zarokanellos, On the relationship between the decadal oscillations of the northern Ionian Sea and the salinity
515 distributions in the eastern Mediterranean, *J. Geophys. Res.*, 116, C12002, doi:10.1029/2011JC007280. 2011
- [Giering, S. L. C., Cavan, E. L., Basedow, S. L., Briggs, N., Burd, A. B., Darroch, L. J., ... & Waite, A. M.. Sinking organic
particles in the ocean—flux estimates from in situ optical devices. *Frontiers in Marine Science*, 6, 834. 2020](#)
- Guidi, L., Stemann, L., Jackson, G. A., Ibanez, F., Claustre, H., Legendre, L., ... & Gorsky a, G. Effects of phytoplankton
community on production, size, and export of large aggregates: A world-ocean analysis. *Limnology and*
520 *Oceanography*, 54(6), 1951-1963. 2009
- Guieu, C., D'Ortenzio, F., Dulac, F., Taillandier, V., Doglioli, A., Petrenko, A., Barrillon, S., Mallet, M., Nabat, P., and
Desboeufs, K.: Introduction: Process studies at the air–sea interface after atmospheric deposition in the Mediterranean
Sea – objectives and strategy of the PEACETIME oceanographic campaign (May–June 2017), *Biogeosciences*, 17,
5563–5585, <https://doi.org/10.5194/bg-17-5563-2020>, 2020.
- 525 Henson, S. A., Sanders, R., Madsen, E., Morris, P. J., Le Moigne, F., & Quartly, G. D. : A reduced estimate of the strength
of the ocean's biological carbon pump. *Geophysical Research Letters*, 38(4). 2011
- Herman, A.W., Beanlands, B., Phillips, E.F., The next generation of Optical Plankton Counter: the Laser-OPC. *Journal of*
Plankton Research, 26, 1135-1145. 2004

- 530 Herman, A.W., Harvey, M., Application of normalized biomass size spectra to laser optical plankton counter net
intercomparisons of zooplankton distributions. *Journal of Geophysical Research*, 111, 1-9. 2006
- Karageorgis, A. P., Gardner, W. D., Georgopoulos, D., Mishonov, A. V., Krasakopoulou, E., & Anagnostou, C. . Particle
dynamics in the Eastern Mediterranean Sea: A synthesis based on light transmission, PMC, and POC archives (1991–
2001). *Deep Sea Research Part I: Oceanographic Research Papers*, 55(2), 177-202. 2008
- 535 Karageorgis, A.P., Georgopoulos, D., Kanellopoulos, T.D., Mikkelsen, O.A., Pagou, K., Kontoyiannis, H., Kontoyianni, A.,
Anagnostou, C. : Spatial and seasonal variability of particulate matter optical and size properties in the eastern
mediterranean sea. *J. Marine Syst.* 105, 123–134. 2012
- Kiko, R : Marine particle size distribution dataset obtained with the Underwater Vision Profiler 5 - Part 1 (2008-2010).
<https://doi.pangaea.de/10.1594/PANGAEA.924374>. 2020
- 540 Laurenceau-Cornec, E.C., Le Moigne, F.A.C., Gallinari, M., Moriceau, B., Toullec, J., Iversen, M.H., Engel, A., De La
Rocha, C.L., New guidelines for the application of Stokes' models to the sinking velocity of marine aggregates. *Limnol.*
Oceanogr. 2019
- Laurenceau-Cornec, E.C., Trull, T.W., Davies, D.M., Bray, S.G., Doran, J., Planchon, F., Carlotti, F., Jouandet, M.P.,
Cavagna, A.J., Waite, A.M., Blain, S., The relative importance of phytoplankton aggregates and zooplankton fecal pellets
to carbon export: insights from free-drifting sediment trap deployments in naturally iron-fertilised waters near the
545 Kerguelen Plateau. *Biogeosciences* 12, 1007–1027. 2015.
- Lavigne, H., Civitarese, G., Gačić, M., & D'Ortenzio, F. . Impact of decadal reversals of the north Ionian circulation on
phytoplankton phenology. *Biogeosciences*, 15(14). 2018
- Le Bot, P., C. Kermabon, P. Lherminier, F. Gaillard: Rapport technique OPS/LPO 11-01. Ifremer, Centre de Brest, France.
<https://www.umr-lops.fr/en/Technology/Software/CASCADE-7.2> 2011
- 550 Le Moigne, F. A. C. . Pathways of organic carbon downward transport by the oceanic biological carbon pump. *Frontiers in*
Marine Science, 6, 634. 2019
- Le Moigne, F.A.C.; Gallinari, M.; Laurenceau, E.; De La Rocha, C.L.. Enhanced rates of particulate organic matter
remineralization by microzooplankton are diminished by added ballast minerals. *Biogeosciences*, 10 (9). 5755-5765.
<https://doi.org/10.5194/bg-10-5755-2013>. 2013
- 555 MacIntyre, S., Alldredge, A.L., Gotschalk, C.C., Accumulation of Marine Snow at Density Discontinuities in the Water
Column. *Limnology and Oceanography*, 40, 449-468. 1995
- Malanotte-Rizzoli, P., Manca, B. B., d'Alcalà, M. R., Theocharis, A., Bergamasco, A., Bregant, D., ... & Sansone, E. . A
synthesis of the Ionian Sea hydrography, circulation and water mass pathways during POEM-Phase I. *Progress in*
Oceanography, 39(3), 153-204. 1997
- 560 Marañón, E., Van Wambeke, F., Uitz, J., Boss, E. S., Pérez-Lorenzo, M., Dinasquet, J., Haëntjens, N., Dimier, C., and
Taillandier, V.: Deep maxima of phytoplankton biomass, primary production and bacterial production in the
Mediterranean Sea during late spring, *Biogeosciences Discuss.*, <https://doi.org/10.5194/bg-2020-261>, in review, 2020.

- Martin, P., Lampitt, R., Perry, M.J., Sanders, R., Lee, C., D'asaro, E. Export and mesopelagic particle flux during a North Atlantic spring diatom bloom. *Deep sea Res. part I* 58, 338–349. 2011
- 565 Mazzocchi, M. G., Nervegna, D., D'elia, G., Di Capua, I., Aguzzi, L., & Boldrin, A.. Spring mesozooplankton communities in the epipelagic Ionian Sea in relation to the Eastern Mediterranean Transient. *Journal of Geophysical Research: Oceans*, 108(C9). 2003
- [McDonnell, A. M., Lam, P. J., Lamborg, C. H., Buesseler, K. O., Sanders, R., Riley, J. S., ... & Bishop, J. K. \(2015\). The oceanographic toolbox for the collection of sinking and suspended marine particles. *Progress in oceanography*, 133, 17-31.](#)
- 570
- Menna, M., Suarez, N. R., Civitarese, G., Gačić, M., Rubino, A., & Poulain, P. M. . Decadal variations of circulation in the Central Mediterranean and its interactions with mesoscale gyres. *Deep Sea Research Part II: Topical Studies in Oceanography*, 164, 14-24. 2019
- Moutin, T., Doglioli, A.M., de Verneil, A., Bonnet, S. —Preface: The Oligotrophy to the Ultra-oligotrophy PACific Experiment (OUTPACE cruise, 18 February to 3 April 2015). *Biogeosciences*, 14, 3207-3220, doi:[10.5194/bg-14-3207-2017](#). 2017
- 575
- Nardelli, B. B., Tronconi, C., Pisano, A., & Santoleri, R.. High and Ultra-High resolution processing of satellite Sea Surface Temperature data over Southern European Seas in the framework of MyOcean project. *Remote Sensing of Environment*, 129, 1-16. 2013
- 580
- Ohman, M. D., Powell, J. R., Picheral, M., & Jensen, D. W.. Mesozooplankton and particulate matter responses to a deep-water frontal system in the southern California Current System. *Journal of plankton research*, 34(9), 815-827. 2012
- Petrenko, A., Doglioli, A.M., Nencioli, F., Kersalé, M., Hu, Z., d'Ovidio, F.. *A review of the LATEX project: mesoscale to submesoscale processes in a coastal environment*. *Ocean Dynam.*, doi: 10.1007/s10236-017-1040-9. 2017
- Picheral, M., Guidi, L., Stemmann, L., Karl, D. M., Iddaoud, G., & Gorsky, G. (2010). The Underwater Vision Profiler 5: An advanced instrument for high spatial resolution studies of particle size spectra and zooplankton. *Limnology and Oceanography: Methods*, 8(9), 462-473.
- 585
- Pinardi, N., Zavatarelli, M., Adani, M., Coppini, G., Fratianni, C., Oddo, P., ... & Bonaduce, A. (2015). Mediterranean Sea large-scale low-frequency ocean variability and water mass formation rates from 1987 to 2007: A retrospective analysis. *Progress in Oceanography*, 132, 318-332.
- 590
- Pisano, A., Nardelli, B. B., Tronconi, C., & Santoleri, R. The new Mediterranean optimally interpolated Pathfinder AVHRR SST Dataset (1982–2012). *Remote Sensing of Environment*, 176, 107-116. 2016
- Prairie, J. C., Ziervogel, K., Camassa, R., McLaughlin, R. M., White, B. L., Dewald, C., & Arnosti, C.. Delayed settling of marine snow: Effects of density gradient and particle properties and implications for carbon cycling. *Marine Chemistry*, 175, 28-38. 2015
- 595
- Rabitti, S., Bianchi, F., Boldrin, A., Daros, L., Socal, G., & Totti, C. Particulate matter and phytoplankton in the Ionian Sea. *Oceanologica Acta*, 17(3), 297-307. 1994

- Ramondenc, S., Madeleine, G., Lombard, F., Santinelli, C., Stemmann, L., Gorsky, G., & Guidi, L. An initial carbon export assessment in the Mediterranean Sea based on drifting sediment traps and the Underwater Vision Profiler data sets. *Deep Sea Research Part I: Oceanographic Research Papers*, 117, 107-119. 2016
- 600 Richardson, T.L., Jackson, G.A.,. Small phytoplankton and carbon export from the surface ocean. *Science* (80-.). 315, 838–840. <https://doi.org/10.1126/science.1133471>. 2007
- Riley, J., Sanders, R., Marsay, C., Le Moigne, F.A.C., Achterberg, E., Poulton, A., The relative contribution of fast and slow sinking particles to ocean carbon export. *Global Biogeochem. Cycles* 26. <https://doi.org/doi:10.1029/2011GB004085>. 2012
- 605 Rousselet, L., Doglioli, A. M., de Verneil, A., Pietri, A., Della Penna, A., Berline, L., ... & Petrenko, A. (2019). Vertical motions and their effects on a biogeochemical tracer in a cyclonic structure finely observed in the Ligurian Sea. *Journal of Geophysical Research: Oceans*, 124(6), 3561-3574.
- 610 Stange, P., Bach, L., Le Moigne, F.A.C., Taucher, J., Boxhammer, T., Riebesell, U.,. Quantifying the time lag between organic matter production and export in the surface ocean: Implications for estimates of export efficiency. *Geophys. Res. Lett.* <https://doi.org/10.1002/2016GL070875>. 2017
- Stemmann L., G., Jackson, G. Gorsky. A vertical model of particle size distributions and fluxes in the midwater column that includes biological and physical processes. II. Application to a three year survey in the NW Mediterranean Sea .*Deep Sea Research I* 51 (7): 865-884. 2004
- 615 Von Schuckmann, K., Le Traon, P. Y., Smith, N., Pascual, A., Djavidnia, S., Gattuso, J. P., ... & Álvarez Fanjul, E.. Copernicus Marine Service Ocean State Report, Issue 3. *Journal of Operational Oceanography*, 12(sup1), S1-S123. 2019
- Waite, A. M., Stemmann, L., Guidi, L., Calil, P. H., Hogg, A. M. C., Feng, M., ... & Gorsky, G. The wineglass effect shapes particle export to the deep ocean in mesoscale eddies. *Geophysical Research Letters*, 43(18), 9791-9800. 2016

Station	Arrival date – local time	Departure date– local time	Depth (m)	Lat N	Long E
SAV	23/05/2017 11:30	23/05/2017 14:17	2945	37°50.4	17°36.4
ST7	23/05/2017 21:10	24/05/2017 07:15	3627	36°39.5	18°09.3
ION	24/05/2017 18:02	28/05/2017 20:54	3054	35°29.4	19°46.57
ION-Tr	28/05/2017 19:41	29/05/2017 14:51	2980-3520	35°21.0 35°38.81	20°03.3 18°52.2
ST8-Tr	30/05/2017 00:43	30/05/2017 01:33	3200-3314	36°10.3 36°12.6	16°47.5 16°37.5
ST8	30/05/2017 03:53	30/05/2017 09:41	3314	36°12.6	16°37.5

Table 1. Sampling in the Ionian Sea (see figure 1). Note that ST8 and ST8-Tr are considered in the study as one single station.

	ION-W (n=13)	ION-C (n=10)	ION-E (n=15)	ST8 (n=8)	SAV (n=15)
5-50 m					
%MEP	0.35 (0.2-0.5)	0.53 (0.2-0.8)	1.23 (0.6-1.8)	0.87 (0.4-1.3)	3.20 (2.1-5.6)
AI	0.05 (0.03-0.16)	0.08 (0.04-0.12)	0.10 (0.07-0.15)	0.15 (0.08-0.24)	0.08 (0.07-0.10)
%detritus	16.08	26.46	47.54	38.87	71.47
50-150 m					
%MEP	1.26 (0.8-2.0)	1.29 (0.9-1.7)	7.76 (4.7-9.1)	6.08 (3.7-7.4)	4.44 (3.4-7.4)
AI	0.05 (0.04-0.11)	0.08 (0.04-0.11)	0.10 (0.09-0.11)	0.10 (0.08-0.14)	0.08 (0.07-0.10)
%detritus	48.14	48.73	93.64	87.53	79.67
150-300 m					
%MEP	1.17 (0.7-1.6)	1.00 (0.6-1.25)	5.68 (2.0-7.5)	2.24 (1.5-3.4)	2.94 (1.8-4.2)
AI	0.05 (0.03-0.16)	0.07 (0.03-0.13)	0.11 (0.07-0.14)	0.16 (0.09-0.25)	0.08 (0.06-0.11)
%detritus	46.28	42.36	85.83	62.54	69.35
5-300 m					
%MEP	0.88 (0.67-1.14)	0.96 (0.63-1.25)	6.10 (3.40-7.33)	4.87 (3.00-5.97)	3.81 (2.9-5.7)
AI	0.05 (0.04-0.13)	0.08 (0.04-0.12)	0.10 (0.09-0.12)	0.11 (0.08-0.15)	0.08 (0.07-0.10)
%detritus	39.16	41.33	87.62	81.98	75.84
Abund m ⁻³ x10 ⁴	0.64 (0.61-0.68)	0.99 (0.65-1.37)	1.82 (1.46-2.01)	3.69 (2.71-6.86)	5.43 (3.30-8.35)
BioV mm ³ m ⁻³ x10 ³	0.33 (0.13-1.24)	0.37 (0.17-1.29)	0.80 (0.38-1.20)	1.47 (0.71-2.25)	2.58 (1.42-3.56)
NBSS	y = -0.85x + 3.71	y = -0.89x + 3.95	y = -0.85x + 4.25	y = -0.93x + 4.58	y = -0.86x + 4.76

Table 2. LOPC particle derived parameters for ION-Tr, ST8 and SAV, for three depth layers. Abund : Total abundance and BioV : total biovolume.




Figure 1. Bathymetry of the Ionian Sea (isobaths 100, 1000, 2000, 3000, 4000m) with the locations of stations (black squares), transects and transect parts (black lines and dots). Schematic circulation (blue lines with arrows) for cyclonic conditions of the North Ionian Gyre redrawn from Menna et al (2019). The blue dashed line indicates the average AW path.





Figure 2 : Time series of A) 8-day satellite Chl-a and B) daily SST prior to sampling, averaged over the northern Ionian Sea (16° to 22°E, 34° to 40°N). Sampling at ION-Tr is marked by the dashed vertical line.



640 **Figure 3.** A) Satellite SST (May 29-30th, night, °C). B) Chl-a (May 30th, mg/m³). On panels A and B, underway surface salinity from TSG is overlaid as dots, darker for saltier water. C) Underway surface abundance (cells.cm⁻³) of nanoeukaryotes1 group from cytometry. D) Underway surface abundance (cells.cm⁻³) of coccolithophore like group from cytometry. On all panels, FSLE ridges are overlaid in light grey (>0,15 day⁻¹). Stations positions are marked as red squares and ION-Tr parts as blue dots as figure 1.




Figure 4. A) Absolute Dynamic Topography (ADT, color) and surface geostrophic circulation from altimetry averaged over period May 22-30th (black arrows). Underway ADCP currents are overlaid (red arrows). ION-Tr parts marked as blue dots as figure 1. B) Velocity along ION-Tr transect derived from altimetry for May 29th and measured with ADCP.




Figure 5. Variables sampled along ION-Tr transect from MVP, plotted as depth vs longitude. a) Salinity, b) ADCP current speed, c) Fluorescence and d) Total particle abundance. Density and current speed are overlaid as respectively black and white contours on all panels. The three parts of the transect where ION-W, ION-C, ION-E pseudo-stations are located are delimited by black vertical lines. ION station position is marked as red dot on top of each panel.






Figure 6. TS-diagram for stations ION, SAV, ST7, ST8 restricted to 0-300m (top) and for ION-Tr parts (bottom), ION-W, ION-C, ION-E.



660

Figure 7. Vertical profiles of variables at ION, SAV, ST7, ST8 (top) and for ION-Tr parts (bottom), ION-W, ION-C, ION-E. Panels A and F Salinity, B and G density, C Nitrate, D and H Chl-a, E and I total particle abundance. Note that particles were not quantified at ST7.



665 | Figure 8. Backward Lagrangian trajectories for particles launched along ION transect and tracked for one month, overlaid on average ADT (colors) and surface currents (black vectors) averaged for period May 1-30th. The particle launch area is indicated by the white polygon. Panels A, B, C, D are trajectories separated according to the four major origins. panel A for Wo, B for No, C for So, D for Eo. Black dots are the particle positions at the end of the backward advection, ie one month before our sampling. E) Final positions of particles for each origin. The pink dots are the stations positions. F) Proportion of each origin along ION-Tr computed for each 0.1° of longitude. The three parts of ION-Tr shown on figure 5 are plotted at the bottom.

670 |

Supplementary material

https://people.mio.osupytheas.fr/~berline.l/movie_ADT.gif

675 | **Appendix A. Absolute Dynamic Topography evolution for period Jan 1st-June 15th 2017. Ship track is marked as thin black line and ION-Tr as thick black line. The anticyclonic eddy sampled during ION-Tr is visible during the whole period.**