



A novel representation of biological nitrogen fixation and competitive dynamics between nitrogen-fixing and non-fixing plants in a land model (GFDL LM4.1-BNF)

Sian Kou-Giesbrecht^{1*}, Sergey Malyshev², Isabel Martínez Cano³, Stephen W. Pacala³, Elena Shevliakova², Thomas A. Bytnerowicz⁴, and Duncan N. L. Menge¹

¹Department of Ecology, Evolution and Environmental Biology, Columbia University, New York, 10027, USA

²Geophysical Fluid Dynamics Laboratory (GFDL), National Oceanic and Atmospheric Administration (NOAA), Princeton, 08540, USA

³Department of Ecology and Evolutionary Biology, Princeton University, Princeton, 08544, USA

10 ⁴Department of Integrative Biology, The University of Texas, Austin, 78712, USA

Correspondence to: Sian Kou-Giesbrecht (sk4220@columbia.edu)

Abstract. Representing biological nitrogen fixation (BNF) is an important challenge for incorporating nitrogen (N) cycling into land models. Initial representations of BNF in land models applied simplified phenomenological relationships. More recent representations of BNF are mechanistic and include the dynamic response of symbiotic BNF to N limitation of plant growth. However, they generally do not include the competitive dynamics between N-fixing and non-fixing plants, which is a key ecological mechanism that determines ecosystem-scale symbiotic BNF. Furthermore, asymbiotic BNF is generally not included in land models. Here, we present LM4.1-BNF, a novel representation of BNF (asymbiotic and symbiotic) and an updated representation of N cycling in the Geophysical Fluid Dynamics Laboratory Land Model 4.1 (LM4.1). LM4.1-BNF incorporates a mechanistic representation of asymbiotic BNF by soil microbes, a representation of the competitive dynamics between N-fixing and non-fixing plants, and distinct asymbiotic and symbiotic BNF temperature responses derived from corresponding observations. LM4.1-BNF makes reasonable estimations of major carbon (C) and N pools and fluxes and their temporal dynamics, in comparison to the previous version of LM4.1 with N cycling (LM3-SNAP) and to previous representations of BNF in land models generally (phenomenological representations and those without competitive dynamics between N-fixing and non-fixing plants and/or asymbiotic BNF). LM4.1-BNF can be applied to project the dynamic response of vegetation to N limitation of plant growth and the degree to which this will constrain the terrestrial C sink under elevated atmospheric CO₂ concentration and other global change factors.

1 Introduction

The terrestrial carbon (C) sink is controlled by the availability of nitrogen (N) for plant growth (Elser et al., 2007; LeBauer and Treseder, 2008; Wright et al., 2018). Land models are applied to project the terrestrial C sink (Arora et al., 2020) and are progressively incorporating representations of N cycling and N limitation of plant growth (Goll et al., 2017; Lawrence et al.,



2019; Medvigy et al., 2009; Smith et al., 2014; Wang et al., 2010; Yang et al., 2009; Zaehle and Friend, 2010). However, the degree to which N limitation of plant growth will constrain the terrestrial C sink under elevated atmospheric CO₂ concentration is unresolved (Terrer et al., 2019), as there is substantial variation between different land models (Wieder et al., 2015b).

The representation of biological N fixation (BNF), the primary natural input of N to terrestrial ecosystems (Fowler et al., 2013; Vitousek et al., 2013), is a key challenge to incorporating N cycling into land models because of its complexity (Davies-Barnard et al., 2020; Meyerholt et al., 2020; Stocker et al., 2016; Thomas et al., 2015; Wieder et al., 2015a). BNF occurs in multiple niches across terrestrial ecosystems: by symbioses between N-fixing bacteria living in root nodules of plants (hereafter, symbiotic BNF) and by a host of other organisms such as soil microbes, bryophytes, and lichens (hereafter, 40 asymbiotic BNF for simplicity although some of these organisms are symbiotic associations; see Reed et al., 2011). Symbiotic and asymbiotic BNF are regulated by a myriad of abiotic and biotic controls, that vary temporally, spatially, and among different niches (Zheng et al., 2019). In particular, symbiotic BNF responds dynamically to N limitation of plant growth: it is up-regulated under N limitation of plant growth and down-regulated under non-N limitation of plant growth 45 (Vitousek et al., 2013). BNF could, as such, be pivotal to overcoming N limitation of plant growth under elevated atmospheric CO₂ concentration (Liang et al., 2016; Terrer et al., 2016, 2018).

Many coupled C-N land models use the empirical relationship of BNF with either net primary production (NPP; Goll et al., 2017) or evapotranspiration (ET; B. Smith et al., 2014; Yang et al., 2009; Zaehle & Friend, 2010) to represent BNF. However, these are simplified phenomenological relationships that are not based on the ecological mechanisms underlying 50 BNF (Cleveland et al., 1999). Furthermore, implementing and comparing a NPP-based and ET-based representation of BNF within a land model (CLM5) resulted in projections of the terrestrial C sink that differed by 50 Pg C in 2100 under the representative concentration pathway 8.5 (RCP8.5; Wieder et al., 2015a). Finally, a recent meta-analysis of BNF found no evidence for the empirical relationship of BNF with either NPP or ET (Davies-Barnard and Friedlingstein, 2020).

Recent coupled C-N land models have simulated symbiotic BNF mechanistically rather than phenomenologically as 55 responding dynamically to N limitation of plant growth. The Geophysical Fluid Dynamics Laboratory (GFDL) Land Model 3 (LM3) can include the Symbiotic Nitrogen Acquisition by Plants (SNAP) model (Sulman et al., 2019), in which plant C allocation to N-fixing bacteria is optimized to maximize plant growth. However, LM3-SNAP as well as other land models that have implemented a mechanistic representation of symbiotic BNF such as CLM5 (Lawrence et al., 2019) and CABLE (Haverd et al., 2018; Peng et al., 2020; Wang et al., 2010), represent a single general plant C pool capable of BNF and 60 cannot represent community dynamics. In observed ecosystems, symbiotic BNF responds dynamically to N limitation of plant growth at both the population scale (via individual-scale regulation of symbiotic BNF rate; Menge, Wolf, & Funk, 2015) and at the community scale (via competitive dynamics between N-fixing and non-fixing plants; Boring and Swank, 1984; Chapin III et al., 1994; Menge and Hedin, 2009): Under strong N limitation, N-fixing plants up-regulate symbiotic BNF rate and have a competitive advantage over non-fixing plants, but, under weak N limitation, N-fixing plants down- 65 regulate symbiotic BNF rate and are competitively excluded by non-fixing plants because of the high C cost of symbiotic



BNF (Gutschick, 1981; Sheffer et al., 2015). As such, the competitive dynamics between N-fixing and non-fixing plants is a key ecological mechanism that could determine ecosystem-scale symbiotic BNF. Finally, the abundance of N-fixing trees is spatially variable (Menge et al., 2019; Staccone et al., 2020), but its representation is not possible in land models that represent a single general plant C pool capable of BNF although is necessary to accurately estimate regional symbiotic BNF.

70 Asymbiotic BNF is generally not included in coupled C-N land models. Phenomenological representations of BNF merge asymbiotic and symbiotic BNF, although they are regulated by different controls (Zheng et al., 2019). Mechanistic representations of BNF either merge asymbiotic and symbiotic BNF (e.g. LM3-SNAP; Sulman et al., 2019), represent asymbiotic BNF as a constant from averaged observations (e.g. CABLE; Wang and Houlton, 2009), or represent asymbiotic BNF phenomenologically as a function of ET (e.g. CLM5; Lawrence et al., 2019). Importantly, although asymbiotic and

75 symbiotic BNF exhibit different temperature responses (Bytnerowicz et al., submitted), the symbiotic BNF temperature response is, when included, derived primarily from asymbiotic BNF observations (Houlton et al., 2008), and the asymbiotic BNF temperature response is omitted.

Here, we present LM4.1-BNF, a novel representation of BNF and an updated representation of N cycling in the GFDL Land Model 4.1 (LM4.1; Shevliakova et al., in prep). LM4.1 includes height-structured competition for light and water between

80 plant cohorts using the perfect plasticity approximation (Martinez Cano et al., 2020; Purves et al., 2008; Strigul et al., 2008; Weng et al., 2015). LM4.1-BNF builds on the framework of LM4.1, including competition for light, water, and N between plant cohorts that associate with N-fixing bacteria and non-fixer plant cohorts. LM4.1-BNF introduces several improvements to the representation of N cycling in LM3-SNAP by incorporating novel representations of the following ecological mechanisms:

- 85 1. *Symbiotic BNF and competitive dynamics between N-fixing and non-fixing plants*: Plant cohorts with a N-fixer vegetation type conduct symbiotic BNF and compete with plant cohorts with a non-fixer vegetation type.
2. *Asymbiotic BNF*: Soil microbes conduct asymbiotic BNF, as well as decomposition, nitrification, and denitrification.
3. *BNF temperature response*: Asymbiotic and symbiotic BNF have different temperature responses derived from
- 90 asymbiotic BNF observations (Houlton et al., 2008) and symbiotic BNF observations (Bytnerowicz et al., submitted) respectively.
4. *N limitation*: N limitation is determined by current stored non-structural N relative to the demand for non-structural N. N limitation increases active root uptake of inorganic N and decreases root N exudation following observations (Canarini et al., 2019; Nacry et al., 2013).
- 95 5. *Dynamic plant C allocation to growth and N uptake*: N limitation decreases the growth of leaves, sapwood, and seeds, proportionally increasing the growth of fine roots following observations (Poorter et al., 2012). N limitation stimulates C allocation to N uptake (including symbiotic BNF) relative to growth. C limitation, which is determined by current stored non-structural C relative to the demand for non-structural C, stimulates C allocation to growth



100 relative to N uptake. Thereby, plant C allocation is optimized to maximize growth following observations (Rastetter and Shaver, 1992).

We focus our analysis on temperate forests which are generally N-limited (Elser et al., 2007; LeBauer and Treseder, 2008). We parameterise a N-fixer vegetation type based on *Robinia pseudoacacia* (black locust), which is the most abundant N-fixing tree species in the coterminous US, accounting for 64% of tree-associated BNF in the coterminous US (Staccone et al., 2020). We compare *Robinia* to a non-fixer vegetation type based on *Acer rubrum* (red maple), which is the most
105 abundant non-fixing tree species in the North region of the coterminous US (Oswalt et al., 2019). We evaluate LM4.1-BNF at Coweeta Hydrologic Laboratory in North Carolina, US, which has observations on symbiotic BNF by *Robinia* (Boring and Swank, 1984).

We conduct three analyses to assess the performance of LM4.1-BNF in estimating major C and N pools and fluxes in comparison to previous representations of BNF in land models generally. In the first analysis, we compare mechanistic and
110 phenomenological representations of BNF; We compare LM4.1-BNF (with BNF represented mechanistically as described above) to LM4.1-BNF with BNF represented as a function of NPP and to LM4.1-BNF with BNF represented as a function of ET. In the second analysis, we examine the role of competitive dynamics between N-fixing and non-fixing plants; We compare LM4.1-BNF simulations with both *Robinia* and *Acer* to LM4.1-BNF simulations with only *Acer* and LM4.1-BNF simulations with only *Acer* that can associate with N-fixing bacteria, which are representative of land models that represent a
115 single general plant C pool capable of BNF and cannot represent community dynamics. In the third analysis, we examine the role of asymbiotic BNF; We compare LM4.1-BNF simulations with asymbiotic BNF to LM4.1-BNF simulations without asymbiotic BNF, which is representative of land models that do not include asymbiotic BNF.

2 Model description

2.1 Overview of a land tile and vegetation types

120 We provide an overview of LM4.1-BNF with a focus on the novel elements relative to LM4.1 (Shevliakova et al., in prep) and LM3-SNAP (Sulman et al., 2019). A complete description of LM4.1-BNF is available in Appendix A.

LM4.1-BNF consists of a grid, in which grid cells are approximately 100 km by 100 km. LM4.1-BNF represents the heterogeneity of the land surface as a mosaic of land tiles within a grid cell. Each land tile represents a fraction of the grid cell area and does not have an associated location within the grid cell. A land tile may represent natural vegetation at a given
125 stage of recovery post-disturbance, urban area, pastureland, rangeland, or cropland. Land tiles are created dynamically due to a disturbance, such as human land use, fire, or natural mortality of vegetation.

A land tile contains multiple plant cohorts that compete for light and water following the Perfect Plasticity Approximation (Martinez Cano et al., 2020; Purves et al., 2008; Strigul et al., 2008; Weng et al., 2015) and compete for N (presented below). Plant cohorts consist of identical individual trees belonging to a vegetation type that occupy a given canopy layer
130 and that have a spatial density (determined by recruitment and mortality). A vegetation type can be associated with



exclusively arbuscular mycorrhizae (AM), exclusively ectomycorrhizae (EM), both AM and N-fixing bacteria, or both EM and N-fixing bacteria. A land tile can contain multiple plant cohorts of the same or of different vegetation types. As such, there is intraspecific competition (among plant cohorts of the same vegetation type within a tile) and interspecific competition (among plant cohorts of different vegetation types within a tile). Growth is based on allometric equations (Eq. (A40-42)) and is modulated by N availability. Recruitment and mortality follow Weng et al., 2015 and Martinez Cano et al., (2020) and are not directly influenced by N availability but are indirectly influenced by N availability via its effect on growth.


There are six plant tissue C and N pools: leaf, fine root, sapwood, heartwood, seed, and non-structural C or N. The C:N ratios of the leaf, fine root, sapwood, heartwood, and seed tissue pools are fixed (for a given vegetation type). There are three soil organic C and N pools (labile plant-derived, labile microbe-derived, and recalcitrant) and two soil inorganic N pools (ammonium (NH_4^+) and nitrate (NO_3^-)) in each soil layer. There are 20 soil layers of varying thickness to a total depth of 10 m. Soil C and N are transferred between soil layers via leaching. Fig. 1 displays a diagram of key C and N pools and fluxes.

We define a N-fixer vegetation type with a parameterisation based on *Robinia pseudoacacia* and a non-fixer vegetation type with a parameterisation based on *Acer rubrum*. Both *Acer* and *Robinia* associate with AM. We used the US Forest Inventory and Analysis (FIA) database (US Forest Service, 2020a), the US FIA Forest Health Monitoring database (US Forest Service, 2020b), and the Biomass and Allometry Database (BAAD; Falster et al., 2015) to parameterise the allometries of these vegetation types (Eq. (A40-42); Appendix B and C). The vegetation types also differed in other key traits (Table 1). In particular, the C:N ratio of leaves differed between vegetation types that associated with AM, EM, and N-fixing bacteria (Adams et al., 2016; Averill et al., 2019). See Table D1 for all vegetation type-specific parameters. Other model parameters are from Weng et al., (2015) or Sulman et al., (2019) or are derived from published observations (Appendix C). Some parameters were not well constrained by available observations and were tuned to fit to observed patterns of C and N cycling in temperate forests (Appendix C). See Table D2 for general parameters.

Table 1: Key parameter differences between vegetation types. See Table D1 for remaining vegetation type-specific parameters.

Vegetation type	Leaf C:N ratio	Maximum rate of carboxylation (V_{cmax}) at 15°C	Wood C density	Leaf mass per area
<i>Acer rubrum</i>	30 kg C kg N ⁻¹	17 $\mu\text{mol m}^{-2} \text{s}^{-1}$	340 kg C m ⁻³	0.0482 kg C m ⁻²
<i>Robinia pseudoacacia</i>	14 kg C kg N ⁻¹	23 $\mu\text{mol m}^{-2} \text{s}^{-1}$	280 kg C m ⁻³	0.0380 kg C m ⁻²

155



160

Figure 1: Diagram of key C and N pools (boxes) and fluxes (arrows). C pools and fluxes are indicated in blue. N pools and fluxes are indicated in orange. NSC represents non-structural C and NSN represents non-structural N. Orange and blue boxes have a fixed C:N ratio. Plant turnover, symbiont turnover, and soil C and N pools and fluxes are not displayed for visual clarity. The black dashed box represents the rhizosphere. a) Vegetation type that associates with arbuscular mycorrhizae (AM). (b) Vegetation type that associates with AM and N-fixing bacteria. (c) Vegetation type that associates with ectomycorrhizae (EM). (d) Vegetation type that associates with EM and N-fixing bacteria.

2.2 Symbiotic BNF and N uptake by roots, AM, and EM

All vegetation types take up inorganic N via passive and active root uptake. Passive root uptake of inorganic N follows LM3-SNAP (Eq. (A1)). Active root uptake of inorganic N follows LM3-SNAP but is modified to increase with N stress following observations (Nacry et al., 2013) (described in further detail below; Eq. (16)). AM take up inorganic N following LM3-SNAP (Eq. (A4)). EM decompose and take up organic C and N following LM3-SNAP but is modified to additionally take up inorganic N following observations (Phillips et al., 2013) (Eq. (A8)).

The symbiotic BNF rate by N-fixing bacteria (N_{Nfix} ; [kg N indiv^{-1} yr $^{-1}$]) is

$$170 \quad N_{Nfix} = r_{Nfix} B_{Nfix} f(T) \quad (1)$$

where r_{Nfix} is a rate constant, B_{Nfix} is the biomass C of the nodule (includes both plant and N-fixing bacteria tissue) [kg C indiv^{-1}], and $f(T)$ is the soil temperature dependence function. For *Robinia*,

$$f(T) = \max \left[0.0, \left(\frac{45.67 - (T - 273.15)}{45.67 - 31.89} \right) \left(\frac{(T - 273.15) + 1.43}{31.89 + 1.43} \right)^{\frac{31.89 + 1.43}{45.67 - 31.89}} \right] \quad (2)$$



175 where T is the average soil temperature across soil layers [K]. This reaches its maximum at 31.9 °C (Fig. D1). This is derived from Bytnerowicz et al., (submitted).

Respiration associated with symbiotic BNF is $cost_{Nfix}N_{Nfix}$, where $cost_{Nfix}$ is the C cost of symbiotic BNF per unit N (Eq. (A52)).

Note that the description of EM is included although both *Acer* and *Robinia* associate with AM.

2.3 Asymbiotic BNF

180 Soil microbes are represented as a single C pool that conducts decomposition, nitrification, denitrification, and asymbiotic BNF. The rates of C and N decomposition, rates of C and N decomposition during denitrification, rates of change of biomass C and N, and maintenance respiration rate of soil microbes follow LM3-SNAP (Eq. (A11-A19)). The N surplus or deficit of soil microbes and C and N growth rates of soil microbes are modified to include asymbiotic BNF (Eq. (A22-24)).

The asymbiotic BNF rate of soil microbes in soil layer k ($N_{Nfix\ asymb}(k)$; [kg N m⁻² yr⁻¹]) is

$$185 \quad N_{Nfix\ asymb}(k) = r_{Nfix\ asymb} C_M(k) f(T(k)) \quad (3)$$

where $r_{Nfix\ asymb}$ is a rate constant, $C_M(k)$ is the biomass C of soil microbes in soil layer k [kg C m⁻²], and $f(T(k))$ is the soil temperature dependence function.

$$f(T(k)) = e^{-2.6+0.21(T(k)-273.15)\left(1-\frac{0.5(T(k)-273.15)}{24.4}\right)} \quad (4)$$

190 which reaches its maximum at 24.4 °C (Fig. D1). This is derived from the observations compiled by Houlton et al., 2008 with the study of symbiotic BNF removed (Schomberg and Weaver, 1992) and is normalized to a maximum of 1.

2.4 N limitation, plant C allocation to growth, and plant C allocation to rhizosphere priming

The non-structural C pool (NSC ; [kg C indiv⁻¹]) gains C from photosynthesis. NSC loses C to respiration and C allocation to growth, symbionts, and root C exudation. The rate of change of NSC ($\frac{dNSC}{dt}$; [kg C indiv⁻¹ yr⁻¹]) is

$$\frac{dNSC}{dt} = P - R - (G_{C,l} + G_{C,r} + G_{C,sw} + G_{C,seed}) - C_{alloc} - L_{C,exudate} \quad (5)$$

195 where P is the photosynthesis rate [kg C indiv⁻¹ yr⁻¹], R is the respiration rate (maintenance and growth) [kg C indiv⁻¹ yr⁻¹], $G_{C,l}$ is the growth rate of the leaf C pool (C_l ; [kg C indiv⁻¹]) [kg C indiv⁻¹ yr⁻¹], $G_{C,r}$ is the growth rate of the fine root C pool (C_r ; [kg C indiv⁻¹]) [kg C indiv⁻¹ yr⁻¹], $G_{C,sw}$ is the growth rate of the sapwood C pool (C_{sw} ; [kg C indiv⁻¹]) [kg C indiv⁻¹ yr⁻¹], $G_{C,seed}$ is growth rate of the seed C pool (C_{seed} ; [kg C indiv⁻¹]) [kg C indiv⁻¹ yr⁻¹], C_{alloc} is the rate of C allocation to symbionts (Eq. (A43-46)), and $L_{C,exudate}$ is the rate of root C exudation (Eq. (A38)). Note that sapwood is converted to
200 heartwood following Martinez Cano et al., (2020).

The non-structural N pool (NSN ; [kg N indiv⁻¹]) gains N from N uptake via roots and symbionts. NSN loses N to N allocation to growth, symbionts, and root N exudation. The rate of change of NSN ($\frac{dNSN}{dt}$; [kg N indiv⁻¹ yr⁻¹]) is



$$\frac{dNSN}{dt} = U - \left(\frac{G_{C,l}}{C:N_l} + \frac{G_{C,r}}{C:N_r} + \frac{G_{C,sw}}{C:N_{sw}} + \frac{G_{C,seed}}{C:N_{seed}} \right) - N_{alloc} - L_{N,exudate} \quad (6)$$

where U is the N uptake rate via roots and symbionts (Eq. (A55)) [$\text{kg N indiv}^{-1} \text{ yr}^{-1}$], $C:N_l$ is the fixed C:N ratio of leaves, $C:N_r$ is the fixed C:N ratio of fine roots, $C:N_{sw}$ is the fixed C:N ratio of sapwood, $C:N_{seed}$ is the fixed C:N ratio of seeds, N_{alloc} is the rate of N allocation to symbionts (Eq. (A47)), and $L_{N,exudate}$ is the rate of root N exudation (Eq. (A39)).

Non-N-limited growth is calculated according to Weng et al., (2015) and Martinez Cano et al., (2020). The total allocation of NSC to growth is determined by the target NSC (NSC_{target} ; [kg C indiv^{-1}]) and minimizes the deviation between NSC and NSC_{target} . NSC_{target} is a multiple of the target C_l ($C_{l,target}$; [kg C indiv^{-1}]), which reflects the ability of a plant to re-
 210 after defoliation (Hoch et al., 2003; Richardson et al., 2013), and is calculated as

$$NSC_{target} = q C_{l,target} \quad (7)$$

where q is a proportionality constant. The allocation of NSC to the growth of each tissue depends on the total allocation of NSC to growth and the target C pool of each tissue, and minimizes the deviation between the C pool of each tissue and the target C pool of each tissue. The target C pool of each tissue is dynamic and is determined by allometry (Eq. (A40-42)),
 215 canopy position, and phenology.

In LM4.1-BNF, $G_{C,l}$, $G_{C,r}$, $G_{C,sw}$, and $G_{C,seed}$ are adjusted to include N limitation and are calculated as

$$G_{C,l} = (1 - N_{stress}) \Delta_l \quad (8)$$

$$G_{C,r} = \Delta_r \quad (9)$$

$$G_{C,sw} = (1 - N_{stress}) \Delta_{sw} \quad (10)$$

$$220 \quad G_{C,seed} = (1 - N_{stress}) \Delta_{seed} \quad (11)$$

where N_{stress} is N stress [unitless] and Δ_l , Δ_r , Δ_{sw} , and Δ_{seed} are the non-N-limited growth rates of C_l , C_r , C_{sw} , and C_{seed} respectively [$\text{kg C indiv}^{-1} \text{ yr}^{-1}$] following Weng et al., (2015) and Martinez Cano et al., (2020). Because plants increase C allocation to fine roots relative to other tissues when N-limited (Poorter et al., 2012), $G_{C,r}$ is not adjusted to include N limitation.

225 In LM4.1-BNF, N_{stress} is the relative difference between NSN and NSN_{target} and is calculated as

$$N_{stress} = \max \left[0, \frac{NSN_{target} - NSN}{NSN_{target}} \right] \quad (12)$$

where NSN_{target} is the target NSN [kg N indiv^{-1}]. N_{stress} is smoothed with a low-pass filter over 30 days to reflect the persisting influence of N stress (Mooney et al., 1991). NSN_{target} is calculated as

$$NSN_{target} = \frac{NSC_{target}}{C:N_l} \quad (13)$$

230 This is similar to LM3-SNAP, which compared the target leaf and root N pools to NSN , but is modified to reflect the treatment of NSC_{target} in LM4.1 by including the target sapwood and seed N pools.

Plant turnover decreases C_l , C_r , and C_{sw} and from N_l , N_r , and N_{sw} at a constant tissue-specific rate. A fraction of the turnover of C_l and N_l is retranslocated into NSC and NSN respectively.



Under N limitation, plants increase root C exudation to stimulate N mineralization in the rhizosphere (rhizosphere priming;
 235 Cheng et al., 2014; Finzi et al., 2015). $L_{C,exudate}$ increases with N_{stress} and is calculated as

$$L_{C,exudate} = r_{leakage,C} NSC N_{stress} \quad (14)$$

where $r_{leakage,C}$ is a rate constant.

Under N limitation, plants decrease root N exudation (Canarini et al., 2019). $L_{N,exudate}$ decreases with N_{stress} and is calculated as

$$240 \quad L_{N,exudate} = r_{leakage,N} NSN (1 - N_{stress}) \quad (15)$$

where $r_{leakage,N}$ is a rate constant.

Under N limitation, plants increase active root uptake of inorganic N (Nacry et al., 2013). The rate of active root uptake of inorganic N in soil layer k ($N_{active}(k)$; [kg N indiv⁻¹ yr⁻¹]) increases with N_{stress} and is calculated as

$$N_{active}(k) = f(NO_3(k), NH_4(k), f_{rhiz}(k)) N_{stress} \quad (16)$$

245 where $f(NO_3(k), NH_4(k), f_{rhiz}(k))$ is a function of the NH_4^+ pool in soil layer k ($NH_4(k)$; [kg N m⁻²]), the NO_3^- pool in soil layer k ($NO_3(k)$; [kg N m⁻²]), and the rhizosphere volume fraction of soil layer k ($f_{rhiz}(k)$; [m³ m⁻³]), and is given in Eq. (A3).

2.5 Plant C allocation to symbionts (AM, EM and N-fixing bacteria)

The rate of C allocation to AM ($C_{alloc,AM}$; [kg C indiv⁻¹ yr⁻¹]) is

$$250 \quad C_{alloc,AM} = f_{alloc,AM} NSC \quad (17)$$

where $f_{alloc,AM}$ is the fraction of NSC allocated to AM per unit time. $C_{alloc,AM}$ is not related to N_{stress} because, although AM increase N uptake, AM is maintained by the plant primarily for phosphorus uptake (Smith and Smith, 2011).

The rate of C allocation to EM ($C_{alloc,EM}$; [kg C indiv⁻¹ yr⁻¹]) is

$$C_{alloc,EM} = f_{alloc,EM} NSC N_{stress} \quad (18)$$

255 where $f_{alloc,EM}$ is the maximum fraction of NSC allocated to EM per unit time. $C_{alloc,EM}$ is a function of N_{stress} because biomass C of EM increases with N limitation (Phillips et al., 2013).

Plants that associate with N-fixing bacteria can regulate symbiotic BNF to different extents, termed their BNF strategy (Menge et al., 2015). For plants with a perfectly facultative BNF strategy, symbiotic BNF increases with N limitation. For plants with an incomplete BNF strategy, symbiotic BNF increases with N limitation but is maintained at a minimum. For
 260 plants with an obligate BNF strategy, symbiotic BNF is constant. For plants with either a facultative or an incomplete BNF strategy, the rate of C allocation by the plant to N-fixing bacteria ($C_{alloc,Nfix}$; [kg C indiv⁻¹ yr⁻¹]) is

$$C_{alloc,Nfix} = \max[f_{alloc,Nfix} NSC N_{stress}, f_{alloc,Nfix,min} NSC] \quad (19)$$

where $f_{alloc,Nfix}$ is the fraction of NSC allocated to N-fixing bacteria per unit time and $f_{alloc,Nfix,min}$ is the minimum fraction of NSC allocated to N-fixing bacteria per unit time. For plants with a perfectly facultative BNF strategy



265 $f_{alloc,Nfix,min} = 0$ and for plants with an incomplete down-regulator BNF strategy $f_{alloc,Nfix,min} > 0$. *Robinia* has an incomplete down-regulator BNF strategy.

For plants with an obligate BNF strategy, $C_{alloc,Nfix}$ is

$$C_{alloc,Nfix} = f_{alloc,Nfix} NSC \quad (20)$$

270 Additionally, plants allocate a small quantity of N to symbionts such that symbiont growth can be initiated. The rate of N allocation by the plant to symbionts ($N_{alloc,j}$; $j = AM, EM, N\text{-fixing bacteria}$; [$\text{kg N indiv}^{-1} \text{ yr}^{-1}$]) is

$$N_{alloc,j} = \frac{C_{alloc,j}}{C:N_{alloc}} \quad (21)$$

where $C:N_{alloc}$ is the C:N ratio of C and N allocated to symbionts by the plant.

Plant C allocation to symbionts increases biomass C of symbionts, which increases plant N uptake via symbionts (Appendix A.1 and A.6).

275 2.6 Dynamic plant C allocation to growth and N uptake

The order of plant C allocation to growth, symbionts, and rhizosphere priming is determined by C limitation relative to N limitation (Cheng et al., 2014; Finzi et al., 2015; Poorter et al., 2012; Treseder, 2004; Zheng et al., 2019). If a plant is more C-limited than N-limited, $NSC < NSN \cdot C:N_{leaf}$, the plant allocates C to growth, then to N-fixing bacteria (if associated) and EM (if associated), then to rhizosphere priming, and finally to AM. If a plant is more N-limited than C-limited, $NSC >$
 280 $NSN \cdot C:N_{leaf}$, the plant allocates C to N-fixing bacteria (if associated) and EM (if associated), then to rhizosphere priming, then to growth, and finally to AM.

2.7 Soil N₂O and NO emissions

Soil N₂O and NO emissions occur during nitrification (aerobic oxidation of NH₄⁺ with oxygen as an electron acceptor, which produces N₂O and NO as by-products) and denitrification (anaerobic oxidation of organic C with NO₃⁻ as an electron
 285 acceptor, which produces N₂O as a by-product).

Following LM3V-N (Huang and Gerber, 2015), soil N₂O emission rate in soil layer k ($N_2O(k)$; [$\text{kg N m}^{-2} \text{ yr}^{-1}$]) is

$$N_2O(k) = \gamma_{N_2O,nit} nit(k) + \frac{1}{1 + \max\left(3.52, 22 \exp\left(-0.8 \frac{NO_3(k)}{HR(k)}\right)\right) \max\left(0.1, 0.015 \frac{\theta(k)}{\theta_{sat}} - 0.32\right)} denit(k) \quad (22)$$

where $\gamma_{N_2O,nit}$ is the proportion of the nitrification rate that is emitted as N₂O, $nit(k)$ is nitrification rate in soil layer k [$\text{kg N m}^{-2} \text{ yr}^{-1}$] (Eq. (A56)), $HR(k)$ is heterotrophic respiration in soil layer k [$\text{kg C m}^{-2} \text{ yr}^{-1}$], $\theta(k)$ is volumetric soil water content of soil layer k [$\text{m}^3 \text{ m}^{-3}$], θ_{sat} is saturation volumetric soil water content, and $denit(k)$ is denitrification rate in soil layer k [$\text{kg N m}^{-2} \text{ yr}^{-1}$] (Eq. (A57)).
 290

Following LM3V-N (Huang and Gerber, 2015), soil NO emission rate in soil layer k ($NO(k)$; [$\text{kg N m}^{-2} \text{ yr}^{-1}$]) is



$$NO(k) = \left(15.2 + \frac{35.5 \tan^{-1} \left(0.68 \pi \left(2.09 \left(1 - \frac{\theta(k)}{\theta_{sat}} \right)^{4/3} - 1.68 \right) \right)}{\pi} \right) \gamma_{N_2O,nit} nit(k) \quad (23)$$

2.8 Phenomenological representations of BNF (LM4.1-BNF_{NPP} and LM4.1-BNF_{ET})

295 To compare mechanistic and phenomenological representations of BNF in our first analysis, we developed LM4.1-BNF_{NPP} and LM4.1-BNF_{ET} which have phenomenological representations of BNF. In LM4.1-BNF_{NPP}, BNF is represented as a function of NPP in LM4.1-BNF. BNF rate (BNF_{NPP} ; [kg N m⁻²]) is

$$BNF_{NPP} = a_{NPP} (1 - e^{b_{NPP} NPP}) \quad (24)$$

where NPP is net primary production [kg C m⁻² yr⁻¹], and a_{NPP} and b_{NPP} are constants from Meyerholt et al., 2016.

300 In LM4.1-BNF_{ET}, BNF is represented as a function of ET in LM4.1-BNF. BNF rate (BNF_{ET} ; [kg N m⁻²]) is

$$BNF_{ET} = a_{ET} ET + b_{ET} \quad (25)$$

where ET is evapotranspiration [mm yr⁻¹] and a_{ET} and b_{ET} are constants from Meyerholt et al., 2016.

305 BNF_{NPP} and BNF_{ET} enter $NH_4(k)$ (distributed across all soil layers proportional to thickness). In LM4.1-BNF_{NPP} and LM4.1-BNF_{ET}, growth and turnover of symbionts and plant C allocation to symbionts do not occur, and asymbiotic BNF does not occur. All other components of C and N cycling in LM4.1-BNF_{NPP} and LM4.1-BNF_{ET} are the same as LM4.1-BNF.

3 Numerical experiments and evaluation description

3.1 Numerical experiments description

We ran numerical experiments for the grid cell containing Coweeta Hydrological Laboratory (CHL) in North Carolina, US (35.05°N, 83.45°W), which is part of the Long-Term Ecological Research Network and has observations on symbiotic BNF
 310 by *Robinia* (Boring and Swank, 1984).

We ran the LM4.1-BNF spin up for 1000 years at pre-industrial atmospheric CO₂ concentration (284.26 ppm) to allow the soil C and N pools to reach an approximate steady state. Then, we initialised LM4.1-BNF, LM4.1-BNF_{NPP}, and LM4.1-BNF_{ET} numerical experiments with seedlings (removed vegetation C and N pools from the spin up) and the LM4.1-BNF spin up soil C and N pools. We ran numerical experiments for another 300 years at current atmospheric CO₂ concentration
 315 (324.53 ppm). See Table D3 for a summary of atmospheric CO₂ concentration (Dlugokencky and Tans, 2020; Meinshausen et al., 2017), meteorological forcings (Sheffield et al., 2006), and N deposition rates (Dentener, 2006) used in the spin up and numerical experiments. We initialised the LM4.1-BNF spin up with *Acer* seedlings, and we initialised LM4.1-BNF numerical experiments with both *Acer* and *Robinia* seedlings. To compare mechanistic and phenomenological representations of BNF in our first analysis (Table 2), we initialised LM4.1-BNF_{NPP} and LM4.1-BNF_{ET} numerical
 320 experiments with only *Acer* seedlings. To examine the role of competitive dynamics between N-fixing and non-fixing plants



in our second analysis (Table 2), we initialised LM4.1-BNF numerical experiments with both *Acer* and *Robinia* seedlings, only *Acer* seedlings or only *Acer* seedlings that can associate with N-fixing bacteria (N-fixer *Acer*). To examine the role of asymbiotic BNF in our third analysis (Table 2), we initialised LM4.1-BNF numerical experiments with both *Acer* and *Robinia* seedlings, only *Acer* seedlings, or only N-fixer *Acer* seedlings. LM4.1-BNF, LM4.1-BNF_{NPP}, and LM4.1-BNF_{ET} simulations are initialised such that all plant cohorts have the same height (0.5 m; Table D4) and dbh is determined from height by allometry (Eq. (A41); Table D4). See Table D4 for a summary of initial density, height, and dbh of seedlings in numerical experiments.

We ran the LM3-SNAP spin up for 1000 years at pre-industrial atmospheric CO₂ concentration to allow the soil C and N pools to reach an approximate steady state. Then, we initialised LM3-SNAP numerical experiments with seedlings and the LM3-SNAP spin up soil C and N pools. We ran numerical experiments for another 300 years at current atmospheric CO₂ concentration. See Table D3 for a summary of atmospheric CO₂ concentration, meteorological forcings, and N deposition rates used in the spin up and numerical experiments. We initialised the LM3-SNAP spin up and the LM3-SNAP numerical experiments with a temperate deciduous vegetation type.

Table 2: Description of numerical experiments. In the first analysis, we compare mechanistic and phenomenological representations of BNF. In the second analysis, we examine the role of competitive dynamics between N-fixing and non-fixing plants. In the third analysis, we examine the role of asymbiotic BNF.

Analysis	BNF representation	Species	Asymbiotic BNF
1. Mechanistic and phenomenological representations of BNF	LM4.1-BNF	<i>Robinia</i> and <i>Acer</i>	represented
	LM4.1-BNF _{NPP}	<i>Robinia</i> and <i>Acer</i>	-
	LM4.1-BNF _{ET}	<i>Robinia</i> and <i>Acer</i>	-
2. Competitive dynamics between N-fixing and non-fixing plants	LM4.1-BNF	<i>Robinia</i> and <i>Acer</i>	represented
	LM4.1-BNF	<i>Robinia</i> and <i>Acer</i>	-
	LM4.1-BNF	<i>Acer</i>	represented
3. Asymbiotic BNF	LM4.1-BNF	N-fixer <i>Acer</i>	represented
	LM4.1-BNF	N-fixer <i>Acer</i>	-

3.2 Evaluation description

We evaluated the LM4.1-BNF numerical experiment initialised with both *Acer* and *Robinia* seedlings. To evaluate LM4.1-BNF at the population scale (individuals of the same species in an ecosystem), we compared the diameter at breast height (dbh) growth rate of each vegetation type (*Acer* and *Robinia*) of the numerical experiments to US FIA database tree data (Fig. 2; US Forest Service, 2020a). To evaluate LM4.1-BNF at the community scale (populations in an ecosystem), we compared the dbh distribution of the numerical experiments to US FIA database tree data (Fig. 3; US Forest Service, 2020a) and we compared the basal area fraction of each vegetation type (*Acer* and *Robinia*) over time of the numerical experiments to US FIA database tree data (Fig. 4; US Forest Service, 2020a). To evaluate LM4.1-BNF at the ecosystem scale, we made four comparisons. First, we compared total plant biomass C to US FIA database tree data (Fig. 5; US Forest Service, 2020a)




and site observations from CHL (Fig. 5; Boring and Swank, 1984). Second, we compared asymbiotic and symbiotic BNF rate to site observations from CHL (Fig. 6; Boring and Swank, 1984; Todd et al., 1978). Third, we compared soil C and N
350 pools and fluxes to US FIA database soil data (Fig. 7; US Forest Service, 2020a), site observations from CHL (Fig. 7; Binkley et al., 1992; Knoepp, 2009a, 2009b, 2018; Swank & Waide, 1988), and a meta-analysis of temperate forests (Fig. 7; Stehfest and Bouwman, 2006). Fourth, we compared ecosystem C fluxes to a meta-analysis of temperate forests (Fig. 8; Anderson-Teixeira et al., 2018). See Appendix B for data availability and processing. See Table D5 for a summary of the validated variables and data sources.

355 **4 Evaluation: comparison of LM4.1-BNF to observations and to LM3-SNAP**

Here we describe the evaluation of LM4.1-BNF, in which we compare LM4.1-BNF simulations to observations and to LM3-SNAP (which represents a single general plant C pool capable of BNF and cannot represent community dynamics (i.e., competitive dynamics between N-fixing and non-fixing plants) and does not represent asymbiotic BNF). The comparisons
360 between LM4.1-BNF simulations and observations at population (Fig. 2), community (Fig. 3-4), and ecosystem scales (Fig. 5-8) show that LM4.1-BNF captures observed forest growth, successional dynamics, and major pools and fluxes of C and N, in particular asymbiotic and symbiotic BNF. Note that parameters were not tuned to fit observations specific either to CHL or to forests in North Carolina, US (Appendix B); Population, community, and ecosystem-scale metrics emerge from the ecological mechanisms represented in LM4.1-BNF.

4.1 Population scale

365 LM4.1-BNF makes reasonable estimates for dbh growth rates but overestimates dbh growth rates for both *Robinia* and *Acer* (mean 5.52 vs. 4.46 mm yr⁻¹ and 3.96 vs. 1.71 mm yr⁻¹ respectively; Fig. 2ab). Note that LM3-SNAP cannot distinguish between *Robinia* and *Acer*.




370 **Figure 2: Simulated dbh growth rate for (a) *Acer* / non-fixer and (b) *Robinia* compared to FIA data (in North Carolina). Simulated data are trees with dbh > 12.7 cm to reflect the dbh range of FIA data and are weighted by the stand age distribution of FIA data (Fig. D6). FIA data of all non-fixing trees are aggregated to represent *Acer*. Simulated and FIA data are scaled to display an equal maximum density.**



4.2 Community scale

LM4.1-BNF makes a reasonable estimate for the dbh distribution but overestimates density (maximum 350 vs. 98 indiv ha⁻¹; Fig. 3). LM4.1-BNF effectively reproduces the successional dynamics of *Robinia* and *Acer* (Fig. 4): *Robinia* is competitively excluded by *Acer* at approximately the same time scale as observations (< 2% basal area fraction at 150 years). Note that LM3-SNAP cannot distinguish between plant cohorts with different dbhs or to distinguish between *Robinia* and *Acer*.



380 **Figure 3: Simulated dbh distribution compared to FIA data (in North Carolina). Simulated data are trees with dbh > 12.7 cm to reflect the dbh range of FIA data and are weighted by the stand age distribution of FIA data (Fig. D6).**




Figure 4: Simulated relative basal area of *Acer* and *Robinia* over time compared to FIA data (in North Carolina). Simulated data are trees with dbh > 12.7 cm to reflect the dbh range of FIA data. FIA data of all non-fixing trees are aggregated to represent *Acer*. Each point represents an FIA plot. See Fig. D7 for absolute basal area.




385 4.3 Ecosystem scale

LM4.1-BNF effectively reproduces the temporal dynamics of total plant biomass C, particularly in comparison to LM3-SNAP which underestimates total plant biomass C (Fig. 5). LM4.1-BNF simulates a reasonable estimate (mean 13.4 vs. 10.8 kg N ha⁻¹ yr⁻¹ over the final 100 years) for asymbiotic BNF rate, which is not represented in LM3-SNAP (Fig. 6a). LM4.1-BNF effectively reproduces the temporal dynamics of symbiotic BNF rate: LM4.1-BNF simulates a symbiotic BNF rate pulse in early succession that reaches 120.8 kg N ha⁻¹ yr⁻¹ at 12 years, then declines to ~ 0 kg N ha⁻¹ yr⁻¹ at 300 years (Fig. 6b). LM3-SNAP simulates higher symbiotic BNF rate in late succession in comparison to LM4.1-BNF (mean 8.3 vs. ~ 0 kg N ha⁻¹ yr⁻¹ over the final 100 years; Fig. 6b) because there is no competitive exclusion of N-fixing plants by non-fixing plants due to weak N limitation of plant growth in late succession in LM3-SNAP, which represents a single general plant C pool capable of BNF and cannot represent community dynamics. Additionally, LM3-SNAP simulates lower total plant biomass C than LM4.1-BNF simulations (mean 44.0 vs. 174.6 Mg C ha⁻¹) because of the persisting high C cost of symbiotic BNF.


LM4.1-BNF makes reasonable estimates for soil C and N pools and fluxes, which are comparable to those of LM3-SNAP. LM4.1-BNF underestimates total soil C and N (mean 13.3 vs. 33.0 Mg C ha⁻¹ and 0.2 vs. 1.8 Mg N ha⁻¹ respectively; Fig. 7ab), and overestimates soil NH₄⁺ and NO₃⁻ (mean 4.7 vs. 1.0 kg N ha⁻¹ and 1.6 vs. 0.1 kg N ha⁻¹ respectively; Fig. 7c). LM4.1-BNF underestimates N mineralization rate and net nitrification rate (mean 14.9 vs. 39.5 kg N ha⁻¹ yr⁻¹ and 7.2 vs. 12.8 kg N ha⁻¹ yr⁻¹ respectively; Fig. 7d). LM4.1-BNF overestimates N₂O emission rate (mean 5.1 vs. 0.9 kg N ha⁻¹ yr⁻¹) and underestimates NO emission rate (mean 0.2 vs. 0.8 kg N ha⁻¹ yr⁻¹; Fig. 7e). Note that LM3-SNAP cannot simulate N₂O and NO emission rates. LM4.1-BNF underestimates dissolved organic N (DON) leaching rate (mean ~0 vs. 0.6 kg N ha⁻¹ yr⁻¹) and NH₄⁺ leaching rate (mean ~0 vs. 0.05 kg N ha⁻¹ yr⁻¹), and overestimates NO₃⁻ leaching rate (mean 0.13 vs. 0.10 kg N ha⁻¹ yr⁻¹; Fig. 7e). LM3-SNAP substantially overestimates NO₃⁻ leaching rate (Fig. 7e).

405 Lastly, LM4.1-BNF makes reasonable estimates for ecosystem C flux rates, particularly in comparison to LM3-SNAP which overestimates gross primary production (GPP), NPP, and heterotrophic respiration (HR) (Fig. 8). LM4.1-BNF overestimates GPP and NPP (mean 15.6 vs. 13.1 Mg C ha⁻¹ yr⁻¹ and 8.1 vs. 7.5 Mg C ha⁻¹ yr⁻¹ respectively). LM4.1-BNF overestimates HR (mean 7.6 vs. 4.7 Mg C ha⁻¹ yr⁻¹) and consequentially underestimates net ecosystem production (NEP) (mean 0.5 vs. 4.8 Mg C ha⁻¹ yr⁻¹).




410

Figure 5: Simulated total plant biomass C over time by LM4.1-BNF and LM3-SNAP compared to FIA data (in North Carolina). Each point represents an FIA plot.




415 **Figure 6: Simulated BNF rate by LM4.1-BNF and LM3-SNAP. (a) Simulated asymbiotic BNF rate compared to CHL site data. Simulated data are averaged over the last 100 yr of the 300 yr simulation to reflect the site data which is from mature forests. Error bars indicate two standard deviations. (b) Simulated symbiotic BNF rate over time compared to CHL site data for a 4 yr, 17 yr, 38 yr and mature forest (plotted at 300 yr).**



420 **Figure 7: Simulated soil C and N pools, soil N fluxes, and soil N loss rates by LM4.1-BNF and LM3-SNAP. (a) Simulated total soil C (depth 0-10 cm) compared to FIA data (in North Carolina) and CHL site data. (b) Simulated total soil N (depth 0-10 cm) compared to FIA data (in North Carolina) and CHL site data. (c) Simulated soil NH₄⁺ and NO₃⁻ (depth 0-10 cm) compared to CHL site data. (d) Simulated N mineralization rate and net nitrification rate (depth 0-10 cm) compared to CHL site data. (e) Simulated N₂O and NO emission rates compared to a meta-analysis estimate for temperate forests, and simulated dissolved organic N (DON), NH₄⁺, and NO₃⁻ leaching rate compared to CHL site data. Simulated data are averaged over the last 100 yr of the 300 yr**



425 simulation to reflect the site data which is from mature forests. Error bars indicate two standard deviations. NA indicates that LM3-SNAP cannot estimate N₂O or NO emissions.



430 Figure 8: Simulated gross primary production (GPP), net primary production (NPP), heterotrophic respiration (HR), and net ecosystem production (NEP) by LM4.1-BNF and LM3-SNAP compared to the ForC database. Simulated data are averaged over the last 100 yr of the 300 yr simulation to reflect the data which is from mature forests. Error bars indicate two standard deviations.




5 LM4.1.-BNF performance relative to previous BNF representations

Here we describe the three analyses we conducted to identify LM4.1-BNF improvements to estimating major C and N pools and fluxes. First, we compare LM4.1-BNF to LM4.1-BNF with BNF represented as a function of NPP (LM4.1-BNF_{NPP}) and
435 LM4.1-BNF with BNF represented as a function of ET (LM4.1-BNF_{ET}) to compare mechanistic and phenomenological representations of BNF. Second, we compare LM4.1-BNF simulations with both *Robinia* and *Acer* to LM4.1-BNF simulations with only *Acer* and LM4.1-BNF simulations with only *Acer* that can associate with N-fixing bacteria (hereafter, N-fixer *Acer*) to examine the role of competitive dynamics between N-fixing and non-fixing plants. Third, we compare
440 LM4.1-BNF simulations (with both *Robinia* and *Acer*, only *Acer*, and only N-fixer *Acer*) with asymbiotic BNF to LM4.1-BNF simulations without asymbiotic BNF to examine the role of asymbiotic BNF.

5.1 Mechanistic and phenomenological representations of BNF

In our first analysis (Table 2), we compare mechanistic and phenomenological representations of BNF, and their implications for C and N cycling. LM4.1-BNF, LM4.1-BNF_{NPP}, and LM4.1-BNF_{ET} simulations estimate different total plant biomass C (Fig. 9a). LM4.1-BNF predicts the largest total plant biomass C (mean 174 Mg C ha⁻¹ over the final 100 years),
445 followed by LM4.1-BNF_{ET} (mean 70 Mg C ha⁻¹ over the final 100 years) and LM4.1-BNF_{NPP} (mean 0 Mg C ha⁻¹ over the final 100 years). This is because, in LM4.1-BNF, BNF responds dynamically to strong N limitation of plant growth in early succession and BNF (mean 57 kg N ha⁻¹ yr⁻¹ over the initial 100 years) supports total plant biomass C accumulation. Conversely, in LM4.1-BNF_{NPP} and LM4.1-BNF_{ET}, BNF does not respond dynamically to strong N limitation of plant growth in early succession and BNF is not sufficient (mean 22 and ~ 0 kg N ha⁻¹ yr⁻¹ over the initial 100 years for LM4.1-BNF_{ET} and
450 LM4.1-BNF_{NPP} respectively) to support total plant biomass C accumulation (Fig. 9b). As such, LM4.1-BNF effectively reproduces the temporal dynamics of symbiotic BNF rate, whereas LM4.1-BNF_{ET} and LM4.1-BNF_{NPP} predicted relatively constant symbiotic BNF rates. In observed ecosystems, strong N limitation of plant growth occurs in early succession. N-fixing trees are generally important pioneer species and can relieve strong N limitation of plant growth in early succession (Chapin III et al., 1994; Cierjacks et al., 2013; Menge et al., 2010). Consequently, symbiotic BNF is highest in early
455 succession (Batterman et al., 2013; Boring and Swank, 1984; Menge and Hedin, 2009; Sullivan et al., 2014). Simulated soil C and N pools, soil N fluxes, soil N loss rates, and ecosystem C fluxes are relatively similar between simulations and are displayed in Fig. D2-3.

A similar result was found by Meyerholt et al., (2020), who compared five alternative representations of BNF within the O-CN model, including a BNF representation based on NPP, a BNF representation based on ET, and a BNF representation
460 responding dynamically to N limitation of plant growth. As with our results, they found that the BNF representation responding dynamically to N limitation of plant growth predicted the largest total plant biomass C. However, their study did not compare these results to simulations that include competitive dynamics between N-fixing and non-fixing plants because the land model (O-CN) represents a single general plant C pool capable of BNF and cannot represent community dynamics.



465 **Figure 9:** Simulated (a) total plant biomass C and (b) symbiotic BNF rate over time from LM4.1-BNF, LM4.1-BNF_{NPP}, and LM4.1-BNF_{ET}.



5.2 Competitive dynamics between N-fixing and non-fixing plants

In our second analysis (Table 2), we examine the role of competitive dynamics between N-fixing and non-fixing plants and its implication for C and N cycling. LM4.1-BNF simulations initialised with only *Acer* seedlings accumulate total plant biomass C slower than LM4.1-BNF simulations initialised with both *Robinia* and *Acer* seedlings (mean 48.6 vs. 74.5 Mg C ha⁻¹ over the initial 100 years respectively; Fig. 10a). In LM4.1-BNF simulations initialised with only *Acer* seedlings, stronger N limitation of plant growth in early succession due to the absence of a N-fixer vegetation type slows total plant biomass C accumulation. Nevertheless, total plant biomass C accumulates due to asymbiotic BNF and high N deposition at CHL (13.9 kg N ha⁻¹ yr⁻¹), reaching a similar level to LM4.1-BNF simulations initialised with both *Robinia* and *Acer* seedlings after 100 years.

LM4.1-BNF simulations initialised with only N-fixer *Acer* seedlings accumulate total plant biomass C slightly slower than LM4.1-BNF simulations initialised with both *Robinia* and *Acer* seedlings (mean 60.0 vs. 74.5 Mg C ha⁻¹ over the initial 100 years respectively; Fig. 10a). However, in LM4.1-BNF simulations initialised with only N-fixer *Acer* seedlings, a higher symbiotic BNF rate persists throughout succession in comparison to LM4.1-BNF simulations initialised with both *Robinia* and *Acer* seedlings (mean 151.4 vs. ~ 0 kg N ha⁻¹ yr⁻¹ over the final 100 years respectively; Fig. 10b). This occurs because there is no competitive exclusion of N-fixing plants by non-fixing plants due to weak N limitation of plant growth in late succession, which occurs in LM4.1-BNF simulations initialised with both *Robinia* and *Acer* seedlings. Additionally, LM4.1-BNF simulations initialised with only N-fixer *Acer* seedlings estimate a lower total plant biomass C than LM4.1-BNF simulations initialised with both *Robinia* and *Acer* seedlings (mean 123.3 vs. 174.5 Mg C ha⁻¹ over the final 100 years respectively; Fig. 10a) because of the high C cost of symbiotic BNF. In LM4.1-BNF simulations initialised with only N-fixer *Acer* seedlings, over-estimated symbiotic BNF causes over-estimated soil inorganic N, soil N flux rates, and soil N loss rates (Fig. D4). In particular, accurately estimating soil N loss rates is important due to their numerous consequences such as eutrophication and the positive radiative forcing of N₂O (Fowler et al., 2013). Simulated ecosystem C fluxes are relatively similar between simulations and are displayed in Fig. D5.

Levy-Varon et al., (2019) conducted a similar study, in which a N-fixer vegetation type was included in the ED2 model. Similarly, they found that simulations without a N-fixer vegetation type accumulate total plant biomass C slower than simulations with a N-fixer vegetation type. However, ED2 differs from LM4.1-BNF in a multitude of processes. In particular, ED2 does not include representations of asymbiotic BNF, mycorrhizae, or rhizosphere priming. Furthermore, the representation of BNF in ED2 assumes instantaneous down-regulation of symbiotic BNF rate (in comparison to a time lag in down-regulation of symbiotic BNF rate due to the time between plant C allocation to symbiotic BNF, the growth of N-fixing bacteria, and symbiotic BNF in LM4.1-BNF following observations (Bytnerowicz et al., submitted)).




Figure 10: Simulated (a) total plant biomass C and (b) symbiotic BNF rate over time from LM4.1-BNF initialised with both *Robinia* and *Acer*, only *Acer*, and only N-fixer *Acer*, with and without asymbiotic BNF. aBNF indicates asymbiotic BNF.



500 5.3 Asymbiotic BNF

In our third analysis (Table 2), we examine the role of asymbiotic BNF and its implications for C and N cycling. LM4.1-BNF simulations initialised with *Acer* without asymbiotic BNF accumulate total plant biomass C slower than LM4.1-BNF simulations initialised with *Acer* with asymbiotic BNF (30.4 vs. 48.6 Mg C ha⁻¹ over the initial 100 years; Fig. 10a). In LM4.1-BNF simulations initialised with *Acer* without asymbiotic BNF, stronger N limitation of plant growth in early
505 succession due to the absence of both asymbiotic BNF and a N-fixer vegetation type (i.e., symbiotic BNF) substantially slows total plant biomass C accumulation. Nevertheless, total plant biomass C accumulates due to high N deposition at CHL (13.9 kg N ha⁻¹ yr⁻¹), reaching a similar level to LM4.1-BNF simulations initialised with both *Robinia* and *Acer* seedlings after 300 years. Simulated soil C and N pools, soil N fluxes, soil N loss rates, and ecosystem C fluxes are relatively similar between simulations and are displayed in Fig. D4-5.

510 6 Discussion

6.1 Limitations

LM4.1-BNF captures the major pools and fluxes of C and N and their temporal dynamics. Importantly, LM4.1-BNF is novel in that it captures both the competitive dynamics between N-fixing and non-fixing plants as well as asymbiotic BNF. However, LM4.1-BNF has limitations.
515 LM4.1-BNF does not explicitly include asymbiotic BNF by bryophytes, lichens, and other organisms beyond soil microbes. This is regulated differently from asymbiotic BNF by soil microbes, specifically by light (Reed et al., 2011). In particular, in boreal forests and arctic tundra, asymbiotic BNF by bryophytes is a significant N flux (DeLuca et al., 2002). Additionally, herbaceous symbiotic BNF in the forest understory could be significant but few studies have quantified its magnitude and controls (Cleveland et al., 1999).
520 The asymbiotic BNF temperature response is heavily biased towards high latitudes; the studies we used in its derivation had a mean latitude of 60° (Chan, 1991; Chapin et al., 1991; Coxson and Kershaw, 1983; Liengen and Olsen, 1997; Roper, 1985). More studies on the asymbiotic BNF temperature response at lower latitudes are necessary.
The symbiotic BNF temperature response could acclimate to changing temperature (Bytnerowicz et al., submitted). The C cost of symbiotic BNF, which we assumed to be constant per unit N, could depend on temperature or other factors. These
525 issues could influence the simulated response of symbiotic BNF and consequently total plant biomass C to increasing temperatures due to climate change. Thus, further empirical work on the effect of temperature on symbiotic BNF is necessary.
Finally, more observations of N cycling in general are necessary to validate N cycling representations in land models (Stocker et al., 2016; Thomas et al., 2015; Vicca et al., 2018). Global observations on N limitation of plant growth, soil N, N



530 gas emission rates, N leaching rates, and, in particular, asymbiotic and symbiotic BNF rates are limited. Constraining these
N pools and fluxes is critical to rigorously validating novel N cycling representations in land models.

6.2 Extensions

Robinia pseudoacacia is the most abundant N-fixing tree species in the coterminous US (Staccone et al., 2020) and is also a
common N-fixing tree across temperate forests; it is also found in Africa, Asia, Australia, Europe, and South America
535 (Cierjacks et al., 2013). As such, *Robinia pseudoacacia* is representative of temperate N-fixing trees.

Although N-fixing trees are generally important pioneer species and can relieve strong N limitation of plant growth in early
succession (Chapin III et al., 1994), N-fixing trees can also be strong competitors. As such, in addition to having a
facilitative effect on neighbouring plant growth (Hulvey et al., 2013), they can also have no effect on neighbouring plant
growth (Lai et al., 2018; Xu et al., 2020) or an inhibitory effect on neighbouring plant growth (Chapin III et al., 2016; Taylor
540 et al., 2017). This depends on abiotic and biotic factors (Staccone et al., 2021) and could be explored further with LM4.1-
BNF.

The LM4.1-BNF representation of BNF, while implemented and validated in a temperate forest, can be expanded to other
terrestrial ecosystems, such as tropical and boreal forests. This will require parameterisation of representative N-fixing and
non-fixing tree species but will not require re-structuring the model equations. Furthermore, this representation of BNF could
545 be incorporated into other land models.

7 Conclusions

Here we present LM4.1-BNF: an updated representation of BNF and other aspects of N cycling in LM4.1, which is the land
component of the GFDL Earth System Model (Zhao et al., 2018a, 2018b). LM4.1-BNF is the first land model to include a
representation of the competitive dynamics between N-fixing and non-fixing plants, a mechanistic representation of
550 asymbiotic BNF, and distinct asymbiotic and symbiotic BNF temperature responses derived from corresponding
observations. Comparisons of simulations with observations show that LM4.1-BNF captures observed forest growth,
successional dynamics, and major pools and fluxes of C and N and their temporal dynamics at population, community, and
ecosystem scales. Furthermore, LM4.1-BNF represents these more accurately than previous representations of BNF in land
models. By incorporating both the competitive dynamics between N-fixing and non-fixing plants, which is a key ecological
mechanism that determines ecosystem-scale symbiotic BNF, as well as asymbiotic BNF, LM4.1-BNF yields accurate
555 ecosystem-scale estimates of BNF and its temporal dynamics. Furthermore, the novel representation of soil NO and N₂O
emissions in LM4.1-BNF enables the estimation of the magnitude of the terrestrial NO and N₂O source, which can be driven
by BNF (Kou-Giesbrecht and Menge, submitted). The representation of these ecological mechanisms in LM4.1-BNF is
general and could be incorporated into other land models.



560 Extending LM4.1-BNF to other biomes and incorporating LM4.1-BNF within the GFDL Earth System Model would allow a
 more accurate assessment of the response of BNF and the terrestrial C sink to elevated atmospheric CO₂ concentration,
 which intensifies N limitation of plant growth (Terrer et al., 2019; Zheng et al., 2020), and elevated N deposition, which
 relieves N limitation of plant growth (Reay et al., 2008; Schulte-Uebbing and de Vries, 2018; Zheng et al., 2020). In
 particular, such an endeavour could address whether BNF and N deposition will provide sufficient N to sustain CO₂
 565 sequestration by terrestrial ecosystems under elevated atmospheric CO₂ concentration.

Appendix A Model Description

A.1 N uptake by roots and symbionts

All vegetation types take up inorganic N via passive and active root uptake. Processes in Section A.1 occur on the fast
 timescale (30 min).

570 A.1.1 Passive root uptake of inorganic N

The rate of passive root uptake of inorganic N in soil layer k ($N_{passive}(k)$; [kg N indiv⁻¹ yr⁻¹]) is

$$N_{passive}(k) = U_{H_2O}(k) \left(\frac{NO_3(k) + NH_4(k)}{H_2O(k)} \right) \quad (A1)$$

where $U_{H_2O}(k)$ is the water uptake flux in soil layer k [kg H₂O m⁻² yr⁻¹], $NO_3(k)$ is the NO₃⁻ pool in soil layer k [kg N m⁻²],
 $NH_4(k)$ is the NH₄⁺ pool in soil layer k [kg N m⁻²], and $H_2O(k)$ is the soil water content of soil layer k [kg H₂O m⁻²].

575 A.1.2 Active root uptake of inorganic N

The rate of active root uptake of inorganic N in soil layer k ($N_{active}(k)$; [kg N indiv⁻¹ yr⁻¹]) is

$$N_{active}(k) = \left(r_{NO_3} \frac{NO_3(k) f_{rhiz}(k) / \Delta z(k)}{NO_3(k) f_{rhiz}(k) / \Delta z(k) + k_{M,NO_3}} \frac{\Delta z(k)}{n_{indiv}} + r_{NH_4} \frac{NH_4(k) f_{rhiz}(k) / \Delta z(k)}{NH_4(k) f_{rhiz}(k) / \Delta z(k) + k_{M,NH_4}} \frac{\Delta z(k)}{n_{indiv}} \right) N_{stress} \quad (A2)$$

where r_{NO_3} and r_{NH_4} are rate constants, $f_{rhiz}(k)$ is the rhizosphere volume fraction of soil layer k [m³ m⁻³], $\Delta z(k)$ is the
 thickness of soil layer k, k_{M,NO_3} and k_{M,NH_4} are half-saturation constants, and n_{indiv} is the spatial density [indiv m⁻²].

580 $f_{rhiz}(k)$ is calculated as

$$f_{rhiz}(k) = \pi((r_{rhiz} + r_{root})^2 - r_{root}^2) C_r(k) SRL \frac{n_{indiv}}{\Delta z(k)} \quad (A3)$$

where r_{rhiz} is the radius of the rhizosphere around fine roots, r_{root} is the radius of fine roots, $C_r(k)$ is the biomass C of fine
 roots in soil layer k [kg C indiv⁻¹], and SRL is the specific root length. N stress of the plant (N_{stress} ; [unitless]) is given in Eq.
 (A34). This follows LM3-SNAP but is modified to increase with N stress.



585 A.1.3 Inorganic N uptake by arbuscular mycorrhizae

The rates of NO_3^- and NH_4^+ uptake by AM in soil layer k ($N_{AM,\text{NO}_3}(k)$ and $N_{AM,\text{NH}_4}(k)$ respectively; [$\text{kg N indiv}^{-1} \text{ yr}^{-1}$]) are

$$N_{AM}(k) = r_{\text{NO}_3,AM} \frac{\frac{\text{NO}_3(k)}{\Delta z(k)}}{\frac{\text{NO}_3(k)}{\Delta z(k)} + k_{AM,\text{NO}_3}} \frac{\frac{B_{AM}(k)n_{\text{indiv}}}{\Delta z(k)}}{\frac{B_{AM}(k)n_{\text{indiv}}}{\Delta z(k)} + k_{AM}}{n_{\text{indiv}}}} \Delta z(k) + r_{\text{NH}_4,AM} \frac{\frac{\text{NH}_4(k)}{\Delta z(k)}}{\frac{\text{NH}_4(k)}{\Delta z(k)} + k_{AM,\text{NH}_4}} \frac{\frac{B_{AM}(k)n_{\text{indiv}}}{\Delta z(k)}}{\frac{B_{AM}(k)n_{\text{indiv}}}{\Delta z(k)} + k_{AM}}{n_{\text{indiv}}}} \Delta z(k) \quad (\text{A4})$$

where $r_{\text{NO}_3,AM}$ and $r_{\text{NH}_4,AM}$ are rate constants, k_{AM,NO_3} and k_{AM,NH_4} are half-saturation constants, $B_{AM}(k)$ is the biomass C of AM in soil layer k [kg C indiv^{-1}], and k_{AM} is a half-saturation constant. This follows LM3-SNAP.

590 A.1.4 Organic and inorganic N uptake by ectomycorrhizae

The rates of C and N decomposition by EM in soil layer k of organic matter type i , where i = labile plant-derived, labile microbe-derived, or recalcitrant, ($D_{C,i,EM}(k)$ and $D_{N,i,EM}(k)$ respectively; [$\text{kg C indiv}^{-1} \text{ yr}^{-1}$] and [$\text{kg N indiv}^{-1} \text{ yr}^{-1}$]) are

$$D_{C,i,EM}(k) = \frac{V_{EM,max,ref,i}}{\exp(-E_{a,i}/RT_{ref})} \exp(-E_{a,i}/RT(k)) \frac{\left(\frac{\theta(k)}{\theta_{sat}}\right)^3 \left(1 - \frac{\theta(k)}{\theta_{sat}}\right)^{2.5}}{f_{\theta,max}} \frac{C_{U,i}(k)}{n_{\text{indiv}}} \frac{B_{EM}(k)/C_{U,i}(k)}{B_{EM}(k)/C_{U,i}(k) + k_{M,EM}} \quad (\text{A5})$$

$$D_{N,i,EM}(k) = \frac{V_{EM,max,ref,i}}{\exp(-E_{a,i}/RT_{ref})} \exp(-E_{a,i}/RT(k)) \frac{\left(\frac{\theta(k)}{\theta_{sat}}\right)^3 \left(1 - \frac{\theta(k)}{\theta_{sat}}\right)^{2.5}}{f_{\theta,max}} \frac{N_{U,i}(k)}{n_{\text{indiv}}} \frac{B_{EM}(k)/C_{U,i}(k)}{B_{EM}(k)/C_{U,i}(k) + k_{M,EM}} \quad (\text{A6})$$

595 where $V_{EM,max,ref,i}$ is the maximum decomposition rate of organic matter type i , $E_{a,i}$ is the activation energy of the decomposition of organic matter type i , R is the ideal gas constant, T_{ref} is reference temperature, $T(k)$ is soil temperature of soil layer k [K], $\theta(k)$ is volumetric soil water content of soil layer k [$\text{m}^3 \text{ m}^{-3}$], θ_{sat} is saturation volumetric soil water content [$\text{m}^3 \text{ m}^{-3}$], $f_{\theta,max}$ is a factor normalizing the dependence on $\theta(k)$ to a maximum value of 1 [unitless], $B_{EM}(k)$ is the biomass C of EM in soil layer k [kg C indiv^{-1}], $C_{U,i}(k)$ is the soil C pool of type i in soil layer k [kg C m^{-2}], $N_{U,i}(k)$ is the soil N pool
 600 type i in soil layer k [kg C m^{-2}], and $k_{M,EM}$ is a half-saturation constant. This follows LM3-SNAP.

The rates of C and N uptake by EM in soil layer k ($C_{EM}(k)$ and $N_{EM}(k)$ respectively; [$\text{kg C indiv}^{-1} \text{ yr}^{-1}$] and [$\text{kg N indiv}^{-1} \text{ yr}^{-1}$]) are

$$C_{EM}(k) = \sum_i \varepsilon_{C,i,EM} D_{C,i,EM}(k) \quad (\text{A7})$$

$$N_{EM}(k) = \sum_i \varepsilon_{N,i,EM} D_{N,i,EM}(k) + r_{\text{NO}_3,EM} \frac{\frac{\text{NO}_3(k)}{\Delta z(k)}}{\frac{\text{NO}_3(k)}{\Delta z(k)} + k_{EM,\text{NO}_3}} \frac{\frac{B_{EM}(k)n_{\text{indiv}}}{\Delta z(k)}}{\frac{B_{EM}(k)n_{\text{indiv}}}{\Delta z(k)} + k_{EM}}{n_{\text{indiv}}}} \Delta z(k) + r_{\text{NH}_4,EM} \frac{\frac{\text{NH}_4(k)}{\Delta z(k)}}{\frac{\text{NH}_4(k)}{\Delta z(k)} + k_{EM,\text{NH}_4}} \frac{\frac{B_{EM}(k)n_{\text{indiv}}}{\Delta z(k)}}{\frac{B_{EM}(k)n_{\text{indiv}}}{\Delta z(k)} + k_{EM}}{n_{\text{indiv}}}} \Delta z(k) \quad (\text{A8})$$

where $\varepsilon_{C,i,EM}$ is the C uptake efficiency of soil C type i by EM, $\varepsilon_{N,i,EM}$ is the N uptake efficiency of soil N type i by EM, $r_{\text{NO}_3,EM}$ and $r_{\text{NH}_4,EM}$ are rate constants, k_{EM,NO_3} and k_{EM,NH_4} are half-saturation constants, and k_{EM} is a half-saturation constant. $\sum_i (1 - \varepsilon_{C,i,EM}) D_{C,i,EM}(k)$ is released as CO_2 . $\sum_i (1 - \varepsilon_{N,i,EM}) D_{N,i,EM}(k)$ enters $\text{NH}_4(k)$. This follows LM3-SNAP but is modified to additionally take up inorganic N.



610 A.1.5 Symbiotic BNF by N-fixing bacteria

The symbiotic BNF rate by N-fixing bacteria (N_{Nfix} ; [kg N indiv^{-1} yr $^{-1}$]) is

$$N_{Nfix} = r_{Nfix} B_{Nfix} f(T) \quad (\text{A9})$$

where r_{Nfix} is a rate constant, B_{Nfix} is the biomass C of the nodule (includes both plant and N-fixing bacteria tissue) [kg C indiv^{-1}], and $f(T)$ is the soil temperature dependence function. For *Robinia*,

$$615 \quad f(T) = \max \left[0.0, \left(\frac{45.67 - (T - 273.15)}{45.67 - 31.89} \right) \left(\frac{(T - 273.15) + 1.43}{31.89 + 1.43} \right)^{\frac{31.89 + 1.43}{45.67 - 31.89}} \right] \quad (\text{A10})$$

where T is the average soil temperature across soil layers [K]. This reaches its maximum at 31.9 °C (Fig. D1). This is derived from Bytnerowicz et al., (submitted).

A.2 Asymbiotic BNF

Soil microbes are represented as a single C pool that conducts decomposition, nitrification, denitrification, and asymbiotic
 620 BNF. The rates of C and N decomposition by soil microbes in soil layer k of organic matter type i , where i = labile plant-derived, labile microbe-derived or recalcitrant, ($D_{C,i}(k)$ and $D_{N,i}(k)$ respectively; [kg C m^{-2} yr $^{-1}$] and [kg N m^{-2} yr $^{-1}$]) are

$$D_{C,i}(k) = \frac{v_{max,ref,i}}{\exp(-E_{a,i}/RT_{ref})} \exp(-E_{a,i}/RT(k)) \frac{\left(\frac{\theta(k)}{\theta_{sat}}\right)^3 \left(1 - \frac{\theta(k)}{\theta_{sat}}\right)^{2.5}}{f_{\theta,max}} C_{U,i}(k) \frac{C_M(k)/C_{U,i}(k)}{C_M(k)/C_{U,i}(k) + k_M} \quad (\text{A11})$$

$$D_{N,i}(k) = \frac{v_{max,ref,i}}{\exp(-E_{a,i}/RT_{ref})} \exp(-E_{a,i}/RT(k)) \frac{\left(\frac{\theta(k)}{\theta_{sat}}\right)^3 \left(1 - \frac{\theta(k)}{\theta_{sat}}\right)^{2.5}}{f_{\theta,max}} N_{U,i}(k) \frac{C_M(k)/C_{U,i}(k)}{C_M(k)/C_{U,i}(k) + k_M} \quad (\text{A12})$$

where $v_{max,ref,i}$ is the maximum decomposition rate of organic matter type i , $C_M(k)$ is the biomass C of soil microbes in soil
 625 layer k [kg C m^{-2}], and k_M is the half-saturation constant. This follows LM3-SNAP.

The potential rates of C and N decomposition during denitrification by soil microbes in soil layer k of organic matter type i ($D_{C,i,denit,pot}(k)$ and $D_{N,i,denit,pot}(k)$ respectively; [kg C m^{-2} yr $^{-1}$] and [kg N m^{-2} yr $^{-1}$]) are

$$D_{C,i,denit,pot}(k) = \frac{v_{denit,max,ref,i}}{\exp(-E_{a,i}/RT_{ref})} \exp(-E_{a,i}/RT(k)) \frac{\left(\frac{\theta(k)}{\theta_{sat}}\right)^{5.5}}{f_{\theta,max}} C_{U,i}(k) \frac{C_M(k)/C_{U,i}(k)}{C_M(k)/C_{U,i}(k) + k_{M,denit}} \quad (\text{A13})$$

$$D_{N,i,denit,pot}(k) = \frac{v_{denit,max,ref,i}}{\exp(-E_{a,i}/RT_{ref})} \exp(-E_{a,i}/RT(k)) \frac{\left(\frac{\theta(k)}{\theta_{sat}}\right)^{5.5}}{f_{\theta,max}} N_{U,i}(k) \frac{C_M(k)/C_{U,i}(k)}{C_M(k)/C_{U,i}(k) + k_{M,denit}} \quad (\text{A14})$$

630 where $v_{denit,max,ref,i}$ is the maximum decomposition rate of organic matter type i during denitrification and $k_{M,denit}$ is the half-saturation constant. This follows LM3-SNAP.

The rates of C and N decomposition during denitrification by soil microbes in soil layer k of organic matter type i ($D_{C,i,denit}(k)$ and $D_{N,i,denit}(k)$ respectively; [kg C m^{-2} yr $^{-1}$] and [kg N m^{-2} yr $^{-1}$]) are

$$D_{C,i,denit}(k) = \frac{D_{C,i,denit,pot}(k) NO_3(k)}{NO_3(k) + k_{denit} f_{denit} \sum_i D_{C,i,denit,pot}(k)} \quad (\text{A15})$$



$$635 \quad D_{N,i,denit}(k) = \frac{D_{N,i,denit,pot}(k)NO_3(k)}{NO_3(k)+k_{denit}f_{denit}\sum_i D_{C,i,denit,pot}(k)} \quad (A16)$$

where k_{denit} is the half-saturation constant, and f_{denit} is the stoichiometric ratio of NO_3^- demand for C decomposition. $D_{C,i,denit}(k)$ and $D_{N,i,denit}(k)$ include NO_3^- limitation of denitrification. This follows LM3-SNAP.

The rate of change of biomass C of soil microbes in soil layer k ($\frac{dC_M(k)}{dt}$; [kg C m⁻² yr⁻¹]) is

$$\frac{dC_M(k)}{dt} = G_{M,C}(k) - \frac{C_M(k)}{\tau_M} \quad (A17)$$

640 where $G_{M,C}(k)$ is the C growth rate of soil microbes in soil layer k (Eq. (A23)) and τ_M is the combined maintenance respiration and turnover time of soil microbes. This follows LM3-SNAP.

The rate of change of biomass N of soil microbes in soil layer k ($\frac{dN_M(k)}{dt}$; [kg N m⁻² yr⁻¹]) is

$$\frac{dN_M(k)}{dt} = G_{M,N}(k) - \frac{C_M(k)\varepsilon_{t,M}}{C:N_M} \quad (A18)$$

645 where $G_{M,N}(k)$ is the N growth rate of soil microbes in soil layer k (Eq. (A24)), $\varepsilon_{t,M}$ is the fraction of maintenance respiration in combined maintenance respiration and turnover, and $C:N_M$ is the C:N ratio of soil microbes. This follows LM3-SNAP.

The maintenance respiration rate of soil microbes in soil layer k ($R_{maint}(k)$; [kg C m⁻² yr⁻¹]) is

$$R_{maint}(k) = \frac{C_M(k)}{\tau_M}(1 - \varepsilon_{t,M}) \quad (A19)$$

This is released as CO₂. This follows LM3-SNAP.

650 The asymbiotic BNF rate of soil microbes in soil layer k ($N_{Nfix\ asymb}(k)$; [kg N m⁻² yr⁻¹]) is

$$N_{Nfix\ asymb}(k) = r_{Nfix\ asymb}C_M(k)f(T(k)) \quad (A20)$$

where $r_{Nfix\ asymb}$ is a rate constant and $f(T(k))$ is the soil temperature dependence function.

$$f(T(k)) = e^{-2.6+0.21(T(k)-273.15)\left(1-\frac{0.5(T(k)-273.15)}{24.4}\right)} \quad (A21)$$

655 which reaches its maximum at 24.4 °C (Fig. D1). This is derived from the observations compiled by Houlton et al., 2008 with the study of symbiotic BNF removed (Schomberg and Weaver, 1992) and is normalized to a maximum of 1.

The N surplus or deficit of soil microbes in soil layer k ($\phi_N(k)$; [kg N m⁻² yr⁻¹]) is

$$\phi_N(k) = \sum_i \varepsilon_{N,i}D_{N,i}(k) + \sum_i \varepsilon_{N,i}D_{N,i,denit}(k) + N_{Nfix,asymb}(k) + \frac{\sum_i \varepsilon_{C,i}D_{C,i}(k) + \sum_i \varepsilon_{C,i}D_{C,i,denit}(k) - R_{maint}(k)}{C:N_M} \quad (A22)$$

660 where $\varepsilon_{N,i}$ is the N uptake efficiency of soil N type i by soil microbes and $\varepsilon_{C,i}$ is the C uptake efficiency of soil C type i by soil microbes. $\phi_N(k) > 0$ indicates net N mineralization (N surplus) and $\phi_N(k) < 0$ indicates net N immobilization (N deficit) by soil microbes in soil layer k. $\sum_i(1 - \varepsilon_{C,i})(D_{C,i}(k) + D_{C,i,denit}(k))$ is released as CO₂. $\sum_i(1 - \varepsilon_{N,i})(D_{N,i}(k) + D_{N,i,denit}(k))$ enters $NH_4(k)$. This follows LM3-SNAP but was modified to include asymbiotic BNF.

$G_{M,C}(k)$ and $G_{M,N}(k)$ depend on whether growth of soil microbes is C-limited ($\phi_N(k) \geq -Imm_{max}(k)$) or N-limited ($\phi_N(k) < -Imm_{max}(k)$) and are calculated as



$$G_{M,C}(k) = \begin{cases} \sum_i \varepsilon_{C,i} D_{C,i}(k) + \sum_i \varepsilon_{C,i} D_{C,i,denit}(k), & \phi_N(k) \geq -Imm_{max}(k) \\ C:N_M [\sum_i \varepsilon_{N,i} D_{N,i}(k) + \sum_i \varepsilon_{N,i} D_{N,i,denit}(k) + N_{Nfix,asymb}(k) + Imm_{max}(k)] + R_{maint}(k), & \phi_N(k) < -Imm_{max}(k) \end{cases} \quad (A23)$$

$$G_{M,N}(k) = \begin{cases} \frac{\sum_i \varepsilon_{C,i} D_{C,i}(k) + \sum_i \varepsilon_{C,i} D_{C,i,denit}(k) - R_{maint}(k)}{C:N_M}, & \phi_N(k) \geq -Imm_{max}(k) \\ \sum_i \varepsilon_{N,i} D_{N,i}(k) + \sum_i \varepsilon_{N,i} D_{N,i,denit}(k) + N_{Nfix,asymb}(k) + Imm_{max}(k), & \phi_N(k) < -Imm_{max}(k) \end{cases} \quad (A24)$$

670 This follows LM3-SNAP but was modified to include asymbiotic BNF.

The maximum N immobilization rate of soil microbes in soil layer k ($Imm_{max}(k)$; [kg N m⁻² yr⁻¹]) is

$$Imm_{max}(k) = \left(V_{max,ref,NH_4} \exp\left(\frac{-E_{a,NH_4}}{RT(k)}\right) NH_4(k) + V_{max,ref,NO_3} \exp\left(\frac{-E_{a,NO_3}}{RT(k)}\right) NO_3(k) \right) \frac{\left(\frac{\theta(k)}{\theta_{sat}}\right)^3 \left(1 - \frac{\theta(k)}{\theta_{sat}}\right)^{2.5}}{f_{\theta,max}} \quad (A25)$$

675 where V_{max,ref,NH_4} is the maximum NH₄⁺ immobilization rate, E_{a,NH_4} is the activation energy of NH₄⁺ immobilization, V_{max,ref,NO_3} is the maximum NO₃⁻ immobilization rate, and E_{a,NO_3} is the activation energy of NO₃⁻ immobilization. This follows LM3-SNAP.

When growth of soil microbes in soil layer k is N-limited ($\phi_N(k) < -Imm_{max}(k)$), there is overflow respiration of excess C. The overflow respiration of excess C in soil layer k ($R_{overflow}(k)$; [kg C m⁻² yr⁻¹]) is

$$R_{overflow}(k) = \begin{cases} 0, & \phi_N(k) \geq -Imm_{max}(k) \\ -(\phi_N(k) + Imm_{max}(k)) C:N_M, & \phi_N(k) < -Imm_{max}(k) \end{cases} \quad (A26)$$

680 This is released as CO₂.

If $\phi_N(k) > 0$, the net N mineralization flux in soil layer k is $\phi_N(k)$. If $-Imm_{max}(k) \leq \phi_N(k) \leq 0$, the net N immobilization flux in soil layer k is $-\phi_N(k)$. If $\phi_N(k) < -Imm_{max}(k)$, the net N immobilization flux in soil layer k is $-Imm_{max}(k)$. The fraction of the net N immobilization flux in soil layer k that is NH₄⁺ immobilization is

$$\frac{\frac{V_{max,ref,NH_4}}{\exp\left(\frac{-E_{a,NH_4}}{RT_{ref}}\right)} \exp\left(\frac{-E_{a,NH_4}}{RT(k)}\right) NH_4(k)}{\frac{V_{max,ref,NH_4}}{\exp\left(\frac{-E_{a,NH_4}}{RT_{ref}}\right)} \exp\left(\frac{-E_{a,NH_4}}{RT(k)}\right) NH_4(k) + \frac{V_{max,ref,NO_3}}{\exp\left(\frac{-E_{a,NO_3}}{RT_{ref}}\right)} \exp\left(\frac{-E_{a,NO_3}}{RT(k)}\right) NO_3(k)}. \quad \text{The fraction of the net N immobilization flux in soil layer k that is } NO_3^- \text{ immobilization is}$$

$$\frac{\frac{V_{max,ref,NO_3}}{\exp\left(\frac{-E_{a,NO_3}}{RT_{ref}}\right)} \exp\left(\frac{-E_{a,NO_3}}{RT(k)}\right) NO_3(k)}{\frac{V_{max,ref,NH_4}}{\exp\left(\frac{-E_{a,NH_4}}{RT_{ref}}\right)} \exp\left(\frac{-E_{a,NH_4}}{RT(k)}\right) NH_4(k) + \frac{V_{max,ref,NO_3}}{\exp\left(\frac{-E_{a,NO_3}}{RT_{ref}}\right)} \exp\left(\frac{-E_{a,NO_3}}{RT(k)}\right) NO_3(k)}. \quad \text{This follows LM3-SNAP.}$$

685 flux in soil layer k that is NO₃⁻ immobilization is



Processes in Section A.2 occur on the fast timescale (30 min).

A.3 Plant growth and N limitation

The non-structural C pool (NSC ; [kg C indiv^{-1}]) gains C from photosynthesis. NSC loses C to respiration and C allocation to growth, symbionts, and root C exudation. The rate of change of NSC ($\frac{dNSC}{dt}$; [kg C indiv^{-1} yr^{-1}]) is

$$\frac{dNSC}{dt} = P - R - (G_{C,l} + G_{C,r} + G_{C,sw} + G_{C,seed}) - C_{alloc} - L_{C,exudate} \quad (\text{A27})$$

where P is the photosynthesis rate [kg C indiv^{-1} yr^{-1}], R is the respiration rate (maintenance and growth) [kg C indiv^{-1} yr^{-1}], $G_{C,l}$ is the growth rate of the leaf C pool (C_l ; [kg C indiv^{-1}]) [kg C indiv^{-1} yr^{-1}], $G_{C,r}$ is the growth rate of the fine root C pool (C_r ; [kg C indiv^{-1}]) [kg C indiv^{-1} yr^{-1}], $G_{C,sw}$ is the growth rate of the sapwood C pool (C_{sw} ; [kg C indiv^{-1}]) [kg C indiv^{-1} yr^{-1}], $G_{C,seed}$ is growth rate of the seed C pool (C_{seed} ; [kg C indiv^{-1}]) [kg C indiv^{-1} yr^{-1}], C_{alloc} is the rate of C allocation to symbionts (Eq. (A43-46)), and $L_{C,exudate}$ is the rate of root C exudation (Eq. (A38)). Note that sapwood is converted to heartwood following Martinez Cano et al., (2020).

The non-structural N pool (NSN ; [kg N indiv^{-1}]) gains N from N uptake via roots and symbionts. NSN loses N to N allocation to growth, symbionts, and root N exudation. The rate of change of NSN ($\frac{dNSN}{dt}$; [kg N indiv^{-1} yr^{-1}]) is

$$\frac{dNSN}{dt} = U - \left(\frac{G_{C,l}}{C:N_l} + \frac{G_{C,r}}{C:N_r} + \frac{G_{C,sw}}{C:N_{sw}} + \frac{G_{C,seed}}{C:N_{seed}} \right) - N_{alloc} - L_{N,exudate} \quad (\text{A28})$$

where U is the N uptake rate via roots and symbionts (Eq. (A55)) [kg N indiv^{-1} yr^{-1}], $C:N_l$ is the fixed C:N ratio of leaves, $C:N_r$ is the fixed C:N ratio of fine roots, $C:N_{sw}$ is the fixed C:N ratio of sapwood, $C:N_{seed}$ is the fixed C:N ratio of seeds, N_{alloc} is the rate of N allocation to symbionts (Eq. (A47)), and $L_{N,exudate}$ is the rate of root N exudation (Eq. (A39)).

Non-N-limited growth is calculated according to Weng et al., (2015). The total allocation of NSC to growth is determined by the target NSC (NSC_{target} ; [kg C indiv^{-1}]) and minimizes the deviation between NSC and NSC_{target} . NSC_{target} is a multiple of the target C_l ($C_{l,target}$; [kg C indiv^{-1}]), which reflects the ability of a plant to re-leaf after defoliation (Hoch et al., 2003; Richardson et al., 2013), and is calculated as

$$NSC_{target} = q C_{l,target} \quad (\text{A29})$$

where q is a proportionality constant. The allocation of NSC to the growth of each tissue depends on the total allocation of NSC to growth and the target C pool of each tissue, and minimizes the deviation between the C pool of each tissue and the target C pool of each tissue. The target C pool of each tissue is dynamic and is determined by allometry (Eq. (A40-42)).

In LM4.1-BNF, $G_{C,l}$, $G_{C,r}$, $G_{C,sw}$, and $G_{C,seed}$ are adjusted to include N limitation and are calculated as

$$G_{C,l} = (1 - N_{stress})\Delta_l \quad (\text{A30})$$

$$G_{C,r} = \Delta_r \quad (\text{A31})$$

$$G_{C,sw} = (1 - N_{stress})\Delta_{sw} \quad (\text{A32})$$

$$G_{C,seed} = (1 - N_{stress})\Delta_{seed} \quad (\text{A33})$$



where N_{stress} is N stress [unitless] and $\Delta_l, \Delta_r, \Delta_{sw}$, and Δ_{seed} are the non-N-limited growth rates of C_l, C_r, C_{sw} , and C_{seed} respectively [kg C indiv⁻¹ yr⁻¹] following Weng et al., (2015). Because plants increase C allocation to fine roots relative to other tissues when N-limited (Poorter et al., 2012), $G_{C,r}$ is not adjusted to include N limitation.

720 In LM4.1-BNF, N_{stress} is the relative difference between NSN and NSN_{target} and is calculated as

$$N_{stress} = \max \left[0, \frac{NSN_{target} - NSN}{NSN_{target}} \right] \quad (A34)$$

where NSN_{target} is the target NSN [kg N indiv⁻¹]. N_{stress} is smoothed with a low-pass filter over 30 days to reflect the persisting influence of N stress (Mooney et al., 1991). NSN_{target} is calculated as

$$NSN_{target} = \frac{NSC_{target}}{C:N_l} \quad (A35)$$

725 This is similar to LM3-SNAP, which compared the target leaf and root N pools to NSN , but is modified to reflect the treatment of NSC_{target} in LM4.1 by including the target sapwood and seed N pools.

N demand for plant growth (N_{demand} ; [kg N indiv⁻¹]) is

$$N_{demand} = \frac{G_{C,l}}{C:N_l} + \frac{G_{C,r}}{C:N_r} + \frac{G_{C,sw}}{C:N_{sw}} + \frac{G_{C,seed}}{C:N_{seed}} \quad (A36)$$

If $N_{demand} > 0.5 NSN$, $G_{C,l}, G_{C,r}, G_{C,sw}$, and $G_{C,seed}$ are reduced to prevent N_{demand} from depleting NSN

$$730 \quad G_{C,t} = \begin{cases} G_{C,t}, & N_{demand} < 0.5 NSN \\ G_{C,t} \frac{NSN}{N_{demand}}, & N_{demand} > 0.5 NSN \end{cases} \quad (A37)$$

where t = leaf, root, sapwood, or seed and 0.5 was set to maintain a baseline NSN .

Plant turnover decreases C_l, C_r , and C_{sw} and from N_l, N_r , and N_{sw} at a constant tissue-specific rate and enters $C_{U,labile\ plant-derived}$ or $C_{U,recalcitrant}$ and $N_{U,labile\ plant-derived}$ or $N_{U,recalcitrant}$ respectively. A fraction of the turnover of C_l and N_l is retranslocated into NSC and NSN respectively.

735 Under N limitation, plants increase root C exudation to stimulate N mineralization in the rhizosphere (rhizosphere priming; Cheng et al., 2014; Finzi et al., 2015). $L_{C,exudate}$ increases with N_{stress} and is calculated as

$$L_{C,exudate} = r_{leakage,C} NSC N_{stress} \quad (A38)$$

where $r_{leakage,C}$ is a rate constant. $L_{C,exudate}$ enters the rhizosphere $C_{U,labile\ plant-derived}$.

740 Under N limitation, plants decrease root N exudation (Canarini et al., 2019). $L_{N,exudate}$ decreases with N_{stress} and is calculated as

$$L_{N,exudate} = r_{leakage,N} NSN (1 - N_{stress}) \quad (A39)$$

where $r_{leakage,N}$ is a rate constant. $L_{N,exudate}$ enters the rhizosphere $N_{U,labile\ plant-derived}$.

745 Plant growth occurs during the growing season. Plant maintenance respiration and plant turnover occur throughout the year. After the transition from the growing season to the non-growing season, turnover of C_l and N_l occur at an elevated rate until $C_l = 0$ and $N_l = 0$. Plant C allocation to symbionts occurs during the growing season. Root exudation occurs during the



growing season. Symbiont growth, maintenance respiration, and turnover occur throughout the year. N uptake by roots and symbionts occurs throughout the year.

Photosynthesis and respiration occur on the fast timescale (30min). Plant turnover occurs on the fast timescale (30 min).

Plant growth, plant C allocation to symbionts, and root exudation occur on the daily timescale.

750 **A.4 Plant allometry**

Plant allometry follows Martinez Cano et al., 2020. Crown area (CA ; [m^2]) is a function of diameter at breast height (power function)

$$CA = \alpha_{CA} D^{\theta_{CA}} \quad (A40)$$

where D is diameter at breast height [m], and α_{CA} and θ_{CA} are allometry parameters.

755 Height (H ; [m]) is a function of diameter at breast height (generalized Michaelis-Menten equation)

$$H = \frac{\alpha_{HT} D^{\theta_{HT}}}{\gamma_{HT} + D^{\theta_{HT}}} \quad (A41)$$

where α_{HT} , θ_{HT} , and γ_{HT} are allometry parameters.

Wood mass (W ; [kg C]) is a function of diameter at breast height and height

$$W = \alpha_{BM} \rho_{wood} D^2 H \quad (A42)$$

760 where α_{BM} is an allometry parameter and ρ_{wood} is wood density.

A.5 Plant C allocation to symbionts (AM, EM and N-fixing bacteria)

The rate of C allocation to AM ($C_{alloc,AM}$; [kg C $indiv^{-1}$ yr^{-1}]) is

$$C_{alloc,AM} = f_{alloc,AM} NSC \quad (A43)$$

765 where $f_{alloc,AM}$ is the fraction of NSC allocated to AM per unit time. $C_{alloc,AM}$ is not related to N_{stress} because, although AM increase N uptake, AM is maintained by the plant primarily for phosphorus uptake (Smith and Smith, 2011).

The rate of C allocation to EM ($C_{alloc,EM}$; [kg C $indiv^{-1}$ yr^{-1}]) is

$$C_{alloc,EM} = f_{alloc,EM} NSC N_{stress} \quad (A44)$$

where $f_{alloc,EM}$ is the maximum fraction of NSC allocated to EM per unit time. $C_{alloc,EM}$ is a function of N_{stress} because biomass C of EM increases with N limitation (Phillips et al., 2013).

770 Plants that associate with N-fixing bacteria can regulate symbiotic BNF to different extents, termed their BNF strategy (Menge et al., 2015). For plants with a perfectly facultative BNF strategy, symbiotic BNF increases with N limitation. For plants with an incomplete BNF strategy, symbiotic BNF increases with N limitation but is maintained at a minimum. For plants with an obligate BNF strategy, symbiotic BNF is constant. For plants with either a facultative or an incomplete BNF strategy, the rate of C allocation by the plant to N-fixing bacteria ($C_{alloc,Nfix}$; [kg C $indiv^{-1}$ yr^{-1}]) is

$$775 \quad C_{alloc,Nfix} = \max[f_{alloc,Nfix} NSC N_{stress}, f_{alloc,Nfix,min} NSC] \quad (A45)$$



where $f_{alloc,Nfix}$ is the fraction of NSC allocated to N-fixing bacteria per unit time and $f_{alloc,Nfix,min}$ is the minimum fraction of NSC allocated to N-fixing bacteria per unit time. For plants with a perfectly facultative BNF strategy $f_{alloc,Nfix,min} = 0$ and for plants with an incomplete down-regulator BNF strategy $f_{alloc,Nfix,min} > 0$. We model *Robinia* with an incomplete down-regulator BNF strategy.

780 For plants with an obligate BNF strategy, $C_{alloc,Nfix}$ is

$$C_{alloc,Nfix} = f_{alloc,Nfix} NSC \quad (A46)$$

Additionally, plants allocate a small quantity of N to symbionts such that symbiont growth can be initiated. The rate of N allocation by the plant to symbionts ($N_{alloc,j}$; $j = AM, EM, N\text{-fixing bacteria}$; [$\text{kg N indiv}^{-1} \text{ yr}^{-1}$]) is

$$N_{alloc,j} = \frac{C_{alloc,j}}{C:N_{alloc}} \quad (A47)$$

785 where $C:N_{alloc}$ is the C:N ratio of C and N allocated to symbionts by the plant.

Processes in Section A.5 occur on the daily timescale.

A.6 Growth and turnover of symbionts

Plant C allocation to symbionts is transferred to an intermediate C pool ($C_{int,j}$; $j = AM, EM, \text{ or } N\text{-fixing bacteria}$; [kg C indiv^{-1}]). The rate of change of $C_{int,j}$ ($\frac{dC_{int,j}}{dt}$; [$\text{kg C indiv}^{-1} \text{ yr}^{-1}$]) is

$$790 \quad \frac{dC_{int,j}}{dt} = C_{alloc,j} - \frac{G_j}{\varepsilon_{symb}} \quad (A48)$$

where ε_{symb} is the proportion of C uptake by a symbiont from $C_{int,j}$ that is assimilated. The C growth rate of a symbiont with strategy j (G_j ; $j = AM \text{ or } N\text{-fixing bacteria}$; [$\text{kg C indiv}^{-1} \text{ yr}^{-1}$]) is

$$G_j = \varepsilon_{symb} r_{growth} C_{int,j} \quad (A49)$$

where r_{growth} is the growth rate of a symbiont.

795 The growth rate of EM (G_{EM} ; [$\text{kg C indiv}^{-1} \text{ yr}^{-1}$]) is

$$G_{EM} = \varepsilon_{symb} (r_{growth} C_{int,EM} + C_{EM}) \quad (A50)$$

$\frac{G_j}{\varepsilon_{symb}} (1 - \varepsilon_{symb})$ ($j = AM, EM, \text{ or } N\text{-fixing bacteria}$) is released as CO_2 (growth respiration).

The rate of change of biomass C of a symbiont with strategy j ($\frac{dB_j}{dt}$; $j = AM \text{ or } EM$; [$\text{kg C indiv}^{-1} \text{ yr}^{-1}$]) is

$$\frac{dB_j}{dt} = G_j - \xi_j B_j - \frac{B_j}{\tau_j} \quad (A51)$$

800 where ξ_j is rate of maintenance respiration of a symbiont with strategy j and τ_j is the turnover time of a symbiont with

strategy j . $\xi_j B_j$ is released as CO_2 (maintenance respiration). $\frac{B_j}{\tau_j}$ enters $C_{U,labile \text{ microbe-derived}}$ and $\frac{B_j}{C:N_j \tau_j}$ enters

$N_{U,labile \text{ microbe-derived}}$, where $C:N_j$ is the C:N ratio of a symbiont with strategy j .

The rate of change of biomass C of N-fixing bacteria ($\frac{dB_{Nfix}}{dt}$; [$\text{kg C indiv}^{-1} \text{ yr}^{-1}$]) is



$$\frac{dB_{Nfix}}{dt} = G_{Nfix} - \xi_{Nfix}B_{Nfix} - cost_{Nfix}N_{Nfix} - \frac{B_{Nfix}}{\tau_{Nfix}} \quad (A52)$$

805 where ξ_{Nfix} is rate of maintenance respiration of N-fixing bacteria, $cost_{Nfix}$ is the C cost of symbiotic BNF per unit N, and τ_{Nfix} is the turnover time of N-fixing bacteria. $\xi_{Nfix}B_{Nfix}$ is released as CO₂ (maintenance respiration) and $cost_{Nfix}N_{Nfix}$ is released as CO₂ (respiration associated with symbiotic BNF). $\frac{B_{Nfix}}{\tau_{Nfix}}$ enters $C_{U,labile\ microbe-derived}$ and $\frac{B_{Nfix}}{C:N_{Nfix}\tau_{Nfix}}$ enters $N_{U,labile\ microbe-derived}$, where $C:N_{Nfix}$ is the C:N ratio of N-fixing bacteria.

N acquired by symbionts is transferred to an intermediate N pool ($N_{int,j}$; [kg N indiv⁻¹]). The rate of change of $N_{int,j}$ ($\frac{dN_{int,j}}{dt}$;
 810 $j = AM$ or EM ; [kg N indiv⁻¹ yr⁻¹]) is

$$\frac{dN_{int,j}}{dt} = \sum_k N_j + \frac{C_{alloc,j}}{C:N_{alloc}} - \frac{1}{C:N_j} (G_j - \xi_j B_j) - r_{up,veg} N_{int,j} \quad (A53)$$

where $r_{up,veg}$ is the rate of plant N uptake from $N_{int,j}$.

The rate of change of $N_{int,Nfix}$ ($\frac{dN_{int,Nfix}}{dt}$; [kg N indiv⁻¹ yr⁻¹]) is

$$\frac{dN_{int,Nfix}}{dt} = N_{Nfix} + \frac{C_{alloc,Nfix}}{C:N_{alloc}} - \frac{1}{C:N_{Nfix}} (G_{Nfix} - \xi_{Nfix}B_{Nfix} - cost_{Nfix}N_{Nfix}) - r_{up,veg} N_{int,Nfix} \quad (A54)$$

815 U is calculated as

$$U = \sum_k N_{passive}(k) + \sum_k N_{active}(k) + \sum_j r_{up,veg} N_{int,j} \quad (A55)$$

Processes in Section A.6 occur on the fast timescale (30 min). The purpose of $C_{int,j}$ and $N_{int,j}$ are to translate between the fast timescale (30min) and the daily timescale. Plant C allocation to symbionts occurs on the daily timescale (alongside plant growth), but plant N uptake occurs on the fast timescale.

820 A.7 Dynamic plant C allocation to N uptake relative to plant growth

The order of plant C allocation to growth, symbionts, and rhizosphere priming is determined by C limitation relative to N limitation (Cheng et al., 2014; Finzi et al., 2015; Poorter et al., 2012; Treseder, 2004; Zheng et al., 2019). If a plant is more C-limited than N-limited, $NSC < NSN \cdot C:N_{leaf}$. The plant allocates C to growth, then to N-fixing bacteria (if associated) and EM (if associated), then to rhizosphere priming, and finally to AM. If a plant more N-limited than C-limited, $NSC >$
 825 $NSN \cdot C:N_{leaf}$. The plant allocates C to N-fixing bacteria (if associated) and EM (if associated), then to rhizosphere priming, then to growth, and finally to AM.

A.8 Soil N₂O and NO emissions

Soil N₂O and NO emissions occur during nitrification (aerobic oxidation of NH₄⁺ with oxygen as an electron acceptor, which produces N₂O and NO as by-products) and denitrification (anaerobic oxidation of organic C with NO₃⁻ as an electron
 830 acceptor, which produces N₂O as a by-product).

Nitrification rate in soil layer k ($nit(k)$; [kg N m⁻² yr⁻¹]) is



$$nit(k) = \frac{V_{nit,max,ref}}{\exp\left(\frac{-E_{a,nit}}{RT_{ref}}\right)} \exp\left(\frac{-E_{a,nit}}{RT(k)}\right) \frac{\left(\frac{\theta(k)}{\theta_{sat}}\right)^3 \left(1 - \frac{\theta(k)}{\theta_{sat}}\right)^{2.5}}{f_{\theta,max}} NH_4(k) \quad (A56)$$

where $V_{nit,max,ref}$ is the maximum nitrification rate and $E_{a,nit}$ is the activation energy of nitrification. This follows LM3-SNAP.

835 Denitrification rate in soil layer k ($denit(k)$; [$\text{kg N m}^{-2} \text{ yr}^{-1}$]) is

$$denit(k) = f_{denit} \sum_i D_{C,i,denit}(k) \quad (A57)$$

This follows LM3-SNAP.

Soil N_2O emission rate in soil layer k ($N_2O(k)$; [$\text{kg N m}^{-2} \text{ yr}^{-1}$]) is

$$N_2O(k) = \gamma_{N_2O,nit} nit(k) + \frac{1}{1 + \max\left(3.52, 22 \exp\left(-0.8 \frac{NO_3(k)}{HR(k)}\right)\right) \max\left(0.1, 0.015 \frac{\theta(k)}{\theta_{sat}} - 0.32\right)} denit(k) \quad (A58)$$

840 where $\gamma_{N_2O,nit}$ is the proportion of the nitrification rate that is emitted as N_2O and $HR(k)$ is heterotrophic respiration in soil layer k (summation of maintenance respiration, overflow respiration, and decomposition respiration) [$\text{kg C m}^{-2} \text{ yr}^{-1}$]. This follows LM3V-N (Huang and Gerber, 2015).

Soil NO emission rate in soil layer k ($NO(k)$; [$\text{kg N m}^{-2} \text{ yr}^{-1}$]) is

$$NO(k) = \left(15.2 + \frac{35.5 \tan^{-1}\left(0.68\pi\left(2.09\left(1 - \frac{\theta(k)}{\theta_{sat}}\right)^{4/3} - 1.68\right)\right)}{\pi} \right) \gamma_{N_2O,nit} nit(k) \quad (A59)$$

845 This follows LM3V-N (Huang and Gerber, 2015).

Processes in Section A.8 occur on the fast timescale (30 min).

A.9 Additional N sources

N deposition enters $N_{U,labile\ plant-derived}$, NH_4 and NO_3 . Processes in Section A.9 occur on the fast timescale (30 min).

Appendix B Data availability and processing

850 The US Forest Inventory and Analysis (FIA) database (US Forest Service, 2020a) was downloaded in 2019 from https://apps.fs.usda.gov/fia/datamart/CSV/datamart_csv.html. For tree data, plots with Plot Design Code 1 (national plot design) were selected, dead and cut trees were excluded, trees measured at the root collar were excluded, trees with visually estimated or modelled heights were excluded, and accessible forest land was selected (excludes agriculture and urban areas). Canopy trees had crown class code open grown, dominant, or codominant. Understory trees had crown class code
 855 intermediate or overtopped. For soil data, mineral soil layers were selected. For seedling data, plots with Plot Design Code 1 (national plot design) were selected, and accessible forest land was selected (excludes agriculture and urban areas).



US FIA Forest Health Monitoring database (US Forest Service, 2020b) was downloaded in 2019 from https://www.fia.fs.fed.us/tools-data/other_data/index.php. Dead trees and trees with damaged crowns were excluded.
860 Timberland and woodland land uses were selected.

The Biomass and Allometry Database (BAAD; Falster et al., 2015) was downloaded from <https://esajournals.onlinelibrary.wiley.com/doi/abs/10.1890/14-1889.1>. Field / wild plants were selected. Temperate forest plants were selected.

TRY database data (Kattge et al., 2020) were downloaded in 2019 from <https://www.try-db.org/TryWeb/Prop0.php>.

865 Total soil C, total soil N, soil NH_4^+ , soil nitrate NO_3^- , N mineralization rate, net nitrification rate, and bulk density from CHL (Knoepp, 2009a, 2009b, 2018) were downloaded from https://coweeta.uga.edu/dbpublic/dataset_details.asp?accession=1124,
https://coweeta.uga.edu/dbpublic/dataset_details.asp?accession=1125, and
https://coweeta.uga.edu/dbpublic/dataset_details.asp?accession=1123.

Soil N_2O and NO emissions are from Stehfest and Bouwman, (2006). Data from deciduous temperate forests were selected.

870 Gross primary production, net primary production, heterotrophic respiration and net ecosystem production are from the ForC database (Anderson-Teixeira et al., 2018). Data from deciduous temperate broadleaf forests were selected.

Appendix C Model parameterization

Eq. (A40) was fit using nonlinear least-squares with the US FIA Forest Health Monitoring database (US Forest Service, 2020b) for each species to calculate α_{CA} and θ_{CA} (Fig. D8-9). Eq. (A41) was fit using nonlinear least-squares with the US
875 FIA database tree data (US Forest Service, 2020a) for each species to calculate α_{HT} , θ_{HT} and γ_{HT} (Fig. D10-11). Eq. (A42) was fit using nonlinear least-squares with the US FIA database tree data (US Forest Service, 2020a) for each species to calculate α_{BM} (Fig. D12-13). Note that wood mass (aboveground and belowground) is calculated by the US FIA using allometric equations of dbh.

880 The following equations from Weng et al., (2015) were utilized with the US FIA database tree data from consecutive censuses for each species (US Forest Service, 2020a) to calculate canopy tree background mortality rate (μ_C) and understory tree background mortality rate (μ_U)

$$n_{dead,C} = n_C(1 - e^{-\mu_C}) \quad (\text{A60})$$

$$n_{dead,U} = n_U \left(1 - e^{-\mu_U \left(1 + \frac{4e^{30(0.025-D_U)}}{1+e^{30(0.025-D_U)}} \right)} \right) \quad (\text{A61})$$

885 where $n_{dead,C}$ is the mean number of dead canopy individuals in a given year [indiv m^{-2}], n_C is the mean number of canopy individuals in a given year [indiv m^{-2}], $n_{dead,U}$ is the mean number of dead understory individuals in a given year [indiv m^{-2}],



n_U is the mean number of understory individuals in a given year [indiv m^{-2}], and D_U is the mean diameter at breast height of an understory individual in a given year [m].

890 The following equation from Martinez Cano et al., (2020) was fit using nonlinear least-squares with the BAAD (Falster et al., 2015) to calculate the proportionality constant between crown area and cross-sectional sapwood area (φ_{CSA}) (Fig. D14-15)

$$\varphi_{CSA} = \frac{SA_{sw}(y_{HT} + D^{\theta_{HT}})}{D^{\theta_{CA}} + \theta_{HT}} \quad (\text{A62})$$

895 where SA_{sw} is the cross-sectional sapwood area at breast height [m^2] and D is the diameter [m] from the BAAD (for deciduous angiosperm).

The following equation from Weng et al., (2015) was utilized with the BAAD (Falster et al., 2015) to calculate the proportionality constant between root area and leaf area (φ_{RL})

900 $\varphi_{RL} = \frac{c_r}{c_l} \cdot LMA \cdot SRA = 0.43 \cdot 0.0375 \left[\frac{\text{kg C}}{\text{m}^2} \right] \cdot 45 \left[\frac{\text{m}^2}{\text{kg C}} \right] = 0.79$, where 0.43 is from the BAAD (for deciduous angiosperm), 0.0375 kg C m^{-2} is from Poorter et al., (2009) (for deciduous trees), and 45 m^2 kg C is from Jackson et al., (1997) (for temperate deciduous forests).

The following equation from Martinez Cano et al., (2020) was utilized with the BAAD (Falster et al., 2015) to calculate the fraction of sapwood in branches (f_{br})

905 $f_{br} = \frac{c_{br}}{c_{sw}} = 0.255$, where 0.255 is from the BAAD. Note that we did not distinguish between deciduous angiosperm and evergreen gymnosperm due to insufficient data.

The following equation (from Eq. (A20)) was used to calculate $r_{Nfix\ asymb}$

910 $r_{Nfix\ asymb} = \frac{N_{Nfix\ asymb}}{C_M} = \frac{12 \left[\frac{\text{kg N}}{\text{ha yr}} \right]}{50\ 000 \left[\frac{\text{kg C}}{\text{ha}} \right] \times 0.01} = 0.024 \left[\frac{\text{kg N}}{\text{kg C yr}} \right]$ where 12 kg N ha^{-1} yr^{-1} is from Reed et al., (2011), 50000 kg C ha^{-1} is from Scharlemann et al., (2014), and 1% (0.01) is from Chapin et al., (2011).

The following equation (from Eq. (A38)) was used to calculate $r_{leakage,C}$

915 $r_{leakage,C} = \frac{L_C}{NSC} = \frac{0.05 \cdot GPP}{q \cdot C_l} = \frac{0.05 \cdot 13050 \left[\frac{\text{kg C}}{\text{ha yr}} \right]}{4 \cdot 3.2 \left[\frac{\text{kg C}}{\text{indiv}} \right] \cdot 390 \left[\frac{\text{indiv}}{\text{ha}} \right]} = 0.131 \left[\frac{1}{\text{yr}} \right]$ where 0.05 is from Jones et al., (2009), 13050 kg C ha^{-1} yr^{-1} is from the ForC database (Anderson-Teixeira et al., 2018), 3.2 kg C indiv^{-1} is from the BAAD (Falster et al., 2015), and ~ 390 indiv ha^{-1} is from Crowther et al., (2015) (for temperate forests). Note that we did not distinguish between deciduous angiosperm and evergreen gymnosperm in the BAAD because other data are from all temperate forests.



r_{growth} from LM3-SNAP was adapted to the units of LM4.1

$$r_{growth} = \frac{0.3 \left[\frac{kg\ C}{yr\ m^2} \right] \left[\frac{10000m^2}{ha} \right]}{8 \left[\frac{kg}{ha} \right] \left[\frac{kg\ C}{2kg} \right], 106 \left[\frac{kg}{ha} \right] \left[\frac{kg\ C}{2kg} \right], 4 \left[\frac{kg}{ha} \right] \left[\frac{kg\ C}{2kg} \right], 250 \left[\frac{kg}{ha} \right] \left[\frac{kg\ C}{2kg} \right]} = 65\ yr^{-1}$$

where 0.3 kg C yr⁻¹ m⁻² is from LM3-SNAP, 8, 106 and 4 kg ha⁻¹ are from Boring and Swank (1984), and 250 kg ha⁻¹ yr⁻¹ is from Binkley et al., (1992).

Parameters related to plant C allocation to symbionts ($f_{alloc,AM}$, $f_{alloc,EM}$, and $f_{alloc,Nfix}$) were estimated by varying the parameter incrementally at different orders of magnitude. Symbiont biomass C was compared to literature estimates (Boring and Swank (1984) and Binkley et al., (1992) for nodule biomass C and Zhu and Miller (2003) for mycorrhizal biomass C.



925 Appendix D Figures and Tables






Figure D1: Soil temperature dependence function of asymbiotic BNF and symbiotic BNF by *Robinia*. Asymbiotic BNF reaches its maximum at 24.4 °C and symbiotic BNF by *Robinia* reaches its maximum at 31.9°C.




930 **Figure D2: Simulated soil C and N pools, soil N fluxes, and soil N loss rates from LM4.1-BNF, LM4.1-BNF_{NPP}, and LM4.1-BNF_{ET}. (a) Simulated total soil C (depth 0-10 cm). (b) Simulated total soil N (depth 0-10 cm). (c) Simulated soil NH₄⁺ and NO₃⁻ (depth 0-10 cm). (d) N mineralization rate and net nitrification rate (depth 0-10 cm). (e) Simulated N₂O and NO emission rate and simulated dissolved organic N (DON), NH₄⁺, and NO₃⁻ leaching rate. Simulated data are averaged over the last 100 yr of the 300 yr simulation to reflect the site data which is from mature forests. Error bars indicate two standard deviations.**



935

Figure D3: Simulated gross primary production (GPP), net primary production (NPP), heterotrophic respiration (HR), and net ecosystem production (NEP) from LM4.1-BNF, LM4.1-BNF_{NPP}, and LM4.1-BNF_{ET}. Simulated data are averaged over the last 100 yr of the 300 yr simulation to reflect the data which is from mature forests. Error bars indicate two standard deviations.



940 **Figure D4: Simulated soil C and N pools, soil N fluxes, and soil N loss rates from LM4.1-BNF initialised with both *Robinia* and**
Acer, only *Acer*, and only N-fixer *Acer*, with and without asymbiotic BNF. (a) Simulated total soil C (depth 0-10 cm). (b) Simulated
 total soil N (depth 0-10 cm). (c) Simulated soil NH₄⁺ and NO₃⁻ (depth 0-10 cm). (d) N mineralization rate and net nitrification rate
 (depth 0-10 cm). (e) Simulated N₂O and NO emission rate and simulated dissolved organic N (DON), NH₄⁺, and NO₃⁻ leaching rate.
 945 Simulated data are averaged over the last 100 yr of the 300 yr simulation to reflect the site data which is from mature forests.
 Error bars indicate two standard deviations.




Figure D5: Simulated gross primary production (GPP), net primary production (NPP), heterotrophic respiration (HR), and net ecosystem production (NEP) from LM4.1-BNF initialised with both *Robinia* and *Acer*, only *Acer*, and only N-fixer *Acer*, with and without asymbiotic BNF. Simulated data are averaged over the last 100 yr of the 300 yr simulation to reflect the data which is from mature forests. Error bars indicate two standard deviations.

950





Figure D6: Histogram of FIA stand age for FIA plots in North Carolina. These distributions were used to weigh the data displayed in Fig. 2 and 3.



955 **Figure D7: Simulated absolute basal area of *Acer* and *Robinia* over time compared to FIA data (in North Carolina). Simulated data are trees with dbh > 12.7 cm to reflect the dbh range of FIA data. FIA data of all non-fixing trees are aggregated to represent *Acer*. Each point represents an FIA plot.**





Figure D8: Crown area model fit to FIA Forest Health Monitoring data for *Acer*.



960

Figure D9: Crown area model fit to FIA Forest Health Monitoring data for *Robinia*.





Figure D10: Height model fit to FIA data for *Acer*.



965 Figure D11: Height model fit to FIA data for *Robinia*.




Figure D12: Wood mass model fit to FIA data for *Acer* (displayed at mean height of *Acer*).





Figure D13: Wood mass model fit to FIA data for *Robinia* (displayed at mean height of *Robinia*).



970

Figure D14: Sapwood area at breast height model fit to BAAD for *Acer*.




Figure D15: Sapwood area at breast height model fit to BAAD for *Robinia*.

975 Table D1: Vegetation type-specific parameters.

Parameter	Vegetation type	Value	Unit	Source
-----------	-----------------	-------	------	--------



$C: N_l$	<i>Acer</i>	30	kg C kg N ⁻¹	TRY database
	<i>Robinia</i>	14		
ρ_{wood}	<i>Acer</i>	340	kg C m ⁻³	TRY database
	<i>Robinia</i>	280		
LMA	<i>Acer</i>	0.0482	kg C m ⁻²	TRY database
	<i>Robinia</i>	0.0380		
V_{cmax} at 15°C	<i>Acer</i>	17	μmol m ⁻² s ⁻¹	TRY database, converted to 15°C using method from Medlyn et al., 2002
	<i>Robinia</i>	23		
α_{HT}	<i>Acer</i>	46.175656	m	see Appendix C
	<i>Robinia</i>	43.87161		
θ_{HT}	<i>Acer</i>	0.782971	unitless	see Appendix C
	<i>Robinia</i>	0.89594		
γ_{HT}	<i>Acer</i>	0.485517	unitless	see Appendix C
	<i>Robinia</i>	0.40675		
α_{BM}	<i>Acer</i>	0.3922393	unitless	see Appendix C
	<i>Robinia</i>	0.3383768		
α_{CA}	<i>Acer</i>	134.18322	unitless	see Appendix C
	<i>Robinia</i>	66.3140		
θ_{CA}	<i>Acer</i>	1.02731	unitless	see Appendix C
	<i>Robinia</i>	0.8128		
φ_{CSA}	<i>Acer</i>	0.21346	m ⁻¹	see Appendix C
	<i>Robinia</i>	0.16983		
μ_C	<i>Acer</i>	0.01810868	yr ⁻¹	see Appendix C
	<i>Robinia</i>	0.03105176		
μ_U	<i>Acer</i>	0.04024044	yr ⁻¹	see Appendix C
	<i>Robinia</i>	0.08785878		
r_{Nfix}	<i>Robinia</i>	6.3	kg N kg C ⁻¹ yr ⁻¹	Bytnerowicz et al., submitted

Table D2: General parameters.

Parameter	Value	Unit	Source
N uptake by roots and symbionts			
r_{NO_3}	0.1	kg N m ⁻³ yr ⁻¹	Sulman et al., 2019
k_{M,NO_3}	0.005	kg N m ⁻³	Sulman et al., 2019
r_{NH_4}	0.1	kg N m ⁻³ yr ⁻¹	Sulman et al., 2019
k_{M,NH_4}	0.005	kg N m ⁻³	Sulman et al., 2019
$\Delta z(k)$	0.02, 0.04, 0.04, 0.05, 0.05, 0.10, 0.10, 0.20, 0.20, 0.20, 0.40, 0.40, 0.40, 0.40, 0.40, 1.0, 1.0, 1.0, 1.5, 2.5	m	Sulman et al., 2019
r_{root}	0.00029	m	Jackson et al., 1997
SRL	24545	m kg C ⁻¹	Jackson et al., 1997
SRA	45	m ² kg C ⁻¹	Jackson et al., 1997
r_{rhiz}	0.001	m	Sulman et al., 2019
$r_{NO_3,AM}$	0.2	kg N m ⁻³ yr ⁻¹	Sulman et al., 2019



k_{AM,NO_3}	0.001	kg N m^{-3}	Sulman et al., 2019
$r_{NH_4,AM}$	0.2	$\text{kg N m}^{-3} \text{ yr}^{-1}$	Sulman et al., 2019
k_{AM,NH_4}	0.001	kg N m^{-3}	Sulman et al., 2019
k_{AM}	0.3	kg C m^{-3}	Sulman et al., 2019
$V_{EM,max,ref,i}$	2, 0.3, 2	yr^{-1}	Sulman et al., 2019
$E_{a,i}$	6000, 40000, 6000	kJ mol^{-1}	Sulman et al., 2019
R	8.314472	$\text{kJ mol}^{-1} \text{ K}^{-1}$	-
T_{ref}	293.15	K	-
$k_{M,EM}$	0.015	unitless	Sulman et al., 2019
$\varepsilon_{C,i,EM}$	0.1, 0.05, 0.1	unitless	Sulman et al., 2019
$\varepsilon_{N,i,EM}$	0.7, 0.7, 0.7	unitless	Sulman et al., 2019
$r_{NO_3,EM}$	0.2	$\text{kg N m}^{-3} \text{ yr}^{-1}$	Sulman et al., 2019
k_{EM,NO_3}	0.001	kg N m^{-3}	Sulman et al., 2019
$r_{NH_4,EM}$	0.2	$\text{kg N m}^{-3} \text{ yr}^{-1}$	Sulman et al., 2019
k_{EM,NH_4}	0.001	kg N m^{-3}	Sulman et al., 2019
k_{EM}	0.3	kg C m^{-3}	Sulman et al., 2019
Asymbiotic BNF			
$V_{max,ref,i}$	18, 0.2, 4.5	yr^{-1}	Sulman et al., 2019
k_M	0.045	unitless	Sulman et al., 2019
θ_{sat}	0.439	$\text{m}^3 \text{ m}^{-3}$	Sulman et al., 2019
$V_{denit,max,ref,i}$	0.018, 0.00025, 0.0045	yr^{-1}	Sulman et al., 2019
$k_{M,denit}$	0.045	unitless	Sulman et al., 2019
k_{denit}	0.0027	yr	Sulman et al., 2019
f_{denit}	0.93	kg N kg C^{-1}	Sulman et al., 2019
τ_M	0.25	yr	Sulman et al., 2019
$\varepsilon_{t,M}$	0.6	unitless	Sulman et al., 2019
$C:N_M$	10.0	kg C kg N^{-1}	Chapin et al., 2011
$r_{Nfix asymb}$	0.024	$\text{kg N kg C}^{-1} \text{ yr}^{-1}$	see Appendix C
$\varepsilon_{N,i}$	0.7, 0.4, 0.7	unitless	Sulman et al., 2019
$\varepsilon_{C,i}$	0.6, 0.05, 0.6	unitless	Sulman et al., 2019
V_{max,ref,NH_4}	365	yr^{-1}	Sulman et al., 2019
E_{a,NH_4}	37000	kJ mol^{-1}	Sulman et al., 2019
V_{max,ref,NO_3}	365	yr^{-1}	Sulman et al., 2019
E_{a,NO_3}	37000	kJ mol^{-1}	Sulman et al., 2019
Plant growth and N limitation			
$C:N_r$	42	kg C kg N^{-1}	Roumet et al., 2016
$C:N_{sw}$	287	kg C kg N^{-1}	Meerts, 2002
$C:N_{hw}$	427	kg C kg N^{-1}	Meerts, 2002
LAI_{max}	5.1	$\text{m}^2 \text{ m}^{-2}$	Asner et al., 2003
φ_{RL}	0.79	unitless	see Appendix C
f_{br}	0.255	unitless	see Appendix C
q	4	unitless	Weng et al., 2015
$r_{leakage,C}$	0.131	yr^{-1}	see Appendix C
$r_{leakage,N}$	0.15	yr^{-1}	Sulman et al., 2019
Plant C allocation to symbionts			
$f_{alloc,AM}$	0.01	yr^{-1}	estimated (see Appendix C)



$f_{alloc,EM}$	0.01	yr^{-1}	estimated (see Appendix C)
$f_{alloc,Nfix}$	1.0	yr^{-1}	estimated (see Appendix C)
$f_{alloc,Nfix,min}$	0.5	$\text{kg C indiv}^{-1} \text{ yr}^{-1}$	D. N. L. Menge, unpublished data
$C:N_{alloc}$	1000	kg C kg N^{-1}	Sulman et al., 2019
Growth and turnover of symbionts			
ϵ_{symb}	0.666	unitless	Sulman et al., 2019 (for plant growth)
r_{growth}	65	yr^{-1}	see Appendix C
ξ_{AM}	1.25	yr^{-1}	Sulman et al., 2019 (for fine roots)
ξ_{EM}	1.25	yr^{-1}	Sulman et al., 2019 (for fine roots)
ξ_{Nfix}	1.25	yr^{-1}	Sulman et al., 2019 (for fine roots)
$cost_{Nfix}$	4.8	kg C kg N^{-1}	Gutschick et al., 1981
τ_{AM}	2.0	yr	Sulman et al., 2019 (for fine roots)
τ_{EM}	2.0	yr	Sulman et al., 2019 (for fine roots)
τ_{Nfix}	2.0	yr	Sulman et al., 2019 (for fine roots)
$C:N_{AM}$	10	kg C kg N^{-1}	Johnson, 2010
$C:N_{EM}$	14	kg C kg N^{-1}	Zhang and Elser, 2017
$C:N_{Nfix}$	6	kg C kg N^{-1}	Boring and Swank, 1984
$r_{up,veg}$	250	yr^{-1}	Sulman et al., 2019
Soil N ₂ O and NO emissions			
$V_{nit,max,ref}$	5	yr^{-1}	Sulman et al., 2019
$E_{a,nit}$	37000	kJ mol^{-1}	Sulman et al., 2019
$\gamma_{N_2O,nit}$	0.0017	unitless	Bateman and Baggs, 2005
LM4.1-BNF _{NPP}			
a_{NPP}	0.0018	$\text{kg N m}^{-2} \text{ yr}^{-1}$	Meyerholt et al., 2016
b_{NPP}	-3	$\text{m}^2 \text{ yr kg C}^{-1}$	Meyerholt et al., 2016
LM4.1-BNF _{ET}			
a_{ET}	2.34e-6	$\text{kg N mm}^{-1} \text{ m}^{-2}$	Meyerholt et al., 2016
b_{ET}	-1.72e-5	$\text{kg N m}^{-2} \text{ yr}^{-1}$	Meyerholt et al., 2016

Table D3: Summary of the spin up and numerical experiments.

	Spin up	Numerical experiments
Atmospheric CO ₂ concentration	Pre-industrial: 284.26 ppm (Meinshausen et al., 2017)	mean 1948 – 1978: 324.53 ppm (Dlugokencky and Tans, 2020)
Meteorological forcing	1948-1978 (Sheffield et al., 2006)	1948-1978 (Sheffield et al., 2006)
N deposition rate	1993 (Dentener, 2006)	1993 (Dentener, 2006)



980

Table D4: Initial densities and heights for simulations. The initial densities of *Robinia* and *Acer* were derived from the US FIA database seedling data for plots with at least one *Robinia* individual in North Carolina. Diameter at breast height is determined from height by allometry (Eq. (A41)).

BNF representation	Initialised species	Density	Height	Diameter at breast height
LM4.1-BNF	<i>Acer</i>	0.54 indiv m ⁻²	0.5 m	0.00160 m
	<i>Robinia</i>	0.13 indiv m ⁻²	0.5 m	0.00336 m
LM4.1-BNF	<i>Acer</i>	0.5 indiv m ⁻²	0.5 m	0.00160 m
LM4.1-BNF _{NPP}	<i>Acer</i>	0.5 indiv m ⁻²	0.5 m	0.00160 m
LM4.1-BNF _{ET}	<i>Acer</i>	0.5 indiv m ⁻²	0.5 m	0.00160 m

985 **Table D5: Validated variables and data sources.**

Variable	Data source for Coweeta Hydrologic Laboratory
dbh growth rate of each vegetation type	FIA database tree data, North Carolina plots with at least one <i>Robinia</i> individual (US Forest Service, 2020a)
dbh distribution	FIA database tree data, North Carolina plots with at least one <i>Robinia</i> individual (US Forest Service, 2020a)
Basal area fraction of each vegetation type	FIA database tree data, North Carolina plots with at least one <i>Robinia</i> individual (US Forest Service, 2020a)
Total plant biomass C	FIA database tree data, North Carolina plots with at least one <i>Robinia</i> individual (US Forest Service, 2020a)
Asymbiotic BNF rate	Todd et al., 1978
Symbiotic BNF rate	Boring and Swank, 1984
Total soil C and N	FIA database soil data, North Carolina plots (US Forest Service, 2020a)
Soil NH ₄ ⁺ and NO ₃ ⁻	Knoepp, 2009a, 2018
N mineralization and net nitrification rates	Knoepp, 2009a, 2018
N ₂ O and NO emission rates	Knoepp, 2009b, 2009a
Stehfest and Bouwman, 2006	
DON, NH ₄ ⁺ , and NO ₃ ⁻ leaching rate	Swank and Waide, 1988
Gross primary production, heterotrophic respiration, net primary production, and net ecosystem production	Anderson-Teixeira et al., 2018

Code availability

Full GFDL land model code is available upon request through the GFDL.

Author contribution

S.K.-G. designed LM4.1-BNF. D.M., S.M., I.M.C., E.S., S.P., and T.B. advised on the model. S.M. implemented LM3-SNAP into LM4.1. S.K.-G. implemented modifications for LM4.1-BNF. S.K.-G. conducted the analysis. S.K.-G. wrote the initial draft. All authors contributed to the writing.

990



Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

995 We thank Ben Sulman for initial assistance with the model. We thank Anika Staccone for assistance with the US FIA database. We thank the Menge lab and the Soper lab for their comments on the manuscript. Data from several databases were used (Appendix B). This study was supported by the National Science Foundation, award NSF DEB-1457650. We acknowledge the support of the Natural Sciences and Engineering Research Council of Canada. Cette recherche a été financée par le Conseil de recherches en sciences naturelles et en génie du Canada.

1000 References

- Adams, M. A., Turnbull, T. L., Sprent, J. I. and Buchmann, N.: Legumes are different: Leaf nitrogen, photosynthesis, and water use efficiency, *Proc. Natl. Acad. Sci.*, 113(15), 4098–4103, doi:10.1073/pnas.1523936113, 2016.
- Anderson-Teixeira, K. J., Wang, M. M. H., McGarvey, J. C., Herrmann, V., Tepley, A. J., Bond-Lamberty, B. and LeBauer, D. S.: ForC: a global database of forest carbon stocks and fluxes, *Ecology*, 99(6), 1507, doi:10.1002/ecy.2229, 2018.
- 1005 Arora, V. K., Katavouta, A., Williams, R. G., Jones, C. D., Brovkin, V., Friedlingstein, P., Schwinger, J., Bopp, L., Boucher, O., Cadule, P., Chamberlain, M. A., Christian, J. R., Delire, C., Fisher, R. A., Hajima, T., Ilyina, T., Joetzjer, E., Kawamiya, M., Koven, C. D., Krasting, J. P., Law, R. M., Lawrence, D. M., Lenton, A., Lindsay, K., Pongratz, J., Raddatz, T., Séférian, R., Tachiiri, K., Tjiputra, J. F., Wiltshire, A., Wu, T. and Ziehn, T.: Carbon–concentration and carbon–climate feedbacks in CMIP6 models and their comparison to CMIP5 models, *Biogeosciences*, 17(16), 4173–4222, doi:10.5194/bg-17-4173-2020,
- 1010 2020.
- Asner, G. P., Scurlock, J. M. O. and Hicke, J. A.: Global synthesis of leaf area index observations: implications for ecological and remote sensing studies, *Glob. Ecol. Biogeogr.*, 12(3), 191–205, 2003.
- Averill, C., Bhatnagar, J. M., Dietze, M. C., Pearse, W. D. and Kivlin, S. N.: Global imprint of mycorrhizal fungi on whole-plant nutrient economics, *Proc. Natl. Acad. Sci. U. S. A.*, 116(46), 23163–23168, doi:10.1073/pnas.1906655116, 2019.
- 1015 Bateman, E. J. and Baggs, E. M.: Contributions of nitrification and denitrification to N₂O emissions from soils at different water-filled pore space, *Biol. Fertil. Soils*, 41(6), 379–388, doi:10.1007/s00374-005-0858-3, 2005.
- Batterman, S. A., Hedin, L. O., van Breugel, M., Ransijn, J., Craven, D. J. and Hall, J. S.: Key role of symbiotic dinitrogen fixation in tropical forest secondary succession, *Nature*, 502(7470), 224–227, doi:10.1038/nature12525, 2013.
- Binkley, D., Sollins, P., Bell, R., Sachs, D. and Myrold, D.: Biogeochemistry of adjacent conifer and alder-conifer stands, *Ecology*, 73(6), 2022–2033, doi:10.2307/1941452, 1992.
- 1020 Boring, L. R. and Swank, W. T.: The Role of Black Locust (*Robinia Pseudo-Acacia*) in Forest Succession, *J. Ecol.*, 72(3),



- 749–766, 1984.
- Bytnerowicz, T. A., Akana, P. R., Griffin, K. L. and Menge, D. N. L.: The temperature sensitivity of woody dinitrogen fixation across species and growing temperatures, n.d.
- 1025 Canarini, A., Kaiser, C., Merchant, A., Richter, A. and Wanek, W.: Root exudation of primary metabolites: Mechanisms and their roles in plant responses to environmental stimuli, *Front. Plant Sci.*, 10(157), 1–19, doi:10.3389/fpls.2019.00157, 2019.
- Chan, Y. K.: Temperature response of an associative N₂-fixing *Pseudomonas* species in pure culture, *Can. J. Microbiol.*, 37(9), 715–718, doi:10.1139/m91-122, 1991.
- Chapin, D. M., Bliss, L. C. and Bledsoe, L. J.: Environmental regulation of nitrogen fixation in a high arctic lowland ecosystem, *Can. J. Bot.*, 69(12), 2744–2755, doi:10.1139/b91-345, 1991.
- 1030 Chapin, F. S., Matson, P. A. and Vitousek, P. M.: *Principles of Terrestrial Ecosystem Ecology*, Springer Science & Business Media., 2011.
- Chapin III, F. S., Walker, L. R., Fastie, C. L. and Sharman, L. C.: Mechanisms of Primary Succession Following Deglaciation at Glacier Bay, Alaska, *Ecol. Monogr.*, 64(2), 149–175, 1994.
- 1035 Chapin III, F. S., Conway, A. J., Johnstone, J. F., Hollingsworth, T. N. and Hollingsworth, J.: Absence of net long-term successional facilitation by alder in a boreal Alaska floodplain, *Ecology*, 97(11), 2986–2997, doi:10.1002/ecy.1529, 2016.
- Cheng, W., Parton, W. J., Gonzalez-Meler, M. A., Phillips, R., Asao, S., McNickle, G. G., Brzostek, E. and Jastrow, J. D.: Synthesis and modeling perspectives of rhizosphere priming, *New Phytol.*, 201(1), 31–44, doi:10.1111/nph.12440, 2014.
- Cierjacks, A., Kowarik, I., Joshi, J., Hempel, S., Ristow, M., von der Lippe, M. and Weber, E.: Biological flora of the british isles: *Robinia pseudoacacia*, *J. Ecol.*, 101(6), 1623–1640, doi:10.1111/1365-2745.12162, 2013.
- 1040 Cleveland, C. C., Townsend, A. R., Schimel, D. S., Fisher, H., Hedin, L. O., Perakis, S., Latty, E. F., Fischer, C. Von, Elseroad, A. and Wasson, M. F.: Global patterns of terrestrial biological nitrogen (N₂) fixation in natural ecosystems, *Glob. Biochem. Cycles*, 13(2), 623–645, doi:10.1029/1999GB900014, 1999.
- Coxson, D. S. and Kershaw, K. A.: Rehydration response of nitrogenase activity and carbon fixation in terrestrial *Nostoc* commune from *Stipa-Bouteola* grassland, *Can. J. Bot.*, 61(10), 2658–2668, doi:10.1139/b83-293, 1983.
- 1045 Crowther, T. W., Glick, H. B., Covey, K. R., Bettigole, C., Maynard, D. S., Thomas, S. M., Smith, J. R., Hintler, G., Duguid, M. C., Amatulli, G., Tuanmu, M.-N., Jetz, W., Salas, C., Stam, C., Piotta, D., Tavani, R., Green, S., Bruce, G., Williams, S. J., Wiser, S. K., Huber, M. O., Hengeveld, G. M., Nabuurs, G.-J., Tikhonova, E., Borchardt, P., Li, C.-F., Powrie, L. W., Fischer, M., Hemp, A., Homeier, J., Cho, P., Vibrans, A. C., Umunay, P. M., Piao, S. L., Rowe, C. W., Ashton, M. S., Crane, P. R. and Bradford, M. A.: Mapping tree density at a global scale, *Nature*, 525(7568), 201–205, doi:10.1038/nature14967, 2015.
- Davidson, E. A., Keller, M., Erickson, H. E., Verchot, L. V. and Veldkamp, E.: Testing a Conceptual Model of Soil Emissions of Nitrous and Nitric Oxides, *Bioscience*, 50(8), 667, doi:10.1641/0006-3568(2000)050[0667:TACMOS]2.0.CO;2, 2000.
- 1055 Davies-Barnard, T. and Friedlingstein, P.: The Global Distribution of Biological Nitrogen Fixation in Terrestrial Natural



- Ecosystems, *Global Biogeochem. Cycles*, 34(3), 1–17, doi:10.1029/2019GB006387, 2020.
- Davies-Barnard, T., Meyerholt, J., Zaehle, S., Friedlingstein, P., Brovkin, V., Fan, Y., Fisher, R. A., Jones, C. D., Lee, H., Peano, D., Smith, B., Warlind, D. and Wiltshire, A.: Nitrogen Cycling in CMIP6 Land Surface Models: Progress and Limitations, *Biogeosciences*, 17, 5129–5148, doi:10.5194/bg-2019-513, 2020.
- 1060 DeLuca, T. H., Zackrisson, O., Nilsson, M.-C. and Sellstedt, A.: Quantifying nitrogen-fixation in feather moss carpets of boreal forests, *Lett. to Nat.*, 419(6910), 917–920, doi:10.1038/nature01051, 2002.
- Dentener, F. J.: Global Maps of Atmospheric Nitrogen Deposition, 1860, 1993, and 2050., 2006.
- Dlugokencky, E. and Tans, P.: Trends in atmospheric carbon dioxide. [online] Available from: https://www.esrl.noaa.gov/gmd/ccgg/trends/gl_data.html, 2020.
- 1065 Elser, J. J., Bracken, M. E. S., Cleland, E. E., Gruner, D. S., Harpole, W. S., Hillebrand, H., Ngai, J. T., Seabloom, E. W., Shurin, J. B. and Smith, J. E.: Global analysis of nitrogen and phosphorus limitation of primary producers in freshwater, marine and terrestrial ecosystems, *Ecol. Lett.*, 10(12), 1135–1142, doi:10.1111/j.1461-0248.2007.01113.x, 2007.
- Falster, D. S., Duursma, R. A., Ishihara, M. I., Barneche, D. R., FitzJohn, R. G., Vårhammar, A., Aiba, M., Ando, M., Anten, N., Aspinwall, M. J., Baltzer, J. L., Baraloto, C., Battaglia, M., Battles, J. J., Lamberty, B. B., Van Breugel, M., Camac, J.,
- 1070 Claveau, Y., Coll, L., Dannoura, M., Delagrangé, S., Domec, J. C., Fatemi, F., Feng, W., Gargaglione, V., Goto, Y., Hagihara, A., Hall, J. S., Hamilton, S., Harja, D., Hiura, T., Holdaway, R., Hutley, L. B., Ichie, T., Jokela, E. J., Kantola, A., Kelly, J. W. G., Kenzo, T., King, D., Kloeppe, B. D., Kohyama, T., Komiyama, A., Laclau, J. P., Lusk, C. H., Maguire, D. A., Le Maire, G., Mäkelä, A., Markesteijn, L., Marshall, J., McCulloh, K., Miyata, I., Mokany, K., Mori, S., Myster, R. W., Nagano, M., Naidu, S. L., Nouvellon, Y., O’Grady, A. P., O’Hara, K. L., Ohtsuka, T., Osada, N., Osunkoya, O. O., Peri, P.
- 1075 L., Petritan, A. M., Poorter, L., Portsmouth, A., Potvin, C., Ransijn, J., Reid, D., Ribeiro, S. C., Roberts, S. D., Rodríguez, R., Acosta, A. S., Santa-Regina, I., Sasa, K., Selaya, N. G., Sillett, S. C., Sterck, F., Takagi, K., Tange, T., Tanouchi, H., Tissue, D., Umehara, T., Utsugi, H., Vadeboncoeur, M. A., Valladares, F., Vanninen, P., Wang, J. R., Wenk, E., Williams, R., De Aquino Ximenes, F., Yamaba, A., Yamada, T., Yamakura, T., Yanai, R. D. and York, R. A.: BAAD: a Biomass And Allometry Database for woody plants, *Ecology*, 96(5), 1445, doi:10.1890/14-1889.1, 2015.
- 1080 Finzi, A. C., Abramoff, R. Z., Spiller, K. S., Brzostek, E. R., Darby, B. A., Kramer, M. A. and Phillips, R. P.: Rhizosphere processes are quantitatively important components of terrestrial carbon and nutrient cycles, *Glob. Chang. Biol.*, 21(5), 2082–2094, doi:10.1111/gcb.12816, 2015.
- Fowler, D., Coyle, M., Skiba, U., Sutton, M. A., Cape, J. N., Reis, S., Sheppard, L. J., Jenkins, A., Grizzetti, B., Galloway, J. N., Vitousek, P., Leach, A., Bouwman, A. F., Butterbach-Bahl, K., Dentener, F., Stevenson, D., Amann, M. and Voss, M.:
- 1085 The global nitrogen cycle in the twenty-first century, *Philos. Trans. R. Soc. B Biol. Sci.*, 368(1621), 20130164, doi:10.1098/rstb.2013.0164, 2013.
- Goll, D. S., Winkler, A. J., Raddatz, T., Dong, N., Colin Prentice, I., Ciais, P. and Brovkin, V.: Carbon-nitrogen interactions in idealized simulations with JSBACH (version 3.10), *Geosci. Model Dev.*, 10(5), 2009–2030, doi:10.5194/gmd-10-2009-2017, 2017.



- 1090 Gutschick, V. P.: Evolved Strategies in Nitrogen Acquisition by Plants, *Am. Nat.*, 118(5), 607–637, 1981.
- Haverd, V., Smith, B., Nieradzik, L., Briggs, P. R., Woodgate, W., Trudinger, C. M., Canadell, J. G. and Cuntz, M.: A new version of the CABLE land surface model (Subversion revision r4601) incorporating land use and land cover change, woody vegetation demography, and a novel optimisation-based approach to plant coordination of photosynthesis, *Geosci. Model Dev.*, 11(7), 2995–3026, doi:10.5194/gmd-11-2995-2018, 2018.
- 1095 Hoch, G., Richter, A. and Körner, C.: Non-structural carbon compounds in temperate forest trees, *Plant, Cell Environ.*, 26(7), 1067–1081, doi:10.1046/j.0016-8025.2003.01032.x, 2003.
- Houlton, B. Z., Wang, Y. P., Vitousek, P. M. and Field, C. B.: A unifying framework for dinitrogen fixation in the terrestrial biosphere, *Nature*, 454(7202), 327–330, doi:10.1038/nature07028, 2008.
- Huang, Y. and Gerber, S.: Global soil nitrous oxide emissions in a dynamic carbon-nitrogen model, *Biogeosciences*, 12(21), 6405–6427, doi:10.5194/bg-12-6405-2015, 2015.
- 1100 Hulvey, K. B., Hobbs, R. J., Standish, R. J., Lindenmayer, D. B., Lach, L. and Perring, M. P.: Benefits of tree mixes in carbon plantings, *Nat. Clim. Chang.*, 3(10), 869–874, doi:10.1038/nclimate1862, 2013.
- Jackson, R. B., Mooney, H. A. and Schulze, E.-D.: A global budget for fine root biomass, surface area, and nutrient contents, *Proc. Natl. Acad. Sci.*, 94(14), 7362–7366, 1997.
- 1105 Johnson, N. C.: Resource Stoichiometry Elucidates the Structure and Function of Arbuscular Mycorrhizas across Scales, *New Phytol.*, 185(3), 631–647, doi:10.1111/j.1469-8137.2009.03110.x, 2010.
- Jones, D. L., Nguyen, C. and Finlay, R. D.: Carbon flow in the rhizosphere: Carbon trading at the soil-root interface, *Plant Soil*, 321, 5–33, doi:10.1007/s11104-009-9925-0, 2009.
- Knoepp, J.: Measurements of Soil Bulk Density at the Coweeta Terrestrial Gradient Sites. [online] Available from: https://coweeta.uga.edu/dbpublic/dataset_details.asp?accession=1123, 2009a.
- 1110 Knoepp, J.: Measurements of Soil Nitrogen Transformations at the Coweeta Terrestrial Gradient Sites. [online] Available from: https://coweeta.uga.edu/dbpublic/dataset_details.asp?accession=1125, 2009b.
- Knoepp, J.: Measurements of Soil %C and %N at the Coweeta Terrestrial Gradient Sites. [online] Available from: https://coweeta.uga.edu/dbpublic/dataset_details.asp?accession=1124, 2018.
- 1115 Kou-Giesbrecht, S. and Menge, D. N. L.: Nitrogen-fixing trees increase soil nitrous oxide emissions: a meta-analysis, n.d.
- Lai, H. R., Hall, J. S., Batterman, S. A., Turner, B. L. and van Breugel, M.: Nitrogen fixer abundance has no effect on biomass recovery during tropical secondary forest succession, *J. Ecol.*, 106(4), 1415–1427, doi:10.1111/1365-2745.12979, 2018.
- Lawrence, D. M., Fisher, R. A., Koven, C. D., Oleson, K. W., Swenson, S. C., Bonan, G., Collier, N., Ghimire, B., van Kampenhout, L., Kennedy, D., Kluzek, E., Lawrence, P. J., Li, F., Li, H., Lombardozzi, D., Riley, W. J., Sacks, W. J., Shi, M., Vertenstein, M., Wieder, W. R., Xu, C., Ali, A. A., Badger, A. M., Bisht, G., van den Broeke, M., Brunke, M. A., Burns, S. P., Buzan, J., Clark, M., Craig, A., Dahlin, K., Drewniak, B., Fisher, J. B., Flanner, M., Fox, A. M., Gentine, P., Hoffman, F., Keppel-Aleks, G., Knox, R., Kumar, S., Lenaerts, J., Leung, L. R., Lipscomb, W. H., Lu, Y., Pandey, A., Pelletier, J. D.,



- Perket, J., Randerson, J. T., Ricciuto, D. M., Sanderson, B. M., Slater, A., Subin, Z. M., Tang, J., Thomas, R. Q., Val Martin,
1125 M. and Zeng, X.: The Community Land Model Version 5: Description of New Features, Benchmarking, and Impact of
Forcing Uncertainty, *J. Adv. Model. Earth Syst.*, 11(12), 4245–4287, doi:10.1029/2018MS001583, 2019.
- LeBauer, D. S. and Treseder, K. K.: Nitrogen Limitation of Net Primary Productivity in Terrestrial Ecosystems is Globally
Distributed, *Ecology*, 89(2), 371–379, doi:10.1016/j.agee.2013.04.020, 2008.
- Levy-Varon, J. H., Batterman, S. A., Medvigy, D., Hall, J. S., Breugel, M. Van and Hedin, L. O.: Tropical carbon sink
1130 accelerated by symbiotic dinitrogen fixation, *Nat. Commun.*, 10(1), doi:10.1038/s41467-019-13656-7, 2019.
- Liang, J., Qi, X., Souza, L. and Luo, Y.: Processes regulating progressive nitrogen limitation under elevated carbon dioxide:
A meta-analysis, *Biogeosciences*, 13(9), 2689–2699, doi:10.5194/bg-13-2689-2016, 2016.
- Liengen, T. and Olsen, R. A.: Seasonal and site-specific variations in nitrogen fixation in a high arctic area, Ny-Alesund,
Spitsbergen, *Can. J. Microbiol.*, 43(8), 759–769, doi:10.1139/m97-109, 1997.
- 1135 Martinez Cano, I., Shevliakova, E., Malyshev, S., Wright, S. J., Detto, M., Pacala, S. W. and Muller-Landau, H. C.:
Allometric constraints and competition enable the simulation of size structure and carbon fluxes in a dynamic vegetation
model of tropical forests (LM3PPA-TV), *Glob. Chang. Biol.*, 26(8), 4478–4494, doi:10.1111/gcb.15188, 2020.
- Medlyn, B. E., Dreyer, E., Ellsworth, D., Forstreuter, M., Harley, P. C., Kirschbaum, M. U. F., Le Roux, X., Montpied, P.,
Strassemeyer, J., Walcroft, A., Wang, K. and Loustau, D.: Temperature response of parameters of a biochemically based
1140 model of photosynthesis. II. A review of experimental data, *Plant, Cell Environ.*, 25(9), 1167–1179, doi:10.1046/j.1365-
3040.2002.00891.x, 2002.
- Medvigy, D., Wofsy, S. C., Munger, J. W., Hollinger, D. Y. and Moorcroft, P. R.: Mechanistic scaling of ecosystem function
and dynamics in space and time: Ecosystem Demography model version 2, *J. Geophys. Res. Biogeosciences*, 114(G01002),
doi:10.1029/2008JG000812, 2009.
- 1145 Meerts, P.: Mineral nutrient concentrations in sapwood and heartwood : a literature review, *Ann. For. Sci.*, 59(7), 713–722,
2002.
- Meinshausen, M., Vogel, E., Nauels, A., Lorbacher, K., Meinshausen, N., Etheridge, D. M., Fraser, P. J., Montzka, S. A.,
Rayner, P. J., Trudinger, C. M., Krummel, P. B., Beyerle, U., Canadell, J. G., Daniel, J. S., Enting, I. G., Law, R. M.,
Lunder, C. R., O’Doherty, S., Prinn, R. G., Reimann, S., Rubino, M., Velders, G. J. M., Vollmer, M. K., Wang, R. H. J. and
1150 Weiss, R.: Historical greenhouse gas concentrations for climate modelling (CMIP6), *Geosci. Model Dev.*, 10(5), 2057–2116,
doi:10.5194/gmd-10-2057-2017, 2017.
- Menge, D. N. L. and Hedin, L. O.: Nitrogen Fixation in Different Biogeochemical Niches along a 120 000-Year
Chronosequence in New Zealand, *Ecology*, 90(8), 2190–2201, 2009.
- Menge, D. N. L., Denoyer, J. L. and Lichstein, J. W.: Phylogenetic constraints do not explain the rarity of nitrogen-fixing
1155 trees in late-successional temperate forests, *PLoS One*, 5(8), e12056, doi:10.1371/journal.pone.0012056, 2010.
- Menge, D. N. L., Wolf, A. A. and Funk, J. L.: Diversity of nitrogen fixation strategies in Mediterranean legumes, *Nat.*
Plants, 1(6), 1–5, doi:10.1038/nplants.2015.64, 2015.



- Menge, D. N. L., Chisholm, R. A., Davies, S. J., Abu Salim, K., Allen, D., Alvarez, M., Bourg, N., Brockelman, W. Y., Bunyavejchewin, S., Butt, N., Cao, M., Chanthorn, W., Chao, W. C., Clay, K., Condit, R., Cordell, S., da Silva, J. B.,
1160 Dattaraja, H. S., de Andrade, A. C. S., de Oliveira, A. A., den Ouden, J., Drescher, M., Fletcher, C., Giardina, C. P., Savitri
Gunatilleke, C. V., Gunatilleke, I. A. U. N., Hau, B. C. H., He, F., Howe, R., Hsieh, C. F., Hubbell, S. P., Inman-Narahari, F.
M., Jansen, P. A., Johnson, D. J., Kong, L. S., Král, K., Ku, C. C., Lai, J., Larson, A. J., Li, X., Li, Y., Lin, L., Lin, Y. C.,
Liu, S., Lum, S. K. Y., Lutz, J. A., Ma, K., Malhi, Y., McMahon, S., McShea, W., Mi, X., Morecroft, M., Myers, J. A.,
Nathalang, A., Novotny, V., Ong, P., Orwig, D. A., Ostertag, R., Parker, G., Phillips, R. P., Abd. Rahman, K., Sack, L.,
1165 Sang, W., Shen, G., Shringi, A., Shue, J., Su, S. H., Sukumar, R., Sun, I. F., Suresh, H. S., Tan, S., Thomas, S. C., Toko, P.
S., Valencia, R., Vallejo, M. I., Vicentini, A., Vrška, T., Wang, B., Wang, X., Weiblen, G. D., Wolf, A., Xu, H., Yap, S.,
Zhu, L. and Fung, T.: Patterns of nitrogen-fixing tree abundance in forests across Asia and America, *J. Ecol.*, 107(6), 2598–
2610, doi:10.1111/1365-2745.13199, 2019.
- Meyerholt, J., Zaehle, S. and Smith, M. J.: Variability of projected terrestrial biosphere responses to elevated levels of
1170 atmospheric CO₂ due to uncertainty in biological nitrogen fixation, *Biogeosciences*, 13(5), 1491–1518, doi:10.5194/bg-13-
1491-2016, 2016.
- Meyerholt, J., Sickel, K. and Zaehle, S.: Ensemble projections elucidate effects of uncertainty in terrestrial nitrogen
limitation on future carbon uptake, *Glob. Chang. Biol.*, 26(7), 3978–3996, doi:10.1111/gcb.15114, 2020.
- Mooney, H. A., Winner, W. E. and Pell, E. J.: Response of plants to multiple stresses, Academic Press., 1991.
- 1175 Nacry, P., Bouguyon, E. and Gojon, A.: Nitrogen acquisition by roots: Physiological and developmental mechanisms
ensuring plant adaptation to a fluctuating resource, *Plant Soil*, 370, 1–29, doi:10.1007/s11104-013-1645-9, 2013.
- Oswalt, S. N., Smith, W. B., Miles, P. D. and Pugh, S. A.: Forest Resources of the United States, 2017., 2019.
- Peng, J., Wang, Y. P., Houlton, B. Z., Dan, L., Pak, B. and Tang, X.: Global Carbon Sequestration Is Highly Sensitive to
Model-Based Formulations of Nitrogen Fixation, *Global Biogeochem. Cycles*, 34(1), e2019GB006296,
1180 doi:10.1029/2019GB006296, 2020.
- Phillips, R. P., Brzostek, E. and Midgley, M. G.: The mycorrhizal-associated nutrient economy: A new framework for
predicting carbon-nutrient couplings in temperate forests, *New Phytol.*, 199(1), 41–51, doi:10.1111/nph.12221, 2013.
- Poorter, H., Niinemets, Ü., Poorter, L., Wright, I. J. and Villar, R.: Causes and consequences of variation in leaf mass per
area (LMA): a meta-analysis, *New Phytol.*, 182(3), 565–588, doi:10.1111/j.1469-8137.2008.02681.x, 2009.
- 1185 Poorter, H., Niklas, K. J., Reich, P. B., Oleksyn, J., Poot, P. and Mommer, L.: Biomass allocation to leaves, stems and roots:
Meta-analyses of interspecific variation and environmental control, *New Phytol.*, 193(1), 30–50, doi:10.1111/j.1469-
8137.2011.03952.x, 2012.
- Purves, D. W., Lichstein, J. W., Strigul, N. and Pacala, S. W.: Predicting and understanding forest dynamics using a simple
tractable model, *Proc. Natl. Acad. Sci.*, 105(44), 17018–17022, doi:10.1073/pnas.0807754105, 2008.
- 1190 Rastetter, E. B. and Shaver, G. R.: A model of multiple-element limitation for acclimating vegetation, *Ecology*, 73(4), 1157–
1174, doi:10.2307/1940666, 1992.



- Reay, D. S., Dentener, F. J., Smith, P., Grace, J. and Feely, R. A.: Global nitrogen deposition and carbon sinks, *Nat. Geosci.*, 1(7), 430–437, doi:10.1038/ngeo230, 2008.
- Reed, S. C., Cleveland, C. C. and Townsend, A. R.: Functional Ecology of Free-Living Nitrogen Fixation: A Contemporary
1195 Perspective, *Annu. Rev. Ecol. Evol. Syst.*, 42(1), 489–512, doi:10.1146/annurev-ecolsys-102710-145034, 2011.
- Richardson, A. D., Carbone, M. S., Keenan, T. F., Czimeczik, C. I., Hollinger, D. Y., Murakami, P., Schaberg, P. G. and Xu,
X.: Seasonal dynamics and age of stemwood nonstructural carbohydrates in temperate forest trees, *New Phytol.*, 197(3),
850–861, doi:10.1111/nph.12042, 2013.
- Roper, M. M.: Straw decomposition and nitrogenase activity (C_2H_2 reduction): Effects of soil moisture and temperature, *Soil*
1200 *Biol. Biochem.*, 17(1), 65–71, doi:10.1016/0038-0717(85)90091-4, 1985.
- Roumet, C., Birouste, M., Picon-Cochard, C., Ghestem, M., Osman, N., Vrignon-Brenas, S., Cao, K. fang and Stokes, A.:
Root structure-function relationships in 74 species: Evidence of a root economics spectrum related to carbon economy, *New*
Phytol., 210(3), 815–826, doi:10.1111/nph.13828, 2016.
- Scharlemann, J. P. W., Tanner, E. V. J., Hiederer, R. and Kapos, V.: Global soil carbon: Understanding and managing the
1205 largest terrestrial carbon pool, *Carbon Manag.*, 5(1), 81–91, doi:10.4155/cmt.13.77, 2014.
- Schomberg, H. H. and Weaver, R. W.: Nodulation, Nitrogen Fixation, and Early Growth of Arrowleaf Clover in Response to
Root Temperature and Starter Nitrogen, *Agron. J.*, 84(6), 1046–1050, doi:10.2134/agronj1992.00021962008400060026x,
1992.
- Schulte-Uebbing, L. and de Vries, W.: Global-scale impacts of nitrogen deposition on tree carbon sequestration in tropical,
1210 temperate, and boreal forests: A meta-analysis, *Glob. Chang. Biol.*, 24(2), e416–e431, doi:10.1111/gcb.13862, 2018.
- Sheffer, E., Batterman, S. A., Levin, S. A. and Hedin, L. O.: Biome-scale nitrogen fixation strategies selected by climatic
constraints on nitrogen cycle, *Nat. Plants*, 1(12), 1–6, doi:10.1038/nplants.2015.182, 2015.
- Sheffield, J., Goteti, G. and Wood, E. F.: Development of a 50-Year High-Resolution Global Dataset of Meteorological
Forcings for Land Surface Modeling, *J. Clim.*, 19(13), 3088–3111, doi:10.1175/JCLI3790.1, 2006.
- 1215 Shevliakova, E., Malyshev, S., Martinez-Cano, I., Milly, P. C. D., Pacala, S. W., Ginoux, P., Dunne, K. A., Dunne, J. P.,
Dupuis, C., Findell, K., Ghannam, K., Horowitz, L. W., John, J. G., Knutson, T. R., Krasting, P. J., Naik, V., Zadeh, N.,
Zeng, F. and Zeng, Y.: The land component LM4.1 of the GFDL Earth System Model ESM4.1: biophysical and
biogeochemical processes and interactions with climate, n.d.
- Smith, B., Wärlind, D., Arneth, A., Hickler, T., Leadley, P., Siltberg, J. and Zaehle, S.: Implications of incorporating N
1220 cycling and N limitations on primary production in an individual-based dynamic vegetation model, *Biogeosciences*, 11(7),
2027–2054, doi:10.5194/bg-11-2027-2014, 2014.
- Smith, S. E. and Smith, F. A.: Roles of Arbuscular Mycorrhizas in Plant Nutrition and Growth: New Paradigms from
Cellular to Ecosystem Scales, *Annu. Rev. Plant Biol.*, 62(1), 227–250, doi:10.1146/annurev-arplant-042110-103846, 2011.
- Staccone, A., Liao, W., Perakis, S., Compton, J., Clark, C. and Menge, D.: A spatially explicit, empirical estimate of tree-
1225 based biological nitrogen fixation in forests of the United States, *Global Biogeochem. Cycles*, 34(2), e2019GB006241,



- doi:10.1029/2019gb006241, 2020.
- Staccone, A. P., Kou-Giesbrecht, S., Taylor, B. N. and Menge, D. N. L.: Nitrogen-fixing trees have no net effect on forest growth in the coterminous United States, *J. Ecol.*, 109(2), 877–887, doi:10.1111/1365-2745.13513, 2021.
- Stehfest, E. and Bouwman, L.: N₂O and NO emission from agricultural fields and soils under natural vegetation:
1230 Summarizing available measurement data and modeling of global annual emissions, *Nutr. Cycl. Agroecosystems*, 74(3), 207–228, doi:10.1007/s10705-006-9000-7, 2006.
- Stocker, B. D., Prentice, I. C., Cornell, S. E., Davies-Barnard, T., Finzi, A. C., Franklin, O., Janssens, I., Larmola, T., Manzoni, S., Näsholm, T., Raven, J. A., Rebel, K. T., Reed, S., Vicca, S., Wiltshire, A. and Zaehle, S.: Terrestrial nitrogen cycling in Earth system models revisited, *New Phytol.*, 210(4), 1165–1168, doi:10.1111/nph.13997, 2016.
- 1235 Strigul, N., Pristinski, D., Purves, D., Dushoff, J. and Pacala, S.: Scaling from trees to forests: Tractable macroscopic equations for forest dynamics, *Ecol. Monogr.*, 78(4), 523–545, doi:10.1890/08-0082.1, 2008.
- Sullivan, B. W., Smith, W. K., Alan, R., Nasto, M. K., Reed, S. C. and Chazdon, R. L.: Spatially robust estimates of biological nitrogen (N) fixation imply substantial human alteration of the tropical N cycle, *Proc. Natl. Acad. Sci.*, 111(22), 8101–8106, doi:10.1073/pnas.1511978112, 2014.
- 1240 Sulman, B. N., Shevliakova, E., Brzostek, E. R., Kivlin, S. N., Malyshev, S., Menge, D. N. L. and Zhang, X.: Diverse mycorrhizal associations enhance terrestrial C storage in a global model, *Global Biogeochem. Cycles*, 33(4), 501–523, doi:10.1029/2018GB005973, 2019.
- Swank, W. T. and Waide, J. B.: Characterization of Baseline Precipitation and Stream Chemistry and Nutrient Budgets for Control Watersheds, in *Forest Hydrology and Ecology at Coweeta.*, 1988.
- 1245 Taylor, B. N., Chazdon, R. L., Bachelot, B. and Menge, D. N. L.: Nitrogen-fixing trees inhibit growth of regenerating Costa Rican rainforests, *Proc. Natl. Acad. Sci.*, 114(33), 8817–8822, doi:10.1073/pnas.1707094114, 2017.
- Terrer, C., Vicca, S., Hungate, B. A., Phillips, R. P. and Prentice, I. C.: Mycorrhizal association as a primary control of the CO₂ fertilization effect., *Science*, 353(6294), 72–74, doi:10.1126/science.aaf4610, 2016.
- Terrer, C., Vicca, S., Stocker, B. D., Hungate, B. A., Phillips, R. P., Reich, P. B., Finzi, A. C. and Prentice, I. C.: Ecosystem
1250 responses to elevated CO₂ governed by plant–soil interactions and the cost of nitrogen acquisition, *New Phytol.*, 217(2), 507–522, doi:10.1111/nph.14872, 2018.
- Terrer, C., Prentice, I., Jackson, R., Keenan, T., Kaiser, C., Vicca, S., Fisher, J., Reich, P., Stocker, B., Hungate, B., Penuelos, J., McCallum, I., Soudzilovskala, N., Cernusak, L., Talhelm, A., Van, S. K., Piao, S., Newton, P., Hovenden, M., Blumenthal, D., Liu, Y., Muller, C., Winter, K., Field, C., Viechtbauer, W., Van, L. C., Hoosbeek, M., Watanabe, M., Koike,
1255 T., Leshyk, V., Polley, W. and Franklin, O.: Nitrogen and phosphorus constrain the CO₂ fertilization of global plant biomass, *Nat. Clim. Chang.*, 9(9), 684–689, doi:10.1038/s41558-019-0545-2, 2019.
- Thomas, R. Q., Brookshire, E. N. J. and Gerber, S.: Nitrogen limitation on land: How can it occur in Earth system models?, *Glob. Chang. Biol.*, 21(5), 1777–1793, doi:10.1111/gcb.12813, 2015.
- Todd, R. L., Meyer, R. D. and Waide, J. B.: Nitrogen Fixation in a Deciduous Forest in the South-Eastern United States,



- 1260 Ecol. Bull., 26, 172–177, 1978.
Treseder, K. K.: A meta-analysis of mycorrhizal responses to nitrogen, phosphorus, and atmospheric CO₂ in field studies, *New Phytol.*, 164(2), 347–355, doi:10.1111/j.1469-8137.2004.01159.x, 2004.
US Forest Service: US Forest Inventory and Analysis Database., 2020a.
US Forest Service: US Forest Inventory and Analysis Forest Health Monitoring Database., 2020b.
- 1265 Vicca, S., Stocker, B. D., Reed, S., Wieder, W. R., Bahn, M., Fay, P. A., Janssens, I. A., Lambers, H., Peñuelas, J., Piao, S., Rebel, K. T., Sardans, J., Sigurdsson, B. D., Van Sundert, K., Wang, Y. P., Zaehle, S. and Ciais, P.: Using research networks to create the comprehensive datasets needed to assess nutrient availability as a key determinant of terrestrial carbon cycling, *Environ. Res. Lett.*, 13(12), 125006, doi:10.1088/1748-9326/aaeae7, 2018.
Vitousek, P. M., Menge, D. N., Reed, S. C. and Cleveland, C. C.: Biological nitrogen fixation: rates, patterns and ecological
1270 controls in terrestrial ecosystems, *Philos. Trans. R. Soc. B Biol. Sci.*, 368(1621), 20130119, doi:10.1098/rstb.2013.0119, 2013.
Wang, Y. P. and Houlton, B. Z.: Nitrogen constraints on terrestrial carbon uptake: Implications for the global carbon-climate feedback, *Geophys. Res. Lett.*, 36(24), 1–5, doi:10.1029/2009GL041009, 2009.
Wang, Y. P., Law, R. M. and Pak, B.: A global model of carbon, nitrogen and phosphorus cycles for the terrestrial biosphere,
1275 *Biogeosciences*, 7(7), 2261–2282, doi:10.5194/bg-7-2261-2010, 2010.
Weng, E., Malyshev, S., Lichstein, J. W., Farrior, C. E., Dybzinski, R., Zhang, T., Shevliakova, E. and Pacala, S. W.: Scaling from individuals to ecosystems in an Earth System Model using a mathematically tractable model of height-structured competition for light, *Biogeosciences*, 11(12), 17757–17860, doi:10.5194/bgd-11-17757-2014, 2015.
Wieder, W. R., Cleveland, C. C., Lawrence, D. M. and Bonan, G. B.: Effects of model structural uncertainty on carbon cycle
1280 projections: biological nitrogen fixation as a case study, *Environ. Res. Lett.*, 10(4), 044016, doi:10.1088/1748-9326/10/4/044016, 2015a.
Wieder, W. R., Cleveland, C. C., Smith, W. K. and Todd-Brown, K.: Future productivity and carbon storage limited by terrestrial nutrient availability, *Nat. Geosci.*, 8(6), 441–444, doi:10.1038/ngeo2413, 2015b.
Wright, S. J., Turner, B. L., Yavitt, J. B., Harms, K. E., Kaspari, M., Tanner, E. V. J., Bujan, J., Griffin, E. A., Mayor, J. R.,
1285 Pasquini, S. C., Sheldrake, M. and Garcia, M. N.: Plant responses to fertilization experiments in lowland, species-rich, tropical forests, *Ecology*, 99(5), 1129–1138, doi:10.1002/ecy.2193, 2018.
Xu, H., Detto, M., Fang, S., Chazdon, R. L., Li, Y., Hau, B. C. H., Fischer, G. A., Weiblen, G. D., Hogan, J. A., Zimmerman, J. K., Uriarte, M., Thompson, J., Lian, J., Cao, K., Kenfack, D., Alonso, A., Bissengou, P., Memiaghe, H. R., Valencia, R., Yap, S. L., Davies, S. J., Mi, X. and Yao, T. L.: Soil nitrogen concentration mediates the relationship between
1290 leguminous trees and neighbor diversity in tropical forests, *Commun. Biol.*, 3(1), 1–8, doi:10.1038/s42003-020-1041-y, 2020.
Yang, X., Wittig, V., Jain, A. K. and Post, W.: Integration of nitrogen cycle dynamics into the Integrated Science Assessment Model for the study of terrestrial ecosystem responses to global change, *Global Biogeochem. Cycles*,



- 23(GB4029), 1–18, doi:10.1029/2009GB003474, 2009.
- 1295 Zaehle, S. and Friend, A. D.: Carbon and nitrogen cycle dynamics in the O-CN land surface model: 1. Model description, site-scale evaluation, and sensitivity to parameter estimates, *Global Biogeochem. Cycles*, 24(GB1006), 1–13, doi:10.1029/2009GB003521, 2010.
- Zhang, J. and Elser, J. J.: Carbon:Nitrogen:Phosphorus stoichiometry in fungi: A meta-analysis, *Front. Microbiol.*, 8, 1–9, doi:10.3389/fmicb.2017.01281, 2017.
- 1300 Zhao, M., Golaz, J.-C., Held, I. M., Guo, H., Balaji, V., Benson, R., Chen, J.-H., Chen, X., Donner, L. J., Dunne, J. P., Dunne, K., Durachta, J., Fan, S.-M., Freidenreich, S. M., Garner, S. T., Ginoux, P., Harris, L. M., Horowitz, L. W., Krasting, J. P., Langenhorst, A. R., Liang, Z., Lin, P., Lin, S.-J., Malyshev, S. L., Mason, E., Milly, P. C. D., Ming, Y., Naik, V., Paulot, F., Paynter, D., Phillipps, P., Radhakrishnan, A., Ramaswamy, V., Robinson, T., Schwarzkopf, D., Seman, C. J., Shevliakova, E., Shen, Z., Shin, H., Silvers, L. G., Wilson, J. R., Winton, M., Wittenberg, A. T., Wyman, B. and Xiang, B.:
1305 The GFDL Global Atmosphere and Land Model AM4.0/LM4.0: 1. Simulation Characteristics With Prescribed SSTs, *J. Adv. Model. Earth Syst.*, 10(3), 691–734, doi:10.1002/2017MS001208, 2018a.
- Zhao, M., Golaz, J.-C., Held, I. M., Guo, H., Balaji, V., Benson, R., Chen, J.-H., Chen, X., Donner, L. J., Dunne, J. P., Dunne, K., Durachta, J., Fan, S.-M., Freidenreich, S. M., Garner, S. T., Ginoux, P., Harris, L. M., Horowitz, L. W., Krasting, J. P., Langenhorst, A. R., Liang, Z., Lin, P., Lin, S.-J., Malyshev, S. L., Mason, E., Milly, P. C. D., Ming, Y., Naik, V.,
1310 Paulot, F., Paynter, D., Phillipps, P., Radhakrishnan, A., Ramaswamy, V., Robinson, T., Schwarzkopf, D., Seman, C. J., Shevliakova, E., Shen, Z., Shin, H., Silvers, L. G., Wilson, J. R., Winton, M., Wittenberg, A. T., Wyman, B. and Xiang, B.:
The GFDL Global Atmosphere and Land Model AM4.0/LM4.0: 2. Model Description, Sensitivity Studies, and Tuning Strategies, *J. Adv. Model. Earth Syst.*, 10(3), 735–769, doi:10.1002/2017MS001209, 2018b.
- Zheng, M., Zhou, Z., Luo, Y., Zhao, P. and Mo, J.: Global pattern and controls of biological nitrogen fixation under nutrient
1315 enrichment: A meta-analysis, *Glob. Chang. Biol.*, 25(9), 3018–3030, doi:10.1111/gcb.14705, 2019.
- Zheng, M., Zhou, Z., Zhao, P., Luo, Y., Ye, Q., Zhang, K., Song, L. and Mo, J.: Effects of human disturbance activities and environmental change factors on terrestrial nitrogen fixation, *Glob. Chang. Biol.*, 26(11), 6203–6217, doi:10.1111/gcb.15328, 2020.
- Zhu, Y.-G. and Miller, R. M.: Carbon cycling by arbuscular mycorrhizal fungi in soil–plant systems, *Trends Plant Sci.*, 8(9),
1320 407–409, doi:10.1016/S1360-1385(03)00192-4, 2003.