

1 **Additional carbon inputs to reach a 4 per 1000 objective in**  
2 **Europe: feasibility and projected impacts of climate change**  
3 **based on Century simulations of long-term arable**  
4 **experiments**

5 Elisa Bruni<sup>1</sup>, Bertrand Guenet<sup>1,2</sup>, Yuanyuan Huang<sup>3</sup>, Hugues Clivot<sup>4,5</sup>, Iñigo Virto<sup>6</sup>,  
6 Roberta Farina<sup>7</sup>, Thomas Kätterer<sup>8</sup>, Philippe Ciais<sup>1</sup>, Manuel Martin<sup>9</sup>, Claire Chenu<sup>10</sup>

7 <sup>1</sup>Laboratoire des Sciences du Climat et de l'Environnement, LSCE/IPSL, CEA-CNRS-UVSQ, Université  
8 Paris-Saclay, F-91191 Gif-sur-Yvette, France

9 <sup>2</sup>LG-ENS (Laboratoire de géologie) - CNRS UMR 8538 - École normale supérieure, PSL University -  
10 IPSL, 75005 Paris France

11 <sup>3</sup>CSIRO Oceans and Atmosphere, Aspendale 3195, Australia

12 <sup>4</sup>Université de Lorraine, INRAE, LAE, 68000, Colmar, France

13 <sup>5</sup>Université de Reims Champagne Ardenne, INRAE, FARE, UMR A 614, 51097 Reims, France

14 <sup>6</sup>Departamento de Ciencias. IS-FOOD, Universidad Pública de Navarra, 31009 Pamplona, Spain

15 <sup>7</sup>CREA - Council for Agricultural Research and Economics, Research Centre for Agriculture  
16 and Environment, 00198 Rome, Italy

17 <sup>8</sup>Swedish University of Agricultural Sciences, Department of Ecology, Box 7044, 75007 Uppsala, Sweden

18 <sup>9</sup>INRA Orléans, InfoSolUnit, Orléans, France

19 <sup>10</sup>Ecosys, INRA-AgroParisTech, Université Paris-Saclay, Campus AgroParisTech, 78850 Thiverval-  
20 Grignon, France

21 *Correspondence to:* Elisa Bruni (elisa.bruni@lscce.ipsl.fr)

22

23

24

25

26

27

28

29

30 **Abstract.** The 4 per 1000 initiative aims to maintain and increase soil organic carbon (SOC) stocks for soil  
31 fertility, food security and climate change adaptation and mitigation. One way to enhance SOC stocks is to  
32 increase carbon (C) inputs to the soil.

33 In this study, we assessed the amount of organic C inputs that are necessary to reach a target of SOC stocks  
34 increase by 4% per year on average, for 30 years, in 14 long-term agricultural sites in Europe. We used the  
35 Century model to simulate SOC stocks and assessed the required level of additional C inputs to reach the 4  
36 per 1000 target at these sites. Then, we analyzed how this would change under future scenarios of  
37 temperature increase. Initial stocks were simulated assuming steady state. We compared modelled C inputs  
38 to different treatments of additional C used on the experimental sites (exogenous organic matter addition  
39 and one treatment with different crop rotations). The model was calibrated to fit the control plots, i.e.  
40 conventional management without additional C inputs from exogenous organic matter or changes in crop  
41 rotations, and was able to reproduce the SOC stocks dynamics.

42 We found that, on average among the selected experimental sites, annual C inputs will have to increase by  
43  $43.15 \pm 5.05$  %, which is  $0.66 \pm 0.23$  MgC ha<sup>-1</sup> per year (mean  $\pm$  standard error), with respect to the initial  
44 C inputs in the control treatment. The simulated amount of C inputs required to reach the 4% SOC increase  
45 was lower or similar to the amount of C inputs actually used in the majority of the additional C input  
46 treatments of the long-term experiments. However, Century might be overestimating the effect of  
47 additional C inputs on SOC stocks. In the experimental sites, we found that treatments with additional C  
48 inputs were increasing by 0.25% on average. This means that the C inputs required to reach the 4 per 1000  
49 target might actually be much higher. Furthermore, we estimated that annual C inputs will have to increase  
50 even more due to climate warming, that is 54% more and 120% more, for a 1°C and 5°C warming,  
51 respectively. We showed that modelled C inputs required to reach the target depended linearly on the initial  
52 SOC stocks, raising concern on the feasibility of the 4 per 1000 objective in soils with a higher potential  
53 contribution on C sequestration, that is soils with high SOC stocks. Our work highlights the challenge of  
54 increasing SOC stocks at large scale and in a future with warmer climate.

## 55 **1 Introduction**

56 Increasing organic carbon (C) stocks in agricultural soils is beneficial for soil fertility and crop production  
57 and for climate change adaptation and mitigation. This consideration was at the basis of the 4 per 1000  
58 (4p1000) initiative, proposed by the French Government during the 21<sup>st</sup> Conference of the Parties (COP21)  
59 on climate change. The 4p1000 initiative aims to promote agricultural practices that enable the  
60 conservation of organic carbon in the soil ([www.4p1000.org](http://www.4p1000.org)). Because soil organic carbon (SOC) stocks  
61 are two to three times higher than those in the atmosphere, even a small increase of the SOC pool can  
62 translate into significant changes in the atmospheric pool (Minasny et al., 2017). To demonstrate the  
63 importance of SOC, the initiative took as an example the fact that increasing global SOC stocks up to 0.4 m  
64 depth by 4p1000 (0.4%) per year of their initial value could offset the net annual carbon dioxide (CO<sub>2</sub>)  
65 anthropogenic emissions to the atmosphere (Soussana, 2017). While increasing SOC stocks by 4p1000

66 annually is not a normative target of the initiative, this value can be taken as a reference to which current  
67 situations and alternative strategies are compared (e.g. Pellerin et al., 2017).

68 Strategies of conservation and expansion of existing SOC pools may be necessary but are not sufficient to  
69 mitigate climate change (Paustian et al., 2016). In this sense, increasing SOC stocks cannot be regarded as a  
70 dispensation to continue business as usual, but rather as a wedge of negative greenhouse gases (GHG)  
71 emissions (Wollenberg et al., 2016), as well as a strategy for improving most soils' resilience to changes in  
72 the climate.

73 The potential to increase SOC stocks is particularly relevant in cropped soils, where the depletion of  
74 organic matter with respect to the original non-cultivated situation has been demonstrated (Clivot et al.,  
75 2019; Goidts and van Wesemael, 2007; Meersmans et al., 2011; Saffih-Hdadi and Mary, 2008; Sanderman  
76 et al., 2017; Zinn et al., 2005) and where straightforward management practices can be implemented to  
77 promote the conservation or increment of SOC (Chenu et al., 2019; Guenet et al., 2020; Paustian et al.,  
78 2016). Moreover, increasing the organic C content in agricultural soils is known to improve their fertility  
79 and water retention capacity (Lal 2008), indirectly enhancing agricultural productivity and food security.

80 SOC stocks are a function of C inputs and C outputs. To increase SOC stocks one can either increase C  
81 inputs to the soil (i.e. adding plant material or organic fertilizers) or reduce C outputs resulting from  
82 mineralization and, in some cases, soil erosion. Increasing SOC stocks can be achieved via agricultural  
83 practices such as retention of crop residues and organic amendments to the soil, cover cropping, diversified  
84 rotations and agroforestry systems (Chenu et al., 2019; Powlson et al., 2011). However, some of these  
85 practices only lead to local *carbon storage* at field scale, rather than a net *carbon sequestration* from the  
86 atmosphere at larger scales (Chenu et al., 2019).

87 Assessing the evolution of SOC stocks over time is important to estimate correctly the potential of SOC  
88 storage in agricultural soils and evaluate management practices in terms of both SOC stocks increase and  
89 sequestration potential. The dynamics of SOC stocks can be either measured in agricultural soils through  
90 long-term experiments (LTEs) and soil monitoring networks or estimated via biogeochemical models  
91 (Campbell and Paustian, 2015; Manzoni and Porporato, 2009). Combining measurements of SOC with  
92 models provides a wider applicability of the information collected in field trials, as it allows SOC stocks  
93 and their future trends to be estimated. However, validity of models in the studied areas has to be assessed  
94 and models need to be initialized. This means that the initial status of SOC has to be set, either for lack of  
95 data on total initial stocks, or to determine the allocation of C among model's compartments that cannot be  
96 measured. This is commonly accomplished by assuming that SOC is at equilibrium at the beginning of the  
97 experiment (Luo et al., 2017; Xia et al., 2012).

98 The feasibility and applicability of a 4‰ increase target depend on biotechnical and socio-economic  
99 factors. As we mentioned earlier, a number of practices are known to increase SOC stocks in agricultural  
100 systems. However, it is still debated whether they will be sufficient to reach the 4p1000 objective. Minasny  
101 et al. (2017) described opportunities and limitations of a 4‰ SOC increase in 20 regions across the world.  
102 Several authors (e.g. Baveye et al., 2018; van Groenigen et al., 2017; VandenBygaart, 2018) argued that

103 some of the examples described in Minasny et al. (2017) were not representative of wide-scale agriculture  
104 and suggested that a 4‰ rate is not attainable in many practical situations (Poulton et al., 2018).  
105 Implementing new agricultural practices that allow the maintenance and increase of SOC stocks might  
106 require structural land management changes that not all farmers will be willing to adopt. Incentivizing and  
107 sustaining virtuous practices to increase SOC stocks should be a strategy for policymakers to overcome  
108 socio-economic barriers (e.g. Lal, 2018; Soussana, 2017) and in order to do that, they need to be correctly  
109 informed. Recent works have assessed the biotechnical limitations of a SOC increase, studying the required  
110 and available biomass to reach a 4p1000 target in European soils (Wiesmeier et al., 2016; Martin et al.,  
111 2021; Riggers et al., 2021).

112 Our work was set up in this context with the objectives to: 1) estimate the amount of C inputs needed to  
113 increase SOC stocks by 4‰ per year; 2) investigate if this amount is attainable with currently implemented  
114 soil practices (i.e. organic amendments and different crop rotations) and 3) study how the required C inputs  
115 are going to evolve in a future driven by climate change. We used the biogeochemistry SOC model  
116 Century, which is one of the most widely used and validated models (Smith et al., 1997), to simulate SOC  
117 stocks in 14 different agricultural LTEs around Europe. We set the target of SOC stocks increase to 4‰ per  
118 year for 30 years, relative to the initial stocks in the reference treatments. With an inverse modeling  
119 approach, we estimated the amount of additional C inputs required to reach a 4p1000 target at these sites.  
120 Finally, we evaluated the dependency of the required additional C inputs to different scenarios of increased  
121 temperature.

## 122 2 Materials and methods

### 123 2.1. Experimental sites

124 We compiled data from 14 LTEs in arable cropping systems across Europe (Fig. 1), where a total of 46  
125 treatments with increased C inputs to the soil were performed and one control plot in each experiment was  
126 implemented (Table 1). The experiments lasted between 11 and 53 years (median value of 16 years) in the  
127 period from 1956 to 2018. Most of the experiments had at least 3 replicates, except for the Italian site  
128 *Foggia*, the French site *Champ Noël 3* and the British site *Broadbalk*, where no replicates were available.  
129 We selected experiments where dry matter (DM) yields and SOC had been measured at several dates. C  
130 inputs in all sites, except for control plots and all plots in *Foggia*, included exogenous organic matter  
131 (EOM) addition, e.g. animal manure, household waste, sewage sludge or compost additions. In *Foggia*,  
132 different rotations without organic matter addition were studied and compared to a wheat-only treatment,  
133 considered as the control plot. The annual C inputs to the soil were substantially higher in the rotations  
134 compared to the control. More information on crop rotations and C inputs for each treatment can be found  
135 in Table 1.

136 Cropping systems in the 60 treatments (14 control plots and 46 additional C input treatments) were mainly  
137 cereal-dominated rotations (wheat, maize, barley and oat). In particular, four were cereal monocultures

138 (silage maize in *Champ Noël 3*, *Le Rheu 1* and *Le Rheu 2* and winter wheat in *Broadbalk*) and four sites  
 139 had rotations of different cereals (winter wheat and silage or grain maize in *Crécom 3 PRO*, *Feucherolles*,  
 140 *La Jaillièrre 2 PRO* and *Avrillé*). The other sites rotated cereal crops with legumes (chickpea, pea) and/or  
 141 root crops (fodder beet, fodder rape and Swedish turnip), oilseed crops (sunflower and oilseed rape), cover  
 142 crops (mustard and rapeseed) and one rotation included tomatoes. Straw residues were systematically  
 143 exported except in French sites, where residues were sometimes incorporated into the soil as accounted for  
 144 in the C input calculations. All LTEs were under conventional tillage, which was performed with a tractor,  
 145 except in the case of *Ultuna*, where it was performed manually. All experiments were rainfed, except for  
 146 *Foggia*, where tomatoes were irrigated in summer. The French sites *Champ Noël 3*, *Crécom 3 PRO*, *La*  
 147 *Jaillièrre 2 PRO*, *Le Rheu 1* and *Trévarez* received optimal amounts of mineral fertilizers both in the control  
 148 plot and in the different organic matter treatments. All other experiments did not receive any mineral  
 149 fertilization. All control plots, apart from *Arazuri*, had decreasing SOC stock trends (SOC approximated  
 150 with a linear regression:  $SOC = m \cdot t + SOC_0$ , with average relative change:  $\frac{m}{SOC_0} \cdot 100 = -0.76 \%$ ,  $R^2 =$   
 151 0.58). Over the 46 treatments of additional C input, 18 exhibited increasing SOC stocks at a higher rate  
 152 than 4‰ per year on average over the experiment length (Table 1). Six treatments had increasing SOC  
 153 stocks, but at a lower ratio than 4p1000. The other 22 treatments with additional C inputs had decreasing  
 154 SOC stocks ( $MgC\ ha^{-1}$ ). However, the decreasing trend was, in these cases, lower than the decreasing trend  
 155 in the respective control plot, on the majority of the treatments.

156 **Table 1: Summary of the agricultural experiments included in the study: crop rotations grown at site, amount of**  
 157 **carbon inputs ( $MgC\ ha^{-1}$  per year) estimated from crop yields as in (Bolinder et al., 2007), type of treatments,**  
 158 **amount of additional organic carbon from organic treatments ( $MgC\ ha^{-1}$  per year) and mean annual SOC stocks**  
 159 **variation (%).**

Site	ID Treatment	Rotations*	Carbon inputs from crop rotations	Treatment type	Additional carbon inputs from organic treatments	SOC annual variation
			$MgC\ ha^{-1}$ year <sup>-1</sup>		$MgC\ ha^{-1}$ year <sup>-1</sup>	%
Champ Noël 3	Min**	sM	1.29	Reference+N*	0	-0.92
(CHNO3)	LP	Silage maize	1.49	Pig manure	0.79	-0.89
Colmar	T0	wW/Mg/sB/S	2.79	Reference	0	-0.78
(COL)	BIO1	wW/Mg/sB/S	3.93	Biowaste	1.01	0.15
	BOUE1	wW/Mg/sB/S	3.96	Sewage sludge	0.49	-0.61
	CFB1	wW/Mg/sB/S	4.04	Cow manure	1.07	-0.01
	DVB1	wW/Mg/sB/S	4.00	Green manure+Sewa ge sludge	1.08	0.18
	FB1	wW/Mg/sB/S	3.93	Cow manure	1.36	-0.01

Crécom 3 PRO (CREC3)	Min	wW/sM	1.84	Reference+N	0	-0.06
	FB2	wW/sM	1.92	Cow manure	1.82	0.49
	FV	wW/sM	1.96	Poultry manure	0.47	-1.46
Feucherolles (FEU)	T0	wW/ Mg	2.22	Reference	0	-0.66
	BIO1	wW/Mg	3.44	Biowaste	2.21	3.60
	DVB1	wW/Mg	3.45	Green manure+Sewa ge sludge	2.45	3.69
	FB1	wW/Mg	3.55	Cow manure	2.28	1.36
	OMR1	wW/Mg	3.45	Household waste	2.11	1.72
Jeu-les-Bois (JEU)	M0	wB/R/wW	2.99	Reference	0	-1.33
	CFB1	wB/R/wW	2.89	Cow manure	1.1	1.61
	CFB2	wB/R/wW	3.06	Poultry manure	1.94	1.52
	FB2	wB/R/wW	3.11	Cow manure	2.43	0.99
La Jaillièrè 2 PRO (LAJA2)	Min	sM/wW	1.59	Reference+N	0	-1.43
	CFB	sM/wW	1.25	Cow manure	1.14	-0.88
	CFP	sM/wW	1.21	Pig manure	1	-1.09
	CFV	sM/wW	1.31	Poultry manure	0.94	-1.60
	FB	sM/wW	1.29	Cow manure	1.44	-0.64
	FP	sM/wW	1.27	Pig manure	1.07	-1.03
	FV	sM/wW	1.40	Poultry manure	0.93	-1.59
Le Rheu 1 (RHEU1)	Min	sM	1.31	Reference+N	0	-1.51
	CFB1	sM	1.31	Cow manure	1.06	-1.21
Le Rheu 2 (RHEU2)	T0	sM	1.03	Reference	0	-1.72
	CFP1	sM	1.20	Pig manure	0.78	-1.28
	FP	sM	1.30	Pig manure	1.62	-0.74
Arazuri (ARAZ)	DO_N0	B/P/W/Sf/O	0.98	Reference	0	1.00
	D1_F1	B/P/W/Sf/O	1.40	Sewage sludge	2.82	0.40
	D1_F2	B/P/W/Sf/O	1.41	Sewage sludge	1.4	1.22
	D1_F3	B/P/W/Sf/O	1.44	Sewage sludge	0.78	1.22
	D2_F1	B/P/W/Sf/O	1.30	Sewage sludge	5.64	0.22
	D2_F2	B/P/W/Sf/O	1.40	Sewage	2.8	2.32

	D2_F3	B/P/W/Sf/O	1.49	sludge Sewage sludge	1.56	0.93
Ultuna	P0_B	O/sT/Mu/sB/FB/OsR/W/F R/M	1.03	Reference	0	-0.52
(ULTU)	S_F	O/sT/Mu/sB/FB/OsR/W/F R/M	1.10	Straw	1.77	-0.09
	GM_H	O/sT/Mu/sB/FB/OsR/W/F R/M	1.82	Green manure	1.76	0.11
	PEAT_I	O/sT/Mu/sB/FB/OsR/W/F R/M	1.14	Peat	1.97	2.17
	FYM_J	O/sT/Mu/sB/FB/OsR/W/F R/M	1.76	Farmyard Manure	1.91	0.69
	SD_L	O/sT/Mu/sB/FB/OsR/W/F R/M	0.82	Sawdust	1.84	0.56
	SS_O	O/sT/Mu/sB/FB/OsR/W/F R/M	2.59	Sewage sludge	1.84	1.36
Broadbalk	3_Nill	wW	0.36	Reference	0	-0.09
(BROAD)	19_Cast	wW	0.65	Castor meal	0.43	0.42
	22_FYM	wW	2.07	Farmyard Manure	3	0.38
Foggia***	T0	W	1.56	Reference	0	-0.86
	Dw-Dw-Fall	W/W/F	2.13	Rotation	0.57	0.01
	Dw-Fall	W/F	1.95	Rotation	0.39	-0.33
	Dw-Oa-Fall	W/O/F	2.20	Rotation	0.64	-0.33
	Dw-Dw-Cp	W/W/C	2.53	Rotation	0.97	-0.15
	Dw-Dw-To	W/W/T	2.57	Rotation	1.01	-0.59
Trévarez	Min	RG/Mg/wW/sM	1.94	Reference+N	0	-0.66
(TREV)	FB	RG/Mg/wW/sM	2.04	Cow manure	1.52	-0.39
	FP	RG/Mg/wW/sM	2.02	Pig manure	1.18	-0.18
Avrillé	T12TR	wW/sM	2.25	Reference	0	-1.18
(AVRI)	T2TR	wW/sM	2.36	Cow manure	1.68	-0.76

\*Crops: sM = silage Maize, Mg= Maize grain, wW = winter Wheat, W = Wheat,

sB = spring Barley, wB = winter Barley, B = barley, S = sugarbeet,


R = Rapeseed, Sf = Sunflower, O = Oats, P = Pea, sT = Swedish Turlip, Mu =

Mustard, DF = Fodder Beet, OsR = Oilseed Rape, FR = fodder Rape,

F = green Fallow, C = Chickpeas, T = Tomato, RG = Ray Grass

\*\*Optimal amounts of mineral fertilizers added to the control plot and to all other treatments in the experiment

\*\*\* In Foggia, additional carbon inputs from organic treatments were calculated for each rotation as the difference between C inputs in the rotation and the reference wheat-only rotation.



161

162 **Figure 1: Location of the 60 field trials distributed among the 14 cropland experiments around Europe.**

163

### 2.1.1. Climate forcing

164

Mean temperature of the sites ranged from a minimum of 5.7 °C to a maximum of 15.5 °C, while mean soil

165

humidity to approximately 20 cm depth ranged between 20.2 and 24.6 kg<sub>H<sub>2</sub>O</sub> m<sup>-2</sup><sub>soil</sub> in the dataset (Table 2).

166

When available, observed daily air temperature was used as an approximation of soil temperature.

167

Otherwise, land-atmosphere model ORCHIDEE was used to simulate soil surface temperature and soil

168

humidity at site-scale (Krinner et al., 2005). ORCHIDEE simulations were run over each site using a 3-

169

hourly global climate dataset at 0.5° (GSWP3 <http://hydro.iis.u-tokyo.ac.jp/GSWP3/>). Plant cover was set

170

to C3 plant functional type (PFT) for agriculture.

171

**Table 2: Information about experimental sites, including: mean annual values of temperature (C°) and soil humidity to approximately 20 cm depth (kg<sub>H<sub>2</sub>O</sub> m<sup>-2</sup><sub>soil</sub>) simulated with the ORCHIDEE model at each experimental site, measured pH, bulk density (g cm<sup>-3</sup>), clay (%) and initial SOC stocks in the control plots (MgC ha<sup>-1</sup>) at the experimental sites. Reference papers for each site are indicated. <sup>1</sup>For *Arazuri*, data were directly provided by the Spanish Mancomunidad de la Comarca de Pamplona.**

172

173

174

175

Sites	Reference paper	Coordinates	Years	Mean annual Temperature °C	Mean annual soil humidity kg H <sub>2</sub> O m <sup>-2</sup> <sub>soil</sub>	pH	Bulk density g cm <sup>-3</sup>	Clay %	Initial SOC stocks MgC ha <sup>-1</sup>
Champ Noël 3*	(Clivot et al., 2019)	48.09° N, 1.78° W	1990 - 2008	12.1	21.6	6.3	1.35	15.1	40.57
Colmar	(Levvasseu	48.11° N,	2000 - 2013	9.6	24.6	8.3	1.3	23.1	54.33



	r et al., 2020)	7.38 ° E				3			
Crécom 3 PRO	(Clivot et al., 2019)	48.32 ° N, 3.16 ° W	1986 - 2008	11.8	22.9	6.1 5	1.36	14.6	62
Feucherolles	(Levavasseau r et al., 2020)	48.88 ° N, 1.96 ° E	1998 - 2013	11.9	21.2	6.7 3	1.32	15.6	39.78
Jeu-les-Bois	(Clivot et al., 2019)	46.68 ° N, 1.79 ° E	1998 - 2008	12.2	22.1	6.2 7	1.52	10	48.53
La Jaillièrè 2 PRO	(Levavasseau r et al., 2020)	47.44 ° N, 0.98 ° W	1995 - 2009	12.7	20.5	6.8	1.37	20.8	32.42
Le Rheu 1*	(Clivot et al., 2019)	48.09 ° N, 1.78 ° W	1994 - 2009	12.2	21.8	5.8 5	1.27	16.4	36.23
Le Rheu 2*	(Clivot et al., 2019)	48.09 N, 1.78 W	1994 - 2009	12.2	21.8	6.0 5	1.28	13.9	36.53
Arazuri <sup>1</sup>	-	42.81 ° N, 1.72 ° W	1993 - 2018	12.7	20.4	8.6	1.67	27.9	55.39
Ultuna	(Kätterer et al., 2011)	59.82 ° N, 17.65 ° E	1956 - 2008	5.7	22.6	6.2 3	1.4	36.5	41.72
Broadbalk	(Powlson et al. 2012)	51.81 ° N, 0.37 ° W	1968 - 2015	10.2	21.5	7.8	1.25	25	24.84
Foggia	(Farina et al., 2017)	41.49 ° N, 15.48 ° E	1992 - 2008	15.5	22.4	8.1	1.32	41	63.22
Trévarez	(Clivot et al., 2019)	48.15 ° N, 3.76 ° W	1986 - 2008	11.8	23.4	6.0 1	1.48	19.2	115.33
Avrillé*	(Clivot et al., 2019)	47.50 ° N, 0.60 ° W	1983 - 1991	12.0	20.2	6.5 9	1.4	17.6	54.46

\*These experiments were part of the initial French database (AIAL) described in Clivot et al. (2019), but they were not selected for the final modelling work of this latter study. For more information, see also Bouthier et al. (2014).

176

### 2.1.2. Soil characteristics

177

The sampling depth of the experiments varied between 20 and 30 cm. SOC stocks were measured in 3 – 4 replicates, apart from *Foggia* and *Champ Noël 3* experiments, where no replicates were available, and *Broadbalk*. In this experiment, SOC was measured in each plot using a semi-cylindrical auger where 10-20 cores were taken from across the plot and bulked together (more details can be found on the e-RA website<sup>1</sup>). The clay content ranged from 10% (*Jeu-les-Bois*) to 41% (*Foggia*). Soil pH varied from a minimum of 5.85 in *Le Rheu 1* to a maximum of 8.33 in *Colmar*. The average bulk density (BD) in the control plots was 1.38 g cm<sup>-3</sup>. SOC stocks (MgC ha<sup>-1</sup>) were calculated at each site using the following equation:

178

179

180

181

182

183

184

<sup>1</sup> www.era.rothamsted.ac.uk

185  $SOC (MgC\ ha^{-1}) = SOC(\%) \cdot BD(g\ cm^{-3}) \cdot sampling\ depth\ (cm),$  (1)


186 where SOC (%) is the concentration of organic C in the soil, BD is the average bulk density of the  
 187 experimental plot. It should be noted that the application of EOMs might induce differences in BD with  
 188 time, which in turn affects the calculations of SOC stocks. No adjustment was made in this sense, since  
 189 data on the evolution of BD was available only for a few sites. This might explain differences between the  
 190 SOC stocks calculated for *Broadbalk* in this paper and those found by Powlson et al. (2012) in the same  
 191 site, by adjusting soil weights to observed decreases in top soil BD due to accumulating farmyard manure  
 192 (FYM). Initial SOC stocks values in the control plot and mean climate variables for each site are reported  
 193 in Table 2.

194 **2.1.3. Carbon inputs**

195 The allocation of C in the aboveground and belowground parts of the plant was estimated with the  
 196 approach first described by Bolinder et al. (2007) for Canadian experiments and then adapted by Clivot  
 197 et al. (2019) to the same French sites we use in this study. This methodology allows splitting C inputs from  
 198 crop residues after harvest into aboveground and belowground C inputs, using measured dry matter yields  
 199 and estimations of the shoot-to-root ratio (S:R) and harvest indexes (HI) of the crops (see Fig. 2). The  
 200 aboveground plant material is estimated as the harvested part of the plant ( $C_P$ ), which is exported from the  
 201 soil, plus the straw and stubble that are left in the soil after harvest ( $C_S$ ). The harvested part consists of the  
 202 measurements of DM yields ( $Y_P$ ), while the straw and stubble are estimated using the HI coefficient of the  
 203 different crops in the rotation (Bolinder et al., 2007). We assumed that the values used in Clivot et al.  
 204 (2019) for the HI compiled from French experimental sites were applicable to all the sites in our dataset,  
 205 which mainly include temperate sites over Europe. When these values were not available for some crops,  
 206 they have been directly derived from Bolinder et al. (2007) or other sources in the literature (S:R ratio for  
 207 fallow from Mekonnen, Buresh, and Jama (1997) and tomato from Lovelli et al. (2012)). When straw was  
 208 exported from the field, we considered that only a fraction of  $C_S$  was left on the soil. This fraction was set  
 209 to 0.4 for all sites and to 0.2 in *Ultuna*, where almost no stubble was left on the soil, since plots were  
 210 harvested by hand and crops were cut at the soil surface. We considered a C content of 0.44 gC gDM<sup>-1</sup> in  
 211 the aboveground plant material (Redin et al., 2014) and 0.4 gC gDM<sup>-1</sup> in the belowground part material  
 212 (Bolinder et al., 2007). We used the asymptotic equation of Gale and Grigal (1987) to determine the  
 213 cumulative BG input fraction from the soil surface to a considered depth:

214  $BG_{F\ depth} = 1 - \beta^{depth},$  (2)

215 where  $\beta$  is a crop-specific parameter determined using the root distributions for temperate agricultural  
 216 crops, reported in Fan et al. (2016) and Clivot et al. (2019). The depth was set to 30 cm, since it was the  
 217 depth at which soil samples were taken in the majority of the sites. For more details on the C inputs  
 218 allocation method and the allometric functions involved, see Bolinder et al. (2007) and Clivot et al. (2019).



219  
220  
221  
222  
223

Figure 2: Adapted from (Bolinder et al., 2007). Representation of the distribution of carbon in the different parts of the plant:  $C_P$  represents the carbon in the harvested product (grain, forage, tuber);  $C_S$  is the carbon in the aboveground residues (straw, stover, chaff);  $C_R$  is the carbon present in roots and  $C_E$  represents all the extra-root carbon (including all root-derived materials not usually recovered in the root fraction).

224

## 2.2. Century model

225

### 2.2.1. Model description

226

For this study, we selected the Century model, which has proved to be well suited to simulate accurately the soil C dynamics in a range of pedoclimatic areas and cropping systems (Bortolon et al., 2011; Cong et al., 2014; Parton et al., 1993), and because we had the full command of the model for fine tuning of parameters. Soil C dynamics in a soil organic matter (SOM) model with first-order kinetics can be mathematically described by the following first-order differential matrix equation:

231

$$\frac{dSOC(t)}{dt} = I + A \cdot \xi_{TWLCl}(t) \cdot K \cdot SOC(t), \quad (3)$$

232

where  $I$  is the vector of the external C inputs to the soil system, with four nonzero elements (Fig. 3). The second term  $A \cdot \xi_{TWLCl}(t) \cdot K \cdot SOC(t)$  of the equation represents organic matter decomposition rates (diagonal matrix  $K$ ), losses through respiration ( $\xi_{TWLCl}(t)$ ), transfers of C among different SOC pools ( $A$ ) and SOC evolution with time ( $SOC(t)$ ) (see Appendix A). We used the daily time-step version of the SOM model Century (Parton et al., 1988) to simulate the amount of C inputs required to reach a 4‰ annual increase of SOC storage over 30 years. In the version used, only SOC is modelled and plant growth is directly accounted as variations of C inputs. The original version of Century simulates the fluxes of SOC depending on soil relative humidity, temperature and texture (as a percentage of clay). As shown in Fig. 3, the model is discretized into 7 compartments that exchange C with each other: 4 pools of litter (aboveground metabolic, belowground metabolic, aboveground structural and belowground structural) and 3 pools of SOC (active, slow and passive). The litter C is partially released to the atmosphere as respired

242


243 CO<sub>2</sub> and partially converted to SOM in the active, slow and passive pools (see Table S1 in the supporting  
 244 information for default Century parameters). The decomposition rate of C in the  $i^{th}$  pool depends on  
 245 climatic conditions, litter and soil characteristics and is calculated using environmental response functions,  
 246 as follows:

$$247 \xi_{TWLCl}(t)_i \cdot K_i = k_i \cdot f_T(t) \cdot f_W(t) \cdot f_{L,i} \cdot f_{Clay,i}, \quad (4)$$

248 where  $i = 1, \dots, 7$  is one of the aboveground (AG) and belowground (BG) metabolic and structural litter  
 249 pools, and the active, slow and passive SOC pools;  $K_i$  is the  $(K)_{ii}$  element of the diagonal matrix  $\mathbf{K}$  in Eq.  
 250 (3);  $k_i$  is the specific mineralization rate of pool  $i$ ,  $f_T(t)$  is a function of daily soil temperature,  $f_W(t)$  is a  
 251 function used as a proxy to describe the effects of soil moisture,  $f_{L,i}$  is a reduction rate parameter acting on  
 252 the AG and BG structural pools only, depending on the lignin concentration in the litter and  $f_{Clay,i}$  is a  
 253 reduction rate function of clay on SOC mineralization in the active pool. The temperature function  $f_T(t)$   
 254 describes the exponential dependence of soil decomposition on surface temperature, through the  $Q_{10}$   
 255 relationship that was first presented by M. J. H. van't Hoff in 1884:

$$256 f_T(t) = Q_{10}^{\frac{(T(t)-T_{ref})}{10}}, \quad (5)$$

257 where  $Q_{10}$  is the temperature coefficient, usually set to 2 and  $T_{ref}$  is the reference temperature of 30 °C. The  
 258  $Q_{10}$  factor is a measure of the soil respiration change rate as a consequence of increasing temperature by  
 259 10°. The other environmental response functions are described in Appendix A.



260

261 **Figure 3: Representation of litter and soil organic carbon (SOC) pools in Century. The model takes as inputs**  
 262 **litter carbon from plants (aboveground metabolic (I<sub>1</sub>), belowground metabolic (I<sub>2</sub>), aboveground structural (I<sub>3</sub>)**  
 263 **and belowground structural (I<sub>4</sub>)). A certain fraction of carbon can be transferred from one pool to another and**  
 264 **each time a transfer occurs, part of this carbon is respired and leaves the system to the atmosphere as CO<sub>2</sub>. The**  
 265 **SOC active pool receives carbon from each litter pool, while only the structural material is transferred to the**  
 266 **SOC slow pool. Litter material never goes directly to the SOC passive pool while the three SOC pools exchange**  
 267 **C within each other.**

268

### 2.2.2. Model initialization

269 The initialization of the model consists of specifying the sizes of the SOC pools at the beginning of the  
 270 experiment. Here, we assumed initial pools are in equilibrium with C inputs before the experiments begin,  
 271 in absence of knowledge about past land use and climate making initial pools different from steady state  
 272 (Sanderman et al., 2017). Then, initialization can be done either by running the model iteratively for  
 273 thousands of years to approximate the steady state solution (numerical spin-up), or semi-analytically by  
 274 solving the set of differential equations that describes the C transfers within model compartments (Xia et  
 275 al., 2012). We solved the matrix equation by inverse calculations for determining pools sizes at steady  
 276 state, as in Xia et al. (2012) and Huang et al. (2018). These authors demonstrated that the matrix inversion  
 277 approach exactly reproduces the steady state and SOC dynamics of the model. By speeding up the  
 278 performance of the simulations, this technique allowed us to perform the optimization of model parameters,  
 279 the sensitivity analysis of SOC to climatic variables and the quantification of model outputs uncertainties  
 280 through Monte-Carlo (MC) iterative procedures. We solved the matrix equation by using its semi-analytical  
 281 solution and the following algorithm: 1) calculating annual averages of matrix items obtained by Century  
 282 simulations, driven by 30 years of climatic forcing; 2) setting Eq. (3) to zero to solve the state vector **SOC**.  
 283 For each agricultural site, the 30 years of climate forcing were set as the 30 years preceding the beginning  
 284 of the experiment, and the litter input estimated from observed vegetation was set to be the average litter  
 285 input in the control plot over the experiment duration.

286

### 2.2.3. Model calibration: optimization of the metabolic:structural fractions of the litter 287 inputs

288 In the Century model, AG and BG carbon inputs are further separated into metabolic and structural  
 289 fractions, according to the lignin to nitrogen (L:N) ratio. Because the L:N ratio was not available for all the  
 290 crops in the database, we fitted model simulations to observed SOC dynamics for the control plot of each  
 291 site, i.e. the reference plot without additional C inputs, in order to get the metabolic:structural (M:S)  
 292 fraction of the AG and BG carbon inputs. We used the sequential least-squares quadratic programming  
 293 function in Python (SciPy v1.5.1, `scipy.optimize` package with `method='SLSQP'`), a nonlinear constrained,  
 294 gradient-based optimization algorithm (Fu et al., 2019). We successfully performed the optimization on 13  
 295 sites, where at least three measures of SOC stocks were available. For *Jeu-les-Bois*, which includes two  
 296 SOC measurements only, we decided to use the same optimized values as for *Feucherolles*, which has  
 297 similar pedoclimatic conditions and crop rotations. The optimization consisted in minimizing the following  
 298 function:

$$299 \quad J_{fit} = \sum_{i=1}^n \frac{(SOC_i^{model} - SOC_i^{obs})^2}{\sigma_i^2 SOC_{obs}^2}, \quad (6)$$

300 where  $i=1, \dots, n$  is the year of the experiment,  $SOC_i^{model}$  (MgC ha<sup>-1</sup>) is the SOC simulated with Century for  
 301 year  $i$ ,  $SOC_i^{obs}$  (MgC ha<sup>-1</sup>) is the observed SOC for year  $i$  in the control plot and  $\sigma_i^2 SOC_{obs}^2$  is the variance of

302 the  $SOC_i^{obs}$  estimated from the different replicates. When replicates were not available, we recalculated  
 303  $\sigma^{2SOC_{obs}}$  as the variance amongst  $SOC^{obs}$  samples of the whole experiment. The optimized M:S values are  
 304 reported in Table 3 and represent the average quality of litter C in the rotating crops along the duration of  
 305 the experiments that match control SOC data at each site.

306 **Table 3: Optimized values of the aboveground metabolic (AM), aboveground structural (AS), belowground**  
 307 **metabolic (BM) and belowground structural (BS) fractions of the litter inputs and the Q10 and reference**  
 308 **temperature (°C) parameters.**

Site	AM	AS	BM	BS	Q <sub>10</sub>	Reference temperature °C
CHNO3	0.85	0.15	0.26	0.74	5.0	21.2
COL	0.85	0.15	0.57	0.43	2.0	30.0
CREC3	0.15	0.85	0.29	0.71	2.0	30.0
FEU	0.85	0.15	0.52	0.48	5.0	21.6
JEU*	0.85	0.15	0.52	0.48	5.0	21.6
LAJA2	0.85	0.15	0.72	0.28	5.0	21.5
RHEU1	0.85	0.15	0.49	0.51	5.0	21.3
RHEU2	0.85	0.15	0.32	0.68	5.0	21.3
ARAZ	0.53	0.47	0.53	0.47	3.0	30.0
ULTU	0.85	0.15	0.85	0.15	2.2	30.0
BROAD	0.42	0.58	0.15	0.85	2.9	30.0
FOGGIA	0.15	0.85	0.15	0.85	5.0	27.1
TREV1	0.15	0.85	0.15	0.85	5.0	23.0
AVRI	0.85	0.15	0.76	0.24	2.0	30.0

#### 309 2.2.4. Model calibration: optimization of temperature dependency parameters

310 We optimized the Q<sub>10</sub> and daily soil reference temperature parameters, which affect SOC decomposition.  
 311 The Q<sub>10</sub> factor is fixed to 2 in Century. However, many authors have shown that Q<sub>10</sub> measurements vary  
 312 with pedoclimatic conditions and vegetation activity (Craine et al., 2010; Lefèvre et al., 2014; Meyer et al.,  
 313 2018; Wang et al., 2010). For this reason, and to reproduce correctly interregional variations among the  
 314 sites in the dataset, we optimized both the Q<sub>10</sub> and reference temperature parameters to better fit the SOC  
 315 dynamics (MgC ha<sup>-1</sup>) of each agricultural site at control plot. We decided to bind the Q<sub>10</sub> between 1 and 5,  
 316 following the variation of Q<sub>10</sub> found by Wang et al. (2010) over 384 samples collected in the Northern  
 317 Hemisphere. The reference temperature ranged between 10 and 30°C. We used the SLSQP optimization  
 318 algorithm and the cost function of Eq. (6) to perform the optimization, which was successful in 13 sites and  
 319 we assigned the values obtained from the optimization of *Feucherolles* to *Jeu-les-Bois*, where SOC  
 320 measurements were too sparse to perform a two-dimensional optimization. Optimized values of Q<sub>10</sub> and  
 321 reference temperature are reported in Table 3.

322 Model performance in the control plot was evaluated using two residual-based metrics. The first one is the  
323 Mean Squared Deviation (MSD), decomposed into its three components to help locating the source of error  
324 of model simulations: the Squared Bias (SB), the Non-Unity slope (NU) and the Lack of Correlation (LC).  
325 The second metrics used is the Normalized Root Mean Squared Deviation (NRMSD) (see Appendix B).

### 326 **2.3. 4p1000 analysis**

#### 327 **2.3.1. Optimization of C inputs to reach the 4p1000 target**

328 After the spin-up to steady state, the model was set to calculate the SOC stocks dynamics of the control plot  
329 and the C inputs for virtual treatments, assuming an average increase of SOC stocks by 4‰ per year over  
330 30 years. 30 years is considered as a period of time over which the variation of SOC can be detected  
331 correctly. During this period length, we supposed the soil was fed with constant amounts of C inputs from  
332 plant material. For the control, we derived C inputs from measurements of DM yields and calculated the  
333 annual mean over the whole experiment length. For the virtual treatments, we used an optimization  
334 algorithm to calculate the required amount of C inputs to reach a linear increase of SOC storage by 4‰ per  
335 year above the SOC stock at the start of the simulation. Mathematically, we minimized the following  
336 function:

$$337 J_{4p1000} = |SOC_0 \cdot (1 + 0.004 \cdot 30) - SOC_{30}^{model}(\mathbf{I})|, \quad (7)$$

338 where  $\mathbf{I}$  is the 1x4 vector of C inputs to minimize over,  $SOC_0$  is the initial SOC stock and  $SOC_{30}^{model}(\mathbf{I})$  is  
339 the SOC stock after 30 years of simulation. During the optimization, the M:S fractions were allowed to  
340 vary to estimate the quality of the optimal C inputs. Instead, we kept the aboveground:belowground ratio of  
341 the C inputs fixed to its initial value, to bind the model in order to represent agronomically plausible C  
342 inputs. In fact, if not bound, the model tends to increase the belowground C fraction to unrealistic values  
343 (assuming the same crop rotations persisted on site). On the other hand, keeping the  
344 aboveground:belowground ratio fixed implies that the simulated additional C inputs will be spread equally  
345 on surface and belowground. As for the previous optimizations, we used the Python function SLSQP to  
346 solve the minimization problem. The outcome of the optimization is a 4x1 vector ( $\mathbf{I}_{opt}$ ) representing the  
347 amount of C in the four litter input pools that matches the 4p1000 rate target.

#### 348 **2.3.2. Uncertainties quantification**

349 Uncertainties of model outcomes were quantified using a Monte-Carlo approach. We initially calculated  
350 the standard error (SE) of the mean C inputs derived from yield measurements for each experimental site:

$$351 SE = \sqrt{\frac{\sigma^2_I}{s}}, \quad (8)$$

352 where  $\sigma^2_I$  is the variance of the estimated C input from yield measurements and  $s$  is the length of the  
353 experiment. If not available, we calculated  $\sigma^2_I$  as the average relative variance of C inputs among the

354 control plots. We therefore randomly generated  $N$  vectors of  $C$  inputs ( $I$ ) around the calculated standard  
355 error and performed the 4p1000 optimization  $N$  times, each time using one of the generated vectors  $I$  as a  
356 prior for the optimization. To correctly assess the uncertainty over the required  $C$  inputs we set  $N$  to 50  
357 (Anderson, 1976). The standard error of model outputs was calculated with Eq. (8), where the variance was  
358 set as the variance of the modelled carbon outputs and the experiment size ( $s$ ) to 50.

### 359 2.3.3. Sensitivity analysis to temperature

360 We tested the sensitivity of model outputs to temperature, running two simulations with increased  
361 temperatures. We considered two representative concentration pathways (RCPs) of global average surface  
362 temperature change projections (IPCC, 2015). The first scenario (RCP2.6) is the one that contemplates  
363 stringent mitigation policies and predicts that average global land temperature will increase by  $1^{\circ}\text{C}$  during  
364 the period 2081-2100, compared to the mean temperature of 1986-2005. The second scenario (RCP8.5)  
365 estimates an average temperature increase of  $+4.8^{\circ}\text{C}$ , compared to the same period of time. We ran two  
366 simulations of increasing temperature scenarios with Century. We considered the same initial conditions as  
367 the standard simulations, hence running the spin-up with the average soil temperature and relative humidity  
368 of the 30 years preceding the experiments. Then, we increased daily temperature by  $1^{\circ}\text{C}$  (AS1) and  $5^{\circ}\text{C}$   
369 (AS5) for the entire simulation length, to assess the sensitivity of modelled  $C$  inputs to increasing  
370 temperatures. Nevertheless, it must be noted that our simulations are running over a 30 years period, not the  
371 entire 21st Century. Thus, the temperature sensitivity analysis should not be considered as a test of climatic  
372 scenarios but as a classical sensitivity analysis where the boundaries were defined following RCP2.6 and  
373 RCP8.5 predictions of increased temperatures.


## 374 3 Results

### 375 3.1. Fit of calibrated model to control SOC values

376 Modelled and measured SOC stocks in the control plot were compared to evaluate the capability of the  
377 calibrated version of Century to reproduce the dynamics of SOC stocks in the selected sites (Fig. 4.c). As  
378 shown in Fig. 4.b, the NRMSD of the control plot SOC stocks is lower than 15% for all the treatments,  
379 indicating that overall model simulations fitted the observed SOC stocks well (observed SOC stocks  
380 variance was 16.3% on average in the control plots). The correlation coefficient between modelled and  
381 observed SOC stocks in the control plots was 0.96 (Fig. 4.c). Figure 4.a provides the values of the three  
382 components of the MSD indicator for each site. It can be noticed that the LC and NU components are the  
383 highest contributors to MSD. This means that the major sources of error are the representation of the data  
384 shape and magnitude of fluctuation among the measurements. The highest NRMSD can be found in *Le*  
385 *Rheu 1* and *Le Rheu 2* (around 12% and 14% respectively). In these sites the model seems to better capture  
386 the shape of the data (low LC compared to the other sites), but it misses the representation of mean SOC  
387 stock (high SB) and data scattering (high NU) of the experimental profiles. We tested the capability of




388 Century to reproduce SOC stocks increase in the additional C input treatments (Fig. 5). Figure 5 shows the  
389 correlation between additional C inputs and SOC stock increase in the C input treatments ( $R^2 = 0.23$ ). In the  
390 same graph, we can appreciate additional C inputs simulated by Century to reach the 4p1000 target being  
391  $0.66 \pm 0.23 \text{ MgC ha}^{-1}$  per year (mean  $\pm$  standard deviation from the mean). This shows that Century is  
392 generally overestimating the effect of additional C inputs on SOC stocks increase. However, the effect of  
393 additional C inputs on observed SOC stock increase varies largely across different treatments.



394  
 395  
 396  
 397  
 398

**Figure 4:** a) Decomposed mean squared deviation (MgC ha<sup>-1</sup>)<sup>2</sup> in control plots for all sites. LC = Lack of Correlation, NU = Non-Unity slope and SB = Squared Bias. b) Normalized root squared deviation (%) in control plots for all sites c) Fit of predicted versus observed SOC stocks (MgC ha<sup>-1</sup>) in control plots for all sites ( $R^2 = 0.96$ ).



399  
400  
401  
402

Figure 5: Correlation between additional carbon inputs ( $\text{MgC ha}^{-1}$  per year) and annual SOC stock increase (%) in the carbon inputs treatments and mean  $\pm$  standard deviation of the additional carbon inputs to reach the 0.4% target in Century.

403

### 3.2. Estimates of additional carbon inputs and SOC changes

404

#### 3.2.1. Virtual C inputs to reach the 4p1000

405

Figure 6 represents the average percentage change of C inputs required to reach the 4% annual increase of

406

SOC stocks, among the whole sites. The increase of C inputs is given for each litter pool. On average, a

407

$43.15 \pm 5.05$  % (mean  $\pm$  SE across sites) increase of total annual C inputs compared to the current situation

408

in the control plot, is required to meet the 4p1000 target. In terms of absolute values, this represents an

409

additional  $0.66 \pm 0.23$   $\text{MgC ha}^{-1}$  inputs per year, i.e.  $2.35 \pm 0.21$   $\text{MgC ha}^{-1}$  total inputs per year (equivalent

410

approximately to  $4.05 \pm 0.36$   $\text{MgDM ha}^{-1}$  per year). What stands out in the graph is that, on average among

411

the studied sites, the AG structural litter pool should be more than doubled, while the other pools need only

412

to increase by about half of their initial value. In terms of absolute values, the structural AG biomass

413

(which was initially  $0.29$   $\text{MgC ha}^{-1}$  per year on average in the control treatments) would need an additional

414

$0.18$   $\text{MgC ha}^{-1}$  per year to reach the 4p1000; the metabolic AG (initially  $0.70$   $\text{MgC ha}^{-1}$  per year on

415

average) needs an additional  $0.14$   $\text{MgC ha}^{-1}$  per year; structural and metabolic BG biomass (initially  $0.65$


416 and 0.52 MgC ha<sup>-1</sup> per year) require an additional C input corresponding to 0.21 and 0.13 MgC ha<sup>-1</sup> per  
 417 year respectively.

418 Analysis of the SOC pools evolution in the runs with optimized C inputs to match the 4p1000 increase rate,  
 419 indicates that the active and slow pools increased by 0.58% and 0.61% per year respectively, while the  
 420 passive pool increased annually by 0.01% (Fig. 7). In absolute values, the slow compartment contributed  
 421 the most to the increase of SOC during the 30 years runs, as it increased by 2.7 MgC ha<sup>-1</sup> on average among  
 422 the sites (against an increase of 0.1 and 0.06 MgC ha<sup>-1</sup> in the active and passive compartments  
 423 respectively). This corresponds to a storage efficiency for the 30 years of simulation of approximately 13.7  
 424 % in the slow pool, compared to a storage efficiency of 0.5% and 0.34% in the active and in the passive  
 425 pools respectively.

426 We found a high linear correlation (R<sup>2</sup>=0.80) between observed initial SOC stocks and optimized C inputs  
 427 (Fig. 8). It is logical and expected that for low initial SOC stocks in steady state, a small increase of C  
 428 inputs is sufficient to reach the 4p1000 target. Conversely, when SOC is high at the beginning of the  
 429 experiment (e.g. *Trévarez*) much higher C inputs must be employed since our target increase rate is a  
 430 relative target. The regression line that emerges from the cross sites' relationship can be written as:


$$431 I^{4p1000} = 0.013 \cdot SOC_0^{obs} + 0.001, \quad (9)$$

432 where  $I^{4p1000}$  are the simulated C inputs needed to reach the 4p1000 target (MgC ha<sup>-1</sup> per year) and  
 433  $SOC_0^{obs}$  (MgC ha<sup>-1</sup>) is the observed initial SOC stock.




434

435 **Figure 6:** Sites average percentage change of carbon inputs needed to reach the 4p1000 (TOT), separated into  
 436 the four litter input pools. AM = aboveground metabolic, BM = belowground metabolic, AS = aboveground  
 437 structural, BS = belowground structural and TOT = total litter inputs. Error bars indicate the standard error.  
 438 N.B: Total change of carbon inputs (TOT) was calculated as the percentage change between the total amount of  
 439 carbon inputs before and after the 4p1000 optimization, averaged across all sites.



440  
441  
442  
443

Figure 7: Sites average soil organic carbon pools (ACT = active, SLOW = slow and PASS= passive) evolution (MgC ha<sup>-1</sup>) over the 30 years of simulation to reach the 4p1000 target. In the graph the mean percentage increase is given for each SOC pool.




444  
445  
446  
447

Figure 8: Correlation between initial observed SOC stocks (MgC ha<sup>-1</sup>) and modelled carbon inputs needed to reach the 4p1000 target (MgC ha<sup>-1</sup> year<sup>-1</sup>). The correlation coefficient ( $R^2$ ) is 0.80 and the regression line is  $y = 0.013 \cdot x + 0.001$ .

448

### 3.2.2. Virtual versus actual C inputs in the experimental carbon treatments

449 In Fig. 9 we compare the C inputs required to reach the 4p1000 target to the actual inputs used across the  
450 46 treatments of additional C. The additional C ( $\text{MgC ha}^{-1}$  per year) shown in the graph for all experimental  
451 treatments refers to exogenous organic amendments, plus additional C due to increased crop yields,  
452 relatively to the control plot. The most striking result emerging from the data is that modelled additional C  
453 inputs are systematically lower or similar to at least one treatment of additional C in all sites, except for  
454 *Foggia*. In *Foggia* experiment, different crop rotations were compared and no additional EOM was  
455 incorporated to the soil. Here, none of the rotations had sufficient additional C content (compared to the  
456 control wheat-only treatment), to meet the required C input level predicted by Century for a 4p1000  
457 increase rate. Overall, 86.91% of the experimental treatments used higher amounts of C inputs compared to  
458 the modelled need of additional C inputs in the same site. For the other treatments, the difference between  
459 simulated and observed additional C input was not significant. In the experimental treatments were applied  
460  $1.52 \text{ MgC ha}^{-1}$  per year on average and SOC stocks were found to be increasing by 0.25% per year relative  
461 to initial stocks. Modelled additional C input to reach a 0.4% increase was  $0.66 \text{ MgC ha}^{-1}$  per year, on  
462 average among the sites.




463

464 **Figure 9: Additional modelled carbon inputs (MgC ha<sup>-1</sup> year<sup>-1</sup>) to reach the 4p1000 (grey bars) compared to**  
 465 **additional carbon input treatments (colored bars) on each experimental site. Additional carbon inputs for field**  
 466 **trials are calculated as the sum of organic fertilizers and the delta carbon inputs from crop yields (compared to**  
 467 **the control plot). Additional carbon treatments are separated into different categories: BIO waste = biowaste**  
 468 **compost, green manure + sewage sludge and household waste, Cow Manure = cow manure and**  
 469 **farmyard manure (in *Broadbalk* and *Ultuna*), Pig Manure, Poultry Manure, Sewage Sludge, Rotations =**  
 470 **different crop rotations, Other organic amendments (OA) = straw, sawdust and peat (in *Ultuna*) and Castor**  
 471 **Meal (in *Broadbalk*). The error bars shown are the standard errors computed with the Monte Carlo method.**

472 **3.3. Carbon input requirements with temperature increase**

473 The temperature sensitivity analysis of the Century model for the 4p1000 target framework is plotted in  
474 Fig. 10. The required amount of C inputs to reach the 4p1000 target is likely to increase with increasing  
475 temperature scenarios. In particular, C inputs will have to increase on average by 54% in the AS1 scenario  
476 of +1°C and 120% in the AS5 scenario of +5°C temperature change, relative to current C inputs in the  
477 control plots. This represents an additional C inputs increase of 11% and 77% respectively, compared to the  
478 business as usual scenario with current temperature setup (CURR). What can be clearly seen in the graph is  
479 the increased amount of C inputs required in *Trévarez*, where C inputs should more than quadruplicate to  
480 reach the 4p1000 objective.



481  
482 **Figure 10: Temperature sensitivity analysis of carbon inputs increase (%) to reach the 4p1000 objective.**  
483 **CURR=business as usual simulation, AS1=RCP2.6 scenario of +1°C temperature increase, AS5=RCP8.5**  
484 **scenario of +5°C temperature change.**

485 **4 Discussion**

486 **4.1. Reliability of the Century model**

487 The Century model has been widely used to simulate SOC stocks dynamics in arable cropping systems  
488 (Bortolon et al., 2011; Cong et al., 2014; Kelly et al., 1997; Xu et al., 2011). Optimizing the



489 metabolic:structural ratio in the reference plots allowed us to initialize the C inputs compartments, since no  
490 measurement of the L:N ratio was available. This allowed us to: 1) take into account the average C quality  
491 of the litter pools in the different crops rotations and 2) estimate correctly the initial values of SOC stocks  
492 on the majority of the sites. On the other hand, this could have influenced the predicted redistribution of C  
493 in the additional C inputs required to reach the 4p1000 (Fig. 6). We suggest that taking into account the  
494 historical site-specific land use could help initialize SOC stocks without requiring any assumption  
495 regarding the M:S ratio (e.g. with historically based equilibrium scenarios as in Lugato et al. (2014)). To  
496 further improve SOC stock simulations, we optimized the  $Q_{10}$  and reference temperature parameters on the  
497 control plots, to account for the different pedo-climatic conditions of the experimental sites and enhance  
498 model predictions of SOC stocks dynamics (Craine et al., 2010; Lefèvre et al., 2014; Meyer et al., 2018;  
499 Wang et al., 2010). Although the dispersion of SOC stocks over time is not perfectly captured in the  
500 majority of the control plots (see the high LC component of the MSD in Fig. 4), the simulations of SOC  
501 dynamics were improved by the optimization of temperature related parameters and the NRMSD was  
502 found to be lower than 15% on all sites. Figure C2 shows that the optimization of temperature sensitive  
503 parameters did not affect significantly the required C input estimation for the current temperature scenario.  
504 This means that, although parameters optimization improved the simulation of SOC stocks in the control  
505 plots, the final results are not affected by it. The capability of Century to simulate SOC stocks in the  
506 simulations of additional C treatments might be a major shortcoming of modeling results. In fact, although  
507 SOC stocks were found to be increasing on average in the additional C treatments (0.25% per year with  
508 1.52 MgC ha<sup>-1</sup> yearly additional C inputs), this increase rate is lower than the 0.4% increase of SOC stocks  
509 predicted by Century with lower amounts of virtual C inputs (0.66 MgC ha<sup>-1</sup> per year). This is pointed out  
510 in Fig. 5, where we can see that predicted additional C inputs to reach the 4‰ are lower than the correlation  
511 line between additional C inputs and SOC stocks increase in field treatments. The overestimation of the C  
512 input effect on SOC stocks in Century might be related to the assumption that SOC stocks are in  
513 equilibrium with C inputs at the onset of the experiment and on the high sensitivity of the model to C  
514 inputs.

## 515 **4.2. Increasing annual SOC stocks by 4p1000**

### 516 **4.2.1. Modelled carbon inputs to reach the 4p1000**

517 Century simulations estimated that annual C inputs should increase by  $43 \pm 5\%$  (SE) on average to reach the  
518 4p1000 target on the selected experimental sites, under the condition that the additional C inputs are  
519 equally distributed among the surface and belowground, in order to maintain the same  
520 aboveground:belowground ratio as at the beginning of the experiment. Martin et al. (2021) found similar  
521 values of required additional C inputs to reach a 4p1000 target in France croplands (i.e. 42%, that is 0.88  
522 MgC ha<sup>-1</sup> per year). This is higher than the values found by Chenu et al. (2019) using default RothC 26.3  
523 parameters, who estimated a relative increase of C inputs in temperate sandy soils by 24% and in temperate  
524 clayey soils by 29%. Riggers et al. (2021) found that in 2095, a minimum increase of C inputs by 45% will

525 be required to maintain SOC stocks of German croplands at the level of 2014. However, they found that to  
526 increase SOC stocks by 4‰ per year, a much higher effort will be required. That is, C inputs in 2095 will  
527 have to increase by 213% relative to current levels.

528 In our study, not only the quantity of C but also the quality will need to change according to Century  
529 predictions. In fact, the predicted aboveground structural litter change was threefold higher than all other  
530 pools on average, representing an additional 0.18 MgC ha<sup>-1</sup> each year. A way for the farmer to increase the  
531 structural fraction of the C inputs is to compost the organic amendments that will be spread on soil surface.  
532 Increasing EOM in large quantities may not be possible everywhere. First of all, the amount of organic  
533 fertilizers is limited at regional scale. If farmers source additional EOMs elsewhere, only those EOMs that  
534 otherwise would be mineralized (e.g. burnt) and not applied to land account as sequestration. Second,  
535 farmers may be prevented from applying high amounts of EOM because of the risk of nitrate and phosphate  
536 pollution (Li et al., 2017; Piovesan et al., 2009). Moreover, producing additional animal manure implies  
537 larger GHG emissions through animal digestion and manure decomposition. Consequently, even if more  
538 manure is returned to the soil, it will not necessarily result in climate change mitigation.

#### 539 **4.2.2. Stability of the additional carbon stored**

540 Another important aspect to take into consideration is the stability of the additional C. In fact, the duration  
541 and persistence of C in the soil might be very different depending on whether or not the proportion of stable  
542 C is important. In the Century model, this translates into questioning whether the fractions of the long  
543 turnover rate pools (the slow and passive SOC pools) have increased. In our simulations, a general pattern  
544 can be detected (Fig. 6) where both passive and slow pools increased, but at very different rates (0.1‰ and  
545 6.1‰ per year respectively). The active pool increased by 5.8‰ annually, with benefits for soil fertility and  
546 hence food security. The additional C is mainly stored in the slow pool (2.7 MgC ha<sup>-1</sup> in 30 years of  
547 simulations), meaning that it will be stored in the soil for around 20 to 30 years. The increase in C inputs  
548 must be sustained to increase SOC stocks at the desired rate, until a new equilibrium will be reached. To  
549 further increase SOC stocks after the new equilibrium, new strategies of additional C could be implemented  
550 later on. For instance, this could be achieved through the implementation of complementary management  
551 options to those considered in the long-term experiments described here, such as residue management,  
552 cover crops, conservation agriculture and agroforestry systems (Chenu et al., 2019; Lal, 1997; Smith et al.,  
553 1997).

#### 554 **4.2.3. Simulated carbon inputs and experimental carbon addition treatments**

555 Different types of organic C treatments were considered in this study and compared to Century simulations  
556 of C inputs required to reach the 4p1000. In all experimental sites with additional EOM inputs, at least one  
557 treatment employed higher amounts of C inputs compared to the simulated C inputs required for a 4‰  
558 annual target. In *Foggia*, C inputs from different crop rotations were studied, but none employed sufficient  
559 amounts of additional C to reach the 4p1000, as predicted by Century. Model results in *Foggia* had a high

560 standard error, mainly due to the fact that the variability of crop yields for this site was not available. Thus,  
561 for this site, we calculated model uncertainty using the average relative variability across the whole dataset,  
562 which could have increased the uncertainty of model outputs.

563 It is important to note that the amount of C inputs simulated by Century was constrained to have the same  
564 AG:BG ratio as at the beginning of the experiment. This means that the additional C inputs should be  
565 distributed equally on soil surface and belowground, not to change the initial allocation of C in the litter  
566 pools. Since all field treatments were performed under conventional tillage, the comparison between  
567 modelled and observed additional C inputs under this constraint holds well.

568 The annual SOC stocks variation (0.25%) estimated in the experimental C treatments across the 14 sites,  
569 indicates that Century might be overestimating the effect of additional C inputs on SOC stocks. In  
570 particular, only 18 out of 46 field treatments (with average additional C inputs of 1.93 MgC ha<sup>-1</sup> per year)  
571 were found to be actually increasing SOC stocks at a higher rate than 4‰ per year, relatively to their initial  
572 SOC stocks. This is similar to the values found by Poulton et al. (2018), who estimated that adding similar  
573 high amounts of C inputs increased SOC stocks at an annual rate higher than 4‰ in 16 long-term  
574 agricultural experiments. Thus, Century seems to be over-predicting the effect of adding C inputs in the  
575 virtual simulations. The overestimation of the Century model might be due to several factors. First of all,  
576 the C inputs prescribed to model simulations were constant through time, while C inputs from plant  
577 material actually vary annually and over the years because of agronomical and climatic factors. Historical  
578 land use and management practices such as tillage were not taken into account, although they affect SOC  
579 stocks (Pellerin et al. 2017). Another factor that the model is not taking into account is N and other  
580 nutrients availability, which might affect the SOC stocks dynamics. This is especially true for treatments  
581 with different frequencies of application (e.g. *Arazuri*), where nutrients depletion is likely to be more  
582 evident when the application is sparser. The method used to estimate C inputs (i.e. the allometric functions  
583 from Bolinder et al. (2007) in our case) also influences the simulation of SOC stocks (Clivot et al., 2019).  
584 However, estimating the increase of C inputs relative to their initial value has likely cancelled out  
585 uncertainties related to the C inputs estimation method in our analysis.

#### 586 **4.2.4. Organic carbon inputs use in Europe**

587 Zhang et al. (2017) estimated that the amount of N inputs from livestock manure applied to European  
588 croplands was 3.9 Tg N in 2014, for a cropland area of 127 Mha in 2015 (Goldewijk et al. 2017). Cattle  
589 manure, which represents the highest proportion of manure produced and applied to croplands, has average  
590 C:N ratio ranging between 10 and 30 (multiple sources from Fuchs et al. (2014) and Pellerin et al. (2017)).  
591 With these data, we can roughly estimate the application of C manure from livestock in European  
592 agricultural soils as ranging between 0.30 and 0.92 MgC ha<sup>-1</sup> each year. Most of the experiments used in  
593 this study used higher amounts of C input (1.52 MgC ha<sup>-1</sup> per year on average). However, the C inputs  
594 requirement predicted by Century, which ranged between 0.24±0.02 and 1.20±1.00 MgC ha<sup>-1</sup> per year,  
595 plus one site with 1.45±0.16 MgC ha<sup>-1</sup> per year, is in line with the average use of livestock manure in

596 Europe. In terms of C sequestration, organic fertilizers coming from animal manure are usually being  
597 applied to the soil at some location, hence they cannot account for additional climate mitigation potential  
598 (Poulton et al., 2018). Rather, they are considered as a business as usual situation that can unlikely be  
599 significantly expanded. However, according to Zhang et al. (2017) estimation, there is room for  
600 improvement since the fraction of livestock manure applied to cropland in the 2010s was approximately  
601 26% of total livestock production in Europe. The estimates from Zhang et al. (2017) refer to livestock  
602 manure only. In our study, we also considered treatments with other types of EOM addition, such as  
603 sewage sludge and household waste. In many countries, a significant proportion of food and urban waste is  
604 currently left on disposal areas, where C is lost to the atmosphere as CO<sub>2</sub> or methane (CH<sub>4</sub>) emissions  
605 (Bijaya et al. 2006). Pellegrini et al. (2016) reported the amounts of sewage sludge disposed on landfill in  
606 Europe (EU26) from Eurostat (2014b). In 2010, this was 0.914 TgDM. Using the Van Bemmelen factor  
607 (1.724) to convert OM to OC (McBratney and Minasny, 2010; Rovira et al., 2015), we estimated that the  
608 sewage sludge disposed on landfill in Europe was around 0.004 MgC ha<sup>-1</sup> per year in 2010. If applied to  
609 cropland, this could potentially increase C inputs to the soil and decrease GHG emissions associated to  
610 landfilled waste. However, in some countries social acceptability of spreading EOM such as sewage sludge  
611 is very low, limiting its actual potential. In Europe, landfilled municipal waste was 0.3 MgC ha<sup>-1</sup> in 2019  
612 (estimated from Eurostat (2020) considering a C content in household waste of 71% (Larsen et al., 2013)).  
613 This is higher than the amount of municipal waste currently composted in Europe (i.e. 0.22 MgC ha<sup>-1</sup> in  
614 2019, according to Eurostat (2020)), showing that additional efforts to improve the reutilization of  
615 municipal waste could help to increase C inputs in agriculture. A contribution to the sequestration of C  
616 from the atmosphere could also come from changing the treatment methods which affect the quality of C in  
617 crop residues and manure, so that their turnover time decreases, e.g. through fermentation or biochar.  
618 However, a full C cycle assessment should be considered to make sure that GHG emissions associated to  
619 such treatments do not exceed additional C storage (Guenet et al., 2020). In general, improving the use  
620 efficiency of EOM to the soil by managing it differently could contribute to some extent to climate change  
621 mitigation, increase soil quality, and reduce mineral fertilizers use (Chadwick et al. 2015). In this study, we  
622 did not include other potentially beneficial management practices, such as cover crops, reduced tillage,  
623 biochar application, improved soil pH, landscape differentiation and mineral amendments. Further research  
624 should investigate if long-term experiments with these management practices would be able to increase  
625 SOC stocks by 4p1000, following Century predictions.

#### 626 **4.2.5. Reaching a 4p1000 target: only a matter of initial SOC stocks?**

627 As we expected, the estimated amount of C inputs to reach the 4p1000 target was linearly correlated to the  
628 initial observed level of SOC stocks (Fig. 7). This result means that site differences in Q<sub>10</sub> and  
629 decomposition rates are less influential than initial SOC in determining the optimal input increase to reach  
630 the 4‰ per year target. The linearity between C inputs and initial SOC stocks is primarily due to the linear  
631 structure of the Century model. In fact, if we consider the stationary solution for which Eq. (2) is equal to 0,

632 SOC depends linearly on the carbon inputs. Therefore, the opposite is also true (i.e. carbon inputs are  
633 linearly dependent to the initial amount of SOC stocks). Moreover, the 4p1000 target itself is defined as the  
634 increase of SOC by 0.4% per year, relatively to its initial value (Minasny et al., 2017). Hence, it implies a  
635 proportional contribution that depends on the initial SOC stocks. Wiesmeier et al. (2016) also observed a  
636 linear relationship between SOC increase and C inputs. This linear relationship means that soils with high  
637 SOC stocks will have to increase their carbon stocks more in absolute terms to meet this quantitative target.  
638 On the other side, smaller amounts of C will have to be employed in sites with low levels of SOC stocks, to  
639 reach a 4p1000 target. However, increasing C inputs where SOC stocks are low might require substantial  
640 changes in the agricultural systems and such quantity of additional OM might not be available at a large  
641 scale. A counterpoint is also that the largest contribution of C sequestration will come from soils with  
642 medium or high SOC stocks (i.e. higher than 50 MgC ha<sup>-1</sup>, such as grasslands and forests). In these soils,  
643 the required additional C inputs will have to be higher according to Century, raising concern on a  
644 compensation of CO<sub>2</sub> emissions through improved SOC stocks at a global scale. This result depends on the  
645 quality of the simulated carbon inputs (i.e. the predicted metabolic:structural ratio) and does not take into  
646 account any notion of soil saturation. Before applying this trend to calculate the required C inputs from  
647 current SOC stocks, we should extend the database to cover different pedo-climatic regions and different  
648 ecosystems of the world. Moreover, inaccuracies in simulations outcomes, such as those found in this  
649 study, need to be reduced. As discussed in subsection 4.2.3, a better representation of C inputs dynamics  
650 and management practices could improve the simulation of SOC stocks.  
651 We suggest to consider multi-model analysis for this type of work in the future (Farina et al., 2021), to  
652 acknowledge different representations of SOC and reduce the effect of single models' uncertainties.  
653 Furthermore, the likely increase of SOC mineralization due to future climate change (Wiesmeier et al.,  
654 2016) needs to be taken into account.

### 655 **4.3. Sensitivity analysis**

656 The predicted need of additional C inputs to reach the 4p1000 target is likely to be higher with future global  
657 warming, as a consequence of modified SOC decomposition rates. Considering the crucial role of soil as a  
658 land-use based option for mitigating climate change, recent studies have shown a growing interest in  
659 temperature sensitivity of SOC stocks decomposition (Dash et al., 2019; Koven et al., 2011; Parihar et al.,  
660 2019; Wiesmeier et al., 2016). We know that the decomposition rate of SOM is affected – generally  
661 increased – with increasing temperatures. However, the magnitude of expected feedbacks is still  
662 surrounded by controversy. In particular, this is mainly due to the diversity of organic compounds in the  
663 soil that are known to have inherent sensitivities to temperature (Davidson and Janssens, 2006). In fact, a  
664 diversity of responses of decomposition rates to future climates can be expected, including increases due to  
665 higher temperature as well as decreases due to water limitation. In this context, the study of the Century  
666 model response to predicted scenarios of temperature increase is of primary importance. We mimicked the  
667 most optimistic (+1°C) and pessimistic (+5°C) RCPs scenarios of the 5th IPCC assessment report.

668 Although these scenarios are calculated over ~100 years, we used these values over a 30 years simulation to  
669 assess the sensitivity of Century to temperature increase. What is striking from our results is that with  
670 increasing temperatures all sites will have to provide considerably higher amounts of C inputs to reach the  
671 4p1000 target (Fig. 9). In particular, the C inputs change needs to more than double in all sites, according to  
672 the worst-case scenario of +5°C. It is important to point out that the optimization of the  $Q_{10}$  and reference  
673 temperature parameters are likely to influence the outcomes of the simulated SOC stocks and therefore the  
674 C inputs need. Nevertheless, comparing the carbon input change simulated with the optimized version of  
675 Century (Fig. 9) to that simulated with the default parameters setting (Fig. C1), shows that the predicted C  
676 inputs change follows the same pattern, even though the intensity of the increase is considerably higher in  
677 the optimized version. These results can be understood in two ways. Either the optimized version of  
678 Century is overestimating the effect of temperature on SOC stocks decomposition, or SOC stocks  
679 decomposition patterns are likely to increase even more intensively when considering the entire range of  
680 possible  $Q_{10}$  values. In either case, further research is needed to reduce the uncertainty around the impact of  
681 climate change on SOC decomposition. Studies should also examine moisture change, which we did not  
682 take into account here. This is likely to be impacted as a consequence of modified precipitations and  
683 temperature (IPCC, 2015), with consequences on root respiration and microbial decomposition (Davidson  
684 and Janssens, 2006). Additionally, increased temperature and CO<sub>2</sub> concentration in the atmosphere, as well  
685 as changes in precipitations are likely to influence net primary production and therefore C inputs to the soil.  
686 All these feedbacks are important and must be taken into account for a comprehensive evaluation of C  
687 cycle effects on climate change.

## 688 5 Conclusion

689 The Century model predicted an average increase of annual C inputs by  $43 \pm 5\%$  to reach a 4p1000 target  
690 over a range of 14 agricultural sites across Europe, with diverse soil types, climates, crop rotations and  
691 practices. The required simulated amount of additional C inputs was found to be systematically lower or  
692 similar to the 46 treatments of C inputs carried out in these sites. However, Century might have  
693 overestimated the predicted effect of additional C inputs on the SOC stocks variation rate, as the only field  
694 treatments that were found increasing SOC stocks by at least 4‰ annually were those using very high  
695 amounts of C inputs (~1.93 MgC ha<sup>-1</sup> per year). The predicted amount of additional C inputs depended  
696 linearly on the initial amount of observed SOC stocks in the control experiments, indicating that lower  
697 amounts of C inputs might be sufficient to reach the 4p1000 target where SOC stocks are low. However,  
698 increasing C inputs might require substantial changes in the agricultural systems and high quantities of  
699 additional organic matter might not be available at a large scale. Furthermore, the required amount of  
700 additional C inputs was found to increase substantially with future scenarios of changes in temperature,  
701 raising concern about the feasibility of a 4p1000 target under climate change and beyond that, the  
702 feasibility of SOC stock preservation. The magnitude of SOC storage potential in agricultural soils depends  
703 largely on site-specific conditions, such as climate, soil type and land use. In this study, we did not take into

704 account the whole life cycle of C at the farm. However, compensating CO<sub>2</sub> emissions from human  
705 activities through SOC sequestration should also comprehend GHG emissions related to the management of  
706 additional EOM. In this study, we considered only temperate, sub-humid and Mediterranean climates. A  
707 broader evaluation of the required C inputs and associated agricultural practices to increase SOC stocks  
708 should be carried out at larger scales. Causes of biases in model simulations should be addressed in future  
709 studies and the representation of C inputs should be improved. We also suggest that future research should  
710 include multiple models, to reduce the influence of extreme model outcomes on the representation of SOC  
711 stocks.

## 712 **Authors contribution**

713 YH provided the initial model code. EB edited and developed the model code, performed the simulations  
714 and prepared the manuscript with contributions from all co-authors. EB, CC, PC and BG designed the  
715 study. HC, IV, RF, TK and MM provided the data. All co-authors participated to the results analysis and  
716 the writing.

## 717 **Competing interests**

718 The authors declare that they have no conflict of interest.

## 719 **Acknowledgements**

720 This work benefited from the French state aid managed by the ANR under the "Investissements d'avenir"  
721 programme with the reference ANR-16-CONV-0003 (CLAND project). We acknowledge Mancomunidad  
722 de la Comarca de Pamplona for maintenance and access to Arazuri site data. Research grant RTA2017-  
723 00088-C03-01 from the Instituto Nacional de Investigación Agraria y Alimentaria, INIA (Spanish Agency).  
724 We acknowledge Margaret Glendining, curator of the electronic Rothamsted Archive (e-RA) for providing  
725 the Broadbalk data. The Colmar and Feucherolles field experiments form part of the SOERE-PRO  
726 (network of long-term experiments dedicated to the study of impacts of organic waste product recycling)  
727 certified by ALLENI (Alliance Nationale de Recherche pour l'Environnement) and integrated as a service  
728 of the 'Investment for future' infrastructure AnaEE-France, overseen by the French National Research  
729 Agency (ANR-11-INBS-0001).

## 730 **Appendix A – Century model description and environmental functions used**

731 The temporal evolution of soil organic carbon is described in the Century model as a first order differential  
732 matrix equation:

$$733 \frac{dSOC(t)}{dt} = I + A \cdot \xi_{TWLCl}(t) \cdot K \cdot SOC(t), \quad (2)$$

734 where  $\mathbf{SOC}(t)$  is the vector describing the SOC state variables. The first term on the right side of the  
 735 equation represents carbon inputs to the soil coming from plant residues and organic material. Carbon  
 736 inputs are allocated into four different litter pools. Hence,  $\mathbf{I}$  is a  $1 \times 7$  matrix with four nonzero elements.  
 737 The second term of the equation represents carbon outputs from the soil, following a first order decay  
 738 kinetics.  $\mathbf{A}$  is a  $7 \times 7$  carbon transfer matrix that quantifies the transfers of carbon among the different pools.  
 739 The diagonal entries of  $\mathbf{A}$  are equal to -1, denoting the entire decomposition flux that leaves each carbon  
 740 pool. The non-diagonal elements represent the fraction of carbon that is transferred from one pool to  
 741 another.  $\mathbf{K}$  is a  $7 \times 7$  diagonal matrix with the diagonal elements representing the potential decomposition  
 742 rate of each carbon pool.  $\xi_{TWLCI}(t)$  is the environmental scalar matrix, a  $7 \times 7$  diagonal matrix with each  
 743 diagonal element denoting temperature ( $f_T(t)$ ), water ( $f_W(t)$ ) lignin ( $f_{L_i}$ ) and clay ( $f_{Clay_i}$ ) scalars, which  
 744 modify the potential decomposition rate. Temperature response function  $f_T(t)$  is described by Eq. (4), the  
 745 others are expressed as follows. The moisture function  $f_W(t)$  is a polynomial function ranging from 0.25  
 746 and 1 and taking the form of:

$$747 \quad f_W(t) = -1.1 \cdot w^2 + 2.4 \cdot w - 0.29, \quad (A1)$$

748 where  $w$  is the daily relative humidity coefficient, which varies between 0 and 1 and was calculated from  
 749 soil moisture ( $m^3_{water} m^{-3}_{soil}$ ), using the following function from (Krinner et al., 2005):

$$750 \quad w = \sum_{texture} \frac{conc_{texture} \cdot moisture - WP_{texture}}{FC_{texture} - WP_{texture}},$$

751 where  $w$  is the estimated relative humidity, ranging between 0 and 1;  $texture =$  sand, silt and clay;  
 752  $conc_{texture}$  is the concentration of the different textures;  $moisture$  is soil moisture ( $m^3_{water} m^{-3}_{soil}$ ),  
 753  $WP_{texture}$  is the wilting point of the different textures (equivalent to 0.0657, 0.0884, 0.1496 for sand, silt  
 754 and clay respectively) and  $FC_{texture}$  is the field capacity of texture (equivalent to 0.1218, 0.1654,  
 755 0.2697 for sand, silt and clay respectively).

756 The decomposition rate of structural litter pools is affected by their lignin content:

$$757 \quad f_{L_i} = e^{-lgc \cdot L}, \quad (A2)$$

758 where  $lgc$  is the coefficient that regulates the lignin effect, while  $L$  is the lignin structural fraction of the  
 759 aboveground and the belowground litter pools.

760 Finally, the fraction of clay in the soil ( $g \text{ clay } g^{-1} \text{ soil}$ ) influences the decomposition rate of the active  
 761 pool:

$$762 \quad f_{Clay_i} = 1 - 0.75 \cdot clay. \quad (A3)$$

## 763 **Appendix B – Model evaluation**

764 Two residual-based metrics were used to evaluate the goodness-of-fit of modelled and observed SOC  
 765 stocks for each site: the Mean Squared Deviation (MSD) and the Normalized Root Mean Squared  
 766 Deviation (NRMSD). The MSD for each site is defined as:

$$767 \quad MSD = \frac{\sum_{i=1}^n (m_i - o_i)^2}{s}, \quad (B1)$$



768 where  $i = 1, \dots, n$  is the year of the experiment,  $m_i$  and  $o_i$  are respectively modelled and observed values of  
 769 SOC stocks and  $s$  is the number of observations in the experiment. Following Gauch et al. (2003), the MSD  
 770 can be decomposed into three components: the Squared Bias (SB), the Non-Unity slope (NU) and the Lack  
 771 of Correlation (LC). SB is calculated as:

$$772 \quad SB = (\bar{m} - \bar{o})^2, \quad (B2)$$

773 where  $\bar{m}$  and  $\bar{o}$  are the mean values of modelled and observed SOC stocks respectively.

774 Calling  $\Delta M_i = (\bar{m} - m_i)$  and  $\Delta O_i = (\bar{o} - o_i)$  we have:

$$775 \quad NU = \left(1 - \frac{\sum_{i=1}^n \Delta M_i \Delta O_i}{\sum_{i=1}^n \Delta M_i^2}\right)^2 \cdot \frac{\sum_{i=1}^n \Delta M_i^2}{s}, \quad (B3)$$

$$776 \quad LC = \left(1 - \frac{\sum_{i=1}^n (\Delta M_i \Delta O_i)^2}{\sum_{i=1}^n \Delta O_i^2 \cdot \sum_{i=1}^n \Delta M_i^2}\right) \cdot \frac{\sum_{i=1}^n \Delta O_i^2}{s}. \quad (B4)$$


777 These three components add up to MSD and help locating the causes of error of model predictions,  
 778 determining areas in the model that require further improvement (Bellocchi et al., 2010). In particular, SB  
 779 provides information about the mean bias of the simulation from measurements, NU indicates the capacity  
 780 of the model to correctly reproduce the magnitude of the fluctuation among the measurements and LC is an  
 781 indication of the dispersion of the points over a scatterplot, i.e. the capacity of the model to reproduce the  
 782 shape of the data (Kobayashi and Salam, 2000).

783 The second statistical measure we used was computed as the squared root of the MSD, normalized by the  
 784 mean observed SOC stocks:

$$785 \quad NRMSD = \frac{\sqrt{MSD}}{\bar{o}} \cdot 100. \quad (B5)$$


786 This indicator is expressed as a percentage and allows to evaluate the model performance independently to  
 787 the units of SOC stocks.

788 **Appendix C – Sensitivity analysis with default Century parameters**



789

790 **Figure C1: Temperature sensitivity analysis of carbon inputs change (%) to reach the 4p1000 objective, using**  
 791 **Century default Q10 and reference temperature parameters. CURR=business as usual simulation, AS1=RCP2.6**  
 792 **scenario of +1°C temperature increase, AS5=RCP8.5 scenario of +5°C temperature change.**



793

794 **Figure C2: Effect of the optimization of the  $Q_{10}$  and reference temperature ( $T_{ref}$ ) parameters on the additional**  
 795 **carbon inputs to reach the 4p1000 predicted by Century (mean  $\pm$  standard deviation).**

796

## References

- 797 Anderson, G. M.: Error propagation by the Monte Carlo method in geochemical calculations, *Geochimica*  
 798 *et Cosmochimica Acta*, 40(12), 1533–1538, doi:10.1016/0016-7037(76)90092-2, 1976.
- 799 Baveye, P. C., Berthelin, J., Tessier, D. and Lemaire, G.: The “4 per 1000” initiative: A credibility issue for  
 800 the soil science community?, *Geoderma*, 309, 118–123, doi:10.1016/j.geoderma.2017.05.005,  
 801 2018.
- 802 Bellocchi, G., Rivington, M., Donatelli, M. and Matthews, K.: Validation of biophysical models: issues and  
 803 methodologies. A review, *Agron. Sustain. Dev.*, 30(1), 109–130, doi:10.1051/agro/2009001, 2010.
- 804 Bijaya, M.: Predicted growth of world urban food waste and methane production, *Waste Management &*  
 805 *Research: The Journal for a Sustainable Circular Economy (WM&R)*, 24(5), 421–433,  
 806 doi:doi.org/10.1177/0734242X06067767, 2006.
- 807 Bolinder, M. A., Janzen, H. H., Gregorich, E. G., Angers, D. A. and VandenBygaart, A. J.: An approach for  
 808 estimating net primary productivity and annual carbon inputs to soil for common agricultural crops  
 809 in Canada, *Agriculture, Ecosystems & Environment*, 118(1–4), 29–42,  
 810 doi:10.1016/j.agee.2006.05.013, 2007.
- 811 Bortolon, E. S. O., Mielniczuk, J., Tornquist, C. G., Lopes, F. and Bergamaschi, H.: Validation of the  
 812 Century model to estimate the impact of agriculture on soil organic carbon in Southern Brazil,  
 813 *Geoderma*, 167–168, 156–166, doi:10.1016/j.geoderma.2011.08.008, 2011.
- 814 Bouthier, A., Duparque, A., Mary, B., Sagot, S., Trochard, R., Levert, M., Houot, S., Damay, N., Denoroy,  
 815 P., Dinh, J.-L., Blin, B., and Ganteil, F.: Adaptation et mise en œuvre du modèle de calcul de bilan  
 816 humique à long terme AMG dans une large gamme de systèmes de grandes cultures et de  
 817 polyculture-élevage, 34, 125–139, 2014.
- 818 Campbell, E. E. and Paustian, K.: Current developments in soil organic matter modeling and the expansion  
 819 of model applications: a review, *Environ. Res. Lett.*, 10(12), 123004, doi:10.1088/1748-  
 820 9326/10/12/123004, 2015.
- 821 Chadwick, Q.: Improving manure nutrient management towards sustainable agricultural intensification in  
 822 China, *Agriculture, Ecosystems and Environment*, 209, 34–46,  
 823 doi:doi.org/10.1016/j.agee.2015.03.025, 2015.

824 Chenu, C., Angers, D. A., Barré, P., Derrien, D., Arrouays, D. and Balesdent, J.: Increasing organic stocks  
825 in agricultural soils: Knowledge gaps and potential innovations, *Soil and Tillage Research*, 188,  
826 41–52, doi:10.1016/j.still.2018.04.011, 2019.

827 Clivot, H., Mouny, J.-C., Duparque, A., Dinh, J.-L., Denoroy, P., Houot, S., Vertès, F., Trochard, R.,  
828 Bouthier, A., Sagot, S. and Mary, B.: Modeling soil organic carbon evolution in long-term arable  
829 experiments with AMG model, *Environmental Modelling & Software*, 118, 99–113,  
830 doi:10.1016/j.envsoft.2019.04.004, 2019.

831 Cong, R., Wang, X., Xu, M., Ogle, S. M. and Parton, W. J.: Evaluation of the CENTURY Model Using  
832 Long-Term Fertilization Trials under Corn-Wheat Cropping Systems in the Typical Croplands of  
833 China, edited by J. Vera, *PLoS ONE*, 9(4), e95142, doi:10.1371/journal.pone.0095142, 2014.

834 Craine, J., Spurr, R., McLauchlan, K. and Fierer, N.: Landscape-level variation in temperature sensitivity of  
835 soil organic carbon decomposition, *Soil Biology and Biochemistry*, 42(2), 373–375,  
836 doi:10.1016/j.soilbio.2009.10.024, 2010.

837 Dash, P. K., Bhattacharyya, P., Roy, K. S., Neogi, S. and Nayak, A. K.: Environmental constraints'  
838 sensitivity of soil organic carbon decomposition to temperature, management practices and climate  
839 change, *Ecological Indicators*, 107, 105644, doi:10.1016/j.ecolind.2019.105644, 2019.

840 Davidson, E. A. and Janssens, I. A.: Temperature sensitivity of soil carbon decomposition and feedbacks to  
841 climate change, *Nature*, 440(7081), 165–173, doi:10.1038/nature04514, 2006.

842 Eurostat: [online] Available from: <http://ec.europa.eu/eurostat/web/products-datasets/-/ten00030> (accessed  
843 September 2015), 2014b.

844 Eurostat: Municipal waste landfilled, incinerated, recycled and composted, EU-27, 1995-2018, [online]  
845 Available from: [https://ec.europa.eu/eurostat/statistics-  
846 explained/index.php?title=Municipal\\_waste\\_statistics](https://ec.europa.eu/eurostat/statistics-explained/index.php?title=Municipal_waste_statistics), 2020.

847 Fan, J., McConkey, B., Wang, H. and Janzen, H.: Root distribution by depth for temperate agricultural  
848 crops, *Field Crops Research*, 189, 68–74, doi:10.1016/j.fcr.2016.02.013, 2016.

849 Farina, R., Marchetti, A., Francaviglia, R., Napoli, R. and Bene, C. D.: Modeling regional soil C stocks and  
850 CO<sub>2</sub> emissions under Mediterranean cropping systems and soil types, *Agriculture, Ecosystems &  
851 Environment*, 238, 128–141, doi:10.1016/j.agee.2016.08.015, 2017.

852 Farina, R., Sándor, R., Abdalla, M., Álvaro-Fuentes, J., Bechini, L., Bolinder, M. A., Brilli, L., Chenu, C.,  
853 Clivot, H., De Antoni Migliorati, M., Bene, C. D., Dorich, C. D., Ehrhardt, F., Ferchaud, F.,  
854 Fitton, N., Francaviglia, R., Franko, U., Giltrap, D. L., Grant, B. B., Guenet, B., Harrison, M. T.,  
855 Kirschbaum, M. U. F., Kuka, K., Kulmala, L., Liski, J., McGrath, M. J., Meier, E., Menichetti, L.,  
856 Moyano, F., Nendel, C., Recous, S., Reibold, N., Shepherd, A., Smith, W. N., Smith, P., Soussana,  
857 J.-F., Stella, T., Taghizadeh-Toosi, A., Tsutsikh, E., and Bellocchi, G.: Ensemble modelling,  
858 uncertainty and robust predictions of organic carbon in long-term bare-fallow soils, 27, 904–928,  
859 [https://doi.org/DOI: 10.1111/gcb.15441](https://doi.org/DOI:10.1111/gcb.15441), 2021.

860 Fu, Z., Liu, G. and Guo, L.: Sequential Quadratic Programming Method for Nonlinear Least Squares  
861 Estimation and Its Application, *Mathematical Problems in Engineering*, 2019, 1–8,  
862 doi:10.1155/2019/3087949, 2019.

863 Fuchs, J., Générmont, S., Houot, S., Jardé, E., Ménasseri, S., Mollier, A., Morel, C., Parnaudeau, V.,  
864 Pradel, M. and Vieublé, L.: Effets agronomiques attendus de l'épandage des Mafor sur les  
865 écosystèmes agricoles et forestiers, 204, 2014.

866 Gale, M. R. and Grigal, D. F.: Vertical root distributions of northern tree species in relation to successional  
867 status, *Can. J. For. Res.*, 17(8), 829–834, doi:10.1139/x87-131, 1987.

868 Gauch, H. G., Hwang, J. T. G. and Fick, G. W.: Model Evaluation by Comparison of Model-Based  
869 Predictions and Measured Values, *Agron. J.*, 95(6), 1442–1446, doi:10.2134/agronj2003.1442,  
870 2003.

871 Goidts, E. and van Wesemael, B.: Regional assessment of soil organic carbon changes under agriculture in  
872 Southern Belgium (1955–2005), *Geoderma*, 141(3–4), 341–354,  
873 doi:10.1016/j.geoderma.2007.06.013, 2007.

874 Goldewijk, K., Beusen, A., Doelman, J. and Stehfest, E.: Anthropogenic land use estimates for the  
875 Holocene – HYDE 3.2, *Earth Syst. Sci. Data*, 9(2), 927–953, doi:10.5194/essd-9-927-2017, 2017.

876 van Groenigen, J. W., van Kessel, C., Hungate, B. A., Oenema, O., Powlson, D. S. and van Groenigen, K.  
877 J.: Sequestering Soil Organic Carbon: A Nitrogen Dilemma, *Environ. Sci. Technol.*, 51(9), 4738–  
878 4739, doi:10.1021/acs.est.7b01427, 2017.

879 Guenet, B., Gabrielle, B., Chenu, C., Arrouays, D., Balesdent, J., Bernoux, M., Bruni, E., Caliman, J.,  
880 Cardinael, R., Chen, S., Ciais, P., Desbois, D., Fouche, J., Frank, S., Henault, C., Lugato, E.,  
881 Naipal, V., Nesme, T., Obersteiner, M., Pellerin, S., Powlson, D. S., Rasse, D. P., Rees, F.,  
882 Soussana, J., Su, Y., Tian, H., Valin, H. and Zhou, F.: Can N<sub>2</sub>O emissions offset the benefits  
883 from soil organic carbon storage?, *Glob. Change Biol.*, gcb.15342, doi:10.1111/gcb.15342, 2020.  
884 Huang, Y., Lu, X., Shi, Z., Lawrence, D., Koven, C. D., Xia, J., Du, Z., Kluzek, E. and Luo, Y.: Matrix  
885 approach to land carbon cycle modeling: A case study with the Community Land Model, *Glob  
886 Change Biol*, 24(3), 1394–1404, doi:10.1111/gcb.13948, 2018.  
887 IPCC, 2014: Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the  
888 Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team,  
889 R.K. Pachauri and L.A. Meyer (eds.)]. IPCC, Geneva, Switzerland, 151, 2015  
890 Kätterer, T., Bolinder, M. A., Andrén, O., Kirchmann, H. and Menichetti, L.: Roots contribute more to  
891 refractory soil organic matter than above-ground crop residues, as revealed by a long-term field  
892 experiment, *Agriculture, Ecosystems & Environment*, 141(1–2), 184–192,  
893 doi:10.1016/j.agee.2011.02.029, 2011.  
894 Kelly, R. H., Parton, W. J., Crocker, G. J., Graced, P. R., Klir, J., Körschens, M., Poulton, P. R. and  
895 Richter, D. D.: Simulating trends in soil organic carbon in long-term experiments using the  
896 century model, *Geoderma*, 81(1–2), 75–90, doi:10.1016/S0016-7061(97)00082-7, 1997.  
897 Kobayashi, K. and Salam, M. U.: Comparing Simulated and Measured Values Using Mean Squared  
898 Deviation and its Components, *AGRONOMY JOURNAL*, 92, 9, 2000.  
899 Koven, C. D., Ringeval, B., Friedlingstein, P., Ciais, P., Cadule, P., Khvorostyanov, D., Krinner, G. and  
900 Tarnocai, C.: Permafrost carbon-climate feedbacks accelerate global warming, *Proceedings of the  
901 National Academy of Sciences*, 108(36), 14769–14774, doi:10.1073/pnas.1103910108, 2011.  
902 Krinner, G., Viovy, N., de Noblet-Ducoudré, N., Ogée, J., Polcher, J., Friedlingstein, P., Ciais, P., Sitch, S.  
903 and Prentice, I. C.: A dynamic global vegetation model for studies of the coupled atmosphere-  
904 biosphere system: DVGM FOR COUPLED CLIMATE STUDIES, *Global Biogeochem. Cycles*,  
905 19(1), doi:10.1029/2003GB002199, 2005.  
906 Lal, R.: Residue management, conservation tillage and soil restoration for mitigating greenhouse effect by  
907 CO<sub>2</sub>-enrichment, *Soil and Tillage Research*, 43(1–2), 81–107, doi:10.1016/S0167-  
908 1987(97)00036-6, 1997.  
909 Lal, R.: Carbon sequestration, *Phil. Trans. R. Soc. B*, 363(1492), 815–830, doi:10.1098/rstb.2007.2185,  
910 2008.  
911 Lal, R.: Digging deeper: A holistic perspective of factors affecting soil organic carbon sequestration in  
912 agroecosystems, *Glob Change Biol*, 24, 3285–3301, <https://doi.org/10.1111/gcb.14054>, 2018.  
913 Larsen, A. W., Fuglsang, K., Pedersen, N. H., Fellner, J., Rechberger, H. and Astrup, T.: Biogenic carbon  
914 in combustible waste: Waste composition, variability and measurement uncertainty, *Waste Manag  
915 Res*, 31(10\_suppl), 56–66, doi:10.1177/0734242X13502387, 2013.  
916 Lefèvre, R., Barré, P., Moyano, F. E., Christensen, B. T., Bardoux, G., Eglin, T., Girardin, C., Houot, S.,  
917 Kätterer, T., van Oort, F. and Chenu, C.: Higher temperature sensitivity for stable than for labile  
918 soil organic carbon - Evidence from incubations of long-term bare fallow soils, *Glob Change Biol*,  
919 20(2), 633–640, doi:10.1111/gcb.12402, 2014.  
920 Levvasseur, F., Mary, B., Christensen, B. T., Duparque, A., Ferchaud, F., Kätterer, T., Lagrange, H.,  
921 Montenach, D., Resseguier, C., and Houot, S.: The simple AMG model accurately simulates  
922 organic carbon storage in soils after repeated application of exogenous organic matter, *Nutr Cycl  
923 Agroecosyst*, 117, 215–229, <https://doi.org/10.1007/s10705-020-10065-x>, 2020.  
924 Li, S., Li, J., Zhang, B., Li, D., Li, G. and Li, Y.: Effect of different organic fertilizers application on  
925 growth and environmental risk of nitrate under a vegetable field, *Sci Rep*, 7(1), 17020,  
926 doi:10.1038/s41598-017-17219-y, 2017.  
927 Lovelli, S., Scopa, A., Perniola, M., Di Tommaso, T. and Sofo, A.: Abscisic acid root and leaf  
928 concentration in relation to biomass partitioning in salinized tomato plants, *Journal of Plant  
929 Physiology*, 169(3), 226–233, doi:10.1016/j.jplph.2011.09.009, 2012.  
930 Lugato, E., Bampa, F., Panagos, P., Montanarella, L. and Jones, A.: Potential carbon sequestration of  
931 European arable soils estimated by modelling a comprehensive set of management practices, *Glob  
932 Change Biol*, 20(11), 3557–3567, doi:10.1111/gcb.12551, 2014.  
933 Luo, Y., Shi, Z., Lu, X., Xia, J., Liang, J., Jiang, J., Wang, Y., Smith, M. J., Jiang, L., Ahlström, A., Chen,  
934 B., Hararuk, O., Hastings, A., Hoffman, F., Medlyn, B., Niu, S., Rasmussen, M., Todd-Brown, K.

935 and Wang, Y.-P.: Transient dynamics of terrestrial carbon storage: mathematical foundation and  
936 its applications, *Biogeosciences*, 14(1), 145–161, doi:10.5194/bg-14-145-2017, 2017.

937 M. J. H. van't Hoff: Etudes de dynamique chimique, Amsterdam, Frederik Muller & C°. [online] Available  
938 from: <https://doi.org/10.1002/recl.18840031003>, 1884.

939 Manzoni, S. and Porporato, A.: Soil carbon and nitrogen mineralization: Theory and models across scales,  
940 *Soil Biology and Biochemistry*, 41(7), 1355–1379, doi:10.1016/j.soilbio.2009.02.031, 2009.

941 Martin, M. P., Dimassi, B., Romàn Dobarco, M., Guenet, B., Arrouays, D., Angers, D. A., Blache, F.,  
942 Huard, F., Soussana, J., and Pellerin, S.: Feasibility of the 4 per 1000 aspirational target for soil  
943 carbon. A case study for France, *Glob Change Biol*, gcb.15547, <https://doi.org/10.1111/gcb.15547>,  
944 2021.

945 McBratney, Alex. B. and Minasny, B.: Comment on “Determining soil carbon stock changes: Simple bulk  
946 density corrections fail” [*Agric. Ecosyst. Environ.* 134 (2009) 251–256], *Agriculture, Ecosystems*  
947 *& Environment*, 136(1–2), 185–186, doi:10.1016/j.agee.2009.12.010, 2010.

948 Meersmans, J., Van WESEMAEL, B., Goidts, E., Van Molle, M., De Baets, S. and De Ridder, F.: Spatial  
949 analysis of soil organic carbon evolution in Belgian croplands and grasslands, 1960–2006: Spatial  
950 analysis of soil organic carbon evolution, *Global Change Biology*, 17(1), 466–479,  
951 doi:10.1111/j.1365-2486.2010.02183.x, 2011.

952 Mekonnen, K., Buresh, R. J. and Jama, B.: Root and inorganic nitrogen distributions in sesbania fallow,  
953 natural fallow and maize fields, 9, 1997.

954 Meyer, N., Welp, G. and Amelung, W.: The Temperature Sensitivity (Q10) of Soil Respiration: Controlling  
955 Factors and Spatial Prediction at Regional Scale Based on Environmental Soil Classes, *Global*  
956 *Biogeochem. Cycles*, 32(2), 306–323, doi:10.1002/2017GB005644, 2018.

957 Minasny, B., Malone, B. P., McBratney, A. B., Angers, D. A., Arrouays, D., Chambers, A., Chaplot, V.,  
958 Chen, Z.-S., Cheng, K., Das, B. S., Field, D. J., Gimona, A., Hedley, C. B., Hong, S. Y., Mandal,  
959 B., Marchant, B. P., Martin, M., McConkey, B. G., Mulder, V. L., O'Rourke, S., Richer-de-  
960 Forges, A. C., Odeh, I., Padarian, J., Paustian, K., Pan, G., Poggio, L., Savin, I., Stolbovoy, V.,  
961 Stockmann, U., Sulaeman, Y., Tsui, C.-C., Vågen, T.-G., van Wesemael, B. and Winowiecki, L.:  
962 Soil carbon 4 per mille, *Geoderma*, 292, 59–86, doi:10.1016/j.geoderma.2017.01.002, 2017.

963 Olson, K. R., Al-Kaisi, M. M., Lal, R. and Lowery, B.: Experimental Consideration, Treatments, and  
964 Methods in Determining Soil Organic Carbon Sequestration Rates, *Soil Science Society of*  
965 *America Journal*, 78(2), 348–360, doi:10.2136/sssaj2013.09.0412, 2014.

966 Pachauri, R. K., Mayer, L. and Intergovernmental Panel on Climate Change, Eds.: Climate change 2014:  
967 synthesis report, Intergovernmental Panel on Climate Change, Geneva, Switzerland., 2015.

968 Parton, W. J., Stewart, J. W. B. and Cole, C. V.: Dynamics of C, N, P and S in grassland soils: a model,  
969 *Biogeochemistry*, 5(1), 109–131, doi:10.1007/BF02180320, 1988.

970 Parton, W. J., Scurlock, J. M. O., Ojima, D. S., Gilmanov, T. G., Scholes, R. J., Schimel, D. S., Kirchner,  
971 T., Menaut, J.-C., Seastedt, T., Garcia Moya, E., Kamnalrut, A. and Kinyamario, J. I.:  
972 Observations and modeling of biomass and soil organic matter dynamics for the grassland biome  
973 worldwide, *Global Biogeochem. Cycles*, 7(4), 785–809, doi:10.1029/93GB02042, 1993.

974 Paustian, K., Lehmann, J., Ogle, S., Reay, D., Robertson, G. P. and Smith, P.: Climate-smart soils, *Nature*,  
975 532(7597), 49–57, doi:10.1038/nature17174, 2016.

976 Pellegrini, M., Saccani, C., Bianchini, A. and Bonfiglioli, L.: Sewage sludge management in Europe: a  
977 critical analysis of data quality, *IJEWM*, 18(3), 226, doi:10.1504/IJEWM.2016.10001645, 2016.

978 Pellerin, S., Bamière, L., Denis, A., Béline, F., Benoit, M., Butault, J.-P., et al.: Stocker du Carbone dans  
979 les sols Français - Quel Potentiel au Regard de L'objectif 4 pour 1000 et à Quel Coût? Synthèse du  
980 rapport d'étude. ADEME., *Environ. Sci. Policy*, 77, 130–139, doi:doi:  
981 10.1016/j.envsci.2017.08.003, 2017.

982 Piovesan, R. P., Favaretto, N., Pauletti, V., Motta, A. C. V. and Reissmann, C. B.: Perdas de nutrientes via  
983 subsuperfície em colunas de solo sob fertilização mineral e orgânica, *Rev. Bras. Ciênc. Solo*,  
984 33(4), 757–766, doi:10.1590/S0100-06832009000400002, 2009.

985 Poulton, P., Johnston, J., Macdonald, A., White, R. and Powlson, D.: Major limitations to achieving “4 per  
986 1000” increases in soil organic carbon stock in temperate regions: Evidence from long-term  
987 experiments at Rothamsted Research, United Kingdom, *Glob Change Biol*, 24(6), 2563–2584,  
988 doi:10.1111/gcb.14066, 2018.

989 Powlson, D. S., Whitmore, A. P., and Goulding, K. W. T.: Soil carbon sequestration to mitigate climate  
990 change: a critical re-examination to identify the true and the false, 62, 42–55,  
991 <https://doi.org/10.1111/j.1365-2389.2010.01342.x>, 2011.

992 Powlson, D. S., W., A. P.: The potential to increase soil carbon stocks through reduced tillage or organic  
993 material additions in England and Wales: A case study., *Agriculture, Ecosystems and*  
994 *Environment*, 146, 23–33, doi:doi:10.1016/j.agee.2011.10.004, 2012.

995 Redin, M., Recous, S., Aita, C., Dietrich, G., Skolaude, A. C., Ludke, W. H., Schmatz, R. and Giacomini,  
996 S. J.: How the chemical composition and heterogeneity of crop residue mixtures decomposing at  
997 the soil surface affects C and N mineralization, *Soil Biology and Biochemistry*, 78, 65–75,  
998 doi:10.1016/j.soilbio.2014.07.014, 2014.

999 Riggers, C., Poeplau, C., Don, A., Frühauf, C., and Dechow, R.: How much carbon input is required to  
1000 preserve or increase projected soil organic carbon stocks in German croplands under climate  
1001 change?, *Plant Soil*, 460, 417–433, <https://doi.org/10.1007/s11104-020-04806-8>, 2021.

1002 Rovira, P., Sauras, T., Salgado, J. and Merino, A.: Towards sound comparisons of soil carbon stocks: A  
1003 proposal based on the cumulative coordinates approach, *CATENA*, 133, 420–431,  
1004 doi:10.1016/j.catena.2015.05.020, 2015.

1005 Saffih-Hdadi, K. and Mary, B.: Modeling consequences of straw residues export on soil organic carbon,  
1006 *Soil Biology and Biochemistry*, 40(3), 594–607, doi:10.1016/j.soilbio.2007.08.022, 2008.

1007 Sanderman, J., Hengl, T. and Fiske, G. J.: Soil carbon debt of 12,000 years of human land use, *Proc Natl*  
1008 *Acad Sci USA*, 114(36), 9575–9580, doi:10.1073/pnas.1706103114, 2017.

1009 Smith, P., Smith, J. U., Powlson, D. S., McGill, W. B., Arah, J. R. M., Chertov, O. G., Coleman, K.,  
1010 Franko, U., Frolking, S., Jenkinson, D. S., Jensen, L. S., Kelly, R. H., Klein-Gunnewiek, H.,  
1011 Komarov, A. S., Li, C., Molina, J. A. E., Mueller, T., Parton, W. J., Thornley, J. H. M., and  
1012 Whitmore, A. P.: A comparison of the performance of nine soil organic matter models using  
1013 datasets from seven long-term experiments, *Geoderma*, 81, 153–225,  
1014 [https://doi.org/10.1016/S0016-7061\(97\)00087-6](https://doi.org/10.1016/S0016-7061(97)00087-6), 1997.

1015 Smith, P., Powlson, D., Glendining, M. and Smith, J.: Potential for carbon sequestration in European soils:  
1016 preliminary estimates for five scenarios using results from long-term experiments, *Global Change*  
1017 *Biology*, 3(1), 67–79, doi:10.1046/j.1365-2486.1997.00055.x, 1997.

1018 Soussana, J.-F.: Matching policy and science\_ Rationale for the ‘4 per 1000 - soils for food security and  
1019 climate’ initiative, 14, 2017.

1020 VandenBygaart, A. J.: Comments on soil carbon 4 per mille by Minasny et al. 2017, *Geoderma*, 309, 113–  
1021 114, doi:10.1016/j.geoderma.2017.05.024, 2018.

1022 Wang, X., Piao, S., Ciais, P., Janssens, I. A., Reichstein, M., Peng, S. and Wang, T.: Are ecological  
1023 gradients in seasonal Q10 of soil respiration explained by climate or by vegetation seasonality?,  
1024 *Soil Biology and Biochemistry*, 42(10), 1728–1734, doi:10.1016/j.soilbio.2010.06.008, 2010.

1025 Wiesmeier, M., Poeplau, C., Sierra, C. A., Maier, H., Frühauf, C., Hübner, R., Kühnel, A., Spörlein, P.,  
1026 Geuß, U., Hangen, E., Schilling, B., von Lützow, M. and Kögel-Knabner, I.: Projected loss of soil  
1027 organic carbon in temperate agricultural soils in the 21st century: effects of climate change and  
1028 carbon input trends, *Sci Rep*, 6(1), 32525, doi:10.1038/srep32525, 2016.

1029 Wollenberg, E., Richards, M., Smith, P., Havlík, P., Obersteiner, M., Tubiello, F. N., Herold, M., Gerber,  
1030 P., Carter, S., Reisinger, A., van Vuuren, D. P., Dickie, A., Neufeldt, H., Sander, B. O.,  
1031 Wassmann, R., Sommer, R., Amonette, J. E., Falcucci, A., Herrero, M., Opio, C., Roman-Cuesta,  
1032 R. M., Stehfest, E., Westhoek, H., Ortiz-Monasterio, I., Sapkota, T., Rufino, M. C., Thornton, P.  
1033 K., Verchot, L., West, P. C., Soussana, J.-F., Baedeker, T., Sadler, M., Vermeulen, S. and  
1034 Campbell, B. M.: Reducing emissions from agriculture to meet the 2 °C target, *Glob Change Biol*,  
1035 22(12), 3859–3864, doi:10.1111/gcb.13340, 2016.

1036 Xia, J. Y., Luo, Y. Q., Wang, Y.-P., Weng, E. S. and Hararuk, O.: A semi-analytical solution to accelerate  
1037 spin-up of a coupled carbon and nitrogen land model to steady state, *Geosci. Model Dev.*, 5(5),  
1038 1259–1271, doi:10.5194/gmd-5-1259-2012, 2012.

1039 Xu, W., Chen, X., Luo, G. and Lin, Q.: Using the CENTURY model to assess the impact of land  
1040 reclamation and management practices in oasis agriculture on the dynamics of soil organic carbon  
1041 in the arid region of North-western China, *Ecological Complexity*, 8(1), 30–37,  
1042 doi:10.1016/j.ecocom.2010.11.003, 2011.

- 1043 Zhang, B., Tian, H., Lu, C., Dangal, S. R. S., Yang, J. and Pan, S.: Global manure nitrogen production and  
1044 application in cropland during 1860–2014: a 5 arcmin gridded global dataset for Earth system  
1045 modeling, *Earth Syst. Sci. Data*, 9(2), 667–678, doi:10.5194/essd-9-667-2017, 2017.  
1046 Zinn, Y. L., Lal, R. and Resck, D. V. S.: Changes in soil organic carbon stocks under agriculture in Brazil,  
1047 *Soil and Tillage Research*, 84(1), 28–40, doi:10.1016/j.still.2004.08.007, 2005.