

# Barium stable isotopes as a fingerprint of biological cycling in the Amazon River Basin

Quentin Charbonnier<sup>1,2</sup>, Julien Bouchez<sup>1</sup>, Jérôme Gaillardet<sup>1,3</sup>, and Éric Gayer<sup>1</sup>

<sup>1</sup>Université de Paris, Institut de physique du globe de Paris, CNRS, F-75005 Paris, France

<sup>2</sup>Institute of Geochemistry and Petrology, Department of Earth Sciences, ETH Zürich, Clausiusstrasse 25, 8092 Zürich, Switzerland

<sup>3</sup>Institut Universitaire de France, Paris, France

**Correspondence:** Quentin Charbonnier (quentin.charbonnier@erdw.ethz.ch)

**Abstract.** Biological cycling of rock-derived nutrients **on the continents** is a major component of element cycles the Earth's surface, but its magnitude still remains elusive **to this day**. The use of **the** stable isotope composition of rock-derived nutrients, which can be fractionated during biological uptake, provides a promising path forward in quantifying biological cycling and its overall contribution to global element cycling. In this paper, **we rely on the nutrient-like behaviour of the trace element barium**  
5 **and use its** elemental and stable isotopic compositions in dissolved and sediment load river samples **to investigate biological cycling in the Amazon Basin**. From these measurements, we show that dissolved Ba derives mainly from silicate rocks while a correlation between dissolved Ba and K abundances suggests a role of biological cycling on the Ba river budget. Further, the isotope composition of Ba ( $\delta^{138}\text{Ba}$ ) in the dissolved load was found to be significantly different from that of the parent silicate rocks implying that dissolved Ba isotopic signatures are affected by i) precipitation of **soil-forming** secondary phases and ii)  
10 biological uptake and release from dead organic matter.

Results from an isotope mass balance applied to the river dissolved load data indicate that after its release to solution by rock weathering, Ba is partitioned between the **river** dissolved load, secondary weathering products such as those found in soils and river sediments, and the biota. In most sub-catchments of the Amazon, river Ba abundances and isotope compositions are significantly affected by biological cycling. Relationships between estimates of **Ba cycled through biota** and independent  
15 metrics of ecosystem dynamics (such as Gross Primary Production and Terrestrial Ecosystem Respiration) allow us to discuss the role of **environmental parameters such as climate or erosion rates** on the **biological cycling of Ba**, and by extension of major rock-derived nutrients.

In addition, catchment-scale mass and isotope budgets of Ba show that the measured riverine export of Ba is lower than the estimated delivery of Ba to the Earth surface through rock alteration. This indicates the existence of a missing Ba component,  
20 that we attribute to the formation of Ba-bearing particulate organics, possibly accumulating as soil organic matter or currently growing biomass within the catchments; and to organic-bound Ba exported as "unsampled" river particulate organic matter.

Given our findings on **the trace element Ba**, we explore whether the river fluxes of most major rock-derived nutrients (K, Mg, Ca) might also be significantly affected by biological uptake or release. A first-order correction of river-derived silicate weathering fluxes from biological cycling shows that, at the Amazon at mouth, the  $\text{CO}_2$  consumption by silicate weathering  
25 could be **several times higher** than the yet-reported value of  $13 \cdot 10^9$  mol of  $\text{CO}_2/\text{yr}$  by Gaillardet et al. (1997).

Overall, our study clearly shows that **in the Amazon - and most likely in other large river basins - the chemical and isotope compositions of rivers bear** a biological imprint, thereby challenging common assumptions made in weathering studies.

## 1 Introduction

A fundamental process on Earth is the chemical weathering of rocks by atmospheric agents such as oxygen, CO<sub>2</sub>, and water. Over the long-term evolution of our planet, chemical weathering reactions and the subsequent formation of secondary mineral phases in the ocean and on the continents have been the principal drivers in forming and sustaining habitable conditions on the planet surface (Langmuir and Broecker, 2012). Continental chemical weathering consists **in** a series of hydrogeochemical processes that transform rocks within the lithosphere into dissolved solutes, into secondary minerals such as clays that form soils, and, most importantly, into the living organisms that compose our biota (*e.g.* Porder, 2019). The complex reactions participating in these transformations occur in what is frequently referred to as the "critical zone" of the planet, a thin layer, almost invisible at the planetary scale, made of soils, surface and underground waters as well as dead and living organic matter (Riebe et al., 2017). These weathering products are then mobilised by the water cycle and harvested by rivers before reaching the ocean. Rivers therefore integrate over the diversity of critical zone processes and offer a quantitative window into the fluxes of matter implied in these processes and their controlling factors (Berner and Berner, 2012).

An important question related to the functioning of the critical zone is the role played by living organisms in its continuous transformation and long-term evolution. Biological activity has been suggested to significantly impact the Earth surface via a wide range of processes (Brantley et al., 2011) that often tend to have a variety of impacts on the critical zone. On the one hand, the decomposition of organic matter and root respiration in soils generates CO<sub>2</sub> and organic acids that acidify the soil solution and consequently trigger chemical weathering reactions (such as dissolution of primary minerals; *e.g.* Drever and Stillings, 1997; Calmels et al., 2014). On the other hand, vegetation and roots can also act as stabilizing agents by preventing physical erosion and thus preserving soil (*e.g.* Marston, 2010). Further, one of the most prominent effects **of** the presence of biota on Earth **follows from the various strategies adopted by living beings** to extract nutrients from rock minerals for their growth and the development of ecosystems (Aerts and Chapin III, 1999). **A major challenge facing ecosystems lies in the acquisition of these nutrients in a manner that is sustainable on the long term. Consequently, many studies documenting ecosystem nutrient acquisition strategies have emphasised that nutrients are strongly "recycled". This means that the nutrients contained in dead organic matter are efficiently taken up again after their release into solution (*e.g.* Jobbágy and Jackson, 2004; Cleveland et al., 2013). Nonetheless, nutrient recycling is never total, such that ecosystems eventually lose a small fraction of their nutrient inventory due to incomplete litter remineralization. Therefore, an alternative pathway has to exist to balance this nutrient loss. In places where Earth surface rejuvenation is relatively inefficient, strong nutrient depletion is compensated for by atmospheric deposition (Chadwick et al., 1999). In areas where erosion is active, the sustained supply of fresh minerals to the Earth surface provides to plants an access to "new" rock-derived nutrients (Porder et al., 2007; Cleveland et al., 2011). However, in such actively eroding areas loss of nutrients is also thought to be accelerated by litter erosion (Uhlir et al., 2017; Uhlir and von Blanckenburg, 2019).**

As a consequence, biological cycling acts as a "traffic officer" for elements and their isotopes through the critical zone, directing their circulation across the Earth surface (e.g. Chaudhuri et al., 2007; Cenki-Tok et al., 2009; Schmitt et al., 2012; Baronas et al., 2018). For simplicity, it has been commonly assumed that from the geochemical point of view element cycling by the biota is at "equilibrium" at the river scale (e.g. Viers et al., 2014) or, in other words that the flux of nutrient uptake is compensated for by an equivalent flux of release from the biota (e.g. Gaillardet et al., 1999a). However, more and more studies are now showing that higher plants are a significant reservoir of major rock-derived nutrients such as Ca, Mg, K or Si at the ecosystem scale (Burgelea et al., 2018) and that nutrient cycling is a key critical zone mechanism for understanding the partitioning of elements at the Earth surface (e.g. Uhlig et al., 2017).

The abundance and isotope signature of elements involved into biological cycling can be used to appraise nutrient uptake and the subsequent fluxes of elements passing through the critical zone. Barium (Ba), an alkali earth element, that has seven stable isotopes ( $^{130}\text{Ba}$ ,  $^{132}\text{Ba}$ ,  $^{134}\text{Ba}$ ,  $^{135}\text{Ba}$ ,  $^{136}\text{Ba}$ ,  $^{137}\text{Ba}$ ,  $^{138}\text{Ba}$ ; although unstable,  $^{130}\text{Ba}$  is considered to be stable given its long radioactive decay time). Besides the formation of secondary phases such as clays or oxides (Gong et al., 2019) and adsorption (Gou et al., 2019) that are considered to be major fractionation processes for a variety of weathering stable isotope tracers, previous studies indicate that Ba is sensitive to uptake by plants and bacteria (Bullen and Chadwick, 2016; Cam et al., 2016; Blondeau et al., 2018). Despite the lack of any known benefit of Ba for plants (Marschner, 2011) and the fact that accumulation of Ba in plants is toxic (Lamb et al., 2013; Myrvang et al., 2016, and reference therein), Ba shares similar chemical properties (ionic radius, charge) with major important rock-derived nutrients such as Mg, Ca, and K (Bullen and Bailey, 2005; Kabata-Pendias and Mukherjee, 2007), leading to significant uptake of Ba in plants and to a "nutrient-like behaviour". In addition to the sizeable amount of Ba usually found in plants, the isotopic composition of Ba has been found to faithfully trace nutrient cycling, to an even better extent than Sr or Ca stable isotopes (Bullen and Chadwick, 2016). This, together with its lack of sensitivity to redox conditions (unlike some of the transition metals or major constituents of the biosphere such as C, N, or S), makes Ba stable isotopes a potentially powerful tool to quantify biological cycling within the critical zone, and leads us to consider in this work that at first order Ba can be used as a proxy for the cycling of major rock-derived nutrients.

The study of Ba stable isotope fractionation at the Earth surface has started only recently (Von Allmen et al., 2010) and most contributions have hitherto focused principally on seawater (Horner et al., 2015; Cao et al., 2016; Hsieh and Henderson, 2017; Bates et al., 2017; Charbonnier et al., 2018), marine sediments (Bridgestock et al., 2018) and sedimentary barite (Crockford et al., 2019). In addition, the isotope composition of Ba dissolved in continental waters has been reported only for a few rivers, such that constraints are still lacking on the global dissolved Ba riverine flux to the ocean as well as on its isotope composition (Cao et al., 2016; Hsieh and Henderson, 2017; Gou et al., 2019).

In this study, we aim to provide estimates of rock-derived nutrient cycling, to reveal its controlling parameters, and to discuss the potential implications thereof through the use of the abundance and isotope signature of Ba in the Amazon, the world largest river basin. We rely on the abundance and isotopic composition of river dissolved and sediment Ba for a series of sub-catchments in the Amazon Basin spanning a variety of different parent rock lithologies, reliefs, climates, and vegetation types. We test different hypotheses to identify the principal drivers of the dissolved Ba isotopic composition of these rivers and demonstrate the significant role of biological cycling on the river Ba isotope signature. Using a set of river-scale elemental


and isotopic mass budgets for Ba in the Amazon River, we further show that biological cycling has a significant impact on the routing of Ba between the different compartments of the critical zone. We examine how this routing is dependent on the **environmental conditions** and in particular how it shifts between the mountainous regions of the Andes and the flatter, more expansive areas of the Amazon Basin. Finally, we explore how the export fluxes of major dissolved species from the Amazon Basin might be impacted by biological cycling and how this might bias estimates of chemical weathering **rates** based on river chemistry. Altogether, our study opens up new perspectives for the use of Ba isotope in sedimentary archives to trace past changes in ecosystem dynamics, provides further constraints on the isotope composition of Ba delivered globally to the oceans, emphasizes the role of biological cycling on the routing of **elements** at the Earth surface, and highlights the role played by life on **riverine** export.

## 2 Material and methods

### 2.1 Geographical and geological setting

The Amazon Basin, the world largest river basin with a drainage **area** of 5,500,000 km<sup>2</sup> and a discharge of 6,300 km<sup>3</sup>/yr (Milliman and Farnsworth, 2013), can be divided into three regions (Fig. 1):

- **The Andean belt** results from the subduction of the Nazca plate below the South American plate. The Northern part of the Amazon Andes is composed of a mixture of igneous rocks such as andesites and sedimentary rocks (*e.g.* shales and carbonates) of Paleozoic, Tertiary and Quaternary ages (Stallard and Edmond, 1983; Putzer, 1984; Moquet et al., 2011) and is drained by the Solimões tributaries, such as the Marañon, Ucayali, Huallaga, Morona, Tigre, or Pastaza. The Southern part of the Amazon Andes is composed almost exclusively of sedimentary rocks deposited during the same geological periods as in the Northern part, and drained by the Madeira tributaries, such as the Beni, Alto Beni, Chepete, Quiquibey, Madre de Dios, Mamoré or Orthon rivers. Most Andean rivers are characterized by high erosion rates (**higher than 500 t/km<sup>2</sup>/yr; Wittmann et al., 2011**) due to steep slopes, making the Andes the quasi-sole source of sediment to the Amazon mainstream (Gibbs, 1972; Allègre et al., 1996). These sediment-laden rivers, called "white waters" (Gibbs, 1967; Stallard and Edmond, 1983) also feature high dissolved loads. Rivers draining the Andean foreland (*e.g.* the Purus river), where Andean sediments have accumulated, are also considered as "white waters", although they have lower sediment and dissolved loads compared to Andean rivers. **In the Andes, precipitation and temperature are around 1,650 mm/yr and 20°C on average, respectively, with more arid and colder climate towards the South (example: around 1,200 mm/yr and 17°C for the Upper Beni).**
- **The Shields** underlie the sedimentary layers of the Amazon plains, and consist in a Precambrian craton composed of plutonic and metamorphic rocks, which crops out North (Guiana Shield) and South (Brazilian Shield) of the Amazon main channel. Amazon tributaries draining the Shields, such as the Tapajós and the Trombetas rivers, are dilute (**Stallard and Edmond, 1983; Gaillardet et al., 1997**) and feature low erosion rates (**about a few tens of t/km<sup>2</sup>/yr; Wittmann et al., 2011**), and are typically of the "clear water"-type (Gibbs, 1972; Stallard and Edmond, 1983), characterized by high



**Figure 1.** Map of the Amazon Basin and location of the sampling sites.

phytoplanktonic photosynthetic activity. There, precipitation rate is higher (around 2,300 mm/yr) and the temperature warmer (26°C) than in the Andes.

130 • **The Amazon Plain** where all tributaries of the Solimões and the Madeira join, and underlain by sediments derived from the Andean belt and the Shields. Rivers draining only the Amazon Plain, such as the upper Negro River, are dilute, sediment-poor, and typically of the "black water"-type, *i.e.* characterized by a high dissolved organic carbon concentration. In the Amazon Plain, Andean-fed rivers carrying sediments (around 200 t/km<sup>2</sup>/yr; Wittmann et al., 2011) have developed floodplains which store river sediments over periods of time up to several 10<sup>6</sup> yrs (Wittmann et al., 2011), before their release to the Amazon hydrographic system. Precipitation (2,130 mm/yr) and temperature (25°C) are relatively similar to those encountered in the Shield basins.

135 Throughout the remainder of the text, we will distinguish our set of sampling sites (Fig. 1) as (1) Northern Andean rivers (Solimões sub-basin); (2) Southern Andean rivers (Madeira sub-basin); (3) "dilute" rivers, lumping together rivers draining the Shields and the Amazon Plain only; (4) the "main tributaries", with sampling sites in the plain, but resulting from the mixing of waters derived from several regions (Andes, Shields, Plain).

## 2.2 Sampling

140 The samples used in this study (Fig. 1) have been collected during different sampling cruises performed from 2001 to 2008 (Dosseto et al., 2006; Bouchez et al., 2011; Dellinger et al., 2015b). Most analyses were performed at the High-Resolution Analytical Platform PARI of IPGP.

River waters were collected in acid-washed polypropylene containers and were filtered on site using Teflon filtration units (0.2- $\mu$ m porosity). Water samples were acidified at pH  $\approx$  2 with ultra pure HNO<sub>3</sub> and stored in a cold room at 4°C after  
145 collection. Major anion and cation concentration were measured using ion chromatography, dissolved silica concentration by UV-VIS spectrophotometry and trace elements by quadrupole ICP-MS.

Sampling procedures for river sediments are reported in Dosseto et al. (2006) and Bouchez et al. (2011). Briefly, at several locations, river sediments were sampled at different depths in the river channel and at the river bottom, in order to access the whole range of river sediment grain size. Major and trace element concentrations were measured by ICP-AES and ICP-MS,  
150 respectively, at the SARM (Service d'Analyse des Roches et des Minéraux, INSU facility, Vandoeuvre-les-Nancy, France; analytical details available at <http://helium.crpq.cnrs-nancy.fr/SARM>). In order to characterize the chemical and isotope composition of typical Andean bedrock, outcropping rocks were collected in 2001 and 2005. These rocks were crushed to obtain homogeneous powders, digested using a mix of HNO<sub>3</sub>/HF and repeatedly taken up in HCl 6 N until full dissolution. Concentration of major and trace elements in rocks were measured using quadrupole ICP-MS at IPGP.

155 In order to investigate the relative solubility of Ba (*i.e.* compared to soluble alkali and alkali-earth elements), we produced a compilation of existing data on the chemical composition of the dissolved load of rivers draining single rock types that are significantly present in the Amazon Basin (andesites, shales, plutonic igneous rocks such as granites, and carbonates), as well as a data compilation for the composition of the rock types themselves. The river data compilation is made of a combination of literature data (Edmond et al., 1995; Louvat and Allègre, 1997; Louvat et al., 2008) and our own new data, while rock data  
160 are derived from the GEOROC database (<http://georoc.mpch-mainz.gwdg.de/georoc/>; see Fig. S1 and data Tables in repository doi:10.5281/zenodo.3698299).

## 2.3 Barium chemical separation and isotope measurements

For each river water sample, a volume of 5 to 15 mL of water was evaporated at 100°C in a Teflon beaker, and the residue was taken up in 0.5 mL HCl 2.5 N. For each solid sample (river sediment or rock), aliquots of homogeneous powders (around 50  
165 mg of crushed powders allowing not only for Ba isotope measurements but also other analyses) were digested using a mix of HNO<sub>3</sub>/HF, evaporated to dryness, and repeatedly taken up in HCl 6 N until full dissolution of the residue. A sample aliquot containing at minimum 250 ng of Ba was evaporated, and the residue was taken up in 0.5 mL HCl 2.5 N.

The chemical separation of Ba was carried out by ion chromatography using the ion-exchange AG50W-X8 resin where the matrix elements such as Na, K, Ca, Mg, Fe, and Al were eluted in a volume of 13 mL of HCl 2.5 N, and Ba was eluted in a  
170 volume of 15 mL of HCl 6 N (Van Zuilen et al., 2016b). To ensure a proper purification of Ba from the matrix, the separation procedure was carried out twice in a row. Ba purification from river waters and sediments does not require the use of carbonate

co-precipitation, a method usually performed for Ba purification in sea water, and which can entail a loss of Ba (*e.g.* Horner et al., 2015). Therefore the double-spike method (*e.g.* Van Zuilen et al., 2016b) was not used here, and we rather corrected the mass instrumental fractionation using the sample-standard-bracketing method (see below). However, to ensure correct and accurate Ba isotope measurements, yield and eluate purity were checked for all samples by quadrupole ICP-MS (Table S.1). The total blank procedure was checked for each separation session, with a maximum of 0.2 ng of Ba, negligible compared to **the typical amount of Ba present in samples**.

The isotope ratios of Ba were measured using MC-ICP-MS (Thermo Fisher Neptune) **using** a SIS spray chamber. The Faraday cups were placed to collect masses 129 (L4), 130 (L3), 131 (L2), 132 (L3), 134 (C), 135 (H1), 136 (H2), 137 (H3) and 138 (H4). The concentration of samples and standards were adjusted to 100  $\mu\text{g/L}$ . Over a session, the reproducibility of the isotope measurement of a pure Ba solution was checked to be in the same range as **the values** reported by Nan et al. (2015). The isobaric interference by Xe on mass 134 was corrected by on-peak zeroes. Although the presence of residual Ce and La in the eluate can in principle lead to interferences on  $^{138}\text{Ba}$ , the very low concentration of these two elements in the river water samples did not significantly impact our measurement, as mass-dependent fractionation was shown for all dissolved load samples by plotting  $^{137}\text{Ba}/^{134}\text{Ba}$  vs.  $^{138}\text{Ba}/^{134}\text{Ba}$  (Fig. S2). For solid samples, the presence of residual yet measurable amounts of Ce and La in the eluate were shown to affect the  $^{138}\text{Ba}$  signal to some extent. Therefore for the whole sample set, mass-dependent fractionation was rather checked using the  $^{137}\text{Ba}/^{135}\text{Ba}$  and  $^{137}\text{Ba}/^{134}\text{Ba}$  ratios (Fig. S2). For the sake of consistency, all data were measured as  $\delta^{137}\text{Ba}$  and are reported here as  $\delta^{138}\text{Ba}$ , using the mass-dependent relation  $\delta^{138}\text{Ba} \approx 1.33 \times \delta^{137}\text{Ba}$ , with the  $\delta^{13x}\text{Ba}$  defined as:

$$\delta^{13x}\text{Ba}_{\text{smpl}} = \left( \frac{^{13x}\text{Ba}/^{134}\text{Ba}_{\text{smpl}}}{^{13x}\text{Ba}/^{134}\text{Ba}_{\text{std}}} - 1 \right) \times 1000 \quad (1)$$

with  $x = 7$  or  $8$ , the subscript *smpl* indicating the sample isotope ratio, and the subscript *std* indicating the reference isotope ratio (NIST SRM 3104a).

Uncertainties on sample  $\delta^{138}\text{Ba}$  values are reported as 95%-confidence interval (CI 95%), calculated as:

$$CI95\% = t_{n-1} \times \frac{S.D.}{\sqrt{n}} \quad (2)$$

where *S.D.* is the standard deviation over the  $n$  measurements of the sample (from 0.02 to 0.15‰), and  $t_{n-1}$  is the student's law factor with  $n - 1$  degrees of freedom at a 95% confidence level. The long-term reproducibility (over one year) and accuracy of the measurements were checked using the reference materials SRM3104a ( $0.00 \pm 0.03\%$  S.D.;  $n = 9$  from 3 chemical separations); SRM3104a spiked with two different river water matrices ( $-0.03 \pm 0.06\%$  S.D.;  $n = 6$  with 1 chemical separation for each matrix); JB-2 ( $0.05 \pm 0.08\%$  S.D.;  $n = 9$  chemical separations from two digestion batches) and BaBe27 ( $-0.81 \pm 0.08\%$  S.D.;  $n = 33$ ) matching well the values reported by Miyazaki et al. (2014) **and** Van Zuilen et al. (2016b) **for two of these two reference materials (BaBe27 = -0.82‰ and JB-2 = 0.07‰)**.

### 3 Results

The concentration and isotope composition of dissolved and particulate Ba in the Amazon river system are available at doi: 10.5281/zenodo.3698299.

#### 205 3.1 Ba abundance

The concentration of river dissolved Ba in the Amazon Basin ranges from 0.049  $\mu\text{mol/L}$  to 0.490  $\mu\text{mol/L}$  with an average of 0.202  $\mu\text{mol/L}$ . As water flow can exert a strong control on elemental concentrations through dilution, normalization to a conservative element such as Na helps comparing dissolved Ba abundances between different water samples. The most Ba-enriched river (highest Ba/Na ratio) is the Negro river. Andean tributaries from the Solimões and Madeira basins show  
210 homogeneous Ba/Na ratios, despite the fact that they drain different rock types. The main tributaries of the Amazon (Solimões and Madeira) show significant Ba enrichment with respect to their Andean tributaries.

Our compilation of Ba abundance (expressed as Ba/X ratios) for rocks and rivers draining single rock types (Fig. S1) highlights that Ba is depleted with respect to other alkali (Li, Na) and alkali Earth (Mg, Ca, Sr) elements during chemical weathering and thus that Ba is **affected by processes in the critical zone after its release from rocks at least to a larger extent**  
215 than these elements **are**.


The concentration of Ba in the suspended sediments of the Amazon River Basin is heterogeneous (from 222 mg/kg to 836 mg/kg). This is likely due to the variable quartz abundance throughout the water column and a subsequent dilution effect (Bouchez et al., 2011). Once normalized to thorium (Th) concentration (Th being an insoluble element with a magmatic compatibility similar to that of Ba), Ba abundance is even more variable, with Ba/Th ratios ranging from 13 to 125. Within  
220 this range, some Ba/Th values differ significantly from the reported ratio for the Upper Continental Crust (UCC) (Ba/Th  $\approx$  50; Taylor and McLennan, 1995; Rudnick and Gao, 2003), suggesting that processes can enrich or deplete Ba in river sediment **compared to source rocks** and/or that Amazon river sediment Ba is at least partially sourced from other rock types.

The fraction of river-borne Ba exported by Amazon rivers as dissolved species is calculated as:

$$w^{Ba} = \frac{[Ba]_{diss}}{[Ba]_{diss} + [Ba]_{spm} \times [spm]} \quad (3)$$

225 with  $[Ba]_{diss}$  the concentration of dissolved Ba in rivers in  $\mu\text{g/L}$ ,  $[Ba]_{spm}$  the Ba concentration in sediment in mg/kg, and  $[spm]$  the concentration of suspended particulate matter in river (obtained by long-term sediment gauging) in g/L. The fraction of river Ba exported as dissolved Ba ranges from 0.03 to 0.85. The lowest values are found for Andean tributaries, the highest values for "dilute" tributaries, whereas the main tributaries show intermediate values. Excluding the "dilute" tributaries, it appears that Ba is mainly transported as solids in rivers. At the mouth of the Amazon, 16 % of Ba is transported in dissolved  
230 forms.






**Figure 2.**  $\delta^{138}\text{Ba}$  of river dissolved load, river sediments, and shale rocks from the Amazon Basin. The composition of vegetation collected over a soil profile in Hawaii (Bullen and Chadwick, 2016) and of the UCC (Upper Continental Crust; Van Zuilen et al., 2016b; Charbonnier et al., 2018; Nan et al., 2018) reported in the literature is also shown.

### 3.2 Ba isotope composition

The isotopic composition of river dissolved Ba shows a wide range of  $\delta^{138}\text{Ba}_{diss}$  values from 0.29‰ to -0.10‰ without any clear difference between the various types of rivers. We acknowledge that, at this stage, we lack time series of  $\delta^{138}\text{Ba}_{diss}$  values. This could be problematic as it has been shown, for example, that  $\delta^{138}\text{Ba}_{diss}$  displays temporal variations in the monsoonal Yellow River Basin (Gou et al., 2019). However, this variability was attributed to variable degrees of Ba sorption onto particles, enabled by the very high concentration of suspended particulate matter in the Yellow River (several g/L; Gou et al., 2019). As the rivers of the Amazon Basin display much lower sediment **concentration** than that of the Yellow River (on the order of 100 mg/L for the largest tributaries; Bouchez et al., 2011), we contend that temporal variability in  $\delta^{138}\text{Ba}_{diss}$  is much less significant.

240 The Ba isotope composition of river sediments,  $\delta^{138}\text{Ba}_{\text{sed}}$  ranges from 0.06‰ to -0.25‰, with no significant difference between river suspended and bed sediments. The reported average values for plutonic rocks and andesites are  $0.00 \pm 0.17\%$  (S.D.) and  $0.07 \pm 0.02\%$  (S.D.), respectively (Charbonnier et al., 2018; Nan et al., 2018). Our own measurements of shale rocks collected in the Andes yield an average value for  $\delta^{138}\text{Ba}_{\text{rock}}$  of  $-0.02 \pm 0.04\%$  (S.D.) (Fig. 2). Ba in secondary weathering products is thus depleted in the "heavy" Ba isotopes with respect to *i.e.* the parent bedrocks that have lower isotope ratios, 245 whereas Ba in the dissolved load is enriched in the "heavy" Ba isotopes (*i.e.* have higher isotope ratios). This observation is consistent with the premise that the dissolved and suspended loads are complementary phases formed during the partitioning of Ba isotopes by chemical weathering (Fig. 2; Gong et al., 2019).

The Ba isotope composition of river sediments does not show any systematic variation with respect to the Al/Si ratio (a tracer for sediment grain size), with Al-rich samples (typically suspended matter **sampled near the river channel surface**) enriched in fine clays and Si-rich samples (typically bedload samples) enriched in coarse quartz (Bouchez et al., 2011, Fig. 3). This observation contrasts with what has been previously reported for other isotope systems such as  $^{87}\text{Sr}/^{86}\text{Sr}$  or  $^7\text{Li}/^6\text{Li}$  (Bouchez et al., 2011; Dellinger et al., 2014) and highlights that Ba isotopes in solid phase weathering products are poorly sensitive to grain size, as shown previously for Ge isotopes (Baronas et al., 2018).



**Figure 3.**  $\delta^{138}\text{Ba}$  of river sediments in the Amazon Basin as a function of their Al/Si ratio, used here as a tracer for sediment grain size (Bouchez et al., 2011).

## 4 Discussion

255 Significant Ba isotope fractionation is observed in the Amazon Basin between the dissolved and particulate loads of rivers (Fig. 2). In the following sections, we discuss the potential causes for this observation and show the influence of biological cycling on Ba isotope signatures.

### 4.1 Influence of critical zone processes on dissolved Ba


Variations in the dissolved Ba abundance and associated isotope composition could be controlled by variations in the sources  
260 of Ba and/or of critical zone processes affecting Ba after its release from rocks.

Although there are several possible sources of solutes contributing to the overall river chemistry, we show in Appendix A that silicate rocks are the main source of dissolved Ba to the river dissolved load in the Amazon Basin, consistent with previous observations from other river systems (Gou et al., 2019). This suggests that processes in the critical zone act to shift the abundance of river dissolved Ba between different basins. The fact that dissolved Ba is depleted with respect to elements  
265 classically considered as soluble (such as Na, Ca, or Mg) compared to rocks further lends support to this inference (Fig. S1).

Although still largely unconstrained, it is suspected that Ba incorporation into secondary weathering products (clays or oxi-hydroxides) occurs in the critical zone (Gong et al., 2019). River dissolved Li abundance and isotope composition can serve as a proxy for the contributions of secondary weathering products to the overall dissolved solute flux (von Strandmann et al., 2012; Dellinger et al., 2015b). When dissolved Ba is compared against dissolved Li, no correlation emerges between  
270  $Ba^*/Na^*$  and  $Li^*/Na^*$  ("**\***" referring to the dissolved element concentration corrected from non-silicate inputs; see Appendix A for details) nor between  $Li^*/Na^*$  and  $\delta^{138}Ba_{diss}$  (Fig. 4a,b) even when these elemental ratios are corrected for the source rock composition (these "corrected ratios" actually correspond to the so-called  $f_{diss}^{Ba}$  and  $f_{diss}^{Li}$  parameters; see below and Fig. S3). This lack of correlation implies that at least one other critical zone process (that is, in addition to the formation of secondary weathering products) scavenges dissolved Ba after its release from rocks. This inference is also supported by the absence of  
275 any significant differences in the Ba isotope compositions between shales (which are clay- and oxide-rich sedimentary rocks), unweathered igneous rocks, and river sediments (Fig. 2), which is in contrast to what is observed for Li (Dellinger et al., 2014).

It has been suggested that the isotope composition of Ba in soils is mainly driven by biological uptake and release (Bullen and Chadwick, 2016). This can be tested using the K abundance, which is known to be strongly impacted by biological activity at the Earth Surface (Chaudhuri et al., 2007; Tripler et al., 2006; Uhlig et al., 2017; Uhlig and von Blanckenburg, 2019) - in  
280 addition to having a magmatic compatibility (Hofmann, 1988) and a high abundance in shales (Taylor and McLennan, 1995; Rudnick and Gao, 2003) similar to those of Ba, **altogether suggesting a similar behaviour of Ba and K at the Earth Surface regarding "inorganic" processes (Kabata-Pendias and Mukherjee, 2007). The significant** positive relationship between river dissolved  $Ba^*/Na^*$  and  $K^*/Na^*$  (Fig. 4c) **suggests that the common behaviour between Ba and K extends to the biological realm.** However, no correlation is observed between  $\delta^{138}Ba_{diss}$  and  $K^*/Na^*$  (Fig. 4d).

285 Overall, the lack of correlation between  $\delta^{138}Ba_{diss}$  and a range of elemental ratios indicates that a combination of several processes is required to explain the Ba isotope signatures throughout the Amazon. In order to quantify the relative role of



**Figure 4.** Scatter plots of  $\text{Ba}^*/\text{Na}^*$  and  $\delta^{138}\text{Ba}_{diss}$  vs.  $\text{Li}^*/\text{Na}^*$  and  $\text{K}^*/\text{Na}^*$  for the dissolved load of the Amazon rivers ( "\*" refers to dissolved **element concentration** corrected from non-silicate inputs; see Appendix A for details).

each of these processes on the river dissolved Ba budget, we first need to calculate the fraction of weathering-derived Ba that remains in solution after Ba uptake by different processes. This can be done using Na as a reference element, *i.e.* under the assumption that dissolved Na is released to rivers only by weathering and is not significantly re-incorporated in solid secondary

290 weathering products during their formation, nor taken up as a nutrient by the biota (Gislason et al., 1996; Georg et al., 2007; Millot et al., 2010; Dellinger et al., 2015b):

$$f_{diss}^{Ba} = \frac{Ba^*/Na^*}{(Ba/Na)_0} \quad (4)$$

with  $Ba^*/Na^*$  the dissolved Ba/Na ratio corrected from non-silicate inputs, and  $(Ba/Na)_0$  the **bedrock ratio** (details about calculation are given in Appendix B). **Assuming** that Ba and Na are not fractionated **with respect to one another** during their  
 295 release from rocks,  $f_{diss}^{Ba}$  is then a proxy for the re-incorporation of Ba into secondary weathering solids (clays, oxides) or into organic matter.

The calculated fraction of initially dissolved Ba left in the dissolved load of rivers,  $f_{diss}^{Ba}$ , ranges from 0.04 in rivers draining only the Andean belt, to 1.63 for the Negro River. This latter value for a parameter which in principle should be lower than < 1, is somewhat surprising, and will be discussed later (section 4.5).

300  $f_{diss}^{Ba}$  does not show any correlation with dissolved  $\delta^{138}Ba_{diss}$  (not shown), supporting the idea that several compartments incorporate Ba in the critical zone following its release from rocks, and **that** the various processes responsible for this incorporation are characterized by different isotopic fractionation factors. In the following, in order to quantitatively tease these effects apart, we **use a mass balance approach** that **accounts for** the two processes likely to fractionate Ba in the critical zone: secondary phase formation and biological uptake.

#### 305 4.2 Assessing the relative role of biological uptake on the Ba cycle in the Amazon Basin

Element fluxes and isotope composition can be used in tandem to estimate the role of various processes on elemental budgets (*e.g.* Bouchez et al., 2013; Baronas et al., 2018). To this aim, we consider the behavior of Ba in the critical zone at the catchment scale under a scenario in which dissolved Ba, originally derived from the partial dissolution of rocks, is partitioned into different net uptake and dissolved export fluxes following the framework developed by Bouchez et al. (2013). The partitioning of Ba  
 310 after its release from rock minerals can be written as follows:

$$F_0^{Ba} = F_{diss}^{Ba} + F_{bio}^{Ba} + F_{sec}^{Ba} \quad (5)$$

with  $F_0^{Ba}$  the dissolution flux of Ba from rocks;  $F_{diss}^{Ba}$  the flux of river dissolved Ba exported from the system;  $F_{bio}^{Ba}$  the net Ba uptake by the biota ("net" meaning Ba uptake minus Ba release during organic matter remineralization);  $F_{sec}^{Ba}$  the Ba flux associated with the net formation (*i.e.* precipitation minus re-dissolution) associated to Ba-bearing secondary weathering  
 315 products such as clays or oxides.

Then dividing both sides of the eq. (5) by  $F_0^{Ba}$ , we obtain the relative proportion of all fluxes with respect to the dissolution flux:

$$1 = f_{diss}^{Ba} + f_{bio}^{Ba} + f_{sec}^{Ba} \quad (6)$$

where the terms  $f_i^{Ba}$  reflect the relative role of each process on Ba fluxes within the critical zone, with respect to the "initial" release flux of Ba from rocks  $F_0^{Ba}$ , and where  $f_{diss}^{Ba}$  represents in particular the "residual" dissolved Ba flux after each process, *i.e.* is equivalent to that in eq. (4). In other words, these terms represent the net formation fluxes of different critical zone compartments:  $f_{bio}^{Ba}$  is the net uptake of Ba by the biosphere and  $f_{sec}^{Ba}$  is the net flux of precipitation of Ba-bearing secondary phases of weathering. It should be emphasized that while these terms reflect the net formation of these pools, in reality the dynamics of these compartments result from the competition between their formation and their destruction. For example, clays formed deep within weathering profiles that are generally in equilibrium **with** solute-rich waters, can be destabilized when exposed to more dilute waters (Dellinger et al., 2015b); and organic matter can be remineralized in reservoirs with long residence times such as soils or floodplains (Bouchez et al., 2014b). A negative net formation ( $f_i^{Ba} < 0$ ) thus indicates that the pool *i* releases more Ba to the dissolved compartment in the critical zone than it incorporates Ba from this dissolved compartment, at least on the time scale over which  $f_i^{Ba}$  is estimated.

Following Bouchez et al. (2013), we can introduce the corresponding isotope mass balance, which importantly is valid only at steady state:

$$\delta_{diss}^{Ba} = \delta_{rock}^{Ba} - f_{bio}^{Ba} \times \Delta_{bio-diss}^{Ba} - f_{sec}^{Ba} \times \Delta_{sec-diss}^{Ba} \quad (7)$$

with  $\delta_{diss}^{Ba}$  the Ba isotope composition of the river,  $\delta_{rock}^{Ba}$  the composition of the dissolved load after rock dissolution (assuming congruent dissolution, this parameter equals the isotopic composition of the average bedrock);  $\Delta_{bio-diss}^{Ba}$  the Ba isotope fractionation factor during biological uptake; and  $\Delta_{sec-diss}^{Ba}$  the Ba isotope fractionation factor associated with the formation of secondary weathering products.

In order to quantify the role of nutrient cycling in the biogeochemical dynamics of Ba at the catchment scale, eq. (7) can be solved for the term  $f_{bio}^{Ba}$ . Replacing  $f_{sec}^{Ba}$  by  $(1 - f_{bio}^{Ba} - f_{diss}^{Ba})$  eq. (6) in eq. (7) and re-arranging leads to:

$$f_{bio}^{Ba} = \frac{\delta_{diss}^{Ba} - \delta_{rock}^{Ba} + \Delta_{sec-diss}^{Ba} - (f_{diss}^{Ba} \times \Delta_{sec-diss}^{Ba})}{(\Delta_{sec-diss}^{Ba} - \Delta_{bio-diss}^{Ba})} \quad (8)$$

Provided that the values of the parameters  $\delta_{rock}^{Ba}$ ,  $\Delta_{sec-diss}^{Ba}$ , and  $\Delta_{bio-diss}^{Ba}$  can be estimated,  $f_{bio}^{Ba}$  can be calculated using our measurements of  $\delta_{diss}^{Ba}$  and  $f_{diss}^{Ba}$  into eq. (8). In Appendix B, we discuss how the values of  $\delta_{rock}^{Ba}$ ,  $\Delta_{sec-diss}^{Ba}$ , and  $\Delta_{bio-diss}^{Ba}$  can be estimated. We also acknowledge that eq. (8) relies on a steady-state assumption, which is also discussed in Appendix B.

Using these constraints, the calculated  $f_{bio}^{Ba}$  values over the Amazon Basin range from  $-0.39_{-0.30}^{+0.25}$  to  $0.54_{-0.20}^{+0.24}$ , and  $f_{bio}^{Ba}$  of the Amazon at mouth is  $0.28_{-0.12}^{+0.25}$  (Table S.2). Only a few rivers show a  $f_{bio}^{Ba}$  close to 0, which challenges the following common assumption that biological cycling operates at steady-state from the point of view of the river dissolved load (*i.e.* that nutrient uptake equals nutrient release) at the catchment scale. Some of the rivers such as the Mamoré show a net release from the biosphere ( $f_{bio}^{Ba} < 0$ ), emphasizing that, locally in time and space, the biota can also act as a source of rock-derived nutrients to the river dissolved load.


The most significant findings of this analysis are the net uptake of Ba ( $f_{bio}^{Ba} > 0$ ) by the biota across many regions of the Amazon Basin and the fact that the extent of this net biological uptake (the exact value of  $f_{bio}^{Ba}$ ) displays significant spatial variation. Therefore, in the following section we explore the geomorphological, climatic, and ecological controls on the extent of Ba biological uptake.

### 4.3 Controls on Ba elemental and isotope partitioning by the weathering regime

The first hypothesis we set out to test regarding the spatial variation in Ba partitioning amongst critical zone compartments ( $f_i^{Ba}$ ) across the Amazon Basin is the role of the Earth surface denudation, which varies significantly between the mountainous Andes and the flatter Amazon lowlands. Denudation rates exert a strong control on the so-called "weathering regime", which in turn influences the behavior of elements and of their isotopes in the critical zone (Bouchez et al., 2013; Dellinger et al., 2015b; Frings et al., 2016). The catchment-scale denudation rate  $D$  is the sum of the chemical weathering  $W$  and physical erosion  $E$  rates measured from river material, typically using estimates of weathering-derived solute fluxes (e.g. Gaillardet et al., 1999b) for weathering and sediment gauging for erosion. Alternatively, denudation rates can also be estimated using cosmogenic nuclides such as  $^{10}\text{Be}$  in sediments (e.g. Wittmann et al., 2011). The three rates  $D$ ,  $E$ , and  $W$  are typically expressed in units of  $\text{t}/\text{km}^2/\text{yr}$  (or in  $\text{mm}/\text{y}$  using a rock-density based conversion). At one end of the spectrum of  $D$ , the "kinetically limited" weathering regime is characterized by high values of  $D$  and short particle residence time in the critical zone (West et al., 2005). Under these conditions, the extent of chemical reactions is limited by the kinetics (or how fast the reaction can proceed), leading to a low share of solute export ( $W$  for weathering) to total export rates (low values of the  $W/D$  ratios, used as an index for weathering intensity; Bouchez et al., 2014a). On the other end of the spectrum, the "supply-limited" regime, in which  $D$  values are low and particles reside for a long time within the weathering reactor, allows for more intense weathering and is characterized by higher  $W/D$  ratios (Riebe et al., 2004). To summarize the relation between these different metrics, low  $W/D$  ratios of mountainous regions are typically associated with both high  $W$  and  $D$  fluxes, and high  $W/D$  ratios characteristic of flatter areas are typically associated with both low  $W$  and  $D$  fluxes.

Weathering intensity, bounded by these "kinetically-limited" and "supply-limited" weathering regime endmembers, also translates into isotope signatures: for example, denudation exerts a strong control on the behavior of Li stable isotopes within the critical zone by modulating the extent of secondary mineral formation (Dellinger et al., 2015b; Frings et al., 2016). Unlike for Li isotopes, there is no direct relationship between  $\delta^{138}\text{Ba}_{diss}$  and  $D$  or  $W/D$  in our dataset (Fig. S4). This is because several critical zone processes act in conjunction to set the river dissolved Ba isotope signature in the Amazon Basin, as explained in section 4.1, while only the formation of secondary products drives dissolved Li isotopes (Lemarchand et al., 2010). However, the individual role of each critical zone process can be evaluated through the examination of the relationships between  $f_i^{Ba}$  (eqs. (4) and (8)) fractions estimated above using our mass balance approach, and  $W/D$  as markers of the weathering regime.

The relative role of the dissolved export to the total Ba input into the critical zone ( $f_{diss}^{Ba}$ ) displays a positive relation with  $W/D$  (Fig. 5a). This observation is in contrast to what has been reported for Li, which shows a lesser extent of Li scavenging from the solution (higher  $f_{diss}^{Li}$ ) when  $W/D$  is high or low, resulting in a so-called "boomerang" relationship (Dellinger et al.,



**Figure 5.** Distribution of Ba amongst different critical zone compartments ( $f_i^{Ba}$ ); dissolved export (*diss*) (panel a), biota (*bio*); eq. (8) (panel b), and secondary weathering products such as clays or oxides (*sec*) (panel c), once released from rocks vs.  $W/D$  (weathering intensity), in the Amazon Basin.  $W/D$  values are taken from Wittmann et al. (2011) and Dellinger et al. (2015b). In panel c,  $f_{sec}^{Ba}$  is calculated as  $1 - f_{diss}^{Ba} - f_{bio}^{Ba}$  (eq. (6)). Uncertainties on  $f_i^{Ba}$  have been evaluated using Monte Carlo error propagation based on the uncertainties on individual parameters (Appendix B; Table S.2.). Note the logarithmic scale used for  $f_{diss}^{Ba}$  in panel a.



2015b). For Ba, it is thus likely that even if, as for Li, Ba scavenging during formation of secondary weathering products is hampered at **low  $W/D$**  because of a kinetic limitation, another process scavenging dissolved Ba is being enhanced at **low  $W/D$** , the nature of which is discussed below.  $f_{bio}^{Ba}$  **does not show any trend with  $W/D$  (Fig. 5b)**, whereas  $f_{sec}^{Ba}$  shows a negative relation with  $W/D$  (Fig. 5c).

As for each river the sum of the three  $f_i^{Ba}$  values is equal to 1 by construction, some of the observed co-variation between  $f_i^{Ba}$  is spurious, as this closure constraint induces a negative covariance between the different  $f$ -terms. In order to independently examine the role of each process, we translate these Ba fractions (equivalent to non-dimensional Ba fluxes) into area-specific, dimensional fluxes  $F_i^{Ba}$  (express as a kg of Ba per year per km<sup>2</sup>) using the fact that by definition from eq. (6):


$$F_i^{Ba} = F_{diss}^{Ba} \times \frac{f_i^{Ba}}{f_{diss}^{Ba}} \quad (9)$$

with  $i = bio, sec, \text{ or } diss$ . In eq. (10) we can calculate  $F_{diss}^{Ba}$  using  $Q$  the river discharge (in m<sup>3</sup>/yr),  $[Ba]^*$  the concentration of silicate-derived river dissolved Ba ( $\mu\text{g/L}$ ), and  $S$  (in km<sup>2</sup>) the drainage area of the catchment (yielding a flux in mg/km<sup>2</sup>/yr; Table S.2):

$$F_{diss}^{Ba} = \frac{[Ba]^* \times Q}{S} \quad (10)$$

The area-specific flux of dissolved Ba remains constant across the range of denudation rates (Fig. 6a) or weathering regime indexed as the  $W/D$  ratio (Fig. 6b). Barium fluxes associated with both biological uptake and the formation of secondary weathering products **is high where denudation is strong**. The flux of Ba associated to secondary phase formation is positively related to the denudation rate and, hence, to the flux of Ba linked to biological uptake. **In settings with the highest denudation rates**, the formation of secondary weathering products is thought to become kinetically limited (Bouchez et al., 2013; Pogge von Strandmann and Henderson, 2015; Dellinger et al., 2015b). However, such limitation is not apparent for the corresponding Ba flux. This indicates again that the relevant processes for scavenging dissolved elements in the critical zone are distinct between Ba and Li.

The absolute flux of dissolved export Ba ( $F_{diss}^{Ba}$ ) being **relatively constant across the Amazon Basin**, combined with the fact that the fluxes of Ba associated to biological uptake and secondary phase formation **are the highest where  $D$  is high**, explains why  $f_{diss}^{Ba}$  (the relative dissolved Ba flux) **increases with  $W/D$  (Fig. 5a)**. Regardless of the explanation for this relatively invariant  $F_{diss}^{Ba}$ , it is critical to note that such a behaviour, observed here for Ba across the whole Amazon Basin, is in stark contrast with what has been reported for Li in the Andean part of the Amazon Basin (Dellinger et al., 2015b). In addition, it should be emphasized that the relationships shown in Figs. 5 and 6 include pure Andean rivers (*e.g.* Alto Beni), pure "dilute" rivers (*e.g.* Tapajós), and mixed rivers with active floodplains (*e.g.* Amazon at Óbidos). Dellinger et al. (2015b) have shown a specific behaviour of Li isotopes in floodplains, expressed as a strong enrichment of heavy Li isotopes in rivers featuring active floodplains, and a loss of dissolved Li between the upstream reaches and the downstream reaches of these rivers. Although we cannot fully exclude a role of processes specific to floodplains on the Ba cycle and isotope fractionation in the Amazon



**Figure 6.** Fluxes of Ba in the critical zone ( $F_i^{Ba}$ : dissolved export (*diss*), formation of secondary weathering products such as clays or oxides (*sec*), and biological uptake (*bio*); eq. (9)) vs.  $D$  (denudation, panel a), and  $W/D$  (weathering intensity, panel b) in the Amazon Basin.  $D$  and  $W/D$  values are taken from Wittmann et al. (2011) and Dellinger et al. (2015b). **Uncertainties on  $F_i^{Ba}$  have been evaluated using Monte Carlo error propagation based on the uncertainties on individual parameters (Appendix B; Table S.2). Negative median values of  $F_i^{Ba}$  are not plotted here because of the logarithmic scale.**

Basin, we note that no such loss of dissolved Ba is observed across the few floodplain reaches where such a calculation can be made (See Supporting text for details and Table S.3 for data; Bouchez et al., 2012; Dellinger et al., 2015b). In most floodplain reaches the **river dissolved Ba flux** actually increases downstream, which can be explained by the relative enrichment of Ba of pure "dilute" tributaries (as shown by their relatively high dissolved Ba/Na ratios; section 4.1) that also contribute water to large rivers featuring active floodplains.

The most striking feature of Fig. 6 remains the increase of the Ba biological uptake flux with denudation rates. Although this finding relies on isotope mass balance equations that inherently bear some uncertainties - in particular related to the steady state assumption itself, or to parameter estimates (Appendix B), such result is consistent with what has been observed at the much smaller soil scale reporting that biological cycling can be influenced by rock supply rate or age (e.g. Vitousek et al., 2003; Porder et al., 2007). In the following section, we explore potential reasons why such a relationship might exist at the

scale of large catchments in the Amazon Basin, in particular using independent estimates of **parameters relevant to ecosystem**  
425 **dynamics**.

#### 4.4 Relationships between biological uptake, weathering regime, and ecosystem dynamics


In the previous section, variations in catchment-scale Ba biological uptake fluxes across the Amazon Basin were examined as a function of  $D$  and  $W/D$  as metrics of the weathering regime. This analysis shows in particular how the partitioning of the nutrient-like element Ba in the critical zone differs from that of the non-nutrient Li. The prominent role of ecosystem dynamics  
430 (especially in terms of ecosystem nutrition) in setting Ba partitioning in the critical zone is thus further explored in the present section.

The fluxes of Ba associated to biological uptake are here compared to: (1) two catchment-scale carbon fluxes, that we believe capture the first-order mass balance dynamics of ecosystems: the Gross Primary Production (GPP; in  $t/km^2/yr$ ) and the Terrestrial Ecosystem Respiration (TER; in  $t/km^2/yr$ ), both retrieved from remote sensing data (Table S.4; Tramontana  
435 et al. (2016); Jung et al. (2019), see Supporting text for details) and (2) two major climatic factors affecting ecosystems, the mean annual temperature (MAT; in  $^{\circ}C$ ) and precipitation (MAP; in  $mm/yr$ ) (Table S.4; <https://www.worldclim.org>, see Supporting text for details). GPP can be envisaged as a measure of the biomass - most catchment-scale biomass estimates being anyway derived from GPP estimates (Tramontana et al., 2016; Jung et al., 2019). Respiration of organic matter leads to the release of nutrients to the ecosystem, such that catchment-scale TER can be rather used as a proxy for the recycling of  
440 rock-derived nutrients (and of elements characterized by a nutrient-like behavior such as Ba) through litter remineralization. Such constraints (GPP, TER, MAP, MAT) do not rely on any geochemical tool, and thus offer a fully independent piece of information on biological activity across the Amazon Basin compared to the Ba isotope approach used in this study until now.

Ecosystem dynamics, in particular as probed by GPP, are also related to rock-derived nutrient cycling as calculated for Ba isotopes (Fig. 7a,b) has shown by the broad difference between Andean and Plain tributaries, lending further confidence to the  
445 validity of our determination of  $F_i^{Ba}$ . In addition,  $F_{bio}^{Ba}$  is broadly negatively related to both MAP and MAT (Fig. 7c and Fig. 7d), suggesting a role of climatic factors on Ba biological cycling.

The relationships of Fig. 7 indicate that in the poorly-erosive, tectonically-quiet, warm and humid areas of the Amazon Basin such as the Shields and Amazon Plain, the elevated GPP (Fig. 7a) suggests that the size of the biota is large (as in particular allowed by high MAP and MAT), while the high TER (Fig. 7b) is indicative of efficient recycling of rock-derived  
450 nutrients from litter, resulting on average in smaller net fluxes associated to Ba biological uptake ( $F_{bio}^{Ba}$ ). On the contrary, in the highly-erosive, colder and drier areas of the Andes, the relatively low GPP (Fig. 7a) suggests a smaller size of the biota reservoir. TER is also relatively low there (Fig. 7b), indicating a lower efficiency in nutrient recycling of rock-derived from litter, thereby leading to a higher net flux of Ba biological uptake by the biosphere. This comparison between independent sets of constraints further strengthens our inference that some catchment-scale relationship exists between weathering regime,  
455 climate, and nutrient cycling across the Amazon basin.

To summarize, biological uptake as indexed by  $F_{bio}^{Ba}$  is higher in the Andes than in the lowland regions of the Amazon Basin. Based on existing knowledge on ecosystem nutrition, a set of mechanistic interpretations can be proposed to explain this



**Figure 7.** Flux of Ba biological uptake ( $F_{bio}^{Ba}$ ; eq. (9)) vs. Gross Primary Production (GPP) (in  $t/km^2/yr$ ; panel a), Terrestrial Ecosystem Respiration (TER) (in  $t/km^2/yr$ ; panel b), Mean Annual Precipitation (MAP) (in  $mm/yr$ ; panel c), and Mean Annual Temperature (MAT) (in  $T^{\circ}C$ ; panel d). GPP and TER were retrieved from remote sensing data (Tramontana et al., 2016; Jung et al., 2019), MAP and MAT are derived from <https://www.worldclim.org>.

finding. First, enhanced nutrient supply to ecosystem through rejuvenation of the Earth surface by denudation and subsequent weathering (high  $W$  and  $D$  in the Andes) might simply allow for more vigorous uptake of rock-derived nutrients (e.g Vitousek et al., 2003; Porder et al., 2007; Hahm et al., 2014; Uhlig and von Blanckenburg, 2019), although we note that no relation exists between  $W$  and  $F_{bio}^{Ba}$  in our dataset (not shown). Second, in mountainous regions lower temperature and precipitation combined with the short residence time of solid in the critical zone might lead to inefficient release of nutrients from litter (Trumbore et al., 1996). In the plains, higher temperature and precipitation increase the energy available for efficient nutrient recycling, with slow denudation and subsequent longer residence time of solid material allowing for a nearly-closed "recycling loop". Other factors influencing biological uptake and litter remineralization that might differ between the Amazon plains and

the Andes include the role of soil humidity (Grubb, 1971) and the residence time of water in the subsurface (Torres et al., 2015).

470 Altogether, a change in nutrient recycling efficiency is a likely explanation for the shift in rock-derived nutrient uptake fluxes between the Andes and the plain, as suggested by Ba isotopes. As a matter of fact, in most ecosystems the major nutrition acquisition pathway relies on the rapid recycling of litter (*e.g.* Cleveland et al., 2013; Wilcke et al., 2017; Uhlig and von Blanckenburg, 2019). Nonetheless, recycling is never complete as a small fraction of litter is exported from soils by erosion before remineralization (Selva et al., 2007; Scalley et al., 2012), as shown by the presence of particulate organic carbon of biogenic origin in river material (Selva et al., 2007; Bouchez et al., 2014b). Therefore, a slow but sustained supply from rock dissolution is needed to balance the loss from un-recycled nutrients by litter erosion, defined as the "geogenic" pathway  
475 (Uhlig and von Blanckenburg, 2019). Our results on the Amazon Basin suggest that a shift occurs in the relative role of these two pathways for acquisition of rock derived-nutrients, from the predominance of the organic pathway in the hot and humid environment of the plains, to an increased importance of the geogenic pathway in the colder, drier, and more erosive Andes (Fig. 7).

480 The above interpretation can further be tested using a so-called "river-scale mass budget", which allows for probing the form under which rock-derived nutrients are exported. Such a test is performed for Ba in the next section.

#### 4.5 The fate of Ba following its biological uptake in the Amazon Basin: clues from Ba elemental and isotope river-scale mass budgets

River-scale mass budgets, based on a hypothesized chemical complementarity between the weathered (chemical export) and eroded (physical export) material exported by rivers, have been used to test whether the chemical composition of the material  
485 exported by rivers can be explained by the composition of the rock undergoing erosion and weathering at the scale of large catchments (Stallard, 1995; Gaillardet et al., 1995, 1999a). A failure in such test can be indicative of issues in estimating the composition of the source rock (Gaillardet et al., 1999a) or in the determination of the composition of the riverine export (Bouchez et al., 2011), of timescale issues related to the typical integration time of the values used for the parameters of the river mass budget (Stallard, 1995; Lemarchand and Gaillardet, 2006), or any combination thereof.

490 In the scope of the present study on Ba isotopes, such river mass budget can be used to test whether all Ba released from rock dissolution is found in the total (that is, dissolved plus particulate) riverine export, in which case the mass budget can be considered to be "equilibrated". A lack of Ba in the riverine export would then mean that Ba is either currently accumulating in the catchment, and/or that the adopted sampling techniques do not capture the entirety of the riverine Ba export, or at least does not integrate the full river Ba export over the time scale that is relevant to establish the Ba river mass budget. As shown  
495 below, the form under which Ba is accumulated in, or exported from the catchment, can also be discussed based on the results of such a river-scale mass budget.

In addition, it is important to note that whereas strong uncertainties affect the different parameters used to compute  $f_{bio}^{Ba}$  above (in particular on the isotope fractionation factors  $\Delta_{sec-diss}^{Ba}$  and  $\Delta_{bio-diss}^{Ba}$ ), river-scale mass budgets as performed in

the following are based only on the relative partitioning of Ba and its isotopes between the dissolved and solid phases of rivers  
500 (Bouchez et al., 2013).

#### 4.5.1 Testing the equilibrium of the Ba river mass budget in the Amazon Basin

To assess whether the Ba river budget is "at equilibrium" in the Amazon, we can compare the fraction of Ba transported as a dissolved species ( $w^{Ba}$ ; eq. (3)) against the relative loss of Ba in river sediments compared to the source rocks. To quantify this loss, we normalise the Ba/Th ratio of river sediments to the Ba/Th ratio of the source rocks (the full derivation of the following  
505 equation is presented in Appendix C).

$$(Ba/Th)_n + w_{fluxes}^{Ba} = 1 \quad (11)$$

The two parameters  $(Ba/Th)_n$  and  $w_{fluxes}^{Ba}$  (as determined from eq. (3)) represent the fraction of dissolved and solid Ba transported in rivers relative to the denudation fluxes, respectively. Eq. (11) means that when the Ba river mass budget is at equilibrium, the sum of  $(Ba/Th)_n$  and  $w_{fluxes}^{Ba}$  is equal to 1; in other words data points should lie on the down-sloping  
510 diagonal in a  $w_{fluxes}^{Ba}$  vs.  $(Ba/Th)_n$  diagram (Fig. S5 and 8a). Rivers plotting below this line are characterized by an export of Ba lower than that expected from the basin-scale denudation rate and the Ba concentration in the source rock.

Such a test for an equilibrated river mass budget can also be performed using isotope ratios (Bouchez et al., 2013; Uhlig et al., 2017):

$$w_{iso}^{Ba} = \frac{\delta_{rock}^{Ba} - \delta_{sed}^{Ba}}{\delta_{diss}^{Ba} - \delta_{sed}^{Ba}} \quad (12)$$

515 In a  $w_{iso}^{Ba}$  vs.  $w_{fluxes}^{Ba}$  diagram, if the river mass budget is at equilibrium the data points should lie along the 1:1 line, while in a  $w_{iso}^{Ba}$  vs.  $(Ba/Th)_n$  diagram, they should lie along the down-sloping diagonal. Note that the  $w_{iso}^{Ba}$  vs.  $(Ba/Th)_n$  evaluation, unlike the  $w_{fluxes}^{Ba}$ - $(Ba/Th)_n$  comparison explained above, does not depend on fluxes. In order to provide an example of an equilibrated river mass budget, in Fig. S5 we re-visit the data provided by Dellinger et al. (2015a), who showed that in the Amazon the sum of the solid and dissolved Li fluxes exported by rivers, as well as their combined isotope composition,  
520 can be accounted for by independent estimate of flux and isotope composition of Li eroded and weathered from rocks. The river mass budget of Li shows that at the scale of the Amazon Basin, time scale issues stemming from the comparison of various parameters integrating Earth surface processes over different timescales (e.g. dissolved fluxes or particulate fluxes) are negligible, at least regarding the "inorganic" part of element cycles in the critical zone. This observation, combined with the fact that estimates of Ba abundance and isotope composition of source rocks can be considered as fairly reliable (Appendix B),  
525 suggests that any disequilibrium that might arise in the Ba river mass budget should find its origin in the nutrient-like behaviour of Ba.

The comparison between  $w_{fluxes}^{Ba}$  and  $(Ba/Th)_n$  (eq. (C5); Fig. 8a) shows that in most of the Amazon rivers, there is less Ba exported by rivers than expected from denudation rates. Notable exceptions are the "dilute" tributaries (Rio Negro, Trombetas and Tapajós). It is possible that in such rivers, the export of Th as a dissolved species induced by organic colloids compromises

530 the use of **the river mass budget (in particular eqs. (C2) and (C3); Appendix C)**. We also note that in these lowland areas, the input from dust (Abouchami et al., 2013; Moran-Zuloaga et al., 2018) can be significant **which could bias river-scale mass budgets**. In the following, we focus our interpretations on the other river systems, which all display a negative offset from the equilibrium line in Fig. 8a. For these rivers, we note that if a dust component was included in the left-hand sides of (C1) or (C2), the observed disequilibrium would go in the same direction and would even be larger, thus leaving this inference unchanged.


535 The observed disequilibrium in the Ba river mass budget also remains when comparing  $w_{iso}^{Ba}$  (eq. (12)) to  $(Ba/Th)_n$  (eq. (C5), Fig. 8b) or  $w_{fluxes}^{Ba}$  to  $w_{iso}^{Ba}$  (Fig. 8c), despite significant uncertainty on the values of  $w_{iso}^{Ba}$  due to both analytical uncertainties and error propagation (for example when  $\delta_{diss}^{Ba}$  and  $\delta_{sed}^{Ba}$  are close; see eq. (12) and discussion in Bouchez et al., 2013).

**From here**, in order to identify the potential causes for the lack of equilibrium in the river-scale mass budgets observed for most tributaries of the Amazon Basin, we can test the different scenarios that could reconcile the isotopic and elemental exports of Ba by Amazon rivers with denudation rates. As there is less Ba exported by rivers than expected from denudation (Fig. 8a), these different scenarios will all consider that to reconcile the two measures, **the** addition of a "missing Ba component", associated with a specific isotopic signature called hereafter  $\delta_{miss}^{Ba}$ , is needed. To illustrate these scenarios, arrows in the panels of Fig. 8 indicate in which directions the values on the three diagram axes will change upon addition of the missing Ba component:

- 1) the missing Ba is under a dissolved form and its isotopic signature ( $\delta_{miss}^{Ba}$ ) is lower than  $\delta_{diss}^{Ba}$
- 2) the missing Ba is under a dissolved form and  $\delta_{miss}^{Ba}$  is higher than  $\delta_{diss}^{Ba}$
- 3) the missing Ba is under a particulate form and  $\delta_{miss}^{Ba}$  is lower than  $\delta_{sed}^{Ba}$
- 4) the missing Ba is under a particulate form and  $\delta_{miss}^{Ba}$  is higher than  $\delta_{sed}^{Ba}$

550 For scenarios 1 and 2, adding the missing (dissolved) Ba would increase  $w_{fluxes}^{Ba}$  (eq. (3)) but leave  $(Ba/Th)_n$  unchanged. Scenarios 3 and 4 would result in a decrease of  $w_{fluxes}^{Ba}$ , while the shift they would induce on  $(Ba/Th)_n$  would depend on the Ba/Th ratio of the missing Ba component. We contend that this ratio is relatively high, as we expect - as discussed below in section 4.5.2 - that this missing Ba solid component would be of biological nature, and as Th is not a nutrient. In this case, scenarios 3 and 4 would result in the addition of Ba-rich material to the river solid load, and thus to an increase in  $(Ba/Th)_n$ , leading to non-vertical vectors in Fig. 8b and Fig. 8c. Finally, following eq. (12),  $w_{iso}^{Ba}$  would increase if  $\delta_{miss}^{Ba}$  is higher than  $\delta_{diss}^{Ba}$  (scenario 1) or than  $\delta_{sed}^{Ba}$  (scenario 3); and decrease if  $\delta_{miss}^{Ba}$  is lower (scenarios 2 and 4).

While by construction all these scenarios bring the data points closer to the equilibrium line in Fig. 8a, it appears that scenario 2 can be immediately excluded because adding dissolved Ba with a high  $\delta_{miss}^{Ba}$  would shift the river mass budget toward stronger disequilibrium (Fig. 8b and c). In addition, the direction given by the vector of scenario 4 is also unlikely to reconcile  $w_{iso}^{Ba}$  and  $(Ba/Th)_n$  for any river in Fig. 8b and  $w_{iso}^{Ba}$  and  $w_{fluxes}^{Ba}$  for most rivers in Fig. 8c. Indeed, some rivers (Beni at Riberalta and Rurrenabaque, Ucayali, and Mamoré) are above the equilibrium line in Fig. 8c, and their Ba river mass budget could be reconciled if the vector 1 had a sub-horizontal direction, which could be the case if the difference between  $\delta_{diss}^{Ba}$  and



**Figure 8.** Test for the equilibrium of the river mass budget of Ba in the Amazon Basin using elemental (eqs. (3) and (C4)) and isotopic (eq. (12)) ratios. Uncertainties on  $w_{fluxes}^{Ba}$  and  $(Ba/Th)_n$  have been calculated through gaussian error propagation. Values of  $w_{fluxes}^{Ba}$  have been calculated following two ways (using river gauging: circles and cosmogenic nuclides: diamonds; for a given river both estimates are linked by a stippled line) allowing us to partially account for potential timescale issues in the estimation of sediment export (see Appendix C). Uncertainties on  $w_{iso}^{Ba}$  have been evaluated using Monte Carlo error propagation based on the uncertainties on individual parameters in eq. (12), and are shown here as 68%CI. In panel c, "B@Rur", "B@Rib", "U", "M", "A" and "S" reflect the sampling sites: Beni @ Rurrenabaque, Beni @ Riberalta, Ucayali, Madeira, Amazon, and Solimões, respectively.

$\delta_{miss}^{Ba}$  was small. However, we first note that for these rivers the uncertainty on  $w_{iso}^{Ba}$  is large. Second, the most important finding regarding this comparison between  $w_{iso}^{Ba}$  and  $w_{fluxes}^{Ba}$  in Fig. 8c is still that the observed disequilibrium cannot be explained by scenario 2 and is unlikely to be explained by scenario 4.

To summarize, ruling out scenarios 2 and 4 allows us to infer that the isotopic composition of the missing Ba ( $\delta_{miss}^{Ba}$ ) is lower than that of Ba exported by rivers. Therefore, the remaining valid scenarios are 1 (the missing Ba is dissolved) and 3 (the



missing Ba is solid). However, the nature of (and the uncertainties encountered in) the analysis performed above prevents a full assessment of the exact nature of this missing Ba (dissolved or solid), and of its isotopic composition.

570 Nevertheless, the difference **between** the Li (Fig. S5) and Ba (Fig. 8) river mass budgets, and the fact that the Ba isotopic composition of the missing reservoir is low (hence consistent with the Ba isotope composition of a biological component; Bullen and Chadwick, 2016, and Fig. 2) are strong arguments towards a role of biological cycling in forming this Ba component missing from the river-scale mass budget. In the following, we further explore this hypothesis by comparing estimates of the amount of missing Ba to other independent estimates of Ba fluxes across the Amazon Basin.

#### 575 4.5.2 What is the nature of the "missing Ba" component in the Amazon riverine export?

**Given the role** of biological uptake on Ba abundance and isotope ratio in the rivers of the Amazon Basin, here we test for the influence of biological uptake on the "missing Ba" inferred from the river mass budget above.

First, we quantify the missing Ba as a relative flux ( $f_{miss}^{Ba}$ ) as the difference between the Ba flux actually exported by rivers and the Ba export that should result from denudation and the concentration of Ba in rocks. This actually corresponds to the graphical offset (vertical or horizontal) between  $(Ba/Th)_n$  (or equivalently  $w_{fluxes}^{Ba}$ ) and the down-sloping diagonal line in Fig. 8:

$$f_{miss}^{Ba} = 1 - w_{fluxes}^{Ba} - (Ba/Th)_n \quad (13)$$


It is important to note that  $f_{miss}^{Ba}$  can be interpreted **either** as an actual **relative** flux of Ba exported from the catchment but missed by our sampling scheme; or as a **relative** rate of Ba build-up into a reservoir within the catchment; or a combination thereof. We return to this distinction later when discussing the interpretation of  $f_{miss}^{Ba}$ , but name this metric a **relative** "flux" for the moment, for the sake of simplicity. To compare the flux of missing Ba ( $f_{miss}^{Ba}$ ) with the amount of Ba cycled by the biota in the Amazon ( $f_{bio}^{Ba}$ ), we first convert  $f_{miss}^{Ba}$  into a dimensional flux:

$$F_{miss}^{Ba} = D \times f_{miss}^{Ba} \times [Ba]_{rock} \quad (14)$$

with  $F_{miss}^{Ba}$  the area-specific flux of missing Ba in rivers, expressed in *e.g.* kg/km<sup>2</sup>/yr. We note that the denudation rates derived from the sum of sediment gauging (*E* **integrating over decades**) and silicate weathering flux (*W*) vs. cosmogenic nuclides which already represents the total denudation flux (*D* **integrating over thousands of years**; Wittmann et al., 2011; Dellinger et al., 2015a) can **differ** (especially in the Beni River system). Therefore, for rivers where both constraints are available, we calculate two values of  $F_{miss}^{Ba}$ : one based on *D* calculated as *E* from sediment gauging, plus the silicate weathering flux; and the other based on *D* determined from cosmogenic nuclides.

595 **Despite significant uncertainty, it is clear that the missing flux  $F_{miss}^{Ba}$  (eq. (14)) is of the same order of magnitude as  $F_{bio}^{Ba}$**  (Table S.6; Fig. 9) across the Amazon Basin. This **observation** further supports that biological uptake **is a viable explanation** for the disequilibrium in the river mass budget of the Amazon Basin. Interestingly, the relationship between  $F_{miss}^{Ba}$  and  $F_{bio}^{Ba}$  is

stronger when  $F_{miss}^{Ba}$  is calculated using cosmogenic isotopes in eq. (14) (Table S.6; Fig. 9). The cosmogenic-derived denudation rates integrate over longer timescales compared to that of sediment gauging (von Blanckenburg, 2005), suggesting that the link between Ba biological uptake and the Ba river mass budget is a long-term feature of the Ba cycle in the Amazon Basin. Altogether, these observations further support that the existence of the missing Ba flux in the Amazon Basin, inferred from the river-scale mass budget performed in section 4.5.1, is indeed linked to biological cycling.




**Figure 9.** Flux of Ba biological uptake ( $F_{bio}^{Ba}$ ; eq. (9)) vs. the missing Ba flux as quantified from the river-scale mass budget ( $F_{miss}^{Ba}$ ; eq. (14)) in the Amazon Basin. **Uncertainties have been propagated using a combination of Monte Carlo simulation and gaussian error propagation.**

A first possibility to account for such a missing Ba flux could derive from groundwater contributions that can export a significant amount of dissolved Ba. Thus, sampling schemes focusing only on river export might miss this potentially large dissolved Ba flux coming from the groundwater. However, with the magnitude of the river Ba dissolved flux and that of  $F_{miss}^{Ba}$  in the Amazon Basin being similar (Tables S.2 and S.6) coupled with observed Ba concentration in groundwater being similar to those measured in the rivers, this explanation would require that the groundwater discharge of the Amazon River is similar to that of the river discharge. Although seasonal groundwater storage in the Amazon Basin appears to be a significant component of the regional hydrological budget, especially in the central part of the Amazon Basin (Frappart et al., 2019), such a high contribution of groundwater to the overall water export of the whole Amazon basin seems unlikely. In addition, we do not see any particular reason why such a groundwater export would affect Ba and not Li (Fig. 8 and Fig. S5). It also appears unlikely that  $F_{miss}^{Ba}$  could be interpreted as a dissolved pool of Ba accumulating in the catchments given the typical residence times of groundwater even in large catchments and the fact that the observed disequilibrium in the Ba river mass budget seems to hold over long time scales (see above).

615 Therefore, the most likely interpretation for the existence of a significant missing Ba is the export (if interpreted as an actual flux) or accumulation (if interpreted as a pool building up) of Ba-bearing solids. Given the relationship observed between  $F_{bio}^{Ba}$  and  $F_{miss}^{Ba}$  (Fig. 9), it is plausible that these particulates are of organic nature, *i.e.* living beings, or dead organic matter (litter and/or river particulate organic carbon). In support of this interpretation, we note that there exists a relationship between TER, MAP, and MAT vs.  $F_{miss}^{Ba}$  (Fig. 10a,b,c), showing a more pronounced Ba missing flux, in basins for which precipitation, temperatures and the efficiency of the remineralization are lower and, thus, where the formation (and possibly export) of litter and dead particulate organic matter is stronger (Section 4.4).

620



**Figure 10.** Missing Ba flux ( $F_{miss}^{Ba}$ ; eq. (14)) vs. Terrestrial Ecosystem Respiration (TER; panel a); Mean Annual Precipitation (MAP; panel b); and Mean Annual Temperature (MAT; panel c). Uncertainties have been propagated using Monte Carlo simulation, and gaussian propagation error.

Accumulation of organic-bound Ba within catchments could explain the existence of the missing Ba. For example, a pool of solid, organic-bound Ba could be forming through accumulation of litter in soils, or through increase in the biomass - this can be the case in areas where forests grow again after important deforestation in the Amazon Basin. Although this hypothesis needs to be tested further, current growth of biomass in some regions of the Andes would support this scenario (*e.g.* Feeley and Silman, 2010).

625

Alternatively, the missing Ba could also reflect the existence of a particulate, organic debris flux. Uhlig et al. (2017) have suggested that the export of Mg and other rock-derived nutrients as particulate organic matter in mountain belts could represent a crucial missing riverine elemental flux, but is traditionally challenging to sample, notably because this export is mostly episodic, and, thus, difficult to constrain resulting in a typical timescale issue in the river mass budget, where the sampling frequency of river material does not allow to capture the entirety of the long-term riverine export. In large river systems such as the Amazon, the effect of episodic export on the total organic matter flux is likely to be smeared out, as suggested by the

630

relative regularity of particulate organic carbon fluxes throughout the hydrological cycle (Bouchez et al., 2014b). However, large woody debris (*e.g.* trunks) or river macrophytes could in principle represent a significant fraction of the particulate export of large rivers and be essentially unaccounted for due to limited amount of water that can be withdrawn with the sampling scheme commonly used (only a few liters with river point samplers) (Bouchez et al., 2011).

To test for the validity of this scenario, we can predict the flux of particulate organic carbon that would be needed to account for the computed  $F_{miss}^{Ba}$ :

$$F_{miss}^C = \frac{F_{miss}^{Ba}}{(Ba/C)_{org}} \quad (15)$$

with  $F_{miss}^C$  the missing flux of organic carbon and  $(Ba/C)_{org}$  the Ba/C ratio of the organic matter exported by rivers.  $F_{miss}^C$  can then be compared with  $E_{org}^C$ , the flux of particulate organic carbon exported by the river:

$$E_{org}^C = E \times POC \quad (16)$$

with POC representing the relative content of particulate organic carbon in the river sediments (Bouchez et al., 2014b), and  $E$  the erosion rate (from sediment gauging) (Table S.6).

Two estimates can be proposed for the value of  $(Ba/C)_{org}$  to calculate  $F_{miss}^C$  (eq. (15)). In the first estimate, we take the average concentration of Ba in organic matter of 15 mg/kg based on our own compilation (Li, 2000; Chiarenzelli et al., 2001; Bullen and Bailey, 2005; Vaganov et al., 2013, Table S.7), and a concentration of C in organic matter around 50%. In the second estimate, we use the ratio between the fluxes of Ba and C escaping litter decay, quantified as:

$$(Ba/C)_{org} = \frac{F_{bio}^{Ba}}{NEE} \quad (17)$$

where NEE is the Net Ecosystem Exchange calculated as GPP-TER, and corresponding to the net accumulation of carbon by the ecosystem. The values of  $(Ba/C)_{org}$  calculated this way differs significantly between the Andes and the plains. In the Andes, the Ba abundance in organic matter estimated using eq. (17) is much lower than that derived from our compilation (around 0.60 mg/kg; Table S.6). In the plains, the calculated Ba concentration in organic matter is even lower (around 0.01 mg/kg; Table. S.6). At face value, the difference between the values obtained for the Andes and the plain from eq. (17) suggests that the more efficient nutrient recycling is, the more weakly Ba is retained in the organic material exported by the river.

However, and regardless of the value used for  $(Ba/C)_{org}$ , the comparison between  $E_{org}^C$  (eq. (16)) and  $F_{miss}^C$  (eq. (15)) suggests that more than 99% of the particulate organic carbon would be missed when estimating the POC flux through a "classical" sampling scheme. We believe that it is unlikely that such a high proportion of the POC flux can be unaccounted for.

It remains thus challenging to identify the exact form of the missing Ba (*i.e.* flux vs. pool, or a combination thereof) inferred from the river-scale mass budget. However, the most likely interpretation for the existence of this missing component is the partitioning of Ba into biological material after uptake. As a consequence, our analysis emphasize the role of biological cycling on the routing and transport of rock-derived nutrients at the large catchment scale.

To summarize the above discussion, this examination of the Ba elemental and isotope budgets for a range of different river catchments throughout the Amazon Basin shows how stable isotope signatures can be used to explore nutrient cycling in the critical zone and, further, opens the possibility for an application to other isotope systems and other locations. Additionally, applying this approach to Ba provides novel information on the dynamics of the rock-derived nutrients, and possible of organic matter, at the scale of large catchments.

#### 4.6 A conceptual model for the influence of biological cycling and weathering regime on the fate of Ba as a nutrient-like element in the Amazon Basin.


To summarize our analysis, we propose the following interpretation for the behavior of Ba as a nutrient-like behavior, broadly reflecting the behaviour of major rock-derived nutrients in the Amazon Basin (Fig. 11):

- Except for some rivers draining the lowlands (see below), in the plain regions (Fig. 11a) litter remineralization outpaces litter erosion, allowing the biosphere to efficiently recycle Ba - and thus possibly rock-derived nutrients - from the litter and to minimize the uptake of "new" Ba derived from rock dissolution (Cleveland et al., 2013). In such conditions, primary production is strong, as promoted by high MAP and MAT, and soil stability allowed by low erosion rates leads to a weak export of particulate organic carbon from the catchment.
- In mountains regions of the Amazon (Fig. 11b), nutrient recycling by vegetation from litter is slower than litter erosion. As a consequence, the biota continuously extracts Ba and possibly major rock-derived nutrients from the surrounding rocks, which compensates for the loss of litter by erosion. Primary production is lower because the growth of biota is limited by the low MAP and MAT, strong denudation rates and short residence times of material, whereas export of rock-derived nutrients by particulate organic matter from the catchment (or accumulation within the catchment) is high.
- Some "dilute" rivers draining the lowlands are enriched in Ba with respect to the source rocks, which implies a net export of dissolved Ba from the biomass / organic matter component of the critical zone and thus suggests that these pools are currently not at steady state.


#### 4.7 Using Ba stable isotopes to refine the river mass budget of major elements

The above analysis shows that the cycling of Ba (and those of other rock-derived nutrients) by land biota is a significant component of terrestrial biogeochemical cycles. As a consequence, biological cycling within a catchment and the routing, or exchange, of rock-derived elements amongst the different compartments of the critical zone could bear consequence on the dissolved riverine export of major nutrients (Ca, Mg, K) that are traditionally used to estimate alkalinity fluxes and CO<sub>2</sub> consumption by chemical weathering at the catchment scale. In other words, nutrients that are taken up by the biota to a significant extent simply "go missing" in the river dissolved export fluxes. In the following, we add to existing inventories of river-borne cations concentrations deriving strictly from silicate weathering a component taken up by the biota, in order to

### a) Plains



### b) Mountains



**Figure 11.** Conceptual sketch for the Ba cycle in the critical zone in the **plains (a)** and **mountains (b)** of the Amazon Basin.  $E_{sec}^{Ba}$  and  $E_{bio}^{Ba}$  correspond to the erosion flux of secondary phase of weathering and organic-bound Ba, respectively. Other terms are defined in the text. For the sake of simplicity, the formed particulate organic carbon pool containing rock-derived nutrients such as Ba is assumed to be exported rather than accumulated in the basin.

update estimates of  $\text{CO}_2$  consumption fluxes by silicate weathering reaction at the basin scale in the light of our findings (see Appendix D for details).

695 First noting that the **biological** uptake flux of **any nutrient X** is:

$$[X]_{bio} = [Ba]_{diss} \times \frac{f_{bio}^{Ba}}{f_{diss}^{Ba}} \times \left( \frac{X}{Ba} \right)_{bio} \quad (18)$$

with  $[X]_{bio}$  the concentration of a major soluble element  $X$  (Ca, K or Mg,  $[Na]_{bio}$  being negligible as Na is not a nutrient) retained by the biosphere, and  $(X/Ba)_{bio}$  the  $X/Ba$  ratio of the organic component. The  $(X/Ba)_{bio}$  ratio is obtained from our own compilation of the chemical composition of the vegetation (Li, 2000; Chiarenzelli et al., 2001; Bullen and Bailey, 2005; Vaganov et al., 2013, ; Table S.7).  $[X]_{bio}$  can then be added to the silicate-derived dissolved river concentration of  $X$  in the calculation of "corrected" silicate weathering fluxes. To that aim, we express the impact of biological cycling on silicate

weathering fluxes as the ratio between cation deriving from silicate weathering added from that deriving from biological cycling ( $W_{silcorr}$ ) and uncorrected cation ( $W_{sil}$ ):

$$R_{(sil+bio)/sil} = \frac{W_{silcorr}}{W_{sil}} = \frac{2 \times ([Ca]_{sil}^{2+} + [Ca]_{bio}^{2+}) + 2 \times ([Mg]_{sil}^{2+} + [Mg]_{bio}^{2+}) + ([K]_{sil}^{+} + [K]_{bio}^{+}) + [Na]_{sil}^{+}}{2 \times [Ca]_{sil}^{2+} + 2 \times [Mg]_{sil}^{2+} + [K]_{sil}^{+} + [Na]_{sil}^{+}} \quad (19)$$

705  $W_{sil}$  is calculated using the sum of the concentration of each X element from silicate weathering and is expressed in meq/L, with  $W_{sil} = 2 \times [Ca]_{sil}^{2+} + 2 \times [Mg]_{sil}^{2+} + [K]_{sil}^{+} + [Na]_{sil}^{+}$  (data from Dellinger et al. (2015b)). As our primary interest is to quantify the effect of biological cycling on CO<sub>2</sub> consumption, we do not consider the SiO<sub>2</sub> component of the weathering flux in our analysis.

710  $R_{(sil+bio)/sil}$  values range from  $1.1_{-0.1}^{+1.7}$  (Huallaga River) to  $4.8_{-1.6}^{+3.0}$  (Trombetas River; Table S.8), while the value for the Amazon at mouth is  $2.1_{-0.5}^{+1.1}$ . These numbers show that the CO<sub>2</sub> consumption by silicate weathering, once corrected for biological uptake, could be higher by a factor of around two across the Amazon Basin compared to previous estimates. Despite significant uncertainties, this analysis shows that river-derived weathering fluxes warrant re-evaluation regarding the potential role played by the biota as a storage of rock-derived nutrients.

## 5 Conclusions

715 In the Amazon, the abundance and isotope composition of Ba - which we show to be sourced **mainly** from silicate rocks - is controlled by a combination of the formation of secondary weathering products and biological uptake, with a stronger isotope effect of the latter. **This "nutrient-like behavior" enables the use of Ba as a surrogate for major rock-derived nutrients, and allows us to examine the relationships between rock-derived nutrients cycling, weathering, and ecosystem dynamics**

Using isotope mass balance and estimates of isotope fractionation factors, we are able to propose a quantitative pattern  
720 for the partitioning of Ba liberated in the critical zone through rock weathering amongst secondary phases of weathering, the dissolved load, and the biota. The comparison of this partitioning to catchment-scale mass and isotope budgets of Ba (showing that the riverine export of Ba does not match the flux of Ba release from rocks) suggests that biological cycling of Ba leads to the formation of a "missing Ba component" in the critical zone of the Amazon Basin. Although the exact nature of this missing Ba component remains elusive (in particular regarding the fact that it could be accumulated within the catchment  
725 and/or exported by rivers), our favored interpretation is that it is locked into dead particulate organic matter (*i.e.* litter or river particulate organic carbon).

The proportion of Ba remaining in solution compared to that "initially" released by mineral dissolution is negatively correlated to denudation rates and positively correlated to weathering intensity, reflecting the combined effects of scavenging by secondary mineral precipitation and biological uptake. Biological uptake **of Ba** is found to be stronger **in the Andes**, possibly  
730 reflecting the need for the biota to compensate for the loss of organic components lost by erosion. **In the plains, where** denudation rate is low **and energy available for litter remineralization is high**, organic-bound Ba can **be released from litter** such that additional biological uptake **from rock weathering** is not required to replace the loss or nutrients bound to solid organic material by the ecosystem.

Finally, using the estimated uptake of Ba by the biosphere, we re-assess silicate weathering fluxes calculated from the cation  
735 loads of rivers with the effect of biological uptake now corrected for. We show that the cation dissolved load at the mouth of the Amazon is **significantly underestimated**, shifting best estimates of CO<sub>2</sub> consumption deduced from the cation dissolved load. This observation should be confirmed by using other rock-derived nutrients to reduce the uncertainties. Further, we believe that the method developed in this study to account for contributions from biological cycling should be extended to other large rivers in order to re-assess the CO<sub>2</sub> consumption at a global scale.

740 To conclude, our work shows that barium and its isotopes can be used to track biological cycling of rock-derived nutrients in large river basins and reveal its impact on the routing of nutrients throughout the critical zone. Such an informative approach warrants extension to other settings and other isotope systems in order to further reveal the role played by Life on chemical weathering process.



## 6 Appendices

### 745 Appendix A: Identification of dissolved Ba sources in the Amazon Basin rivers

Spatial variation in dissolved Ba abundance and isotope composition throughout the Amazon Basin could be due to source effects. For this reason, it is necessary to **assess** the relative contribution of atmospheric deposition and of release from different rock sources to the Ba river dissolved load (Gou et al., 2019).

Rain water is most likely not a significant source of Ba given the low concentration of dissolved Ba in surface seawater (a  
750 common source of solutes to rainwater) compared to that of dissolved Ba in rivers (Gaillardet et al., 2003). It is possible to further quantify the contribution of rain-derived Ba following:

$$[Ba]_{riv}^* = [Ba]_{riv} - [Cl]_{atm} \times \left( \frac{Ba}{Cl} \right)_{atm} \quad (A1)$$

where  $[Ba]_{riv}^*$  is the dissolved concentration of **dissolved** Ba in the river corrected from rain inputs,  $[Ba]_{riv}$  the measured concentration of Ba of the river,  $[Cl]_{atm}$  the concentration of dissolved Cl derived from the atmosphere following Gaillardet  
755 et al. (1997), and  $(Ba/Cl)_{atm}$  the seawater Ba/Cl ratio  $\approx 2 \times 10^{-7}$  **assuming that sea salts are the sole Ba contributor to rain**. Following this calculation, Ba derived from rain inputs is less than 1% of the total dissolved Ba for all rivers analyzed here.

Carbonate rocks formed in ancient marine environments are one of the most important sources of dissolved alkali-earth elements to rivers. However, the abundance of Ba in modern oceans is controlled by barite formation, rather than by carbonate precipitation (Dehairs et al., 1980). None of the  $Ca^*/Na^*$  vs.  $Ba^*/Na^*$  and  $\delta^{138}Ba$  vs.  $Ca^*/Na^*$  plots (Fig. S6a,b) show a  
760 correlation that could indicate mixing between a silicate and a carbonate weathering end member as a control on dissolved Ba abundance. We can assess the proportion of river dissolved Ba derived from carbonate weathering using the  $\alpha_{Ca}^{carbonate}$  (proportion of riverine dissolved Ca derived from carbonate weathering) from Dellinger et al. (2015b) using:

$$[Ba]_{riv}^{**} = [Ba]_{riv}^* - [Ca]_{riv}^* \alpha_{Ca}^{carb} \times \left( \frac{Ba}{Ca} \right)_{carb} \quad (A2)$$

where  $[Ba]_{riv}^{**}$  is the dissolved concentration of Ba corrected from both rain and carbonate inputs;  $[Ca]_{riv}$  is the measured  
765 concentration of dissolved Ca in the river; and  $(Ba/Ca)_{carb}$  the  $Ba^*/Ca^*$  ratio of rivers draining only carbonate rocks (**see Table S.2**), corrected from rain inputs, equal to  $5 \times 10^{-5}$  (median of  $n = 12$  measurements; Fig. S1). The highest proportion of dissolved Ba derived from carbonate rocks was found for the Madre de Dios river (about 10%), whereas carbonate weathering inputs to the Ba dissolved load for other tributaries do not exceed 5%.

In aqueous solutions, Ba shows a high affinity for sulfate ions, which leads to the formation of barite ( $BaSO_4$ ) in oceans  
770 and marine sediments. In particular, pre-concentration of Ba and  $SO_4$  induced by organic matter decay in micro-environments leads to the precipitation of  $BaSO_4$  (Griffith and Paytan, 2012). It has been also shown that the weathering of organic-rich sedimentary rocks such as black shales can be an important input of dissolved Ba to rivers (Dalai et al., 2002) which could be due to presence of barite in these rocks (Arndt et al., 2009). In addition, as an alkali-earth element, Ba could be contained in trace but significant amounts in gypsum, which could also contribute to the river dissolved Ba budget. To test for both

775 the influence gypsum and barite dissolution,  $\text{SO}_4^*/\text{Na}^*$  vs.  $\text{Ba}^*/\text{Na}^*$  and  $\delta^{138}\text{Ba}$  vs.  $\text{SO}_4^*/\text{Na}^*$  are plotted in Fig. S6c,d. No correlation can be observed between these parameters, implying that neither gypsum nor barite represent significant sources of river dissolved Ba in the Amazon Basin. To conclude, dissolved Ba in the Amazon Basin is almost exclusively derived from silicate rocks, and in our analysis we use our estimates of  $[\text{Ba}]^*$  of eq. (A1) as reflecting silicate-derived inputs.

## Appendix B: Parameters and assumptions of the Ba isotope mass balance

780 The estimation of  $f_{bio}^{Ba}$  from eq. (8) requires the estimation of several parameters, which we review hereafter.

### Source rock Ba abundance ( $(\text{Ba}/\text{Na})_0$ )

In order to compute the values of  $f_{diss}^{Ba}$ , the source ratio  $(\text{Ba}/\text{Na})_0$  first needs to be constrained. In the case of lithologically mixed basins, it can be calculated as:

$$(\text{Ba}/\text{Na})_0 = \sum_{i=1} \alpha_{rocki}^{Na} \times \left( \frac{\text{Ba}}{\text{Na}} \right)_0^i \quad (\text{B1})$$

785 with  $\alpha_{rocki}^{Na}$  the proportion of Na and  $\left( \frac{\text{Ba}}{\text{Na}} \right)_0^i$  the Ba/Na ratio of the rock  $i$ , and where  $i$  = shales, andesites, and plutonic rocks, as carbonate and evaporite rocks do not significantly influence the dissolved Ba and Na budgets, as shown Appendix A. The sources of dissolved Na within the Amazon catchment have been identified by Dellinger et al. (2015b). Values and uncertainties for each end member are reported in Table S.2.

### Source rock Ba isotope composition ( $\delta^{138}\text{Ba}_{rock}$ )

790 The value of  $\delta_{rock}^{Ba}$  to be used in eq. (8) depends on the relative contribution of each source rock type to the Ba dissolved flux. Three main silicate rocks have to be considered as potential sources of Ba for the Amazon basin: shales, andesites and plutonic rocks. First, using the relative contribution of Na ( $\alpha_i^{Na}$ ) calculated by Dellinger et al. (2015b), we can estimate the relative Ba contribution from each rock following:

$$\alpha_{rocki}^{Ba} = \frac{(\text{Ba}/\text{Na})_i}{(\text{Ba}/\text{Na})_0} \times \alpha_{rocki}^{Na} \quad (\text{B2})$$

795 Then, the average  $\delta_{rock}^{Ba}$  value can then be calculated as:

$$\delta_{rock}^{Ba} = \sum_{i=1} \alpha_{rocki}^{Ba} \times \delta_{rocki}^{Ba} \quad (\text{B3})$$

with  $\delta_{rock}^{Ba}$  the Ba isotope composition of each source rock  $i$ . We use our own measurements for shales ( $\delta_{rockshales}^{Ba} = -0.02 \pm 0.04\%$ ;  $n = 4$ ) and data from literature (Van Zuilen et al., 2016b; Nan et al., 2018) for andesites ( $\delta_{rockandesite}^{Ba} = 0.07 \pm 0.02\%$ ;  $n = 7$ ) and plutonic rocks ( $\delta_{rockplutonic}^{Ba} = 0.00 \pm 0.04\%$ ;  $n = 71$ ).

800 **Ba isotope fractionation factor associated to the formation of secondary weathering products ( $\Delta_{sec-diss}^{Ba}$ )**

As discussed above in section 4.1, secondary phases, *e.g.* clays or oxy-hydroxides, can at least partially drive the isotope composition of dissolved Ba in rivers (Gong et al., 2019). No experimental study on Ba isotope fractionation during precipitation of secondary weathering phases has been carried out to date, except for sorption on silica hydrogel which did not result in

a strong Ba isotope fractionation (Van Zuilen et al., 2016a). Nevertheless, we can try to assess a Ba isotope fractionation factor during formation of clays and oxides using our own measurements and previously published data. Marine clays do not show large variations in  $\delta^{138}\text{Ba}$ , with an isotope composition slightly depleted (by  $\sim -0.05\text{‰}$ ) in heavy isotopes with respect to igneous and sedimentary rocks (Bridgestock et al., 2018). Barium isotope fractionation during formation of secondary weathering products can also be estimated from data on soils. The difference between soil water and bulk solid soil Ba isotope composition over a latosol profile in China ranges from  $-0.01\text{‰}$  to  $-0.18\text{‰}$  (average value of  $-0.09\text{‰}$ ), likely induced by the difference in mineralogical composition, and by the varying respective role of sorption and precipitation over the profile (Gong et al., 2019). The large river approach undertaken here likely mitigates such effects of mineralogical heterogeneity, allowing for considering one "lumped" isotope fractionation factor (Bouchez et al., 2013). Our own measurement of Ba isotopes in sediments from the Amazon rivers show the same slight depletion in heavy isotopes compared to shales from the Amazon Basin (this study), or to published data on andesites (Van Zuilen et al., 2016b), with  $\delta^{138}\text{Ba}$  values ranging from  $-0.05$  to  $-0.10\text{‰}$ , consistent with the estimates reached above. Therefore, taking into account the above constraints and the associated uncertainty, we estimate that  $\Delta_{sec-diss}^{Ba} = -0.09 \pm 0.05\text{‰}$ .

#### **Ba isotope fractionation factor associated to biological uptake ( $\Delta_{bio-diss}^{Ba}$ )**

Bullen and Chadwick (2016) showed that the isotopic composition of dissolved Ba in the critical zone is strongly impacted by biological processes. However, no proper estimate of the Ba isotope fractionation factor associated to biological uptake is available yet. In order to represent the poor constraints on this parameter we consider  $\Delta_{bio-diss}^{Ba} = -0.25$  to  $-0.75$  using three  $\delta^{138}\text{Ba}$  values reported by Bullen and Chadwick (2016) for vegetation samples, and propagated uncertainty on this parameter in the following calculations assuming a uniform distribution between these values.

#### **Discussing the steady-state assumption**

Eq. (7) assumes that the Ba-bearing materials exported by rivers (excluding primary minerals, *i.e.* dissolved Ba, organics, and secondary weathering phases) correspond to all Ba derived from rock dissolution, thereby achieving Ba steady-state for all critical zone compartments. In such case, the net formation of each secondary phase can be envisaged to be compensated for by an erosion flux, as proposed in Bouchez et al. (2013).

In first "non-steady state scenario", the organic reservoir (biota + soil solid organics) grows simply because litter fall and erosion of organic matter are outpaced by biological uptake. In such a scenario, the isotopic composition of the dissolved Ba exported by rivers would not change. Therefore, in this case, the strict steady state assumption can be relaxed for the organic (biota plus soil organic matter) reservoir without undermining the validity of the steady state isotope mass balance of eq. (7).

Another possibility for non-steady state dynamics is the scenario where litter remineralization releases Ba to waters faster than Ba is taken up by vegetation (implying that the organic reservoir decreases in size), in which case the steady state assumption of the dissolved Ba reservoir is not met, compromising the use of eq. (7) above which derives from a steady-state framework (Bouchez et al., 2013). We first note that negative  $f_{bio}^{Ba}$  values, which should result from such cases where remineralization is outpaced by uptake, are only observed for three rivers from our dataset. Second, if  $f_{bio}^{Ba} < 0$ , over sufficiently short time scales the isotope composition of dissolved Ba can then rather be interpreted as a mixture between i) a "rock-derived" end-member, possibly affected by the formation of secondary phases of weathering and ii) an organic-derived end-member.

## Appendix C: River mass budget assumption and derivation

840 Here we present in details the different assumption and derivation required to performed river mass budget in section 4.5.

Quantitatively speaking, for an "equilibrated" mass budget, the amount of Ba delivered to the Earth surface by rock uplift and conversion into critical zone material, should be equal to the sum of Ba dissolved ( $F_{diss}^{Ba}$ ) and solid ( $F_{sed}^{Ba}$ ) exports by rivers:

$$F_{diss}^{Ba} + F_{sed}^{Ba} = D \times [Ba]_{rock} \quad (C1)$$

Using term  $D \times [Ba]_{rock}$  (with  $[Ba]_{rock}$  the abundance of Ba in the source rock) in eq. C1 is equivalent to assuming that  
 845 the denudation rate  $D$  is a proxy for the supply of rock material to the Earth surface. We acknowledge that  $D$  can be seen as an imperfect proxy for the supply of rock material to the Earth surface because its validity requires that the export of material by the river is equal to the supply of rock at depth, hence that some form of steady state exists for the regolith covering the Earth surface. However, we take the fact that in the Amazon the Li river mass budget (using eq. (C1)) is at equilibrium (Dellinger et al., 2015a) as a good indicator that this approach is reliable. We also note that this estimate of Ba supply to the Earth surface  
 850 neglects inputs from rock-derived nutrients from global dust (Abouchami et al., 2013; Moran-Zuloaga et al., 2018), which can be important in the Amazon lowlands (see discussion in section 4.5).

The same equation can be written for Th, which is an insoluble element with a magmatic compatibility close to that of Ba (Gaillardet et al., 1999a), thus minimizing the influence of the crustal composition when using rock Ba/Th ratios (see below):

$$F_{sed}^{Th} = D \times [Th]_{rock} \quad (C2)$$

855 with  $[Th]_{rock}$  the abundance of Th in the source rock and  $F_{sed}^{Th}$  the flux of Th transported as solids (*i.e.* as sediments) by the river. Then dividing eq. (C2) by eq. (C1) side by side:

$$\left(\frac{Ba}{Th}\right)_{rock} = \frac{F_{diss}^{Ba} + F_{sed}^{Ba}}{F_{sed}^{Th}} \quad (C3)$$

Then dividing the Ba/Th ratio of river sediments  $(Ba/Th)_{sed}$  (Table S.5) by the two sides of eq. (C3), and noting that

$$\left(\frac{Ba}{Th}\right)_{sed} = \frac{F_{sed}^{Ba}}{F_{sed}^{Th}}$$

$$860 \frac{\left(\frac{Ba}{Th}\right)_{sed}}{\left(\frac{Ba}{Th}\right)_{rock}} = \left(\frac{Ba}{Th}\right)_n = \frac{F_{sed}^{Ba}}{F_{sed}^{Ba} + F_{diss}^{Ba}} \quad (C4)$$

The term  $(Ba/Th)_n$  in eq. (C4) correspond to the inverse of the  $\alpha$ -depletion factor of soluble elements in river sediments, introduced by Gaillardet et al. (1999a); and is conceptually equivalent to  $1 + \tau_{Ba,Th}$  where  $\tau_{Ba,Th}$  is the mass transfer coefficient developed for quantify the depletion in soluble elements in soils (Brimhall and Dietrich, 1987). Finally, given the

definition of  $w^{Ba}$  (eq. (3), called hereafter " $w_{fluxes}^{Ba}$ " (for better distinction with " $w_{iso}^{Ba}$ " calculated below), an equilibrated river  
 865 mass budget should thus result in:

$$\frac{F_{sed}^{Ba}}{F_{sed}^{Ba} + F_{diss}^{Ba}} + \frac{F_{diss}^{Ba}}{F_{sed}^{Ba} + F_{diss}^{Ba}} = (Ba/Th)_n + w_{fluxes}^{Ba} = 1 \quad (C5)$$

Both parameters should in principle take values between 0 and 1, and we note that in our dataset, as Ba is always depleted over Th in the measured river sediment material compared to rocks,  $(Ba/Th)_n$  is between 0 and 1. Regarding  $w_{fluxes}^{Ba}$ , it is important to note that eq. (3) can be used based on river sediment gauging (yielding the averaged [ $spm$ ]) as in (3), or through:

$$870 \quad w_{fluxes}^{Ba} = \frac{Q \times [Ba]_{diss}}{D \times [Ba]_{rock}} \quad (C6)$$

Using (3),  $w_{fluxes}^{Ba}$  is by construction between 0 and 1, whereas if eq. (C6) is used, it is possible that  $w_{fluxes}^{Ba}$  takes values higher than 1 because the numerator and the denominator of the right-hand side of eq. C6 are evaluated in independent ways, and using metrics reflecting different time scales of critical zone dynamics. However, in our dataset, regardless of the way  $w_{fluxes}^{Ba}$  is calculated all values calculated for  $w_{fluxes}^{Ba}$  in the Amazon Basin are between 0 and 1.

875  $(Ba/Th)_{rock}$  in  $(Ba/Th)_n$  is estimated for each river using the rock contributions given by Dellinger et al. (2017), and the Ba/Th ratios of individual rock types (equal to 48 for igneous plutonic and shale rocks - which have a similar Ba/Th ratio - and of 140 for andesites; Table S.6).

## Appendix D: Calculation of dimensional, catchment-scale Ba biological uptake fluxes

The isotope mass balance model presented in section 4.3 allows to estimate  $f_{bio}^{Ba}$ , which a relative, hence non-dimensional flux of Ba biological uptake, with respect to the flux of Ba release from rock partial dissolution. In section 4.7, our analysis requires a dimensional values equivalent for  $f_{bio}^{Ba}$ , but for other rock-derived, major nutrients.

Recalling the definition of  $f_{bio}^{Ba}$ :

$$f_{bio}^{Ba} = \frac{F_{bio}^{Ba}}{F_0^{Ba}} \quad (D1)$$

and applying the same definition for another element X:

$$f_{bio}^X = \frac{F_{bio}^X}{F_0^X} \quad (D2)$$

Now recalling the definition of  $f_{diss}^{Ba}$ :

$$f_{diss}^{Ba} = \frac{F_{diss}^{Ba}}{F_0^{Ba}} \quad (D3)$$

and applying the same definition for another element X :

$$f_{diss}^X = \frac{F_{diss}^X}{F_0^X} \quad (D4)$$

we can re-arrange these equations and obtain:

$$\frac{f_{bio}^{Ba}}{f_{diss}^{Ba}} = \frac{F_{bio}^{Ba}}{F_{diss}^{Ba}} \quad (D5)$$

such that the net, dimensional flux of X biological uptake is:

$$F_{bio}^X = Q \times [Ba]_{diss} \times \frac{f_{bio}^{Ba}}{f_{diss}^{Ba}} \times \left( \frac{X}{Ba} \right)_{bio} = F_{diss}^{Ba} \times \frac{F_{bio}^{Ba}}{F_{diss}^{Ba}} \times \left( \frac{X}{Ba} \right)_{bio} \quad (D6)$$

895 *Author contributions.* QC performed new analytical work, interpreted data, and wrote text. JB and JG designed the study, conducted field work and performed previous analytical work used in the present contribution, interpreted data, and wrote text. EG performed the analysis of remote sensing data and wrote text.

*Competing interests.* The authors declare that they have no conflict of interest.

*Data availability.* Supplementary data tables are available in the online version of the paper.

900 *Acknowledgements.* The authors are grateful to Pascale Louvat, Thibaud Sontag, Jessica Dallas, Caroline Gorge, and Pierre Burckel for analytical support. Nicole Fernandez is thanked for English corrections. Geochemical analyses presented in this study were enabled by the IPGP multidisciplinary program PARI and by the Region Île-de-France SESAME Grant no. 12015908, and by the grant "Émergences" awarded by the City of Paris to Julien Bouchez.

## References

- Abouchami, W., Nätke, K., Kumar, A., Galer, S. J., Jochum, K. P., Williams, E., Horbe, A. M., Rosa, J. W., Balsam, W., Adams, D., et al.:  
905 Geochemical and isotopic characterization of the Bodélé Depression dust source and implications for transatlantic dust transport to the  
Amazon Basin, *Earth and Planetary Science Letters*, 380, 112–123, 2013.
- Aerts, R. and Chapin III, F. S.: The mineral nutrition of wild plants revisited: a re-evaluation of processes and patterns, in: *Advances in  
ecological research*, vol. 30, pp. 1–67, Elsevier, 1999.
- Allègre, C. J., Dupré, B., Négrel, P., and Gaillardet, J.: Sr, Nd, Pb isotope systematics in Amazon and Congo River systems: constraints about  
910 erosion processes, *Chemical Geology*, 131, 93–112, 1996.
- Arndt, S., Hetzel, A., and Brumsack, H.-J.: Evolution of organic matter degradation in Cretaceous black shales inferred from authigenic  
barite: A reaction-transport model, *Geochimica et Cosmochimica Acta*, 73, 2000–2022, 2009.
- Baronas, J. J., Torres, M. A., West, A. J., Rouxel, O., Georg, B., Bouchez, J., Gaillardet, J., and Hammond, D. E.: Ge and Si isotope signatures  
in rivers: A quantitative multi-proxy approach, *Earth and Planetary Science Letters*, 503, 194–215, 2018.
- 915 Bates, S. L., Hendry, K. R., Pryer, H. V., Kinsley, C. W., Pyle, K. M., Woodward, E. M. S., and Horner, T. J.: Barium isotopes reveal role of  
ocean circulation on barium cycling in the Atlantic, *Geochimica et Cosmochimica Acta*, 204, 286–299, 2017.
- Berner, E. K. and Berner, R. A.: *Global environment: water, air, and geochemical cycles*, Princeton University Press, 2012.
- Blondeau, M., Benzerara, K., Ferard, C., Guigner, J.-M., Poinot, M., Coutaud, M., Tharaud, M., Cordier, L., and Skouri-Panet, F.: Impact  
of the cyanobacterium *Gloeomargarita lithophora* on the geochemical cycles of Sr and Ba, *Chemical Geology*, 483, 88–97, 2018.
- 920 Bouchez, J., Gaillardet, J., France-Lanord, C., Maurice, L., and Dutra-Maia, P.: Grain size control of river suspended sediment geochemistry:  
Clues from Amazon River depth profiles, *Geochemistry, Geophysics, Geosystems*, 12, 2011.
- Bouchez, J., Gaillardet, J., Lupker, M., Louvat, P., France-Lanord, C., Maurice, L., Armijos, E., and Moquet, J.-S.: Floodplains of large  
rivers: Weathering reactors or simple silos?, *Chemical Geology*, 332, 166–184, 2012.
- Bouchez, J., von Blanckenburg, F., and Schuessler, J. A.: Modeling novel stable isotope ratios in the weathering zone, *American Journal of  
925 Science*, 313, 267–308, 2013.
- Bouchez, J., Gaillardet, J., and von Blanckenburg, F.: Weathering intensity in lowland river basins: from the Andes to the Amazon mouth,  
*Procedia Earth and Planetary Science*, 10, 280–286, 2014a.
- Bouchez, J., Galy, V., Hilton, R. G., Gaillardet, J., Moreira-Turcq, P., Pérez, M. A., France-Lanord, C., and Maurice, L.: Source, transport  
and fluxes of Amazon River particulate organic carbon: insights from river sediment depth-profiles, *Geochimica et Cosmochimica Acta*,  
930 133, 280–298, 2014b.
- Brantley, S. L., Megonigal, J. P., Scatena, F. N., Balogh-Brunstad, Z., Barnes, R. T., Bruns, M. A., Van Cappellen, P., Dontsova, K., Hartnett,  
H. E., Hartshorn, A. S., et al.: Twelve testable hypotheses on the geobiology of weathering, *Geobiology*, 9, 140–165, 2011.
- Bridgestock, L., Hsieh, Y.-T., Porcelli, D., Homoky, W. B., Bryan, A., and Henderson, G. M.: Controls on the barium isotope compositions  
of marine sediments, *Earth and Planetary Science Letters*, 481, 101–110, 2018.
- 935 Brimhall, G. H. and Dietrich, W. E.: Constitutive mass balance relations between chemical composition, volume, density, porosity, and strain  
in metasomatic hydrochemical systems: results on weathering and pedogenesis, *Geochimica et Cosmochimica Acta*, 51, 567–587, 1987.
- Bullen, T. and Bailey, S. W.: Identifying calcium sources at an acid deposition-impacted spruce forest: a strontium isotope, alkaline earth  
element multi-tracer approach, *Biogeochemistry*, 74, 63–99, 2005.



- Bullen, T. and Chadwick, O.: Ca, Sr and Ba stable isotopes reveal the fate of soil nutrients along a tropical climosequence in Hawaii, *Chemical Geology*, 422, 25–45, 2016.
- 940 Burghelea, C. I., Dontsova, K., Zaharescu, D. G., Maier, R. M., Huxman, T., Amistadi, M. K., Hunt, E., and Chorover, J.: Trace element mobilization during incipient bioweathering of four rock types, *Geochimica Et Cosmochimica Acta*, 234, 98–114, 2018.
- Calmels, D., Gaillardet, J., and François, L.: Sensitivity of carbonate weathering to soil CO<sub>2</sub> production by biological activity along a temperate climate transect, *Chemical Geology*, 390, 74–86, 2014.
- 945 Cam, N., Benzerara, K., Georgelin, T., Jaber, M., Lambert, J.-F., Poinot, M., Skouri-Panet, F., and Cordier, L.: Selective uptake of alkaline earth metals by cyanobacteria forming intracellular carbonates, *Environmental Science & Technology*, 50, 11 654–11 662, 2016.
- Cao, Z., Siebert, C., Hathorne, E. C., Dai, M., and Frank, M.: Constraining the oceanic barium cycle with stable barium isotopes, *Earth and Planetary Science Letters*, 434, 1–9, 2016.
- Cenki-Tok, B., Chabaux, F., Lemarchand, D., Schmitt, A.-D., Pierret, M.-C., Viville, D., Bagard, M.-L., and Stille, P.: The impact of water–rock interaction and vegetation on calcium isotope fractionation in soil-and stream waters of a small, forested catchment (the Strengbach case), *Geochimica et Cosmochimica Acta*, 73, 2215–2228, 2009.
- 950 Chadwick, O. A., Derry, L. A., Vitousek, P. M., Huebert, B. J., and Hedin, L. O.: Changing sources of nutrients during four million years of ecosystem development, *Nature*, 397, 491–497, 1999.
- Charbonnier, Q., Moynier, F., and Bouchez, J.: Barium isotope cosmochemistry and geochemistry, *Science Bulletin*, 63, 385–394, 2018.
- 955 Chaudhuri, S., Clauer, N., and Semhi, K.: Plant decay as a major control of river dissolved potassium: a first estimate, *Chemical Geology*, 243, 178–190, 2007.
- Chiarenzelli, J., Aspler, L., Dunn, C., Cousens, B., Ozarko, D., and Powis, K.: Multi-element and rare earth element composition of lichens, mosses, and vascular plants from the Central Barrenlands, Nunavut, Canada, *Applied Geochemistry*, 16, 245–270, 2001.
- Cleveland, C. C., Townsend, A. R., Taylor, P., Alvarez-Clare, S., Bustamante, M. M., Chuyong, G., Dobrowski, S. Z., Grierson, P., Harms, K. E., Houlton, B. Z., et al.: Relationships among net primary productivity, nutrients and climate in tropical rain forest: a pan-tropical analysis, *Ecology letters*, 14, 939–947, 2011.
- 960 Cleveland, C. C., Houlton, B. Z., Smith, W. K., Marklein, A. R., Reed, S. C., Parton, W., Del Grosso, S. J., and Running, S. W.: Patterns of new versus recycled primary production in the terrestrial biosphere, *Proceedings of the National Academy of Sciences*, 110, 12 733–12 737, 2013.
- 965 Crockford, P. W., Wing, B. A., Paytan, A., Hodgskiss, M. S., Mayfield, K. K., Hayles, J. A., Middleton, J. E., Ahm, A.-S. C., Johnston, D. T., Caxito, F., et al.: Barium-isotopic constraints on the origin of post-Marinoan barites, *Earth and Planetary Science Letters*, 519, 234–244, 2019.
- Dalai, T. K., Krishnaswami, S., and Sarin, M. M.: Barium in the Yamuna River System in the Himalaya: Sources, fluxes, and its behavior during weathering and transport, *Geochemistry, Geophysics, Geosystems*, 3, 1–23, 2002.
- 970 Dehairs, F., Chesselet, R., and Jedwab, J.: Discrete suspended particles of barite and the barium cycle in the open ocean, *Earth and Planetary Science Letters*, 49, 528–550, 1980.
- Dellinger, M., Gaillardet, J., Bouchez, J., Calmels, D., Galy, V., Hilton, R. G., Louvat, P., and France-Lanord, C.: Lithium isotopes in large rivers reveal the cannibalistic nature of modern continental weathering and erosion, *Earth and Planetary Science Letters*, 401, 359–372, 2014.
- 975 Dellinger, M., Bouchez, J., Gaillardet, J., and Faure, L.: Testing the Steady State Assumption for the Earth’s Surface Denudation Using Li Isotopes in the Amazon Basin, *Procedia Earth and Planetary Science*, 13, 162–168, 2015a.

- Dellinger, M., Gaillardet, J., Bouchez, J., Calmels, D., Louvat, P., Dosseto, A., Gorge, C., Alanoca, L., and Maurice, L.: Riverine Li isotope fractionation in the Amazon River basin controlled by the weathering regimes, *Geochimica et Cosmochimica Acta*, 164, 71–93, 2015b.
- Dellinger, M., Bouchez, J., Gaillardet, J., Faure, L., and Moureau, J.: Tracing weathering regimes using the lithium isotope composition of detrital sediments, *Geology*, 45, 411–414, 2017.
- 980 Dosseto, A., Bourdon, B., Gaillardet, J., Maurice-Bourgoin, L., and Allegre, C. J.: Weathering and transport of sediments in the Bolivian Andes: Time constraints from uranium-series isotopes, *Earth and Planetary Science Letters*, 248, 759–771, 2006.
- Drever, J. and Stillings, L.: The role of organic acids in mineral weathering, *Colloids and Surfaces A: Physicochemical and Engineering Aspects*, 120, 167–181, 1997.
- 985 Edmond, J., Palmer, M., Measures, C., Grant, B., and Stallard, R.: The fluvial geochemistry and denudation rate of the Guayana Shield in Venezuela, Colombia, and Brazil, *Geochimica et Cosmochimica Acta*, 59, 3301–3325, 1995.
- Feeley, K. J. and Silman, M. R.: Land-use and climate change effects on population size and extinction risk of Andean plants, *Global change biology*, 16, 3215–3222, 2010.
- Frappart, F., Papa, F., Güntner, A., Tomasella, J., Pfeffer, J., Ramillien, G., Emilio, T., Schiatti, J., Seoane, L., da Silva Carvalho, J., et al.: The spatio-temporal variability of groundwater storage in the Amazon River Basin, *Advances in water resources*, 124, 41–52, 2019.
- 990 Frings, P. J., Clymans, W., Fontorbe, G., Christina, L., and Conley, D. J.: The continental Si cycle and its impact on the ocean Si isotope budget, *Chemical Geology*, 425, 12–36, 2016.
- Gaillardet, J., Dupré, B., and Allègre, C. J.: A global geochemical mass budget applied to the Congo Basin rivers: erosion rates and continental crust composition, *Geochimica et Cosmochimica Acta*, 59, 3469–3485, 1995.
- 995 Gaillardet, J., Dupre, B., Allegre, C. J., and Négrel, P.: Chemical and physical denudation in the Amazon River Basin, *Chemical geology*, 142, 141–173, 1997.
- Gaillardet, J., Dupré, B., and Allègre, C. J.: Geochemistry of large river suspended sediments: silicate weathering or recycling tracer?, *Geochimica et Cosmochimica Acta*, 63, 4037–4051, 1999a.
- Gaillardet, J., Dupré, B., Louvat, P., and Allegre, C.: Global silicate weathering and CO<sub>2</sub> consumption rates deduced from the chemistry of large rivers, *Chemical geology*, 159, 3–30, 1999b.
- 1000 Gaillardet, J., Viers, J., and Dupré, B.: Trace elements in river waters, *Treatise on geochemistry*, 5, 605, 2003.
- Georg, R. B., Reynolds, B. C., West, A., Burton, K., and Halliday, A. N.: Silicon isotope variations accompanying basalt weathering in Iceland, *Earth and Planetary Science Letters*, 261, 476–490, 2007.
- Gibbs, R. J.: The geochemistry of the Amazon River system: Part I. The factors that control the salinity and the composition and concentration of the suspended solids, *Geological Society of America Bulletin*, 78, 1203–1232, 1967.
- 1005 Gibbs, R. J.: Water chemistry of the Amazon River, *Geochimica et Cosmochimica Acta*, 36, 1061–1066, 1972.
- Gislason, S. R., Arnorsson, S., and Armannsson, H.: Chemical weathering of basalt in Southwest Iceland; effects of runoff, age of rocks and vegetative/glacial cover, *American Journal of Science*, 296, 837–907, 1996.
- Gong, Y., Zeng, Z., Zhou, C., Nan, X., Yu, H., Lu, Y., Li, W., Gou, W., Cheng, W., and Huang, F.: Barium isotopic fractionation in latosol developed from strongly weathered basalt, *Science of The Total Environment*, 687, 1295–1304, 2019.
- 1010 Gou, L.-F., Jin, Z., Galy, A., Gong, Y.-Z., Nan, X.-Y., Jin, C., Wang, X.-D., Bouchez, J., Cai, H.-M., Chen, J.-B., et al.: Seasonal riverine barium isotopic variation in the middle Yellow River: Sources and fractionation, *Earth and Planetary Science Letters*, p. 115990, 2019.
- Griffith, E. M. and Paytan, A.: Barite in the ocean—occurrence, geochemistry and palaeoceanographic applications, *Sedimentology*, 59, 1817–1835, 2012.

- 1015 Grubb, P.: Interpretation of the 'Massenerhebung' effect on tropical mountains, *Nature*, 229, 44–45, 1971.
- Hahm, W. J., Riebe, C. S., Lukens, C. E., and Araki, S.: Bedrock composition regulates mountain ecosystems and landscape evolution, *Proceedings of the National Academy of Sciences*, 111, 3338–3343, 2014.
- Hofmann, A. W.: Chemical differentiation of the Earth: the relationship between mantle, continental crust, and oceanic crust, *Earth and Planetary Science Letters*, 90, 297–314, 1988.
- 1020 Horner, T. J., Kinsley, C. W., and Nielsen, S. G.: Barium-isotopic fractionation in seawater mediated by barite cycling and oceanic circulation, *Earth and Planetary Science Letters*, 430, 511–522, 2015.
- Hsieh, Y.-T. and Henderson, G. M.: Barium stable isotopes in the global ocean: Tracer of Ba inputs and utilization, *Earth and Planetary Science Letters*, 2017.
- Jobbágy, E. G. and Jackson, R. B.: The uplift of soil nutrients by plants: biogeochemical consequences across scales, *Ecology*, 85, 2380–2389, 1025 2004.
- Jung, M., Koirala, S., Weber, U., Ichii, K., Gans, F., Camps-Valls, G., Papale, D., Schwalm, C., Tramontana, G., and Reichstein, M.: The FLUXCOM ensemble of global land-atmosphere energy fluxes, *Scientific data*, 6, 74, 2019.
- Kabata-Pendias, A. and Mukherjee, A. B.: Trace elements from soil to human, Springer Science & Business Media, 2007.
- Lamb, D. T., Matanitobua, V. P., Palanisami, T., Megharaj, M., and Naidu, R.: Bioavailability of barium to plants and invertebrates in soils 1030 contaminated by barite, *Environmental science & technology*, 47, 4670–4676, 2013.
- Langmuir, C. H. and Broecker, W.: How to Build a Habitable Planet: The Story of Earth from the Big Bang to Humankind-Revised and Expanded Edition, Princeton University Press, 2012.
- Lemarchand, D. and Gaillardet, J.: Transient features of the erosion of shales in the Mackenzie basin (Canada), evidences from boron isotopes, *Earth and Planetary Science Letters*, 245, 174–189, 2006.
- 1035 Lemarchand, E., Chabaux, F., Vigier, N., Millot, R., and Pierret, M.-C.: Lithium isotope systematics in a forested granitic catchment (Strengbach, Vosges Mountains, France), *Geochimica et Cosmochimica Acta*, 74, 4612–4628, 2010.
- Li, Y. H.: A compendium of geochemistry: from solar nebula to the human brain, Princeton University Press, 2000.
- Louvat, P. and Allègre, C. J.: Present denudation rates on the island of Reunion determined by river geochemistry: basalt weathering and mass budget between chemical and mechanical erosions, *Geochimica et Cosmochimica Acta*, 61, 3645–3669, 1997.
- 1040 Louvat, P., Gislason, S. R., and Allègre, C. J.: Chemical and mechanical erosion rates in Iceland as deduced from river dissolved and solid material, *American Journal of Science*, 308, 679–726, 2008.
- Marschner, H.: Marschner's mineral nutrition of higher plants, Academic press, 2011.
- Marston, R. A.: Geomorphology and vegetation on hillslopes: interactions, dependencies, and feedback loops, *Geomorphology*, 116, 206–217, 2010.
- 1045 Milliman, J. D. and Farnsworth, K. L.: River discharge to the coastal ocean: a global synthesis, Cambridge University Press, 2013.
- Millot, R., Vigier, N., and Gaillardet, J.: Behaviour of lithium and its isotopes during weathering in the Mackenzie Basin, Canada, *Geochimica et Cosmochimica Acta*, 74, 3897–3912, 2010.
- Miyazaki, T., Kimura, J.-I., and Chang, Q.: Analysis of stable isotope ratios of Ba by double-spike standard-sample bracketing using multiple-collector inductively coupled plasma mass spectrometry, *Journal of Analytical Atomic Spectrometry*, 29, 483–490, 2014.
- 1050 Moquet, J. S., Crave, A., Viers, J., Seyler, P., Armijos, E., Bourrel, L., Chavarri, E., Lagane, C., Laraque, A., Casimiro, W. S. L., et al.: Chemical weathering and atmospheric/soil CO<sub>2</sub> uptake in the Andean and Foreland Amazon basins, *Chemical Geology*, 287, 1–26, 2011.

- Moran-Zuloaga, D., Ditas, F., Walter, D., Saturno, J., Brito, J., Carbone, S., Chi, X., de Angelis, I. H., Baars, H., Godoi, R. H., et al.: Long-term study on coarse mode aerosols in the Amazon rain forest with the frequent intrusion of Saharan dust plumes, *Atmospheric Chemistry and Physics*, 18, 10 055–10 088, 2018.
- 1055 Myrvang, M. B., Hillersøy, M. H., Heim, M., Bleken, M. A., and Gjengedal, E.: Uptake of macro nutrients, barium, and strontium by vegetation from mineral soils on carbonatite and pyroxenite bedrock at the Lillebukt Alkaline Complex on Stjernøy, Northern Norway, *Journal of Plant Nutrition and Soil Science*, 179, 705–716, 2016.
- Nan, X., Wu, F., Zhang, Z., Hou, Z., Huang, F., and Yu, H.: High-precision barium isotope measurements by MC-ICP-MS, *Journal of Analytical Atomic Spectrometry*, 30, 2307–2315, 2015.
- 1060 Nan, X.-Y., Yu, H.-M., Rudnick, R. L., Gaschnig, R. M., Xu, J., Li, W.-Y., Zhang, Q., Jin, Z.-D., Li, X.-H., and Huang, F.: Barium isotopic composition of the upper continental crust, *Geochimica et Cosmochimica Acta*, 233, 33–49, 2018.
- Pogge von Strandmann, P. A. and Henderson, G. M.: The Li isotope response to mountain uplift, *Geology*, 43, 67–70, 2015.
- Porder, S.: How plants enhance weathering and how weathering is important to plants, *Elements: An International Magazine of Mineralogy, Geochemistry, and Petrology*, 15, 241–246, 2019.
- 1065 Porder, S., Vitousek, P. M., Chadwick, O. A., Chamberlain, C. P., and Hilley, G. E.: Uplift, erosion, and phosphorus limitation in terrestrial ecosystems, *Ecosystems*, 10, 159–171, 2007.
- Putzer, H.: The geological evolution of the Amazon Basin and its mineral resources, in: *The Amazon*, pp. 15–46, Springer, 1984.
- Riebe, C. S., Kirchner, J. W., and Finkel, R. C.: Erosional and climatic effects on long-term chemical weathering rates in granitic landscapes spanning diverse climate regimes, *Earth and Planetary Science Letters*, 224, 547–562, 2004.
- 1070 Riebe, C. S., Hahm, W. J., and Brantley, S. L.: Controls on deep critical zone architecture: A historical review and four testable hypotheses, *Earth Surface Processes and Landforms*, 42, 128–156, 2017.
- Rudnick, R. L. and Gao, S.: Composition of the continental crust, *Treatise on geochemistry*, 3, 659, 2003.
- Scalley, T. H., Scatena, F., Moya, S., and Lugo, A.: Long-term dynamics of organic matter and elements exported as coarse particulates from two Caribbean montane watersheds, *Journal of Tropical Ecology*, pp. 127–139, 2012.
- 1075 Schmitt, A.-D., Vigier, N., Lemarchand, D., Millot, R., Stille, P., and Chabaux, F.: Processes controlling the stable isotope compositions of Li, B, Mg and Ca in plants, soils and waters: A review, *Comptes Rendus Geoscience*, 344, 704–722, 2012.
- Selva, E. C., Couto, E. G., Johnson, M. S., and Lehmann, J.: Litterfall production and fluvial export in headwater catchments of the southern Amazon, *Journal of Tropical Ecology*, pp. 329–335, 2007.
- Stallard, R. F.: Tectonic, environmental, and human aspects of weathering and erosion: a global review using a steady-state perspective, *Annual Review of Earth and Planetary Sciences*, 23, 11–39, 1995.
- 1080 Stallard, R. F. and Edmond, J. M.: Geochemistry of the Amazon: 2. The influence of geology and weathering environment on the dissolved load, *Journal of Geophysical Research: Oceans*, 88, 9671–9688, 1983.
- Taylor, S. R. and McLennan, S. M.: The geochemical evolution of the continental crust, *Reviews of Geophysics*, 33, 241–265, 1995.
- Torres, M. A., West, A. J., and Clark, K. E.: Geomorphic regime modulates hydrologic control of chemical weathering in the Andes–Amazon, *Geochimica et Cosmochimica Acta*, 166, 105–128, 2015.
- 1085 Tramontana, G., Jung, M., Camps-Valls, G., Ichii, K., Ráduly, B., Reichstein, M., Schwalm, C. R., Arain, M. A., Cescatti, A., Kiely, G., et al.: Predicting carbon dioxide and energy fluxes across global FLUXNET sites with regression algorithms, *Biogeosciences Discussions*, 2016.

- 1090 Tripler, C. E., Kaushal, S. S., Likens, G. E., and Todd Walter, M.: Patterns in potassium dynamics in forest ecosystems, *Ecology Letters*, 9, 451–466, 2006.
- Trumbore, S. E., Chadwick, O. A., and Amundson, R.: Rapid exchange between soil carbon and atmospheric carbon dioxide driven by temperature change, *Science*, 272, 393–396, 1996.
- Uhlir, D. and von Blanckenburg, F.: How slow rock weathering balances nutrient loss during fast forest floor turnover in montane, temperate forest ecosystems, *Frontiers in Earth Science*, 7, 159, 2019.
- 1095 Uhlir, D., Schuessler, J. A., Bouchez, J., Dixon, J. L., and von Blanckenburg, F.: Quantifying nutrient uptake as driver of rock weathering in forest ecosystems by magnesium stable isotopes, *Biogeosciences*, 14, 3111, 2017.
- Vaganov, E. A., Grachev, A. M., Shishov, V. V., Panyushkina, I. P., Leavitt, S. W., Knorre, A. A., Chebykin, E. P., and Menyailo, O. V.: Elemental composition of tree rings: A new perspective in biogeochemistry, in: *Doklady Biological Sciences*, vol. 453, pp. 375–379, Springer, 2013.
- 1100 Van Zuilen, K., Müller, T., Nägler, T. F., Dietzel, M., and Küsters, T.: Experimental determination of barium isotope fractionation during diffusion and adsorption processes at low temperatures, *Geochimica et cosmochimica acta*, 186, 226–241, 2016a.
- Van Zuilen, K., Nägler, T. F., and Bullen, T. D.: Barium Isotopic Compositions of Geological Reference Materials, *Geostandards and Geo-analytical Research*, 40, 543–558, 2016b.
- Viers, J., Oliva, P., Dandurand, J.-L., Dupré, B., and Gaillardet, J.: Chemical weathering rates, CO<sub>2</sub> consumption, and control parameters deduced from the chemical composition of rivers, 2014.
- 1105 Vitousek, P., Chadwick, O., Matson, P., Allison, S., Derry, L., Kettley, L., Luers, A., Mecking, E., Monastra, V., and Porder, S.: Erosion and the rejuvenation of weathering-derived nutrient supply in an old tropical landscape, *Ecosystems*, 6, 762–772, 2003.
- Von Allmen, K., Böttcher, M. E., Samankassou, E., and Nägler, T. F.: Barium isotope fractionation in the global barium cycle: First evidence from barium minerals and precipitation experiments, *Chemical geology*, 277, 70–77, 2010.
- 1110 von Blanckenburg, F.: The control mechanisms of erosion and weathering at basin scale from cosmogenic nuclides in river sediment, *Earth and Planetary Science Letters*, 237, 462–479, 2005.
- von Strandmann, P. A. P., Opfergelt, S., Lai, Y.-J., Sigfússon, B., Gislason, S. R., and Burton, K. W.: Lithium, magnesium and silicon isotope behaviour accompanying weathering in a basaltic soil and pore water profile in Iceland, *Earth and Planetary Science Letters*, 339, 11–23, 2012.
- 1115 West, A. J., Galy, A., and Bickle, M.: Tectonic and climatic controls on silicate weathering, *Earth and Planetary Science Letters*, 235, 211–228, 2005.
- Wilcke, W., Velescu, A., Leimer, S., Bigalke, M., Boy, J., and Valarezo, C.: Biological versus geochemical control and environmental change drivers of the base metal budgets of a tropical montane forest in Ecuador during 15 years, *Biogeochemistry*, 136, 167–189, 2017.
- 1120 Wittmann, H., von Blanckenburg, F., Maurice, L., Guyot, J.-L., Filizola, N., and Kubik, P. W.: Sediment production and delivery in the Amazon River basin quantified by in situ-produced cosmogenic nuclides and recent river loads, *Bulletin*, 123, 934–950, 2011.