

# Leaf-scale quantification of the effect of photosynthetic gas exchange on $\Delta^{17}\text{O}$ of atmospheric $\text{CO}_2$

Getachew Agmuas Adnew<sup>1</sup>, Thijs L. Pons<sup>2</sup>, Gerbrand Koren<sup>3</sup>, Wouter Peters<sup>3,4</sup>, Thomas Röckmann<sup>1</sup>

<sup>1</sup>Institute for Marine and Atmospheric research Utrecht (IMAU), Utrecht University, The Netherlands

5 <sup>2</sup>Institute of Environmental Biology, Utrecht University, The Netherlands

<sup>3</sup>Department of Meteorology and Air Quality, Wageningen University, The Netherlands

<sup>4</sup>Centre for Isotope Research, University of Groningen, The Netherlands

*Correspondence to:* Getachew Agmuas Adnew (g.a.adnew@uu.nl)

10

## Abstract

Understanding the processes that affect the triple oxygen isotope composition of atmospheric  $\text{CO}_2$  during gas exchange can help constrain the interaction and fluxes between the atmosphere and the biosphere. We conducted leaf cuvette experiments under controlled conditions, using three plant species. The experiments were conducted at two different light intensities and using  $\text{CO}_2$  with different  $\Delta^{17}\text{O}$ . We directly quantify for the first time the effect of photosynthesis on  $\Delta^{17}\text{O}$  of atmospheric  $\text{CO}_2$ . Our results demonstrate the established theory for  $\delta^{18}\text{O}$  is applicable to  $\Delta^{17}\text{O}$ - $\text{CO}_2$  at leaf-level and we confirm the two key factors determine the effect of photosynthetic gas exchange on the  $\Delta^{17}\text{O}$  of atmospheric  $\text{CO}_2$ . The relative difference between  $\Delta^{17}\text{O}$  of the  $\text{CO}_2$  entering the leaf and the  $\text{CO}_2$  in equilibrium with leaf water, and the back-diffusion flux of  $\text{CO}_2$  from the leaf to the atmosphere, which can be quantified by the  $c_m/c_a$  ratio where  $c_a$  is the  $\text{CO}_2$  mole fraction in the surrounding air and  $c_m$  the one at the site of oxygen isotope exchange between  $\text{CO}_2$  and  $\text{H}_2\text{O}$ . At low  $c_m/c_a$  ratio the discrimination is governed mainly by diffusion into the leaf, and at high  $c_m/c_a$  ratio by back-diffusion of  $\text{CO}_2$  that has equilibrated with the leaf water. Plants with a higher  $c_m/c_a$  ratio modify the  $\Delta^{17}\text{O}$  of atmospheric  $\text{CO}_2$  more strongly than plants with a lower  $c_m/c_a$  ratio. Based on the leaf cuvette experiments, the global value for discrimination against  $\Delta^{17}\text{O}$  of atmospheric  $\text{CO}_2$  during photosynthetic gas exchange is estimated to be  $-0.57 \pm 0.14$  ‰ using  $c_m/c_a$  values of 0.3 and 0.7 for  $\text{C}_4$  and  $\text{C}_3$  plants, respectively. The main uncertainties in this global estimate arise from variation in  $c_m/c_a$  ratios among plants and growth conditions.

30

35

## 1. Introduction

40 Stable isotope measurements of CO<sub>2</sub> provide important information on the magnitude of the CO<sub>2</sub> fluxes  
between atmosphere and biosphere, which are the largest components of the global carbon cycle  
(Farquhar et al., 1989; 1993; Ciais et al., 1997a; 1997b; Flanagan and Ehleringer, 1998; Yakir and  
Sternberg, 2000; Gillon and Yakir, 2001; Cuntz et al., 2003a; 2003b). A better understanding of the  
45 terrestrial carbon cycle is essential for predicting future climate and atmospheric CO<sub>2</sub> mole fractions  
(Booth et al., 2012). Gross primary productivity (GPP), the total carbon dioxide uptake by vegetation  
during photosynthesis, can only be determined indirectly and remains poorly constrained (Cuntz, 2011;  
Welp et al., 2011). For example, Beer et al. (2010) estimated global GPP to be 102-135 PgC yr<sup>-1</sup> (85 %  
confidence interval, CI) using machine learning techniques by extrapolating from a database of eddy-  
50 covariance measurements of CO<sub>2</sub> fluxes. This estimate has since then been widely used as target for  
terrestrial vegetation models (Sitch et al., 2015), and replicated based on cross-consistency checks with  
atmospheric inversions, sun-induced fluorescence (SIF) and global vegetation models (Jung et al., 2020).  
As an alternative, Welp et al. (2011) estimated global GPP to be 150-175 PgC yr<sup>-1</sup> using variations in  
δ<sup>18</sup>O of atmospheric CO<sub>2</sub> after the 1997/98 El Nino event; see equation 1 for definition of the δ value.

55 The concept behind the latter study was that atmospheric CO<sub>2</sub> exchanges oxygen isotopes with leaf and  
soil water, and this isotope exchange mostly determines the observed variations in δ<sup>18</sup>O of CO<sub>2</sub> (Francey  
and Tans, 1987; Yakir, 1998). Following the 97/98 ENSO event, the anomalous δ<sup>18</sup>O signature imposed  
on tropical leaf and soil waters was transferred to atmospheric CO<sub>2</sub>, before slowly disappearing as a  
function of the lifetime of atmospheric CO<sub>2</sub>. This in turn is governed by the land vegetation uptake of  
60 CO<sub>2</sub> during photosynthesis, as well as soil invasion of CO<sub>2</sub> (Miller et al., 1999; Wingate et al., 2009). For  
the photosynthesis term, the equilibration of CO<sub>2</sub> with water is an uncertain parameter in this calculation,  
partly because the δ<sup>18</sup>O of water at the site of isotope exchange in the leaf is not well defined. Importantly,  
a significant δ<sup>18</sup>O variation can occur in leaves due to the preferential evaporation of H<sub>2</sub><sup>16</sup>O relative to  
H<sub>2</sub><sup>18</sup>O (Gan et al., 2002; Farquhar and Gan, 2003; Gan et al., 2003; Cernusak et al., 2016), which induces  
65 a considerable uncertainty in estimating δ<sup>18</sup>O of CO<sub>2</sub>. Similar considerations for the transfer of the δ<sup>18</sup>O  
signature of precipitation into the soils, and then up through the roots, stems, and leaves make <sup>18</sup>O of CO<sub>2</sub>  
a challenging measurement to interpret (Peylin et al., 1999; Cuntz et al., 2003a; 2003b).

Classical isotope theory posits that oxygen isotope distributions are modified in a mass-dependent way.  
This means that the <sup>17</sup>O/<sup>16</sup>O ratio changes by approximately half of the corresponding change in <sup>18</sup>O/<sup>16</sup>O  
70 (equation 2), and it applies to the processes involved in gas exchange between atmosphere and plants.  
However, in 1983 Thiemens and co-workers (Heidenreich and Thiemens, 1983; Thiemens, 1983;  
Heidenreich and Thiemens, 1986) reported a deviation from mass-dependent isotope fractionation in  
ozone (O<sub>3</sub>) formation called mass-independent isotope fractionation (Δ<sup>17</sup>O, equation 3). In the  
stratosphere, the Δ<sup>17</sup>O of O<sub>3</sub> is transferred to CO<sub>2</sub> via isotope exchange of CO<sub>2</sub> with O(<sup>1</sup>D) produced from  
75 O<sub>3</sub> photolysis (Yung et al., 1991; Yung et al., 1997; Shaheen et al., 2007), which results a large Δ<sup>17</sup>O in  
stratospheric CO<sub>2</sub> (Thiemens et al., 1991; 1995; Lyons, 2001; Lämmerzahl et al., 2002; Thiemens, 2006;  
Kawagucci et al., 2008; Wiegel et al., 2013).

80 Once  $\Delta^{17}\text{O}$  has been created in stratospheric  $\text{CO}_2$ , the only process that modify its signal is isotope exchange with leaf water, soil water and ocean water at the Earth's surface, after  $\text{CO}_2$  has re-entered the troposphere (Boering, 2004; Thiemens et al., 2014; Liang and Mahata, 2015; Hofmann et al., 2017). Isotope exchange with leaf water is more efficient relative to ocean water due to the presence of the enzyme carbonic anhydrase (CA), which effectively catalyzes the conversion of  $\text{CO}_2$  and  $\text{H}_2\text{O}$  to  $\text{HCO}_3^-$  and  $\text{H}^+$  and vice versa (Francey and Tans, 1987; Friedli et al., 1987; Badger and Price, 1994; Gillon and 85 Yakir, 2001). The isotope exchange in the atmosphere is negligible due to lower liquid water content, lower residence time and the absence of carbonic anhydrase (Mills and Urey, 1940; Miller et al., 1971; Johnson, 1982; Silverman, 1982; Francey and Tans, 1987).

$\Delta^{17}\text{O}$  of  $\text{CO}_2$  has been suggested as additional independent tracer for constraining global GPP (Hoag et al., 2005; Thiemens et al., 2013; Hofmann et al., 2017; Liang et al., 2017b; Koren et al., 2019), because 90 the processes involved in plant-atmosphere gas exchange are all mass-dependent. Therefore,  $\Delta^{17}\text{O}$  at the  $\text{CO}_2$ - $\text{H}_2\text{O}$  exchange site in the leaf will vary much less than  $\delta^{18}\text{O}$ . Nevertheless, mass-dependent isotope fractionation processes with slightly different three-isotope fractionation slopes are involved, which have been precisely established in the past years. Figure 1 shows how the different processes affect  $\Delta^{17}\text{O}$  of the  $\text{H}_2\text{O}$  and  $\text{CO}_2$  reservoirs involved. The triple isotope slope of oxygen in meteoric waters is taken as 95 reference slope,  $\lambda_{Ref}=0.528$  (Meijer and Li, 1998; Barkan and Luz, 2007; Landais et al., 2008; Luz and Barkan, 2010; Uemura et al., 2010) and we assume that soil water is similar to meteoric water. Due to transpiration and diffusion in the leaf,  $\Delta^{17}\text{O}$  of leaf water gets modified following a humidity dependent three-isotope slope  $\theta_{trans}=0.522-0.008 \times h$  (Landais et al., 2006). Exchange of oxygen isotopes between leaf water and  $\text{CO}_2$  follows  $\lambda_{\text{CO}_2\text{-H}_2\text{O}}=0.5229$  (Barkan and Luz, 2012) which determines the  $\Delta^{17}\text{O}$  of  $\text{CO}_2$  100 inside the leaf at the  $\text{CO}_2$ - $\text{H}_2\text{O}$  exchange site. Finally, the  $\Delta^{17}\text{O}$  of the  $\text{CO}_2$  is modified when  $\text{CO}_2$  diffuses into and out of the leaf with  $\lambda_{diff}=0.509$  (Young et al., 2002).

In the first box model study of Hoag et al. (2005), the small deviations in  $\Delta^{17}\text{O}$  of  $\text{CO}_2$  due to differences in three-isotope slopes were neglected and exchange with water was assumed to reset  $\Delta^{17}\text{O}$  to 0. Hofmann et al. (2017) included the different isotope effects shown in Figure 1 in their box model. Koren et al. 105 (2019) incorporated all the physico-chemical processes affecting  $\Delta^{17}\text{O}$  of  $\text{CO}_2$  in a 3D atmospheric model and investigated the spatiotemporal variability of  $\Delta^{17}\text{O}$  and its use as tracer for GPP. Using these and other similar models, numerous measurements of  $\Delta^{17}\text{O}$  in atmospheric  $\text{CO}_2$  from different locations have been performed and used to estimate GPP (Liang et al., 2006; Barkan and Luz, 2012; Thiemens et al., 2014; Liang and Mahata, 2015; Laskar et al., 2016; Hofmann et al., 2017). The three-isotope slopes of 110 the processes involved in the gas exchange (Figure 1) have been precisely determined in idealized experiments. In the advanced models mentioned above it is assumed that when all the pieces are put together they results in a realistic overall modification of  $\Delta^{17}\text{O}$  of  $\text{CO}_2$  in the atmosphere surrounding the leaf. However, this has not been confirmed by measurements previously.

115 In this study we report the effect of photosynthesis on  $\Delta^{17}\text{O}$  of  $\text{CO}_2$  in the surrounding air at the leaf scale.  
 We measured  $\Delta^{17}\text{O}$  of  $\text{CO}_2$  entering and leaving a leaf cuvette to calculate the isotopic fractionation  
 associated with photosynthesis for three species that are representative for three different biomes. The  
 fast-growing annual herbaceous  $\text{C}_3$  species *Helianthus annuus* (sunflower) has a high photosynthetic  
 capacity ( $A_N$ ) and high stomatal conductance ( $g_s$ ) and is representative for temperate and tropical crops  
 120 (Fredeen et al., 1991). The slower growing perennial evergreen  $\text{C}_3$  species *Hedera hybernica* (ivy) is  
 representative of forests and other woody vegetation and stress subjected habitats (Pons et al., 2009). The  
 fast-growing, agronomically important crop *Zea mays* (maize) is an herbaceous annual  $\text{C}_4$  species with a  
 high  $A_N$  and a low  $g_s$ , typical for savanna type vegetation (Weijde et al., 2013). The mole fraction of  $\text{CO}_2$   
 at the  $\text{CO}_2$ - $\text{H}_2\text{O}$  exchange site ( $c_m$ ) is an important parameter to determine the effect of photosynthesis on  
 125  $\Delta^{17}\text{O}$  of  $\text{CO}_2$ . In  $\text{C}_3$  plants, the  $\text{CO}_2$ - $\text{H}_2\text{O}$  exchange can occur anywhere between the plasma membrane  
 and the chloroplast since the catalyzing enzyme CA has been found in the chloroplast, cytosol,  
 mitochondria and plasma membrane (Fabre et al., 2007; DiMario et al., 2016). For  $\text{C}_4$  plants, CA is mainly  
 found in the cytosol and the  $\text{CO}_2$ - $\text{H}_2\text{O}$  exchange occurs there (Badger and Price, 1994). In our  
 experiments, sunflower and ivy are used to cover the wide  $c_m/c_a$  ratio range among  $\text{C}_3$  plants and maize  
 130 represents the  $c_m/c_a$  ratio for the  $\text{C}_4$  plants. Using our results from the leaf scale experiments, we estimated  
 the effect of terrestrial vegetation on  $\Delta^{17}\text{O}$  of  $\text{CO}_2$  in the global atmosphere.

## 1. Theory

### 135 1.1. Notation and definition of $\delta$ values

Isotopic composition is expressed as the deviation of the heavy to light isotope ratio in a sample relative  
 to a reference ratio and it is denoted as  $\delta$ , expressed in per mill (‰). In the case of oxygen isotopes, the  
 isotope ratios are  $^{18}\text{R} = [^{18}\text{O}]/[^{16}\text{O}]$  and  $^{17}\text{R} = [^{17}\text{O}]/[^{16}\text{O}]$  and the reference material is Vienna Standard  
 Mean Ocean Water (VSMOW):

$$140 \quad \delta^n\text{O} = \frac{^n\text{R}_{\text{sample}}}{^n\text{R}_{\text{VSMOW}}} - 1, \quad n \text{ refers to } 17 \text{ or } 18 \quad (1)$$

For most processes, isotope fractionation depends on mass, and therefore the fractionation against  $^{17}\text{O}$  is  
 approximately half of the fractionation against  $^{18}\text{O}$  (equation 3).

$$\ln(\delta^{17}\text{O} + 1) = \lambda \times \ln(\delta^{18}\text{O} + 1) \quad (2)$$

145 The mass-dependent isotope fractionation factor  $\lambda$  ranges from 0.5 to 0.5305 for different molecules and  
 process (Matsuhisa et al., 1978; Thieme, 1999; Young et al., 2002; Cao and Liu, 2011).  $\Delta^{17}\text{O}$  is used to  
 quantify the degree of deviation from equation (2) (see equation 3). Note that  $\Delta^{17}\text{O}$  changes not only by  
 mass-independent isotope fractionation processes, but also by mass-dependent isotope fractionation  
 processes with a different  $\lambda$  value from the one used in the definition of  $\Delta^{17}\text{O}$  (Barkan and Luz, 2005;  
 150 Landais et al., 2006; 2008; Luz and Barkan, 2010; Barkan and Luz, 2011; Pack and Herwartz, 2014).

$$\Delta^{17}\text{O} = \ln(\delta^{17}\text{O} + 1) - \lambda \times \ln(\delta^{18}\text{O} + 1) \quad (3)$$

The choice of  $\lambda$  is in principle arbitrary and in this study, we use  $\lambda = 0.528$ , which was established for meteoric waters (Meijer and Li, 1998; Landais et al., 2008; Brand et al., 2010; Luz and Barkan, 2010; Barkan and Luz, 2012; Sharp et al., 2018). Equation 3 can be linearized to (Miller, 2002)  $\Delta^{17}\text{O} = \delta^{17}\text{O} - \lambda \times \delta^{18}\text{O}$ , but this approximation causes an error that increases with  $\delta^{18}\text{O}$  (Miller, 2002; Bao et al., 2016).

## 1.2. Discrimination against $\Delta^{17}\text{O}$ of $\text{CO}_2$

The overall isotope fractionation associated with the photosynthesis of  $\text{CO}_2$  is commonly quantified using the term discrimination as described in (Farquhar and Richards, 1984; Farquhar et al., 1989; Farquhar and Lloyd, 1993). We use the symbol  $\Delta_A$  for discrimination due to assimilation in this manuscript since the commonly used  $\Delta$  is already used for the definition of  $\Delta^{17}\text{O}$  (see equation 3).  $\Delta_A$  quantifies the enrichment or depletion of carbon and oxygen isotopes of  $\text{CO}_2$  in the surrounding atmosphere relative to the  $\text{CO}_2$  that is assimilated (Farquhar and Richards, 1984). It can be calculated from the isotopic composition of the  $\text{CO}_2$  entering and leaving the leaf cuvette (Evans et al., 1986; Gillon and Yakir, 2000a; Barbour et al., 2016) as:

$$\Delta_A^{n\text{O}} = \frac{n_{\text{R}_a}}{n_{\text{R}_e}} - 1 = \frac{\delta^{n\text{O}_a} - \delta^{n\text{O}_e}}{1 + \delta^{n\text{O}_a}} = \frac{\zeta \times (\delta^{n\text{O}_a} - \delta^{n\text{O}_e})}{1 + \delta^{n\text{O}_a} - \zeta \times (\delta^{n\text{O}_a} - \delta^{n\text{O}_e})} \quad (4)$$

where the indices  $e$ ,  $a$  and  $A$  refer to  $\text{CO}_2$  entering and leaving the cuvette and being assimilated, respectively.  $\zeta = \frac{c_e}{c_e - c_a}$ , where  $c_e$  and  $c_a$  are the mole fractions of  $\text{CO}_2$  entering and leaving the cuvette. For quantifying the effect of photosynthesis on  $\Delta^{17}\text{O}$  in our experiments, the  $\Delta_A \Delta^{17}\text{O}$  is calculated from  $\Delta_A^{17}\text{O}$  and  $\Delta_A^{18}\text{O}$  using the three-isotope slope  $\lambda_{\text{RL}} = 0.528$ , similar to equation 3. In previous studies slightly, different formulations have been used to define the effect of photosynthesis on  $\Delta^{17}\text{O}$ , and a comparison of the different definitions is provided in the supplementary material (equation S37-S40).

It is important to note that when the logarithmic definition of  $\Delta^{17}\text{O}$  or  $\Delta_A \Delta^{17}\text{O}$  is used, values are not additive (Kaiser et al., 2004). In linear calculations, the error gets larger when the relative difference in  $\delta^{18}\text{O}$  between the two  $\text{CO}_2$  gases increases regardless of the  $\Delta^{17}\text{O}$  of the individual  $\text{CO}_2$  gases (Figure S1). Therefore,  $\Delta_A \Delta^{17}\text{O}$  values have to be calculated from the individual  $\Delta_A^{17}\text{O}$  and  $\Delta_A^{18}\text{O}$  values, and not by linear combinations of the  $\Delta^{17}\text{O}$  of air entering and leaving a plant chamber.

## 2. Materials and methods

### 2.1. Plant material and growth conditions

Sunflower (*Helianthus annuus* L. cv “sunny”) was grown from seeds in 0.6 L pots with potting soil (Primasta, the Netherlands) for about four weeks. All leaves appearing above the first leaf pair were removed to avoid shading. Established juvenile ivy (*Hedera hybernica* L.) plants were pruned and planted

in 6 L pots for 6 weeks. Ivy leaves that had developed and matured were used for the experiments. Maize (*Z. mays* L. cv “saccharate”) was grown from seed in 1.6 L pots for at least 7 weeks. For maize, the 4<sup>th</sup> or higher leaf number was used for the experiments when mature. A section of the leaf at about 1/3 from the tip was inserted in the leaf cuvette. They were placed on a sub-irrigation system that provided water during the growth period in a controlled environment growth chamber, air temperature 20°C, relative humidity 70 % and CO<sub>2</sub> mole fraction of about 400 ppm. The photosynthetic photon flux density (PPFD) was about 300  $\mu\text{mol m}^{-2} \text{s}^{-1}$  during a daily photoperiod of 16 hours measured with a PPF meter (Licor LI-250A, Li-Cor Inc, Nebraska, USA).

195

## 2.2. Gas exchange experiments

Gas exchange experiments were performed in an open system where a controlled flow of air enters and leaves the leaf cuvette similar to the setup used by (Pons and Welschen, 2002). A schematic for the gas exchange experimental setup is shown in Figure 2. The leaf cuvette had dimensions of 7 x 7 x 7 cm<sup>3</sup> (lxwxh) and the top part of the cuvette was transparent. The temperature of the leaf was measured with a K type thermocouple. The leaf chamber temperature was controlled by a temperature-controlled water bath kept at 20°C (Tamson TLC 3, The Netherlands). A halogen lamp (PRADOVIT 253, ERNST LEITZ WETZLAR GMBH, Germany) in a slide projector was used as a light source. Infrared was excluded by reflection from a cold mirror. The light intensity was varied with spectrally neutral filters (PRADOVIT 253, ERNST LEITZ WETZLAR GMBH, Germany).

200

205

The CO<sub>2</sub> mole fraction of the incoming and outgoing air was measured with an infrared gas analyzer (IRGA, model LI-6262, LI-COR Inc., Nebraska, USA). The isotopic composition and mole fraction of the incoming and outgoing water vapor were measured with a triple water vapor isotope analyzer (WVIA, model 911-0034, Los Gatos Research, USA). Compressed air (ambient outside air without drying) was passed through soda lime to scrub the CO<sub>2</sub>. The CO<sub>2</sub> free air could be humidified depending on the experiment conditions (see Figure 2). The humidity of the inlet air was monitored continuously with a dewpoint meter (HYGRO-M1, General Eastern, Watertown, MA, USA). Pure CO<sub>2</sub> (either normal CO<sub>2</sub> or isotopically enriched CO<sub>2</sub>) was mixed with the incoming air to produce a CO<sub>2</sub> mole fraction of 500 ppm. The isotopically enriched CO<sub>2</sub> was prepared by photochemical isotope exchange between CO<sub>2</sub> and O<sub>2</sub> under UV irradiation (Adnew et al., 2019).

210

215

An attached leaf or part of it was inserted into the cuvette, the composition of the inlet air was measured, and both IRGA and WVIA were switched to measure the outlet air. Based on the CO<sub>2</sub> mole fraction of the outgoing air the flow rate of the incoming air to the cuvette was adjusted to establish a drawdown of 100 ppm CO<sub>2</sub> due to photosynthesis in the plant chamber. The water vapor content entering the cuvette was adjusted depending on the transpiration rate relative to CO<sub>2</sub> uptake to avoid condensation (Figure 2). The outgoing air was measured continuously until a steady state was reached for CO<sub>2</sub> and H<sub>2</sub>O mole fractions and  $\delta\text{D}$  and  $\delta^{18}\text{O}$  of the water vapor. After a steady state was established, the air was directed to the sampling flask while the IGRA and WVIA were switched back to measure the inlet air. The air passed through a Mg(ClO<sub>4</sub>)<sub>2</sub> dryer before entering the sampling flask.

220

225

After sampling, the leaf area inside the cuvette was measured with a LI-3100C area meter (Li-COR, Inc. USA). Immediately afterward, the leaf was placed in a leak tight 9 mL glass vial and kept in a freezer at -20°C until leaf water extraction.

### 2.3. Calibration of the Water Vapor Isotope Analyzer (WVIA) and leaf water analysis

The WVIA was calibrated using five water standards provided by IAEA (Wassenaar et al., 2018) for both  $\delta^{18}\text{O}$  and  $\delta\text{D}$  (Figure S2). We did not calibrate the WVIA for  $\delta^{17}\text{O}$ , so the  $\delta^{17}\text{O}$  data are not used in the quantitative evaluation. The isotopic composition of the water standards ranged from -50.93 to 3.64‰ and -396.98 ‰ to 25.44 ‰ for  $\delta\text{D}$  and  $\delta^{18}\text{O}$ , respectively. The detailed characterization and calibration of the water WVIA is provided in the supplementary material (Figure S2 to S4).

Leaf water was extracted by cryogenic vacuum distillation for 4 h at 60°C following a well-established procedure as shown in Figure S5 (Wang and Yakir, 2000; Landais et al., 2006; West et al., 2006). Details are provided in the supplementary material. The  $\delta^{17}\text{O}$  and  $\delta^{18}\text{O}$  of leaf water was determined at the Laboratoire des Sciences du Climat et de l'Environnement laboratory using a fluorination technique as described in (Barkan and Luz, 2005; Landais et al., 2006; 2008).

### 2.4. Carbon dioxide extraction and isotope analysis

$\text{CO}_2$  was extracted from the air samples in a system made from electropolished stainless steel (Supplementary Figure S6). Our system used four commercial traps (MassTech, Bremen, Germany). The first two traps were operated at dry ice temperature (-78°C) to remove moisture and some organics. The other two traps were operated at liquid nitrogen temperature (-196°C) to trap  $\text{CO}_2$ . The flow rate during extraction was 55 mL  $\text{min}^{-1}$ , controlled by a mass flow controller (Brooks Instruments, Holland). The reproducibility of the extraction system was 0.030 ‰ for  $\delta^{18}\text{O}$  and 0.007 ‰ for  $\delta^{13}\text{C}$  determined on 14 extractions (1 $\sigma$  standard deviation, Supplementary Table S1).

The  $\Delta^{17}\text{O}$  of  $\text{CO}_2$  was determined using the  $\text{CO}_2$ - $\text{O}_2$  exchange method (Mahata et al., 2013; Barkan et al., 2015; Adnew et al., 2019). The  $\text{CO}_2$ - $\text{O}_2$  exchange system used at Utrecht University is described in (Adnew et al., 2019). In short, equal amounts of  $\text{CO}_2$  and  $\text{O}_2$  were mixed in a quartz reactor containing a platinum sponge catalyst and heated at 750°C for 2hrs. After isotope equilibration, the  $\text{CO}_2$  was trapped at liquid nitrogen temperature, while the  $\text{O}_2$  was collected with 1 pellet of 5Å molecular sieve (1.6 mm, Sigma Aldrich, USA) at liquid nitrogen temperature. The isotopic composition of the isotopically equilibrated  $\text{O}_2$  was measured with a Delta<sup>plus</sup>XL isotope ratio mass spectrometer in dual inlet mode with reference to a pure  $\text{O}_2$  calibration gas that has been assigned values of  $\delta^{17}\text{O} = 9.254$  ‰ and  $\delta^{18}\text{O} = 18.542$  ‰ by E. Barkan at the Hebrew University of Jerusalem. The reproducibility of the  $\Delta^{17}\text{O}$  measurement was better than 0.01 ‰ (Supplementary Table S1).

### 2.5. Leaf cuvette model

265 We used a simple leaf cuvette model to evaluate the dependence of  $\Delta_A \Delta^{17}\text{O}$  on key parameters. In this  
model, the leaf is partitioned into three different compartments: the intercellular air space, the mesophyll  
cell, and the chloroplast. In the leaf cuvette model, we used a 100 ppm drawdown of  $\text{CO}_2$ , similar to the  
leaf exchange experiments, i.e., the  $\text{CO}_2$  mole fraction decreases from 500 ppm in the entering air ( $c_e$ ) to  
400 ppm in the outgoing air ( $c_o$ ), which is identical to the air surrounding the leaf ( $c_a$ ) as a result of  
thorough mixing in the cuvette. The assimilation rate is set to  $20.0 \mu\text{mol m}^{-2}\text{s}^{-1}$ . The leaf area and flowrate  
270 of air are set to  $30 \text{ cm}^2$  and  $0.7 \text{ L min}^{-1}$ , respectively. The isotope composition of leaf water at the site  
where the  $\text{H}_2\text{O}-\text{CO}_2$  exchange occurs is  $\delta^{17}\text{O} = 5.39 \text{ ‰}$  and  $\delta^{18}\text{O} = 10.648 \text{ ‰}$ , which is the mean of the  
measured  $\delta^{17}\text{O}$  and  $\delta^{18}\text{O}$  values of bulk leaf water in our experiments. The leaf water temperature is set  
to  $22^\circ\text{C}$  (similar to the experiment). In the model, the  $\delta^{18}\text{O}$  of the  $\text{CO}_2$  entering the cuvette is set to  $30.47 \text{ ‰}$   
for all the simulations, as in the normal  $\text{CO}_2$  experiments, but the assigned  $\Delta^{17}\text{O}$  values ranges from -  
275  $0.5 \text{ ‰}$  to  $0.5 \text{ ‰}$  which encompasses both the stratospheric intrusion and combustion components. The  
corresponding  $\delta^{17}\text{O}$  of the  $\text{CO}_2$  entering the cuvette is calculated from the assigned  $\delta^{18}\text{O}$  value ( $30.47 \text{ ‰}$ )  
and  $\Delta^{17}\text{O}$  values ( $-0.5 \text{ ‰}$  to  $0.5 \text{ ‰}$ ). For the calculations with this model, we assumed an infinite boundary  
layer conductance. The leaf cuvette model is illustrated in the supplementary material (Figure S7) and the  
detailed code and description is available at [https://git.wur.nl/leaf\\_model/D17O](https://git.wur.nl/leaf_model/D17O).

280

### 3. Results

#### 3.1. Gas exchange parameters

285 Table 1 summarizes the isotopic composition and mole fraction of the  $\text{CO}_2$  used in this study for  
sunflower, ivy and maize. The  $\Delta^{17}\text{O}$  of  $\text{CO}_2$  used in this study varies from  $-0.215 \text{ ‰}$  to  $0.44 \text{ ‰}$  while the  
 $\delta^{18}\text{O}$  value is close to  $30 \text{ ‰}$  for all the experiments. For all the experiments, the mole fraction of  $\text{CO}_2$   
entering the leaf ( $c_a$ ) is 400 ppm whereas the mole fraction of the  $\text{CO}_2$  in the intercellular air space ( $c_i$ ), at  
the  $\text{CO}_2-\text{H}_2\text{O}$  exchange site ( $c_m$ ) and in the chloroplast ( $c_c$ ) varies depending on the assimilation rate and  
290 metabolism type of the plants. Estimating the mesophyll conductance is described in the companion  
paper. A detailed description for estimating  $c_m$  and  $c_c$  is provided in the supplementary material. A list of  
variables and parameters used in this study are summarized in Table 2.

#### 3.2. Discrimination against $^{18}\text{O}$ of $\text{CO}_2$

295

Figure 3a shows discrimination against  $^{18}\text{O}$  associated with photosynthesis ( $\Delta_A^{18}\text{O}$ ) for sunflower, ivy,  
and maize as a function of the  $c_m/c_a$  ratio.  $\Delta_A^{18}\text{O}$  varies with  $c_m/c_a$ , as found in previous studies (Gillon  
and Yakir, 2000a; Barbour et al., 2016). For sunflower, we observe  $\Delta_A^{18}\text{O}$  values between  $29 \text{ ‰}$  and  $64 \text{ ‰}$   
for  $c_m/c_a$  between 0.54 and 0.86. Ivy shows a relatively little variation of  $\Delta_A^{18}\text{O}$  around a mean of  $22 \text{ ‰}$   
300  $\text{‰}$  for  $c_m/c_a$  between 0.48 and 0.58. For maize,  $\Delta_A^{18}\text{O}$  is lower than for the  $\text{C}_3$  plants measured in this  
study, with values between  $10 \text{ ‰}$  and  $20 \text{ ‰}$  for  $c_m/c_a$  between 0.15 and 0.37.



For sunflower changing the irradiance from 300  $\mu\text{mol m}^{-2}\text{s}^{-1}$  (low light, hereafter LL) to 1200  $\mu\text{mol m}^{-2}\text{s}^{-1}$  (high light, hereafter HL) leads to a clear decrease in  $\Delta_A^{18}\text{O}$  (average 22 ‰). For maize, the  $\Delta_A^{18}\text{O}$  change is only 4.4 ‰ on average. For ivy, changing the light intensity does not significantly change the observed  $\Delta_A^{18}\text{O}$ . The solid lines in Figure 3a show results of leaf cuvette model calculations, where the dependence of  $\Delta_A^{18}\text{O}$  on  $c_m/c_a$  is explored for a set of calculations with otherwise fixed parameters. The model agrees well with the experimental results except for ivy, where the model overestimates the discrimination.

310

### 3.3. Discrimination against $\Delta^{17}\text{O}$ of $\text{CO}_2$

The discrimination of photosynthesis against  $\Delta^{17}\text{O}$  of  $\text{CO}_2$  ( $\Delta_A\Delta^{17}\text{O}$ ) is shown in Figure 3b.  $\Delta_A\Delta^{17}\text{O}$  is negative for all experiments and it depends strongly on the  $c_m/c_a$  ratio and  $|\Delta_A\Delta^{17}\text{O}|$  increases with  $c_m/c_a$  ratio. For instance, for  $\Delta^{17}\text{O}$  of  $\text{CO}_2$  entering the cuvette of -0.215 ‰,  $\Delta_A\Delta^{17}\text{O}$  is -0.25 ‰ for maize with  $c_m/c_a$  ratio of 0.3, -0.3 ‰ for ivy with  $c_m/c_a$  ratio of 0.5 ‰ and -0.5 ‰ for sunflower with  $c_m/c_a$  ratio of 0.7 (Figure 3b). For sunflower and ivy,  $\Delta_A\Delta^{17}\text{O}$  is also strongly dependent on the  $\Delta^{17}\text{O}$  of  $\text{CO}_2$  supplied to the cuvette, whereas no significant dependence is found for maize. For an increase in  $\Delta^{17}\text{O}$  of  $\text{CO}_2$  entering the cuvette from -0.215 ‰ to 0.435 ‰,  $\Delta_A\Delta^{17}\text{O}$  increases from -0.3 ‰ to -0.9 ‰ at  $c_m/c_a$  ratio of 0.5 for ivy. For sunflower, an increase in  $\Delta^{17}\text{O}$  of  $\text{CO}_2$  entering the cuvette from -0.215 ‰ to 0.31 ‰ increases  $\Delta_A\Delta^{17}\text{O}$  from -0.8 ‰ to -1.7 ‰ at  $c_m/c_a$  ratio of 0.8. The leaf cuvette model results illustrate the shape of the dependence on the  $c_m/c_a$  ratio and agree well with the experiments. For the leaf cuvette model, the  $\Delta^{17}\text{O}$  value of the water is assigned a constant value of -0.122 ‰ (average  $\Delta^{17}\text{O}$  value for the bulk leaf water).

Figure 4b shows the same values of  $\Delta_A\Delta^{17}\text{O}$  as a function of the difference between  $\Delta^{17}\text{O}$  of  $\text{CO}_2$  entering the leaf and the calculated  $\Delta^{17}\text{O}$  of leaf water at the evaporation site where  $\text{CO}_2\text{-H}_2\text{O}$  exchange takes place ( $\Delta^{17}\text{O}_a - \Delta^{17}\text{O}_{\text{wes}}$ ), for different  $c_m/c_a$  ratios. The leaf cuvette model results (solid lines in Figure 4b) suggest a linear dependence between  $\Delta_A\Delta^{17}\text{O}$  and ( $\Delta^{17}\text{O}_a - \Delta^{17}\text{O}_{\text{wes}}$ ). The experimental results agree with the hypothesis that  $\Delta_A\Delta^{17}\text{O}$  is linearly dependent on  $\Delta^{17}\text{O}_a - \Delta^{17}\text{O}_{\text{wes}}$  at a certain  $c_m/c_a$  ratio. Figure 4a shows the corresponding relation where  $\Delta_A\Delta^{17}\text{O}$  is divided by  $\Delta^{17}\text{O}_a - \Delta^{17}\text{O}_m$ . All the values follow the same relationship as function of the  $c_m/c_a$  ratio, which can be approximated quite well by an exponential function (equation 5). This function quantifies the dependence of  $\Delta_A\Delta^{17}\text{O}$  on  $c_m/c_a$ , and thus the effect of the diffusion of isotopically exchanged  $\text{CO}_2$  back to the atmosphere, which increases with increasing  $c_m/c_a$  ratio.

335

$$\frac{\Delta_A\Delta^{17}\text{O}}{\Delta^{17}\text{O}_a - \Delta^{17}\text{O}_m} = -0.150 \times \exp(3.707 \times c_m/c_a) + 0.028 \quad (5)$$

Figure 5 a and c show results from the leaf cuvette model that illustrates in more detail how  $\Delta^{17}\text{O}_e$  and  $\Delta^{17}\text{O}_{\text{wes}}$  affect  $\Delta^{17}\text{O}_a$  and  $\Delta_A\Delta^{17}\text{O}$  and their dependence on  $c_m/c_a$ . At lower  $c_m/c_a$ , only a very small fraction of  $\text{CO}_2$  that has undergone isotopic equilibration in the mesophyll diffuses back to the atmosphere, and therefore  $\Delta^{17}\text{O}_a$  stays close to the incoming  $\Delta^{17}\text{O}_e$ , modified by the fractionation during  $\text{CO}_2$  diffusion

340

through the stomata (Figure 5a). Figure 5c confirms that indeed at low  $c_m/c_a$ ,  $\Delta_A\Delta^{17}\text{O}$  approaches the fractionation constant expected for diffusion,  $-0.170\text{‰}$ . This diffusional fractionation is independent of the isotopic composition of the  $\text{CO}_2$  entering the leaf, and therefore at low  $c_m/c_a$ , the  $\Delta_A\Delta^{17}\text{O}$  curves for the different values of the anomaly of the  $\text{CO}_2$  entering the leaf converge. For a high  $c_m/c_a$  ratio, the back-diffusion flux of  $\text{CO}_2$  that has equilibrated with water becomes the dominant factor, and in this case, the isotopic composition of the outgoing  $\text{CO}_2$  converges towards this isotope value, independent of the isotopic composition of the incoming  $\text{CO}_2$  (Figure 5a). This can lead to a very wide range of values for the discrimination against  $\Delta^{17}\text{O}$ , because now the effect on  $\Delta^{17}\text{O}$  of the ambient  $\text{CO}_2$  depends strongly on the difference in isotopic composition between incoming  $\text{CO}_2$  and  $\text{CO}_2$  in isotopic equilibrium with the leaf water.

In the model calculations shown in Figure 5b and d, the isotopic composition of the water was changed from  $\Delta^{17}\text{O}_{\text{wes}} = -0.122\text{‰}$  to  $0.300\text{‰}$ , whereas all other parameters were kept the same. The value of  $\Delta^{17}\text{O}_e$  for which  $\Delta^{17}\text{O}_a$  does not depend on  $c_m/c_a$  is shifted accordingly, again being similar to  $\Delta^{17}\text{O}_m$ . At low  $c_m/c_a$   $\Delta_A\Delta^{17}\text{O}$  converges to the same value as in Figure 5c, confirming the role of diffusion into the stomata as discussed above.

Figure 6 shows how  $\delta^{18}\text{O}$  and  $\Delta^{17}\text{O}$  vary in key compartments of the leaf cuvette system that determine the oxygen isotope effects associated with photosynthesis, based on the previously established three-isotope slopes of the various processes (Figure 1). The irrigation water has a  $\Delta^{17}\text{O}$  value of  $0.017$ . The measured bulk leaf water is  $6\text{--}16\text{‰}$  enriched in  $^{18}\text{O}$  and its  $\Delta^{17}\text{O}$  value is lower by  $-0.075$  to  $-0.200\text{‰}$  (mean value  $-0.121\text{‰}$ ) than the irrigation water, calculated using a three-isotope slope of  $\lambda_{\text{trans}} = 0.516$  at  $80\%$  humidity (Landais et al., 2006).  $\Delta^{17}\text{O}$  of leaf water at the evaporation site, calculated from the transpired water, has slightly lower  $\Delta^{17}\text{O}$ , with values between  $-0.119\text{‰}$  and  $-0.237$  (average  $-0.184\text{‰}$ ). Note that the bulk leaf water was not measured for all the experiments. For the experiments where the bulk leaf water is measured,  $\Delta^{17}\text{O}$  of leaf water at the evaporation site ranges from  $-0.160\text{‰}$  to  $-0.231$  with an average value of  $-0.190 \pm 0.020\text{‰}$ . The calculated isotopic composition of water at the exchange site was thus similar, but slightly lower in  $\Delta^{17}\text{O}$  than the values measured for bulk leaf water.  $\text{CO}_2$  exchanges with the water in the leaf with a well-established fractionation constant (see equation S17, supplementary material) and a three-isotope slope of  $\lambda_{\text{CO}_2\text{-H}_2\text{O}} = 0.5229$  (Barkan and Luz, 2012), leading to the lower  $\Delta^{17}\text{O}$  values of the equilibrated  $\text{CO}_2$ . In our experiments, the  $\Delta^{17}\text{O}$  value of  $\text{CO}_2$  in equilibrium with leaf water is lower than the  $\Delta^{17}\text{O}$  value of  $\text{CO}_2$  entering the leaf. The  $\Delta^{17}\text{O}$  of the  $\text{CO}_2$  in the intercellular air space is a mixture between two end members, the  $\Delta^{17}\text{O}$  of the  $\text{CO}_2$  entering the leaf and  $\Delta^{17}\text{O}$  of the  $\text{CO}_2$  in equilibrium with leaf water. This explains why the observed values of  $\Delta_A\Delta^{17}\text{O}$  are negative for the experiments performed in this study.

## 4. Discussion

### 4.1. Discrimination against $^{18}\text{O}$ of $\text{CO}_2$

380 The higher  $\Delta_A^{18}\text{O}_{\text{obs}}$  values for sunflower compared to maize and ivy (Figure 3a) are mainly due to a  
 higher back-diffusion flux ( $c_m/(c_a-c_m)$ ). The back-diffusion flux is higher for the C<sub>3</sub> plants sunflower and  
 ivy than for the C<sub>4</sub> plant maize, a consequence of the lower stomatal conductance and higher assimilation  
 385 in a lower back-diffusion flux. The increase of assimilation rate with higher light intensity decreases the  
 $c_m/c_a$  ratio and thus leads to a lower back-diffusion flux, which explains the decreases of  $\Delta_A^{18}\text{O}_{\text{obs}}$  for  
 maize and most clearly for sunflower. A similar trend of increase in  $\Delta_A^{18}\text{O}_{\text{obs}}$  with an increase in  $c_m/c_a$   
 ratio has been reported in previous studies (Gillon and Yakir, 2000b, a; Osborn et al., 2017). For ivy,  
 $\Delta_A^{18}\text{O}_{\text{obs}}$  and  $\Delta_A^{17}\text{O}_{\text{obs}}$  do not decrease with an increase in irradiance, because the change in assimilation  
 390 rate with irradiance is small. Thus,  $c_m$  will not decrease strongly and the effect on the back diffusion is  
 smaller than the variability in  $\Delta_A^{18}\text{O}_{\text{obs}}$  of different leaves of the same plant.

In our experiments, photosynthesis causes enrichment  $\delta^{18}\text{O}$  of atmospheric CO<sub>2</sub> for both C<sub>3</sub> and C<sub>4</sub> plants,  
 i.e. positive values  $\Delta_A^{18}\text{O}$ . In principle,  $\Delta_A^{18}\text{O}$  can also be negative if the  $\delta^{18}\text{O}_m$  are depleted relative to  
 395 the ambient CO<sub>2</sub>. This is in contrast to  $\Delta_A^{13}\text{C}$ , which will always be positive since it is determined by the  
 fractionation due to the PEPC and RuBisCO enzyme activity (Figure S8 and S9, supplementary material).  
 In general, in our experiments, the  $\Delta_A^{18}\text{O}_{\text{obs}}$  values are about five times larger than  $\delta^{18}\text{O}_a - \delta^{18}\text{O}_e$ , the  
 $\delta^{18}\text{O}$  difference between CO<sub>2</sub> entering and leaving the cuvette (Figure S10 to S12 supplementary  
 material). This is easy to understand from the definition of  $\Delta_A$ . Taking  $\Delta_A^{18}\text{O}$  as an example,  $\Delta_A^{18}\text{O}_{\text{obs}} =$   
 400  $\frac{\zeta(\delta^{18}\text{O}_a - \delta^{18}\text{O}_e)}{1 + \delta^{18}\text{O}_a - \zeta(\delta^{18}\text{O}_a - \delta^{18}\text{O}_e)} \approx \zeta(\delta^{18}\text{O}_a - \delta^{18}\text{O}_e)$  and in our experiments,  $\zeta = c_e / (c_e - c_a) \approx 500 / (500 - 400) = 5$ .

#### 4.2. Discrimination against the $\Delta^{17}\text{O}$ of CO<sub>2</sub>

The leaf cuvette model includes the isotope fractionations of all the individual processes that have been  
 quantified in dedicated experiments previously (Figure 1). The good agreement of the model results with  
 405 the measurements (Figure 3a) demonstrates that when all these processes are combined in the quantitative  
 description of a gas exchange experiment, they actually result in a correct quantification of the isotope  
 effects associated with photosynthesis. This has already been demonstrated before for  $\Delta_A^{18}\text{O}_{\text{obs}}$  but has  
 now been confirmed for  $\Delta_A\Delta^{17}\text{O}$ .

410 Unlike ivy and sunflower, maize does not show a significant change in  $\Delta_A\Delta^{17}\text{O}$  when CO<sub>2</sub> gases with  
 different  $\Delta^{17}\text{O}$  are supplied to the plant. The C<sub>4</sub> plant maize has a small back-diffusion flux due to its high  
 assimilation rate and low stomatal conductance, leading to a low  $c_m/c_a$  ratio. At low  $c_m/c_a$  ratios,  $\Delta_A\Delta^{17}\text{O}$   
 is expected to be close to the weighted fractionation due to diffusion through boundary layer and stomata.  
 In general, the effect of diffusion on  $\Delta^{17}\text{O}$  of atmospheric CO<sub>2</sub> can be expressed as follows:

415

$$\Delta^{17}\text{O}_{\text{Modified}} = \Delta^{17}\text{O}_a + (\lambda_{\text{RL}} - \lambda_{\text{diffusion}}) \times \ln\alpha_{\text{diffusion}} \quad (6)$$

where  $\Delta^{17}\text{O}_a$  is the  $\Delta^{17}\text{O}$  of the  $\text{CO}_2$  surrounding the leaf,  $\Delta^{17}\text{O}_{\text{modified}}$  is the  $\Delta^{17}\text{O}$  of the  $\text{CO}_2$  modified due to diffusional fractionation and  $\lambda_{\text{diffusion}}$ ,  $\lambda_{\text{RL}}$  and  $\alpha_{\text{diffusion}}$  are the oxygen three-isotope relationships during diffusion from the  $\text{CO}_2\text{-H}_2\text{O}$  exchange site to the atmosphere, the reference slope used and the fractionation against  $^{18}\text{O}$  for  $\text{CO}_2$  during diffusion through the stomata. Using the values  $\lambda_{\text{RL}} = 0.528$ ,  
420  $\lambda_{\text{diffusion}} = 0.509$  (Young et al., 2002) and  $\alpha_{\text{diffusion}} = 0.9912$  (Farquhar and Lloyd, 1993), the effect of diffusional fractionation on the  $\Delta^{17}\text{O}$  of atmospheric  $\text{CO}_2$  is  $-0.168\text{‰}$  regardless of the anomaly of the  $\text{CO}_2$  entering the leaf, and the model results confirm this at low  $c_m/c_a$  ratio (Figure 5 c and d, inset).

At a high  $c_m/c_a$  ratio,  $\Delta^{17}\text{O}_a$  is dominated by the back-diffusion flux of  $\text{CO}_2$  that has equilibrated with water. As a consequence,  $\Delta^{17}\text{O}_a$  converges to a common value that is independent of the anomaly of the  
425  $\text{CO}_2$  entering the cuvette and is determined by the isotopic composition of leaf water. Figure 5 confirms that the end member is equal to the  $\Delta^{17}\text{O}$  of  $\text{CO}_2$  in equilibrium with leaf water,  $\Delta^{17}\text{O}_m$ . In fact, when  $\Delta^{17}\text{O}_a = \Delta^{17}\text{O}_m$ ,  $\Delta^{17}\text{O}_a$  does not change with  $c_m/c_a$ , indicating that in this case the  $\Delta^{17}\text{O}$  of the  $\text{CO}_2$  diffusing back from the leaf is the same as the  $\Delta^{17}\text{O}(\text{CO}_2)$  entering the leaf.

$\bar{a}_{18}$  is the overall discrimination occurring during the diffusion of  $^{12}\text{C}^{18}\text{O}^{16}\text{O}$  from the ambient air  
430 surrounding the leaf to the  $\text{CO}_2\text{-H}_2\text{O}$  exchange site (see Table 2 for the list of variables). In our study  $\bar{a}_{18}$  ranges from  $5\text{‰}$  to  $7.2\text{‰}$ , lower than the literature estimate of  $7.4\text{‰}$  (Farquhar et al., 1993).  $\bar{a}_{18}$  depends on the ratio of stomatal conductance, which is associated with a strong fractionation of  $8.8\text{‰}$ , to mesophyll conductance with an associated fractionation of only  $0.8\text{‰}$ . Therefore, the higher the ratio ( $g_s/g_{m18}$ ) the lower the  $\bar{a}_{18}$  (Table S2, supplementary material). The difference in  $\bar{a}_{18}$  of  $2.4\text{‰}$  between  
435 the literature value of  $7.4\text{‰}$  and the lowest  $\bar{a}_{18}$  estimate in this study will introduce an error of only  $0.046\text{‰}$  in the  $\Delta^{17}\text{O}$  value (see equation 6). The uncertainty  $\bar{a}_{18}$  has lower influence on the  $\Delta_A\Delta^{17}\text{O}$  of  $\text{C}_3$  plants compared to  $\text{C}_4$  plants since the diffusional fractionation is less important at the higher  $c_m/c_a$  ratio where  $\text{C}_3$  plants operate.

### 440 4.3. Global average value of $\Delta_A\Delta^{17}\text{O}$ and $\Delta^{17}\text{O}$ isoflux

We can use the established relationship between  $\Delta_A\Delta^{17}\text{O}$  and  $\Delta^{17}\text{O}_a - \Delta^{17}\text{O}_{\text{wes}}$  for a certain  $c_m/c_a$  ratio to provide a bottom-up estimate for the global effect of photosynthesis on  $\Delta^{17}\text{O}$  in atmospheric  $\text{CO}_2$ , based on data obtained in real gas exchange experiments. For this, we use results from a recent modeling study, which provides global average values for  $\text{CO}_2$  and leaf water ( $\Delta^{17}\text{O}(\text{CO}_2) = -0.168\text{‰}$ ,  $\Delta^{17}\text{O}(\text{H}_2\text{O}_{\text{-leaf}}) = -$   
445  $0.067\text{‰}$ ; (Koren et al., 2019); Figure S13 and 14, supplementary material). The  $\Delta^{17}\text{O}(\text{CO}_2)$  values agree well with the limited amount of available measurements (Table 3).

To extrapolate  $\Delta_A\Delta^{17}\text{O}$  determined in the leaf scale experiments to the global scale, global average  $c_m/c_a$  ratios of  $0.7$  and  $0.3$  are used for  $\text{C}_3$  and  $\text{C}_4$  plants, respectively, similar to previous studies (Hoag et al.,  
450 2005; Liang et al., 2017b). From SIBCASA model results we obtained an annual variability of  $c_i/c_a$  values with a standard deviation of  $0.12$  and  $0.17$  for  $\text{C}_4$  and  $\text{C}_3$  plants respectively (Figure S15, supplementary material) (Schaefer et al., 2008; Koren et al., 2019). We use this variability as upper limit of the error estimate for  $c_m/c_a$  as shown in the light orange and light pink shaded areas in Figure 4b. This error is converted to an error in  $\Delta_A\Delta^{17}\text{O}$  using the relation with  $c_m/c_a$ . Based on the linear dependency of  $\Delta_A\Delta^{17}\text{O}$

455 and  $\Delta^{17}\text{O}_a - \Delta^{17}\text{O}_{\text{wes}}$ , we estimate the  $\Delta_A \Delta^{17}\text{O}$  for tropospheric  $\text{CO}_2$  based on the  $\Delta^{17}\text{O}$  of leaf water and  $c_m/c_a$  ratio. In Figure 4b, the dashed black vertical line indicates  $\Delta^{17}\text{O}_a - \Delta^{17}\text{O}_{\text{wes}}$  obtained from the 3D global model (Koren et al., 2019). The results of the global estimate and parameters used for the extrapolation of leaf scale study to the global scale are summarized in Table 3.

460 The  $\delta^{17}\text{O}$  value of atmospheric  $\text{CO}_2$  (21.53 ‰) is calculated from the global  $\delta^{18}\text{O}$  and  $\Delta^{17}\text{O}$  values (41.5 ‰ and -0.168 ‰, respectively) (Koren et al., 2019). The  $\delta^{17}\text{O}$  and  $\delta^{18}\text{O}$  values of global mean leaf water are calculated from the soil water. A global mean  $\delta^{18}\text{O}$  value of soil water is -8.4 ‰ assuming soil water to be similar to precipitation (Bowen and Revenaugh, 2003; Koren et al., 2019). The  $\delta^{17}\text{O}$  value of soil water is -4.4 ‰, calculated using equation 7 (Luz and Barkan, 2010).

465

$$\ln(\delta^{17}\text{O}_{\text{soil}} + 1) = 0.528 \times \ln(\delta^{18}\text{O}_{\text{soil}} + 1) + 0.033 \quad (7)$$

$\delta^{17}\text{O}$  and  $\delta^{18}\text{O}$  of leaf water are calculated from  $\delta^{17}\text{O}$  and  $\delta^{18}\text{O}$  of soil water with fractionation factors of 1.0043 and 1.0084, respectively (Hofmann et al., 2017; Koren et al., 2019). The fractionation factor for  $\delta^{17}\text{O}$  is calculated using  $\alpha^{17} = (\alpha^{18})^{\lambda_{\text{trans}}}$  with  $\lambda_{\text{trans}} = 0.516$ , assuming relative humidity to be 75 % (Landais et al., 2006). The  $\delta^{17}\text{O}$  and  $\delta^{18}\text{O}$  values of global mean leaf water are then -0.136 ‰ and -0.131 ‰, respectively. Thus, the difference between global atmospheric  $\text{CO}_2$  and leaf water is  $\delta^{17}\text{O}_{\text{CO}_2 - \text{water}} = 21.666$  ‰ and  $\delta^{18}\text{O}_{\text{CO}_2 - \text{water}} = 41.631$  ‰. This yields  $\Delta^{17}\text{O}_{\text{CO}_2 - \text{water}} = -0.101$  ‰, and this value is indicated as dashed black line in Figure 4. The grey shaded area indicates the propagated error using the standard deviation of the relevant parameters in 180 x 360 grid boxes for 12 months of leaf water and 45 x 60 grid boxes for 24 months for  $\text{CO}_2$  (Koren et al., 2019). In Figure 4b, the intersection between the dashed black vertical line and the discrimination lines for the representative  $c_m/c_a$  ratios of  $\text{C}_3$  and  $\text{C}_4$  plants corresponds to the  $\Delta_A \Delta^{17}\text{O}$  value of  $\text{C}_3$  and  $\text{C}_4$  plants. For  $\text{C}_4$  plants ( $c_m/c_a = 0.3$ ) this yields  $\Delta_A \Delta^{17}\text{O} = -0.3$  ‰ (gray dashed line in Figure 4b) and for  $\text{C}_3$  plants ( $c_m/c_a = 0.7$ ),  $\Delta_A \Delta^{17}\text{O} = -0.65$  ‰ (black dashed line in Figure 4b).

480 Three main factors contribute to the uncertainty of the extrapolated  $\Delta_A \Delta^{17}\text{O}$  value. The first is the measurement error, which contributes 0.25 ‰ (standard error for individual experiments). The second factor is the uncertainty in the difference between  $\Delta^{17}\text{O}$  of atmospheric  $\text{CO}_2$  and leaf water, and we use results from the global model to estimate an error. For  $\Delta^{17}\text{O}$  of atmospheric  $\text{CO}_2$ , statistics for all 45 x 60 grid boxes for 24 months (2012-2013) show a range of -0.218 ‰ to -0.151 ‰, with a mean of -0.168 ‰ and a standard deviation of 0.013 ‰ (Figure S13, supplementary material). For  $\Delta^{17}\text{O}$  of the leaf water statistics for all 180 x 360 grid boxes for 12 months show a range of -0.236 ‰ and -0.027 ‰ (Figure S14, supplementary material). The mean is -0.067 ‰ with a standard deviation of 0.041 ‰. From the combined errors we estimate the error in  $(\Delta^{17}\text{O}_a - \Delta^{17}\text{O}_{\text{wes}})$  to be 0.043 ‰. The third uncertainty in the extrapolation of  $\Delta^{17}\text{O}$  comes from the uncertainty in the  $c_m/c_a$  ratio. For  $\text{C}_3$  and  $\text{C}_4$  plants, these errors are indicated by the light orange and light blue shadings in Figure 4b.

490 Taking these uncertainties into account leads to a mean value of  $\Delta_A \Delta^{17}\text{O} = -0.3 \pm 0.18$  ‰ for  $\text{C}_4$  plants and  $\Delta_A \Delta^{17}\text{O} = -0.65 \pm 0.18$  ‰ for  $\text{C}_3$  plants. The leaf scale discrimination against  $\Delta^{17}\text{O}$  is then extrapolated to

495 global vegetation using these representative values of  $\Delta_A\Delta^{17}\text{O}$  and the relative fractions of photosynthesis by  $\text{C}_4$  and  $\text{C}_3$  plants, respectively as:

$$\Delta_A\Delta^{17}\text{O}_{\text{global}} = f_{\text{C}_4} \times \Delta_A\Delta^{17}\text{O}_{\text{C}_4} + f_{\text{C}_3} \times \Delta_A\Delta^{17}\text{O}_{\text{C}_3} \quad (8)$$

500 where  $f_{\text{C}_4}$  and  $f_{\text{C}_3}$  are the photosynthesis weighted global coverage of  $\text{C}_4$  and  $\text{C}_3$  vegetation.  $\Delta_A\Delta^{17}\text{O}_{\text{C}_4}$  and  $\Delta_A\Delta^{17}\text{O}_{\text{C}_3}$  quantify the discrimination against  $\Delta^{17}\text{O}$  by  $\text{C}_4$  and  $\text{C}_3$  plants, which are calculated using estimated values of  $c_m/c_a$  from a model. Using assimilation weighted fractions of 23 % for  $\text{C}_4$  and 77 % for  $\text{C}_3$  vegetation (Still et al., 2003), the global mean value of  $\Delta_A\Delta^{17}\text{O}$  obtained from equation 8 is -0.57±0.14 ‰.

505 Isoflux is the product of isotope composition and gross mass flux of the molecule. In the case of assimilation, the net flux  $F_A = F_{AL} - F_{LA}$  is multiplied with the discrimination associated with assimilation (Ciais et al., 1997a).  $F_{LA}$  and  $F_{AL}$  are total  $\text{CO}_2$  fluxes from leaf to the atmosphere and from atmosphere to leaf, respectively. The global scale  $\Delta^{17}\text{O}_A$  isoflux is calculated by multiplying the discrimination with the assimilation flux as:

$$F_A \times \Delta_A\Delta^{17}\text{O} = A \times (f_{\text{C}_4} \times \Delta_A\Delta^{17}\text{O}_{\text{C}_4} + f_{\text{C}_3} \times \Delta_A\Delta^{17}\text{O}_{\text{C}_3}) \quad (9)$$

510 where,  $A=0.88 \times \text{GPP}$  is the terrestrial assimilation rate. The factor 0.88 accounts for the fraction of  $\text{CO}_2$  released due to autotrophic respiration (Ciais et al., 1997a). The  $\Delta_A\Delta^{17}\text{O}$  isoflux due to photosynthesis is calculated using a GPP value of  $120 \text{ PgCyr}^{-1}$  (Beer et al., 2010) and  $A=0.88 \times \text{GPP}$ , resulting in an isoflux of  $-60 \pm 15 \text{ ‰ PgCyr}^{-1}$  globally. This is the first global estimate of  $\Delta_A\Delta^{17}\text{O}$  based on direct measurements of the discrimination during assimilation. Our value is in good agreement with previous model estimates. 515 Hofmann et al. (2017) estimated an isoflux ranging from -42 to -92 ‰PgCyr<sup>-1</sup> (converted to a reference line with  $\lambda=0.528$ ) using an average  $c_m/c_a$  ratio of 0.7 for both  $\text{C}_4$  and  $\text{C}_3$  plants and  $\Delta^{17}\text{O}$  of -0.147 ‰ for atmospheric  $\text{CO}_2$ . A model-estimated value from (Hoag et al., 2005) is -47 ‰PgCyr<sup>-1</sup> (converted to our reference slope of  $\lambda=0.528$ ), derived with a more simple model and using  $\Delta^{17}\text{O}$  of -0.146 ‰ with  $c_m/c_a$  ratio of 0.33 and 0.66 for  $\text{C}_4$  and  $\text{C}_3$  plants, respectively.

520 The main uncertainty in the extrapolation of  $\Delta_A\Delta^{17}\text{O}$  from the leaf experiments to the global scale is the uncertainty in the  $c_m/c_a$  ratio. The error from the uncertainty in  $c_m/c_a$  ratio increases when the relative difference in  $\Delta^{17}\text{O}$  between  $\text{CO}_2$  and leaf water increases (Figure 5b). It is difficult to determine a single representative  $c_m$  value for different plants because this value would need to be properly weighted with temperature, irradiance,  $\text{CO}_2$  mole fraction and other environmental factors (Flexas et al., 2008; 2012; Shrestha et al., 2019). Recent developments in laser spectroscopy techniques (McManus et al., 2005; Nelson et al., 2008; Tuzson et al., 2008; Kammer et al., 2011) might enable more and easier measurements of  $c_m/c_a$  both in the laboratory and under field conditions. This could lead to a better understanding of variations in the  $c_m/c_a$  ratio among plant species and, temporally, spatially and environmentally.

530

## 5. Conclusions

In order to directly quantify the effect of photosynthetic gas exchange on the  $\Delta^{17}\text{O}$  of atmospheric  $\text{CO}_2$ , gas exchange experiments were carried out in leaf cuvettes using two  $\text{C}_3$  plants (sunflower and ivy) and one  $\text{C}_4$  plant (maize) with isotopically normal and slightly anomalous ( $^{17}\text{O}$ -enriched)  $\text{CO}_2$ . Results for  $^{18}\text{O}$  agree with results reported in the literature previously. Our results for  $\Delta^{17}\text{O}$  confirm that the formalism developed by Farquhar and others for  $\delta^{18}\text{O}$  is also applicable to the evaluation of  $\Delta^{17}\text{O}$ . In particular, our experiments confirm that two parameters determine the effect of photosynthesis on  $\text{CO}_2$ : 1) the  $\Delta^{17}\text{O}$  difference between the incoming  $\text{CO}_2$  and  $\text{CO}_2$  in equilibrium with leaf water and 2) the  $c_m/c_a$  ratio, which determines the degree of back-flux of isotopically exchanged  $\text{CO}_2$  from the mesophyll to the atmosphere. At low  $c_m/c_a$  ratios,  $\Delta_A\Delta^{17}\text{O}$  is mainly influenced by the diffusional fractionation. Under our experimental conditions, the isotopic effect increased with  $c_m/c_a$ , e.g.  $\Delta_A\Delta^{17}\text{O}$  was  $-0.3\text{‰}$  and  $-0.65\text{‰}$  for maize and sunflower with  $c_m/c_a$  ratios of 0.3 and 0.7, respectively. However, experiments with mass independently fractionated  $\text{CO}_2$  demonstrate that the results depend strongly on the  $\Delta^{17}\text{O}$  difference between the incoming  $\text{CO}_2$  and  $\text{CO}_2$  in equilibrium with leaf water. This is supported by calculations with a leaf cuvette model.

$\delta^{18}\text{O}$  is largely affected by kinetic and equilibrium processes between  $\text{CO}_2$  and leaf water, and also leaf water isotopic inhomogeneity and dynamics. The  $\Delta^{17}\text{O}$  variation is much smaller compared to  $\delta^{18}\text{O}$  and is better defined since conventional bio-geo-chemical processes that modify  $\delta^{17}\text{O}$  and  $\delta^{18}\text{O}$  follow a well-defined three-isotope fractionation slope. Results from the leaf exchange experiments were upscaled to the global atmosphere using modeled values for  $\Delta^{17}\text{O}$  of leaf water and  $\text{CO}_2$ , which results in  $\Delta_A\Delta^{17}\text{O} = -0.57 \pm 0.14\text{‰}$  and a value for the  $\Delta^{17}\text{O}$  isoflux of  $-60 \pm 15\text{‰ PgCyr}^{-1}$ . This is the first study that provides such an estimate based on direct leaf chamber measurements, and the results agree with previous  $\Delta^{17}\text{O}$  calculations. The largest contribution to the uncertainty originates from uncertainty in the  $c_m/c_a$  ratio and the largest contributions to the isoflux come from  $\text{C}_3$  plants, which have both a higher share of the total assimilation and higher discrimination.  $\Delta_A\Delta^{17}\text{O}$  is less sensitive to  $c_m/c_a$  ratios at lower values of  $c_m/c_a$ , for instance for  $\text{C}_4$  plants, maize.

$\Delta^{17}\text{O}$  of tropospheric  $\text{CO}_2$  is controlled by photosynthetic gas exchange, respiration, soil invasion, and stratospheric influx. The stratospheric flux is well established and the effect of photosynthetic gas exchange can now be quantified more precisely. To untangle the contribution of each component to the  $\Delta^{17}\text{O}$  atmospheric  $\text{CO}_2$  we recommend measuring the effects of foliage respiration and soil invasion both in the laboratory and at the ecosystem scale.

565

Code and data availability.

570 The data used in this study are included in the paper either with figures or tables. The python code for  
the cuvette model is available at [https://git.wur.nl/leaf\\_model/D170](https://git.wur.nl/leaf_model/D170).

#### Author contributions.

GAA and TR designed the main idea of the study. GAA and TP designed the leaf cuvette setup. TP  
monitors plant growth. GAA and TR designed the CO<sub>2</sub> extraction and CO<sub>2</sub>-H<sub>2</sub>O exchange system. GAA  
575 conducted all the measurements. GK provided the leaf cuvette model. WP enabled the work within the  
ASICA project. All authors discussed the results at different steps of the project. GAA and TR prepared  
the manuscript with contributions from all the co-authors.

#### Competing interests.

The authors declare that they have no conflict of interest

580

### Acknowledgments

The authors thank Leonard I. Wassenaar and Stefan Terzer-Wassmuth from the International Atomic and  
Energy Agency, Vienna for supplying water standards. The authors thank Eugeni Barkan and Rolf Vieten  
585 from the Hebrew University of Jerusalem for calibration of our O<sub>2</sub> and CO<sub>2</sub> working gases. We are  
grateful to Amaelle Landais from Laboratoire des Sciences Du Climat et de l'Environnement Université  
Paris-Saclay for measuring the  $\Delta^{17}\text{O}$  of leaf water samples for our study. The authors thank Amzad  
Laskar for useful discussion during the design of the experiment. This work is funded by the EU ERC  
project ASICA.

590

595

### Reference

- Adnew, G. A., Hofmann, M. E. G., Paul, D., Laskar, A., Surma, J., Albrecht, N., Pack, A., Schwieters, J., Koren, G., Peters,  
W., and Röckmann, T.: Determination of the triple oxygen and carbon isotopic composition of CO<sub>2</sub> from atomic ion fragments  
600 formed in the ion source of the 253 Ultra High-Resolution Isotope Ratio Mass Spectrometer, *Rapid Commun. Mass. Sp.*, 33,  
17, 2019.
- Badger, M. R., and Price, G. D.: The role of carbonic anhydrase in photosynthesis, *Annu. Rev. Plant Biol.*, 45, 23, 1994.
- Bao, H., Cao, X., and Hayles, J. A.: Triple oxygen isotopes: fundamental relationships and applications, *Annu. Rev. Earth  
Planet. Sci.*, 44, 29, 2016.
- 605 Barbour, M. M., Evans, J. R., Simonin, K. A., and von Caemmerer, S.: Online CO<sub>2</sub> and H<sub>2</sub>O oxygen isotope fractionation  
allows estimation of mesophyll conductance in C<sub>4</sub> plants, and reveals that mesophyll conductance decreases as leaves age in  
both C<sub>4</sub> and C<sub>3</sub> plants, *New Phytol.*, 14, 2016.




- Barkan, E., and Luz, B.: High precision measurements of  $^{17}\text{O}/^{16}\text{O}$  and  $^{18}\text{O}/^{16}\text{O}$  ratios in  $\text{H}_2\text{O}$ , *Rapid Commun. Mass. Sp.*, 19, 3737-3742, 10.1002/rcm.2250, 2005.
- 610 Barkan, E., and Luz, B.: Diffusivity fractionations of  $\text{H}_2^{16}\text{O}/\text{H}_2^{17}\text{O}$  and  $\text{H}_2^{16}\text{O}/\text{H}_2^{18}\text{O}$  in air and their implications for isotope hydrology, *Rapid Commun. Mass. Sp.*, 21, 6, 2007.
- Barkan, E., and Luz, B.: The relationships among the three stable isotopes of oxygen in air, seawater and marine photosynthesis, *Rapid Commun. Mass. Sp.*, 25, 2, 2011.
- 615 Barkan, E., and Luz, B.: High-precision measurements of  $^{17}\text{O}/^{16}\text{O}$  and  $^{18}\text{O}/^{16}\text{O}$  ratios in  $\text{CO}_2$ , *Rapid Commun. Mass. Sp.*, 26, 2733-2738, 10.1002/rcm.6400, 2012.
- Barkan, E., Musan, I., and Luz, B.: High-precision measurements of  $\delta^{17}\text{O}$  and  $^{17}\text{O}$ -excess of NBS19 and NBS18, *Rapid Commun. Mass. Sp.*, 29, 2219-2224, 10.1002/rcm.7378, 2015.
- 620 Beer, C., Reichstein, M., Tomelleri, E., Ciais, P., Jung, M., Carvalhais, N., Rodenbeck, C., Arain, M. A., Baldocchi, D., Bonan, G. B., Bondeau, A., Cescatti, A., Lasslop, G., Lindroth, A., Lomas, M., Luyssaert, S., Margolis, H., Oleson, K. W., Rouspard, O., Veenendaal, E., Viovy, N., Williams, C., Woodward, F. I., and Papale, D.: Terrestrial gross carbon dioxide uptake: global distribution and covariation with climate, *Science*, 329, 834-838, 10.1126/science.1184984, 2010.
- Boering, K. A.: Observations of the anomalous oxygen isotopic composition of carbon dioxide in the lower stratosphere and the flux of the anomaly to the troposphere, *Geophysical Research Letters*, 31, 10.1029/2003gl018451, 2004.
- 625 Booth, B. B. B., Jones, C. D., Collins, M., Totterdell, I. J., Cox, P. M., Sitch, S., Huntingford, C., Betts, R. A., Harris, G. R., and Lloyd, J.: High sensitivity of future global warming to land carbon cycle processes, *Environmental Research Letters*, 7, 10.1088/1748-9326/7/2/024002, 2012.
- Bowen, G. J., and Revenaugh, J.: Interpolating the isotopic composition of modern meteoric precipitation, *Water Resour. Res.*, 39, 2003.
- 630 Brand, W. A., Assonov, S. S., and Coplen, T. B.: Correction for the  $^{17}\text{O}$  interference in  $\delta^{13}\text{C}$  measurements when analyzing  $\text{CO}_2$  with stable isotope mass spectrometry (IUPAC Technical Report), *Pure Appl. Chem.*, 82, 1719-1733, 10.1351/pac-rep-09-01-05, 2010.
- Cao, X., and Liu, Y.: Equilibrium mass-dependent fractionation relationships for triple oxygen isotopes, *Geochimica et Cosmochimica Acta*, 75, 7435-7445, 10.1016/j.gca.2011.09.048, 2011.
- 635 Cernusak, L. A., Barbour, M. M., Arndt, S. K., Cheesman, A. K., English, N. B., Feild, T. S., Helliker, B. R., Holloway-Phillips, M. M., Holtum, J. A., Kahmen, A., McInerney, F. A., Munksgaard, N. C., Simonin, K. A., Song, X., Stuart-Williams, H., West, J. B., and Farquhar, G. D.: Stable isotopes in leaf water of terrestrial plants, *Plant Cell Environ.*, 39, 1087-1102, 10.1111/pce.12703, 2016.
- Ciais, P., Denning, A. S., Tans, P. P., Berry, J. A., Randall, D. A., Collatz, G. J., Sellers, P. J., White, J. W. C., Trolrier, M., Meijer, H. A. J., Francey, R. J., Monfray, P., and Heimann, M.: A three-dimensional synthesis study of  $\delta^{18}\text{O}$  in atmospheric  $\text{CO}_2$ : 1. Surface fluxes, *J. Geophys. Res. Atmos.*, 102, 5857-5872, 10.1029/96jd02360, 1997a.
- 640 Ciais, P., Tans, P. P., Denning, A. S., Francey, R. J., Trolrier, M., Meijer, H. A. J., White, J. W. C., Berry, J. A., Randall, D. A., and Collatz, G. J.: A three-dimensional synthesis study of  $\delta^{18}\text{O}$  in atmospheric  $\text{CO}_2$ : 2. Simulations with the TM2 transport model, *J. Geophys. Res. Atmos.*, 102, 10, 1997b.
- 645 Cuntz, M., Ciais, P., Hoffmann, G., Allison, C. E., Francey, R. J., Knorr, W., Tans, P. P., White, J. W. C., and Levin, I.: A comprehensive global three-dimensional model of  $\delta^{18}\text{O}$  in atmospheric  $\text{CO}_2$ : 2. Mapping the atmospheric signal *J. Geophys. Res. Atmos.*, 108, 2003a.
- Cuntz, M., Ciais, P., Hoffmann, G., and Knorr, W.: A comprehensive global three-dimensional model of  $\delta^{18}\text{O}$  in atmospheric  $\text{CO}_2$ : 1. Validation of surface processes, *J. Geophys. Res. Atmos.*, 108, 24, 2003b.
- 650 Cuntz, M.: A dent in carbon's gold standard, *Nature*, 477, 547-548, 2011.
- DiMario, R. J., Quebedeaux, J. C., Longstreth, D. J., Dassanayake, M., Hartman, M. M., and Moroney, J. V.: The cytoplasmic carbonic anhydrases  $\beta\text{CA}_2$  and  $\beta\text{CA}_4$  are required for optimal plant growth at low  $\text{CO}_2$ , *Plant Physiol.*, 171, 13, 2016.
- Evans, J. R., Sharkey, T. D., Berry, J. A., and Farquhar, G. D.: Carbon isotope discrimination measured concurrently with gas exchange to investigate  $\text{CO}_2$  diffusion in leaves of higher plants, *Funct. Plant Biol.*, 13, 11, 1986.
- 655 Fabre, N., Reiter, I. M., Becuwe-Linka, N., Genty, B., and Rumeau, D.: Characterization and expression analysis of genes encoding alpha and beta carbonic anhydrases in *Arabidopsis*, *Plant Cell Environ.*, 30, 617-629, 10.1111/j.1365-3040.2007.01651.x, 2007.

- Farquhar, D. G., and Lloyd, J.: Carbon and oxygen isotope effects in the exchange of carbon dioxide between terrestrial plants and the atmosphere, *Stable isotopes and plant carbon-water relations*, Academic Press Inc., London, 47 pp., 1993.
- 660 Farquhar, G. D., and Richards, R. A.: Isotopic composition of plant carbon correlates with water-use efficiency of wheat genotypes, *Functional Plant Biology*, 11, 11, 1984.
- Farquhar, G. D., Ehleringer, J. R., and Hubic, K. T.: Carbon isotope discrimination and photosynthesis, *Annu. Rev. Plant Physiol. Plant Mol. Biol.*, 40, 34, 1989.
- Farquhar, G. D., Lloyd, J., Taylor, J. A., Flanagan, L. B., Syvertsen, J. P., Hubick, K. T., Wong, S. C., and Ehleringer, J. R.: Vegetation effects on the isotope composition of oxygen in atmospheric CO<sub>2</sub>, *Nature*, 363, 4, 1993.
- 665 Farquhar, G. D., and Gan, K. S.: On the progressive enrichment of the oxygen isotopic composition of water along a leaf, *Plant, Cell & Environment*, 26, 18, 2003.
- Flanagan, L. B., and Ehleringer, J. R.: Ecosystem-atmosphere CO<sub>2</sub> exchange: interpreting signals of change using stable isotope ratios, *Trends Ecol. Evol.*, 13, 4, 1998.
- 670 Flexas, J., Ribas-Carbo, M., Diaz-Espejo, A., Galmes, J., and Medrano, H.: Mesophyll conductance to CO<sub>2</sub>: current knowledge and future prospects, *Plant Cell Environ.*, 31, 19, 2008.
- Flexas, J., Barbour, M. M., Brendel, O., Cabrera, H. M., Carriquí, M., Díaz-Espejo, A., Douthe, C., Dreyer, E., Ferrio, J. P., and Gago, J.: Mesophyll diffusion conductance to CO<sub>2</sub>: an unappreciated central player in photosynthesis *Plant Sci.*, 193, 14, 2012.
- Francey, R. J., and Tans, P. P.: Latitudinal variation in oxygen-18 of atmospheric CO<sub>2</sub>, *Nature*, 327, 2, 1987.
- 675 Fredeen, A. L., Gamon, J. A., and Field, C. B.: Responses of photosynthesis and carbohydrate-partitioning to limitations in nitrogen and water availability in field-grown sunflower *Plant Cell Environ.*, 14, 7, 1991.
- Friedli, H., Siegenthaler, U., Rauber, D., and Oeschger, H.: Measurements of concentration, <sup>13</sup>C/<sup>12</sup>C and <sup>18</sup>O/<sup>16</sup>O ratios of tropospheric carbon dioxide over Switzerland, *Tellus B*, 8, 1987.
- 680 Gan, K. S., Wong, S. C., Yong, J. W. H., and Farquhar, G. D.: <sup>18</sup>O spatial patterns of vein xylem water, leaf water, and dry matter in cotton leaves, *Plant Physiol.*, 130, 13, 2002.
- Gan, K. S., Wong, S. C., Yong, J. W. H., and Farquhar, G. D.: Evaluation of models of leaf water <sup>18</sup>O enrichment using measurements of spatial patterns of vein xylem water, leaf water and dry matter in maize leaves, *Plant Cell Environ.*, 26, 16, 2003.
- 685 Gillon, J., and Yakir, D.: Influence of carbonic anhydrase activity in terrestrial vegetation on the <sup>18</sup>O content of atmospheric CO<sub>2</sub>, *Science*, 291, 3, 2001.
- Gillon, J. S., and Yakir, D.: Internal conductance to CO<sub>2</sub> diffusion and C<sup>18</sup>OO discrimination in C<sub>3</sub> leaves, *Plant Physiol.*, 123, 13, 2000a.
- Gillon, J. S., and Yakir, D.: Naturally low carbonic anhydrase activity in C<sub>4</sub> and C<sub>3</sub> plants limits discrimination against C<sup>18</sup>OO during photosynthesis, *Plant Cell Environ.*, 23, 12, 2000b.
- 690 Heidenreich, J. E., and Thiemens, M. H.: A non-mass-dependent isotope effect in the production of ozone from molecular oxygen *J. CHEM. Phys.*, 78, 3, 1983.
- Heidenreich, J. E., and Thiemens, M. H.: A non-mass-dependent oxygen isotope effect in the production of ozone from molecular oxygen: The role of molecular symmetry in isotope chemistry *J. Chem. Phys.*, 84, 5, 1986.
- 695 Hoag, K. J., Still, C. J., Fung, I. Y., and Boering, K. A.: Triple oxygen isotope composition of tropospheric carbon dioxide as a tracer of terrestrial gross carbon fluxes, *Geophys. Res. Lett.*, 32, 10.1029/2004gl021011, 2005.
- Hofmann, M. E. G., Horváth, B., Schneider, L., Peters, W., Schützenmeister, K., and Pack, A.: Atmospheric measurements of Δ<sup>17</sup>O in CO<sub>2</sub> in Göttingen, Germany reveal a seasonal cycle driven by biospheric uptake, *Geochim. Cosmochim. Ac.*, 199, 143-163, 10.1016/j.gca.2016.11.019, 2017.
- 700 Johnson, K. S.: Carbon dioxide hydration and dehydration kinetics in seawater 1, *Limnology and Oceanography*, 27, 6, 1982.
- Jung, M., Schwalm, C., Migliavacca, M., Walther, S., Camps-Valls, G., Koirala, S., Anthoni, P., Besnard, S., Bodesheim, P., and Carvalhais, N.: Scaling carbon fluxes from eddy covariance sites to globe: synthesis and evaluation of the FLUXCOM approach, *Biogeosciences*, 17, 22, 2020.
- Kaiser, J., Röckmann, T., and Brenninkmeijer, C. A. M.: Contribution of mass-dependent fractionation to the oxygen isotope anomaly of atmospheric nitrous oxide, *J. Geophys. Res.*, 109, D03305, 2004.

- 705 Kammer, A., Tuzson, B., Emmenegger, L., Knohl, A., Mohn, J., and Hagedorn, F.: Application of a quantum cascade laser-based spectrometer in a closed chamber system for real-time  $\delta^{13}\text{C}$  and  $\delta^{18}\text{O}$  measurements of soil-respired  $\text{CO}_2$ , *Agr. Forest. Meteorol.*, 151, 9, 2011.
- Kawagucci, S., Tsunogai, U., Kudo, S., Nakagawa, F., Honda, H., Aoki, S., Nakazawa, T., Tsutsumi, M., and Gamo, T.: Long-term observation of mass-independent oxygen isotope anomaly in stratospheric  $\text{CO}_2$ , *Atmos. Chem. Phys.*, 8, 8, 2008.
- 710 Koren, G., Schneider, L., Velde, I. R. v. d., Schaik, E. v., Gromov, S. S., Adnew, G. A., D.J.Mrozek, Hofmann, M. E. D., Liang, M.-C., Mahata, S., Bergamaschi, P., Laan-Luijckx, I. T. v. d., Krol, M. C., Röckmann, T., and Peters, W.: Global 3-D Simulations of the Triple Oxygen Isotope Signature  $\Delta^{17}\text{O}$  in Atmospheric  $\text{CO}_2$  *J. Geophys. Res. Atmos.*, 124, 28, 2019.
- Lämmerzahl, P., Röckmann, T., and Brenninkmeijer, C. A. M.: Oxygen isotope composition of stratospheric carbon dioxide, *Geophys. Res. Lett.*, 29, 10.1029/2001gl014343, 2002.
- 715 Landais, A., Barkan, E., Yakir, D., and Luz, B.: The triple isotopic composition of oxygen in leaf water, *Geochim. Cosmochim. Ac.*, 70, 4105-4115, 10.1016/j.gca.2006.06.1545, 2006.
- Landais, A., Barkan, E., and Luz, B.: Record of  $\delta^{18}\text{O}$  and  $^{17}\text{O}$ -excess in ice from Vostok Antarctica during the last 150,000 years, *Geophys. Res. Lett.*, 35, 10.1029/2007gl032096, 2008.
- 720 Laskar, A. H., Mahata, S., and Liang, M.-C.: Identification of anthropogenic  $\text{CO}_2$  using triple oxygen and clumped isotopes, *Environ. Sci. Technol.*, 50, 18, 10.1021/acs.est.6b02989, 2016.
- Liang, M.-C., Blake, G. A., Lewis, B. R., and Yung, Y. L.: Oxygen isotopic composition of carbon dioxide in the middle atmosphere, *PNAS*, 104, 4, 2006.
- Liang, M.-C., and Mahata, S.: Oxygen anomaly in near surface carbon dioxide reveals deep stratospheric intrusion, *Sci. Rep.*, 5, 11352, 10.1038/srep11352, 2015.
- 725 Liang, M.-C., Mahata, S., Laskar, A. H., and Bhattacharya, S. K.: Spatiotemporal variability of oxygen isotope anomaly in near surface air  $\text{CO}_2$  over urban, semi-urban and ocean areas in and around Taiwan, *Aerosol Air Qual. Res.*, 17, 24, 2017a.
- Liang, M.-C., Mahata, S., Laskar, A. H., Thiemens, M. H., and Newman, S.: Oxygen isotope anomaly in tropospheric  $\text{CO}_2$  and implications for  $\text{CO}_2$  residence time in the atmosphere and gross primary productivity, *Sci Rep*, 7, 13180, 10.1038/s41598-017-12774-w, 2017b.
- Luz, B., and Barkan, E.: Variations of  $^{17}\text{O}/^{16}\text{O}$  and  $^{18}\text{O}/^{16}\text{O}$  in meteoric waters *Geochim. Cosmochim. Ac.*, 74, 10, 2010.
- Lyons, J. R.: Transfer of mass-independent fractionation in ozone to other oxygen-containing radicals in the atmosphere, *Geophys. Res. Lett.*, 28, 3231-3234, 10.1029/2000gl012791, 2001.
- 735 Mahata, S., Bhattacharya, S. K., Wang, C. H., and Liang, M.-C.: Oxygen isotope exchange between  $\text{O}_2$  and  $\text{CO}_2$  over hot platinum: an innovative technique for measuring  $\Delta^{17}\text{O}$  in  $\text{CO}_2$ , *Anal. Chem.*, 85, 6894-6901, 10.1021/ac4011777, 2013.
- Matsuhisa, Y., Goldsmith, J. R., and Clayton, R. N.: Mechanisms of hydrothermal crystallization of quartz at 250 °C and 15 kbar, *Geochimica et Cosmochimica Acta*, 42, 9, 1978.
- McManus, J. B., Nelson, D. D., Shorter, J. H., Jimenez, R., Herndon, S., Saleska, S., and Zahniser, M.: A high precision pulsed quantum cascade laser spectrometer for measurements of stable isotopes of carbon dioxide, *J. Mod. Optic.*, 52, 12, 2005.
- 740 Meijer, H., and Li, W.: The use of electrolysis for accurate  $^{17}\text{O}$  and  $^{18}\text{O}$  isotope measurements in water isotopes, *Isotopes Environ. Health Stud.*, 34, 20, 1998.
- Miller, J. B., Yakir, D., White, J. W. C., and Tans, P. P.: Measurement of  $^{18}\text{O}/^{16}\text{O}$  in the soil-atmosphere  $\text{CO}_2$  flux, *Global Biogeochem. Cy.*, 13, 13, 1999.
- Miller, M. F.: Isotopic fractionation and the quantification of  $^{17}\text{O}$  anomalies in the oxygen three-isotope system: an appraisal and geochemical significance, *Geochim. Cosmochim. Acta*, 66, 8, 2002.
- 745 Miller, R. F., Berkshire, D. C., Kelley, J. J., and Hood, D. W.: Method for determination of reaction rates of carbon dioxide with water and hydroxyl ion in seawater, *Environ. Sci. Tech.*, 5, 6, 1971.
- Mills, G. A., and Urey, H. C.: The kinetics of isotopic exchange between carbon dioxide, bicarbonate ion, carbonate ion and water, *J. AM. Chem. Soc.*, 62, 7, 1940.
- 750 Nelson, D. D., McManus, J. B., Herndon, S., Zahniser, M. S., Tuzson, B., and Emmenegger, L.: New method for isotopic ratio measurements of atmospheric carbon dioxide using a 4.3  $\mu\text{m}$  pulsed quantum cascade laser *Appl. Phys. B-Lasers. O.*, 90, 8, 2008.

- Osborn, H. L., Alonso-Cantabrana, H., Sharwood, R. E., Covshoff, S., Evans, J. R., Furbank, R. T., and von Caemmerer, S.: Effects of reduced carbonic anhydrase activity on CO<sub>2</sub> assimilation rates in *Setaria viridis*: a transgenic analysis, *J. Exp. Bot.*, 68, 11, 2017.
- 755 Pack, A., and Herwartz, D.: The triple oxygen isotope composition of the Earth mantle and understanding  $\Delta^{17}\text{O}$  variations in terrestrial rocks and minerals *Earth. Planet. Sc. Lett.*, 390, 7, 2014.
- Peylin, P., Ciais, P., Denning, A. S., Tans, P. P., Berry, J. A., and White, J. W.: A 3-dimensional study of  $\delta^{18}\text{O}$  in atmospheric CO<sub>2</sub>: contribution of different land ecosystems, *Tellus B*, 51, 25, 1999.
- 760 Pons, T. L., and Welschen, R. A. M.: Overestimation of respiration rates in commercially available clamp-on leaf chambers. Complications with measurement of net photosynthesis, *Plant, Cell and Environment*, 25, 5, 2002.
- Pons, T. L., Flexas, J., von Caemmerer, S., Evans, J. R., Genty, B., Ribas-Carbo, M., and Brugnoli, E.: Estimating mesophyll conductance to CO<sub>2</sub>: methodology, potential errors, and recommendations *J. Exp. Bot.*, 60, 7, 2009.
- 765 Schaefer, K., Collatz, G. J., Tans, P., Denning, A. S., Baker, I., Berry, J., Prihodko, L., Suits, N., and Philpott, A.: Combined simple biosphere/Carnegie-Ames-Stanford approach terrestrial carbon cycle model, *J. Geophys. Res.-Biogeo.*, 113, 2008.
- Shaheen, R., Janssen, C., and Röckmann, T.: Investigations of the photochemical isotope equilibrium between O<sub>2</sub>, CO<sub>2</sub> and O<sub>3</sub>, *Atmos. Chem. Phys.*, 7, 495-509, 2007.
- 770 Sharp, Z. D., Wostbrock, J. A. G., and Pack, A.: Mass-dependent triple oxygen isotope variations in terrestrial materials, *Geochemical Perspectives Letters*, 27-31, 10.7185/geochemlet.1815, 2018.
- Shrestha, A., Song, X., and Barbour, M. M.: The temperature response of mesophyll conductance, and its component conductances, varies between species and genotypes, *Photosynth. Res.*, 18, 2019.
- Silverman, D. N.: Carbonic anhydrase: Oxygen-18 exchange catalyzed by an enzyme with rate-contributing Proton-transfer steps *Methods in enzymology*, Elsevier, 1982.
- 775 Sitch, S., Friedlingstein, P., Gruber, N., Jones, S. D., Murray-Tortarolo, G., Ahlström, A., Doney, S. C., Graven, H., Heinze, C., and Huntingford, C.: Recent trends and drivers of regional sources and sinks of carbon dioxide, *Biogeosciences*, 12, 26, 2015.
- Still, C. J., Berry, J. A., Collatz, G. J., and DeFries, R. S.: Global distribution of C<sub>3</sub> and C<sub>4</sub> vegetation: Carbon cycle implications, *Global Biogeochem. Cy.*, 17, 6-1-6-14, 10.1029/2001gb001807, 2003.
- 780 Thiemens, M. H., Jackson, T., Mauersberger, K., Schüler, B., and Morton, J.: Oxygen isotope fractionation in stratospheric CO<sub>2</sub>, *Geophys. Res. Lett.*, 18, 669-672, 1991.
- Thiemens, M. H., Jackson, T., Zipf, E. C., Erdman, P. W., and Egmond, C. v.: Carbon dioxide and oxygen isotope anomalies in the mesosphere and stratosphere, *Science* 270, 3, 1995.
- 785 Thiemens, M. H.: Mass-independent isotope effects in planetary atmospheres and the early solar system, *Science*, 283, 4, 1999.
- Thiemens, M. H.: History and Applications of Mass-Independent Isotope Effects, *Annu. Rev. Earth Planet. Sci.*, 34, 62, 10.1146/, 2006.
- Thiemens, M. H., Chakraborty, S., and Jackson, T. L.: Decadal  $\Delta^{17}\text{O}$  record of tropospheric CO<sub>2</sub>: Verification of a stratospheric component in the troposphere, *J. Geophys. Res. Atmos.*, 119, 8, 10.1002/2013JD020317, 2013.
- 790 Thiemens, M. H., and Heidenreich, J.E. III: Mass-independent fractionation of oxygen: a novel isotope effect and its possible cosmochemical implications, *Science* 10.1126/science.219.4588.1073, 1983.
- Thiemens, M. K., Chakraborty, S., and Jackson, T. L.: Decadal  $\Delta^{17}\text{O}$  record of tropospheric CO<sub>2</sub>: Verification of a stratospheric component in the troposphere, *J. Geophys. Res. Atmos.*, 119, 8, 2014.
- 795 Tuzson, B., Mohn, J., Zeeman, M. J., Werner, R. A., Eugster, W., Zahniser, M. S., Nelson, D. D., McManus, J. B., and Emmenegger, L.: High precision and continuous field measurements of  $\delta^{13}\text{C}$  and  $\delta^{18}\text{O}$  in carbon dioxide with a cryogen-free QCLAS, *Appl. Phys. B-Lasers O.*, 92, 7, 2008.
- Uemura, R., Barkan, E., Abe, O., and Luz, B.: Triple isotope composition of oxygen in atmospheric water vapor, *Geophys. Res. Lett.*, 37, L04402, 2010.
- Wang, X.-F., and Yakir, D.: Using stable isotopes of water in evapotranspiration studies, *Hydrol. Process.*, 14, 14, 2000.
- 800 Wassenaar, L. I., Terzer-Wassmuth, S., Douence, C., Araguas-Araguas, L., Aggarwal, P. K., and Coplen, T. B.: Seeking excellence: An evaluation of 235 international laboratories conducting water isotope analyses by isotope-ratio and laser-absorption spectrometry, *Rapid Commun Mass Spectrom*, 32, 393-406, 10.1002/rcm.8052, 2018.

- Weijde, T. v. d., Kamei, K. C. L. A., Torres, A. F., Vermerris, W., Dolstra, O., Visser, R. G. F., and Trindade, L. M.: The potential of C<sub>4</sub> grasses for cellulosic biofuel production, *Front. Plant. Sci.*, 4, 107, 2013.
- 805 Welp, L. R., Keeling, R. F., Meijer, H. A. J., Bollenbacher, A. F., Piper, S. C., Yoshimura, K., Francey, R. J., Allison, C. A., and Wahlen, M.: Interannual variability in the oxygen isotopes of atmospheric CO<sub>2</sub> driven by El Niño, *Nature*, 477, 579-582, 10.1038/nature10421, 2011.
- West, A. G., Patrickson, S. J., and Ehleringer, J. R.: Water extraction times for plant and soil materials used in stable isotope analysis, *Rapid Commun Mass Spectrom*, 20, 1317-1321, 10.1002/rcm.2456, 2006.
- 810 Wiegel, A. A., Cole, A. S., Hoag, K. J., Atlas, E. L., Schauffler, S. M., and Boering, K. A.: Unexpected variations in the triple oxygen isotope composition of stratospheric carbon dioxide, *PNAS*, 110, 17680-17685, 10.1073/pnas.1213082110, 2013.
- Wingate, L., Ogée, J., Cuntz, M., Genty, B., Reiter, I., Seibt, U., Yakir, D., Maseyk, K., Pendall, E. G., Barbour, M. M., Mortazavi, B., Burlett, R., Peylin, P., Miller, J., Mencuccini, M., Shim, J. H., Hunt, J., and Grace, J.: The impact of soil microorganisms on the global budget of δ<sup>18</sup>O in atmospheric CO<sub>2</sub>, *Proceedings of the National Academy of Sciences*, 106, 4, 2009.
- 815 Yakir, D.: Oxygen-18 of leaf water: a crossroad for plant-associated isotopic signals, *Stable isotopes: integration of biological, ecological and geochemical processes*, 21, 1998.
- Yakir, D., and Sternberg, L. S. L.: The use of stable isotopes to study ecosystem gas exchange, *Oecologia*, 123, 4, 2000.
- 820 Young, E. D., Galy, A., and Nagahara, H.: Kinetic and equilibrium mass-dependent isotope fractionation laws in nature and their geochemical and cosmochemical significance, *Geochim. Cosmochim. Ac.*, 66, 9, 2002.
- Yung, Y. L., DoMore, W. B., and Pinto, J. P.: Isotopic exchange between carbon dioxide and ozone via O (<sup>1</sup>D) in the stratosphere, *Geophys. Res. Lett.*, 18, 3, 1991.
- Yung, Y. L., Lee, A. Y. T., Irion, F. W., DeMore, W. B., and Wen, J.: Carbon dioxide in the atmosphere: Isotopic exchange with ozone and its use as a tracer in the middle atmosphere, *J. Geophys. Res.*, 102, 9, 1997.
- 825




830 Figure 1 Schematic for mass-dependent isotope fractionation process that affects the  $\Delta^{17}\text{O}$  of the CO<sub>2</sub> and H<sub>2</sub>O during the photosynthetic gas exchange (not to scale). The triple oxygen isotope relationship for the individual isotope fractionation processes (both kinetic and equilibrium fractionation) are assigned with  $\theta$ .


835  $\theta_{trans}=0.522-0.008 \times h$ , where  $h$  is relative humidity (Landais et al., 2006), in this study the humidity is 75 %,  $\theta_{trans}=0.516$ .  $\theta_{CO_2-H_2O}$  (Barkan and Luz, 2012),  $\theta_{CO_2-diff}$  (Young et al., 2002),  $\theta_{H_2O(v)-H_2O(l)}$  (Barkan and Luz, 2005) and  $\theta_{H_2O(v)-diff}$  (Barkan and Luz, 2007). Where  $v$  and  $l$  for vapor and liquid water, respectively;  $\epsilon^{18O}$  is enrichment or depletion in  $^{18}O$  isotope composition due to the corresponding isotope fractionation process; *diff* and *trans* stand for diffusion and transpiration, respectively.

840

845




850 **Figure 2** Schematic diagram of the leaf cuvette experimental setup. IRGA stands for the infrared gas analyzer, WVSS is the water vapor standard source, WVIA is the water vapor isotope analyzer, N-CO<sub>2</sub> is normal CO<sub>2</sub>, E-CO<sub>2</sub> is <sup>17</sup>O-enriched CO<sub>2</sub>.



855


Figure 3 a)  $\Delta_A^{18}\text{O}_{\text{obs}}$  during photosynthesis for two  $\text{C}_3$  plants, sunflower (circles) and ivy (triangles) and  $\text{C}_4$  plant maize (stars) as a function of  $c_m/c_a$ . The solid lines show results from the leaf cuvette model, where  $\delta^{18}\text{O}$  of the  $\text{CO}_2$  entering the cuvette is 30.47‰. b)  $\Delta_A \Delta^{17}\text{O}$  of  $\text{CO}_2$  as a function of  $c_m/c_a$  for isotopically different  $\text{CO}_2$  gases entering the cuvette (color bar shows  $\Delta^{17}\text{O}_e$ ) for sunflower (circles), ivy (triangles) and maize (stars).  $\Delta_A \Delta^{17}\text{O}$  values calculated using the leaf cuvette model are shown as solid lines in corresponding colors ( $\Delta^{17}\text{O}_e$  values given in the legend). The shaded areas indicate the  $c_m/c_a$  ranges for  $\text{C}_4$  and  $\text{C}_3$  plants and the vertical dashed lines indicate the mean  $c_m/c_a$  ratio used for extrapolating from the leaf scale to the global scale. Solid line are leaf cuvette model results for the corresponding  $c_m/c_a$  ratio.

865






870 **Figure 4** a) Dependency of  $\Delta_A \Delta^{17}\text{O}$  on the relative difference on the  $\Delta^{17}\text{O}$   $\text{CO}_2$  entering the leaf and the  
 875  $\Delta^{17}\text{O}$  of  $\text{CO}_2$  in equilibrium with leaf water against  $c_m/c_a$  ratio. b) dependency of  $\Delta_A \Delta^{17}\text{O}$  on the difference  
 between the  $\Delta^{17}\text{O}$  of  $\text{CO}_2$  entering the cuvette and the  $\Delta^{17}\text{O}$  of leaf water at the evaporation site color  
 coded for different  $c_m/c_a$  ratios. The solid lines are results of the leaf cuvette model for different  $c_m/c_a$   
 ratios stated in the legend. The dashed vertical black line indicates the difference between the global  
 average  $\Delta^{17}\text{O}$  value for  $\text{CO}_2$  (-0.168 ‰) and leaf water (-0.067 ‰) (Koren et al., 2019). The gray and  
 yellow horizontal dashed lines indicate  $\Delta_A \Delta^{17}\text{O}$  of  $\text{C}_4$  and  $\text{C}_3$  plants for  $c_m/c_a$  ratio of 0.3 and 0.7,  
 respectively globally.



880 **Figure 5** a) and b)  $\Delta^{17}\text{O}_a$  as a function of  $c_m/c_a$  for various values of  $\Delta^{17}\text{O}_e$  (see legend) for  $\Delta^{17}\text{O}_{\text{wes}} = -$   
 0.122 ‰ in a) and  $\Delta^{17}\text{O}_{\text{wes}} = 0.300$  ‰ in b). c) and d) show the corresponding values for  $\Delta_A \Delta^{17}\text{O}$ .  $\Delta^{17}\text{O}_{\text{global}}$   
 is the global average  $\Delta^{17}\text{O}$  value for atmospheric  $\text{CO}_2$  (Koren et al., 2019). When  $\Delta^{17}\text{O}$  of  $\text{CO}_2$  entering  
 the cuvette is approximately 0.2 ‰ lower than the  $\Delta^{17}\text{O}$  of leaf water at the  $\text{CO}_2$ - $\text{H}_2\text{O}$  exchange site,  $\Delta^{17}\text{O}$   
 of the  $\text{CO}_2$  leaving the cuvette does not change when  $c_m/c_a$  vary.



885

Figure 6 Isotopic composition of various relevant oxygen reservoirs that affect the  $\Delta^{17}\text{O}$  of atmospheric  $\text{CO}_2$  during photosynthesis: irrigation water (grey triangle), calculated leaf water at the evaporation site (brown circles), measured bulk leaf water (brown star),  $\text{CO}_2$  entering the cuvette (black circles),  $\text{CO}_2$  leaving the leaf cuvette (green circles),  $\text{CO}_2$  equilibrated with leaf water at the evaporation site (blue circles),  $\text{CO}_2$  equilibrated with bulk leaf water (blue stars).  $\Delta^{17}\text{O}$  is calculated with  $\lambda=0.528$ .

890

895

900

905

910 Table 1: Summary of gas exchange parameters and isotopic composition of maize, sunflower and ivy. Mole fraction at the site of exchange ( $c_m$ ) is calculated assuming complete isotopic equilibrium with the water at the CO<sub>2</sub>-H<sub>2</sub>O exchange site. The water at the CO<sub>2</sub>-H<sub>2</sub>O exchange site is assumed the same as the isotopic composition at the site of evaporation. Number in the parenthesis are the standard deviation of the mean ( $1\sigma$ ).

Parameter		Sunflower	Ivy	Maize	Irradiance ( $\mu\text{mol m}^{-2} \text{s}^{-1}$ )
$A_n$	$\mu\text{mol mol}^{-1} \text{m}^{-2} \text{s}^{-1}$	18(0.7)	12(0.7)	17(2)	300
		29(2)	15(2)	32(2)	1200
$g_s$	$\text{mol m}^{-2} \text{s}^{-1}$	0.45(0.14)	0.11(0.02)	0.08(0.01)	300
		0.40(0.04)	0.15(0.03)	0.16(0.02)	1200
$\delta^{18}\text{O}_e$	‰	27.26 to 31.80	28.28 to 30.48	27.26 to 30.48	
$\Delta^{17}\text{O}_e$	‰	-0.227 to 0.409	-0.215 to 0.435	-0.215 to 0.310	
$\delta^{18}\text{O}_a$	‰	33.25 to 43.87	32.64 to 35.86	34.04 to 29.764	
$\Delta^{17}\text{O}_a$	‰	-0.333 to 0.163	-0.276 to 0.327	-0.270 to 0.296	
$\Delta_A^{18}\text{O}_{\text{obs}}$	‰	57.12(4.70)	22.20(1.32)	17.23(1.32)	300
		34.48(3.25)	24.35(3.09)	12.78(0.83)	1200
$\Delta_A \Delta^{17}\text{O}_{\text{obs}}$	‰	-2.61 to -0.43	-1.03 to -0.19	-0.36 to -0.09	
$\delta^{18}\text{O}_m$	‰	52.02(1.24)	47.17(1.17)	52.62(0.52)	300
		52.62(1.42)	51.09(1.76)	55.15(1.55)	1200
$\Delta^{17}\text{O}_m$	‰	-0.41(0.001)	-0.35(0.001)	-0.40(0.01)	300
		-0.41(0.01)	-0.38(0.02)	-0.42(0.02)	1200
$c_a$	ppm	402 (3)	403 (3)	403 (3)	
$c_i$	ppm	357(10)	284(0.1)	194(20)	300
		323(10)	301(13)	194(15)	1200
$c_c$	ppm	277(15)	188(30)		300
		201(42)	163(21)		1200
$c_m$	ppm	320(10)	220(10)	134(15)	300
		252(27)	214(12)	88 (17)	1200

915

Table 2 List of symbols and variables

Symbol	description	Unit/calculation/value
<b>Gas exchange</b>		
$A_n$	Rate of CO <sub>2</sub> assimilation	$\frac{u_e}{s} \left( c_e - c_a \left( \frac{1-w_e}{1-w_a} \right) \right)$ , mol m <sup>-2</sup> s <sup>-2</sup>

$E$	Transpiration rate	$\frac{u_e}{s} \left( \frac{w_a - w_e}{1 - w_a} \right)$ , mol m <sup>-2</sup> s <sup>-2</sup>
$w_i$	Mole fraction of water vapour inside leaf	$\frac{613.65 \times e^{\left( \frac{17.502 \times T_{leaf}}{240.97 + T_{leaf}} \right)} \times 10^{-5}}{P}$ , mol mol <sup>-1</sup>
$w_a$	Mole fraction of water vapour leaving the cuvette /leaf surrounding	WVIA / IRGA, mol mol <sup>-1</sup>
$w_e$	Mole fraction of water vapour entering the cuvette	WVIA/IRGA, mol mol <sup>-1</sup>
$c_e$	Mole fraction of CO <sub>2</sub> entering the cuvette	IRGA, mol mol <sup>-1</sup>
$c_a$	Mole fraction of CO <sub>2</sub> in the leaf surrounding/ leaving the cuvette	IRGA, mol mol <sup>-1</sup>
$u_e$	Flow rate of air entering the cuvette	mol s <sup>-1</sup>
$s$	Surface area of the leaf inside the cuvette	m <sup>2</sup>
$P$	Atmospheric pressure	bar
$T_{leaf}$	Leaf temperature	°C
$g_{s(H_2O)}$	Stomatal conductance for water vapour	$\frac{g_{H_2O}^t \times g_b(H_2O)}{g_b(H_2O) - g_{H_2O}^t}$
$g_{b(H_2O)}$	Boundary layer conductance for water vapour	Calibrated for the cuvette we used
$g_{H_2O}^t$	Conductance for water vapor through the boundary layer and stomata	$E \left( \frac{1 - \left( \frac{w_i + w_a}{2} \right)}{w_i - w_a} \right)$ , mol m <sup>-2</sup> s <sup>-1</sup>
$g_s$	Stomatal conductance for CO <sub>2</sub>	$\frac{g_s(H_2O)}{1.6}$
$g_b$	Boundary conductance for CO <sub>2</sub>	$\frac{g_b(H_2O)}{1.37}$
$g_{CO_2}^t$	Conductance for CO <sub>2</sub> through the boundary layer and stomata	$\frac{g_s \times g_b}{g_s + g_b}$
$\Gamma^*$	CO <sub>2</sub> compensation point	45 μmol m <sup>-2</sup> s <sup>-1</sup>
$g_{m13}$	CO <sub>2</sub> conductance from intercellular air space to the site of carboxylation calculated using ΔA <sup>13</sup> C (for C <sub>3</sub> plants only)	mol m <sup>-2</sup> s <sup>-1</sup> bar <sup>-1</sup>
$g_{m18}$	CO <sub>2</sub> conductance from intercellular air space to CO <sub>2</sub> -H <sub>2</sub> O exchange site calculated using ΔA <sup>18</sup> O	mol m <sup>-2</sup> s <sup>-1</sup> bar <sup>-1</sup>
$g_{m17}$	CO <sub>2</sub> conductance from intercellular air space to CO <sub>2</sub> -H <sub>2</sub> O exchange site calculated using ΔA <sup>17</sup> O	mol m <sup>-2</sup> s <sup>-1</sup> bar <sup>-1</sup>
$g_{m\Delta 17}$	CO <sub>2</sub> conductance from intercellular air space to CO <sub>2</sub> -H <sub>2</sub> O exchange site calculated using ΔAΔ <sup>17</sup> O	mol m <sup>-2</sup> s <sup>-1</sup> bar <sup>-1</sup>
$c_i$	Mole fraction of CO <sub>2</sub> in the intercellular air space	$\frac{\left( \frac{g_{CO_2}^t}{2} \right) c_a - A_n}{\left( \frac{g_{CO_2}^t}{2} \right)}$ mol mol <sup>-1</sup>
$c_s$	Mole fraction of CO <sub>2</sub> at the leaf surface	$c_a - \frac{A_n}{g_b}$
$c_m$	Mole fraction of CO <sub>2</sub> at the site of CO <sub>2</sub> -H <sub>2</sub> O exchange	mol mol <sup>-1</sup>
$c_c$	Mesophyll conductance to the chloroplast (for C <sub>3</sub> plants)	$c_i - \frac{A_n}{g_{m13}}$ mol mol <sup>-1</sup>
$i^{13}$	Ternary correction for <sup>13</sup> CO <sub>2</sub>	$\frac{(1 + a_{13bs})E}{2g_{CO_2}^t}$
$i^{18}$	Ternary correction for C <sup>18</sup> OO	$\frac{(1 + a_{18bs})E}{2g_{CO_2}^t}$
$i^{17}$	Ternary correction for C <sup>17</sup> OO	$\frac{(1 + a_{17bs})E}{2g_{CO_2}^t}$
$R_D$	Dark respiration rate	0.8 μmol m <sup>-2</sup> s <sup>-1</sup>
$R_L$	Day respiration rate	0.5 × R <sub>D</sub> μmol m <sup>-2</sup> s <sup>-1</sup>
<b>Oxygen and carbon isotope effects</b>		
$\epsilon_k^{18}$	Kinetic fractionation of water vapour in air	$\frac{28g_b + 19g_s}{g_b + g_s}$ , ‰
$\epsilon_{equ}^{18}$	Equilibrium fractionation between liquid and gas phase of water vapor	$2.644 - 3.206 \left( \frac{10^3}{T_{leaf}} \right) + 1.534 \left( \frac{10^6}{T_{leaf}^2} \right)$ , ‰
$a_{13bs}$	Weighted fractionation for <sup>13</sup> COO as CO <sub>2</sub> diffuses through the boundary layer and stomata	$\frac{(c_s - c_i)a_{13s} + (c_a - c_s)a_{13b}}{c_a - c_i}$ ‰

$a_{17bs}$	Weighted fractionation for $C^{17}OO$ as $CO_2$ diffuses through the boundary layer and stomata	$\frac{(c_s - c_i)a_{17s} + (c_a - c_s)a_{17b}}{c_a - c_i}$ ‰
$a_{18bs}$	Weighted fractionation for $C^{18}OO$ as $CO_2$ diffuses through the boundary layer and stomata	$\frac{(c_s - c_i)a_{18s} + (c_a - c_s)a_{18b}}{c_a - c_i}$ ‰
$a_{13bs}$	Weighted fractionation for $^{13}COO$ as $CO_2$ diffuses through the boundary layer and stomata	$\frac{(c_s - c_i)a_{13s} + (c_a - c_s)a_{13b}}{c_a - c_i}$ , ‰
$a_{18bs}$	Weighted fractionation for $C^{18}OO$ as $CO_2$ diffuses through the boundary layer and stomata	$\frac{(c_s - c_i)a_{18s} + (c_a - c_s)a_{18b}}{c_a - c_i}$ , ‰
$a_{17bs}$	Weighted fractionation for $C^{17}OO$ as $CO_2$ diffuses through the boundary layer and stomata	$\frac{(c_s - c_i)a_{17s} + (c_a - c_s)a_{17b}}{c_a - c_i}$ , ‰
$\bar{a}_{17}$	Weighted fractionation of $C^{17}OO$ as it diffuses through the boundary layer, stomata and liquid phase in series	$\frac{(c_i - c_m)a_{17w} + (c_s - c_i)a_{17s} + (c_a - c_s)a_{17b}}{c_a - c_m}$ , ‰
$\bar{a}_{18}$	Weighted fractionation of $C^{18}OO$ as it diffuses through the boundary layer, stomata and liquid phase in series	$\frac{(c_i - c_m)a_{18w} + (c_s - c_i)a_{18s} + (c_a - c_s)a_{18b}}{c_a - c_m}$ , ‰
$a_{13b}$	Fractionation in $^{13}CO_2$ as $CO_2$ diffuses through the boundary layer	2.9‰
$a_{13s}$	Fractionation in $^{13}CO_2$ as $CO_2$ diffuses through the stomata	4.4‰
$a_m$	Fractionation factor for dissolution and diffusion through water	1.8‰
$f$	Fractionation factor for photorespiration (decarboxylation of glycine)	16‰
$e$	Fractionation factor for day respiration	$R_p + e^*$ , ‰
$e^*$	Apparent fractionation for day respiration	$\delta^{13}C_p - \Delta_A^{13}C - \delta^{13}C_{substrate}$ , ‰
$b$	Fractionation factor for uptake by RubisCO	29‰
$\alpha_f$	Fractionation due to photorespiration (decarboxylation of glycine)	$1+f$
$\alpha_e$	Fractionation due to day respiration	$1+e$
$\alpha_b$	Fractionation due to uptake by RubisCO	$1+b$
$a_{17b}$	Fractionation of $C^{17}OO$ as $CO_2$ diffuses through the boundary layer	2.9‰
$a_{17s}$	Fractionation in $C^{17}OO$ as $CO_2$ diffuses through stomata	4.4‰
$a_{18b}$	Fractionation of $C^{18}OO$ as $CO_2$ diffuses through the boundary layer	5.8‰
$a_{18s}$	Fractionation in $C^{18}OO$ as $CO_2$ diffuses through stomata	8.8‰
$a_{17w}$	Fractionation in $C^{17}OO$ due to diffusion and dissolution in water	0.382‰
$a_{18w}$	Fractionation in $C^{18}OO$ due to diffusion and dissolution in water	0.8‰
$\epsilon^{18}w$	Equilibrium fractionation of $CO_2$ and water for $C^{18}OO$	$\frac{17604}{T_{leaf}} - 17.93$ , ‰
$\epsilon^{18}k$	kinetic fractionation of water vapor in air	$\frac{28 \times g_b + 19 \times g_s}{g_b + g_s}$
$\epsilon^{18}equ$	equilibrium fractionation between liquid and gas phase water	$2.644 - 3.206 \times \left(\frac{10^3}{T}\right) + 1.534 \times \left(\frac{10^6}{T}\right)$
<b>Isotopic composition</b>		
$\delta^{17}O_A$	$\delta^{17}O$ of the assimilated $CO_2$	$\frac{\delta^{17}O_a - \Delta_A^{17}O}{\Delta_A^{17}O + 1} = \delta^{17}O_a - \frac{c_e}{c_e - c_a} (\delta^{17}O_a - \delta^{17}O_e)$
$\delta^{18}O_A$	$\delta^{18}O$ of the assimilated $CO_2$	$\frac{\delta^{18}O_a - \Delta_A^{18}O}{\Delta_A^{18}O + 1} = \delta^{18}O_a - \frac{c_e}{c_e - c_a} (\delta^{18}O_a - \delta^{18}O_e)$
$\delta^{17}O_{io}$	$\delta^{17}O$ of $CO_2$ in the intercellular air space ignoring ternary correction	$\delta^{17}O_a \left(1 - \frac{c_a}{c_i}\right) (1 + a_{17bs}) + \frac{c_a}{c_i} (\delta^{17}O_a - a_{17bs}) + a_{17bs}$ , ‰

$\delta^{18}O_{io}$	$\delta^{18}O$ of CO <sub>2</sub> in the intercellular air space ignoring ternary correction	$\delta^{18}O_A \left(1 - \frac{c_a}{c_i}\right) (1 + a18bs) + \frac{c_a}{c_i} (\delta^{18}O_a - a18bs) + a18bs$ , %
$\delta^{17}O_i$	$\delta^{17}O$ of CO <sub>2</sub> in the intercellular air space	$\frac{\delta^{17}O_{io} + t^{17} \left( \delta^{17}O_A \left( \frac{c_a+1}{c_i} \right) - \delta^{17}O_a \frac{c_a}{c_i} \right)}{1+t^{17}}$ , ‰
$\delta^{18}O_i$	$\delta^{18}O$ of CO <sub>2</sub> in the intercellular air space	$\frac{\delta^{18}O_{io} + t^{18} \left( \delta^{18}O_A \left( \frac{c_a+1}{c_i} \right) - \delta^{18}O_a \frac{c_a}{c_i} \right)}{1+t^{18}}$ , ‰
$\delta^{18}O_{trans}$	$\delta^{18}O$ of transpired water vapour	$\left( \frac{w_a}{w_a - w_e} \right) (\delta^{18}O_{wa} - \delta^{18}O_{we}) + \delta^{18}O_{we}$ , ‰
$\delta^{18}O_{wes}$	$\delta^{18}O$ of water at the evaporation site	$\delta^{18}O_{wes} = \delta^{18}O_{trans} + \varepsilon^{18}_k + \varepsilon^{18}_{equ} + \frac{w_a}{w_i} \times (\delta^{18}O_{wa} - \varepsilon^{18}_k + \delta^{18}O_{trans})$
$\delta^{17}O_m$	$\delta^{17}O$ of CO <sub>2</sub> at the site of CO <sub>2</sub> -H <sub>2</sub> O exchange	$(\delta^{17}O_{wes} + 1) \times (1 + \varepsilon^{17}_w) - 1$ , ‰
$\delta^{18}O_m$	$\delta^{18}O$ of CO <sub>2</sub> at the site of CO <sub>2</sub> -H <sub>2</sub> O exchange	$(\delta^{18}O_{wes} + 1) \times (1 + \varepsilon^{18}_w) - 1$ , ‰
$\delta^{13}C_{substrate}$	Isotope ( <sup>13</sup> C) ratio of substrate used for dark respiration	$\frac{\delta^{13}C_a - \Delta A^{13}C}{\Delta A^{13}C + 1}$ , ‰
$\Delta A^{13}C$	<sup>13</sup> C-photosynthetic discrimination	$\frac{\zeta(\delta^{13}C_a - \delta^{13}C_e)}{1 + \delta^{13}C_a - \zeta(\delta^{13}C_a - \delta^{13}C_e)}$ , ‰
$\Delta A^{13}C_{obs}$	<sup>13</sup> C-photosynthetic discrimination (Farquhar model)	$\left( \frac{1}{1-t} \right) \left[ a_{13bs} \frac{c_a - c_i}{c_a} \right] + \left( \frac{1+t}{1-t} \right) \left[ a_m \frac{c_i - c_c}{c_a} + b \frac{c_c}{c_a} - \frac{\alpha_b}{\alpha_e} e \frac{R_D}{R_D + A_n} \frac{c_c - \Gamma^*}{c_a} - \frac{\alpha_b}{\alpha_f} f \frac{\Gamma^*}{c_a} \right]$
$\Delta A^{13}C_i$	<sup>13</sup> C-photosynthetic discrimination (assuming no mesophyll conductance, i.e. c <sub>i</sub> =c <sub>c</sub> )	$\left( \frac{1}{1-t} \right) \left[ \bar{a} \frac{c_a - c_i}{c_a} \right] + \left( \frac{1+t}{1-t} \right) \left[ b \frac{c_i}{c_a} - \frac{\alpha_b}{\alpha_e} e \frac{R_D}{R_D + A_n} \frac{c_i - \Gamma^*}{c_a} - \frac{\alpha_b}{\alpha_f} f \frac{\Gamma^*}{c_a} \right]$
$\Delta A^{18}O$	<sup>18</sup> O-photosynthetic discrimination	$\frac{\zeta(\delta^{18}O_a - \delta^{18}O_e)}{1 + \delta^{18}O_a - \zeta(\delta^{18}O_a - \delta^{18}O_e)}$ , ‰
$\Delta A^{17}O$	<sup>17</sup> O-photosynthetic discrimination	$\frac{\zeta(\delta^{17}O_a - \delta^{17}O_e)}{1 + \delta^{17}O_a - \zeta(\delta^{17}O_a - \delta^{17}O_e)}$ , ‰
$\Delta A^{17}O_{FM}$	Farquhar model for <sup>17</sup> O-photosynthetic discrimination	$\frac{\bar{a}_{17} + \frac{c_m}{c_a - c_m} \delta^{17}O_{ma}}{1 - \frac{c_m}{\alpha - c_m} \delta^{17}O_{ma}}$ ‰
$\Delta A^{18}O_{FM}$	Farquhar model for <sup>18</sup> O-photosynthetic discrimination	$\frac{\bar{a}_{18} + \frac{c_m}{c_a - c_m} \delta^{18}O_{ma}}{1 - \frac{c_m}{\alpha - c_m} \delta^{18}O_{ma}}$ ‰
$\delta^{17}O_e$	$\delta^{17}O$ of CO <sub>2</sub> entering the cuvette	‰
$\delta^{17}O_a$	$\delta^{17}O$ of CO <sub>2</sub> leaving the cuvette	‰
$\delta^{18}O_e$	$\delta^{18}O$ of CO <sub>2</sub> entering the cuvette	‰
$\delta^{18}O_a$	$\delta^{18}O$ of CO <sub>2</sub> leaving the cuvette	‰
$\delta^{17}O_{ma}$	$\delta^{17}O$ of CO <sub>2</sub> equilibrated with the leaf water at the evaporating site relative to the CO <sub>2</sub> leaving the cuvette	$\frac{\delta^{17}O_m - \delta^{17}O_a}{1 - \delta^{18}O_a}$ , ‰
$\delta^{18}O_{ma}$	$\delta^{18}O$ of CO <sub>2</sub> equilibrated with the leaf water at the evaporating site relative to the CO <sub>2</sub> leaving the cuvette	$\frac{\delta^{18}O_m - \delta^{18}O_a}{1 - \delta^{18}O_a}$ , ‰
$\delta^{18}O_{we}$	$\delta^{18}O$ of water vapour entering the cuvette	WVIA, ‰
$\delta^{18}O_{wa}$	$\delta^{18}O$ of water vapour leaving the cuvette/leaf surrounding	WVIA, ‰

920

925

930

Table 3: Summary of the parameters used for the extrapolation of leaf scale experiments to the global scale and the results obtained. Bottom part: overview of available  $\Delta^{17}\text{O}$  measurements.

<b>Parameters and values used for global estimation</b>		
Parameter	Value	ref
GPP	120 PgCyr <sup>-1</sup>	(Beer et al., 2010)
$f_{C4}$	23 %	(Still et al., 2003)
$f_{C3}$	77 %	(Still et al., 2003)
$c_m/c_a$ (C <sub>3</sub> )	0.7	(Hoag et al., 2005)
$c_m/c_a$ (C <sub>4</sub> )	0.3	(Hoag et al., 2005)
$\Delta^{17}\text{O}$ leaf water (global mean, modelled)	-0.067±0.04 ‰	(Koren et al., 2019)
$\Delta^{17}\text{O}$ CO <sub>2</sub> (global mean, modelled)	-0.168±0.013 ‰	(Koren et al., 2019)
$\Delta_A\Delta^{17}\text{O}$ (global mean for C <sub>4</sub> )	-0.3±0.18 ‰	(Figure 5b, for $c_m/c_a$ ratio of 0.3)
$\Delta_A\Delta^{17}\text{O}$ (global mean for C <sub>3</sub> )	-0.65±0.18 ‰	(Figure 5b, for $c_m/c_a$ ratio of 0.7)
$\Delta_A\Delta^{17}\text{O}$ (global mean for whole vegetation)	-0.57±0.14 ‰	(Equation 13)
$\Delta_A\Delta^{17}\text{O}$ -isoflux (global mean for C <sub>4</sub> )	-7.3±4 ‰PgCyr <sup>-1</sup>	(Equation 14, only for C <sub>4</sub> )
$\Delta_A\Delta^{17}\text{O}$ -isoflux (global mean for C <sub>3</sub> )	-53±15 ‰PgCyr <sup>-1</sup>	(Equation 14, only for C <sub>3</sub> )
$\Delta_A\Delta^{17}\text{O}$ -isoflux (global mean for whole vegetation)	-60±15 ‰PgCyr <sup>-1</sup>	(equation 14)
$\Delta_A\Delta^{17}\text{O}$ -isoflux (global mean for whole vegetation)	-47 ‰PgCyr <sup>-1</sup>	(Hoag et al., 2005)
$\Delta_A\Delta^{17}\text{O}$ -isoflux (global mean for whole vegetation)	-42 to -92 ‰PgCyr <sup>-1</sup>	(Hofmann et al., 2017)
<b><math>\Delta^{17}\text{O}</math> value of tropospheric CO<sub>2</sub></b>		
$\Delta^{17}\text{O}(\text{CO}_2)$ for CO <sub>2</sub> samples collected in La Jolla-UCSD (California, USA) (1990 to 2000)	-0.173±0.046 ‰	(Thiemens et al., 2014)
$\Delta^{17}\text{O}(\text{CO}_2)$ for CO <sub>2</sub> samples collected in Israel	0.034±0.010 ‰	(Barkan and Luz, 2012)
$\Delta^{17}\text{O}(\text{CO}_2)$ for CO <sub>2</sub> samples collected in South china sea (2013-2014)	-0.159±0.084 ‰	(Liang et al., 2017b;Liang et al., 2017a)
$\Delta^{17}\text{O}(\text{CO}_2)$ for CO <sub>2</sub> samples collected in Taiwan (2012-2015)	-0.150±0.080 ‰	(Liang et al., 2017b;Liang et al., 2017a)
$\Delta^{17}\text{O}(\text{CO}_2)$ for CO <sub>2</sub> samples collected in California (USA) (2015)	-0.177±0.029 ‰	(Liang et al., 2017b;Liang et al., 2017a)
$\Delta^{17}\text{O}(\text{CO}_2)$ for CO <sub>2</sub> samples collected in Göttingen (Germany) (2010-2012)	-0.122±0.065 ‰	(Hofmann et al., 2017)

935

940

945

