

Plant genotype determines biomass response to flooding frequency in tidal wetlands

Svenja Reents¹, Peter Mueller², Hao Tang¹, Kai Jensen¹, Stefanie Nolte^{3,4}

¹Applied Plant Ecology, Institute of Plant Science and Microbiology, Universität Hamburg, Hamburg, 22609, Germany

²Smithsonian Environmental Research Center, Edgewater, Maryland, 21087, United States

5 ³School of Environmental Sciences, University of East Anglia, Norwich, NR4 7TJ, United Kingdom

⁴Centre for Environment, Fisheries and Aquaculture Science, Lowestoft, NR33 0HT, United Kingdom

Correspondence to: Svenja Reents (svenja.reents@uni-hamburg.de)

10 **Abstract.** The persistence of tidal wetland ecosystems like salt marshes is threatened by human interventions and climate change. Particularly the threat of accelerated sea level rise (SLR) has recently gained increasing attention by the scientific community. However, studies investigating the effect of SLR on plants and vertical marsh accretion are usually restricted to the species or community level and do not consider phenotypic plasticity or genetic diversity. To investigate the response of genotypes within the same salt-marsh species to SLR, we used two known genotypes of *Elymus athericus* (Link) Kerguelen
15 (low-marsh and high-marsh genotypes). In a factorial marsh organ experiment we exposed both genotypes to different flooding frequencies and quantified plant growth parameters. With increasing flooding frequency, the low-marsh genotype showed higher aboveground biomass production compared to the high-marsh genotype. Additionally, the low-marsh genotype generally formed longer rhizomes, shoots and leaves, regardless of flooding frequency. Belowground biomass of both
20 genotypes decreased with increasing flooding frequency. We conclude that the low-marsh genotype is better adapted to higher flooding frequencies through its ability to allocate resources from below- to aboveground biomass. Given the strong control of plant biomass production on salt-marsh accretion, we argue that these findings yield important implications for our understanding of ecosystem resilience to SLR as well as plant-species distribution in salt marshes.

1 Introduction

25 Salt marshes are wetland ecosystems predominantly found along coastlines where they form a transition zone between the marine and the terrestrial environment. Salt marshes provide important ecosystem services like protection of coastlines against storm surges by wave attenuation (Möller et al., 2014), supply of nursery grounds for commercially important fish (Bolle et al., 2009) and mitigation of climate change by long-term carbon sequestration (McLeod et al., 2011). However, human interventions such as land reclamation, eutrophication and climate change threaten the persistence of salt marshes, causing loss rates of 1-2 % of the global area per year (Duarte et al., 2008). Particularly the threat of accelerated sea level rise (SLR)
30 has recently gained increasing attention by the scientific community (Fitzgerald et al., 2008; Kirwan and Megonigal, 2013; Schuerch et al., 2018).

Generally, up to a certain locally varying threshold of SLR, salt marshes are able to keep up with rates of SLR through their ability to accrete vertically (Kirwan and Megonigal, 2013). During this process, salt-marsh plants act as ‘ecosystem engineers’ because their aboveground biomass reduces water flow velocity and hydrodynamic forces, which results in a decrease in the sediment-loading capacity of the water and an increase in sediment settlement (Morris et al., 2002; Yang, 1998). Furthermore, a high belowground biomass production and decreased decomposition rates in oxygen-deficient soils lead to an accumulation of organic matter and thereby expansion of soil volume. Whether above- or belowground processes predominantly contribute to vertical accretion, depends on the marsh type (i.e. minerogenic vs. organogenic) and often depends on local tidal amplitude (Allen, 2000; Kirwan and Megonigal, 2013; Nolte et al., 2013b). Yet, in both cases, biomass production of the marsh vegetation is an important driver of accretion. When accretion rates are too low and threshold values of SLR are exceeded, plant growth is negatively affected, leading to a negative feedback loop, which ensues marsh submergence and finally marsh loss (Chmura, 2013; Kirwan and Guntenspergen, 2012).

Recently, more studies have therefore been focusing on salt-marsh resilience to projected rates of SLR by studying the vegetation response. To examine how vegetation responds to changes in flooding regimes, so called ‘marsh organ’ experiments have been proven as convenient and insightful approaches. A marsh organ consists of several mesocosms arranged next to each other and along an elevational gradient. Kirwan and Guntenspergen (2012), for example, placed a marsh organ in a brackish marsh to examine the effect of varying sea levels on plant growth of two marsh species and what possible implications this may have on accretion processes. They observed that marsh elevation within the tidal frame determines whether root biomass increases or decreases with SLR. In contrast to belowground productivity, aboveground biomass response was species dependent. Numerous other studies confirmed that responses of marsh plants to sea level changes are highly species-specific (Eleuterius and Eleuterius, 1979; Kirwan and Guntenspergen, 2015; Langley et al., 2013; Morris et al., 2013). Therefore, the effect of SLR on plant biomass production, and thus the ability of the ecosystem to accrete vertically, can only be understood if SLR-induced shifts in plant species composition also are taken into account.

However, such studies on the effect of SLR on plants are usually restricted to the species or community level (Kirwan and Guntenspergen, 2015; Langley et al., 2013; Morris et al., 2013). They usually do not consider the high degree of phenotypic plasticity and genetic diversity within many species, as for instance evident between different locally adapted populations (Valladares et al., 2014). Yet, understanding a species’ adaptive genetic plasticity can be crucial to evaluate its response to environmental change (Razgour et al., 2019). Indeed, in some cases environmental change can in fact induce larger variability in plant biomass production within than between species (Beierkuhnlein et al., 2011). We therefore argue, that intraspecific differences in the biomass response to SLR of salt-marsh plants are likely relevant and require more attention by the scientific community to understand ecosystem resilience.

To investigate the response of genotypes within the same salt-marsh species to SLR, we used two known genotypes of the grass *Elymus athericus* (Link) Kerguélen. This species is widely distributed in NW European salt marshes and usually grows in high-elevated and therefore less-frequently flooded zones of the salt marsh (= high marsh) (Nolte et al., 2019). In the high marsh, the tall grass forms dense, monospecific stands with a low local plant species diversity (Bakker et al., 2003). Recently,

however, populations of *E. athericus* have been observed spreading into lower and thus more frequently flooded zones of the marsh (= low marsh) (Oloff et al., 1997; Veeneklaas et al., 2013). Low- and high-marsh genotypes are visually distinguishable, as the low-marsh genotype develops a specific phenotype different from the high-marsh genotype (Bockelmann et al., 2003). If genotypes respond differently to increased flooding frequencies, for instance in biomass productivity, it might affect salt-marsh responses to SLR.

The aim of this study was to investigate potential adaptations of the low-marsh genotype to increased flooding frequencies, to improve current evaluations of salt-marsh resilience to SLR. To test the hypothesis that the low-marsh genotype performs better at increased flooding frequencies than the high-marsh genotype, which would be reflected in above- and belowground growth parameters like biomass production and shoot, leaf and rhizome lengths, we conducted a factorial marsh organ experiment. We exposed both genotypes of *E. athericus* to three different flooding frequencies and quantified plant growth parameters to compare their performance and assess morphological adaptations of the low-marsh genotype.

2 Material and methods

2.1 Plant collection and culture

Plants were collected in April 2015 from a salt marsh on the Dutch Island Schiermonnikoog (53°30'N, 6°16'E) from stands that have previously been identified to be dominated by genetically distinct populations of *E. athericus*, i.e. high-marsh (HM) and low-marsh (LM) genotypes (Bockelmann et al., 2003). On Schiermonnikoog, *Elymus athericus* can be found in higher and lower elevated sites, which are inundated 20 – 125 and 90 – 270 times per year, respectively (Bockelmann et al., 2003). Soil salinities range from 22 to 29 ppt (Bolhuis et al., 2013). Plants and soil were extracted in the form of intact sods to keep them alive during transport. In Hamburg, soil was removed and roots were rinsed before both genotypes were planted separately in trays with standardised potting soil. Until the start of the experiment (i.e. for 24 months), plants were kept under identical environmental conditions in a common garden at the Institute of Plant Science and Microbiology. Ramets of these plants were used for this study. In July 2017, single plants of similar biomass were selected based on visual assessment (no obvious outliers), transplanted to separate pots and randomly assigned to the flooding treatments (described in 2.2). Initial shoot length and shoot number was tested for differences between genotypes and flooding frequencies to ensure that results were not biased by unequal plant size at the beginning of the experiment. There were no significant differences regarding shoot length (genotype: $F = 0.787$, $p = 0.380$; flooding frequency: $F = 0.127$, $p = 0.881$; genotype*flooding frequency: $F = 0.231$, $p = 0.795$) and number of shoots (genotype: Wald = 2.203, $p = 0.137$; flooding frequency: Wald = 0.357, $p = 0.837$; genotype*flooding frequency: Wald = 0.005, $p = 0.997$). The pots were 15 cm in diameter, 17 cm in height and had holes in the bottom to facilitate drainage. They were filled with salt marsh soil taken from the salt marsh at Sönke-Nissen-Koog, Germany (54°36'N, 8°49'E) which was sieved (with a 1 cm mesh) and homogenised beforehand (see Nolte et al. (2013a) for soil properties). Eight replicates (i.e. single plants in separate pots) per genotype were assigned to one of three flooding treatments, so that a total number of 48 plants were used in this study.

2.2 Experimental set-up

Plants were placed onto three steps (step height: 20 cm) within a tidal-tank facility (Hanke et al., 2015), to create three different
100 flooding frequencies. The tidal tank is located outdoors at the Institute of Plant Science and Microbiology and has a total
volume of 6.75 m³ (dimensions: 3 x 1.5 x 1.5 m). A pump was used to fill and empty the tidal tank at regular intervals to mimic
tides by alternating between three different maximum water levels. Pots were fully drained between flooding events. Flooding
with the respective maximum water level reached 3 cm above soil surface, lasted two hours and took place twice a day. The
lowest step was flooded every day, which represented the highest frequency. The flooding of the middle step (moderate
105 flooding frequency) happened weekly, while plants on the highest step were flooded only every two weeks (lowest flooding
frequency). Highest and lowest flooding frequency reflect the natural flooding gradient between pioneer zone and high marsh
in many NW European salt marshes, including the site where our plants were collected (Bockelmann et al., 2002). A CTD
diver combined with a baro diver (Van Essen Instruments, Delft, The Netherlands) was used to monitor flooding cycles.
Artificial sea salt (AB Aqua Medic GmbH, Germany) was suspended in tap water to create a salinity of about 20-22 ppt. To
110 minimise the impact of other effects than flooding frequency and genotype, the pots were circulated on each step, other
seedlings and algae were removed once a week. Concurrently, water level and salinity were checked as well. The experiment
ran for approx. 12 weeks from mid-July to early October 2017.

2.3 Measurements

2.3.1 Biomass

115 At the end of the experiment (2nd of October 2017), the plants were harvested, separated into above- and belowground
biomass, dried for two days at 70°C and weighed. Belowground biomass was divided into rhizomes and roots and weighed.
The length of rhizomes was also recorded. Additionally, above- and belowground biomass were used to calculate the
belowground:aboveground ratio.

2.3.2 Plant growth

120 At the beginning and the end of the experiment, plant shoot and leaf length as well as number of shoots and leaves were
measured. The difference between both measurements was calculated and designated as Δ . Only living plant material was
taken into account and length measurements (leaf and shoot length) were carried out on the longest leaves or shoots.

2.4 Statistical analysis

Data were tested for normality by applying the Shapiro-Wilk-Test. Except count data (e.g. number of leaves), all parameters
125 were normally distributed and therefore further analysed applying factorial ANOVAs. Due to the well-balanced study design,
potential moderate deviations from homogeneity of variance between groups were considered unimportant for ANOVA testing
(Box, 1954; McGuinness, 2002). Each analysis included genotype and flooding frequency as well as their interaction as

explanatory variables. To detect significant differences between treatments (flooding frequency and genotype), post-hoc tests (Tukey's HSD) were conducted. To analyse count data, i.e. number of shoots and number of leaves, generalized linear models (GLM) were applied assuming a Poisson distribution and including the explanatory variables genotype and flooding frequency, as well as two-way interaction effects. Each GLM was checked for overdispersion (Pearson Chi2 dispersion parameter) and was refitted afterwards if necessary, using the standard procedure of the applied program. All statistical analyses were performed using STATISTICA 13 (StatSoft Inc., Tulsa, OK, USA).

3 Results

135 3.1 Biomass

Total biomass, defined as the sum of dry above- and belowground plant biomass, differed significantly between genotypes and flooding frequencies (Table 1). In addition, the interaction of both factors showed a significant effect on total biomass. Total biomass production of the high-marsh genotype decreased steadily with increasing flooding frequency, whereas the total biomass of the low-marsh genotype decreased less distinctly (Fig. 1a). The difference between low-marsh and high-marsh genotypes was most pronounced at the highest flooding frequency. In fact, the low-marsh genotype produced almost twice as much total biomass at highest flooding frequency as the high-marsh genotype (LM: 4.61 ± 0.70 g and HM: 2.66 ± 0.52 g, mean \pm standard deviation).

Genotype and flooding frequency as well as their interaction had a significant effect on the aboveground biomass production. Aboveground biomass of the high-marsh genotype decreased with increasing flooding frequency from 3.31 ± 0.57 g to 2.03 ± 0.38 g (Fig. 1b). However, aboveground biomass production of the low-marsh genotype remained constant at about 3.46 ± 0.45 g on all flooding frequencies.

In contrast to the genotype-specific aboveground biomass response to flooding, belowground biomass of both genotypes decreased with increasing flooding frequency (Fig. 1c). Results indicate a more pronounced effect of flooding frequency on belowground biomass production compared to the factor genotype (Table 1). Under all flooding frequencies, the low-marsh genotype produced slightly more belowground biomass than the high-marsh genotype (LM: 1.63 ± 0.78 g, HM: 1.30 ± 0.76 g).

Root biomass production (belowground biomass without rhizomes) was significantly affected by flooding frequency ($F = 10.69$, $p < 0.001$), but did not differ between genotypes. Root biomass decreased with increasing flooding frequency for both genotypes (Fig. 1d). In contrast, biomass of rhizomes was significantly affected by both genotype and flooding frequency (Table 1). Mean rhizome biomass of the low-marsh genotype was higher than of the high-marsh genotype (LM: 0.55 ± 0.50 g, HM: 0.26 ± 0.28 g) with the most pronounced differences on lowest and highest flooding frequency (Fig. 1e). In some cases, the low-marsh genotype formed very long rhizomes (up to 166 cm length, coiled around the soil). On average, rhizomes of the low-marsh genotype were nearly twice as long as those of the high-marsh genotype (LM: 51.43 ± 41.11 cm, HM: 26.63 ± 27.23 cm, Table 2). Genotypes significantly differed in the length of rhizomes ($F = 6.102$, $p < 0.05$, Table 1).

160 Belowground:aboveground-ratio was significantly affected by flooding frequency (Table 1) and decreased with increasing flooding frequency (Fig. 1f).

3.2 Δ Leaf and shoot length

The increase in shoot and leaf length significantly differed between genotypes (Table 1). Regarding leaf length, the high-marsh genotype showed approximately the same increase on all flooding frequencies (1.9 ± 4.45 cm). The low-marsh genotype had similar increases of leaf lengths at the lowest flooding frequency but showed pronounced increases of leaf length with increasing flooding frequency (7.03 ± 2.17 cm) (Fig. 2a). Increase in shoot length of the low-marsh genotype was twice as high as that of the high-marsh genotype (LM: 10.78 ± 6.18 cm, HM: 5.57 ± 6.58 cm, Fig. 2b).

3.3 Δ Number of leaves and shoots

Neither genotype nor flooding frequency had a significant effect on the increase in number of shoots (Table 1). However, for the increase in number of leaves, a significant effect of flooding frequency was detected (Wald = 19.69, $p < 0.001$). With highest flooding frequency, both genotypes produced the lowest number of new leaves (LM: 10.4 ± 4.0 , HM: 8.9 ± 2.1 ; Table 2).

4 Discussion

Assessments of plant responses to changed hydrological conditions (e.g. SLR) have thus far focused mainly on comparisons on species level. However, variability in plant responses within species can be considerably higher than between species (Beierkuhnlein et al., 2011). In this study, we therefore investigated differences in plant response between genotypes of the same species (*Elymus athericus*) to better understand the importance of intraspecific variability for evaluations of future ecosystem functionality and resilience. We found, in line with our hypothesis on biomass production, that the low-marsh genotype performs better than the high-marsh genotype under increased flooding frequency. Additionally, the low-marsh genotype generally formed longer rhizomes, shoots and leaves, regardless of flooding frequency (Fig. 1 & 2). We argue that these findings yield important implications for our understanding of ecosystem resilience to SLR as well as plant-species distribution in salt marshes.

We found a higher total biomass of the low-marsh genotype, which was particularly pronounced under high flooding frequency (Fig. 1a). This result indicates a better adaptation of the low-marsh genotype to lower elevated, more frequently flooded conditions. When separating above- and belowground biomass, the high-marsh genotype showed a decrease of both biomass parameters with increasing flooding frequency. Aboveground biomass is important for maintaining photosynthesis (Johnson, 2016), so that its reduction can be interpreted as reduction in performance as well. In contrast to the high-marsh genotype showing a marked reduction in aboveground biomass, the low-marsh genotype maintained aboveground biomass across all flooding treatments.

190 We also found a decrease of belowground biomass with increasing flooding frequency in both genotypes. This can be interpreted as an adaptive trait, because a reduction of belowground biomass reduces the number of respiring roots and thereby improves the diffusion of oxygen to the roots (Naidoo and Naidoo, 1992; Voesenek et al., 1988). In line with these results, an increased aboveground biomass production while belowground biomass decreased was found for other flooding adapted plant species such as *Taxodium distichum*, *Danthonia montevidensis* and *Paspalum dilatatum* (Megonigal and Day, 1992; Rubio et al., 1995).

The difference in aboveground biomass response between the two genotypes seems to be mainly explained by genotype-specific increases in leaf and shoot length, whereas the number of both remained similar (Fig. 2, Table 1). Likewise, Voesenek et al. (1988) found a marked increase in leaf length in the flooding adapted *Rumex palustris* under waterlogging, but no increase in number of leaves. The distinct increase in leaf and shoot length in addition to the simultaneous reduction of belowground biomass of the low-marsh genotype of *Elymus athericus* found in our study, indicate resource allocation as response to flooding. For other species it was found that reallocated resources fuel elongation of shoots and leaves to maintain gas exchange and avoid light dissipation through water (Blanch et al., 1999; Grace, 1989). Our results suggest that this response may be also present in *E. athericus*, which could improve its chances of survival under higher flooding frequencies e.g. due to accelerated SLR.

205 Vertical accretion in the minerogenic salt marshes of the Wadden Sea is primarily driven by sedimentation (Allen, 2000; Nolte et al., 2013b), which is strongly controlled by the sediment-trapping capacity of the aboveground biomass (Morris et al., 2002; Yang, 1998). The strong aboveground biomass response to increased flooding frequencies of the low-marsh genotype found in our study may therefore have a positive effect on vertical accretion rates and thereby marsh resilience to rising sea levels. *E. athericus* is not the only salt-marsh species characterised by a high degree of genetic diversity. In previous studies, genotypes of several salt-marsh grasses have been described and tested for intraspecific differences in plant response to changing environmental conditions, including *Puccinellia maritima*, *Phragmites australis* and *Spartina alterniflora* (Gray, 1985; Mozdzer and Megonigal, 2012; Proffitt et al., 2003; Seliskar et al., 2002). They showed high genotypic variations affecting colonisation success, species composition and even ecosystem function.

Compared to the root biomass of both genotypes, which responded similarly and decreased with increasing flooding frequency, rhizome length differed significantly between the genotypes (Table 1). The formation of longer rhizomes by the low-marsh genotype, especially under high flooding frequency, could serve as an escape strategy to expand into more favourable habitats (Hartnett and Bazzaz, 1983; Lovett-Doust, 1981). In previous studies, it was reported that *E. athericus* usually expands via a ‘phalanx’ growth strategy, which means that parental plants invest in many but rather short rhizomes to utilise resources in a favourable habitat (Bockelmann and Neuhaus, 1999). Field observations of the same authors and results of our study, however, indicate that *E. athericus* is able to alter its strategy to the ‘guerrilla’ form, by producing longer rhizomes. The ‘guerrilla’ strategy is usually found in plants characteristic for early successional stages as it enables plants to spread quickly and exploit new favourable areas (Lovett-Doust, 1981). However, overall rhizome length of our study should be interpreted with caution because of potential edge effects caused by the experimental mesocosms.

225 The change of expansion strategy together with a better adaptation to higher flooding frequencies may lead to a displacement of the high-marsh genotype under accelerated SLR. However, until now, the Wadden Sea salt marshes are able to cope with current rates of sea level rise due to high accretion rates (Esselink et al., 2017; Nolte et al., 2013a). If rates of SLR remain stable, the low-marsh genotype of the tall grass *E. athericus* has the potential to expand further into the low marsh and outcompete other species via light competition, potentially reducing local species diversity.

5 Methodological considerations

230 We suggest that the experimental setup including a tidal tank and steps proved suitable to investigate the effects of different flooding frequencies on salt-marsh vegetation. Nevertheless, we recommend repeating this experiment in situ, for example as a transplant experiment, to estimate actual effect size under more natural conditions, as drainage and plant-soil interactions might have been different in the tidal tank and could have affected biomass production.

6 Conclusion

235 The present work revealed marked differences in the plant biomass response to changes in flooding frequency between two genotypes of the dominant European salt-marsh grass *Elymus athericus*. Furthermore, we observed large differences in rhizome production between genotypes, which is interpreted as a change in growth strategy. The alteration of its growth strategy and the higher aboveground biomass productivity of the low-marsh genotype implies a larger potential of the low-marsh genotype to invade and establish at lower elevations of the tidal frame. Considering the generally low plant species
240 diversity of salt marshes (e.g. Silliman, 2014; Wanner et al., 2014) and the strong feedbacks between plant growth and accelerated SLR (Kirwan and Megonigal, 2013), it is possible that intraspecific variation and adaptive capacity in salt-marsh plants acts as an important but overlooked mediator of ecosystem resilience.

7 Data availability

All data presented in this paper is available from the corresponding author upon request.

245 8 Author contribution

All authors contributed to the design of the experiments. SR and HT conducted the experiment and performed the measurements. SR analysed the data and wrote the manuscript. All authors contributed to the discussion of results and the final manuscript.

9 Competing interests

250 The authors declare that they have no conflict of interest.

10 Acknowledgements

We would like to thank Chris Smit and his colleagues from the University of Groningen for the provision of the plants. Furthermore, we would like to acknowledge Max Beiße, Marion Klötzl, and Maren Winnacker for their assistance during the preparation of the experiment. Christoph Reisdorff provided advice on experimental setup and measurements.

255 References

- Allen, J. R. L.: Morphodynamics of Holocene salt marshes: a review sketch from the Atlantic and Southern North Sea coasts of Europe, *Quat. Sci. Rev.*, 19, 1155–1231, 2000.
- Bakker, J. P., Bos, D. and De Vries, Y.: To graze or not to graze, that's the question, in *Proceedings of the 10th International Scientific Wadden Sea Symposium*, vol. 88., 2003.
- 260 Beierkuhnlein, C., Thiel, D., Jentsch, A., Willner, E. and Kreyling, J.: Ecotypes of European grass species respond differently to warming and extreme drought, *J. Ecol.*, 99, 703–713, doi:10.1111/j.1365-2745.2011.01809.x, 2011.
- Blanch, S. J., G. Ganf, G. and Walker, K. F.: Growth and resource allocation in response to flooding in the emergent sedge *Bolboschoenus medianus*, *Aquat. Bot.*, 63(2), 145–160, doi:10.1016/S0304-3770(98)00109-0, 1999.
- Bockelmann, A. C. and Neuhaus, R.: Competitive exclusion of *Elymus athericus* from a high-stress habitat in a European salt
265 marsh, *J. Ecol.*, 87(3), 503–513, 1999.
- Bockelmann, A. C., Bakker, J. P., Neuhaus, R. and Lage, J.: The relation between vegetation zonation, elevation and inundation frequency in a Wadden Sea salt marsh, *Aquat. Bot.*, 73(3), 211–221, doi:10.1016/S0304-3770(02)00022-0, 2002.
- Bockelmann, A. C., Reusch, B. H., Bijlsma, R. and Bakker, J. P.: Habitat differentiation vs. isolation-by-distance: The genetic population structure of *Elymus athericus* in European salt marshes, *Mol. Ecol.*, 12(2), 505–515, doi:10.1046/j.1365-
270 294X.2003.01706.x, 2003.
- Bolhuis, H., Fillinger, L. and Stal, L. J.: Coastal microbial mat diversity along a natural salinity gradient, *PLoS One*, 8(5), doi:10.1371/journal.pone.0063166, 2013.
- Bolle, L. J., Neudecker, T., Vorberg, R., Damm, U., Diederichs, B., Jager, Z., Scholle, J., Daenhardt, A., Lüerßen, G. and Marencic, H.: Trends in Wadden Sea fish fauna., 2009.
- 275 Chmura, G. L.: What do we need to assess the sustainability of the tidal salt marsh carbon sink?, *Ocean Coast. Manag.*, 83, 25–31, doi:10.1016/j.ocecoaman.2011.09.006, 2013.
- Duarte, C. M., Dennison, W. C., Orth, R. J. W. and Carruthers, T. J. B.: The charisma of coastal ecosystems: Addressing the imbalance, *Estuaries and Coasts*, 31(2), 233–238, doi:10.1007/s12237-008-9038-7, 2008.
- Eleuterius, L. N. and Eleuterius, C. K.: Tide levels and salt marsh zonation, *Bull. Mar. Sci.*, 29(3), 394–400, 1979.
- 280 Esselink, P., Duin, W. E. Van, Bunje, J., Cremer, J., Folmer, E. O., Frikke, J., Glahn, M., Groot, A. V. De, Hecker, N., Hellwig, U., Jensen, K., Körber, P., Petersen, J. and Stock, M.: Salt marshes, *Wadden Sea Qual. Status Rep.*, Eds.: Kloepper S. et al., Common Wadden Sea Secret, 2017.

- Fitzgerald, D. M., Fenster, M. S., Argow, B. A. and Buynevich, I. V.: Coastal impacts due to sea-level rise, *Annu. Rev. Earth Planet. Sci.*, 36(601–647), 2008.
- 285 Grace, J. B.: Effects of water depth on *Typha latifolia* and *Typha domingensis*, *Am. J. Bot.*, 76(5), 762–768, 1989.
- Gray, A. J.: Adaptation in perennial coastal plants - with particular reference to heritable variation in *Puccinellia maritima* and *Ammophila arenaria**, *Vegetatio*, 61, 179–188, 1985.
- Hanke, J. M., Ludewig, K. and Jensen, K.: Effects of water level and competition on the endangered river corridor plant *Cnidium dubium* in the context of climate change, *Wetl. Ecol. Manag.*, 23(2), 215–226, doi:10.1007/s11273-014-9371-5, 290 2015.
- Hartnett, D. C. and Bazzaz, F. .: Physiological integration among intracolonial ramets in *Solidago Canadensis*, *Ecology*, 64(4), 779–788, 1983.
- Johnson, M. P.: Photosynthesis, *Essays Biochem.*, 60(3), 255–273, doi:10.1042/EBC20160016, 2016.
- Kirwan, M. L. and Guntenspergen, G. R.: Feedbacks between inundation, root production, and shoot growth in a rapidly 295 submerging brackish marsh, *J. Ecol.*, 100(3), 764–770, doi:10.1111/j.1365-2745.2012.01957.x, 2012.
- Kirwan, M. L. and Guntenspergen, G. R.: Response of plant productivity to experimental flooding in a stable and a submerging marsh, , 18(5), 903–913, 2015.
- Kirwan, M. L. and Megonigal, J. P.: Tidal wetland stability in the face of human impacts and sea-level rise, *Nature*, 504, 53–60, doi:10.1038/nature12856, 2013.
- 300 Langley, J. A., Mozdzer, T. J., Shepard, K. A., Hagerty, S. B. and Megonigal, J. P.: Tidal marsh plant responses to elevated CO₂, nitrogen fertilization, and sea level rise, *Glob. Chang. Biol.*, 19, 1495–1503, doi:10.1111/gcb.12147, 2013.
- Lovett-Doust, L.: Population dynamics and local specialization in a clonal perennial (*Ranunculus Repens*): II . The dynamics of leaves, and a reciprocal transplant-replant experiment, *J. Ecol.*, 69(3), 757–768, doi:10.2307/2259634, 1981.
- McLeod, E., Chmura, G. L., Bouillon, S., Salm, R., Björk, M., Duarte, C. M., Lovelock, C. E., Schlesinger, W. H. and Silliman, 305 B. R.: A blueprint for blue carbon: Toward an improved understanding of the role of vegetated coastal habitats in sequestering CO₂, *Front. Ecol. Environ.*, 9(10), 552–560, 2011.
- Megonigal, J. P. and Day, F. P.: Effects of flooding on root and shoot production of Bald Cypress in large experimental enclosures, *Ecology*, 73(4), 1182–1193, 1992.
- Möller, I., Kudella, M., Rupprecht, F., Spencer, T., Paul, M., Van Wesenbeeck, B. K., Wolters, G., Jensen, K., Bouma, T. J., 310 Miranda-Lange, M. and Schimmels, S.: Wave attenuation over coastal salt marshes under storm surge conditions, *Nat. Geosci.*, 7(10), 727–731, doi:10.1038/NGEO2251, 2014.
- Morris, J. T., Sundareshwar, P. V., Nietch, C. T., Kjerfve, B. and Cahoon, D. R.: Responses of coastal wetlands to rising sea level, *Ecology*, 83(10), 2869–2877, 2002.
- Morris, J. T., Sundberg, K. and Hopkinson, C. S.: Salt marsh primary production and its responses to relative sea level and 315 nutrients, *Oceanography*, 26(3), 78–84, 2013.
- Mozdzer, T. J. and Megonigal, J. P.: Jack-and-Master trait responses to elevated CO₂ and N: A comparison of native and

- introduced *Phragmites australis*, PLoS One, 7(10), doi:10.1371/journal.pone.0042794, 2012.
- Naidoo, G. and Naidoo, S.: Waterlogging responses of *Sporobolus-Virginicus* (L) Kunth, Oecologia, 90(3), 445–450, doi:10.1007/BF00317704, 1992.
- 320 Nolte, S., Müller, F., Schuerch, M., Wanner, A., Esselink, P., Bakker, J. P. and Jensen, K.: Does livestock grazing affect sediment deposition and accretion rates in salt marshes?, Estuar. Coast. Shelf Sci., 135, 296–305, doi:10.1016/j.ecss.2013.10.026, 2013a.
- Nolte, S., Koppelaar, E. C., Esselink, P., Dijkema, K. S., Schuerch, M., De Groot, A. V., Bakker, J. P. and Temmerman, S.: Measuring sedimentation in tidal marshes: A review on methods and their applicability in biogeomorphological studies, J. Coast. Conserv., 17(3), 301–325, doi:10.1007/s11852-013-0238-3, 2013b.
- 325 Nolte, S., Wanner, A., Stock, M. and Jensen, K.: *Elymus athericus* encroachment in Wadden Sea salt marshes is driven by surface elevation change, Appl. Veg. Sci., 22, 454–464, doi:10.1111/avsc.12443, 2019.
- Oloff, H., De Leeuw, J., Bakker, J. P., Platerink, R. J. and van Wijnen, H. J.: Vegetation succession and herbivory in a salt marsh: Changes induced by sea level rise and silt deposition along an elevational gradient, J. Ecol., 85(6), 799–814, 1997.
- 330 Proffitt, C. E., Travis, S. E. and Edwards, K. R.: Genotype and elevation influence *Spartina alterniflora* colonization and growth in a created salt marsh, Ecol. Appl., 13(1), 180–192, 2003.
- Razgour, O., Forester, B., Taggart, J. B., Bekaert, M., Juste, J., Ibáñez, C., Puechmaille, S. J., Novella-Fernandez, R., Alberdi, A. and Manel, S.: Considering adaptive genetic variation in climate change vulnerability assessment reduces species range loss projections, PNAS, 116(21), 10418–10423, doi:10.1073/pnas.1820663116, 2019.
- 335 Rubio, G., Casasola, G. and Lavado, R. S.: Adaptations and biomass production of two grasses in response to water logging and soil nutrient enrichment, Oecologia, 102, 102–105, 1995.
- Schuerch, M., Spencer, T., Temmerman, S., Kirwan, M. L., Wolff, C., Lincke, D., Mcowen, C. J., Pickering, M. D., Reef, R., Vafeidis, A. T., Hinkel, J., Nicholls, R. J. and Brown, S.: Future response of global coastal wetlands to sea-level rise, Nature, 561, 231–234, doi:10.1038/s41586-018-0476-5, 2018.
- 340 Seliskar, D. M., Gallagher, J. L., Burdick, D. M. and Mutz, L. A.: The regulation of ecosystem functions by ecotypic variation in the dominant plant: a *Spartina alterniflora* salt-marsh case study, J. Ecol., 90, 1–11, 2002.
- Silliman, B. R.: Quick guide: Salt marshes, Curr. Biol., 24(9), 348–350, doi:10.1016/j.cub.2014.03.001, 2014.
- Valladares, F., Matesanz, S., Guilhaumon, F., Araújo, M. B., Balaguer, L., Benito-Garzón, M., Cornwell, W., Gianoli, E., van Kleunen, M., Naya, D. E., Nicotra, A. B., Poorter, H. and Zavala, M. A.: The effects of phenotypic plasticity and local adaptation on forecasts of species range shifts under climate change, Ecol. Lett., 17, 1351–1364, doi:10.1111/ele.12348, 2014.
- 345 Veeneklaas, R. M., Dijkema, K. S., Hecker, N. and Bakker, J. P.: Spatio-temporal dynamics of the invasive plant species *Elytrigia atherica* on natural salt marshes, Appl. Veg. Sci., 16(2), 205–216, doi:10.1111/j.1654-109X.2012.01228.x, 2013.
- Voeselek, L. A. C. J., Blom, C. W. P. M. and Pouwels, R. H.: Root and shoot development of *Rumex* species under waterlogged conditions, Can. J. Bot., 67, 1865–1869, doi:10.1139/b89-236, 1988.
- 350

Wanner, A., Suchrow, S., Kiehl, K., Meyer, W., Pohlmann, N., Stock, M. and Jensen, K.: Scale matters: Impact of management regime on plant species richness and vegetation type diversity in Wadden Sea salt marshes, *Agric. Ecosyst. Environ.*, 182, 69–79, doi:10.1016/j.agee.2013.08.014, 2014.

355 Yang, S. L.: The role of *Scirpus* marsh in attenuation of hydrodynamics and retention of fine sediment in the Yangtze estuary, *Estuar. Coast. Shelf Sci.*, 47, 227–233, 1998.





Figure 1: Total biomass (a), aboveground biomass (b), belowground biomass (c), root biomass (d), rhizome biomass (e) and belowground-aboveground ratio (f) of both genotypes on three different flooding frequencies (mean + standard errors). Stars show significant differences between low-marsh (LM) and high-marsh (HM) genotypes within the same flooding treatment based on

365 Tukey's HSD post-hoc test (* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$). Capital letters indicate significant differences between flooding frequencies. For statistics see Table 1.



370 **Figure 2: Delta leaf length (a) and delta shoot length (b) of both genotypes on three different flooding frequencies (mean + standard errors). Capital letters indicate significant differences between flooding frequencies. For statistics see Table 1.**

375

380

385

Table 1: Summary statistics of main response variables in the experiment testing for effects of flooding and genotype of *Elymus athericus* on its biomass and growth. Count variables (number of shoots, number of leaves) are analysed through GLM, all other variables by a two-way factorial ANOVA. Significant p-values are shown in bold letters. Means and standard errors are shown in Figures 1 and 2.

Response variable	statistical test	genotype (df = 1)		flooding frequency (df = 2)		genotype * flooding frequency (df = 2)	
		test statistic	p-value	test statistic	p-value	test statistic	p-value
<i>Total biomass [g]</i>	factorial ANOVA	F = 13.18	≤ 0.001	F = 18.24	≤ 0.001	F = 3.36	≤ 0.05
<i>Aboveground biomass [g]</i>	factorial ANOVA	F = 13.84	≤ 0.001	F = 6.38	≤ 0.01	F = 4.77	≤ 0.05
<i>Belowground biomass [g]</i>	factorial ANOVA	F = 4.31	≤ 0.05	F = 21.93	≤ 0.001	F = 0.53	0.59
<i>Δ Shoot length [cm]</i>	factorial ANOVA	F = 7.77	≤ 0.01	F = 0.32	0.73	F = 1.03	0.37
<i>Δ Leaf length [cm]</i>	factorial ANOVA	F = 5.09	≤ 0.05	F = 1.08	0.35	F = 1.42	0.25
<i>Δ Number of shoots</i>	GLM	Wald = 0.00	0.95	Wald = 5.87	0.05	Wald = 0.78	0.68
<i>Δ Number of leaves</i>	GLM	Wald = 0.44	0.51	Wald = 19.69	≤ 0.001	Wald = 1.60	0.45
<i>Root biomass [g]</i>	factorial ANOVA	F = 0.03	0.86	F = 10.69	≤ 0.001	F = 0.07	0.93
<i>Rhizome biomass [g]</i>	factorial ANOVA	F = 6.49	≤ 0.05	F = 3.84	≤ 0.05	F = 0.51	0.60
<i>Rhizome length [cm]</i>	factorial ANOVA	F = 6.10	≤ 0.05	F = 1.67	0.20	F = 0.45	0.64
<i>Belowground:Aboveground</i>	factorial ANOVA	F = 0.34	0.56	F = 14.56	≤ 0.001	F = 0.13	0.88

Table 2: Mean and standard error (SE) of rhizome length [cm], delta number of leaves and delta number of shoots for different combinations of factors genotype and flooding frequency. LM = low-marsh genotype, HM = high-marsh genotype.

			Rhizome length [cm]		Δ Number of leaves		Δ Number of shoots	
genotype	flooding frequency	N	mean	SE	mean	SE	mean	SE
LM	low	8	63.38	15.13	15.88	2.43	7.00	0.82
HM	low	8	37.13	9.85	18.00	3.19	8.13	1.52
LM	moderate	8	45.28	10.35	21.75	2.45	9.50	0.78
HM	moderate	8	32.81	11.76	16.88	3.42	8.50	1.65
LM	high	8	45.63	18.10	10.43	1.53	6.25	1.18
HM	high	8	9.94	2.91	8.86	0.80	5.88	1.08