

Experimental production of charcoal morphologies to discriminate fuel source and fire type: an example from Siberian taiga

Angelica Feurdean¹

5 ¹ Department of Physical Geography, Goethe University, Altenhöferallee 1, 60438 Frankfurt am Main, Germany

Correspondence to: angelica.feurdean@gmail.com; Feurdean@em.uni-frankfurt.de

10

15

20

25

30

Abstract

The analysis of charcoal fragments in peat and lake sediments is the most widely used approach used to reconstruct past biomass burning. With a few exceptions, this method typically relies on the quantification of the total charcoal content of the sediment. To enhance charcoal analyses for the reconstruction of past fire regimes and make the method more relevant to studies of both plant evolution and fire management, the extraction of more information from charcoal particles is critical. Here, I used a muffle oven to burn seven fuel types comprising 17 species from boreal Siberia (near Teguldet village), which are also commonly found in the Northern Hemisphere, and built on published schemes to develop morphometric and finer diagnostic classifications of the experimentally charred particles. I then combined these results with those from fossil charcoal from a peat core taken from the same location (Ulukh-Chayahk) to demonstrate the relevance of these experiments to the fossil charcoal records. Results show that graminoids, Sphagnum, and wood (trunk) lose the most mass at low burn temperatures (<300°C), whereas heathland shrub leaves, brown moss, and ferns at high burn temperatures. This suggests that species with low mass retention in high-temperature fires are likely to be underrepresented in the fossil charcoal record. The charcoal particle aspect ratio appeared to be the strongest indicator of the fuel type burnt. Graminoid charcoal particles are most elongate (6.7-11.5), with a threshold above 6 that may be indicative of wetland graminoids. Leaves are the shortest and bulkiest (2.1-3.5) while twigs and wood are intermediate (2.0-5.2). Further, the use of fine diagnostic features was more successful in separating wood, graminoids, and leaves, but it was difficult to further differentiate these fuel types due to overlapping morphometrics and morphologies. High aspect ratio particles, dominated by graminoid and Sphagnum morphologies, may be robust indicators of low temperature, surface fires, whereas abundant wood and leaf morphologies, and low aspect ratio particles are indicative of higher-temperature fires. However, the overlapping morphologies of leaves and wood from trees and shrubs make it hard to distinguish between high-intensity surface fires, combusting living shrubs and dead wood and leaves, and high-intensity crown fires which have burnt living trees. Distinct particle shape may also influence charcoal transportation, with elongated particles (graminoids) potentially having a more heterogeneous distribution and be deposited farther away from the origin of fire than the rounder, polygonal leaf particles. Despite these limitations, the combined use of charred-particle aspect ratios and fuel morphotypes can aid the more robust interpretation of fuel source and fire type changes. Lastly, I highlight the further investigation needed to refine the histories of past wildfires.

1 Introduction

Wildfires are the most common disturbance type in boreal forests, triggering gap dynamics or stand-scale forest replacement depending on intensity (temperature of fire) and frequency (Goldammer and Furayev, 1996). Ongoing and anticipated increases in the intensity and frequency of wildfire in boreal forests is raising concerns about its impact forest composition as well as climate (Jones et al., 2020). Although the largest area of boreal forest globally is in Siberia (Goldammer and Furayev, 1996), its extent and access restrictions mean there are only a few datasets recording changes in wildfire activity, especially from a longer-term perspective (Marlon et al., 2016). Long-term records of wildfire activity are vital to understanding how fire regimes vary with changes in climate and human-vegetation interaction, as well as the impacts of fires on boreal forests.

Wildfires can reach temperatures up to 1800 °C. Charcoal, however, is an inorganic carbon compound resulting from the incomplete combustion of plant tissues, which typically occurs at temperatures between 280 and 500 °C (Rein, 2014). Charcoal particles vary in size and form, characteristics such as edge aspects, surface features, cleavage, lustre, or anatomical details (tracheids with border pits, leaf veins, cuticles, etc.) that can be used to determine their origin are often preserved (Ward and Hardy, 1991; MacDonald et al., 1991; Scott 2010; Enache and Cumming, 2006; Jensen et al., 2007; Courtney-Mustaphi and Pizaric, 2014a; Hubau et al., 2012, 2013; Prince et al., 2018). Although macroscopic charcoal analysis, typically counting charcoal pieces or charcoal area per unit sediment volume from a small sediment volume (1-2 cm³), is widely done, there has been less focus on the reconstruction of fuel sources. Yet this is a crucial factor in determining fire type, i.e., the burning of surface fuels in low or high intensity fires or distinguishing between surface and crown fires. At a minimum, this requires a better characterisation of charcoal morphology to indicate the nature of the plant material burnt (Courtney-Mustaphi and Pizaric, 2014 a,b; Feurdean et al., 2017; Hawthorne et al., 2018). The determination of fire type is not only helpful for palaeofire reconstructions, but could provide an accessible tool for ecosystem managers and modellers, and for assessing and mitigating the risks of fires that might impact settlements and infrastructures (Moritz et al., 2014).

Ongoing efforts have advanced the utility of charcoal morphological analyses for fuel type identification and fire regime reconstruction. Umbanhowar and McGrath (1998), Crawford and Belcher (2014), and Pereboom et al. (2020) conducted morphometric measurements of the length, aspect ratio (length/width), and surface area of charcoal particles by burning in the laboratory known plant materials originating from American prairie, tropical and arctic environments. They concluded that longer fragments correspond to graminoids, whereas shorter fragments originate from wood, shrubs, and leaves. Nichols (2000) and Crawford and Belcher (2014) additionally found that charcoal morphometry is generally preserved during transportation by water. Other studies have focused on the effects of burning conditions, i.e., open-flame ignition, muffle furnace experiments (Umbanhowar and McGrath, 1998; Orvis et al., 2005), and combustion calorimetry (Hudspith et al., 2018) on charcoal production. Belcher et al. (2005, 2015) investigated in the laboratory whether fire can be ignited by thermal radiation and be the reason for major extinction events in deep geological times, with results giving little support for this hypothesis. Jensen et al. (2005) and Courtney-Mustaphi and Pizaric (2014a) examined subtler diagnostic features (morphology, surface features, lustre) of laboratory-produced charcoal morphotypes of a small number of North American grasses and leaves of coniferous and deciduous trees. Enache and Cumming (2006, 2007) and Mustaphi and Pizaric (2014a) classified charcoal morphologies in Canadian lake sediments based on particle shape, aspect ratio, and surface features, and linked these morphometric characteristics to fuel types. Courtney Mustaphi and Pizaric (2014a) also discussed the potential for categorising charcoal morphologies to explore relationships to taphonomic processes and fuel types. A number of recent studies have attributed fossil charred particles to specific fuel and fire types based on published morphotype categorisations (Walsh et al., 2010, Daniau et al., 2013; Aleman et al., 2013; Leys et al., 2015; Courtney-Mustaphi and Pizaric, 2014a,b, 2018; Feurdean et al., 2017, 2019a,b, 2020; Unkelbach et al., 2018).

This paper presents the first results of laboratory-produced (muffle oven) charcoal morphologies spanning a range of fuel types from 17 Siberian species, with the aim of characterising the diversity of charcoal morphologies produced by boreal understory

and forest vegetation to facilitate more robust interpretations of fuel sources from this region. Specifically, (i) it evaluates whether there are morphological distinctions (morphometrics and finer anatomical features) between species or fuel types, and (ii) the effect of burning temperature on the mass, morphometrics, and finer anatomical features of charred plant material. This combination of factors has not been widely tested in the laboratory, so this study has the potential to advance our understanding of the link between sedimentary charcoal morphologies and fire types and, since the species occur across most of the Northern Hemisphere, refine wildfire histories in boreal regions.

2 Material and Methods

2.1 Laboratory analysis

Plant materials used for laboratory burning experiments were identified in the field, stored in plastic bags for transportation, and air-dried. Selected materials include a range of fuel types (graminoid, trunk wood, twigs from trees and shrubs, leaves from coniferous and deciduous trees, shrubs, forbs, and ferns, moss, and fern stems with leaves) from the most common tree, shrub, herb, fern, and moss species around a forested bog near Teguldet village, Tomsk district, Russia. This light boreal taiga forest is primarily composed of *Pinus* and *Betula*. Additionally, needles and twigs of *Picea abies* were collected from Taunus, near Frankfurt am Main, Germany (Table 1). All plant material was collected from living plants, except trunk wood, which was taken from a dead tree.

To determine the mass, morphometrics, and finer diagnostic features of residual charred plant material, as well the effect of increasing temperatures on all these characteristics, the dried remains of individual plant species were placed in ceramic crucibles, weighed, covered with a lid to limit oxygen availability and avoid mixing of the charred particles, and heated for 2 h in a muffle oven (preheated for 1 h) then roasted at 250, 300, 350, 400, or 450 °C (File S1). No burning experiments were conducted at higher temperatures because all plant material turned to ash i.e., a solid residue mostly composed of minerals that crumbled apart into soot and flyash (Rein et al., 2014). The effect of mixing plant material in known ratios on charred mass and morphometrics at an intermediate temperature (300 °C) was also tested. For this experiment, the plant material was combined in volumes to approximate the predominant fuel mixtures for low-intensity surface fires (75% graminoid and moss: 25% shrub; 50% graminoid: 50% moss and fern), intermediate- to high-intensity surface fires (25% graminoid and moss: 75% shrub; 50% graminoid and moss: 50% shrub), and high-intensity crown fires (50% graminoid, shrub, and moss: 50% wood and leaf). The experimental temperatures were chosen based on the range of temperatures reported in the literature (250–500 °C; Umbanhower and McGrath, 1998; Orvis et al., 2005; Jensen et al., 2007; Pereboom et al., 2020). Although dry roasting in a muffle oven approximates some aspects of the heating conditions of vegetation in a natural fire, it does not explore the impact of how long the material was at specific burning temperature and oxygen conditions (Belcher et al., 2015; Hudspith et al., 2017). Dry roasting also reduces the influence of flame dynamics and turbulent airflow, and therefore plant tissue is more rapidly reduced to ash than in natural fires. After cooling, the remaining charred mass of each sample was weighed and the ratio of charred to pre-combustion mass was calculated. Charred samples were then split into two subsets. The first was left intact and stored as reference material. The second sub-sample was gently disaggregated with a mortar and pestle to mimic the

135 natural breakage that charcoal particles incur over time in sediment (Umbanhowar and McGrath, 1998; Crawford and Belcher,
2014; Belcher et al., 2015), then washed through a 125- μm sieve to remove smaller fragments. This second sub-sample was
used to make morphometric measurements and to characterize finer diagnostic features. Morphometric measurements of
individual charred particles were obtained from photographs taken at 4 \times magnification with a digital camera (Kern DXM
1200F). On average, more than 100 charcoal particles larger than 150 μm were automatically detected in most samples, except
for those burnt at higher temperatures, where particles were more prone to breaking up. The major (L) and minor (W) axes,
140 along with surface area (A) of each particle were measured following the algorithm given in Appendix A1, and the aspect ratio
 L/W was calculated. Finer diagnostic features such as shape, surface features (reticulates, tracheids with border pits, leaf veins,
the arrangement of epidermal cells, cuticles with stomata, etc), and cleavage were characterized at 4 \times by inspection of
microphotographs or observations of the charred particles themselves under a microscope or stereomicroscope.
To demonstrate the applicability of these experiments to fossil records, seven samples with high charcoal content were selected
145 from a sediment core from Ulukh-Chayakh mire near Teguldet village (Feurdean et al., in prep). Sample preparation followed
Feurdean et al. (2020) and included bleaching overnight and washing in a 150- μm sieve. The results were compared to pollen
and plant macrofossils data from the same core.

2.2. Numerical analysis

150 The medians and standard deviations of charcoal morphometrics (L , L/W , A) were aggregated for each species, fuel type, and
burn temperature, and are displayed as box plots. A two-tailed Mann-Whitney test was used to test whether the medians of the
charcoal morphometrics of various fuel types were equal (File S2). This test does not assume a normal distribution, only similar
distributions in both groups.

155 3. Results

3.1 The influence of temperature on charred mass production

Only needles and shrub leaves were greenish or brownish in colour at 250 $^{\circ}\text{C}$, but plant materials of all species were black
with a typical charcoal appearance after burning at 300 and 350 $^{\circ}\text{C}$. A few fuel types (graminoid, *Sphagnum*, and some twigs)
turned to ash at 400 $^{\circ}\text{C}$, whereas all other types of plant tissue became ash at 450 $^{\circ}\text{C}$ (File S1). Most of the charred materials
160 remained intact and retained all their morphological characteristics. However, samples burnt at higher temperatures tended to
break easily during sample manipulation.

The average percentage of charred mass retained at 300 $^{\circ}\text{C}$ decreased as follows (Table 2; Fig. 1): brown moss and fern (50%)
> shrub twigs (46%) > shrub leaf (44%) > forb leaf (42%) > needles (41%) > tree twigs (40%) > graminoid (29%) > *Sphagnum*
(22%) > trunk wood (11%). This trend in mass loss was similar at all temperatures, although charcoal mass showed a marked
165 decline from 38-84 % at 250 $^{\circ}\text{C}$ to 0.2-23% at 400 $^{\circ}\text{C}$ across all fuel types. The charred mass of mixed-fuel samples at 300 $^{\circ}\text{C}$
was lowest for samples with high contents of graminoid and *Sphagnum* (33-35%) and highest for samples with greater
proportions of shrub material (38%).

3.2 Fuel-dependent variations in length, aspect ratio, and surface area

170 Graminoid charcoals burnt at 300 °C ($L/W = 11.5$, Fig. 2b, f, Table 2) were consistently more elongate than those of twigs
(shrub, 5.2; tree, 3.8), moss and fern stems (4.6), and leaves (2.7). Charred needles were more elongate (3.1) than those of
leaves from heathland shrubs (2.4) and broadleaf trees (2.1). The Mann-Whitney test confirmed that (i) the median aspect ratio
of graminoids was significantly different from those of all other fuel types ($p < 0.001$), (ii) those of all types of wood were
175 different from those of leaves ($p < 0.001$) and moss (except at 350 °C) and (iii) those of leaves were different from those of
moss ($p < 0.001$; Table S2). The lengths (major axis, L) of charred particles from different fuel types, however, were less
clearly differentiated (Figs. 3a, c, Files S2, S3). The surface area (A) varied greatly between individual taxa and fuel types,
however, fragments of shrub leaves tended to be larger than all other fuel types (Figs. 3b, d, Table 2, Files S2, S4). The
morphometrics of mixed-fuel samples showed that charcoal from samples with abundant graminoids and moss were more
elongate (higher L/W) than those with higher proportions of shrubs, wood, and/or leaves (Fig. 2h). Similarly, the longest
180 charcoal particles (higher L) were from samples with greater proportions of graminoids and moss (Fig. 3e), whereas the
charcoals with the largest surface areas were from samples with more abundant shrubs and leaves (Fig. 3f). The aspect ratios
and lengths of individual taxa and fuel types changed slightly with temperature, but the general trends were similar across all
temperatures (Figs. 2, 3a, c, File S3). In contrast, relative surface areas varied more with temperature changes (Figs. 3b, d, File
S4).

185 3.3 Finer diagnostic features of the charcoal morphologies of various fuel types

3.3.1 Graminoid charcoal

Graminoid (*Carex*, *Calamagrostis*, *Eriophorum vaginatum*) charred particles were consistently flat, rectangular, and elongated
(Files S5a, S6a). They mostly broke parallel to the long axis when pressured, resulting in highly elongated pieces with straight
margins. They can also appear as featureless long, thin filaments. Charcoal produced at higher temperatures (350 °C) often
190 had more irregular, zig-zag, or denticulate margins. The most commonly preserved surface features were rectangular epidermal
cells or contained oval voids, reticulated or mesh patterns, and/or isolated veins.

3.3.2 Wood charcoal (trunk, tree and shrub twigs)

Wood charcoal pieces from tree trunk (*Pinus sylvestris*) were blocky and quadrilateral with corner angles of 90° (Files S5b,
195 S6b). Wood charcoal from tree (*P. sylvestris*, *P. sibirica*, *Picea abies*, *Betula pendula*) and shrub (Ericaceae) twigs showed
both quadrilateral and polygonal shapes. The edges of both trunk and twig charcoal were smooth, serrated, or denticulate, and
surface textures were smooth, foliated, or striated (File S5b). Trunk charcoal of *P. sylvestris* showed rows of brown, open pits
in the tracheid walls. Under the microscope, trunk charcoal fragments were shinier and darker than twig charcoals. Large
charcoal pieces often broke parallel to the long axis, producing many tiny, elongated pieces (trunks) or pieces of various forms
200 (twigs).

3.3.3 Leaf charcoal (needles, deciduous tree and shrub, forb, and fern)

Charred needle fragments were elongated and rectangular (corner angles of 90°; Files S5c, S6a). Their edges were smooth but became serrated and denticulate when broken. Surface features included visible venation and ridges. Charcoals from the leaves of deciduous trees (*Betula*), heathland shrubs (*Oxyccocum*, *Ledum*, *Camadaphne*, *Vaccinium*), herbaceous plants (*Rubus*), and ferns (Polypodiaceae) were polygonal. Only those of *Cnidium* leaves were elongated, reflecting their needle shape. Edges were mostly undulate, but sometimes smooth or denticulate. Surface textures were generally smooth (featureless), but sometimes included visible venation and ridges. When broken, they showed voids, reticulated mesh patterns, and curly fibres. Birch leaves produced visible charred veins with three branches diverging from a node. When pressured and broken, small leaf pieces had fracture lines radiating out at a variety of angles.

3.3.4 Moss and fern stems

Sphagnum produced two types of charcoal morphologies. Stems produced elongate particles with ramifications (scars) where leaves branched from the stem. Leaves preserved the anatomical features of the unburned leaves, i.e., a mesh-like appearance (Files S5a, S6a). *Polytrichum* produced several charcoal morphologies (quadrilateral, polygonal, or curved with angular edges) with generally featureless surfaces, although some showed mesh patterns. This charcoal type often splits along the main axis. *Equisetum* leaves and stems were generally quadrilateral with straight, undulate, or denticulate margins, and oval voids and reticulated mesh patterns on their surfaces.

3.4 Morphometrics and finer diagnostic features of fossil charcoal

The average aspect ratio of the seven sediment samples varied between 2.8 and 11.1, whereas the average length between 403 and 742 (Table 3). Samples with higher aspect ratios contained abundant morphologies of graminoids, *Equisetum* and moss (Table 3). The average surface area varied from 16.565 to 248.287, where samples with greater surface area have higher number of leaves.

4 Discussion

These laboratory burning experiments provide information on charcoal morphology i.e., morphometrical aspects, fine diagnostic features, and charcoal production for 17 plant species belonging to seven fuel types from boreal Siberia. This dataset expands the geographical coverage of fuel types researched, demonstrates the applicability of charcoal morphology assemblages to fossil records, and improves the interpretation of fire types based on charcoal morphologies.

4.1 The influence of combustion temperature on charcoal production: implications for charcoal-based fire reconstructions

Knowledge of the charred mass is critical for determining biases in charcoal production to biomass quantity and fire temperature (Walsh and Li, 1964). Results from these burning experiments clearly show that the effect of temperature on

charcoal production is fuel dependent. Graminoid, *Sphagnum*, and trunk wood produce the lowest amounts of charcoal per unit biomass and lost their mass more rapidly with increasing burning temperature i.e., from 40-63% at 250°C to 0.2-3% at 400°C (Fig. 1; Table 2). Leaves of heathland shrubs, forbs, and ferns (Polypodiaceae), as well as fern stems with leaves (*Equisetum*), produced the most charcoal per unit biomass and retained the greatest mass at higher temperatures from 50-84% at 250°C to 4-24% at 400°C (Fig. 1). The charred mass of mixed-fuel samples burnt at 300 °C also changed according to the dominant fuel type. Peerboom et al. (2020) burned plant tissue from similar taxa occurring in the Alaskan tundra in a muffle oven (at 500 °C) and likewise found that graminoids have a lower charred-mass (25-27%) retention than shrubs (up to 33%). However, they did not test the leaves and wood of shrubs separately and they did not investigate the effect of different temperatures on charred mass. Burning experiments on American forest steppe plants in a muffle oven (at 350°C) and under open flame conditions showed that the mass retention of grass and deciduous leaf charcoal decreased more rapidly with temperature compared to wood charcoal (Umbanhower and McGrath, 1998), in partial agreement with the findings from this study. A calorimetric combustion study of various fuel types from mostly tropical plants found the charred mass of wood, needles, and *Equisetum* was greater than that of other leaf types, because of higher bulk densities and fuel load (Hudspith et al., 2017).

Although oven experiments do not fully replicate the burn conditions of natural wildfires, these experimental findings have practical implications for charcoal-based fire reconstructions. First, fuel types with low charred-mass retention at higher temperatures are likely to be under-represented in the sedimentary charcoal record. Cyperaceae (sedges) are the most common graminoids in fens and meso- and eutrophic bogs worldwide, and *Eriophorum* (sedge) and *Sphagnum* (moss) are common in oligotrophic bogs. These fuel types will turn to ash even in relatively low-intensity fires (<300°C) and thus may leave little or no trace of charcoal in sediments. Second, the woody biomass of typical heathland shrubs from oligotrophic bogs such as *Vaccinium*, *Camadaphne*, *Oxycoccus*, and *Ledum*, as well as brown moss (*Polytrichum commune*), which is common in all boreal habitats, preserve almost half of their biomass when burnt at temperatures up to 300 °C but their mass declines strongly (11-15%) at higher temperatures (>350°C). This suggests that charcoal from these fuel types is likely to be preserved only when fires are of low- to intermediate-intensity. Third, the leaves of shrubs, forbs, and ferns, and stems with leaves of *Equisetum* are more likely to persist as charcoal (27-38%) after high-temperature fires and thus may contribute disproportionately to sedimentary charcoal. Mineral constituents can slow pyrolysis (thermal decomposition of plant material producing volatile products and a solid charred residue) and this may explain why *Equisetum* stems, with high silica content, produce more charcoal. Fuel with higher lignin content such as wood might also be expected to produce more charcoal than fuels higher in cellulose and hemicellulose such as leaves (Yang et al., 2007). However, this did not appear to be the case in the current experiments where leaves retained a higher charred mass than the wood with increasing burn temperature. This suggests there is a need for further research on the quantitative relationship between temperature and charcoal mass retention to fuels with various structures, chemistry, and bulk density.

4.2 Fuel-dependent variability in charcoal morphometrics: implications for the reconstruction of fuel-type and transportation by air and water

270
275
280
285
290
295
300

The consistency of the morphometrics of charred fragments between species in the same genus or family suggests that these measurements are useful for fuel-type identification. Graminoid charcoal particles are at least twice more elongated (6.7-11.5) than all other charcoal types and differ the most from leaf charcoal across all temperatures (Fig. 2; Table 2; File S7). Highly elongate and narrow graminoid charcoal are thought to reflect the occurrence of conspicuous veins parallel to the long axis (Umbanhowar and McGrath, 1998; Crawford and Belcher, 2014). Charred fragments of leaves (2.0- 2.7 broadleaves; 3.1-3.5 needles) are also markedly more circular than those of other fuel types. However, there is some degree of overlap between the aspect ratios of twigs (2.5-5.2), and moss and fern stems (3.5-4.75; Table 2). In agreement with Crawford and Belcher (2014) and Umbanhowar and McGrath (1998), these experiments showed that smaller particles have a lower aspect ratio (more circular particles). Although larger charcoal fragments may be more suitable to categorise fuel type, there is no obvious threshold for determining what size particles should be used. Charcoal fragments from mixed-fuel samples preserve the aspect ratio of the dominant fuel type; particles with highest aspect ratios (3.5) were found in samples with greater proportions of graminoids and moss. Length and mean surface area do not appear to distinguish between fuel types reliably, except for the slight tendency that charred shrub leaf particles are larger than those of all other fuel types (Figs. 3, File S4; Table 2). The larger shrub leaf fragments may be explained by the arrangement of leaf venation, with fragments breaking along the three branching veins that diverge from nodes (Umbanhowar and McGrath, 1998; Jenssen et al., 2007). The measured aspect ratio of graminoids and shrub and forb leaves from this study are quite similar to those from the Alaskan arctic, where graminoids (*Eriophorum vaginatum* and *Carex bigelowii*) show aspect ratios ranging from 5.46 to 8.09 (mean 6.77), and shrubs (*Ledum palustre*, *Salix pulchra*, *Betula nana*, *Rubus chamaemorus*, *Vaccinium vitis-idaea*) from 2.09 to 2.50 (mean 2.42; Pereboom et al., 2020; Table 4). Considerably shorter graminoid particles (3.62) were obtained from American steppe forests (Umbanhowar and McGrath, 1998), although the aspect ratios obtained for leaves (1.91) and wood (2.13) were closer to those obtained in this study. Crawford and Belcher (2014) produced charcoal under laboratory conditions with an aspect ratio of 3.7 for graminoids, 2.23 for leaves, 1.97 for wood, and 2.8 for *Pinus sylvestris* needles (Table 4). Fossil charcoal assemblages from tropical African forests and grasslands were used to separate graminoids (aspect ratio <2.0) from shrubs (>2.0; Aleman et al., 2013). However, Daniau et al. (2013) interpreted an increase in the aspect ratio as an indication of the increased proportion of burning of the grass fuel. Mustaphi and Pisaric (2014a) also observed that burning monocotyledons from boreal Canada in the laboratory generally produced more elongated charcoal morphologies than other fuels. In terms of surface area, Umbanhowar and McGrath (1998), show a surface area of 65.630 μm^2 (56.737 herein) for graminoids, 50.150 μm^2 (103.138 herein) for wood and 64.946 μm^2 (versus 114.952 herein) for leaves, comparable to Pereboom et al. (2020) who found little differentiation between average surface area for shrub (88.246 μm^2), and graminoid (87.474 μm^2) species. The present study suggests that the aspect ratio is generally preserved for all fuel types over the range of temperatures explored, but the length and surface area of fuel types changed less consistently with increasing temperature. Umbanhowar and McGrath, (1998) found that burn

temperature did not significantly change the aspect ratios of graminoid and leaf charcoals, but marginally reduced those of wood.

305 The combined results from this study and the published literature suggest that, despite some variability in morphometrics of charcoal assemblages from similar fuel types, the aspect ratio decreases from graminoids to wood and leaves. Differences in the aspect ratio might allow the distinction of graminoids from other fuel types in a consistent way. Although there is a wide range of individual measurements, the mean aspect ratios of the three graminoid species (6.7-11.5) suggests that a threshold aspect ratio of 6 could be used to discriminate graminoids. This threshold value appears to be most consistent for wetland graminoids (mean 6.77 for arctic Alaska) but may be too high for graminoids from temperate grasslands (3.8-4.66; Table 4; 310 File S7). Although there is also a good consistency in the aspect ratio of laboratory-produced wood (2.1-4.5) and leaf (2.0-3.5) charcoal particles across studies, these values overlap, suggesting that it is not possible to specify a threshold value at which charcoal particles are indicative of wood or leaves. The use of charcoal morphologies in fuel type identification, therefore, requires the use of fine anatomical features (see section 4.3) or validation from other sources such as anthracological analysis as employed in archaeobotanical studies (Hubau et al., 2012; 2013; 2015; de Melo Júnior, 2017).

315 Particle shape affects the behaviour of charcoal during transportation by air (Clark and Hussey, 1996; Clark, 1998) and water (Nicols et al., 2000). Models, assuming a uniform spherical particle shape, and empirical data of transportation by fume indicate that the amount of charcoal particles is greatest near the fire source (Clark et al., 1998; Clark and Royall, 1995; Tinner et al., 2006; Higuera et al., 2007; Peters and Higuera, 2007). However, recent models accounting for different shapes, sizes, and densities of charcoal show that non-spherical particles have lower settling velocities than spherical particles and produce a 320 spatially more extensive and heterogeneous particle-size distribution pattern, i.e, dispersal distances for spherical and aspherical particles greater than 150 μm could be up to 20 km apart (Vacula and Richter 2018). Similarly, Clark and Hussey (1996) derived a velocity index for sedimentary charcoal particles and found that non-spherical particles have lower settling velocities and higher residence time into the atmosphere than the elongated particles. Based on these studies, it appears that non-spherical charcoal particles (elongated) such as those of graminoids, moss, and ferns are likely to have a more 325 heterogeneous distribution and be deposited farther away from the origin of a fire than the rounder, polygonal leaf particles. Erosion during hydrological transportation can also change the shape of buried (sedimentary) charcoal and can be an important consideration when interpreting charcoal morphometrics (Patterson et al., 1988; Nichols et al., 2000; Scott et al., 2000).

Laboratory experiments simulating fluvial transportation found that the surface area of leaf charcoal decreases and circularity increases with transportation, whereas changes in the shape of woody particles is less evident with transportation, and grassy 330 charcoal preserves a high aspect ratio during transportation (Crawford and Belcher, 2014). However, Nichols et al (2000), found a slight rounding of sharp-angled edges of wood and a greater propensity for breakage of charcoal produced at higher temperatures, the latter also found in this study. These findings give further support that the typical appearance of graminoids as elongated particles and of leaves as circular is preserved during transportation. Nevertheless, other studies using sedimentary charcoal records suggest that erosion during transport accounts for the rounding (degree angles are eroded) of robust charcoal 335 types such as wood, whereas fragile pieces of leaves and grass may break (Vanniery et al., 2003; Mustaphi and Pisaric, 2014b;

Mustaphi, et al., 2015). The differential transportation by air and fragility of sedimentary charcoal morphotypes calls for investigations for the influence of particle shape on charcoal transportation and strategies targeting coring locations for generating robust quantitative data for palaeofire interpretations.

4.3 Finer diagnostic features of the charcoal morphologies for fuel type identification

340 Results from fine diagnostic features on charcoal particles show that these can be used to attribute charcoal particles to certain
fuel types with some confidence. Apart from the extremely elongated shape that differentiates graminoid charred particles
from all other fuel types, graminoids are further distinguished under both microscope and stereomicroscope by their flat
appearance and breakage into thin filaments (Files S5a, S6a). Rectangular epidermal cells, reticulate meshes, oval voids of
former epidermal stomata are also good diagnostic features of graminoids (Grosse-Brauckman, 1974). The graminoid
345 charcoals produced in this study are most similar to types C4, C6, D1, D2, and D3 described by Courtney-Mustaphi and Pisaric
(2014a) and Enache and Cumming (2006). Comparative studies on graminoid charcoal originating from the Poaceae (grass)
and Cyperaceae (sedge) would further improve the identification of fuel types given the ecological differences of the two
groups, with sedges growing on wetlands and grass on drier habitats.

A distinct feature of woody charcoal is that they are layered with foliated or striated textures and break into many tiny particles
350 when pressured (Files S5a, S6b). This is due to the abundance of fibres and xylem, which leads to charcoals splitting at various
angles (Vaughan and Nichols, 1995). Additionally, conifer wood charcoal presents distinct rows of open pits in the tracheid
walls (Schweingruber, 1978). Attempts to distinguish between charred trunk and twig particles were less successful, although
charred trunk particles are blockier. Foliated charred wood fragments also share appearance with moss and fern stems. These
woody charcoals are most similar to types A1, B1, B2, and B3 (Mustaphi and Pisaric, 2014a; Enache and Cumming, 2006).

355 Typical features of charred deciduous leaves are their polygonal shapes with surfaces characterised by void spaces or undulated
surfaces (Files S5c, S6a). Netted venation is also sometimes visible, mostly with three branches diverging from a node. In
contrast, conifer needles are elongated, often show ramification, and can have a wood-like appearance. The deciduous leaf
charcoals found here are most similar to morphologies A2, A3, A4, A5, and A46, and conifer needle charcoals to C1, C2, and
C3 (Courtney-Mustaphi and Pisaric, 2014a; Enache and Cumming, 2006).

360 Charred *Sphagnum* leaves preserve the meshed pattern of fresh plant material (Grosse-Brauckman 1972). Often, stems contain
ramification, likely scars of former leaves (Files. S5a, S6a). Both *Sphagnum* and *Polytrichum* charcoals present curvy
fragments not seen in other fuel types. However, stems of *Sphagnum* and *Polytrichum* can be easily mistaken for shrub twigs.
Burnt *Equisetum* can resemble graminoid charcoal. Charred moss is similar to morphologies C4 and C7 (Courtney-Mustaphi
and Pisaric, 2014a; Enache and Cumming, 2006).

365

4.4 The morphometrics and morphologies of fossil charcoal particles: implications for fuel type identification

Charcoal fragments from Holocene samples ranging from 6700 to 180 cal yr BP at the Ulukh-Chayakh mire preserved the
aspect ratio of the dominant morphologies i.e., particles with the highest aspect ratios (4-11) were found in samples with a

greater proportion of graminoids, *Equisetum*, and moss (Table 3). Likewise, a greater surface area was found in samples with
370 a higher number of leaves. Comparative results from fossil charcoal morphologies and morphometrics to those from pollen
and plant macrofossils from the same depths show a partial agreement. For example, the pollen record indicates that
percentages of tree were >90% and shrub up to 3% during the entire period, whereas the abundance of woody charcoal
morphologies increased infrequently. This suggests that although there was a continuous source of woody fuel to burn, high
intensity fire, producing wood charcoal occurred only occasionally. There is, however, a better agreement between samples
375 with greater aspect ratios and morphologies of understory vegetation i.e., graminoids, *Equisetum*, and moss, and the proportion
of these plants in the pollen and plant macrofossil records (Table 3). These findings are in line with the Siberian wildfire
behaviour of predominantly low intensity, surface fires fuelled by graminoids, forbs, ferns, and mosses, or intermediate
intensity surface fires (shrubs) and only infrequently as high intensity crown fires (Anderson, 1982). Another practical
application of this finding is that the morphometrical and morphological characterisation of fossil charcoal is more
380 representative of fuel type (what plant types were burning), whereas the pollen data and plant macrofossils reflect plant types
growing regionally and locally.

4.5 Applications for fuel and fire type reconstructions

The physical and chemical characteristics of fuel are key factors influencing ignition and fire propagation. Major chemical
385 components of fuels are cellulose, hemicellulose, and lignin, and minor ones include terpenes, resins, and minerals (Plana and
Pastor, 2014). Fuels rich in cellulose and hemicellulose (i.e., leaves) pyrolyse at a lower and narrow temperature range (200
and 400 °C), whereas those rich in lignin (i.e., wood) pyrolyse at a higher and over a wider range of temperatures (160-900°C;
Yang et al., 2007). Fuel types rich in terpene and resins (conifer wood, needles, Ericaceae) burn faster and at higher
temperatures, whereas those rich in mineral components (graminoids) burn at lower temperatures (200 °C) and less efficiently
390 (Plana and Pastor, 2014). Results from the current burning experiments and fossil charcoal samples suggests that the combined
use of morphometric and morphological features and charred mass can help distinguish some of the predominant fuel source.
Knowledge of the fuel source may in turn provide clues on fire type, i.e., the combination of fire intensity (temperature) and
severity (effect on vegetation). Fossils samples dominated by graminoid morphotypes show a high aspect ratio (4 -11), in line
with the elongated shape of graminoid charcoal found in burning experiments (>6.7-11.5). As graminoid charcoal typically
395 preserves at lower temperatures, it likely suggests a graminoid fuel source, and therefore a lower-intensity fire (Fig.4). Fossil
samples with abundant leaves and wood morphologies showed considerably lower aspect ratios (3-3.2), in agreement with
values from laboratory-derived morphologies of leaves (2.1-3.5) and wood (2.0-5.2). Thus, shorter, and bulkier charcoal
particles likely indicate the increased prevalence of leaves and wood as a fuel source (Figs. 2, 4). Because the morphometrical
and morphological characteristics of leaves, and wood from trees and shrubs overlap, it is hard to distinguish between high-
400 intensity surface fires, combusting living shrubs and dead wood and leaves, and high-intensity crown fires which have burnt
living trees. Nevertheless, the fact that the past fires may have been of higher intensity at times of leaves and wood charcoal

dominance than during the graminoid charcoal dominance is additionally suggested by increased abundance of morphologies of *Equisetum* and *Polytrichum*, taxa found to remain as charcoal after burning at high temperature.

405 In summary, the consistency of results from this study with those from the literature on various vegetation types (boreal, temperate, and tropical woodlands, and grasslands) suggests the potential of charcoal morphometrics and morphologies in palaeoecology. For example, the expansion of open habitats during deep geological times or with human impact, the recession of latitudinal and elevational treelines, or the predominant occurrence of surface fires is likely to be reflected in an increase in aspect ratio and graminoid morphologies relative to total biomass burning. Conversely, the closing up of the forests, shrub encroachment, or the predominance of crown fires may show itself in a decreased aspect ratio of particles and increased bulky morphologies derived from leaves and wood. Results could also provide forest managers with the range of fire types that key 410 boreal species experienced in the past, useful when aiming to make choices for prescribed burning to remove fuel and prevent large fires or select species that will be fit to cope with future fire regimes. Answering all questions, however, will need further investigations to relate the proportion of charcoal morphotypes to the quantity of biomass and extend the morphometric and morphological characterisation to key species of interest.

415

5. Conclusions and recommendations

This study presents the first results on the morphometry and other diagnostic features of charred particles produced in the laboratory from seven fuel types comprising 17 plant species from boreal Siberia and demonstrates the applicability of these experiments to interpreting fossil charcoal records. The use of a higher number of fuel types from species with broad 420 geographical coverage combined with an exploration of various combustion temperatures improves the link between charcoal morphologies, fuel types, and fire characteristics. Results show a distinct effect of temperature on fuel types, suggesting that species with low mass retention (graminoid, *Sphagnum*, and trunk wood) during high fire temperature are likely to be underrepresented in the fossil charcoal record. The aspect ratio was the strongest indicator of fuel type. Graminoid charcoal particles are more elongate (6.7-11.5) than all other fuel types and a threshold above 6 that may be indicative of wetland 425 graminoids, leaves are the shortest and bulkiest (2.1-3.5), and twigs and wood are intermediate (2.0-5.2). Other diagnostic features can be used to separate wood, graminoids, and leaves, but not to make further distinctions within these fuel types. Distinct particle shape may influence charcoal transportation, with elongated particles (graminoids, moss, and ferns) potentially having a more heterogeneous distribution and be deposited farther away from the origin of a fire than the rounder, polygonal particles (leaf).

430 Despite these limitations, the combined use of particle aspect ratio and charred morphotypes should allow more robust interpretations of changes in fuel source and fire type from charcoal records. Future efforts to determine fuel sources based on analyses of small charcoal fragments will require: i) a more detailed examination of plant anatomy; ii) investigation of the proportion of particular charcoal morphotype to the quantity of biomass; iii) quantifying the relationship between the chemical composition of fuels, combustion temperature, and charcoal production; iv) determining the influence of particle shape on differential transportation and fragility; and v) using image-recognition software to collect data on charcoal characteristics 435

such as roundedness, reflectance and others features that could improve the estimation of fire temperature and erosion during transportation.

Data sets: A limited amount of burnt plant material can be made available upon request.

440

Author contribution: AF designed the burning experiments and carried them out. AF performed the morphometrical and morphological analyses; AF performed numerical analysis and data presentation; AF wrote the manuscript; AF acquired the financial support for the project leading to this publication.

Competing interests: The author declare that they have no conflict of interest.

445 **Special issue statement:** The role of fire in the Earth system: understanding interactions with the land, atmosphere, and society (ESD/ACP/BG/GMD/NHESS inter-journal SI)

Acknowledgements I would like to thank Markus Rosenstihl for the help to develop the code for automatic detection of charred particles and drawing the pictograms in Fig 4; Dagmar Fritzsich for initial brainstorming on the burning experiments; Doris Schneider for help with burning plant material in muffle oven; and Sergey Kirpotin for help with identification of plant species in field. Simon Hutchinson and Mirjam Pfeiffer provided some linguistic suggestions.

450

Financial support: This work was supported from the Deutsche Forschungsgemeinschaft grant number FE_1096/6.

455 **References**

- Anderson, R.C.: The eastern prairie-forest transition – an overview. In Ballard, H.E. Jr., Brewer, L.S. and Fox, C., editors, Proceedings of the eighth North American Prairie Conference, Kalamazoo, Michigan: Western Michigan University, 86–92, 1992.
- Aleman, J.C., Blarquez, O., and Bentaleb, I., Bonté, P., Brossier B, Carcaillet, C., Gond, V., Gourlet-Fleury, S., Kpolita, A., Lefèvre, I., and Oslisly, R.: Tracking landcover changes with sedimentary charcoal in the Afrotropics. *The Holocene*, 23, 1853–1862, <https://doi.org/10.1177/0959683613508159>, 2013.
- 460 Belcher, C.M., Collinson, M.E., and Scott, A.C.: Constraints on the thermal energy released from the Chicxulub impactor: new evidence from multi-method charcoal analysis. *Journal of the Geological Society*, 162, 591-602, <https://doi.org/10.1144/0016-764904-104>, 2005.
- 465 Belcher, C.M., Hadden, R.M., Rein, G., Morgan, J.V., Artemieva, N., and Goldin, T.: An experimental assessment of the ignition of forest fuels by the thermal pulse generated by the Cretaceous–Palaeogene impact at Chicxulub. *Journal of the Geological Society*, 172, 175-185. <https://doi.org/10.1144/jgs2014-082>, 2015.

- Clark, J. S.: Particle motion and the theory of charcoal analysis: source area, transport, deposition, and sampling. *Quaternary research*, 30, 67-80. [https://doi.org/10.1016/0033-5894\(88\)90088-9](https://doi.org/10.1016/0033-5894(88)90088-9), 1988.
- 470 Clark, J.S. and Hussey, T.C.: Estimating the mass flux of charcoal from sedimentary records: effects of particle size, morphology, and orientation. *The Holocene*, 6, 129–44. <https://doi.org/10.1177/095968369600600201>, 1996.
- Courtney-Mustaphi, C.J. and Pisaric, M.F.: A classification for macroscopic charcoal morphologies found in Holocene lacustrine sediments. *Progress in Physical Geography*, 38, 734-754. <https://doi.org/10.1177/0309133314548886>, 2014a.
- 475 Courtney-Mustaphi, C.J. and Pisaric M.F.J.: Holocene climate-fire-vegetation interactions at a subalpine watershed in southeastern British Columbia, Canada. *Quaternary Research*, 81, 228–239. <https://doi.org/10.1016/j.yqres.2013.12.002>, 2014b.
- Courtney Mustaphi, C.J., Davis, E.L., Perreault, J.T., and Pisaric, M.F.J.: Spatial variability of recent macroscopic charcoal deposition in a small montane lake and implications for reconstruction of watershed-scale fire regimes. *Journal of Paleolimnology*, 54, 71–86. <https://doi.org/10.1007/s10933-015-9838-2>, 2015.
- 480 Courtney-Mustaphi C.J. and Pisaric, M.F.J.: Forest vegetation change and disturbance interactions over the past 7500 years at Sasquatch Lake, Columbia Mountains, western Canada. *Quaternary International*, 488, 95-106, <https://doi.org/10.1016/j.quaint.2017.03.045>, 2018.
- Crawford, A.J. and Belcher, C.M.: Charcoal morphometry for paleoecological analysis: the effects of fuel type and transportation on morphological parameters. *Applications in Plant Sciences*, 2, 1400004. <https://doi.org/10.3732/apps.1400004>, 2014.
- 485 Daniau, A.-L., Goñi, M.F.S., Martinez, P., Urrego, D.H.V., Bout-Roumazelles, V., Desprat, S., and Marlon, J.R.: Orbital-scale climate forcing of grassland burning in southern Africa. *Proceedings of the National Academy of Sciences, USA*, 110, 5069– 5073. <https://doi.org/10.1073/pnas.1214292110>, 2013.
- de Melo Júnior, J.C.F.: A new archaeobotanical protocol for collecting concentrated wood charcoal from archaeological bonfire sites. *International Journal of Development Research*, 7, 14241-14247, 2017.
- 490 Enache, M.D. and Cumming, B.F.: Tracking recorded fires using charcoal morphology from the sedimentary sequence of Prosser Lake, British Columbia (Canada). *Quat Res.*, 65, 282-292. <https://doi.org/10.1016/j.yqres.2005.09.003>, 2006.
- Enache, M.D. and Cumming, B.F.: Charcoal morphotypes in lake sediments from British Columbia (Canada): an assessment of their utility for the reconstruction of past fire and precipitation. *J. Paleolimnol.* 38, 347-363. <https://doi.org/10.1007/s10933-006-9084-8>, 2007.
- 495 Feurdean, A., Veski, S., Florescu, G., Vannièrè, B., Pfeiffer, M., O'Hara, R.B., Stivrins, N., Amon, L., Heinsalu, A., Vassiljev, J., and Hickler, T.: Broadleaf deciduous forest counterbalanced the direct effect of climate on Holocene fire regime in hemiboreal/boreal region (NE Europe). *Quaternary Science Reviews*, 169, 378-390. <https://doi.org/10.1016/j.quascirev.2017.05.024>, 2017.
- 500 Feurdean, A. and Vasiliev, I.: The contribution of fire to the late Miocene spread of grasslands in eastern Eurasia (Black Sea region). *Scientific reports*, 9, 1-7. <https://doi.org/10.1038/s41598-019-43094-w>, 2019a.

- Feurdean, A., Tonkov, S., Pfeiffer, M., Panait, A., Warren, D., Vanni re, B., and Marinova, M.: Fire frequency and intensity associated with functional traits of dominant forest type in the Balkans during the Holocene. *European Journal of Forest Research* 138, 1049–1066. <https://doi.org/10.1007/s10342-019-01223-0>, 2019b.
- 505 Feurdean, A., Florescu, G., Tan au, I., Vanni re, B., Diaconu, A.C., Pfeiffer, M., Warren, D., Hutchinson, S.M., Gorina, N., Ga ka, M., and Kirpotin, S.: 2020. Recent fire regime in the southern boreal forests of western Siberia is unprecedented in the last five millennia. *Quaternary Science Reviews*, 244, 106495. <https://doi.org/10.1016/j.quascirev.2020.106495>, 2020.
- 510 Goldammer, J.G. and Furyaev, V.V. (Eds): *Fire in Ecosystems of Boreal Eurasia*, Springer, Dordrecht, Germany, 1996.
- Grosse-Brauckmann, G.:  ber pflanzliche Makrofossilien mitteleurop ischer Torfe. *TELMA*, Band 4. 51-117, Hannover, 1974.
- Higuera, P. E., Peters, M. E., Brubaker, L. B., and Gavin, D. G.: Understanding the origin and analysis of sediment-charcoal records with a simulation model. *Quaternary Science Reviews*, 26, 1790-1809. <https://doi.org/10.1016/j.quascirev.2007.03.010>, 2007.
- 515 Hawthorne, D., Mustaphi, C.J.C., Aleman, J.C., Blarquez, O., Colombaroli, D., Daniau, A.L., Marlon, J.R., Power, M., Vanni re, B., Han, Y. and Hantson, S., and GMCD.: A tool for exploring proxy-fire linkages and spatial patterns of biomass burning. *Quaternary International*, 488, 3-17. <https://doi.org/10.1016/j.quaint.2017.03.046>, 2018.
- Hudspith, V.A., Hadden, R.M., Bartlett, A.I., and Belcher, C.M.: Does fuel type influence the amount of charcoal produced in wildfires? Implications for the fossil record. *Palaeontology*, 61, 159-171. <https://doi.org/10.1111/pala.12341>, 2018.
- 520 Hubau, W., Van den Bulcke, J., Mees, F., Van Acker, J., and Beeckman, H.: Charcoal identification in species rich biomes: a protocol for Central Africa optimised for the Mayumbe forest. *Review of Palaeobotany and Palynology*, 171, 164– 178, <https://doi.org/10.1016/j.revpalbo.2011.11.002>, 2012.
- Hubau, W., Van den Bulcke, J., Kitin, P., Brabant, L., Van Acker, J., and Beeckman, H.: Complementary imaging techniques for charcoal examination and identification. *IAWA Journal*, 34, 147– 168. <https://doi.org/10.1163/22941932-00000013>, 2013.
- 525 Hubau, W., Van den Bulcke, J., Van Acker, J., and Beeckman, H.: Charcoal-inferred Holocene fire and vegetation history linked to drought periods in the Democratic Republic of Congo. *Glob Change Biol*, 21, 2296-2308. <https://doi.org/10.1111/gcb.12844>, 2015.
- 530 Jensen, K., Lynch, E., Calcote, R., and Hotchkiss, S.C.: Interpretation of charcoal morphotypes in sediments from Ferry Lake, Wisconsin, USA: do different plant fuel sources produce distinctive charcoal morphotypes? *Holocene* 17, 907-915. <https://doi.org/10.1177/0959683607082405>, 2007.
- Jones, M.W., Smith, A., Betts, R., Canadel, J.G., I , Prentice, I.C., and Le Qu er , C.: Climate change increases the risk of wildfires. <https://sciencebrief.org/briefs/wildfires>, 2020.

- 535 Leys, B., Brewer, S.C., McConaghy, S., Mueller, J., and McLauchlan, K.K.: Fire history reconstruction in grassland ecosystems: amount of charcoal reflects local area burned. *Environmental Research Letters*, 10,114009, 2015.
- Marlon, J. R., Kelly, R., Daniau, A.-L., Vanni re, B., Power, M. J., Bartlein, P., Higuera, P., Blarquez, O., Brewer, S., Br ucher, T., Feurdean, A., Romera, G. G., Iglesias, V., Maezumi, S. Y., Magi, B., Courtney Mustaphi, C. J., and Zhihai, T.: Reconstructions of biomass burning from sediment-charcoal records to improve data–model comparisons, *Biogeosciences*, 13, 3225–3244. <https://doi.org/10.5194/bg-13-3225-2016>, 2016.
- 540 Moritz, M.A., Batllori, E., Bradstock, R.A., Gill, A.M., Handmer, J., Hessburg, P.F., Leonard, J., McCaffrey, S., Odion, D.C., Schoennagel, T., and Syphard, A.D.: Learning to coexist with wildfire. *Nature*, 515, 58–66. <https://doi.org/10.1038/nature13946>, 2014.
- MacDonald, G.M., Larsen, C.P.S., Szeicz, J.M., and Moser, K.A.: The reconstruction of boreal forest fire history from lake sediments: a comparison of charcoal, pollen, sedimentological, and geochemical indices. *Quaternary Science Reviews*, 10, 53–71 . [https://doi.org/10.1016/0277-3791\(91\)90030-X](https://doi.org/10.1016/0277-3791(91)90030-X), 1991.
- 545 Nichols, G.J., Cripps, J.A., Collinson, M.E., and Scott, A.S.: Experiments in waterlogging and sedimentology of charcoal: results and implications, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 164, 43-56, [https://doi.org/10.1016/S0031-0182\(00\)00174-7](https://doi.org/10.1016/S0031-0182(00)00174-7), 2000.
- 550 Orvis, K.H., Lane, C.S., and Horn, S.P.: Laboratory production of vouchered reference charcoal from small wood samples and non-woody plant tissues. *Palynology* 29, 1–11. <https://www.jstor.org/stable/3687800>, 2005.
- Patterson, W.A. III, Edwards, K.J., and Maguire, D.J.: Microscopic charcoal as an indicator of fire. *Quaternary Science Reviews* 6, 3–23, 1987.
- Pereboom, E.M., Vachula, R.S., Huang, Y., and Russell, J.: The morphology of experimentally produced charcoal distinguishes fuel types in the Arctic tundra. *The Holocene*, 7, 1-6. <https://doi.org/10.1177/0959683620908629>, 2020.
- 555 Peters, M. and Higuera, P.: Quantifying the source area of macroscopic charcoal with a particle dispersal model. *Quaternary Research*, 67, 304-310. doi:10.1016/j.yqres.2006.10.004, 2007.
- Planas, E. and Pastor, E.: Wildfire behaviour and danger rating. In: Belcher, C., editor, *Fire phenomena and the Earth system: an interdisciplinary guide to fire science*, 53-76. doi:10.1002/9781118529539, John Wiley & Sons, Ltd, 2014.
- 560 Prince, T.J., Pisaric, M.F., and Turner, K.W.: Postglacial reconstruction of fire history using sedimentary charcoal and pollen from a small lake in southwest Yukon Territory, Canada. *Frontiers in Ecology and Evolution*, 6, 209. <https://www.frontiersin.org/article/10.3389/fevo.2018.00209>, 2018.
- Rein, G.: Smoldering fire and natural hazard. In: Belcher, C., editor, *Fire phenomena and the Earth system: an interdisciplinary guide to fire science*,15-34. doi:10.1002/9781118529539, John Wiley & Sons, Ltd, 2014.
- 565 Scott, A.C.: Charcoal recognition, taphonomy and uses in palaeoenvironmental analysis. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 291, 11– 39. <https://doi.org/10.1016/j.palaeo.2009.12.012>, 2010.
- Schweingruber, F.H.: *Mikroskopische Holzanatomie Formenspektren mitteleurop ischer Stamm- und Zweigh olzer zur Bestimmung von rezentem und subfossilem*. Z rcher AG, CH-6301, Zug, 1978.

- 570 Umbanhowar, C.E. and McGrath, M.J.: Experimental production and analysis of microscopic charcoal from wood, leaves and
grasses. *The Holocene*, 8, 341–346. <https://doi.org/10.1191/095968398666496051>, 1998.
- Unkelbach, J., Dulamsuren, C., Punsalpaamuu, G., Saindovdon, D., and Behling, H.: Late Holocene vegetation, climate,
human and fire history of the forest-steppe-ecosystem inferred from core G2-A in the ‘Altai Tavan Bogd’ conservation
area in Mongolia. *Veget Hist Archaeobot*, 27, 665–677 DOI: 10.1007/s00334-017-0664-5, 2018.
- 575 Vachula, R. S. and Richter, N.: Informing sedimentary charcoal-based fire reconstructions with a kinematic transport model.
The Holocene, 28, 173-178, <https://doi.org/10.1177/0959683617715624>, 2018.
- Vannière, B., Bossuet, G., Walter-Simonnet, A.V., Gauthier, E., Barral, P., Petit, C., Buatier, M., and Daubigny, A.: Land
use change, soil erosion and alluvial dynamic in the lower Doubs Valley over the 1st millenium AD (Neublans, Jura,
France). *Journal of Archaeological Science*, 1283-1299, 2003.
- 580 Vaughan, A. and Nichols, G.: Controls on the deposition of charcoal: implications for sedimentary accumulations of fusain.
*Journal of Sedimentary Research A*65, 130–135. <https://doi.org/10.1306/D426804A-2B26-11D7-8648000102C1865D>,
1995.
- Walsh, M.K., Whitlock, C., and Bartlein, P.J.: A 14,300- year-long record of fire–vegetation–climate linkages at Battle Ground
Lake, southwestern Washington. *Quaternary Research* 70: 251–264. <https://doi.org/10.1016/j.yqres.2008.05.002>, 2008.
- 585 Walsh, P.M. and Li, T.: Fragmentation and attrition of coal char particles undergoing collisions during combustion at
temperatures from 900–1100 K . *Combustion and Flame* 99, 749–757 . [https://doi.org/10.1016/0010-2180\(94\)90070-1](https://doi.org/10.1016/0010-2180(94)90070-1),
1994.
- Ward, D.E. and Hardy, C.C.: Smoke emissions from wildland fires. *Environment International* 17, 117–134, 1991.
- Whitlock, C., Larsen, C.: Charcoal as a fire proxy. In Smol, JP, Birks, HJB and Last, WM., editors, *Tracking environmental
change using lake sediments. Volume 3: terrestrial, algal, and siliceous indicators*. Kluwer Academic Publishers, 75-97,
590 10.1007/0-306-47668-1, https://doi.org/10.1007/0-306-47668-1_5, 2002.
- Yang, H., Yan, R., Chen, H., Lee, D.H., and Zheng, C.: Characteristics of hemicellulose, cellulose and lignin
pyrolysis. *Fuel*, 86, 1781– 1788. <https://doi.org/10.1016/j.fuel.2006.12.013>. 2007

595

600

Figure legends and embedded figures

605 **Figure 1.** The percent of charred mass retained after burning known plant species from Siberia in a muffle oven at 250, 300, 350, and 400 °C. Abbreviations: L, leaf; N, needles; t, twig; w, wood. The median mass retained for similar fuel types (identified by the same colour) are reported as black diamonds.




Figure 3. The median lengths (μm) and surface areas (μm^2) of charred particles from (a, b) individual taxa, (c, d) fuel types, and (e, f) fuel mixtures at 300 °C. Abbreviations as in Figure 1. See Figure 2 for description of box plots and colour coding.

625





Figure 4. Schematic representation of fire types and the potential link with fuel types burnt and predominant charcoal morphometrics (aspect ratio) and morphologies as well as charcoal production.



Tables

650 **Table 1.** List of plant materials burned. All plants are from Siberia, except for *Picea abies*, originating from Taunus, Germany. All plant material was dried before the combustion in the muffle oven. Leaves of deciduous trees and shrubs include veins and petioles. The term twig was used only for woody species i.e., deciduous and coniferous trees and shrubs and no distinction was made between soft young wood.

655	Plant type	Scientific name	Family	Common name	Plant burned
	Trees				
	Conifer tree	<i>Pinus sylvestris</i>	Pinaceae	Scots pine	Needles
	Conifer tree	<i>Pinus sylvestris</i>	Pinaceae	Scots pine	Twigs
	Conifer tree	<i>Pinus sylvestris</i>	Pinaceae	Scots pine	Dead wood
660	Conifer tree	<i>Pinus sibirica</i>	Pinaceae	Siberian pine	Needles
	Conifer tree	<i>Pinus sibirica</i>	Pinaceae	Siberian pine	Twigs
	Conifer tree	<i>Picea abies</i>	Pinaceae	Norway spruce	Needles
	Conifer tree	<i>Picea abies</i>	Pinaceae	Norway spruce	Twigs
	Deciduous tree	<i>Betula pendula</i>	Betulaceae	Silver birch	Leaves
665	Deciduous tree	<i>Betula pendula</i>	Betulaceae	Silver birch	Twigs
	Shrubs				
	Shrub	<i>Vaccinium myrtillus</i>	Ericaceae	Bilberry	Leaves
	Shrub	<i>Vaccinium myrtillus</i>	Ericaceae	Bilberry	Twigs
670	Shrub	<i>Oxycoccus palustre</i>	Ericaceae	Bilberry	Leaves
	Shrub	<i>Oxycoccus palustre</i>	Ericaceae	Bilberry	Twigs
	Shrub	<i>Empetrum nigrum</i>	Ericaceae	Bilberry	Leaves
	Shrub	<i>Empetrum nigrum</i>	Ericaceae	Bilberry	Twigs
	Shrub	<i>Ledum palustre</i>	Ericaceae	Bilberry	Leaves
675	Shrub	<i>Ledum palustre</i>	Ericaceae	Bilberry	Twigs
	Shrub	<i>Chamaedaphne calyculata</i>	Ericaceae	Bilberry	Leaves
	Shrub	<i>Chamaedaphne calyculata</i>	Ericaceae	Bilberry	Twigs
	Herbaceous				
680	Graminoid	<i>Eriophorum vaginatum</i>	Cyperaceae	Cotton grass	Leaves
	Graminoid	<i>Calmagrostis</i>	Poaceae	Reed grass	Leaves
	Graminoid	<i>Carex</i> spp.	Cyperaceae	Sedge	Leaves
	Forb	<i>Cnidium dubium</i>	Apiaceae		Leaves
	Forb	<i>Rubus</i> spp.	Rosaceae	Raspberry	Leaves
685	Fern	<i>Polypodium</i>	Polypodiaceae	Fern	Leaves
	Fern	<i>Equisetum palustre</i>	Equisetaceae	Horsetail	Stem
	Moss	<i>Sphagnum</i> spp.	Sphagnaceae	Peat moss	Stem +leaves
	Moss	<i>Polytrichum commune</i>	Polytrichiaceae	Hair moss	Stem +leaves

690

695

Table 2. Summary of the mean aspect ratio, length (μm), surface area (μm^2), and mass retained (%) of charcoal produced in the muffle oven for individual plant species and fuel types from Siberia (SD=standard deviation). For the full taxa name see Table 1.

Fuel types	Species	Aspect ratio at T/°C				Length at T/°C				Surface area at T/°C				Mass retained at T/°C			
		250	300	350	400	250	300	350	400	250	300	350	400	25	30	35	40
Graminoid	<i>Eriophorum</i>	7.2±5.0	11.1±6.4	5.8±3.7	N/A	521±545	797±496	635±74	N/A	37665	42744	51820	N/A	42.6	29	3.6	2.9
	<i>Calmagrostis</i>	7.4±4.6	12.9±11.8	6.0±3.8	N/A	620±622	951±712	440±370	N/A	82058	99009	30244	N/A	54.7	29.2	8.1	1.7
	<i>Carex</i>	7.4±5.1	10.6±7.0	8.3±6.0	N/A	487±499	841±646	690±709	N/A	32165	82947	51982	N/A	50.9	29.5	7.9	1.2
	Mean	7.3±0.1	11.5±1.2	6.7±1.4		543±64	862±79	588±131	N/A	50629	74900	44682	N/A	49.4	29.2	6.5	1.9
Moss & ferns	<i>Equisetum</i>	4.8±3.3	4.5±2.8	3.1±2.2	4.8±3.1	506±501	286±233	655±510	530±424	71381	25955	166741	75098	54.4	49.4	31.0	21.4
	<i>Polytrichum</i>	4.8±3.1	4.2±2.6	4.2±2.6	4.7±2.9	673±665	461±408	459±476	572±535	123311	84632	68170	99522	77.0	55.5	11.4	3.6
	<i>Sphagnum</i>	3.4±1.8	5.2±6.2	3.2±1.6	N/A	612±574	524±636	319±217	N/A	109185	77224	24724	N/A	38.3	21.7	8.3	0.2
	Mean	4.3±0.8	4.6±0.5	3.5±0.6	4.7±0.1	598±84	423±123	477±168	551±29	97346	62739	86545	87310	56.6	42.2	16.6	8.4
Wood (trunk)	<i>Pinus sylvestris</i>	2.0±0.9	4.9±2.8	2.5±0.9	N/A	408±347	391±300	482±384	N/A	105057	57028	97673	N/A	63.5	11.1	1.8	1.4
Wood (tree twig)	<i>Betula pendula</i>	2.8±1.4	4.5±3.1	2.0±0.8	N/A	459±402	361±236	347±85	N/A	120539	37630	54180	N/A	50.8	40.2	10.9	0.3
	<i>Picea abies</i>	2.3±1.2	3.1±1.9	2.5±1.3	N/A	435±371	439±303	598±575	N/A	92767	92107	199556	N/A	56.5	40.1	16.8	3.4
	<i>Pinus sibirica</i>	2.4±1.3	3.1±1.9	2.6±1.3	N/A	450±379	318±248	427±343	N/A	119815	15058	83461	N/A	56.0	43.3	11.0	1.3
	<i>Pinus sylvestris</i>	3.2±1.7	3.5±2.0	2.9±1.7	2.9±1.5	593±558	379±318	407±295	407±350	140004	74797	74929	88944	55.2	38.1	5.7	1.4

	<i>Mean</i>	2.5 ±0.4	3.8 ±0.8	2.5 ±0.3		469 ±72	377 ±44	452 ±95	N/A	11563 9	55051	10204 9	N/A	54. 5	40. 2	11. 2	1.5
Wood (shrub twig)	<i>Chamae daphne</i>	3.5±1 .9	6.3 ±3.5	3.7 ±2.1	N/A	387±34 0	347±28 8	525±52 8	521±55 5	61732	73382	13106 3	N/A	56. 8	41. 3	11. 4	1.6
	<i>Oxycocc us</i>	3.8±3 .0	6.2 ±3.5	4.7 ±3.3	4.1± 2.3	674±66 9	590±41 9	591±58 0	1053±7 00	15787 3	51906	94418	24919 7	78. 8	49. 4	35. 9	14. 8
	<i>Ledum</i>	4.4±2 .4	5.4 ±2.8	2.9 ±1.3	4.0± 2.7	333±19 7	458±28 8	342±26 6	617±58 0	57191	56836	25469 6	13186 0	56. 6	52. 5	15. 6	0.9
	<i>Vacciniu m</i>	5.0±3 .1	3.0 ±2.6	3.7 ±2.5	N/A	461±44 6	818±71 4	441±30 6	N/A	72214	29102 3	75212	N/A	60. 8	42. 2	16. 9	2.5
	Mean	4.2 ±0.7	5.2 ±1.5	3.8 ±0.7	4.0± 0.7	463±15 0	553±20 2	474±10 7	730±28 3	87252	11806 1	13659 7	N/A	62. 7	46	20	4.8
Needles	<i>Picea abies</i>	2.6±1 .6	2.2±1 .1	2.3 ±0.9	N/A	549±53 5	342±27 6	608±69 2	N/A	11837 5	86728	24236 4	N/A	67. 2	52. 3	26. 2	4.9
	<i>Pinus sibirica</i>	4.0±2 .3	3.6 ±1.8	4.7 ±2.9	N/A	606±58 7	385±29 2	432±41 4	N/A	11106 8	46629	52880	N/A	62. 3	34. 0	14. 3	2.4
	<i>Pinus sylv.</i>	4.0±2 .5	3.7 ±4.0	3.5 ±2.9	N/A	690±66 0	445±37 6	492±36 8	N/A	13450 4	86639	83649	N/A	59. 7	37. 3	10. 6	0.6
	Mean	3.5 ±0.8	3.1 ±0.8	3.5 ±0.1	N/A	613 ±71	390 ±52	510 ±90	N/A	12130 3	73332	12629 7	N/A	58. 3	35. 6	20. 3	3.5
Broadlea f (tree)	<i>Betula pendula</i>	2.1±1 .3	2.1 ±0.9	2.0 ±0.8	N/A	493±49 8	335±23 4	354±18 5	N/A	15659 1	67692	57934	N/A	58. 3	35. 6	20. 3	3.5
Broadlea f (shrub)	<i>Chamae daphne</i>	2.4±1 .3	2.3 ±1.7	1.8 ±0.7	N/A	633±49 3	347±28 9	571±39 5	N/A	17662 0	73382	21647 3	N/A	61. 8	44. 7	27. 2	4.3
	<i>Oxycocc us</i>	2.1±0 .9	2.3 ±1.4	2.2 ±1.0	1.9± 0.8	730±48 6	590±41 9	410±34 3	260±10 8	22208 1	18419 6	94939	31142	69. 1	52. 8	32. 0	6.1
	<i>Ledum</i>	2.3±1 .0	2.7 ±1.4	2.7 ±1.6	N/A	728±69 8	393±32 2	568±58 6	N/A	28175 1	67913	21978 8	N/A	84. 5	47. 4	31. 9	8.3
	<i>Vacciniu m</i>	2.6±2 .0	2.1 ±0.9	3.8 ±2.8	2.1± 1.4	441±33 4	414±30 9	442±30 4	401±25 0	88832	10554 7	88333	75360	59. 1	43. 4	21. 4	3.9
	Mean	2.3 ±0.1	2.4 ±0.2	2.6 ±0.8	2.0± 0.4	563 ±98	385 ±71	426±10 1	330±10 0	17463 6	87725	10640 8	53251	68. 5	46. 7	28	5.6
Leaf (forb)	<i>Rubus</i>	1.9±0 .7	2.2 ±1.1	2.1 ±1.0	N/A	398±35 5	354±23 7	368±31 5	N/A	99109	79877	88927	N/A	49. 8	40	35. 2	12. 1

	<i>Cnidium</i>	2.3±1 .2	4.9 ±2.6	2.6 ±2.2	N/A	466±38 4	372±26 3	549±59 8	N/A	10861 0	34207	17682 8	N/A	50. 3	41. 1	37. 3	23. 6
	<i>Polypodium</i>	2.1±1 .0	2.9 ±2.0	2.5 ±1.3	N/A	521±39 9	464±29 2	383±28 4	N/A	15556 1	10624 1	60473	N/A	61. 4	50. 6	29. 4	18. 2
	Mean	2.1 ±0.2	3.3 ±1.4	2.4 ±0.3	N/A	466 ±61	418 ±65	466±11 7	N/A	13208 5	70224	11865 0	N/A	56. 6	44. 3	33. 6	17. 6

Table 3. The mean aspect ratio, length, surface area, and the number of each charcoal morphotypes in the Holocene samples ranging from 185 to 6750 cal yr BP (35-303 cm) from Ulukh-Chayakh mire. The local to regional vegetation is represented by the percentages of the main pollen types, whereas the local vegetation is represented by the plant macrofossils (values in brackets presented as numbers except for the wood remains presented as percentages).

705	Depth (cm)	35	84	85	172	248	268	303
	Aspect ratio	3.2	3.0	3.0	4.0	11.1	4.3	2.8
	Length (μm)	742	555	723	488	403	515	431
	Surface (μm^2)	248.287	122.599	219.413	49.673	16.565	53.354	53.698
710	Charcoal morphologies (plant macrofossil)							
	Wood	23 (6%)	13(5%)	6 (5%)	11 (15%)	1 (0%)	8 (0%)	6 (0%)
	Leaf	11 (0)	7 (0)	3 (0)	2 (0)	0 (0)	4 (0)	4 (0)
	Needle	0 (7)	1 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
715	Graminoid	21 (30)	9 (70)	1 (70)	11 (45)	4 (65)	19 (70)	3 (0)
	<i>Equisetum</i>	15 (10)	3(5)	1 (0)	0 (15)	0 (10)	5 (5)	0 (90)
	Moss	10 (30)	0 (5)	0 (5)	1 (5)	1 (5)	3 (5)	1 (0)
	Pollen (%)							
720	Trees	90	92	92	94	97	92	97
	Shrub	1.1	3.4	3.4	3.5	0.2	0.5	0
	Graminoid	5	6.3	6.3	3	13	8.6	9.5
	<i>Equisetum</i>	0.5	3.4	3.4	1.2	6.7	9	18
	Moss	0	0.2	0.2	0	0	0.3	0
725								

Table 4. Comparative results of the aspect ratio from plant species analysed in this study with those from literature. *Pinus sylvestris* wood sums the mean aspect ratio of wood from trunk and twig; wood total sums the mean aspect ratio of wood from trees and shrubs (trunk and twig); broadleaf sums the mean aspect ratio of leaf from trees and shrubs; whereas leaf total averages the mean aspect ratio of all leaf types.

730	Fuel type	250°C	300°C	350°C	400°C	500/550°C	open flame	References
	Graminoid (boreal)	7.3	11.5	6.7	-	-	-	This study
735	Graminoid (artic) -	-	-	-	-	6.7	-	Pereboom et al. (2020)
	Graminoid (forest steppe)	-	-	3.6	-	-	4.8	Umbanhowar & McGrath (1998)
	Graminoid (grass)	-	-	-	-	3.7	-	Crawford & Belcher (2014)
	<i>Pinus sylvestris</i> (wood)	2.7	4.1	2.7	2.9	-	-	This study
740	<i>Pinus sylvestris</i> (wood)	-	-	-	-	2.8	-	Crawford and Belcher (2014)
	Wood (total)	3.4	4.5	3.1	4.0	-	-	This study
	Wood (forest steppe)	-	-	2.1	-	-	2.3	Umbanhowar & McGrath (1998)
	Shrubs (wood and leaf)	-	-	-	-	2.4	-	Pereboom et al. (2020)
	Broadleaf	2.2	2.2	2.3	2.0	-	-	This study
745	Needles	3.5	3.1	3.5	-	-	-	This study
	Leaf (total)	2.5	2.7	2.6	2.0	-	-	This study
	Leaf (forest steppe)	-	-	1.9	-	-	2.1	Umbanhowar & McGrath (1998)

750

Appendix**Appendix A.** The watershed algorithm used to calculate morphometrics of charred particles.

755 The algorithm for automatic detection of morphometrics is based on functions from the Python module skimage (watershed
algorithm). First, the picture is converted to a grey scale. A Sobel gradient of the picture is then calculated, which results in an
elevation map. To use the watershed algorithm to detect the charcoal particles, a map of markers in the grey picture with grey
values higher than 140 was then create. These are the starting points of the watershed region fill algorithm. Finally, any holes
in the watershed regions were filled with the help of a binary fill method (Soille and Vincent, 1990). The detected particles
were subject to the calculation of morphometrics such as surface area and lengths along the major and minor axis via supported
760 functions. Particles with the length of major axis smaller than 150 μm were excluded from these calculations. The pixel area
has been calibrated with a micrometre scale and the results scaled accordingly.

Reference

765 Soille, P., Vincent, L. M.: Determining watersheds in digital pictures via flooding simulations. Proc. SPIE 1360: 240-250.
doi:10.1117/12.24211, 1990.

770