

EThe identification stimation of Evaluation of Phosphorus the natural background of phosphate estimation in the Scheldt a lowland river using tidal marsh sediment cores

Florian Laurysen¹, Philippe Crombé², Tom Maris³, Elliot Van Maldegem², Marijn Van de Broek⁴,
5 Stijn Temmerman³, Erik Smolders¹

¹Division of Soil and Water Management, Department of Earth and Environmental Sciences, KU Leuven, Kasteelpark Arenberg 20 bus 2459, 3001 Leuven, Belgium

²Department of Archaeology, Ghent University, Sint-Pietersnieuwstraat 35, 9000, Ghent, Belgium

10 ³University of Antwerp, Ecosystem Management Research Group, Campus Drie Eiken, D.C.120, Universiteitsplein 1, 2610 Wilrijk, Belgium

⁴Sustainable Agroecosystems group, Department of Environmental Systems Science, Swiss Federal Institute of Technology, ETH Zürich, Zürich, Switzerland

Correspondence to: Florian Laurysen (florian.laurysen@kuleuven.be)

15 **Abstract.** Elevated phosphate (PO₄) concentrations can harm the ecological status in water by eutrophication. In the majority of surface waters in lowland regions such as Flanders (Belgium), the local PO₄ levels exceed the limits defined by environmental policy and fail to decrease, despite decreasing total phosphorus (P) emissions. In order to underpin the definition of currents limits, this study was set up to identify the pre-industrial background PO₄ concentration in surface water of the Scheldt river, a tidal river in Flanders. We used the sedimentary records preserved in tidal marsh sediment cores as an archive
20 for reconstructing historical changes in surface water PO₄. For sediment samples at sequential depths below the sediment surface, we dated the time of sediment deposition and analysed the extractable sediment-P. The resulting time series of sediment-P was linked to the time series of measured surface water-PO₄ concentrations (data 1967-present). By combining those datasets, the sorption characteristics of the sediment could be described using a Langmuir type sorption model. The calibrated sorption model allowed us to estimate a pre-industrial background surface water PO₄ levels, based on deeper
25 sediment-P that stabilised at concentrations smaller than the modern. In three out of the four cores, the sediment-P peaked around 1980, coinciding with the surface water PO₄. The estimated pre-industrial (~1800) background PO₄-concentration in the Scheldt river water was 62 [57; 66 (95%CI)] µg PO₄-P L⁻¹. That concentration exceeds the previously estimated natural background values in Flanders (15-35 µg TP L⁻¹) and is about half of the prevailing limit in the Scheldt river (120 µg PO₄-P L⁻¹). In the 1930s, river water concentrations were estimated at 140 [128; 148] µg PO₄-P L⁻¹, already exceeding the current
30 limit. The method developed here proved useful for reconstructing historical, background PO₄ concentrations of a lowland tidal river. A similar approach can apply to other lowland tidal rivers to provide a scientific basis for local, catchment specific PO₄ backgrounds.

1 Introduction

Elevated Excess phosphorus (P) concentrations in surface waters is a global problem (Azevedo et al., 2015; Dodds and Smith, 2016; Elser et al., 2007). Eutrophication by excess nutrients, including Pphosphate (PO_4), and nitrogen-nitrogen (N) (N), can lead to hypoxia, acidification, and harmful algal blooms (Azevedo et al., 2015; Correll, 1998; Watson et al., 2018). Therefore, limiting P concentrations in the surface water is crucial to ensuring a good ecological status, human health and well-being. LowlandDownstream river systems are at higher risk for nutrient-stimulated eutrophication than upland streams (Watson et al., 2018). As a result, eutrophication of lowland rivers is on the international agenda, and P is considered the limiting nutrient (Jarvie et al., 2006; Mainstone and Parr, 2002; Reynolds, 2000). This study focuses on dissolved orthophosphate (PO_4), almost identical to the reactive P determined by a colour reaction. Other P forms present in surface water include organic P fractions, and P adsorbed to mineral colloids. Total P (TP) refers to all P forms together. The environmental limits for P are either expressed as reactive P (equated to PO_4 -P limits), as TP limits or both.

The lowland rivers of Like Flanders (North of Belgium), densely populated regions many lowland regions (The Netherlands, Germany) do not achieve good water quality mainly due to the excess of nutrients (Bitschofsky and Nausch, 2019; Huet, 1990; Van Der Molen et al., 1998; Van Puijenbroek et al., 2014; Rönspieß et al., 2020; Schulz and Herzog, 2004). For example, since 2004, the average PO_4 concentration in Flemish waterways has stabilised since 2004 at $290 \mu\text{g PO}_4\text{-P L}^{-1}$. That concentration is well above the limits varying between $70\text{-}140 \mu\text{g PO}_4\text{-P L}^{-1}$ for different river types (Smolders et al., 2017; VMM, 2018). Despite the current net-zero P-balance in agricultural soils in that region, the situation did not improve since 2004 (VMM, 2017). The question arises when, or even if these limits can be achieved.

However, the natural background concentrations of lowland river PO_4 -P may be higher than those of upland rivers because of biogeochemical processes typical for such waters. For example, diatom assemblages revealed natural eutrophic conditions in The Spree river in Germany, with total P (TP) concentrations of $80 \mu\text{g L}^{-1}$, compared to recent data of $120 \mu\text{g TP L}^{-1}$ (Schönfelder and Steinberg, 2004; Zak et al., 2006).

Therefore, estimating the background is essential for developing nutrient limits, as it provides a baseline for water quality assessments. It is crucial to ensure the P form when interpreting surface water P concentrations because different P forms occur in surface waters. Phosphorus is either present in solution (dissolved P) or associated with the suspended matter. This study focused on dissolved orthophosphate (PO_4), almost identical to the reactive P determined by a colour reaction. Other P forms present in surface water include organic P fractions, and P adsorbed to mineral colloids. Total P refers to all P forms together. The environmental limits are either expressed as reactive P (equated to PO_4 -P limits), as TP limits or both.

Since 2000, the European Union has regulated surface water quality with the Water Framework Directive 2000/60/EC (WFD), which does not prescribe limits but provides a framework for local regulations. ~~The WFD does not prescribe limits for P in rivers and lakes but provides a framework for local regulations.~~ ~~The WFD has identified a high ecological status of a river if nutrient conditions remain within the range associated with undisturbed conditions, i.e. natural background levels~~ (European Commission, 2000). ~~The WFD does not prescribe limits for P in rivers and lakes but provides a framework for local regulations.~~ ~~The local maximum P concentrations identify pristine environments with minimal anthropogenic disturbance, i.e. the natural background~~ (European Commission, 2000). However, the definition of the natural background has been subject to debate for many river basins (Matschullat et al., 2000; van Raaphorst et al., 2000). ~~Here we define, the~~ ~~The~~ natural background can be defined as the situation (ose concentration range)s found in the environment without any human activity, reflecting only natural geochemical processes (Laane, 1992; Reimann and Garrett, 2005). This definition implies that concentrations ~~have to~~ ~~must~~ be estimated before human activity, which is not always feasible. Therefore, a pre-industrial background can be defined instead, inferred from samples dating before the industrial revolution (Reimann and Garrett, 2005). The pre-industrial background ~~is can~~ logically ~~be~~ affected by anthropogenic processes. ~~Alternatively, natural background concentrations can be estimated by sampling regions with an expected minimal anthropogenic influence, i.e. reference lakes and rivers~~ (Cardoso et al., 2007). ~~Natural background concentrations have been established for different chemical elements in rivers in Europe, however that did not include TP or PO₄-P~~ (Salminen et al., 2005).

~~For P in densely populated regions such as Flanders, Defining the natural background concentrations for P in densely populated regions such as~~ ~~For example, in Belgium, the industrial revolution started around 1800 with three million inhabitants~~ (Vanhaute, 2003). ~~Before that, large scale agriculture dates back to the middle ages and the Roman period. However, the most substantial increase in nutrient emissions occurred after the 1950s due to sewer infrastructure, mineral fertilisers and P-loaded detergents~~ (Billen et al., 2005; van Raaphorst et al., 2000).

~~Flanders can only be inferred indirectly. Natural background TP concentrations for Flanders were set at 15-35 µg P L⁻¹, have been inferred from reference lakes sampled in Central and Baltic states in Europe~~ (Cardoso et al., 2007) ~~and were set at 15-35 µg P L⁻¹~~ (Cardoso et al., 2007). However, the natural background PO₄-P concentrations of lowland rivers ~~PO₄-P may be higher than those of upland rivers. For example, diatom assemblages revealed natural eutrophic conditions in The Spree river in Germany, with TP concentrations of 80 µg L⁻¹, compared to recent data of 120 µg TP L⁻¹~~ (Schönfelder and Steinberg, 2004; Zak et al., 2006).

~~Lowland rivers can be significantly loaded with P derived from the river bed sediment, defined as internal loading~~ (Froelich, 1988). ~~The internal loading of P better better explains temporal and spatial trends in river P concentrations than the P emissions of the lowland river system of Flanders~~ (Smolders et al., 2017). ~~That is explained by biogeochemical processes~~ ~~Biogeochemical processes in the sediment explain that result in the sediment.~~ Ferric iron- (Fe(III)) and aluminium-oxyhydroxides have a high affinity for PO₄-anions and limit the PO₄ in solution (Borgegaard, 1990; Holtan et al., 1988). However, anoxic conditions lead

Formatted: Subscript

Formatted: Not Highlight

100 to the reductive dissolution of those Fe(III) minerals releasing the associated P to the overlying water when the sediment is
strongly reduced (Baken et al., 2015; van Dael et al., 2020). ~~The S~~small rivers and ditches in lowland ~~river~~regions have a low
water velocity and are nutrient-rich, leading to anoxic conditions during summer and fall. ~~Those anoxic events, explaining~~
the typical summer peaks in PO₄ in small rivers. ~~Moreover, -and-~~regional differences in sediment P/Fe concentration ratios
explain regional differences in surface water PO₄ concentrations (Smolders et al., 2017).

105 Sediment analysis is linked to surface water P and can be a valuable tool ~~F~~for the assessment ~~of~~assessing historical river water
quality, ~~sediment analysis is valuable~~. In surface waters, sediments can ~~both~~ serve as a sink or a source of PO₄, depending on
the sediment surface chemistry and water concentrations (Froelich, 1988; House and Denison, 1998; Simpson et al., 2021; van
der Zee et al., 2007). For example, P storage on fine bed sediments can amount to 60% ~~of the total P export~~ of a catchment
110 nutrient budget (Ballantine et al., 2009; Svendsen and Kronvang, 1993). The essential processes for PO₄ are adsorption and
desorption from Fe oxy-hydroxides, present in the suspended matter or bed sediments (Froelich, 1988; van Raaphorst and
Kloosterhuis, 1994; van der Zee et al., 2007). Ferric iron (Fe(III)) and aluminium oxyhydroxides have a high affinity for PO₄-
anions and limit the PO₄ in solution (Borggaard, 1990; Holtan et al., 1988). As a result, the surface water PO₄ concentration
depends on the sorption capacity of Fe oxy-hydroxides, which decreases with increasing pH and salinity (van der Zee et al.,
115 2007). However, anoxic conditions lead to the reductive dissolution of those Fe(III) minerals releasing. As a result, the
associated P is released to the overlying water when the sediment is strongly reduced (Baken et al., 2015). It is now well
established that such reducing conditions explain the typical summer peaks in PO₄ and that regional differences in sediment
Fe concentrations explain regional differences in surface water PO₄ concentrations in Flanders (Smolders et al., 2017a).

120 Several authors in Belgium and the Netherlands described the relation between soil characteristics and pore water PO₄
concentrations. They used soil analysis to identify agricultural areas sensitive to phosphate leaching as the soil P content
showed a good correlation with pore water P concentrations (Breeuwsma et al., 1995; Lookman et al., 1995; Schoumans and
Chardon, 2015; Schoumans and Groenendijk, 2000; van der Zee, 1988). Unfortunately, a similar relation between sediments
deposited by rivers and surface waters has not yet been described. ~~However, s~~Sediments-P concentrations can likely predict
surface water P concentrations ~~and has been relevant for the long term reconstruction of P in the environment~~ because the P
125 adsorption characteristics of sediments have been related to surface water concentrations (Wang et al., 2009; Zhou et al., 2005).
~~In addition, s~~Streams sediment can buffer P concentrations, and by sediment analysis, we can identify if stream sediment acts
as a source or a sink for dissolved P (Jarvie et al., 2005; McDowell, 2015). Thus, the existing literature shows that analysis of
sediment P content could be related to surface water P concentrations. ~~A similar relation to what has been observed in soils~~
and pore water.

130 In addition, sediment P analysis has shown to be relevant for the long term reconstruction of P in the environment. For example,
Boyle et al. (2015) used P profiles from lake sediments in the UK to infer ~~catchment P inputs over the last 10.000 years and~~
~~linked that to~~ the historical evolution in population density ~~over 10.000 years in catchments~~. Similarly, banded iron formations
in deep oceanic waters allowed ~~to infer~~inferring oceanic P concentrations ~~of~~ over two billion years ago (Bjerrum and Canfield,

Field Code Changed

Field Code Changed

2002). Likewise, the sediments deposited by rivers or lakes react with surface water PO_4 and are deposited in regularly flooded areas. Thus, those sediments can serve as an archive for reconstructing historical P emissions trends and provide useful information on historical P concentrations in adjacent water bodies (Birch et al., 2008).

In lowland rivers with tidal influence, like the Scheldt, vegetated tidal marshes develop along the river banks, vegetated tidal marshes develop. Tidal marshes directly adjacent to tidal rivers are regularly flooded during high tides; these flooding events deposit, so river sediments and associated elements like P on the densely vegetated marsh surface are deposited on these densely vegetated marshes (Friedrichs and Perry, 2001; De Swart and Zimmerman, 2009; Temmerman et al., 2004a). As a result, the net accumulation of sediments increases the elevation of tidal marshes increases over time due to their net accumulation of sediments (Temmerman et al., 2003a). Therefore, researchers have used tidal marshes as sediment archives of deposited substances other than P, such as organic carbon (Van de Broek et al., 2019) and silicon (Struyf et al., 2007). However, it remains to be investigated to what extent P concentrations measured in tidal marsh sediment archives can be used to reconstruct historical changes in PO_4 concentrations in the adjacent estuary.

In Flanders, environmental pressure is high, and eutrophication affects most water bodies (European Commission, 2019). The average PO_4 concentration in Flemish waterways stabilised since 2004 at $290 \mu\text{g PO}_4\text{-P L}^{-1}$. That concentration is well above the limits varying between $70\text{--}140 \mu\text{g PO}_4\text{-P L}^{-1}$ for different river types (Smolders et al., 2017; VMM, 2018). However, despite the current net-zero P balance in agricultural soils, the situation did not improve since 2004 (MIRA, 2017). Therefore, the question arises when or even if these limits can be achieved. Surprisingly, the prevailing limits in Flanders lack a scientific basis and are not adapted to the local situation (Amery, 2015; Schneiders, 2007). Flanders is densely populated, and it was considered impossible to locate pristine or reference lakes in the current environment to develop nutrient limits. Instead, natural background TP concentrations for Flanders have been inferred from reference lakes sampled in Central and Baltic states in Europe identified for the WFD (Cardoso et al., 2007). Based on that study, background TP concentrations in lakes of Flanders were set at $15\text{--}35 \mu\text{g P L}^{-1}$, selected based on lakes with representative depth and alkalinity. Until now, no TP or PO_4 natural background has been established for rivers in Europe (Salminen et al., 2005).

This study tested and evaluated a methodology to estimate the pre-industrial background water PO_4 concentrations based on the analysis of tidal marsh sediment. Those sediments, which have been deposited over multiple centuries on the banks of Flanders largest tidal river Flanders largest tidal river banks over multiple centuries. Using our results, sediment analysis and a sorption model we could provide the provided the first estimate of pre-industrial PO_4 levels in the Scheldt in, a large lowland river, the Scheldt. First, we described the tidal marsh sediment sorption characteristics by linking the P concentration in dated of tidal marsh sediments to historical measurements of PO_4 in the Scheldt river water. Those sorption characteristics allowed us to an estimation of historical river water PO_4 concentrations. This estimate was based on an analysis of sediments deposited in the 1800s or before industrialisation. The underlying assumption is that sediment-P remains immobile and that the sediments

depth profile reflects the historical trend of PO₄ in the Scheldt river. Accordingly, we argue that the sediment P-composition in deeper sediment layers of tidal marshes provides an [estimate-archive](#) of the historic PO₄ concentration of the adjacent river.

170 A database containing measurements of the PO₄ concentration in the Scheldt river's surface water (1967-current) verified this assumption. [We-This study hypothesis](#) that the previously estimated natural background P of this major lowland river is larger than [that](#) estimated earlier for lakes (15-35 µg P L⁻¹).

2 Materials and methods

175 2.1 Study area

Freshwater tidal marshes were sampled at four locations along the Scheldt river (Fig. 1, Table S1). The Scheldt estuary is located in northern Belgium and the south-western Netherlands and flows into the North Sea. The river basin of the Scheldt covers a large part of Flanders (71%) and the adjacent region of Northern France; the total catchment area is approximately 22.000 km². The population living in the river basin is about 10 million (Meire et al., 2005). The tidal wave extends from the mouth (Vlissingen) to 160 km upstream near Ghent, where sluices stop the tidal wave. The estuary's freshwater tidal zone reaches from Ghent to Rupelmonde (Fig. 1). This research focused on freshwater tidal marshes, i.e. located in this freshwater tidal zone of the estuary. Brackish waters experience the mixing of seawater, making it difficult to distinguish the anthropogenic sources from seawater influence. Furthermore, saltwater in the North Sea has PO₄ concentrations about a factor ten lower than fresh water in the Scheldt river (Burson et al., 2016). This research was focused on freshwater, lowland river systems and the human influence on the P concentrations, saltwater environments were beyond the scope of this study.

180
185




Figure 1: Map of the Scheldt Estuary, triangles indicate the locations of the sampled tidal marshes, Old1 and Young1 were only 250 m apart, and on the scale of the map, they overlap

190 Sediment accreting in tidal marshes originates from the deposition of riverine suspended matter, including inorganic mineral sediment and organic matter (Callaway et al., 1996). We discriminate between old and young tidal marshes, hereafter referred to as marshes. Old marshes have a higher elevation compared to young marshes. As a general mechanism, young marsh surfaces accumulate sediments quickly and increase their elevation asymptotically up to an equilibrium level, **which is** around the mean high water level (MHWL) (Pethick, 1981; Temmerman et al., 2003a). Temmerman et al. (2003a) defined an old marsh as visible on topographic maps of Ferraris (1774 - 1777), so it was formed before the 19th century (Temmerman et al., 195 2003a). Young marshes in the Scheldt estuary were formed more recently, by the natural establishment of pioneer marsh vegetation on formerly bare tidal mudflats, generally after 1944. During the last decades, the young marshes had a surface elevation below MHWL. **As a result,** ~~young~~ marshes experienced more frequent inundations and **therefore** had larger sediment accretion rates than old marshes. For example, between 1931 and 1951, young marshes accumulated at rates of 1.6 to 3.2 cm yr⁻¹, ~~and during the subsequent period 1955-2002, accumulation was slower at 0.4-1.8 cm yr⁻¹.~~ In contrast, ~~to the young marshes,~~ the elevation of old marshes **is at any time** was very close to the yearly MHWL increase rate of 0.3 to 0.6 cm yr⁻¹ in the Western Scheldt (Temmerman et al., 2003a).

This study analysed depth profiles of sediment cores originating from tidal marshes along the freshwater Scheldt river. The analysis contained two old and two young marshes, ~~referred to as~~ Old1, Old2, Young1 and Young2 (locations indicated in Fig. 1). The coordinates of sampling locations can be found in (Van de Broek et al., 2018; Van De Broek et al., 2016) and supplementary information (SI.I). Marshes Old1 and Young1 originated from the Tidal marsh named the Notelaer, Old 2 from Grembergen and Young2 from Mariekerke. ~~In total, we analysed eight cores; three replicate cores for both sites Old1 and Young 1 and one core each~~ Eight cores were analysed; three replicate cores for both sites Old1 and Young 1 and one core for Old2 and Young2.

2.2 PO₄ concentration in surface waters

The ~~IMIS~~ (Flanders Marine Institute) ~~(IMIS)~~ provided surface water phosphate (PO₄) data measured colourimetrically on a filtered water sample and total phosphorus (TP) by acid digestion and a segmented flow analyser. ~~Concentration D~~ data of PO₄ ~~concentrations~~ in Scheldt river were available from 1967 ~~up to~~ 2019, ~~originating from different sources and~~ compiled by the ~~Research program environmental effects Sigmaplan~~ ~~(OMES)~~ ~~monitoring~~. ~~The OMES program~~ ~~who~~ did additional quality controls on the data (ECOBE - UA; The Flemish Waterway, 2019). ~~The PO₄ data originated from~~ different sources described in the supplementary information (SI.V) (De Pauw, 2007; ECOBE - UAntwerpen, 2007; Institute for Hygiene en Epidemiology (IHE), 2007; *OMES: Monitoring physical-chemical water quality in the Zeeschelde*, 2016; Van Meel, 1958).

The open-source software R (R Core Team, 2020) was used to compile all available datasets for PO₄ closest to the study sites (Temse) and to calculate annual means by averaging all observations within ~~each a year for which data was available~~. The annual means of PO₄ were used to visualise the evolution of PO₄ in the Scheldt river (Fig. 2). ~~The~~ ~~Because~~ ~~P~~ ~~emissions~~ ~~of~~ ~~P~~ mainly originate from point sources due to domestic loading ~~(Billen et al., 2005)~~. ~~As a result,~~ the ~~increasing~~ surface water P concentration ~~increased~~ ~~between~~ 1950 and 1975 ~~can be~~ related to the rise in the number of households connected to sewer systems. ~~At first,~~ ~~without~~ ~~no~~ ~~appropriate~~ wastewater treatment ~~being~~ ~~was~~ in place, ~~(Billen et al., 2005)~~ ~~resulting in a sharp~~ ~~increase in nutrient loads to the river~~. ~~However,~~ ~~s-~~ Since 1985, wastewater treatment has significantly improved the situation (Billen et al., 2005).




Figure 2. Concentrations of phosphate in the Scheldt River at Temse, annual means and standard deviation (error bar) around the annual mean. Samples were taken in Temse close to tidal marsh sites. (data sources: (ECOBE-UA and De Vlaamse Waterweg, 2016; ECOBE - UA Antwerpen, 2007; Institute voor Hygiëne en Epidemiologie (IHE), 2007; Van Meel, 1958; De Pauw, 2007)

2.3 Sediment sampling

The sediment samples used here had been collected during a previous study about carbon sequestration in tidal marsh sediments in the Scheldt estuary (Van de Broek et al., 2018; Van De Broek et al., 2016). Collection of undisturbed sediment profiles on the tidal marshes took place between July and September 2016 (Old1, Young1, Old2, Young2; Fig. 1). Undisturbed sediment cores were taken using a gauge auger (0.06 m diameter) at each sampling location. Undisturbed sediment cores were taken at each sampling location using a gauge auger (0.06 m diameter). In the field, the cores were divided into subsamples. The cores were divided into subsamples with a 0.03 m interval. The sediment samples were dried at a maximum temperature of 50 °C for 48 hours, crushed and sieved to a <2 mm grain size. Macroscopic vegetation residues were removed manually using tweezers (Van de Broek et al., 2018). Bulk density, grain size distribution and organic carbon (OC) content were analysed by Van de Broek et al. (2018). We refer to Van de Broek et al. (2016, 2018) for further information about sample collection and processing. sample collection and processing information.

2.4 Sediment analysis

The dried sediment samples were analysed for oxalate-extractable P, Fe, Al and Mn (P_{ox} , Fe_{ox} , Al_{ox} , Mn_{ox} ; Schwertmann, 1964). The preparation of extraction solution and dilutions were made with ultrapure water (Milli-Q®), and all glassware used was acid-soaked overnight in a 1% HCl acid bath to prevent P contamination. That acid oxalate extractant, a mixture of ammonium oxalate (0.2 M) and oxalic acid (0.2 M) at pH = 3, targets poorly crystalline oxyhydroxides of Fe, Al and Mn and the associated P (Schwertmann, 1964). Those P-poorly crystalline oxyhydroxides are the most reactive due to their large specific surface area (Hiemstra et al., 2010). The extraction was done with 1 g of dry sediment in 50 ml extraction solution over two hours in an end-over-end shaker at 20°C (26 rpm). The suspension was filtered through a 0.45 µm membrane filter (CHROMAFIL® Xtra PET - 45/25). Analytical blanks, internal reference samples, and duplicate samples were included in every batch to ensure the analysis's quality, purity, and reproducibility. The extracts were diluted 20 times and measured by inductively coupled plasma optical emission spectrometry (ICP-OES). The degree of P-saturation (DPS; %) was calculated as in Eq. (1). The DPS represents the ratio of extractable (P_{ox}) to the P sorption capacity of the sediment. This P sorption capacity is estimated as half of the sum of oxalate Fe_{ox} and Al_{ox} , because not all the Fe and Al in the soil is available for P sorption with Fe_{ox} , Al_{ox} and P_{ox} in molar units.

$$DPS = \frac{P_{ox}}{0.5(Fe_{ox} + Al_{ox})} 100\%, \quad (1)$$

The DPS is expressed in percentage and can be interpreted as the ratio of sorption sites on the sediment occupied by P. Previous research used the DPS to identify agricultural areas sensitive to phosphate leaching and showed a good correlation with pore water P concentrations (Breeuwsma et al., 1995; Lookman et al., 1995; Schoumans and Chardon, 2015; Schoumans and Groenendijk, 2000; van der Zee, 1988). The DPS relation was verified for porewater-soil systems and was developed and verified by (van der Zee et al., 1990). Lexmond et al. (1982) illustrated that the maximal sorbed P was about half the pool available after a long-term precipitation experiment. Therefore, the factor 0.5 is an empirical value representing the soil's sorption capacity, an empirical value based on pore water measurements of non-calcareous sandy soils and is considered the sorption capacity of the soil. Lexmond et al. (1982) illustrated that the maximal sorbed P was about half the pool available after a long-term precipitation experiment. The parameter α primarily affects the maximum sorption capacity. So they set α at 0.5 ± 0.1 . However, even among soils, this parameter varied between 0.3 and 0.6 (Lexmond et al., 1982). We are interested for this research, in low background concentrations for this research are most important, so maximal sorption, occurring at high PO_4 concentrations, is less relevant.

2.5 Age-depth model

The sediment analysis and the surface water PO_4 data had to be linked by a corresponding date and location to fit a sorption model. Therefore, an age-depth model was used to calculate the time since deposition of each sediment sample. Temmerman et al. (2004b, 2004a) developed a time-stepping marsh sedimentation model (MARSED). That model estimates

275 sediment deposition rates and the resulting evolution of the ~~sediment surface~~tidal marsh elevation ~~in the tidal marshes of in~~ the Scheldt estuary. Hence, we could ~~also use that model~~MARSED to determine the time since deposition of sediments throughout the sampled sediment profiles. The MARSED model simulates the tidal supply of suspended sediments and the settling ~~of these sediments~~ to the marsh surface during tidal inundation cycles integrated over the years. The model was calibrated and validated against measured ~~data on~~ sediment deposition rates on the Scheldt estuary tidal marshes from 1945 until 2002 (Temmerman et al. 2003; 2004). The empirical data on sediment deposition rates were derived from radiometric and paleoenvironmental dating of sediment cores at the ~~same exact~~ locations sampled for the present study (Temmerman et al., 280 2004a, 2004b).

For our current study, ~~we extrapolated we extrapolated~~ the MARSED model simulations of sediment accretion from 2002 until 2016, the sampling date of the sediment cores (Van de Broek et al. 2018). ~~However, We found that the MARSED simulations model overestimated the observed marsh surface elevation in 2016 (observed by RTK GPS surveying; Van de Broek et al. 2018; Poppelmonde, 2017) by 25 cm for sampling location Old1, 29 cm for Young1, 19 cm for Old2, and 8 cm for Young2~~ 285 ~~(observed by RTK GPS surveying; Van de Broek et al. 2018; Poppelmonde, 2017).~~ The MARSED model was initially designed to simulate the overall sediment accretion and surface elevation changes in tidal marshes in response to sea-level rise scenarios, for which those errors were acceptable. ~~However~~In contrast, ~~the most important was the time of sediment deposition throughout the sediment profile for the present study or the present study, we wanted to estimate the time of sediment deposition throughout the sediment profiles as good as possible.~~ Therefore, the original age-depth relation calculated by MARSED was recalibrated ~~by using~~ observed age-depth points. The observed age-depth points originated ~~d~~ from GPS measurements of marsh elevation in 2016 (M. Van de Broek, unpublished data) and previously published radiometric and paleoenvironmental dating (Temmerman et al. 2003; 2004). This rescaling procedure is explained in the Supporting Information (Fig. S1, S2, S3, S4). ~~The observed age depth points were available starting from 1958 for sampling site Old1, 1947 for Young1, 1963 for Old2 and 1968 for Young2 (Temmerman et al. 2004).~~ 290

295 An approximate extrapolation procedure was used to estimate the sediment deposition time from depths below the oldest measured age-depth points (mentioned in the previous sentence). ~~The observed age-depth points were available from 1958 for sampling site Old1, 1947 for Young1, 1963 for Old2 and 1968 for Young2 (Temmerman et al. 2004).~~ ~~This~~This extrapolation procedure ~~was could~~ only ~~be~~ applied for old marshes, which were defined as marshes that existed at least since the end of the 18th century (Temmerman et al. 2003a; 2004). Two sediment cores originated from ~~such~~ old tidal marshes (Old1 and Old2). 300 ~~We know from~~Based on observed age-depth points, ~~it has been proven~~ that ~~the~~ old marshes reached equilibrium with the MHWL before 1944. ~~After 1944, and that they~~Old marshes ~~have~~ built up their elevation ~~after 1944~~ at a rate comparable to ~~the rate of~~ local MHWL rise (Temmerman et al., 2003a). Here, we assumed that ~~also~~ between 1800 and 1944, these old marshes ~~also~~ accreted at a rate comparable to the MHWL rise.

305 Historical tide gauge data of MHWL rise was available from 1901 for site Old1 and 1930 for site Old2 (ScheldeMonitor Team and VNSC, 2020; Temmerman et al., 2003a) and linear regression of the MHWL against time was used to estimate the marsh surface elevation before 1944 (Fig. S6, S7). However, the ~~accuracy of the dating~~dating accuracy will be lower going further

back in time. ~~Such~~ Furthermore, such extrapolation to earlier dates is not appropriate for young marshes, as they were only formed after 1950 by pioneer vegetation establishment on formerly bare mudflats (Temmerman et al. 2003a; 2004). Those mudflat sediment profiles do not have continuous sedimentary records as tidal marshes and are likely to be disturbed by erosion and sedimentation alternations (Belliard et al., 2019). Therefore, the sediment deposition time could not be extrapolated for the young marsh sampling locations ~~before the oldest available measured age-depth points, dating back to 1947 for site Young1 and 1968 for site Young2.~~

2.6 Relating surface water PO₄ with sediment P: the sediment-water model

The age-depth model and linear regression of MHWL provided a deposition year for each sediment sample. ~~Through that age-depth relation~~ Thereby, the dataset of water PO₄ between 1967-2016 was linked to the sediment DPS for each core. The resulting dataset contained all available surface water PO₄ readings between 1967 and 2016, closest to the tidal marshes in Temse (n = 1932) and a corresponding DPS value. ~~The DPS value~~ of a sediment sample originates from ~~a specific layer of~~ one sediment core or, ~~when available for Old1 and Young1,~~ a mean DPS of the replicate sediment samples. This dataset allowed to fit a sorption model further termed the sediment-water model. Schoumans and Groenendijk (2000) presented a Langmuir-type sorption model to predict PO₄ concentration leaching from a soil layer based on the DPS Eq. (2).

$$[PO_4] = K^{-1} \frac{DPS}{100 - DPS}, \quad (2)$$

With [PO₄] phosphate concentration in (kg L⁻¹), K the sorption constant (L kg⁻¹), DPS (degree of P-saturation; %). This model adequately described P sorption in soil across a wide range of pH values, including the Scheldt river pH (Schoumans and Groenendijk, 2000; Warrinnier et al., 2018). The model relies on surface complexation between PO₄ and Fe-, Al-oxyhydroxides in the sediment. ~~That complexation is,~~ determined by a chemical equilibrium between solid (adsorbed) and dissolved PO₄ phase (Warrinnier et al., 2019). The parameter K of existing soil models has been calibrated for soil - pore water system, and the sediment-water parameter (K) is unlikely equal. Therefore, the model was calibrated by fitting parameter K (Eq. 2) on sediment DPS measurements and recent Scheldt water PO₄ measurements. ~~The parameter K (Eq. 2) was calibrated to the most recent Scheldt water data.~~ As a result, the fitted K-value is adapted to the local geochemistry of tidal marsh sediments and the surface water.

We explored 16 different scenarios to fit the sediment-water model Eq. (2). These scenarios illustrate the statistical uncertainty surrounding the estimated PO₄ concentrations. The model was fitted ~~separately for each site sediment core separately for each site sediment core~~ or on the combined replicate cores for Old1 and Young1 (SI.VI). Every sediment sample had between one and three replicates, depending on the depth and the site. ~~We entered either the~~ The average value of these replicates ~~was used~~ or the individual replicates DPS values. One sediment sample covered several deposition years, so multiple PO₄ observations corresponded with each sediment sample. Again, the average of all corresponding PO₄ readings was taken, or all available values were used separately. ~~The combination of mean or individual DPS and PO₄, resulting in~~ 16 models (Table S2). For each of these, the parameter ~~K was fitted estimation of Eq. (2) was fitted~~ by non-linear ~~least squares~~ regression with ~~R using~~

Rstudio interface (R Core Team, 2020; RStudio Team, 2015). ~~Non-linear least-squares regression to the PO₄-DPS data was used to estimate the model parameter (K), yielding the lowest sum of squared errors.~~

2.7 Evaluation Model Performance

The predictions of the sediment-water model were evaluated based on several parameters; the Residual Standard Error (RSE), the Nash Sutcliffe Model Efficiency (E) and by plotting the measured surface water PO₄ against predicted PO₄ between 2007 and 2016 (Table S2; Fig. S10). Additionally, the ~~percentage~~ bias (Pbias) was calculated for data points between 2007 and 2016. The Pbias measures the average tendency of the simulated data to be larger or smaller than their observed counterparts. That difference is expressed as a percentage of deviation from the observations (Moriassi et al., 1983; Eq. 3). The predictions of recent years is interesting to evaluate the model's performance ~~because of~~ two reasons. First, the most recent surface water PO₄-concentrations are relatively low and more representative of background concentrations. Second, the monitoring data have a high temporal resolution, and the age-depth model is more accurate at shallow depths.

$$PBias = \frac{\sum_{i=1}^n (y_i^{obs} - y_i^{sim})}{\sum_{i=1}^n y_i^{obs}} \quad (3)$$

3 Results

3.1 History of surface water PO₄ concentrations

The Scheldt PO₄-concentrations varied greatly over the past decades, with the peak in surface water PO₄-concentrations between 1975 and 1985 (Fig. 2). In Temse, the annual mean concentrations rose from 410 µg PO₄-P L⁻¹ in 1967 and peaked in 1980 with 1570 µg PO₄-P L⁻¹. Between 1990 and 2003, ~~a decrease followed the peak and over the last decade,~~ concentrations ~~decreased and~~ stabilised between 160 and 200 µg PO₄-P L⁻¹ in Temse. ~~The C~~current PO₄-levels are a factor two lower than in 1967 and almost a factor ten lower than the peak in 1980 (Fig. 2; Table 1).

3.2 Sediment cores

The P_{ox} in the sediments ranged between 370 mg P kg⁻¹ and 13,000 mg P kg⁻¹, while the DPS ranged between 13% and 94% (Table 1). In all soil cores starting at the surface, the DPS and P_{ox} increased with depth and peaked at about 0.5 m below the surface (Fig. S7, Fig. S8). In deeper (>1.0 m) sediment layers, P_{ox} and DPS decreased and stabilised for Old1, Young1 and Young2 (Table 1). Overall, the P_{ox} increased by ~~on average, a an average~~ factor of 3.5 between the surface and the maximum concentrations (Fig. S8, Table1). ~~The sediments with The~~these peak DPS were deposited between 1960 and 1985 in three of the four sediment cores (Fig. 2). Only the core Old2 ~~reached the peakpeaked~~ earlier (ca. 1940-1950). Most importantly for this work, DPS for Old1 showed an apparent stabilisation in deeper or older layers, which indicated undisturbed sediment layers (Fig. 3, Fig. S7).

Table 1: The sediment oxalate extractable P (P_{ox}) and its Degree of Phosphate Saturation (DPS) of the top, bottom, peak sediment layers at four different tidal marsh locations. Top layers are the sediments closest to the surface, peak layers had maximal P_{ox} and DPS, and bottom layers are those sediments sampled at the largest depth. Values of P_{ox} and DPS are means (\pm standard deviation) of N sediment samples, between top and bottom (cm) depth.

Location		N	Top – Bottom (cm)	P_{ox} (mg kg ⁻¹)	Fe_{ox} (mg kg ⁻¹)	Al_{ox} (mg kg ⁻¹)	DPS (%)
Old1	Top	4	0 - 9	2300 (\pm 2400)	21000 (\pm 1400)	1200 (\pm 130)	36 (\pm 3)
	Peak	7	27 - 57	5400 (\pm 1300)	24000 (\pm 4700)	1800 (\pm 300)	70 (\pm 8)
	Bottom	8	147 - 180	540 (\pm 110)	8500 (\pm 750)	660 (\pm 41)	20 (\pm 4)
Young1	Top	4	0 - 9	2700 (\pm 320)	24000 \pm 1200	1400 (\pm 91)	37 (\pm 2)
	Peak	6	27 - 57	8500 (\pm 3200)	31000 (\pm 7500)	2000 (\pm 120)	85 (\pm 15)
	Bottom	6	129 - 144	910 (\pm 440)	7000 (\pm 3900)	650 (\pm 120)	40 (\pm 4)
Old2	Top	3	0 - 9	2800 (\pm 90)	19000 (\pm 230)	1700 (\pm 36)	45 (\pm 1)
	Peak	3	54 - 69	8000 (\pm 1600)	25000 (\pm 6500)	2600 (\pm 360)	94 (\pm 2)
	Bottom	3	132 - 147	1700 (\pm 620)	11000 (\pm 3900)	1700 (\pm 300)	43 (\pm 6)
Young2	Top	3	0 - 9	2700 (\pm 410)	39000 (\pm 7800)	1300 (\pm 240)	23 (\pm 1)
	Peak	3	48 - 63	7000 (\pm 1200)	42000 (\pm 5400)	1900 (\pm 57)	55 (\pm 7)
	Bottom	3	144 - 183	3200 (\pm 110)	34000 (\pm 770)	1400 (\pm 78)	31 (\pm 2)

Within the first meter, Fe_{ox} was stable in the three soil cores (Old1, Young1, Old2) with concentrations around $20,000 \text{ mg kg}^{-1}$, except for Young2 for which Fe_{ox} was a factor two larger (Fig. S9). For Young1 and Young2, a decrease in Fe_{ox} concentration occurred. Fe_{ox} concentration decreased at depths $> 1 \text{ m}$. For Old1, Fe_{ox} showed a steady decline from $20,000 \text{ mg kg}^{-1}$ at the surface to $10,000 \text{ mg kg}^{-1}$ at the bottom of the profile (Fig. S9). The Al_{ox} concentrations showed a similar trend as the P_{ox} concentrations, with an initial increase followed by a decrease with depth. The strong correlations of Al_{ox} and Fe_{ox} with P_{ox} ($r_{Al} = 0.73$ and $r_{Fe} = 0.65$) illustrate the positive effect of Fe and Al oxyhydroxides on P sorption.

Formatted: Subscript




Figure 3: The Degree of Phosphate Saturation (DPS) timeline based on four tidal marsh sediment sites. Each dot represents a sediment analysis. The year assigned to each sediment analysis was calculated with the age-depth model. Before 1930, no model dates were available. Therefore a linear regression of the MHWL was used to extrapolate the dates for the old marshes. Dates before 1930 are increasingly uncertain going further back in time. For young marshes, such extrapolation was not possible. The points before the formation of the marshes are indicated with a question mark.

390

3.3 Sediment core selection

Under the assumption that PO_4 does not migrate, the tidal marsh sediment cores can provide an archive for river water PO_4 . ~~However, that assumption may be most violated at two locations, Old2 and Young2.~~ Considering P-migration, ~~it is crucial to evaluate~~ the distance from a creek within the tidal marsh ~~is crucial~~. That distance is essential for two reasons. First, within 10 to 20 m of the creeks, the groundwater table fluctuates largely with the tides (Van Putte et al., 2020), which can induce vertical P-migration (Van Putte et al., 2020). Secondly, sediment accretion is more difficult to predict at closer distances to the creeks and can affect the age-depth relation (Temmerman et al., 2003b). ~~The distance from the sediment cores to the nearest creek for the tidal marshes is was within~~ 21 m for Old1, 56 m for Young1, 35 m for Young2 and 5 m for Old2.

~~The assumption that PO_4 does not migrate may be most violated at Old2 and Young2. Furthermore, the~~ The profile of Old2 ~~indicated P-migration because it~~ had a peak of P_{ox} at an earlier date (1950) than was expected from surface water data (1980), ~~indicating P migration~~ (Fig. 2). Consequently, Old2 was not taken up ~~for interpretation of to interpret~~ the relation between DPS and PO_4 . For core Young2, deeper sediment layers had a larger DPS than the surface layers (Table 2). Additionally, the age estimation of sediments older than 1968 was ~~not impossible~~ due to this tidal marsh's young age, ~~which hampers the interpretation of DPS values from deeper layers as representative for background values.~~ Furthermore, Fe_{ox} concentrations were a factor two larger than the other cores (Fig. S9) and a factor two larger than the average sediment Fe-concentration of the Upper-Scheldt basin (VMM, 2019). ~~Finally, the~~ The local enrichment in iron lowers the DPS values and makes the core less representative of the average situation in the Scheldt. These observations made Young2 inappropriate to fit the relation between DPS and PO_4 .

The two remaining soil cores, Old1 and Young1, originated from the same tidal marsh area named "The Notelaer", located near the city of Temse (Fig. 1). ~~That marsh and~~ has been the topic of multiple studies on sediment accretion (Temmerman et al., 2004b, 2003a) and soil OC stocks (Van de Broek et al., 2018; Van De Broek et al., 2016). The sediment profiles of both sites Old1 and Young1 rise and fall in DPS comparable to dynamics in surface water PO_4 -concentrations (Fig. 2, Fig. 3). In deeper sediment layers, DPS and P_{ox} stabilise below levels of recent deposits (Fig. S7, S8). The time series of Old1 displayed a DPS peak around 1960, indicating a shift of 20 years (Figure 2-). However, the core Old1 was taken up ~~for the prediction of~~ for the model fitting ~~because it. That core is essential to predict the background because the core~~ dates back to 1800 at the deepest levels ~~and is essential to predict the background~~. Furthermore, the DPS concentrations stabilised before 1920, indicating that P has not migrated to these depths, making it suitable for background prediction. These observations suggested a well-preserved P_{ox} and DPS profile, essential for the DPS- PO_4 relation. Therefore, Old1 and Young1 are considered the best profiles for applying the sediment-water model and interpretation of background concentrations.

420

Table 2: The predicted concentrations of phosphate ($\text{PO}_4\text{-P } \mu\text{g L}^{-1}$) in the Scheldt river based on the Degree of Phosphate Saturation (DPS) in the sediment layers of marsh Old1, dating back to 1800 (pre-industrial), where DPS values stabilised with depth at 20%. The predicted concentration dated to 1930, where DPS stabilised at 36%. The Pbias is the mean difference of the simulated data and the observed data between 2007 and 2016, expressed as a percentage of the observed data. The conversion of DPS to river phosphate concentration is based on the association of DPS with $\text{PO}_4\text{-P}$. That association was calibrated to data 1967-2016, thereby using different calibrations for sediment-water models; the details of models are in Table S2. Model 3b (in bold), is proposed as the most accurate one (see text).

Model #	K (L kg^{-1}) [95% CI]	pre-industrial background		Pbias (2007 -2016)
		$\mu\text{g PO}_4\text{-P L}^{-1}$ [95% CI]	$\mu\text{g PO}_4\text{-P L}^{-1}$ [95% CI]	
1b	2.1×10^6 [2.0×10^6 ; 2.3×10^6]	120 [110; 130]	270 [245; 281]	62
2b	4.9×10^6 [4.6×10^6 ; 5.2×10^6]	51 [49; 54]	120 [109; 122]	-28
3b	4.1×10^6 [3.8×10^6; 4.4×10^6]	62 [57; 66]	140 [128; 148]	-15

3.4 Sediment-water model fit

The sediment-water model Eq. (2) was fitted on DPS- PO_4 data from the different sediment cores (Table S2). Two observations were omitted because the DPS values were too large (0.99 – 1.02) and produced artefacts in the results. The Nash-Sutcliffe model efficiency (E) ranged between 0.04 and 0.85 depending on the input data (Table S2; Nash and Sutcliffe, 1970). The sediment-water model was fitted on each core's data separately and for the combination of combined the data from Old1 and Young1, as they came from the same tidal marsh location. The models fitted on data from sites Old2 and Young2 were not considered as migration likely affected those cores (crf. section 3.3).


The models fitted on an average DPS (across replicates) associated with individual PO_4 readings were considered most suitable (Models 1b, 2b, 3b; Table S2). A single sediment sample analysis represents an average P signal over the sediment's deposition period. However, due to variation in the marsh surface elevation, the age-depth relation can vary slightly the age-depth relation can vary slightly due to the marsh surface elevation variation. By taking an average DPS from replicate cores, we reduced the variation in the independent variable was reduced to predict PO_4 in water. Furthermore, in most models, the prediction error increased the prediction error increased in most models by relating individual rather than mean DPS values with individual PO_4 measurements (Table S2). Models using unique DPS associated with single PO_4 -data duplicated or even triplicated the PO_4 -data, artificially creating more degrees of freedom (Model 1c, 2c, 3c; Table S2). Using mean PO_4 -values artificially reduced the degrees of freedom, compromising the model predictions by increasing RSE and widening confidence intervals (Models

445 1a, 2a, 3a; Table S2). The fitted parameter K (L kg^{-1}) ranged between 1.0×10^6 and 5.4×10^6 for the different input datasets, with the 95% confidence intervals ranging between 0.8×10^6 and 7.2×10^6 . The variation of parameter K for the various input datasets was larger than the individual confidence limits variation (Table S2). Thus, the uncertainty was more pronounced due to the variability in sediment samples than due to the model fit.

3.5 Model performance

450 The sediment-water model performance was evaluated with several parameters (RSE, Pbias, and E), ~~and by comparing~~ the actual by predicted PO_4 concentrations over the last decade. ~~Those recent~~ ~~those~~ PO_4 concentrations are more comparable to the background (Fig. S10; Table 2; mean Temse [2007-2016] = $170 \mu\text{g PO}_4\text{-PL}^{-1}$). Model 3b was considered the most suitable for predicting background concentrations. The Pbias was the lowest for recent observations for model 3b. The average tendency of simulated data compared to the observations was only -14.9 %, which is within the acceptable range of $\pm 25\%$ (Moriassi et al., 1983). Model 2b had an underestimation of observed data of more than 28%, and model 1b overestimated recent observations by more than 60%. ~~Such an overestimation~~ ~~which~~ is unwanted for calculating the background, ~~and were~~ ~~therefore~~, both ~~were~~ considered unsuitable (Table 2). The actual by predicted plots illustrate a similar message (Fig. S10). Based on these observations, model 3b was considered the best model, although the residual standard error (RSE) was lower for model 2b (Table S2). The selected model 3b successfully reconstructs the rise and fall in surface water PO_4 -concentrations based on the sediment characteristic DPS (Fig. 4).

465 Maxima of monitored and predicted PO_4 -concentrations coincide in time and have a similar size (Fig. 4). For example, in 1973, the average PO_4 concentration predicted by the model was $1200 \mu\text{g PO}_4\text{-P } \mu\text{g L}^{-1}$ and measured concentrations was on average $1300 \mu\text{g PO}_4\text{-P } \mu\text{g L}^{-1}$. The maximal predicted PO_4 concentration was $2200 \mu\text{g PO}_4\text{-PL}^{-1}$, while the maximal observed was $3000 \mu\text{g PO}_4\text{-PL}^{-1}$. Predictions for recent years are within 15% of the observed data (e.g. 2015: Model: $133 \mu\text{g PO}_4\text{-PL}^{-1}$, Measured $155 \mu\text{g PO}_4\text{-PL}^{-1}$). Between 1940 and 1990, the modelled PO_4 -concentrations ~~show~~ ~~had~~ more variation. Likewise, monitored PO_4 -data are spread more between 1967 and 1990 (Fig. 2). Before 1930, modelled PO_4 -concentrations stabilised at levels below current observations (Fig. 4).



470

Figure 4: Measured (grey crosses; +) and predicted (black points) of PO₄-P concentrations (µg L⁻¹) in the Scheldt river in Temse. The concentrations are calculated from the sediment phosphate saturation (DPS) of the tidal marshes at Old1 and Young1, using sediment-water model 3b.

3.6 Estimating background PO₄ concentrations in the Scheldt river

475

The deepest sediment layers are most suitable for predicting background PO₄ concentrations of the Scheldt river water. These layers are the oldest and expected to have experienced the lowest impact of P additions from anthropogenic sources. The Old1 marsh site was appropriate for this purpose as it developed before 1774, before the industrial revolution in Belgium. The average DPS for the bottom sediments, dated between 1800 and 1840, was 20% for core Old1 (Table 1; Fig. 2); these samples are considered to represent the pre-industrial background. That DPS value produced PO₄-concentrations of 62 µg PO₄-P L⁻¹ [95%CI (57; 66)] for the pre-industrial background, using sediment-water model 3b (Table 2). The sediment dated to 1930 had a DPS of 36%. For that value, the same sediment-water model predicted a PO₄-concentration of 140 µg PO₄-P L⁻¹ [95%CI (128; 148)] (Table 2).

480

4 Discussion

4.1 Pre-industrial background vs ambient PO₄ concentrations.

485 This work presents a novel approach to reconstruct background surface water PO₄ concentration in a tidal river using the DPS of adjacent tidal marsh sediments. The background concentration is essential in the context of developing local nutrient limits. The predicted pre-industrial background concentration (62 µg PO₄-P L⁻¹; Table 2) is about half of the current surface limit of the Scheldt (120 µg PO₄-P L⁻¹; Flemish Government, 1995). Remarkably, the predicted background concentrations are about a factor two larger than the background estimates of lake waters for Flanders today (15-35 µg PO₄-P L⁻¹; Cardoso et al., 2007).

490 ~~Our analysis suggests that~~ That the pre-industrial PO₄ concentration was is about three times lower than the current concentrations in the Scheldt. For example, between 2007 and 2016, the mean PO₄ concentration of the Scheldt in Temse was 170 µg PO₄-P L⁻¹. However, in the 1930s, the concentration was estimated at 140 µg PO₄-PL⁻¹ and larger than current limits, at a time before widespread connection to sewer systems, P-loaded detergents, and application of mineral fertilisers.

495 ~~Those results, suggesting that the sediment internal loading related to summer anoxia in lowland rivers contributes naturally triggered by summer anoxia in lowland rivers contributes~~ to larger PO₄ concentrations ~~than estimated before in lowland rivers~~ (see introduction). The summer ~~PO₄ peak of PO₄ concentration is present for last about five months of the year in Flanders, and can thereby have and largely affect the~~ a significant influence on the mean P concentrations in the rivers ~~about five months per year in Flanders and largely affects the rivers' mean P concentrations~~ (Smolders et al., 2017). ~~Summer anoxia can occur in eutrophic lakes or sometimes in oligotrophic brown water lakes (Nürnberg, 1995). For example, in 75 lakes in the US, anoxia occurred between zero and 83 days a year, which is less than the five months or 150 days in Flemish rivers (Nürnberg, 1995; Smolders et al., 2017b).~~

500 ~~Also~~ Additionally, lowland rivers in Flanders are primarily groundwater-fed, ~~and, as on average,~~ 73% of streamflow can be attributed to base flow. ~~The and n.~~ Natural groundwater in Belgium has a median P concentration between 150 - 320 µg P L⁻¹ (Edmunds and Shand, 2009). ~~Groundwater. Therefore, groundwater~~ feeding the river waters ~~will increase the P levels logically affects the river P concentrations.~~ In contrast, ~~mostly primarily rain-rain-~~ fed lakes will have lower P concentrations, with rain P ranging between 1.5 and 120 µg P L⁻¹ (Migon and Sandroni, 1999).

505 ~~Summer anoxia also occurs in eutrophic lakes, or sometimes oligotrophic brown water lakes (Nürnberg, 1995). For example, in 75 lakes in the US, anoxia occurred between zero and 83 days a year, which is less than the five months or 150 days in Flemish rivers (Nürnberg, 1995; Smolders et al., 2017b).~~

510 ~~Our analysis suggests that the pre-industrial PO₄ concentration was about three times lower than the current concentrations in the Scheldt. For example, between 2007 and 2016, the mean PO₄ concentration of the Scheldt in Temse was 170 µg PO₄-PL⁻¹. However, in the 1930s, the concentration was estimated at 140 µg PO₄-PL⁻¹ and larger than current limits, at a time before widespread connection to sewer systems, P-loaded detergents, and application of mineral fertilisers.~~

515 4.2 Limitations of the model

Care needs to be taken with background extrapolations to ensure that post-depositional processes have not modified the biogeochemical patterns and that the area represents the area of interest (Reimann and Garrett, 2005). Several factors can obscure the reconstructed background concentrations. First, vertical migration of P can enrich deeper sediment layers, causing an overestimation of the background. Second, the sediment profiles at the tidal marshes are almost permanently saturated, ~~and~~ ~~so~~ the intrusion of P-rich groundwater could affect the P concentrations in the tidal marsh sediment. Moreover, ~~depending on the tidal marsh elevation,~~ periodic flooding occurs at an approximate range of 300-350 inundations per year, ~~depending on the tidal marsh elevation~~ (Temmerman et al., 2003b). These conditions could favour ~~phosphorus-P~~ migration due to the reductive dissolution of Fe (oxy)hydroxides (Baken et al., 2015; van Dael et al., 2020).

~~We removed t~~Two cores with indications of PO₄ migration ~~were removed from the analysis~~ to address the issue (Old2 and Young2).

These cores were identified by ~~comparing the peak in the~~ DPS age profile ~~with the known peak in surface water PO₄ concentrations in the 1980s~~ and considering the distance from the nearby creeks (Fig. 2; Fig. 1). Additionally, the DPS levels of the deepest sediment layers were compared with layers at the surface. The surface layers had lower DPS levels than the deepest layers for one core (Young 2). The two remaining cores (Old1, Young1) had lower DPS levels in deeper sediment layers (Fig. S7). More importantly, the modelled peak in PO₄ concentrations based on the cores Old1, Young1 were found within two years of the monitored peak and had a similar magnitude (Fig. 4). The coinciding peaks illustrate little migration of PO₄ in Old1 and Young1, thereby justifying these cores as an archive for water-PO₄.

The limited migration is also logical: at the average DPS of 90 % in sediment showing at the peak; the sorption models predict that the solid-liquid P concentration ratio is 2900 L kg⁻¹ ~~with~~ the average K value of models of Table 2. That solid-liquid ratio can be converted to a dimensionless retardation factor ~~representing, representing~~ the ratio of the distance migrated by the PO₄ compared to the distance travelled by percolating water. ~~For example, the retardation was calculated to be 7500 with a bulk density (pb) of 1.3, of 7500 with a bulk density (p_b) of 1.3 and porosity (θ) of 0.5 and~~ With a net vertical annual water percolation of about 2 meters. ~~That, the~~ retardation corresponds to a net vertical P migration rate of 2.5 cm over 100 years, i.e. vanishingly small (calculation details not shown).

Secondly, there is uncertainty on the age-depth estimation of the sampled sediment profiles. ~~We expect that t~~The age-depth model is ~~expected to be~~ most reliable for the Young1 sediment core, as it is based on a fitting of a modelled age-depth relation to four observed age-depth points, while we only had two observed age-depth points available for the other cores (Temmerman et al., 2004a). Additionally, observed age-depth points were not older than 1944. Hence, the extrapolation of the age-depth model to periods before the older available age-depth points is increasingly uncertain.

545 4.3 Pre-industrial and natural background values

The population increase between 1800 and 1930 can provide a first, very crude estimate of the population-DPS relation in the Scheldt basin. In 1800 the population in Belgium was around 3 million. ~~Later, in~~ 1930, this number ~~has~~ more than doubled to 7 million (Vanhaute, 2003). A linear relation between both suggests that the DPS is 8% for the pre-anthropogenic pristine environment, corresponding with a PO_4 -concentration 19-41 $\mu\text{g PO}_4\text{-P L}^{-1}$, i.e. close to what researchers have indicated for
550 pristine lakes. Such predictions need to be corroborated with older sediment observations and other archaeological information. The Scheldt river is logically more aerated than smaller lowland rivers where summer anoxia is naturally more present, i.e. the pristine $\text{PO}_4\text{-P}$ values will be higher.

Formatted: Subscript

5 Conclusions

555 Our study illustrated that tidal marsh sediments could ~~estimate-evaluate~~ pre-industrial background PO_4 -concentrations of the freshwater rivers like the Scheldt river. A sediment assessment can record ~~and time-integrated environmental events, which provide~~ ~~time-integrated environmental events, providing~~ useful spatial and temporal information. Our data estimated the pre-industrial background concentration at 62 $\mu\text{g PO}_4\text{-P L}^{-1}$ [95% CI (57; 66)], about half of the environmental limits set for surface waters in Flanders and neighbouring countries. Around 1930, the PO_4 levels were only about 20% lower than today, which is
560 a remarkably large concentration at a time before the massive application of mineral fertilisers, with lower population density and limited connection to sewer systems. The current PO_4 concentrations decreased ~~by a factor ten times~~ from the peak found 40 years ago, reflecting wastewater treatment efforts and reducing diffuse P emission. It is also clear from this study that the pristine, pre-anthropogenic $\text{PO}_4\text{-P}$ concentrations in the Scheldt river are well below the current ambient ones.

Data availability

565 The ~~results of the sediment data analysis and age depth model~~ ~~supplement provides the sediment data analysis and age depth model results in~~ ~~are provided in the supplement as~~ csv format. ~~Results~~ ~~In addition, results~~ of surface water data are available upon request at ~~the~~ the IMIS (Flanders Marine Institute).

Author contribution

570 FL, ES and PC designed the research. FL conducted the investigation process, and developed the methodology under supervision of ES. MVDB carried out the fieldwork and conceptualised the use of the samples. ST provided the methodology for the age-depth model and software. TM validated the use of the surface water data. EVM and FL placed the results in

perspective with historical data. All the authors contributed to discussion and data interpretations, review and editing of the work.

Competing interests

575 The authors declare that they have no conflict of interest

Acknowledgements

This project was supported by the Research Fund Flanders (FWO), project G089319N. The results of this research greatly depended on the data collected by the OMES-monitoring and The Flemish Waterway. Many years of intensive data collection and quality assessment of the Scheldt river resulted in a unique and valuable phosphate time series. We have the utmost respect
580 for their work and are thankful we could apply the dataset for this research. We acknowledge Dries Grauwels and Kristin Coorevits for technical assistance. We recognise the efforts from the unanimous reviewers for their constructive comments on the work, which improved the quality of the result. Finally, thanks to the Scheldt for providing this beautiful sediment archive to travel back in time and explore environmental history.

585 References

- Azevedo, L. B., Van Zelm, R., Leuven, R. S. E. W., Hendriks, A. J. and Huijbregts, M. A. J.: Combined ecological risks of nitrogen and phosphorus in European freshwaters, *Environ. Pollut.*, 200, 85–92, doi:10.1016/J.ENVPOL.2015.02.011, 2015.
- Baken, S., Verbeeck, M., Verheyen, D., Diels, J. and Smolders, E.: Phosphorus losses from agricultural land to natural waters are reduced by ~~immobilization~~–immobilisation in iron-rich sediments of drainage ditches, *Water Res.*, 71, 160–170, doi:10.1016/j.watres.2015.01.008, 2015.
590
- Ballantine, D. J., Walling, D. E., Collins, A. L. and Leeks, G. J. L.: The content and storage of phosphorus in fine-grained channel bed sediment in contrasting lowland agricultural catchments in the UK, *Geoderma*, 151(3–4), 141–149, doi:10.1016/j.geoderma.2009.03.021, 2009.
- Belliard, J. P., Silinski, A., Meire, D., Kolokythas, G., Levy, Y., Van Braeckel, A., Bouma, T. J. and Temmerman, S.: High-resolution bed level changes in relation to tidal and wave forcing on a narrow fringing macrotidal flat: Bridging intra-tidal, daily and seasonal sediment dynamics, *Mar. Geol.*, doi:10.1016/j.margeo.2019.03.001, 2019.
- Billen, G., Garnier, J. and Rousseau, V.: Nutrient fluxes and water quality in the drainage network of the Scheldt basin over the last 50 years, *Hydrobiologia*, doi:10.1007/s10750-004-7103-1, 2005.
- Birch, G. F., McCready, S., Long, E. R., Taylor, S. S. and Spyrikis, G.: Contaminant chemistry and toxicity of sediments in

- 600 Sydney Harbour, Australia: Spatial extent and chemistry-toxicity relationships, *Mar. Ecol. Prog. Ser.*, doi:10.3354/meps07445, 2008.
- Bitschofsky, F. and Nausch, M.: Spatial and seasonal variations in phosphorus speciation along a river in a lowland catchment (Warnow, Germany), *Sci. Total Environ.*, 657, 671–685, doi:10.1016/J.SCITOTENV.2018.12.009, 2019.
- Bjerrum, C. J. and Canfield, D. E.: Ocean productivity before about 1.9 Gyr ago limited by phosphorus adsorption onto iron oxides, *Nature*, doi:10.1038/417159a, 2002.
- 605 Borggaard, O. K.: Dissolution and adsorption properties of soil iron oxides., Royal Veterinary and Agricultural University., 1990.
- Breeuwsma, A., Reijerink, J. G. A. and Schoumans, O. F.: Impact of manure on accumulation and leaching of phosphate in areas of intensive livestock farming, in *Animal waste and the land-water interface.*, 1995.
- 610 Van de Broek, M., Vandendriessche, C., Poppelmonde, D., Merckx, R., Temmerman, S. and Govers, G.: Long-term organic carbon sequestration in tidal marsh sediments is dominated by old-aged allochthonous inputs in a macrotidal estuary, *Glob. Chang. Biol.*, doi:10.1111/gcb.14089, 2018.
- Van de Broek, M., Baert, L., Temmerman, S. and Govers, G.: Soil organic carbon stocks in a tidal marsh landscape are dominated by human marsh embankment and subsequent marsh progradation, *Eur. J. Soil Sci.*, doi:10.1111/ejss.12739, 2019.
- 615 Van De Broek, M., Temmerman, S., Merckx, R. and Govers, G.: Controls on soil organic carbon stocks in tidal marshes along an estuarine salinity gradient, *Biogeosciences*, doi:10.5194/bg-13-6611-2016, 2016.
- Burson, A., Stomp, M., Akil, L., Brussaard, C. P. D. and Huisman, J.: Unbalanced reduction of nutrient loads has created an offshore gradient from phosphorus to nitrogen limitation in the North Sea, *Limnol. Oceanogr.*, 61(3), 869–888, doi:10.1002/LNO.10257, 2016.
- 620 Callaway, J. C., Nyman, J. A. and DeLaune, R. D.: Sediment accretion in coastal wetlands: A review and a simulation model of processes, *Curr. Top. Wetl. Biogeochem.*, 1996.
- Cardoso, A. C., Solimini, A., Premazzi, G., Carvalho, L., Lyche, A. and Rekolainen, S.: Phosphorus reference concentrations in European lakes, in *Hydrobiologia*, vol. 584, pp. 3–12., 2007.
- Correll, D. L.: The Role of Phosphorus in the Eutrophication of Receiving Waters: A Review, *J. Environ. Qual.*, 27(2), 261–266, doi:10.2134/jeq1998.00472425002700020004x, 1998.
- 625 van Dael, T., De Cooman, T., Verbeeck, M. and Smolders, E.: Sediment respiration contributes to phosphate release in lowland surface waters, *Water Res.*, 168, 115168, doi:10.1016/j.watres.2019.115168, 2020.
- Dodds, W. K. and Smith, V. H.: Nitrogen, phosphorus, and eutrophication in streams, *Int. Waters*, 6(2), 155–164, doi:10.5268/IW-6.2.909, 2016.
- 630 ECOBE-UA and De Vlaamse Waterweg: OMES: Monitoring fysical-chemical water quality in the Zeeschelde. [online] Available from: <http://www.vliz.be/en/imis?module=dataset&dasid=1069> (Accessed 28 September 2020), 2016.
- ECOBE - UA; The Flemish Waterway: OMES monitoring data Zeeschelde since 1995., 2019.
- ECOBE - UAntwerpen: AZ monitoring water quality of the Scheldt. [online] Available from:

<http://www.vliz.be/en/imis?module=dataset&dasid=1468> (Accessed 28 September 2020), 2007.

635 Edmunds, W. and Shand, P.: Natural groundwater quality., 2009.

Elser, J. J., Bracken, M. E. S., Cleland, E. E., Gruner, D. S., Harpole, W. S., Hillebrand, H., Ngai, J. T., Seabloom, E. W., Shurin, J. B. and Smith, J. E.: Global analysis of nitrogen and phosphorus limitation of primary producers in freshwater, marine and terrestrial ecosystems, *Ecol. Lett.*, 10(12), 1135–1142, doi:10.1111/j.1461-0248.2007.01113.x, 2007.

European Commission: DIRECTIVE 2000/60/EC OF THE EUROPEAN PARLIAMENT AND OF THE COUNCIL of 23
640 October 2000 establishing a framework for Community action in the field of water policy., 2000.

Flemish Government: VLAREM II, Vlarem II, EMIS Navig. [online] Available from: <https://navigator.emis.vito.be/mijn-navigator?woId=263> (Accessed 23 December 2020), 1995.

Friedrichs, C. T. and Perry, J. E.: Tidal Salt Marsh Morphodynamics: A Synthesis, *J. Coast. Res.*, 2001.

645 Froelich, P. N.: Kinetic control of dissolved phosphate in natural rivers and estuaries: A primer on the phosphate buffer mechanism I, *Limnol. Oceanogr.*, 33(4part2), 649–668, doi:10.4319/lo.1988.33.4part2.0649, 1988.

Hiemstra, T., Antelo, J., Rahnamaie, R. and Riemsdijk, W. H. va.: Nanoparticles in natural systems I: The effective reactive surface area of the natural oxide fraction in field samples, *Geochim. Cosmochim. Acta*, 74(1), 41–58, doi:10.1016/j.gca.2009.10.018, 2010.

650 Holtan, H., Kamp-Nielsen, L. and Stuanes, A. O.: Phosphorus in soil, water and sediment: an overview, *Hydrobiologia*, 170(1), 19–34, doi:10.1007/BF00024896, 1988.

House, W. A. and Denison, F. H.: Phosphorus dynamics in a lowland river, *Water Res.*, 32(6), 1819–1830, doi:10.1016/S0043-1354(97)00407-7, 1998.

Huet, H. J. W. J. Van: Phosphorus eutrophication research in the lake district of south western Friesland, The Netherlands. Preliminary results of abiotic studies, *Trophic Relationships Inl. Waters*, 75–85, doi:10.1007/978-94-009-0467-5_10, 1990.

655 Institute voor Hygiëne en Epidemiologie (IHE): Scheldt water quality data. [online] Available from: <http://www.vliz.be/en/imis?module=dataset&dasid=1438> (Accessed 28 September 2020), 2007.

Formatted: Dutch (Belgium)

Jarvie, H. P., Neal, C. and Withers, P. J. A.: Sewage-effluent phosphorus: A greater risk to river eutrophication than agricultural phosphorus?, *Sci. Total Environ.*, 360(1–3), 246–253, doi:10.1016/j.scitotenv.2005.08.038, 2006.

660 Laane, R. W. P. M.: Background concentrations of natural compounds in rivers, sea water, atmosphere and mussels, The Hague. [online] Available from: <http://publicaties.minienm.nl/documenten/background-concentrations-of-natural-compounds-in-rivers-sea-wat> (Accessed 19 October 2020), 1992.

Lexmond, T. M., Riemsdijk, W. H. van and Haan, F. A. M. de: Onderzoek naar fosfaat en koper in de bodem in het bijzonder in gebieden met intensieve veehouderij, L.H. [online] Available from: <https://research.wur.nl/en/publications/onderzoek-naar-fosfaat-en-koper-in-de-bodem-in-het-bijzonder-in-g> (Accessed 15 September 2021), 1982.

Formatted: Dutch (Belgium)

665 Lookman, R., Vandeweert, N., Merckx, R. and Vlaskak, K.: Geostatistical assessment of the regional distribution of phosphate sorption capacity parameters (FeOX and AlOX) in northern Belgium, *Geoderma*, 66(3–4), 285–296, doi:10.1016/0016-7061(94)00084-N, 1995.

- Mainstone, C. P. and Parr, W.: Phosphorus in rivers - Ecology and management, *Sci. Total Environ.*, 282–283, 25–47, doi:10.1016/S0048-9697(01)00937-8, 2002.
- 670 Matschullat, J., Ottenstein, R. and Reimann, C.: Geochemical background - Can we calculate it?, *Environ. Geol.*, 39(9), 990–1000, doi:10.1007/s002549900084, 2000.
- Van Meel, L.: hydrobiology of the Sea-Scheldt near Liefkenshoek. [online] Available from: <http://www.vliz.be/en/imis?module=dataset&david=1412> (Accessed 28 September 2020), 1958.
- Meire, P., Ysebaert, T., Van Damme, S., Van Den Bergh, E., Maris, T. and Struyf, E.: The Scheldt estuary: A description of a changing ecosystem, *Hydrobiologia*, 540(1–3), 1–11, doi:10.1007/s10750-005-0896-8, 2005.
- 675 Migon, C. and Sandroni, V.: Phosphorus in rainwater: Partitioning inputs and impact on the surface coastal ocean, *Limnol. Oceanogr.*, 44(4), 1160–1165, doi:10.4319/lo.1999.44.4.1160, 1999.
- Van Der Molen, D. T., Portielje, R., Boers, P. C. M. and Lijklema, L.: Changes in sediment phosphorus as a result of eutrophication and oligotrophication in Lake Veluwe, The Netherlands, *Water Res.*, 32(11), 3281–3288, doi:10.1016/S0043-680 1354(98)00117-1, 1998.
- Moriasi, D. N., Arnold, J. G., Liew, M. W. Van, Bingner, R. L., Harmel, R. D. and Veith, T. L.: MODEL EVALUATION GUIDELINES FOR SYSTEMATIC QUANTIFICATION OF ACCURACY IN WATERSHED SIMULATIONS, *Trans. ASABE*, 50(3), 1983.
- Nash, J. E. and Sutcliffe, J. V.: River flow forecasting through conceptual models part I - A discussion of principles, *J. Hydrol.*, doi:10.1016/0022-1694(70)90255-6, 1970.
- 685 Nürnberg, G. K.: Quantifying anoxia in lakes, *Limnol. Oceanogr.*, 40(6), 1100–1111, doi:10.4319/LO.1995.40.6.1100, 1995.
- De Pauw, C.: The environment and plankton of the WesterScheldt estuary, Ghent. [online] Available from: <http://www.vliz.be/en/imis?module=dataset&david=1390>, 2007.
- Pethick, J. S.: Long-term accretion rates on tidal salt marshes., *J. Sediment. Petrol.*, doi:10.1306/212F7CDE-2B24-11D7-690 8648000102C1865D, 1981.
- Poppelmonde, D.: Organic carbon dynamics in tidal marshes of the Scheldt estuary A combined field and modelling approach, KULeuven, VUB., 2017.
- Van Puijenbroek, P. J. T. M., Cleij, P. and Visser, H.: Aggregated indices for trends in eutrophication of different types of fresh water in the Netherlands, *Ecol. Indic.*, 36, 456–462, doi:10.1016/J.ECOLIND.2013.08.022, 2014.
- 695 R Core Team: R: A language and environment for statistical computing, [online] Available from: <https://www.r-project.org/>, 2020.
- van Raaphorst, W. and Kloosterhuis, H. T.: Phosphate sorption in superficial intertidal sediments, *Mar. Chem.*, 48(1), 1–16, doi:10.1016/0304-4203(94)90058-2, 1994.
- van Raaphorst, W., de Jonge, V. N., Dijkhuizen, D. and Frederiks, B.: Natural background concentrations of phosphorus and nitrogen in the Dutch Wadden Sea, *Rapp. voor Kust en Zee*, 53pp., 2000.
- 700 Reimann, C. and Garrett, R. G.: Geochemical background - Concept and reality, *Sci. Total Environ.*,

doi:10.1016/j.scitotenv.2005.01.047, 2005.

Reynolds, C. S.: Phosphorus recycling in lakes: Evidence from large limnetic enclosures for the importance of shallow sediments, *Freshw. Biol.*, 35(3), 623–645, doi:10.1111/j.1365-2427.1996.tb01773.x, 2000.

705 Rönspieß, L., Dellwig, O., Lange, X., Nausch, G. and Schulz-Bull, D.: Spatial and seasonal phosphorus dynamics in a eutrophic estuary of the southern Baltic Sea, *Estuar. Coast. Shelf Sci.*, 233, 106532, doi:10.1016/J.ECSS.2019.106532, 2020.

RStudio Team: RStudio: Integrated Development for R., [online] Available from: <http://www.rstudio.com/>, 2015.

Salminen, R., Batista, M. J., Bidovec, M. D., Demetriades, A., De Vivo, B., De Vos, W., Duris, M., Gilucis, A., Gregorauskiene, V., Halamic, J., Heitzmann, P., Lima, A., Jordan, G., Klaver, G., Klein, P., Lis, J., Locutura, J., Marsina, K.,

710 Mazreku, A., ~~O'Connor~~O'Connor, P. J., Olsson, S. Å., Ottesen, R.-T., Petersell, V., Plant, J. a., Reeder, S., Salpeteur, I., Sandström, H., Siewers, U., Steenfelt, A. and Tarvainen, T.: Part 1- Background information, methodology and maps., in *Geochemical Atlas of Europe.*, 2005.

ScheldeMonitor Team and VNSC, R. & M.: Data downloaded from ScheldeMonitor: a data portal with information, data and products on the Scheldt Estuary, Data downloaded from ScheldeMonitor a data portal with information, data Prod. Scheldt Estuary [online] Available from: <https://rshiny.scheldemonitor.org/waterniveauschelde/> (Accessed 12 April 2021), 2020.

715 Schönfelder, I. and Steinberg, C. E. W.: How did the nutrient concentrations change in northeastern German lowland rivers during the last four millennia?-A paleolimnological study of floodplain sediments, [online] Available from: <https://doi.org/10.18452/9393> (Accessed 8 October 2021), 2004.

Schoumans, O. F. and Chardon, W. J.: Phosphate saturation degree and accumulation of phosphate in various soil types in The Netherlands, *Geoderma*, 237, 325–335, doi:10.1016/j.geoderma.2014.08.015, 2015.

720 Schoumans, O. F. and Groenendijk, P.: Modeling Soil Phosphorus Levels and Phosphorus Leaching from Agricultural Land in the Netherlands, *J. Environ. Qual.*, 29(1), 111–116, doi:10.2134/jeq2000.00472425002900010014x, 2000.

Schulz, M. and Herzog, C.: The influence of sorption processes on the phosphorus mass balance in a eutrophic German lowland river, *Water, Air, Soil Pollut.*, doi:10.1023/B:WATE.0000026535.27164.56, 2004.

725 Schwertmann, U.: Differenzierung der Eisenoxide des Bodens durch Extraktion mit Ammoniumoxalat-Lösung, *Zeitschrift für Pflanzenernährung, Düngung, Bodenkd.*, 105(3), 194–202, doi:10.1002/jpln.3591050303, 1964.

Simpson, Z. P., McDowell, R. W., Condon, L. M., McDaniel, M. D., Jarvie, H. P. and Abell, J. M.: Sediment phosphorus buffering in streams at baseflow: A meta-analysis, *J. Environ. Qual.*, 50(2), 287–311, doi:10.1002/JEQ2.20202, 2021.

Smolders, E., Baetens, E., Verbeeck, M., Nawara, S., Diels, J., Verdievel, M., Peeters, B., De Cooman, W. and Baken, S.: 730 Internal Loading and Redox Cycling of Sediment Iron Explain Reactive Phosphorus Concentrations in Lowland Rivers, *Environ. Sci. Technol.*, 51(5), 2584–2592, doi:10.1021/acs.est.6b04337, 2017.

Struyf, E., Temmerman, S. and Meire, P.: Dynamics of biogenic Si in freshwater tidal marshes: Si regeneration and retention in marsh sediments (Scheldt estuary), *Biogeochemistry*, doi:10.1007/s10533-006-9051-5, 2007.

735 Svendsen, L. M. and Kronvang, B.: Retention of nitrogen and phosphorus in a Danish lowland river system: implications for the export from the watershed, *Nutr. Dyn. Retent. Land/Water Ecotones Lowl. Temp. Lakes Rivers*, 123–135,

Formatted: German (Germany)

doi:10.1007/978-94-011-1602-2_15, 1993.

De Swart, H. E. and Zimmerman, J. T. F.: Morphodynamics of tidal inlet systems, *Annu. Rev. Fluid Mech.*, doi:10.1146/annurev.fluid.010908.165159, 2009.

740 Temmerman, S., Govers, G., Meire, P. and Wartel, S.: Modelling long-term tidal marsh growth under changing tidal conditions and suspended sediment concentrations, Scheldt estuary, Belgium, *Mar. Geol.*, doi:10.1016/S0025-3227(02)00642-4, 2003a.

Temmerman, S., Govers, G., Wartel, S. and Meire, P.: Spatial and temporal factors controlling short-term sedimentation in a salt and freshwater tidal marsh, scheldt estuary, Belgium, SW Netherlands, *Earth Surf. Process. Landforms*, doi:10.1002/esp.495, 2003b.

745 Temmerman, S., Govers, G., Wartel, S. and Meire, P.: Modelling estuarine variations in tidal marsh sedimentation: Response to changing sea level and suspended sediment concentrations, *Mar. Geol.*, doi:10.1016/j.margeo.2004.10.021, 2004a.

Temmerman, S., Govers, G., Meire, P. and Wartel, S.: Simulating the long-term development of levee-basin topography on tidal marshes, *Geomorphology*, 63(1–2), 39–55, doi:10.1016/j.geomorph.2004.03.004, 2004b.

Vanhaute, E.: en arbeid in België in de ' lange negentiende eeuw ', , 118(2001), 153–178, 2003.

VMM: Milieurapport Vlaanderen - Systeembalans 2017. [online] Available from: 750 http://www.milieurapport.be/Upload/main/0_topicrapporten/361312_Systeembalans2017_nieuw.pdf, 2017.

Wang, Y., Shen, Z., Niu, J. and Liu, R.: Adsorption of phosphorus on sediments from the Three-Gorges Reservoir (China) and the relation with sediment compositions, *J. Hazard. Mater.*, 162(1), 92–98, doi:10.1016/j.jhazmat.2008.05.013, 2009.

755 Warrinnier, R., Goossens, T., Braun, S., Gustafsson, J. P. and Smolders, E.: Modelling heterogeneous phosphate sorption kinetics on iron oxyhydroxides and soil with a continuous distribution approach, *Eur. J. Soil Sci.*, 69(3), 475–487, doi:10.1111/ejss.12549, 2018.

Warrinnier, R., Goossens, T., Amery, F., Vanden Nest, T., Verbeeck, M. and Smolders, E.: Investigation on the control of phosphate leaching by sorption and colloidal transport: Column studies and multi-surface complexation modelling, *Appl. Geochemistry*, 100, 371–379, doi:10.1016/j.apgeochem.2018.12.012, 2019.

760 Watson, S. J., Cade-Menun, B. J., Needoba, J. A. and Peterson, T. D.: Phosphorus Forms in Sediments of a River-Dominated Estuary, *Front. Mar. Sci.*, 0(SEP), 302, doi:10.3389/FMARS.2018.00302, 2018.

Zak, D., Kleeberg, A. and Hupfer, M.: Sulphate-mediated phosphorus ~~mobilization~~-mobilisation in riverine sediments at increasing sulphate concentration, River Spree, NE Germany, *Biogeochemistry*, 80(2), 109–119, doi:10.1007/s10533-006-0003-x, 2006.

765 van der Zee, C., Roelvros, N. and Chou, L.: Phosphorus speciation, transformation and retention in the Scheldt estuary (Belgium/The Netherlands) from the freshwater tidal limits to the North Sea, *Mar. Chem.*, 106(1-2 SPEC. ISS.), 76–91, doi:10.1016/j.marchem.2007.01.003, 2007.

van der Zee, S. E. A. T. M.: Transport of reactive contaminants in heterogeneous soil systems., 1988.

van der Zee, S. E. A. T. M., van Riemsdijk, W. H. and de Haan, F. A. M.: HET PROTOKOL FOSFAATVERZADIGDE GRONDEN., 1990.

Formatted: Dutch (Belgium)

Formatted: Dutch (Belgium)

770 Zhou, A., Tang, H. and Wang, D.: Phosphorus adsorption on natural sediments: Modeling and effects of pH and sediment composition, *Water Res.*, 39(7), 1245–1254, doi:10.1016/j.watres.2005.01.026, 2005.