

1 **Temporal dynamics of surface ocean carbonate chemistry in response**
2 **to natural and simulated upwelling events during the 2017 coastal El**
3 **Niño near Callao, Peru**

4 Shao-Min Chen^{1, 2}, Ulf Riebesell¹, Kai G. Schulz³, Elisabeth von der Esch¹, Eric P. Achterberg¹, and
5 Lennart T. Bach⁴

6 ¹GEOMAR Helmholtz Centre for Ocean Research Kiel, Kiel, Germany

7 ²Department of Earth and Environmental Sciences, Dalhousie University, Halifax, Canada

8 ³Centre for Coastal Biogeochemistry, School of Environment, Science and Engineering, Southern Cross University, Lismore,
9 Australia

10 ⁴Institute for Marine and Antarctic Studies, University of Tasmania, Tasmania, Australia

11

12 *Correspondence to:* Shao-Min Chen (shaomin.chen@dal.ca)

13 **Abstract.** Oxygen minimum zones (OMZs) are characterized by enhanced carbon dioxide (CO₂) levels and low pH and are
14 being further acidified by uptake of anthropogenic atmospheric CO₂. With ongoing intensification and expansion of OMZs
15 due to global warming, carbonate chemistry conditions may become more variable and extreme, particularly in the Eastern
16 Boundary Upwelling Systems. In austral summer (Feb-Apr) 2017, a large-scale mesocosm experiment was conducted in the
17 coastal upwelling area off Callao (Peru) to investigate the impacts of on-going ocean deoxygenation on biogeochemical
18 processes, coinciding with a rare coastal El Niño event. Here we report on the temporal dynamics of carbonate chemistry in
19 the mesocosms and surrounding Pacific waters over a continuous period of 50 days with high temporal resolution observations
20 (every 2nd day). The mesocosm experiment simulated an upwelling event in the mesocosms by addition of nitrogen (N)-
21 deficient and CO₂-enriched OMZ water. Surface water in the mesocosms was acidified by the OMZ water addition, with pH_T
22 lowered by 0.1-0.2 and pCO₂ elevated to above 900 µatm. Thereafter, surface pCO₂ quickly dropped to near or below the
23 atmospheric level (405.22 µatm in 2017, NOAA/GML) mainly due to enhanced phytoplankton production with rapid CO₂
24 consumption. Further observations revealed that the dominance of dinoflagellate *Akashiwo sanguinea* and contamination of
25 bird excrements played important roles in the dynamics of carbonate chemistry in the mesocosms. Compared to the simulated
26 upwelling, natural upwelling events in the surrounding Pacific waters occurred more frequently with sea-to-air CO₂ fluxes of
27 4.2-14.0 mmol C m⁻² d⁻¹. The positive CO₂ fluxes indicated our site was a local CO₂ source during our study, which may have
28 been impacted by the coastal El Niño. However, our observations of DIC drawdown in the mesocosms suggests that CO₂
29 fluxes to the atmosphere can be largely dampened by biological processes. Overall, our study characterized carbonate
30 chemistry in near-shore Pacific waters that are rarely sampled in such temporal resolution and hence provided unique insights
31 into the CO₂ dynamics during a rare coastal El Niño event.

32 **1 Introduction**

33 One of the most extensive oxygen minimum zones (OMZs) in the global ocean can be found off central/northern Peru (4 - 16°
34 S, Chavez and Messié, 2009). High biological productivity is stimulated by permanent upwelling of cold, nutrient-rich water
35 to the surface supporting a remarkable fish production off Peru (Chavez et al., 2008; Montecino and Lange, 2009; Albert et
36 al., 2010). The high primary production also leads to enhanced remineralization of sinking organic matter in subsurface waters
37 which depletes dissolved oxygen (O₂) and creates an intense and shallow OMZ (Chavez et al., 2008). The depletion of O₂ in
38 OMZs plays an important role in the global nitrogen (N) cycle, accounting for 20 - 40% N loss in the ocean (Lam et al., 2009;
39 Paulmier and Ruiz-Pino, 2009). Denitrification and anammox processes that occur in O₂ depleted waters remove **biologically**
40 **available** N from the ocean and produce an N deficit and hence phosphorus (P) excess with respect to the Redfield ratio (C:N:P
41 = 106:16:1) in the water column (Redfield, 1963; Deutsch et al., 2001; Deutsch et al., 2007; Hamersley et al., 2007; Galán et
42 al., 2009; Lam et al., 2009). Upwelling of this N-deficient water has been found to control the surface-water nutrient
43 stoichiometry and thus influence phytoplankton growth and community compositions (Franz et al., 2012; Hauss et al., 2012).

44 Apart from being N-deficient, the OMZ waters are also characterized by enhanced carbon dioxide (CO₂) concentrations and
45 low pH from respiratory processes and are further acidified by **increasing uptake of** anthropogenic atmospheric CO₂ (Feely et
46 al., 2008; Friederich et al., 2008; Paulmier et al., 2008; Paulmier et al., 2011). Accordingly, surface water carbonate chemistry
47 is influenced by upwelling of CO₂-enriched OMZ water (Van Geen et al., 2000; Capone and Hutchins, 2013). The upwelled
48 CO₂-enriched OMZ water can give rise to surface CO₂ levels >1,000 µatm, pH values as low as 7.6, and under-saturation for
49 the calcium carbonate mineral aragonite (Feely et al., 2008; Hauri et al., 2009). As a result, there is a significant flux of CO₂
50 from the ocean to the atmosphere off Peru, which is further facilitated by surface ocean warming, making the Peruvian
51 upwelling region a year-round CO₂ source to the atmosphere (Friederich et al., 2008). In contrast, rapid utilization of upwelled

52 CO₂ and nutrients by phytoplankton can occasionally deplete surface CO₂ below atmospheric equilibrium and dampen the CO₂
53 outgassing (Van Geen et al., 2000; Friederich et al., 2008; Loucaides et al., 2012). The enhanced primary production in turn
54 contributes to increasing export of organic matter, enhanced bacterial respiration, O₂ consumption and CO₂ production at
55 depth. Such a positive feedback may determine the intensity of the underlying OMZ and promote carbon (C) preservation in
56 marine sediments (Dale et al., 2015).

57 In response to reduced O₂ solubility and enhanced stratification induced by global warming, OMZs have been intensifying and
58 expanding over the past decades (Stramma et al., 2008; Fuenzalida et al., 2009; Stramma et al., 2010). Based on regional
59 observations and model projections, a decline in dissolved O₂ concentrations has been reported for most regions of the global
60 ocean (Matear et al., 2000; Matear and Hirst, 2003; Whitney et al., 2007; Stramma et al., 2008; Keeling et al., 2009; Bopp et
61 al., 2013; Schmidtko et al., 2017; Oschlies et al., 2018). The vertical expansion of OMZs represents shoaling of CO₂-enriched
62 seawater, which has become further enriched by oceanic uptake of anthropogenic CO₂ (Doney et al., 2012; Gilly et al., 2013;
63 Schulz et al., 2019). Since biogeochemical processes in OMZs are directly linked to the C cycle and control surface nutrient
64 stoichiometry, with on-going ocean warming and acidification, the deoxygenation may have cascading effects on plankton
65 productivity and composition, C uptake, and food web functioning (Keeling et al., 2009; Gruber, 2011; Doney et al., 2012;
66 Gilly et al., 2013; Levin and Breitburg, 2015). Therefore, it is important to monitor the changes in CO₂ when investigating the
67 effects of deoxygenation on marine ecosystems.

68 To investigate the potential impacts of upwelling on pelagic biogeochemistry and natural plankton communities in the Peruvian
69 OMZ, a large-scale *in situ* mesocosm study was carried out in the coastal upwelling area off Peru. An upwelling event was
70 simulated in the mesocosms by addition of OMZ waters collected from two different locations where the OMZ was considered
71 to contain different nutrient concentrations and N:P ratios. The ecological and biogeochemical responses in the mesocosms
72 were monitored and compared with those influenced by natural upwelling events in the ambient coastal water surrounding the
73 mesocosms. As part of this collaborative research project, questions specific to the present paper were: (1) How does surface
74 water carbonate chemistry respond to an upwelling event?; and (2) How does upwelled OMZ water with different chemical
75 signatures modulate surface water carbonate chemistry? The current study will mainly focus on the temporal changes in surface
76 water carbonate chemistry within the individual mesocosms, including observations made in the ambient Pacific water and a
77 local estimate of air-sea CO₂ exchange, together with the influence by a rare coastal El Niño event (Garreaud, 2018). This
78 provides first insights into how inorganic C cycling links to chemical signatures of OMZ waters in a natural plankton
79 community and its implications for ongoing environmental changes.

80 **2 Material and methods**

81 **2.1 Study site**


82 The experiment was conducted in the framework of the Collaborative Research Center 754 "Climate-Biogeochemistry
83 Interactions in the Tropical Ocean" (www.sfb754.de/en) and in collaboration with the Instituto del Mar del Peru (IMARPE)
84 in Callao, Peru (Fig. 1a). The coastal area off Callao lies within the Humboldt Current System and is influenced by wind-
85 induced coastal upwelling (Bakun and Weeks, 2008).

86 **2.2 Mesocosm setup**

87 Eight "Kiel Off-Shore Mesocosms for Future Ocean Simulations" (KOSMOS) units (M1-M8), extending 19 m below the sea
88 surface, were deployed by the research vessel *Buque Armada Peruana (BAP) Morales* and moored at 12.06° S, 77.23° W in

89 the coastal upwelling area off Callao, Peru (Fig. 1a) on February 23rd, 2017 (late austral summer). The technical design of
90 these sea-going mesocosms is described by Riebesell et al. (2013). For a more detailed description of the mesocosm
91 deployment and maintenance in this study, please refer to Bach et al. (2020^a).

92 The mesocosm bags were filled with surrounding seawater through the upper and lower openings. Both openings were covered
93 by screens with a mesh size of 3 mm to avoid enclosing larger organisms such as fish. The mesocosm bags were left open
94 below the water surface for two days, allowing free exchange with surrounding coastal water. On February 25th, mesocosm
95 bags were closed with the screens removed, tops pulled above the sea surface and bottoms sealed with 2-m long conical
96 sediment traps (Fig. 1b). The experiment started with the closure of the mesocosms (day 0) and lasted for 50 days. Each
97 mesocosm bag enclosed a seawater volume of ~54 m³. After the bags were closed, daily or every-2nd-day sampling was
98 performed to monitor the initial conditions of the enclosed water before simulating an upwelling event on day 11 and 12 (see
99 Sect. 2.4 for details).



100
101 **Figure 1** The study site of the mesocosm experiment (a) created and modified using Ocean Data View (Schlitzer, Reiner, Ocean
102 Data View, odv.awi.de, 2021) and a schematic illustration of a KOSMOS mesocosm unit (b). We acknowledge reprint permission
103 from the AGU as parts of this drawing was used for a publication by Bach et al. (2016). The star symbol marks the approximate
104 location of mesocosm deployment.

105 2.3 Simulated upwelling and salt addition

106 To simulate an upwelling event in the mesocosms, OMZ-influenced waters were collected from the nearby coastal area and
107 added to the mesocosms. Two OMZ water masses were collected at Station 1 (12° 01.70' S, 77° 13.41' W) at a depth of ~30
108 m and at Station 3 (12° 02.41' S, 77° 22.50' W) at a depth of ~70 m respectively using a deep-water collection system as
109 described by Taucher et al. (2017). These two water masses were sampled for chemical and biological variables as the
110 mesocosms (see Sect. 2.4). The OMZ water collected from Station 3 had a dissolved inorganic nitrogen (DIN) concentration
111 of 4.3 $\mu\text{mol L}^{-1}$ (denoted as “Low DIN” in this paper) and was added to M2, M3, M6 and M7. The OMZ water from Station 1
112 had a DIN of 0.3 $\mu\text{mol L}^{-1}$ (denoted as “Very low DIN” in this paper) and was added to M1, M4, M5 and M8. Before OMZ
113 water addition, approximately 9 m³ of seawater were removed from 11-12 m of each mesocosm on March 5th (day 8). During
114 the night of March 8th (day 11), ~10 m³ of OMZ water were added to 14-17 m of each mesocosm. On March 9th (day 12), ~10
115 m³ of seawater were removed from 8-9 m followed by an addition of ~12 m³ OMZ water to 0-9 m of each mesocosm.

116 To maintain a low-O₂ bottom layer in the mesocosms and avoid convective mixing induced by heat exchange with the
117 surrounding Pacific, 69 L of a concentrated sodium chloride (NaCl) brine solution were added to the bottom of each mesocosm
118 (10-17 m) on day 13, which increased the bottom salinity by ~0.7 units. Since then, turbulent mixing induced by sampling
119 activities continuously interrupted the artificial halocline. Hence, on day 33, 46 L of the NaCl brine solution were added again
120 to the bottom of each mesocosm (12.5-17 m), which increased the bottom salinity by ~0.5 units. At the end of the experiment
121 after the last sampling (day 50), 52 kg of NaCl brine was added again to each mesocosm to calculate the enclosed seawater
122 volume from a measured salinity change by ~0.2 units (see Czerny et al., 2013a and Schulz et al., 2013 for details). The average
123 final volume for each mesocosm bag was calculated at ~54 m³. With known sampling volumes and deep-water addition
124 volumes during the experiment, the enclosed volumes of each mesocosm on each sampling day could be calculated. The NaCl
125 solution for the halocline establishments had been prepared in Germany by dissolving 300 kg of food industry grade NaCl
126 (free of anti-caking agents) in 1000 L deionized water (Milli-Q, Millipore) and purified with ion exchange resin (LewawitTM
127 MonoPlus TP260®, Lanxess, Germany) to minimize potential contaminations with trace metals (Czerny et al., 2013a). The
128 NaCl solution for the volume determination was produced on site using locally purchased table salt. For a more detailed
129 description of OMZ water and salt additions, please refer to Bach et al. (2020a).

130 **2.4 Sampling procedures and CTD operations**

131 Sampling was carried out in the morning (7 a.m.-11 a.m. local time) daily or every 2nd day throughout the entire experimental
132 period. Depth-integrated samples were taken from the surface (0-10 m for day 3-28) and bottom layer (10-17 m for day 3-28)
133 of the mesocosms and the surrounding coastal water (named “Pacific”) using a 5-L integrating water sampler (IWS, HYDRO-
134 BIOS, Kiel). Due to the deepening of the oxycline as observed from the CTD profiles, the sampling depth for the surface was
135 adjusted to 0-12.5 m while that for the bottom was changed to 12.5-17 m from day 29 until the end of the experiment (day 50).

136 For gas-sensitive variables such as pH and dissolved inorganic carbon (DIC), 1.5 L of seawater from each integrated depth in
137 each mesocosm were taken directly from the fully-filled 5-L integrating water sampler. Clean polypropylene sampling bottles
138 (rinsed with deionized water in the laboratory; Milli-Q, Millipore) were pre-rinsed with sample water immediately prior to
139 sampling. Bottles were filled from bottom to top using pre-rinsed Tygon tubing with overflow of at least one sampling bottle
140 volume (1.5 L) to minimize the impact of CO₂ air-water gas exchange. Nutrient samples were collected into 250 ml
141 polypropylene bottles using pre-rinsed Tygon tubing (see Bach et al., 2020a for details). Sample containers were stored in cool
142 boxes for ~3 hours, protected from sunlight and heat before being transported to the shore. Once in the lab, sample water was
143 sterile-filtered by gentle pressure using syringe filters (0.2 µm pore size), Tygon tubing and a peristaltic pump to remove
144 particles that may cause changes to seawater carbonate chemistry (Bockmon and Dickson, 2014). For DIC measurements, the
145 water was filtered from the bottom of the 1.5-L sample bottle into 100-ml glass-stoppered bottles (DURAN) with an overflow
146 of at least 100 ml to minimize contact with air. Once the glass bottle was filled with sufficient overflow, it was immediately
147 sealed without headspace using a round glass stopper. This procedure was repeated to collect a second bottle (100 ml) of
148 filtered water for pH measurements. The leftover seawater was directly filtered into a 500 ml polypropylene bottle for total
149 alkalinity (TA) measurements (non-gas-sensitive). Filtered DIC and pH samples were stored at 4 °C in the dark and TA samples
150 were at room temperature in the dark until further analysis. Samples were analysed for DIC and pH on the same day of
151 sampling, while TA was determined overnight (see Sect. 2.5 for analytical procedures).

152 CTD casts were performed with a multiparameter logging probe (CTD60M, Sea and Sun Technology) in the mesocosms and
153 Pacific on every sampling day. From the CTD casts, profiles of salinity, temperature, pH, dissolved O₂, chlorophyll *a* (chl *a*)
154 and photosynthetically active radiation were obtained (see Schulz and Riebesell, 2013 and Bach et al., 2020a for details).

155 2.5 Carbonate chemistry and nutrient measurements

156 Total alkalinity was determined at room temperature (22-32°C) by a two-stage open-cell potentiometric titration using a
157 Metrohm 862 Compact Titrosampler, Aquatrode Plus (Pt1000) and a 907 Titrand unit in the IMARPE laboratory following
158 Dickson et al. (2003). The acid titrant was prepared by preparing a 0.05 mol kg⁻¹ hydrochloric acid (HCl) solution with an
159 ionic strength of ca. 0.7 mol kg⁻¹ (adjusted by NaCl). Approximately 50-grams of sample water from each sample was weighed
160 into the titration cell with the exact weight recorded (precision: 0.0001 g). After the two-stage titration, the titration data
161 between a pH of ~3.5 and 3 was fitted to a modified non-linear Gran approach described in Dickson et al. (2007) using
162 MATLAB (The MathWorks). The results were calibrated against certified reference materials (CRMs) batch 142 (Dickson,
163 2010) measured on each measurement day. In this paper, measured TA values refer to the measured values that have been
164 calibrated against the CRM.

165 Seawater pH_T (total scale) was determined spectrophotometrically by measuring the absorbance ratios after adding the
166 indicator dye m-cresol purple (mCP) as described in Carter, et al. (2013). Before measurements, samples were acclimated to
167 25.0°C in a thermostatted bath. The absorbance of samples with mCP was determined on a Varian-Cary 100 double-beam
168 spectrophotometer (Varian), scanning between 780 and 380 nm at 1-nm resolution. During the spectrophotometric
169 measurement, the temperature of the sample was maintained at 25.0°C by a water-bath connected to the thermostatted 10-cm
170 cuvette. The pH_T values were calculated from the baseline-corrected absorbance ratios and corrected for *in situ* salinity
171 (obtained from CTD casts) and pH change caused by dye addition (using the absorbance at the isosbestic point, i.e. 479 nm)
172 as described in Dickson et al. (2007). To minimize potential CO₂ air-water gas exchange, a syringe pump (Tecan Cavro XLP)
173 was used for sample/dye mixing and cuvette injection (see Schulz et al. 2017 for details). For the dye correction, a batch of
174 sterile filtered seawater of known salinity was prepared. The pH_T was determined once for an addition of 7 ul of dye and once
175 of 25 ul at five pH levels (raised to 7.95 with NaOH and lowered to 7.74, 7.58, 7.49 and 7.36 with HCl stepwise). The pH
176 change resulting from the dye correction addition was calculated from the change in measured absorbance ratio for each pair
177 of dye additions (see Clayton and Byrne, 1993 and Dickson et al., 2007 for details). The dye-corrected pH_T values measured
178 at 25.0°C and atmospheric pressure were then re-calculated for *in situ* temperature and pressure as determined by CTD casts
179 (averaged over 0-10/12.5 m for surface and 10/12.5-17 m for bottom). For carbonate chemistry speciation calculations (see
180 Sect. 2.6), the dye-corrected pH_T values were used as one of the input parameters.

181 Dissolved inorganic carbon was measured by infrared absorption using a LI-COR LI-7000 on an AIRICA system
182 (MARIANDA, Kiel, see Taucher et al., 2017 and Gafar and Schulz, 2018 for details). The results were calibrated against
183 CRMs batch 142 (Dickson, 2010). Unfortunately, due to a malfunctioning of the AIRICA system, we obtained measured DIC
184 data only up to March 7th (day 10). Therefore, measured TA and pH_T were used for calculations of carbonate system
185 parameters at *in situ* temperature and salinity but we used DIC measurements from day 3-10 for consistency checks of
186 calculated carbonate chemistry parameters. In this paper, measured DIC values refer to the measured values that have been
187 calibrated against the CRM.

188 Inorganic nutrients were analyzed colorimetrically (NO₃⁻, NO₂⁻, PO₄³⁻ and Si(OH)₄) and fluorimetrically (NH₄⁺) using a
189 continuous flow analyzer (QuAatro AutoAnalyzer with integrated photometers, SEAL Analytical) connected to a fluorescence
190 detector (FP-2020, JASCO). All colorimetric methods were conducted according to Murphy and Riley (1962), Mullin and
191 Riley (1955a, b) and Morris and Riley (1963) and corrected following the refractive index method developed by Coverly et al.
192 (2012). For details of the quality control procedures, see Bach et al. (2020^a).

193 2.6 Carbonate chemistry speciation calculations and propagated uncertainties

194 Calculations of carbonate chemistry parameters (*in situ* pH_T, DIC, pCO₂, and calcium carbonate saturation state for calcite and
 195 aragonite) were performed with the Excel version of CO2SYS (Version 2.1, Pierrot et al., 2006) using K1 and K2 dissociation
 196 constants from Mehrbach et al., (1973) which were refitted by Lueker et al. (2000). The dissociation constant for KHSO₄ from
 197 Dickson (1990) and for total boron from Uppström (1974) were applied in the calculations (see Orr et al., 2015 for details).
 198 The observed pH_T and TA as well as inorganic nutrient concentration (phosphate and silicic acid) were used as input CO₂
 199 system parameters. *In situ* salinity and temperature were obtained by CTD casts and averaged over surface (0-10 m or 0-12.5
 200 m) and bottom (10-17 m or 12.5-17 m) waters for each sampling day. In situ pressure was approximated for surface (5 dbar)
 201 and bottom (13.5 or 14.75 dbar) waters. For details of calculation procedures and choices of constants, see Lewis et al. (1998)
 202 and Orr et al. (2015).

203 To evaluate the performance of pH_T and TA measurements, quality control procedures were performed. First, standard
 204 deviations of pH_T measurements were graphed over time. Measured TA values of a control sample (CRM batch 142, Dickson,
 205 2010) were plotted over time, compared to the warning and control limits calculated from their mean and standard deviation
 206 (for details please see Dickson et al., 2007) as well as the certified value of the CRM. To compute a range control chart for the
 207 evaluation of measurement repeatability, the absolute difference between duplicate measurements of CRMs on each sampling
 208 day was calculated and plotted over time, compared to the warning and control limits calculated from their mean and standard
 209 deviation (for details see Dickson et al., 2007).

210 We used the R package *seacarb* with a Gaussian approach and an input variable pair (pH_T, TA) to calculate uncertainties for
 211 calculated CO₂ system parameters (Orr et al. 2018; Gattuso et al. 2020). The contribution of input uncertainties in nutrient
 212 concentrations and *in situ* salinity and temperature to the uncertainties in the CO2SYS-based calculations are often small (<
 213 0.1%; Orr et al. 2018) so they were not considered in our propagation. The input uncertainties of pH_T and TA were estimated
 214 based on our measurements (Table 1). Standard uncertainties include random and systematic errors. For TA, systematic errors
 215 were removed by calibrating the measured results using CRMs (see Sect. 2.5). Hence, the random error of TA, estimated by
 216 the averaged standard deviation of all the CRM measurements (4.4 μmol kg⁻¹; n = 62), was used as the standard uncertainty.
 217 For pH_T, an uncertainty of 0.01 was used as the standard uncertainty. Due to the unavailability of CRMs that correct for
 218 systematic error in pH measurements, the standard deviations of repeated measurements (0.0012; n = 377) only accounted for
 219 the random components of standard uncertainties (Orr et al. 2018). Therefore, we used 0.01 in our uncertainty propagation as
 220 an approximation of the total standard uncertainty for pH_T, which has been used in previous assessments (Orr et al. 2018).

221 **Table 1: Standard uncertainties of pH_T and TA estimated based on our measurements are denoted by u(pH_T) and u(TA). Based**
 222 **on u(pH_T) and u(TA), propagated uncertainties were estimated for each data point in R and averaged for each reported variable**
 223 **(μ), with Standard deviation (σ), minimum (min) and maximum (max) values presented. The relative percentage (%) of**
 224 **propagated standard uncertainties were calculated by dividing the propagated uncertainty by the corresponding data point and**
 225 **averaged for each reported variable (μ), with σ, min and max values presented.**

u(pH _T)	u(TA) μmol kg ⁻¹	ΔpCO ₂		ΔDIC		ΔΩ _{Ar}		ΔΩ _{Ca}	
		μatm	%	μmol kg ⁻¹	%	%	%		
0.01	μ	35.94	3.8	6.63	0.3	0.08	5.1	0.13	5.1
	σ	12.60	0.3	0.80	0.0	0.03	0.3	0.05	0.3
	min	15.07	3.2	5.88	0.3	0.04	4.4	0.07	4.4
	max	62.84	4.8	8.72	0.4	0.16	5.8	0.24	5.8

226 The air-sea flux of CO₂ (FCO₂, mmol C m⁻² d⁻¹) in the Pacific was determined based on

$$227 \quad FCO_2 = k K_0 \Delta pCO_2 \quad (1)$$

228 where k is the gas transfer velocity parameterized as a function of wind speed, K_0 is the solubility of CO_2 in seawater dependent
229 on *in situ* salinity and temperature (Weiss, 1974), and $\Delta p\text{CO}_2$ is the difference between $p\text{CO}_2$ in the surface water and in the
230 atmosphere (Wanninkhof 2014). Wind data were averaged over 2 sampling days for the sampling location from a satellite-
231 derived gridded dataset (GLDAS Model, near surface wind speed, 0.25 x 0.25 degrees, 3 hour temporal resolution, 12.375° to
232 11.875°S, 77.375° to 76.875°W), obtained from NASA Giovanni (Rodell et al., 2004; Beaudoin and Rodell, 2020). *In situ*
233 salinity and temperature were obtained from the CTD casts (see Sect. 2.4). Calculated $p\text{CO}_2$ based on (pH_T , TA) and an
234 estimated atmospheric $p\text{CO}_2$ of 405.22 μatm (referenced to year 2017, NOAA/GML) were used in the air-sea flux estimation.

235 **3 Results**

236 **3.1 Responses of surface layer nutrient concentrations**










237 The OMZ-influenced water masses were collected from two locations and added to the mesocosms to simulate an upwelling
238 event (see Sect. 2.3). The two water masses were named “Low DIN” and “Very low DIN” respectively based on their DIN
239 concentrations (Table 2). Both water masses shared similar silicic acid (Si) and phosphate (PO_4^{3-}) concentrations but differed
240 in DIN concentration. The “Low DIN” water had a DIN concentration of 4.3 $\mu\text{mol L}^{-1}$, 14 times as high as that of the “Very
241 low DIN” water (0.3 $\mu\text{mol L}^{-1}$; Table 2).

242 **Table 2: Inorganic nutrient concentrations of the two collected deep-water masses. Please note that DIN is the sum of nitrate, nitrite**
 243 **and ammonium. P is phosphate. Si is silicic acid. Color codes denote the two water masses and are applied to the mesocosms treated**
 244 **with respective water masses in the following figures and tables.**


Water mass	Si ($\mu\text{mol L}^{-1}$)	DIN ($\mu\text{mol L}^{-1}$)	PO_4^{3-} ($\mu\text{mol L}^{-1}$)	N: P ratio (mol: mol)
Low DIN	19.6	4.3	2.5	1.7
Very low DIN	17.4	0.3	2.6	0.1

245 On day 10 before OMZ water addition, the average surface DIN concentration of the two treatment groups were similar (3.4
 246 $\mu\text{mol L}^{-1}$), but lower than that in the Pacific (9.8 $\mu\text{mol L}^{-1}$; Table 3). Surface layer DIN concentration in the mesocosms ranged
 247 between 2.0 and 6.0 $\mu\text{mol L}^{-1}$ before OMZ water addition (Fig. 2a). The addition of OMZ water elevated surface DIN in the
 248 “Low DIN” mesocosms to 3.6-6.4 $\mu\text{mol L}^{-1}$ but lowered that in the “Very low DIN” to 0.9-2.0 $\mu\text{mol L}^{-1}$. The average surface
 249 DIN concentration in the “Very low DIN” decreased to 1.6 $\mu\text{mol L}^{-1}$ while the “Low DIN” slightly increased to 4.7 $\mu\text{mol L}^{-1}$
 250 (Table 3), followed by a sharp depletion on day 16 except for M3. M3 received the highest input of DIN (6.4 $\mu\text{mol L}^{-1}$) and
 251 was not depleted until day 24. Despite several small peaks in M3, M4, M5 and M6 ($\leq 1.6 \mu\text{mol L}^{-1}$), surface DIN concentration
 252 in the mesocosms were at around limits of detection (LODs: $\text{NH}_4^+ = 0.063 \mu\text{mol L}^{-1}$, $\text{NO}_2^- = 0.054 \mu\text{mol L}^{-1}$, $\text{NO}_3^- = 0.123$
 253 $\mu\text{mol L}^{-1}$) most of the time after depletion. A slight rise could be observed from day 44 towards the last sampling day (day 48).
 254 In the Pacific, surface layer DIN concentration was mostly greater than 5 $\mu\text{mol L}^{-1}$ (except on day 16 and 18) and became
 255 considerably higher during the second half of the experiment ($> 10 \mu\text{mol L}^{-1}$ for day 26-44; Fig. 2a).

256 **Table 3: DIN concentration ($\mu\text{mol L}^{-1}$) in the surface layer of each mesocosm (M1-M8) and the average DIN concentration ($\mu\text{mol L}^{-1}$)**
 257 **for each treatment (“Low DIN” and “Very Low DIN”, n = 4) before (t10) and after deep water addition (t13). The DIN**
 258 **concentration in the surface Pacific water is also shown. Color codes and symbols denote the respective mesocosm in the following**
 259 **figures.**

	M1	M2	M3	M4	M5	M6	M7	M8	Low DIN	Very Low DIN	Pacific
t10	3.7	2.2	5.0	3.3	3.9	3.4	3.2	2.6	3.4 \pm 1.2	3.4 \pm 0.5	9.8
t13	1.8	3.6	6.4	2.0	1.6	4.7	4.0	0.9	4.7 \pm 1.3	1.6 \pm 0.5	9.2
											

260 Surface layer PO_4^{3-} concentrations in the mesocosms initially ranged between 1.1 and 1.5 $\mu\text{mol L}^{-1}$ and were elevated by OMZ
 261 water addition to around 1.9 $\mu\text{mol L}^{-1}$ (Fig. 2b). Thereafter, PO_4^{3-} exhibited a slow but steady decline until the end of the study
 262 with a slightly higher decrease in “Low DIN” mesocosms (blue symbols; Fig. 2b). Throughout the study, PO_4^{3-} in the
 263 mesocosms was never lower than 1.1 $\mu\text{mol L}^{-1}$. Surface layer PO_4^{3-} in the Pacific was generally higher, fluctuating between
 264 1.4 and 2.9 $\mu\text{mol L}^{-1}$. In the mesocosms, enhanced chl *a* concentrations were observed at depths shallower than 5 m and below
 265 15 m before OMZ water addition (Fig. 2c). Following OMZ water addition, a chl *a* maximum occurred at ~ 10 m and persisted
 266 until day 40, except for M3 and M4 with a 1-week delayed increase in the former and a lack of bloom in the latter (Fig. 2c).
 267 After day 40, chl *a* concentrations in all mesocosms (except for M4) increased to 12-38 $\mu\text{g L}^{-1}$ with a bloom occurring in 0-10
 268 m (Fig. 2c) Throughout the study, a chl *a* maximum was continuously observed above 10 m in the Pacific (Fig. 2c).



269

270 **Figure 2** Temporal dynamics of depth-integrated surface DIN concentration (a), PO_4^{3-} concentration (b) and vertical distribution of
 271 chl a concentration determined by CTD casts (c). The black solid lines on top of the coloured contours represent the average values
 272 over the entire water column, with the corresponding additional y-axes on the right. The vertical white lines represent the day when
 273 OMZ water was added to the mesocosms. Abbreviation: OWA, OMZ
 274 water addition. SA, salt addition. Dataset is available at <https://doi.pangaea.de/10.1594/PANGAEA.923395> (Bach et al., 2020b).

275 3.2 Temporal dynamics of carbonate chemistry

276 Before OMZ water addition, surface layer pH_T in the mesocosms ranged between 7.80-7.94 with a slight decline by ~ 0.1 over
 277 time (Fig. 3a). The initial surface layer TA ranged between 2,310 and 2,330 $\mu\text{mol kg}^{-1}$ (Fig. 3b; day 3-12). Surface layer pCO_2
 278 and DIC ranged from 541 to 749 μatm and 2,119 to 2,180 $\mu\text{mol kg}^{-1}$, respectively (Fig. 3c, d).

279 The two collected OMZ-water masses shared similar carbonate chemistry properties despite the differences in DIN
 280 concentrations. In both water masses, pH_T was ~ 7.48 , DIC was $\sim 2,305\text{-}2,310 \mu\text{mol kg}^{-1}$, TA was $\sim 2,337 \mu\text{mol kg}^{-1}$, and pCO_2
 281 was between 1,700 and 1,780 μatm (Table 4).


282 **Table 4: The *in situ* pH_T , TA, DIC, pCO_2 , Ω_{Ar} and Ω_{Ca} of the two collected OMZ-water masses.**

Water mass	pH_T	TA ($\mu\text{mol kg}^{-1}$)	DIC ($\mu\text{mol kg}^{-1}$)	pCO_2 (μatm)	Ω_{Ar}	Ω_{Ca}
Low DIN	7.49	2336.5	2305.4	1707.5	0.90	1.38
Very low DIN	7.47	2338.2	2312.1	1775.3	0.87	1.34

283 Surface DIC and pCO_2 were elevated from $\sim 2,150 \mu\text{mol kg}^{-1}$ and $\sim 600 \mu\text{atm}$ to $\sim 2,200 \mu\text{mol kg}^{-1}$ and $\sim 900 \mu\text{atm}$ (except M7)
 284 by OMZ water addition, respectively, without distinct differences between the two treatments (Mann-Whitney U-Test, $p >$
 285 0.05 ; Fig. 3c). Following OMZ water addition, surface pCO_2 in the mesocosms decreased quickly and reached minima at 340-
 286 500 μatm (except M3 and M4) on day 24 and 26. These minima corresponded with DIC minima at 2,040-2,110 $\mu\text{mol kg}^{-1}$ and
 287 pH_T maxima at 7.9-8.1 (except M3 and M4; Fig. 3c, d). After reaching the minima, surface layer pCO_2 exhibited a steady
 288 increase to 410- 680 μatm from day 24 to day 38 and later declined in M3, M5, and M7 while the rest remained relatively
 289 stable until day 42 (Fig. 3c). Interestingly, and unlike the other mesocosms, after OMZ water addition, pCO_2 in M3 steadily
 290 declined from 928 to 342 μatm until the end of the experiment while that in M4 remained constantly higher than the other
 291 mesocosms ($> 700 \mu\text{atm}$), with a slightly decreasing trend to 645 μatm towards the end of the study (Fig. 3c).


292 In the Pacific, much lower surface pH_T and higher surface pCO_2 and DIC were observed compared to the mesocosms, with an
 293 average of 7.7 (7.6-7.8), 1,078 μatm (775 – 1358 μatm) and 2,221 $\mu\text{mol kg}^{-1}$ (2173 – 2269 $\mu\text{mol kg}^{-1}$; minimum to maximum
 294 range in parenthesis; Fig. 3c, d), respectively. TA in the Pacific was initially similar to that in the mesocosms, fluctuating
 295 between 2,310 and 2,330 $\mu\text{mol kg}^{-1}$, and later decreased to $\sim 2,310 \mu\text{mol kg}^{-1}$ for the rest of the study.

296 Surface waters in the mesocosms and the Pacific were always saturated with respect to calcite and aragonite throughout the
 297 entire experimental period, with lower values observed in the Pacific (Fig. 4a, c). Bottom waters in the mesocosms and Pacific
 298 were always saturated with respect to calcite during the experiment (Fig. 4b) while bottom waters in the Pacific were
 299 undersaturated with respect to aragonite before day 13 (0.88-0.99) and had Ω_{Ar} values slightly above 1.0 for the rest of the
 300 study period (Fig. 4d).



301

302 **Figure 3** Temporal dynamics of measured depth-integrated surface pH_r (a) and TA (b), and calculated pCO₂ (c) and DIC (d). The
 303 error ribbons present measurement and propagated standard uncertainties of the calculations, respectively. Color codes and
 304 symbols denote the respective mesocosm. Abbreviation: OWA, OMZ water addition. SA, salt addition.




305

306 **Figure 4** Temporal dynamics of depth-integrated surface calcite saturation state (a), bottom calcite saturation state (b), surface
 307 aragonite saturation state (c), and bottom aragonite saturation state (d) in the mesocosms and the surrounding Pacific. The error
 308 ribbons present the propagated standard uncertainties of the calculations. When $\Omega > 1$ (above red dashed line), seawater is
 309 supersaturated for calcium carbonate. When $\Omega < 1$ (below red dashed line), seawater is under-saturated for calcium carbonate.
 310 Color codes and symbols denote the respective mesocosm. Abbreviation: OWA, OMZ water addition. SA, salt addition.

311 3.3 Air-sea CO₂ fluxes in the Pacific

312 Positive FCO₂ values indicate CO₂ outgassing from the surface waters to the atmosphere, while negative values indicate a CO₂
 313 flux from the atmosphere to the ocean. The air-sea CO₂ flux in the Pacific was constantly positive throughout our study,
 314 fluctuating from 4.2 to 14.0 mmol C m⁻² d⁻¹ over time (Fig. 5a). The minima of FCO₂ occurred on day 26 and 30, while the
 315 maximum occurred on day 32 when near surface wind was the highest (2.89 m s⁻¹; Fig. 5b), corresponding to the minima and
 316 maxima of surface pCO₂. Co-occurring with a decrease in surface temperature to below 19°C after day 36 (Fig 5c), FCO₂
 317 slightly declined from ~10 to ~6 mmol C m⁻² d⁻¹ (Fig. 5a). FCO₂ was positively correlated with near surface wind speed (R² =
 318 0.4). No correlation was found between FCO₂ and temperature (R² = 0).



319

320 **Figure 5** Temporal dynamics of surface air-sea CO_2 flux (a), near surface wind speed (b) and surface temperature (c) in the Pacific.
 321 $FCO_2 > 0$ (above red dashed line) indicates CO_2 outgassing from the sea surface to the atmosphere. $FCO_2 < 0$ (below red dashed
 322 line) indicates a CO_2 flux from the atmosphere to the sea.

323 4 Discussion

324 4.1 Quality control and propagated uncertainties


325 To compare the sensitivity of different calculated variables to uncertainties in the input variables, the propagated uncertainties
 326 were averaged for each calculated variables, reported in numerical values and percentages relative to the calculated values of
 327 each variable (Table 1). Among the 4 reported variables, Ω_{Ca} and Ω_{Ar} were the most sensitive to uncertainties in pH_T and TA
 328 with an average uncertainty of 5.1%. This adds ambiguity to whether the bottom water (10-17 m for day 3-28; 12.5-17 m for
 329 day 29-50) in the Pacific was undersaturated with respect to aragonite when Ω_{Ar} was oscillating near 1 (Fig. 4d). The
 330 propagated uncertainty in pCO_2 was slightly lower (3.8%) while DIC was the least sensitive (0.3%).

331 We examined the internal consistency between DIC measurements and calculations. DIC was measured from day 3 until the
 332 malfunction of the instrument on day 10. In total, 53 sets of measured DIC and calculated DIC (from measured pH_T and TA)
 333 values were obtained from day 3 to day 10 and compared to test their consistency (Fig. 6a). The calculated DIC values were

334 generally in agreement with the measured values ($R^2 = 0.985$, $p < 0.005$), showing that the calculations made an overall good
335 prediction for the measured DIC values. The average of the residuals (calculated DIC– measured DIC) was $-8.27 \pm 6.9 \mu\text{mol}$
336 kg^{-1} , indicating an underestimation of calculated DIC. This result is consistent with a previous observation of underestimated
337 calculated DIC (pH_T , TA) compared with measured DIC when applying the same set of constants ($-6.6 \pm 7.9 \mu\text{mol kg}^{-1}$;
338 Raimondi et al., 2019). The reasons for such underestimation have not been addressed in previous studies and remain unclear.
339 No significant relationships with input variables pH_T and TA ($R^2 = 0.12$ for both) and temperature ($R^2 = 0.30$) were found in
340 the DIC residuals (salinity remained the same from day 3 to day 10). The lack of correlation with pH_T and TA indicated that
341 the underestimation in calculated DIC was not a result from changes in pH_T and TA. Although dissociation constants are
342 known to be salinity- and temperature-dependent, the lack of correlation between DIC residuals and temperature may be
343 attributed to the relatively narrow ranges of temperature in the mesocosms ($17.9\text{-}20.9^\circ\text{C}$ from day 3-10). The offsets were
344 typically larger at lower temperatures (e.g., samples from the Arctic, Chen et al. 2015).

345 To assess the quality of carbonate chemistry measurements in this study, the stability and performance of measurements were
346 evaluated. The standard deviation of triplicate pH_T measurements varied up to 0.003 with an average of 0.0012 throughout the
347 whole experiment (Fig. 6b). The average standard deviation was in agreement with reported analytical precisions of pH (0.003,
348 Orr et al. 2018; 0.002, Raimondi et al., 2019; Ma et al., 2019).

349 For TA, triplicate measurements of CRM distributed to before and after the sample measurements were carried out on each
350 measuring day to monitor the stability of the measurement process and the performance of the system. Based on the offsets, a
351 correction factor was applied to the measured values of samples on each sampling day to calibrate for instrument drift. As
352 shown in Fig. 6c, 90.5% of the measured TA values of CRM fell between warning limits (UWL and LWL) with one data point
353 falling outside the control limits (UCL and LCL), overall suggesting a relatively stable measurement system. The average
354 measured TA was $2209.9 \mu\text{mol kg}^{-1}$, which was $17.69 \mu\text{mol kg}^{-1}$ lower than the certified concentration of the CRM (2227.59
355 $\mu\text{mol kg}^{-1}$), indicating a relatively poor accuracy (compared to the suggested bias of less than $2 \mu\text{mol kg}^{-1}$; Dickson et al., 2003;
356 Dickson et al., 2007). The poor accuracy could be attributed to the fact that the concentration of the acid titrant was not checked
357 after being prepared, as suggested in the protocol (Dickson et al., 2003). A range control chart was computed based on duplicate
358 measurements of CRM made prior to the sample measurements on each sampling day to evaluate the consistency of the offset
359 between measured and certified TA values over the course of the study (Fig. 6d; Dickson et al., 2007). The absolute difference
360 (range) between the repeated CRM measurements was on average $1.4 \mu\text{mol kg}^{-1}$. All the range values fell below the UWL
361 ($3.50 \mu\text{mol kg}^{-1}$; Fig. 6d), suggesting a relatively good precision of the measurement system.



362

363 **Figure 6** Comparison of calculated values of DIC (pH_T , TA) and measured values (a). The black line is the regression line, with the
 364 corresponding equation and R^2 shown in the top-left corner. The blue dashed line shows the regression line forced through the
 365 origin. Standard deviations of all the triplicate pH_T measurements on each sampling day over the study period. Orange dashed line
 366 shows the average ($n = 377$) of the standard deviations (b). TA values of CRM measurements on each sampling day over the study
 367 period. Orange dashed line shows the average ($n = 62$) of the measured values and green dashed line indicates the certified value of
 368 the CRM (c). The absolute difference in TA values between duplicate CRM measurements (range) on each sampling day over the
 369 study period. Orange dashed line shows the average ($n = 21$) of the ranges (d). Abbreviation: UCL, upper control limit. UWL, upper
 370 warning limit. LWL, lower warning limit. LCL, lower control limit.

371 4.2 CO_2 responses to the simulated upwelling event

372 At the beginning of the experiment, surface pCO_2 levels in the mesocosms were $>500 \mu\text{atm}$ (Fig. 3c). This suggests that we
 373 initially enclosed an upwelled water mass that was enriched with respiratory CO_2 . The addition of OMZ water with high
 374 concentrations of CO_2 to the mesocosms reduced the surface pH_T by 0.1-0.2 and increased the surface pCO_2 to $>900 \mu\text{atm}$
 375 (except for M7, which was $819.4 \mu\text{atm}$ on day 13). The simulated upwelling substantially reduced the variability in CO_2
 376 between mesocosms because OMZ water addition replaced $\sim 20 \text{ m}^3$ of seawater in each mesocosm (out of $\sim 54 \text{ m}^3$). The
 377 enhanced pCO_2 level is comparable with our observations in the ambient Pacific water ($>775 \mu\text{atm}$; Fig. 3c). These values also

378 agree with reported observations for our study area in 2013 ($>1,200 \mu\text{atm}$ in the upper 100 m and $> 800 \mu\text{atm}$ at the surface;
379 Bates, 2018).

380 In the days after OMZ water addition, surface pCO_2 in the mesocosms dropped near or below the atmospheric level (405.22
381 μatm , NOAA/GML) with a decline in DIC by $\sim 100 \mu\text{mol kg}^{-1}$ (except M3 and M4; Fig. 3c, d). The declining pCO_2 could be
382 partially attributed by CO_2 outgassing due to a high CO_2 gradient from the sea surface to the air. Due to a rare coastal El Nino
383 event (Garreaud, 2018), the CO_2 loss process may have been enhanced by a rapid surface warming ($19.8\text{-}21.0^\circ\text{C}$ from day 14
384 to 36; Fig. 5) which reduced surface CO_2 solubility (Zeebe and Wolf-Gladrow, 2001). However, air-sea gas exchange could
385 not explain surface CO_2 under-saturation in relation to the atmosphere, as observed in response to OMZ water addition in some
386 mesocosms (Van Geen et al., 2000; Friederich et al., 2008; Fig. 3c). Biological production has typically one to four times
387 greater impacts on CO_2 drawdown than air-sea gas exchange in the equatorial Pacific where surface waters are exposed to
388 local wind stress (Feely et al., 2002). This interpretation is supported by the continuously high DIC in M4 where photosynthetic
389 biomass build-up was substantially lower (Fig. 3d). Hence, the depletion of nutrients (Fig. 2a, b) and increase in chl *a*
390 concentration (Fig 2c; Bach et al., 2020a) strongly suggest that the loss of DIC (except M4) was primarily driven by biological
391 uptake and phytoplankton growth. Nevertheless, it is difficult to dissect how much CO_2 was outgassed and how much was
392 taken up photosynthetically as we did not measure air-sea gas exchange in the mesocosms (please note that equations from
393 Wanninkhof, (2014) are not applicable for mesocosms; Czerny et al., 2013b). In previous mesocosm studies, N_2O addition has
394 been a common practice to monitor air-sea gas exchange in the mesocosms (Czerny et al., 2013b). However, this was not
395 carried out in our study because it would have interfered with ^{15}N label incubations to determine N loss processes (Schulz et
396 al., 2021). Due to high variability of DOC data and the poorly constrained gas exchange of CO_2 , C budgeting often comes with
397 high uncertainties and large errors, even for a relatively simple dataset (Czerny et al., 2013b; Boxhammer et al., 2018). It
398 becomes even more difficult for the current dataset because the water column was not homogeneously mixed like in previous
399 studies. With the inability to estimate CO_2 gas exchange, it was impossible to calculate a reasonable C budget for this study.

400 Before OMZ water addition, dissolved inorganic N:P ratios in the mesocosms ranged from 1.6 to 3.5 (data not shown),
401 indicating N is the limiting nutrient in the water column (Bach et al., 2020a). Not surprisingly, the uptake of DIC was higher
402 in the “Low DIN” mesocosms which received more input of DIN from OMZ water addition, with on average $41.0 \mu\text{mol kg}^{-1}$
403 higher drawdown compared to the “Very Low DIN” from day 13 to day 24 (excluding M3 and M4; Mann-Whitney U-Test, p
404 $= 0.05$; Table S1). This observation agrees with the general expectations that addition of limiting nutrients to water column
405 should enhance biological biomass build-up. Such differences in DIC uptake, however, were not reflected in the build-up of
406 particulate organic carbon (POC) in the mesocosms (excluding M3 and M4; Mann-Whitney U-Test, $p > 0.1$). As mentioned
407 above, the differences in OMZ-water DIN between the two treatments were minor and hence, their potential to trigger treatment
408 difference were small. Also, *A. sanguinea* was persistent in the water column in the mesocosms (except M4 where it never
409 bloomed), retaining the biomass in the water column as, not sinking out until the very end of the experiment (Bach et al.,
410 2020). Due to the developing N-limitation after the biomass build-up there much of the consumed DIC could have been
411 channelled to dissolved organic carbon (DOC) pool. Indeed, we observed a pronounced increase in DOC following OMZ
412 water addition (except for M4; Igarza et al., in prep, 2021). The increase in DOC may be attributed to extracellular release by
413 phytoplankton due to nutrient limitation, or cellular lysis of phytoplankton cells by bacteria (Mykkestad 2000; Igarza et al., in
414 prep, 2021).

415 After day 24, variability in carbonate chemistry between individual mesocosms increased, with a general trend of recovering
416 from CO_2 -undersaturated conditions during the peak of the bloom (except for M3 and M4; Fig. 3c). One factor that may have
417 controlled the differences in CO_2 increase are the mesocosm-specific phytoplankton succession patterns. A shift from a diatom-
418 dominated community to a dominance of dinoflagellates (in particular *Akashiwo sanguinea*) occurred when DIN was

419 exhausted, which was absent in M3 and M4 (Bach et al., 2020a). The different succession patterns in the plankton community
420 are the most likely explanation why M3 and M4 behaved differently from the others in terms of surface layer productivity, and
421 hence carbonate chemistry. Although the rate of DIN depletion in M3 and M4 were similar to the others, the reduction in pCO₂
422 in M3 experienced a 1-week delay which is consistent with the delayed build-up of chl *a* biomass (Fig. 2c, 3c). On the other
423 hand, the pCO₂ level in M4 remained constantly elevated throughout the experiment, as of a lack of a phytoplankton bloom
424 (Fig. 2c, 3c). M4 was the only mesocosm where a *A. sanguinea* remained undetectable, whereas a delayed and reduced
425 contribution by *A. sanguinea* was observed in M3. This strongly suggests that *A. sanguinea* was a key factor driving the trend
426 of carbonate chemistry in the mesocosms.

427 Near the end of the experiment, a slight decline in pCO₂ became apparent in the mesocosms which co-occurred with a second
428 phytoplankton bloom observed in the uppermost layer of the water column (Fig. 2c, 3c). This bloom was likely fuelled by
429 surface eutrophication due to defecating sea birds. During the last part of our experiment, Inca terns (*Larosterna inca*) were
430 frequently observed to rest on the roofs and the edges of the mesocosms (Bach et al., 2020a). Bird excrements, dropped into
431 the mesocosms, are known to be enriched in inorganic nutrients, ~~especially ammonium~~ (Bedard et al., 1980). They generally
432 contain 60% water, 7.3% N and 1.5% P and the main form of N is uric acid and ammonium which makes them slightly acidic
433 (De La Peña- Lastra, 2021). Therefore, the droppings may lower the pH of the surface water. However, this should have been
434 visible in decreasing alkalinity, which was not the case. The excrements may also be high in dissolved organic nitrogen (DON),
435 evidenced by a substantial increase in DON concentrations in the mesocosm surface from day 38 onward (Igarza et al., in prep,
436 2021). The triggered surface eutrophication and phytoplankton blooms were noticeable from an accumulation of chl *a* biomass
437 above the mixed layer in the mesocosms near the end of the study (Fig. 2c). As a result, another drawdown of DIC could be
438 observed in the mesocosms except for M4, M6 and M8. While the build-up of chl *a* was comparable with that triggered by
439 OMZ water addition, the drawdown in DIC was less pronounced, potentially counteracted by the release of CO₂ by enhanced
440 respiration and remineralization following the previous bloom. Also, the second bloom occurred in the top 2 meter in the
441 mesocosms (Fig. 2c) where gas exchange can quickly replete the DIC drawdown during photosynthesis and biomass build up.

442 There are potential complications when monitoring the carbonate chemistry dynamics in an enclosed mesocosm. First, the
443 brine addition to the mesocosms could have influenced the community composition which played an important role in driving
444 the carbonate chemistry. However, the effects of the brine on the enclosed organisms have been discussed in past studies and
445 considered negligible for a salinity increase of less than 1 (~0.7 and ~0.5 increase for both salt additions in our study
446 respectively; Czerny et al., 2013a). The difference of salinity in the mesocosms from the Pacific was less than 1 throughout
447 the study so we believe salinity was not a stressor to the system (Bach et al., 2020). Second, multiple factors may introduce
448 variabilities to air-sea gas exchange in the mesocosms. With the water surface sheltered from direct wind forcing by the two-
449 meter high plastic walls of the mesocosm bag, air-sea gas exchange could be very low (but not zero due to other energy inputs
450 such as thermal convection and surrounding wave movements; Czerny et al., 2013b). On the other hand, the extensive daily
451 sampling that actively perturbing the mesocosm surface may enhance the gas exchange (Czerny et al., 2013b). With a high
452 variability of DOC data and a lack of direct measurements of CO₂ gas exchange, it is impossible to estimate a reasonable C
453 budget for our dataset (Czerny et al., 2013b; Boxhammer et al., 2018). Other complications include heterogenous initial
454 conditions in the mesocosms and light limitation due to self-shading (please see Bach et al., 2020 for an extensive discussion).
455 It would have been ideal to have control mesocosms that were treated the same way except the OMZ water addition to rule
456 out the effects induced by mesocosm manipulations and changes in hydrodynamics. This has been compromised to ensure
457 enough replicate numbers for both treatments despite the enormous cost of mesocosm experimentation. Nevertheless, a
458 previous study has examined impacts of different mixing techniques in outdoor mesocosms and found no effects on
459 phytoplankton biomass and minor effects on phytoplankton and zooplankton community composition (Striebel et al., 2013).
460 In our study, various measures were also taken to minimize the mixing (brine additions and slow casting of CTD).

461 4.3 Temporal changes of carbonate chemistry in the coastal Pacific near Callao

462 According to estimations by Takahashi et al. (2009) of global air-sea CO₂ fluxes, our study site in the equatorial Pacific (14°N-
463 14°S) is a major source of CO₂ to the atmosphere. Our near-coastal location showed high pCO₂ levels over the study period
464 (with an average of 1,078 μatm), with a sea-to-air CO₂ flux of 4.2-14.0 mmol C m⁻² d⁻¹ (Fig. 5). Compared to the criterion of
465 high CO₂ fluxes (5 mmol C m⁻² d⁻¹ or more) as proposed by Paulmier et al. (2008), our study site was a strong CO₂ source to
466 the atmosphere most of the time. These results of air-sea CO₂ fluxes were slightly higher than observations by Friederich et al.
467 (2008) along the coast of Peru in February, 2004-2006 (0.85-4.54 mol C m⁻² yr⁻¹; spatially averaged for 5-15°S along the coast
468 of Peru). This is not surprising because Friederich et al. averaged the air-sea CO₂ fluxes for 0-200 km from shore where much
469 lower pCO₂ were observed offshore (< 600 μatm), compared to our nearshore study site. The decline in pCO₂ with increasing
470 distance from shore was driven by biological uptake and outgassing to the atmosphere (Friederich et al., 2008; Loucaides et
471 al., 2012). However, when compared to the magnitude of DIC drawdown triggered by upwelling events in the mesocosms, the
472 flux of CO₂ to the atmosphere was insignificant. Assuming a 10 m mixed layer in the Pacific with a DIC concentration of
473 2,200 μmol kg⁻¹, the DIC content below 1 m² surface area would be ~22 mol m⁻². With an upper bound outgassing of 14.2
474 mmol C m⁻² d⁻¹ over 10 days (day 13-24), the loss of CO₂ would only be 0.142 mol m⁻². On the other hand, the average DIC
475 drawdown of 118.2 μmol kg⁻¹ in the “Very Low DIN” and 160.3 μmol kg⁻¹ in the “Low DIN” mesocosms (M3 and M4
476 excluded) during this period accounts for 1.18 mol m⁻² and 1.60 mol m⁻², respectively, over the same water column. This shows
477 that biological processes, drawing down CO₂, is stronger than loss by air-sea gas exchange.

478 During our study, we experienced a coastal El Niño event, which has been the strongest on record (compared to those recorded
479 in 1891 and 1925) and induced rapid sea surface warming of ~1.5°C and enhanced stratification (Garreaud, 2018). Previous
480 investigations showed that the impact of reduced upwelling on CO₂ fluxes is pronounced for upwelling areas (Feely et al.,
481 1999; Feely et al., 2002). A decline in upwelling of CO₂-enriched OMZ water results in a decrease in sea-to-air CO₂ fluxes.
482 For example, during the 1991-94 El Niño year, a total reduction in CO₂ fluxes to the atmosphere was reported for the equatorial
483 Pacific. They were only 30-80% of that of a non-El-Niño year (Feely et al., 1999; Feely et al., 2002). This is likely to be the
484 case for our study location. Most studies investigated air-sea CO₂ fluxes at larger time and regional scales (Feely et al., 1999;
485 Friederich et al., 2008; Takahashi et al., 2009). Therefore, it is difficult to conclude the magnitude of the coastal El Niño
486 influence on the local CO₂ fluxes in our study by comparing our results with previous observations. Nevertheless, our
487 observations can serve as a first evidence of carbonate chemistry dynamics in the coastal Peruvian upwelling system during a
488 coastal El Niño event. Observations of sea surface carbonate chemistry with a high temporal resolution (every-2nd-day) in near-
489 shore waters are scarce, as rarely covered by typical research expeditions in the open ocean (Takahashi et al., 2009; Franco et
490 al., 2014), especially during such an extremely rare coastal El Niño event. Comparisons of our data with previous or future
491 observations may enhance our understanding of how inorganic carbon cycling interact with extreme climate events in
492 upwelling systems.

493 CO₂-enriched OMZ water has been occasionally reported to be under-saturated with respect to aragonite (Feely et al., 2008;
494 Fassbender et al., 2011). In our study, calcite under-saturation did not occur in the mesocosms or in the Pacific (Fig. 4).
495 Aragonite under-saturation, however, was observed below the surface (10-17 m for day 3-28; 12.5-17 m for day 29-50) of the
496 Pacific at the start of the experiment (Fig. 4d), when pCO₂ was the highest (pCO₂ > 1100 μatm; Fig. 3c). Aragonite under-
497 saturation was also observed in the two deep water masses collected at deeper depths (30 m and 70 m) in the Pacific (Table
498 4). Throughout the study period, the aragonite saturation state fluctuated close to around 1 below the surface (Fig. 4d).
499 Considering the water column we sampled in the Pacific still belonged to the upper surface ocean, we could expect deeper and
500 more CO₂-enriched water in the underlying OMZ to be most likely under-saturated with respect to calcite and aragonite. Hence,
501 our observations of aragonite under-saturation in the Pacific suggest a potential risk of dissolution for marine calcifiers in

502 response to the on-going intensification and expansion of acidified OMZ water (Comeau et al., 2009; Lischka et al., 2011;
503 Maas et al., 2012).

504 **5 Conclusion**

505 Our observations in the mesocosms revealed that, following the addition of two OMZ water masses with different nutrient
506 signatures, there was a higher drawdown of DIC in response to slightly more DIN input from the OMZ water addition but no
507 difference in the build-up of POC and chl *a* (Fig. 2a, 2c, 3d). The timing of the first phytoplankton bloom was consistent with
508 a shift from a diatom-dominated community to *A. sanguinea* dominance in most mesocosms, indicating that *A. sanguinea* was
509 a key factor driving the changes in carbonate chemistry under N-limited conditions. A second phytoplankton bloom was
510 triggered by defecations of Inca terns, which eased the N limitation in the mesocosms (Fig. 2c). These findings provide
511 improved insights into the links between upwelling-induced N limitation, phytoplankton community shifts and carbonate
512 chemistry dynamics in the Peruvian upwelling system.

513 The surrounding Pacific waters at the study site were characterized by constantly high pCO₂ levels (with an average of 1,078.1
514 µatm). Most CO₂ flux estimates have been conducted in the open ocean and few studies surveyed coastal regions (Takahashi
515 et al., 2009; Franco et al., 2014). Our study site was a strong CO₂ source to the atmosphere most of the time (4.2-14.2 mmol
516 C m⁻² d⁻¹), despite a rare coastal El Niño event. However, evidence from our mesocosm experiment suggests biological
517 responses that draw down DIC can quickly turn a CO₂ source into a sink in the upwelling system. The influence of the co-
518 occurring coastal El Niño event on the local CO₂ fluxes remains unclear. Nevertheless, future carbonate chemistry fluctuations
519 are expected to be enhanced by expanding and intensifying ocean deoxygenation, as well as reducing buffer factors (Schulz et
520 al., 2019). Hence, it is essential to improve our understanding of the mechanisms driving the inorganic carbon cycling in
521 upwelling systems. As a unique dataset that characterized near-shore carbonate chemistry with a high temporal resolution
522 during a rare coastal El Niño event, our study gives important insights into the carbonate chemistry responses to extreme
523 climate events in the Peruvian upwelling system.

524 **Data availability**

525 All data will be made available on the permanent repository www.pangaea.de after publication.

526 **Author contribution**

527 UR, KGS, and LTB designed the experiment. All authors contributed to the sampling. S-MC measured, calculated, and
528 analyzed carbonate chemistry. LTB and KGS supervised the carbonate chemistry analysis. KGS carried out the CTD casts and
529 data analyses. EvdE and EPA measured and analyzed nutrients. S-MC wrote the manuscript with input from all the co-authors.

530 **Competing interests**

531 The authors declare that they have no conflict of interests.

532 **Acknowledgements**

533 This project was supported by the Collaborative Research Centre SFB 754 Climate-Biogeochemistry Interactions in the
534 Tropical Ocean financed by the German Research Foundation (DFG). Additional funding was provided by the EU project
535 AQUACOSM and the Leibniz Award 2012 granted to U.R. We thank all participants of KOSMOS Peru 2017 experiment for

536 mesocosm maintenance and sample collection and analysis. Special thanks go to the staff of IMARPE, the captains and crews
537 of *Bap Morales*, *IMARPE VI* and *B.I.C. Humboldt*, and Marina de Guerra del Perú, in particular the submarine section of the
538 Navy of Callao, and the Dirección General de Capitanías y Guardacostas for their support and assistance planning and carrying
539 out the experiment. We are thankful to Club Náutico Del Centro Naval for hosting our laboratories, office space, and support.
540 This work is a contribution in the framework of the Cooperation agreement between the IMARPE and GEOMAR through the
541 German Ministry for Education and Research (BMBF) project ASLAEL 12-016 and the national project Integrated Study of
542 the Upwelling System off Peru developed by the Direction of Oceanography and Climate Change of IMARPE, PPR 137
543 CONCYTEC. Analyses and visualizations used in this paper were produced with the Giovanni online data system, developed
544 and maintained by the NASA GES DISC.

545 References

- 546 Albert, A., Echevin, V., Lévy, M. and Aumont, O.: Impact of nearshore wind stress curl on coastal circulation and primary
547 productivity in the Peru upwelling system, *J. Geophys. Res. Oceans*, 115(C12), <https://doi.org/10.1029/2010JC006569>, 2010.
- 548 Bach, L.T., Boxhammer, T., Larsen, A., Hildebrandt, N., Schulz, K.G. and Riebesell, U.: Influence of plankton community
549 structure on the sinking velocity of marine aggregates, *Global Biogeochem. Cycles*, 30(8), 1145-1165, 2016.
- 550 Bach, L. T., Paul, A. J., Boxhammer, T., von der Esch, E., Graco, M., Schulz, K. G., Achterberg, E., Aguayo, P., Arístegui, J.,
551 Ayón, P., Baños, I., Bernales, A., Boegeholz, A. S., Chavez, F., Chavez, G., Chen, S.-M., Doering, K., Filella, A., Fischer, M.,
552 Grasse, P., Haunost, M., Hennke, J., Hernández-Hernández, N., Hopwood, M., Igarza, M., Kalter, V., Kittu, L., Kohnert, P.,
553 Ledesma, J., Lieberum, C., Lischka, S., Löscher, C., Ludwig, A., Mendoza, U., Meyer, J., Meyer, J., Minutolo, F., Ortiz Cortes,
554 J., Piiparinen, J., Sforza, C., Spilling, K., Sanchez, S., Spisla, C., Sswat, M., Zavala Moreira, M., and Riebesell, U.: Factors
555 controlling plankton community production, export flux, and particulate matter stoichiometry in the coastal upwelling system
556 off Peru, *Biogeosciences*, 17, 4831–4852, <https://doi.org/10.5194/bg-17-4831-2020>, 2020a.
- 557 ~~Bach, L. T., Paul, A., Boxhammer, T., von der Esch, E., Graco, M., Schulz, K. G., Achterberg, E. P., Aguayo, P., Arístegui,~~
558 ~~J., Ayón, P., Baños, I., Bernales, A., Boegeholz, A. S., Chavez, F. P., Chen, S. M., Doering, K., Filella, A., Fischer, M. A.,~~
559 ~~Grasse, P., Haunost, M., Hennke, J., Hernandez Hernandez, N., Hopwood, M., Igarza, M., Kalter, V., Kittu, L., Kohnert, P.,~~
560 ~~Ledesma, J., Lieberum, C., Lischka, S., Löscher, C., Ludwig, A., Mendoza, U., Meyer, J., Meyer, J., Minutolo, F., Ortiz Cortes,~~
561 ~~J., Piiparinen, J., Sforza, C., Spilling, K., Sanchez, S., Spisla, C., Sswat, M., Zavala Moreira, M., and Riebesell, U.: KOSMOS~~
562 ~~2017 Peru mesocosm study: overview data, PANGAEA, <https://doi.org/10.1594/PANGAEA.923395>, 2020b.~~
- 563 Bakun, A. and Weeks, S.J.: The marine ecosystem off Peru: What are the secrets of its fishery productivity and what might its
564 future hold?, *Prog. Oceanogr.*, 79(2-4), 290-299, <https://doi.org/10.1016/j.pocean.2008.10.027>, 2008.
- 565 Bates, N.R.: Seawater carbonate chemistry distributions across the Eastern South Pacific Ocean sampled as part of the
566 GEOTRACES project and changes in marine carbonate chemistry over the past 20 years, *Front Mar. Sci.*, 5(398), 1-18,
567 <https://doi.org/10.3389/fmars.2018.00398>, 2018.
- 568 Beaudoin, H. and M. Rodell, NASA/GSFC/HSL: GLDAS Noah Land Surface Model L4 3 hourly 0.25 x 0.25 degree V2.1,
569 Greenbelt, Maryland, USA, Goddard Earth Sciences Data and Information Services Center (GES DISC), Accessed: [2021-04-
570 08], 10.5067/E7TYRXPJKWOQ, 2020
- 571 Bedard, J., Therriault, J.C. and Berube, J.: Assessment of the importance of nutrient recycling by seabirds in the St. Lawrence
572 Estuary, *Can. J. Fish. Aquat. Sci.* 37(4), 583-588, <https://doi.org/10.1139/f80-074>, 1980.
- 573 Bockmon, E.E. and Dickson, A.G.: A seawater filtration method suitable for total dissolved inorganic carbon and pH analyses,
574 *Limnol. Oceanogr. Methods*, 12(4), 191-195, <https://doi.org/10.4319/lom.2014.12.191>, 2014.

575 Bopp, L., Resplandy, L., Orr, J.C., Doney, S.C., Dunne, J.P., Gehlen, M., Halloran, P., Heinze, C., Ilyina, T., Seferian, R. and
576 Tjiputra, J.: Multiple stressors of ocean ecosystems in the 21st century: projections with CMIP5 models, *Biogeosciences*, 10,
577 6225-6245, <https://doi.org/10.5194/bg-10-6225-2013>, 2013.

578 Capone, D.G. and Hutchins, D.A.: Microbial biogeochemistry of coastal upwelling regimes in a changing ocean, *Nat. Geosci.*,
579 6(9), 711, <https://doi.org/10.1038/ngeo1916>, 2013.

580 Carter, B.R., Radich, J.A., Doyle, H.L. and Dickson, A.G.: An automated system for spectrophotometric seawater pH
581 measurements, *Limnol. Oceanogr. Methods*, 11(1), 16-27, <https://doi.org/10.4319/lom.2013.11.16>, 2013.

582 Chavez, F.P. and Messié, M.: A comparison of eastern boundary upwelling ecosystems, *Prog. Oceanogr.*, 83(1-4), 80-96,
583 <https://doi.org/10.1016/j.pocean.2009.07.032>, 2009.

584 Chavez, F.P., Bertrand, A., Guevara-Carrasco, R., Soler, P. and Csirke, J.: The northern Humboldt Current System: Brief
585 history, present status and a view towards the future, *Prog. Oceanogr.*, 79, 95-105,
586 <https://doi.org/10.1016/j.pocean.2008.10.012>, 2008.

587 Chen, B., Cai, W.J. and Chen, L.: The marine carbonate system of the Arctic Ocean: assessment of internal consistency and
588 sampling considerations, summer 2010, *Mar. Chem.*, 176, 174-188, <https://doi.org/10.1016/j.marchem.2015.09.007>, 2015.

589 Clayton, T.D. and Byrne, R.H.: Spectrophotometric seawater pH measurements: total hydrogen ion concentration scale
590 calibration of m-cresol purple and at-sea results, *Deep Sea Res. Part I Oceanogr. Res. Pap.*, 40(10), 2115-2129,
591 [https://doi.org/10.1016/0967-0637\(93\)90048-8](https://doi.org/10.1016/0967-0637(93)90048-8), 1993.

592 Comeau, S., Gorsky, G., Jeffree, R., Teyssié, J.L. and Gattuso, J.P.: Impact of ocean acidification on a key Arctic pelagic
593 mollusc (*Limacina helicina*). *Biogeosciences*, 6(9), <https://doi.org/10.5194/bg-6-1877-2009>, 2009.

594 Cooperative Global Atmospheric Data Integration Project: Multi-laboratory compilation of atmospheric carbon dioxide data
595 for the period 1957-2018; obspack_co2_1_GLOBALVIEWplus_v5.0_2019-08-12; NOAA Earth System Research
596 Laboratory, Global Monitoring Division. <http://dx.doi.org/10.25925/20190812>, 2019.

597 Correa, D., Tam, J., Pasapera, J., Saavedra, M. and Ingunza, A.: Modeling marine circulation and hypothetical discharges in
598 Callao Bay, Peru, *Inf. Inst. Mar Peru*, 35(3), 181-192, 2008.

599 Coverly, S., Kérouel, R. and Aminot, A.: A re-examination of matrix effects in the segmented-flow analysis of nutrients in sea
600 and estuarine water, *Anal. Chim. Acta*, 712, 94-100, 2012.

601 Cyronak, T., Schulz, K.G. and Jokiel, P.L.: The Omega myth: what really drives lower calcification rates in an acidifying
602 ocean, *ICES J. Mar. Sci.*, 73(3), 558-562, <https://doi.org/10.1093/icesjms/fsv075>, 2016.

603 Czerny, J., Schulz, K.G., Krug, S.A., Ludwig, A. and Riebesell, U.: The determination of enclosed water volume in large
604 flexible-wall mesocosms "KOSMOS", *Biogeosciences*, 10(3), 1937-1941, <https://doi.org/10.5194/bg-10-1937-2013>, 2013a.

605 [Czerny, J., Schulz, K. G., Ludwig, A., and Riebesell, U.: Technical Note: A simple method for air-sea gas exchange
606 measurements in mesocosms and its application in carbon budgeting, *Biogeosciences*, 10, 1379-1390,
607 <https://doi.org/10.5194/bg-10-1379-2013>, 2013b.](https://doi.org/10.5194/bg-10-1379-2013)

608 Dale, A.W., Sommer, S., Lomnitz, U., Montes, I., Treude, T., Liebetrau, V., Gier, J., Hensen, C., Dengler, M., Stolpovsky, K.,
609 Bryant, L.D., and Wallmann, K.: Organic carbon production, mineralisation and preservation on the Peruvian margin,
610 *Biogeosciences*, 12, 1537-1559, <https://doi.org/10.5194/bg-12-1537-2015>, 2015.

611 [De La Peña-Lastra, S.: Seabird droppings: Effects on a global and local level, *Science of The Total Environment*, Feb
612 1:754:142148, <https://doi.org/10.1016/j.scitotenv.2020.142148>, 2021](https://doi.org/10.1016/j.scitotenv.2020.142148)

613 Deutsch, C., Gruber, N., Key, R.M., Sarmiento, J.L. and Ganachaud, A.: Denitrification and N₂ fixation in the Pacific Ocean,
614 *Global Biogeochem. Cycles*, 15(2), 483-506, <https://doi.org/10.1029/2000GB001291>, 2001.

615 Deutsch, C., Sarmiento, J.L., Sigman, D.M., Gruber, N. and Dunne, J.P.: Spatial coupling of nitrogen inputs and losses in the
616 ocean, *Nature*, 445(7124), 163-167, <https://doi.org/10.1038/nature05392>, 2007.

617 Dickson, A.G.: Standards for ocean measurements, *Oceanogr.*, 23(3), 34-47, <https://doi.org/10.5670/oceanog.2010.22>, 2010.

618 Dickson, A.G., Wesolowski, D.J., Palmer, D.A. and Mesmer, R.E.: Dissociation constant of bisulfate ion in aqueous sodium
619 chloride solutions to 250. degree, C, J. Phys. Chem., 94(20), 7978-7985, <https://doi.org/10.1021/j100383a042>, 1990.

620 Dickson, A.G., Afghan, J.D. and Anderson, G.C.: Reference materials for oceanic CO₂ analysis: a method for the certification
621 of total alkalinity, Mar. Chem., 80(2-3), 185-197, [https://doi.org/10.1016/S0304-4203\(02\)00133-0](https://doi.org/10.1016/S0304-4203(02)00133-0), 2003.

622 Dickson, A.G., Sabine, C.L. and Christian, J.R.: Guide to best practices for ocean CO₂ measurements, North Pacific Marine
623 Science Organization, 2007.

624 Dlugokencky, E. and Tans, P.: NOAA/GML (www.esrl.noaa.gov/gmd/ccgg/trends/)

625 Doney, S.C., Ruckelshaus, M., Duffy, J.E., Barry, J.P., Chan, F., English, C.A., Galindo, H.M., Grebmeier, J.M., Hollowed,
626 A.B., Knowlton, N. and Polovina, J.: Climate change impacts on marine ecosystems, Ann. Rev. Mar. Sci., 4, 11-37,
627 <https://doi.org/10.1146/annurev-marine-041911-111611>, 2012.

628 Douglas, N.K. and Byrne, R.H.: Achieving accurate spectrophotometric pH measurements using unpurified meta-cresol purple
629 Mar. Chem., 190, pp.66-72, 2017.

630 Echevin, V., Aumont, O., Ledesma, J. and Flores, G.: The seasonal cycle of surface chlorophyll in the Peruvian upwelling
631 system: A modelling study, Prog. Oceanogr., 79(2-4), 167-176, <https://doi.org/10.1016/j.pocean.2008.10.026>, 2008.

632 Fassbender A.J., Sabine C.L., Feely R.A., Langdon C., Mordy C.W.: Inorganic carbon dynamics during northern California
633 coastal upwelling, Cont. Shelf Res., 31(11), 1180-1192, <https://doi.org/10.1016/j.csr.2011.04.006>, 2011.

634 Fassbender, A.J., Sabine, C.L. and Feifel, K.M.: Consideration of coastal carbonate chemistry in understanding biological
635 calcification, Geophys. Res. Lett., 43(9), 4467-4476, <https://doi.org/10.1002/2016GL068860>, 2016.

636 Feely, R.A., Wanninkhof, R., Takahashi, T. and Tans, P.: Influence of El Niño on the equatorial Pacific contribution to
637 atmospheric CO₂ accumulation, Nature, 398(6728), 597-601, <https://doi.org/10.1038/19273>, 1999.

638 Feely, R.A., Sabine, C.L., Hernandez-Ayon, J.M., Ianson, D. and Hales, B.: Evidence for upwelling of corrosive " acidified"
639 water onto the continental shelf, Science, 320(5882), 1490-1492, <https://doi.org/10.1126/science.1155676>, 2008.

640 Franco, A.C., Hernández-Ayón, J.M., Beier, E., Garçon, V., Maske, H., Paulmier, A., Färber-Lorda, J., Castro, R. and Sosa-
641 Ávalos, R.: Air-sea CO₂ fluxes above the stratified oxygen minimum zone in the coastal region off Mexico. J. Geophys. Res.
642 Oceans, 119(5), 2923-2937, 2014.

643 Franz, J., Krahnemann, G., Lavik, G., Grasse, P., Dittmar, T. and Riebesell, U.: Dynamics and stoichiometry of nutrients and
644 phytoplankton in waters influenced by the oxygen minimum zone in the eastern tropical Pacific, Deep Sea Res. Part I
645 Oceanogr. Res. Pap., 62, 20-31, <https://doi.org/10.1016/j.dsr.2011.12.004>, 2012.

646 Friederich, G.E., Ledesma, J., Ulloa, O. and Chavez, F.P.: Air-sea carbon dioxide fluxes in the coastal southeastern tropical
647 Pacific, Prog. Oceanogr., 79(2-4), 156-166, <https://doi.org/10.1016/j.pocean.2008.10.001>, 2008.

648 Fuenzalida, R., Schneider, W., Garcés-Vargas, J., Bravo, L. and Lange, C.: Vertical and horizontal extension of the oxygen
649 minimum zone in the eastern South Pacific Ocean, Deep Sea Res. Part II Top. Stud. Oceanogr., 56(16), 992-1003,
650 <https://doi.org/10.1016/j.dsr2.2008.11.001>, 2009.

651 Gafar, N.A. and Schulz, K.G.: A three-dimensional niche comparison of *Emiliania huxleyi* and *Gephyrocapsa oceanica*:
652 Reconciling observations with projections, Biogeosciences, 15(11), 3541-3560, <https://doi.org/10.5194/bg-15-3541-2018>,
653 2018.

654 Galán, A., Molina, V., Thamdrup, B., Woebken, D., Lavik, G., Kuypers, M.M. and Ulloa, O.: Anammox bacteria and the
655 anaerobic oxidation of ammonium in the oxygen minimum zone off northern Chile, Deep Sea Res. Part II Top. Stud.
656 Oceanogr., 56(16), 1021-1031, <https://doi.org/10.1016/j.dsr2.2008.09.016>, 2009.

657 Garreaud, R.D.: A plausible atmospheric trigger for the 2017 coastal El Niño. Int. J. Climatol., 38, e1296-e1302,
658 <https://doi.org/10.1002/joc.5426>, 2018.

659 Gattuso, J.P., Epitalon, J.M., Lavigne, H., Orr, J., Gentili, B., Hagens, M., Hofmann, A., Mueller, J.D., Proye, A., Rae, J. and
660 Soetaert, K.: Package 'seacarb', <http://CRAN.R-project.org/package=seacarb>, Accessed 5 Jun 2020, 2020.

661 Gilly, W.F., Beman, J.M., Litvin, S.Y. and Robison, B.H.: Oceanographic and biological effects of shoaling of the oxygen
662 minimum zone, *Ann. Rev. Mar. Sci.*, 5, 393-420, <https://doi.org/10.1146/annurev-marine-120710-100849>, 2013.

663 Gruber, N.: Warming up, turning sour, losing breath: ocean biogeochemistry under global change, *Philos. Trans. A Math.*
664 *Phys. Eng. Sci.*, 369(1943), 1980-1996, <https://doi.org/10.1098/rsta.2011.0003>, 2011.

665 Hamersley, M.R., Lavik, G., Woebken, D., Rattray, J.E., Lam, P., Hopmans, E.C., Damsté, J.S.S., Krüger, S., Graco, M.,
666 Gutiérrez, D. and Kuypers, M.M.: Anaerobic ammonium oxidation in the Peruvian oxygen minimum zone, *Limnol. Oceanogr.*,
667 52(3), 923-933, <https://doi.org/10.4319/lo.2007.52.3.0923>, 2007.

668 Haugan, P.M. and Drange, H.: Effects of CO₂ on the ocean environment, *Energy Convers. Manag.*, 37(6-8), 1019-1022,
669 [https://doi.org/10.1016/0196-8904\(95\)00292-8](https://doi.org/10.1016/0196-8904(95)00292-8), 1996.

670 Hauri, C., Gruber, N., Plattner, G.K., Alin, S., Feely, R.A., Hales, B. and Wheeler, P.A.: Ocean acidification in the California
671 current system, *Oceanogr.*, 22(4), 60-71, 2009. www.jstor.org/stable/24861024. Accessed 22 June 2020.

672 Hauss, H., Franz, J.M. and Sommer, U.: Changes in N: P stoichiometry influence taxonomic composition and nutritional
673 quality of phytoplankton in the Peruvian upwelling, *J. Sea Res.*, 73, 74-85, <https://doi.org/10.1016/j.seares.2012.06.010>, 2012.

674 Hofmann, G.E., Barry, J.P., Edmunds, P.J., Gates, R.D., Hutchins, D.A., Klinger, T. and Sewell, M.A.: The effect of ocean
675 acidification on calcifying organisms in marine ecosystems: an organism-to-ecosystem perspective, *Annu. Rev. Ecol. Evol.*
676 *Syst.*, 41, 127-147, <https://doi.org/10.1146/annurev.ecolsys.110308.120227>, 2010.

677 Igarza, M., Sánchez, S., Bernales, A., Gutiérrez, D., Meyer, J., Riebesell, U., Graco, M., Bach, L., Dittmar, T., and Niggemann,
678 J.: Dissolved organic matter production during an artificially-induced red tide off central Peru. *Biogeosciences*, in preparation,
679 2021.

680 Keeling, R.F., Körtzinger, A. and Gruber, N.: Ocean deoxygenation in a warming world, *Ann. Rev. Mar. Sci.*, 2, 199-229,
681 <https://doi.org/10.1146/annurev.marine.010908.163855>, 2010.

682 Koeve, W. and Oschlies, A.: Potential impact of DOM accumulation of fCO₂ and carbonate ion computations in ocean
683 acidification experiments, *Biogeosciences (BG)*, 9, 3787-3798, <https://doi.org/10.5194/bg-9-3787-2012>, 2012.

684 Lam, P., Lavik, G., Jensen, M.M., van de Vossenberg, J., Schmid, M., Woebken, D., Gutiérrez, D., Amann, R., Jetten, M.S.
685 and Kuypers, M.M.: Revising the nitrogen cycle in the Peruvian oxygen minimum zone, *Proc. Natl. Acad. Sci.*, 106(12), 4752-
686 4757, <https://doi.org/10.1073/pnas.0812444106>, 2009.

687 Lefèvre, N., Aiken, J., Rutllant, J., Daneri, G., Lavender, S. and Smyth, T.: Observations of pCO₂ in the coastal upwelling off
688 Chile: Spatial and temporal extrapolation using satellite data, *J. Geophys. Res. Oceans*, 107(C6), 8-1,
689 <https://doi.org/10.1029/2000JC000395>, 2002.

690 Levin, L.A. and Breitburg, D.L.: Linking coasts and seas to address ocean deoxygenation, *Nat. Clim. Chang.*, 5(5), 401-403,
691 <https://doi.org/10.1038/nclimate2595>, 2015.

692 Lischka, S., Büdenbender, J., Boxhammer, T. and Riebesell, U.: Impact of ocean acidification and elevated temperatures on
693 early juveniles of the polar shelled pteropod *Limacina helicina*: mortality, shell degradation, and shell
694 growth, *Biogeosciences*, 8, 919-932, <http://dx.doi.org/10.5194/bg-8-919-2011>, 2011.

695 Loucaides, S., Tyrrell, T., Achterberg, E.P., Torres, R., Nightingale, P.D., Kitidis, V., Serret, P., Woodward, M. and Robinson,
696 C.: Biological and physical forcing of carbonate chemistry in an upwelling filament off northwest Africa: Results from a
697 Lagrangian study, *Global Biogeochem. Cycles*, 26(3), <https://doi.org/10.1029/2011GB004216>, 2012.

698 Lueker, T.J., Dickson, A.G. and Keeling, C.D.: Ocean pCO₂ calculated from dissolved inorganic carbon, alkalinity, and
699 equations for K₁ and K₂: validation based on laboratory measurements of CO₂ in gas and seawater at equilibrium, *Mar. Chem.*,
700 70(1-3), 105-119, [https://doi.org/10.1016/S0304-4203\(00\)00022-0](https://doi.org/10.1016/S0304-4203(00)00022-0), 2000.

701 Ma, J., Shu, H., Yang, B., Byrne, R.H. and Yuan, D.: Spectrophotometric determination of pH and carbonate ion concentrations
702 in seawater: Choices, constraints and consequences, *Anal. Chim. Acta*, 1081, 18-31, <https://doi.org/10.1016/j.aca.2019.06.024>,

703 2019. Maas, A.E., Wishner, K.F. and Seibel, B.A.: The metabolic response of pteropods to acidification reflects natural CO₂-
704 exposure in oxygen minimum zones. *Biogeosciences*, 9(2), 747-757, <https://doi.org/10.5194/bg-9-747-2012>, 2012.

705 Matear, R.J. and Hirst, A.C.: Long-term changes in dissolved oxygen concentrations in the ocean caused by protracted global
706 warming. *Global Biogeochem. Cycles*, 17(4), <https://doi.org/10.1029/2002GB001997>, 2003.

707 Matear, R.J., Hirst, A.C. and McNeil, B.I.: Changes in dissolved oxygen in the Southern Ocean with climate change. *Geochem.*
708 *Geophys. Geosyst.*, 1(11), <https://doi.org/10.1029/2000GC000086>, 2000.

709 McNeil, C.L. and Merlivat, L.: The warm oceanic surface layer: Implications for CO₂ fluxes and surface gas measurements,
710 *Geophys. Res. Lett.*, 23(24), 3575-3578, <https://doi.org/10.1029/96GL03426>, 1996.

711 Mehrbach, C., Culbertson, C.H., Hawley, J.E. and Pytkowicz, R.M.: Measurement of the apparent dissociation constants of
712 carbonic acid in seawater at atmospheric pressure 1, *Limnol. Oceanogr.*, 18(6), 897-907,
713 <https://doi.org/10.4319/lo.1973.18.6.0897>, 1973.

714 Messié, M. and Chavez, F.P.: Nutrient supply, surface currents, and plankton dynamics predict zooplankton hotspots in coastal
715 upwelling systems. *Geophys. Res. Lett.*, 44(17), 8979-8986, <https://doi.org/10.1002/2017GL074322>, 2017.

716 Mogollón, R. and Calil, P.H.: Modelling the mechanisms and drivers of the spatiotemporal variability of pCO₂ and air-sea
717 CO₂ fluxes in the Northern Humboldt Current System, *Ocean Modelling*, 132, 61-72,
718 <https://doi.org/10.1016/j.ocemod.2018.10.005>, 2018.

719 Montecino, V. and Lange, C.B.: The Humboldt Current System: Ecosystem components and processes, fisheries, and sediment
720 studies. *Prog. Oceanogr.*, 83(1-4), 65-79, <https://doi.org/10.1016/j.pocean.2009.07.041>, 2009.

721 Morris, A.W. and Riley, J.P.: The determination of nitrate in sea water, *Anal. Chim. Acta*, 29, 272-279,
722 [https://doi.org/10.1016/S0003-2670\(00\)88614-6](https://doi.org/10.1016/S0003-2670(00)88614-6), 1963.

723 Mullin, J. and Riley, J.P.: The colorimetric determination of silicate with special reference to sea and natural waters, *Anal.*
724 *Chim. Acta*, 12, 162-176, [https://doi.org/10.1016/S0003-2670\(00\)87825-3](https://doi.org/10.1016/S0003-2670(00)87825-3), 1955.

725 Murphy, J.A.M.E.S. and Riley, J.P.: A modified single solution method for the determination of phosphate in natural waters,
726 *Anal. Chim. Acta*, 27, 31-36, [https://doi.org/10.1016/S0003-2670\(00\)88444-5](https://doi.org/10.1016/S0003-2670(00)88444-5), 1962.

727 Mykkestad, S.M.: Dissolved organic carbon from phytoplankton, In *Mar. Chem.*, 111-148, Springer, Berlin, Heidelberg, 2000.

728 Orr, J.C., Epitalon, J.M. and Gattuso, J.P.: Comparison of ten packages that compute ocean carbonate chemistry, *Biogeosci.*
729 *Discuss.*, 12(5), 1483-1510, <https://doi.org/10.5194/bg-12-1483-2015>, 2015.

730 Orr, J.C., Epitalon, J.M., Dickson, A.G. and Gattuso, J.P.: Routine uncertainty propagation for the marine carbon dioxide
731 system, *Mar. Chem.*, 207, 84-107, <https://doi.org/10.1016/j.marchem.2018.10.006>, 2018.

732 Oschlies, A., Brandt, P., Stramma, L. and Schmidtko, S.: Drivers and mechanisms of ocean deoxygenation, *Nat. Geosci.*, 11,
733 467-473, <https://doi.org/10.1038/s41561-018-0152-2>, 2018.

734 Paulmier, A., Ruiz-Pino, D. and Garçon, V.: The oxygen minimum zone (OMZ) off Chile as intense source of CO₂ and N₂O,
735 *Cont. Shelf Res.*, 28(20), 2746-2756, <https://doi.org/10.1016/j.csr.2008.09.012>, 2008.

736 Paulmier, A., Ruiz-Pino, D. and Garçon, V.: CO₂ maximum in the oxygen minimum zone (OMZ), *Biogeosciences*, 8, 239-
737 252, <https://doi.org/10.5194/bg-8-239-2011>, 2011.

738 Pierrot, D., Lewis, E. and Wallace, D.W.R.: MS Excel program developed for CO₂ system calculations, ORNL/CDIAC-105a,
739 Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, US Department of Energy, Oak Ridge,
740 Tennessee, 2006.

741 Raimondi, L., Matthews, J.B.R., Atamanchuk, D., Azetsu-Scott, K. and Wallace, D.W.: The internal consistency of the marine
742 carbon dioxide system for high latitude shipboard and in situ monitoring, *Mar. Chem.*, 213, 49-70,
743 <https://doi.org/10.1016/j.marchem.2019.03.001>, 2019.

744 Redfield, A.C.: The influence of organisms on the composition of seawater, *The Sea*, 2, 26-77, 1963.

745 Riebesell, U., Czerny, J., Bröckel, K.V., Boxhammer, T., Büdenbender, J., Deckelnick, M., Fischer, M., Hoffmann, D., Krug,
746 S.A., Lentz, U. and Ludwig, A.: A mobile sea-going mesocosm system—new opportunities for ocean change research,
747 *Biogeosciences*, 10(3), 1835-1847, <https://doi.org/10.5194/bg-10-1835-2013>, 2013.

748 Rodell, M., P.R. Houser, U. Jambor, J. Gottschalck, K. Mitchell, C. Meng, K. Arsenault, B. Cosgrove, J. Radakovich, M.
749 Bosilovich, J.K. Entin, J.P. Walker, D. Lohmann, and D. Toll: The Global Land Data Assimilation System, *Bull. Amer.*
750 *Meteor. Soc.*, 85, 381-394, doi:10.1175/BAMS-85-3-381, 2004

751 RStudio Team (2015). RStudio: Integrated Development for R. RStudio, Inc., Boston, MA URL <http://www.rstudio.com/>.

752 Schmidtko, S., Stramma, L. and Visbeck, M.: Decline in global oceanic oxygen content during the past five decades, *Nature*,
753 542(7641), 335-339, 2017.

754 Schulz, K.G. and Riebesell, U.: Diurnal changes in seawater carbonate chemistry speciation at increasing atmospheric carbon
755 dioxide, *Mar. Biol.*, 160(8), 1889-1899, <https://doi.org/10.1007/s00227-012-1965-y>, 2013.

756 Schulz, K.G., Bellerby, R.G.J., Brussaard, C.P.D., Büdenbender, J., Czerny, J., Engel, A., Fischer, M., Koch-Klavnsen, S.,
757 Krug, S., Lischka, S. and Ludwig, A.: Temporal biomass dynamics of an Arctic plankton bloom in response to increasing
758 levels of atmospheric carbon dioxide, *Biogeosciences*, 10, 161-180, <https://doi.org/10.5194/bg-10-161-2013>, 2013.

759 Schulz, K.G., Bach, L.T., Bellerby, R.G., Bermúdez, R., Büdenbender, J., Boxhammer, T., Czerny, J., Engel, A., Ludwig, A.,
760 Meyerhöfer, M. and Larsen, A.: Phytoplankton blooms at increasing levels of atmospheric carbon dioxide: experimental
761 evidence for negative effects on prymnesiophytes and positive on small picoeukaryotes, *Front. Mar. Sci.*, 4, 64,
762 <https://doi.org/10.3389/fmars.2017.00064>, 2017.

763 Schulz, K.G., Hartley, S. and Eyre, B.: Upwelling amplifies ocean acidification on the East Australian Shelf: implications for
764 marine ecosystems, *Front. Mar. Sci.*, 6, 636, <https://doi.org/10.3389/fmars.2019.00636>, 2019.

765 [Schulz, K. G., Achterberg, E. P., Arístegui, J., Bach, L. T., Baños, I., Boxhammer, T., Erler, D., Igarza, M., Kalter, V., Ludwig,](#)
766 [A., Löscher, C., Meyer, J., Meyer, J., Minutolo, F., von der Esch, E., Ward, B. B., and Riebesell, U.: Nitrogen loss processes](#)
767 [in response to upwelling in a Peruvian coastal setting dominated by denitrification – a mesocosm approach, *Biogeosciences*,](#)
768 [18, 4305–4320, https://doi.org/10.5194/bg-18-4305-2021, 2021.](#)

769 Stramma, L., Johnson, G.C., Sprintall, J. and Mohrholz, V.: Expanding oxygen-minimum zones in the tropical oceans, *Science*,
770 320(5876), 655-658, <https://doi.org/10.1126/science.1153847>, 2008.

771 Stramma, L., Schmidtko, S., Levin, L.A. and Johnson, G.C.: Ocean oxygen minima expansions and their biological impacts,
772 *Deep Sea Res. Part I Oceanogr. Res. Pap.*, 57(4), 587-595, <https://doi.org/10.1016/j.dsr.2010.01.005>, 2010.

773 [Striebel, M., Kirchmaier, L. and Hingsamer, P.: Different mixing techniques in experimental mesocosms—does mixing affect](#)
774 [plankton biomass and community composition?, *Limnology and Oceanography: Methods*, 11\(4\), 176-186,](#)
775 [https://doi.org/10.4319/lom.2013.11.176, 2013.](#)

776 Takahashi, T., Sutherland, S.C., Wanninkhof, R., Sweeney, C., Feely, R.A., Chipman, D.W., Hales, B., Friederich, G., Chavez,
777 F., Sabine, C. and Watson, A.: Climatological mean and decadal change in surface ocean pCO₂, and net sea–air CO₂ flux over
778 the global oceans, *Deep Sea Res. Part II Top. Stud. Oceanogr.*, 56(8-10), 554-577, <https://doi.org/10.1016/j.dsr2.2008.12.009>,
779 2009.

780 Taucher, J., Bach, L.T., Boxhammer, T., Nauendorf, A., Achterberg, E.P., Algueró-Muñiz, M., Arístegui, J., Czerny, J.,
781 Esposito, M., Guan, W. and Haunost, M.: Influence of ocean acidification and deep water upwelling on oligotrophic plankton
782 communities in the subtropical North Atlantic: insights from an in situ mesocosm study, *Front Mar. Sci.*, 4, 85,
783 <https://doi.org/10.3389/fmars.2017.00085>, 2017.

784 Trenberth, K.E.: The definition of el nino, *Bull. Am. Meteorol. Soc.*, 78(12), 2771-2778, <https://doi.org/10.1175/1520->
785 [0477\(1997\)078<2771:TDOENO>2.0.CO;2, 1997.](#)

786 Uppström, L.R.: The boron/chlorinity ratio of deep-sea water from the Pacific Ocean, *Deep Sea Res. and Oceanogr. Abstr.*,
787 21, 161-162, [https://doi.org/10.1016/0011-7471\(74\)90074-6](https://doi.org/10.1016/0011-7471(74)90074-6), 1974.

788 Van Geen, A., Takesue, R.K., Goddard, J., Takahashi, T., Barth, J.A. and Smith, R.L.: Carbon and nutrient dynamics during
789 coastal upwelling off Cape Blanco, Oregon, Deep Sea Res. Part II Top. Stud. Oceanogr., 47(5-6), 975-1002,
790 [https://doi.org/10.1016/S0967-0645\(99\)00133-2](https://doi.org/10.1016/S0967-0645(99)00133-2), 2000.

791 Volk, T. and Hoffert, M.I.: Ocean carbon pumps: Analysis of relative strengths and efficiencies in ocean-driven atmospheric
792 CO₂ changes. The carbon cycle and atmospheric CO₂: natural variations Archean to present, 32, 99-110,
793 <https://doi.org/10.1029/GM032p0099>, 1985.

794 Wanninkhof, R.: Relationship between wind speed and gas exchange over the ocean revisited, Limnol. Oceanogr. Methods,
795 12(6), 351-362, <https://doi.org/10.4319/lom.2014.12.351>, 2014.

796 Weiss, R.: Carbon dioxide in water and seawater: the solubility of a non-ideal gas, Mar. Chem., 2(3), 203-215, 1974.

797 Whitney, F.A., Freeland, H.J. and Robert, M.: Persistently declining oxygen levels in the interior waters of the eastern subarctic
798 Pacific, Prog. Oceanogr., 75(2), 179-199, <https://doi.org/10.1016/j.pocean.2007.08.007>, 2007.

799 Zeebe, R.E. and Wolf-Gladrow, D.: CO₂ in seawater: equilibrium, kinetics, isotopes (No. 65), Gulf Professional Publishing,
800 2001.