

Geochemical consequences of oxygen diffusion from the oceanic crust into overlying sediments and its significance for biogeochemical cycles based on sediments of the NE Pacific

Gerard J. M. Versteegh^{1,2,3}, Andrea Koschinsky^{2,3}, Thomas Kuhn⁴, Inken Preuss^{2*}, Sabine Kasten^{1,3,5}

5 ¹Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Bremerhaven, 27570, Germany

²Jacobs University Bremen, Department of Physics and Earth Sciences, Bremen, 28759, Germany

³MARUM – Zentrum für Marine Umweltwissenschaften, Universität Bremen, Bremen, 28359, Germany

10 ⁴Bundesanstalt für Geowissenschaften und Rohstoffe, Hannover, 30655, Germany

⁵University of Bremen, Faculty of Geosciences, Bremen, 28359, Germany

*present address: GEOMAR Helmholtz Centre for Ocean Research, Kiel, 24148, Germany

Correspondence to: Gerard J. M. Versteegh (gerard.versteegh@awi.de)

Abstract. Exchange of dissolved substances at the sediment–water interface provides an important link
15 between the short–term and long–term geochemical cycles in the ocean. A second, as yet poorly
understood sediment–water exchange is supported by low–temperature circulation of seawater through
the oceanic basement underneath the sediments. From the basement, upwards diffusing oxygen and
other dissolved species modify the sediment whereas reaction products diffuse from the sediment down
into the basement, where they are transported by the basement fluid and released to the ocean. Here, we
20 investigate the impact of this ‘second’ route with respect to transport, release and consumption of
oxygen, nitrate, manganese, nickel, and cobalt on the basis of sediment cores retrieved from the Clarion
Clipperton Zone (CCZ) in the equatorial Pacific Ocean. We show that in this abyssal ocean region
characterised by low organic–carbon burial and sedimentation rates vast areas exist where the
downward and upward directed diffusive fluxes of oxygen meet so that the sediments are oxic
25 throughout. This is especially the case where sediments are thin, or in the proximity of faults. Oxygen
diffusing upward from the basaltic crust into the sediment contributes to the degradation of sedimentary
organic matter. Where the sediments are entirely oxic, nitrate produced in the upper sediment by
nitrification is lost both by upward diffusion into the bottom water and by downward diffusion into the

Gerard Versteegh 3-6-21 10:06

Gelöscht: Where the oxygen profiles do not meet, they are separated by a suboxic sediment interval characterised by Mn^{2+} in the pore–water.

fluids circulating within the basement. Where the oxygen profiles do not meet, they are separated by a suboxic sediment interval characterised by Mn^{2+} in the porewater. Where pore-water Mn^{2+} in the suboxic zones remains low, nitrate consumption is low and the sediment continues to deliver nitrate to the ocean bottom waters and basement fluid. We observe that at elevated pore-water manganese concentrations, nitrate consumption exceeds production and nitrate diffuses from the basement fluid into the sediment. Within the suboxic zone, not only manganese but also cobalt and nickel are released into the pore water by reduction of Mn oxides, diffuse towards the oxic/suboxic fronts above and below where they precipitate, effectively removing these metals from the suboxic zone and concentrating them at the two oxic/suboxic redox boundaries. We show that not only diffusive fluxes in the top part of deep-sea sediments modify the geochemical composition over time, but also diffusive fluxes of dissolved constituents from the basement into the bottom layers of the sediment. Hence, paleoceanographic interpretation of sedimentary layers should carefully consider such deep secondary modifications in order to prevent misinterpretation as primary signatures.

45 1 Introduction

The heat flux from the mantle through the lithosphere into the ocean is high where the ocean crust is young and sediment cover sparse such as at spreading zones, and the heat flux typically decreases with age as the crust cools (e.g., Williams and von Herzen, 1974; Sclater et al., 1980; Davis et al., 1999). Ventilation of the basaltic crust by seawater is an important aspect of this cooling and is predicted to occur in off-axis settings up to an age of 65 Ma (e.g., Stein and Stein, 1992; Fisher et al., 2003; Hutnak et al., 2006). Most of the ocean floor has cooled over time, and conversely, most ventilation of the basaltic crust occurs at low temperatures (Hasterok et al., 2011; Coogan and Gillis, 2018). Water percolating through the crust changes the chemistry of the basement and vice versa (Fisher and Wheat, 2010; Orcutt et al., 2013). Moreover, sediments overlying the basement may be modified by substances diffusing from the crust upwards such as oxygen whereas sedimentary products may diffuse down and interact with the crust and the fluid circulating therein (Ziebis et al., 2012; Orcutt et al., 2013; Mewes et al., 2016; Kuhn et al., 2017). Due to the vastness of the abyssal ocean floor, this low temperature ventilation may have a large impact on the global geochemical and biogeochemical cycles (e.g., Baker et al., 1991). This impact is however poorly constrained. This is to a large extent due to sediments accumulating on the basement, making it less accessible and due to low sediment permeability, they progressively insulate the crust, both thermally and hydrodynamically (Anderson et al., 1979). As a result, water can enter or leave the basement only where sediment cover remains thin or absent such as at seamounts, ridges and faults. Finally, diffusion of dissolved species from the basement into

Gerard Versteegh 26-7-21 15:39

Formatiert: Hochgestellt

Gerard Versteegh 3-6-21 10:07

Gelöscht: the basement becomes a nitrate source

Gerard Versteegh 3-6-21 10:08

Gelöscht: long periods

Gerard Versteegh 26-7-21 15:41

Gelöscht: still

Gerard Versteegh 3-6-21 10:09

Gelöscht: poor

65 | the sediments and vice versa modifies the composition of both fluid and sediment and adds an extra aspect to the geochemical and biogeochemical impact of basement ventilation (e.g., Wheat and Mottl, 2000; Fisher et al., 2003; Wheat and Fisher, 2008; Mewes et al., 2016; Kuhn et al., 2017; Hulme et al., 2019).

1.1 Sediment geochemistry.

70 | The ocean floor is subject to a continuous rain of reduced material, notably organic matter (OM). Sedimentary life harvests the energy stored in this reduced material by exploiting and catalysing the redox reactions possible, giving priority to the energetically most favourable redox pair (e.g., Froelich et al., 1979; Berner, 1980). In oxic sediments, the dominant redox process is aerobic respiration. Oxygen entering the sediment by diffusion is consumed on its way down and if sufficient OM is available dissolved oxygen becomes exhausted resulting in an oxic/suboxic interface at depth. In a steady state situation, this interface keeps pace with sedimentation and its depth relative to the sediment surface remains constant (e.g., de Lange et al., 1983; Kasten et al., 2003). From this interface downwards successively energetically less favourable redox reactions follow, provided sufficient electron donors remain (e.g., Canfield and Thamdrup, 2009; Kasten et al., 2003). However, due to successively reduced activation energies, oxidation of OM will be increasingly less complete (e.g., Tegelaar et al., 1989; Armarson and Keil, 2007; Zonneveld et al., 2010). Dissolved redox products with remaining oxidation potential may diffuse to a zone with energetically more favourable redox reaction conditions and become further oxidised at the corresponding redox front (e.g., the oxic/suboxic interface). Oxidation of immobile components requiring a high activation energy slows down or even stops, leading to long-term preservation as is the case of the more refractory OM below the oxic/suboxic interface (e.g., Arndt et al., 2013; Zonneveld et al., 2010).

85 | Less input of reactive OM or lower sedimentation rates shift the balance between oxygen diffusion and its benthic consumption deeper into the sediment (e.g. Emerson et al., 2003). The result is increased oxygen exposure time of the sediments and a deeper oxic/suboxic interface, which can reach depths of up to several meters and more (Fischer et al., 2009; Rühlemann et al., 2011; Ziebis et al., 2012; Orcutt et al., 2013; Mewes et al., 2014; d'Hondt et al., 2015; Mewes et al., 2016; Mogollón et al., 2016; Kuhn et al., 2017; Volz et al., 2018). The higher degree of OM degradation results in a more refractory residue. Conversely, OM degradation rates decrease more rapidly and to a larger extent with distance into the sediment as compared to regions with higher productivity and/or sedimentation rates (Mogollón et al., 2016).

90 | In sediment-filled basins of the western flank of the Mid-Atlantic ridge at "North Pond" and in the deep-sea sediments of the Clarion Clipperton Zone (CCZ) in the Northeast Pacific Ocean oxygen was shown to not only diffuse down across the sediment-water interface but also upward into the basal sediments from the underlying oceanic basement (Ziebis et al., 2012; Orcutt et al., 2013; Mewes et al., 2016; Kuhn et al., 2017). For this upward diffusion both oxygen concentration profiles and redox dynamics involved differ from those near the sediment surface. Oxygen concentrations of the fluids circulating in the permeable crust were demonstrated to be lower than those in the bottom water overlying the sediment surface due to partial consumption as a consequence of oxidation of reduced mineral phase contained in the crustal basalt

Gerard Versteegh 26-7-21 15:42

Gelöscht: s

Gerard Versteegh 26-7-21 15:42

Gelöscht: Mewes et al., 2016;

Gerard Versteegh 3-6-21 10:21

Gelöscht: it

Gerard Versteegh 21-7-21 08:46

Gelöscht: into the sediment and

(Fisher and Wheat, 2010; Orcutt et al., 2013; Mewes et al., 2016). The oxygen entering the sediment from below also encounters old, refractory OM and meets electron donors diffusing down from the redox-zones above. In this zone of upward diffusing oxygen, oxygen exposure time of OM decreases upward from the sediment-basement interface and may have been very long, up to equalling the age of the sediment where this upward diffusive supply of oxygen never ceased.

105 Although the upward diffusion of oxygen from the basement has only been documented for a few ocean environments, its impact on the sediment composition, OM preservation/burial and biogeochemical cycles is becoming better understood (e.g. D'Hondt et al., 2015, 2019a; Kuhn et al., 2017; Mewes et al., 2016; Morono et al., 2020; Ziebis et al., 2012). Still material and information are relatively limited. This is to a large extent a result of coring practice and/or list of parameters typically measured onboard ship. Mostly, coring devices such as piston and gravity corers are too short, or the sediment cover involved too thick to reach beyond the redox zones in the upper meters sediments and reach the deep oxic redox zone 110 induced by the upward diffusion of oxygen from seawater circulating within the basement. Furthermore, although there is a long history of deeper drilling into the ocean sediments, and through them into the ocean crust (such as in the DSDP, ODP and IODP frameworks), the instances during which ex situ oxygen measurements were performed on these long sediment cores are rare (e.g. Fischer et al., 2009; Ziebis et al., 2012; Orcutt et al., 2013; d'Hondt et al., 2013). To overcome this 115 problem, we went to the NE Pacific to the region of the CCZ in the framework of RV *Sonne* expedition SO240 (Kuhn et al., 2015) and among other pore-water and solid-phase components performed ex situ oxygen measurements on whole-round gravity cores (Fig. 1). Sedimentation rates in the CCZ are low ranging between 0.2 and 1.2 cm/ka (Mewes et al., 2014; Volz et al., 2018), thus combining a relatively thin sediment cover with a long period of sedimentation and seawater circulation within the upper basaltic crust that delivered dissolved oxygen to the overlying sediment which penetrated the entire 120 sediment column or at least to the deepest sediment recovered. Sediments with such low sedimentation rates represent about 70% of the ocean floor (Bowles et al., 2014; Mewes et al., 2016) and conversely it has been suggested that extensive and long periods of oxidation are a widespread phenomenon in abyssal ocean deposits (D'Hondt et al., 2015, 2019b). Therefore, understanding the processes associated with upward oxygen supply from the basement into overlying sediments is crucial for our understanding of global biogeochemical cycles and for properly interpreting marine sedimentary records/archives. To 125 shed more light on the exchange between the basaltic basement and the ocean as well as the role of the oceanic sediments herein, the oceanic basement and overlying sediments of the abyssal plain in the CCZ were investigated seismically, thermally and geochemically during and after RV *Sonne* cruise SO240 in the framework of the project FLUM (FLUM project Kuhn et al., 2015; Fig1).

1.2 Study area

130 The oceanic basement underlying the sediments between the Clarion and the Clipperton Fracture Zones was formed from the late Cretaceous to the Miocene (Eittrheim et al., 1992). The region hosts many seamounts and faults, which due to their steep topography are often barren of sediment on their slopes (e.g., Rühlemann et al., 2011). This allows seawater to circulate

Gerard Versteegh 3-6-21 16:29

Gelöscht: in

Gerard Versteegh 3-6-21 16:29

Gelöscht: down

Gerard Versteegh 3-6-21 16:33

Gelöscht: involved in case

Gerard Versteegh 23-6-21 11:00

Gelöscht: As yet

Gerard Versteegh 22-6-21 16:15

Gelöscht: abyssal

Gerard Versteegh 22-6-21 16:22

Gelöscht:

Gerard Versteegh 23-6-21 11:01

Gelöscht: and

Gerard Versteegh 23-6-21 11:02

Gelöscht: largely under-explored and poorly

Gerard Versteegh 22-6-21 16:19

Gelöscht:

Gerard Versteegh 22-6-21 16:20

Gelöscht: extremely

Gerard Versteegh 5-6-21 15:11

Gelöscht: and oxygen penetration from below. As such, the lower oxidation front had ample time to move up and to get into the reach of our coring devices.

Gerard Versteegh 23-6-21 11:30

Gelöscht: we hypothesise

Gerard Versteegh 23-6-21 11:30

Gelöscht: must be

Gerard Versteegh 23-6-21 11:42

Gelöscht: may be

Gerard Versteegh 5-6-21 15:29

Gelöscht: have been

Gerard Versteegh 5-6-21 15:14

Gelöscht: has been

through the oceanic crust to enter or leave (Fisher and Wheat, 2010; Kuhn et al., 2017; Mewes et al., 2016). Sediment accretion is slow (typically a few mm/ka) due to the great distance from land, relatively great water depth, a low carbon flux to the sea floor of 1.5–1.8 mg C_{org} m⁻²d⁻¹ (Lutz et al., 2007) and a sea floor below the **current** carbonate compensation depth.

155 Meter-scale oxygen penetration into the sediment results in extensive organic matter mineralisation (TOC typically < 0.2%, and mostly below 0.4% at the surface) due to oxygen exposure times in the order of hundreds of thousands of years. (e.g., Müller and Mangini, 1980; Mewes et al., 2016; Volz et al., 2018).

In this paper we concentrate on the impact dissolved **inorganic** substances migrating from the **basaltic** crust have on the overlying sediment and vice versa sediment interactions to better understand the impact of seawater ventilation of oceanic
160 crusts on global biogeochemical cycles. We build upon recent insights on the low-temperature exchange of dissolved components between the basement and the overlying sediment and role of the sedimentary record (Mewes et al., 2016; Mogollón et al., 2016; Mewes et al., 2014; Fisher and Wheat, 2010; Kuhn et al., 2017; Volz et al., 2018). We pay special attention to the upward diffusion of oxygen from the basement to obtain a better insight on its impact on sediment composition, preservation/burial of OM and its relevance for global biogeochemical cycles.

165 2 Material and Methods

The oceanic crust in the region of investigation between 11.5°N, 116.3°W and 13.1°N, 119.6°W (Fig.1) is of middle to late Eocene age (17–21 Ma) (Barckhausen et al., 2013). **Sediment thicknesses are based on sediment echo sounding (PARASOUND) and single channel seismics using a 100 m streamer chain for signal detection and GI airgun.** (Kuhn et al., 2015; Table 1). Sediments **were** recovered during RV *Sonne* cruise SO240 (Kuhn et al., 2015) by means of box-, multiple-,
170 gravity- and piston corer (Table 1; Fig 1). Cores consist generally of stiff and compact brown clays with **colour** depending on the MnO₂ contents. **For example, less MnO₂ (< concentration) resulted in a lighter brown colour and more MnO₂ (> concentration) resulted in a darker colour.** (Kuhn et al., 2015). Of these, cores 22KL, 42SL, 72SL, 81SL and 96SL were taken close by or above a fault, cores 5SL, 35SL, 51SL, 53SL and 117SL were taken close to a seamount, core 69SL was taken near both a fault and a seamount and cores 9KL, 58SL, 65SL and 108SL were taken near neither a fault nor a seamount.

175 2.1 Oxygen analyses

Pore-water oxygen in the retrieved sediments was determined ex situ onboard using amperometric Clark-type oxygen sensors according to the procedure described by **Revsbech (1989)**, Ziebis et al. (2012), Mewes et al. (2014, 2016) and Volz et al. (2018). For conversion of the measured mV to mmol/l we used Eq. (1):

$$180 \quad O_2 \text{ (mmol/l)} = V_d \times (V_{sat} - V_{Ar}) \times O_{2(sol)} \quad (1)$$

Gerard Versteegh 23-6-21 13:05

Gelösch: Sediment cover is between 0 and 90 meters thick

Gerard Versteegh 5-6-21 15:16

Gelösch: have been

Gerard Versteegh 5-6-21 15:18

Gelösch: lighter brown (

Gerard Versteegh 5-6-21 15:18

Gelösch:)

Gerard Versteegh 5-6-21 15:19

Gelösch: and darker (

Gerard Versteegh 5-6-21 15:19

Gelösch:)

Gerard Versteegh 5-6-21 15:19

Gelösch: sections

190 where V = sensor voltage at depth (V_d), under O_2 saturation (V_{sat}) and in argon (V_{ar}). $O_{2(sol)}$ is the oxygen solubility at the
given temperature (measured), salinity (measured) and pressure conditions (atmospheric pressure) and was calculated using
the equation of García and Gordon (1992). Measurements were performed in an acclimatised room kept at 4 to 6°C.

2.2 Pore–water sampling and analyses

195 Upon recovery of the sediment cores, pore water was sampled using rhizons according to the procedure described by
Seeberg–Elverfeldt et al. (2005) with a typical average pore diameter of 0.1 μm (Mewes et al., 2014). For NO_3^- analyses a
QuAAtro Continuous Segmented Flow Analyser was used (Seal Analytical). Pore–water Mn^{2+} Ni^{2+} and Co^{2+} were
determined after sample acidification with distilled HNO_3 by inductively coupled plasma optical emission spectrometry
(ICP–OES; IRIS Intrepid ICP–OES Spectrometer, Thermo Elemental) using the NIST2702 standard as reference and
following the procedures described by Mewes et al., (2014).

200 2.3 Sediment analyses

For solid–phase element analysis, 50 mg of freeze–dried and homogenised bulk sediment were digested with a mixture of
HF (0.5 ml, 40%, suprapur), HNO_3 (3 ml, 65%) and HCl (1.5ml, 30%, suprapur) at 220°C using a MARS Express
microwave system (CEM) as described by Nöthen and Kasten (2011). Major and trace elements were quantified in the
laboratories of the Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research (AWI) in Bremerhaven and
205 the Jacobs University Bremen using inductively coupled plasma mass spectrometry (ICP–MS) and inductively coupled
plasma optical emission spectrometry (ICP–OES), and at the Federal Institute for Geosciences and Natural Resources (BGR)
in Hanover by quantitative XRF (X–ray Fluorescence Spectroscopy) (Rietveld, 1967). Cross calibration, achieved by
quantifying the same elements with ICP–MS, ICP–OES and XRF for a few cores showed that the XRF measurements
reliably reproduce the changes in the elemental composition of the sediment.

210 The total carbon content (TC), the content of organic carbon (C_{org}) and the total sulphur (TS) content were determined using
LECO CS 230 Carbon–Sulfur–Analyzer at the BGR. The samples were burned in a high–frequency–oven at 1800°–2200°C
and the carbon and sulphur contents were determined using infrared–detection. Untreated samples were used to determine
TC and TS. Another sample aliquot was treated with 2N HCl at 80°C to remove carbonate–carbon and subsequently
measured for C_{org} . The difference between TC and C_{org} equals to carbonate–carbon.

215 Porosity was calculated from the water content of samples with a known volume, corrected for salinity and using a bulk
mineral density of 2.53 g cm^{-3} measured on core 22KL.

Gerard Versteegh 5-6-21 15:23

Gelöscht: has been

Gerard Versteegh 5-6-21 15:24

Gelöscht: has been

3 Results

220 The oxygen profiles of the cores analysed which are oxic throughout are highly asymmetric. From core top to bottom, oxygen shows a strong decrease in the upper 30 centimetres and below a less pronounced decrease with depth leading to an interval of minimum oxygen concentrations between 2–3 m (Fig. 2). From there in most of these cores, a **nearly** linear re-increase in oxygen concentrations is observed towards the bottom of the cores (51, 53SL, 117SL, 35SL, 81SL, 69SL, 96SL, Fig. 2). Core 5SL does not entirely follow these general patterns but shows an S-shaped oxygen profile (Fig. 2) we explain
225 by assuming non-equilibrium from recent mass transport adding sediment at the location of this core, and/or by lateral transport of oxygen and for that reason we do not take this core into further consideration.

At two sites, these upper and lower oxic zones are separated by an intermediate suboxic interval as indicated by the presence of Mn^{2+} in the pore-water (Fig. 3, 22KL, 42SL). There are five cores where the suboxic zone continues to the core bottom (9KL, 58SL, 65SL, 72SL 108SL). Of these seven cores that are partly suboxic, none has been retrieved close to a seamount, and none except for 22KL and 42SL has been retrieved from above a fault. Except for 58SL, all cores which are partly suboxic show convex pore-water Mn^{2+} profiles whereby the profile from the upper oxic/suboxic interface downward is steeper than the upward profile from the lower (extrapolated) oxic/suboxic interface. For core 22KL pore-water oxygen concentrations decrease linearly with depth. For site 58SL a downward decrease in Mn^{2+} concentration is absent, and concentrations increase to the core bottom.

235 In all but two cores pore-water NO_3^- concentrations decrease downward from the core top (Figs 2, 3). For cores 22KL and 35SL the maximum is not at the sediment surface but, slightly below in the top meter and from this maximum the downward decrease commences (Figs 2, 3). In the cores with a suboxic zone the downward decrease in NO_3^- has a concave shape in the upper part, which is followed in the suboxic zone by a less steep linear decrease (9KL, 42SL, 58SL) or linear increase (22KL). The entirely oxic cores (35SL, 53SL, 117SL, 96SL) all display a linear downward decrease in NO_3^- . Extrapolation
240 of the NO_3^- concentrations to the sediment–basement interface (see Table 1 for sediment thicknesses) provides values between 43 and 48 $\mu\text{mol/l}$ at the interface for all oxic cores and for cores 22KL and 108SL. Core 5SL, and the remaining cores with a suboxic interval (lower extrapolated NO_3^- concentrations) do not follow this general pattern.

Pore-water trace metals that co-vary in their concentrations with Mn are Co and to a lesser extent Ni (Fig. 4). In combination with the solid-phase data of these elements, we see from top to bottom (if present, Fig. 4) an oxic zone (with
245 for 22KL clearly decreasing TOC values), without free pore-water Mn and Co and with low pore-water Ni and elevated solid-phase Mn, Co, and Ni values, often with considerable scatter. Below this, a ‘transition zone’ (Fig. 4) is present with low oxygen, (and for core 22KL TOC low and constant) and solid-phase Mn, Co, Ni at the same level as in the oxic zone directly above whereas in the pore-water these elements mostly steeply decrease downward. Below this ‘transition zone’, solid-phase trace metal values are low and pore-water values are high. Below this, there is a sharp transition to oxic

250 conditions without a transition zone. Like in the upper oxic zone, pore-water Mn and Co are below detection limit and dissolved Ni is low again, while the solid-phase values of all three elements increase (Fig. 4). Dissolved Fe has not been detected in any of the pore-waters, neither in the oxic nor in the suboxic sections. Solid-phase manganese for core 22KL shows an interval with low values in the suboxic interval.

4 Discussion

255 For 63-91% of the sea floor, oxygen is high at the sediment-water interface and decreases to zero at depth (e.g., Wenzhöfer and Gludd, 2002; D'Hondt et al., 2015) due to mineralisation of organic matter (OM) or oxidation of reduced pore-water and sediment compounds. The oxygen profile is concave since oxygen is consumed over the entire upper oxic zone (e.g., Froelich et al., 1979; Cai, and Sayles, 1996; Wenzhöfer and Gludd, 2002). The cores analysed in this study show the typical features of abyssal, carbon-starved ocean sediments with sufficient deep-water ventilation. OM oxygen exposure times are long, OM degradation extensive, the remaining OM refractory and conversely sedimentary oxygen consumption rates low. As expected, we observe oxygen penetrating several meters into the sediments (see also e.g., Fischer et al., 2009). Despite this, for some cores, sufficient degradable OM remains available to exhaust oxygen at depth. Below the oxic zone and with increasing sediment depth, we observe zones of denitrification and manganese reduction. (Froelich et al., 1979; Berner, 1980; Mogollón et al., 2016; Volz et al., 2020). However, iron reduction and sulphate reduction were not observed due to the low content and reactivity of buried OM.

4.1 Pore-water oxygen

270 As a result of core retrieval by gravity- and piston-corers, the surface sediment proper is usually lost, or at least heavily disturbed. Oxygen concentrations in the bottom waters are close to 160 $\mu\text{mol/l}$ (Mewes et al., 2015, Mogollón et al., 2016, Garcia et al., 2018) and rapidly decrease to values identical to those found for the gravity- and piston-cores discussed here within the upper 5 cm (not shown). The presence of a second, upward directed diffusive flux of oxygen in many of the sediments investigated in our study and in some others suggests that oxygen is supplied from low-temperature seawater ventilation of the underlying basement. This seems to be a widespread phenomenon (Wheat and Fisher, 2008; Fisher and Wheat, 2010; Ziebis et al., 2012; Orcutt et al., 2013; Mewes et al., 2016; Kuhn et al., 2017). The interaction between the upward and downward directed oxygen profiles through time shapes the pore-water and solid-phase element profiles in the investigated sediments. Two types of geochemical settings were encountered in the investigated sediments. One type is represented by entirely oxic sediments with concave oxygen profiles (Fig. 2) and the other by cores with a suboxic zone with a second oxic zone below where oxygen concentrations increase towards the core bottom (Fig. 3). A compilation of the extension of the upward and downward pore-water oxygen and manganese profiles (Fig. 5) shows that the oxygen pore-

Gerard Versteegh 21-7-21 10:20

Gelöscht: an

Gerard Versteegh 6-6-21 13:00

Gelöscht: most

Gerard Versteegh 6-6-21 12:58

Gelöscht: oceanic sediments

Gerard Versteegh 6-6-21 12:58

Gelöscht: steeply

Gerard Versteegh 26-7-21 16:11

Gelöscht: its entire oxic length

Gerard Versteegh 5-6-21 15:26

Gelöscht: have

Gerard Versteegh 5-6-21 15:26

Gelöscht: been

Gerard Versteegh 5-6-21 15:26

Gelöscht: have been

Gerard Versteegh 6-6-21 08:07

Gelöscht: of which some show below this

290 water profiles from the core top downward are restricted to the upper three meters. The relatively constant thickness of the
upper **oxic zone of the sediments**, (mostly 2–3 m), for the oxic cores suggests that the combination of OM flux to the
sediment and benthic mineralisation, oxygen diffusion and sedimentation rate is relatively constant so that the system would
turn into suboxic conditions somewhere below the upper 2–3 m. Although sedimentation rates in the area vary (Müller and
Mangini, 1980; Mewes et al., 2014, 2016; Volz et al., 2018), they are significantly lower than the rate at which oxygen
diffuses into the sediments and thus have no significant influence on the oxygen penetration depth. This would imply that the
thickness of the sediment cover and the diffusive oxygen delivery from the basement into the overlying sediments, i.e., the
295 steepness of the upward oxygen profile in the end determines if, and to what extent, a suboxic zone is present in the
sediment.

In the case where oxygen consumption exceeds supply from above and below, two oxic/suboxic fronts are formed with the
intermediate suboxic zone having free Mn^{2+} in pore water. In a steady-state situation, both fronts represent metastable
equilibria between diffusion of reduced compounds produced in the suboxic zone to the oxidation front and consumption and
diffusion of oxygen in the oxic zones. Whereas in the upper oxic zone the concave oxygen profile indicates that oxygen is
consumed over the entire oxic zone due to the presence of more reactive OM close to the sediment surface, in the lower oxic
zone the linear oxygen profiles indicate that oxygen consumption along the profile is insignificant compared to that at the
front (Fig. 3). In combination with sediment thickness, this linearity allows us to extrapolate the profiles to the basement and
calculate basement oxygen concentrations in several occasions (Table 2, using the seismic-based estimates of sediment
300 thicknesses; Table 1). The thus obtained values, that vary strongly (from 29 $\mu\text{mol/l}$ @ 21m in 96SL to 156 $\mu\text{mol/l}$ @ 51m in
69SL), an observation also made for North Pond at the Mid–Atlantic Ridge (Ziebis et al., 2012; Orcutt et al., 2013). The
differences may be caused e.g., by differences in fluid advection velocities, fluid path–length in the basement in combination
with consumption of oxygen by reduced compounds in the basement or proximity to subsurface faults (Kuhn et al., 2017).

305 For core 69SL the projected value is very close to that of the bottom waters (160 $\mu\text{mol/l}$) and, therefore, it is suggested that
this core is taken closely to an active site of bottom water uptake by the basaltic basement. To a lesser extent this is true for
Site 51SL (131 $\mu\text{mol/l}$). For sites with lower projected O_2 concentrations at the bottom of the core we infer a longer distance
to the site of inlet or recharge and/or a more sluggish fluid circulation through the basaltic basement.

315 For the partly suboxic sites the upper oxic/suboxic front is (logically) always shallower than the depth in the entirely oxic
cores where the downward and upward O_2 fluxes meet. The sea floor in the investigated region is not a smooth abyssal plain
but characterized by abundant seamounts of different heights. The variable and dynamic interaction of laterally flowing deep
water masses with these seamounts results in bottom water currents of variable speed. As such deposition of OM raining to
the seafloor will be uneven as well, with pockets enriched and areas impoverished in OM flux. We infer that this patchiness
is the main factor causing the differences in the depth of the upper oxic/suboxic front, as also has already been discussed by
Mewes et al. (2014) and Volz et al. (2018). In the investigated region bioturbation depth is usually < 7 cm and occasionally
320 reaches twice as deep (Volz et al., 2018). This is supported by the exponential decrease in oxygen concentrations, which is

Gerard Versteegh 26-7-21 16:11

Gelösch: oxygen pore–water profile

Gerard Versteegh 26-7-21 15:49

Gelösch: ; Mewes et al., 2016;

Gerard Versteegh 26-7-21 17:58

Gelösch: water. In a steady-

Gerard Versteegh 5-7-21 14:16

Gelösch: for cores 35, 69 and 96

Gerard Versteegh 5-7-21 14:16

Gelösch: (

Gerard Versteegh 5-7-21 13:56

Gelösch: SL35, 66 $\mu\text{mol/l}$ @ 18m; 69SL,

Gerard Versteegh 5-7-21 13:53

Gelösch: 204

Gerard Versteegh 5-7-21 13:56

Gelösch: 70

Gerard Versteegh 5-7-21 13:58

Gelösch: ; 96SL, 25 $\mu\text{mol/l}$ @ 15m

Gerard Versteegh 5-7-21 13:58

Gelösch: are considerably different

restricted to the upper 4-5 cm of the accompanying multicores. As such, bioturbation is much shallower than the depth of the front (50-300 cm) and as such plays no significant role in shaping the pore-water profiles. The exception is 42SL for which suboxic conditions already prevail at the core top. However, in the accompanying multicore, the upper 25 cm measured are all oxic whereas the rapid and exponential decrease in oxygen concentrations is restricted to the upper 3.5 cm. We thus infer that for site 42SL, an unusually thick section of the upper sediments has become lost upon core recovery and also here the oxic/suboxic front was located > 25 cm below the surface, well below the zone of bioturbation.

The influence of the sediment thickness on pore-water oxygen profiles is evident from the cores from areas with a thin sediment cover, namely cores/sites 96SL, 5SL, 51SL, 53SL 117SL which are all entirely oxic (Table 1; Fig. 5) whereas longer cores appear oxic or partly suboxic. Cores 9KL, 65SL, 58SL and 108SL are the only cores from areas characterised by sediment thicknesses of > 15 m, that are neither above a fault and except for 58SL nor close to a seamount, and they all have an intermediate suboxic interval. Oxic core 81SL and partly suboxic core 108SL have similar lengths, and were taken close to each other (Fig. 1) whereby core 81SL has been taken above a fault in contrast to core 108SL. Considering the different lengths of the oxygen diffusion paths from below, we suggest that subsurface faults may facilitate upward oxygen diffusion, thus explaining the absence of a suboxic zone for core 81SL as already suggested by Kuhn et al. (2017). The presence of a fault may also explain why oxygen penetrates further upwards at site 72SL (near fault) compared to 65SL (no fault) despite both cores having been taken close to each other (Fig. 1) and having a similar sediment cover (73 and 72 m respectively; Fig 5). The presence of an underlying fault may also explain why core 69SL is entirely oxic, despite the thickness of the sediment cover. Finally, it may explain why in core 22KL the lower oxic/suboxic interface is located at a shallower sediment depth than in core 9SL. At extensional faults, pore volumes may increase, so that it is conceivable that faults increase sediment permeability. However, for these cores (> 10 m thickness), fluid advection may be neglected due to the low permeability of these sediments (Spinelli et al., 2004; Kuhn et al., 2017).

Apart from sediment thickness and the effect of faults, the oxygen concentration at the sediment-basement interface is a third variable determining the oxygen profile into the overlying sediments. This third variable depends a.o. on the flow rate, length of way of the fluids in the basement, the amount of reducible compounds in the basin, and basement temperature, whereby the latter two variables also depend on basement age (e.g., Coogan and Gillis 2018). Near sites where oxygen-rich seawater enters the basement, such as near larger seamounts, oxygen concentrations at the sediment-basement interface will be highest (e.g., Fischer et al., 2003; Wheat et al., 2004). Since oxygen is consumed within the basement, and oxygen diffuses from the basement into the sediment, further away from these sites of inflow, oxygen concentrations at the sediment-basement interface will be lower. The oxygen diffusion is not necessarily vertical only. For instance, 69SL is taken close to a ridge system with a steep slope and sediment cover may be sufficiently thin (Fig. 1) to allow fluid venting from the basement into the ocean. As such it is conceivable that lateral oxygen diffusion, perpendicular to this venting, contributes to the upward oxygen profile for 69SL resulting in sediment columns in proximity of these venting systems that may be entirely oxic, possibly overruling the effect of sediment thickness.

- Gerard Versteegh 6-7-21 10:14
Gelöscht: close to a seamount nor
- Gerard Versteegh 5-6-21 15:27
Gelöscht: have been
- Gerard Versteegh 5-7-21 14:22
Gelöscht: from sediments with similar thickness
- Gerard Versteegh 21-7-21 08:58
Gelöscht: 2
- Gerard Versteegh 21-7-21 08:57
Gelöscht: 1
- Gerard Versteegh 22-6-21 15:32
Gelöscht: a
- Gerard Versteegh 30-6-21 11:27
Gelöscht: S
- Gerard Versteegh 21-7-21 09:02
Gelöscht: , despite the thicker sediment cover for core 22KL
- Gerard Versteegh 21-7-21 09:16
Gelöscht: Considering the differences between core 72SL (partly suboxic) and core 69SL (entirely oxic) which are both taken above a fault we need to take into account
- Gerard Versteegh 21-7-21 09:10
Gelöscht: that the degree of facilitated diffusion by subsurface faults may differ so that sediments may differ in the degree to which they are suboxic
- Gerard Versteegh 22-6-21 15:34
Gelöscht: Especially i
- Gerard Versteegh 26-7-21 17:21
Gelöscht: n combination with extension
- Gerard Versteegh 22-6-21 15:35
Gelöscht: F
- Gerard Versteegh 6-6-21 16:12
Gelöscht: . This decrease in oxygen concentration per unit distance from the site of entrance will be faster if flow rates through the basement are low. Over time, the reactivity of the basement decreases as does temperature, and both may lead to a reduction of oxygen loss per unit distance through the sediment, provided a simultaneous reduction in fluid flow rates doesn't compensate this
- Gerard Versteegh 22-6-21 15:20
Gelöscht: b

Only core 58SL shows no downward decrease in its pore-water Mn^{2+} profile. It has been taken from the thickest sediment cover encountered like cores 65SL and 72SL but in contrast to these two latter cores, there are no faults nearby. So, of all cores this one has the best prerequisites for having a suboxic zone that extends deep into the sediment. We therefore suppose that also for this location upward oxygen diffusion modifies the sediment but we are not able to observe this since the pore-water Mn^{2+} maximum and subsequent downward decrease are found below the depth of the retrieved sediments/bottom of the core.

4.2 Pore-water nitrate

Usually in oxic sediments, nitrification of ammonium and diffusive loss of nitrate to the bottom water results in a nitrate maximum in the subsurface (Hensen et al., 2006). This was observed in our associated multi core data (not shown) where values increase from 36 $\mu\text{mol/l}$ at the sediment surface to reach maxima between 40 and 60 $\mu\text{mol/l}$ at 10 to 50 cm depth, and which agree with earlier studies in the region (Mewes et al., 2014, Volz et al., 2018). The absence of such a subsurface nitrate maximum for all piston- and gravity-cores but 22KL and 35SL (Fig. 2) is explained by the loss of surface sediments as a result of the core recovery procedure. OM mineralisation slows down with increasing sediment depth so does nitrate production. Furthermore, nitrate is diffusively lost to the overlying ocean bottom water (e.g., Hensen and Zabel 2000, Hensen et al., 2006), and to the deeper sediment. The subsurface pore-water nitrate maximum in the upper oxic zone of the cores reflects the balance between these three processes. For the entirely oxic cores, deeper in the sediment the linearly downward decreasing nitrate concentrations (Fig. 2) suggest the profiles to be driven by diffusion only, whereby nitrate is delivered to the basement fluid. Whereas the MUC core-tops indicate nitrate maximum concentrations of 35–40 $\mu\text{mol/l}$ for ocean bottom waters (e.g., Hensen et al., 2006), extrapolation of the deeper linear nitrate profiles to the basement-sediment interface (from seismics; Kuhn et al., 2015) results in higher concentrations (43–48 $\mu\text{mol/l}$) for the sediment-basement interface. These values differ at most only 5 $\mu\text{mol/l}$ suggesting the basement fluid to have a relatively constant nitrate concentration. It also implies that the seismic-based estimates of sediment thicknesses must be relatively accurate. Since the bottom-water nitrate concentrations are lower than those in the basement fluids we infer that there is a net flux of nitrate from the sediment to the basement. This stability of nitrate concentrations at the sediment-basement interface contrasts with the variability of inferred oxygen concentrations at this interface suggesting very different dynamics such as a much stronger influence of processes in the basement on oxygen concentrations.

The cores with a suboxic zone (Fig. 3) have a more complex nitrate chemistry. From the nitrate maximum in the upper oxic zone well into the suboxic zone, the curvature of the profile becomes increasingly less (as also observed in some cores by e.g. Mewes et al., 2014; Mogollón et al., 2016 for this region), indicating a decreasing contribution of aerobic nitrification and suboxic heterotrophic nitrification with depth. Extrapolation of the nitrate profile of core 108SL down to the anticipated

Gerard Versteegh 21-7-21 09:32

Gelöscht: The o

Gerard Versteegh 21-7-21 09:32

Gelöscht: which

Gerard Versteegh 21-7-21 09:33

Gelöscht: shows no downward decrease and as such no signs of influence from below has

Gerard Versteegh 21-7-21 09:26

Gelöscht: (82 m) and as such the sediment is 10 m thicker than for the other sites with thick sediments (65SL, 69SL, 72SL). Moreover

Gerard Versteegh 21-7-21 09:30

Gelöscht: or seamounts

Gerard Versteegh 6-6-21 13:22

Gelöscht: see

Gerard Versteegh 9-6-21 08:34

Gelöscht: We find this back

Gerard Versteegh 9-6-21 08:35

Gelöscht: s

Gerard Versteegh 9-6-21 08:36

Gelöscht: As

435 depth of the basement–sediment interface provides values estimated between 42 and 47 $\mu\text{mol/l}$, which agrees with the values
inferred from the entirely oxic cores. As such, the sediments represented by core 108SL are a source of nitrate for the
basement fluid. For core 22KL, nitrate values reach below those inferred for the basement fluid. Here, the change from a
gentle decrease to a linear increase at 9 m core depth in core 22KL marks the point where upward nitrate diffusion from the
basement equals downward diffusion and nitrate consumption. Linear extrapolation of this downward increase provides for
440 | cores/sites. However, in contrast to the oxic cores and core 108SL, the sediments represented by 22KL are a sink for
basement–fluid nitrate. For the remaining cores with a suboxic zone (9KL, 42SL, 58SL, 65SL, and 72SL), denitrification is
even more substantial and minimum nitrate values always reach below those inferred for the basement fluid. We infer for the
sediments below these cores a nitrate profile analogous to that seen in 22KL. However, whereas the nitrate minimum occurs
within 22KL, we infer that these other cores did not reach sufficiently deep to include the nitrate minimum. From this
445 | minimum, (anticipated to be located below the maximum core depth), we infer a linear downward nitrate increase towards
the basement — suggesting that the basement fluid represents a nitrate source.

There appears to be a clear linear relationship between the minimum pore–water nitrate concentration measured and the
maximum pore–water manganese concentration (Fig. 6), which roughly suggests that for each $\mu\text{mol/l}$ increase in manganese,
nitrate decreases by also one $\mu\text{mol/l}$ and this suggests a coupling between the manganese and nitrate cycles. The ratio is very
450 | different from the manganese mediated denitrification where 2.5 manganese are needed to consume one nitrate to produce
0.5 nitrogen molecule (e.g., Mogollón et al., 2016). We note that such an opposite relation can be induced by two processes:
(1) Oxidation by O_2 increases NO_3^- (from NH_4^+) and decreases Mn^{2+} (producing MnO_2) whereas (2) OM degradation
consumes NO_3^- (producing NH_4^+) and produces Mn^{2+} (from MnO_2) (see e.g., Mogollón et al., 2016). Apparently, the long–
term dynamic equilibria between these processes result in the observed net relation between the minimum pore–water nitrate
455 | concentration measured and the maximum pore–water manganese concentration in the sediments of this region.

Thus, we observe that some sediments deliver nitrate to the basement whereas others take up nitrate from the basement. We
also observe that the inferred nitrate values in the basaltic basement are higher than those in the ocean bottom waters, which
implies that after entering the basement, the basement fluids get enriched in nitrate. We infer, therefore, that overall, the
nitrate uptake from the basement by the sediment is less than the nitrate delivery to the basement by the sediment so that the
460 | basement gets enriched in nitrate relative to the ocean bottom waters. This nitrate enrichment can also be observed for the
Cocos Ridge, (Wheat and Fischer, 2008) and North Pond, Mid Atlantic Ridge (Wankel et al., 2015, Wheat et al., 2020).

465 | For the nitrate profiles in marine sediments there seem to be three situations. In the more eutrophic ocean and coastal
margins, sediments get suboxic at shallow sediment depth and in the suboxic zone nitrate is completely removed from the
sediment by denitrification (e.g., Froelich et al., 1979). As nitrate is absent from the deeper sediment, a flux of nitrate will
occur from the basaltic basement upwards, into the sediment where it gets consumed. Towards increasingly oligotrophic

Gerard Versteegh 30-6-21 11:27

Gelöscht: S

Gerard Versteegh 9-6-21 08:46

Gelöscht: reach

Gerard Versteegh 23-6-21 15:48

Gelöscht: and Fisher, 2008

470 conditions, and lower sedimentation rates, oxygen can diffuse deeper into the sediment and the depth at which nitrate is completely consumed increases as well e.g., just outside the South Pacific Gyre (SPG) core SPG-12 (D'Hondt et al., 2009), IODP 329 Site U1371 (D'hondt et al., 2015) and Western N. Atlantic KN223 Site 10 (Buchwald et al., 2018).

475 A second situation occurs where nitrate remains present throughout the sediment. It seems that this situation occurs with further decreasing productivity/sedimentation rate and denitrification in the suboxic zone such that it is insufficient to create a zone without nitrate. This occurs at e.g., at the Dorado Outcrop on the East Pacific Rise (Wheat et al., 2019). Here, manganese reduction starts when nitrate falls below 10-20 $\mu\text{mol/kg}$. In such cases, the redox succession probably remains suboxic with manganese reduction, and doesn't enter the zone of iron reduction. In the CCZ cores we observe a negative linear 1:1 relation between nitrate and pore-water manganese concentrations (Fig 6). In the East Pacific Rise (Wheat et al., 2019) dissolved manganese appears if nitrate falls below 25 $\mu\text{mol/kg}$ and nitrate becomes depleted if pore-water manganese is near 70 $\mu\text{mol/kg}$. As a consequence every μmol increase in nitrate results in a 2.8 μmol decrease in pore-water manganese. 480 As long as sedimentary nitrate remains below the values in the basaltic basement, nitrate diffuses from this basement into the sediment, a situation we observe for most of our sediments with a suboxic zone. However, as maximum dissolved manganese concentrations decrease, minimum nitrate concentrations increase and eventually, nitrate concentrations stay higher in the sediment than in the basaltic basement, establishing a net nitrate flux from the sediment into the fluid circulating in underlying basaltic basement. This is the case at site108SL and also occurs at IODP 329 Site U1370 at the edge of the SPG (D'Hondt et al., 2015). 485

A third situation evolves if the sediments become entirely oxic. The zone with dissolved manganese disappears and denitrification ceases. In the CCZ, we find for this situation nitrate concentrations up to 52 $\mu\text{mol/l}$. Mewes et al (2014) found a slightly higher threshold of 60 $\mu\text{mol/l}$ (Fig. 6 green dots). In all these cases, sediments are a nitrate source to the basaltic basement. This is also observed at North Pond, Western Flank of the Mid Atlantic Ridge, at IODP 336 sites U1382B, U1383D, U1384A (Wankel et al., 2015). In the latter study maximum nitrate concentrations increase with decreasing denitrification rates. Towards the even more oligotrophic conditions such as towards the centre of the gyre SPG, sedimentation rates decrease and the minimum oxygen concentrations increase. We infer that towards such increasingly carbon-poor settings maximum sedimentary nitrate concentrations decrease as well and so does the nitrate flux to the basaltic basement. This situation is encountered in the SPG, cores SPG 1-11 (D'Hondt et al., 2009) and IODP 329 SPG sites U1365 to U1369 (D'hondt et al., 2015) where maximum NO_3^- concentrations decrease towards the SPG centre. 495

On a global scale, from the eutrophic continental margins and upwelling zones towards the highly oligotrophic central ocean gyres, export production and sedimentation rate decrease and oxygen penetration into the sediments (from above and below) increases. As a result of nitrification in the upper sediment, there is always a net flux of nitrate from the sediment to the bottom waters and further down into the sediment. However, nitrate exchange with the basement varies. In more eutrophic sediments, denitrification is so extensive that nitrate concentrations deeper in the sediment are lower than in the basaltic basement and – as a consequence - the basement delivers nitrate to the sediment. However, towards the center of the 500

Gerard Versteegh 19-7-21 09:26

Formatiert: Tiefgestellt

Gerard Versteegh 19-7-21 09:26

Formatiert: Schriftart:Schriftfarbe:
Schwarz, Hochgestellt, Rahmen:: (Kein
Rahmen)

gyres (more oligotrophic settings), denitrification rates decrease and nitrate becomes present throughout the sediment. If denitrification in the suboxic zone further declines, minimum nitrate levels eventually exceed those in the basaltic basement and a net flux of nitrate into this basement is induced. Eventually, also the suboxic zone disappears, denitrification ceases and the sediments become entirely oxic. Towards even more oligotrophic settings, maximum nitrate levels in the sediments decrease and so does the nitrate flux from the sediment to the basaltic basement.

4.3 Pore–water manganese and iron

While in sediments with sufficient reactive OM the manganese reduction zone is followed by iron reduction, there is no evidence of this latter process in the sediments investigated in the framework of this study — i.e. the abundance of electron donors in the form of OM is too low to induce Fe reduction. In all cases where dissolved manganese is present in the pore water, we observe a maximum followed by a decrease with depth (Fig. 3). However, in those cases where we can follow this decrease further to zero, it is followed by oxic sediments and not by free iron. Consequently, we infer that in the complete region of this study insufficient reductive power is available to induce iron reduction. This has also been observed for the North Pond at the Mid Atlantic Ridge where only Hole 1382B shows pore-water dissolved manganese but no free/dissolved Fe^{2+} (see Janus Database: www-odp.tamu.edu/database/).

4.4 Nickel and cobalt in pore water and solid phase

Both pore–water and solid–phase nickel and cobalt in the suboxic cores tend to behave similar as/to manganese (Fig. 4), which agrees with the assumption that both metals are preferentially associated with the redox cycling of Mn (Aplin and Cronan, 1985; Koschinsky et al., 2001). In the suboxic zone, Mn oxide reduction also releases dissolved Ni^{2+} and Co^{2+} into the pore–water. Especially cobalt behaves similar to manganese, both clearly showing the ‘transition zone’ between the upper oxic zone and the suboxic zone (upper front). We call this a ‘transition zone’ since these elements are still present in the pore–water but already show elevated solid–phase contents. Nickel is much less bound to the oxidation fronts and seems to remain more mobile in the pore waters at higher oxygen levels than is the case for manganese and cobalt. The reason for this may be oxidation of pore–water Co^{2+} to Co^{3+} when bound to the Mn oxide phase in the sediment, which is a much more efficient scavenging process compared to simple adsorption, such as for Ni^{2+} (Murray and Dillard, 1979). Experiments with manganese oxide rich sediments also showed that the bonding of Ni is on average weaker and more easily reversible than for Co, and oxidation of Co during sorption on Mn oxides was confirmed (Kay et al., 2001).

4.5 Stability of the upper and lower redox fronts

The absence of a ‘transition zone’ for cobalt and manganese between the suboxic zone and the lower oxic zone (lower front) illustrates that the upper front is much more dynamic than the lower front. The lower front is the result of a very long route

of oxygen transport from the ocean into and through the basement. As such, changes in ocean water oxygen arrive at the lower front delayed and with lower amplitude and as a consequence of oxidation of reduced mineral phases in the crust also with lower concentrations. As a result, the lower front is much more stable and probably foremost moves slowly but steadily upward, oxidising the OM residue left behind when the upper front moved upwards with ongoing sedimentation.

535 4.6 Different aspects of the top and bottom redox zonations

Considering the above, we may add geochemical details to the conceptual model of low-temperature circulation of seawater/fluids and their components through ocean crust and diffusive exchange with overlying sediments that was presented by Fisher and Wheat (2010, Fig. 7) and schematically visualise the observed diagenetic profiles for sediments taking up nitrate from the basement (Fig. 8). The classical model of a succession of redox zones from the ocean floor into the sediment has to be extended by adding a similar reversed zonation scheme from the basement-sediment interface into the sediment for those regions where basement fluids are oxidic. However, there are some marked differences between the redox succession from the sediment surface downwards, and that from the basement and lower/overlying sediments upwards.

At the sediment surface the OM is fresh, highly reactive and subject to aerobic degradation first (Mewes et al., 2014; Mogollón et al., 2016; Volz et al., 2018). With ongoing sedimentation and concurrent upward movement of the oxidic zone, progressively less energetically favourable redox processes follow. Generally, in such situations, the reaction time is often too short for a given redox process to consume all available substrate. Reduced species that can't be oxidised by the next, energetically less favourable redox pair will remain, resulting in a surplus of reduced material that becomes fossilised at depth.

In contrast to this, oxygen penetrating from the basement into the sediment encounters the oldest sediment first. Compared to the sediment surface, reaction time is long and at the basalt-sediment interface aerobic conditions may have prevailed since the onset of sedimentation. Here, oxygen exposure time is not limiting the degree to which the substrate (OM) can be oxidised but the reactivity of the substrate. From the basement, the successively energetically less favourable redox zones follow each other upward so that at a particular depth, with time, progressively stronger oxidants may be expected, each consuming the fossilised reduced species (such as OM) that were left by the upwards migrating redox zones near the sediment surface. It should be noted that the fossilisation process may have changed the reactivity and degradability of the fossilised species over time (e.g., in sediments where sulphate reduction occurs; however, not in our sediments), natural sulphurisation may have cross-linked residual organic matter reducing its degradability by microorganisms (Zonneveld et al., 2010)). The order of redox processes upwards thus mirrors that from the sediment-water interface downwards. However, in this lower oxidic zone (from the basement upwards), oxidation times are long so that only the most refractory components remain. As such, reaction rates should be negligible so that the oxygen profile will approach linearity up to the lower oxidic/suboxidic interface above. Thus, the front position is primarily a balance between the downward migration/burial of

Gerard Versteegh 23-6-21 19:24

Gelöscht: The oxygen from the upper front has a much shorter route from its oceanic source and is thus much more subject to changes in deep-water oxygen content and/or marine production.

Gerard Versteegh 26-7-21 16:01

Gelöscht: /diffusion

Gerard Versteegh 26-7-21 16:01

Gelöscht:)

Gerard Versteegh 26-7-21 17:22

Gelöscht: Here,

Gerard Versteegh 23-6-21 13:37

Gelöscht: reaction

Gerard Versteegh 23-6-21 13:37

Gelöscht: may equal sediment

Gerard Versteegh 23-6-21 13:38

Gelöscht: age

Gerard Versteegh 9-6-21 10:15

Gelöscht: not

Gerard Versteegh 9-6-21 10:14

Gelöscht:

Gerard Versteegh 22-6-21 09:27

Gelöscht: Considering the large amount of time available, consumption in the lower oxidic zone (from the basement upwards)

Gerard Versteegh 21-7-21 10:20

Gelöscht: an

Gerard Versteegh 19-7-21 09:29

Gelöscht: In sediments that are carbon-starved and have virtually no oxidisable organic material left in the solid phase.

oxidisable species and upward oxygen flux from the basement. This is what we observe at site 22KL. Here, the downward directed profile of Mn, Co, Ni and other reduced species become linear already in the suboxic zone and meet a counter-directed linear upward oxygen diffusion profile (Fig. 8). If oxygen supply from below increases, the oxygen flux from below may meet the upper oxic zone. Thus, in the CCZ, the entirely oxic cores display a configuration where the balance between supply of reduced species to the sediment, oxygen supply and sedimentation rate is such that a suboxic zone is absent as a pool of reduced fossilised material. Here, the upward oxygen flux meets the downward flux at the point of no flux. The lower profile contributes here to the aerobic degradation at the lower end of the upper profile.

If in such entirely oxic sediments the upward oxygen flux becomes less, the oxygen minimum and the downward directed oxygen diffusion will extend downward and on its way will meet sediments with progressively increasing oxygen exposure time. Where reaction rates become very low, this downward profile will become quasilinear. Despite very low oxygen consumption in such sediments, if they are sufficiently thick, the system may become suboxic eventually. Since this suboxic zone is formed where previously the oxygen from below had exhaustively removed electron donors, manganese remobilisation may be very slow relative to diffusion, or even absent due to lack of electron donors and thus, we expect to encounter a suboxic sediment where porewater manganese is undetectable, a situation that may apply to core 81SL (Fig. 2).

The widespread interaction between basement fluid and the overlying sediment through diffusion of dissolved substances adds an important dynamic component to the geochemical interaction of sediment, basement and deep ocean (as compared to direct exchange between the surface sediment and the ocean bottom waters). The deep oligotrophic and fully oxidized ocean sediments possibly account for 30-50% of the ocean floor (D'Hondt et al., 2015). Most of the rest of the deep ocean is covered by sediments with oxic surface sediments and probably also oxic bottom sediments. All these sediments collect oxidized substances and store them for long periods of time. Fluid circulation through the basaltic basement increases the exchange between sediment and the bottom water and thus the buffer capacity.

5 Conclusions

Analyses of the concentrations of oxygen, nitrate, manganese, nickel and cobalt in pore waters and the solid-phase of sediments from the CCZ were summarised in a conceptual graph (Fig. 8) and show:

1. An upper oxygen profile of relatively constant length (2–3 m) resulting from oxygen penetrating the sediments from the ocean bottom water downwards and a lower oxygen profile of variable length resulting from oxygen penetrating from the basement upwards.
2. Where the upper and lower oxygen profiles meet, sediments are entirely oxic and organic carbon mineralisation is very low. In the absence of denitrification, nitrate produced in the sediment diffuses into the ocean bottom water and basement fluid.

Gerard Versteegh 22-6-21 09:29

Gelöscht: equal

Gerard Versteegh 24-6-21 14:58

Gelöscht: sediment–water interface

Gerard Versteegh 24-6-21 15:11

Gelöscht: where it

Gerard Versteegh 24-6-21 15:05

Gelöscht: ly refractory carbon since

Gerard Versteegh 24-6-21 15:12

Gelöscht: by the lower oxygen profile increases towards the basement

Gerard Versteegh 24-6-21 17:24

Gelöscht: the residual carbon is very refractory

Gerard Versteegh 19-7-21 09:30

Gelöscht: .

Gerard Versteegh 19-7-21 09:30

Gelöscht: M

Gerard Versteegh 26-7-21 17:33

Formatiert: Einzug: Erste Zeile: 1 cm

Gerard Versteegh 5-6-21 15:28

Gelöscht: have been

Gerard Versteegh 24-6-21 18:34

Gelöscht: due to

Gerard Versteegh 24-6-21 18:35

Gelöscht: due to

Gerard Versteegh 24-6-21 18:37

Gelöscht: .
3

Gerard Versteegh 24-6-21 18:37

Gelöscht: 3

Gerard Versteegh 22-6-21 09:32

Gelöscht: exhaustive

Gerard Versteegh 22-6-21 09:39

Gelöscht: N

Gerard Versteegh 22-6-21 09:38

Gelöscht: production is larger than denitrification so that the sediments are a source of nitrate for

630 3. Where the upper and lower oxygen profiles do not meet, they are separated by an intermediate suboxic zone with free pore-water manganese and cobalt. Here denitrification continues and in some cases may be in excess of nitrate production so that the basement fluid delivers nitrate to the sediment.

635 4. The inferred nitrate concentrations at the basement-sediment interface are relatively constant (43–48 $\mu\text{mol/l}$) and higher than in the ocean bottom waters so that in the sampled region overall more nitrate goes from the sediment to the basement than vice versa.

640 6. The behaviour of the upper and lower oxic/suboxic transitions is very different. The upper oxic/suboxic redox boundary is at the end of a curved oxygen profile. The redox boundary moves up with sedimentation, leaving unoxidised OM behind in the manganese reduction zone below. Moreover, due to the relatively short distance between bottom water and the front, the position of the upper front is sensitive to changes in bottom water oxygen content and as such relatively unstable. The lower oxic/suboxic redox boundary is at the end of a linear oxygen profile, and it moves up as a result of lacking (sufficiently labile) oxidisable material further down. It leaves behind an 'energetic desert' without further oxidation potential. The long transport route of bottom water oxygen, via basement fluid (advection) into the sediment and to the lower oxic/suboxic transition (diffusion), strongly buffers amplitude and frequency of changes in oxygen content, so that the lower transition is much more stable than the upper front.

645 7. Over time, manganese, nickel and cobalt diffuse from the suboxic zone to the upper and lower redox fronts/transitions where they are oxidised and precipitate. At the upper transition they add to the manganese, cobalt and nickel that were deposited by sedimentation. With this transition moving upward with sedimentation, these elements re-enter the suboxic zone, remobilise and move up to the oxic/suboxic transition again, leading to a continuous increasing in solid-phase contents of these elements near the redox front. With the lower transition moving upwards, these elements remain fixed in the oxidised sediment below, resulting in a permanent loss of these elements from the suboxic zone.

650 8. Since in the region between the Clarion and Clipperton Fracture Zones sediment cover is generally thin and sedimentation rates are generally low, we are mostly able to observe the upper oxygen profile and accompanying redox zone as well as in reverse the lower redox zone and oxygen profile, each with its associated set of phenomena such as nitrate diffusion into the basement fluid. In many other ocean regions sediments are thicker and productivity higher so that the lower oxygen profile will be below the reach of the gravity- or piston-coring device due to a more complete redox zonation reaching deeper into the sediment and drilling is needed to reach the successive redox zones as well as the reversed redox zonation followed by the deep oxygen profile to the basement (provided sufficient oxygen is provided by the basement). We are just beginning to understand the impact of this deep redox zonation on the biogeochemical cycles of carbon and nitrogen as well as the distribution of carbon and other elements (such as manganese and associated metals such as nickel and cobalt and, in organically richer sediments, iron and sulphur) in the deep sediments. Fortunately, deep drilling (e.g., through international consortia like DSDP, ODP and IODP) logistically and financially enable access to these deeper sediments to investigate to

Gerard Versteegh 24-6-21 18:40

Gelöscht: 4

Gerard Versteegh 24-6-21 18:40

Gelöscht: 5

Gerard Versteegh 24-6-21 18:39

Gelöscht: Extrapolation of nitrate profiles indicates that the

Gerard Versteegh 24-6-21 18:39

Gelöscht: and we conclude that

Gerard Versteegh 24-6-21 18:40

Gelöscht: (see conclusion 3)

Gerard Versteegh 24-6-21 18:40

Gelöscht: (see conclusion 4)

Gerard Versteegh 21-7-21 10:21

Gelöscht: an

Gerard Versteegh 21-7-21 10:21

Gelöscht: an

what extent theories and hypotheses erected on the basis of shallower settings are applicable for the majority of the ocean floor. Considering the potential area involved this warrants a substantial research effort.

675 9. The vast amount of oxygen and oxidised materials in sediments below the oligotrophic ocean buffer the redox environment of the ocean bottom waters. Fluid circulation through the basaltic basement increases the exchange between sediment and the bottom water and thus the buffer capacity. Our study has pointed out that not only diffusive fluxes in the top part of deep-sea sediments modify the geochemical composition over time, but also diffusive transfer from basement fluids into the bottom layers of the sediment, and vice versa. Hence, paleoceanographic interpretation of sedimentary archives should carefully consider such secondary modifications and adjust rates of (bio)geochemical processes in order to prevent misinterpretation as primary signatures

680

6 Code and Data Availability

Data are available at <https://doi.pangaea.de/10.1594/PANGAEA.932959>.

7 Author Contributions

685 Gerard J. M. Versteegh interpreted the data and wrote the manuscript. Inken Preuss carried out the porewater analyses, Thomas Kuhn performed the solid phase analyses. Sabine Kasten and Andrea Koschinsky, together with Thomas Kuhn, designed and supervised the project. All authors discussed data and contributed to the manuscript at different stages.

8 Competing Interests

The authors declare they have no competing interests.

690 9 Acknowledgments

We thank captain Lutz Mallon and his crew for the excellent co-operation and the pleasant atmosphere during RV *Sonne* cruise SO240. We also appreciate the enthusiasm and engagement of the scientific participants of the expedition. We thank the reviewers Geoff Wheat and Beth Orcutt for their detailed and constructive reviews. This work was funded by the German Federal Ministry for Education and Science (BMBF) under grant no. 03G0240A-C and by internal funding of the Federal

Gerard Versteegh 14-7-21 12:07

Gelöscht: xxxxxx

Gerard Versteegh 14-7-21 12:08

Gelöscht: A doi has been applied but has not yet been issued.

Institute for Geosciences and Natural Resources (BGR) under the project no. A-0203002.A. We acknowledge further financial support from the Helmholtz Association (Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research). The working areas are within the area contracted by the International Seabed Authority to the German Federal Institute for Geosciences and Natural Resources for exploration of polymetallic nodules.

References

Anderson, R. N., Hobart, M. A., Langseth, M. G.: Geothermal convection through oceanic crust and sediments in the Indian Ocean, *Science*, 204, 828–832, 1979, doi: 10.1126/science.204.4395.828.

Aplin, A. C., Cronan, D. S.: Ferromanganese oxide deposits from the Central Pacific Ocean, II. Nodules and associated sediments, *Geochim. Cosmochim. Ac.* 49, 437–451, 1985, doi: 10.1016/0016-7037(85)90035-3.

Armarson, T. S., Keil, R. G.: Changes in organic matter–mineral interactions for marine sediments with varying oxygen exposure times, *Geochim. Cosmochim. Ac.* 71, 3545–3556, 2007, doi: 10.1016/j.gca.2007.04.027.

Arndt, S., Jørgensen, B. B., LaRowe, D. E., Middelburg, J. J., Pancost, R. D., Regnier, R.: Quantifying the degradation of organic matter in marine sediments: A review and synthesis, *Earth–Sci. Rev.*, 123, 53–86, 2013, doi: 10.1016/j.earscirev.2013.02.008.

Baker, P. A., Stout, P. M., Kastner, M., Elderfield, H.: Large–scale lateral advection of seawater through oceanic crust in the central equatorial Pacific, *Earth Planet. Sci. Lett.*, 105, 522–533, 1991, doi: 10.1016/0012-821X(91)90189-O.

Barckhausen, U., Bagge, M., Wilson, D. S.: Seafloor spreading anomalies and crustal ages of the Clarion–Clipperton Zone, *Mar. Geophys. Res.*, 34, 79–88, 2013, doi: 10.1007/s11001-013-9184-6.

Berner, R. A.: *Early Diagenesis. A Theoretical Approach*. Princeton University Press, Princeton NJ, 1980.

Bowles, M. W., Mogollón, J. M., Kasten, S., Zabel, M., Hinrichs, K.–U.: Global rates of marine sulfate reduction and implications for sub–sea–floor metabolic activities, *Science*, 344, 889–891, 2014, doi: 10.1126/science.1249213.

Cai, W. –J., Sayles, F. L.: Oxygen penetration depths and fluxes in marine sediments, *Mar. Chem.*, 52, 123–131, 1996, doi: 10.1016/0304-4203(95)00081-X.

Canfield, D. E., Thamdrup, B.: Towards a consistent classification scheme for geochemical environments, or, why we wish the term 'suboxic' would go away, *Geobiol.*, 7, 385–392, 2009, doi: 10.1111/j.1472-4669.2009.00214.x.

Coogan, L. A. Gillis, K. M. Low–Temperature Alteration of the Seafloor: Impacts on Ocean Chemistry, *Annu. Rev. Earth Planet. Sci.*, 46, 21–45, 2018, doi: 10.1146/annurev-earth-082517-010027.

Gerard Versteegh 23-6-21 17:50

Gelöscht: ta

Gerard Versteegh 23-6-21 17:49

Gelöscht: ta

- D'Hondt, S., Spivack, A. J., Pockalny, R., Ferdelman, T. G., Fischer, J. P., Kallmeyer, J., Abrams, L. J., Smith, D. C., Graham, D., Hasiuk, F., Schrum, H., Stancin, A. M.: Subseafloor sedimentary life in the South Pacific Gyre, *PNAS*, 106, 11651–11656, 2009, doi: 10.1073/pnas.0811793106.
- 730 D'Hondt, S., Inagaki, F., Alvarez Zarikian, C., IODP Expedition 329 Scientific Party: IODP Expedition 329: Life and Habitability Beneath the Seafloor of the South Pacific Gyre. *Sci. Drilling*, 15, 4–10, 2013, doi: 10.2204/iodp.sd.15.01.2013.
- D'Hondt, S., Inagaki, F., Zarikian, C. A., Abrams, L. J., Dubois, N., Engelhardt, T., Evans, H., Ferdelman, T., Gribsholt, B., Harris, R. N., Hoppie, B., Hyun, J. -H., Kallmeyer, J., Kim, J., Lynch, J. E., McKinley, C., Mitsunobu, S., Morono, Y., Murray, R. W., Pockalny, R., Sauvage, J., Shimono, T., Shiraishi, F., Smith, D., C. Smith-Duque, C., Spivack, A. J., Steinsbu, B. O., Suzuki, Y., Szpak, M., Toffin, L., Uramoto, G., Yamaguchi, Y. T., Zhang, G. -L., Zhang, X. -H., Ziebis, W.: Presence of oxygen and aerobic communities from sea floor to basement in deep-sea sediments, *Nat. Geosci.*, 8, 299–304, 2015, doi: 10.1038/NGEO2387.
- 735 D'Hondt, S., Inagaki, F., Orcutt, B. N., Hinrichs K.-U.: IODP advances in the understanding of subseafloor life. *Oceanogr.*, 32, 198-207, 2019, doi: 10.1038/s41467-019-11450-z.
- D'Hondt, S., Pockalny, R., Fulfer, V. M., Spivack, A. J.: Subseafloor life and its biogeochemical impacts, *Nat. Commun.*, 10, 3519, 2019, doi: 10.1038/s41467-019-11450-z.
- Davis, E. E., Chapman, D. S., Wang, K., Villinger, H., Fisher, A. T., Robinson, S. W., Grigel, J., Pribnow, D., Stein, J., Becker: Regional heat flow variations across the sedimented Juan de Fuca Ridge eastern flank: Constraints on lithospheric cooling and lateral hydrothermal heat transport, *J. Geophys. Res. – Atm.*, 104, 17675–17688, 1999, doi: 10.1029/1999JB900124.
- 745 Eittrheim, S. L., Ragozin, N., Gribbidenko, H. S., Helsley, C. E.: Crustal age between the Clipperton and Clarion Fracture Zones, *Geophys. Res. Lett.*, 19, 2365–2368, 1992, doi:10.1029/92GL02928.
- Emerson, S., Hedges, J.: Sediment diagenesis and benthic flux. In: Holland H. D., Turekian K. K. (eds.) *Treatise on Geochemistry vol. 6*, Elsevier, Amsterdam, pp. 293–319, 2003, doi: 10.1016/B0-08-043751-6/06112-0
- 750 Fischer, J. P., Ferdelman, T. G., D'Hondt, S., Roy, H., Wenzhöfer, F.: Oxygen penetration deep into the sediment of the South Pacific gyre, *Biogeosci.*, 6, 1467–1478, 2009, doi: 10.5194/bg-6-1467-2009.
- Fisher, A. T., Davis, E. E., Hutnak, M., Spiess, V., Zühlsdorff, L., Cherkaoui, A., Christiansen, L., Edwards, K., Macdonald, R., Villinger, H., Mottl, M. J., Wheat, C. G., Becker, K.: Hydrothermal recharge and discharge across 50 km guided by seamounts on a young ridge flank, *Nature*, 421, 618–621, 2003, doi: 10.1038/nature01352.
- 755 Fisher, A. T., Wheat, C. G.: Seamounts as conduits for massive fluid, heat, and solute fluxes on ridge flanks, *Oceanogr.*, 23, 74–87, 2010, doi: 10.5670/oceanog.2010.63.

Gerard Versteegh 13-7-21 16:34

Formatiert: Standard, Einzug: 0 cm, Hängend: 1 cm, Tabstopps bei: Nicht an 0 cm

Gerard Versteegh 23-6-21 17:49

Gelöscht: entific

Gerard Versteegh 23-6-21 17:49

Gelöscht: ure

Gerard Versteegh 23-6-21 17:40

Gelöscht: ence

765 Froelich, P. N., Klinkhammer, G. P., Bender, M. L., Luedtke, N. A., Heath, G. R., Cullen, D., Dauphin, P., Hammond, D., Hartman, B., Maynard, V.: Early oxidation of organic matter in pelagic sediments of the eastern equatorial Atlantic: suboxic diagenesis, *Geochim. Cosmochim. Ac.* **43**, 1075–1090, 1979, doi: 10.1016/0016-7037(79)90095-4.

770 García, H. E., Gordon, L. I.: Oxygen solubility in seawater: Better fitting equations, *Limnol. Oceanogr.*, **37**, 1307–1312, 1992, doi: 10.4319/lo.1992.37.6.1307.

Garcia, H. E., Weathers, K., Paver, C. R., Smolyar, I., Boyer, T. P., Locarnini, R. A., Zweng, M. M., Mishonov, A. V., Baranova, O. K., Seidov, D., Reagan, J. R. *World Ocean Atlas 2018, Volume 4: Dissolved Inorganic Nutrients (phosphate, nitrate and nitrate+nitrite, silicate)*. Mishonov, A., Technical Ed, NOAA Atlas NESDIS 84, 35pp, 2018.

775 Hasterok, D., Chapman, D. S., Davis, E. E.: Oceanic heat flow: Implications for global heat loss, *Earth Planet. Sci. Lett.*, **311**, 386–395, 2011, doi: 10.1016/j.epsl.2011.09.044.

Heller, C., Kuhn, T., Versteegh, G. J. M., Wegorzewski, A. V., Kasten, S.: The geochemical behavior of metals during early diagenetic alteration of buried manganese nodules, *Deep Sea Res. Pt. I*, **142**, 16–33, 2018, doi: 10.1016/j.dsr.2018.09.008.

780 Hensen, C., Zabel, M., Schulz, H. D.: A comparison of benthic nutrient fluxes from deep-sea sediments off Namibia and Argentina. *Deep-Sea Res. Pt. II* **47**, 2029–2050, 2000, doi: 10.1016/S0967-0645(00)00015-1.

Hensen, C., Zabel, M., Schulz, H. N., Early Diagenesis at the Benthic Boundary Layer: Oxygen, Nitrogen, and Phosphorus in Marine Sediments, In: Schulz H. D., Zabel M. (eds.), *Marine Geochemistry*, Springer, Heidelberg, pp. 207–240, 2006, doi: 10.1007/3-540-32144-6_6.

Hulme, S. M. Wheat C. G.: Subseafloor fluid and chemical fluxes along a buried-basement ridge on the eastern flank of the Juan de Fuca Ridge. *Geochem, Geophys. Geosys.*, **20**, 4922–4938, 2019, doi: 10.1029/2019GC008408.

785 Hutnak, M., Fisher, A. T., Zühlsdorff, L., Spiess, V., Stauffer, P. H., Gable, C. W.: Hydrothermal recharge and discharge guided by basement outcrops on 0.7–3.6 Ma seafloor east of the Juan de Fuca Ridge: Observations and numerical models, *Geochem. Geophys. Geosyst.*, **7**, 2006, doi: 10.1029/2006GC001242.

Kasten, S., Zabel, M., Heuer, V., Hensen, C.: Processes and signals of non-steady state diagenesis in deep-sea sediments, in: Wefer G., Mulitza S., Rathmeyer V. (eds.), *The South Atlantic in the Late Quaternary*, Springer, Berlin, pp. 431–459, 2003, doi: 10.1007/978-3-642-18917-3_20.

Kay, J., Conklin, M. H., Fuller, Christopher C, O'Day, P. A.: Processes of nickel and cobalt uptake by a manganese oxide forming sediment in Pinal Creek, Globe Mining District, Arizona, *Environmental Sci. Technol.*, **35**, 4719–4725, 2001, doi: 10.1021/es010514d.

Gerard Versteegh 23-6-21 17:48
Gelöscht: ta

Gerard Versteegh 23-6-21 17:48
Gelöscht: earch

Gerard Versteegh 23-6-21 17:47
Gelöscht:

Gerard Versteegh 3-6-21 10:16
Formatiert: Schriftart:Schriftfarbe:
 Schwarz, Englisch (USA), Rahmen:: (Kein
 Rahmen)

Gerard Versteegh 23-6-21 17:41
Gelöscht: istry

Gerard Versteegh 23-6-21 17:41
Gelöscht: ics

Gerard Versteegh 23-6-21 17:41
Gelöscht: ems

Gerard Versteegh 23-6-21 17:41
Gelöscht: ence and

Gerard Versteegh 23-6-21 17:41
Gelöscht: ogy

- 800 Koschinsky, A., Fritsche, U., Winkler, A.: Sequential leaching of Peru Basin surface sediment for the assessment of aged and fresh heavy metal associations and mobility, *Deep Sea Res.* Pt. II: 48, 3683–3699, 2001, doi: 10.1016/S0967-0645(01)00062-5.
- 805 Kuhn, T., Bösel, J., Dohrmann, I., Filmair, C., Fronzek, J., Gerken, J., Goergens, R., Hartmann, J. F., Heesemann, B., Heller, C., Heyde, I., Janssen, A., Kasten, S., Kaul, N., Kevel, O., Kleint, C., Lückge, A., Müller, P., Preuss, I. –M., Purkiani, K., Ritter, S., Rühlemann, C., Schwab, A., Sing, R., Stegger, U., Sturm, S., Uhlenkott, K., Villinger, H., Vink, A., Wedemeyer, H., Węgorzewski, A.: SO240 – Flum: Low-temperature. Fluid. Circulation. at Seamounts and Hydrothermal Pits: Heat Flow Regime, Impact on Biogeochemical Processes and its Potential Influence on the Occurrence and Composition of Manganese Nodules in the Equatorial Eastern Pacific, 1–185, 2015, doi: 10.2312/cr_so240.
- 810 Kuhn, T., Versteegh, G. J. M., Villinger, H., Dohrmann, I., Heller, C., Koschinsky, A., Kaul, N., Ritter, S., Węgorzewski, A. V., Kasten, S.: Widespread sea-water circulation in 18–22 Ma oceanic crust: Impact on heat flow and sediment geochemistry, *Geology*, 45, 799–802, 2017, doi:10.1130/G39091.1.
- de Lange, G. J.: Geochemical evidence of a massive slide in the southern Norwegian Sea. *Nature* 305, 420–422, 1983, doi: 10.1038/305420a0.
- 815 Lutz, M. J., Caldeira, K., Dunbar, R. B., Behrenfeld, M. J.: Seasonal rhythms of net primary production and particulate organic carbon flux describe the biological pump efficiency in the global ocean. *J. Geophys. Res. Oceans*, 112, C10011, 2007, doi: 10.1029/2006JC003706.
- Mewes, K., Mogollón, J. M., Picard, A., Rühlemann, C., Kuhn, T., Nöthen, K., Kasten, S.: Impact of depositional and biogeochemical processes on small scale variations in nodule abundance in the Clarion-Clipperton Fracture Zone, *Deep Sea Res. Pt. I*, 91, 125–141, 2014, doi:10.1016/j.dsr.2014.06.001.
- 820 Mewes, K., Mogollón, J. M., Picard, A., Rühlemann, C., Eisenhauer, A., Kuhn, T., Ziebis, W., Kasten, S.: Diffusive transfer of oxygen from seamount basaltic crust into overlying sediments: An example from the Clarion-Clipperton Fracture Zone, *Earth Planet. Sci. Lett.*, 433, 215–225, 2016, doi:10.1016/j.epsl.2015.10.028.
- Mogollón, J. M., Mewes, K., Kasten, S.: Quantifying manganese and nitrogen cycle coupling in manganese-rich, organic carbon-starved marine sediments: Examples from the Clarion-Clipperton fracture zone, *Geophys. Res. Lett.*, 43, 7114–7123, 2016, doi:10.1002/2016gl069117.
- 825 Morono, Y., Ito, M., Hoshino, T., Terada, T., Hori, T., Ikehara, M., D’Hondt, S., Inagaki, F.: Aerobic microbial life persists in oxic marine sediment as old as 101.5 million years, *Nature Commun.*, 11, 3626, 2020, doi: 10.1038/s41467-020-17330-1.

Gerard Versteegh 23-6-21 17:41

Gelöscht: earch

Gerard Versteegh 23-6-21 17:47

Gelöscht: :

- 830 | Murray, J. W., Dillard, J. G.: The oxidation of cobalt(II) adsorbed on manganese dioxide, *Geochim. Cosmochim. Ac.* **43**, 781–787, 1979, doi:10.1016/0016-7037(79)90261-8.
- Müller, P. J., Mangini, A.: Organic carbon decomposition rates in sediments of the Pacific manganese nodule belt dated by ^{230}Th and ^{231}Pa , *Earth Planet. Sci. Lett.*, **51**, 94–114, 1980, doi: 10.1016/0012-821X(80)90259-9.
- Nöthen, K., Kasten, S.: Reconstructing changes in seep activity by means of pore water and solid phase Sr/Ca and Mg/Ca ratios in pockmark sediments of the Northern Congo Fan, *Mar. Geol.*, **287**, 1–13, 2011, doi: 10.1016/j.margeo.2011.06.008.
- 835 | Orcutt, B. N., Wheat, C. G., Rouxel, O., Hulme, S., Edwards, K. J., Bach, W.: Oxygen consumption rates in subseafloor basaltic crust derived from a reaction transport model, *Nat. Commun.*, **4**, 2539, 2013, doi:10.1038/ncomms3539.
- Rietveld, H. M.: Line profiles of neutron powder-diffraction peaks for structure refinement, *Acta Crystallogr.*, **22**, 151–152, 1967, doi: 10.1107/S0365110X67000234.
- 840 | Revsbech, N. P.: An oxygen microsensor with a guard cathode. *Limnol. Oceanogr.* **34**, 474–478, 1989, doi: 10.4319/lo.1989.34.2.0474.
- Rühlemann, C., Kuhn, T., Wiedicke, M., Kasten, S., Mewes, K. and Picard, A. (2011) Current status of manganese nodule exploration in the German license area. Proceedings of the Ninth (2911) ISOPE Ocean Mining Symposium, Maui, Hawaii, USA, June 19–24, 2011, 168–173.
- 845 | Sclater, J. G., Jaupart, C., Galson, D.: The heat flow through oceanic and continental crust and the heat loss of the Earth, *Rev. Geophys. Rev. Geophys.*, **18**, 269–311, 1980, doi: 10.1029/RG018i001p00269.
- Seeberg-Elverfeldt, J., Schlüter, M., Feseker, T., Kölling, M.: Rhizon sampling of porewaters near the sediment–water interface of aquatic systems, *LOM3. Limnol. Oceanogr.: Methods.*, **3**, 361–371, 2005, doi: 10.4319/lom.2005.3.361.
- 850 | Spinelli, G. A., Giambalvo, E. R., Fisher, A. T.: Sediment permeability, distribution, and influence on fluxes in oceanic basement. In: Davis E. E., Elderfield H. (eds.), *Hydrogeology of the oceanic lithosphere*, Cambridge University Press, Cambridge, 151–188, 2004.
- Stein, C. A., Stein, S.: A model for the global variation in oceanic depth and heat flow with lithospheric age, *Nature*, **359**, 123–129, 1992, doi: 10.1038/359123a0.
- 855 | Tegelaar, E. W., de Leeuw, J. W., Derenne, S., Largeau, C.: A reappraisal of kerogen formation, *Geochim. Cosmochim. Ac.* **53**, 3103–3106, 1989, doi: 10.1016/0016-7037(89)90191-9.
- Volz, J. B., Mogollón, J. M., Geibert, W., Martínez Arbizu, P., Koschinsky, A., Kasten, S.: Natural spatial variability of depositional conditions, biogeochemical processes and element fluxes in sediments of the eastern Clarion–Clipperton Zone, Pacific Ocean, *Deep Sea Res. Pt. I*, **140**, 159–172, 2018, doi: 10.1016/j.dsr.2018.08.006.

Gerard Versteegh 23-6-21 17:42

Gelöscht: ta

Gerard Versteegh 23-6-21 17:42

Gelöscht: ta

Gerard Versteegh 23-6-21 17:43

Gelöscht: earch

Gerard Versteegh 23-6-21 17:46

Gelöscht: ar

- 865 Volz, J. B. , Liu, B. , Köster, M. , Henkel, S. , Koschinsky, A. and Kasten, S.: Post-depositional manganese mobilization during the last glacial period in sediments of the eastern Clarion-Clipperton Zone, Pacific Ocean , *Earth Planet. Sci. Lett.* 532, 116012, 2020. doi: 10.1016/j.epsl.2019.116012.
- Wenzhöfer, F. and Gludd, R. N.: Benthic carbon mineralization in the Atlantic: a synthesis based on in situ data from the last decade. *Deep Sea Res.* Pt. I: Oceanographic Research Papers 49, 1255–1279, 2002, doi: 10.1016/S0967-0637(02)00025-0.
- 870 Wheat, C. G., Fisher, A. T.: Massive, low-temperature hydrothermal flow from a basaltic outcrop on 23 Ma seafloor of the Cocos Plate: Chemical constraints and implications, *Geochem. Geophys. Geosyst.* 9, 1–16, 2008, doi: 10.1029/2008GC002136.
- Wheat, C. G., Mottl, M. J.: Composition of pore and spring waters from Baby Bare: global implications of geochemical fluxes from a ridge flank hydrothermal system, *Geochim. Cosmochim. Ac.* 64, 629–642, 2000, doi: 10.1016/s0016-7037(99)00347-6.
- 875 Wheat, C. G., Hartwell, A. M., McManus, J., Fisher, A. T., Orcutt, B. N., Schlicht, L. E. M., Niedenzu, S., and Bach, W.: Geology and fluid discharge at Dorado Outcrop, a low temperature ridge-flank hydrothermal system, *Geochem. Geophys. Geosyst.*, 20, 487–504, 2019, doi: 10.1029/2018GC007933.
- 880 Wankel, S. D., Buchwald, C., Ziebis, W., Wenk, C. B., Lehmann, M. F.: Nitrogen cycling in the deep sedimentary biosphere: nitrate isotopes in porewaters underlying the oligotrophic North Atlantic, *Biogeosci.*, 12, 7483–7502, 2015, doi: 10.5194/bg-12-7483-2015.
- Wheat, C. G., Becker, K., Villinger, H., Orcutt, B. N., Fournier, T., Hartwell, A., Paul, C.: Subseafloor cross-hole tracer experiment reveals hydrologic properties, heterogeneities, and reactions in slow-spreading oceanic crust, *Geochem. Geophys. Geosyst.*, 21, 1–15, 2020, doi: 10.1029/2019GC008804.
- 885 Williams, D. L., von Herzen, R. P.: Heat Loss from the Earth: New Estimate, *Geology*, 2, 327–328, 1974, doi: 10.1130/0091-7613(1974)2<327:HLFTEN>2.0.CO;2.
- Ziebis, W., McManus, J., Ferdelman, T., Schmidt-Schierhorn, F., Bach, W., Muratli, J., Edwards, K. J., Villinger, H.: Interstitial fluid chemistry of sediments underlying the North Atlantic gyre and the influence of subsurface fluid flow, *Earth Planet. Sci. Lett.*, 323–324, 79–91, 2012, 10.1016/j.epsl.2012.01.018.
- 890 Zonneveld, K. A. F., Versteegh, G. J. M., Kasten, S., Eglinton, T. I., Emeis, K. C., Huguet, C., Koch, B. P., de Lange, G. J., de Leeuw, J. W., Middelburg, J. J., Mollenhauer, G., Prahl, F. G., Rethemeyer, J., Wakeham, S. G.: Selective preservation of organic matter in marine sediments – processes and impact on the fossil record, *Biogeosci.*, 7, 483–511, 2010, doi:10.5194/bg-7-483-2010.

Gerard Versteegh 23-6-21 17:43

Gelöscht: earch

Gerard Versteegh 23-6-21 17:46

Gelöscht: ar

Gerard Versteegh 23-6-21 17:43

Gelöscht: istry

Gerard Versteegh 23-6-21 17:43

Gelöscht: ics

Gerard Versteegh 23-6-21 17:43

Gelöscht: tems

Gerard Versteegh 23-6-21 17:43

Gelöscht: ta

Table 1: Sediment cores analysed (adapted from Kuhn et al., 2015).

Working Area	Core	Latitude (N)	Longitude (W)	Water depth (m)	Length (cm)	Heat flow ($mW \cdot m^{-2}$)	Sediment thickness (m)	Length upper oxic zone (m)	Lower ox. front (m)	Length lower oxic zone (m)
1	SSL F, SM ¹	13° 10.525'	118° 06.705'	4287	756	-	<10	2	n.a. ²	8?
1	9KL	13° 10.524'	118° 10.104'	4335	1187	4.1 _↓	23	0.5	14 ³	3
1	22KL F	13° 10.527'	118° 08.184'	4302	1301	10	20 _↓	1.5	10	14
1	51SL F, SM	13° 10.526'	118° 06.584'	4286	537	42	10 _↓	3	n.a.	7
1	53SL SM	13° 10.508'	118° 06.110'	4273	482	54.8	8	2.5	n.a.	5
1	117SL SM	13° 11.103'	118° 05.992'	4271	600	181.1 _↓	10 _↓	2	n.a.	n.a.
2	35SL SM	12° 54.128'	118° 24.791'	4319	982	9	18	2.5	n.a.	15.5
2	42SL F	12° 51.249'	118° 23.976'	4290	1066	37.4	29	<0.5	7	22
3	58SL SM	12° 53.216'	119° 08.351'	4309	1244	48.9 _↓	73	3?	?	?
3	65SL	12° 56.107'	119° 08.884'	4293	1275	63.4 _↓	72	3?	?	?
3	69SL F	12° 39.855'	119° 13.374'	4275	1265	135.5 _↓	51	3	n.a.	67
3	72SL F	12° 55.597'	119° 08.833'	4294	853	85.3	73	2	?	?
4	81SL F	11° 50.064'	116° 32.890'	4355	1346	91.7	20	2(4?)	n.a.	17
4	96SL F, SM	11° 49.260'	117° 13.195'	4145	980	80.4	21 _↓	3	n.a.	<12
4	108SL	11° 48.796'	116° 31.767'	4326	1038	54 _↓	20	3	16 ²	4

¹ Core names, SL = gravity core, KL = piston core, F = core taken above a fault, SM = core taken near a sea mountain; ² n.a. = not available; _↓ Extrapolated from porewater Mn profile.

Gerard Versteegh 1-7-21 11:06

Gelöscht: ¹

Gerard Versteegh 1-7-21 10:58

Gelöscht: 5

Gerard Versteegh 30-6-21 15:39

Gelöscht: 17

Gerard Versteegh 1-7-21 11:07

Gelöscht: ²

Gerard Versteegh 30-6-21 15:39

Gelöscht: 24

Gerard Versteegh 30-6-21 15:28

Gelöscht: <

Gerard Versteegh 1-7-21 10:58

Gelöscht: 95

Gerard Versteegh 30-6-21 15:30

Gelöscht: -

Gerard Versteegh 21-7-21 08:48

Gelöscht: K

Gerard Versteegh 1-7-21 10:59

Gelöscht: 50

Gerard Versteegh 30-6-21 15:39

Gelöscht: 83

Gerard Versteegh 1-7-21 10:59

Gelöscht: 5

Gerard Versteegh 30-6-21 15:39

Gelöscht: 71

Gerard Versteegh 1-7-21 11:00

Gelöscht: 0

Gerard Versteegh 30-6-21 15:39

Gelöscht: 70

Gerard Versteegh 30-6-21 15:39

Gelöscht: 72

Gerard Versteegh 30-6-21 15:31

Gelöscht: <

Gerard Versteegh 30-6-21 15:31

Gelöscht: 15

Gerard Versteegh 1-7-21 11:01

Gelöscht: 60

Gerard Versteegh 21-7-21 10:23

Formatiert: Zeilenabstand: einfach

Gerard Versteegh 1-7-21 11:08

Gelöscht: ¹

Gerard Versteegh 1-7-21 11:08

Gelöscht: ²

Table 2: Linear regressions and extrapolated NO_3^- concentrations at the sediment base. Regressions are based on the lower, linear part of the O_2 and NO_3^- profiles. Whereby: Concentration in $\mu\text{mol} / \text{L} = a * (\text{sediment depth in meters}) + b$

	Species	a	b	r^2	concentration ($\mu\text{mol} / \text{L}$) ¹	Depth (m)	
	5 SL	NO_3^-			45.5 _v	7.5	
	22 KL	NO_3^-	0.266	37.88	0.65	43.2 _v	20 _v
	35 SL	NO_3^-	-0.415	52.84	0.85	45.4	18
	51 SL	O_2	19.89	-68.05	0.84	131	10
	51 SL	O_2	2.90	8.923	0.83	37.9	10
	53 SL	NO_3^-			46 ¹	5	
	53 SL	O_2	4.84	27.31	0.32	66	8
	81 SL	NO_3^-			50 ?	20	
	81 SL	O_2	6.89	-45.98	0.96	91.82	20
	96 SL	NO_3^-	-0.153	51.18	0.38	47.9 _v	21 _v
	96 SL	O_2	0.71	14.45	0.85	29.36	21
	35 SL	O_2	3.71	0.903 _v	0.95	65.9	18
	69 SL	O_2	2.98	4.48	0.95	156 _v	51 _v

¹ Estimated from the values and trend at the core base.

Gerard Versteegh 2-7-21 10:31

Gelöscht: 1

Gerard Versteegh 30-6-21 15:33

Gelöscht: 5

Gerard Versteegh 30-6-21 15:33

Gelöscht: 2

Gerard Versteegh 30-6-21 15:38

Gelöscht: 8

Gerard Versteegh 30-6-21 15:38

Gelöscht: 15

Gerard Versteegh 1-7-21 13:43

Gelöscht: 29

Gerard Versteegh 30-6-21 15:35


Gelöscht: 204

Gerard Versteegh 30-6-21 15:36

Gelöscht: 70

Unknown

Formatiert: Schriftart:(Standard) Times New Roman, 9 pt, Fett, Unterstreichungsfarbe: Automatisch, Schriftfarbe: Automatisch, Rahmen:: (Kein Rahmen)



945 **Figure 1: Maps (adapted from Kuhn et al., 2015). Top panel A. Geographical setting. B. Bathymetric map of the region of investigation with the four working areas indicated. Lower four panels cut-outs of bathymetric maps of the four working areas, with core locations indicated by red dots. Intervals between isobaths for working areas 1 to 3 are 50 m, and 100 m for working area 4. Star, core taken above a fault; circle, core taken near a seamount; pentagon, core taken above a fault and near a seamount; square core taken neither close to a seamount, nor above a fault.**

Gerard Versteegh 30-6-21 10:40

Gelöscht: , bathymetric maps of the working areas

Fig. 2 Entirely oxic cores





Figure 2: Dissolved oxygen, manganese and nitrate profiles of cores without free porewater Mn^{2+} . For all cores, nitrate profiles do not display significant denitrification beyond the uppermost meter (e.g., in 35SL). NB: core 5SL may reflect disequilibrium conditions due to recent slumping. NB: Nitrate increase with depth for 81SL is aberrant as is the absence of free manganese. EST denotes estimated sediment thickness for each core (from Table 1).

955

Fig. 3 Partly suboxic cores



960 **Figure 3: Dissolved oxygen, manganese and nitrate profiles of cores with free porewater manganese in order of maximum manganese concentrations (decreasing minimum nitrate values). For all cores nitrate values reach below the inferred value for basement fluid of 45 $\mu\text{mol/l}$, suggesting the sediments at these locations are nitrate sinks. Core 22KL illustrates this with a nitrate minimum at 7 m core depth, indicating the depth where denitrification equals the nitrate fluxes from above and below. EST denotes estimated sediment thickness for each core (from Table 1).**




Fig 4a




Fig 4b




Fig 4c

000

005





Figure 4a–d: Combined porewater and solid–phase profiles for oxygen, Mn, Co, Ni, and nitrate in the cores with suboxic zones.

010

015

020



Unknown
Formatiert: Schriftart:(Standard) Times
 New Roman, Unterstreichungsfarbe:
 Automatisch, Schriftfarbe: Automatisch,
 Rahmen:: (Kein Rahmen)

025

Figure 5: Compilation of oxygenation states of cores for the entire sediment column (sediment surface to basement) taken during RV *Sonne* Cruise SO240. Sediment thickness is based on seismic and parasound data (Kuhn et al., 2015); oxidation state based on core porewater oxygen and manganese concentrations. Note that for the totally oxic cores (blue only) the oxygen minimum (where the downward dark blue, and upward light blue oxygen profiles meet) is always in the upper few meters. The upper oxic/suboxic transition also occurs in the upper few meters. The black line at 3 m depth marks the lower limit of the upper oxic zone for all cores except for 108SL.

030




Figure 6: Relationship between maximum Mn^{2+} and minimum NO_3^- concentrations for the cores with a suboxic zone.

Unknown
Formatiert: Schriftart:(Standard) Times New Roman, 9 pt, Fett, Unterstreichungsfarbe: Automatisch, Schriftfarbe: Automatisch, Rahmen:: (Kein Rahmen)

035

040

Circulation through oceanic crust




Figure 7: Model of fluid circulation in the basaltic basement, diffusive exchange with the sediment and the sediment oxidation state. Note the influence of faults. Representative cores taken during SO240 are placed such that their geochemical profiles match with the model (modified from Fisher and Wheat, 2010).

045




Figure 8: Conceptual model summarising the diagenetic processes discussed in this paper as a result of oxygen entering the sediment from the ocean bottom waters and from the oceanic basement. Left panel: porewater profiles, Right panel solid fraction. Numbers refer to concentrations in $\mu\text{mol/L}$. The model is based on the profiles of inorganic species and TOC in core.