

Mixed layer depth dominates over upwelling in regulating the seasonality of ecosystem functioning in the Peruvian Upwelling System

Tianfei Xue¹, Ivy Frenger¹, A.E. Friederike Prowe¹, Yonss Saranga José¹, and Andreas Oschlies^{1,2}

¹GEOMAR Helmholtz Centre for Ocean Research Kiel, Kiel, Germany

²Christian-Albrechts-University Kiel, Germany

Correspondence: Tianfei Xue (txue@geomar.de)

Abstract. The Peruvian Upwelling System hosts a marine ecosystem with extremely high productivity. Observations show that the Peruvian Upwelling System is the only Eastern Boundary Upwelling System (EBUS) with an out-of-phase relationship between seasonal surface chlorophyll concentrations and upwelling intensity. This "seasonal paradox" triggers the following questions: (1) What are the unique characteristics of the Peruvian Upwelling System, compared with other EBUS, that lead to the out-of-phase relationship; and (2) How does the seasonal paradox influence ecosystem functioning? Using observational climatologies for four EBUS, we diagnose that the Peruvian Upwelling System is the only one to reveal that intense upwelling coincides with deep mixed layers. We then apply a coupled regional ocean circulation-biogeochemical model (CROCO-BioEBUS) to assess how the interplay between mixed layers and upwelling regulates the seasonality of surface chlorophyll in the Peruvian Upwelling System. Our model reproduces the "seasonal paradox" within 200 km off the Peruvian coast. We confirm previous findings regarding the main contribution of mixed layer depth to the seasonality of chlorophyll, relative to upwelling. Deep mixed layers in austral winter cause vertical dilution of phytoplankton and strong light limitation, impacting growth. The effect of advection ~~on the buildup of phytoplankton biomass~~, though second-order, is consistent with previous findings for the Peruvian system and other EBUS, with enhanced offshore export opposing the coastal buildup of biomass. In addition, we find that the relatively colder temperatures of upwelled waters slightly dampen phytoplankton productivity and further slow the buildup of phytoplankton biomass. This impact from the combination of deep mixed layers and upwelling propagates through the ecosystem, from primary production to export and export efficiency. Our findings emphasise the crucial role of the interplay between mixed layer depth and upwelling and suggest that surface chlorophyll may increase, along with a weakened seasonal paradox, in response to shoaling mixed layers under climate change.

Copyright statement.

20 1 Introduction

The Peruvian Upwelling System (PUS) hosts a disproportionately productive ecosystem, supporting 10% of the world's fishing yield while covering only 0.1% of the ocean area (Chavez et al., 2008). As one of the Eastern Boundary Upwelling Systems (EBUS), winds favouring upwelling raise cool, nutrient-rich waters to the surface, supporting high primary production and fish yield. Simultaneously, high primary production, together with subsequent export and remineralisation contributes to the formation of a sub-surface oxygen-deficient zone which is particularly shallow and intense in the PUS (Fuenzalida et al., 2009; Stramma et al., 2010; Getzlaff et al., 2016). Particularly due to its high productivity, the response of the PUS to climate change is of great social and economic interest (Pauly et al., 1998; Bakun, 1990; Bakun et al., 2010), and a variety of studies have investigated how physical and biogeochemical processes influence the production of phytoplankton as well as its potential links to ecosystem functioning in the PUS.

30 While the PUS has been frequently compared to other EBUS (e.g., the Benguela, California, and Canary Systems), it is set apart by how surface chlorophyll responds to the variation of upwelling on a seasonal scale. The high productivity of EBUS primarily benefits from the upwelling of nutrient-rich waters, driven by alongshore equatorward winds. Hence, it is commonly assumed that the magnitude of phytoplankton biomass in EBUS is directly correlated with the wind-driven upwelling intensity (Bakun, 1973). However, in the PUS, upwelling intensity and surface chlorophyll are not correlated on a seasonal scale (here-
35 after referred to as "seasonal paradox"; Chavez, 1995; Thomas et al., 2001; Echevin et al., 2008; Chavez and Messié, 2009). Instead, they are out of phase, with the lowest surface chlorophyll concentration in austral winter corresponding to maximum upwelling intensity (Calienes et al., 1985). Echevin et al. (2008) used a regional coupled physical-biogeochemical model to simulate the "seasonal paradox" and found that deep mixed layers caused dilution of surface phytoplankton, reduced growth due to limited light, and subsequently reduced iron levels (as phytoplankton require more iron under low-light conditions).
40 This ultimately leads to low chlorophyll under strong upwelling conditions. Results from Messié and Chavez (2015) corroborated iron and light limitations found in Echevin et al. (2008), showing additionally that relatively strong offshore advection in austral winter regulated the buildup of phytoplankton and thus also contributed to the seasonal paradox. Guillen and Calienes (1981) suggested that lower surface irradiation in winter might amplify light limitation and further limit phytoplankton growth, while insolation was found not to play a major role in Echevin et al. (2008). Additionally, Echevin et al. (2008) concluded that
45 temperature played no role in regulating phytoplankton growth. Despite previous research on surface chlorophyll seasonality, uncertainty still remains regarding why the seasonal paradox occurs only in the PUS and not in the other EBUS, and it is unexplored how the seasonal paradox affects ecosystem functioning.

This study addresses the following key questions: (1) what are the unique characteristics of the PUS, compared to other EBUS, that lead to this seasonal paradox; (2) what are the mechanisms that cause low surface phytoplankton in winter; and (3)
50 how do these mechanisms affect ecosystem functioning.

2 Data and Methods

2.1 Regional ocean circulation-biogeochemical model: setup and simulation

We use a climatological simulation of the three-dimensional regional ocean circulation model CROCO (Coastal and Regional Ocean COmmunity model; Debreu et al., 2012) coupled with the biogeochemical model BioEBUS (Biogeochemical model for the Eastern Boundary Upwelling Systems; Gutknecht et al., 2013) for this study.

The same technical setup, including the model grid, is used as in José et al. (2017), along with an updated version of the ocean circulation model CROCO. CROCO is the next generation of the ROMS AGRIF model (Tedesco et al., 2019), and is a free-surface and split-explicit regional ocean model system (ROMS; Shchepetkin and McWilliams, 2005). We employ a two-way nesting approach, with the larger coarser-resolution domain covering the Southeast Pacific and the smaller higher-resolution domain focusing on the PUS. The larger domain has a $1/4^\circ$ resolution, spanning from 69°W to 120°W and from 18°N to 40°S . The embedded "child" domain has a resolution of $1/12^\circ$ and extends from 5°N to 31°S and from 69°W to 102°W (Fig. [1a-b & A1](#)) and is used in this study. Both the coarse- and fine-resolution domains use 32 sigma levels in the vertical direction, with finer resolution towards the surface and shallower regions. The surface layer thickness ranges from 0.5 m in the coastal region (water depth around 50 m) to around 3 m in the offshore region, ~~corresponding to a~~ (water depth of more than 4000 m). Initial and boundary conditions are provided by the monthly climatological SODA reanalysis (Simple Ocean Data Assimilation; Carton and Giese, 2008) from 1990–2010. Surface forcing is based on the monthly climatological heat and freshwater fluxes from COADS (Comprehensive Ocean-Atmosphere Data Set; Worley et al., 2005), along with wind data from QuikSCAT (Quick Scatterometer; Liu et al., 1998). The physical setup is the same as in José et al. (2017) and has been evaluated therein, showing that the model reproduces the circulation of the region reasonably well.

The biogeochemical BioEBUS model used in this study was developed explicitly for applications to EBUS and oxygen minimum zones (Gutknecht et al., 2013). BioEBUS is a nitrogen-based model, originating from the $\text{N}_2\text{P}_2\text{Z}_2\text{D}_2$ model by Koné et al. (2005). It simulates two phytoplankton and two zooplankton groups: small and large phytoplankton, along with micro- and mesozooplankton. Furthermore, there are two detritus pools, categorised by size. BioEBUS resolves the N species (nitrate, nitrite and ammonium) and simulates processes under oxic, hypoxic and suboxic conditions (e.g., remineralisation, nitrification, denitrification and anammox). The BioEBUS model was first used to study the Peruvian marine biogeochemistry by Montes et al. (2014), and is capable of producing a realistic simulation of the oxygen distribution. Initial and boundary conditions for nitrate and oxygen are taken from CARS (CSIRO (Commonwealth Scientific and Industrial Research Organisation) Atlas of Regional Seas; Ridgway et al., 2002), and initial conditions for phytoplankton are based on monthly climatological SeaWiFS (Sea-viewing Wide Field-of-view Sensor; O'Reilly et al., 1998) estimates. A detailed description of these biogeochemical processes can be found in Gutknecht et al. (2013). The parameter settings are the same as in José et al. (2017), except for a few adjustments ~~to of~~ biological parameters (Table. A1) to improve the fit between the simulated ecology, in particular phytoplankton and zooplankton ~~biomass and seasonality~~, and the observations.

CROCO-BioEBUS is run in coupled mode from the beginning of the simulation. The time-stepping of the physical model is the same as the coupling time step, with a duration of 1200 seconds. The time-stepping of the biogeochemical model has a

85 duration of 400 seconds. The coupled model is run for a 25-year spin-up period. Physical and biogeochemical fields are spun
up after one year for the upper 10 m, while waters in the depth range of upwelling ~~source~~ (100 m) require 3–10 years longer to
reach a statistical quasi-equilibrium (Fig. C1). We run the model for a total of 30 climatologically-forced years, using the last
five years for the analyses. As we observe from the surface ecology, our results are not sensitive to deep ocean spin-up. This
study focuses on the 200 km band off the Peruvian coast (white line region in Fig. 1a-b), which shows clear seasonal variation
90 as well as strong upwelling.

2.2 Analysis approaches

To assess the seasonal variance of phytoplankton biomass concentration in each grid box (C), we analysed the budget of the
phytoplankton biomass and how its tendency is driven by physical versus biological processes:

$$\frac{\partial C}{\partial t} = PHY(C) + BIO(C) \quad (1)$$

95 with [$BIO = PP - GRAZ - MORT - EXU - SINK$; $PHY = MIX + ADV$]

PHY represents the physical processes, including advection ~~and mixing~~ ADV and mixing MIX, whereas BIO represents the
biological processes, namely primary production PP , consumptive mortality $GRAZ$, natural mortality $MORT$, exudation
 EXU and sinking $SINK$. All biological and physical fluxes ~~except for entrainment~~ were saved monthly from the simulation,
100 with units of $\text{mmol N m}^{-3} \text{ s}^{-1}$, and we integrated the terms offline over the mixed layer depth (MLD) using the croco-tools
provided for post-processing (<https://www.croco-ocean.org/download/croco-project/>). ~~We calculated the monthly entrainment
as a residual, following the calculation of entrainment for physical variables in the CROCO code. That is, entrainment was
calculated as the difference between the MLD-integrated biomass tendency and the sum of the biological and physical fluxes
integrated over the MLD of the previous month. The mixing term also includes entrainment from varying MLD as a minor~~
105 contribution.

We analysed in detail the drivers of PP , which was ~~determined by~~ calculated online by multiplying phytoplankton concen-
tration (C) and the growth factors ($L_{(PAR)}$, $L_{(T)}$, $L_{(N)}$)

$$PP = C \cdot L_{(PAR)} \cdot L_{(T)} \cdot L_{(N)} \quad (2)$$

where $L_{(PAR)}$, $L_{(T)}$, $L_{(N)}$ represent the light-, temperature- or nitrogen-related growth factors, respectively. Here, the phy-
110 toplankton growth rate was defined as a multiplicative function of the light-, temperature- or nitrogen-related growth factors.
The limitation experienced by phytoplankton within the mixed layer L_{mld} is calculated offline from each growth factor ($L_{(PAR)}$,
 $L_{(T)}$ and $L_{(N)}$), using phytoplankton concentration (C) within the mixed layer as a weight (Eq. 3). Light-, temperature- and
nitrogen-related growth factors that each phytoplankton cell experienced were computed online.

$$L_{\text{mld}} = \frac{\sum_0^{\text{mld}} L_{(PAR)} \cdot L_{(T)} \cdot L_{(N)} \cdot C}{\sum_0^{\text{mld}} C} \quad (3)$$

115 For the analysis, we attributed the seasonal change of the average phytoplankton biomass concentration (ΔC_{mld}) within the
mixed layer to the change in the integrated phytoplankton content within the mixed layer (ΔB_{mld}), as well as the change in

volume of the mixed layer (ΔV_{mld}). Using the chain rule and the condition that $V^2 \gg V\Delta V$, we approximated a discrete change in the mixed layer tracer concentration (ΔC_{mld}) as follows:

$$\Delta C_{\text{mld}} = \frac{1}{V_{\text{mld}}} \Delta B_{\text{mld}} - B_{\text{mld}} \frac{\Delta V_{\text{mld}}}{V_{\text{mld}}^2} = \frac{B_{\text{mld}}}{V_{\text{mld}}} \frac{\Delta B_{\text{mld}}}{B_{\text{mld}}} - \frac{B_{\text{mld}}}{V_{\text{mld}}} \frac{\Delta V_{\text{mld}}}{V_{\text{mld}}} \quad (4)$$

120 To assess the relative contributions, we then divided by $C_{\text{mld}} = B_{\text{mld}} \cdot V_{\text{mld}}^{-1}$ to obtain

$$\frac{\Delta C_{\text{mld}}}{C_{\text{mld}}} = \frac{\Delta B_{\text{mld}}}{B_{\text{mld}}} - \frac{\Delta V_{\text{mld}}}{V_{\text{mld}}} \quad (5)$$

which allowed us to attribute decreased concentration of phytoplankton in the mixed layer $\downarrow C_{\text{mld}}$ to a decrease in the phytoplankton biomass B_{mld} or an increase in the mixed-layer volume V_{mld} , and vice versa.

2.3 Observational data and model assessment

125 For EBUS comparisons, we digitised SeaWIFS climatological surface chlorophyll and upwelling (~~a combination of Ekman transport and Ekman pumping, Messié et al. (2009)~~), (a combination of Ekman transport and Ekman pumping; Messié et al., 2009)
estimated based on winds from QuikSCAT, from Chavez and Messié (2009). Additionally, we used surface nitrate data from the World Ocean Atlas (WOA; Garcia et al., 2019), the gridded ARGO mixed layer dataset (~~Holte et al. (2017)~~), (http://mixedlayer.ucsd.edu/monthly-climatology (http://mixedlayer.ucsd.edu/; Holte et al., 2017), monthly climatologies of MODIS sea surface temperature (SST) and chlorophyll data (https://oceancolor.gsfc.nasa.gov/data/aqua/) to analyse and evaluate the model results.

The model was evaluated based on averages over the focus region, with monthly observational data. The correlation coefficient between the model simulation and observations, the root mean square error (RMSE) and the normalised standard deviation (SD) of the observations relative to the model results are shown in a Taylor diagram as a summary of the evaluation (~~Fig. 1e; Taylor et al. (1991)~~); (a comparison of the spatial pattern and the seasonal cycles of variables is provided in the appendix, see
135 Figs. B1-B4). (Fig. 1c; Taylor et al., 1991, a comparison of the spatial pattern and the seasonal cycles of variables is provided in the appendix)
Model results fit the observational data reasonably well. The model effectively simulated sea surface temperature (SST). SST
with $R > 0.95$, $1 < \sigma^* < 1.2$ and $RMSE^* < 0.4$ (R : correlation coefficient, σ^* : normalised SD, and $RMSE^*$: normalised RMSE). It also captured the observed seasonal cycle well, though it produced slightly stronger seasonal variations compared to those from the observational data. Although the seasonal variation was somewhat overestimated, the simulated MLD (defined based on a
140 0.2°C temperature difference criterion) remained mostly largely within the observed range of ARGO-based MLD (Fig. B3). As for biogeochemical variables, the model effectively simulated surface nitrate, with $R > 0.95$, $0.6 < \sigma^* < 1$ and $RMSE^* < 0.4$, but overestimated the nitrate compared to WOA. Cruise ~~However, cruise~~ data (Fig. B2c-d) show that the overestimation could have arisen from WOA failing to capture the high-surface nitrate concentration in the coastal region under strong upwelling. A comparison of the simulated and observed seasonal cycle of surface chlorophyll in the focus region (Fig. 1d) revealed that
145 modelled chlorophyll generally followed the seasonal trend of satellite and *in situ* data, with the amplitude of the seasonal cycle in between amplitudes from satellite and *in situ* data. Overall, the model showed reasonably good agreement with observational data on a seasonal scale, sufficiently supporting an investigation of the seasonal paradox with CROCO-BioEBUS.




Figure 1. (a) Spatial distribution of the amplitude of the annual cycle of surface chlorophyll in log scale ($\log(\text{chl} \text{ (mg m}^{-3}\text{)}^{-1})$); (b) Map of annual mean upwelling velocity $w-w$ (m d^{-1}) at the bottom of the mixed layer, with contour lines indicating MLD (m). White lines highlight the focus area; (c) Taylor diagram for seasonal SST (red), MLD (yellow), surface nitrate concentrations (purple) and chlorophyll concentrations (green). The black dot indicates the model simulation as a reference. The radial distance from the origin is proportional to the standard deviation, normalised by the standard deviation of the data. The green dashed lines show the RMSE. The correlations between model and observations are given by the azimuthal position; (d) Seasonal cycles of surface chlorophyll concentration from model simulation (black solid line), satellite data (dotted line; SeaWiFS (diamond) and MODIS (square)) and in-situ data (digitised from Pennington et al. (2006, star, 250 km band off the coast) and Echevin et al. (2008, cross)). Error bars indicate the standard deviation.

3 Results

3.1 Anticorrelation of chlorophyll and upwelling: The seasonal paradox only appears in the Peruvian upwelling system

150

Compared with other EBUS (spatial extent of EBUS regions indicated in Fig. A2), the Peruvian system is unique in that it shows a clear anticorrelation between surface chlorophyll concentration and upwelling intensity on a seasonal scale, with lowest chlorophyll concentrations when upwelling is most intense (Fig. 2a, $R^2 = 0.71$; Chavez and Messié, 2009). While the surface chlorophyll in the Benguela system does not feature a strong seasonality, surface chlorophyll closely follows upwelling intensity in the California ($R^2 = 0.92$) and Canary ($R^2 = 0.88$) systems, suggesting that upwelling of nutrient-rich waters fuels the increase in chlorophyll. Indeed, comparatively low surface nitrate concentrations indicate that nitrate is depleted, potentially

155

limiting phytoplankton growth throughout the year in the California system and for approximately half the year in the Canary system (Fig. 2b). In contrast, the Benguela ($R^2 = 0.63$) and Peruvian systems ($R^2 = 0.90$) feature replete surface nitrate over most of the year. Because higher nitrate concentrations correlate with lower chlorophyll in these cases, nitrate is not observed
160 to be a limiting factor.

In the Peruvian system, a strong relationship exists between deepening mixed layers and decreasing chlorophyll (Fig. 2c, $R^2 = 0.91$), which supports the notion that dilution of phytoplankton over a deeper mixed layer and/or light limitation plays a role, as found by Echevin et al. (2008). The California system shows a similar response to mixed layer variations ($R^2 = 0.62$), suggesting that the same process may play a role there as well. Additionally, surface chlorophyll shows significant correlation
165 with SST in the Peruvian (Fig. 2d, $R^2 = 0.73$) and California systems ($R^2 = 0.65$), suggesting that increasing temperatures stimulate phytoplankton growth.

Strikingly, the Peruvian system is the only one of the four EBUS where high-strong upwelling coincides with deep MLD (Fig. 2e, $R^2 = 0.79$). The Canary and Benguela systems exhibit pronounced seasonality either in upwelling or in mixed layer depth, respectively. In the California system, the relationship of upwelling and mixed layer depth is opposite to that of the
170 Peruvian system, with the highest-strongest upwelling occurring in the shallowest mixed layers. Given the paradox that strong upwelling in the Peruvian system occurs at the time of the yearly chlorophyll minimum, it is intuitive that the concurrent deep mixed layers offset the positive impact of upwelled nutrients. Nutrient enrichment would only stimulate higher productivity if the region was nutrient limited. If concentrations are already elevated, adding more nutrients would have a weak impact. We will further investigate the interplay of the seasonality of mixed layers and upwelling in the Peruvian system in the following
175 sections.

3.2 Modelled phytoplankton biomass, dissolved inorganic nitrogen, upwelling and the MLD in the Peruvian system

We used a regional ocean circulation model, coupled to a marine biogeochemical model (CROCO-BioEBUS), to further analyse the Peruvian system (see the “Data and Methods” section). The model effectively reproduced the observed estimate of the seasonal out-of-phase relationship between surface phytoplankton biomass and upwelling intensity as well as nitrate concentrations (Fig. 2, open circles). Over the course of the year, surface chlorophyll, surface nitrogen concentrations, upwelling intensity and MLD varied by 40 - 60% relative to their annual mean values. Surface phytoplankton biomass concentration reached its maximum from late austral summer to early autumn (March to April), when upwelling was relatively weak (Fig. 3a,b). During this time window, less nitrogen is available within a shallow MLD compared with the rest of the year. In austral winter (July to September), when upwelling introduces ample nitrogen into the deep mixed layer, surface phytoplankton
185 concentration reaches a minimum.

3.3 Biomass dilution by the deepening mixed layer

Dilution of phytoplankton in deepening winter mixed layers is a key driver behind the seasonality of surface phytoplankton concentration. Within the research area, the MLD showed a seasonal variation with the shallowest mixed layer in austral summer, around 10 m, and the deepest mixed layer in austral winter, around 45 m. Phytoplankton were vertically well mixed




Figure 2. Correlations of surface chlorophyll (SeaWiFS climatology, in mg m⁻³) with (a) upwelling (a combination of Ekman transport and Ekman pumping, estimated based on winds from QuikSCAT, in Sv, digitised from Chavez and Messié (2009), for calculations see Messié et al. (2009)); (b) surface nitrate concentration (WOA, in mmol N m⁻³); (c) MLD (ARGO, in m); (d) SST (MODIS, in °C) and (e) correlation of MLD and upwelling transport among four eastern boundary upwelling systems (EBUS). For the Peruvian system, we also show the model (CROCO-BioEBUS) results. Lines and R² values are displayed for correlations with R² > 0.5.

190 within the mixed layer throughout the year (Fig. 3c). In austral winter, within the ‘deep-mixing’ regime, phytoplankton were evenly distributed over a relatively deep mixed layer, diluting phytoplankton biomass. Accordingly, phytoplankton biomass




Figure 3. Seasonal cycles of (a) upwelling intensity (in Sv, solid line) and surface dissolved inorganic nitrogen (DIN) concentration (in mmol N m^{-3} , dotted line); (b) surface (in mmol N m^{-3} , solid line) and mixed layer depth (MLD)-integrated phytoplankton biomass (in mmol N m^{-2} , dotted line); (c) phytoplankton depth-month distribution, showing the seasonal cycle of MLD (in m, solid line) within the focus region. The shaded area indicates the decline phase of MLD-integrated phytoplankton biomass.

concentrations in the mixed layer as well as at the surface decreased. Hence, we infer that seasonal mixed layer deepening and shoaling alone is an important factor in driving phytoplankton concentrations at the ocean surface, as observed for instance from satellite images.

195 While dilution caused a decrease in winter surface phytoplankton biomass, it explained only part of the observed biomass decrease. The decline persisted, even though attenuated, when integrating phytoplankton over the mixed layer (Fig. 3b). The phytoplankton concentration at the surface and within the mixed layer declined by around 70%, while it declined by around 30% for MLD-integrated biomass between late April and late July (shaded area in Fig. 3 & 4, hereafter referred to as the decline phase). The decline of surface phytoplankton concentrations can be attributed to the decline due to the increase in
 200 mixed layer volume ΔV (dilution effect, see Eq. 5) and the decrease in biomass ΔB within the mixed layer through local biological and physical processes (see Eq. 5). During the decline phase, ΔV contributed by slightly more than half to the concentration change, while ΔB contributed slightly less than half. That is, the dilution effect due to the deepening mixed layer in the decline phase amplified the decline of surface biomass concentrations by approximately a factor of two. However,

dilution could not fully explain the low phytoplankton biomass in conditions where the supply of nitrogen is ample; in such
205 conditions, MLD-integrated biomass still declined by around 30%.

3.4 Biological and physical processes change the total biomass within the mixed layer

3.4.1 Disentangling physical and biological processes

In addition to causing dilution due to the deepening mixed layer, the imbalance of a series of biological and physical processes
during the decline phase diminished phytoplankton concentrations. To disentangle their contributions to the decline of phyto-
210 plankton concentration without the complicating factor of the dilution effect, we next analysed the change of phytoplankton
biomass integrated over the mixed layer (Fig. 4a) and its drivers; that is, the mixed layer budget of phytoplankton biomass
(Eq. 1). We separated biological processes (e.g., primary production, grazing from zooplankton, natural mortality, exudation
and sinking) and physical processes (mixing ~~-, advection and entrainment~~ and advection) that affect the integrated biomass.
Throughout the year, the net biological flux (the sum of all biological fluxes) was positive ("biological gain", Fig. 4b), thus
215 supporting an increase in biomass. In contrast, the net physical flux, the sum of all physical fluxes, was negative ("physical
loss"), therefore supporting a decrease in biomass. The time point t1 marks the seasonal maximum of the MLD-integrated
phytoplankton biomass, and t2 marks the minimum at the end of the decline phase. At t1 and t2, the net biological and physical
fluxes balanced (Fig. 4b, eb-c) and the tendency of the mixed layer phytoplankton biomass was zero (Fig. 4a). Between t1
and t2 (Fig. 4b), the net biomass supply due to biological fluxes decreased more quickly than the net biomass removal due to
220 physical fluxes, resulting in an imbalance of the fluxes and the decrease in biomass between t1 and t2.

To determine which terms from Eq. 1 mostly drove the decrease of the biomass between t1 and t2 (Fig. 5a), we integrated the
change of each term over time (that is, the derivatives) between t1 and t2. Therefore, in Fig. 5b and c, if a bar was positive(red),
the change of the term during the decline phase (t1 - t2) promoted an increase of the phytoplankton biomass, mostly as a result
of reduced grazing pressure and reduced downward mixing. If the bar was negative(grey), the change of the term during the
225 decline phase (t1 - t2) opposed an increase in phytoplankton biomass. The "opposing terms" that acted to reduce phytoplankton
biomass were the ones that contributed to the seasonal paradox; that is, decline in biomass despite increased supply of nutri-
ents due to upwelling. These terms mostly referred to the reduced primary production, with a secondary contribution from the
~~reduced convergence~~ increased divergence due to advection, ~~as well as a small effect due to detrainment~~. Details regarding the
two major contributors, primary production and advection, are presented in the following sections.

230

3.4.2 Factors limiting primary production

Primary production changed due to variations in both the growth factor and the biomass (Eq. 2). The growth factor (calculated
as in Eq. 3, Fig. 6a) combined the effects of light, temperature and nitrogen on phytoplankton growth. It showed a clear decrease
of around 30% during the decline phase. Optimal phytoplankton growth conditions were reached in March, despite the low




Figure 4. Seasonal cycles of (a) total phytoplankton biomass change (ΔB ; in $\text{mmol N m}^{-2} \text{d}^{-1}$). Grey shading indicates the decline phase, with t1 and t2 marking the beginning and end of the decline phase; (b) phytoplankton fluxes resulting from net biological gain (bio) and net physical loss (phy, in $\text{mmol N m}^{-2} \text{d}^{-1}$, as shown in Eq. 1 plus the entrainment and detrainment introduced by the temporal variation of the MLD); (c) Stacked bar plots displaying balancing budgets of phytoplankton fluxes at t1 and t2 (PP stands for primary production; GRAZ for: consumptive mortality; MORT for: natural mortality; EXU for: exudation; SINK for: sinking; MIX for: mixing; ADV for: advection and ENTR for entrainment). Fluxes are distinguished as sources (positive values) and sinks (negative values) of phytoplankton biomass, with the sum of all source and sink terms balancing at times t1 and t2. All fluxes are integrated over the MLD.

235 dissolved inorganic nitrogen (DIN) conditions, within the warmest and brightest environment. The lowest growth rate occurred just after the decline phase, despite relatively high nitrogen concentrations, due to limiting light and temperature conditions.

Strong light limitation experienced by phytoplankton, in combination with low temperatures, slowed growth during the decline phase. Light conditions for phytoplankton growth were optimal in March when the water was rather stratified and worsened over the decline phase to a minimum in August when the water column was most deeply mixed (Fig. 6a). The light-related growth factor declined by 17% during the decline phase and would decrease the growth factor by approximately 60% in the absence of other limiting factors, estimated from (estimated from the product rule for differentiation and the multiplicative relation of growth factors shown in Eq. 32 & 3). Decreasing temperature was the second most important contributor in slowing the growth rate during the decline phase. The temperature-related growth factor reached its maximum by March, similar to the light-related growth factor, and reached its minimum by October. The temperature-related growth factor declined by 12% and would decrease the growth factor by around 40% during the decline phase, in the absence of other limiting factors. In contrast, the seasonality of the growth factor due to nitrogen showed the opposite seasonality compared to the total growth factor. Clearly, light and temperature regulated primary production and overrode the effect of enhanced nitrogen supply during the decline phase. Therefore, while light was the dominant mechanism that reduced productivity towards winter, we found that temperature played a relevant secondary role.

245




Figure 5. (a) Seasonal cycle of phytoplankton source and sink processes, as well as phytoplankton biomass change (ΔB multiplied by a factor of 10; solid line); bar plots of the integrated change over the decline phase due to (b) biological fluxes and (c) physical fluxes averaged over the focus region (PP: primary production; GRAZ: consumptive mortality; MORT: natural mortality; EXU: exudation; SINK: sinking; MIX: mixing; ADV: advection). Red and grey shading indicate a positive or negative sign of the bars here designates fluxes promoting and opposing an increase in MLD-integrated phytoplankton biomass, respectively. The magnitude (including the sign) of the integrated change is given as numbers above and below the bars. PP stands for primary production; GRAZ for consumptive mortality; MORT for natural mortality; EXU for exudation; SINK for sinking; MIX for mixing; ADV for advection and ENTR for entrainment. All fluxes are integrated over the MLD.




Figure 6. (a) Seasonal cycles of the normalised total (black) growth factor and light (yellow)-, temperature (red)- and nitrogen (blue)-related growth factors for phytoplankton over the mixed layer. The grey shading indicates the decline phase. (b) Seasonal correlation of MLD and mixed layer-averaged light-related growth factor; (c) correlation of upwelling intensity and mixed layer-averaged temperature-related growth factor. Colours indicate the time of the year (months), with black edges indicating the months of the decline phase. The R^2 values of the correlations are shown in the right-hand sides within the panels.

250 Stronger light and temperature limitation during the decline phase were due to deeper mixing and stronger upwelling of cold waters, respectively (Fig. 6b-c). While upwelling intensity was approximately correlated with MLD, the maximum upwelling occurred just after the deepest mixed layers. The variation of MLD-averaged light limitation was correlated ($R^2 = 0.92$) with the change of MLD. As phytoplankton were evenly distributed within the mixed layer, deeper MLD indicated that, on average, more phytoplankton were exposed to lower light conditions during the decline phase, with a minimum in August when mixed
 255 layers were deepest. The change of the temperature-related growth factor within the mixed layer was closely related with the seasonal variation of upwelling intensity ($R^2 = 0.83$), with the lowest values occurring in September and October when upwelling intensity reached its maximum. During the decline phase, cold waters were upwelled into the mixed layer at a higher rate, further damping phytoplankton growth in addition to the effects of limiting light conditions. Reduced winter surface solar

radiation and heat loss to the atmosphere also played a role in the seasonality of the light and temperature growth factors, respectively (Fig. C2), though to a much smaller extent (not shown) [which agrees with findings in Echevin et al. \(2008\)](#).

3.4.3 Enhanced upwelling and offshore transport of phytoplankton

An enhanced advective loss of mixed layer phytoplankton is a second-order process, promoting the decrease of MLD-integrated phytoplankton biomass during the decline phase [\(Fig. 5c\)](#). Similar to the effects of nutrients, phytoplankton biomass is affected by the seasonality of upwelling and offshore export of waters. Relatively dilute concentrations of phytoplankton growing below the base of the mixed layer are upwelled into the mixed layer, while waters with mixed-layer-averaged phytoplankton concentrations are pushed offshore. During the decline phase, the upwelling and offshore transport of water increased and a greater volume of what was produced in coastal waters was exported offshore: 4% of primary production was lost via advection by the end of the decline phase compared to 2% gained at the beginning. This greater loss of biomass due to divergent lateral advection was mainly caused by stronger upwelling during the decline phase (Fig. C3).

3.5 Seasonal Paradox: from phytoplankton to export

Small and large zooplankton exhibit the same "seasonal paradox" pattern as phytoplankton, and so does the export of organic material to the deeper ocean. Similar to phytoplankton, both small and large zooplankton are vertically well mixed within the mixed layer throughout the year (contours and colours in Fig. 7a, respectively); ~~biomass~~ [Biomass](#) concentrations are high in austral summer and low in austral winter, in opposition to the upwelling trend. Additionally, the particulate organic matter, the sum of plankton biomass and other organic particles, follows the same pattern, with large amounts of particulate organic matter concentrated in a shallow mixed layer during the productive summer (Fig. 7b). The pattern of organic matter in the water column is then reflected in the export pattern of sinking organic material, composed of large phytoplankton as well as small and large detritus (Fig. 7c). Export below 100 m depth is high during the productive summer, when the mixed layer is shallow and particulate organic matter is large, and low in winter (Fig. 7b, black line). That is, the model's ecosystem is affected by the seasonal variation of the MLD.

Finally, export efficiency also follows the seasonal cycle of the MLD. Export efficiency is defined as the export of sinking organic material through the 100 m depth level, relative to primary production in the upper 100 m. It reaches a maximum in austral summer, when MLD is shallow, and a minimum in austral winter, when MLD is deep (Fig. 7d). As both export and primary production show the same seasonal trend as phytoplankton biomass, export must overcompensate the change in primary production and vary even more, in order to allow export efficiency to reveal the same seasonal trend. Export largely consists of large detritus originating from large zooplankton faecal pellets and mortality (Fig. 7c). Since large detritus possesses the fastest sinking speed compared to other components, it sinks the most efficiently. The relative contribution of fast-sinking large detritus to total export is largest in summer, close to 100 %, which may partially explain the higher export efficiency. In addition to changes in composition of the sinking organic material, other processes may cause export to be amplified relative to phytoplankton production. These include: (1) changes in structure and trophic transfer efficiency of the plankton food web, and (2) a varying degradation of sinking organic matter in the upper 100 m, that is, differences in the remineralisation. The

detailed mechanisms behind the seasonality of export efficiency are beyond the focus of this paper and will be investigated in a separate study.

4 Discussion

295 4.1 Mixed layer depth drives surface phytoplankton biomass seasonality in the Peruvian upwelling system

The regional ocean circulation-biogeochemical model that we used successfully reproduced the "seasonal paradox", defined as the seasonal out-of-phase surface chlorophyll concentration and upwelling intensity, as derived from observations. As shown in the results, the low surface chlorophyll concentration in high-strong upwelling conditions during austral winter was constrained by a combined effect of MLD-driven and upwelling-driven processes. Under strong upwelling conditions during austral winter, 300 phytoplankton were-was diluted over a deeper mixed layer, leading to a decrease within the mixed layer. Likewise, surface phytoplankton concentrations decreased by over 50%. Also, phytoplankton growth was slowed due to deteriorating light and temperature conditions, as well as strong upwelling pushing phytoplankton offshore.

Several previous studies have also focused on the possible reasons behind the seasonal paradox in the PUS. Echevin et al. (2008) also used a regional ocean circulation-biogeochemical model to examine the reasons for the relatively low surface 305 chlorophyll concentration off the Peruvian coast in austral winter. Based on a series of model sensitivity experiments regarding vertical mixing, surface temperature, iron limitation and insolation cycle, Echevin et al. (2008) concluded that the low surface chlorophyll in austral winter is mainly generated by the combined effect of dilution and deteriorating light with deepening mixed layers. Additionally, the iron sensitivity experiment confirmed the existence of iron limitation in austral winter, which corroborated the findings in Messié and Chavez (2015). Messié and Chavez (2015) pointed out that more severe iron limitation 310 under low light could also be one of the reasons behind low primary production under strong upwelling conditions. According to results from culture experiments, phytoplankton iron demand would increase under light limitation (Sunda and Huntsman, 1997). Based on observations, Friederich et al. (2008) suggested that winds in the high-upwelling-strong-upwelling winter conditions favour curl-driven offshore upwelling, which would draw more offshore iron-deficient waters to the surface. On the contrary, a model study (Albert et al., 2010) found that stronger wind-curl-driven upwelling actually recruits more nutrient- 315 rich water from a shoaling coastal undercurrent, thus enhancing surface chlorophyll concentrations. We could not assess the role of iron in regulating the seasonality of phytoplankton biomass because our biogeochemical model did not simulate iron. Nevertheless, our study confirmed the importance of vertical redistribution of biomass and light limitation due to vertical mixing.

4.2 Upwelling into deep mixed layers: A unique feature of the Peruvian upwelling system and its implications

320 As stated in the previous paragraphs, based on the differences in the seasonalities of MLD and upwelling in the Peruvian system, upwelling of nutrient-rich waters occurs when growth conditions are least optimal, in particular when light availability is lowest due to deep mixed layers. In contrast, within the California system, nutrients are upwelled into the shallowest mixed




Figure 7. Monthly depth distribution of (a) large and small zooplankton (in colours and contour lines, respectively) and (b) organic matter with the seasonal cycle of vertical export through the 100 m depth horizon (black line); (c) Seasonal cycle of the relative contributions to sinking organic matter from large phytoplankton (green) and small (blue) and large (purple) detritus; (d) correlation of export efficiency with mixed layer depth (MLD) within the focus region. The export efficiency is defined as the ratio of export through the 100 m depth level to primary production in the upper 100 m. Colours indicate the month of the year.

layers. While this nutrient supply coincides with shoaling mixed layers, associated with improved light conditions and reduced dilution, it does not result in as high a level of phytoplankton concentrations as for the Peruvian system. This supports that
325 nutrient limitation contributes substantially to processes in the California system, as the supply of nutrients to shallow mixed
layers through upwelling appears to be insufficient to relieve nutrient limitation. Additionally, if nutrients are upwelled into
deep mixed layers [as in the Peruvian system](#) and allow the onset of a bloom, zooplankton standing stocks might be low and
require more time to catch up, eventually reducing phytoplankton biomass. On the contrary, if nutrients are upwelled into
shallow mixed layers, zooplankton standing stocks are likely already elevated, allowing zooplankton to immediately limit any
330 increase in phytoplankton biomass.

While the Canary and Benguela systems lack a pronounced seasonality in MLD and phytoplankton, respectively, we point
out a few aspects that may elucidate the role of MLD in these systems. Given that the Canary system does not feature a sub-
stantial seasonal MLD variability, it is intuitive that phytoplankton follows the seasonality of upwelling intensity more strongly
compared to the other EBUS. While mixed layer conditions do not modulate the seasonality of phytoplankton, they may con-
335 tribute to high phytoplankton concentrations in the Canary system insofar as mixed layers are shallow throughout the year,
creating favourable light conditions. Finally, the Benguela system features a rather constant upwelling throughout the year
into varying mixed layers. The unresponsiveness of phytoplankton to the varying MLD could hypothetically be due to com-
pensating effects of deepening mixed layers that dilute phytoplankton and deteriorate light conditions, but are simultaneously
accompanied by an enhanced supply of nutrients that ~~is~~ [are](#) mixed up from below.

340 Other factors may also contribute to regulating phytoplankton in the EBUS aside from nutrients, dilution and light associ-
ated with upwelling and MLD(~~also see the Results section; Messié and Chavez, 2015~~), including the advection of biomass and
regulation by temperature that varies with upwelling (~~see Results~~)-([also see the Results section; Messié and Chavez, 2015](#)).
[Eddies have been found to favour offshore export and subduction of phytoplankton and nutrients \(Lathuilière et al., 2010; Gruber et al., 201](#)
In addition, Lachkar and Gruber (2011) suggest that a longer residence time because of a wide shelf and weak mesoscale
345 activity may ~~also~~ promote phytoplankton growth in the Canary system. Next to iron supply from the shelves and upwelling of
source waters, Fung et al. (2000) also found that atmospheric deposition of iron varies between EBUS.

4.3 Seasonal paradox and ecosystem functioning

The interplay of mixed layer depth and upwelling that leads to the seasonal paradox in the PUS propagates further up the food
chain, modulating the trophodynamics. In austral summer, the shallow mixed layer along with the add-on effect from upwelling
350 supports the highest phytoplankton biomass and primary production, providing an ideal feeding place for zooplankton. In
contrast, during the winter zooplankton face a food shortage, less efficient grazing due to dilution, and transport offshore due
to enhanced upwelling. Similar to the spatial match-mismatch observed for phytoplankton and top predators in the Benguela
system (Grémillet et al., 2008), mesozooplankton with ~~their slower growth rates~~ [its slower growth rate](#) may also be negatively
affected by enhanced upwelling.

355 In our model, mesozooplankton ~~are~~ [is](#) responsible for the major part of the export in the coastal upwelling region. During
the productive season, the faecal material of mesozooplankton accounts for close to 100% of the sinking matter, which is in

good agreement with what Stukel et al. (2013) observed for the California system. ~~Both~~ We found that both primary production and export can be determined from the mixed layer dynamics and food web structures (consistent with Ducklow et al., 2001; Turner, 2015; Steinberg and Landry, 2017). The efficiency of the export, defined as the ratio of export to primary production, is also related to trophodynamics. We find that export efficiency is positively correlated with MLD on a seasonal scale. As mentioned ~~above~~ in the "Results" section, it partially depends on the composition of the exported material. Mesozooplankton produce fast-sinking large detritus, which enhances the export efficiency during the productive season. Kelly et al. (2018) observed that export efficiency is negatively correlated with net primary productivity in the California system. They suggested that the negative correlation in the California system arises from a seasonal decoupling of export and particle production through long-lived particles that introduce a temporal lag of mesozooplankton production and export to depth. Henson et al. (2019) also identify a negative correlation between export efficiency and primary productivity on a global scale. They imply in their study that not just the phytoplankton community, but also the food web structure, is important to export efficiency. Currently, it is not entirely clear why the PUS export efficiency behaves differently in our model. We suggest that the interplay of the mixed layer and upwelling in EBUS and ecosystem functioning are closely linked, warranting further examination.

370 5 Conclusions and potential implications

In summary, CROCO-BioEBUS performs well with respect to observational data and successfully reproduces the "seasonal paradox" with an out-of-phase relationship between surface chlorophyll and upwelling intensity in the Peruvian coastal waters. In agreement with an earlier model study (Echevin et al., 2008), the seasonal cycle of surface chlorophyll concentration in our simulations is driven mostly by MLD-related processes, specifically dilution and light limitation. Furthermore, our model results provide evidence for secondary contributions from upwelling-related processes such as temperature limitation and advection. This is consistent with Lachkar and Gruber (2011) and Messié and Chavez (2015), who suggested that advection is relevant to the seasonal cycle, but in contrast to Echevin et al. (2008), who found that temperature was not important. Differences in results from Echevin et al. and our results likely originate in the different biogeochemical model components (e.g., different parameterisations of temperature dependencies of phytoplankton growth). Given the disparity of the models, the role of temperature limitation in the PUS warrants further investigations in order to better constrain second-order drivers of the seasonal paradox. The sensitivity of the different processes within the plankton ecosystem to temperature, as well as their interplay, are topics of active research (e.g. Thomas et al., 2017; Chen and Laws, 2017; Morán et al., 2018; Marañón et al., 2018; Barton and Yvon-Durocher, 2019) and relevant particularly in light of global warming.

We find that the seasonal variability of phytoplankton propagates up the food chain and is reflected in trophodynamics and ecosystem functioning. In particular, zooplankton and organic matter within the water column mirror the seasonal cycle of phytoplankton. Finally, export and export efficiency are well-correlated with the MLD over the course of the annual cycle. Given that changes in MLD are correlated to many ecosystem components related to plankton ecosystem functioning, we argue for a more thorough understanding of the ~~interactions behind the mixed layer and upwelling dynamics, along with food web processes.~~ A impact of the seasonal paradox on the ecosystem. In particular effects on the trophic transfer of energy through the

390 plankton food web to higher trophic levels such as fish will determine ecosystem functions like trophic transfer efficiency, fish
production, and ultimately potentially fisheries yields. Thus, a better understanding of how the interplay of MLD and upwelling
impacts the ecosystem in the contemporary PUS will ultimately help to better project how coastal upwelling ecosystems, and
in particular the Peruvian system, may vary under climate change.

Phytoplankton will inevitably be influenced by climate change, responding to changes in the biotic and abiotic environment.
395 Impacts in a changing climate will arise from changes in stratification and upwelling, that further lead to shifting growth
conditions due to changes of light, temperature and nutrients (Behrenfeld, 2014). A recent regional modelling study (Echevin
et al., 2020) projects a weak decrease in upwelling along with increasing stratification in the PUS due to climate change.
Our results suggest that the decreasing upwelling and increasing stratification will both contribute to an increase in surface
phytoplankton, in agreement with the findings of Echevin et al. (2020). While a reduction of upwelling might lead to a reduced
400 supply of nutrients, the region is far from being nutrient limited. Therefore, a reduction in upwelling could rather have an effect
via temperature, reducing the cooling effect of upwelled waters. We hypothesise that the coastal region would experience more
phytoplankton growth and biomass buildup with a reduction of upwelling, due to warmer surface waters and weaker offshore
advection compared to the current environmental situation. ~~Aecording~~ Moreover, according to our results, shoaling of the
mixed layer will be more relevant than a decrease in upwelling intensity, reducing the dilution of phytoplankton and the light
405 limitation in austral winter. This could possibly lead to an attenuation of the seasonal paradox in the future. As export and
export efficiency are also regulated by MLD dynamics, we expect not only enhanced export but also an increase in the fraction
of primary production that is transported to the deep ocean under global warming.

Code availability. CROCO and BioEBUS models are available at <http://www.croco-ocean.org>

Data availability. The model data used in this paper are available via the corresponding author

410 **Appendix A: Methods**

A1 Two-way nesting approach

Figure A1 visualises the coarser-resolution parent domain and nested finer-resolution child domain that contains the focus
region. The variables in section B are shown for the child domain.

A2 Adjustment of biogeochemical model parameters

415 The parameter setting is the same as in José et al. (2017), with only a few biological parameters adjusted to make the ecol-
ogy (phyto- and zooplankton biomasses, productivity) better fit observational data. The changed parameters along with value
ranges from literature are listed in Table A1 and will be further explained below.




Figure A1. Bathymetry of the "parent" ($1/4^\circ$ resolution) and "child" ($1/12^\circ$ resolution) domains. White lines near the coast highlight the focus region.

Here, we assign a higher mortality rate for large phytoplankton to simulate the potential impact of virus infection during bloom conditions (Suttle, 2005). Simulated phytoplankton biomass and its seasonality has been calibrated and evaluated against chlorophyll concentration data from MODIS monthly climatology data (<https://oceancolor.gsfc.nasa.gov/>). Nitrate has been evaluated based on [WOA-World Ocean Atlas \(WOA; Garcia et al., 2019\)](#) and cruise data ([M92 and M93 Thomsen et al., 2016](#)) while simulated MLD has been validated against the ARGO mixed layer database ([Holte et al. \(2017\)](#), <http://mixedlayer.uesd.edu/>).

425 Appendix B: Model evaluation

B1 Surface chlorophyll concentration

The large-scale spatial pattern of annual average surface chlorophyll of the monthly climatology of MODIS data and CROCO-BioEBUS are similar (Fig. B1), with higher chlorophyll concentrations in coastal regions and lower concentrations offshore (note that chlorophyll is shown in log-scale). The satellite data features a higher cross-shore chlorophyll concentration gradient compared to the model simulation. The model's overestimation of the low offshore chlorophyll and hence weaker cross-shore




Figure A2. Map of annual mean surface chlorophyll ($mg\ chl\ m^{-3}$) with white lines highlight the regions that we average over in our analyses in Fig.2. Coastal EBUS regions picking here are the same as Chavez and Messié (2009)

gradient potentially is due to the lack of iron limitation in the model. Apart from that, the model is also not able to correctly capture the alongshore pattern (Fig. B1), i.e. it misses two observed high surface chlorophyll concentration patches between 8°S to 10°S and 12°S to 14°S (Bruland et al., 2005). Within a 200 km band near the coast, both satellite data and the model simulation show a similar seasonality with maximum chlorophyll concentrations exceeding 4 $mg\ /m^3$ from March to April and minimum concentrations around 2 $mg\ /m^3$ in August. In general, simulated surface chlorophyll concentrations agree reasonably well with satellite data.

Table A1. Adjusted biological ~~Parameters~~ parameters and range of published parameter values

Parameters	Symbols	Units	Value	Range
Max growth rate of P_L	a_{P_L}	d^{-1}	0.6	0.6^a - 3.0^b
Mortality rate of P_L	μ_{P_L}	d^{-1}	0.15	0.027^c - 0.2^d
Preference of Z_S for P_S	$e_{Z_S P_S}$	-	0.65	see references ^e
Preference of Z_S for P_L	$e_{Z_S P_L}$	-	0.35	see references ^e
Preference of Z_L for P_S	$e_{Z_L P_S}$	-	0.1	see references ^{f,g}
Preference of Z_L for P_L	$e_{Z_L P_L}$	-	0.4	see references ^{f,g}
Preference of Z_L for Z_S	$e_{Z_L Z_S}$	-	0.5	see references ^{f,g}
Excretion rate of Z_S	γ_{Z_S}	d^{-1}	0.1	0.03^h - 0.1^i
Excretion rate of Z_L	γ_{Z_L}	d^{-1}	0.1	0.05^h - 0.1^i
Mortality rate of Z_L	μ_{Z_L}	$\text{mmol N m}^{-3} \text{d}^{-1}$	0.135	0.05^a - 0.25^j

The values for diet preferences were picked based on a combination of calibrating the model against observations of plankton biomasses and observed qualitative diet preferences in the references.

^a Gutknecht et al. (2013)

^b Andersen et al. (1987)

^c Koné et al. (2005)

^d Taylor et al. (1991)

^e Bohata (2016)

^f Kleppel (1993)

^g Schukat et al. (2014)

^h Aumont (2005)

ⁱ Fennel et al. (2006)

^j Lima and Doney (2004)




Figure B1. Annual mean surface chlorophyll concentration (in $\log(\text{chl} (\text{mg m}^{-3})^{-1})$) distribution of (a) MODIS and (b) CROCO-BioEBUS. White lines highlight the focus region.

B2 Surface nitrate concentration

The simulated surface nitrate distribution shows the same seasonality as observations from the World Ocean Atlas (WOA; Garcia et al., 2019) (Fig. B2). The simulated surface nitrate concentration in the coastal region is biased high compared to the




Figure B2. Spatial distribution of surface nitrate concentration based on (a) WOA and (b) CROCO-BioEBUS; (c) January and (d) February as simulated by CROCO-BioEBUS. Dots indicate measurements from the cruises M92 (January) and M93 (February); (e) ~~seasonal~~Seasonal cycle of surface nitrate concentration from WOA (cross), CROCO-BioEBUS (line) and cruises (pentagram, hexagram) within the focus region. White lines highlight the focus region. The black box indicates the maps of panel c-d.

WOA data. This may be partly due to the WOA data failing to capture high-nitrate concentrations due to coastal upwelling. This notion is supported by nitrate concentration data from ~~a cruise~~two cruises (Thomsen et al., 2016, M92 and M93) in austral summer that show nitrate concentrations in the coastal region are high compared to the model data.




Figure B3. Annual average spatial distribution of mixed layer depth (MLD) from (a) ARGO and (b) CROCO-BioEBUS; (c) Seasonal variation of average mixed layer depth from ARGO (crosses blue), with lines indicating the standard deviation of Boyer Montégut climatology (purple) and model simulation (line black) with error bar indicating the standard deviation within focus region. White lines highlight the focus region.

B3 Mixed layer depth

445 We validate the simulated MLD against both the gridded ARGO mixed layer dataset (Holte et al. (2017), <http://mixedlayer.ucsd.edu/>) both in terms of spatial pattern and seasonal variability and the de Boyer Montégut climatology mixed layer data available from the IFREMER/LOS Mixed Layer Depth Climatology website (www.ifremer.fr/cerweb/deboyer/mld) within the research area (Fig. B3). All observational data and simulated MLD are calculated as the depth where a 0.2°C difference to the surface temperature is reached. The annually averaged spatial distribution of MLD within the research area presents the same features as ARGO: shallower MLD in the coastal region (around 20 m) and deeper MLD in the offshore region (around 80 m). The simulated seasonal variability of MLD within the research region generally follows the seasonal trend of the Argo-ARGO and the Boyer Montégut climatology data. The water column within the research region is most stratified in February to March and most deeply mixed in August. Although simulated MLD in austral winter is somewhat deeper than Argo and the de Boyer Montégut climatology data, the simulated MLD and the de Boyer Montégut climatology data are largely within the range of the ARGO data. Deeper simulated MLD compare to observation could partially come from not including chlorophyll shading effect on water cooling (Echevin et al., 2021).

450

455




Figure B4. Annual average spatial distribution of sea surface temperature (SST, in °C) from (a) MODIS and (b) CROCO-BioEBUS; (c) Seasonal variation of average sea surface temperature (SST) from MODIS (cross) and model simulation (line) within focus region. White lines highlight the focus region.

B4 Sea surface temperature

The simulated SST has been validated against monthly climatological MODIS data in terms of both spatial pattern and seasonal variability within the research area (Fig. B4). The annually averaged spatial distribution of SST is well simulated by the model. The model successfully captures the cold coastal upwelled water as well as slightly warmer water masses further offshore. The simulated SST seasonality within the research region generally follows the seasonal trend of the observations, with a cool bias of less than 1°C. The surface waters within the research region are warmest in February to March matching the modelled/observed shallowest mixed layers and coldest from August to October. In general, the simulated SST matches the observations well both in terms of spatial pattern and seasonal variation.

465 B5 Mesozooplankton distribution

In addition, we calibrated zooplankton in the BioEBUS model against observational estimates (Fig. B5). ~~It is something that is~~ [Calibration and assessment of simulated zooplankton is](#) often omitted, despite the central role of zooplankton parameterisations on plankton dynamics (Anderson et al., 2010; Prowe et al., 2012). While the observations show a large spread, the simulated large-scale spatial distribution of mesozooplankton generally follows the observed pattern, with high mesozooplankton biomass




Figure B5. Annual average spatial distribution of integrated mesozooplankton biomass over upper 200 m based on (in mmol N m^{-2}) : Dots indicate mesozooplankton observational data from observations (Moriarty and O'Brien, 2013, <https://doi.org/10.5194/essd-5-45-2013>) and (b) model simulation. White lines highlight (c) Scatter plot of observed and simulated integrated mesozooplankton biomass over upper 200 m (in mmol N m^{-2}). The dashed line indicates the focus region 1:1 line.

470 in the upwelling region and low biomass further offshore. The model overestimated simulated zooplankton biomass compared to the observational data in the offshore region is likely partially related to the overestimated offshore phytoplankton biomass, which in turn presumably results from the lack of iron limitation in the model. As shown in Fig. B5c, most data points fall close to the 1:1 line. However, the model is not able to capture the few data points with very high zooplankton biomass. The model simulates a stripe of low zooplankton biomass concentrations in the focus region near the coast (due to offshore advection combined with slow mesozooplankton growth) that is difficult to assess, as observations near the coast are sparse.

475 Though, the feature is This feature may be apparent to some extent in the observations in the southern focus region. Note that observational zooplankton biomass estimates are based on a wide range of methods and accordingly have a large uncertainty that is difficult to quantify (O'Brien, 2007). An agreement of model and observations in magnitude and large scale pattern is therefore a meaningful result.




Figure C1. Time series of temperature T (at 10 m & 100 m depth), nitrate N (at 10 m & 100 m depth), mixed layer depth MLD and phytoplankton phyto (at 10 m) over 30 years of simulation in the focus region.

480 Appendix C: Additional Figures

The whole time series of temperature and nitrate concentration at 10 m and 100 m are shown in Fig. C1a-b. Surface fields are spun up after one year while water at 100 m takes 3-10 years longer to reach a steady state. In the meanwhile, mixed layer and surface layer chlorophyll are also spun up after one year (Fig. C1c-d).




Figure C2. (a) Correlation between surface short-wave radiation (Wm^{-2}) and the averaged light-related growth factor within mixed layer; (b) Correlation between the surface heat forcing (in $^{\circ}\text{Cd}^{-1}$) and averaged temperature-related growth factor within mixed layer. Colour indicates the time of the year and black edges the decline phase.




Figure C3. Correlation between upwelling intensity and phytoplankton convergence of advection over the mixed layer. A negative convergence equals a divergence of phytoplankton biomass due to the combined effect of upwelling and lateral transports. Color indicates the time of the year and black edges the decline phase. The correlation coefficient ($R^2=0.81$) is shown for the decline phase.

Apart from above mentioned mixed layer depth and upwelling intensity, short-wave surface radiation and surface net heat flux are of second-order importance to light- and temperature-related variance during the decline phase respectively (Fig.C2).

Phytoplankton net advection flux over the mixed layer closely follows the upwelling intensity during the decline phase (Fig.C3, $R^2 = 0.81$). When the mixed layer depth is relatively shallow, the correlation between upwelling intensity and phytoplankton convergence of advection over the mixed layer is insignificant.

Author contributions. IF and AO designed the study. TX and YSJ carried out the simulations. TX, IF and FP conducted the analysis. All
490 authors discussed the results and wrote the manuscript.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. This work is financially supported by the China Scholarship Council (TX, grant no.201808460055). Further support for this work was provided by the BMBF funded projects Coastal Upwelling System in a Changing Ocean CUSCO (IF, AO) and Humboldt Tipping (YJ).

495 **References**

- Albert, A., Echevin, V., Lévy, M., and Aumont, O.: Impact of nearshore wind stress curl on coastal circulation and primary productivity in the Peru upwelling system, *Journal of Geophysical Research: Oceans*, 115, 2010.
- Andersen, V., Nival, P., and Harris, R. P.: Modelling of a planktonic ecosystem in an enclosed water column, *Journal of the Marine Biological Association of the United Kingdom*, 67, 407–430, 1987.
- 500 Anderson, T. R., Gentleman, W. C., and Sinha, B.: Influence of grazing formulations on the emergent properties of a complex ecosystem model in a global ocean general circulation model, *Progress in Oceanography*, 87, 201–213, 2010.
- Aumont, O.: PISCES biogeochemical model, Unpublished report, 2005.
- Bakun, A.: Coastal upwelling indices, west coast of North America, 1946-71, 1973.
- Bakun, A.: Global climate change and intensification of coastal ocean upwelling, *Science*, 247, 198–201, 1990.
- 505 Bakun, A., Field, D. B., Redondo-Rodriguez, A., and Weeks, S. J.: Greenhouse gas, upwelling-favorable winds, and the future of coastal ocean upwelling ecosystems, *Global Change Biology*, 16, 1213–1228, 2010.
- Barton, S. and Yvon-Durocher, G.: Quantifying the temperature dependence of growth rate in marine phytoplankton within and across species, *Limnology and Oceanography*, 64, 2081–2091, 2019.
- Behrenfeld, M. J.: Climate-mediated dance of the plankton, *Nature Climate Change*, 4, 880–887, 2014.
- 510 Bohata, K.: Microzooplankton of the northern Benguela Upwelling System, PhD thesis, 2016.
- Bruland, K. W., Rue, E. L., Smith, G. J., and DiTullio, G. R.: Iron, macronutrients and diatom blooms in the Peru upwelling regime: brown and blue waters of Peru, *Marine Chemistry*, 93, 81–103, 2005.
- Calienes, R., Guillén, O., Lostaunau, N., et al.: Variabilidad espacio-temporal de clorofila, producción primaria y nutrientes frente a la costa peruana, *Boletín Instituto del Mar del Perú*, 10, 1–44, 1985.
- 515 Carton, J. A. and Giese, B. S.: A reanalysis of ocean climate using Simple Ocean Data Assimilation (SODA), *Monthly Weather Review*, 136, 2999–3017, 2008.
- Chavez, F. P.: A comparison of ship and satellite chlorophyll from California and Peru, *Journal of Geophysical Research: Oceans*, 100, 24 855–24 862, 1995.
- Chavez, F. P. and Messié, M.: A comparison of eastern boundary upwelling ecosystems, *Progress in Oceanography*, 83, 80–96, 2009.
- 520 Chavez, F. P., Bertrand, A., Guevara-Carrasco, R., Soler, P., and Csirke, J.: The northern Humboldt Current System: Brief history, present status and a view towards the future, *Progress in Oceanography*, 79, 95–105, 2008.
- Chen, B. and Laws, E. A.: Is there a difference of temperature sensitivity between marine phytoplankton and heterotrophs?, *Limnology and Oceanography*, 62, 806–817, 2017.
- Debreu, L., Marchesiello, P., Penven, P., and Cambon, G.: Two-way nesting in split-explicit ocean models: Algorithms, implementation and validation, *Ocean Modelling*, 49, 1–21, 2012.
- 525 Ducklow, H. W., Steinberg, D. K., and Buesseler, K. O.: Upper ocean carbon export and the biological pump, *OCEANOGRAPHY-WASHINGTON DC-OCEANOGRAPHY SOCIETY*, 14, 50–58, 2001.
- Echevin, V., Aumont, O., Ledesma, J., and Flores, G.: The seasonal cycle of surface chlorophyll in the Peruvian upwelling system: A modelling study, *Progress in Oceanography*, 79, 167–176, 2008.
- 530 Echevin, V., Gévaudan, M., Espinoza-Morriberón, D., Tam, J., Aumont, O., Gutierrez, D., and Colas, F.: Physical and biogeochemical impacts of RCP8.5 scenario in the Peru upwelling system, *Biogeosciences*, 17, 3317–3341, 2020.

- Echevin, V., Hauschildt, J., Colas, F., Thomsen, S., and Aumont, O.: Impact of Chlorophyll Shading on the Peruvian Upwelling System, *Geophysical Research Letters*, p. e2021GL094429, 2021.
- 535 Fennel, K., Wilkin, J., Levin, J., Moisan, J., O'Reilly, J., and Haidvogel, D.: Nitrogen cycling in the Middle Atlantic Bight: Results from a three-dimensional model and implications for the North Atlantic nitrogen budget, *Global Biogeochemical Cycles*, 20, 2006.
- Friederich, G. E., Ledesma, J., Ulloa, O., and Chavez, F. P.: Air–sea carbon dioxide fluxes in the coastal southeastern tropical Pacific, *Progress in Oceanography*, 79, 156–166, 2008.
- Fuenzalida, R., Schneider, W., Garcés-Vargas, J., Bravo, L., and Lange, C.: Vertical and horizontal extension of the oxygen minimum zone in the eastern South Pacific Ocean, *Deep Sea Research Part II: Topical Studies in Oceanography*, 56, 992–1003, 2009.
- 540 Fung, I. Y., Meyn, S. K., Tegen, I., Doney, S. C., John, J. G., and Bishop, J. K.: Iron supply and demand in the upper ocean, *Global Biogeochemical Cycles*, 14, 281–295, 2000.
- Garcia, H., Weathers, K., Paver, C., Smolyar, I., Boyer, T., Locarnini, M., Zweng, M., Mishonov, A., Baranova, O., Seidov, D., et al.: *World Ocean Atlas 2018. Vol. 4: Dissolved Inorganic Nutrients (phosphate, nitrate and nitrate+ nitrite, silicate)*, 2019.
- Getzlaff, J., Dietze, H., and Oschlies, A.: Simulated effects of southern hemispheric wind changes on the Pacific oxygen minimum zone, *Geophysical Research Letters*, 43, 728–734, 2016.
- 545 Grémillet, D., Lewis, S., Drapeau, L., van Der Lingen, C. D., Huggett, J. A., Coetzee, J. C., Verheye, H. M., Daunt, F., Wanless, S., and Ryan, P. G.: Spatial match–mismatch in the Benguela upwelling zone: should we expect chlorophyll and sea-surface temperature to predict marine predator distributions?, *Journal of Applied Ecology*, 45, 610–621, 2008.
- Gruber, N., Lachkar, Z., Frenzel, H., Marchesiello, P., Münnich, M., McWilliams, J. C., Nagai, T., and Plattner, G.-K.: Eddy-induced reduction of biological production in eastern boundary upwelling systems, *Nature geoscience*, 4, 787–792, 2011.
- 550 Guillen, O. and Calienes, R.: Upwelling off chimbote, *Coastal upwelling*, 1, 312–326, 1981.
- Gutknecht, E., Dadou, I., Le Vu, B., Cambon, G., Sudre, J., Garçon, V., Machu, E., Rixen, T., Kock, A., Flohr, A., et al.: Coupled physical/biogeochemical modeling including O₂-dependent processes in the Eastern Boundary Upwelling Systems: application in the Benguela, *Biogeosciences*, 10, 3559–3591, 2013.
- 555 Henson, S., Le Moigne, F., and Giering, S.: Drivers of carbon export efficiency in the global ocean, *Global biogeochemical cycles*, 33, 891–903, 2019.
- Holte, J., Talley, L. D., Gilson, J., and Roemmich, D.: An Argo mixed layer climatology and database, *Geophysical Research Letters*, 44, 5618–5626, 2017.
- José, Y. S., Dietze, H., and Oschlies, A.: Linking diverse nutrient patterns to different water masses within anticyclonic eddies in the upwelling system off Peru, *Biogeosciences*, 14, 1349–1364, 2017.
- 560 Kelly, T. B., Goericke, R., Kahru, M., Song, H., and Stukel, M. R.: CCE II: Spatial and interannual variability in export efficiency and the biological pump in an eastern boundary current upwelling system with substantial lateral advection, *Deep Sea Research Part I: Oceanographic Research Papers*, 140, 14–25, 2018.
- Kleppel, G.: On the diets of calanoid copepods, *Marine Ecology-Progress Series*, 99, 183–183, 1993.
- 565 Koné, n. V., Machu, E., Penven, P., Andersen, V., Garçon, V., Fréon, P., and Demarcq, H.: Modeling the primary and secondary productions of the southern Benguela upwelling system: A comparative study through two biogeochemical models, *Global Biogeochemical Cycles*, 19, 2005.
- Lachkar, Z. and Gruber, N.: What controls biological production in coastal upwelling systems? Insights from a comparative modeling study, *Biogeosciences*, 8, 2961–2976, 2011.

- 570 Lathuilière, C., Echevin, V., Lévy, M., and Madec, G.: On the role of the mesoscale circulation on an idealized coastal upwelling ecosystem, *Journal of Geophysical Research: Oceans*, 115, 2010.
- Lima, I. D. and Doney, S. C.: A three-dimensional, multnutrient, and size-structured ecosystem model for the North Atlantic, *Global Biogeochemical Cycles*, 18, 2004.
- Liu, W. T., Tang, W., and Polito, P. S.: NASA scatterometer provides global ocean-surface wind fields with more structures than numerical
575 weather prediction, *Geophysical Research Letters*, 25, 761–764, 1998.
- Marañón, E., Lorenzo, M. P., Cermeño, P., and Mouriño-Carballido, B.: Nutrient limitation suppresses the temperature dependence of phytoplankton metabolic rates, *The ISME journal*, 12, 1836–1845, 2018.
- Messié, M. and Chavez, F. P.: Seasonal regulation of primary production in eastern boundary upwelling systems, *Progress in Oceanography*, 134, 1–18, 2015.
- 580 Messié, M., Ledesma, J., Kolber, D. D., Michisaki, R. P., Foley, D. G., and Chavez, F. P.: Potential new production estimates in four eastern boundary upwelling ecosystems, *Progress in Oceanography*, 83, 151–158, 2009.
- Montes, I., Dewitte, B., Gutknecht, E., Paulmier, A., Dadou, I., Oschlies, A., and Garçon, V.: High-resolution modeling of the Eastern Tropical Pacific oxygen minimum zone: Sensitivity to the tropical oceanic circulation, *Journal of Geophysical Research: Oceans*, 119, 5515–5532, 2014.
- 585 Morán, X. A. G., Calvo-Díaz, A., Arandia-Gorostidi, N., and Huete-Stauffer, T. M.: Temperature sensitivities of microbial plankton net growth rates are seasonally coherent and linked to nutrient availability, *Environmental microbiology*, 20, 3798–3810, 2018.
- Moriarty, R. and O’Brien, T.: Distribution of mesozooplankton biomass in the global ocean, *Earth System Science Data*, 5, 45–55, 2013.
- O’Brien, T. D.: COPEPOD, a global plankton database: A review of the 2007 database contents and new quality control methodology, 2007.
- O’Reilly, J. E., Maritorena, S., Mitchell, B. G., Siegel, D. A., Carder, K. L., Garver, S. A., Kahru, M., and McClain, C.: Ocean color
590 chlorophyll algorithms for SeaWiFS, *Journal of Geophysical Research: Oceans*, 103, 24 937–24 953, 1998.
- Pauly, D., Christensen, V., Dalsgaard, J., Froese, R., and Torres, F.: Fishing down marine food webs, *Science*, 279, 860–863, 1998.
- Pennington, J. T., Mahoney, K. L., Kuwahara, V. S., Kolber, D. D., Calienes, R., and Chavez, F. P.: Primary production in the eastern tropical Pacific: A review, *Progress in oceanography*, 69, 285–317, 2006.
- Prowe, A. F., Pahlow, M., Dutkiewicz, S., Follows, M., and Oschlies, A.: Top-down control of marine phytoplankton diversity in a global
595 ecosystem model, *Progress in Oceanography*, 101, 1–13, 2012.
- Ridgway, K., Dunn, J., and Wilkin, J.: Ocean interpolation by four-dimensional weighted least squares—Application to the waters around Australasia, *Journal of atmospheric and oceanic technology*, 19, 1357–1375, 2002.
- Schukat, A., Auel, H., Teuber, L., Lahajnar, N., and Hagen, W.: Complex trophic interactions of calanoid copepods in the Benguela upwelling system, *Journal of Sea Research*, 85, 186–196, 2014.
- 600 Shchepetkin, A. F. and McWilliams, J. C.: The regional oceanic modeling system (ROMS): a split-explicit, free-surface, topography-following-coordinate oceanic model, *Ocean modelling*, 9, 347–404, 2005.
- Steinberg, D. K. and Landry, M. R.: Zooplankton and the ocean carbon cycle, *Annual review of marine science*, 9, 413–444, 2017.
- Stramma, L., Schmidtko, S., Levin, L. A., and Johnson, G. C.: Ocean oxygen minima expansions and their biological impacts, *Deep Sea Research Part I: Oceanographic Research Papers*, 57, 587–595, 2010.
- 605 Stukel, M. R., Ohman, M. D., Benitez-Nelson, C. R., and Landry, M. R.: Contributions of mesozooplankton to vertical carbon export in a coastal upwelling system, *Marine Ecology Progress Series*, 491, 47–65, 2013.

- Sunda, W. G. and Huntsman, S. A.: Interrelated influence of iron, light and cell size on marine phytoplankton growth, *Nature*, 390, 389–392, 1997.
- Suttle, C. A.: Viruses in the sea, *Nature*, 437, 356–361, 2005.
- 610 Taylor, A., Watson, A., Ainsworth, M., Robertson, J., and Turner, D.: A modelling investigation of the role of phytoplankton in the balance of carbon at the surface of the North Atlantic, *Global Biogeochemical Cycles*, 5, 151–171, 1991.
- Tedesco, P., Gula, J., Ménesguen, C., Penven, P., and Krug, M.: Generation of submesoscale frontal eddies in the Agulhas Current, *Journal of Geophysical Research: Oceans*, 124, 7606–7625, 2019.
- Thomas, A., Carr, M.-E., and Strub, P. T.: Chlorophyll variability in eastern boundary currents, *Geophysical Research Letters*, 28, 3421–3424, 615 2001.
- Thomas, M. K., Aranguren-Gassis, M., Kremer, C. T., Gould, M. R., Anderson, K., Klausmeier, C. A., and Litchman, E.: Temperature–nutrient interactions exacerbate sensitivity to warming in phytoplankton, *Global Change Biology*, 23, 3269–3280, 2017.
- Thomsen, S., Kanzow, T., Krahnemann, G., Greatbatch, R. J., Dengler, M., and Lavik, G.: The formation of a subsurface anticyclonic eddy in the Peru-Chile Undercurrent and its impact on the near-coastal salinity, oxygen, and nutrient distributions, *Journal of Geophysical Research: Oceans*, 121, 476–501, 2016. 620
- Turner, J. T.: Zooplankton fecal pellets, marine snow, phytodetritus and the ocean’s biological pump, *Progress in Oceanography*, 130, 205–248, 2015.
- Worley, S. J., Woodruff, S. D., Reynolds, R. W., Lubker, S. J., and Lott, N.: ICOADS release 2.1 data and products, *International Journal of Climatology: A Journal of the Royal Meteorological Society*, 25, 823–842, 2005.