

# Net soil carbon balance in afforested peatlands and separating autotrophic and heterotrophic soil CO<sub>2</sub> effluxes

Renée Hermans<sup>15</sup>, Rebecca McKenzie<sup>26</sup>, Roxane Andersen<sup>2</sup>, Yit Arn Teh<sup>3</sup>, Neil Cowie<sup>4</sup> and Jens-Arne Subke<sup>1</sup>

5 <sup>1</sup>Department of Biological and Environmental sciences, University of Stirling, Stirling, UK

<sup>2</sup>Environmental Research Institute, University of Highlands & Islands, Thurso, UK

<sup>3</sup>School of Natural and Environmental Sciences, Newcastle University, Newcastle, UK

<sup>4</sup>Centre for Conservation Science, Royal Society for the Protection of Birds Scotland, Edinburgh, UK

10 <sup>5</sup>IUCN UK Peatland Programme, Edinburgh, UK

<sup>6</sup>Geography Department, Loughborough University, Loughborough, UK

*Correspondence to:* renee.kerkvliet-hermans@iucn.org.uk

**Abstract.** Peatlands are a significant global carbon (C) store, which can be compromised by drainage and afforestation. Quantifying the rate of C loss from peat soils under forestry is challenging, as soil CO<sub>2</sub> efflux includes both CO<sub>2</sub> produced from heterotrophic peat decomposition and CO<sub>2</sub> produced by tree roots and associated fungal networks (autotrophic respiration). We experimentally terminated autotrophic belowground respiration in replicated forest plots by cutting through all living tree roots (“trenching”), and measured soil surface CO<sub>2</sub> flux, litter input, litter decay rate and soil temperature and moisture over two years. Decomposition of cut roots was measured and CO<sub>2</sub> fluxes were corrected for this, this resulted in a large change in the fraction heterotrophic : autotrophic flux, suggesting that even two years after trenching decaying root biomass make significant contributions to the CO<sub>2</sub> flux. Annual peat decomposition (heterotrophic CO<sub>2</sub> flux) was  $115 \pm 16$  g C m<sup>-2</sup> y<sup>-1</sup>, representing c. 40% of total soil respiration. Decomposition of needle litter is accelerated in the presence of an active rhizosphere, indicating a priming effect by labile C inputs from roots. This suggests that our estimates of peat mineralization in our trenched plots are conservative, and underestimate overall rates of peat C loss. Considering also input of litter from trees, our results indicate that the soils in these 30 year-old drained and afforested peatlands are a net sink for C, since substantially more C enters the soil as organic matter, than is decomposed heterotrophically. This study doesn’t account for fluvial C fluxes, which represents a small flux compared to the CO<sub>2</sub> soil efflux; further, root litter and exudate deposition could be a significant C source that is only partially sampled by our approach, adding to these plantations being a potential carbon sink. However, the C balance for these soils should be taken over the lifespan of the trees, in order to determine if the soils under these drained and afforested peatlands are a sustained sink of C, or become a net source over longer periods of forestry.

## 30 **1 Introduction**

Large peatland areas in the boreal and temperate zone have been drained and afforested, in Western Europe, especially in the UK, Ireland and the Fennoscandia region, with conifers replacing native peatland vegetation (Andersen et al., 2016). In the UK alone, >800,000 ha (approximately 20%) were affected by this land-use change (Anderson et al., 2016). As well as causing

habitat loss, drainage and afforestation of peatlands influences peatland hydrology, biogeochemistry, and carbon (C) storage.

35 In Fennoscandia, drainage to improve tree growth on naturally treed peatlands has been shown to cause significant changes in decomposition and greenhouse gas (GHG) fluxes (Ojanen et al., 2013). However, the effects of drainage and afforestation on temperate peatlands that were previously dominated by smaller-statured vegetation (e.g. blanket bogs, moorland, heathland) are more uncertain (Sloan et al., 2018). While changes in hydrology and soil redox potential are anticipated to accelerate soil C loss and alter the composition of GHG emissions, there is very little quantitative data on how this shift in land-use changes

40 soil C dynamics and GHG emissions in temperate peatlands (Hermans et al., 2018). Since northern hemisphere peatlands are estimated to store a third of global terrestrial carbon (Gorham, 2010), changes linked to drainage and afforestation of temperate peatlands could have significant impacts on regional C dynamics and GHG balances.

There is a broad consensus that peatland drainage accelerates the loss of endogenous peat C stocks, but the impacts of drainage and afforestation on total soil C stocks and soil respiration are less certain (Hargreaves et al., 2003; Mayer et al.,

45 2020; Simola et al., 2012; Vanguelova et al., 2019; Zerva & M. Mencuccini, 2005). Afforestation of previously open peatlands in the British Isles (e.g. Mazzola et al., 2020) and Fennoscandia (e.g. Tolvanen et al., 2020) differ from other types of drained peatlands because this land-use change involves a wholesale shift in the functional composition of the plant community (i.e. replacement of short-statured vegetation with trees), leading to potential interactions or synergistic effects arising from changes to both soil hydrology and plant community composition.

50 The comparatively high rates of net primary production and larger stature of the trees on drained and afforested peatlands can represent a significant net ecosystem C sink, and consequently represent a large source of detrital material to soil (Yamulki et al., 2013). Thus, it is possible that this input of C from more highly productive trees could partially or wholly offset or outstrip some of the losses of C derived from mineralization of the peat. Moreover, changes to soil organic matter (SOM) chemistry due to a shift towards inputs of more recalcitrant and nutrient-poor coniferous litter could further enhance soil C storage in

55 afforested peatlands in the British Isles and Fennoscandia. This is because the trees planted in these regions (e.g. Sitka spruce) produce tissues that are less readily decomposed than the organic matter (OM) generated by the short-statured vegetation that they have replaced (i.e. larger proportion of woody debris generated with higher C:N ratios and greater proportion of recalcitrant compounds like lignin) (Hermans et al., 2018). This may affect soil C stocks by inhibiting or slowing overall rates of decay due to the tree litter's recalcitrance (Liski et al., 2002), but has not been studied in tree plantations on deep peat so

60 far.

In order to close these knowledge gaps, we need to determine the C balance of drained and afforested peatland soils, tracking C inputs from tissue turnover (e.g. litterfall and rhizodeposition), as well as C losses from SOM decay. However, separating the direct and indirect effects of trees on peat mineralization is methodologically challenging (Subke et al., 2006). Soil CO<sub>2</sub> efflux includes both CO<sub>2</sub> released from peat mineralization (heterotrophic CO<sub>2</sub> flux) and CO<sub>2</sub> produced from root-

65 rhizosphere (i.e. autotrophic) respiration; experimental manipulation of autotrophic C supply to the rhizosphere allows a separation of these two main component fluxes, but introduces a number of potential artefacts (see Subke et al., 2006 for a methods overview including method-related assumptions and artefacts). Root trenching, where roots are severed throughout

the depth of the rooting zone to create areas of forest floor with no recent C input from trees to roots or associated mycorrhizas, has been used across many forest sites, and provides useful insights into relative contributions of autotrophic and heterotrophic soil CO<sub>2</sub> efflux and the respective temporal dynamics. This disruption of root exudation and continuous root turnover could influence peat decay through processes such as microbial priming effects (Kuzyakov, 2006; Subke et al., 2006), where microbial activity (linked with an active rhizosphere) in the soil is stimulated by the addition of easily accessible C from roots (Kuzyakov, 2006). Further experiments are required to evaluate the effects of these other plant-facilitated processes on peat decay, to better constrain estimates of heterotrophic activity in soil.

In this study, we experimentally determined the C budget of a drained and afforested peat soil. Carbon inputs were tracked by quantifying litterfall, while C outputs were determined using a trenching technique to partition root-rhizosphere and heterotrophic respiration. In addition, we conducted a root decomposition study in rhizosphere and living root-free (i.e. trenched) soil, to constrain artefacts associated with the trenching and in order to determine if rhizosphere-linked processes (e.g. priming effects) facilitated OM decay. We hypothesised that 1) the soils under these forestry plantations are a net C source, 2) soil CO<sub>2</sub> efflux is dominated by autotrophic respiration and 3) interactions between C supply to the rhizosphere by trees and surface litter decomposition result in higher decomposition rates of the litter due to rhizosphere priming effects.

## 2 Methods

### 2.1 Study site

The research took place in RSPB's Forsinard Flows National Nature Reserve in the north of Scotland (58° 22' N, 3° 53' W). Four paired plots were established in the beginning of June 2014 in three separate forestry plantation blocks of identical age containing a mixture, 3:5 (on average), of Sitka Spruce (*Picea sitchensis*) and Lodgepole Pine (*Pinus contorta*). The plantations were drained, around 30 years old and very dense (about 5000 trees per ha), with no vascular plant understory, but sporadic patches of moss, predominantly feather moss, e.g. *Hypnum jutlandicum*, *Hylocomium splendens* and in some instances, *Sphagnum fallax* and *S. capillifolium* in furrows. The average diameter at breast height (DBH) for Sitka Spruce was 13.3 cm and for Lodgepole Pine 17.9 cm. The average canopy cover was 76.3% (Smith and Hancock, 2016). Peat depths in these three forestry blocks varied between 30 and 260 cm, with depths at research plots between 137 and 204 cm (Smith et al., 2014). The average annual precipitation in the research area between 1981-2010 was 970.5 mm with an average maximum air temperature of 11.4°C and average minimum air temperature of 3.3°C measured at the Kinbrace weather station approximately 20 km from the plots (Location: 58°13'89"N, 3°55'1.2"W; Altitude: 103 m above mean sea level; Met Office, 2018). Seasonal averaged water table relative to ground surface is -350 (±20) mm in spring (March-May), -457 (±34) mm in summer (June-August), -404 (±49) mm in autumn (September-November) and -244 (±14) mm in winter (December-February).

## 2.2 Experimental set up

Candidate locations for the four paired trenched and control areas in each plot were initially identified at random and soil surface respiration measured. Based on respiration results, two closely matched plots were selected, and randomly allocated a treatment (trenching or control). Paired plots were no more than 10 metres apart from each other.

The double-mouldboard ploughing of the peat at the time of afforestation created a regular micro topography with low-lying furrows (c. 1.5 m wide) flanked by high ridges (plough throws; c. 0.75 m wide) on either side. In between two plough throws, there is an area of c. 0.50 m width of the original (unploughed) surface (Figure 1). The height from the bottom of the furrow to the top of the plough throw is on average 0.5 m and from the original surface to the plough throw is about 0.15 - 0.2 m. In general, conifer seedlings were planted on the plough throws because of the improved drainage compared to the original surface. For our study, each plot encompassed the three types of micro-topography.




Figure 1 Micro topography in forest plantations, with location of measurements for flux chambers (not to scale).

110

## 2.3 Trenching

A 40 cm deep and 30 cm wide trench was cut to just below the main rooting depth of the trees, cutting through all roots present. The trench was double-lined using polypropylene gardening cloth, and re-filled with peat soil in between the two layers of cloth to prevent in-growth of roots (Figure 2). The dimensions of each trench plot were about 3.5 x 1.5 meters and included

115 all three micro topographic forms. These dimensions maximised the space between trees, but did not represent ground within 60 cm of trees.



Figure 2 Left: trenching plot dug and all living tree roots are sawn trough. Right: finished trench plot; trenches double lined with polypropylene gardening cloth and filled in with peat.

## 120 **2.4 CO<sub>2</sub> measurements**

Three pairs of PVC collars of 10 cm height with a diameter of 20 cm were installed to a depth of three cm within the three microforms (Figure 1) of both trenched and control plots. CO<sub>2</sub> measurements were taken using custom-built dark dynamic closed chamber with a height of five cm and a diameter of 20 cm, which were placed on the permanent collars for three minutes. Elasticated rubber material placed over the joint of chambers and collars for the duration of the measurement provided an air-tight seal. The chamber was connected to an EGM 4 Infrared Gas Analyser (PP-Systems, Amesbury, MA, USA), recording CO<sub>2</sub> concentrations every 4-5 seconds. Fluxes were calculated from increases in CO<sub>2</sub> concentration within the chamber over 3 minutes. Measurements were carried out ten times between August 2014 and July 2016.

125

## **2.5 Litter**

Six litter traps (0.07 m<sup>2</sup> each) were located close to each plot, and litter (falling needles and twigs) collected at each sampling visit. Dry weight of all litter was recorded as area-based averages for each plot separately.

130

Litter was allowed to fall onto the soil surface within collars for the duration of the experiment. To be able to distinguish peat respiration from litter respiration, surface litter was removed manually from one (always the same) of the two paired collars in each microform before measuring respiration. The litter present in the “collar with litter” was weighed after a respiration measurement and then placed back in the collar. Litter adjacent to the collar was also collected and weighed in the field, then  
135 taken back to the lab, dried and weighed again to establish the wet to dry mass ratio of litter and calculate litter dry mass within each collar.

## 2.6 Roots

Root biomass was determined from soil cores (0-20 cm deep and 6.5 cm diameter) taken from each microform in all plots, at the start (June 2014) and end (July 2016) of the experiment. Both dead and living roots from each core were carefully separated  
140 and sorted into three root diameter classes: smaller than 2 mm, 2 to 5 mm, and greater than 5 mm. All roots and the root-free soil were dried at 50°C for 7 days, and weighed to establish percentage roots per gram of soil. Root depth was found to be 25 cm when digging trenches and root density for 0-20 cm was assumed to be representative for 0-25 cm.

To estimate root decomposition directly, roots were taken from soil collected in each plot, dried (50°C for 7 days) and separated into the same size classes as described previously. Between 0.36 and 0.69 g of dried root material (separately  
145 for each size class) were placed in polyester mesh bags (10 x 10 cm; mesh size of 0.5 mm) for field incubations. Bags were soaked in water for 2 days prior to field placement, to mimic field conditions. Four replicate bags of each size class were buried at 5-10 cm depth in all three microforms in all plots four weeks after trenching. To account for any weight loss that may have occurred prior to field incubation, five bags of each size class were taken into the field and not buried, but taken back to the lab; the proportional mass loss of litter in these bags was used to correct the initial root mass of all other bags.  
150 One bag per root class per microform was collected from all sites in November 2014, March 2015, July 2015 (except root class >5 mm, since there was not enough material for four bags) and July 2016. After retrieval, bags were dried for seven days at 50°C, and root dry mass recorded.

Root decay was fitted to an exponential decay function:

$$M_t = M_0 e^{-kt} \quad (1)$$

155 With  $M_t$  the remaining amount of root biomass after collection from the field,  $M_0$  the initial root biomass,  $t$  time and  $k$  the decay constant. Data fits were performed separately for root size and microform.

## 2.7 Soil moisture and temperature

Between June 2014 and July 2016, soil moisture and soil temperature at 5 and 20 cm soil depth were recorded at 30-minute intervals in all three microforms in a nearby plot, using 12-bit smart temperature sensors, S-TMB-M002 (Onset Computer  
160 Corporation, Bourne, MA, USA) and 10HS soil moisture smart sensors, S-SMD-M005 (Decagon Devices, Inc., Pullman, WA,

USA combined with Onset’s smart sensor technology, measuring  $m^3/m^3$  (volumetric water content)) connected to a Hobo micro station (Onset Computer Corporation, Bourne, MA, USA).

In addition to this, soil temperature (10 cm thermistor) and moisture (HH2 moisture meter, Delta-T Devices, Cambridge, measuring  $m^3/m^3$  (volumetric water content)) were measured at 5 cm depth next to each collar during sampling.

165 Air temperature was also measured one meter above the ground during sampling.

## 2.8 Statistical analyses and flux calculations

Data were analysed using R (RStudio Team, 2016). All  $CO_2$  data was log transformed to meet the criteria of normality, and a linear mixed-effect model (LMM) using the *lme4* package in R (Bates et al., 2015) was used to predict  $CO_2$  fluxes based on environmental drivers. All numerical predictors were standardized to one standard deviation prior to statistical analyses, to

170 allow relative effect sizes of predictors to be compared directly (Nakagawa & Schielzeth, 2010). Model selection was done

based on information theory (Burnham & Anderson, 2002). First a model of maximum complexity was built, with interactions between soil moisture, soil temperature, trench treatment and microform plus interactions between trench treatment, microform and litter treatment, with plot as a random effect ( $lmer(CO_2 \text{ flux} \sim (\text{soil moisture} * \text{soil temperature} * \text{treatment} * \text{microform} + \text{treatment} * \text{microform} * \text{litter treatment}) + (1|plot))$ ). Then, all possible combinations of this model were identified using the

175 ‘dredge’ function in the *MuMIn* package (Barton, 2017). Goodness of model fit was assessed with the small-sample size corrected Akaike’s Information Criterion (AICc), which is calculated using the number of parameters and either the maximum likelihood estimate for the model or the residual sum of squares. “Likelihood” here is a measure of the extent to which a sample

provides support for particular values of a parameter in a parametric model. AICc values of different models can be compared and the model with the lowest AICc is selected as the ‘best approximating model’, hereafter called ‘top model’ (Burnham &

180 Anderson, 2002). Any of the models with a delta AICc of less than 2 are considered to be as good as the best model (Richards, 2005). ‘Dredge’ also gives a weight to the models it produces, ranging between 0 and 1; with for example a weight of 0.7

meaning that there is a 70% chance that that model is the best approximating model of the models considered. If the weight of the best model is low, it is not possible to say that that model really is the best model, meaning other models also have to be considered. In this study, the model with the best AICc and highest weight was used. Significance (p-values) for the mixed

185 effect model were calculated using the package *lmerTest* (Kuznetsova et al., 2016).

Annual fluxes were calculated using the predict function over the mixed effects model from library *lme4* in R (Bates et al., 2015) and predictions and error terms were back transformed. The predictions were made over half-hourly measurements of soil moisture and soil temperature at 5 cm soil depth in all three microforms just outside the plots.

From these predictions, partitioned fluxes were calculated from the collars without litter as:

$$190 F_a + \varepsilon = (F_{control} + \varepsilon) - ((F_{trench} + \varepsilon) - (F_{dead\ roots} + \varepsilon)) \quad (2)$$


Where  $F_a$  is autotrophic  $\text{CO}_2$  flux,  $F_{\text{control}}$  is the  $\text{CO}_2$  flux from the control plots,  $F_{\text{trench}}$  is the  $\text{CO}_2$  flux from trenched plots,  $F_{\text{dead roots}}$  is the  $\text{CO}_2$  flux coming from the dead roots in the trenched plots created by the trenching technique and  $\epsilon$  the associated error terms, error terms where propagated.

195 The annual flux from litter was calculated from the difference in modelled annual fluxes between collars with and without litter.

### 3 Results

#### 3.1 Temporal trends in soil $\text{CO}_2$ fluxes

200 Trenching initially led to an increase in soil respiration, followed by a significant reduction in soil  $\text{CO}_2$  flux. Soil respiration fluxes from both control and trenched plots showed a clear annual cycle, with highest fluxes in summer. After the initial perturbation fluxes from trenched plots are significantly lower than fluxes from control plots ( $p < 0.001$ ) and this difference is greater in the summer (Figure 3).



205 Figure 3 Mean  $\text{CO}_2$  fluxes from control (grey squares) and trenched (black circles) plots over time, averaging across all microforms ( $n = 12$ ) and both litter treatments. Error bars are standard errors and are in most instances smaller than symbols.

Soil  $\text{CO}_2$  fluxes were best explained with a combination of soil moisture, soil temperature, trenching treatment, microform and litter treatment, with an interaction between soil moisture and soil temperature, including 'plot' as a random



210 effect. Table 1 shows model estimates for each variable, with their standard error and p-value. The marginal  $R^2$  was 0.40 and conditional  $R^2$  was 0.41 on the log scale, since predictions are made in the original units all uncertainties reflate. The set of models with a  $\Delta AICc$  of less than 2 is shown in Table 2. The predictors of the ‘top model’ were present in over half of the models in the top model set, so this model was used.

215 Table 1 Model estimates with standard errors and p-value for best-fit model. All numerical predictors were standardized to one standard deviation prior to analysis.

<b>Fixed effect</b>	<b>Estimate</b>	<b>Std. Error</b>	<b>p-value</b>
Intercept	-0.22	0.10	0.05
Trenched	-0.50	0.07	<0.001
Microform - Original surface	0.42	0.12	<0.001
Microform - Plough throw	0.35	0.13	0.006
Soil moisture	-0.12	0.06	0.03
Soil temperature	0.35	0.03	<0.001
No Litter	-0.17	0.06	0.008
Soil moisture x Soil temperature	-0.11	0.04	0.008

Table 2 Model selection summary, showing the 4 best ranked models with a delta AICc of less than 2. Models are ranked by AICc and weight, where higher weighted models have more statistical support. Df= degrees of freedom, Loglik=Log likelihood, and  $\Delta$ AICc is the difference in AICc to the 'top model'.

Candidate models	Df	LogLik	AICc	$\Delta$ AICc	Weight
Trenched + Microform+ No Litter + Soil moisture + Soil temperature + Soil moisture x Soil temperature	10	-403	826	0.00	0.34
Trenched + Microform+ No Litter + Soil moisture + Soil temperature	9	-404	826	0.37	0.29
Trenched + Microform+ Soil moisture + Soil temperature + Soil moisture x Soil temperature	9	-404	827	1.21	0.19
Trenched + Microform+ Soil moisture + Soil temperature	8	-405	827	1.30	0.18

### 3.2 Spatial trends and litter contributions to soil CO<sub>2</sub> flux

In both control and trenched plots, fluxes from plough throw (control:  $1.23 \pm 0.10$ ; trenched:  $0.85 \pm 0.07 \mu\text{mol m}^{-2} \text{s}^{-1}$ ;  $p=0.01$ ) and original surface (control:  $1.48 \pm 0.10$ ; trenched:  $0.83 \pm 0.06 \mu\text{mol m}^{-2} \text{s}^{-1}$ ;  $p<0.001$ ) were significantly higher than fluxes from furrow (control:  $0.90 \pm 0.06$ ; trenched:  $0.47 \pm 0.04 \mu\text{mol m}^{-2} \text{s}^{-1}$ ). Across all microforms fluxes from collars with litter ( $1.03 \pm 0.05 \mu\text{mol m}^{-2} \text{s}^{-1}$ ) were significantly higher than fluxes from collars without litter ( $0.91 \pm 0.05 \mu\text{mol m}^{-2} \text{s}^{-1}$ ,  $p=0.008$ ).

### 3.3 Role of environmental drivers in modulating CO<sub>2</sub> flux

Observed soil CO<sub>2</sub> efflux values correlated positively with soil temperature, whilst soil moisture showed an inconsistent correlation with flux values; a significant ( $p=0.008$ ) interaction between soil temperature and soil moisture means that at high temperatures CO<sub>2</sub> flux decreases with increasing soil moisture, but at low temperatures flux increases when soil moisture increases, with an inflection between positive and negative moisture dependence between 6 and 7 °C (Figure 4).

There was no difference in soil temperature between trenched and control plots, but soil moisture was significantly higher in trenched plots than in control plots ( $p<0.001$ ). To account for the artificially elevated soil moisture in trenched plots, CO<sub>2</sub> fluxes were corrected using the global relationship between soil moisture and CO<sub>2</sub> flux (Figure 4) to calculate fluxes expected to occur under moisture conditions in control plots.




Figure 4 Combined effect of soil temperature and soil moisture at 5 cm depth on soil CO<sub>2</sub> flux from the control sites, estimated from the 'top model'.

### 240 3.4 Partitioned fluxes

Soil CO<sub>2</sub> efflux was partitioned into component fluxes for all measuring dates from August 2014 onwards to remove disturbance related artefacts observed immediately after trenching. The model prediction of heterotrophic respiration is calculated by subtracting the autotrophic flux from the total soil flux and includes emissions from decomposition of cut roots. Flux simulations based on the soil model details indicate significantly lower autotrophic fluxes than heterotrophic fluxes (p=0.01, Figure 5). Across all microforms, heterotrophic fluxes represented 61% and autotrophic fluxes represented 39% of the total fluxes. From these predictions, annual sums for autotrophic and heterotrophic fluxes have been calculated, giving an average peat decomposition flux of  $184 \pm 21 \text{ g C m}^{-2} \text{ y}^{-1}$ . Total soil respiration without needle litter is  $301.3 \pm 34.2 \text{ g C m}^{-2} \text{ y}^{-1}$ , with needle decomposition adding  $41.2 \pm 53.5 \text{ g C m}^{-2} \text{ y}^{-1}$  to annual fluxes, giving a total soil respiration including needle litter of  $343 \pm 35 \text{ g C m}^{-2} \text{ y}^{-1}$ .

250




Figure 5 Modelled and measured fluxes of heterotrophic (black) and autotrophic (grey) soil CO<sub>2</sub> efflux from the three topographic microforms. Closed symbols are average fluxes with error bars (n = 4 plots). Connecting lines are the predicted fluxes using soil temperature and moisture at 5 cm depth. a) Original surface, b) Plough throw and c) Furrow.

### 3.5 Impact of litter and roots


Litter fall was  $719 \pm 71.3$  grams of litter per  $\text{m}^2 \text{y}^{-1}$ , with no detectable difference between trenched and control plots. Assuming a C fraction of biomass of 50% (Mathews, 1993), this represents an input to the soil of  $359 \text{ g C m}^{-2} \text{y}^{-1}$  via litter fall.

260  $\text{CO}_2$  flux from surface litter is calculated from the difference in the modelled annual fluxes between the collars with and without litter, with the fluxes from collars with litter significantly higher than the  $\text{CO}_2$  flux from the collars without litter ( $p = 0.008$ , Table 1). C emitted by litter in the control plots appears to be higher than in trenched plots and the  $\text{CO}_2$  flux per gram of litter is 1.7 to 3.6 times higher from the control plots than from the trenched plots (Table 3).

265 Table 3 Mean amount of C emitted as  $\text{CO}_2$  by surface needle litter in 'litter collars', for both years (standard error in brackets).

Microform	Litter [g $\text{m}^{-2}$ ]		$\text{CO}_2$ flux [g C $\text{m}^{-2} \text{y}^{-1}$ ]		$\text{CO}_2$ flux per g litter [mg C (g litter) $^{-1}$ ]	
	Trench	Control	Trench	Control	Trench	Control
<b>Original surface</b>	472.4 (174.0)	237.8 (61.0)	34.4 (32.4)	62.5 (50.4)	72.8 (73.6)	263 (222)
<b>Plough throw</b>	351.1 (125.2)	263.5 (85.4)	36.0 (40.8)	60.1 (61.3)	103 (122)	228 (244)
<b>Furrow</b>	589.8 (288.8)	558.4 (256.2)	26.4 (28.8)	43.2 (40.8)	44.8 (53.5)	77.4 (81.2)

270 Root biomass per  $\text{m}^2$  with an assumed rooting depth of 25 cm in August 2014 was  $761 \pm 324 \text{ g}$ ,  $603 \pm 110 \text{ g}$  and  $715 \pm 257 \text{ g}$  for  $< 2 \text{ mm}$  roots,  $2\text{-}5 \text{ mm}$  roots and  $> 5 \text{ mm}$  roots, respectively. For both the control and the trench plots, roots smaller than 2 mm declined in total biomass from the start of the experiment to the end of the experiment, but there were no significant difference between the control and trenched plots at the start and end of the experiment. Despite an apparent trend towards higher root biomass in control plots in July 2016, these differences were not statistically significant. There was also no significant differences between the beginning and end of the experiment for root classes  $2\text{-}5 \text{ mm}$  and  $>5 \text{ mm}$  (Figure 6).



275 Figure 6 Root biomass per square meter to 25 cm in control (grey) and trenched (black) plots at the beginning of the experiment (June 2014) and end of experiment (July 2016), split into three root size classes, per microform. a) root size <2 mm, b) root size 2-5 mm, c) root size >5 mm. Symbols are average fluxes with error bars representing standard errors (n = 4 plots).

### 3.6 Root decomposition

280 Root mass in decomposition bags showed a consistent decline over the duration of the experiment (Figure 7). The calculated decay constant ( $k$ ) showed systematic differences by microform, with highest decay rates tending to occur for incubations at “original surface” (Table 4).




Figure 7 Root mass remaining in root bags over number of days buried, for each root size class averaged across all microforms.

a) root size <2 mm, b) root size 2-5 mm, c) root size >5 mm.

285

Table 4 Root decay constants and associated C emissions in trenched plots by microform and root size classes (standard error in brackets).

	<b>Root diameter</b>	<b>Decay constant</b>	<b>C emitted year 1</b>	<b>C emitted year 2</b>
	<b>mm</b>	<b>y<sup>-1</sup></b>	<b>g C m<sup>-2</sup> y<sup>-1</sup></b>	<b>g C m<sup>-2</sup> y<sup>-1</sup></b>
Original surface	<2	0.106 (0.019)	57 (18)	52 (17)
	2-5	0.180 (0.027)	28 (7)	23 (6)
	>5	0.190 (0.027)	50 (25)	41 (21)
	<b>Total</b>		<b>135 (32)</b>	<b>116 (27)</b>
Plough throw	<2	0.097 (0.015)	39 (12)	35 (11)
	2-5	0.153 (0.026)	33 (6)	28 (5)
	>5	0.100 (0.022)	10 (1)	9 (1)
	<b>Total</b>		<b>82 (13)</b>	<b>73 (12)</b>
Furrow	< 2	0.068 (0.015)	10 (3)	9 (3)
	2-5	0.097 (0.018)	11 (4)	10 (3)
	>5	0.065 (0.023)	3 (/)	3 (/)
	<b>Total</b>		<b>24 (5)</b>	<b>22 (5)</b>

### 3.7 C flux from dead roots

The amount of C emitted from the decaying roots is calculated using the exponential decay function (1), with the biomass of roots per m<sup>2</sup> to a depth of 20 cm in the trenched plots at the beginning of the experiment as M<sub>0</sub>. It is assumed that all biomass lost is emitted as CO<sub>2</sub> and that 50% of roots is C (Mathews, 1993), as conservative assumptions, meaning that estimates are maximum possible CO<sub>2</sub> flux from decaying roots in trenched plots. To correct for soil CO<sub>2</sub> efflux generated in trenched plots as an artefact of creating dead root biomass, annual estimated CO<sub>2</sub> emissions were corrected by subtracting root-decay based estimates from trenched plot CO<sub>2</sub> emissions (Table 5).

295



Table 5 Corrected for dead root decay in trenched plots, heterotrophic soil CO<sub>2</sub> efflux (excluding litter) (F<sub>h</sub>) and autotrophic (F<sub>a</sub>) fluxes (standard error in brackets) in g C m<sup>-2</sup> y<sup>-1</sup> for both first (August 2014 – August 2015) and second year (August 2015 – August 2016) of the study.

	Year 1		Year 2		Average		
	F <sub>h</sub>	F <sub>a</sub>	F <sub>h</sub>	F <sub>a</sub>	F <sub>h</sub>	F <sub>a</sub>	F <sub>soil</sub>
Original surface	86.5 (37.7)	276 (35)	80.2 (33.2)	242 (30)	83.3 (35.4)	259 (33)	342 (48)
Plough throw	118 (29)	212 (20)	124.1 (28.0)	199 (19)	121 (28)	206 (19)	327 (34)
Furrow	134 (21)	125 (10)	107 (17)	105 (9)	120 (19)	115 (9)	235 (21)

300

Heterotrophic fluxes represents approximately 24% and autotrophic fluxes 76% of the total soil fluxes in the original surface, 37% and 63% respectively in the plough throw, 51% and 49% respectively, in the furrow and 38% and 62% respectively averaged over all microforms.

### 3.8 Weighted average for soil CO<sub>2</sub> flux in Flow Country forest plantations

305 In order to scale soil CO<sub>2</sub> fluxes (excluding litter) from different microforms to the level of an entire forest stand, fluxes from individual microforms were weighted by their fractional area (Table 6). This results in a slight shift in proportion of heterotrophic and autotrophic CO<sub>2</sub> flux sources to 40% and 60% respectively and a total area weighted soil CO<sub>2</sub> flux of 289.4 ± 18.6 g C m<sup>-2</sup> y<sup>-1</sup>.

310 Table 6 Area-weighted heterotrophic peat decomposition (F<sub>h</sub>) and autotrophic (F<sub>a</sub>) fluxes (standard error in brackets) in g C m<sup>-2</sup> y<sup>-1</sup> averaged over both years measured.

Microform	Fractional area	Area-weighted F <sub>h</sub>	Area-weighted F <sub>a</sub>
Original surface	0.14	12 (5)	36 (5)
Plough throw	0.43	52 (13)	88 (8)
Furrow	0.43	52 (8)	49 (4)
Total	1	115 (16)	174 (10)

## 4 Discussion

Our results indicate that the soils in these 30 year old drained and afforested peatlands are a net sink for C, as substantially more C enters the soil as organic matter than is decomposed heterotrophically. The soil surface C balance of the soil under these forest plantations is visualised in Figure 8, with the annual CO<sub>2</sub> fluxes of the forest plantation based on the area-weighted fluxes. Based on these fluxes, we tentatively estimate the soil C sink (the difference between litter C input and total heterotrophic soil CO<sub>2</sub> efflux) to be  $191 \pm 64$  g C m<sup>-2</sup> y<sup>-1</sup> for our site.





Figure 8 Summary of soil C budget (in g C m<sup>-2</sup> y<sup>-1</sup>) under the canopy of a Sitka spruce and Lodgepole pine plantation on deep peat. Downward arrows indicate litterfall, whilst upward fluxes show autotrophic (F<sub>a</sub>) (black), peat decomposition flux (grey) and surface needle litter (white) across the three main microforms. Total forest soil fluxes with area weighted values are visualised in the top right corner, including the estimated soil C sink (grey and white diagonal striped).

We found a C input of 359 g C m<sup>-2</sup> y<sup>-1</sup> via litter fall, similar to other Sitka Spruce forests of similar age to these forest plantations in the UK, which range from 273 to 573 g C m<sup>-2</sup> y<sup>-1</sup> (Morison et al. 2012). This is balanced by total soil efflux including litter-derived CO<sub>2</sub> of  $342.5 \pm 34.7$  g C m<sup>-2</sup> y<sup>-1</sup>, i.e. the amount of C entering the soil as surface litter alone falls within a similar range to C leaving as CO<sub>2</sub>. Missing in the C budget in Figure 8 are the input of C from root exudates that remain in the soil C pool (net rhizodeposition), potential losses through aquatic C export, and root turnover. Gaffney, et al. (2020) have measured the aquatic C flux (DOC, DIC, POC) from similar forest plantations in the north of Scotland and have found a flux of 13.9 g C m<sup>-2</sup> y<sup>-1</sup>.

$^2 \text{ y}^{-1}$ , which is a small flux compared to the  $\text{CO}_2$  soil efflux. Root:shoot allocation in forest ecosystems is usually in the order of 1:3 (Laiho & Laine, 1997), so it is possible that belowground litter and exudate deposition could account for a significant C input only partially sampled by our approach (i.e. excluding large root stocks), which means that our tentative soil C sink estimate is conservative.

335 The rate of peat decomposition in these drained and afforested peatlands is substantial, but overall soil  $\text{CO}_2$  efflux and ratio of heterotrophic/autotrophic respiration falls within boreal averages for forested ecosystems (Figure 9; Bond-Lamberty and Thomson, 2010).



340 Figure 9 Heterotrophic annual flux against soil respiration annual flux ( $\text{g C m}^{-2} \text{ y}^{-1}$ ) in Boreal forests in black dots, Boreal forests on peat soils in grey triangles, from Bond-Lamberty and Thomson (2010) dynamic database downloaded on 19/04/2020. Our study is included in the red square. See Table 7 in Appendix for references to studies used to create this graph.

Results from other drained temperate peatland sites under conifer plantation vary from around  $260 \text{ g C m}^{-2} \text{ y}^{-1}$  at a  
345 39-year old drained Sitka spruce plantation in Ireland on naturally treeless blanket bog (Byrne & Farrell, 2005) to values of  $972 \text{ g C m}^{-2} \text{ y}^{-1}$  on average across 18 to 44 year old Sitka spruce plantations on poorly drained Dystric Histosols in the south of Ireland (Jovani-Sancho et al., 2018), with similar annual fluxes of  $453 \text{ g C m}^{-2} \text{ y}^{-1}$  reported from under a 43-year old lodgepole pine plantation in central Scotland (Yamulki et al., 2013). Afforested boreal peatlands also show generally higher soil  $\text{CO}_2$  flux values, ranging from  $577$  to  $705 \text{ g C m}^{-2} \text{ y}^{-1}$  (e.g. Mäkiranta et al., 2008; Minkkinen et al., 2018; Ojanen et al.,  
350 2010). Also taking into account possible measurement bias associated with the chemical absorption method used by Byrne & Farrell (2005), which generally underestimates fluxes compared to more reliable infrared gas analysis based methods (Janssens & Ceulemans, 1998), our results of  $342.5 \text{ g C m}^{-2} \text{ y}^{-1}$  sit at the low end of total soil respiration fluxes. These low rates are corroborated by independent chamber flux measurements across forest sites in the Flowcountry (Hermans et al., unpublished), and are likely to relate to poor site fertility and specific biochemical properties of the peat at these sites (Hermans et al., 2019).

355 Our peat oxidation rates of  $115 \pm 16 \text{ g C m}^{-2} \text{ y}^{-1}$  are also low compared to temperate afforested peatlands, which have been reported to vary between 211 and  $407 \text{ g C m}^{-2} \text{ y}^{-1}$  for Sitka spruce sites in Ireland (Jovani-Sancho et al., 2018), whilst boreal peatlands show an even larger range of heterotrophic flux sums (145 to  $670 \text{ g C m}^{-2} \text{ y}^{-1}$ ; Ojanen et al., 2010). Our low estimates result in part from the root decomposition correction, which reduced heterotrophic flux estimates by around  $60 \text{ g C m}^{-2} \text{ y}^{-1}$ , and is not routinely applied across other studies.

360 Minkkinen et al (2018) measured carbon fluxes of a drained, naturally forested peatland (Kalevansuo peatland forest) in the south of Finland over 4 years. They found that the ecosystem was a strong sink of  $\text{CO}_2$  in all studied years, with an average NEE for the 4 years of  $-234 \text{ g C m}^{-2} \text{ y}^{-1}$ . By subtracting the carbon sink of the tree stand from eddy-covariance derived NEE the authors show that the soil was a carbon sink of  $-60 \text{ g C m}^{-2} \text{ y}^{-1}$ . However, modelling their forest soil respiration fluxes from chamber measurements, they found the peat respiration alone made up 53% of the total forest floor respiration flux, litter  
365 22%, roots 16% and autotrophic respiration of above-ground vegetation 8%, with soils representing a weak C source. Our results show a lower percentage of peat respiration of 38% of the total soil respiration minus the litter flux. The higher percentage found by Minkkinen et al (2018) could possibly be explained by them not applying a dead root correction to their fluxes, which as we show could lead to a big difference in fluxes. For forestry-drained boreal peatlands, Ojanen et al (2013) report a range of soil C balances from weak source to sinks depending on site fertility. Accounting for belowground litter  
370 production and fine root turnover is an important source of uncertainty in soil C balance estimates (Ojanen et al., 2014), and whilst our soil C balance focuses on surface litter only, our calculations indicate a weak C sink even before belowground litter input is considered.

Trenching is likely to underestimate heterotrophic respiration and rates of peat mineralization since this approach does not account for rhizosphere effects such as priming (Walker et al., 2016). Therefore, it is likely that the estimated rate of  
375 peat oxidation from trenching is conservative. The observed difference of 1.7 to 3.6 times higher  $\text{CO}_2$  flux per gram of litter from the control plots than from the trenched plots (Table 3) indicates that heterotrophic processes are reduced under trenching. In presence of an active rhizosphere (control plots), decomposition of needle litter and/or soil organic matter (SOM) appears to be faster than when the rhizosphere is not active (trenched plots). This priming of organic matter decomposition is likely to be the result of microbial activity in the soil stimulated by the addition of easily accessible C from roots (Kuzyakov, 2006).  
380 Friggens et al. (2020) recently demonstrated the accelerated loss of organic soil under coniferous and deciduous forest plantations on peaty podzols in Scotland, which is consistent with soil priming through the elevated input of C from trees in the rhizosphere. Several studies have shown that mycorrhizal fungi are involved in soil priming (Kuyper, 2017; Paterson et al., 2016), further underlining the interaction between plant-associated biotic activity and decomposition in underlying peat. Therefore, in the control plots a slightly larger proportion of the total  $\text{CO}_2$  flux could be heterotrophic decomposition than the  
385 fluxes from the trenched plots suggest, which means there could be an underestimation of heterotrophic flux in our results, in line with results from literature (Subke et al., 2004; Subke et al., 2011).

To calculate the root biomass at the start of the experiment, one soil core per microform was taken and since trees were standing close to each other (1.5 to 2 m apart) this was assumed to be representative for the whole microform. It was not

possible to distinguish between living and dead roots in the soil cores, so our results might overestimate the living root biomass.  
390 The dead root emission correction made a big difference to the ratio of heterotrophic to autotrophic flux, going from 61% and  
39% respectively over all microforms to 38% and 62% respectively (without area-weighting of fluxes), so a decrease in  
heterotrophic flux of 23%. This is in line with the corrections used in other studies; a comparison of corrections used in  
trenching studies indicates a range from 2% to 24%, with an average of 12% (Subke et al., 2006). This big difference in the  
fraction heterotrophic : autotrophic flux suggests that even two years after trenching, decaying root biomass make significant  
395 contributions to the CO<sub>2</sub> flux.

That soils under a 30-year old conifer plantation on deep peat are a C sink is an interesting finding. However, in order  
to determine if these soils are a long-term sink or source of carbon, the balance between soil C input from roots and litter and  
C loss via oxidation should be quantified over the lifespan of the plantation. The average peat depth in these forest plots is 126  
(±16) cm, with 0.47 kg C m<sup>-2</sup> per centimetre depth (Cannell et al., 1993), which represents ca. 59.3 (±7.3) kg C m<sup>-2</sup> stored in  
400 the peat under these plantations. The largest carbon losses most likely occurred in the initial planting phase of afforestation  
(Simola et al., 2012; Vanguelova et al., 2019; Zerva & M. Mencuccini, 2005), but fluxes have not been measured through this  
phase and cannot be quantified as new policy prevents planting on deep peat (Forestry Commission, 2016). In other parts of  
the world, peatland drainage is still actively happening and studies from these sites show very high rates of peat oxidation  
during the first 5-10 years of conversion (e.g. Cooper et al., 2020; McCalmont et al., 2021; Minkkinen et al., 2018; Prananto  
405 et al., 2020)(McCalmont et al., 2021). In the UK forest plantations on deep peat sometimes end in clear felling of the site and  
restoration of the peat. The results of this study could also help inform what the CO<sub>2</sub> fluxes will be when timber is harvested  
and large quantities of felling residues are left above ground as well as whole root systems below ground. However, we note  
that changes in water table and soil moisture conditions created by the soil rewetting associated with clear-felling will have  
significant and separate impacts beyond the conditions of active drainage under which we took our measurements.

410

## 5 Code and data availability

The main dataset used for the analysis in this paper are publicly available on: ... The R code used for the analysis is not  
publicly available. Requests to access the code should be directed to renee.kerkvliet-hermans@iucn.org.uk.

## 6 Author contribution

415 Funding acquisition and initial conceptualization of the whole PhD project was done by J-A Subke, R. Andersen, Y. Teh and  
N. Cowie. Further in-depth conceptualization of this particular work was done by R. Hermans and J-A Subke, with support  
from R. Andersen, Y Teh and N. Cowie. Investigation, methodology, project administration, data curation, formal analysis,  
software, visualization and writing was done by R. Hermans with supervision and guidance of J-A Subke, R. Andersen, Y.  
Teh and N. Cowie.

## 420 7 Competing interests

The authors declare that they have no conflict of interest.

## 8 Acknowledgements

We are grateful for the PhD studentship that enabled this work, jointly funded by the University of Stirling and the Royal Society for the Protection of Birds. Trenching of plots was supported by volunteers working at the RSPB Forsinard reserve and the Environmental Research Institute of the University of the Highlands and Islands. We thank Rebecca McKenzie, Paul Gaffney, Peter Gilbert and Nathalie Triches for help with field measurements. Further we gratefully acknowledge Daniela Klein, Norrie Russell, Trevor Smith (RSPB), Graeme Findlay (Forestry Commission Scotland), Fountains Forestry and Donald MacLennan (Brook Forestry) for guidance and land access.

## 9 References

- 430 Andersen, R., Farrell, C., Graf, M., Muller, F., Calvar, E., Frankard, P., Caporn, S., & Anderson, P. (2016). An overview of the progress and challenges of peatland restoration in Western Europe. *Restoration Ecology*, 1–12. <https://doi.org/10.1111/rec.12415>
- Anderson, R., Vasander, H., Geddes, N., Laine, A., Tolvanen, A., O’Sullivan, A., & Aapala, K. (2016). Afforested and forestry-drained peatland restoration. In A. Bonn, T. Allott, M. Evans, H. Joosten, & R. Stoneman (Eds.), *Peatland restoration and ecosystem services - science, policy and practise* (pp. 213–233). Cambridge University press.
- 435 Barton, K. (2017). MuMIn: Multi-Model Inference. *R Package Version 1.40.0*. <https://cran.r-project.org/package=MuMIn>
- Bates, D., Maechler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software*, 67(1), 1–48. <https://doi.org/10.18637/jss.v067.i01>
- Burnham, K. P., & Anderson, D. R. (2002). *Model Selection and Multimodel Inference: A Practical Information-Theoretic Approach (2nd ed)*. Springer, New York, USA.
- 440 Byrne, K. A., & Farrell, E. P. (2005). The effect of afforestation on soil carbon dioxide emissions in blanket peatland in Ireland. *Forestry*, 78(3), 217–227. <https://doi.org/10.1093/forestry/cpi020>
- Cannell, M. G. R., Dewar, R. C., & Pyatt, D. G. (1993). Conifer plantations on drained peatlands in Britain: A net gain or loss of carbon? *Forestry*, 66(4), 353–369. <https://doi.org/10.1093/forestry/66.4.353>
- 445 Cooper, H. V., Evers, S., Aplin, P., Crout, N., Dahalan, M. P. Bin, & Sjoergersten, S. (2020). Greenhouse gas emissions resulting from conversion of peat swamp forest to oil palm plantation. *Nature Communications*, 11(1). <https://doi.org/10.1038/s41467-020-14298-w>
- Forestry Commission. (2016). Supplementary guidance to support the FC forests and peatland habitats guideline note (2000). *Scottish Forestry Publications*, 1–5.

- 450 Friggens, N. L., Hester, A. J., Mitchell, R. J., Wookey, J. S. P. A., & Parker, T. C. (2020). Tree planting in organic soils does not result in net carbon sequestration on decadal timescales. *Global Change Biology, April*, 1–11. <https://doi.org/10.1111/gcb.15229>
- Gorham, E. (2010). Northern Peatlands: Role in the Carbon Cycle and Probable Responses to Climatic Warming. *Ecological Applications, 1*(2), 182–195.
- 455 Hargreaves, K. J., Milne, R., & Cannell, M. G. R. (2003). Carbon balance of afforested peatland in Scotland. *Forestry, 76*(3), 299–317.
- Hermans, R., Zahn, N., Andersen, R., Teh, Y. A., Cowie, N., & Subke, J. A. (2018). An incubation study of GHG flux responses to a changing water table linked to biochemical parameters across a peatland restoration chronosequence. *Mires and Peat, 23*, 1–18. <https://doi.org/10.19189/MaP.2018.DW.354>
- 460 Janssens, I. A., & Ceulemans, R. (1998). Spatial variability in forest soil CO<sub>2</sub> efflux assessed with a calibrated soda lime technique. *Ecology Letters, 1*(2), 95–98.
- Jovani-Sancho, A. J., Cummins, T., & Byrne, K. A. (2018). Soil respiration partitioning in afforested temperate peatlands. *Biogeochemistry, 141*(1), 1–21. <https://doi.org/10.1007/s10533-018-0496-0>
- Kuyper, T. W. (2017). Carbon and Energy Sources of Mycorrhizal Fungi: Obligate Symbionts or Latent Saprotrophs? In *Mycorrhizal Mediation of Soil: Fertility, Structure, and Carbon Storage* (pp. 357–374). Elsevier Inc. <https://doi.org/10.1016/B978-0-12-804312-7.00020-6>
- 465 Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2016). lmerTest: Tests in Linear Mixed Effects Models. *R Package Version 2.0-33*. <https://cran.r-project.org/package=lmerTest>
- Kuzyakov, Y. (2006). Sources of CO<sub>2</sub> efflux from soil and review of partitioning methods. *Soil Biology and Biochemistry, 38*(3), 425–448. <https://doi.org/10.1016/j.soilbio.2005.08.020>
- 470 Laiho, R., & Laine, J. (1997). Tree stand biomass and carbon content in an age sequence of drained pine mires in southern Finland. *Forest Ecology and Management, 93*(1–2), 161–169. [https://doi.org/10.1016/S0378-1127\(96\)03916-3](https://doi.org/10.1016/S0378-1127(96)03916-3)
- Liski, J., Perruchoud, D., & Karjalainen, T. (2002). Increasing carbon stocks in the forest soils of western Europe. *Forest Ecology and Management, 169*(1–2), 159–175. [https://doi.org/10.1016/S0378-1127\(02\)00306-7](https://doi.org/10.1016/S0378-1127(02)00306-7)
- 475 Mäkiranta, P., Minkinen, K., Hytönen, J., & Laine, J. (2008). Factors causing temporal and spatial variation in heterotrophic and rhizospheric components of soil respiration in afforested organic soil croplands in Finland. *Soil Biology and Biochemistry, 40*(7), 1592–1600. <https://doi.org/10.1016/j.soilbio.2008.01.009>
- Mathews, G. (1993). *The carbon content of trees. Forestry Commission Technical Paper 4*. 21.
- Mayer, M., Prescott, C. E., Abaker, W. E. A., Augusto, L., Cécillon, L., Ferreira, G. W. D., James, J., Jandl, R., Katzensteiner, K., Laclau, J. P., Laganière, J., Nouvellon, Y., Paré, D., Stanturf, J. A., Vanguelova, E. I., & Vesterdal, L. (2020). Influence of forest management activities on soil organic carbon stocks: A knowledge synthesis. *Forest Ecology and Management, 466*, 118127. <https://doi.org/10.1016/j.foreco.2020.118127>
- 480 Mazzola, V., Perks, M. P., Smith, J., Yeluripati, J., & Xenakis, G. (2020). Seasonal patterns of greenhouse gas emissions from

- a forest-to-bog restored site in northern Scotland: Influence of microtopography and vegetation on carbon dioxide and methane dynamics. *European Journal of Soil Science*, August, 1–22. <https://doi.org/10.1111/ejss.13050>
- 485 McCalmont, J., Kho, L. K., Teh, Y. A., Lewis, K., Chocholek, M., Rumpang, E., & Hill, T. (2021). Short- and long-term carbon emissions from oil palm plantations converted from logged tropical peat swamp forest. *Global Change Biology*, 27(11), 2361–2376. <https://doi.org/10.1111/gcb.15544>
- Met Office. (2018). *Kinbrace climate*. <https://www.metoffice.gov.uk/public/weather/climate/gfm5qbgxz>
- 490 Minkkinen, K., Ojanen, P., Penttilä, T., Aurela, M., Laurila, T., Tuovinen, J. P., & Lohila, A. (2018). Persistent carbon sink at a boreal drained bog forest. *Biogeosciences*, 15(11), 3603–3624. <https://doi.org/10.5194/bg-15-3603-2018>
- Morison, J., Matthews, R., Miller, G., Perks, M., Randle, T., Vanguelova, E., White, M., & Yamulki, S. (2012). Understanding the carbon and greenhouse gas balance of forests in Britain. *Research Report - Forestry Commission, UK*.
- Nakagawa, S., & Schielzeth, H. (2010). Repeatability for Gaussian and non-Gaussian data: A practical guide for biologists. *Biological Reviews*, 85(4), 935–956. <https://doi.org/10.1111/j.1469-185X.2010.00141.x>
- 495 Ojanen, P., Lehtonen, A., Heikkinen, J., Penttilä, T., & Minkkinen, K. (2014). Soil CO<sub>2</sub> balance and its uncertainty in forestry-drained peatlands in Finland. *Forest Ecology and Management*, 325, 60–73. <https://doi.org/10.1016/j.foreco.2014.03.049>
- Ojanen, P., Minkkinen, K., Alm, J., & Penttilä, T. (2010). Soil–atmosphere CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O fluxes in boreal forestry-drained peatlands. *Forest Ecology and Management*, 260(3), 411–421. <https://doi.org/10.1016/j.foreco.2010.04.036>
- 500 Ojanen, P., Minkkinen, K., & Penttilä, T. (2013). The current greenhouse gas impact of forestry-drained boreal peatlands. *Forest Ecology and Management*, 289, 201–208. <https://doi.org/10.1016/j.foreco.2012.10.008>
- Paterson, E., Sim, A., Davidson, J., & Daniell, T. J. (2016). Arbuscular mycorrhizal hyphae promote priming of native soil organic matter mineralisation. *Plant and Soil*, 408(1–2), 243–254. <https://doi.org/10.1007/s11104-016-2928-8>
- 505 Prananto, J. A., Minasny, B., Comeau, L.-P., Rudyanto, R., & Grace, P. (2020). Drainage increases CO<sub>2</sub> and N<sub>2</sub>O emissions from tropical peat soils. *Global Change Biology*, 26(8), 4583–4600. <https://doi.org/https://doi.org/10.1111/gcb.15147>
- Richards, S. A. (2005). Testing ecological theory using the information-theoretic approach: examples and cautionary results. *Ecology*, 86(10), 2805–2814.
- RSPB unpublished data. (n.d.). *No Title*.
- 510 RStudio Team. (2016). *RStudio: Integrated Development Environment for R* (1.0.136). RStudio, Inc. <http://www.rstudio.com/>
- Simola, H., Pitkänen, A., & Turunen, J. (2012). Carbon loss in drained forestry peatlands in Finland, estimated by re-sampling peatlands surveyed in the 1980s. *European Journal of Soil Science*, 63(6), 798–807. <https://doi.org/10.1111/j.1365-2389.2012.01499.x>
- Sloan, T. J., Payne, R. J., Anderson, A. R., Bain, C., Chapman, S., Cowie, N., Gilbert, P., Lindsay, R., Mauquoy, D., Newton, A. J., & Andersen, R. (2018). Peatland afforestation in the UK and consequences for carbon storage. *Mires and Peat*, 23, 1–17. <https://doi.org/10.19189/MaP.2017.OMB.315>
- Smith, T., & Hancock, M. (2016). *Interim report on monitoring of forest to bog restoration, Dyke and Forsinain forests:*



*baseline forest characteristics and vegetation.*

- 520 Smith, T., Hancock, M., & Stagg, P. (2014). *The RSPB Forest to Bog Restoration Project Forsinard Flows Interim Report – July 2012 – Sept 2013. April.*
- Subke, Inglima, I., & Francesca Cotrufo, M. (2006). Trends and methodological impacts in soil CO<sub>2</sub> efflux partitioning: A metaanalytical review. *Global Change Biology*, 12(6), 921–943. <https://doi.org/10.1111/j.1365-2486.2006.01117.x>
- 525 Subke, J.-A., Hahn, V., Battipaglia, G., Linder, S., Buchmann, N., & Cotrufo, M. F. (2004). Feedback interactions between needle litter decomposition and rhizosphere activity. *Oecologia*, 139(4), 551–559. <https://doi.org/10.1007/s00442-004-1540-4>
- Subke, Voke, N. R., Leronni, V., Garnett, M. H., & Ineson, P. (2011). Dynamics and pathways of autotrophic and heterotrophic soil CO<sub>2</sub> efflux revealed by forest girdling. *Journal of Ecology*, 99(1), 186–193. <https://doi.org/10.1111/j.1365-2745.2010.01740.x>
- 530 Tolvanen, A., Tarvainen, O., & Laine, A. M. (2020). Soil and water nutrients in stem-only and whole-tree harvest treatments in restored boreal peatlands. *Restor Ecol*, 28(6), 1357–1364.
- Vanguelova, E. I., Crow, P., Benham, S., Pitman, R., Forster, J., Eaton, E. L., & Morison, J. I. L. (2019). Impact of Sitka spruce (*Picea sitchensis* (Bong.) Carr.) afforestation on the carbon stocks of peaty gley soils- A chronosequence study in the north of England. *Forestry*, 92(3), 242–252. <https://doi.org/10.1093/forestry/cpz013>
- 535 Walker, T. N., Garnett, M. H., Ward, S. E., Oakley, S., Bardgett, R. D., & Ostle, N. J. (2016). Vascular plants promote ancient peatland carbon loss with climate warming. *Global Change Biology*, 22(5), 1880–1889. <https://doi.org/10.1111/gcb.13213>
- Yamulki, S., Anderson, R., Peace, a., & Morison, J. I. L. (2013). Soil CO<sub>2</sub> CH<sub>4</sub> and N<sub>2</sub>O fluxes from an afforested lowland raised peatbog in Scotland: implications for drainage and restoration. *Biogeosciences*, 10(2), 1051–1065. <https://doi.org/10.5194/bg-10-1051-2013>
- 540 Zerva, A., & M. Mencuccini. (2005). Carbon stock changes in a peaty gley soil profile after afforestation with Sitka spruce (*Picea sitchensis*). *Annals of Forest Science*, 62, 873–880. <https://doi.org/10.1051/forest>

## 10 Appendix

545 **Table 7 List of articles used in** Error! Reference source not found.

<b>Authors</b>	<b>Year</b>	<b>Title</b>
Bond-Lamberty, B, C Wang, and ST Gower	2004	Contribution of Root Respiration to Soil Surface CO <sub>2</sub> Flux in a Boreal Black Spruce Chronosequence
Bond-Lamberty, Ben, Dustin Bronson, Emma Bladyka, and Stith T. Gower	2011	A Comparison of Trenched Plot Techniques for Partitioning Soil Respiration
Gaumont-Guay, D., T. A. Black, A. G. Barr, T. J. Griffis, R. S. Jassal, P. Krishnan, N. Grant, and Z. Nestic.	2014	Eight Years of Forest-Floor CO <sub>2</sub> Exchange in a Boreal Black Spruce Forest: Spatial Integration and Long-Term Temporal Trends
Gaumont-Guay, David, T. Andrew Black, Tim J. Griffis, Alan G. Barr, Rachhpal S. Jassal, and Zoran Nestic	2006	Interpreting the Dependence of Soil Respiration on Soil Temperature and Water Content in a Boreal Aspen Stand
Gaumont-Guay, David, T. Andrew Black, Tim J. Griffis, Alan G. Barr, Kai Morgenstern, Rachhpal S. Jassal, and Zoran Nestic	2006	Influence of Temperature and Drought on Seasonal and Interannual Variations of Soil, Bole and Ecosystem Respiration in a Boreal Aspen Stand
Gaumont-Guay, David, T Andrew Black, Alan G Barr, Rachhpal S Jassal, and Zoran Nestic	2008	Biophysical Controls on Rhizospheric and Heterotrophic Components of Soil Respiration in a Boreal Black Spruce Stand
Hermle, Sandra, Michael B. Lavigne, Pierre Y. Bernier, Onil Bergeron, and David Paré	2010	Component Respiration, Ecosystem Respiration and Net Primary Production of a Mature Black Spruce Forest in Northern Quebec
Högberg, Peter, Anders Nordgren, Nina Buchmann, Andrew F. S. Taylor, Alf Ekblad, Mona N. Högberg, Gert Nyberg, Mikael Ottosson-Löfvenius, and David J. Read	2001	Large-Scale Forest Girdling Shows That Current Photosynthesis Drives Soil Respiration

- Howard, Erica A., Stith T. Gower, 2004 Effects of Logging on Carbon Dynamics of a Jack Pine Forest  
Jonathan A. Foley, and in Saskatchewan, Canada  
Christopher J. Kucharik
- Laganière, Jérôme, David Paré, 2012 The Effect of Boreal Forest Composition on Soil Respiration Is  
Yves Bergeron, and Han Y.H. Mediated through Variations in Soil Temperature and C  
Chen Quality
- Ma, Y., Piao, S., Sun, Z., Lin, X., 2014 Stand ages regulate the response of soil respiration to  
Wang, T., Yue, C., and Yang, Y. temperature in a *Larix principis-rupprechtii* plantation
- Mäkiranta, Päivi, Kari Minkkinen, 2008 Factors Causing Temporal and Spatial Variation in  
Jyrki Hytönen, and Jukka Laine Heterotrophic and Rhizospheric Components of Soil  
Respiration in Afforested Organic Soil Croplands in Finland
- Mahli, Y, D D Baldocchi, and P G 1999 The Carbon Balance of Tropical, Temperate and Boreal Forests  
Jarvis
- Molchanov, A. G. 2009 CO<sub>2</sub> Emission from the Surface of Dark Gray Forest Soils of  
the Forest Steppe and Sandy Soddy-Podzolic Soils of the  
Southern Taiga
- Mustamo, P., Maljanen, M., 2016 Respiration and emissions of methane and nitrous oxide from a  
Hyvärinen, M., Ronkanen, A. K., boreal peatland complex comprising different land-use types.  
and Kløve, B.
- O'Connell, K. E. B., Gower, S. T., 2003 Net Ecosystem Production of Two Contrasting Boreal Black  
& Norman, J. M. Spruce Forest Communities
- Ojanen, Paavo, Kari Minkkinen, 2010 Soil-Atmosphere CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O Fluxes in Boreal Forestry-  
Jukka Alm, and Timo Penttil Drained Peatlands
- Ojanen, Paavo, Kari Minkkinen, 2012 Chamber Measured Soil Respiration: A Useful Tool for  
Annalea Lohila, Tiina Badorek, Estimating the Carbon Balance of Peatland Forest Soils?  
and Timo Penttilä
- Pumpanen, J., Kulmala, L., 2015 Seasonal dynamics of autotrophic respiration in boreal forest  
Lindén, A., Kolari, P., Nikinmaa, soil estimated by continuous chamber measurements  
E., and Hari, P.
- Russell, C A, and R P Voroney 1998 Carbon Dioxide Efflux from the Floor of a Boreal Aspen  
Forest. I. Relationship to Environmental Variables and  
Estimates of C Respired

- Ryan, M. G., M. B. Lavigne, and S. T. Gower 1997 Annual Carbon Cost of Autotrophic Respiration in Boreal Forest Ecosystems in Relation to Species and Climate
- Sawamoto, Takuji, Ryusuke Hatano, Masato Shibuya, Kunihide Takahashi, Alexander P. Isaev, Roman V. Desyatkin, and Trofim C. Maximov 2003 Changes in Net Ecosystem Production Associated with Forest Fire in Taiga Ecosystems, near Yakutsk, Russia
- Vogel, JG;, DW; Valentine, and RW Ruess 2005 Soil and Root Respiration in Mature Alaskan Black Spruce Forests That Vary in Soil Organic Matter Decomposition Rates
- Widén, Britta, and Hooshang Majdi 2001 Soil CO<sub>2</sub> Efflux and Root Respiration at Three Sites in a Mixed Pine and Spruce Forest: Seasonal and Diurnal Variation
- Zha, Tianshan, Zisheng Xing, Kai Yun Wang, Seppo Kellomaki, and Alan G. Barr 2007 Total and Component Carbon Fluxes of a Scots Pine Ecosystem from Chamber Measurements and Eddy Covariance