

# Photosynthetic activity in Devonian Foraminifera

Zofia Dubicka<sup>1,2</sup>, Maria Gajewska<sup>1</sup>, Wojciech Kozłowski<sup>1</sup>, Pamela Hallock<sup>3</sup>, Johann Hohenegger<sup>4</sup>

<sup>1</sup>Faculty of Geology, University of Warsaw, Warsaw, Poland

<sup>2</sup>Ecological Chemistry, Alfred-Wegener-Institut Helmholtz-Zentrum für Polar- und Meeresforschung, Bremerhaven,

5 Germany

<sup>3</sup>College of Marine Science, University of South Florida, St. Petersburg, USA

<sup>4</sup>Department of Palaeontology, Universität Wien, Wien, Austria

Correspondence to: Maria Gajewska (mi.gajewska@uw.edu.pl)

**Abstract.** Photosynthetically-active foraminifera are ~~prolific important~~ carbonate producers ~~in warm, sunlit, surface waters of the oceans, contributing nearly 5% of the reef and nearly 1% of the total global calcium carbonate budgets. The abilities to be photosynthetically active, Foraminifera have repeatedly developed mixotrophic strategies (i.e., the ability of an organism or holobiont to both feed and photosynthesize) realize~~ by ~~facultative or obligate~~ endosymbiosis with microalgae, or by sequestering plastids (kleptoplasts) of ~~digested-ingested~~ algae. ~~These ecological behaviors are a great advantage. Mixotrophy provides access to essential nutrients (e.g., N, P) through feeding, while providing carbohydrates and lipids produced through photosynthesis, resulting in substantial energetic advantage in warm, sunlit environments where food and dissolved nutrients are scarce for the continued growth, survival of the hosts and enhance of foraminiferal calcification. Our morphological and isotopic data provide concurrent pieces of evidence for, as of now, the earliest (mid-Devonian) evidence for photosymbiosis photosynthetic activity in protists observed in the first true~~ advanced, multichambered, calcareous foraminifera, ~~which are Semitextularia, from the tropical shelf of the Laurussia paleocontinent. from the tropical shelf of the Laurussia paleocontinent.~~ This adaptation ~~likely might have had a significant impact on~~ the evolutionary radiation of calcareous Foraminifera in the Devonian (“Givetian revolution”), ~~which was~~ one of the most important evolutionary events in foraminiferal history, ~~that. The observed phenomenon~~ coincided with the worldwide development of diverse calcifying marine communities inhabiting shelf environments linked with Devonian stromatoporoid-coral reefs.

## 25 1 Introduction

Among photosynthetically-active protists, symbiont-bearing benthic Foraminifera (SBBF), ~~which compose less than 10% of all living families (Lee and Anderson, 1991),~~ are important carbonate producers (e.g., Prazeres and Renema, 2019, ~~and references therein~~) contributing nearly 1% of ~~the total~~ global calcium carbonate budgets (Hohenegger, 2006; Langer ~~et al., 1997~~79). The occurrence of modern SBBF is limited to tropical and warm temperate zones and is related to the light requirements of the hosted symbiotic microalgae (e.g., Hallock, 1979; Larsen and Drooger, 1977; Renema, 2018). Living in warm, shallow, euphotic environments enables hosts to use their tests as “glass houses” suitable for the husbandry of endosymbiotic algae (e.g., Hohenegger, 2009; Hottinger, 1982). Hence, SBBF inhabit mainly upper euphotic shallow-reef and lagoon environments, ~~however, though~~ some groups hosting diatoms can be found along reef slopes in ~~highly~~ transparent water down to 130 m depth of the lower euphotic zone (e.g., Hallock, 1999; Hohenegger, 2000). ~~This ability Photosymbioses are is a great ecological energetically and ecologically highly~~ advantageous in warm, clear, nutrient-poor environments

(e.g., Hallock, 1981a, 1987; Lee et al., 2010; Schmidt et al., 2004). ~~as t~~ Through photosynthesis, endosymbionts can provide their hosts with the energy for much of their metabolic needs (i.e., mixotrophy; Dubinsky and Berman-Frank, 2001; Hallock, 1981b; Selose et al., 2017). ~~while and can also~~ enhancing the process of foraminiferal calcification (e.g., Hallock, 1999; Kinoshita et al., 2017; McConnaughey and Whelan, 1997; ter Kuile, 1991). ~~The most important b~~ Benefits for the algal symbionts ~~are may include~~ protection afforded by the host tests (Hohenegger, 2000, 2018), ~~if the symbionts actually do benefit~~ (Wooldridge, 2010) and uptake of inorganic nutrients from host metabolites (Hallock, 1999).

~~Apart from endosymbiosis, another~~ A more direct ecological adaptation that enables foraminifersforaminifera to be photosynthetically active is by kleptoplasty (Cesbron et al., 2017; Goldstein et al., 2004; Pillet et al., 2011). ForaminifersForaminifera sequester plastids from microalgal prey (e.g., ~~diatoms~~), ~~and those plastids can remain~~ photosynthetically active for days to weeks ~~within the foraminifera~~ (Jaufrais et al., 2016, 2018). Products of photosynthesis, such as glucose, can be especially important during periods of food scarcity. In dysphotic and aphotic habitats like the deeper bathyal benthos, sequestered plastids that originate from planktic diatoms can assimilate inorganic carbon, nitrate and sulfur (Jaufrais et al., 2019). Feeding upon microalgae, and harvesting and maintaining active plastids, is an evolutionarily more direct way to function as a mixotrophic organism compared to the more complicated array of adaptations associated with obligate symbioses.

Considering the important roles played by SBBF and kleptoplastic foraminifersforaminifera in carbonate production of present coral reefs and shallow-marine ecosystems (e.g., Baccaert, 1986; Fujita and Fujimura, 2008; Prazeres and Renema, 2019), as well as the advantage of endoplasmic photosynthesis in foraminiferal evolution, we are seeking the first evidence of photosynthetic activity in foraminifersforaminifera in the past. Presumptive candidates are the first multi-chambered calcareous foraminifersforaminifera with leaf-like tests such as *Semitextularia* (Eifelian–Frasnian), which are morphological counterparts to some recent SBBF, notably peneroplids (Figs. 1, 2). *Semitextulariids* were globally distributed on Devonian inner shelves, in shallow-marine, well-illuminated habitats such as tropical reefs and lagoons (see Table S1). They, and other well-known Devonian groups such as *Nanicellidae* and *Eonodosariidae*, died out following the Kellwasser event of the Frasnian–Famennian crisis (Vachard et al., 2010) that triggered the rapid collapse of the prolific stromatoporoid-coral reef ecosystems. In principle, the origin of *Semitextularia* occurred at the beginning of the Middle Devonian foraminiferal radiation, which is the so-called “Givetian revolution” (Vachard et al., 2014) and is considered to be the most important Paleozoic evolutionary event for these protists. This event is associated with ‘the replacement of the primitive agglutinated wall of the Textulariata by the more advanced secreted wall of the Fusulinata’ (Vachard et al., 2014, p. 217; see also BouDagher-Fadel, 2018).

In this paper, we hypothesize that the Devonian semitextularids were not only among the first true multi-chambered calcareous foraminifersforaminifera, but may also have been the first algal symbiont-bearing or chloroplast-sequestering foraminifersforaminifera. To test this hypothesis, we compared paleo-populations of *Semitextularia* derived from facies of coeval horizons (upper Eifelian, Middle Devonian; Skaly Beds, HCM, Central Poland) characterized by contrasting paleo-depths with different levels of light intensity representing the tropical southern shelf of the Laurussia paleo-continent

70 (Szulczewski, 1995). Samples were compared in terms of a possible photosynthesis-related microhabitat effect recorded in the carbon-isotopic data from fossil shells, since photosynthesis by algal symbionts ~~of—in~~ modern larger benthic ~~foraminifers~~foraminifera has a significant impact on  $\delta^{13}\text{C}$  composition. Enrichment of  $^{13}\text{C}$  in the microenvironment occurs when  $^{12}\text{CO}_2$  is ~~sequestered—preferentially taken up~~ by algae during photosynthesis (Ravelo and Hillaire-Marcel, 2007). Moreover, our findings are supported by ecological similarities and a functional-morphological comparison between  
75 *Semitextularia* and recent SBBFpeneroplids.

## 2 Materials and Methods

The studied samples were collected in the lower part of the Skały Formation (upper Eifelian–lower Givetian) outcrop in the Holy Cross Mountains inlier (HCM, Central Poland). The Skały Formation represents an epicratonic facies of the southern (so-called Fennosarmatian) tropical shelf of the Laurussia paleocontinent (Szulczewski, 1995) close to the northern part of the  
80 East European Platform (referred to as the Łysogóry unit) located in the HCM. The Skały outcrop (50°53'044.69"N 21°9'33.75"E) is part of the well-known Grzegorzowice-Skały (G-S) succession exposed along the Dobruchna brook (Pajchłowa, 1957; Zeuschner, 1869), which is a key section for the Devonian of the Łysogóry facies region (Szulczewski, 1995). The Miłoszów outcrop (50°54'09.5"N 21°07'14.5"E; Samosonowicz, 1936), located 3 km west of Skały, complements the key succession by virtue of the closely-correlated Miłoszów limestone complex (see faunal-summary tables in Biernat,  
85 1966; Pajchłowa, 1957; Stasińska, 1958), which was recently exposed in Skały. For detailed geological, stratigraphic, and sedimentary environment background, see Supplement Discussion/Note).

We examined a total of 25 samples from the upper Eifelian (Middle Devonian) Skały beds of the Holy Cross Mountains in Central Poland. The specimens studied were *Semitextularia thomasi* Miller and Carmer, 1933 ~~Miller and Carmer, 2015~~, including the isotopic signatures of the tests and the enclosing rock matrix. The latter included: (1) a marly  
90 intercalation in Miłoszów coral-bearing biostromes (two intervals Miłoszów 11 and 12), representing an upper euphotic (well-illuminated) environment (10 samples); (2) Skały brachiopod shales (SBS), containing a mesophotic (lower euphotic) coral assemblage (Skały 11 and A+11A/Skały brachiopod shales, 14 samples); and (3) marls with abundant solitary corals (Skały C sample), representing a proximal foreereef environment characterized by intermediate photic conditions. All samples were derived from very soft marly  
95 clays with low permeability and very low paleoheating rate (Dubicka et al., in press), that ensured favorable conditions for excellent fossil preservation. Very well-preserved, three-dimensional specimens, with no sediment or mineral infillings, with original foliated test microstructure, and with no evidence of any dissolution or recrystallization, were analyzed (Narkiewicz and Malec, 2005). Lack of diagenetic alteration of the Skały samples can be additionally supported by uniquely preserved brachiopods with preserved brachidia, primary shell structures (Biernat, 1966), and in some cases also relicts of the original  
100 color pattern (Biernat, 1984). Also, the porous stereom of crinoids includes original (metastable) high-Mg calcite, which is the first phase to be affected during the diagenetic alteration of calcareous fossils (Gorzelał et al., 2011).

During maceration, samples were mechanically disintegrated (crumbled into ca. 1–2 cm<sup>3</sup> pieces), dried and repeatedly soaked in water and detergent, then heated and dried again following decantation. To intensify the rock disintegration process, some samples were also subjected to a maceration technique using Rewoquat (Jarochovska et al., 2013). After being washed  
105 in an ultrasonic cleaner, the residuum was sieved using a mesh size of 0.053 mm. From the residue, foraminifera were hand-picked and studied using a Nikon SMZ 18 stereoscopic microscope. More detailed taxonomic observations and photographic documentation were made using a Zeiss Sigma VP scanning electron microscope at the Faculty of Geology, University of Warsaw.

For oxygen and carbon isotopic analyses, 35 foraminiferal samples and 25 bulk-rock samples were prepared.  
110 Foraminifera were manually chosen from the residue prior to undergoing multiple additional cleanings in an ultrasonic cleaner. Sediment-free *Semitextularia thomasi* ~~Miller and Carner~~ specimens were placed in Eppendorf tubes (ca. 10 large adult specimens in each tube). Bulk rock samples were prepared by grinding a portion of rock material of ca. 2 g to powder using a mortar. The isotopic analysis was performed at the GeoZentrum at the University of Erlangen, Germany. Samples were reacted with 100% phosphoric acid at 70°C using a Gasbench II connected to a Thermo Fisher Delta V Plus mass spectrometer.  
115 Reproducibility of the analyses was 0.06 and 0.06‰ (±1 s.d.) for  $\delta^{18}\text{O}$  and  $\delta^{13}\text{C}$  values of laboratory standard Sol 2 (n = 16) and 0.07 and 0.06 ‰ (±1 s.d.) for  $\delta^{18}\text{O}$  and  $\delta^{13}\text{C}$  values of laboratory standard Erl 5 (n = 22), respectively. All isotopic data are reported in per mil relative to the VPDB scale and presented in Tables S1 and S2 for Miłoszów and Skały, respectively.

All statistical analyses were carried out using the program package PAST 4.02 (Hammer et al., 2001).

### 3 Results

#### 120 3.1 Morphometrics of the *Semitextularia* tests

*Semitextularia* was the only plurilocular foraminifera in the Skały and Miłoszów assemblages. *Semitextularia* is the oldest and pioneering multichambered form. Apart from *Semitextularia*, in the studied sections, there are some bilocular and tubular forms of the family Moravamminidae family were found, as well as and many different microproblematica, for which the systematic position and paleoecology are unknown . (e.g., Langer, 1979)

125 Differences in test morphology of *Semitextularia* between specimens from Miłoszów, representing an upper euphotic environment (see appendix Discussion), and Skały, interpreted as lower euphotic (i.e., mesophotic) environment (Zapalski et al., 2017), were tested for differences in size and shape. Size represented by *test height* (i.e., maximum diameter) did not differ significantly among samples [ANOVA:  $p(H_0) = 0.499$ ; Tables S2, S3], followed by pairwise comparisons. Only specimens from Skały 1 were, on average, slightly smaller compared to the other samples. Shape, represented by the  
130 ~~size independent~~ parameters  $\sqrt{\text{area}}/\text{height}$  and  $\text{width}/\text{height}$ , also did not differ among samples [ANOVA;  $p(H_0) = 0.697$ ,  $p(H_0) = 0.122$ ; Table S4], again with some smaller  $\text{width}/\text{height}$  ratios for Skały 1. The complete overlapping of samples is supported by regression functions of *test area* to *test height* (Fig. 3).

### 3.2 $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ values in Devonian foraminiferal shells along with rock-matrix samples

135 Carbon isotopic data from *Semitextularia* specimens, as well as from bulk-rock samples were compared (Tables S5, S6). For *Semitextularia*,  $\delta^{13}\text{C}$ -values were significantly higher in specimens from Miłoszów (2.86, 2.71) compared to Skały [1.47, 0.84, 1.60;  $p(H_0) = 1.05\text{E-}06$ ]. In contrast, in bulk-rock samples,  $\delta^{13}\text{C}$ -values were significantly lower in Miłoszów (-0.05, -2.24) compared to Skały 11 (1.71) and Skały C (0.96) specimens, with intermediate values in Skały A+11A rock samples [0.10;  $p(H_0) = 3.07\text{E-}11$ ].

140 Oxygen isotopic values in *Semitextularia* tests also resulted in significant differences among samples [ $p(H_0) = 3.56\text{E-}08$ ]. Averaged  $\delta^{18}\text{O}$ -values were significantly lower in Miłoszów (-6.35, -6.44) compared to Skały C (-5.68) and Skały A+11A (-5.49), while Skały 11 values fell in between (-5.95). Values for  $\delta^{18}\text{O}$  of bulk-rock samples also differed significantly [ $p(H_0) = 9.44\text{E-}08$ ], with Miłoszów samples showing significantly lower values (-6.62, -6.91) compared to Skały 11 (-5.56) and Skały C (-5.43) samples; with values for Skały A+11A samples falling in between (-6.17).

145 Comparisons among stable-isotopic values of *Semitextularia* specimens and bulk-rock samples differed (Table S7). In Miłoszów 11 [samples](#), the  $\delta^{13}\text{C}$ -values in *Semitextularia* (2.86) were much higher than for the bulk-rock sample (-0.05). Similar differences were found at Miłoszów 12, with high values for *Semitextularia* (2.71) compared to the bulk-rock sample (-0.05). Skały samples are quite different from Miłoszów. In [material from](#) both Skały 11 and Skały A+11A, the  $\delta^{13}\text{C}$ -values of bulk-rock samples were significantly higher (both samples 1.71) compared to *Semitextularia* (1.47 in Skały 11, 0.84 in Skały A+11A). At Skały C, however, the carbon isotopes are statistically similar, 1.60 for *Semitextularia* versus 0.96 for bulk rock.

Comparing oxygen-isotopic data between *Semitextularia* and bulk-rock samples produced different results. The  $\delta^{18}\text{O}$ -values do not differ between *Semitextularia* specimens and bulk rock in either Miłoszów sample (Table S7). Similar congruence was found in Skały C samples, with  $\delta^{18}\text{O}$ -values of -5.68 for *Semitextularia* specimens and -5.43 for bulk rock. 155 Although oxygen-isotopic values for Skały 11 (-5.95 for *Semitextularia*, -5.56 for bulk rock) were similar, because the variance in both groups was extremely low (0.003, 0.005), the differences were significant. In contrast, the  $\delta^{18}\text{O}$ -values of *Semitextularia* specimens (-5.49) and bulk-rock samples (0.10) from Skały A+11A were very different.

Comparisons of  $\delta^{13}\text{C}$  and  $\delta^{18}\text{O}$  values based on [foraminifers/foraminifera](#) and bulk-rock samples are shown as scatter diagrams for Miłoszów and Skały samples (Fig. 4).

## 160 4 Discussion

[A simplified depositional-biofacies model for the uppermost Eifelian Skały Formation in the Łysogóry basin \(HCM\) is provided in Figure 5, presenting our interpretation of the paleoenvironments represented by \*Semitextularia\*-bearing samples are indicated.](#) The  $^{13}\text{C}$  enrichment of *Semitextularia* foraminiferal tests compared to the associated bulk rock (matrix) was interpreted as related to the fractionation of isotopes due to photosynthetic activity in [foraminifers/foraminifera](#) that either 165 hosted photosynthetic endosymbionts or were kleptoplastic.

This hypothesis of photosynthetic activity in *Semitextularia* is based on many previous observations that intracellular photosynthesis influences the  $\delta^{13}\text{C}$  of foraminiferal tests (e.g., Ravelo and Fairbanks, 1995; Wefer et al., 1981; Wefer and Berger, 1991). Fractionation of carbon isotopes during photosynthesis can produce  $\delta^{13}\text{C}$  differences between foraminiferal tests and the  $\delta^{13}\text{C}_{\text{DIC}}$  of the ambient seawater in which the foraminifers live, as it strongly influences the foraminiferal microenvironment, i.e., the “internal carbon pool” (ICP) (e.g., Ravelo and Hillaire-Marcel, 2007, Wefer and Berger, 1991; Zeebe et al., 1999). The probability that *Semitextularia* was photosynthetically active is supported by the apparently optically-transparent lamellar test structure (Fig. 1 h–j) that enabled light penetration (Dubicka et al., in press). Furthermore, the test morphology of *Semitextularia* is comparable to representatives of some smaller SBBF to the point that *Semitextularia* seems to be the fossil morphological counterpart of peneroplids. Specimens of both foraminiferal taxa can be characterized by fan-like, bilaterally flat tests with a high surface-to-volume ratios (Hohenegger, 2009), and multiple apertures forming a series of small holes along the entire last chamber. This kind of test represents a way to increase the surface area allowing more symbionts/plastids to be positioned just beneath the test wall to gather light (e.g., Haynes, 1965; Hansen and Buchhardt, 1979), while multiple apertures are morphological adaptations to resist hydrodynamic forces by strengthening attachments through bundles of pseudopodia, as exemplified by the *Peneroplis* life strategy (Hohenegger, 2011). Specifically, peneroplids hide in the uppermost layer of sand grains or fix with a net of pseudopods to algal filaments (Fig. 2b; Hohenegger, 2009, 2011).

However, differences in  $\delta^{13}\text{C}$  values in *Semitextularia* compared to the bulk-rock samples are much smaller in Skąły A11+A samples than those recorded in specimens from Miłoszów and are very minimal in Skąły 11 and Skąły C specimens. Therefore, there is no clear evidence for photosynthetic activity in the *Semitextularia* tests from the Skąły settings, which likely were characterized by lower light intensities (Zapalski et al., 2017). The paleodepth of Skąły is estimated as ca. 30–40 meters (see Supplement Geological Setting), which is within the range of the water depth of Recent mixotrophic foraminifers, primarily those housing diatoms, which did not evolve until the Jurassic (Sims et al., 2006). Species hosting dinoflagellates, chlorophytes or rhodophytes tend to be shallower dwelling, especially in shelf-margin environments with reduced water transparency such as the Florida reef tract (Baker et al., 2009), compared to much clearer oceanic waters (Hohenegger, 2006). Thus, reduced photosynthetic activity at the seafloor at the Skąły location was likely related to input of nutrients and fine sediments into the basin, thereby limiting light penetration required for sufficient photosynthetic activity to influence carbon-isotopic ratios in the foraminiferal tests (e.g., Hansen and Buchhardt, 1979). Light penetration declines exponentially with water depth and so declines rapidly with depth even with relatively small changes in water transparency (e.g., Hallock and Schlager, 1986; Hohenegger, 2000, 2004).

Another point to discuss is the consistency in test morphology of the studied *Semitextularia* as shown by regression analyses of the  $\text{test\_width}/\text{test height}$  and  $\sqrt{\text{area}}/\text{test height}$  ratios (Fig. 3). These ratios can indicate light requirements when harboring microalgae (Hohenegger, 2004), as has been observed in the surface/volume-ratios of algal symbiont-bearing rotaliids (Röttger and Hallock, 1982) and some porcelanous species whose distributions are well correlated with depth (39, 42 e.g., Eder et al., 2018; Hallock, 1979; Hohenegger, 2000). Because light intensities would have been higher in the upper

200 euphotic Miłoszów location compared to the mesophotic Skały sites, one could expect differences in test parameters. For example, in hyaline amphisteginid species, which are characterized by highly transparent lamellar test structure, the maximum test diameters are relatively consistent within their depth range, though their minimum diameters (also referred to as “test thickness”) decline with depth (e.g., Hallock and Hansen, 1979; Hohenegger et al., 1999; Larsen and Drooger, 1977). In ~~this case~~ *Amphistegina*, optimal light penetration into the test is facilitated by ~~thinning of the~~ changes in lamellar thickness, which could not be demonstrated for *Semitextularia* because possessing nonlamellar walls like miliolid foraminifera (e.g. peneroplids). ~~However, the micro-erosion of the test surface of the specimen shown in Figure 1a, as well as irregularities in chamber walls shown in thin section (Fig. 1h1), indicate at least partial internal subdivision of the individual chambers, analogous to that seen in modern *Archaias angulatus* (e.g., Cottey and Hallock, 1988).~~

205 ~~If we suspect that *Semitextularia* was photosynthetically active, at least in quite shallow water, the question arises: were these foraminifera kleptoplastic, as seen in some modern taxa (Jaufrais et al., 2016, 2018), or were they indeed symbiotic with some kind of microalgae? And if *Semitextularia* were photosymbiotic, was the relationship a facultative symbioses or an obligate symbiosis? And in the latter case, were the algal cells essentially organelles, as in peneroplids in which the red algal cells are not bound by a host membrane (e.g., Lee and Anderson, 1991, and references therein).~~ was kleptoplastic, maintaining plastids that photosynthesized most actively in the upper euphotic zone. ~~These questions cannot be answered with our current specimens and isotopic data.~~ Photosynthetic activity sufficient to influence carbon-isotopic ratios ~~is known to declines~~ with depth in ~~foraminifers~~ foraminifera with diatom endosymbionts (Hansen and Buchardt, 1979). Moreover, obligate endosymbiosis is ~~likely~~ a more complex co-evolutionary process, especially obligate photosymbioses in which the host is unable to survive without symbionts. Facultative algal symbioses, which occur in some planktic taxa, are unknown among Recent SBBF (e.g., Hallock, 1999; Lee and Anderson, 1991; Prazeres and Renema, 2019). ~~Because photosymbiosis in benthic foraminifera may have been somewhat unstable at the beginning, *Semitextularia* might have been benefitting from also mixotrophyic most of the times and just by sporadically sequestering host and maintaining chloroplasts off from their prey algal food is a much simpler adaptation and therefore a parsimonious hypothesis, as this photosymbiotic strategy could have been rather unstable at the beginning. Mixotrophic strategy might be additionally~~ Evidence for photosynthetic activity, whether kleptoplastic or symbiotic, is supported by our isotopic data, specifically different the isotopic contrast differences between ~~foraminiferal tests and bulk rock samples, which seems to be~~ appears to have been related to the paleodepth/illumination intensity.

225 ~~The basic life history strategy of the Foraminifera involves alternation of generations (Goldstein, 1997), and can be assumed for *Semitextularia* based both on phylogeny and by the possible triadimorphism apparent in fossilized tests. For example, the initial chambers of the specimen in Figure 1e appears to be much smaller than those of other specimens shown, possibly indicating in Figure 1. This difference is consistent with a sexually-produced agamont; (microspheric-microsphere, (B form) individuals. Diameters of initial chambers in specimens in Figure 1b, d, f, are nearly double that of 1e, while diameter of initial chambers of the specimen in Figure 1c is about three times larger. The specimen shown in Figure 1a is poorly preserved, but does resemble that of Figure 1c. These differences are similar to the size range demonstrated for *Amphistegina*~~

235 *gibbosa* by Harney et al. (1998) for ~~microspheric agamonts (microspheric, sexually-produced, B form) specimens,~~  
megalospheric specimens produced by megalospheric parents (schizonts, A1 forms), and megalospheric specimens that  
produced gametes (gamonts, A2 form), compared to generally more common asexually produced, megalospheric individuals.  
As ~~hypothesized in previous studies~~ hypothesized by ~~(Dettmering et al., 1998) and~~ Krüger et al. (1996), the transfer of  
endosymbionts during sexual reproduction is certainly more complicated than direct transfer during asexual reproduction.  
240 Thus, successive asexual generations (schizogony) as a key strategy for local population increase, while sexual reproduction  
facilitates dispersal, is highly advantageous in foraminiferal taxa that host algal endosymbionts (e.g., Hallock and Seddighi,  
2021; Pappazoni and Seddighi, 2018). Future studies that examine initial chamber-size distributions in fossil populations of  
*Semitextularia* or other suspected photosymbiotic taxa could provide additional evidence for symbiosis rather than  
kleptoplasty.

The evolution of foraminiferal lineages characterized by algal endosymbiosis has occurred repeatedly and  
245 independently numerous times since the late Paleozoic, involving unrelated algal groups as well. Previously, the earliest  
calcareous ~~foraminifers~~ foraminifera postulated to host algal endosymbionts were the large, complex fusulinids (e.g.,  
BouDagher-Fadel, 2018, and references therein). A true symbiotic relationship between *Semitextularia* and an algal taxon  
cannot be ruled out from the observations reported here. Nevertheless, our observations are also consistent with kleptoplasty,  
which is ~~the a~~ the simplest hypothesis to explain photosynthetic activity in *Semitextularia*.

## 250 **5 Conclusions**

~~Summarizing~~, *Semitextularia*, an extinct and pioneering multi-chambered calcareous foraminiferal genus (Eifelian,  
Middle Devonian), shows morphological homology and comparable habitat with ~~the~~ recent photosymbiont-bearing  
~~peneroplids~~, *Peneroplis*. The  $\delta^{13}\text{C}$  enrichment, up to 2.86‰ in relation to rock-matrix samples, indicates that *Semitextularia*  
255 ~~could have utilized photosynthetic activity, either been through~~ kleptoplasty ~~ie by~~, acquiring and maintaining plastids from ~~their~~  
microalgal ~~prey food, or by benefit directly from photosynthesis. Another possibility is that these foraminifers maintain~~ ing  
algal endosymbionts, which would have involved co-evolution with a microalgal taxon. This beneficial ecological behavior  
could have had an impact on the evolutionary radiation of calcareous multi-chambered Foraminifera during the mid-Devonian  
radiation of carbonate-producing biotas.

## 260 **Author Contributions**

ZD developed the original idea presented in the manuscript. ZD and WK designed research. MG performed foraminiferal  
extraction and prepared samples for isotopic studies. ZD, MG and WK carried out fieldwork. ZD, MG, WK and JH prepared  
figures; JH provided statistical analyses. All authors analyzed data and ~~wrote~~ contributed to writing the paper.

## 265 **Competing Interest Statement**

The authors declare no competing interests.



### **Financial support**

270

This research was supported by NCN UMO (2018/29/B/ST10/01811). Z.D. thanks the Alexander von Humboldt Foundation for supporting her research stay at the Alfred Wegener Institute.

## References

- ~~Miller, B. B. and Carter, C.: The test article, J. Sci. Res., 12, 135–147, doi:10.1234/56789, 2015.~~
- 275 ~~Smith, A. A., Carter, C., and Miller, B. B.: More test articles, J. Adv. Res., 35, 13–28, doi:10.2345/67890, 2014.~~
- ~~van Edig, X., Schwarze, S., and Zeller, M.: The robustness of indicator based poverty assessment tools in changing environments – empirical evidence from Indonesia, in: Tropical Rainforests and Agroforests under Global Change, Environmental Science and Engineering (Environmental Engineering), edited by: Tschardtke, T., Leuschner, C., Veldkamp, E., Faust, H., Gubardja, E., and Bidin, A., Springer, Berlin, Heidelberg, Germany, 191–211, [https://doi.org/10.1007/978-3-642-00493-3\\_9](https://doi.org/10.1007/978-3-642-00493-3_9), 2010.~~
- 280 Baccaert, J.: Foraminiferal bio- and thanatocoenoses of reef flats, Lizard Island, Great Barrier Reef, Australia. Nature of Substrate, Annales de la Société royale zoologique de Belgique, 116, 3–14, 1986.
- Baker, R. D., Hallock, P., Moses, F. E., Williams, D. E., and Ramirez, A.: Larger foraminifers of the Florida Reef Tract, USA: Distribution patterns on reef-rubble habitats, J. Foram. Res., 39, 267–277, 2009.
- 285 Biernat, G.: Middle Devonian brachiopods from the Bodzentyn Syncline (Holy Cross Mountains, Poland), Acta Palaeontol. Pol., 17, 1–162, 1966.
- Biernat, G.: Colour pattern in Middle Devonian rhynchonellid brachiopods from the Holy Cross Mountains, Acta Geol. Pol., 34, 63–72, 1984.
- [BouDagher-Fadel MK. Evolution and Geological Significance of Larger Benthic Foraminifera, 2<sup>nd</sup> ed. London: UCL Press; 704pp, 2018.](#)
- 290 ~~BouDagher Fadel, M.: The Palaeozoic Larger Benthic Foraminifera, in: Evolution and Geological Significance of Larger Benthic Foraminifera, edited by BouDagher Fadel, M, Elsevier Science, 45–160, 2008.~~
- Cesbron, E. Geslin, C. Le Kieffre, T., and Jauffrais, T.: Sequestered chloroplasts in the benthic foraminifer *Haynesina germanica*: Cellular organization, oxygen fluxes and potential ecological implications, J. Foram. Res., 47, 268–278, 2017.
- 295 [Cotter, T. L., and Hallock, P.: Test surface degradation in \*Archaias angulatus\*, J. Foram. Res., 18, 187–202.](#)
- Dettmering, C., Röttger, R., Hohenegger, J., and Schmaljohann, R.: The trimorphic life cycle in foraminifera: Observations from cultures allow new evaluation, Eur. J. Protistol., 34, 363–368, 1998.
- Dubicka, Z., Gajewska, M., Kozłowski, W., and Mikhalevich, V.: Test structure in some pioneer multichambered Paleozoic Foraminifera, Proc. Natl. Acad. Sci. U.S.A (in press).
- 300 [Dubinsky, Z., and Berman-Frank, I.: Uncoupling primary production from population growth in photosynthesizing organisms in aquatic ecosystems, Aquat. Sci., 63, 4–17, 2001.](#)
- [Eder, W., Hohenegger, J., and Briguglio, A.: Test flattening in the larger foraminifer \*Heterostegina depressa\*: Predicting bathymetry from axial sections, Paleobiology, 44, 76–88, 2018.](#)

- 305 Fujita, K. and Fujimura, H.: Organic and inorganic carbon production by algal symbiont-bearing foraminifera on northwest Pacific coral-reef flats, *J. Foram. Res.*, 38, 117–126, 2008.
- Goldstein, S. T.: Gametogenesis and the antiquity of reproductive pattern in the Foraminiferida, *J. Foram. Res.*, 27, 319–328, 1997.
- Goldstein, S. T., Bernhard, J. M., and Richardson, E. A.: Chloroplast sequestration in the foraminifer *Haynesina germanica*: Application of high pressure freezing and freeze substitution, *Microsc. Microanal.*, 10, 1458–1459, 2004.
- 310 Gorzelak, P., Stolarski, J., Dubois, P., Kopp, C., and Meibom, A.: <sup>26</sup>Mg labeling of the sea urchin regenerating spine: Insights into echinoderm biomineralization process, *J. Struct. Biol.*, 176, 119–126, 2011.
- Hallock, P.: Trends in test shape in large, symbiont-bearing Foraminifera, *J. Foram. Res.*, 9, 61–69, 1979.
- Hallock, P.: Light dependence in *Amphistegina*, *J. Foram. Res.*, 11, 40–46, 1981a.
- 315 [Hallock, P.: Algal symbiosis: a mathematical analysis. \*Mar. Biol.\* 6, 249–255, 1981b.](#)
- [Hallock, P.: Fluctuations in the trophic resource continuum: a factor in global diversity cycles?, \*Paleoceanography\*, 2, 457–471, 1987.](#)
- Hallock, P.: Symbiont-bearing Foraminifera, in: *Modern Foraminifera*, edited by Sen Gupta, B. K., Springer, 123–39, 1994.
- Hallock, P. and Hansen, H. J.: Depth adaptation in *Amphistegina*: Change in lamellar thickness, *Bull. Geol. Soc.*, 27, 99–104, 1979.
- 320
- Hallock, P. and Schlager, W.: Nutrient excess and the demise of coral reefs and carbonate platforms, *Palaios*, 1, 389–398, 1986.
- [Hallock, P., and Seddighi, M. Why did some larger benthic foraminifera become so large and flat? \*Sedimentology\*, doi: 10.1111/sed.12837, 2021.](#)
- 325
- Hammer, Ø., Harper, D., and Paul, D. R.: Past: Paleontological Statistics Software Package for Education and Data Analysis, *Palaeontologia Electronica*, 4, 1–9, 2001.
- Hansen, H. J. and Buchhardt, B.: Depth distribution of *Amphistegina* in the Gulf of Elat, *Utrecht Micropaleontol. Bull.*, 30, 205–244, 1979.
- [Harney, J., Hallock, P., and Talge, H.K.: Observations on a trimorphic life cycle in \*Amphistegina gibbosa\* populations from the Florida Keys. \*J. Foram. Res.\*, 28, 141–147, 1998.](#)
- 330
- Haynes, J.: Symbiosis, wall structure and habitat in foraminifera, *Contrib. Cushman Found. Foram. Res.*, 16, 40–44, 1965.
- Hohenegger, J.: Coenoclines of larger foraminifera: *Micropaleontol.*, 46, 127–151, 2000.
- Hohenegger, J.: Depth coenoclines and environmental considerations of western Pacific larger foraminifera, *J. Foram. Res.*, 34, 9–34, 2004.
- 335
- Hohenegger, J.: The importance of symbiont-bearing benthic foraminifera Symbiont-Bearing Benthic Foraminifera for West Pacific Carbonate Beach Environments, *Mar. Micropaleontol.*, 61, 4–39, 2006.
- Hohenegger, J.: Functional Shell Geometry of Symbiont-bearing Benthic Foraminifera, *Galaxea JCRS*, 11, 81–89, 2009.


- Hohenegger, J.: Large Foraminifera: Greenhouse constructions and gardeners in the oceanic microcosm, The Kagoshima University Museum, Kagoshima, Japan, 2011.
- 340 Hohenegger, J.: Foraminiferal growth and test development, *Earth-Sci. Rev.*, 185, 140–162, 2018.
- Hohenegger, J., Yordanova, E., Nakano, Y., and Tatzreiter, F.: Habitats of larger foraminifera on the upper reef slope of Sesoko Island, Okinawa, Japan, *Mar. Micropaleontol.* 36, 109–168, 1999.
- [Hottinger, L.: Larger Foraminifera, giant cells with a historical background. \*Naturwissenschaften\*, 69, 361–371, 1982.](#)
- Jarochovska, E., Tonarová, P., Munnecke, A., Ferrová, L., Sklenář, J., and Vodrážková, S.: An acid-free method of microfossil  
345 extraction from clay-rich lithologies using the surfactant Rewoquat, *Palaeontol. Electron.*, 16, 1-16, 2013.
- Jauffrais, T., Jesus, B., Metzger, E., Mouget, J. L., Jorissen, F., and Geslin, E.: Effect of light on photosynthetic efficiency of sequestered chloroplasts in intertidal benthic foraminifera (*Haynesina germanica* and *Ammonia tepida*), *Biogeosciences*, 13, 2715–2726, 2016.
- Jauffrais, T., LeKieffre, C., Koho, K. A., Tsuchiya, M., Schweizer, M., Bernhard, J. M., Meibom, A., and Geslin, E.:  
350 Ultrastructure and distribution of kleptoplasts in benthic foraminifera from shallow-water (photic) habitats, *Mar. Micropaleontol.*, 138, 46–62, 2018.
- Jauffrais, T., LeKieffre, C., Schweizer, M., Geslin, E., Metzger, E., Bernhard, J. M., Jesus, B., Filipsson, H. L., Maire, O., and Meibom, A.: Kleptoplastidic benthic foraminifera from aphotic habitats: Insights into assimilation of inorganic C, N and S studied with sub-cellular resolution, *Environ. Microbiol.*, 21, 125–141, 2019.
- 355 [Kinoshita, S., Eder, W., Wo, J., Hohenegger, J., and Briguglio, A.: Growth, chamber building rate and reproduction time of \*Palaeonummulites venosus\* \(Foraminifera\) under natural conditions. \*Coral Reefs\*, 36, 1097–1109, 2017.](#)
- Krüger, R. Röttger, R. Lietz, R., and Hohenegger, J.: Biology and reproductive process of the large foraminiferan *Cycloclypeus carpenteri* (Protozoa, Nummulitidae), *Archives for Protist Studies*, 147, 307–321, 1996.
- ~~Langer, W.: New calcareous microproblematica from the Devonian of Western 269 Germany (in German), *Neues Jahrb. Geol. Palaontol. Abh.*, 723–733, 1979.~~  
360 [Langer, M. R., Silk, M. T. and Lipps, J. H.: Global ocean carbonate and carbon dioxide production: the role of reef foraminifera. \*J. Foram. Res.\*, 27, 271–277, 1997.](#)
- Larsen, A. R. and Drooger, C. W.: Relative thickness of the test in the *Amphistegina* species of the Gulf of Elat, *Utrecht Micropal. Bull.*, 15, 225–240, 1977.
- 365 Lee, J. J. and Anderson, O. R.: Symbiosis in Foraminifera, in: *Biology of Foraminifera*, edited by Lee, J. J. and Anderson, O. R., LA Press, London, 157–220, 1991.
- Lee, J. J., Cervasco, M., Morales, J., Billik, M., and Fine, M.: Symbiosis ~~d~~Drove ~~c~~Cellular ~~e~~Evolution: Symbiosis ~~f~~Fueled ~~e~~Evolution of ~~l~~Lineages of Foraminifera (~~e~~Eukaryotic ~~c~~Cells) into ~~e~~Exceptionally ~~c~~Complex ~~g~~Giant ~~p~~Protists, *Symbiosis*, 51, 13–25, 2010.
- 370 McConnaughey, T. A. and Whelan, J. F.: Calcification generates protons for nutrient and bicarbonate uptake, *Earth-Sci. Rev.*, 42, 95-117, 1997.


- Miller, A.K. and Carmer, A.M.: Devonian Foraminifera from Iowa. *Journal of Paleontology*, 7, p. 423–431, 1933.
- Miller, B. B. and Carter, C.: The test article, *J. Sci. Res.*, 12, 135–147, doi:10.1234/56789, 2015.
- Smith, A. A., Carter, C., and Miller, B. B.: More test articles, *J. Adv. Res.*, 35, 13–28, doi:10.2345/67890, 2014.
- 375 van Edig, X., Schwarze, S., and Zeller, M.: The robustness of indicator based poverty assessment tools in changing environments — empirical evidence from Indonesia, in: *Tropical Rainforests and Agroforests under Global Change, Environmental Science and Engineering (Environmental Engineering)*, edited by: Tschardtke, T., Leuschner, C., Veldkamp, E., Faust, H., Guhardja, E., and Bidin, A., Springer, Berlin, Heidelberg, Germany, 191–211, <https://doi.org/10.1007/978-3-642-00493-3-9>, 2010.
- 380 Narkiewicz, K. and Malec, J.: New conodont CAI database (CAI) [in Polish], *Przegląd Geologiczny*, 53, 33–37, 2005.
- Pajchłowa, M.: Dewon w profilu Grzegorzowice-Skały (in Polish with English summary), *Biuletyn Instytutu Geologicznego*, 122, 145–254, 1957.
- Papazzoni, C.A. and Seddighi, M.: What, if anything, is a Nummulite Bank? *J. Foram. Res.*, 48, 276–287, 2018.
- Pillet, L., de Vargas, C., and Pawłowski, J.: Molecular identification of sequestered diatom chloroplasts and kleptoplastidy in  
385 foraminifera, *Protist*, 162, 394–404, 2011.
- Prazeres, M. and Renema, W.: Evolutionary sSignificance of the mMicrobial aAssemblages of lLarge bBenthic fForaminifera, *Biol. Rev.*, 94, 828–848, 2019.
- Ravelo, A. C. and Fairbanks, R. G.: Carbon isotopic fractionation in multiple species of planktonic foraminifera from core-tops in the tropical Atlantic, *J. Foram. Res.*, 25, 53–74, 1995.
- 390 Ravelo, A. C. and Hillaire-Marcel, C.: The Use of Oxygen and Carbon Isotopes of Foraminifera in Paleooceanography, in: *Proxies in Late Cenozoic Paleooceanography*, edited by Hillaire-Marcel, C and De Vernal, A., Elsevier Science, 735–764, 2007.
- Renema, W.: Terrestrial Influence as a Key Driver of Spatial Variability in Large Benthic Foraminiferal Assemblage Composition in the Central Indo-Pacific, *Earth-Sci. Rev.*, 177, 514–544, 2018.
- 395 Röttger, R. and Hallock, P.: Shape trends in *Heterostegina depressa* (Protozoa, Foraminiferida), *J. Foram. Res.*, 12, 197–204, 1982.
- Samsonowicz, J.: Sprawozdanie z badań w r. 1935 na północ od kopalni Staszic między Pokrzywianką. Psarką i Swiśliną (in Polish), *Posiedzenie naukowe Państwowego Instytutu Geologicznego*, 44, 41–45, 1936.
- Schmidt, D. N., Renaud, S., Bollmann, J., Schiebel, R., and Thierstein, H. R.: Size distribution of Holocene planktic foraminifer  
400 assemblages: biogeography, ecology and adaption, *Mar. Micropaleontol.*, 50, 319–338, 2004.
- Selosse, M. A., Charpin, A., and Not, F.: Mixotrophy everywhere on land and in water. The grand écart hypothesis, *Ecol. Lett.*, 20, 246–263, 2017.
- Sims, P. A., Mann, D. G., and Medlin, L. K.: Evolution of the diatoms: Insights from fossil, biological and molecular data, *Phycologia*, 45, 361–402, 2006.

- 405 Stasińska, A.: Tabulata. Heliolitida et Chaetetida du Devonien moyen des monts de Sainte-Croix, *Acta Palaeontol. Pol.*, 3, 161–282, 1958.
- Szulcowski, M.: Depositional evolution of the Holy Cross Mts. (Poland) in the Devonian and Carboniferous – a review, *Geological Quarterly*, 39, 471-488, 1995.
- ter Kuile, B.: Mechanisms for calcification and carbon cycling in algal symbiont-bearing foraminifera, in: *Biology of Foraminifera*, edited by Lee, J. J. and Anderson, O. R., London Academic Press, London, UK, 73-89, 1991.
- 410 Vachard, D., Haig, D. W, and Mory, A. J.: Lower Carboniferous (middle Visean) foraminifera and algae from an interior sea, Southern Carnarvon Basin, Australia, *Geobios*, 47, 57–74, 2014.
- Vachard, D., Pille, L., and Gaillot, J.: Palaeozoic Foraminifera: Systematics, Palaeoecology and Responses to Global Changes, *Rev. de Micropaleontol.*, 53, 209-254, 2010.
- 415 Wefer, G. and Berger, W. H.: Isotope paleontology: growth and composition of extant calcareous species, *Mar. Geol.*, 100, 207-248, 1991.
- Wefer, G., Killingley, J. S., and Lutze, G. F.: Stable isotopes in recent larger foraminifera, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 33, 253-270, 1981.
- [Wooldridge, S.A.: Is the coral-algae symbiosis really ‘mutually beneficial’ for the partners? \*Bioessays\*, 32, 615–625, 2010.](#)
- 420 Zapalski, M. K., Wrzolek, T., Skompski, S., and Berkowski, B.: Deep in shadows, deep in time: The oldest mesophotic coral ecosystems from the Devonian of the Holy Cross Mountains (Poland), *Coral Reefs*, 36, 847–860, 2017.
- Zeebe, R. E., Bijma, J., and Wolf-Gladrow, D. A.: A diffusion-reaction model of carbon isotope fractionation in foraminifera, *Marine Chemistry*, 64, 199-227, 1999.
- 425 Zeuschner, L.: Geognostische Beschreibung der mittleren devonischen Schichten zwischen Grzegorzowice und Skaly-Zagaje bei Nowa Slupia, *Zeitschrift der Deutschen Geologischen Gesellschaft*, 21, 263–274, 1869.

Figure captions

430






435 **Figure 1. Images of *Semitextularia* (Miller and Carmer, 1933) specimens: (a–g) SEM images of the *Semitextularia* tests; (h) internal test texture in a conventional light-microscope image (h1) and using polarized-light microscopy (h2); (i) image of *Semitextularin* thin section showing transparent test (a–i; scale bars are equal to 100  $\mu\text{m}$ ); (j–l) SEM images showing lamellar-foliated fibrous structure of the test wall (a) Miłoszów, sample B1, MWGUW ZI/67/MG7.02. (b) Miłoszów, sample 11, MWGUW ZI/67/MG5.03. (c) Miłoszów, sample 11, MWGUW ZI/67/MG5.04. (d) Miłoszów, sample 12, MWGUW ZI/67/MG5.30. (h) Miłoszów, sample 12, MWGUW ZI/67/MG5.27. (i) Miłoszów, sample MO, MWGUW ZI/67/MG5.40. (j) Miłoszów, sample 12R, MWGUW ZI/67/MG9.23. In image c, x indicates test height (i.e., maximum diameter) while y is test width.**

440


**Figure 1. SEM images of *Semitextularia* (Miller and Carmer, 1933) specimens (a–g; scale bars are equal to 100  $\mu\text{m}$ ) and cross sections of the *Semitextularia* test (h–j) showing internal lamellar test texture as well as conventional light microscope image (h1) and polarizing light microscopy (under crossed nicols) (h2, i) images of *Semitextularin* thin**



445 ~~section showing transparent test (a-i; scale bars are equal to 100  $\mu$ m). (a) Miłoszów, sample B1, MWGUW ZI/67/MG7.02. (b) Miłoszów, sample 11, MWGUW ZI/67/MG5.03. (c) Miłoszów, sample 11, MWGUW ZI/67/MG5.04. (d) Miłoszów, sample 12, MWGUW ZI/67/MG5.30. (e) Miłoszów, sample 12, MWGUW ZI/67/MG5.27. (f) Miłoszów, sample MO, MWGUW ZI/67/MG5.40. (g) Miłoszów, sample 12R, MWGUW ZI/67/MG9.23.~~




450 **Figure 2.** (a) Living peneroplids and *Neorotalia calcar* from algal mats on reef crests in Belau. (b) *Peneroplis planatus* attached to filamentous microalgae within sand grains from the reef crest at Sesoko Jima, Japan, together with *Calcarina gaudichaudii* and *Baculogypsina sphaerulata*.




**Figure 3.** Relationships between test area and test height fitted by power functions.

455



**Figure 4.** Variations in  $\delta^{13}\text{C}$  and  $\delta^{18}\text{O}$  values of *Semitextularia* tests and bulk-rock samples from Miłoszów and Skały.



460 **Figure 5.** Simplified depositional-biofacies model for the uppermost Eifelian Skaly Formation in the Lysogóry basin (HCM), with *Semitextularia*-bearing samples localization.