

SI of The Cretaceous physiological adaptation of angiosperms to a declining pCO₂ : a trait-oriented modelling approach

Julia Bres¹, Pierre Sepulchre¹, Nicolas Viovy¹, and Nicolas Vuichard¹

¹Laboratoire des Sciences du Climat et de l'Environnement, LSCE/IPSL, CEA-CNRS-UVSQ, Université Paris-Saclay, 91191 Gif-sur-Yvette, France

Correspondence: Julia Bres (julia.bres@lsce.ipsl.fr)




Figure S 1. Annual zonal mean of (a) LAI ($\text{m}^2[\text{leaf}] \text{m}^{-2}[\text{ground}]$) and (b) soil moisture stress for transpiration (unitless) for ANGIO(1120) (black line), ANGIO(280) (black dashed line), NOANGIOhp(1120) (gray line) and NOANGIOhp(280) (gray dashed line).




Figure S 2. Annual mean GPP ($\text{kgC m}^{-2} \text{y}^{-1}$) for a) ANGIO(1120), b) ANGIO(280), c) NOANGIOh(1120), d) NOANGIOh(280), e) NOANGIOp(1120), f) NOANGIOp(280), g) NOANGIOhp(1120) and h) NOANGIOhp(280).




Figure S 3. Annual canopy stomatal conductance ($\text{mol m}^{-2}[\text{ground}] \text{s}^{-1}$) for a) ANGIO(1120) and b) ANGIO(280), anomalies of annual mean canopy stomatal conductance ($\text{mol m}^{-2}[\text{ground}] \text{s}^{-1}$) for c) NOANGIOh(1120) vs ANGIO(1120), d) NOANGIOh(280) vs ANGIO(280), e) NOANGIOp(1120) vs ANGIO(1120), f) NOANGIOp(280) vs ANGIO(280), g) NOANGIOhp(1120) vs ANGIO(1120) and h) NOANGIOhp(280) vs ANGIO(280). The t-test 95 % confidence level anomalies are given by dots.




Figure S 4. Annual mean potential evaporation (mm d^{-1}) for a) ANGIO(1120) and b) ANGIO(280), anomalies of annual mean potential evaporation (mm d^{-1}) for c) NOANGIOh(1120) vs ANGIO(1120), d) NOANGIOh(280) vs ANGIO(280), e) NOANGIOp(1120) vs ANGIO(1120), f) NOANGIOp(280) vs ANGIO(280), g) NOANGIOhp(1120) vs ANGIO(1120) and h) NOANGIOhp(280) vs ANGIO(280). The t-test 95 % confidence level anomalies are given by dots.