

Quantification of Blue Carbon in Salt Marshes of the Pacific Coast of Canada

Stephen G. Chastain¹, Karen E. Kohfeld^{1,2}, Marlow G. Pellatt^{1,3}, Carolina Olid^{4,5}, Maija Gailis⁶

¹School of Resource & Environmental Management, Simon Fraser University, Burnaby, Canada V5A 1S6. Coast Salish Territories, xʷməθkʷəy̓əm (Musqueam), Skwxwú7mesh (Squamish) & səlilwətaɬ (Tsleil-Waututh)

²School of Environmental Science, Simon Fraser University, Burnaby, Canada, V5A 1S6

³Parks Canada, Protected Areas Establishment and Conservation Directorate, Vancouver, British Columbia, Canada V6B 6B4

⁴Department of Forest Ecology and Management, Swedish University of Agricultural Science, Umeå, Sweden

⁵UB-Geomodels Research Institute, Departament de Dinàmica de la Terra i l'Oceà, Facultat de Ciències de la Terra, Universitat de Barcelona, 08028 Barcelona, Spain

⁶Environment and Climate Change Canada, Climate Change Branch, Ottawa, Ontario, Canada

Correspondence to: Karen E Kohfeld (kohfeld@sfu.ca)

Abstract. Tidal salt marshes are known to accumulate “blue carbon” at high rates relative to their surface area, which render these systems among the Earth’s most efficient carbon (C) sinks. However, the potential for tidal salt marshes to mitigate global warming remains poorly constrained because of the lack of representative sampling of tidal marshes from around the globe, inadequate areal extent estimations, and inappropriate dating methods for accurately estimating C accumulation rates. Here we provide the first estimates of organic C storage and accumulation rates in salt marshes along the Pacific Coast of Canada, within the Clayoquot Sound UNESCO Biosphere Reserve and Pacific Rim National Park Reserve, a region currently underrepresented in global compilations. Within the context of other sites from the Pacific Coast of North America, these young Clayoquot Sound marshes have relatively low C stocks but are accumulating C at rates that are higher than the global average, with pronounced differences between high and low marsh habitats. The average C stock calculated during the past 30 years is $54 \pm 5 \text{ Mg C ha}^{-1}$ (mean \pm standard error), which accounts for 81 % of the C accumulated to the base of the marsh peat layer ($67 \pm 9 \text{ Mg C ha}^{-1}$). The total C stock is just under one-third of previous global estimates of salt marsh C stocks, likely due to the shallow depth and young age of the marsh. In contrast, the average C accumulation rate (CAR) ($184 \pm 50 \text{ g C m}^{-2} \text{ yr}^{-1}$ to the base of the peat layer) is higher than both CARs from salt marshes along the Pacific coast ($112 \pm 12 \text{ g C m}^{-2} \text{ yr}^{-1}$) and global estimates ($91 \pm 7 \text{ g C m}^{-2} \text{ yr}^{-1}$). This difference was even more pronounced when we considered individual marsh zones: CARs were significantly greater in high marsh ($303 \pm 45 \text{ g C m}^{-2} \text{ yr}^{-1}$) compared to the low marsh sediments ($63 \pm 6 \text{ g C m}^{-2} \text{ yr}^{-1}$), an observation unique to Clayoquot Sound among NE Pacific Coast marsh studies. We attribute low CARs in the low marsh zones to shallow-rooting vegetation, reduced terrestrial sediment inputs, negative relative sea level rise in the region, and enhanced erosional processes. Per-hectare, CARs in Clayoquot Sound marsh soils are approximately 2-7 times greater than C uptake rates based on net ecosystem productivity in Canadian boreal forests, which highlights their potential importance as C reservoirs and the need to consider their C accumulation capacity as a climate mitigation co-benefit when conserving for other salt marsh ecosystem services.

35 1 Introduction

Coastal, vegetated ecosystems, such as seagrass meadows, mangroves, and tidal salt marshes, have recently been recognized for their ability to store large amounts of “blue carbon” within their soils and sediments (Kennedy et al., 2013; Howard et al. 2014). While blue carbon ecosystems cover approximately 0.2 % of the ocean surface, they have been estimated to be responsible for up to 50 % of total coastal ocean carbon (C) burial (including estuaries and continental platforms) (Duarte et al. 2005), and their per-area C sequestration rate is substantially greater than that of terrestrial forest soils (McLeod et al. 2011). Globally, blue C ecosystems have been estimated to sequester between 75 and 224 Tg C yr⁻¹ (Duarte et al. 2013). For comparison, deep ocean organic carbon burial rates were recently estimated to be $20 \pm 6 \text{ Tg C yr}^{-1}$ (Hayes et al. 2021), with an additional burial of 222 Tg C yr⁻¹ estimated for shelves and platforms shallower than 1000 m (Burdige, 2007). Due to this high C storage and accumulation rate capacity per unit area, coastal vegetated ecosystems have been suggested to play an important role in climate warming mitigation (Howard et al. 2017). However, when the ecosystem is degraded, the stored C can be released, and annual C uptake by the ecosystem ceases, resulting in losses of ecosystem services (McLeod et al. 2011, Pendleton et al 2012). Thus, to better inform policies that identify priority areas for conservation, more precise measurement of C stocks and accumulation potential are needed (Howard et al. 2017).

Global estimates of salt marsh area, C stocks, and C accumulation rates (CAR) are subject to large uncertainties. Duarte et al. (2013) noted a 20-fold uncertainty in global estimates of salt marsh area (ranging from 22,000-to 400,000 km²) associated with ambiguous classification schemes for wetlands. For example, some classification systems consider freshwater and saltwater marshes in the same category (Duarte et al. 2013). Similarly, the estimated, global soil C stock of all salt marshes ranges

Deleted: IPCC 2014


between 0.4 and 6.5 Pg C, a 16-fold range (Duarte et al. 2013). The average global CAR for salt marshes was estimated as 91
55 $\pm 19 \text{ g C m}^{-2} \text{ yr}^{-1}$ by the Intergovernmental Panel on Climate Change (IPCC) (Kennedy et al., 2013) and then as $245 \pm 26 \text{ g C m}^{-2} \text{ yr}^{-1}$ in a subsequent global compilation (mean \pm standard deviation) (Ouyang and Lee 2014). Furthermore, these reviews
of salt marsh CAR estimates note disproportionate representation from certain areas of the world (Ouyang and Lee 2014;
Chmura et al. 2003). Some areas, such as Europe and eastern North America, have dozens of CAR data points, while others,
such as western North America, East Asia, and Australia are underrepresented. Regions such as Africa, India, and South
60 America have fewer or no data at all. The high variability in CAR from site to site combined with the 20-fold uncertainty in
global marsh area estimates result in global salt marsh CAR estimates ranging from 0.9 to 31.4 Tg C yr⁻¹ (Ouyang and Lee
2014). This 35-fold range is seven times greater than the global range for mangroves (Ouyang and Lee 2014; Donato et al.
2011).

The extensive use of ¹³⁷Cs radioisotope or a marker horizon method for sediment dating also limits comparisons of CAR
65 between studies. The difficulties of quantifying low concentrations of ¹³⁷Cs in coastal sediments, and its low retention in high
organic content sediments (Davis et al. 1984), limits the applicability of this technique to estimate CAR in salt marshes and
leads to overestimates of sediment accumulation rates when compared to other dating methods such as ²¹⁰Pb dating (e.g.
70 Callaway et al. 2012; Johannessen and MacDonald 2016). Unlike ¹³⁷Cs, the natural radionuclide ²¹⁰Pb is highly retained in
organic matter, which makes it suitable for establishing the chronology of sedimentary deposits accumulated over
approximately the past 100 years (Krishnaswamy et al. 1971; Arias-Ortiz et al. 2018). Also, the lower detection limits of ²¹⁰Pb
reduce analytical errors in the derived chronologies below those obtained by using ¹³⁷Cs providing, thus, better constrained
recent CAR for salt marshes (e.g., Corbett and Walsh, 2015).

The Commission for Environmental Cooperation (CEC), a tri-national governmental organization promoting scientific
cooperation between Canada, the United States, and Mexico, identified the Pacific coast of Canada as a significant data gap in
75 terms of our knowledge of the aerial extent and quantification of blue C storage and accumulation. Additionally, a review of
global salt marsh CAR data identified only eight sites on the entire Pacific coast of the continent, none of which were north of
38.2 °N (Ouyang and Lee 2014). This study aims to address this data gap by providing C stock and CAR from the Pacific
Coast of Canada as a part of the Government of Canada's contribution to a continent-wide assessment of blue C mitigation
potential. We sampled seven salt marshes within the United Nations Educational, Scientific, and Cultural Organization
80 (UNESCO) Clayoquot Sound Biosphere Reserve, British Columbia's Tofino Mudflats Wildlife Management Area, and Pacific
Rim National Park Reserve of Canada on Vancouver Island, British Columbia (Figure 1). These mesotidal estuarine marshes,
often constrained in size by surrounding topography, are typical of the marshes found on the Pacific coast of British Columbia,
and therefore provide a good representation of many of the tidal wetland ecosystems found here (Ryder et al. 2007). We
calculated soil C stocks (per unit area) across the high and low marsh zones of each marsh and used ²¹⁰Pb dating in a subset of
85 the collected cores to quantify CAR. We then used the extent of high and low marsh areas from aerial imagery to estimate total
C storage and total annual C accumulation. Finally, we place these new data within the context of CAR estimates from salt
marshes in the NE Pacific region.

Deleted: IPCC

Deleted: Fig. 1



90

Figure 1. Site locations. Cypress River Flats (CRF), Cannery Bay East (CBE), Cannery Bay West (CBW), Kennedy Cove South (KCS), Shipwreck Cove (SWC), Grice Bay at Kootowis Creek (GBK), and Tofino Mudflats (TMF). Pacific Rim National Park and Reserve (blue crosshatching) covers the southern portion of the map and the Tofino Mudflats Wildlife Management Area (pink crosshatching) covers portions of the southwestern area. The study region lies within the Clayoquot Sound UNESCO Biosphere Reserve, British Columbia, Canada (inset, outlined).

95

2. Methods

2.1 Study Area

Clayoquot Sound is a complex system of inlets on the west coast of Vancouver Island, British Columbia, Canada. The sparsely-populated area includes several protected area designations, including the Long Beach Unit of Pacific Rim National Park and Reserve of Canada, the Province of British Columbia's Tofino Mudflats Wildlife Management Area, and the UNESCO Clayoquot Sound Biosphere Reserve, which protects 366,000 hectares of the west coast of Vancouver Island (Figure 1). The region is part of the temperate rainforest biome with high annual rainfall (3270 mm y^{-1}) and average annual temperature of 9.5

Deleted: Fig. 1

105 °C (Environment Canada, 1981-2010 averages for Tofino). While we were not able to measure the precise tidal range at each location, the mean tidal range in Tofino is approximately 2.7 m (DFO 2022).

These sites are typical of salt marshes along Canada's Pacific coast because they include small, pocket marshes encompassing an enclosed, semi-circular area of coastline as well as larger, estuarine marshes. Unlike geomorphological settings in Atlantic Canada (e.g. van Ardenne et al. 2018), we do not see extensive formation of spits and lagoons; many of our sites were in enclosed bays but were not cut off by spits. All sites were somewhat close to fluvial sources of varying size. Surface water salinity in the surrounding waters ranged from 5.9 at KCS to 24 in Grice Bay, and 29 at Roberts Point six km south of CRF (Postlethwaite et al. 2018).

2.2 Field Sampling

Within each marsh (n = 7), vegetation composition was noted in 50 x 50 cm quadrats before sediment cores (n = 34) were extracted during summer 2016 along linear transects perpendicular to the low tide shoreline, following the methodology of Howard et al. (2014). Coring spots were approximately evenly spaced along the transect (between 9 and 24 meters apart) from land to sea and spanned the low and high marsh zones. Core locations were chosen to avoid ditches and channels without organic soil accumulation; these ditches and channels made up a relatively small portion of total marsh surface area (<5 %).

We note that, within this system, the partitioning between high and low marsh zones appears very closely linked with associated vegetation. Therefore, vegetation composition was recorded as an indicator of 'low' vs 'high' marsh zones (Porter 1982; Weinmann et al. 1984) around each coring spot. Coring spots were considered low marsh if the species *Triglochin maritima*, *Salicornia* spp., *Fucus* spp. or *Ditschilis spicata* were present, and high marsh if it included *Plantago maritima*, *Deschampsia caespitosa*, *Grindelia integrifolia*, *Potentilla anserina*, *Lysimachia maritima*, or *Eleocharis* spp. If a spot contained a mixture of these species, the majority percent cover of high or low marsh species was used to determine whether the spot was high or low marsh. *Carex lyngbyei* were often found throughout both strata and so were not considered unique to one zone. The high marsh species' ranges align approximately with the mean extreme high-water line of estuarine marshes in Clayoquot Sound, while low marsh encompasses elevations between the mean lower high water and the mean extreme high-water lines (Jefferson 1973 as cited in Deur 2000; Weinmann et al. 1984). This method was verified using detrended correspondence analysis, which showed that vegetation assemblages reflected distinct low and high marsh zones (Hill and Gauch 1980; Appendix B). This tight coupling between vegetation type and marsh zone has also been observed in other studies on the west coast of Vancouver Island (Deur et al. 2000) and studies in nearby Boundary Bay (Gailis et al. 2021). Vegetation associations related to salinity and inundation are well documented and commonly used in salt marsh delineation (MacKenzie and Moran, 2004).

Sediment cores were collected using a simple percussion coring technique in which a length of two-inch (57 mm) diameter, PVC vacuum tubing fitted with a plastic core catcher (AMS Inc.) was hammered into the ground until the depth of refusal (DoR). The DoR was reached between depths of 7 and 62 cm. In 19 of the 34 cores, the DoR was reached when the corer penetrated approximately 5-15 cm into layers of sand or gravel at the base of the marshes (see Supplemental Information, Table A2). A similar layer is reached in the eelgrass meadows that form seaward of these marshes (Postlethwaite et al., 2018).

Deleted: T

Deleted: 14

Deleted: Fisheries and Oceans 2016

Deleted: V

Moved down [2]: This method was verified using detrended correspondence analysis, which showed that vegetation assemblages reflected distinct low and high marsh zones (Hill and Gauch 1980; Appendix B). This method was verified using detrended

Moved (insertion) [2]

At these locations, our cores were able to sample the depth of initiation of marsh-related organic C accumulation. At eight of the 34 locations, the cores reached depths where peat mixed substantially with sand or clay; the remaining seven cores ended in peat layers. With three exceptions, the DoR occurred at minimum %C.

At one site (GBK) a steel sledge corer (AMS Inc.) was used to extract four cores, but mechanical problems required switching 150 to the simpler method described above. These four cores penetrated approximately 15-30 cm into the sand/gravel layer below the marsh, which contained minimal amounts of C. Marsh characteristics (e.g., average percent carbon, dry bulk density, soil C density, C stocks, and C accumulation rates) were determined by considering only the upper part of the profile containing peat and overlying sand, gravel, or clay, and these cores were excluded from our average estimates of the DoR.

While in the field, all cores were stored upright in portable coolers with ice packs until their return to the laboratory where 155 they were photographed, logged, and stored long-term under refrigeration at 4°C at the Parks Canada laboratory in Vancouver, British Columbia. Carbon and ²¹⁰Pb lab work was completed 2-4 months after sample collection, between July and October 2016. Additional dating lab work was completed by approximately 30 months after collection.

Use of the percussion corer resulted in sediment compaction during sample collection, which averaged about 20% across all 160 cores (range 0-55%) (Table A1). Nevertheless, we opted to use a percussion corer instead of a gouge corer because the percussion corer had a closed chamber with internal PVC sleeves. Our experience with these sediments has shown that a gouge corer would have been susceptible to disturbance and sediment mixing due to the nature of the open chamber of the corer. We also did not use a Russian corer because compaction would have been similar to what we experienced with the percussion corer, and we did not want to introduce increased contamination through the pivoting nature of the sampling chamber with the Russian corer. Digging pits with a shovel was not an option as this study took place in a national park and biosphere reserve.

165 We note below that correction for compaction was not necessary for estimation of C stocks because the C stocks were estimated directly from sediment cores and not from the overall depth of marsh soils (thus all carbon in the peat layer, regardless of compression, is included in the calculation). However, when we do need to account for compaction (e.g. Figure 3), we use a compaction factor (Howard et al. 2014; Gailis et al. 2021), estimated for each core by dividing the length of core penetration by the length of core recovered (Table A1).

170 2.3 Marsh Area Estimations

ArcMap 10.3 tools were used with 50 x 50 cm resolution aerial orthophotos taken in July 2014 (Government of British 175 Columbia) to estimate the area of high and low marsh zones. The difference between high marsh and low marsh was delineated by eye between darker-green, denser high marsh vegetation and lighter-green, salt-tolerant, and less-dense low marsh vegetation. This method was verified using the detrended correspondence analysis (e.g. Hill and Gauch 1980) of vegetation survey data and was found to accurately categorize 94 % of the cores into the correct marsh zone (see Supplemental Information, Appendix B). Similar approaches in Boundary Bay marsh in BC, Canada have resulted in a high degree of correspondence between marsh color and low vs high marsh delineation (Gailis et al., 2021).

2.4 Soil Carbon Content

Organic C content (%C = g C g⁻¹ marsh sediment) was determined using loss-on-ignition (%LOI) validated with CN Elemental 180 and coulometric analysis (Froelich 1980). LOI analysis was performed on every 1 cm subsample by homogenizing samples with a mortar and pestle, combusting them at 550°C for four hours, weighing, and combusting again at 1000°C for two hours (Heiri et al. 2001). The percentage mass loss-on-ignition (%LOI) was estimated as:

$$\%LOI = \frac{DW_i - DW_f}{DW_i} \times 100 \quad (1)$$

where DW_i is initial dry weight [g] and DW_f is the dry weight [g] after burning. For comparison, the %C was also estimated 185 by measuring total C (%TC) and inorganic C (%IC) on a subset of 93 samples (see supplemental data). %TC was measured on these homogenized subsamples using dry combustion elemental analysis with an Elementar Elemental Analyzer for CN analysis at the University of British Columbia's Department of Earth, Ocean, and Atmospheric Sciences. The same subsamples were then analyzed for %IC using a UIC CM5014 CO₂ coulometer connected to a UIC CM5130 acidification module in the Climate, Oceans, and Paleo-Environments (COPE) laboratory at Simon Fraser University. Measurements of %IC were 190 subtracted from the %TC measurements to estimate %C (Hodgson and Spooner 2016; Hedges and Stern 1984; Schumacher 2002; Howard et al. 2014). Inorganic C was negligible in all 93 of the subsamples analysed (max: 0.015 %) and assumed to be zero for all C calculation purposes. The strong correlation ($R^2 = 0.96$, $p < 0.05$) observed between %LOI and %C for this set of samples allowed us to use this relationship to convert %LOI to %C for all samples using [the following equation, setting any negative %C value resulting from the use of a negative intercept equal to zero](#):

$$195 \quad \%C = 0.44(\%LOI) - 1.80 \quad (2)$$

Soil C density (SCD [g C cm⁻³]) was obtained as the product of the C content and the dry bulk density (DBD [g C cm⁻³]):

$$200 \quad SCD \left(\frac{g C}{cm^3} \right) = \left(\frac{\%C}{100} \right) \times DBD \quad (3)$$

where DBD was estimated after drying the sediment for no less than 72 hours at 60°C. Specifically, DBD was measured by bisecting the core while still inside its casing with a clean, serrated knife to avoid cross-contamination or inaccurate volume 205 measurement that might be caused by extruding the core. Volume was precisely measured immediately afterward using a custom-designed, brass implement designed to remove a single cubic centimeter of soil at a time from the flat face of the bisected core.

2.5 Marsh profiles chronology

The naturally-occurring radionuclide ²¹⁰Pb was analysed in sediment from a subset of eight cores to determine recent C 205 accumulation rates (CARs) (Appleby et al. 1992). One high marsh and low marsh core, each, from CBE, CRF, GBK, and TMF were chosen to represent the two marsh zones in various sites of different sizes. [We used these vegetation communities to establish representativeness of the high and low marsh sites that we sampled for ²¹⁰Pb analysis](#). A set of subsamples (n = 6-7

for cores shorter than 20 cm, n=11-17 for longer cores) was sent to Core Scientific International (Winnipeg, Canada), Flett Research (Winnipeg, Canada), and MyCore Scientific (Dunrobin, Canada) to determine ^{210}Pb content. Activities of ^{210}Pb were determined by α -spectrometry through its granddaughter ^{210}Po , assumed in secular equilibrium. The atmospheric or excess ^{210}Pb ($^{210}\text{Pb}_{\text{ex}}$) fraction used to derive the age-depth model was determined as the difference between the total ^{210}Pb activity and its parent nuclide ^{226}Ra activity (Table A2). ^{226}Ra activities were determined by α -spectrometry at Flett Research using calibrated geometries in a glass vessel, Spectech UCS 30 Alpha Scintillation Spectrometer purged with helium, and sealed for at least 11 days (Mathieu et al. 1988). Samples were sealed and stored for two hours before counting to ensure secular equilibrium of ^{226}Ra daughters. ^{226}Ra was determined through counting ^{222}Rn activity for 60,000 seconds (Minimum Detectable Activity (MDA) = 0.0167 Bq kg $^{-1}$).

Because most of the cores did not show complete excess ^{210}Pb profile (see Results), sediment chronologies were estimated using the Constant Flux: Constant Sedimentation (CF:CS) model (Krishnaswamy et al. 1971; Equation 4), which assumes a constant atmospheric deposition of ^{210}Pb to the marsh surface and a constant mass accumulation rate (MAR, [g cm $^{-2}$ yr $^{-1}$]).

Under these assumptions, the MAR can be obtained from the slope [cm 2 g $^{-1}$] of the linear best-fit line of the relationship between the natural log of the excess ^{210}Pb activity against the cumulative mass [g cm $^{-2}$] using:

$$\text{MAR (g cm}^{-2}\text{yr}^{-1}\text{)} = \frac{0.0311\text{ (yr}^{-1}\text{)}}{\text{slope (cm}^2\text{g}^{-1}\text{)}} \quad (4)$$

where 0.0311 yr $^{-1}$ is the radioactive decay constant of ^{210}Pb .

2.6 Carbon stocks

Organic C stocks [g cm $^{-2}$] are used as a proxy for total organic C storage in marshes and are calculated by integrating all peat layers (n) of the profile [cm]:

$$\text{C stock (g C cm}^{-2}\text{)} = \sum_{i=0}^n \text{SCD}_i \times 1 \text{ cm} \quad (5)$$

As most cores penetrated to a sand or gravel layer found at the base of the marsh (see Sect. 2.2), we are confident that the estimated C stocks (“based of peat C stocks”, hereafter) capture all marsh-associated C emplaced since the formation of the marsh. Because C stocks are estimated directly from sediment cores and not from the overall depth of marsh soils, correction of core compaction was not necessary.

To facilitate comparison with other studies, we also report the mean C stock (Mg C ha $^{-1}$) to a depth of 20 cm for the region (20-cm C stocks, hereafter). Here we estimated the accumulated C to the corrected (uncompacted) depth of 20 cm in each core, with the caveat that 3 of the 34 cores did not reach a depth of 20 cm and therefore were not included in this calculation (Table A1).

Measurements of C stocks based on the accumulation of organic C up to certain depth, however, might give an incorrect impression of C sequestration capacity when comparing sites with different accumulation rates. The dating of the cores, however, allows one to circumvent this problem by normalizing C stocks to a certain age horizon. We therefore used a horizon

Deleted: h

240 of 30 years before time of sampling to calculate these stocks (hereafter 30-year C stock). We chose this age-depth horizon because it represented the oldest ^{210}Pb date shared between the eight dated cores.

Finally, we estimated the total C storage (Mg C), in each marsh and for all seven marshes combined, by multiplying the average C stocks by the aerial extent of each marsh (Table 1). The total C storage was also estimated for low marsh and high marsh areas separately (Mg C). These estimates were made both to the depth of the base of the deepest peat (base of peat) layer and 245 for the 20-cm horizon carbon stock estimates. Note that total C storage was not estimated for the 30-years horizon because we only had ages in four of the seven marshes, and thus there were three marshes where C storage could not be estimated to 30 years.

2.7 Carbon Accumulation Rates

We calculated C accumulation rates (CARs) in two ways: (1) to the “base of peat” depth of the deepest peat-containing layer 250 overlying sand, gravel, or clay, and (2) to the greatest, common age horizon shared by all cores to allow comparisons between cores over equivalent timespans (30 years), which we call the “30-year CAR.”. We include the whole core CAR calculation to assess any potential bias in the results due to higher overall C content in more recent accumulation and for easier comparison with other studies. This organic C fraction is calculated as:

$$\%C = \frac{\sum_{t=0}^{t=i \text{ yrs}} c \text{ (g cm}^2\text{)}}{\sum_{t=0}^{t=i \text{ yrs}} m \text{ (g cm}^2\text{)}} \quad (6)$$

255 where i is either 30 years or the oldest peat-containing layer with detectable excess ^{210}Pb activity for the core. The CAR is the product of this %C and the MAR from Eq. 4:

$$CAR \text{ (g C cm}^2 \text{ yr}^{-1}\text{)} = \%C_{core} \cdot MAR \text{ (g cm}^{-2} \text{ yr}^{-1}\text{)} \quad (7)$$

Average CARs were calculated for the high marsh, the low marsh, and the entire region, using the eight cores with ^{210}Pb dating, both for the whole cores as well as to the 30-year horizon (Table 2). Marsh-wide C accumulation rates *C Accumulation* were 260 calculated by multiplying CAR by the marsh area, for the high marsh, low marsh, and then for the full marsh (Table 2):

$$C \text{ Accumulation (g C yr}^{-1}\text{)} = [CAR_{high \text{ marsh}} \text{ (g C m}^{-2} \text{ yr}^{-1}\text{)} \times area_{high \text{ marsh}} \text{ (m}^2\text{)} + CAR_{low \text{ marsh}} \text{ (g C m}^{-2} \text{ yr}^{-1}\text{)} \times area_{low \text{ marsh}} \text{ (m}^2\text{)}] \quad (8)$$

Marsh-wide annual carbon accumulation was also estimated for the whole study area, by multiplying the average CARs (high marsh, low marsh, and total) by the area of the seven marshes (Table 2).

265 **2.8 Statistical Analysis**

We use a simple t-test to test for significant differences in marsh characteristics (depth of peat base, carbon content, DBD, and SCDs), C stocks, and CARs between high and low marsh zones. We also compare base of peat C stocks, 20-cm C stocks, 30-yr C stocks, as well as for CARs estimated to 30 years and to the base of peat layer.

3 Results

270 **3.1 Soil Properties**

With a few exceptions, marsh soils in Clayoquot Sound consisted of three layers separated by defined horizons: peat, and sand/clay layers (See Supplemental Information, Tables A2 and A3, Fig. A1). In all cores, organic C concentrations were highest in the surface layers (10-45 %) and decreased to lowest values (~ 0 %) in the deepest parts of the cores ([Error! Reference source not found.](#)).

Deleted: topsoil,

275 Depths of refusal in the 34 cores ranged from 7 cm in a low marsh core of CBW to a maximum of 62 cm at CBE. The average (\pm standard error (SE)) corrected depth of the deepest peat layer overlying a sand, gravel, or clay was 29 ± 3 cm for the high marsh sites and 17 ± 3 cm for the low marsh sites, with an overall average base of peat depth of 23 ± 2 cm (Fig. 3a). Differences in the deepest peat layer depths were statistically significant between the high and low marsh zones (p -value < 0.05). We note, however, that the 10 cores which did not penetrate into the sand, clay, or gravel layer were not used in these calculations.

Deleted: Fig. 2

280 While the average base of peat depth of high marsh cores is significantly higher than that of low marsh cores, no significant differences were found for average DBD, average % C, or average SCD ($p > 0.05$) (Fig. 3b-d). Carbon contents averaged $21 \pm 2\%$ in the high marsh and $18.5 \pm 2\%$ in the low marsh cores and $20 \pm 1\%$ overall (Fig. 3b) and did not vary significantly between the high and low marsh zones. DBDs and SCDs also showed similar average values and distributions in the high and low marsh cores (Fig. 3c and 3d). When corrected for compaction, soil carbon densities averaged $0.027 \pm 0.002 \text{ g C cm}^{-3}$ for

285 all sites and ranged from 0.008 to $0.067 \text{ g C cm}^{-3}$.




Figure 2. Percent Carbon (vertical axis) versus depth (horizontal axis) for high marsh (black) and low marsh (grey) cores at all sites. Marsh name abbreviations are defined in Fig. 1. Depths have been corrected for core compaction.

290




Figure 3. Distributions of (a) deepest depth of peat overlying sand, gravel, or clay (cm), (b) average percent carbon, (c) average dry bulk density (corrected for compaction), and (d) average soil carbon density (corrected for compaction). (b), (c), and (d) are estimated to the deepest peat depth for cores in the high marsh zones, low marsh zones, and all cores combined. Horizontal bar is the median; ends of the boxes represent the upper and lower quartiles; whiskers indicate minimum and maximum values while dots indicate outliers. Differences between depths of deepest peat layer for high and low marshes are statistically significant ($t = 2.86$, p -value = 0.0086). Unpaired t-tests show that the high and low marsh values are not statistically different for %C ($t = 1.03$, p -value = 0.31), DBD ($t = 0.25$, p -value = 0.80), and SCD ($t = 0.91$, p -value = 0.37).


3.2 Carbon Stocks and Total Carbon Storage

295 The seven marshes ranged in size from 0.5 to 27 ha, with a total area of 47 ha (Table 1). The high marsh comprised 19-63 % of each individual marsh area and 58 % of the area of all seven marshes combined (Table 1).
 300 The average C stock to the base peat layer for all cores was $67 \pm 9 \text{ Mg C ha}^{-1}$ (mean \pm SE). The base of peat C stock estimates for the high marshes ($80 \pm 14 \text{ Mg C ha}^{-1}$) were 1.5-fold higher than those found in low marsh cores ($52 \pm 8 \text{ Mg C ha}^{-1}$, unpaired t test, $p < 0.05$) (Fig. 4 and 5, Table 1).


305 Similar results were obtained when C stocks were calculated only for the upper 20 cm (Fig. 4, Table 1). The average 20-cm C stock was $54 \pm 5 \text{ Mg C ha}^{-1}$. High marshes averaged $64 \pm 6 \text{ Mg C ha}^{-1}$, which was statistically higher than the low marsh average of $43 \pm 7 \text{ Mg C ha}^{-1}$ ($p < 0.05$). An average of 81 % of the total C stocks in a core accumulates in the top 20 cm of soil. Summing C stocks to the 30-year horizon of the eight dated cores results in an average of $56 \pm 14 \text{ Mg C ha}^{-1}$ (Table 1; Fig. 4). Using the 30-year horizon accentuates the statistically significant difference in C stocks accumulating in the high vs low marsh 310 zones ($p < 0.05$), with high marsh C stocks ($87 \pm 16 \text{ Mg C ha}^{-1}$) up to ~4 times higher than those found in low marshes ($24 \pm 2 \text{ Mg C ha}^{-1}$).

When combined with our estimates of marsh area, we calculate that the total C storage of these seven marshes was $4189 \pm 303 \text{ Mg C}$, 69 % of which was stored in the high marsh (Table 2; [Error! Reference source not found.](#) 5c). For only the top 20 cm, this total is $2764 \pm 149 \text{ Mg C}$, 63 % of which accumulated in the high marsh zones. Finally, over the last 30 years, $2852 \pm 438 \text{ Mg C}$ have accumulated in these marshes, 84% of which accumulated in the high marsh zones (Table 1).

Deleted: Fig.



320 **Figure 4.** Carbon stock estimates for Clayoquot Sound for the high marsh zone (green), low marsh zone (blue), and combined (grey) for all cores. Estimates are made to the base of peat layer, for the surface 20 cm, and for the most recent 30 years of accretion. Note that 30-yr values are based on 4 cores each for high and low marshes (see Methods). Unpaired t-tests show statistically significant differences between the high and low marsh C stocks calculated to the base of peat depth ($t = 2.91$, p -value = 0.0065), 20-cm C stocks ($t = 2.20$, p -value = 0.036) and 30-year C stocks ($t = 3.95$, p -value = 0.0074).



325 **Figure 5.** (a) High and low marsh carbon stocks (Mg C ha^{-1}) measured to the base of peat layer and for the last 30 years; (b) High
326 and low marsh carbon accumulation rates (CARs, $\text{g C m}^{-2} \text{y}^{-1}$) measured to the base of peat layer and for the last 30 years; (c) total
327 marsh carbon (Mg C) estimate to the base of peat layer for high and low marsh zones of each marsh, and (d) marsh wide annual C
328 accumulation (Mg C y^{-1}) estimated to the base of peat layer and for the last 30 years. Paired differences between high and low marsh
329 CARs are statistically significant for CARs calculated to the base of peat layer ($t=4.283$, p -value = 0.0234) and for 30-yr CARs
330 ($t=4.043$, p -value = 0.0272). (Statistical comparisons of C stocks are in Fig. 4).

Table 1. Carbon Stocks

Site	Marsh Area (ha)	Carbon Stock (Mg C ha ⁻¹), Ave ± SE			Total Carbon (Mg C), Ave ± SE	
		To Peat Base	Top 20 cm	30-yr	To Peat Base	Top 20 cm
<i>High Marsh</i>						
CBE	1.42	129 ± 11	74 ± 4	108	183 ± 16	106 ± 6
CBW	0.27	41 ± 13	47 ± 0.4		11 ± 3	13 ± 0.1
CRF	17.31	131 ± 16	70 ± 7	120	2266 ± 283	1214 ± 121
GBK	6.9	51 ± 6	47 ± 9	59	353 ± 41	322 ± 63
KCS	0.47	66	66		31	31
SWC	0.2	88 ± 46	89 ± 45		18 ± 9	18 ± 9
TMF	0.82	54 ± 9	52 ± 9	61.5	44 ± 7	42.5 ± 7.5
Ave ± SE	---	80 ± 14	64 ± 6	87 ± 16	---	---
Sum ± SE	27.4	---	---	---	2906 ± 287	1746 ± 137
<i>Low Marsh</i>						
CBE	2.57	46 ± 16	38 ± 16	22	118 ± 41	99 ± 40
CBW	0.23	51 ± 34	23		12 ± 8	5
CRF	10.11	89 ± 8	66 ± 4	27	899 ± 85	663 ± 43
GBK	4.79	42	42	19	199	199
KCS	0.32	25 ± 2	24 ± 4		8 ± 0.8	7.5 ± 1
SWC	0.83	---	---		---	---
TMF	0.69	59 ± 6	63 ± 6	29	41 ± 4	44 ± 4
Ave ± SE	---	52 ± 8	43 ± 7	24 ± 2	---	---
Sum ± SE	19.5	---	---	---	1283 ± 97	1018 ± 59
<i>Total Marsh</i>						
Ave ± SE	---	67 ± 9	54 ± 5	56 ± 14	---	---
Sum ± SE	46.9	---	---	---	4189 ± 303	2764 ± 149

Base of peat C stock was estimated by summing the mass of carbon from each 1-cm sample interval to the base of each core. Base of peat C stock values are averages of multiple cores, except where only one core was collected (e.g., high marsh KCS and low marsh GBK). The 30-yr high and low marsh average C stocks are based only on CBE, CRF, GBK, and TMF. 30-yr marsh carbon (Mg C) is estimated by multiplying high marsh areas by average high marsh 30-yr C stocks (2384 ± 438 Mg C), low marsh area by average low marsh 30-yr C stock (468 ± 19 Mg C), which is summed to get a total of 2852 ± 438 Mg C.

3.3 Age-depth relationships and mass accumulation rates over time

340 Activities of ^{210}Pb ranged from 2 to 372 Bq kg⁻¹, with maximum activities found at the surface. Activities of ^{226}Ra ranged from 1.6-4.3 Bq kg⁻¹ and were subtracted from the $^{210}\text{Pb}_{\text{total}}$ to obtain the excess ^{210}Pb ($^{210}\text{Pb}_{\text{xs}}$). For most of the cores, the $^{210}\text{Pb}_{\text{sup}}$ level was not reached. Only GBK 1-4 and TMF 1-2 showed a complete $^{210}\text{Pb}_{\text{xs}}$ profile.

Activities of $^{210}\text{Pb}_{\text{xs}}$ showed an almost monotonic decline with depth, suggesting that processes such as bioturbation, mixing, or disconformities have not disturbed the records, and the logarithmic $^{210}\text{Pb}_{\text{xs}}$ profile plotted against the cumulative mass 345 resulted in a linear profile (Fig. 6).

3.4 Mass Accumulation Rates and Carbon Accumulation Rates

The “basal age”, defined as the age at the base of the deepest peat layer, ranged from 13 to 140 years (Table 2). At all sites, the basal ages are consistently older in low marsh cores (106 ± 17 yrs) when compared to high marsh cores (33 ± 9 yrs). As a result, the MARs were an order of magnitude higher in high marsh zones compared to low ones, with average MAR 350 of 3978 ± 1231 and 399 ± 67 $\text{g m}^{-2} \text{ yr}^{-1}$ in the low marsh sediment, respectively, averaging 2188 ± 885 $\text{g m}^{-2} \text{ yr}^{-1}$ (range 262 - 6977 $\text{g m}^{-2} \text{ yr}^{-1}$) for the whole marsh (Table 2).

The average base of peat CAR for the dated cores was 184 ± 50 $\text{g C m}^{-2} \text{ yr}^{-1}$ (range 47 to 386 $\text{g C m}^{-2} \text{ yr}^{-1}$, Table 2). Similar value was obtained by using the 30-year average CARs, with an average of 178 ± 46 $\text{g C m}^{-2} \text{ yr}^{-1}$ (range 59 to 394 $\text{g C m}^{-2} \text{ yr}^{-1}$) (Table 2; Fig. 5b). In the base of peat estimates, CARs in the four high marsh cores (average 303 ± 45 $\text{g C m}^{-2} \text{ yr}^{-1}$) were 355 significantly higher than the low marsh core average (63 ± 7 $\text{g C m}^{-2} \text{ yr}^{-1}$; $p < 0.05$). This difference was also apparent when comparing the 30-year average CARs (280 ± 52 $\text{g C m}^{-2} \text{ yr}^{-1}$ and 76 ± 7 $\text{g C m}^{-2} \text{ yr}^{-1}$ for high and low marsh, respectively, $p < 0.05$).

When we use the high and low marsh areas with the average of the CARs calculated to the base of the peat in each core, we find that the high marsh accumulates 83 ± 12 Mg C yr^{-1} , and the low marsh accumulates 13 ± 1.4 Mg C yr^{-1} (Fig. 5c, Table 2). 360 Combining these estimates produces an annual marsh-wide C accumulation of 96 ± 12 Mg C yr^{-1} , 86% of which is accumulating annually in the high marsh zones. Similar results are obtained when using the 30-yr CARs, with high marshes accumulating 85 % of marsh-wide annual C accumulation (77 ± 14 Mg C yr^{-1}) (Table 2).

Table 2. Core and Marsh Wide Carbon Accumulation Rates

Site	Area (ha)	Carbon Accumulation Rate (CAR, g C m ⁻² y ⁻¹)		Marsh wide Carbon Accumulation ^b (Mg C y ⁻¹), Ave ± SE		Sedimentation Characteristics, Ave ± SE ^c		
		30-yr	Peat Base	30-yr	Peat Base	Basal Age (years before 2016) ^d	LSR (cm yr ⁻¹)	MAR (g cm ⁻² yr ⁻¹)
High Marsh								
CBE 1-1	1.42	342 ± 25	293 ± 21	4.9 ± 0.4	4.2 ± 0.3	54.1 ± 3.9	0.97 ± 0.56	0.156 ± 0.011
CRF 2-1	17.31	394 ± 41	352 ± 37	68.2 ± 7.1	60.9 ± 6.4	26.7 ± 2.8	1.08 ± 0.55	0.242 ± 0.025
GBK 1-2	6.9	189 ± 9	386 ± 18	13.0 ± 0.6	26.6 ± 1.2	13.3 ± 0.6	1.80 ± 1.40	0.697 ± 0.033
TMF 2-1	0.82	193 ± 8	182 ± 8	1.6 ± 0.1	1.5 ± 0.1	38.5 ± 1.7	1.00 ± 0.80	0.496 ± 0.022
Ave ± SE	27.4 ^a	279.5 ± 52	303 ± 45	76.6 ± 14.3	83 ± 12	33.2 ± 8.7	1.21 ± 0.20	0.40 ± 0.12
Low Marsh								
CBE 1-5	2.57	69 ± 21	65 ± 20	1.8 ± 0.5	1.9 ± 0.6	131.4 ± 39.4	0.175 ± 0.060	0.0346 ± 0.0104
CRF 3-2	10.11	87 ± 20	62 ± 14	8.8 ± 2.0	6.3 ± 1.4	139.5 ± 32.5	0.167 ± 0.075	0.0262 ± 0.0061
GBK 1-4	4.79	59 ± 12	47 ± 10	2.8 ± 0.6	2.3 ± 0.5	87.8 ± 18.5	0.133 ± 0.19	0.0408 ± 0.086
TMF 1-2	0.69	88 ± 10	80 ± 9	0.6 ± 0.1	0.6 ± 0.1	66.7 ± 7.6	0.36 ± 0.27	0.0579 ± 0.0066
Ave ± SE	19.5 ^a	76 ± 8.2	63.5 ± 7	14.8 ± 1.4	13 ± 1.4	106.3 ± 17.4	0.21 ± 0.051	0.040 ± 0.0067
Total Marsh								
Ave ± SE	46.9 ^a	178 ± 46	184 ± 50	83 ± 21 ^e	86 ± 23 ^e	69.7 ± 16.5	0.711 ± 0.212	0.219 ± 0.089

^aMarsh areas represent the sums for the high, low, and total marsh.

^bMarsh-wide Carbon Accumulation rates are estimated as the average (high, low, total) CAR multiplied by the marsh area (first column) for all seven marshes.


^c Associated DBDs, %C, and SCD values are in Table A1.

^dBasal Age reflects age determined in deepest sediment layer containing peat overlying sand, clay, or gravel.

^eThe total marsh-wide carbon accumulation values stated here are based on using the average total marsh CAR multiplied by the total marsh area. The values stated in the text are slightly higher because they are based on adding the low marsh and high marsh annual carbon accumulation values together.

365

370



375

Figure 6. Natural log of Excess ^{210}Pb activity versus cumulative mass from the surface. The equation in each graph shows the results from the regression analyses used for the calculation of the mean accumulation rate ($\text{g cm}^{-2} \text{y}^{-1}$) using the CF:CS model. Grey shaded area represents 95% confidence intervals.

4 Discussion

380 4.1 Carbon Stocks

We calculate that the total C storage from salt marshes in Clayoquot Sound is $4189 \pm 303 \text{ Mg C}$, with only 31% accumulated in the low marsh (see Sect. 3.2). Similar, small contributions from the low marsh zone have been found in other estuarine systems (Connor et al. 2001; Adams et al. 2012). In nearby Boundary Bay, BC, Gailis et al. (2021) also found that the low marsh contributed a disproportionately small fraction (15%) to the total marsh-wide C stocks relative to their area. As with 385 Boundary Bay, we found that the small low marsh contribution was driven in part by the smaller proportion of low marsh area

(42% of total) in Clayoquot Sound. Gailis et al. (2021) also attributed lower C stocks in low marsh sediments to vegetation type and associated rooting depths (which are shallower in low marsh *Salicornia* spp). Similar differences in vegetation are observed in the Clayoquot Sound marshes. Furthermore, when normalized to a constant depth horizon (20 cm), the C stocks in Clayoquot Sound low marsh zones are 28% lower than those in the high marsh zones (Fig. 4). However, the main drivers

390 of the smaller C stocks are significantly shallower depths of the base of the peat layer in low marsh relative to the high marsh cores in Clayoquot Sound (Fig. 3a). High marsh peat layers are on average 71% thicker than those found in the low marsh zones, suggesting that low marsh sediments are simply too shallow to accumulate as much C as the high marsh zones.

The C stocks in Clayoquot Sound marshes are less than one-third of the globally averaged estimate of 250 Mg C ha⁻¹ for salt marsh C stocks in the upper 50 cm (Chmura et al. 2003; Pendleton et al. 2012). This is true whether we consider the base of

395 peat, 20-cm, or 30-yr estimates (Table 1). One possible contributor to these discrepancies could be the use of different formulas relating %C to LOI. Chmura et al. (2003) utilized the formula of Craft et al. (1991) to convert LOI measurements to %C. In contrast, we used an empirical relationship based on measurements collected from Clayoquot Sound marsh samples. Comparison of these two regression equations suggest that estimates of %C are very similar for values of %C equal to or less than 30%, but the %C estimates diverge for percentages above 30% (Fig. A3), with calculated differences in %C exceeding 400 20% at %LOIs above 80%. However, we note that the interquartile (Q1:Q3) range of our %C values fall between 4.4 and 28.8%, suggesting that most of our samples have %C values of less than 30%, where the two equations produce similar results. To test the potential impact of the different equations, we compared C stocks (estimated to peat base) that were calculated using the two different %C-LOI relationships for the eight cores that were ²¹⁰Pb dated (Fig. A4). Using the Craft et al. (1991) regression inflates our C stock values by about 30%, but this increase is not sufficiently large to account for the full difference 405 between our C stocks and the global values in Chmura et al. (2003), which are three times greater than our estimated C stocks.

The C stocks in these seven Clayoquot Sound marshes are consistent with regional values found in Boundary Bay, BC (83 ± 30 and 39 ± 24 Mg C ha⁻¹ for high and low marsh, respectively, Gailis et al. 2021), in Stillaguamish, Washington (60 ± 4 Mg C ha⁻¹; Poppe and Rybczyk, 2019), and Snohomish estuary, Washington (72 Mg C ha⁻¹ and 54 Mg C ha⁻¹ for the top 30 cm of two marshes, Crooks et al., 2014). However, these C stocks are an order of magnitude lower than estimates of total soil C mass

410 in high and low marshes along the Pacific coast of the USA (543 ± 47 and 411 ± 70 Mg C ha⁻¹, respectively (Kauffman et al. 2020)).

The lower overall C stocks in these Pacific marshes may in part be attributable to a lower overall C content: the average soil C density of the seven Clayoquot Sound marshes is 0.027 ± 0.002 g C cm⁻³, which is lower than regional estimates for the Oregon Coast (0.034 ± 0.011 g C cm⁻³; Peck et al. 2020) and the global mean of 0.039 ± 0.003 g C cm⁻³ (Chmura et al. 2003).

415 However, low overall C content is not likely to be the main cause of the lower C stocks overall because these SCDs are comparable to the means for the US (0.028 ± 0.013 g C cm⁻³; Holmquist et al. 2018) and Pacific Northwest marshes (0.023 ± 0.002 g C cm⁻³) in which C stocks are an order of magnitude higher than Clayoquot Sound (Kauffman et al. 2020).

A more likely cause is simply the shallow thickness of the Clayoquot Sound marshes, with the depth to the base of the deepest peat layer averaging 23 ± 2 cm (17 cm for low marsh and 29 for high marsh). Similarly shallow depths of refusal (penetrating

Deleted:

into sand or clay layers) have been found in seagrass beds of Clayoquot Sound (Postlethwaite et al. 2018) as well as in Boundary Bay marsh (Gailis et al., 2021). In contrast, a sampling of seagrass and salt marsh systems along the US Pacific Coasts suggest that these ecosystems have substantially thicker soil C depths, with > 85% having mean soil profile depths greater than 2 m (Kauffman et al. 2020). Because data represented in many syntheses come from regions with marshes that 425 have depths of one meter or greater (e.g. Macreadie et al. 2017), extrapolating the results to the total marsh area likely overestimates total C storage. Thus, we recommend that ongoing work to assess blue C contents globally begin to consider shallower depth horizons and/or age horizons as a means for creating a global comparison.

4.2 Carbon Accumulation Rates, Comparisons of Marsh Zones and Pacific Coast Marshes

While marsh C stocks in Clayoquot Sound are substantially lower than the global and Pacific coast averages, the average CARs 430 are actually higher than those found in other parts of the Pacific Coast, especially when we separate measurements by marsh zone (Fig. 7). When all ^{210}Pb -dated sites are combined, we obtain an average CAR for the NE Pacific region of $112 \pm 12 \text{ g C m}^{-2} \text{ yr}^{-1}$ (mean \pm SE, $n=18$). While this value is comparable to the global average CAR of $91 \pm 2.3 \text{ g C m}^{-2} \text{ yr}^{-1}$ (Kennedy et al. 435 2013), both are substantially lower than the regionally averaged CAR of $184 \pm 50 \text{ g C m}^{-2} \text{ yr}^{-1}$ for Clayoquot Sound (Fig. 7c). When we consider only high marsh cores, the average CAR for Clayoquot Sound further increases to $303 \pm 45 \text{ g C m}^{-2} \text{ yr}^{-1}$, 440 substantially larger than the NE Pacific high-marsh average of $145 \pm 26 \text{ g C m}^{-2} \text{ yr}^{-1}$ (Fig. 7a). In fact, the NE Pacific high marsh average is driven substantially by the inclusion of Clayoquot Sound marshes and drops to $96 \pm 12 \text{ g C m}^{-2} \text{ yr}^{-1}$ when Clayoquot Sound marshes are excluded. This comparison suggests that while the overall C storage in Clayoquot Sound marshes is low, the current rates of accumulation are much higher than the regional average, particularly in the high marsh zones. Thus, assuming these rates of C accumulation continue, high marsh zones of Clayoquot Sound have the potential to play a more important role as C sinks in the future.

One effort has been made to assemble a global compilation and synthesis of CARs within salt marsh ecosystems (Ouyang and Lee, 445 2014). Here we compare our results with this compilation with the caveats that (a) Ouyang and Lee (2014) relied heavily on ^{137}Cs dating or marker horizon methods for their estimates, and (b) appears to have some instances of double-counting of sites (e.g., averages of 3 sites were included as a 4th site) and minor issues with attribution of some geographic locations. That said, we note that our new average estimate for NE Pacific CARs of $112 \pm 12 \text{ g C m}^{-2} \text{ yr}^{-1}$ is substantially lower than estimates compiled by Ouyang and Lee (2014), whether considering the global average ($245 \text{ g C m}^{-2} \text{ yr}^{-1}$), the NE Pacific average ($167 \text{ g C m}^{-2} \text{ yr}^{-1}$, $n=6$), the NW Atlantic region ($172 \text{ g C m}^{-2} \text{ yr}^{-1}$; $n=64$; $35.0\text{--}47.4^\circ\text{N}$), or the subset of cores from Atlantic Canada ($188 \text{ g C m}^{-2} \text{ yr}^{-1}$; $n=40$; $43.6\text{--}47.4^\circ\text{N}$). For the NE Pacific region, this compilation predominantly included sites where ^{137}Cs -dating was used to estimate CARs (Ouyang and Lee, 2014). The remaining NE Pacific CAR studies ($n=2$) used marker horizon methods, which, when included, increase the average CAR to $174 \pm 128 \text{ g C m}^{-2} \text{ yr}^{-1}$. Marker horizon and ^{137}Cs dating methods are shown to produce CARs higher than those calculated from ^{210}Pb (Chmura et al. 2003; Callaway et al. 2012). This is due to gradual decomposition in the sediment column, averaging CAR over a shorter time period, and to a shallower depth, oversampling high C soils vis-à-vis lower C soils at depth. As most of the CAR data points in the NW Atlantic region in

Deleted: IPCC
Deleted: (
Deleted:)

Ouyang and Lee's 2014 summary were also calculated using ^{137}Cs dating or marker horizon methods (n=62), these also represent roughly equivalent values to Clayoquot Sound if the observed bias is considered.

Differences between Clayoquot Sound and regionally averaged CARs are not apparent for low marsh zone, for which the 460 Clayoquot Sound average of $63 \pm 7 \text{ g C m}^{-2} \text{ yr}^{-1}$ is statistically similar to the regional estimate of $74 \pm 6 \text{ g C m}^{-2} \text{ yr}^{-1}$ (Fig. 7b). Within the context of Clayoquot Sound, the low marsh zone exhibits substantially lower C stocks and accumulation rates than the high marsh zones. Basal ages (age of the deepest peat layer overlying sand) are substantially older in low marsh sediments when compared to the high marsh, resulting in significantly lower sedimentation rates and CARs.

Our observation of significantly lower CARs in the low marsh compared to the high marsh could result from several factors.

465 As mentioned previously, the vegetation in the low marsh zone is dominated by *Salicornia* spp, whose lower productivity and shallow root systems could be producing and trapping less C (Kelleway et al. 2017; Gailis et al. 2021). A second, potential contributor to slow accumulation rates of C could be the relatively low sediment discharge from terrestrial sources such as the Kennedy River (Nuwer and Kiel, 2005). Previous work in Clayoquot Sound attributed low carbon accumulation rates in eelgrass, in part, to relatively clear waters (secchi disc measurements) and C/N ratios that reflect marine rather than terrestrial 470 sediment contributions (Postlethwaite et al. 2018).

A third important feature of the Clayoquot Sound marshes is that the region is experiencing vertical land motion that counteracts the effects of sea level rise (Mazzotti et al. 2007, 2008; Montillet et al. 2018). Previous work suggests that, in many cases, marshes are responding to rising sea levels by migrating inland (Kirwan et al. 2016). In contrast, in our study region the nearby town of Tofino experiences uplift of 2.86 mm y^{-1} , such that relative sea level is dropping by $1.08\text{--}1.15 \text{ mm y}^{-1}$ (Montillet 475 et al. 2018). In fact, sea level change along the Pacific Coast of Canada where these marshes are developing averages $-0.76 \pm 1.32 \text{ mm/y}$ (James et al. 2014), suggesting that, while some areas along the Pacific Canadian coast are exposed to sea level rise, average conditions for regional salt marshes are likely comparable to what is being experienced in Clayoquot Sound.

Recent work along the Oregon coast suggests that relative sea level rise is a dominant control on vertical accretion rates in tidal marsh systems, and areas experiencing negative sea level rise (or land uplift) have the lowest rates of C accumulation 480 (Peck et al. 2020). The negative sea level rise and relatively low C accumulation rates, particularly in the low marsh settings of Clayoquot Sound and Pacific Rim National Park, are consistent with this result.

A final discriminating characteristic between the high and low marsh cores in Clayoquot Sound are the ^{210}Pb inventories (Fig. 485 A5), which suggest that the high and low marsh zones are subjected to different erosional processes. Given that these cores all come from the same area one would assume that all the cores should have similar inputs of ^{210}Pb (Olid et al. 2010; Arias-Ortiz et al. 2018). However, a comparison of the ^{210}Pb inventories shows that the high marshes accumulate much more ^{210}Pb than the low marshes. This difference, together with the lower concentrations of the ^{210}Pb in the upper layers of the low marsh cores, could result from focusing /soil formation causing greater ^{210}Pb accumulation in the high marshes and/or reduced accumulation in the low marshes due to erosion and winnowing processes. One possible explanation here is that the extensive, shallow flats in the low marsh regions of Clayoquot Sound are subjected to more intensive tidal erosion and therefore removal of soil organic


Deleted:

Deleted: A3

Deleted: humus

Deleted: humus

material, which results in lower accumulation rates relative to the high marsh zones. Further studies are needed to estimate the
495 significance of these drivers within the region.



4

Figure 7. Comparison of Clayoquot Sound CARs with other salt marshes with ^{210}Pb dating, for whole core (black circles) and 30-year (red triangle) estimates. Individual Clayoquot Sound marshes are indicated with open symbols followed by a regional mean (\pm SE). Additional site data taken from: [1] Boundary Bay, BC, Canada (Gailis et al., 2021); [2] Stillaguamish, WA, USA (Poppe and

500 Rybczyk, 2019); [3] Snohomish Estuary, WA, USA (Crooks et al., 2014); [4] Animal Slough (AS) in southern Puget Sound, WA, USA (Note that red triangle is 50-yr average; Drexler et al., 2019); [5] coastal OR, USA (Peck et al., 2020); [6] San Francisco Bay, CA, USA (Callaway et al., 2012). Global and other regional averages are taken from [7] Kennedy et al. (2013) and [8] Ouyang and Lee (2014); only a fraction of data (<10%) in [8] were dated using ^{210}Pb . Horizontal line represents Kennedy et al. (2013) global average.

Deleted: IPCC

Deleted: IPCC

505 3. Comparison of Whole-Core and 30-year CARs

Estimating CARs to a consistent age horizon of 30 years allowed us to compare C accumulation between marshes and marsh zones, but it also reveals interesting information about differences in CAR depending on the approach used to estimate these values. Previous comparisons in nearby Boundary Bay marsh showed that CARs were 20% higher during the last 30 years when compared to the whole core estimates (Gailis et al. 2021). Because %C decreases with depth in all 34 cores from 510 Clayoquot Sound, we might anticipate that CARs for the most recent years will always result in a higher CAR than a measurement of C stock to the depth of minimum ^{210}Pb detectable activity. This can be partially attributed to remineralization of soil C over time, which would decrease overall %C in the deeper portions of the sediment cores (Callaway et al. 2012; Johannessen and MacDonald 2016). In Clayoquot Sound, however, the CARs estimated for the last 30 years were not substantially different from our estimates made to the base of the peat layer (Table 2), particularly in the high marsh zone 515 where the average age for the base of the peat layer was 33 ± 9 years. Slightly larger differences were seen in the slowly-accumulating low marsh zones, where the 30-year CAR was about 15% higher. Overall, the young ages of these marshes result in small differences between the two estimates.

The examination of 30-year CARs as well as CARs in relatively young marshes could have interesting implications for future blue C assessments. Many marsh restoration projects in which C is being measured are currently on time scales of 10 to 30 520 years past the initiation of restoration (e.g. Abbott et al. 2019; Drexler et al. 2019). Vegetational succession and C storage in a restored tidal marsh can take over 10 years (Havens et al., 2002; Craft et al. 2003), and even a 30-year monitoring period would capture a period of gradually improving C sequestration potential in restored tidal marsh soils (e.g. Howe et al. 2009). However, the inflation of CARs in younger sediments in the Clayoquot Sound and Boundary Bay sediments could provide a caution that 525 CARs measured from relatively young marker horizons are at risk of overestimating total C in sediments, especially if these CARs are used to project C sequestration over longer time periods. Several recent studies have shown substantially higher CARs in restored vs natural marshes after approximately 10 years (e.g., Poppe and Rybczyk, 2019; Drexler et al. 2019). The causes of these inflated CARs can be very site-specific, depending on the types of colonizing vegetation (Kelleway et al. 2017) and the relative contributions of allochthonous mineral materials (Drexler et al., 2019). In the case of Clayoquot Sound, it is possible that these high CARs will be sustained, making this marsh extremely important as a regional C sink. However, we 530 also cannot rule out the possibility that longer-term processes, such as remineralization and sediment redistribution, could reduce the actual amount of C stored as the marshes age, such that the accumulation rates become more comparable to those found on other parts of the Pacific coast (Fig. 7).

535 4.4 Comparison with carbon storage in boreal forest ecosystems

The relevance of blue C storage potential to climate change mitigation depends on the scale over which it is considered. In the literature, the carbon accumulation potential of blue carbon is often compared with terrestrial ecosystems, and it is often stated that, per unit area, blue C ecosystems accumulate carbon at higher estimated rates than terrestrial forest soils (McLeod et al. 2011).

540 A first-order comparison for Canadian ecosystems shows the same pattern of higher carbon uptake rates per unit area in tidal wetlands when compared with measurements of net ecosystem productivity of terrestrial forests. Forested areas in Canada have been estimated to take up carbon at rates ranging from $35 \text{ g C m}^{-2} \text{ yr}^{-1}$ (Canada-wide estimate, Stinson et al., 2011) to $63 \text{ g C m}^{-2} \text{ yr}^{-1}$ (British Columbia, Peng et al., 2014). The 30-year CAR for salt marshes in Clayoquot Sound are approximately 2-7 times higher, averaging $178 \pm 46 \text{ g C m}^{-2} \text{ yr}^{-1}$. Note that we considered forests' net ecosystem productivity, which is higher

545 than C uptake rates of $4.6 \pm 2.1 \text{ g C m}^{-2} \text{ yr}^{-1}$ for boreal forest soils cited by McLeod et al. (2011). Consideration of the areal extent of each of these ecosystems is extremely important for contextualizing blue C uptake among Canadian ecosystems. Estimates of total salt marsh area in Canada range from 44,000 ha (Bridgman et al., 2006) to 111,274 ha (Mcown et al., 2017). In contrast, the boreal forest ecosystem is one of Canada's largest terrestrial biomes and encompasses approximately 270 million ha (Kurz et al. 2013). Our Clayoquot Sound data represent only a small area of a single region of

550 the west coast, but if we assume our CAR estimate of $184 \pm 50 \text{ g C m}^{-2} \text{ yr}^{-1}$ from Clayoquot Sound approximates the average for all tidal salt marshes in Canada, Canada's marshes would accumulate somewhere between 81,000 and 205,000 Mg C yr⁻¹. This annual carbon accumulation from Canadian salt marshes is between 0.4 and 2.9 % of the carbon estimated to accumulate annually by Canadian boreal forests ($6,750,000\text{--}18,090,000 \text{ Mg C yr}^{-1}$).

555 Our back-of-the-envelope comparison is not intended to provide a formal estimate of carbon sequestration in Canadian salt marshes but instead highlights several important assumptions and uncertainties that are important to large-scale assessments of blue carbon sequestration. The first is that regional-to-global estimates of blue C accumulation rates depend on regionally specific measurements of salt marsh C stocks and accumulation rates, and extrapolation of blue C accumulation rates based on globally derived averages (or averages from regions other than the one in question) can lead to large uncertainties in blue C accumulation rates. Our calculation above assumes that the CAR estimates from Clayoquot Sound are representative of the

560 pan-Canadian average for CARs in salt marshes. Using the published value for Atlantic Canada ($188 \text{ g C m}^{-2} \text{ yr}^{-1}$, Ouyang and Lee, 2014) would not substantially change our calculation. However, if the average pan-Canadian CAR is closer to the global estimate of $91 \pm 2.3 \text{ g C m}^{-2} \text{ yr}^{-1}$ (Kennedy et al., 2013), our estimate of annual carbon accumulation would halve. Thus, a pan-Canadian estimate of CAR (and its associated uncertainty) would be useful for improving the estimated carbon sequestration rates in Canadian salt marshes.

565 Second, our comparison with C sequestration in Canadian forests is based on net ecosystem productivity estimates from 2011 and 2013. With climate change, the increased prevalence of natural disturbances such as forest fires and insect outbreaks is projected to result in episodic losses of carbon from the above-ground and soil carbon pools (e.g., Stinson et al., 2011, Walker

Deleted: are

Deleted: ,

Deleted: However, c

Moved (insertion) [1]

Deleted: which will be affected by several processes; and (b) large uncertainties associated with the areal extent of salt marshes is 1,207 ha (CEC, 2016a) field research in Boundary Bay suggests that actual marsh area is only one-quarter the size at 275 ha (Gailis et al., 2021). This suggests that (a) estimated annual carbon accumulation for salt marshes in Canada.

Deleted: With

Deleted: would

Deleted: the

580 *et al., 2019). Between 1990-2008, 24 ± 19 Tg C yr⁻¹ was removed from the forest carbon pool by fire (Kurz et al., 2013), with British Columbia forests becoming a sustained source of CO₂ since 2003 (e.g. Giles-Hansen and Wei, 2022). The degree to which climate change will impact forest carbon sequestration in future will depend on several factors, including increases in disturbance frequency and intensity (Walker et al. 2019), subsequent recovery and shifts in forest species dominance (e.g. Mack et al. 2021), as well as management practices (Price et al. 2013). At the same time, as sea level rise is currently a minimal threat to the Canadian Pacific salt marshes, they are likely to continue to function as efficient C sinks despite global warming.*

585 *Finally, in addition to the regionally-specific nature of processes controlling blue C accumulation, we note that estimation of the areal extent of these marshes varies by an order of magnitude within Canada alone, and determinations of the relative proportions and high-to-low marsh settings have not been quantified. We note, for example, that while large-scale databases suggest that the areal extent of salt marsh in Boundary Bay, BC, is 1,207 ha (CEC, 2016a), our field research in Boundary Bay*

590 *has determined that the actual marsh area is only one-quarter the size at 275 ha (Gailis et al., 2021). This suggests that estimation of blue C potential in policy-specific contexts requires a more detailed determination of the areal extent of these marshes, especially when considering provincial-to-national scales. We further suggest that the small area over which blue C accumulates likely influences the scale over which it is best considered for natural land carbon sequestration and accounting. Blue C accumulation occurs in only 0.016-0.1 % the land area of boreal forests nationwide. On a global, national, and even*

595 *provincial scale, total blue C sequestration in salt marshes is relatively small compared to the large areal expanses of forest. However, the very high, per-unit area sequestration of carbon in salt marsh ecosystems can increase its importance at local-to-regional scales and should be considered as an important co-benefit to restoration activities and conservation for other purposes.*

4.5 Future Work

600 While this study has provided a useful first step in quantifying blue C stocks, storage, and accumulation rates, we note the following considerations in estimating the potential of blue C within this region. First, our study has estimated blue C in 47 ha of high and low marsh areas of Clayoquot Sound. We expect that these mesotidal estuarine marshes, often constrained in size by surrounding topography, are typical of the marshes found on the Pacific coast of British Columbia, the total estimated area of which is 60 square kilometres, or 6000 ha (Ryder et al. 2007). However, given variability in riverine discharge and tectonic

605 settings along the coast, we anticipate that additional measurements (additional marshes, with differing riverine and sedimentary inputs, and variable uplift setting) will be necessary to understand the variability in carbon accumulation along the Pacific Coast of Canada. Second, our study has focused on estimation of carbon stocks and accumulation rates within these marshes, and full estimation of blue C potential will ultimately require estimation of greenhouse gas fluxes of carbon dioxide, methane, and nitrous oxide within these marshes (e.g. Chmura et al. 2011). The mesotidal nature of some of these marsh

610 locations could mean that some of these marshes emit substantial contributions of methane, which may counter their effects as C sinks (e.g. Abdul-Aziz et al. 2018; Huang et al. 2019; Huertas et al. 2019; Li et al. 2021; Wei et al. 2020). Finally, this study has revealed the importance of classifying high vs low marsh zones to best quantify carbon stocks and accumulation

Deleted: B

Deleted: , an amount that may increase substantially later in the 21st century as climate change increases fire frequency.

Deleted:

Deleted: marshes

Moved up [1]: However, even with this large range, this annual carbon accumulation from Canadian salt marshes is between 0.4 and 2.9 % of the carbon estimated to accumulate annually by Canadian boreal forests (6,750,000-18,090,000 Mg C yr⁻¹). ¶

Deleted: This rough calculation reveals three important points. The first is that regional-to-global estimates of blue C accumulation rates depend on regionally specific measurements of salt marsh C stocks and accumulation rates. Extrapolation of blue C accumulation rates based on globally derived averages (or averages from regions other than the one in question) can lead to large uncertainties in blue C accumulation rates. Second, in addition to the regionally-specific nature of processes controlling blue C accumulation, we note that estimation of the areal extent of these marshes varies by an order of magnitude within Canada alone, and determinations of the relative proportions and high-to-low marsh settings have not been quantified. Estimation of blue C potential in policy-specific contexts requires a more detailed determination of the areal extent of these marshes, especially when considering provincial-to-national scales.

Deleted: Finally, w

Deleted: posit

Deleted: Poffenbarger et al. 2011

rates, as the different vegetation growth and sedimentary processes operating in these zones can critically influence CARs, and
640 therefore estimates of C sequestration. Future work thus needs to involve better quantification of the areal extent and
classification of salt marshes, both within Clayoquot Sound, along the Pacific coast, and in Canada as a whole.

5 Conclusion

Our work helps address the data gap for North American blue C by producing CAR estimates using ^{210}Pb -derived mass
accumulation rates of salt marsh sediments, showing CAR comparable to other salt marsh locations in the Northeast Pacific
645 coast. Importantly, this work demonstrates that the marshes of Clayoquot Sound along the Pacific Coast of Canada exhibit
substantially smaller C stocks when compared with other regional salt marshes to the south, along the Pacific Coast of the
United States. We attribute these lower stocks to substantially shallower (< 50 cm) depths of marsh initiation in all high and
low marsh settings. At the same time, while carbon stocks are lower than regional and global averages, the rates of carbon
accumulation in these marshes are comparable and even higher than regional estimates. We show that both 30-year CARs and
650 CARs estimated to the peat base are substantially greater than regional averages generated using only ^{210}Pb -derived mass
accumulation rates. However, significant variability exists between marshes, and different radioisotope dating methods in CAR
measurement create uncertainty. Our results support existing findings that blue C soils accumulate carbon at high rates, limited
only by the spatial extent of such ecosystems. These results provide a natural analogue for monitoring carbon accretion over
timescales relevant for restoration activities.

Table A1. Core Locations, Soil Carbon Characteristics, and Carbon Stocks

Core ID	Lat. (49. $_{-}^{+}$ N)	Long. (125. $_{-}^{+}$ W)	Marsh Zone	Compacted (Complete) Core Length (cm)	*Average Uncorrected DBD \pm SE to Peat Base (g cm $^{-3}$)	Average %C \pm SE To Peat Base	*Average SCD \pm SE to Peat Base (g C cm $^{-3}$)	C Stock to Peat Base (Mg C ha $^{-1}$)	**C Stock to 20 cm (Mg C ha $^{-1}$)	30-yr C Stock (Mg C ha $^{-1}$)
<i>Cannery Bay East</i>										
CBE 1-1	0.14139	0.6662	high	47 (62)	0.320 \pm 0.037	23.4 \pm 1.8	0.031 \pm 0.0013	158.7	62.9	107.7
CBE 1-2	0.14142	0.66629	high	38 (40)	0.309 \pm 0.030	17.0 \pm 1.9	0.032 \pm 0.0019	129.9	78.0	
CBE 1-3	0.14147	0.66636	high	24 (24)	0.148 \pm 0.012	28.3 \pm 1.6	0.038 \pm 0.0008	90.3	79.2	
CBE 1-4	0.14152	0.66639	low	14 (14)	0.154 \pm 0.010	23.5 \pm 1.5	0.035 \pm 0.0018	48.4	-	
CBE 1-5	0.14155	0.66644	low	20 (30)	0.226 \pm 0.021	18.7 \pm 1.4	0.029 \pm 0.004	85.7	43.6	21.9
CBE 2-2	0.1414	0.66618	high	30 (30)	0.16 \pm 0.014	30.3 \pm 1.0	0.045 \pm 0.0020	133.8	86.7	
CBE 2-3	0.14142	0.66614	high	40 (40)	0.119 \pm 0.006	28.4 \pm 1.0	0.032 \pm 0.0006	132.1	65.7	
CBE 2-4	0.14144	0.66609	low	20 (30)	0.179 \pm 0.024	28.6 \pm 2.0	0.033 \pm 0.0036	39.8	62.5	
CBE 2-5	0.14151	0.66602	low	30 (30)	0.767 \pm 0.056	1.9 \pm 0.5	0.013 \pm 0.0029	9.0	9.2	
<i>Cannery Bay West</i>										
CBW 1	0.14115	0.66983	high	16 (20)	0.158 \pm 0.017	20.7 \pm 0.6	0.026 \pm 0.0017	28.7	46.7	
CBW 2	0.14115	0.66982	low	16 (23)	0.266 \pm 0.061	15.9 \pm 3.3	0.025 \pm 0.0029	17.8	23.5	
CBW 3	0.14113	0.66962	low	7 (7)	0.165 \pm 0.004	32.3 \pm 1.8	0.053 \pm 0.0031	84.9	-	
CBW 4	0.14112	0.66955	high	24 (30)	0.157 \pm 0.023	17.8 \pm 1.7	0.019 \pm 0.0020	53.6	47.4	
<i>Cypress River Flats</i>										
CRF 1-1	0.27905	0.90754	high	38 (46)	0.168 \pm 0.008	26.9 \pm 1.1	0.036 \pm 0.0013	163.2	81.7	
CRF 1-2	0.27896	0.90758	low	16 (30)	0.168 \pm 0.009	28.7 \pm 1.0	0.025 \pm 0.0013	76.1	57.8	
CRF 2-1	0.27935	0.90932	high	37 (51)	0.207 \pm 0.021	22.5 \pm 1.5	0.028 \pm 0.0014	118.7	57.5	120.3
CRF 2-2	0.27916	0.90926	low	26 (38)	0.205 \pm 0.025	25.3 \pm 1.9	0.028 \pm 0.0009	104.8	66.7	
CRF 3-1	0.2789	0.911	high	32 (40)	0.225 \pm 0.045	27.1 \pm 2.3	0.032 \pm 0.0016	110.6	105.9	

Core ID	Lat. (49. ⁻ N)	Long. (125. ⁻ W)	Marsh Zone	Compacted (Complete) Core Length (cm)	*Average Uncorrected DBD \pm SE to Peat Base (g cm ⁻³)	Average %C \pm SE To Peat Base	*Average Corrected SCD \pm SE to Peat Base (g C cm ⁻³)	C Stock to Peat Base (Mg C ha ⁻¹)	**C Stock to 20 cm (Mg C ha ⁻¹)	30-yr C Stock (Mg C ha ⁻¹)
CRF 3-2	0.27882	0.91087	low	23 (30)	0.192 \pm 0.023	27.4 \pm 2.3	0.035 \pm 0.0021	85.8	94.8	27.2
<i>Grice Bay – Kootowis Creek</i>										
GBK 1-1	0.08754	0.73238	high	16 (31)***	0.232 \pm 0.032	18.3 \pm 2.3	0.016 \pm 0.0012	41.0	35.2	
GBK 1-2	0.08756	0.73261	high	19 (32)***	0.440 \pm 0.060	10.0 \pm 1.8	0.015 \pm 0.0016	51.2	40.0	59.2
GBK 1-3	0.08763	0.73271	high	23 (26)***	0.257 \pm 0.027	15.7 \pm 1.4	0.032 \pm 0.0028	60.9	64.5	
GBK 1-4	0.08771	0.73283	low	13 (20)***	0.298 \pm 0.017	12.0 \pm 0.8	0.023 \pm 0.0009	41.6	41.6	19.1
<i>Kennedy Cove South</i>										
KCS 1-1	0.13696	0.67082	high	24 (31)	0.227 \pm 0.041	21.4 \pm 2.1	0.030 \pm 0.0045	66.2	66.2	
KCS 1-2	0.13707	0.67085	low	14 (21)	0.248 \pm 0.074	15.0 \pm 6.9	0.018 \pm 0.0012	27.4	-	
KCS 1-3	0.13714	0.67093	low	16 (36)	0.427 \pm 0.097	11.1 \pm 9.5	0.009 \pm 0.0016	25.6	24.0	
KCS 1-4	0.13719	0.67096	low	10 (20)	0.296 \pm 0.085	14.7 \pm 2.2	0.014 \pm 0.0012	28.9	28.9	
KCS 1-5	0.1372	0.67107	low	10 (12)	0.509 \pm 0.125	7.6 \pm 2.1	0.015 \pm 0.0029	17.8	17.8	
<i>Shipwreck Cove</i>										
SWC 1-1	0.12995	0.69943	high	29 (29)	0.248 \pm 0.093	27.6 \pm 3.4	0.033 \pm 0.0030	42.3	43.9	
SWC 2-1	0.13014	0.69908	high	18 (20)	0.541 \pm 0.124	21.0 \pm 2.6	0.067 \pm 0.0093	133.4	133.4	
<i>Tofino Mud Flats</i>										
TMF 1-1	0.13014	0.88689	high	27 (45)	0.326 \pm 0.068	18.6 \pm 3.4	0.019 \pm 0.0012	45.4	42.8	
TMF 1-2	0.1302	0.88688	low	26 (30)	0.241 \pm 0.008	14.0 \pm 0.8	0.038 \pm 0.0012	53.0	57.5	28.9
TMF 2-1	0.12989	0.88661	high	28 (31)	0.751 \pm 0.066	5.6 \pm 1.5	0.022 \pm 0.0041	62.4	61.0	61.5
TMF 2-2	0.13017	0.88665	low	27 (30)	0.231 \pm 0.010	17.2 \pm 0.7	0.035 \pm 0.0010	65.9	69.5	

*Ave DBD values were not corrected for compaction (see Table A3 for corrected DBDs), but SCD values have been corrected for compaction.

** Cores that do not extend to 20 cm and do not penetrate a sand/gravel/clay layer at the base were excluded from the 20 cm calculation.

***GBK cores were extracted using a steel sledge corer that penetrated 15-30 cm into the sand and clay layers. The total core lengths were 46 (88), 60 (100), 59 (66) and 24 (37) cm for GBK 1-1, 1-2, 1-3, and 1-4, respectively. Depths provided here are just below the transition from peat to sand, two cm into the minimal %C layer.

Table A2. Downcore distribution of ^{210}Pb and ^{226}Ra in eight cores used for ^{210}Pb dating.

Upper Depth (cm)	Sediment Type	Supported ^{210}Pb (^{226}Ra , Bq/kg)	^{210}Pb total (Bq/kg)	+/- 1 SD (Bq/kg)	$^{210}\text{Pb}_{\text{exs}}$ (Bq/kg)	+/- 1 SD (Bq/kg)	Slope (cm 2 /g)	Age (yr) before 2016	Age SD (yr)
CBE 1-1 (HM)^a									
0	humus	2.03	196.2	7.2	194.2	7.2	0.20 ± 0.014	0.51	0.04
4	humus	2.03	187.6	7.1	185.6	7.1		2.70	0.19
5	humus	2.03	245.7	9.4	243.7	9.4		3.27	0.24
6	humus	2.03	172.5	6.8	170.5	6.8		3.79	0.27
7	peat	2.03	158.5	6.5	156.5	6.5		4.43	0.32
8	peat	2.03	149.1	6.3	147.0	6.3		5.14	0.37
10	peat	2.03	137.4	6.0	135.3	6.0		6.68	0.48
11	peat	2.03	124.2	5.5	122.2	5.5		7.38	0.53
13	peat	2.03	112.7	5.2	110.7	5.3		9.12	0.66
16	peat	2.03	102.5	4.8	100.4	4.8		12.33	0.89
21	peat	2.03	86.2	4.5	84.2	4.5		21.57	1.55
24	peat	2.03	76.3	4.1	74.3	4.1		29.21	2.10
25*	peat	2.03					31.5	2.27	
27	peat	2.03	62.0	3.8	59.9	3.8		35.31	2.54
32	peat	2.03	53.5	3.6	51.4	3.6		44.37	3.19
34	peat	2.03	46.6	3.2	44.6	3.2		47.51	3.42
37	peat	2.03	36.7	2.8	34.7	2.8		51.56	3.71
38**	peat	2.03					54.1	3.90	
42	sand	2.03	28.1	2.6	26.0	2.7			
46	sand	2.03	24.6	0.5	22.6	0.5			
14 (^{226}Ra)			1.90		0.55				
44 (^{226}Ra)			2.16		0.46				
CBE 1-5 (LM)^b									
0	humus+peat	1.74	374.0	9.5	372.2	9.5	0.90 ± 0.30	4.04	1.21
3	humus+peat	1.74	322.3	9.8	320.5	9.9		16.46	4.93
6*	humus+peat	1.74	83.9	5.1	82.2	5.1		31.77	9.52
10	humus+peat	1.74	19.2	2.0	17.4	2.1		58.04	17.39
14	humus+peat	1.74	12.7	1.9	11.0	2.0		79.12	23.70
18	humus+peat	1.74	20.7	2.2	19.0	2.2		118.11	35.38

Upper Depth (cm)	Sediment Type	Supported					Age (yr)	
		^{210}Pb (^{226}Ra , Bq/kg)	^{210}Pb total (Bq/kg)	+/- 1 SD (Bq/kg)	$^{210}\text{Pb}_{\text{exs}}$ (Bq/kq)	+/- 1 SD (Bq/kg)	Slope (cm^2/g)	before 2016
19**	humus+peat	1.74					131.39	39.36
6 (^{226}Ra)		1.38	0.94					
18 (^{226}Ra)		2.10	0.46					
CRF 2-1 (HM)^a								
0	humus	1.58	225.2	17.5	223.6	17.5	0.13±0.013	0.33
4	humus	1.58	194.2	16.4	192.6	16.4		1.86
8	humus	1.58	169.3	15.4	167.7	15.4		3.44
13	peat	1.58	149.3	14.5	147.8	14.5		7.20
16	peat	1.58	129.9	13.7	128.4	13.8		10.23
19	peat	1.58	114.1	12.9	112.5	13.0		13.58
22	peat	1.58	101.1	12.0	99.5	12.0		16.27
25	peat	1.58	91.8	11.4	90.2	11.5		18.84
28	peat	1.58	80.0	11.1	78.4	11.1		22.36
30**	sand+peat						26.7	2.79
31*	sand	1.58	72.1	10.2	70.5	10.3	30.52	3.18
34	sand	1.58	61.6	9.7	60.1	9.8		
CRF 3-2 (LM)^b								
1	humus+peat	1.58	326.6	14.3	325.0	14.4	1.19±0.28	9.93
4	peat	1.58	212.6	7.1	211.0	7.2		27.89
5*	peat						31.3	7.30
8	peat	1.58	182.8	7.5	181.2	7.6		44.70
11	peat	1.58	130.5	5.6	129.0	5.7		61.90
14	peat	1.58	27.0	2.5	25.5	2.8		85.20
18**	peat						139.5	32.49
20	peat+sand	1.58	2.2	1.2				
8 (^{226}Ra)		2.50	1.17					
14 (^{226}Ra)		0.67	0.67					
GBK 1-2 (HM)^a								
0	humus	2.63	191.6	7.1	189.0	7.1	0.045±0.0021	0.24
4	humus+peat	2.63	183.0	7.0	180.3	7.0		1.20
								0.06

Upper Depth (cm)	Sediment Type	Supported					Age (yr)	
		^{210}Pb (^{226}Ra , Bq/kg)	^{210}Pb total (Bq/kg)	+/- 1 SD (Bq/kg)	$^{210}\text{Pb}_{\text{exs}}$ (Bq/kq)	+/- 1 SD (Bq/kg)	Slope (cm ² /g)	before 2016
5	humus+peat	2.63	175.1	6.8	172.5	6.8		1.48 0.07
7	humus+peat	2.63	169.7	6.7	167.1	6.7		1.99 0.09
9	peat	2.63	157.6	6.5	155.0	6.5		3.00 0.14
12	peat	2.63	145.0	6.2	142.4	6.2		5.03 0.24
16	peat+sand	2.63	131.8	5.9	129.2	5.9		8.11 0.39
19	peat+sand	2.63	117.8	5.2	115.2	5.2		11.90 0.57
20**	peat+sand						13.26	0.63
22	sand	2.63	101.4	4.9	98.7	4.9		16.08 0.76
27	sand	2.63	89.3	4.5	86.6	4.5		23.50 1.12
33*	sand+clay	2.63	75.9	3.9	73.3	3.9		31.26 1.49
37	clay	2.63	57.4	0.5	54.8	0.5		36.58 1.74
GBK 1-4 (LM) ^c								
0	peat	2.63	244.9	5.9	242.3	5.9	0.76 ± 0.16	6.13 1.29
2	peat	2.63	221.8	6.0	219.1	6.0		19.37 4.08
4*	peat	2.63	175.5	6.1	172.9	6.1		32.11 6.76
8	peat	2.63	49.3	2.4	46.6	2.4		61.29 12.90
10	peat-sand	2.63	7.0	0.8	4.3	0.8		77.71 16.36
11**	peat+sand						87.76	18.47
12	Sand	2.63	8.4	0.8	5.8	0.9		
14	Sand	2.63	6.8	0.7	4.2	0.7		
16	Sand	2.63	6.5	0.7	3.8	0.7		
18	Sand	2.63	5.4	0.7	2.8	0.7		
20	Sand	2.63	7.4	0.8	4.8	0.8		
22	Sand	2.63	116.0	4.3	113.4	4.3		
23	Sand	2.63	7.9	1.0	5.3	1.0		
3 (^{226}Ra)			2.58		0.38			
20 (^{226}Ra)			2.68		0.38			

TMF 1-2 (LM) ^b

1	humus+peat	4.33	269.3	6.9	264.9	6.9	0.54 ± 0.061	6.91 0.79
3	peat+mud	4.33	173.9	6.4	169.5	6.4		14.51 1.65
5	peat+mud	4.33	184.0	5.9	179.7	5.9		23.66 2.69

Upper Depth (cm)	Sediment Type	Supported					Age (yr)	
		^{210}Pb (^{226}Ra , Bq/kg)	^{210}Pb total (Bq/kg)	+/- 1 SD (Bq/kg)	$^{210}\text{Pb}_{\text{exs}}$ (Bq/kq)	+/- 1 SD (Bq/kg)	Slope (cm^2/g)	before 2016
7*	peat+mud							33.0 3.76
8	peat+mud	4.33	86.6	4.3	82.2	4.3		36.78 4.19
11	peat+mud	4.33	65.9	3.9	61.6	3.9		49.22 5.61
14	peat+mud	4.33	54.0	3.2	49.7	3.3		61.82 7.04
15**	peat+mud							66.7 7.59
18	sand	4.33	7.1	1.5	2.8	1.5		
18								
<i>(^{226}Ra)</i>		4.33	0.33					

TMF 2-1 (HM)^b

0	humus	3.50	194.4	7.2	190.9	7.2	0.063 \pm	
4	peat+mud	3.50	184.9	7.0	181.4	7.0		0.0027 0.34 0.01
5	peat+mud	3.50	177.8	6.9	174.3	6.9		2.92 0.13
6	peat+mud	3.50	173.1	6.8	169.6	6.8		4.19 0.18
7	peat+mud	3.50	159.9	6.5	156.4	6.5		5.40 0.24
8	peat+mud	3.50	148.2	6.3	144.7	6.3		7.04 0.31
10	peat+mud	3.50	133.5	6.0	130.0	6.0		8.71 0.38
11	sand+peat	3.50	120.9	5.4	117.4	5.4		11.29 0.49
13	sand+peat	3.50	109.3	5.1	105.8	5.1		12.76 0.56
16	sand+peat	3.50	98.1	4.6	94.6	4.7		15.65 0.68
21*	sand+peat	3.50	80.3	4.3	76.8	4.3		20.97 0.92
24	sand+peat	3.50	67.9	3.8	64.4	3.8		31.84 1.39
25**	sand+peat							37.69 1.65
27	sand+gravel	3.50	52.7	0.5	49.2	0.6		
<i>6 (^{226}Ra)</i>		3.50	0.33					

^a measurements from Core Scientific International

^b measurements from Flett Research Ltd

^c measurements from My Core Scientific Laboratory

* Depth of 30-yr horizon

** Depth basal age (age determined in deepest sediment layer containing peat overlying sand, clay, or gravel)

665 670 Note that the 30-yr horizon can occur below the depth of the basal age.

Figure A1. Core stratigraphies. Dated high marsh cores (asterisks) were CBE1-1, CRF2-1, GBK1-2, and TMF2-1; Dated low marsh cores were CBE1-5, CRF3-2, GBK1-4, and TMF1-2. Corrected depth (cm) reflects correction for core compaction. Surface organic layer overlying peat in several cores was black, fluffy, highly organic material overlying the brown peat layer below, which was contained extensive plant roots that had not entirely decayed yet. The transition between these two layers was visually very distinct and also reflected by changes in dry bulk density.

675




Figure A2. %LOI-to-%C Relationship. Relationship between measured LOI values and calculated %C, using elemental analyser EA data on set of 93 subsamples. Measurements from core CBE 1-5 were not used for calculating this relationship due to suspected measurement error.





Figure A3. Comparison of %OC calculation using the empirical regression determined for our study area (blue) and the model of Craft et al. (1991) (orange).



690 **Figure A4. Comparison of C stocks (Mg C/ha) estimated using the Craft et al. (1991) equation versus C stocks estimated using this study's empirical relationship for southern BC, for 8 cores from our study region. Comparison suggests that using the Craft et al. (1991) relationship would produce C stocks that are ~32% greater than our estimates. This difference, while substantial, would not account for global C stocks that are 3 times larger than those found in Clayoquot Sound salt marshes.**





Figure A5. ^{210}Pb inventory (Bq m^{-2}) in high and low marsh cores.

Deleted: 3



700 **Appendix B Groundtruthing Marsh Zone Designations: Detrended Correspondence Analysis**

We used colour orthophotos to visually determine the areas of marsh zones using ArcMap 10.5.1 area measurement tools. A vegetation survey also determined the marsh zone of each core sample site, and vegetation data were examined using a detrended correspondence analysis to verify that vegetation assemblages reflected distinct low and high marsh zones identified from the orthophotos (Hill and Gauch 1980).

705 Our comparison showed that differentiating high and low marsh using orthophotos matched with field vegetation data for 32 of 34 (94 %) of cores. Both CRF 1-2 and CRF 2-2 were classified as low marsh by vegetation survey but fell within the high marsh using the visual orthophotography method. These cores lie 16 m (CRF 1-2) and 12 m (CRF 2-2) away from the low marsh boundary as measured using orthophotos, which is less than their distances from the nearest high marsh cores (17 m and 23 m, respectively). All other cores fell within the correct marsh zone.

710 We also conducted a detrended correspondence analysis (Hill and Gauch 1980) of vegetation data to determine that vegetation encountered in the vicinity of the core samples reflected distinct marsh zones using Canoco v4.5 software. We classified of marsh zones by presence/absence and percent cover of low marsh or high marsh vegetation (e.g. Deur 2000). The analysis showed accurate fit of low marsh cores with low marsh vegetation and high marsh cores with high marsh vegetation, plus the addition of a somewhat indistinct, third cluster of vegetation possibly representing the upland vegetation. This square root-

715 transformed model accounted for 33.2% of all variance in the vegetation dataset (sum of eigenvalues = 3.29). Cores with low marsh vegetation clustered together while high marsh cores clustered separately. An additional, slightly distinct third cluster of upland vegetation indicates that some high marsh cores may have been extracted from near the boundary with a freshwater-dominated upland vegetation or salt-tolerant meadow. The distinction between a salt marsh and a bordering freshwater area has complicated efforts to classify marshes by salinity (Duarte et al. 2013), but this result shows that clustering of vegetation

720 type corresponds reasonably well with each site's designation as high or low marsh.

Deleted: backshore

Deleted: backshore

Deleted: close to

Deleted: backshore

Figure B1 Detrended correspondence analysis results for Marsh vegetation data. Low marsh cores (top, purple squares) corresponded reasonably well with vegetation identified as low marsh, and high marsh cores corresponded with a distinct cluster of high marsh vegetation. The far bottom-right may indicate a population of less salt-tolerant, upland vegetation but it is indistinct from the high marsh.

Deleted: backshore





Figure B2 Reference used for determining marsh stratum based on vegetation. Source: Duer 2000.



Data Availability

Data produced as part of this project are currently being submitted to the Pangaea database with the following doi:
<https://doi.pangaea.de/10.1594/PANGAEA.889454>

740 Author contribution

MGP and KEK acquired funding for the project. SC, MGP, and KEK designed the experiments and SC and MGP carried them out. SC conducted laboratory measurements and SC, CO, MG, MGP and KEK contributed to data analysis. KEK, SC, and CO were responsible for data curation. SC and KEK prepared the manuscript with contributions from all co-authors.

Competing interests

745 The authors declare that they have no conflict of interest.

Acknowledgements

The authors extend their gratitude to the following individuals for their contributions to this research, alphabetically: Dr. Richard Atleo, Celeste Barlow, Hasini Basnayake, Dr. Douglas Deur, Dan Harrison, Hannah Jensen, Victoria Lamothé, Dr. Dana Lepofsky, Gemma MacFarland, Aimee McGowan, Bryn Montgomery, Yiga Phuntsok, Ellie Simpson, Maureen Soon, **750** and Dr. Nancy Turner. We thank Parks Canada for in-kind support at both Pacific Rim National Park Reserve and the Vancouver office, the Raincoast Education Society for their expertise in the field and access to their grounds to reach our study sites, and the Ahousaht First Nation for allowing us to collect samples from their territory.

References

Abbott, K. M., Elsey-Quirk, T., and DeLaune, R. D.: Factors influencing blue carbon accumulation across a 32-year **755** chronosequence of created coastal marshes, *Ecosphere*, 10, 10.1002/ecs2.2828, 2019.

Abdul-Aziz, O. I., Ishtiaq, K. S., Tang, J., Moseman-Valtierra, S., Kroeger, K. D., Gonanea, M. E., et al. (2018). [Environmental controls, emergent scaling, and predictions of greenhouse gas \(GHG\) fluxes in coastal salt marshes](#). *Journal of Geophysical Research: Biogeosciences*, 123, 2234–2256. <https://doi.org/10.1029/2018JG004556>

Adams, C. A., Andrews, J. E., and Jickells, T.: Nitrous oxide and methane fluxes vs. carbon, nitrogen and phosphorous burial in new intertidal and saltmarsh sediments, *Science of The Total Environment*, 434, 240-251, <https://doi.org/10.1016/j.scitotenv.2011.11.058>, 2012.

765 Arias-Ortiz, A., Masqué, P., García-Orellana, J., Serrano, O., Mazarrasa, I., Marbà, N., Lovelock, C. E., Lavery, P. S., and Duarte, C. M.: Reviews and syntheses: ^{210}Pb -derived sediment and carbon accumulation rates in vegetated coastal ecosystems – setting the record straight, *Biogeosciences*, 15, 6791-6818, 10.5194/bg-15-6791-2018, 2018.

Bridgman, S. D., Megonigal, J. P., Keller, J. K., Bliss, N. B., and Trettin, C.: The carbon balance of North American wetlands, *Wetlands*, 26, 889-916, 10.1672/0277-5212(2006)26[889:TCBONA]2.0.CO;2, 2006.

Callaway, J.C., Borgnis, E.L., Turner, R.E., and Milan, C.S.: Carbon sequestration and sediment accretion in San Francisco Bay tidal wetlands, *Estuarine, Coastal and Shelf Science*, 35, 1163–1181, 2012.

770 Chmura, G. L., Anisfeld, S. C., Cahoon, D. R., and Lynch, J. C.: Global carbon sequestration in tidal, saline wetland soils, *Global Biogeochem. Cycles*, 17, doi:10.1029/2002GB001917, 2003.

Chmura, G. L., Kellman, L., and Guntenspergen, G. R.: The greenhouse gas flux and potential global warming feedbacks of a northern macrotidal and microtidal salt marsh, *Environmental Research Letters*, 6, 044016, 10.1088/1748-9326/6/4/044016, 2011.

775 Connor, R. F., Chmura, G. L., and Beecher, C. B.: Carbon accumulation in bay of fundy salt marshes: Implications for restoration of reclaimed marshes, *Global Biogeochemical Cycles*, 15, 943-954, <https://doi.org/10.1029/2000GB001346>, 2001.

Corbett, D. R. and Walsh, J. P.: $^{210}\text{Lead}$ and $^{137}\text{Cesium}$: establishing a chronology for the last century. In: *Handbook of Sea-Level Research*, Shennan, I., Long, A. J., and Horton, B. P. (Eds.), John Wiley & Sons, Ltd., Chichester, UK, 2015.

780 [Craft, C. & Seneca, E. & Broome, Stephen. Loss on Ignition and Kjeldahl Digestion for Estimating Organic Carbon and Total Nitrogen in Estuarine Marsh Soils: Calibration with Dry Combustion. *Estuaries*. 14. 175-179. 10.2307/1351691, 1991.](#)

Craft, C., Megonigal, P., Broome, S., Stevenson, J., Freese, R., Cornell, J., Zheng, L., and Sacco, J.: The pace of ecosystem development of constructed *Spartina alterniflora* marshes, *Ecological Applications*, 13, 1417-1432, <https://doi.org/10.1890/02-5086>, 2003.

785 Crooks, S., Rybczyk, J., O'Connell, K., Devier, D.L., Poppe, K., and Emmett-Mattox, S.: Coastal Blue Carbon Opportunity Assessment for the Snohomish Estuary: The Climate Benefits of Estuary Restoration, Report by Environmental Science Associates, Bellingham, WA, USA, pp. 19-43, 2014.

Davis, R. B., Hess, C. T., Norton, S. A., Hanson, D. W., Hoagland, K. D., and Anderson, D. S.: ^{137}Cs and ^{210}Pb dating of sediments from soft-water lakes in New England (U.S.A.) and Scandinavia, a failure of ^{137}Cs dating, *Chemical Geology*, 44, 151-185, [https://doi.org/10.1016/0009-2541\(84\)90071-8](https://doi.org/10.1016/0009-2541(84)90071-8), 1984.

Deur, D.: A Domesticated Landscape: Native American plant cultivation on the Northwest coast of North America, PhD dissertation, Louisiana State University, Baton Rouge, LA, USA, pp. 69-251, 2000.

[DFO. Marine Environmental Data Section Archive, <https://meds-sdmm.dfo-mpo.gc.ca>, Ecosystem and Oceans Science, Department of Fisheries and Oceans Canada. Data obtained on 2022/06/02.](#)

Donato, D. C., Kauffman, J. B., Murdiyarso, D., Kurnianto, S., Stidham, M., and Kanninen, M.: Mangroves among the most carbon-rich forests in the tropics, *Nature Geoscience*, 4, 293-297, 10.1038/geo1123, 2011.

Drexler, J. Z., Woo, I., Fuller, C. C., and Nakai, G.: Carbon accumulation and vertical accretion in a restored versus historic salt marsh in southern Puget Sound, Washington, United States, *Restoration Ecology*, 27, 1117-1127, 800 <https://doi.org/10.1111/rec.12941>, 2019.

Duarte, C. M., Middelburg, J. J., and Caraco, N.: Major role of marine vegetation on the oceanic carbon cycle, *Biogeosciences*, 2, 1-8, 10.5194/bg-2-1-2005, 2005.

Duarte, C. M., Losada, I. J., Hendriks, I. E., Mazarrasa, I., and Marbà, N.: The role of coastal plant communities for climate change mitigation and adaptation, *Nature Climate Change*, 3, 961-968, 10.1038/nclimate1970, 2013.

805 2016. Canadian Climate Normals 1981-2010 Station Data: Tofino A.http://climate.weather.gc.ca/climate_normals/results_1981_2010_e.html?searchType=stnName&txtStationName=Tofino&searchMethod=contains&txtCentralLatMin=0&txtCentralLatSec=0&txtCentralLongMin=0&txtCentralLongSec=0&stnID=277&dispBack=1. Last access: 13 November, 2016.

810 Froelich, P. N.: *Analysis of organic carbon in marine sediments, Limnology and Oceanography*, 25, 564-572, <https://doi.org/10.4319/lo.1980.25.3.0564>, 1980.

Gailis, M., Kohfeld, K. E., Pellatt, M. G., and Carlson, D.: Quantifying blue carbon for the largest salt marsh in southern British Columbia: implications for regional coastal management, *Coastal Engineering Journal*, <https://doi.org/10.1080/21664250.2021.1894815>, in press 2021.

815 Giles-Hansen, K. and Wei, X.: *Cumulative disturbance converts regional forests into a substantial carbon source, Environmental Research Letters*, 17, 044049, 10.1088/1748-9326/ac5e69, 2022.

Havens, K., Varnell, L., and Watts, B.: Maturation of a constructed tidal marsh relative to two natural reference tidal marshes over 12 years, *Ecological Engineering*, 18, 305-315, 2002.

820 Heiri, O., Lotter, A. F., and Lemcke, G.: Loss on ignition as a method for estimating organic and carbonate content in sediments: reproducibility and comparability of results, *Journal of Paleolimnology*, 25, 101-110, 10.1023/A:1008119611481, 2001.

Hedges, J.I. and Stern, J.H.: Carbon and nitrogen determinations of carbonate-containing solids, *Limnology and Oceanography* 29(3), 657-663, 1984.

Hill, M. O. and Gauch, H. G.: Detrended correspondence analysis: An improved ordination technique, *Vegetatio*, 42, 47-58, 10.1007/BF00048870, 1980.

825 Hodgson, C. and Spooner, A.: The K'omoks and Squamish Estuaries: A Blue Carbon Pilot Project, Final Report to the North American Partnership for Environmental Community Action (NAPECA), (Grant 2014-1362), Comox Valley Watershed Society, Comox, BC, Canada, 2016.

Howard, J., Hoyt, S., Isensee, K., Pidgeon, E., and Telszewski, M.: (eds.) *Coastal Blue Carbon: Methods for assessing carbon stocks and emissions factors in mangroves, tidal salt marshes, and seagrass meadows*, Conservation International,

Deleted: Fisheries and Oceans Canada TOFINO – Tides, Currents, and Water Levels. http://www.tides.gc.ca/eng/data/table/2016/wlev_ref/8615. Last Access: 13 June, 2017, 2016.¶

835 Intergovernmental Oceanographic Commission of UNESCO, International Union for Conservation of Nature,
Arlington, Virginia, USA, 15-24; 149-150, 2014.

Howard, J., Sutton-Grier, A., Herr, D., Kleypas, J., Landis, E., McLeod, E., Pidgeon, E., and Simpson, S.: Clarifying the role
of coastal and marine systems in climate mitigation, *Frontiers in Ecology and the Environment*, 15(1): 42-50,
doi:10.1002/fee.1451, 2017.

840 Howe, A. J., Rodríguez, J. F., and Saco, P. M.: Surface evolution and carbon sequestration in disturbed and undisturbed
wetland soils of the Hunter estuary, southeast Australia, *Estuarine, Coastal and Shelf Science*, 84, 75-83,
<https://doi.org/10.1016/j.ecss.2009.06.006>, 2009.

[Huang J, Luo M, Liu Y, Zhang Y, Tan J. Effects of tidal scenarios on the methane emission dynamics in the Subtropical tidal marshes of the Min River Estuary in Southeast China. *Int J Environ Res Public Health*, 16\(15\):2790, doi:10.3390/ijerph16152790, 2009.](#)

845 Huertas, I. E., de la Paz, M., Perez, F. F., Navarro, G., and Flecha, S.: Methane emissions from the salt marshes of Doñana
wetlands: spatio-temporal variability and controlling factors, *Frontiers in Ecology and Evolution*, 7, 2019.

James, TS, JA Henton, LJ Leonard, A Darlington, DL Forbes, M Craymer, Relative sea-level projections in Canada and the
adjacent mainland United States, *Geological Survey of Canada Open File* 7737, 72 p. doi: 10.4095/295574, 2014.

850 Jefferson, C.: Some Aspects of Plant Succession in Oregon Estuarine Salt Marshes, (unpublished Ph.D. dissertation, Oregon
State University, Corvallis, OR, USA, 1973.

Johannessen, S. C. and Macdonald, R. W.: Geoengineering with seagrasses: is credit due where credit is given?, *Environmental
Research Letters*, 11, 113001, 10.1088/1748-9326/11/11/113001, 2016.

855 Kauffman, J. B., Giovanonni, L., Kelly, J., Dunstan, N., Borde, A., Diefenderfer, H., Cornu, C., Janousek, C., Apple, J., and
Brophy, L.: Total ecosystem carbon stocks at the marine-terrestrial interface: Blue carbon of the Pacific Northwest
Coast, United States, *Global Change Biology*, 26, 5679-5692, <https://doi.org/10.1111/gcb.15248>, 2020.

Kelleway, J. J., Saintilan, N., Macreadie, P. I., Baldock, J. A., and Ralph, P. J.: Sediment and carbon deposition vary among
vegetation assemblages in a coastal salt marsh, *Biogeosciences*, 14, 3763-3779, 10.5194/bg-14-3763-2017, 2017.

[Kennedy HA, Alongi DM, Karim A, Chen G, Chmura GL, Crooks S, Kairo JG, Liao B, Lin G. 2013. Chapter 4 Coastal
Wetlands In: *Supplement to the 2006 IPCC Guidelines on National Greenhouse Gas Inventories: Wetlands*](#)

860 Kirwan, M. L., Walters, D. C., Reay, W. G., and Carr, J. A.: Sea level driven marsh expansion in a coupled model of marsh
erosion and migration, *Geophysical Research Letters*, 43, 4366-4373, <https://doi.org/10.1002/2016GL068507>, 2016.

Krishnaswamy, S., Lal, D., Martin, J. M., and Meybeck, M.: Geochronology of lake sediments, *Earth and Planetary Science
Letters*, 11, 407-414, [https://doi.org/10.1016/0012-821X\(71\)90202-0](https://doi.org/10.1016/0012-821X(71)90202-0), 1971.

865 Kurz, W.A., Shaw, C.H., Boisvenue, C., Stinson, G., Metsaranta, J., Leckie, D., Dyk, A., Smyth, C. and Neilson, E.T.:
Carbon in Canada's boreal forest—a synthesis *Environmental Reviews*, 21, 260-292, 10.1139/er-2013-0041, 2013.

[Li Y, Wang D, Chen Z, Chen J, Hu H, Wang R. Methane emissions during the tide cycle of a Yangtze Estuary salt marsh.
Atmosphere, 12\(2\):245. <https://doi.org/10.3390/atmos12020245>, 2021.](#)

Deleted: IPCC, Hiraishi, T., Krug, T., Tanabe, K., Srivastava, N., Baasansuren, J., Fukuda, M., and Troxler, T.G. (Eds.): Supplement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories: Wetlands, Intergovernmental Panel on Climate Change, Switzerland, 2014.¶

875 [Mack, M. C., Walker, X. J., Johnstone, J. F., Alexander, H. D., Melvin, A. M., Jean, M., and Miller, S. N.: Carbon loss from boreal forest wildfires offset by increased dominance of deciduous trees, Science, 372, 280-283, doi:10.1126/science.abf3903, 2021.](#)

[MacKenzie, W.H. and J.R. Moran. Wetlands of British Columbia: a guide to identification. Res. Br., B.C. Minisitry of Forests, Victoria, B.C. Land Management Handbook No. 52, 2004.](#)

880 Mathieu, G. G., Biscaye, P. E., Lupton, R. A., and Hammond, D. E.: System for measurement of 222Rn at low levels in natural waters, *Health Phys.*, 55, 989-992, 1988.

Mazzotti, S., Jones, C., and Thomson, R.E.: Relative and absolute sea level rise in western Canada and northwestern United States from a combined tide gauge-GPS analysis, *Journal of Geophysical Research*, 113:1-19, C11019, doi:10.1029/2008JC004835, 2008.

885 Mazzotti, S., Lambert, A., Courtier, N., Nykolaishen, L., and Dragert, H.: Crustal uplift and sea level rise in northern Cascadia from GPS, absolute gravity, and tide gauge data, *Geophysical Research Letters*, 34, L15306. <https://doi.org/10.1029/2007GL030283>, 2007.

McLeod, E., Chmura, G.L., Bouillon, S., Salm, R., Björk, M., Duarte, C.M., Lovelock, C.E., Schlesinger, W.H., and Silliman, B.R.: A blueprint for blue carbon: toward an improved understanding of the role of vegetated coastal habitats in sequestering CO₂, *Frontiers in Ecology and the Environment*, 9(10):552–560, 2011.

890 Mcowen, C.J., Weatherdon L., Van Bochove, J.W., Sullivan, E., Blyth, S., Zockler, C., Stanwell-Smith, D., Kingston, N., Martin C.S., Spalding M., and Fletcher, S.: A global map of saltmarshes, *Biodiversity Data Journal* 5: e11764, doi: 10.3897/BDJ.5.e11764, 2017.

Montillet, J.-P., Melbourne, T. I., and Szeliga, W. M.: GPS Vertical Land Motion Corrections to Sea-Level Rise Estimates in the Pacific Northwest, *Journal of Geophysical Research: Oceans*, 123, 1196-1212, <https://doi.org/10.1002/2017JC013257>, 2018.

895 Nuwer, J. M. and Keil, R. G.: Sedimentary organic matter geochemistry of Clayoquot Sound, Vancouver Island, British Columbia, *Limnology and Oceanography*, 50, 1119-1128, <https://doi.org/10.4319/lo.2005.50.4.1119>, 2005.

Olid, C., Garcia-Orellana, J., Martínez-Cortizas, A., Masqué, P., Peiteado-Varela, E., and Sanchez-Cabeza, J.-A.: Multiple site study of recent atmospheric metal (Pb, Zn and Cu) deposition in the NW Iberian Peninsula using peat cores, *Sci Total Environ.*, 408, 5540-5549, 10.1016/j.scitotenv.2010.07.058, 2010.

900 Ouyang, X., and Lee, S.Y.: Updated estimates of carbon accumulation rates in coastal marsh sediments, *Biogeosciences*, 11(18): 5057–5071, <http://doi.org/10.5194/bg-11-5057-2014>, 2014.

Pendleton, L., Donato, D., Murray, B., Crooks, S., Jenkins, W., Sifleet, S., Craft, C., Fourqurean, J. W., Kauffman, J. B., Marba', N. r., Megonigal, P., Pidgeon, E., Herr, D., Gordon, D., and Baldera, A.: Estimating global blue carbon emissions from conversion and degradation of vegetated coastal ecosystems, *PLoS One*, 7, e43542, doi:10.1371/journal.pone.0043542, 2012.

Poffenbarger, H. J., Needelman, B. A., and Megonigal, J. P.: Salinity Influence on Methane Emissions from Tidal Marshes, *Wetlands*, 31, 831-842, 10.1007/s13157-011-0197-0, 2011.

Poppe, K.L., and J.M. Rybczyk. 2019. A blue carbon assessment for the Stillaguamish River estuary: Quantifying the climate
910 benefits of tidal marsh restoration. In Summary report prepared by Western Washington University for Washington
Sea Grant and The Nature Conservancy, 19. Bellingham, WA, USA: Western Washington University.

Porter, G.L.: Vegetation-environment relationships in the tidal marshes of the Fraser River Delta, British Columbia. Masters,
The University of British Columbia Vancouver, BC, Canada, 1982.

Postlethwaite, V. R., McGowan, A. E., Kohfeld, K. E., Robinson, C. L. K., and Pellatt, M. G.: Low blue carbon storage in
915 eelgrass (*Zostera marina*) meadows on the Pacific Coast of Canada, *PLoS One*, 13, e0198348,
10.1371/journal.pone.0198348, 2018.

[Price, D. T., Alfaro, R. I., Brown, K. J., Flannigan, M. D., Fleming, R. A., Hogg, E. H., Girardin, M. P., Lakusta, T., Johnston, M., McKenney, D. W., Pedlar, J. H., Stratton, T., Sturrock, R. N., Thompson, I. D., Trofymow, J. A., and Venier, L. A.: Anticipating the consequences of climate change for Canada's boreal forest ecosystems. Environmental Reviews, 21, 322-365, 10.1139/er-2013-0042, 2013](#)

Ryder, J.L., Kenyon, J.K., Buffet, D., Moore, K., Ceh, M. and Stipek, K.: An integrated biophysical assessment of estuarine
habitats in British Columbia to assist regional conservation planning, No. 476, Technical Report Series, 2007.

Schumacher, Brian A.: Methods for the Determination of Total Organic Carbon in Soils and Sediments, US Environmental
920 Protection Agency Environmental Sciences Division- National Exposure Research Laboratory, Las Vegas, NV, USA,
2-11, 2002.

[Stinson, G., Kurz, W. A., Smyth, C. E., Neilson, E. T., Dymond, C. C., Metsaranta, J. M., Boisvenue, C., Rampley, G. J., Li, Q., White, T. M., and Blain, D.: An inventory-based analysis of Canada's managed forest carbon dynamics, 1990 to
2008, Global Change Biology, 17, 2227-2244, https://doi.org/10.1111/j.1365-2486.2010.02369.x, 2011.](#)

[van Ardenne, L. B., Jolicouer, S., Bérubé, D., Burdick, D., and Chmura, G. L.: The importance of geomorphic context for
930 estimating the carbon stock of salt marshes, *Geoderma*, 330, 264-275,
https://doi.org/10.1016/j.geoderma.2018.06.003, 2018.](#)

[Walker, X. J., Baltzer, J. L., Cumming, S. G., Day, N. J., Ebert, C., Goetz, S., Johnstone, J. F., Potter, S., Rogers, B. M.,
Schuur, E. A. G., Turetsky, M. R., and Mack, M. C.: Increasing wildfires threaten historic carbon sink of boreal forest
soils, *Nature*, 572, 520-523, 10.1038/s41586-019-1474-y, 2019.](#)

[Wei, S., Han, G., Chu, X., Song, W., He, W., Xia, J., and Wu, H.: Effect of tidal flooding on ecosystem CO₂ and CH₄ fluxes
935 in a salt marsh in the Yellow River Delta, *Estuarine, Coastal and Shelf Science*, 232, 106512, 2020.](#)

Weinmann, F.M., Boule, M., Brunner, K., Malek, J., Yoshino V.: Wetland Plants of the Pacific Northwest, US Army Corps
of Engineers, Seattle, WA, USA, 1984.