

**Mass concentration measurements of autumn bioaerosol using low cost sensors in a mature temperate woodland Free Air Carbon Dioxide Enrichment (FACE) experiment: investigating the role of meteorology and carbon dioxide levels**

Aileen B. Baird<sup>1,2</sup>, Edward J. Bannister<sup>1,2</sup>, A. Robert MacKenzie<sup>1,2</sup>, and Francis D. Pope<sup>1,2\*</sup>

<sup>1</sup>School of Geography, Earth & Environmental Sciences, University of Birmingham, Birmingham, B152TT, UK

<sup>2</sup>Birmingham Institute of Forest Research, University of Birmingham, Birmingham, B15 2TT, UK

\*Correspondence to: Francis Pope [f.pope@bham.ac.uk](mailto:f.pope@bham.ac.uk)

**Abstract.**

10 Forest environments contain a wide variety of airborne biological particles (bioaerosols), including pollen, fungal spores, bacteria, viruses, plant detritus and soil particles. Forest bioaerosol plays a number of important roles related to plant and livestock health, human disease and allergenicity, forest and wider ecology, and are thought to influence biosphere/atmosphere interactions via warm and cold cloud formation. Despite the importance of bioaerosols, there are few measurements of forest aerosol, and there is a lack of understanding of how climate change will affect forest bioaerosol in the future.

15 We installed low-cost optical particle counters (OPCs) to measure particles in the size range between 1 and 10  $\mu\text{m}$  ( $\text{PM}_{10}\text{-PM}_1$ ) for a period of two months in Autumn 2018 at the Birmingham Institute of Forest Research (BIFoR) Free Air Carbon Dioxide Enrichment (FACE) facility. In the paper, we propose that the  $\text{PM}_{10}\text{-PM}_1$  metric is a good proxy for bioaerosols because of the bioaerosol representative size range, the location of the study site (a woodland in a rural location), the field measurement taking place during the season of peak fungal activity, and the low hygroscopicity of the particles measured. The BIFoR FACE  
20 facility fumigates three 700  $\text{m}^2$  areas of the forest with an additional 150 ppm  $\text{CO}_2$  above ambient with minimal impacts on other potential environmental drivers such as temperature, humidity, and wind. This experimental set-up enabled us to investigate the effect of environmental variables, including elevated  $\text{CO}_2$  ( $\text{eCO}_2$ ), on bioaerosol proxy concentrations, and to evaluate the performance of the low-cost OPCs in a forested environment.

Operating the low-cost OPCs during Autumn 2018, we aimed to capture predominantly the fungal bioaerosol season. Across  
25 the experimental duration, the OPCs captured both temporal and spatial variation in bioaerosol concentrations. Aerosol concentrations were affected by changing temperatures and wind speeds, but, contrary to our initial hypothesis, not by relative humidity. We detected no effect of the  $\text{eCO}_2$  treatment on total bioaerosol concentrations, but a potential suppression of high concentration bioaerosol events was detected under  $\text{eCO}_2$ . In-canopy atmospheric dispersion modelling indicates that the median spore dispersion distance is sufficiently small that there is little mixing between treatment and control experiments.

30 Our data demonstrate the suitability of low-cost OPCs, interpreted with due caution, for use in forests, and so opens the possibility of forest bioaerosol monitoring in a wider range of habitats, to a wider range of researchers at a modest cost.

## 1 Introduction

Aerobiology investigates the production, transport, and removal of airborne biological particles such as bacterial, fungal, and viral microbes, plant pollen, and soil and plant debris. Fungal spores form a large proportion of the bioaerosol population, with ground level concentrations typically 10,000–50,000 spores m<sup>-3</sup> (compared with 1000–2000 pollen grains m<sup>-3</sup>) (Sesartic and Dallafior, 2011, and references therein). Modelling suggests that fungal spores form approximately 23% of bioaerosol mass globally (Heald and Spracklen (2009). Understanding the dispersal of fungal spores is not only essential for understanding fungal ecology and their direct relevance to the forest plant and soil communities, but also for understanding human disease and allergens, plant and crop health, livestock health, as well as the role played by bioaerosols in ice nucleation and cloud condensation (Pöschl, 2005; Reimmuth-Selzle et al., 2017; Sesartic and Dallafior, 2011a). The significant influence of airborne fungal particles on forest biology is often overlooked.

Spores are produced by fungi as propagules during asexual and sexual reproduction (Læssøe and Petersen, 2019). Fungal spores are varied in shape and size, with typical diameters of 1–30 µm, although their diameters can vary from 1–1000 µm (Fröhlich-Nowoisky et al., 2009; Halbwachs and Bässler, 2015; Jones and Harrison, 2004). Spore dispersal is a complex process, affected by size and shape of the spores, meteorological and ecological factors, and the life history of the fungus. Fungi which produce below-ground fruit bodies, and some mycorrhizal fungi without sporocarps, rarely have spores in significant concentrations in the air, and are more commonly spread by animals. Above-ground fungal fruiting bodies with active release mechanisms or tall stipites may release spores higher into the air (Biedermann and Vega, 2020; Dressaire et al., 2016; Halbwachs and Bässler, 2015; Horton, 2017; Kivlin et al., 2014; Lilleskov and Bruns, 2005; Stephens and Rowe, 2020). Spore size, shape, and ornamentation influence their dispersal, with smaller and/or less dense spores able to travel further, although even very large spores may be dispersed by the air (Chaudhary et al., 2020; Norros et al., 2014, 2012). In addition to the spore and fungal characteristics, the meteorology and ecology of the environment both significantly influence spore dispersal (Halbwachs and Bässler, 2015; Oneto et al., 2020). Mature forests generally have complex, multi-layered canopies which affect airflow through the forest, in addition to altering temperature and relative humidity (RH) variability (Bannister et al., 2021; Gilbert and Reynolds, 2005; Norros et al., 2014).

Meteorological factors have also been shown to impact airborne fungal communities significantly. Several studies have measured concentrations of airborne fungal spores in UK (largely in cities), investigating diurnal and seasonal variables as well as meteorological variables on fungal spores. The effects are almost always species-specific, with a broader seasonal trend of fungal spore concentrations being highest in late summer and autumn (Hollins et al., 2004; O'Connor et al., 2014; Sadyś et al., 2016a, 2014). Larger European studies (e.g. Grinn-Gofroń et al., 2019) and reviews have shown that the two most influential meteorological variables for airborne fungal spore concentrations are temperature and RH (Jones and Harrison, 2004; Moore-Landecker, 2011).

As well as being affected on shorter time scales by meteorological factors, airborne fungal concentrations can also be affected by climate change, both due to increases in CO<sub>2</sub> concentrations as well as the corresponding increase in temperature, although

65 effects on fungi may vary depending on the species (Burge, 2002). Wolf et al. (2003) tested the response of 11 arbuscular  
mycorrhizal (AM) fungi to eCO<sub>2</sub> at the BioCON FACE grassland experiment, with only a single *Glomus* species producing  
additional spores in the soil. Wolf et al. (2010) demonstrated an increase in spore production from *Alternaria alternata* (a  
common airborne allergen) under eCO<sub>2</sub> in a growth chamber. Similarly, in an Aspen tree (*Populus tremuloides*) open-topped  
chamber eCO<sub>2</sub> experiment, Klironomos et al. (1997) found that airborne fungal spore concentrations increased, which they  
70 suggested was due to corresponding increases in spore concentrations in the leaf litter.

In addition to direct changes in spore production, it is possible that fungal sporocarp production, and therefore spore production,  
could also increase due to climate change. Ecological measurements show the autumn fungal fruiting seasons in Europe have  
become longer over the last 50 years. The fungal fruiting season starts earlier and finishes later due to climate change, in  
addition to more fungi having an additional spring fruiting season (Gange et al., 2007; Kausserud et al., 2012). This climate-  
75 induced phenology change has also been demonstrated in a corresponding increase in the airborne fungal spore season (Sadyš  
et al., 2016b). These large changes in the fungal season were mainly attributed to temperature increases; however, at the Aspen  
FACE experiment, it was found that ectomycorrhizal fungi sporocarp production increased under eCO<sub>2</sub>, so it seems possible  
that CO<sub>2</sub> impacts fungal sporocarp production independently of temperature (Andrew and Lilleskov, 2009). All of  
these demonstrated changes in fungal phenology, sporocarp production, and sporulation suggest that bioaerosol concentrations  
80 are also likely to change under eCO<sub>2</sub>. Even if these findings are fungal species-specific, they have potentially wide-ranging  
effects for forested habitats.

The studies described above demonstrate that fungi are likely to change spore and sporocarp production under eCO<sub>2</sub> ranging  
from 192–600 ppm, in addition to the significant changes witnessed in fungal growing seasons under the current anthropogenic  
increases and broader climate effects. Even if these findings are fungal species-specific, they indicate that fungal bioaerosols  
85 concentrations can be expected to change under eCO<sub>2</sub>, with potentially wide-ranging effects for forested habitats (Baird and  
Pope, 2021). However, none of the above studies were completed in complex mature woodlands, with the experiments being  
completed in laboratories, open-topped chambers or, in the case of Aspen FACE, a young plantation forest. Mature and ancient  
woodlands represent a more complex and diverse environment, which are likely to respond differently to eCO<sub>2</sub> than the  
plantations of young trees on agricultural soil studied in earlier FACE experiments (Norby et al., 2016). There are also few  
90 studies of forest airborne spore concentration responses to eCO<sub>2</sub>, with the majority of studies focussing on sporocarp  
production.

In order to study airborne forest fungal bioaerosols under eCO<sub>2</sub> in a mature temperate woodland, we installed low-cost Optical  
Particle Counters (OPCs) into the BIFoR FACE experiment during Autumn 2018. We assume that the bioaerosols can be  
represented by measured difference between PM<sub>10</sub> and PM<sub>1</sub> mass concentrations, as detailed in the methods section.

95 Our hypotheses were:

- A) Hourly fungal bioaerosol concentrations will correlate with hourly weather conditions (wind, RH, temperature).
- B) Fungal bioaerosol concentrations will increase in woodland patches treated with eCO<sub>2</sub> for two years.

## 2 Methods

### 2.1 BIFoR FACE

100 The BIFoR FACE experiment is located in a 19.1 hectare mature oak forest in Staffordshire, UK. The overall aim of the FACE  
experiment is to mimic the effects of anthropogenic climate change by increasing the CO<sub>2</sub> concentrations in areas of the forest  
by 150 ppm above ambient (~400 ppm) (MacKenzie et al., 2020; Norby et al., 2016). This set-up is one of only two FACE  
experiments in mature forests worldwide and it provides a unique experiment to study mature temperate woodland. In the  
context of bioaerosols, this experiment allows the investigation of the direct and indirect effects of eCO<sub>2</sub> in a minimally  
105 disturbed forest environment (Hart et al., 2019).

As shown in the site map in [Figure 1](#), the CO<sub>2</sub> enrichment set-up consists of nine roughly circular “arrays” of three types, each  
array being approximately 30 m in diameter. Arrays 1, 4, and 6 are fumigated with additional CO<sub>2</sub>, increasing the atmospheric  
CO<sub>2</sub> by 150 ppm above ambient (to approximately 550 ppm). The CO<sub>2</sub>, pre-mixed with air, is released into the tree canopies,  
using pipes running the height of 25 m tall towers around the perimeter of the array. Arrays 2, 3, and 5 have the same tower  
set-up, but fumigate only with air. Arrays 7, 8 and 9 are “ghost” arrays without any fumigation infrastructure. Arrays 1–6 are  
110 grouped into three treatment pairs, based on pre-fumigation vegetation and soil biochemistry analysis, each with a single  
elevated CO<sub>2</sub> and single ambient CO<sub>2</sub> array. The array pairs are as follows: 1 and 3; 2 and 4; 5 and 6. The present study used  
Arrays 1–6 only, measuring in each pair of arrays consecutively. Fumigation occurs during the oak growing season,  
approximately 1<sup>st</sup> April to 31<sup>st</sup> October during daylight hours. In 2018, CO<sub>2</sub> fumigation ended on the 31<sup>st</sup> October 2018 at  
115 sundown.

The predominant tree species in the woodland is *Quercus robur* (English oak), with *Corylus avellana* (Hazel) forming a distinct  
understory layer. Other tree species present include *Acer pseudoplatanus* (Sycamore), *Crataegus monogyna* (Common  
Hawthorn), and *Ilex spp.* (Holly). A recent vegetation survey, in 2019, was completed to determine the major ground plant  
species: *Rubus fruticosus* (Bramble), *Phegopteris connectilis* (Beech fern), a *Hedera sp.* (Ivy), and grasses where the canopy  
120 has been opened for access rides (G. Platt, private communication, 2019). Hanging and fallen deadwood is left in place except  
where it poses a direct health-and-safety risk.


In order to focus on the fungal bioaerosol component, we took measurements during the autumn, when pollen and non-fungal  
spore counts from the dominant vegetation were likely to be low. For example, the pollen counts are highest in spring for the  
two dominant tree species at BIFoR FACE. UK oak pollen season can range from March to July and lasts for 4–8 weeks, with  
125 peak concentrations usually occurring in May (Grundström et al., 2019). Hazel pollen season falls earlier in the year, starting  
as early as January, peaking in February or March (National Pollen and Aerobiology Research Unit, 2012). For ground cover  
species, the grass pollen season peaks in the summer months, sometimes extending into early September (National Pollen and  
Aerobiology Research Unit, 2012), and *P. connectilis* sporulation peaking mid-July until mid-September (Page, 1997). None  
of these pollen and fern sporulation seasons coincide with our measurement period. Our measurement period coincides with

Deleted: Figure 1

the fungal fruiting season at BIFoR (see below), as well as previously measured peaks in UK fungal spore concentrations (Davies et al., 1963).

Monthly macro-fungi surveys were completed during the OPC measurement period. The protocol for the macrofungal survey can be found in Van Norman et al., 2008. These surveys revealed the annual peak of fungal fruiting in 2018 occurred across

135 September and October at BIFoR FACE. Species fruiting in October 2018 included, but were not limited to: *Lactarius quietus* (oakbug milkcap), at least one *Russula* (Brittlelegill) species, *Lycoperdon perlatum* (Common puffball), *Mycena rosea* (Rosy Bonnet) and other *Mycena* species, *Auricularia auricula* (Jelly ear), *Hypholoma fasciculare* (Sulphur tuft), *Xylaria hypoxylon* (Candlesnuff Fungus), and a number of resupinate fungi, including several *Stereum* species. All of these fungi produce spores in the range of 3–9  $\mu\text{m}$ , matching the measurement capabilities of the OPCs (Læssøe and Petersen, 2019). *Auricularia auricula*  
140 is an exception; it was observed but produces larger, 16–18  $\mu\text{m}$ -long spores likely outside the OPC size window.



145 **Figure 1.** Map of the BIFoR forest and FACE facility located in Staffordshire (UK) adapted from Hart et al., 2019. The green area shows the whole 19.1 ha Mill Haft woodland, 7.3 ha of which is controlled by the University of Birmingham. The access road is shown as a thick black line. Circles represent scientific research “arrays”. Orange arrays 1, 4, and 6 are surrounded with 25 m tall towers. CO<sub>2</sub> levels inside the array are raised by fumigation to 150 ppm above ambient (from ~400 ppm to ~550 ppm). Blue arrays

2, 3, and 5 have similar towers, but are an infrastructure control, spraying out air with current ambient concentrations of CO<sub>2</sub>. Grey arrays 7, 8, and 9, are non-infrastructure controls, with no towers or fumigation. Treatment and control arrays 1–6 are paired as follows: 1 and 3, 2 and 4, 5 and 6. The landscape surrounding BIFoR FACE is predominantly young tree plantations of mixed broadleaves and conifers (MacKenzie et al., 2020).

## 150 2.2 Instrumentation

In this study, we use optical particle counters, OPC-N2 sensors (Alphasense, Essex), which count and size particles between 0.3 μm and 10 μm in diameter. The sensors are described in detail by Crilley et al. (2018) and Sousan et al. (2016). In brief, the OPCs count the number of particles, and use particle light scattering to determine particle size using Mie scattering approximations (an de Hulst, 1957). To calculate particle mass concentration, particles are assumed to be of uniform spherical shape, the density is assumed to be 1.65 g cm<sup>-3</sup> and the refractive index is assumed to be 1.5 + i0. The values for density and refractive index are the taken from the standard settings of the OPC. The choice of particle density and refractive index has implications for the derived mass concentrations, however comparison between measurements taken within the woodland should be self-consistent. It is noted that fungal spores, the target of this study, are often ellipsoidal in shape and can be defined by a short and long axis diameter. Different fungal species generate spores with different shapes, sizes, and density. The assumptions used by the OPC are typically wrong on a single particle (or bioaerosol) basis but should approach reality on an ensemble averaged basis.

The sensors do not explicitly discriminate between particle types, so in order to discriminate between fungi and other smaller particles (bacteria and anthropogenic aerosols), we excluded data from particles smaller than 1 μm in diameter, measuring from 1 μm up to the maximum 10 μm measuring capacity of the OPCs. This size discrimination, in conjunction with the experimental location and seasonal timing of the experiment make it highly likely the majority of bioaerosols being captured are predominantly of fungal origin.

One of the key concerns raised for the sensing of any aerosol type using low-cost OPCs is that they can report artificially high readings under high RH conditions. Hygroscopic aerosols take up water from the surrounding gaseous environment as a function of RH, with greater water uptake at higher RH. To compare the abundance of aerosol under different environmental conditions, it is preferable to use a ‘dry’ aerosol concentration, with the hygroscopic growth due to water removed. To parameterise particle hygroscopicity, κ-Köhler theory is used, which describes the particle hygroscopicity using a single hygroscopicity parameter kappa (κ) (Petters and Kreidenweis, 2007). Pope (2010) details how κ-Köhler theory can be used to link particle mass to hygroscopicity by Eq. (1).

$$a_w = \frac{\left(\frac{m}{m_0} - 1\right)}{\left(\frac{m}{m_0} - 1\right) + \frac{\rho_w \kappa}{\rho_p}} \quad (1)$$

175  $a_w$  is equilibrium RH expressed as a decimal,  $m$  is the wet aerosol mass,  $m_0$  is the dry aerosol mass, and  $\rho_w$  and  $\rho_p$  the density of water and the dry particulate, respectively. The value of  $\kappa$  can be found by curve fitting of the pollen humidograms ( $m/m_0$  versus  $a_w$ ).

Deleted:

180 Biological particles such as fungal spores and pollen have been demonstrated to be hygroscopic, although they typically have  
a low kappa ( $\kappa$ ) value, with pollen  $\kappa$  typically around 0.05–0.1 as opposed to a  $\kappa$  of approximately 0.3 for average anthropogenic  
aerosols (Griffiths et al., 2012; Pope, 2010). Due to the hygroscopicity of fungal aerosols, high relative humidity can result in  
a high mass concentration reading in sensors, such as the low-cost OPCs, which do not warm or dry the air. This hygroscopic  
effect has been explored using low-cost OPCs in urban environments by Crilley et al. 2020 and Crilley et al. 2018, where they  
used a calibration factor to correct for RH effects.

### 185 2.3 Instrument installation, bias-checking, and calibration

We used a pair of low-cost OPCs to study bioaerosols at BIFoR FACE. The two OPCs used were controlled to take  
measurements every 60 seconds using a Raspberry Pi computer (Crilley et al., 2018, based on Hagan, 2017). The OPCs and  
accompanying instrumentation were installed at 2 m height in a pair of BIFoR FACE arrays at a time (according to the pairing  
detailed in Section 2.1), between 9<sup>th</sup> November and 13<sup>th</sup> December 2018, as per Table 1. The air inlet for each Optical Particle  
190 Counter was orientated towards the south-west in order to face the predominant wind direction for the site (Hart et al., 2019).  
The air flows through the instrument at a rate of 5.5 L/min.

Two Tinytag Plus 2 TGP-4500 (Gemini data loggers, Chichester, UK) units were used to measure RH and temperature  
alongside the low-cost OPCs. The units measure temperature from -25°C to +85°C using an internally mounted thermistor,  
with the manufacturers stating an accuracy of 0.01°C or better. Under field conditions, the TinyTag sensors closely follow  
195 temperature measurements taken by a weather station located at 2 m height in Array 1, with the mean temperature measured  
by the TinyTags measuring within 0.4°C of the weather station. RH is measured using an externally mounted capacitive sensor,  
from 0–100% RH, with a manufacturer stated accuracy of +/- 3.0% RH at 25°C. Tinytag units were placed on the top of the  
OPCs, with the RH sensor facing in the same direction as the OPC inlet.

Above-canopy wind speed was measured using two-D ultrasonic anemometers (WMT700, Vaisala) ~1 m above the canopy  
200 (~25 m height) on the northernmost tower of array 1. Below-canopy wind speed and direction (~2 m height) were measured  
using 03002-L Wind Sentry set (Campbell Scientific, Loughborough, UK) located to the right of the metal walkway near the  
entrance of Array 1. The manufacturer stated minimum threshold wind speed was 0.8 m s<sup>-1</sup> for the Campbell anemometer,  
with 2 m height wind speeds below this minimum threshold considered as zero for the purposes of this experiment.

205 **Table 1. Table listing the dates (dd.mm.yyyy) and locations of equipment installation across the experimental duration at BIFoR  
FACE. Both optical particle counters (OPCs) and TinyTag relative humidity and temperature sensors were installed simultaneously.**

Date range (All dates in 2018)	Tiny Tag unit	OPC unit	Array number	eCO <sub>2</sub> or Ambient
9 <sup>th</sup> Nov to 21 <sup>st</sup> Nov	A	1	A2	Ambient
	B	2	A4	eCO <sub>2</sub>

Deleted:

Deleted: Table 1

Deleted: Error! Reference source not found.

21 <sup>st</sup> Nov to 30 <sup>th</sup> Nov	A	1	A6	eCO <sub>2</sub>
	B	2	A5	Ambient
5 <sup>th</sup> Dec to 13 <sup>th</sup> Dec	A	1	A1	eCO <sub>2</sub>
	B	2	A3	Ambient

210

Both the pair of OPCs and TinyTags were installed for a six day side-by-side intercomparison period at 1.5 m height in Array 1 of BIFoR FACE from 30<sup>th</sup> November 2018 until 05<sup>th</sup> December 2018. Using the data from this intercomparison week, bias calibration factors were calculated.

215 No bias factor was applied to the temperature data from the TinyTag units. RH data from the TinyTags was first filtered to remove any measurements lower than 50% and greater than 99%, and a calibration bias factor of 1.03 applied to TinyTag Unit B to make the two units consistent with each other. A similar calibration was performed on the particulate matter (PM) data collected by the OPCs, with the data from OPC 1 being increased by a bias factor of 1.45. Bias factors were calculated from the six-day side-by-side period, with the factor applied to the data from the full measurement period of 37 days. This bias correction ensured the two OPCs were consistent with each other.

220 For both the TinyTag and OPC pairs, the consistency between the two instruments, rather than absolute calibration, was more important to allow for observation of differences between two different locations. Hence one instrument of each pair was bias-corrected to the other instrument of the pair. However, a comparison outside of the forest environment was completed between the low-cost OPCs and a TSI 3330 (a reference grade OPC), with the low-cost OPCs measuring within 25% of the TSI without any humidity corrections, which is considered a good performance for the low-cost sensors (Crilley et al., 2020, 2018).

#### 225 2.4 Spore dispersal model

We developed a conceptual model to help interpret the OPC results from BIFoR FACE. Within dense forests, the mean horizontal wind speed  $U$  inflects around the tops of the trees, increasing approximately exponentially with height below the inflection and logarithmically with height above it (Bannister et al., 2021; Cionco, 1965; Finnigan, 2000; Raupach et al., 1996). Mean horizontal wind velocity  $U$  at each height  $z$  inside the canopy can, therefore, be approximated via Eq. (2):

230 
$$U(z) = U_{h_c} e^{-a(1-\frac{z}{h_c})} \quad (2)$$

where  $h_c$  is the mean height of the trees and  $U_{h_c}$  is  $U$  at  $h_c$ . The attenuation coefficient,  $a$ , accounts for the flow's response to the density of the forest and tends to increase with increasing leaf area index (LAI) and flexibility of the plant elements (Cionco, 1978; Kaimal and Finnigan, 1994) The exponential relationship in Eq. (2) does not hold in forests with a sparse trunk space and open edges, e.g. many pine plantations, for which a secondary wind velocity maximum occurs below the main crown  
235 (Baldocchi and Meyers, 1988). However, it provides a reasonable first approximation for a forest such as that at the BIFoR FACE site, with extensive understorey growth and closed edges during the leaf-on season.



We adapt the model of Nathan et al., 2002 by using this exponential velocity profile to obtain a rough estimate of the horizontal distance over which spores disperse in a dense forest,  $D$ . We consider spores evenly distributed within small air parcels within the canopy airspace.  $D$  equals the distance an air parcel carrying spores travels between the time of release ( $t_0 = 0$ ) and the time at which the spores in the parcel settle on the ground ( $t_1$ ). As a simplification, here we consider spore ‘release’ to include both detachment by the mean wind and the point at which spores begin to settle back to the ground after being swept upwards by short intense gusts (Aylor, 1978). For a spore falling at an average settling velocity  $v_s$ ,  $t_1 = h_r/v_s$ , where  $h_r$  is the height of spore release. The vertical position of spores within an air parcel during settling at time  $t$  is  $z(t) = h_r - v_s t$ . Substituting this expression into Eq. (2), we generate Eq. (3):

$$U(t) = U_{hc} e^{-\frac{a}{h_c}(h_c - h_r + v_s t)} \quad (3)$$

The horizontal distance over which the spores disperse is therefore:

$$D = \int_0^{t_1} U(t) dt \quad (4a)$$

which, taking  $t_1 = h_r/v_s$  and  $U(t)$  from Eq. (3), gives:

$$D = \frac{U_{hc} \cdot h_c}{av_s} \left[ e^{-a} \left( e^{\frac{ah_r}{h_c}} - 1 \right) \right] \quad (4b)$$

This model assumes the mean wind profile has already adjusted to the presence of the forest, for example, after passing into the forest from surrounding rural areas (see below).

We specified the mean height of the trees as 25 m to reflect those at the BIFoR FACE site. Reported values of  $a$  for forests fall within a relatively narrow range of around 1.5–4.8 (Amiro, 1990; Cionco, 1978; Su et al., 1998). We took  $a = 2.5$ , as measured for forests of similar LAI to BIFoR FACE (Su et al., 1998). Sensitivity testing (not shown) indicated that using higher and lower values of  $a$  respectively decreased and increased the occurrence of spores travelling long distances, but had little effect on the modal value of  $D$ . i.e. using a lower  $a$  thickened the right tail of the probability density function of  $D$  but affected its peak relatively little.

We performed a stochastic ensemble of model runs ( $n = 1,000,000$ ) in R (version 4.0.3, <http://cran.r-project.org>) with low (0–2 m s<sup>-1</sup>), low-medium (*med-l*) (2–3 m s<sup>-1</sup>), high-medium (*med-h*) (3–4 m s<sup>-1</sup>), and high (4–6 m s<sup>-1</sup>) mean wind speeds at the top of the canopy,  $U_{hc}$ . We considered only spores in the bottom 10 m of the forest airspace, i.e.  $h_r$  in the range 0–10 m. Strong turbulent fluxes of momentum, and scalar quantities such as spores, occur in the upper region of forest canopies (Belcher et al., 2012; Finnigan, 2000), which this simplified model cannot capture. Mean settling velocity,  $v_s$ , was specified in the range 0.001–0.005 m s<sup>-1</sup>, taking reported values for fungal spores in the size range 1–10 μm (Di-Giovanni et al., 1995; Tesmer and Schnittler, 2007). For each model run, for we specified a random value of  $U_{hc}$ ,  $h_r$  and  $v_s$  from a uniform distribution within each of these ranges (using the *runif* function in R).

## 2.5 Comparison of measured bioaerosol proxy with regional scale data from Copernicus Atmosphere Monitoring Service (CAMS) global reanalysis (EAC4) data

270 To test the assumption that PM<sub>10</sub>-PM<sub>1</sub> mass concentration is representative of bioaerosols, the measured PM<sub>10</sub>-PM<sub>1</sub> time series  
is compared with Copernicus Atmosphere Monitoring Service (CAMS) global reanalysis (EAC4) data as a measure of regional  
aerosol activity influencing the study site. The Copernicus Atmosphere Monitoring Service (CAMS) reanalysis is the latest  
global reanalysis dataset of atmospheric composition produced by the European Centre for Medium-Range Weather Forecasts  
(ECMWF), consisting of three-dimensional time-consistent atmospheric composition fields, including aerosols and chemical  
species (Inness et al., 2019). The nearest available grid point from the EAC4 dataset (longitude = -2.50, latitude = 52.75) to  
275 the BIFoR site (longitude = -2.32, latitude = 52.81) and use the same time period as the Ambient and eCO<sub>2</sub> measurements.  
The spatial resolution of the reanalysis data is 80 km. The PM<sub>10</sub>-PM<sub>1</sub> metric is calculated from the EAC4 estimates of PM<sub>10</sub>  
and PM<sub>1</sub>. In addition to the EAC4 PM<sub>10</sub>-PM<sub>1</sub> estimate, we also use the aerosol optical depth (AOD) estimates from EAC4,  
including total AOD, the AOD attributed to dust aerosol, the AOD attributed to sea salt aerosol, and the AOD attributed to  
sulphate aerosol (Benedetti et al., 2009; Morcrette et al., 2009). All AODs were for the wavelength of 550 nm.

280

## 2.6 Data processing, visualisation, and analysis

All data analysis was completed in R version 4.0.3 (R Core Team, 2020), with figures created using *openair* and *ggplot*  
(Carslaw and Ropkins, 2012; Wickham, 2016). Relationships between PM<sub>10</sub>-PM<sub>1</sub> concentrations and RH, temperature, and  
wind speed (Figure 2, Figure 3, and Figure 6) were visualised using scatter plots and smoothed loess curves, generated in *ggplot*.  
285 Box plots with mean PM<sub>10</sub>-PM<sub>1</sub> (and interquartile ranges) were generated to visualise differences in bioaerosol concentrations  
between eCO<sub>2</sub> and ambient arrays. Scatter plots with regression lines were generated for Figure 7 and for the supplementary  
figures.  
Data was non-normally distributed, therefore Kruskal-Wallis tests were used to test for statistically significant differences  
between the eCO<sub>2</sub> and Ambient arrays in Section 3.2.

290

## 3 Results

### 3.1 Hypothesis A: Hourly fungal bioaerosol concentrations will correlate with hourly weather conditions (wind, RH, temperature).

During the experimental period, there was a total of 46.3 mm of precipitation, with a median temperature of 7.7°C within the  
295 forest. The lowest recorded temperature was -1.8°C, and the highest 15.8°C. The average daily temperature peak (and the  
lowest RH point) occurred at approximately 13:00 local time. RH throughout the measurement period was typically high, with

Deleted: Figure 3

the median and mean measured between 89–91% RH across the duration of the experiment, with the lowest recorded measurement at 63% RH. Wind speed patterns followed each other closely above and below the canopy, with below canopy measurements significantly lower. The mean wind speed above the canopy was  $2.7 \text{ m s}^{-1}$ , whereas below the canopy it was  $0.4 \text{ m s}^{-1}$ .

Figure 2 Panel A shows the relationship between RH and bioaerosol concentrations at BIFoR FACE: the curve is almost horizontal from 75–90% RH, where at 90% it rapidly increases, most likely indicating the particles becoming exponentially hygroscopic at very high humidities. This sudden increase in particle size at very high humidities demonstrates a clear hygroscopic effect; we therefore performed a correction factor to the data from Crilley et al., 2018, using a  $\kappa$  values of either 0.3 or 0.1 (Griffiths et al., 2012; Pope, 2010). Before correction, the median hourly  $\text{PM}_{10}$ – $\text{PM}_1$  concentration under  $\text{eCO}_2$  conditions was  $15.7 \mu\text{g m}^{-3}$  and  $16.7 \mu\text{g m}^{-3}$  under ambient conditions (Figure 2 Panel A). After the application of the Crilley et al. correction factor using a  $\kappa$  of 0.3, this decreased to a median of  $5.3 \mu\text{g m}^{-3}$  under  $\text{eCO}_2$  and  $5.9 \mu\text{g m}^{-3}$  at ambient  $\text{CO}_2$  concentrations (Figure 2 Panel B), and using a  $\kappa$  of 0.1, the median was  $9.3 \mu\text{g m}^{-3}$  under  $\text{eCO}_2$ , and  $10.1 \mu\text{g m}^{-3}$  under ambient conditions (Figure 2 Panel C). The ratio between the mean concentrations obtained from the  $\text{eCO}_2$  and ambient plots depend upon whether the Crilley correction factor is used and what value is used for  $\kappa$ . The values are 94%, 90%, and 92% for the uncorrected ratio, the ratio using  $\kappa = 0.3$  and the ratio using  $\kappa = 0.1$ , respectively. A comparison of the data before and after correction is shown in Figure 2 panels D, E, and F. Due to the decrease in PM concentrations, with respect to RH, shown using a  $\kappa$  of 0.3, the data corrected using a  $\kappa$  of 0.1 was used for all further analyses, as this was deemed more appropriate given the likely particle composition (low hygroscopicity bioaerosols). After correction to remove the instrument artefacts, there was no clear effect of RH on bioaerosol concentrations. If the observed median concentrations of  $\text{PM}_{10}$ – $\text{PM}_1$ , using the Crilley correction with a  $\kappa$  of 0.1, were solely composed of idealized spherical fungal spores with radius of  $3 \mu\text{m}$  and density of  $1.65 \text{ g/ml}$  this would equate to spore concentration of approximately  $50,000 \text{ spores m}^{-3}$ , which falls at the high end of Sesartic and Dallafor (2011b) estimate of ground level spore concentrations of  $10,000$ – $50,000 \text{ m}^{-3}$ .


Deleted: Error! Reference source not found.

Deleted: Error! Reference source not found.

Deleted: Error! Reference source not found.

Deleted: Error! Reference source not found.

Deleted: Error! Reference source not found.



325 **Figure 2. Relative humidity and bioaerosol concentrations ( $PM_{10}-PM_1$ ) at BIFoR FACE in Autumn 2018. Panel A shows  $PM_{10}-PM_1$  concentrations with increasing RH without any correction applied. Panels B and C plots the same data as Panel A, however a correction factor from Crilley et al., 2018 has been applied to the bioaerosols data to correct for the hygroscopic effect of the particles, Panel B using a  $\kappa$  of 0.3, and Panel C a  $\kappa$  of 0.1. Panels A, B, and C, all have loess curves fitted (Wickham et al., 2019).  
 330 Panels D, E, and F compare the data before and after corrections. All the plots show  $PM_{10}-PM_1$  concentrations under ambient conditions (Arrays 2, 3, and 5) in blue colours, and  $eCO_2$  conditions (Arrays 1, 4, and 6) in orange colours. Data pre-correction is shown in dark orange and blue, corrected data using a  $\kappa$  of 0.3 mid-blue and mid-orange, and the corrected data using a  $\kappa$  of 0.1 uses the lightest blue and orange. All panels use hourly averages of  $PM_{10}-PM_1$  concentrations.**

335 The effects of wind speed and direction on the bioaerosol concentrations were also investigated in depth. Anemometers were located both above (25 m height) and below (2 m height) the canopy, with below-canopy measurements broadly following those above the canopy but measuring significantly lower at the lower height (Figure 3, Panel A). Above the canopy, the hourly wind speed never dropped below  $0.8 \text{ m s}^{-1}$ , whereas recorded speeds of  $0 \text{ m s}^{-1}$  (i.e. below anemometer threshold) were common below canopy. The maximum hourly wind speed above canopy was  $7.05 \text{ m s}^{-1}$ , compared with  $2.59 \text{ m s}^{-1}$  below canopy.

340 Bioaerosol concentrations demonstrated high variability under changing wind speeds; however, a possible U-shaped curve was observed, with higher bioaerosol concentrations at the lowest and highest wind speeds measured, as shown in Figure 3, Panels B and C. This effect was shown using the wind speeds taken from both above (25m) and below (2m) the canopy. Figure

Deleted: Figure 3

Deleted: Error! Reference source not found.

Deleted: Figure 3

Deleted: Error! Reference source not found.


4 shows the effect of wind speed and direction on bioaerosol concentrations using *openair* polar bivariate plots (Carslaw and Ropkins, 2012). Observed wind speed patterns follow broadly similar trends between each pair of arrays, with the majority of medium concentration (medium concentration (green-yellow) bioaerosol activity occurring in the SW and SE quadrants, and only low (blue) concentrations under northerly winds. Peaks in bioaerosol concentration (presumed high sporulation events) are visible in red, with some events being replicated across both eCO<sub>2</sub> treatment and control (e.g. the SW quadrant event in A5 and A6), and other high PM events only occurring in a single array (e.g. the SE quadrant event in A4). By detecting high PM events in a single array at a distinct time shows that the OPCs can detect differences between the BIFoR FACE arrays.

To investigate the relationship between spore dispersal and wind speed further, we used the model outlined in Section 2.4 to investigate the horizontal distance over which spores disperse,  $D$ . The model was run ( $n = 1,000,000$ ) for spore release heights ( $h_r$ ) of 2 m, 5 m, and 10 m. Figure 5 shows the probability density functions of  $D$  under four wind speed scenarios: low, low-medium, medium-high, and high. As would be expected, generally, the lower the spore release height and wind speed, the shorter the overall distance travelled by spores. Most wind conditions experienced at BIFoR FACE fall under the “low” wind speed scenario (black lines), for which the modal distance travelled by spores were less than 20 m for all release heights (Figure 5). With the arrays having diameters of 25–30 m, this means that under typical conditions, we could expect spores to stay within the array they were released in or, at least, very unlikely to be transported into neighbouring arrays (i.e. a spore released under eCO<sub>2</sub> is unlikely to be measurable in an ambient array in low wind conditions). However, at higher wind speeds, a heavy tail is present on each of the plots, indicating that spores are more likely to travel much greater distances at higher wind speeds, potentially causing mixing between the arrays. Finally, we looked at PM<sub>10</sub>-PM<sub>1</sub> concentrations under changing temperature (Figure 6). There was a small positive relationship between temperature and bioaerosol concentrations, with PM<sub>10</sub>-PM<sub>1</sub> concentrations of around 5  $\mu\text{g m}^{-3}$  at the lowest temperatures (1-3 °C), increasing up to 10  $\mu\text{g m}^{-3}$  at the highest temperatures of 12-13 °C.

Deleted: Figure 4

370


Deleted: linear



375


380 *Figure 3. Wind speed and bioaerosol concentrations at BIFoR FACE in Autumn 2018. Panel A shows the hourly average wind speeds above (25 m height) and below (2 m height) the canopy across the experimental duration. Wind speed data at 25 m height is in dark green, and data from 2 m height in light green. Panels B compares hourly average  $PM_{10}$ - $PM_1$  concentrations with hourly average wind speeds measured at 25 m height, and Panel C the same  $PM_{10}$ - $PM_1$  concentrations using wind speeds measured at 2 m height. Panels B and C both have loess lines of fit (Wickham et al., 2019), with data from ambient arrays (Arrays 2, 3, and 5) in blue, and data from  $eCO_2$  arrays (Arrays 1, 4, and 6) in orange.*

385



390 Figure 4. Openair polar bivariate plots displaying bioaerosol concentrations ( $PM_{10}-PM_1$ ), wind speed, and wind direction across the  
 395 Autumn 2018 experimental duration at BIFoR FACE (Carslaw and Ropkins, 2012). Each row of plots shows the data collected in  
 each of the pairs of arrays, one eCO<sub>2</sub>, one ambient) in which the pair of OPCs were located. Dates of OPC installation in each array  
 pair are detailed in [Error! Reference source not found.](#) Bioaerosols were measured using OPCs at 2 m height, and wind data was  
 taken from anemometers at 25 m height. Colour gradients display the concentrations of bioaerosols detected, with low  
 concentrations shown in blue/green colours, mid concentrations in yellow, and high concentrations in orange/red.



- Deleted: showing
- Deleted: with
- Deleted: Plots are displayed in the three array pairs (
- Deleted: Table 1Error! Reference source not found.
- Deleted: ing
- Deleted: the
- Deleted: the



405 **Figure 5. Model outputs of probability density functions for distance travelled by spores in a forested environment. Panels A, B, and C had different spore release heights ( $h_r$ ) inputted into the model- 2, 5, and 10 m respectively. Solid lines are in different colours for each of the low (black), low-medium(blue), medium-high(yellow) and high (red)  $U_{bc}$  (wind speed) cases. Dashed lines show modal D (distance) travelled by the spores.  $n = 1,000,000$  for each case.**

410





Moved down [2]:

Moved (insertion) [2]

Figure 6. Temperature and bioaerosol (PM<sub>10</sub>-PM<sub>1</sub>) concentrations across the experimental duration at BIFoR FACE. Panel A shows temperature data, with mean temperature measured by TinyTag sensors shown in dark green, and light green showing data collected by the Array 1 weather station. Panel B compares temperature data with bioaerosol concentrations, with data from ambient arrays (Arrays 2,3, and 5) in blue, and data from eCO<sub>2</sub> arrays (Arrays 1,4, and 6) in orange, with both treatment types fitted with a loess line (Wickham et al., 2019).

### 3.2 Hypothesis B: Fungal bioaerosol concentrations will increase in woodland patches treated with eCO<sub>2</sub> for two years.

Figure 7 shows six plots comparing PM<sub>10</sub>-PM<sub>1</sub> concentrations between the eCO<sub>2</sub> (orange) and ambient (blue) arrays. When comparing total concentrations across the entire measurement period (Figure 7, Panels A, B, and C), there was no significant difference ( $p = 0.489$ ) between the eCO<sub>2</sub> treatment and ambient, however a heavy tail is present in the ambient arrays. To investigate this relationship further, we separated out the data into low bioaerosol concentration conditions ( $<10 \mu\text{g m}^{-3}$ ) and high bioaerosol concentrations ( $\geq 10 \mu\text{g m}^{-3}$ ) (Figure 7, Panels D, E, and F). When bioaerosol concentration are low (Figure 7, Panels D and E), there was no significant difference between the eCO<sub>2</sub> and ambient treatments ( $p = 0.689$ ). These low concentrations likely represent background levels of aerosols that are consistent throughout the forest. However, when selecting for high concentration events, there was a significant effect seen, whereby eCO<sub>2</sub> treatment suppressed concentrations of high bioaerosol events ( $p = 0.023$ ) (Figure 7, Panels D and F).

Deleted: Figure 7

Deleted: Figure 7

Deleted: Figure 7


Deleted: Figure 7

Deleted: Error! Reference source not found.

Deleted: Figure 7

The boxplot in Panel A shows that there is no significant difference between the medians of eCO<sub>2</sub> treatment versus the control; however, there were slightly higher bioaerosol concentrations shown under ambient conditions. Panel B shows a time series of the data, demonstrating that bioaerosol concentrations match extremely closely between the two treatment groups. The small

440 difference in the overall medians can be largely attributed to the  $PM_{10}$ - $PM_1$  concentration difference between the  $eCO_2$  and ambient during the largest bioaerosol event around the 22<sup>nd</sup> of November (during the second measurement period, shown shaded in grey), where the ambient array (shown in blue) measures higher than the  $eCO_2$  array in orange.





445 **Figure 7. Bioaerosol concentrations across the experimental duration in ambient (Arrays 2,3, and 5) and  $eCO_2$  (Arrays 1, 4, and 6) arrays.  $eCO_2$  treatment is shown in orange, ambient control in blue. Panels A, B, and C include the full dataset, with no significant difference shown between the  $eCO_2$  treatment and control. Panels D, E, and F split the dataset into low ( $<10 \mu g m^{-3}$ ) and high ( $\geq 10 \mu g m^{-3}$ ) bioaerosol concentrations. When the data is split in this way, there was no significant difference between treatments under low bioaerosol concentrations. However, when bioaerosol concentrations were high, concentrations were significantly lower under  $eCO_2$  treatment.**

### 450 3.3 Comparison of the measured bioaerosol proxy with regional scale data from Copernicus Atmosphere Monitoring Service (CAM5) global reanalysis (EAC4) data

455 It can be seen in Figure 8 that the EAC4 estimate of regional  $PM_{10}$ - $PM_1$  does not correlate with the  $PM_{10}$ - $PM_1$  measured within the forest. Hence, we believe the measured  $PM_{10}$ - $PM_1$  is local to woodland and not representative of the regional air mass. The Pearson correlation between the measured  $PM_{10}$ - $PM_1$  in the Ambient and  $eCO_2$  arrays and the EAC4 estimates is **-0.03** and **0.03**, respectively, highlighting negligible correlation between the measured and EAC4 timeseries. To look for further possible correlations between the measured  $PM_{10}$ - $PM_1$  and estimated aerosol properties, correlative analysis was conducted on the measured  $PM_{10}$ - $PM_1$  with the aerosol optical depth (AOD) estimates from EAC4 consisting of the total AOD, the AOD

attributed to dust aerosol, the AOD attributed to sea salt aerosol, and the AOD attributed to sulphate aerosol. Similar to the EAC4  $PM_{10}$ - $PM_{1}$  product, there were no non-negligible correlations with the AOD products, with Pearson correlation values ranging from  $-0.16$  to  $0.23$ . See supplementary material for more information the correlation plots provided in Figures S1 to

460 S11.



Moved down [1]:

Moved (insertion) [1]

465 **Figure 8.** Time series data of the  $PM_{10}$ - $PM_{1}$  mass concentrations measured in the Ambient and  $eCO_2$  arrays and provided by the Copernicus Atmosphere Monitoring Service (CAMS) global reanalysis (EAC4) dataset. The BIFoR data contains only data collected when  $RH < 99\%$ .

#### 4. Discussion

470 In support of hypothesis A, we found that various meteorological variables affected bioaerosol concentrations. The RH before correction was of particular importance, especially as the low-cost OPCs do not have any warming or drying capacity, and therefore are susceptible to recording higher mass concentrations of aerosols under high humidity conditions as particles swell with additional water (Crilley et al., 2018). We found that the RH % threshold for significant particle swelling was 90–95%,

475 which is a much higher RH value than would be expected for anthropogenic aerosols, which typically contain more  
hygroscopic components including salts, and therefore provides evidence that the measured  $PM_{10}$ – $PM_1$  fraction represents a  
predominantly biological source. In their study investigating effects of RH on fungal spore swelling, Reponen et al. (1996)  
demonstrated a similar effect whereby a significant swelling of fungal spores was seen, but only at very high humidities (greater  
than 90%). We believe this threshold for particle swelling further demonstrates that we are recording a biological source. This  
hygroscopic evidence is in addition to the ecological and phenological evidence for spores being the dominant source within  
480 the  $PM_{10}$ – $PM_1$  size fraction during the measurement period.

After correction, we did not see any effect of RH on fungal spore concentrations. The evidence in the literature for the  
relationship between spore concentrations and RH is mixed, probably largely due to species-specific effect. Some fungal  
species are dependent on high RH to develop spores (and then release spores by rain droplets), whereas other species release  
spores in response to a drop in RH (Jones and Harrison, 2004; Li and Kendrick, 1995). The high diversity of fungal species at  
485 BIFoR (each with possibly varying responses to RH) could result in this flat line. Another possible explanation is that the fungi  
never experience low RH conditions due to the constant high RH at BIFoR FACE, potentially with the RH never dropping  
low enough to trigger a burst of spore release. Finally, it is important to mention that although the Crilley correction is an  
essential part of using low-cost sensors in a high-humidity environment, the high swelling threshold of fungal spores means  
that the  $\kappa$  value of 0.3 (or even 0.1) may be inappropriately high for biological particles such as these. As shown in Figure  
490 **2Error! Reference source not found.** Panel B and C, after correcting the data, the  $PM_{10}$ – $PM_1$  concentrations decrease after  
95% RH, although this effect is smaller when using a  $\kappa$  of 0.1. For future studies, calibrating the low-cost OPCs against a  
reference grade instrument in the experimental environment in order to provide a calibrated  $\kappa$ -value would be an appropriate  
measure, and enable a more thorough investigation of the RH dependence, and improve the accuracy of the Crilley correction  
for bioaerosols (Crilley et al., 2020).

495 We measured an approximately U-shaped curve of bioaerosol concentrations in response to wind speed, with the highest  $PM_{10}$ –  
 $PM_1$  concentrations being displayed at low and high wind speeds. This likely indicates a balance between spore release  
mechanisms and mixing of low-aerosol air. At low wind speeds, there is minimal movement of air through the forest, so any  
spores released do not travel significant distances. At high wind speeds, there is much higher movement of air, which decreases  
concentrations by carrying spores further; however, spore release by wind could also be increased (Dressaire et al., 2016). At  
500 medium wind speeds, a balance between these two effects occurs, maintaining bioaerosol concentrations at a lower level (Jones  
and Harrison, 2004).

Our model demonstrated that under low wind-speed conditions, spores are most likely to travel a relatively short distance,  
suggesting that between-array mixing was unlikely under the meteorological conditions experienced during the measurement  
period. Particle release height ( $h_r$ ) had a significant impact on the distance travelled by a particle, which is an important  
505 consideration given the source of bioaerosols. For example, fungal spores are likely to have a significantly lower  $h_r$  than pollen  
which, based on our model, would significantly impact the distance travelled by the different bioaerosol types. Due to the  
complexity of the forest environment, there are a number of aspects which the model did not capture which could alter the

Deleted: Error! Reference source not found.

distance travelled by a spore. Complex wind dynamics such as ejections (infrequent but strong upward gusts of air) and  
510 turbulence around forest edges could increase the distance travelled by a spore, as well as varied spore release dynamics  
between fungi (Dressaire et al., 2016; Norros et al., 2014). The model also assumes a spatially representative wind speed  
profile, which does not account for local velocity effects induced by gaps, clearings, and changes in canopy density. These  
local effects may be important near the ground, where wind speeds are generally low. However, the distance travelled could  
also be shorter than modelled, for example, the model does not capture spore deposition onto forest surfaces other than the  
515 ground, spores being swept downwards by gusts, and changes with wet versus dry deposition.

The data demonstrated a decrease in bioaerosol concentrations at lower temperatures, which is the expected response within  
the range of temperatures we observed in a woodland of this type (Gange et al., 2007). The number of fungal sporocarps  
present across the duration of the experiment decreased, and it is therefore logical to expect that spore production would also  
consequently decrease. For future studies, it would be interesting to begin measurements earlier in the fungal season, allowing  
520 us to capture the peak of sporocarp production, as well as the end of the active eCO<sub>2</sub> fumigation at the FACE experiment.

Regarding Hypothesis B, to our knowledge, this is the first assessment of bioaerosols in any forest FACE experiment to date,  
and therefore provides valuable data contributing to the understanding of the forest environment at BIFoR FACE as well as  
more broadly in the context of forests, and FACE experiments. We did not see a significant effect of eCO<sub>2</sub> on total aerosol  
concentrations. However, when data was split into low and high aerosol concentrations, two differing responses were  
525 demonstrated. Under low concentrations, there was no significant effect of eCO<sub>2</sub> treatment, we therefore think these low  
concentrations represent the background concentrations of aerosols present both inside and in the wider environment around  
the forest. However, when high concentrations of bioaerosols were detected, eCO<sub>2</sub> treatment resulted in a suppression of PM<sub>10</sub>-  
PM<sub>1</sub> concentrations. This is demonstrated in a number of larger sporulation events during the experimental period, for example  
the large sporulation event peaking on 22<sup>nd</sup> November, where although both arrays showed the sporulation event, the magnitude  
530 of this event was significantly different between eCO<sub>2</sub> and ambient.

This evidence, as well as the eCO<sub>2</sub> dispersal data from Hart et al., 2019, and the modal spore travel distances shown in our  
model suggests to us that this experimental set-up is capable of detecting differences between the eCO<sub>2</sub> and ambient arrays. It  
is apparent that at background concentrations these differences are not present between arrays, however there may be a  
535 response to eCO<sub>2</sub> during large sporulation events. This could be for a number of reasons, the primary reason being that the  
experiment was in the very early stages of fumigation (year two of ten total years planned), and there might be a lag expected  
in fungal responses, as they are most likely responding indirectly to eCO<sub>2</sub> via changes in leaf litter and soil. Although the  
literature states a small number of cases of where individual fungal species do respond directly to eCO<sub>2</sub>, it is likely that the  
main effects would be secondary, for example, competition between autotrophs and heterotrophs for nutrients, or an increase  
540 in leaf litter production resulting in an increase the population of decomposer fungi. There is evidence for increased autotrophic  
productivity under eCO<sub>2</sub> at BIFoR FACE in the form of increased leaf-scale photosynthesis (Gardner et al., 2020) and fine  
root production (Ziegler et al., 2021). Whether this increased autotrophic productivity primes, or competes with, fungal activity

requires further work. Continuing to monitor bioaerosol concentrations throughout the 10+ year experimental duration of the BIFoR FACE experiment, along with monitoring in other FACE experiments, will be key in understanding how eCO<sub>2</sub> affects bioaerosol concentrations long term.

Another possible reason for not picking up an eCO<sub>2</sub> response in fungal bioaerosols in the full dataset is that the maximum diameter of the particles that can be detected by the OPCs is 10 μm, which does exclude several fungal species, including those known to be present in the BIFoR forest during the experimental duration. However, we note that many common woodland spore species are smaller than 10 μm, including the following species observed in the forest: *Lactarius quietus*, *Russula* species, *Lycoperdon perlatum*, *Mycena rosea* other *Mycena* species, *Auricularia auricula*, *Hypholoma fasciculare*, *Xylaria hypoxylon*, and various *Stereum* species. In addition to the observed species, many spores commonly observed to be airborne in the UK, including *Cladosporium*, *Ganoderma*, and *Aspergillus* species (Sadyś et al., 2016), have diameters less than 10 μm. If a response to eCO<sub>2</sub> is species-specific, then it is possible that we are missing an effect in fungi with larger spores. It is also possible that although the total and background aerosol concentration was stable under eCO<sub>2</sub>, the aerosol composition could have been different (e.g. altered ratios of fungal species present), which we were not able to detect using the OPCs. Varying fungal species between the ambient and eCO<sub>2</sub> arrays could therefore be responsible for the differing response we detected during sporulation events. In order to determine definitive particle composition, other techniques such as detectors using fluorescence, or DNA sequencing of biological material is required (Fröhlich-Nowoisky et al., 2016; Gosselin et al., 2016; Healy et al., 2012). Finally, our experimental measurement period occurred outside of the eCO<sub>2</sub> fumigation season, and therefore the main growing season of the trees, which could have reduced any direct effects of the eCO<sub>2</sub>, although, given the likely cumulative effect of the eCO<sub>2</sub> treatment over the growing season, and the fact that our measurements were focused on the period of peak sporulation, it seems more likely that our measurements were well-timed to observe any treatment effects.

## 5 Conclusions

We have demonstrated that low-cost OPCs are suitable for measuring PM<sub>10</sub>-PM<sub>1</sub> concentrations in forests, or other high-humidity environments. We demonstrate that the PM<sub>10</sub>-PM<sub>1</sub> metric is a good proxy for bioaerosols because of the bioaerosol representative size range, the location of the study site (a woodland in a rural location), the field measurement taking place during the season of peak fungal activity, and the low hygroscopicity of the particles measured. Through comparison with the EAC4 estimate of PM<sub>10</sub>-PM<sub>1</sub> we highlight that the woodland measurements do not follow regional air pollution trends and that the observed PM<sub>10</sub>-PM<sub>1</sub> concentrations are likely from woodland sources. The findings from this study have consequences for other research into the aerobiology of forests, and also opens up the bioaerosol research field to a wider array of locations and researchers. The low-cost sensors measured significant swelling in PM<sub>10</sub>-PM<sub>1</sub> sized particles at very high RH, which we corrected for using a calibration factor from Crilley et al. (2018). For future work, generating a κ value using a reference-grade instrument in situ would improve accuracy (Crilley et al., 2020). Temperature, wind speed, and wind direction were all shown to affect bioaerosol concentrations; however, we did not see any effect of RH (post-correction). Treatment with eCO<sub>2</sub> may

575 repress concentrations of bioaerosols during high sporulation events ( $p = 0.023$ ), however this significant difference was not  
detected across all concentrations of aerosols ( $p = 0.489$ ). Therefore, further investigation later in the 10+ year experimental  
duration is warranted, as well as investigating forest bioaerosols in other forest FACE experiments globally. The use of the  
PM<sub>10</sub>-PM<sub>1</sub> metric as a proxy for bioaerosols, in woodland and other settings, should be further evaluated through future  
experiments that unambiguously measure bioaerosol concentrations.

#### 580 **6 Data availability**

Data supporting this publication are openly available from the UBIRA eData repository at  
<https://doi.org/10.25500/edata.bham.00000688>. The data provided is the hourly data with the inter-unit calibration bias factor  
applied.

#### 585 **7 Author contribution**

ABB and FDP designed the bioaerosol study as part of the FACE programme designed by ARMK. ABB collected and analysed  
the OPC data and prepared the manuscript. EJB wrote the model. All co-authors discussed the results and contributed to writing  
the manuscript.

#### **8 Competing interests**

590 The authors declare that they have no conflicts of interest.

#### **9 Acknowledgements**

The BIFoR FACE facility is a research infrastructure project supported by the JABBS Foundation and the University of  
Birmingham. ABB is supported by a Natural Environment Research Council (NERC) studentship through the DREAM  
programme (grant NE/M009009/1), EJB is supported by a NERC studentship through the CENTA programme (grant  
595 NE/L002493/1). ARMK gratefully acknowledges support from NERC through grants NE/S015833/1 and NE/S002189/1. FDP  
gratefully acknowledges support from EPSRC through grant EP/T030100/1. With thanks to the British Pteridological Society  
for their literature recommendations and advice on fern phenology.

## 10 References

- 600 Amiro, B.D., 1990. Comparison of turbulence statistics within three boreal forest canopies. *Boundary-Layer Meteorology* 51, 99–121. <https://doi.org/10.1007/BF00120463>
- an de Hulst, H.C., 1957. *Light scattering by small particles*. John Wiley and Sons, New York.
- Andrew, C., Lilleskov, E.A., 2009. Productivity and community structure of ectomycorrhizal fungal sporocarps under increased atmospheric CO<sub>2</sub> and O<sub>3</sub>. *Ecology Letters* 12, 813–822. <https://doi.org/10.1111/j.1461-0248.2009.01334.x>
- 605 Aylor, D.E., 1978. Dispersal in Time and Space: Aerial Pathogens, in: Horsfall, J.G., Cowling, E.B. (Eds.), *Plant Disease: An Advanced Treatise*. Academic Press, New York, pp. 159–179. [https://doi.org/10.1016/0048-3575\(79\)90101-9](https://doi.org/10.1016/0048-3575(79)90101-9)
- Baird, A.B., Pope, F.D., 2021. “Can’t see the forest for the trees”: The importance of fungi in the context of UK tree planting. *Food and Energy Security*.
- Baldocchi, D.D., Meyers, T.P., 1988. Turbulence structure in a deciduous forest. *Boundary-Layer Meteorology* 43, 345–364.
- 610 Bannister, E., MacKenzie, A.R., Cai, X., 2021. Realistic forests and the modeling of forest-atmosphere exchange [WWW Document]. *Earth and Space Science Open Archive*. <https://doi.org/10.1002/essoar.10506854.1>
- Belcher, S.E., Harman, I.N., Finnigan, J.J., 2012. The Wind in the Willows: Flows in Forest Canopies in Complex Terrain. *Annual Review of Fluid Mechanics* 44, 479–504. <https://doi.org/10.1146/annurev-fluid-120710-101036>
- Benedetti, A., Morcrette, J.-J., Boucher, O., Dethof, A., Engelen, R.J., Fisher, M., Flentje, H., Huneeus, N., Jones, L., Kaiser, J.W., Kinne, S., Mangold, A., Razinger, M., Simmons, A.J., Suttie, M., 2009. Aerosol analysis and forecast in the European Centre for Medium-Range Weather Forecasts Integrated Forecast System: 2. Data assimilation. *Journal of Geophysical Research: Atmospheres* 114. <https://doi.org/10.1029/2008JD011115>
- 615 Biedermann, P.H.W., Vega, F.E., 2020. Ecology and Evolution of Insect–Fungus Mutualisms. *Annual Review of Entomology* 65, 431–455. <https://doi.org/10.1146/annurev-ento-011019-024910>
- 620 Burge, H.A., 2002. An update on pollen and fungal spore aerobiology. *Journal of Allergy and Clinical Immunology* 110, 544–552. <https://doi.org/10.1067/mai.2002.128674>
- Carslaw, D.C., Ropkins, K., 2012. openair --- an R package for air quality data analysis. *Environmental Modelling & Software*. 27–28, 52–61.
- Chaudhary, V.B., Nolimal, S., Sosa-Hernández, M.A., Egan, C., Kastens, J., 2020. Trait-based aerial dispersal of arbuscular mycorrhizal fungi. *New Phytologist* n/a. <https://doi.org/10.1111/nph.16667>
- 625 Cionco, R.M., 1978. Analysis of canopy index values for various canopy densities. *Boundary-Layer Meteorology* 15, 81–93. <https://doi.org/10.1007/BF00165507>
- Cionco, R.M., 1965. A Mathematical Model for Air Flow in a Vegetative Canopy. *Journal of Applied Meteorology*. [https://doi.org/10.1175/1520-0450\(1965\)004<0517:ammfaf>2.0.co;2](https://doi.org/10.1175/1520-0450(1965)004<0517:ammfaf>2.0.co;2)



- 630 Crilley, L.R., Shaw, M., Pound, R., Kramer, L.J., Price, R., Young, S., Lewis, A.C., Pope, F.D., 2018. Evaluation of a low-cost optical particle counter (Alphasense OPC-N2) for ambient air monitoring. *Atmos. Meas. Tech* 11, 709–720. <https://doi.org/10.5194/amt-11-709-2018>
- Crilley, L.R., Singh, A., Kramer, L.J., Shaw, M.D., Alam, M.S., Apte, J.S., Bloss, W.J., Hildebrandt Ruiz, L., Fu, P., Fu, W., Gani, S., Gatari, M., Ilyinskaya, E., Lewis, A.C., Ng'ang'a, D., Sun, Y., Whitty, R.C.W., Yue, S., Young, S., Pope, F.D., 2020.
- 635 Effect of aerosol composition on the performance of low-cost optical particle counter correction factors. *Atmospheric Measurement Techniques* 13, 1181–1193. <https://doi.org/10.5194/amt-13-1181-2020>
- Davies, R.R., Denny, M.J., Newton, L.M., 1963. A Comparison Between the Summer and Autumn Air-Sporas at London and Liverpool. *Allergy* 18, 131–147. <https://doi.org/10.1111/j.1398-9995.1963.tb03156.x>
- Di-Giovanni, F., Kevan, P.G., Nasr, M.E., 1995. The variability in settling velocities of some pollen and spores. *Grana* 34, 39–
- 640 44. <https://doi.org/10.1080/00173139509429031>
- Dressaire, E., Yamada, L., Song, B., Roper, M., 2016. Mushrooms use convectively created airflows to disperse their spores. *Proc Natl Acad Sci U S A* 113, 2833–2838. <https://doi.org/10.1073/pnas.1509612113>
- Finnigan, J., 2000. Turbulence in Plant Canopies. *Annual Review of Fluid Mechanics* 32, 519–571.
- Fröhlich-Nowoisky, J., Kampf, C.J., Weber, B., Huffman, J.A., Pöhlker, C., Andreae, M.O., Lang-Yona, N., Burrows, S.M.,
- 645 Gunthe, S.S., Elbert, W., Su, H., Hoor, P., Thines, E., Hoffmann, T., Després, V.R., Pöschl, U., 2016. Bioaerosols in the Earth system: Climate, health, and ecosystem interactions. *Atmospheric Research* 182, 346–376. <https://doi.org/10.1016/j.atmosres.2016.07.018>
- Fröhlich-Nowoisky, J., Pickersgill, D.A., Despres, V.R., Pöschl, U., 2009. High diversity of fungi in air particulate matter. *Proceedings of the National Academy of Sciences* 106, 12814–12819. <https://doi.org/10.1073/pnas.0811003106>
- 650 Gange, A.C., Gange, E.G., Sparks, T.H., Boddy, L., 2007. Rapid and Recent Changes in Fungal Fruiting Patterns. *Science* 316, 71–71. <https://doi.org/10.1126/science.1137489>
- Gardner, A., Ellsworth, D.S., Crous, K.Y., Pritchard, J., MacKenzie, A.R., 2020. Is photosynthetic enhancement sustained through three years of elevated CO<sub>2</sub> exposure in 175-year old *Quercus robur*? *bioRxiv* 2020.12.16.416255. <https://doi.org/10.1101/2020.12.16.416255>
- 655 Gilbert, G.S., Reynolds, D.R., 2005. Nocturnal Fungi: Airborne Spores in the Canopy and Understory of a Tropical Rain Forest. *Biotropica* 37, 462–464.
- Gosselin, M.I., Rathnayake, C.M., Crawford, I., Pöhlker, C., Fröhlich-Nowoisky, J., Schmer, B., Després, V.R., Engling, G., Gallagher, M., Stone, E., Pöschl, U., Huffman, J.A., 2016. Fluorescent bioaerosol particle, molecular tracer, and fungal spore concentrations during dry and rainy periods in a semi-arid forest. *Atmos. Chem. Phys.* 16, 15165–15184. <https://doi.org/10.5194/acp-16-15165-2016>
- 660 Griffiths, P.T., Borlace, J.-S., Gallimore, P.J., Kalberer, M., Herzog, M., Pope, F.D., 2012. Hygroscopic growth and cloud activation of pollen: a laboratory and modelling study. *Atmospheric Science Letters* 13, 289–295. <https://doi.org/10.1002/asl.397>

- Grinn-Gofroñ, A., Nowosad, J., Bosiacka, B., Camacho, I., Pashley, C., Belmonte, J., De Linares, C., Ianovici, N., Manzano, J.M.M., Sadyś, M., Skjøth, C., Rodinkova, V., Tormo-Molina, R., Vokou, D., Fernández-Rodríguez, S., Damialis, A., 2019. Airborne *Alternaria* and *Cladosporium* fungal spores in Europe: Forecasting possibilities and relationships with meteorological parameters. *Science of The Total Environment* 653, 938–946. <https://doi.org/10.1016/j.scitotenv.2018.10.419>
- Grundström, M., Adams-Groom, B., Pashley, C.H., Dahl, Å., Rasmussen, K., de Weger, L.A., Thibaudon, M., Fernández-Rodríguez, S., Silva-Palacios, I., Skjøth, C.A., 2019. Oak pollen seasonality and severity across Europe and modelling the season start using a generalized phenological model. *Science of The Total Environment* 663, 527–536. <https://doi.org/10.1016/j.scitotenv.2019.01.212>
- Hagan, D., 2017. *py-opc*, available at <https://github.com/dhhagan/py-opc> [WWW Document]. URL <https://github.com/dhhagan/py-opc> (accessed 1.27.21).
- Halbwachs, H., Bässler, C., 2015. Gone with the wind – a review on basidiospores of lamellate agarics. *Mycosphere* 6, 78–112. <https://doi.org/10.5943/mycosphere/6/1/10>
- Hart, K.M., Curioni, G., Blaen, P., Harper, N.J., Miles, P., Lewin, K.F., Nagy, J., Bannister, E.J., Cai, X.M., Thomas, R.M., Krause, S., Tausz, M., MacKenzie, A.R., 2019. Characteristics of Free Air Carbon Dioxide Enrichment of a Northern Temperate Mature Forest. *Global Change Biology* 0. <https://doi.org/10.1111/gcb.14786>
- Heald, C.L., Spracklen, D.V., 2009. Atmospheric budget of primary biological aerosol particles from fungal spores. *Geophysical Research Letters* 36. <https://doi.org/10.1029/2009GL037493>
- Healy, D.A., O'Connor, D.J., Sodeau, J.R., 2012. Measurement of the particle counting efficiency of the “Waveband Integrated Bioaerosol Sensor” model number 4 (WIBS-4). *Journal of Aerosol Science* 47, 94–99. <https://doi.org/10.1016/j.jaerosci.2012.01.003>
- Hollins, P.D., Kettlewell, P.S., Atkinson, M.D., Stephenson, D.B., Corden, J.M., Millington, W.M., Mullins, J., 2004. Relationships between airborne fungal spore concentration of *Cladosporium* and the summer climate at two sites in Britain. *Int J Biometeorol* 48, 137–141. <https://doi.org/10.1007/s00484-003-0188-9>
- Horton, T.R., 2017. Spore Dispersal in Ectomycorrhizal Fungi at Fine and Regional Scales, in: Tedersoo, L. (Ed.), *Biogeography of Mycorrhizal Symbiosis*, Ecological Studies. Springer International Publishing, Cham, pp. 61–78. [https://doi.org/10.1007/978-3-319-56363-3\\_3](https://doi.org/10.1007/978-3-319-56363-3_3)
- Inness, A., Ades, M., Agustí-Panareda, A., Barré, J., Benedictow, A., Blechschmidt, A.-M., Dominguez, J.J., Engelen, R., Eskes, H., Flemming, J., Huijnen, V., Jones, L., Kipling, Z., Massart, S., Parrington, M., Peuch, V.-H., Razinger, M., Remy, S., Schulz, M., Suttie, M., 2019. The CAMS reanalysis of atmospheric composition. *Atmospheric Chemistry and Physics* 19, 3515–3556. <https://doi.org/10.5194/acp-19-3515-2019>
- Jones, A.M., Harrison, R.M., 2004. The effects of meteorological factors on atmospheric bioaerosol concentrations—a review. *Science of The Total Environment* 326, 151–180. <https://doi.org/10.1016/j.scitotenv.2003.11.021>
- Kaimal, J.C., Finnigan, J.J., 1994. *Atmospheric boundary layer flows—their structure and measurement*. Oxford University Press, New York. <https://doi.org/10.16085/j.issn.1000-6613.2012.02.016>

- Kausserud, H., Heegaard, E., Buntgen, U., Halvorsen, R., Egli, S., Senn-Irlet, B., Krisai-Greilhuber, I., Damon, W., Sparks, T., Norden, J., Hoiland, K., Kirk, P., Semenov, M., Boddy, L., Stenseth, N.C., 2012. Warming-induced shift in European mushroom fruiting phenology. *Proceedings of the National Academy of Sciences* 109, 14488–14493. <https://doi.org/10.1073/pnas.1200789109>
- Kivlin, S.N., Winston, G.C., Goulden, M.L., Treseder, K.K., 2014. Environmental filtering affects soil fungal community composition more than dispersal limitation at regional scales. *Fungal Ecology, Fungal ecology beyond boundaries: from communities to the globe* 12, 14–25. <https://doi.org/10.1016/j.funeco.2014.04.004>
- 705 Klironomos, J.N., Rillig, M.C., Allen, M.F., Zak, D.R., Pregitzer, K.S., Kubiske, M.E., 1997. Increased levels of airborne fungal spores in response to *Populus tremuloides* grown under elevated atmospheric CO<sub>2</sub>. *Can. J. Bot.* 75, 1670–1673. <https://doi.org/10.1139/b97-880>
- Læssøe, T., Petersen, J.H., 2019. *Fungi of Temperate Europe*. Princeton University Press.
- Li, D.-W., Kendrick, B., 1995. A year-round study on functional relationships of airborne fungi with meteorological factors. *Int J Biometeorol* 39, 74–80. <https://doi.org/10.1007/BF01212584>
- 710 Lilleskov, E.A., Bruns, T.D., 2005. Spore dispersal of a resupinate ectomycorrhizal fungus, *Tomentella subulilacina*, via soil food webs. *Mycologia* 97, 762–769. <https://doi.org/10.1080/15572536.2006.11832767>
- MacKenzie, A.R., Krause, S., Hart, K., Thomas, R., Blaen, P., Hamilton, R., Curioni, G., Quick, S., Kourmouli, A., Hannah, D., Warner, S.C., Brekenfeld, N., Ullah, S., Press, M., 2020. BIFoR FACE: Water-soil-vegetation-atmosphere research in a temperate deciduous forest catchment, including under elevated CO<sub>2</sub> (preprint). *Preprints*. <https://doi.org/10.22541/au.160157598.86879557>
- 715 Moore-Landecker, E., 2011. Fungal Spores, in: *ELS*. American Cancer Society. <https://doi.org/10.1002/9780470015902.a0000378.pub2>
- Morcrette, J.-J., Boucher, O., Jones, L., Salmond, D., Bechtold, P., Beljaars, A., Benedetti, A., Bonet, A., Kaiser, J.W., 720 Razinger, M., Schulz, M., Serrar, S., Simmons, A.J., Sofiev, M., Suttie, M., Tompkins, A.M., Untch, A., 2009. Aerosol analysis and forecast in the European Centre for Medium-Range Weather Forecasts Integrated Forecast System: Forward modeling. *Journal of Geophysical Research: Atmospheres* 114. <https://doi.org/10.1029/2008JD011235>
- Nathan, R., Horn, H.S., Chave, J., Levin, S.A., 2002. Mechanistic models for tree seed dispersal by wind in dense forests and open landscapes., in: Levey, D.J., Silva, W.R., Galetti, M. (Eds.), *Seed Dispersal and Frugivory: Ecology, Evolution and Conservation*. Third International Symposium-Workshop on Frugivores and Seed Dispersal, São Pedro, Brazil, 6-11 August 2000. CABI, Wallingford, pp. 69–82. <https://doi.org/10.1079/9780851995250.0069>
- 725 National Pollen and Aerobiology Research Unit, 2012. . *Pollen Calendars by area - University Of Worcester*. URL <https://www.worcester.ac.uk/about/academic-schools/school-of-science-and-the-environment/science-and-the-environment-research/national-pollen-and-aerobiology-research-unit/pollen-calendar.aspx> (accessed 7.16.20).
- 730 Norby, R.J., De Kauwe, M.G., Domingues, T.F., Duursma, R.A., Ellsworth, D.S., Goll, D.S., Lapola, D.M., Luus, K.A., MacKenzie, A.R., Medlyn, B.E., Pavlick, R., Rammig, A., Smith, B., Thomas, R., Thonicke, K., Walker, A.P., Yang, X.,

- Zaehle, S., 2016. Model-data synthesis for the next generation of forest free-air CO<sub>2</sub> enrichment (FACE) experiments. *New Phytologist* 209, 17–28. <https://doi.org/10.1111/nph.13593>
- Norros, V., Penttilä, R., Suominen, M., Ovaskainen, O., 2012. Dispersal may limit the occurrence of specialist wood decay fungi already at small spatial scales. *Oikos* 121, 961–974. <https://doi.org/10.1111/j.1600-0706.2012.20052.x>
- Norros, V., Rannik, Ü., Hussein, T., Petäjä, T., Vesala, T., Ovaskainen, O., 2014a. Do small spores disperse further than large spores? *Ecology* 95, 1612–1621. <https://doi.org/10.1890/13-0877.1>
- Norros, V., Rannik, Ü., Hussein, T., Petäjä, T., Vesala, T., Ovaskainen, O., 2014b. Do small spores disperse further than large spores? *Ecology* 95, 1612–1621. <https://doi.org/10.1890/13-0877.1>
- 740 O'Connor, D.J., Sadyś, M., Skjøth, C.A., Healy, D.A., Kennedy, R., Sodeau, J.R., 2014. Atmospheric concentrations of *Alternaria*, *Cladosporium*, *Ganoderma* and *Didymella* spores monitored in Cork (Ireland) and Worcester (England) during the summer of 2010. *Aerobiologia* 30, 397–411. <https://doi.org/10.1007/s10453-014-9337-3>
- Oneto, D.L., Golan, J., Mazzino, A., Pringle, A., Seminara, A., 2020. Timing of fungal spore release dictates survival during atmospheric transport. *PNAS* 117, 5134–5143. <https://doi.org/10.1073/pnas.1913752117>
- 745 Page, C.N., 1997. *The Ferns of Britain and Ireland*, 2nd ed. CUP.
- Petters, M.D., Kreidenweis, S.M., 2007. A single parameter representation of hygroscopic growth and cloud condensation nucleus activity. *Atmospheric Chemistry and Physics* 7, 1961–1971. <https://doi.org/10.5194/acp-7-1961-2007>
- Pope, F.D., 2010. Pollen grains are efficient cloud condensation nuclei. *Environ. Res. Lett.* 5, 044015. <https://doi.org/10.1088/1748-9326/5/4/044015>
- 750 Pöschl, U., 2005. *Atmospheric Aerosols: Composition, Transformation, Climate and Health Effects*. *Angewandte Chemie International Edition* 44, 7520–7540. <https://doi.org/10.1002/anie.200501122>
- R Core Team, 2020. *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria.
- Raupach, M.R., Finnigan, J.J., Brunet, Y., 1996. Coherent Eddies and Turbulence in Vegetation Canopies: The Mixing-Layer Analogy. *Boundary-Layer Meteorology 25th Anniversary Volume, 1970–1995* 351–382. [https://doi.org/10.1007/978-94-017-0944-6\\_15](https://doi.org/10.1007/978-94-017-0944-6_15)
- Reinmuth-Selzle, K., Kampf, C.J., Lucas, K., Lang-Yona, N., Fröhlich-Nowoisky, J., Shiraiwa, M., Lakey, P.S.J., Lai, S., Liu, F., Kunert, A.T., Ziegler, K., Shen, F., Sgarbanti, R., Weber, B., Bellinghausen, I., Saloga, J., Weller, M.G., Duschl, A., Schuppan, D., Pöschl, U., 2017. Air Pollution and Climate Change Effects on Allergies in the Anthropocene: Abundance, Interaction, and Modification of Allergens and Adjuvants. *Environ Sci Technol* 51, 4119–4141. <https://doi.org/10.1021/acs.est.6b04908>
- 760 Reponen, T., Willeke, K., Ulevicius, V., Reponen, A., Grinshpun, S.A., 1996. Effect of relative humidity on the aerodynamic diameter and respiratory deposition of fungal spores. *Atmospheric Environment* 30, 3967–3974. [https://doi.org/10.1016/1352-2310\(96\)00128-8](https://doi.org/10.1016/1352-2310(96)00128-8)

- 765 Sadyś, M., Adams-Groom, B., Herbert, R.J., Kennedy, R., 2016a. Comparisons of fungal spore distributions using air sampling at Worcester, England (2006–2010). *Aerobiologia* 32, 619–634. <https://doi.org/10.1007/s10453-016-9436-4>
- Sadyś, M., Kennedy, R., West, J.S., 2016b. Potential impact of climate change on fungal distributions: analysis of 2 years of contrasting weather in the UK. *Aerobiologia* 32, 127–137. <https://doi.org/10.1007/s10453-015-9402-6>
- 770 Sadyś, M., Skjøth, C.A., Kennedy, R., 2014. Back-trajectories show export of airborne fungal spores (*Ganoderma* sp.) from forests to agricultural and urban areas in England. *Atmospheric Environment* 84, 88–99. <https://doi.org/10.1016/j.atmosenv.2013.11.015>
- Sesartic, A., Dallafior, T.N., 2011a. Global fungal spore emissions, review and synthesis of literature data. *Biogeosciences* 8, 1181–1192. <https://doi.org/10.5194/bg-8-1181-2011>
- 775 Sesartic, A., Dallafior, T.N., 2011b. Global fungal spore emissions, review and synthesis of literature data. *Biogeosciences* 8, 1181–1192. <https://doi.org/10.5194/bg-8-1181-2011>
- Sousan, S., Koehler, K., Hallett, L., Peters, T.M., 2016. Evaluation of the Alphasense optical particle counter (OPC-N2) and the Grimm portable aerosol spectrometer (PAS-1.108). *Aerosol Science and Technology* 50, 1352–1365. <https://doi.org/10.1080/02786826.2016.1232859>
- Stephens, R.B., Rowe, R.J., 2020. The underappreciated role of rodent generalists in fungal spore dispersal networks. *Ecology* 780 101, e02972. <https://doi.org/10.1002/ecy.2972>
- Su, H.B., Shaw, R.H., Pawu, K.T., Moeng, C.H., Sullivan, P.P., 1998. Turbulent statistics of neutrally stratified flow within and above a sparse forest from large-eddy simulation and field observations. *Boundary-Layer Meteorology* 88, 363–397. <https://doi.org/10.1023/A:1001108411184>
- Tesmer, J., Schnittler, M., 2007. Sedimentation velocity of myxomycete spores. *Mycological Progress* 6, 229–234. <https://doi.org/10.1007/s11557-007-0539-8>
- 785 Wickham, H., 2016. *ggplot2: Elegant Graphics for Data Analysis*.
- Wickham, H., Averick, M., Bryan, J., Chang, W., D'Agostino McGowan, L., François, R., Grolemund, G., Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T.L., Miller, E., Milton Bache, S., Müller, K., Ooms, J., Robinson, D., Seidel, D.P., Spinu, V., Takahashi, K., Vaughan, D., Wilke, C., Woo, K., Yutani, H., 2019a. Welcome to the Tidyverse [WWW Document]. URL <https://tidyverse.tidyverse.org/articles/paper.html> (accessed 5.11.21).
- 790 Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L.D., François, R., Grolemund, G., Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T.L., Miller, E., Bache, S.M., Müller, K., Ooms, J., Robinson, D., Seidel, D.P., Spinu, V., Takahashi, K., Vaughan, D., Wilke, C., Woo, K., Yutani, H., 2019b. Welcome to the Tidyverse. *Journal of Open Source Software* 4, 1686. <https://doi.org/10.21105/joss.01686>
- 795 Wolf, J., Johnson, N.C., Rowland, D.L., Reich, P.B., 2003. Elevated CO<sub>2</sub> and plant species richness impact arbuscular mycorrhizal fungal spore communities. *New Phytologist* 157, 579–588.

Wolf, J., O'Neill, N.R., Rogers, C.A., Muilenberg, M.L., Ziska, L.H., 2010. Elevated atmospheric carbon dioxide concentrations amplify *Alternaria alternata* sporulation and total antigen production. *Environmental Health Perspectives* 118, 1223–1228. <https://doi.org/10.1289/ehp.0901867>

800 Ziegler, C., Kulawska, A., Kourmouli, A., Hamilton, L., Shi, Z., MacKenzie, A.R., Dyson, R.J., Johnston, I.G., 2021. Quantification and uncertainty of root growth stimulation by elevated CO<sub>2</sub> in mature temperate deciduous forest. *bioRxiv* 2021.04.15.440027. <https://doi.org/10.1101/2021.04.15.440027>