

Phosphorus stress strongly reduced plant physiological activity, but only temporarily, in a mesocosm experiment with *Zea mays* colonized by arbuscular mycorrhizal fungi

Melanie S. Verlinden¹, Hamada AbdElgawad^{2,3}, Arne Ven¹, Lore T. Verryckt¹, Sebastian Wieneke^{1,4}, Ivan A. Janssens¹, Sara Vicca¹

¹Plant and Vegetation Ecology (PLECO), Department of Biology, University of Antwerp, Wilrijk, 2610, Belgium

²Botany and Microbiology Department, Faculty of Science, Beni-Suef University, Beni-Suef, 62521, Egypt

³Integrated Molecular Plant Physiology Research (IMPRES), Department of Biology, University of Antwerp, Antwerp, 2020, Belgium

⁴Remote Sensing Centre for Earth System Research (RSC4Earth, Faculty of Physics and Earth Sciences, University of Leipzig, Leipzig, 04109, Germany

Correspondence to: Melanie S. Verlinden (Melanie.Verlinden@uantwerpen.be); Sara Vicca (Sara.Vicca@uantwerpen.be)

Abstract. Phosphorus (P) is an essential macronutrient for plant growth and one of the least available nutrients in soil. P limitation is often a major constraint for plant growth globally. Although P addition experiments have been carried out to study the long-term effects on yield, data on P addition effects on seasonal variation of leaf-level photosynthesis are scarce. Arbuscular mycorrhizal fungi (AMF) can be of major importance for plant nutrient uptake, and AMF growth may be important for explaining temporal patterns in leaf physiology. In a nitrogen (N) and P fertilization experiment with *Zea mays*, we investigated the effect of P limitation on leaf pigments and leaf enzymes, how these relate to leaf-level photosynthesis, and how these relationships change during the growing season. Previous research indicated that N addition did not affect plant growth and also the leaf measurements in the current study were unaffected by N addition. Contrary to N addition, P addition strongly influenced plant growth and leaf-level measurements. At low soil P availability, leaf-level photosynthetic and respiratory activity were strongly decreased and this was associated with reduced chlorophyll and photosynthetic enzymes. Contrary to the expected increase in P stress over time following gradual soil P depletion, plant P-limitation decreased over time. For most leaf-level processes, pigments and enzymes under study, the fertilization effect had even disappeared two months after planting. Our results point towards a key role for the AMF-symbiosis and consequent increase of P uptake in explaining the vanishing P stress.

30

1 Introduction

Phosphorus (P) is a crucial element in natural ecosystems. It is present in the structure of DNA, in cell membranes, in molecules storing and supplying energy and in several enzymes. As a consequence, P plays a crucial role in plant and soil processes, it regulates productivity and ecosystem functions and influences organisms from the individual to the community level (Elser et al., 2000; Vitousek et al., 2010; Peñuelas et al., 2013). The importance of P for the functioning of the Earth's biogeochemical cycles, especially the carbon cycle, is therefore being

increasingly recognized (Vitousek et al., 2010; Wieder et al., 2015; Vicca et al., 2018) and this is reflected in the recent efforts to include P in terrestrial biosphere models (Wang et al., 2010; Gong et al., 2012; Thum et al., 2019). In plants, P plays a role in most developmental and biochemical processes. Structurally, P is a component of RNA
40 and membrane phospholipids, while metabolically, P functions in the storage and transfer of energy and in energizing of binding sites for metabolic turnover (Schulze et al., 2005; Veneklaas et al., 2012). However, P is one of the least available macronutrients in soils, and P limitation is often a major constraint for plant growth (Augusto et al., 2017). On more than one third of the arable land worldwide, plant productivity is considered to be limited by P (Calderón-Vázquez et al., 2009).

45 Various experiments have been conducted to study the effect of P addition to crops, thereby mainly focusing on the long-term effect on yield (Khan et al., 2018; Johnston and Poulton, 2019). However, data on seasonal variation in leaf-level photosynthesis, especially in crops, are scarce (Rodríguez et al., 2000; Rogers, 2014), while accurate seasonal estimates of photosynthetic capacity are critical for modelling the time course of carbon fluxes (Miner and Bauerle, 2019). The majority of studies investigating effects of nutrients on photosynthesis focus on nitrogen
50 (N) and much less on P and other nutrients (e.g., Brooks, 1986; Brooks et al. 1988; Rodríguez and Goudriaan, 1995; Rodríguez et al., 1998). In addition, it is unclear whether leaf traits, such as leaf nutrients, pigments and enzymes, change seasonally in relation to leaf-level photosynthesis.

Among others, plant P limitation typically results in reduced photosynthesis and plant growth, especially aboveground. P is required for adenosine triphosphate (ATP) synthesis (Veneklaas et al., 2012), which is needed
55 to regenerate Ribulose 1,5-bisphosphate (RuBP) in the Calvin cycle of photosynthesis. Inorganic phosphate (Pi) directly affects the activity of Calvin cycle enzymes through the level of activation. For instance, Pi is required for light activation of Rubisco (Parry et al., 2008). It also directly affects maximum rate of CO₂-limited carboxylation (v_{cmax}) and triose phosphate utilization (Lewis et al., 1994) and RuBP-regeneration-limited rates of electron transport (Loustau et al., 1999). P-deficiency therefore leads to a decrease in RuBP pool size and insufficient ATP,
60 and consequently to a decrease in photosynthetic C assimilation. The concentration and specific activity of Rubisco, the primary CO₂ fixing enzyme in photosynthesis, are generally little affected by P stress (Brooks, 1986; Paul and Stitt, 1993; Pieters et al., 2001, but see Jacob and Lawlor, 1991; Pieters et al., 2001).

Pi can also indirectly affect photosynthesis through the changes in stromal pH (Bhagwat, 1981), where the consumption of Pi as a substrate of photosynthesis could decrease photosynthesis by a direct effect of low stromal
65 Pi concentration on Rubisco. Moreover, the effect of P on photosynthesis depends on the dynamic interactions between sink and source tissues. Low P can reduce carbon export to sinks, and thus decrease sink strength, thereby limiting photosynthesis (Pieters et al., 2001). Concomitantly, leaf starch can increase with P stress (Zhang et al., 2014) due to low availability of P for triphosphate translocation, although decreases of leaf starch have also been observed (Halsted and Lynch, 1996). Moreover, low sink strength restricts the recycling of Pi back to the
70 chloroplast, further reducing photosynthesis (Paul and Foyer, 2001).

In a mesocosm nutrient manipulation experiment setup (previously described in Verlinden et al., 2018), maize (*Zea mays* L.) was planted at different soil N and P availabilities. As demonstrated in Verlinden et al. (2018), this resulted in a strong P, but no N effect on plant growth or photosynthesis at mesocosm scale. In that study, also arbuscular mycorrhizal fungi (AMF) played an important role in explaining plant carbon uptake and allocation.
75 AMF are important for nutrient uptake in maize (Hartnett and Wilson, 1999; Hoeksema et al., 2010), especially for P, and hence AMF growth may also be important for explaining variation in leaf physiology. The objective of

the current study is to test the effect of P limitation on leaf pigments, sugars and photosynthetic enzymes, how they relate to leaf-level photosynthesis, and how these relationships change during the growing season. At low soil P availability, we expected low leaf-level photosynthetic and respiratory activity, associated with reduced chlorophyll and photosynthetic enzymes. Furthermore, P-stress was expected to increase over time, as plants were expected to gradually deplete the soil P.

2 Material and methods

2.1 Experimental design

For this study, we used the first of two mesocosm fertilization experiments. While the first applied a full-factorial NxP fertilization approach and was first described in Verlinden et al. (2018), the second applied a P gradient. Results for the latter are reported in Ven et al. (2020b). The mesocosm experiment consisted of 20 (1 m × 1.2 m, 0.6 m high) insulated boxes was set up in a greenhouse in Sint-Katelijne-Waver, Belgium (51°04'38" N, 4°32'05" E). To each mesocosm we added soil, which was a homogenized mixture of sand originating from a pine forest in a nature reserve in Flanders, white river sand and a minority of compost (details of the experimental setup are described in Verlinden et al. (2018)). On 20 May 2016, 12 seedlings of maize (*Zea mays* L., variety 'Tom Thumb') were planted per mesocosm. Different treatments (set up in five replicates) were distinguished in the level of nutrients added: the +N treatment was fertilized with calcium nitrate at a rate of 95.5 kg N ha⁻¹ (YaraLiva® Calcinit®), the +P treatment received 20 kg P ha⁻¹ as triple superphosphate (Janssens-Smeets®), the combined +N and +P treatment (+NP) received both amounts together. The control treatment received, as all other treatments, only a basic level of micronutrients (Fertigreen® Patentkali® and GroGreen® containing in kg ha⁻¹: 79 Potassium, 19 Magnesium, 53 Sulfur, 0.4 Boron, 0.1 Copper, 2.4 Iron, 1.1 Manganese, 0.1 Molybdenum, 0.4 Zinc). Spores-based inoculum of AMF (species *Rhizophagus irregularis*, Symplanta®) was added to all 20 (4 treatments × 5 replicates) mesocosms. Soil moisture was monitored and kept at a non-limiting (field capacity) level, similar in all plots.

2.2 Measurements and analyses

2.2.1 Leaf C, N and P concentration and Specific Leaf Area

Carbon (C) and N concentrations were determined using an elemental analyzer - model FLASH 2000 (Thermo Fisher Scientific, Waltham, USA). Total leaf P concentration was determined by digestion in tubes with H₂SO₄-salicylic acid- H₂O₂ and selenium (Temminghoff and Houba, 2004). Specific Leaf Area (SLA; m² kg⁻¹) was determined as the ratio of the fresh leaf area and dry leaf mass.

2.2.2 Leaf Photosynthesis

A portable gas exchange system LI-6400 (LI-COR, Lincoln, NE, USA) was used for leaf scale CO₂ gas exchange measurements, operating as an open system (e.g. Verlinden et al., 2013). Leaf-scale measurements were performed during two weeks late June (campaign 1, C1) and repeated end of July (campaign 2, C2), allowing to study the seasonal development. Mean daily photosynthetically active radiation (PAR) during C1 and C2 were respectively 17.1 and 17.7 mol·m⁻² and average temperature respectively 21.7 and 23.3 °C.

In each plot photosynthetic CO₂-response curves (i.e. photosynthesis (A, assimilation) responses to the CO₂ concentration inside leaf air spaces (c_i)) were measured on a recently matured leaf. Leaves were allowed to equilibrate at a CO₂ concentration of 400 μmol mol⁻¹ in the leaf cuvette, after which the net CO₂ assimilation rate at a sequence of different CO₂ concentrations (i.e. 400, 30, 50, 80, 110, 150, 250, 350, 500 and 1000 μmol mol⁻¹) was measured. Photosynthetic photon flux density (PPFD) was fixed at a saturating value of 1200 μmol s⁻¹ m⁻². The resulting A-c_i data were fitted to the biochemical model of C₄ photosynthesis as presented by von Caemmerer (2000) using the package 'Plantecophys' (Duursma, 2015) in R version 3.3.1 (R Foundation for Statistical Computing, Vienna, Austria). The CO₂ assimilation rate is approximated by the minimum of the expressions of an enzyme-limited and an electron-transport-limited CO₂ assimilation rate. The parameters J_{max} (maximum electron transport rate), v_{cmax} (maximal rubisco carboxylation rate) and v_{pmax} (maximum PEP carboxylation rate) were calculated through curve fitting based on minimum least-squares.

Photosynthetic 'light response curves' were obtained by measurements of the net CO₂ assimilation rate at PPFD's of 1200, 500, 250, 100, 80, 60, 40, 30, 25, 20, 15, 10, 5 and 0 μmol m⁻² s⁻¹ (blue-red LED source type 6400-02B, 13% blue light). Leaves were allowed to equilibrate at each step before logging the data. The CO₂ concentration in the cuvette was maintained at 400 μmol mol⁻¹ and the block temperature at 25°C. From the 'light response curves', the net CO₂ assimilation rate at light saturation (A_{max}) and leaf dark respiration (R_{dark}, net CO₂ exchange at zero light) were derived. In addition, light-induced inhibition of leaf respiration was estimated from the 'light response curves' (for PPFD's 0 to 80 μmol m⁻² s⁻¹) from the intersections of the fitted lines above and below the light compensation point the y-axis, giving respectively R_{light} and R_{dark} (Kok, 1948). All selected leaves were harvested and stored at -80°C for later analyses.

2.2.3 Chemical analyses of leaf material

Rubisco activity was analyzed according to Sulpice et al. (2007). It was expressed as the conversion rate of glycerate kinase (3-PGA) of extracted leaf samples, in μmol 3-PGA m⁻² min⁻¹. The activity of Rubisco was determined directly ('direct rubisco'), without incubation of the extract in the presence of 10 mM HCO₃³⁻ and 20 mM Mg²⁺ to convert the non-carbamylated Rubisco into the carbamylated form. The assay of phosphoenolpyruvate carboxylase (PEPC) was coupled with the malate dehydrogenase reaction, the resulting rate of PEPC activity was expressed in μmol HCO₃ m⁻² min⁻¹.

Mono- and oligosaccharides in leaves were analyzed chromatographically according to AbdElgawad et al. (2014). Soluble sugar concentrations were measured by high performance anion exchange chromatography of extracted leaf samples with pulsed amperometric detection (HPAEC-PAD) and the total soluble sugar concentration was calculated as their sum. The remaining pellet of soluble sugars extraction was treated with a mixture of α-amylase and amyloglucosidase to extract starch.

High-performance liquid chromatography (HPLC) was used to analyze leaf pigments. The detection of the carotenoids and xanthophylls was done by a diode array detector (Shimadzu SPD-M10Avp, Kyoto, Japan) at four wavelengths (420, 440, 462, 660 nm) and integrated via the software program (Shimadzu Lab Solutions Lite, Kyoto, Japan) in which the concentration was determined using a calibration curve.

2.2.4 Mycorrhizal fungi

Because AMF growth is potentially crucial for explaining patterns in the leaf response to P limitation, we determined the time course of AMF abundance in each of the mesocosms. To this end, five mesh bags filled with white river sand – and permeable for fungi but not for roots (30 μm mesh size) – were buried vertically into the top soil of each mesocosm one week before planting. They were harvested consecutively 31 (corresponding to C1, *v.i.*) and 61 days (right before C2) after planting. Hyphae were extracted from 4 g mesh bag sand using the method of Rillig et al. (1999). After suspending, processing and staining the sample, hyphal intersects were counted at a magnification of 40×10 using a grid in the microscope ocular. Hyphal length density was calculated following Eq. (1) (Tennant, 1975; Rillig et al., 1999):

$$\text{HLD} = (\pi \cdot n \cdot a \cdot d) \cdot (h \cdot w)^{-1}, \quad (1)$$

where HLD = hyphal length density (mm hyphae g^{-1} soil), n = number of intersects containing AMF hyphae, a = filter area (mm^2) examined, d = dilution factor, h = total length of raster lines projected on filter (mm), and w = soil weight (g).

Mycorrhizal colonization was examined in C1 and C2 by sampling roots from two plants per mesocosm. Per plant, 20 cm of one lateral root containing root hair, was excavated, cut, and stored. Mycorrhizal colonization was quantified by counting arbuscules, vesicles, and hyphae applying the gridline intersection method (Vierheilig et al., 2005). The methodology on root colonization determination is described more elaborately in Verlinden et al. (2018).

2.2.5 Statistical Analyses

Data normality and homoscedasticity were checked using the Shapiro-Wilk and Levene's test, respectively. A three-way mixed analyses of variance (ANOVA) was applied to test if the quantified variables differed between the treatments and between C1 and C2. N addition and P addition were both considered as between-subject variables and time (campaign) as a within-subject variable. Non-significant interactions terms, and further, non-significant factors were removed from the model. In case of significant interaction between factors, the analysis included their multiplied factor levels. A Tukey post-hoc test was applied for pairwise comparison in case of significant factor effects.

3 Results


The addition of P-fertilizer increased soil P availability (Verlinden et al., 2018), as well as leaf P concentration (Table 1). At the time of C1, leaf P concentration was three to four times higher in the +P and +NP treatments than in the non-P-fertilized control and +N treatments. Leaf N:P ratio was higher in the non-P-fertilized treatments than in the P-fertilized treatments (an average N:P ratio of 19.8 versus 37.2 for the non-P-fertilized treatments). However, in C2, the leaf P concentration had increased in all treatments to a similar level (Table 1), as well as the N:P ratio, which decreased for all treatments to a similar level with a mean of 13.8. Leaves in the non-P-fertilized mesocosms were thinner and/or had a lower density than in the P-fertilized mesocosms (Table 1) during C1, with mean SLA values of respectively 52.9 ± 0.9 and 33.9 ± 1.9 $\text{m}^2 \text{kg}^{-1}$. Towards later in the season, SLA decreased in all mesocosms and the difference between non-P-fertilized and P-fertilized mesocosms had disappeared at the time of C2.

185 The majority of leaf physiology parameters differed considerably between C1 versus C2 for the non-P-fertilized
treatments, while for the P-fertilized treatments differences between C1 and C2 were much less pronounced.
During C1, photosynthetic activity was very low in the non-P-fertilized treatments, with a mean A_{\max} of $6.2 (\pm 4.1)$
 $\mu\text{mol m}^{-2} \text{s}^{-1}$ for the control and +N treatments. In contrast, the +P and +NP treatments showed a mean A_{\max} more
than four times higher than in the non-P-fertilized mesocosms (Fig. 1A). A similarly high A_{\max} -level was reached
190 for all treatments in C2 (Fig. 1A). Also R_{dark} was smaller in the non-P-fertilized treatments in C1 (Fig. 1B) and
reached a similar level as the +P and +NP treatments in C2. Photosynthetic parameters J_{\max} , v_{cmax} and v_{pmax} were
all lower in the non-P-fertilized treatments than in the P-fertilized treatments during C1 (Figs. 1C-E), but by the
time of C2, J_{\max} had increased in the non-P-fertilized mesocosms to the level of the P-fertilized mesocosms. v_{cmax}
in the non-P-fertilized mesocosms had even increased to a level of about 45% higher than the P-fertilized
195 mesocosms, while the P-fertilized mesocosms showed very similar J_{\max} , v_{cmax} and v_{pmax} for C1 and C2. Light-
induced inhibition of respiration (Fig. 1F) was variable amongst the mesocosms, though on average it tended to
be higher in the non-P-fertilized mesocosms during C1, whereas no trend was observed during C2. The light
compensation point was initially lower in the non-P-fertilized plants (i.e., in the stressed plants photosynthetic
activity occurred at a lower light availability than in the P-fertilized treatments), whereas during C2 no differences
200 were observed between the mesocosms (Fig. 1G).

Similar to the gas exchange measurements, the leaf chemistry showed a strong difference between non-P-fertilized
and P-fertilized plots during C1, but not during C2. Direct rubisco concentration was initially lower in the non-P-
fertilized mesocosms (Table 1), which was also true for the enzyme PEP-carboxylase (Table 1). A P- and campaign
effect was observed for total chlorophyll (Table 1, similarly for chlorophyll_a and chlorophyll_b, data not shown), its
205 concentration was four times higher in the P-fertilized mesocosms during C1. Also beta-carotene concentration
was initially higher in the P-fertilized mesocosms (Table 1). Zeaxanthin was only detected in the non-P-fertilized
leaves during C1 (Table 1). For both lutein and violaxanthin no differences among the treatments were observed
during C1. There was a tendency of lower starch in the P-stressed mesocosms as compared to the P-fertilized
mesocosms during C1 although there was no P effect, whereas the campaign effect and interactions P x campaign
and N x P x campaign were significant.
210

During C2, direct rubisco concentration increased in the non-P-fertilized mesocosms to the same level as in P-
fertilized mesocosms, while PEP-carboxylase concentration increased in all mesocosms to reach a similar level in
C2. Chlorophyll concentration increased more than 12 times for the non-P-fertilized mesocosms from C1 to C2;
for the P-fertilized mesocosms almost four times. A similar trend was observed for beta-carotene (Table 1), of
215 which concentrations increased five- and threefold respectively. Also lutein and violaxanthin were present in
higher concentrations during C2 (Table 1). Zeaxanthin was not detected during C2. The leaf starch concentration
differed over time, leaves contained much less starch during C1 than during C2 (Table 1).

One month after establishing the experimental setup (during C1), no AMF were detected in plant roots or in the
meshbags (Fig. 2). One month later, i.e. during C2, however, AMF had clearly established, with a mean hyphal
220 length density of 760 mm per gram of soil in all treatments. The percentage of roots colonized was higher in the
non-P-fertilized treatments than in the P-fertilized plots (67% vs. 40% on average) (Fig. 2; Ven et al., 2020a).




225 **Figure 1 A-G:** Means of parameters deduced from leaf CO₂ exchange measurements per treatment and campaign. Error bars indicate standard error. C1: campaign 1, end of June; C2: campaign 2, end of July; control treatment: not fertilized, +N treatment: nitrogen fertilized, +P treatment: phosphorus fertilized, +NP treatment: both nitrogen and phosphorus fertilized. Letters above bars indicate significant differences. Significant effects are given with p- value below the plots. A_{max} = maximal assimilation rate; R_{dark} = leaf dark respiration, R_{dark}/A_{max} = ratio of leaf dark respiration to maximal assimilation rate; J_{max} = maximum electron transport rate; V_{cmax} = maximal rubisco carboxylation rate; V_{pmax} = maximum PEP carboxylation rate;

230

Table 1. Means (and standard errors) of chemically analyzed leaf traits, per campaign and per treatment.

Notes: C1: campaign 1 end of June; C2: campaign 2 end of July. SE: standard error, Superscript letters indicate homogeneous groups as results from post-hoc analysis of statistical analyses of variance.

campaign	treat- ment	leaf N g m ⁻²	leaf P g m ⁻²	leaf N:P -	SLA μmol 3- PGA/m ² min	direct rubisco μmol 3- PGA/m ² min	PEP- carboxylase μmol HCO ₃ /m ² .min	total chlorophyll mg m ⁻²	beta carotene mg m ⁻²	zeaxanthin mg m ⁻²	violaxanthin mg m ⁻²	lutein mg m ⁻²	starch g m ⁻²	insoluble sugars g m ⁻²	soluble sugars mg m ⁻²
<i>p</i> -values:															
P		< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	3.36*10 ⁻²	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01
campaign		< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	
P x campaign		< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	2.78*10 ⁻²	< 0.01	< 0.01	
P x N		3.93*10 ⁻²			2.25*10 ⁻²										
P x N x campaign		4.99*10 ⁻²			3.24*10 ⁻²								4.22*10 ⁻²		
		mean (SE)	mean (SE)	mean (SE)	mean (SE)	mean (SE)	mean (SE)	mean (SE)	mean (SE)	mean (SE)	mean (SE)	mean (SE)	mean (SE)	mean (SE)	mean (SE)
C1	control	0.66 ^a (0.03)	0.018 ^a (0.001)	37.1 ^a (1.7)	50.8 ^a (1.5)	39 ^a (8)	66 ^a (4)	82 ^a (41)	33.0 ^a (8.6)	10.3 ^a (3.1)	1.3 ^a (0.8)	6.4 ^a (3.7)	0.33 ^a (0.06)	0.49 ^a (0.03)	52 ^a (3)
	+N	0.66 ^a (0.02)	0.018 ^a (0.001)	37.3 ^a (0.6)	53.2 ^a (0.9)	35 ^a (5)	54 ^a (9)	71 ^a (21)	25.7 ^a (9.2)	7.1 ^a (2.8)	0.3 ^a (0.1)	2.6 ^a (1.3)	0.27 ^a (0.06)	0.45 ^a (0.04)	56 ^a (8)
	+P	1.01 ^b (0.04)	0.056 ^b (0.008)	19.2 ^b (2.0)	36.4 ^b (2.2)	92 ^b (6)	182 ^b (27)	231 ^b (63)	48.9 ^b (3.8)	0 ^b 0	3.5 ^a (1.2)	5.5 ^a (2.6)	0.49 ^{ab} (0.11)	1.29 ^a (0.13)	73 ^a (12)
	+NP	1.32 ^{bc} (0.10)	0.066 ^{bc} (0.008)	20.5 ^b (1.2)	31.3 ^{bc} (2.9)	112 ^b (6)	218 ^b (21)	351 ^b (79)	52.6 ^b (13.6)	0 ^b 0	4.7 ^a (1.4)	6.5 ^a (2.0)	0.47 ^a (0.06)	1.41 ^a (0.20)	88 ^a (10)
C2	control	1.37 ^c (0.02)	0.103 ^d (0.003)	13.4 ^c (0.3)	26.4 ^{cd} (0.7)	95 ^b (9)	275 ^c (28)	944 ^c (41)	137.5 ^c (8.2)	0 ^b 0	20.3 ^b (2.0)	38.2 ^{ab} (9.0)	1.08 ^{bc} (0.09)	3.89 ^b (0.63)	212 ^b (15)
	+N	1.49 ^c (0.32)	0.102 ^d (0.022)	14.5 ^c (0.5)	25.8 ^{cd} (5.2)	98 ^b (22)	290 ^c (68)	995 ^c (21)	139.0 ^c (30.1)	0 ^b 0	22.9 ^b (6.5)	39.6 ^{ab} (11.8)	1.39 ^c (0.36)	2.86 ^b (0.65)	192 ^b (47)
	+P	1.36 ^c (0.07)	0.092 ^{cd} (0.005)	14.8 ^c (0.4)	20.5 ^d (1.0)	121 ^b (22)	307 ^c (34)	1119 ^d (63)	153.4 ^d (12.7)	0 ^b 0	28.9 ^b (7.5)	67.4 ^b (6.6)	1.28 ^c (0.21)	3.97 ^b (0.62)	209 ^b (22)
	+NP	1.38 ^c (0.08)	0.111 ^d (0.008)	12.5 ^c (0.7)	22.5 ^d (0.9)	123 ^b (10)	296 ^c (11)	1083 ^d (79)	146.2 ^d (15.7)	0 ^b 0	24.6 ^b (3.1)	63.6 ^b (11.4)	0.86 ^{abc} (0.06)	3.36 ^b (0.29)	261 ^b (56)



240

Figure 2: Means of mycorrhizal hyphal length (A) and root colonization (B) by time of C1 and C2. Error bars indicate standard error. Letters above bars indicate statistical differences. Significant effects are given with p-value below the plots

4 Discussion

The unfertilized soil in our experiment was clearly P-impoverished; addition of P increased plant productivity, whereas N addition did not. End-of-season dry biomass reached 81 (\pm 7) and 510 (\pm 24) g m⁻² for the non-P-fertilized and P-fertilized treatments, respectively (Verlinden et al., 2018). N addition had no effect on the leaf-scale measurements, therefore we focus on effects of P.

Leaf photosynthetic parameters and most leaf chemistry parameters showed clear changes throughout the season, as verified by the significant P x campaign interaction effects (Fig. 1, Table 1). During C1, leaf P concentrations in the non-P-fertilized plants were three times lower than in the P-fertilized plants, whereas leaf P concentrations were similar for non-P-fertilized and P-fertilized treatments during C2. Since growth of plants with leaf N:P ratios higher than 16 (Koerselman and Meuleman, 1996) up to 20 (Güsewell, 2004) is considered to be P-limited, the high leaf N:P ratios of about 37 illustrate a clear P-limitation of plant-growth for the non-P-fertilized treatments in C1, while P-fertilized treatments were close to P-limitation. In C2, plants seemed to have reached a favorable allocation of N and P, as shown by the favorable N:P ratio (i.e. between 9 and 18, Beauchamp and Hamilton, 1970) in all treatments. In accordance, the leaves of the non-P-fertilized plants turned yellow in the first weeks of the experiment, but greened up later.

The initial P-limitation present during C1, strongly limited leaf-level photosynthesis as A_{max} , J_{max} and v_{cmax} were three to four times lower in non-P-fertilized than in P-fertilized plants. This inhibitory effect can be attributed to the decrease in the pool size of ribulose-1,5-bisphosphate (RuBP) and its regeneration (Jacob and Lawlor, 1992; Pieters et al., 2001; Calderón-Vázquez et al., 2009), or by feedback inhibition of photosynthesis, but the latter was not specifically tested. Feedback inhibition of photosynthesis can be induced by elevated soluble sugar levels decreasing the gene expression of photosynthetic enzymes (e.g. PEPC, malic enzyme and RuBisCo) (Jeannette et al., 2000; AbdElgawad et al., 2020). This was not likely the case here, since during C1 sugar levels tended to be lower in the non-P-fertilized than in the P-fertilized treatments. Lower starch and soluble sugar synthesis, like in the non-P-fertilized treatments, can slow Pi regeneration, limit ATP production and eventually the functioning of the Calvin cycle, which is known as short-term feedback regulation of photosynthesis (Griffin and Seemann, 1996).

Also Rubisco levels were about three times lower in the non-P-fertilized plants than in the P-fertilized plants (Table 1). Insufficient P restricts the conversion of adenosine diphosphate (ADP) to ATP, limiting the RuBP regeneration (Rao and Terry, 1989; Calderón-Vázquez et al., 2009). C₄ plants can maintain adequate levels of P in the bundle cells, and their growth is therefore generally less constrained by P limitation as compared to C₃ plants (Calderón-

Vázquez et al., 2009). This indicates that in our experiment, plants with absent P fertilization must have experienced extreme P limitation early in the season in our experiment. Nonetheless, during C2, photosynthetic parameters reached similar values for all treatments.

Total chlorophyll can drop drastically in case of P deprivation (Jacob and Lawlor, 1991; Usuda and Shimogawara, 1991). In our experiment, chlorophyll concentration was initially lower in the non-P-fertilized mesocosms as compared to the P-fertilized mesocosms. During C2, however, chlorophyll concentration strongly increased in all treatments, both in the initially non-P-fertilized plants where chlorophyll increase was accompanied by increased photosynthesis, and in the P-fertilized plants. In the latter ones A_{max} did not differ between C1 and C2, indicating that photosynthesis did not increase despite the increase in chlorophyll concentration. Zeaxanthin was only detected in the non-P-fertilized plants during C1. Schlüter et al. (2013) showed the enhancement of protective pigments, such as zeaxanthin, in maize leaves when growing at low P availability. Zeaxanthin plays a key role in the protection of photosynthetic organisms against excess light, minimizing the over-excitation (Jahns and Holzwarth, 2012; Kuczyńska et al., 2012; Ashraf and Harris, 2013). The xanthophyll violaxanthin is reversibly de-epoxidized to zeaxanthin in the xanthophyll cycle when the light absorbed exceeds the capacity of photosynthesis. Zeaxanthin synthesis thus acts as a rescuing mechanism in strongly photo-oxidizing conditions (Dall'Osto et al., 2010) and increased zeaxanthin concentrations imply a decrease of light harvesting. In our experiment, no zeaxanthin was detected later in the season, indicating that stress, likely due to P-limitation, was relieved and plant growth recovered, as also indicated by the increased net photosynthetic rate.

P deprivation has been found to increase the leaf starch concentration in maize (Zhang et al., 2014), although decreases in starch levels under low P conditions have also been reported (Schlüter et al., 2013). In our experiment, P addition had no clear effect on the leaf starch concentration (Table 1), indicating that reduced photosynthetic rates were not due to reduced sink strength. The starch concentration did show a significant campaign effect and more than doubled from C1 to C2. Both sucrose and starch synthesis play important roles in the cellular recycling of phosphate for photosynthesis (Schlüter et al., 2013). A decrease in sugars and starch might lead to lower vitality and productivity of plants, as was previously observed in stressed C_4 leaves (da Silva and Arrabaça, 2004). In our experiment, while there was no effect of P for both sugars and starch, the campaign effect illustrated an increase of sugars and starch from C1 to C2, possibly suggesting that plants in all treatments experienced nutrient stress during C1. Moreover, the increasing sugar and starch levels between C1 and C2 further confirm that the low photosynthetic rates for the low P treatments were not due to reduced sink strength.

Foliar respiration rate is suppressed in the light. The abrupt decline in quantum yield of net CO_2 assimilation that occurs at very low light, often near the photosynthetic light compensation point, is also known as the 'Kok effect' (Kok, 1948). This light-induced inhibition of foliar respiration is reported to vary between 25%-100% (see references in Heskell et al., 2013) and is a source of uncertainty in current models of global terrestrial carbon cycling (Heskell and Tang, 2018). It can be impacted by environmental conditions such as temperature and soil nutrient availability (Heskell et al., 2012; Atkin et al., 2013). Here, the light-induced inhibition of respiration was highly variable among measured plants largely ranging from 0.3 to 0.5 with high uncertainty levels. Several studies showed that increased soil nutrient availability can relax the degree of light induced respiration, which was not confirmed in our experiment (Heskell et al., 2012; Atkin et al., 2013; Shapiro et al., 2004).

We applied 20 kg P ha^{-1} for the P treatment at which A_{max} reached its maximum value of about $27 \mu\text{mol m}^{-2} \text{ s}^{-1}$. Zhang et al. (2014) showed that the critical level of P application for maximal net photosynthetic rate of maize

(i.e. $30.3 \mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$) is between 15 and 28 kg ha^{-1} , which is in agreement with our study. Higher P application rates did not result in higher net photosynthetic rates. In our experiment the non-P-fertilized plants reached similar net photosynthetic rates, but only after colonization by AMF during C2. The campaign effect revealed in our experiment, i.e. the remarkable difference in P effect between C1 and C2, was associated with the (slow) establishment of AMF, which may suggest that increased plant P uptake following mycorrhization caused a recovery of the non-P-fertilized plants, and was beneficial for productivity in the P-fertilized plants as well (Verlinden et al., 2018). In the same experiment, we found that the partitioning to roots and to AMF was larger in the non-P-fertilized mesocosms as compared to the P-fertilized mesocosms (Verlinden et al., 2018). Interestingly, in the absence of AMF, plants that did not receive extra P died prematurely in pasteurized mesocosms not included in this study (but reported in Verlinden et al., 2018).

The similar leaf P concentrations in all treatments during C2 further supports our assumption of a strong stimulation of P-acquisition through mycorrhizae in the non-P-fertilized plants. The establishment of mycorrhizal symbioses is believed to be one of the most successful strategies to maximize the access of plant roots to available P and thus overcome P stress (Smith and Read, 2008; Sánchez-Calderón et al., 2010). The hyphal network of mycorrhizae extends over a very large surface area, increasing prominently the absorbing area of roots. Their extraradical hyphae extend beyond the P depletion zone, absorbing P that is otherwise not accessible for the plant (Plenchette et al., 2005; Roy-Bolduc and Hijri, 2011). Besides, mycorrhizal fungi improve phosphate solubility because they produce exudates that liberate P from the minerals (a.o. Smith et al., 2011; Burghelea et al., 2015; Kobae, 2019; Etesami et al., 2021; Jansa et al., 2021). For example glomalin, a glycoprotein secreted by AMF, aids the uptake of nutrients such as Fe and P that are difficult to dissolve (Miransari, 2010; Emran et al., 2017; Begum et al., 2019). Mycorrhizae thus significantly contribute to plant nutrition and to P uptake in particular (Wright et al., 2005), which in turn can positively affect leaf gas exchange rates (Smith and Read, 2008; Augé et al., 2016). In our experiment, the photosynthetic parameters increased, coinciding the mycorrhization-induced improved P nutrition in the non-P-fertilized plants.

The leaf-scale responses reported here correspond well to the ecosystem-scale GPP measurements reported for the same experiment in Verlinden et al. (2018). In the first weeks, both were (very) low in the absence of P addition, but showed a sudden increase about 6 weeks after planting. Although ecosystem level GPP remained lower for the non-P fertilized treatments, the photosynthesis system seemed to have fully recovered, as indicated by similar levels of leaf photosynthesis among all treatments during C2. These results are in line with the study by Řezáčová et al. (2018), who reported photosynthetic upregulation following upon the establishment of mycorrhizal symbiosis. Also our follow-up experiment with a P gradient confirmed the important stimulating role of AMF for plant productivity and photosynthesis (see Ven et al., 2020b).

To conclude, low P availability significantly decreased photosynthetic capacity, associated with reduced concentrations of photosynthetic enzymes and pigments. In contrast to the expected increase in nutrient stress because of further depletion of the soil as the growing season progressed, nutrient stress decreased over time and for most leaf processes, pigments and enzymes under study, the fertilization effect had disappeared two months after planting. Our results point towards a key role for the AMF-symbiosis and consequent increase of P uptake in explaining the vanishing P stress.

Data availability

Data will be published on Zenodo upon acceptance.

Author contributions

S.V. designed the experiment and research; M.S.V., H.A. and A.V. conducted the field- and lab work; M.S.V.
355 made first data analysis and draft of the manuscript; all authors provided expert advice and critically reviewed the
manuscript.

Competing interests

Sara Vicca is a member of the editorial board of Biogeosciences. The peer-review process was guided by an
independent editor, and the authors have also no other competing interests to declare.

360 Acknowledgements

This research was supported by the Research Foundation—Flanders (FWO) [grant numbers G0D5415N,
12U8918N, 12K0316N]; the European Research Council [grant number ERC-SyG-610028 IMBALANCE-P];
ClimMani COST Action [grant number ES1308]; the European Union Horizon 2020 Research and Innovation
program under the Marie Skłodowska-Curie grant (ReSPeC) [grant number 795299] and Methusalem funding of
365 the Research Council UA. We gratefully acknowledge Fred Kockelbergh and Marc Wellens for technical support;
Bart Hellemans, Lin Leemans, and Eddy De Smet for logistic support at the field site.

References

- AbdElgawad, H., Avramova, V., Baggerman, G., Van Raemdonck, G., Valkenburg, D., Van Ostade, X., Guisez, Y., Prinsen, E., Asard, H., Van den Ende, W. and Gerrit T. S. Beemster, G. T. S.: Starch biosynthesis contributes to the maintenance of photosynthesis and leaf growth under drought stress in maize. *Plant Cell Environ.*, 43, 2254–2271, doi: 10.1111/pce.13813, 2020.
- AbdElgawad, H., Peshev, D., Zinta, G., Van den Ende, W., Janssens, I. A. and Asard, H.: Climate extreme effects on the chemical composition of temperate grassland species under ambient and elevated CO₂: a comparison of fructan and non-Fructan accumulators. *Plos One*, 9, e92044, doi: 10.1371/journal.pone.0092044, 2014.
- Ashraf, M. and Harris, P. J. C.: Photosynthesis under stressful environments: An overview. *Photosynthetica*, 51, 163–190, doi: 10.1007/s11099-013-0021-6, 2013.
- Atkin, O. K., Turnbull, M. H., Zaragoza-Castells, J., Fyllas, N. M., Lloyd J., Meir, P. and Griffin, K. L.: Increased light inhibition of respiration as soil fertility declines along a post-glacial chronosequence in New Zealand. *Plant Soil*, 367, 163–182, doi: 10.1007/s11104-013-1686-0, 2013.
- Augé, R. M., Toler, H. D. and Saxton, A. M.: Mycorrhizal stimulation of leaf gas exchange in relation to root colonization, shoot size, leaf phosphorus and nitrogen: a quantitative analysis of the literature using meta-regression. *Front. Plant Sci.*, 7, 1084, doi: 10.3389/fpls.2016.01084, 2016.
- Augusto, L., Achat, D. L., Jonard, M., Vidal, D. and Ringeval, B.: Soil parent material—A major driver of plant nutrient limitations in terrestrial ecosystems. *Glob. Change Biol.*, 23, 3808–3824, doi: 10.1111/gcb.13691, 2017.
- Beauchamp, E. G. and Hamilton, H. A.: Optimum ratios of nitrogen and phosphorus fertilizers for corn determined by Homes' method of systematic variation. *Can. J. Plant Sci.*, 50, 141–150, doi: 10.4141/cjps70-027, 1970.
- Begum, N., Qin, C., Ahanger, M. A., Raza, S., Khan, M. I., Ashraf, M., Ahmed, N. and Zhang, L.: Role of arbuscular mycorrhizal fungi in plant growth regulation: implications in abiotic stress tolerance. *Front. Plant Sci.*, 10, 1068, doi: 10.3389/fpls.2019.01068, 2019.
- Bhagwat, A. S.: Activation of spinach ribulose 1,5- biphosphate carboxylase by inorganic phosphate. *Plant Sci. Lett.*, 23, 197–206, 1981.
- Brooks, A.: Effects of phosphorus nutrition on ribulose 1,5-bisphosphate carboxylase activation, photosynthetic quantum yield and amounts of some Calvin-cycle metabolites in spinach leaves. *Aust. J. Plant Physiol.* 13, 221–237, doi: 10.1071/PP9860221, 1986.
- Brooks, A., Woo, K. C. and Wong, S. C.: 1988 Effects of phosphorus nutrition on the response of photosynthesis to CO₂ and O₂, activation of ribulose bisphosphate carboxylase and amounts of ribulose bisphosphate and 3-phosphoglycerate in spinach leaves. *Photosyn. Res.*, 15, 133-141, doi: 10.1007/bf00035257, 1988.
- Burghilea, C., Zaharescu, D. G., Dontsova, K., Maier, R., Huxman, T. and Chorover, J.: Mineral nutrient mobilization by plants from rock: influence of rock type and arbuscular mycorrhiza. *Biogeochemistry*, 124, 187-203, doi: 10.1007/s10533-015-0092-5, 2015.
- Calderón-Vázquez, C., Alatorre-Cobos, F., Simpson-Williamson, J. and Herrera-Estrella, L.: Maize under phosphate limitation, in: *Handbook of Maize: Its Biology*, edited by: Bennetzen, J. L. and Hake, S. C., Springer, New York, USA, 381–404, doi: 10.1007/978-0-387-79418-1_19, 2009.

- Dall'Osto, L., Cazzaniga, S., Havaux, M. and Bassi, R.: Enhanced photoprotection by protein-bound vs free xanthophyll pools: a comparative analysis of chlorophyll b and xanthophyll biosynthesis mutants. *Mol. Plant.*, 3, 576-593, doi: 10.1093/mp/ssp117, 2010.
- 410 da Silva, J. M. and Arrabaça, M. C.: Contributions of soluble carbohydrates to the osmotic adjustment in the C₄ grass *Setaria sphacelata*: A comparison between rapidly and slowly imposed water stress. *J. Plant Physiol.*, 161, 551–555, doi: 10.1078/0176-1617-01109, 2004.
- Duursma, R.A: Plantecophys - An R Package for Analysing and Modelling Leaf Gas Exchange Data. *Plos One*, 415 10, e0143346, doi: 10.1371/journal.pone.0143346, 2015.
- Elser, J. J., Fagan, W. F., Denno, R. F., Dobberfuhl, D. R., Folarin, A., Huberty, A., Interlandi, S., Kilham, S. S., McCauley, E., Schulz, K. L., Siemann E. H. and Sterner, R. W.: Nutritional constraints in terrestrial and freshwater food webs. *Nature*, 408, 578-580, doi:10.1038/35046058, 2000. Emran, M., Rashad, M., Gispert, M., and Pardini, G.: Increasing soil nutrients availability and sustainability by glomalin in 420 alkaline soils. *Agricul. Biosystems Eng.*, 2, 74–84, 2017.
- Etesami, H., Jeong, B. R., and Glick, B. R.: Contribution of arbuscular mycorrhizal fungi, phosphate-solubilizing bacteria, and silicon to P uptake by plant. *Front. Plant Sci.*, 12, 1355, doi: 10.3389/fpls.2021.699618, 2021.
- Griffin, K. L. and Seemann, J. R.: Plants, CO₂ and photosynthesis in the 21st century. *Chem. Biol.*, 3, 245-254, 425 doi: 10.1016/S1074-5521(96)90104-0, 1996.
- Güsewell, S.: N:P ratios in terrestrial plants: Variation and functional significance. *New Phytol.*, 164, 243-266, doi: 10.1111/j.1469-8137.2004.01192.x, 2004.
- Halsted, M. and Lynch, J.P.: Phosphorus responses of C₃ and C₄ species. *J. Exp. Bot.*, 47, 497–505, doi: 10.1093/jxb/47.4.497, 1996.
- 430 Hartnett, D. C. and Wilson, G. W. T.: Mycorrhizae influence plant community structure and diversity in tallgrass prairie. *Ecology*, 80, 1187–1195, doi: 10.1890/0012-9658(1999)080[1187:MIPCSA]2.0.CO;2, 1999.
- Heskel, M. A., Anderson, O. R., Atkin, O. K., Turnbull, M. H. and Griffin, K. L.: Leaf- and cell-level carbon cycling responses to a nitrogen and phosphorus gradient in two Arctic tundra species. *Am. J. Bot.*, 99, 1702-1714, doi: 10.3732/ajb.1200251, 2012.
- 435 Heskel, M. A., Atkin, O. K., Turnbull, M. H., Griffin, K. L.: Bringing the Kok effect to light: A review on the integration of daytime respiration and net ecosystem exchange. *Ecosphere*, 4, 98, doi: 10.1890/ES13-00120.1, 2013.
- Heskel, M. A. and Tang, J.: Environmental controls on light inhibition of respiration and leaf and canopy daytime carbon exchange in a temperate deciduous forest. *Tree Physiol.*, 38, 1886–1902, doi: 440 10.1093/treephys/tpy103, 2018.
- Hoeksema, J. D., Chaudhary, V. B., Gehring, C. A., Johnson, N. C., Karst, J., Koide, R. T., Pringle, A., Zabinski, C., Bever, J. D., Moore, J. C., Wilson, G. W. T., Klironomos, J. N., Umbanhowar, J.: A meta-analysis of context-dependency in plant response to inoculation with mycorrhizal fungi. *Ecol. Lett.*, 13, 394–407, doi: 10.1111/j.1461-0248.2009.01430.x, 2010.
- 445 Jacob, J. and Lawlor, D. W.: Stomatal and mesophyll limitations of photosynthesis in phosphate deficient sunflower, maize and wheat plants. *J. Exp. Bot.*, 42, 1003–1011, doi: 10.1093/jxb/42.8.1003, 1991.

- Jacob, J. and Lawlor, D. W.: Dependence of photosynthesis of sunflower and maize leaves on phosphate supply, ribulose-1,5-bisphosphate carboxylase/oxygenase activity, and ribulose-1,5-bisphosphate pool size. *Plant Physiol.*, 98, 801-807, doi: 10.1104/pp.98.3.801, 1992.
- 450 Jahns, P. and Holzwarth, A. R.: The role of the xanthophyll cycle and of lutein in photoprotection of photosystem II. *Biochim. Biophys. Acta*, 1817, 182–193, doi: 10.1016/j.bbabi.2011.04.012, 2012.
- Jansa, J., Finlay, R., Wallander, H., Smith, F. A. and Smith, S. E.: Role of mycorrhizal symbioses in phosphorus cycling, in: *Phosphorus in Action. Soil Biology*, vol 26, edited by: Bünemann, E., Oberson, A. and Frossard, E., Springer, Berlin, Heidelberg, Germany, 137-168, doi: 10.1007/978-3-642-15271-9_6, 2011.
- 455 Jeannette, E., Reyss, A., Gregory, N., Gantet, P., and Prioul, J. L.: Carbohydrate metabolism in a heat-girdled maize source leaf. *Plant Cell Environ.*, 23, 61-69, doi: 10.1046/j.1365-3040.2000.00519.x, 2000.
- Johnston, A. E. and Poulton, P. R.: Phosphorus in Agriculture: A Review of Results from 175 Years of Research at Rothamsted, UK. *J. Environ. Qual.*, 48, 1133–1144, doi: 10.2134/jeq2019.02.0078, 2019.
- Khan, A., Lu, G., Ayaz, M., Zhang, H., Wang, R., Lv, F., Yang, X., Sun, B. and Zhang, S.: Phosphorus efficiency, soil phosphorus dynamics and critical phosphorus level under long-term fertilization for single and double cropping systems. *Agr. Ecosyst. Environ.*, 256, 1-11, doi: 10.1016/j.agee.2018.01.006, 2018.
- 460 Kobae, Y.: Dynamic phosphate uptake in arbuscular mycorrhizal roots under field conditions. *Front. Environ. Sci.*, 6, 159, doi: 10.3389/fenvs.2018.00159, 2019.
- Koerselman, W. and Meuleman, A. F. M.: The vegetation N:P ratio: a new tool to detect the nature of nutrient limitation. *J. Appl. Ecol.*, 33, 1441-1450, doi: 10.2307/2404783, 1996.
- 465 Kok, B.: A critical consideration of the quantum yield of *Chlorella*-photosynthesis. *Enzymologia*, 13, 1-56, 1948.
- Kuczyńska, P., Latowski, D., Niczyporuk, S., Olchawa-Pajor, M., Jahns, P., Gruszecki, W. I. and Strzałka, K.: Zeaxanthin epoxidation - an in vitro approach. *Acta Biochim. Pol.* 59, 105–107, doi: 10.18388/abp.2012_2182, 2012.
- 470 Lewis, J. D., Griffin, K. L., Thomas, R. B. and Strain, B. R.: Phosphorus supply affects the photosynthetic capacity of loblolly pine grown in elevated carbon dioxide. *Tree Physiol.*, 14, 1229-1244, doi: 10.1093/treephys/14.11.1229, 1994.
- Loustau, D., Brahim, M. B., Gaudillère, J. P. and Dreyer, E.: Photosynthetic responses to phosphorus nutrition in two-year-old maritime pine seedlings. *Tree Physiol.*, 19, 707-715, doi: 10.1093/treephys/19.11.707, 1999.
- 475 Miner, G. L. and Bauerle, W. L.: Seasonal responses of photosynthetic parameters in maize and sunflower and their relationship with leaf functional traits. *Plant Cell Environ.*, 42, 1561-1574, doi: 10.1111/pce.13511, 2019.
- Parry, M. A., Keys, A. J., Madgwick, P. J., Carmo-Silva, A. E. and Andralojc, P. J.: Rubisco regulation: a role for inhibitors. *J. Exp. Bot.*, 59, 1569-1580, doi: 10.1093/jxb/ern084, 2008.
- 480 Paul, M. J. and Foyer, C. H.: Sink regulation of photosynthesis. *J. Exp. Bot.*, 52, 1383-1400, doi: 10.1093/jexbot/52.360.1383, 2001.
- Paul, M. J. and Stitt, M.: Effects of nitrogen and phosphorus deficiencies on levels of carbohydrates, respiratory enzymes and metabolites in seedlings of tobacco and their response to exogenous sucrose. *Plant Cell Environ.*, 16, 1047-1057, doi: 10.1111/j.1365-3040.1996.tb02062.x, 1993.
- 485 Peñuelas, J., Poulter, B., Sardans, J., Ciais, P., van der Velde, M., Bopp, L., Boucher, O., Godderis, Y., Hinsinger, P., Llusia, J., Nardin, E., Vicca, S., Obersteiner, M. and Janssens, I. A.: Human-induced nitrogen-

- phosphorus imbalances alter natural and managed ecosystems across the globe. *Nat. Commun.*, 4 , 2934, doi:10.1038/ncomms3934, 2013. Pieters, A. J., Paul, M. J. and Lawlor, D. W.: Low sink demand limits photosynthesis under P(i) deficiency. *J. Exp. Bot.*, 52, 1083-1091, doi: 10.1093/jexbot/52.358.1083, 2001.
- 490 Plenchette, C., Clermont-Dauphin, C., Meynard, J. M. and Fortin, J. A.: Managing arbuscular mycorrhizal fungi in cropping systems. *Can. J. Plant Sci.*, 85, 31-40, doi: 10.4141/P03-159, 2005.
- Rao, I. M. and Terry, N.: Leaf phosphate status, photosynthesis, and carbon partitioning in sugar beet I. Changes in growth, gas exchange, and Calvin cycle enzymes. *Plant Physiol.*, 90, 814-819, doi: 10.1104/pp.90.3.814, 1989.
- 495 Řezáčová, V., Slavíková, R., Zemková, L., Konvalinková, T., Procházková, V., Št'oviček, V., Hršelová, H., Beskid, O., Hujšlová, M., Gryndlerová, H., Gryndler, M., Püschel, D. and Jansa, J.: Mycorrhizal symbiosis induces plant carbon reallocation differently in C₃ and C₄ Panicum grasses. *Plant Soil*, 425, 441–456, doi: 10.1007/s11104-018-3606-9, 2018.
- 500 Rillig, M. C., Field, C. B., Allen, M. F.: Soil biota responses to long-term atmospheric CO₂ enrichment in two California annual grasslands. *Oecologia*, 119, 572–577, doi: 10.1007/s004420050821, 1999.
- Rodríguez, D., Andrade, F. H. and Goudriaan, J.: Does assimilate supply limit leaf expansion in wheat grown in the field under low phosphorus availability? *Field Crops Res.*, 67, 227-238, doi: 10.1016/s0378-4290(00)00098-8, 2000.
- 505 Rodríguez, D. and Goudriaan, J.: Effects of phosphorus and drought stresses on dry-matter and phosphorus allocation in wheat. *J. Plant Nutr.*, 18, 2501-2517, doi: 10.1080/01904169509365080, 1995.
- Rodríguez, D., Keltjens, W. G. and Goudriaan, J.: Plant leaf area expansion and assimilate production in wheat (*Triticum aestivum* L.) growing under low phosphorus conditions. *Plant Soil*, 200, 227-240. doi:10.1023/a:1004310217694, 1998.
- 510 Rogers, A.: The use and misuse of V_{c,max} in Earth System Models. *Photosynth. Res.*, 119, 15-29, doi: 10.1007/s11120-013-9818-1, 2014.
- Roy-Bolduc, A. and Hijri, M.: The use of mycorrhizae to enhance phosphorus uptake: A way out the phosphorus crisis. *J. Biofertil. Biopestici.*, 2, 104, doi: 10.4172/2155-6202.1000104, 2011.
- Sánchez-Calderón, L., Chacon-López, A., Perez-Torres, C. and Herrera-Estrella, L.: Phosphorus: Plants strategies to cope with its scarcity, in: *Cell Biology of Metals and Nutrients. Plant Cell Monographs*, vol 17, edited by: Hell, R. and Mendel, R.R., Springer, Berlin, Heidelberg, Germany, 173–198, doi: 10.1007/978-3-642-10613-2_8, 2010.
- 515 Schlüter, U., Colmsee, C., Scholz, U., Bräutigam, A., Weber, A. P. M., Zellerhoff, N., Bucher, M., Fahnenstich, H. and Sonnewald, U.: Adaptation of maize source leaf metabolism to stress related disturbances in carbon, nitrogen and phosphorus balance. *BMC Genomics* 14, 442, doi: 10.1186/1471-2164-14-442, 2013.
- 520 Schulze, E.-D., Beck, E. and Müller-Hohenstein, K. (Eds.): *Plant Ecology*, Springer, Berlin, Heidelberg, Germany, 2005.
- Shapiro, J. B., Griffin, K. L., Lewis, J. D. and Tissue, T. D.: Response of *Xanthium strumarium* leaf respiration in the light to elevated CO₂ concentration, nitrogen availability and temperature. *New Phytol.*, 162, 377–386, doi: 10.1111/j.1469-8137.2004.01046.x, 2004.
- 525

- Smith, S. E., Jakobsen, I., Grønlund, M. and Smith, F. A.: Roles of arbuscular mycorrhizas in plant phosphorus nutrition: Interactions between pathways of phosphorus uptake in arbuscular mycorrhizal roots have important implications for understanding and manipulating plant phosphorus acquisition. *Plant Physiol.*, 156, 1050-1057, doi: 10.1104/pp.111.174581, 2011.
- 530
- Smith, S. E. and Read, D. J.: *Mycorrhizal Symbiosis* (Third Edition), Academic Press, London, UK, 2008.
- Sulpice, R., Tschoep, H., von Korff, M., Bussis, D., Usadel, B., Hohne, M., Witucka-Wall, H., Altmann, T., Stitt, M. and Gibon, Y.: Description and applications of a rapid and sensitive non-radioactive microplate-based assay for maximum and initial activity of D-ribulose-1,5-bisphosphate carboxylase/oxygenase. *Plant Cell Environ.*, 30, 1163-1175, doi: 10.1111/j.1365-3040.2007.01679.x, 2007.
- 535
- Temminghoff, E. E. J. M. and Houba, V. J. G.: *Plant Analysis Procedures*. Springer Netherlands, Dordrecht, doi: 10.1007/978-1-4020-2976-9, 2004.
- Tennant, D.: A test of a modified line intersect method of estimating root length. *J. Ecol.*, 63, 995-1001, doi: 10.2307/2258617, 1975.
- 540
- Thum, T., Caldararu, S., Engel, J., Kern, M., Pallandt, M., Schnur, R., Yu, L. and Zaehle, S.: A new model of the coupled carbon, nitrogen, and phosphorus cycles in the terrestrial biosphere (QUINCY v1.0; revision 1996). *Geosci. Model Dev.*, 12, 4781-4802, doi:10.5194/gmd-12-4781-2019, 2019.
- Usuda, H. and Shimogawara, K.: Phosphate deficiency in maize. I. Leaf phosphate status, growth, photosynthesis and carbon partitioning. *Plant Cell Physiol.*, 32, 497-504, doi: 10.1093/oxfordjournals.pcp.a078107, 1991.
- 545
- Ven, A., Verbruggen, E., Verlinden, M. S., Olsson, P. A., Wallander, H. and Vicca, S.: Mesh bags underestimated arbuscular mycorrhizal abundance but captured fertilization effects in a mesocosm experiment. *Plant Soil*, 446, 563–575, doi: 10.1007/s11104-019-04368-4, 2020a.
- Ven, A., Verlinden, M. S., Fransen, E., Olsson, P. A., Verbruggen, E., Wallander, H. and Vicca, S.: Phosphorus addition increased carbon partitioning to autotrophic respiration but not to biomass production in an experiment with *Zea mays*. *Plant Cell Environ.*, 43, 2054–2065, doi: 10.1111/pce.13785, 2020b.
- 550
- Veneklaas, E. J., Lambers, H., Bragg, J., Finnegan, P. M., Lovelock, C. E., Plaxton, W. C., Price, C., Scheible, W.-R., Shane, M. W., White, P. J. and Raven, J. A.: Opportunities for improving phosphorus-use efficiency in crop plants. *New Phytol.*, 195, 306-320, doi: 10.1111/j.1469-8137.2012.04190.x., 2012.
- 555
- Verlinden, M. S., Broeckx, L. S., Zona, D., Berhongaray, G., De Groote, T., Camino Serrano, M., Janssens, I. A. and Ceulemans, R.: Net ecosystem production and carbon balance of an SRC poplar plantation during its first rotation. *Biomass Bioenerg.* 56, 412-422, doi: 10.1016/j.biombioe.2013.05.033, 2013.
- Verlinden, M. S., Ven, A., Verbruggen, E., Janssens, I. A., Wallander, H. and Vicca, S.: Favorable effect of mycorrhizae on biomass production efficiency exceeds their carbon cost in a fertilization experiment. *Ecology*, 99, 2525–2534, doi: 10.1002/ecy.2502, 2018.
- 560
- Vicca, S., Stocker, B., Reed, S. C., Wieder, W. R., Bahn, M., Fay, P. A., Janssens, I. A., Lambers, H., Peñuelas, J., Piao, S., Rebel, K., Sardans, J., Sigurdsson, B. D., Van Sundert, K., Wang, Y., Zaehle, S., Ciais, P.: Using research networks to create the comprehensive datasets needed to assess nutrient availability as a key determinant of terrestrial carbon cycling. *Environ. Res. Lett.*, 13, 125006, doi:10.1088/1748-9326/aaeae7, 2018.
- 565

- Vierheilig, H., Schweiger, P. and Brundrett, M.: An overview of methods for the detection and observation of arbuscular mycorrhizal fungi in roots. *Physiol. Plantarum*, 125, 393-404, doi: 10.1111/j.1399-3054.2005.00564.x, 2005.
- 570 Vitousek, P. M., Porder, S., Houlton, B. Z. and Chadwick, O. A.: Terrestrial phosphorus limitation: mechanisms, implications, and nitrogen-phosphorus interactions. *Ecol. Appl.*, 20, 5-15, doi:10.1890/08-0127.1, 2010.
- von Caemmerer, S. (Ed.): *Biochemical models of leaf photosynthesis*. CSIRO Publishing, Melbourne, Australia, 2000.
- Wang, Y. P., Law, R. M., and Pak, B.: A global model of carbon, nitrogen and phosphorus cycles for the terrestrial biosphere. *Biogeosciences*, 7, 2261-2282, doi:10.5194/bg-7-2261-2010, 2010.
- 575 Wieder, W. R., Cleveland, C. C., Smith, W. K., and Todd-Brown, K.: Future productivity and carbon storage limited by terrestrial nutrient availability. *Nat. Geosci.*, 8, 441-444, doi:10.1038/ngeo2413, 2015.
- Winter, H. and Huber, S. C.: Regulation of sucrose metabolism in higher plants: localization and regulation of activity of key enzymes. *Crit. Rev. Biochem. Mol. Biol.*, 35, 253-289, doi: 10.1080/10409230008984165, 2000.
- 580 Wright, D. P., Scholes, J. D., Read, D. J. and Rolfe, S. A.: European and African maize cultivars differ in their physiological and molecular responses to mycorrhizal infection. *New Phytol.*, 167, 881-896, doi: 10.1111/j.1469-8137.2005.01472.x, 2005.
- Zhang, K., Liu, H., Tao, P. and Chen, H.: Comparative proteomic analyses provide new insights into low phosphorus stress responses in maize leaves. *Plos One*, 9, e98215, doi: 10.1371/journal.pone.0098215, 585 2014.