

Climatic Variation Drives Loss and Restructuring of Carbon and Nitrogen in Boreal Forest Wildfire

Johan A. Eckdahl^{1,2}, Jeppe A. Kristensen³, and Daniel B. Metcalfe²

¹Department of Physical Geography and Ecosystem Science, Lund University, Lund, Sweden

²Department of Ecology and Environmental Science, Umeå University, Umeå, Sweden

³School of Geography and the Environment, University of Oxford, Oxford, United Kingdom

Correspondence: Johan Eckdahl (Johan.Eckdahl@nateko.lu.se)

Abstract. The boreal forest landscape covers approximately 10% of the earth's land area and accounts for almost 30% of the global annual terrestrial sink of carbon (C). Increased emissions due to climate change-amplified fire frequency, size and intensity threaten to remove elements such as C and nitrogen (N) from forest soil and vegetation at rates faster than they accumulate. This may result in large areas within the region becoming a net source of greenhouse gases creating a positive feedback loop with a changing climate. Meter-scale estimates of area-normalized fire emissions are ~~regionally limited~~ limited in Eurasian boreal forests and knowledge of their relation to climate and ecosystem properties is sparse. This study sampled 50 separate Swedish wildfires, which occurred during an extreme fire season in 2018, providing quantitative estimates of C and N loss due to fire along a climate gradient. Mean annual precipitation had strong positive effects on total fuel, which was the strongest driver for increasing C and N losses. Mean annual temperature (MAT) influenced both pre- and postfire organic layer soil bulk density and C:N ratio which had mixed effects on C and N losses. Significant fire-induced loss of C estimated in the 50 plots was comparable to estimates in similar Eurasian forests but approximately a quarter of those found in typically more intense North American boreal wildfires. N loss was insignificant though a large amount of fire affected fuel was converted to a low C:N surface layer of char in proportion to increased MAT. These results reveal large quantitative differences in C and N losses between global regions and their linkage to the broad range of climate conditions within Fennoscandia. A need exists to better incorporate these factors into models to improve estimates of global emissions of C and N due to fire in future climate scenarios. Additionally, this study demonstrated a linkage between climate and the extent of charring of soil fuel and discusses its potential for altering C and N dynamics in postfire recovery.

1 Introduction

Worldwide, boreal forests cover approximately 10% of land area (Keenan et al., 2015) and account for a net ~~E~~ carbon (C) sink into plants and soil of 0.31 ± 0.19 ~~Pg of~~ Pg C per year, equivalent to $27.3 \pm 16.7\%$ of the planet's terrestrial C sink (Tagesson et al., 2020). This sink plays a pivotal role in the greenhouse gas content of the atmosphere (Lemprière et al., 2013). Low temperatures and often waterlogged soil conditions reduce the rate of decomposition relative to litter additions resulting in the buildup of thick layers of soil organic material where the majority of C is stored (Malhi et al., 1999; Rapalee et al., 1998). The balance of C transferred between atmospheric and terrestrial stocks on the yearly timescale is dictated by rates of

25 terrestrial net primary production and respiration which are themselves controlled by temperature, moisture and nitrogen (N) availability (Deluca and Boisvenue, 2012).

The predominant disturbance to this C balance in the boreal region is outbreaks of wildfires which reoccur in individual forest stands at the centurial timescale (Bond-Lamberty et al., 2007). Among the immediate effects of fire are a substantial release of C to the atmosphere from soil and vegetation (Schultz et al., 2008) as well as physical restructuring of the habitat
30 through varying degrees of over- and understory removal and changes in soil properties such as bulk density (Certini, 2005). Further, bioavailability of energy sources and nutrients is substantially affected as elements such as C and N are lost and their chemical structure is altered by heating, i.e. charred (Pellegrini et al., 2021; Neff et al., 2005). Along with a changing climate, these effects influence community structure and processes such as soil respiration and nutrient cycling, which can shape future forest C and N cycles on decadal to centurial timescales (Johnstone et al., 2020; Mekonnen et al., 2019). Changing patterns of
35 temperature and precipitation in recent decades have caused increases in frequency, intensity and size of fires across the global boreal region with further amplification predicted in the future (de Groot et al., 2013b; Gillett et al., 2004; Kelly et al., 2013; de Groot et al., 2003). Increasing emissions of C to the atmosphere due to fire and changing C cycling in recovering ecosystems may accelerate climate change (Li et al., 2017).

Both long (~~variation measured interannually~~interannual variability) and short-term (intraannual variability) processes have
40 been identified as drivers of the dynamics of fire events. Particularly, in North American boreal forests, the strongest driver of area-normalized emissions of C ~~in boreal from~~ wildfires appears to be total fuel (i.e. potentially combustible organic material) (Walker et al., 2018, 2020a, b). ~~The accumulation of fuel itself~~ Fuel accumulation has been found to be strongly related to stand age and longer-term moisture levels (a factor of drainage conditions, evapotranspiration and inputs through precipitation and lateral water flow) (Walker et al., 2018, 2020a, b). However, in order for this fuel to be available to be
45 ignited and sustain fire, it must be both adequately dried and located within sufficient proximity to high heat and oxygen exposure during an active fire. This instantaneous proportion of total fuel that is readily combustible is termed fuel availability. Therefore, boreal wildfire models often incorporate short-term fire weather variables (e.g. drought indices, temperature, wind speed, relative humidity) and separate soil fuel loads into distinct compartments such as surface litter (influencing ignition and rate of spread) and the more compactly arranged layers below (acting as a heat reservoir that supports
50 extended smoldering over days to weeks) (de Groot et al., 2003; Van Wagner, 1987; Rabin et al., 2017; Kasischke et al., 2005; Wiggins et al., 2020). Composition of tree species, with their associated fire adaptation strategies, has also been shown to have a strong impact on fire severity and intensity and distinguishes the boreal wildfire regimes of the North American and Eurasian continents (Rogers et al., 2015). Furthermore, climate has been observed to have a conditioning effect on fuel chemical composition through its control over vegetation characteristics and the decomposition state of their detrital inputs, which are often represented by the C:N weight ratio in soils (Vanhala et al., 2008; Kohl et al., 2018). Fuel
55 chemical composition, arrangement, moisture content, applied heat and oxygen availability in turn have all been related to the efficiency of the combustion reaction during fire and therefore emission chemistry and the charring of remaining fuel, which can form a surface pyrogenic layer with C and N weight concentrations that differ from the original material (Santín et al., 2016; Dymov et al., 2021; Schmidt and Noack, 2000). Production of charred material is an innately

60 fire driven process and a representative measure of these interrelated effects. Because additions of char to soils have been observed to have strong impacts on C storage and nutrient cycling ([Schmidt and Noack, 2000](#); [Preston and Schmidt, 2006](#)) ([Pellegriani et al., 2021](#); [Schmidt and Noack, 2000](#); [Preston and Schmidt, 2006](#)), this fire-induced transformation of remaining fuel is valuable to study alongside its loss from the ecosystem during wildfire.

A general approach to estimating boreal C emissions due to a single wildfire can be taken by multiplying total area burned by estimates of area-normalized C emissions (French et al., 2004; van der Werf et al., 2017). While total area burned may be evaluated directly through remote sensing (Giglio et al., 2018; Ruiz et al., 2012), estimating area-normalized C emissions entails a variety of approaches. These approaches are most often derived from the results of labor intensive [wildfire](#) field sampling or controlled burns which are extrapolated to the larger scale either directly or through weighting by remotely sensed data (e.g. topography, vegetation cover, aerosol density) or poorly constrained parameters such as total fuel load (French et al., 2004; Soja et al., 2004; van der Werf et al., 2017; Veraverbeke et al., 2015; Kaiser et al., 2012). This field sampling has been regionally limited and biased towards a few high intensity burn complexes in North America ([Veraverbeke et al., 2021](#)) which may in turn bias global emission estimates ([Veraverbeke et al., 2021](#); [van Leeuwen et al., 2014](#); [Akagi et al., 2011](#)) ([van Leeuwen et al., 2014](#); [Akagi et al., 2011](#)). For example, the Eurasian boreal region is dominated by relatively fire resistant overstory vegetation that avoids excessive heating by promoting lower intensity ground and surface fires than that in boreal North America, which is more prone to spread rapid flaming combustion throughout the canopy (Rogers et al., 2015; de Groot et al., 2013a). C losses in surface fires in Siberian boreal pine forests were found to be 0.88 and 1.69 kg C m⁻² under experimental and wild conditions, respectively (Ivanova et al., 2011; Kukavskaya et al., 2017). These are between one quarter to a half of what is typical in North American wildfire (3.3 kg C m⁻²) (Boby et al., 2010; Walker et al., 2020a). Although Eurasia contains over 70% of the boreal global land area (de Groot et al., 2013a) and about 50% (20 Mha yr⁻¹) of its yearly burnt area (Rogers et al., 2015), methodologies for estimating global and regional C emissions are severely lacking ground validation and meter scale assessments of drivers of C loss variability from this continent (Veraverbeke et al., 2021; van der Werf et al., 2017; Kaiser et al., 2012). Additionally, field measurements of total N loss from wildfire are rare in all boreal regions despite its recognized role as a limiting nutrient and evidence of its immediate removal in similar percentages to C (Boby et al., 2010). Lastly, direct field sampling of boreal wildfire has often focused on individual or small groups of fires located relatively near to each other, with little information about the representativeness of the observations or context of the results within the broader spectrum of fire impacts across the wider region, especially those relating to variation in climate. This knowledge gap has thus far been addressed with conglomerated studies spanning different fire seasons, ecosystem types and research methodologies (Walker et al., 2020a; Gaboriau et al., 2020). Therefore, widely replicated, ~~simultaneous and~~ systematic field measurements of [single-season](#) fire processes in under sampled regions, with particular attention to climate, are needed to derive more robust, generalizable conclusions about boreal forest responses to wildfire.

This study sampled 50 separate fire complexes spanning broad gradients of mean annual temperature (MAT) and precipitation (MAP) which ignited in Sweden during summer 2018 (Fig. 1). This summer, along with that of 2014, were two of the most extreme fire seasons within Sweden in recent history, driven by severe drought conditions (Wilcke et al., 2020). The goal of this study is to distinguish the effects of climate on fire-induced changes in C and N stocks in an undersampled region with

95 in situ, fine scale measurements that both provide insight into local processes and allow for global comparison. Space-for-time substitution (De Frenne et al., 2013) along with a paired-sample design (Boby et al., 2010; Granath et al., 2021) provided insight into possible future conditions of Fennoscandian forests under a changing climate and fire regime. Specifically, it was hypothesized that:

1. Fire significantly reduced C and N stocks across forest compartments.
- 100 2. Fire restructured organic layer C and N stocks by increasing overall bulk density and adjusting their weight concentrations across residual compartments and a newly formed pyrogenic layer.
3. Loss of soil and understory C and N stocks and their transfer to the surface pyrogenic layer were correlated to prefire total fuel amount, composition and distribution amongst forest compartments.
4. A direct relation between climate variables and fire-induced C and N stock changes exists.
- 105 5. The relation between climate and fire driven C and N stock changes is mediated by long-term ecosystem properties that affect the amount of forest fuel removed during fire.

2 Materials and methods

2.1 Experimental design and field site selection

50 burnt plots were selected from a pool of 325 fires identified during the summer 2018 period that had perimeters manually mapped by the Swedish Forest Agency. Perimeters were drawn around burn scars identified using Normalized Burn Ratio (NBR) values derived from Sentinel-2 ~~bottom-of-the-atmosphere~~bottom-of-atmosphere corrected bands 8 and 12. Each of the 20 × 20 m² plots were located within distinct burn scars (greater than 2 km separation) to reduce potential for pseudoreplication or spatial autocorrelation (Bataineh et al., 2006) and allow for increased spread across the climate gradients (Schweiger et al., 2016). Constraints were placed on plot selection using spatial data within the QGIS (QGIS Development Team, 2019) and ArcGIS (Esri Inc., 2019) software environments. Plot-wide values for raster data were taken as the average pixel value within a 20 m diameter circle centered on the plot.

The first constraints on site selection were to avoid wetland or steeply sloping areas. Topo-edaphic derived soil moisture data (TEM) was provided at 10 m resolution by the Swedish Environmental Protection Agency (Naturvårdsverket, 2018). TEM was given as integer values ranging from 0 to 240 (in order of increasing moisture potential) and was based on the 120 Soil Topographic Wetness Index (Buchanan et al., 2014) in areas where soil type information was available and on the two topographic indices Depth to Water (Murphy et al., 2007) and the Topographic Wetness Index (Beven and Kirkby, 1979) where soil information was unavailable. These measures gave an estimate of soil drainage, which can be predictive of long-term soil moisture patterns (Walker et al., 2020b). Elevation data was provided by the Swedish Mapping, Cadastral and Land Registration Authority from a 50 m resolution digital elevation model (Lantmäteriet, 2021). Slope was calculated using the

125 “slope” function within the ArcGIS software environment. Using these datasets all plots were selected to have below both 150
plot-averaged TEM units and 15° slope. This restricted the study to non-wetland ecosystem types, since wetlands tend to have
markedly differing ecosystem functioning than relatively drier forested regions, and to retain focus on climate driven effects
and their space-for-time substitution by reducing the effects of exogenous variables such as topography on models. To further
isolate fire effects in response to variation within a single ecosystem type, Swedish Forest Agency data on tree species was
130 used to select only Scots pine (*Pinus sylvestris*) dominated stands. These stands were determined to be at least 30 years of age
using data from the Swedish University of Agricultural Sciences (Dept. of Forest Resource Management, 2015). The relatively
dry Swedish forested area is predominantly made up of podzols with distinct separation of a low C mineral layer from a high C
organic layer above (Olsson et al., 2009). It is assumed the dominant properties of these layers that affect fire severity are soil
drainage (TEM) and fuel loading (organic layer C). Additional soil property data sources were not available at a high enough
135 resolution and ~~therefore~~, therefore, no further classification of soil properties between and across plot-pairs was performed to
determine site selection.

Next, sites with postfire salvage logging were omitted using recent visual imagery along with Swedish Forest Agency
records. Burnt plots were selected to maximize spacing over latitude, MAT and MAP gradients (Schweiger et al., 2016).
MAT and MAP values were derived from daily records averaged over the period 1961-2017 that were extracted from a 4 km
140 resolution hydrological modeling dataset provided by the Swedish Meteorological and Hydrological Institute (Olsson et al.,
2013). The above mentioned Sentinel-2 imagery was used to locate burnt plots as close to the highest NBR pixel values within
each of their mapped final burn scar perimeters while satisfying all other plot selection requirements. This method intended to
add consistency to plot selection ~~within the larger burn scars~~ by placing sampled regions within the most ~~developed portions of~~
~~each fire~~ highly fire-impacted regions as measured by NBR. Thereby, the potential for observed differences in fire effect between
145 burn scars due to varied spatial proximity to ~~the fire periphery~~ these highest impacted regions is assumed to be reduced. NBR
values were otherwise not used for comparison across separate burn scars. Visual and infrared image processing was performed
in May 2019 through the brandkarta web application provided by the Swedish Forest Agency. No criteria for fire severity were
used for plot selection outside of their detectability using the above methods and data sources.

In order to estimate the effects of fire, prefire properties of each burnt plot were approximated using measurements from
150 an identically-sized, adjacent control plot centered between 15 and 150 m (58 m average) outside the fire boundaries (100
plots total, i.e. 50 plot pairs). A major limitation to this approximation is that observed differences within plot pairs may
be skewed through inaccurate or imprecise representation of prefire burnt plot properties by control plot measurements. An
attempt to reduce these errors was made by incorporating a large sample size ($n = 50$) and strict controls on the matching of
important ecosystem variables. In order to reduce disparities between control and prefire burnt plot properties the following
155 restrictions were placed on control plot selection. Control plot locations were selected to minimize elevation, slope and TEM
differences from the adjacent burnt plot. Swedish Forest Agency data regarding tree species, overstory biomass, and basal
area (collected during 2014) were used to best match properties of control and burnt plot pairs. As a complement to the
quantitative overstory data, aerial and satellite images provided by the Swedish National Land Survey were examined. This

gave quick visual verification that plot pairs, and the area between them, had not been disturbed by fire or logging over the past 30 years and that stand structure appeared homogeneous and uninterrupted by roads or other artificial structures.

Due to their documented effects on C emissions (Walker et al., 2018), metrics related to both short-term and long-term soil moisture conditions were considered in this study. Long-term moisture conditions were separated into a topo-edaphic component (i.e. TEM) and a climatic component (i.e. MAP and MAT). TEM was considered to be representative of the soil drainage which has been observed to relate to C emissions in separate study (Walker et al., 2018). MAP and MAT were considered to have an additional effect on moisture balance through their controls on direct water input and evapotranspiration. The shorter term moisture balance of summer 2018 was assessed with the Standardized Precipitation-Evapotranspiration Index (SPEI) due to its observed relationship to fuel drying and fire activity in boreal Eurasia (Chen et al., 2021; Ponomarev et al., 2016). SPEI was calculated over the first 6 months of 2018 at 0.5° spatial resolution within the SPEIBase data source to capture the extended desiccation process leading up to each fire (Beguería et al., 2019). Due to limited temporal information on 2018 fire activity, common fire weather metrics were unavailable and were instead approximated by summer 2018 anomalies in temperature (Δ MAT) and precipitation (Δ MAP), i.e. the difference in the 2018 June, July, and August average of these values from those during the same months averaged over the period from 1961 to 2017. These two metrics were used in regression for direct comparison to SPEI and to explain C and N losses.

2.2 Sampling

Site visits occurred approximately 1 year postfire ~~over the dates August between 5 to August and 20 August~~ in 2019. This 1 year delay intended to capture the more immediate effects of fire due to potential rapid spikes in nutrient losses and tree mortality, which generally occur within the first year, but avoid the accumulation of discrepancies in C and N stocks relative to control plots due to any longer term differences in rates of decomposition, leaching and litter addition (Granath et al., 2021; Certini, 2005; Sidoroff et al., 2007).

Sampling and analysis were broken into six forest compartments. The compartments included four soil layers ~~(i.e. mineral, duff, moss/litter and char)~~, as well as the two aboveground compartments ~~(i.e. understory and overstory vegetation)~~. The organic layer was defined as the duff, moss/litter and char layers grouped together while the soil category was considered as the organic layer grouped with the mineral layer. The category called “total” refers to the grouping of the soil and understory, but excludes the overstory due to its ~~low observation of minimal~~ combustion (as judged by ~~amounts observed scarcity~~ of canopy blackening). ~~While larger dead wood lying on top of the forest floor (woody Woody~~ debris mixed in the moss/litter layer was sampled in this study) ~~), though not the coarse woody debris laying on top of this layer. While larger dead wood lying on top of the forest floor~~ can contribute to C and N stocks and their losses due to fire, this material is typically of low prevalence in Sweden (Jonsson et al., 2016), difficult to accurately measure and ~~lacks~~ standard methodology to estimate its consumption by fire ~~(Jonsson et al., 2016)~~ is only beginning to be developed (Dieleman et al., 2020). Therefore, focus was retained on the variation of the larger and more precisely readily measurable ecosystem C and N pools in soil and living vegetation.

2.2.1 Soil

Soil horizon depths (i.e. the distance from bottom to top of each individual layer) of the mineral, duff, moss/litter, and char layers were measured at 20 points per plot from 10 equally spaced excavations along each plot diagonal (Kristensen et al., 2015). The mineral layer was measured from its highest rock obstruction to the bottom of the duff layer. The duff layer was considered the grouping of the F (partially decomposed material) and H (humic material) layers in accordance with the Canadian system of soil classification (Canada Soil Survey Committee, 1978), as is common in boreal wildfire literature. The moss/litter layer was all unburnt material on top of the duff layer, including visually identifiable detritus and living moss. In all burnt sites, a layer of conglomerated char formed a clear boundary on top of the moss/litter allowing for distinct measurement. Here, char is defined as fully blackened, brittle material with apparent high heat exposure due to fire. This separation was made based on large observed differences in C and N concentrations in surface pyrogenic layers compared to lower residual layers in similar ecosystems (Santín et al., 2016; Dymov et al., 2021; Bodí et al., 2014)

Samples were acquired for all four soil layers. Four mineral soil samples were taken using a 3 cm diameter, 40 cm long gouge auger corer at four corners of a square each 15 m from the plot center. Where feasible, at least 10 cm vertical mineral cores were taken, however in shallower layers a minimum depth of 5 cm each was collected. Duff samples were collected at the same plot corners as the mineral cores by excavating four soil volumes of approximately $25 \times 25 \text{ cm}^2$ area and at least the full depth of the organic layer. This volume was trimmed to discard the mineral and moss/litter layers off the bottom and top of the volumes, respectively. Right angles were then gently cut with sharp scissors and the 3 dimensions were measured in millimeters (collected samples were at least 400 cm^3 each and aimed to sample the entire in situ depth). Duff and mineral soils were kept frozen until portions were freeze dried for separate analysis. Moss/litter samples were collected by cutting squares, with attention to preservation of the natural in situ volume, until filling a 553 cm^3 steel container. Char layer samples were similarly collected in a 112 cm^3 container. At least one sample each of moss/litter and char were acquired from each plot quadrant, though more were taken at equal spacing along a transect to fill the containers if the layer was thin. On the upper surface of the char layer were small portions of dry, unburnt material, which were likely postfire additions of litter to the forest floor. This material was discarded from the char collection and was not included in C and N stock estimates.

2.2.2 Vegetation

Individual tree bole diameter (sampled at 130 cm height above the forest floor) and species were recorded within each plot perimeter for all trees of at least 5 cm diameter at measurement height. Trees less than 5 cm were uncommon and assumed to contribute negligibly to biomass and C emissions. A tree was recorded as living if standing upright and having any proportion of green needles (Sidoroff et al., 2007). If a fallen tree was charred only on its lower (in standing orientation) portions, it was deemed standing during fire ignition and its measurements were included if its base was within plot boundaries. In burnt plots, the percentage of brown and black needles in each tree canopy was visually approximated as 0%, 25%, 50%, 75%, or 100% with these individual values averaged to give an estimate of total plot canopy browning and blackening. Overstory biomass was calculated by entering bole diameters into allometric equations for Scots pine (*Pinus sylvestris*), Norway spruce (*Picea*

abies), silver birch (*Betula pendula*), and downy birch (*Betula pubescens*) (Marklund, 1987). Birch allometric equations were
225 applied to 7 of the 1716 trees sampled in burnt plots that were not one of these four species (which were unidentified and
deciduous). From the results of the equations, aboveground biomass was considered to be the categories stem wood, stem bark,
living branches, dead branches, stump and needles while belowground was coarse roots (≥ 5 cm) and fine roots (< 5 cm).
Stump, needle, coarse root and fine root categories were inapplicable for birch and its biomass was calculated only from the
remaining categories. To estimate overstory total aboveground C and N, C_R (ratio of sample C weight to total weight) for
230 all components was set to 0.5 and N_R (ratio of sample N weight to total weight) to 0.01 for needles and 0.004 for all other
parts (Boby et al., 2010). Bole diameters from the burnt plots were used to investigate the influence of overstory vegetation
C, N and composition on C and N loss. Bole diameters from adjacent control plots provided plot-pair estimates of increased
canopy blackening/browning and mortality as well as differences in species dominance, C and N to be tested for their influence
on C and N losses over the 50 plot network.

235 Understory samples were taken from control plots by cutting all non-moss, non-tree plant material at the surface of the soil
within four 40×40 cm² patches, each from a separate plot quadrant. To reduce sampling error due to low plot coverage, the
sample patches were chosen by performing four quadrant-wide surveys noting visual estimates of coverage and proportions
of plant functional groups (i.e. graminoids, forbs, shrubs, and pteridophytes). These were applied in selecting representative
patches for the portion of the quadrant that was vegetated, which was always all non-bare rock surface. The conglomerated
240 biomass density and composition for the four samples were applied to the visually estimated non-bare rock surface area of an
entire burnt plot to approximate its prefire understory coverage.

2.3 Sample processing

During sample processing each ecosystem compartment was sorted by weight into categories to produce rough metrics of their
composition and structure, here called compartment composition variables (CCVs). This sorting is described in the immediately
245 following and summarized in Table 1. All samples were dried at 40 °C for at least 3 days. Dry moss/litter samples were weighed
and visual estimates for percentage volume of needles, broad leaves, woody material, moss and lichen were multiplied by total
weight to form CCVs. This broadly categorized, visual estimation, along with the assumption of equal category density, is
meant to test for general effects of variation in proportions of surface fuel types on total soil fuel buildup and fire severity. CCVs
for understory were determined by sorting the sampled understory plant material and measuring dried weights of the functional
250 groups graminoid, forb, shrub, and pteridophyte. Mineral and duff samples were sieved to 2 mm and 4 mm respectively. The
weights of the coarse and fine fractions formed a pair of CCVs for each of the layers. CCVs for the overstory were collected
as the individual aboveground and belowground categories from the allometric equations.

Bulk density of each soil layer per plot was calculated as the total dry weight of its samples divided by their total volume as
measured during collection. All samples were pulverized, except the mineral soil where only the fine earth fraction (< 2 mm)
255 was analyzed (C_R and N_R were set to 0 for the coarse fraction), and packed in tin capsules. The capsules were combusted in
a Costech ECS 4010 elemental analyzer, equipped with a 2 m packed chromatographic column for gas separation, to produce
 C_R and N_R values. After every 10 samples, standardized acetanilide (provided by the company Elemental Microanalysis,

Okehampton, United Kingdom) was run to calibrate the machine within 1%. Duff and mineral layer elemental weight ratios were recalculated by the sum of C or N in each of their fine and coarse fractions and divided by total compartment weight. The
260 C:N ratio for each ecosystem compartment was calculated by dividing its total weight of C by total weight N.

2.4 Data analysis

2.4.1 C and N stock calculation

The measurable properties used in C and N stock calculations within soil compartments are the depth, bulk density and C_R or N_R . Total C and N stocks per soil compartment were calculated as a product of these properties using the equation

$$265 \text{ Stock}_Z = d \cdot \rho \cdot Z_R \quad (1)$$

where subscript Z is substituted with C or N for reference to C or N stocks, d is the soil layer depth in meters, ρ is the layer bulk density (kg m^{-3}) and Z_R is C_R or N_R . Understory compartment C and N stock calculations were performed with the equation

$$\text{Stock}_Z = \frac{m}{A} \cdot Z_R \cdot F \quad (2)$$

270 where m is the sampled mass in kilograms, A is sampled area (m^2), Z_R is C_R or N_R , and F is the estimated fractional vegetation coverage of the $20 \times 20 \text{ m}^2$ plot.

2.4.2 Plot-pair differences and confidence intervals

Unless otherwise noted, all measured changes between control and burnt plots were first calculated by subtracting control plot values of a variable from those of its burnt pair thereby forming a single distribution of 50 elements for statistical testing. These
275 distributions were approximated as normal and all confidence intervals were constructed at the 95% level, unless otherwise noted, using the formula

$$I = \bar{x} \pm z \cdot \frac{\sigma}{\sqrt{n}} \quad (3)$$

where \bar{x} is the sample mean, z is always 1.96 for the 95% interval, σ its standard deviation and n the sample size (always 50). Differences between control and burnt plots were significant when their interval did not include zero. When described as
280 losses, C and N stock differences between control and burnt plots were expressed as a positive number.

2.4.3 Statistical model construction

All regression analyses used the ordinary least squares approach to estimate a function for a single response variable based on linear combinations of the predictor variables and an intercept term. Simple regression was performed using the stats.linregress method from SciPy (Virtanen et al., 2020) providing significance (p), correlation (r), and slope (b). Multiple regression was
285 carried out with the OLS class in the Python 3 statsmodels package (Seabold and Perktold, 2010) with models evaluated in

order of increasing Akaike information criterion (AIC) (Akaike, 1974). Standardized regression coefficients (β) were produced by normalizing all variables (i.e. converting to z scores) before regression. When multicollinearity of explanatory variables in multiple regression was present, the condition number of the model provided by the corresponding OLS object was required to be less than 10 (Alin, 2010). Regression variables were also assessed in their squared, square-root, natural log, exponential and reciprocal transformations.

Multiple regression results were organized into path diagrams based on the assumption that regional climate (i.e. MAT and MAP at the km scale) and soil drainage (TEM) ~~had~~ could have a causal influence on the development of measured forest and fire attributes with negligible reverse effect. A clear potential causal direction of prefire forest properties influencing time-of-fire processes, which together influence C and N losses, is established by the temporal division of these measures across and within a discrete fire event. Therefore, variables in path diagrams are grouped into a causal order of climate/drainage to prefire to time-of-fire to postfire (i.e. C and N losses) properties. The main variables used in model construction are presented in Table 2.

Once causal models using the main variables were established, CCVs were added to the prefire variable category to test for their significance ($p < 0.05$) and increased explanatory power (decreased AIC) of C and N losses in multiple regression as well as their direct correlation to all model variables. This was done by using the original variables ~~and again but replacing~~ from each compartment as well as additional testing of the moss/litter, understory and overstory compartments ~~each~~ with their first two principal components produced by the PCA class in statsmodels. This same analysis was performed using ratios of C and N stocks present in the sampled control plot compartments. These ratios were aboveground (overstory, understory) to belowground (soil), moss/litter to duff, understory plus overstory to duff and organic to mineral.

305 3 Results

3.1 Survey of burnt plot vegetation

The 50 burnt plot overstories were largely dominated by pine with a percentage of spruce stems between 25-50% in five plots, between 50-75% in three plots and two plots with greater than 75%. Birch stems were less than 25% in 44 plots and between 25-50% in six plots, of which only one ~~of the six~~ was spruce dominant. All plots showed some visible charring of tree boles though only three plots had greater than 1% plot-wide canopy blackening. These plots were pine dominant, with two having less than 1% spruce stems while the other had ~~6 spruce~~ six spruce stems of the 27 stems within the sampled area. Prefire aboveground overstory C and N were estimated as 4.46 ± 0.74 kg C m^{-2} and 0.0385 ± 0.0062 kg N m^{-2} , respectively, with 5.31% of C (0.237 ± 0.032 kg C m^{-2}) and 12.3% of N (0.00474 ± 0.00064 kg N m^{-2}) coming from pine and spruce needles. The 50 burnt plots had a large percentage of tree mortality ($45.0 \pm 8.8\%$) compared to control plots ($4.21 \pm 1.63\%$). Total C and N loss, as well as char layer mass, were not correlated to canopy browning, blackening nor increased mortality in burnt plots relative to control plots.

Understory coverage was reduced to $10.2 \pm 5.2\%$ of its estimated prefire values. This left a surface layer of charred material exposed in all plots. The char layer was conglomerated and easily separable from lower layers and a thin clumps of litter

additions on top which were mostly needles. Upon breaking apart the layer, it was found to be completely blackened throughout.
320 Example photographs of four burnt plots are provided in Fig. 2.

Control plot soil C was not correlated to its overstory aboveground C ($p = 0.959$), understory C ($p = 0.285$) or their sum ($p = 0.962$). ~~Replacing C with~~ These relationships regarding N also showed no significant correlations. However, percentage spruce stems in control plots was correlated to organic layer C ($p = 0.033$, $r = 0.302$) and N ($p = 0.007$, $r = 0.378$), though with noticeable heteroscedasticity in the regression fit. Regardless, individual plot-pair differences of estimated over-
325 story C ($p = 0.824$), understory prefire plot coverage ($p = 0.384$), and the percentage of spruce stems ($p = 0.971$) were not related to C losses. The same is true for N. Therefore, there is likely no significant impact of plot-pair vegetation mismatch on the estimates made in this study over the 50 plot network.

3.2 C and N stock losses and restructuring

Averaging across all sites sampled, fire caused significant restructuring of C and N stocks particularly through changing soil
330 depth and bulk density which increased the mass per volume of both C and N. Significant differences in total C stocks of -0.815 ± 0.652 kg C m⁻² in burnt plots from their paired ~~controls~~ control plots were observed, but differences in N were insignificant (0.00208 ± 0.01840 kg N m⁻²). Fire clearly converted large amounts of C and N from lower soil layers to the high N_R surface layer of char. Organic layer C_R was unaffected by fire, however strong changes in N_R were measured resulting in an overall significant increase in the C:N ratio of this layer. Mean values and confidence intervals of the differences
335 between control to burnt plots highlighted in this section for all measured compartments are shown in Table 3 with soil specific properties shown in Table 4.

3.2.1 Total C

The largest total loss of C in burnt plot compartments due to fire was from the duff layer (Fig. 3a). 72.0% of the moss/litter C was removed from burnt plots, comprising an equivalent of 51.7% the total amount of C that was removed from the duff
340 layer. Understory C removal due to fire was near complete but had a relatively small contribution to overall C stocks and their changes. Char layer C averaged across the 50 burnt plots was equivalent to 54.3% of the average C lost due to fire from the combined understory and organic compartments. Burnt plot mineral layers had no significant overall change in C between control and burnt plots.

3.2.2 Total N

345 Fire adjusted the proportions of total N stored in burnt plot forest compartments despite having no overall significant effect on its total amount (Fig. 3b). Similar percentages of N and C were lost from the duff, moss/litter and understory compartments. Char layer N was equivalent to 100.8% of average combined N loss from these combined compartments. Change in N in the mineral layer from control to burnt plots was insignificant.

3.2.3 C:N ratio

350 The greater proportional reduction of C relative to N from control to burnt plots caused significant decreases in the C:N ratio in all compartments except the duff layer, which was unchanged (Fig. 3c). The low C:N ratio in the char layer (29.78 ± 1.70) made a strong contribution to the overall reduction in this value in burnt plot organic layers compared to ~~control~~those in control plots.

3.2.4 C_R and N_R

355 The duff layer C_R and N_R did not change significantly, though the moss/litter layer showed a significant increase in both values in burnt plots compared to their paired ~~controls~~control plots (Fig. 4c, 4d). These layers together with the char layer C_R (0.498 ± 0.019) and N_R (0.0173 ± 0.0011) resulted in the organic layer having no change in C_R but a significant increase in N_R in burnt plots. The mineral layer had significantly lower C_R and N_R in burnt plots both overall and in the fine fraction C_R (-0.0210 ± 0.0145 , -33.7%) and N_R (-0.000509 ± 0.000415 , -24.1%) compared to the ~~controls~~control plots.

360 3.2.5 Soil layer depths and bulk density

Fire had a strong effect on reducing soil layer depths with removal of nearly the entire moss/litter layer and about one third of the duff thickness (Fig. 4a). Together with the formation of the char layer and insignificant mineral layer depth changes, fire removed about a quarter of total soil depth and nearly 40% of the organic layer depth in burnt plots. Bulk density of both the duff and moss/litter layers increased significantly and, along with producing a dense char layer, fire had a strong densifying
365 effect on the organic layer (Fig. 4b). No significant change in bulk density occurred in the mineral layer due to fire.

3.2.6 Statistical contribution of measured changes to C and N losses

To quantify the relative statistical contributions of the variation of fire-induced changes in organic layer depth, bulk density and elemental weight ratios, they were used as predictor variables in multiple regression to explain organic layer C and N losses. The C loss regression produced a model of fit of $R^2 = 0.865$ and standardized regression coefficients for changes in
370 depth ($\beta = -0.670$), bulk density ($\beta = -0.633$) and C_R ($\beta = -0.583$). N loss produced a model fit of $R^2 = 0.777$ and coefficients for loss of depth ($\beta = -0.599$), bulk density ($\beta = -0.398$) and N_R ($\beta = -0.382$). This shows that changes of these variables due to fire all had a strong effect on C and N stock loss estimates. Measured change in organic layer depth is the strongest determinant of losses of N. However, for C, bulk density and elemental weight ratios are nearly as important as depth.

3.3 Forest level drivers of fire-induced C and N loss

375 The strongest correlator to total C and N losses among long-term ecosystem properties was total paired control plot C ($p < 0.001$, $r = 0.703$, $b = 0.744$) and N stocks ($p < 0.001$, $r = 0.585$, $b = 0.574$), respectively. An even stronger correlation was found between control plot organic layer C and N stocks (here abbreviated C_O and N_O) and estimated losses of C_O ($p < 0.001$, $r = 0.736$, $b = 0.762$) and N_O ($p < 0.001$, $r = 0.653$, $b = 0.665$) (Supplementary Fig. 3, 4). Due to this increased explanatory

power, and because the majority of fire affected C and N stocks were located in organic soil layers, the focus of analysis was placed on this C pool. Variables used in regression with percentage changes in C and N stocks tended to have less explanatory power than total C and N stock losses and also were sensitive to outliers having erratic changes in model fit with removal of data points. Therefore only total C and N stock losses were assessed.

The interaction-effect-ratio of C_O and N_O (i.e. its-C:N-ratio $C_O:N_O$) was added to improve model fit. C_O and its-C:N-ratio $C_O:N_O$ strongly explained C_O losses ($p < 0.001$, $R^2 = 0.588$) while N_O and its-C:N-ratio $C_O:N_O$ explained N_O losses with slightly less strength ($p < 0.001$, $R^2 = 0.519$). Multicollinearity between the-organic-layer-C:N-ratio $C_O:N_O$ and C_O ($p = 0.003$, $r = -0.411$, $b = -1.96 \text{ kg C m}^{-2}$) and N_O ($p < 0.001$, $r = -0.578$, $b = -92.2 \text{ kg N m}^{-2}$) did not produce a high condition number in these models (1.55 for C, 1.93 for N) suggesting they are robust to these covariations (Alin, 2010).

Total char layer C was not significantly related to loss of C_O ($p = 0.137$) however a significant correlation using simple regression was found between char layer N and losses of N_O ($p = 0.011$, $r = -0.359$, $b = -0.838$).

CCVs and ratios of C and N stocks amongst control plot compartments could not improve these models explaining C_O and N_O losses in multiple regression with C_O and N_O respectively nor could they significantly explain the buildup of organic layer fuel in control plots or production of char C or N. Relations either did not suitably meet the basic assumptions of regression, were deemed to be confounding or lacked supporting causal mechanism and were at a high risk of omitted-variable bias.

3.4 Climatic drivers of fire-induced C and N loss

MAP had a directly proportional relation to both C_O ($p < 0.001$, $R^2 = 0.465$, $b = 0.0194 \text{ kg C m}^{-2} \text{ mm}^{-1}$) and N_O ($p = 0.012$, $R^2 = 0.352$, $b = 0.000416 \text{ kg N m}^{-2} \text{ mm}^{-1}$) losses (Supplementary Fig. 5, 6). In multiple regression, MAT and MAT^2 explained C_O ($p = 0.008$, $R^2 = 0.186$) and N_O ($p = 0.002$, $R^2 = 0.233$) loss. Using these model fits, C_O and N_O losses were plotted as functions of MAT as negative parabolas both with a peak near 4°C (Supplementary Fig. 7, 8). In multiple regression of the 3 variables, the MAT and MAT^2 terms lost significance and MAP was the dominant explaining factor of C_O and N_O losses. MAT and MAP were not significantly related in simple regression ($p = 0.829$) however MAT and MAT^2 in multiple regression explained MAP ($p < 0.001$, $R^2 = 0.407$). Using these results, plotting MAP as a function of MAT again showed a negative parabola peaking near 4°C . (Supplementary Fig. 9). This suggests that the direct climate dependence of C_O and N_O losses were driven by MAP, with MAT relating indirectly through its association to MAP.

Control plot C_O and N_O were broken down into the three measured aspects of depth, bulk density, and elemental weight ratio and entered as a linear combination in multiple regression within path diagrams (Fig. 5a, b). MAT and MAP enhanced fire-induced C_O and N_O loss by promoting development of a denser, more voluminous fuel load with additional direct effects on losses by MAT. MAT mitigates C_O loss by reducing C_R , however variation in this value does not affect C_O loss as strongly as depth and bulk density. Conversely, MAT promotes N_O loss through a higher N_R fuel but exerts a stronger direct effect on reduction of N_O removal due to fire than in the C_O model. Accordingly, a separate multiple regression showed strong negative effects of MAT and MAP on organic-layer-C:N-ratio $C_O:N_O$ (Fig. 6a) along with a direct effect of MAT ($p < 0.001$, $r = -0.568$, $b = -2.50^\circ\text{C}^{-1}$). CCVs could not strongly link MAT or MAP with C_R , N_R , or the-C:N-ratio-in-control-plot-organic-layers $C_O:N_O$.

Total char layer mass ($p < 0.001$, $r = 0.453$, $b = 0.316 \text{ kg m}^{-2} \text{ }^\circ\text{C}^{-1}$) and char C_R ($p = 0.002$, $r = -0.435$, $b = -0.012 \text{ }^\circ\text{C}^{-1}$) were correlated with MAT but not the total mass of prefire fuel. This means that warmer regions produced larger amounts of lower C_R material irrespective of total fuel amount. In a multiple regression using C_O , the organic layer C:N ratio $C_O:N_O$, MAT, MAP and total char layer C production to explain C_O loss, direct effects of MAT lost significance and overall model fit was improved (Fig. 6a). This suggested that, while controlling for C_O and the organic layer C:N ratio $C_O:N_O$, C_O loss from this layer was reduced by MAT through the creation of char. Similarly, N_O loss is further explained by additions of char layer N to the climate model but a large direct effect of MAT remains (Fig. 6b). Adding the organic layer C:N ratio $C_O:N_O$ to this model increased the AIC and inflated the resulted in the individual p values of itself and MAT to $C_O:N_O$ and MAT reaching greater than 0.15, which is suggestive of a confounding influence of the organic layer C:N ratio $C_O:N_O$ on MAT and N_O loss. Again, CCVs and fuel distribution ratios could not improve either model.

3.5 Moisture and summer 2018 anomalies

TEM was not directly related to C_O ($p = 0.248$) or N_O ($p = 0.259$) stocks in control plots though it slightly improved model fit when joined with MAT and MAP in explaining these variables (Fig. 6a, b). TEM differences between paired burnt and control plots were observed to increase both with control and burnt plot TEM levels however not along gradients of MAT ($p = 0.198$), MAP ($p = 0.771$), C_O ($p = 0.302$) or N_O ($p = 0.423$). TEM differences between paired burnt and control plots were entered into simple regression with losses in C_O ($p = 0.088$, $r = -0.244$, $b = -0.0132 \text{ kg C m}^{-2}$) and N_O ($p = 0.035$, $r = -0.299$, $b = 0.00415 \text{ kg N m}^{-2}$). Although, correction for these slopes by using non-standardized coefficients of regression for TEM, MAT and MAP against C_O and N_O produced no significant change in losses of C_O or N_O .

SPEI was related to MAT ($p < 0.001$, $r = -0.869$), ΔMAT ($p < 0.001$, $r = -0.892$), ΔMAP ($p = 0.002$, $r = -0.433$) but not to MAP ($p = 0.725$), or losses of C_O ($p = 0.712$) and N_O ($p = 0.644$) due to fire. When included in multiple regression with MAP, MAT, char C or N, and C_O or N_O to explain fire-induced C_O or N_O stock losses, respectively, SPEI either did not improve model fit or promoted high uncertainty of variables (increased p values). SPEI was directly related to char C ($p = 0.013$, $r = -0.348$) and N ($p = 0.012$, $r = -0.351$) but had increased p value when explaining these variables in multiple regression with MAT, which suggests SPEI is a confounding variable. Additionally, ΔMAT ($p < 0.001$, $r = 0.956$) and ΔMAP ($p < 0.001$, $r = -0.525$) were strongly correlated to their respective long-term values with ΔMAT offering improved explanation of total char production ($p < 0.001$, $r = 0.477$, $b = 0.179 \text{ kg m}^{-2} \text{ }^\circ\text{C}^{-1}$) over MAT.

4 Discussion

4.1 Trends in ecosystem C and N stocks

Significant overall reduction in C stocks were found in burnt plots relative to their paired controls, with the largest removals from the duff layer. Averaged total C loss was relatively low at $0.815 \pm 0.652 \text{ kg C m}^{-2}$ (15.6%) compared to estimates from inland Alaskan black spruce stands (3.3 kg C m^{-2}) (Boby et al., 2010). However, they were comparable to averaged losses

from experimental Scots pine stands in Siberia ($0.992 \text{ kg C m}^{-2}$) (Ivanova et al., 2011) though half that of surface wildfire in
445 Siberian pine (*Pinus sibirica*) stands (1.69 kg C m^{-2}) (Kukavskaya et al., 2017). Interestingly, this study found average C stock
reduction to be nearly equivalent to that estimated in a Scots pine crown fire, selected for its high severity, within the largest,
most extreme wildfire occurring in Sweden during 2018 (0.80 kg C m^{-2}) (Kelly et al., 2021). In contrast, the largest wildfire
occurring in Sweden during 2014 released an estimated 4.50 kg C m^{-2} from Scots pine dominated stands, which lies just above
the 93rd percentile of C loss estimates within the 50 plot network of the current study (Granath et al., 2021). The extreme value
450 of the 2014 fire was produced from an averaging of regularly spaced ground samples across large portions of the burn scar
and found no relationship between extent of crown fire and C emission levels (Granath et al., 2021; Gustafsson et al., 2019).
These combined observations from Sweden exhibit an emission heterogeneity encompassing the entirety of typical pan-boreal
variation that cannot be explained by presence of crown fire, overstory species, fire size, and perhaps their combined effect
on observable fire intensity. This emphasizes the need to increase the context of studies of individual fire events by further
455 sampling over broader environmental gradients, which may thereby give greater predictive power to known, and yet to be
observed, sets of drivers of boreal wildfire C emissions (Veraverbeke et al., 2021).

Unlike C, however, N stocks were not significantly different overall in burnt plots compared to controls. This contrasts with
the [above-mentioned](#) Alaskan study which estimated percentage removal from soils of N (49.8%) to be similar to C (52.9%)
at an average loss of 0.09 kg N m^{-2} (Boby et al., 2010). Averaging over the 50 burnt plots, fire clearly reduced N in large
460 amounts within the residual duff and moss/litter layers but captured it within the high N_R char layer, preventing differences
when considering the overall soil profile. Proximity to the soil surface suggest a portion of the char layer was likely always
derived from fire interacting with the understory and moss/litter layer, however averaged char layer C and N stocks were
greater than losses from the two layers combined suggesting there were large contributions also from the duff. In burnt plots
with residual moss/litter an upwards mixing of mobilized duff C and N (e.g. in the form of [volatiles-vapor](#) or soot) may have
465 occurred due to heating of lower portions of the fuel bed. Because the char layer was conglomerated and completely blackened,
it is unlikely that material was incorporated postfire. However, material may have been added from downward movement of
overstory components during the time of fire or deposition of aerosols coming from outside the plot. By selecting plots well
within the final fire perimeter it was assumed that incoming and outgoing aerosols during the fire would be approximately
equal and that extended aerosol deposition from more remote sources would accumulate equally on control and burnt plots.
470 The above mentioned study in Alaskan black spruce forests, which are known for their great extent of canopy damage (Walker
et al., 2020a; Boby et al., 2010), showed C and N loss from the canopy to be about an order of magnitude lower than losses
from soil while also assuming that losses from the tree bole are negligible and that a large fraction of these overstory losses
were released to the atmosphere (Boby et al., 2010). An experimental high intensity crown fire in a Canadian Jack pine (*Pinus
banksiana*) forest with relatively low soil C stores (1.97 kg C m^{-2}) had minimal C losses from tree boles at 0.08 kg C m^{-2} ,
475 though needle C losses were substantial at 0.51 kg C m^{-2} (94.4% needle C) (Santín et al., 2015). In the current study, despite
a substantial portion of C and N stored in needles, low levels of overstory blackening and its lack of correlation with char
layer mass suggests that the large majority of C and N stock changes between control and burnt plots were captured within the
sampled soil and understory compartments.

The lack of change in total N stock due to fire is consistent with available evidence from existing study in Fennoscandian
480 forests where fire had only slight effects on total N over extended periods (Palviainen et al., 2017). N losses in non-boreal
forests have been related to fire temperature during time of fire with lower intensity fires transferring a greater proportion of
pools of organic N into soil ammonium and nitrate rather than removing N in gaseous forms (Neary et al., 1999). Labora-
tory studies have linked the amount of N transferred from organic to inorganic forms during heat exposure to both applied
temperature and fuel type (Gundale and DeLuca, 2006; Makoto et al., 2011). Therefore, the N cycle in boreal systems may
485 be highly dependent on active fire properties, fuel type and resulting fuel transformation and the greater N losses in Alaska
compared to Eurasia could be explained by its dissimilar fuel and the characteristically more intense crown fires across the
North American boreal zone (de Groot et al., 2013a; Wooster and Zhang, 2004). Fire intensity and temperature have also been
linked to both C and overall fuel transformation in experimental forest fire (Santín et al., 2016), so it is of interest to compare
remote measurements (i.e. satellite data) of these time-of-fire properties to on-site measured ecosystem changes. This can lead
490 to a more complete predictive understanding of wildfire in the entire Fennoscandian region and beyond.

4.2 Soil restructuring

In addition to removals, C and N were densified by fire in the organic layers due to their significant drops in depth and
increased bulk density in burnt plots. The lower mean C_R in burnt plot organic layers relative to control [plots](#) is assumed to
be due to the increased ratio of remaining incombustible material to remaining organic material. However, large variability
495 of duff C_R in burnt plots contributed to the statistical insignificance of this change and appeared to be related to extreme
volume reductions which reduced the duff layer to exceptionally ashy, rocky material in some plots. The large, significant
increase in N_R in the organic layer can be attributed to the formation of the high N_R char layer. The strong alteration of char
layer C:N relative to prefire fuel is comparable to results from studies that have incorporated similar pyrogenic layers that
are observed to be a mixture of organic or inorganic material types across broad ranges of combustion completeness (Bodí
500 et al., 2014). For example, the C:N ratio in a pyrogenic layer 1 year after a low intensity Siberian Larch (*Larix sibirica*) forest
surface fire in Russia was 31.4 (prefire 49.1), which is much lower than the 43.8 C:N ratio (prefire 39.4) observed the day after
an experimental, high intensity jack pine crown fire in Canada (Dymov et al., 2021; Santín et al., 2016), further suggesting
general differences in thermolability of soil C and N under regionally varied characteristic heating regimes.

Fire-induced morphological change of fuel has been shown to play a strong role in N retention where highly porous char
505 material adsorbs inorganic N preventing its leaching loss from the system until its reuptake into organic forms by plants or
microbes (Makoto et al., 2012). This sorptive power has been observed to fade over the interval between fire events suggesting
newly produced char is required for this retention effect (Zackrisson et al., 1996). The high N_R of the char layer may therefore
be due in part to adsorption of fire mineralized N or preserved, prefire mineral N and act as a steady source of bioavailable
nutrients to plant and microbial communities during succession. In addition to these inorganic forms, N may be stored in
510 this layer in partially combusted or depolymerized organic forms (Certini, 2005). This study employed coarse-scale sampling
of char based on soil horizon identification with separation in the field and a more rigorously defined assessment of char
production in all layers may provide further detail relating it to climate and soil processes.

4.3 Climate linked effects of fire

Both MAP and MAT had significant direct relations to total C and N removals from plots with the strongest mediator being estimated prefire C and N stocks in the organic layer. MAP had a stronger effect on the buildup of control plot fuel, namely through a positive correlation with total organic layer depth. MAT affected C and N losses through increasing bulk density, reducing C_R and increasing N_R thus reducing the C:N ratio in the organic layer in control plots, suggesting warmer conditions had a fuel conditioning effect through greater decomposition of organic soils (Callesen et al., 2007). When controlling for control plot organic layer C and N stocks and their ratios using multiple regression, MAT had a direct negative effect on C and N losses from this layer. This direct effect was largely mediated by the incorporation of measures of fire-induced fuel transformation into the models, i.e. production of char layer C or N. These models suggest that warmer regions tended to conserve larger pools of fire affected fuel as charred material rather than release it from the ecosystem either in gaseous, particulate or dissolved forms over the 1 year postfire period. This fuel restructuring in turn may have extended effects on C and N turnover through links to nutrient availability and the biotic community which in turn affect process rates such as primary production and soil decomposition (Makoto and Koike, 2021; Schmidt and Noack, 2000).

4.4 Considering representativeness and prediction of future wildfire impacts

A caveat of the pair plot matching methodology is that burnt plots may have had a greater tendency to ignite due to specific properties that heighten their fire susceptibility relative to [controlscontrol plots](#). As a result, underestimation of C and N losses may have occurred via burnt plots being biased to a greater prefire fuel load than their paired [controlscontrol plots](#) (systematic error) rather than these differences being approximately random (random error). This was evident in the fact that N losses were centered near 0 by their mean yet had a strong correlation with control plot total N despite the improbability that fire actually increased total N within nearly half of the burnt plot sample pool. Contrarily, control plots were biased towards higher TEM, which was observed to be related to greater fuel loading in measured control plots, though the magnitude of this bias did not increase along gradients of climate or fuel loading nor did attempts to correct for it significantly affect C and N loss estimates. Therefore, control plot matching appears to have been performed adequately within the scope of data collected, with any potential bias coming from unknown parameters. Further investigation of these unknowns is merited in order to improve control plot matching methodology and better constrain emission estimates in this region.

Further support of trends identified within the pair plot methodology was provided by separating analysis of the pyrogenic char layer, which was unique to burnt plots, from the soil profile as a whole. Counter to expectation, the large shifts in N_R and the C:N ratio in the char layer relative to lower layers in burnt plots show a clear preferential removal of C relative to N due to fire. This C:N ratio reduction was opposite to the C:N increases observed in a pyrogenic layer in more intense North American crown fire, which may be due to differences in fire temperature and heating duration (Santín et al., 2016). Therefore, until more predictive knowledge is gained over the proportional losses of C and N, accurate estimates of their stock changes due to fire may require measurements of densities of these individual elements throughout the residual soil profile in burnt plots.

545 Variation of many ecosystem properties could not be statistically linked to C and N losses, though that does not mean they are
without role in determining extent of emissions and ecosystem change. Cross regional boreal wildfire study has shown variables
such as the abundance of spruce to strongly affect C emissions across regions, but within a single region the abundance may be
too homogeneous to show an effect (Walker et al., 2020a). Furthermore, ratios of fine to heavy fuel loads have been manipulated
in experimental burns to produce varying fire severity (Alexander et al., 2018; Ludwig et al., 2018). Accordingly, CCVs and
550 fuel distribution amongst compartments in this study may have been simply too homogeneous to produce significant results
but may nevertheless still provide valuable statistical signals for understanding drivers of fire processes across regions and fire
severities. Of particular interest would be further investigation of the potential effect of dead wood structure on C and N stocks
and fire severity, which remains understudied in boreal wildfire research.

Previous studies have demonstrated strong effects of moisture on boreal wildfire C emissions. For example, a study including
555 several large North American fire complexes found C emissions to increase along gradients of topo-edaphic derived soil mois-
ture (due to its positive relation to total fuel) until reaching high moisture sites where the trend inverted and began to decrease
due to the inhibiting effects of this increased moisture on fuel availability (Walker et al., 2018). The position of this point of
inflection along the long-term moisture curve is likely dictated by short-term moisture levels, which are in turn controlled by
the extremity of drying during a fire season. Accordingly, fuel availability, and therefore C emissions, are strongly dependent
560 on drying processes specific to individual stand composition and structure and its local fire weather. By incorporating measures
of long (TEM, MAT, MAP) and short-term moisture balance (SPEI), climate based path models gained only slight improve-
ments in explaining fuel buildup and no increased explanatory power regarding C and N losses. Therefore, the study design
distinguished climate driven effects on fire severity with only minimal restrictions on site selection (non-sloping, non-wetland)
thereby providing results that are generally representative for Fennoscandian, non-wetland forests under similar drought con-
565 ditions of summer 2018. However, with its strong correlations to drought indices, anomalies of temperature and precipitation
as well as fuel charring and remaining direct effects on emissions, more remains to be understood about how MAT (and in
addition, intraannual distribution of MAP) relates to the fire regime across the conditions of fire seasons of differing extremity.

Fuel loading has been found to have a varied strength of control on boreal wildfire C loss globally. This study found total C
loss to relate to belowground C in simple regression with an R^2 of 0.494 and to aboveground C insignificantly. These results
570 are within the broad range found in Walker et al. (2020a) where total C losses across four North American boreal ecoregions
were directly related to prefire belowground C with R^2 values of 0.024 (insignificant), 0.07, 0.051, and 0.579 and to respective
prefire aboveground C with R^2 values of 0.229, 0.005 (insignificant), 0.101, and 0.336. Little is known about which factors
dictate the strength of these controls and what portions of unexplained variation can be attributed to either additional measurable
factors, methodological error or stochastic fire processes for a particular wildfire event. This study attempted to address these
575 issues by testing many measurable ecosystem properties across several ecosystem C storage compartments finding the organic
layer C stock along with its climate related prefire fuel conditioning and combustion susceptibility to be most predictive
of C loss (Fig. 5a). These trends were demonstrated using a consistent methodology that incorporates high replication and
broad spatial coverage, thereby offering better constraints on remaining unexplained variation across the region than might be
provided by sampling a single burn scar or comparing results from different study designs.

580 Boreal wildfire literature has tended to focus on highly flammable forest ecosystems with intense burning of several hundred hectares or more despite the vast majority of fires in the boreal region being less than 200 ha (Stocks et al., 2002; Valendik, 1996). This research bias may limit knowledge to a particular population by studying fire events only at their utmost extremes and effectively masking important signals of ecosystem heterogeneity on fire severity at differing intensities. The only restriction this study placed on wildfire intensity and size was that the fire activity could be remotely confirmed using Sentinel-2
585 infrared data. This has the potential to affect comparability of results to separate studies both in terms of total C and N losses as well as their drivers which may exhibit differing patterns and strengths over ranges of fire intensity. It may therefore be beneficial for wildfire studies to be examined and compared within categories of absolute severity and intensity. This methodology might be particularly useful in gaining understanding of the drivers of fire severity as it increases in traditionally fire protected ecosystems such as wetlands (Zoltai et al., 1998), which tend to store vastly larger amounts of C per area than their
590 dryer forested counterparts in the boreal region (Deluca and Boisvenue, 2012).

5 Conclusions

This study measured wildfire impacts across current climatic gradients of precipitation and temperature to show that climate controls total releases of C and N during fire events mainly due to its effect on increasing organic layer fuel load. The role of MAP is focused on the total quantity of this fuel load whereas MAT has a more qualitative effect by influencing bulk density, C_R
595 and N_R in the organic layer. When controlling for total organic layer fuel, increasing MAT, and to a lesser extent MAP, reduces C loss due to fire through preconditioning of the organic layer as measured by a lowered C:N ratio. Additionally, both C and N losses are mitigated by increased MAT through the sequestering of fire affected fuel into a surface layer of charred material. This conservation effect is stronger for N, which had no overall significant loss in stocks due to fire, and which also had stronger unexplained direct mitigating effects of MAT on its loss that were hypothesized to be related to time-of-fire properties such
600 as fire intensity and temperature. While remaining ecosystem variables regarding fuel composition and distribution could not be strongly linked to total C and N losses it is of interest to analyze their role in cross-regional comparisons and to investigate whether they influence other fire related properties such as ignition likelihood, fire propagability and intensity. Advancing knowledge of the intricate ties between instantaneous processes of fire events and their long-term effects on C and N cycling demands comprehensive research approaches that pay particular attention to climate sensitivity. This knowledge is imperative
605 for producing accurate predictions of boreal forest functioning under future climate scenarios.

Data availability. All data used to produce the results in this document are original unless otherwise stated in the text. It is found freely available through Eckdahl et al. (2021).

Author contributions. Author contributions using the CRediT contributor roles taxonomy are as follows: conceptualization (JAE, JAK, DBM), data curation (JAE), formal analysis (JAE), funding acquisition (JAK, DBM), investigation (JAE, JAK, DBM), methodology (JAE, JAK, DBM), project administration (JAE, JAK, DBM), resources (JAE, DBM), software (JAE), supervision (JAE, JAK, DBM), validation (JAE, JAK, DBM), visualization (JAE), writing – original draft preparation (JAE), and writing – review & editing (JAE, JAK, DBM).

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. JAE was supported by the strategic research area Biodiversity and Ecosystems in a Changing Climate, BECC, at Lund University. JAK was supported by the Carlsberg Foundation (grant CF20-0238). C and N analyses were performed in the Department of Geology, Lund University under supervision of Karl Ljung. Valuable information was provided by representatives from the Swedish Forest Agency and Swedish Civil Contingencies Agency. Thank you to Geerte Fälthammar-de Jong for dedicated assistance in field work planning and execution. Crucial field and lab assistance was provided by Rieke Madsen, Femke Pijcke, Lotte Wendt, Julia Iwan, and Henni Yläänne. Guidance from Karl Ljung, Åsa Wallin, Patrik Vestin and Michael Gundale in sample collection and processing was critical and highly appreciated. Thanks are given to three anonymous reviewers who provided helpful suggestions for improvement during manuscript development. Friends, family, and colleagues were vital sources of support and inspiration throughout this work.

References

- Akagi, S. K., Yokelson, R. J., Wiedinmyer, C., Alvarado, M. J., Reid, J. S., Karl, T., Crouse, J. D., and Wennberg, P. O.: Emission factors for open and domestic biomass burning for use in atmospheric models, *Atmospheric Chemistry and Physics*, 11, 4039–4072, <https://doi.org/10.5194/acp-11-4039-2011>, 2011.
- 625 Akaike, H.: A new look at the statistical model identification, *IEEE Transactions on Automatic Control*, 19, 716–723, <https://doi.org/10.1109/TAC.1974.1100705>, 1974.
- Alexander, H. D., Natali, S. M., Loranty, M. M., Ludwig, S. M., Spektor, V. V., Davydov, S., Zimov, N., Trujillo, I., and Mack, M. C.: Impacts of increased soil burn severity on larch forest regeneration on permafrost soils of far northeastern Siberia, *Forest Ecology and Management*, 417, 144–153, <https://doi.org/10.1016/j.foreco.2018.03.008>, 2018.
- 630 Alin, A.: Multicollinearity, *WIREs Computational Statistics*, 2, 370–374, <https://doi.org/10.1002/wics.84>, 2010.
- Bataineh, A. L., Oswald, B. P., Bataineh, M., Unger, D., Hung, I.-K., and Scognamillo, D.: Spatial autocorrelation and pseudoreplication in fire ecology, *Fire Ecology*, 2, 107–118, <https://doi.org/10.4996/fireecology.0202107>, 2006.
- Beguería, S., Latorre, B., Reig, F., and Vicente-Serrano, S. M.: Global SPEI database, <https://spei.csic.es/database.html>, 2019.
- Beven, K. J. and Kirkby, M. J.: A physically based, variable contributing area model of basin hydrology, *Hydrological Sciences Bulletin*, 24, 43–69, <https://doi.org/10.1080/02626667909491834>, 1979.
- 635 Bobby, L. A., Schuur, E. A. G., Mack, M. C., Verbyla, D., and Johnstone, J. F.: Quantifying fire severity, carbon, and nitrogen emissions in Alaska's boreal forest, *Ecological Applications*, 20, 1633–1647, <https://doi.org/10.1890/08-2295.1>, 2010.
- Bodí, M. B., Martín, D. A., Balfour, V. N., Santín, C., Doerr, S. H., Pereira, P., Cerdà, A., and Mataix-Solera, J.: Wildland fire ash: Production, composition and eco-hydro-geomorphic effects, *Earth-Science Reviews*, 130, 103–127, <https://doi.org/https://doi.org/10.1016/j.earscirev.2013.12.007>, 2014.
- 640 Bond-Lamberty, B., Peckham, S. D., Ahl, D. E., and Gower, S. T.: Fire as the dominant driver of central Canadian boreal forest carbon balance, *Nature*, 450, 89–92, <https://doi.org/10.1038/nature06272>, 2007.
- Buchanan, B. P., Fleming, M., Schneider, R. L., Richards, B. K., Archibald, J., Qiu, Z., and Walter, M. T.: Evaluating topographic wetness indices across central New York agricultural landscapes, *Hydrology and Earth System Sciences*, 18, 3279–3299, <https://doi.org/10.5194/hess-18-3279-2014>, 2014.
- 645 Callesen, I., Raulund-Rasmussen, K., Westman, C., and Tau-Strand, L.: Nitrogen pools and C:N ratios in well-drained Nordic forest soils related to climate and soil texture, *Boreal Environment Research*, 12, 681–692, 2007.
- Canada Soil Survey Committee: The Canadian system of soil classification, Research Branch, Canada Department of Agriculture, 1978.
- Certini, G.: Effects of fire on properties of forest soils: a review, *Oecologia*, 143, 1–10, <https://doi.org/10.1007/s00442-004-1788-8>, 2005.
- 650 Chen, D., Zhang, P., Seftigen, K., Ou, T., Giese, M., and Barthel, R.: Hydroclimate changes over Sweden in the twentieth and twenty-first centuries: a millennium perspective, *Geografiska Annaler: Series A, Physical Geography*, 103, 103–131, <https://doi.org/10.1080/04353676.2020.1841410>, 2021.
- De Frenne, P., Graae, B. J., Rodríguez-Sánchez, F., Kolb, A., Chabrierie, O., Decocq, G., De Kort, H., De Schrijver, A., Diekmann, M., Eriksson, O., Gruwez, R., Hermy, M., Lenoir, J., Plue, J., Coomes, D. A., and Verheyen, K.: Latitudinal gradients as natural laboratories to infer species' responses to temperature, *Journal of Ecology*, 101, 784–795, <https://doi.org/10.1111/1365-2745.12074>, 2013.
- 655 de Groot, W., Bothwell, P., Carlsson, D., and Logan, K.: Simulating the effects of future fire regimes on western Canadian boreal forests, *Journal of Vegetation Science*, 14, 355–364, <https://doi.org/10.1111/j.1654-1103.2003.tb02161.x>, 2003.

- de Groot, W. J., Cantin, A. S., Flannigan, M. D., Soja, A. J., Gowman, L. M., and Newbery, A.: A comparison of Canadian and Russian boreal forest fire regimes, *Forest Ecology and Management*, 294, 23–34, <https://doi.org/10.1016/j.foreco.2012.07.033>, 2013a.
- 660 de Groot, W. J., Flannigan, M. D., and Cantin, A. S.: Climate change impacts on future boreal fire regimes, *Forest Ecology and Management*, 294, 35–44, <https://doi.org/10.1016/j.foreco.2012.09.027>, 2013b.
- DeLuca, T. H. and Boisvenue, C.: Boreal forest soil carbon: distribution, function and modelling, *Forestry*, 85, 161–184, <https://doi.org/10.1093/forestry/cps003>, 2012.
- Dept. of Forest Resource Management, S. U. o. A.: SLU Forest Map, <https://www.slu.se/centrumbildningar-och-projekt/riksskogstaxeringen/statistik-om-skog/slu-skogskarta/om-slu-skogskarta/>, 2015.
- 665 Dieleman, C. M., Rogers, B. M., Potter, S., Veraverbeke, S., Johnstone, J. F., Laflamme, J., Solvik, K., Walker, X. J., Mack, M. C., and Turetsky, M. R.: Wildfire combustion and carbon stocks in the southern Canadian boreal forest: Implications for a warming world, *Global Change Biology*, 26, 6062–6079, <https://doi.org/https://doi.org/10.1111/gcb.15158>, 2020.
- Dymov, A., Startsev, V., Milanovsky, E., Valdes-Korovkin, I., Farkhodov, Y., Yudina, A., Donnerhack, O., and Guggenberger, G.: Soils and soil organic matter transformations during the two years after a low-intensity surface fire (Subpolar Ural, Russia), *Geoderma*, 404, 115–128, <https://doi.org/https://doi.org/10.1016/j.geoderma.2021.115278>, 2021.
- 670 Eckdahl, J., Kristensen, J., and Metcalfe, D.: Dataset for Boreal Forest Wildfire and Climate Linked Drivers of Carbon and Nitrogen Loss, <https://doi.org/10.5281/zenodo.5078669>, 2021.
- Esri Inc.: ArcGIS Pro, Esri Inc., <https://www.esri.com/en-us/arcgis/products/arcgis-pro/>, 2019.
- 675 French, N. H., Goovaerts, P., and Kasischke, E. S.: Uncertainty in estimating carbon emissions from boreal forest fires, *Journal of Geophysical Research: Atmospheres*, 109, <https://doi.org/10.1029/2003JD003635>, 2004.
- Gaboriau, D. M., Remy, C. C., Girardin, M. P., Asselin, H., Hély, C., Bergeron, Y., and Ali, A. A.: Temperature and fuel availability control fire size/severity in the boreal forest of central Northwest Territories, Canada, *Quaternary Science Reviews*, 250, 106–119, <https://doi.org/10.1016/j.quascirev.2020.106697>, 2020.
- 680 Giglio, L., Boschetti, L., Roy, D. P., Humber, M. L., and Justice, C. O.: The Collection 6 MODIS burned area mapping algorithm and product, *Remote Sensing of Environment*, 217, 72–85, <https://doi.org/10.1016/j.rse.2018.08.005>, 2018.
- Gillett, N., Weaver, A., Zwiers, F., and Flannigan, M.: Detecting the effect of climate change on Canadian forest fires, *Geophysical Research Letters*, 31, <https://doi.org/10.1029/2004GL020876>, 2004.
- Granath, G., Evans, C. D., Strengbom, J., Fölster, J., Grelle, A., Strömqvist, J., and Köhler, S. J.: The impact of wildfire on biogeochemical fluxes and water quality in boreal catchments, *Biogeosciences*, 18, 3243–3261, <https://doi.org/10.5194/bg-18-3243-2021>, 2021.
- 685 Gundale, M. J. and DeLuca, T. H.: Temperature and source material influence ecological attributes of ponderosa pine and Douglas-fir charcoal, *Forest ecology and management*, 231, 86–93, <https://doi.org/10.1016/j.foreco.2006.05.004>, 2006.
- Gustafsson, L., Berglund, M., Granström, A., Grelle, A., Isacson, G., Kjellander, P., Larsson, S., Lindh, M., Pettersson, L. B., Strengbom, J., Stridh, B., Sävström, T., Thor, G., Wikars, L.-O., and Mikusiński, G.: Rapid ecological response and intensified knowledge accumulation following a north European mega-fire, *Scandinavian Journal of Forest Research*, 34, 234–253, <https://doi.org/10.1080/02827581.2019.1603323>, 2019.
- 690 Ivanova, G., Conard, S., Kukavskaya, E., and McRae, D.: Fire impact on carbon storage in light conifer forests of the Lower Angara region, Siberia, *Environmental Research Letters*, 6, 045–053, <https://doi.org/10.1088/1748-9326/6/4/045203>, 2011.
- Johnstone, J., Celis, G., Chapin III, F., Hollingsworth, T., Jean, M., and Mack, M.: Factors shaping alternate successional trajectories in burned black spruce forests of Alaska, *Ecosphere*, 11, e03–129, <https://doi.org/10.1002/ecs2.3129>, 2020.
- 695

- Jonsson, B. G., Ekström, M., Esseen, P.-A., Grafström, A., Ståhl, G., and Westerlund, B.: Dead wood availability in managed Swedish forests – Policy outcomes and implications for biodiversity, *Forest Ecology and Management*, 376, 174–182, <https://doi.org/https://doi.org/10.1016/j.foreco.2016.06.017>, 2016.
- 700 Kaiser, J. W., Heil, A., Andreae, M. O., Benedetti, A., Chubarova, N., Jones, L., Morcrette, J.-J., Razinger, M., Schultz, M. G., Suttie, M., and van der Werf, G. R.: Biomass burning emissions estimated with a global fire assimilation system based on observed fire radiative power, *Biogeosciences*, 9, 527–554, <https://doi.org/10.5194/bg-9-527-2012>, 2012.
- Kasischke, E. S., Hyer, E. J., Novelli, P. C., Bruhwiler, L. P., French, N. H., Sukhinin, A. I., Hewson, J. H., and Stocks, B. J.: Influences of boreal fire emissions on Northern Hemisphere atmospheric carbon and carbon monoxide, *Global Biogeochemical Cycles*, 19, <https://doi.org/10.1029/2004GB002300>, 2005.
- 705 Keenan, R. J., Reams, G. A., Achard, F., de Freitas, J. V., Grainger, A., and Lindquist, E.: Dynamics of global forest area: Results from the FAO Global Forest Resources Assessment 2015, *Forest Ecology and Management*, 352, 9–20, <https://doi.org/10.1016/j.foreco.2015.06.014>, 2015.
- Kelly, J., Ibáñez, T. S., Santín, C., Doerr, S. H., Nilsson, M.-C., Holst, T., Lindroth, A., and Kljun, N.: Boreal forest soil carbon fluxes one year after a wildfire: Effects of burn severity and management, *Global Change Biology*, 27, 4181–4195, <https://doi.org/https://doi.org/10.1111/gcb.15721>, 2021.
- 710 Kelly, R., Chipman, M. L., Higuera, P. E., Stefanova, I., Brubaker, L. B., and Hu, F. S.: Recent burning of boreal forests exceeds fire regime limits of the past 10,000 years, *Proceedings of the National Academy of Sciences*, 110, 13 055–13 060, <https://doi.org/10.1073/pnas.1305069110>, 2013.
- Kohl, L., Philben, M., Edwards, K. A., Podrebarac, F. A., Warren, J., and Ziegler, S. E.: The origin of soil organic matter controls its composition and bioreactivity across a mesic boreal forest latitudinal gradient, *Global Change Biology*, 24, e458–e473, <https://doi.org/10.1111/gcb.13887>, 2018.
- 715 Kristensen, T., Ohlson, M., Bolstad, P., and Nagy, Z.: Spatial variability of organic layer thickness and carbon stocks in mature boreal forest stands—implications and suggestions for sampling designs, *Environmental monitoring and assessment*, 187, 1–19, <https://doi.org/10.1007/s10661-015-4741-x>, 2015.
- 720 Kukavskaya, E., Buryak, L., Kalenskaya, O., and Zarubin, D.: Transformation of the ground cover after surface fires and estimation of pyrogenic carbon emissions in the dark-coniferous forests of Central Siberia, *Contemporary problems of ecology*, 10, 62–70, <https://doi.org/10.1134/S1995425517010073>, 2017.
- Lantmäteriet: Markhöjdmodell Nedladdning, grid 50+, <https://www.lantmateriet.se/sv/Kartor-och-geografisk-information/geodataprodukter/produktlista/markhojdmodell-nedladdning-grid-50/#steg=1>, 2021.
- 725 Lemprière, T., Kurz, W., Hogg, E., Schmoll, C., Rampley, G., Yemshanov, D., McKenney, D., Gilseman, R., Beatch, A., Blain, D., Bhatti, J., and Krčmar, E.: Canadian boreal forests and climate change mitigation, *Environmental Reviews*, 21, 293–321, <https://doi.org/10.1139/er-2013-0039>, 2013.
- Li, F., Lawrence, D. M., and Bond-Lamberty, B.: Impact of fire on global land surface air temperature and energy budget for the 20th century due to changes within ecosystems, *Environmental Research Letters*, 12, 044 014, <https://doi.org/10.1088/1748-9326/aa6685>, 2017.
- 730 Ludwig, S. M., Alexander, H. D., Kielland, K., Mann, P. J., Natali, S. M., and Ruess, R. W.: Fire severity effects on soil carbon and nutrients and microbial processes in a Siberian larch forest, *Global Change Biology*, 24, 5841–5852, <http://dx.doi.org/10.1111/gcb.14455>, 2018.
- Makoto, K. and Koike, T.: Charcoal ecology: Its function as a hub for plant succession and soil nutrient cycling in boreal forests, *Ecological Research*, 36, 4–12, <https://doi.org/https://doi.org/10.1111/1440-1703.12179>, 2021.

- Makoto, K., Choi, D., Hashidoko, Y., and Koike, T.: The growth of *Larix gmelinii* seedlings as affected by charcoal produced at two different temperatures, *Biology and Fertility of Soils*, 47, 467–472, <https://doi.org/10.1007/s00374-010-0518-0>, 2011.
- Makoto, K., Shibata, H., Kim, Y., Satomura, T., Takagi, K., Nomura, M., Satoh, F., and Koike, T.: Contribution of charcoal to short-term nutrient dynamics after surface fire in the humus layer of a dwarf bamboo-dominated forest, *Biology and Fertility of Soils*, 48, 569–577, <https://doi.org/10.1007/s00374-011-0657-y>, 2012.
- Malhi, Y., Baldocchi, D., and Jarvis, P.: The carbon balance of tropical, temperate and boreal forests, *Plant, Cell & Environment*, 22, 715–740, <https://doi.org/10.1046/j.1365-3040.1999.00453.x>, 1999.
- Marklund, L.: Biomass functions for Norway spruce (*Picea abies* (L.) Karst.) in Sweden [biomass determination, dry weight], Rapport-Sveriges Lantbruksuniversitet, Institutionen foer Skogstaxering (Sweden), 1987.
- Mekonnen, Z. A., Riley, W. J., Randerson, J. T., Grant, R. F., and Rogers, B. M.: Expansion of high-latitude deciduous forests driven by interactions between climate warming and fire, *Nature plants*, 5, 952–958, <https://doi.org/10.1038/s41477-019-0495-8>, 2019.
- Murphy, P. N., Ogilvie, J., Connor, K., and Arp, P. A.: Mapping wetlands: a comparison of two different approaches for New Brunswick, Canada, *Wetlands*, 27, 846–854, [https://doi.org/10.1672/0277-5212\(2007\)27\[846:MWACOT\]2.0.CO;2](https://doi.org/10.1672/0277-5212(2007)27[846:MWACOT]2.0.CO;2), 2007.
- Naturvårdsverket: Markfuktighetsindex producerat som del av Nationella marktäckedata, NMD, <https://metadatakatalogen.naturvardsverket.se/metadatakatalogen/GetMetaDataById?id=cae71f45-b463-447f-804f-2847869b19b0>, 2018.
- Neary, D. G., Klopatek, C. C., DeBano, L. F., and Ffolliott, P. F.: Fire effects on belowground sustainability: a review and synthesis, *Forest ecology and management*, 122, 51–71, [https://doi.org/10.1016/S0378-1127\(99\)00032-8](https://doi.org/10.1016/S0378-1127(99)00032-8), 1999.
- Neff, J., Harden, J., and Gleixner, G.: Fire effects on soil organic matter content, composition, and nutrients in boreal interior Alaska, *Canadian Journal of Forest Research*, 35, 2178–2187, <https://doi.org/10.1139/x05-154>, 2005.
- Olsson, J., Södling, J., and Wetterhall, F.: Högupplösta nederbördsdata för hydrologisk modellering: en förstudie, <https://www.smhi.se/publikationer/hogupplosta-nederbordsdata-for-hydrologisk-modellering-en-forstudie-1.32746>, 2013.
- Olsson, M. T., Erlandsson, M., Lundin, L., Nilsson, T., Nilsson, Å., Stendahl, J., et al.: Organic carbon stocks in Swedish Podzol soils in relation to soil hydrology and other site characteristics, *Silva Fennica*, 43, 209–222, <https://doi.org/10.14214/sf.207>, 2009.
- Palviainen, M., Pumpanen, J., Berninger, F., Ritala, K., Duan, B., Heinonsalo, J., Sun, H., Köster, E., and Köster, K.: Nitrogen balance along a northern boreal forest fire chronosequence, *PLoS one*, 12, <https://doi.org/10.1371/journal.pone.0174720>, 2017.
- Pellegrini, A. F., Harden, J., Georgiou, K., Hemes, K. S., Malhotra, A., Nolan, C. J., and Jackson, R. B.: Fire effects on the persistence of soil organic matter and long-term carbon storage, *Nature Geoscience*, pp. 1–9, <https://doi.org/10.1038/s41561-021-00867-1>, 2021.
- Ponomarev, E. I., Kharuk, V. I., and Ranson, K. J.: Wildfires Dynamics in Siberian Larch Forests, *Forests*, 7, <https://doi.org/10.3390/f7060125>, 2016.
- Preston, C. M. and Schmidt, M. W.: Black (pyrogenic) carbon: a synthesis of current knowledge and uncertainties with special consideration of boreal regions, *Biogeosciences*, 3, 397–420, <https://doi.org/10.5194/bg-3-397-2006>, 2006.
- QGIS Development Team: QGIS Geographic Information System, QGIS Association, <https://www.qgis.org>, 2019.
- Rabin, S. S., Melton, J. R., Lasslop, G., Bachelet, D., Forrest, M., Hantson, S., Kaplan, J. O., Li, F., Mangeon, S., Ward, D. S., Yue, C., Arora, V. K., Hickler, T., Kloster, S., Knorr, W., Nieradzik, L., Spessa, A., Folberth, G. A., Sheehan, T., Voulgarakis, A., Kelley, D. I., Prentice, I. C., Sitch, S., Harrison, S., and Arneth, A.: The Fire Modeling Intercomparison Project (FireMIP), phase 1: experimental and analytical protocols with detailed model descriptions, *Geoscientific Model Development*, 10, 1175–1197, <https://doi.org/10.5194/gmd-10-1175-2017>, 2017.

- Rapalee, G., Trumbore, S. E., Davidson, E. A., Harden, J. W., and Veldhuis, H.: Soil carbon stocks and their rates of accumulation and loss in a boreal forest landscape, *Global Biogeochemical Cycles*, 12, 687–701, <https://doi.org/10.1029/98GB02336>, 1998.
- Rogers, B. M., Soja, A. J., Goulden, M. L., and Randerson, J. T.: Influence of tree species on continental differences in boreal fires and climate feedbacks, *Nature Geoscience*, 8, 228–234, <https://doi.org/10.1038/ngeo2352>, 2015.
- 775 Ruiz, J. A. M., Riaño, D., Arbelo, M., French, N. H., Ustin, S. L., and Whiting, M. L.: Burned area mapping time series in Canada (1984–1999) from NOAA-AVHRR LTDR: A comparison with other remote sensing products and fire perimeters, *Remote Sensing of Environment*, 117, 407–414, <https://doi.org/10.1016/j.rse.2011.10.017>, 2012.
- Santín, C., Doerr, S. H., Preston, C. M., and González-Rodríguez, G.: Pyrogenic organic matter production from wildfires: a missing sink in the global carbon cycle, *Global Change Biology*, 21, 1621–1633, <https://doi.org/https://doi.org/10.1111/gcb.12800>, 2015.
- 780 Santín, C., Doerr, S. H., Merino, A., Bryant, R., and Loader, N. J.: Forest floor chemical transformations in a boreal forest fire and their correlations with temperature and heating duration, *Geoderma*, 264, 71–80, <https://doi.org/https://doi.org/10.1016/j.geoderma.2015.09.021>, 2016.
- Schmidt, M. W. and Noack, A. G.: Black carbon in soils and sediments: analysis, distribution, implications, and current challenges, *Global biogeochemical cycles*, 14, 777–793, <https://doi.org/10.1029/1999GB001208>, 2000.
- 785 Schultz, M. G., Heil, A., Hoelzemann, J. J., Spessa, A., Thonicke, K., Goldammer, J. G., Held, A. C., Pereira, J. M. C., and van het Bolscher, M.: Global wildland fire emissions from 1960 to 2000, *Global Biogeochemical Cycles*, 22, <https://doi.org/10.1029/2007GB003031>, 2008.
- Schweiger, A. H., Irl, S. D. H., Steinbauer, M. J., Dengler, J., and Beierkuhnlein, C.: Optimizing sampling approaches along ecological gradients, *Methods in Ecology and Evolution*, 7, 463–471, <https://doi.org/10.1111/2041-210X.12495>, 2016.
- Seabold, S. and Perktold, J.: statsmodels: Econometric and statistical modeling with python, in: 9th Python in Science Conference, <http://dx.doi.org/10.25080/Majora-92bf1922-011>, 2010.
- 790 Sidoroff, K., Kuuluvainen, T., Tanskanen, H., and Vanha-Majamaa, I.: Tree mortality after low-intensity prescribed fires in managed *Pinus sylvestris* stands in southern Finland, *Scandinavian Journal of Forest Research*, 22, 2–12, <https://doi.org/10.1080/02827580500365935>, 2007.
- Soja, A. J., Cofer, W. R., Shugart, H. H., Sukhinin, A. I., Stackhouse, P. W., McRae, D. J., and Conard, S. G.: Estimating fire emissions and disparities in boreal Siberia (1998–2002), *Journal of Geophysical Research: Atmospheres*, 109, <https://doi.org/10.1029/2004JD004570>, 2004.
- 795 Stocks, B. J., Mason, J. A., Todd, J. B., Bosch, E. M., Wotton, B. M., Amiro, B. D., Flannigan, M. D., Hirsch, K. G., Logan, K. A., Martell, D. L., and Skinner, W. R.: Large forest fires in Canada, 1959–1997, *Journal of Geophysical Research: Atmospheres*, 107, FFR 5–1–FFR 5–12, <https://doi.org/10.1029/2001JD000484>, 2002.
- 800 Tagesson, T., Schurgers, G., Horion, S., Ciais, P., Tian, F., Brandt, M., Ahlström, A., Wigneron, J.-P., Ardö, J., Olin, S., Fan, L., Wu, Z., and Fensholt, R.: Recent divergence in the contributions of tropical and boreal forests to the terrestrial carbon sink, *Nature Ecology and Evolution*, 4, <https://doi.org/10.1038/s41559-019-1090-0>, 2020.
- Valendik, E.: Temporal and spatial distribution of forest fires in Siberia, in: *Fire in Ecosystems of Boreal Eurasia*, pp. 129–138, Springer, https://doi.org/10.1007/978-94-015-8737-2_9, 1996.
- 805 van der Werf, G. R., Randerson, J. T., Giglio, L., van Leeuwen, T. T., Chen, Y., Rogers, B. M., Mu, M., van Marle, M. J. E., Morton, D. C., Collatz, G. J., Yokelson, R. J., and Kasibhatla, P. S.: Global fire emissions estimates during 1997–2016, *Earth System Science Data*, 9, 697–720, <https://doi.org/10.5194/essd-9-697-2017>, 2017.

- van Leeuwen, T. T., van der Werf, G. R., Hoffmann, A. A., Detmers, R. G., Rucker, G., French, N. H. F., Archibald, S., Carvalho Jr., J. A., Cook, G. D., de Groot, W. J., Hély, C., Kasischke, E. S., Kloster, S., McCarty, J. L., Pettinari, M. L., Savadogo, P., Alvarado, E. C.,
810 Boschetti, L., Manuri, S., Meyer, C. P., Siegert, F., Trollope, L. A., and Trollope, W. S. W.: Biomass burning fuel consumption rates: a field measurement database, *Biogeosciences*, 11, 7305–7329, <https://doi.org/10.5194/bg-11-7305-2014>, 2014.
- Van Wagner, C.: Development and structure of the canadian forest fire weather index system, in: *Can. For. Serv., Forestry Tech. Rep.*, Citeseer, 1987.
- Vanhala, P., Karhu, K., Tuomi, M., Björklöf, K., Fritze, H., and Liski, J.: Temperature sensitivity of soil organic matter decomposition in
815 southern and northern areas of the boreal forest zone, *Soil biology and biochemistry*, 40, 1758–1764, <https://doi.org/10.1016/j.soilbio.2008.02.021>, 2008.
- Veraverbeke, S., Rogers, B. M., and Randerson, J. T.: Daily burned area and carbon emissions from boreal fires in Alaska, *Biogeosciences*, 12, 3579–3601, <https://doi.org/10.5194/bg-12-3579-2015>, 2015.
- Veraverbeke, S., Delcourt, C. J., Kukavskaya, E., Mack, M., Walker, X., Hessilt, T., Rogers, B., and Scholten, R. C.: Direct and longer-term
820 carbon emissions from arctic-boreal fires: A short review of recent advances, *Current Opinion in Environmental Science Health*, 23, 100277, <https://doi.org/https://doi.org/10.1016/j.coesh.2021.100277>, 2021.
- Virtanen, P., Gommers, R., Oliphant, T. E., Haberland, M., Reddy, T., Cournapeau, D., Burovski, E., Peterson, P., Weckesser, W., Bright, J., van der Walt, S. J., Brett, M., Wilson, J., Millman, K. J., Mayorov, N., Nelson, A. R. J., Jones, E., Kern, R., Larson, E., Carey, C. J., Polat, İ., Feng, Y., Moore, E. W., VanderPlas, J., Laxalde, D., Perktold, J., Cimrman, R., Henriksen, I., Quintero, E. A., Harris, C. R., Archibald,
825 A. M., Ribeiro, A. H., Pedregosa, F., van Mulbregt, P., and SciPy 1.0 Contributors: SciPy 1.0: Fundamental Algorithms for Scientific Computing in Python, *Nature Methods*, 17, 261–272, <https://doi.org/10.1038/s41592-019-0686-2>, 2020.
- Walker, X., Rogers, B., Veraverbeke, S., Johnstone, J., Baltzer, J., Barrett, K., Bourgeau-Chavez, L., Day, N., Groot, W., Dieleman, C., Goetz, S., Hoy, E., Jenkins, L., Kane, E., Parisien, M.-A., Potter, S., Schuur, E., Turetsky, M., Whitman, E., and Mack, M.: Fuel availability not fire weather controls boreal wildfire severity and carbon emissions, *Nature Climate Change*, pp. 1–7, <https://doi.org/10.1038/s41558-020-00920-8>, 2020a.
- Walker, X. J., Rogers, B. M., Baltzer, J. L., Cumming, S. G., Day, N. J., Goetz, S. J., Johnstone, J. F., Schuur, E. A. G., Turetsky, M. R., and Mack, M. C.: Cross-scale controls on carbon emissions from boreal forest megafires, *Global Change Biology*, 24, 4251–4265, <https://doi.org/10.1111/gcb.14287>, 2018.
- Walker, X. J., Baltzer, J. L., Bourgeau-Chavez, L., Day, N. J., Dieleman, C. M., Johnstone, J. F., Kane, E. S., Rogers, B. M., Turetsky, M. R.,
835 Veraverbeke, S., and Mack, M. C.: Patterns of Ecosystem Structure and Wildfire Carbon Combustion Across Six Ecoregions of the North American Boreal Forest, *Frontiers in Forests and Global Change*, 3, <https://doi.org/10.3389/ffgc.2020.00087>, 2020b.
- Wiggins, E. B., Andrews, A., Sweeney, C., Miller, J. B., Miller, C. E., Veraverbeke, S., Commane, R., Wofsy, S., Henderson, J. M., and Randerson, J. T.: Evidence for a larger contribution of smoldering combustion to boreal forest fire emissions from tower observations in Alaska, *Atmospheric Chemistry and Physics Discussions*, 2020, 1–26, <https://doi.org/10.5194/acp-2019-1067>, 2020.
- 840 Wilcke, R. A. I., Kjellström, E., Lin, C., Matei, D., Moberg, A., and Tyrlis, E.: The extremely warm summer of 2018 in Sweden – set in a historical context, *Earth System Dynamics*, 11, 1107–1121, <https://doi.org/10.5194/esd-11-1107-2020>, 2020.
- Wooster, M. J. and Zhang, Y. H.: Boreal forest fires burn less intensely in Russia than in North America, *Geophysical Research Letters*, 31, <https://doi.org/10.1029/2004GL020805>, 2004.
- Zackrisson, O., Nilsson, M.-C., and Wardle, D. A.: Key ecological function of charcoal from wildfire in the Boreal forest, *Oikos*, pp. 10–19,
845 <https://doi.org/10.2307/3545580>, 1996.

Zoltai, S., Morrissey, L., Livingston, G., and Groot, W. d.: Effects of fires on carbon cycling in North American boreal peatlands, *Environmental Reviews*, 6, 13–24, <https://doi.org/10.1139/a98-002>, 1998.




Figure 1. Using climate data averaged over the period 1961-2017 provided by the Swedish Meteorological and Hydrological Institute (SMHI) for plot selection, MAT had a range of 0.43-7.77 °C and MAP of 539-772 mm over an approximately 57-67° latitudinal change. The 50 plot pairs ~~are pictured~~ were added as points upon to a figure provided by SMHI of MAT and MAP gradients over the last normal period 1961-1990.



Figure 2. ~~Four~~ Field pictures from four separate burnt plots ~~were photographed taken~~ approximately 1 year after fire. Trees show charring of boles and enhanced exposure of roots due to fire-reduced organic layer height. Upon a surface char layer are new litter additions composed of mostly needles. Patchy regrowth of understory was present in many plots.




Figure 3. Mean C (a) and N (b) stocks and their ratio (c) between burnt and control plots amongst forest compartments. The organic category is considered the grouping of the duff, moss/litter and char layers while the total category is the grouping of the organic and mineral soil layers along with the understory. Error bars are the bootstrapped 95% confidence interval of the mean.




Figure 4. Mean soil compartment depths (a), bulk density (b), C_R (c), and N_R (d) in burnt and control plots. The organic category is considered the grouping of the duff, moss/litter and char layers while the soil category is the grouping of the organic and mineral soil layers. Error bars are the bootstrapped 95% confidence interval of the mean.




Figure 5. Diagram of proposed pathways for the effects of MAP and MAT on C (a) and N (b) loss. Non-climate variables regard the organic layer. Each variable node is labeled with the R^2 from simple or multiple regression using explanatory variables represented by all incoming arrows. Arrows are labeled by and sized in proportion to the magnitude of their standardized regression coefficients and are significant. Green arrows represent positive relationships while red represent negative relationships. Omitted for simplicity are direct correlations between bulk density to C_R ($p = 0.010$, $r = 0.356$) and depth to N_R ($p = 0.010$, $r = -0.363$).




Figure 6. Path diagrams for C (a) and N (b) loss including char and C:N variables. Prefire C, N, and C:N as well as losses of C and N are regarding the organic layer. Each variable node is labeled with the R^2 from simple or multiple regression using explanatory variables represented by all incoming arrows. Arrows are labeled by and sized in proportion to the magnitude of their standardized regression coefficients and are significant. Green arrows represent positive relationships while red represent negative relationships.

Table 1. CCV categories for ecosystem compartments given in dry weight (kg).

| Compartment | Categories |
|-------------------------|--|
| mineral | coarse (≥ 2 mm), fine (< 2 mm) |
| duff | coarse (≥ 4 mm), fine (< 4 mm) |
| moss/litter | needles, broad leaves, woody material, moss and lichen |
| understory | graminoids, forbs, shrubs, pteridophytes |
| overstory (aboveground) | stem wood, stem bark, living branches, dead branches, stump, needles |
| overstory (belowground) | coarse roots (≥ 5 cm), fine roots (< 5 cm) |

Table 2. ~~The main~~ Main variables used in path diagram construction ~~are~~ listed and grouped by category. Each category row is assumed to have a potential causal effect on lower rows.

| Category | Variables |
|------------------|--|
| climate/drainage | MAT, MAP, TEM |
| prefire soil | C, N, C:N, C_R , N_R , bulk density, depth |
| time-of-fire | SPEI, Δ MAT, Δ MAP, char C, char N |
| postfire | C loss, N loss |

Table 3. Mean values of distributions ($n = 50$) formed by subtracting control plot variables from their paired burnt plot. In parentheses are first the 95% confidence interval of the distribution and then the percentage change calculated from the overall means of control and burnt plots. Statistically significant fire-induced changes are in bold.

| | C ($kg\ C\ m^{-2}$) | N ($kg\ N\ m^{-2}$) | C:N |
|-------------|--------------------------------|----------------------------------|----------------------------|
| Total | -0.815 (0.652, -15.6) | 0.0021 (0.0184, 1.6) | -7.01 (2.58, -16.8) |
| Understory | -0.0717 (0.0126, -90.6) | -0.00127 (0.00022, -90.6) | – (–, –) |
| Soil | -0.744 (0.651, -14.5) | 0.0034 (0.0184, 2.6) | -6.88 (2.61, -16.6) |
| Organic | -0.747 (0.633, -17.8) | 0.0016 (0.0187, 1.6) | -9.46 (2.97, -19.8) |
| Char | 0.971 (0.182, –) | 0.0357 (0.0080, –) | – (–, –) |
| Moss/Litter | -0.584 (0.258, -72.0) | -0.00985 (0.00490, -68.2) | -9.09 (4.01, -15.6) |
| Duff | -1.13 (0.58, -33.5) | -0.0242 (0.0173, -29.6) | -1.24 (3.32, -2.7) |
| Mineral | 0.003 (0.248, 0.4) | 0.00179 (0.00834, 5.2) | -3.11 (2.46, -10.2) |

Table 4. Mean values of distributions ($n = 50$) formed by subtracting control plot variables from their paired burnt plot. In parentheses are first the 95% confidence interval of the distribution and then the percentage change calculated from the overall means of control and burnt plots. Statistically significant fire-induced changes are in bold.

| | Depth (cm) | Bulk Density ($g\ cm^{-3}$) | C_R | N_R |
|-------------|----------------------------|-------------------------------|--------------------------------|------------------------------------|
| Soil | -3.92 (1.47, -22.0) | 0.0665 (0.0305, 30.7) | -0.0326 (0.0395, -14.7) | 0.00029 (0.00104, 5.4) |
| Organic | -4.60 (1.19, -39.9) | 0.0358 (0.0110, 45.0) | -0.0040 (0.0193, -0.9) | 0.00243 (0.00080, 24.3) |
| Char | 1.91 (0.29, -) | - (-, -) | - (-, -) | - (-, -) |
| Moss/Litter | -3.98 (0.76, -85.6) | 0.0162 (0.0057, 42.8) | 0.0215 (0.0083, 4.7) | 0.00185 (0.00073, 21.9) |
| Duff | -2.53 (1.04, -36.8) | 0.0270 (0.0190, 24.9) | -0.0212 (0.0239, -4.6) | -0.000071 (0.000714, -0.7) |
| Mineral | 0.68 (1.32, 10.9) | 0.0057 (0.0352, 0.3) | -0.0199 (0.0127, -34.2) | -0.000505 (0.000369, -25.8) |