

Human and livestock faecal biomarkers at the prehistorical encampment site of Ullafelsen in the Fotsch Valley, Stubai Alps, Austria - potential and limitations

5 Marcel Lerch^{1,2}, Tobias Bromm², Clemens Geitner³, Jean Nicolas Haas⁴, Dieter Schäfer⁵, Bruno Glaser², Michael Zech^{1,2}

¹ Heisenberg Chair of Physical Geography with focus on paleoenvironmental research, Institute of Geography, Technische Universität Dresden, Helmholtzstraße 10, 01069 Dresden, Germany

10 ² Soil Biogeochemistry Group, Institute of Agricultural and Nutritional Sciences, Martin-Luther-University Halle-Wittenberg, Von-Seckendorff-Platz 3, 06120, Halle (Saale), Germany

³ Institute of Geography, University of Innsbruck, Innrain 52f, 6020 Innsbruck, Austria

⁴ Institute of Botany, University of Innsbruck, Sternwartestraße 15, 6020 Innsbruck, Austria

⁵ Institute of Geology, University of Innsbruck, Innrain 52f, 6020 Innsbruck, Austria

Correspondence to: Marcel Lerch (marcel.lerch@tu-dresden.de)

15 **Abstract.** The Ullafelsen at 1869 m a.s.l. in the Tyrolean Stubai Alps next to Innsbruck is an important (geo-)archaeological reference site for the Mesolithic period. Buried fireplaces on the Ullafelsen plateau were dated at 10.9 - 9.5 cal. kyrs BP and demonstrate together with thousands of flint stone artifacts the presence of hunter-gatherers during the Early Holocene. Grazing livestock has been a predominant anthropozoological impact in the Fotsch Valley presumably since the Bronze Age (4.2-2.8 kyrs BP). In order to study the human and/or livestock faeces input on the Ullafelsen, we carried out steroid analyses
20 on 2 modern ruminant faeces samples from cattle and sheep, 37 soil samples from seven archaeological soil profiles and 9 soil samples from five non-archaeological soil profiles from the Fotsch Valley used as reference sites. The dominance of 5 β -stigmastanol and deoxycholic acid in modern cattle and sheep faeces can be used as markers for the input of ruminant faeces in soils. The OA_h horizons, which have accumulated and developed since the Mesolithic, revealed high contents of steroids (sterols, stanols, stanones and bile acids), the E (LL) horizon coinciding with the Mesolithic living floor is characterized by
25 medium contents of steroids. By contrast, the subsoil horizons Bh, Bs and BvCv contain low contents of faecal biomarkers indicating that leaching of steroids into the podsolc subsoils is not an important factor. High content of 5 β -stigmastanol and deoxycholic acid in all soil samples give evidence for faeces input of ruminants. The steroid patterns and ratios indicate a negligible input of human faeces on the Ullafelsen. In conclusion, our results reflect a strong faeces input by livestock, rather than by humans as found for other Anthrosols such as Amazonian Dark Earths. Further studies need to focus on the question
30 of the exact timing of faeces deposition.

Keywords. Livestock faeces, Steroids, Bile acids, Mesolithic, Neolithic, Prehistorical animal husbandry, Anthrosol, Geoarchaeology

1 Introduction

35 Archaeological research in high mountain regions received increasing attention during the last decades. Based on the finding
of the copper age mummy called "Ötzi" at the Tisenjoch in the Ötztaler Alps in 1991, archaeological research projects were
also launched in the Central Alps (Schäfer, 2011b). Mesolithic hunter-gatherers lived in the Alpine regions since the
beginning of the Holocene (Fontana et al., 2016). For instance, Schäfer et al. (2016) and Cornelissen and Reitmaier (2016)
40 provided evidence for the presence of Mesolithic people at the upper subalpine or alpine zones in the central and south-
eastern Swiss Alps.

Concerning the Tyrolean Alps, the Mesolithic site of Ullafelsen (1869 m a.s.l.) in the Fotsch Valley was discovered by the
archaeologist Dieter Schäfer in 1994 and became an important archaeological reference site (Schäfer, 2011a; 2011b) (Figs. 1
and 2). At this site, thousands of archaeological artifacts and many buried fireplaces were found. This provides clear
evidence for the presence and the human environment interaction of our ancestors (Schäfer, 2011a; 2011b). Previous
45 archaeological research demonstrates that the Ullafelsen and its surrounding was used as a summer camp for hunting by
Mesolithic hunter-gatherers during the Preboreal and Boreal from around 10.9 to 9.5 kyrs BP (Schäfer, 2011a). The soils on
the Ullafelsen are strongly influenced by this Mesolithic impact and latest since the Bronze Age alpine pastoralism changed
the vegetation of the Fotsch Valley dramatically (Schäfer, 2011a; Zech et al., 2021).

50

- Figure 1 -

From a pedological point of view, a striking and frequently occurring light layer (LL) below the topsoils was described for
the Ullafelsen and was a focus of previous investigations (Geitner and Schäfer, 2010; Geitner et al., 2011; Geitner et al.,
2014). Similar light horizons are typical for soils developed in the subalpine zone of the Central Alps and are usually
55 interpreted as eluvial horizons (E) horizons of podzols (Zech and Wilke, 1977; Egli et al., 2008), or as eventual loess deposit
(Geitner et al., 2011; Schäfer, 2011a). At the Ullafelsen, most artifacts and Mesolithic fire places were found within and
directly on the top of the E (LL) horizon (Schäfer, 2011a). Therefore, the E (LL) horizon is regarded as Mesolithic living
floor, the humic-rich subsoil below the E (LL) horizon was considered as Late Glacial buried former topsoil (2Ahb horizon)
(Geitner et al., 2011; Schäfer, 2011a).

60 Most recently, Zech et al. (2021) demonstrated the great potential of *n*-alkane and black carbon biomarkers for contributing
to a better understanding of pedogenesis and landscape evolution. Black carbon results based on benzene polycarboxylic
acid (BPCA) analyses corroborated fire-induced human impact on the E (LL) horizon. The absence of leaf wax-derived *n*-
alkane biomarkers in the subsoils together with the absence of Late Glacial radiocarbon ages challenge the existence of a
Late Glacial buried topsoil (2Ahb horizon) and rather point to a humus-enriched podzolic Bh horizon (Zech et al., 2021).

65 In the context of alpine pastoralism (grazing, dairying) since the Neolithic period in different parts of the Alps,
(cattle)husbandry and agriculture became increasingly important for human society (Reitmaier et al., 2018; Gilck and

Poschlod, 2019). Grazing livestock such as cattle and sheep have been a predominant anthropozoological impact for the Ullafelsen and its surrounding presumably since the Bronze Age (4.2-2.8 kyrs. BP) as suggested by previous analyses (Zech et al., 2021).

70 Faecal biomarker analyses have become an attractive tool in palaeoenvironmental and archaeological research during the last decades (Baeten et al., 2012; Glaser and Birk, 2012; Prost et al., 2017). The respective molecules are considered as diagnostic markers for detecting ancient faecal inputs in soils (Bull et al., 1999b), whereby steroids (Δ^5 -sterols, 5α -stanols, 5β -stanols, epi- 5β -stanols, stanones and bile acids) are the relevant compound classes (Bull et al., 2005). These provide insights into ancient agricultural practices and the former presence of animals or humans (Prost et al., 2017).

75 Previous studies prove the specific steroids signal for various animals (Bull et al., 2005; Birk et al., 2011; Prost et al., 2017; Haurrault et al., 2019). Accordingly, faecal biomarker analyses allow to distinguish between faeces of herbivores/ruminants, pigs and humans. Based on their specific steroids signal, origin of faeces input can be detected in soils and sediments (von der Lühne et al., 2013; Haurrault et al., 2019). However, plants show also a specific steroids signal, which has to be considered during interpretations (Evershed et al., 1997; Hartmann, 1998). A finer differentiation between faeces of different
80 livestock can be achieved by the combination of several steroids (Δ^5 -sterols, 5α -stanols, 5β -stanols, epi- 5β -stanols, stanones and bile acids) (Prost et al., 2017).

Faecal steroids as organic compounds/lipid molecules can accumulate and persist in sediments and soils for more than thousands of years (Bull et al., 2001; Schroeter et al., 2020). These have a low water solubility and are thus usually neither leached into deeper soil horizons (Bull et al., 2002) nor detectable in soil leachates (Lloyd et al., 2012). It remains to be
85 investigated whether this also holds true for very low soil pH values < 4 like in our study area, where organic substances can become mobile via complexation with metal ions.

The 5β -stanols coprostanol and 5β -stigmastanol are products of anaerobic microbial reduction of Δ^5 -sterols. In mammals, this reduction is performed by gut bacteria and results in different ratios of 5β -stanols depending on food intake (Bull et al., 1999a). The Δ^5 -sterol cholesterol is the precursor molecule of coprostanol, while the Δ^5 -sterols β -sitosterol and stigmasterol
90 are the precursor molecules of 5β -stigmastanol and epi- 5β -stigmastanol. (Schroeter et al., 2020). Stanones are intermediate products of the transformation of Δ^5 -sterols to 5β -stanols, epi- 5β -stanols and 5α -stanols, occurs in the gut of animals and also in the environment (Prost et al., 2017). Epimerization of 5β -stanols, which occurs in soils due to microbial and diagenetic transformation, has to be considered when applying steroid ratios (Bull et al., 1999a; von der Lühne et al., 2018).

Up to now, analyses of livestock faeces show 5β -stigmastanol and deoxycholic acid (DCA) as the dominant steroid
95 compounds for ruminants (cattle and sheep), whereas coprostanol is a faecal marker for omnivores such as humans and pigs (Glaser and Birk, 2012; Prost et al., 2017; Haurrault et al., 2019). In contrast, plants contain high amounts of β -sitosterol and stigmasterol (Δ^5 -sterols) in roots and litter (Piironen et al., 2000; Verma and Gupta, 2013).

Bile acids are formed from cholesterol in the liver and transformed via the bile into the gut of mammals as primary bile acids (e.g. cholic acid (CA) or chenodeoxycholic acid (CDCA)). These primary bile acids are further mediated to secondary bile
100 acids (e.g. lithocholic acid (LCA) or deoxycholic acid (DCA)) in the intestine by microorganisms (Bull et al., 2002; Kuhajda

et al., 2006; Prost et al., 2017). Human faeces show high abundance of LCA, whereas ruminant faeces have a dominance of DCA (Prost et al., 2017; Shillito et al., 2020). Hence, steroid compounds can be useful markers for reconstructing settlement history of a site based on past faecal inputs.

Faecal biomarkers are currently used in various scientific disciplines all over the world. Glaser et al. (2001), Glaser and Birk
105 (2012) and Wiedner et al. (2015) investigated Anthropogenic Dark Earths, also known as terra preta de Índio, in Central Amazonia. By applying steroid markers, they provided evidence for settlement activities in this part of the tropical rainforest. High nutrient contents induced by the deposition of human and livestock faeces clearly demonstrated the anthropogenic origin of terra preta de Índio (Birk et al., 2011; Birk et al., 2012; Glaser and Birk, 2012). Another study used steroids for identification of temporary mass graves of concentration camp prisoners at the end of World War II (von der Lühe et al.,
110 2020). Findings revealed elevated faecal steroid contents and thus corroborate the former input of human decomposition products as well as faecal and tissue constituents of buried bodies (von der Lühe et al., 2020).

The aim of our here presented geoarchaeological study was to contribute to a better understanding of human and livestock impact at the prehistorical encampment site of Ullafelsen with the use of faecal biomarkers. More specifically, the following questions are addressed: (i) Do faecal biomarker patterns of modern ruminant faeces around the Ullafelsen reflect the steroid
115 patterns reported in literature and is a clear distinction from human faecal biomarker patterns possible? (ii) Do the steroid contents and patterns of the soil profiles at the Ullafelsen allow discrimination between human and livestock faeces input? (iii) Do the faecal steroid ratios of the soil profiles on the Ullafelsen allow the reconstruction of the faecal input history during the Holocene? We hypothesize that human faeces input is detectable in the E horizon representing the Mesolithic living floor (LL), whereas livestock faeces input dominates in the overlaying OAh horizon. (iv) Is there any evidence for
120 leaching of steroid biomarkers in soils with very low pH values such as in our study area?

2 Material and methods

2.1 Study area: The Ullafelsen as prehistorical encampment site in the Fotsch Valley, Stubai Alps, Austria

The prehistorical encampment site of Ullafelsen, also called "Riegelschrofen", is located in the Fotsch Valley at an altitude
of 1869 m above the sea level (a.s.l.). The 13 km long Fotsch Valley belongs to the Stubai Alps southwest of Innsbruck,
125 the capital of the Austrian state Tyrol. The Ullafelsen is a round hump at the eastern site of the Fotsch Valley and is located in the subalpine vegetation zone (Fig. 1). This rock ledge lies 40 m above the level of the adjacent creek, called "Fotscherbach" (Schäfer, 2011a). The geographic coordinates of the archaeological excavation area at the Ullafelsen are N 47.14702°, E 11.21475° (WGS84).

As a part of the transition zone between the wetter Northern and the drier Central Alps, our study area is characterized by a
130 temperate climate with a mean annual temperature of 10°C in the summertime (July) and -3°C in the wintertime (January). The mean annual precipitation is approx. 1500 mm (Schäfer, 2011a; Schlosser, 2011).

The vegetation is predominated by Swiss stone pine (*Pinus cembra*) and Juniper (*Juniperus communis* ssp. *alpina*). Furthermore, there are scattered European larch (*Larix decidua*), Norway spruce (*Picea abies*), Green alder (*Alnus viridis*) and Birch (*Betula pendula*). Alpine rose (*Rhododendron ferrugineum*), Lingonberry (*Vaccinium vitis-idaea*), European blueberry (*Vaccinium myrtillus*) and Ling heather (*Calluna vulgaris*) are occurring as alpine dwarf shrubs. The vegetation cover also consists of several herbs and grasses (Kemmer, 2011; Zech et al., 2021).

From a geological point of view, the Fotsch Valley represents a part of the "Öztal-Stubai-cristalline-complex". Typical rocks for this study area are the metamorphic rocks mica slate and paragneiss. In addition, there also exist a variety of unconsolidated quaternary sediments (Nittel, 2011). The basic material under the anthropogenically-influenced soils at the Ullafelsen consists, amongst others, of weathered till (Nittel, 2011). Despite human influence, these soils were mainly formed by podsolization during the Holocene (Zech et al., 2021). Typical soils in the alpine and subalpine zone of the Fotsch Valley are Cambisols and Leptosols. Under alpine dwarf shrub vegetation, Podzols have frequently developed, whereas in flatter valley floors and on some slope positions also Histosols can be found (Geitner et al., 2014).

2.2 Sampling of modern faeces, archaeological and reference soil profiles

As part of fieldwork in July and August 2017/2018, we collected 37 soil samples from seven profile walls of archaeological soil profiles on the Ullafelsen, 9 soil samples from five non-archaeological soil profiles from the Fotsch Valley as reference sites and 2 faeces samples from cattle and sheep (Table 1). Four profile walls are directly from the archaeological excavation area (1.1 C4w, 1.1 B5s, 1.1 B5w, 1.1 G5n). Three profile walls are from a close-by trench (1.9 NW, 1.9 NE, 1.9 SW) (Fig. 2). The latter is located two meters below the archaeological excavation area at an altitude of approximately 1867 m a.s.l. Samples of reference sites were collected from the soil profiles 4.4, 4.11, 4.12, 5.5 and 5.6w, which are primarily located along pathways and river section on hillsides above 2000 m a.s.l. in the Fotsch Valley (Table 1). Sampling was conducted by soil horizons, which were classified according to the WRB (IUSS Working Group WRB, 2015).

Samples of cattle faeces were collected on the Ullafelsen, while samples of sheep faeces were collected from the Fotsch Valley above 2000 m a.s.l. We chose these sampling sites for collecting faeces samples depending on a high density of grazing livestock mainly consisting of sheep. The food of grazing livestock from the Ullafelsen and surroundings consists of local vegetation, mostly grasses and alpine dwarf shrubs. After drying and grinding our faeces samples, we stored these separately in snap cap vials.

- Figure 2 -

160

For data evaluation, soil samples from soil profiles on the Ullafelsen were sorted by horizons (n=37): OAh1 (n=6), OAh2 (n=4), OAh3 (n=3), E (LL) (n=8), Bh (n=6), Bs (n=5), BvCv (n=5). The here investigated samples from reference soil profiles were from the OAh (n=5) and Bh (n=4) horizons (Table 1). Zech et al. (2021) carried out Total organic carbon (TOC) and TOC/N analyses for all these 46 soil samples. Figure 3 illustrates the soil profile 1.9 NW with the horizons

165 OAh1-OAh2-OAh3-E(LL)-Bh-Bs-BvCv. All other investigated soil profiles have a similar sequence of horizons. Results of grain size analyses for the soil profiles at the Ullafelsen show a dominance of sand (Geitner et al., 2011). In comparison to the other soil horizons, the E (LL) horizon is characterized by remarkably higher amounts of silt (Geitner et al., 2011).

- Figure 3 -

170

TOC values of soil samples from archaeological soil profiles on the Ullafelsen ranged from 0.3 to 28.8 % (Table 1). The maximum TOC content was observed in the OAh3 horizon, being in accordance with darker color and higher density of charcoal particles. Total nitrogen contents of the investigated soil profiles ranged from 0.0 to 1.2 %. The TOC/N ratios ranged from 12.4 to 37.2 with the highest ratios in the OAh3 horizon coinciding with charcoal particles in this soil horizon, related to former fireplaces, and high amounts of other organic material (Zech et al., 2021). High TOC/N ratios in the Bh horizons reflect podsolization processes (Glaser and Birk, 2012; Zech et al., 2014). TOC and TN values of soil samples from reference soil profiles from the Fotsch Valley ranged from 4.9 to 29.4 % and 0.3 to 2.2 %, respectively. The highest TOC and TN values were measured in the OAh horizons (Table 1).

175

Radiocarbon-dated Mesolithic charcoal yielded ¹⁴C ages ranging from 10.9 to 9.5 cal. kyrs BP (Schäfer, 2011a). More recently, Zech et al. (2021) yielded ¹⁴C ages for bulk *n*-alkanes ranging from 8.2 to 4.9 cal. kyrs BP. This discrepancy suggests that a *n*-alkane producing vegetation cover, consisting of herbs, grasses and alpine dwarf-shrubs, did not predominate immediately after the Mesolithic abandonment. Rather, it must be assumed that non-*n*-alkane producing conifers, such as *Pinus cembra* or *Picea abies*, predominated the vegetation cover after the Mesolithic life on the Ullafelsen (Zech et al., 2021).

180

In addition to the 37 soil samples at the archaeological site on the Ullafelsen and the 9 soil samples from reference soil profiles from the Fotsch Valley, we analyzed 2 mixed faeces samples from cattle and sheep, which belong to the typical livestock at the Ullafelsen and surroundings. TOC contents of cattle and sheep ranged from 42.6 to 43.5 % (Table 1).

185

- Table 1 -

190 2.3 Faecal biomarker analyses

Firstly, all 46 soil and 2 faeces samples were air-dried, sieved (≤ 2 mm) and finely ground. Using an elemental analyzer coupled to an isotope ratio mass spectrometer (EA-IRMS), total carbon (TC) and total nitrogen (TN) of soils samples were determined by Zech et al. (2021). Due to the non-carbonate parent rock material as well as the low pH values (< 4 in CaCl₂) of the soils and sediments on the Ullafelsen (Geitner et al., 2011), the measured TC values can be considered as TOC values.

195

From these data, the TOC/N ratio was calculated.

Steroid (sterol, stanol, stanone and bile acid) analyses were performed according to Birk et al. (2012), Wiedner et al. (2015) and von der Lüche et al. (2020). The weight of sample depended on the TOC content of the sample and ranged for the 46

analyzed soil samples from 1.0 to 5.0 g of finely ground material. For faeces, it is necessary to use a much smaller weight due to the higher TOC contents. In case of our two faeces (cattle and sheep), the weight of sample taken was ~60 mg. Prior to extraction, 100 µg of α -preganol and 100 µg of iso-deoxycholic acid were added to each sample as recovery standard (internal standard 1, IS1) for sterol/stanol and bile acid fractions, respectively. We used no internal standard for the stanone fractions. Regarding to data analysis and interpretation, we consider no stanones in our calculated ratios. The total lipid extract (TLE) was obtained by soxhlet extraction for 36 hours using a solvent mixture of dichloromethane/methanol (2:1, v/v, 180 ml).

After extraction, the solvent was removed using rotary evaporator and the TLEs were saponified using 3.5 ml of 0.7 M KOH in methanol at room temperature overnight for approx. 12 hours. Neutral lipids (sterols, stanols, stanones) as well as acidic lipids (bile acids, fatty acids) were separated by sequential liquid-liquid extraction. To obtain the neutral lipid fraction, the extracts were spiked with 10 ml H₂O and afterwards separated from the aqueous phase with 3 x 15 ml chloroform. For gaining the acidic lipid fraction, the aqueous solution was acidified with 6 M HCl (pH \leq 3-4) and the bile acids were extracted with 3 x 15 ml chloroform. Both fractions were separately collected in flasks and evaporated under nitrogen.

Before purification by solid phase extraction (SPE), acidic fraction were methylated by adding 1 ml of 1.25 M HCl in methanol and heating at 80°C for 2 hours. Bile acids and fatty acid methyl esters were extracted with 1 ml H₂O and 3 x 1 ml *n*-hexane. SPE was performed using glass columns (\emptyset 11 mm) containing 1.5 cm activated silica gel (Mesh: 70-230; pore size: 100 Å; type: Merck) in *n*-hexane. After pre-conditioning with 5 ml dichloromethane/*n*-hexane (2:1, v/v), the extracts were transferred with dichloromethane/*n*-hexane (2:1, v/v) on the silica-column. After elution of the fatty acid methyl esters to waste with 5 ml dichloromethane/*n*-hexane (2:1, v/v), bile acid methyl esters were eluted with 5 ml of dichloromethane/methanol (2:1, v/v) and collected in reactivals. The dried bile acid methyl esters were redissolved in 50 µl toluene and silylated with 98 µl BSTFA (N,O-Bis-(trimethylsilyl)-trifluoroacetamid; puriss; Sigma-Aldrich) and 2 µl TSIM (1-(Trimethylsilyl)imidazol; puriss; Sigma-Aldrich) at 80°C for 1 hour. After cooling, 50 µl 5 α -cholestane (10 ng µl⁻¹ in toluene) were added as internal standard 2 (IS2). All substances were transferred into GC-vials afterwards.

For the SPE of the neutral lipid fractions, activated silica gel (Mesh: 70-230; pore size: 100 Å; type: Merck) was deactivated with 5 % H₂O. The neutral lipid extract was transferred with *n*-hexane to the SPE glass columns (\emptyset 11 mm) containing 1.5 cm deactivated silica gel. Preconditioning was carried out with 5 ml *n*-hexane. By adding 5 ml *n*-hexane, aromatic and aliphatic fractions were eluted, but not used for further analyses. The sterol, stanol and stanone-containing fraction was eluted with 3 ml dichloromethane and 2 ml dichloromethane/acetone (2:1, v/v). The eluates were collected in reactivals before drying under nitrogen (N₂). Subsequently, the fraction of sterols, stanols and stanones were silylated by adding 100 µl Sylon HTP (a mixture of HMDS+TMCS+Pyridine (3:1:9, v:v:v); puriss; Supelco) and derivatized at 70°C for 1 hour. The eluates were dried with N₂ after cooling. 50 µl of 5 α -cholestane (10 ng µl⁻¹ in toluene) as internal standard 2 (IS2) and 100 µl of toluene were added to the dried eluates and transferred into GC-vials.

For external calibration, six concentrations from a stock solution (10 ng µl⁻¹ in hexane) containing all target sterols, stanols and stanones as well as the recovery standard (IS1) were prepared ranging from 50 to 2500 ng per vial. After derivatization

500 ng of 5 α -cholestane (dissolved in dry toluene, 50 μ l, 10 ng μ l⁻¹) were added as internal standard 2 (IS2). Peak areas of the analytes were divided by the peak area of IS2 and calibration curves for each substance were calculated using these ratios. Calibration curves for the bile acids were prepared accordingly. All calibration curves had coefficient of determination (R^2) > 0.98. Limit of detection was defined as signal-to-noise ratio of 3:1 and varied for the analytes between 0.6 ng g⁻¹ soil for all Δ^5 -sterols, stanols and stanones and 4 ng g⁻¹ soil for all bile acids. At last, recoveries of the first internal standard (IS1) were calculated for all samples. Recovery of IS1 of all soil samples ranged between 51-116% for sterols, stanols and stanones as well as 50-122% for bile acids. All results for the samples were corrected for the losses during extraction and purification with those individually calculated recoveries.

240 Quantification of all steroid substances took place using gas chromatography-mass spectrometry (GC-MS) with a 5971A quadrupole mass spectrometer connected to a HP5090 gas chromatograph (both made from Hewlett Packard) with a DB-5 MS 30 m fused silica column (25 mm ID and 0.25 μ m film thickness, Agilent Technologies).

All measurements of steroids were conducted with the following settings for GC-MS: injection volume: 1 μ l, carrier gas: helium (purity of 5), injector temperature: 290°C, injection in splitless mode, interface temperature: isotherm at 280°C. The column temperature program of the gas chromatography was held at 80°C for 1.5 min and then raised at 12°C min⁻¹ to 265°C. Further steps are the increase of the temperature at 0.8°C min⁻¹ to 288°C and at 10°C min⁻¹ to 300°C afterwards, whereas it was kept for 12 min.

2.4 Data analysis

For a detailed interpretation of the measured results in terms of degradation effects and distinguishing between omnivore-, herbivore- or plant-derived faecal biomarkers, the following ratios were calculated and plotted as box plot diagrams:

$$\text{Ratio 1} = \frac{\text{Coprostanol} + \text{Epicoprostanol}}{\text{Coprostanol} + \text{Epicoprostanol} + 5\alpha\text{-Cholestanol}} \quad (\text{Bull et al., 1999a})$$

$$\text{Ratio 2} = \frac{\text{Coprostanol} + \text{Epicoprostanol}}{5\beta\text{-Stigmastanol} + \text{Epi-}5\beta\text{-Stigmastanol}} \quad ((\text{Prost et al., 2017}), \text{ edited by Lerch, M.})$$

$$\text{Ratio 3} = \frac{\beta\text{-Sitosterol}}{\beta\text{-Sitosterol} + 5\beta\text{-Stigmastanol} + \text{Epi-}5\beta\text{-Stigmastanol}}$$

$$\text{Ratio 4} = \frac{5\beta\text{-Stigmastanol}}{\text{Epi-}5\beta\text{-Stigmastanol}}$$

260 According to Bull et al. (1999a) and Schroeter et al. (2020), we used ratio 1 to estimate microbial degradation of steroids (5 β -stanols, epi-5 β -stanols and 5 α -stanols) and to identify human faeces input (Wiedner et al., 2015). It is known that soil microorganisms contribute to degradation of steroids in soils (Bull et al., 2001). 5 α -Cholestanol is a degradation product of

the Δ^5 -sterol precursor cholesterol, transformed by soil microorganisms, and occurs naturally in the environment (Prost et al., 2017). This ratio considers the input and preservation of stanols. High values for ratio 1 (0.7-1) indicate an increased faeces deposition, while values < 0.7 indicate low faeces input in soils (Bull et al., 1999a; Schroeter et al., 2020).

For distinguishing between human and herbivore faeces, ratio 2 can be applied. A dominant 5β -stanol marker in omnivore faeces such as humans or pigs is coprostanol, whereas 5β -stigmastanol is a typical 5β -stanol marker for herbivore faeces. To account for ongoing epimerization of 5β -stanols in soils, both epimers are included in this ratio. Major input of human faeces results in values > 1 , whereas values < 1 indicate an input of herbivore faeces (Ossendorf et al., 2019).

Ratio 3 was calculated for differentiation between plant-derived steroids and livestock-derived steroids. According to Prost et al. (2017), β -sitosterol belongs to the typical Δ^5 -sterols, which are characteristic for plant biomass and thus normally occur at high abundance in soils. Values between 0 and 0.5 suggest low input of β -sitosterol and values between 0.5 and 1 point at a high occurrence of this plant Δ^5 -sterol. It has to be considered that faeces of ruminants can also contain high amounts of β -sitosterol due to their plant-dominated diet (Haurrault et al., 2019; Schroeter et al., 2020).

Ratio 4 was calculated for all soil samples and can be used as proxy for degradation of 5β -stigmastanol. Epi- 5β -stigmastanol is a transformation product (epimer) of 5β -stigmastanol and is often found in anaerobic environments and soils (von der Lühse et al., 2018). With increasing degradation, ratio 4 gets lower due to the higher proportion of epi- 5β -stigmastanol.

The following equation was applied for calculating a bile acid ratio:

280

$$\text{Ratio 5} = \frac{\text{Deoxycholic acid (DCA)}}{\text{Lithocholic acid (LCA)}} \quad (\text{Prost et al., 2017})$$

Based on the bile acids DCA and LCA, ratio 5 can be used for distinguishing ruminant from human faeces. Prost et al. (2017) published reference values, which are characteristic for ruminant species and humans respectively. Human faeces contain not only high amounts of coprostanol, but also of LCA. In contrast, ruminant faeces show a dominance of DCA (Shillito et al., 2020). A small ratio 5 (3-5) indicates a dominance of human faeces, whereas high values (5-21) show a dominance of ruminant faeces such as cattle or sheep (Prost et al., 2017).

285

3 Results and discussion

3.1 Biomarker patterns of modern livestock faeces

290 The total sterol, stanol and stanone contents of modern cattle and sheep faeces ranged from 2401.6 to 2671.6 $\mu\text{g g}^{-1}$ and 2109.8 to 2421.9 $\mu\text{g g}^{-1}$, respectively (Table S3). The total bile acid contents of modern cattle and sheep faeces ranged from 55.9 to 86.8 $\mu\text{g g}^{-1}$ and 78.6 to 216.9 $\mu\text{g g}^{-1}$, respectively (Table S4).

For checking the reproducibility of our results, we repeated the analyses of our faeces samples twice (Table S3; Table S4). Laboratory replicate analyses show that the method of faecal biomarker analyses works correctly and our results are reproducible. Figure 4 illustrates the steroid (sterols, stanols, stanones and bile acids) contents and their patterns in modern cattle and sheep faeces. The following results refer to measurement 1 (Table S3; Table S4).

The predominating steroid in cattle faeces is 5 β -stigmastanol (930.9 $\mu\text{g g}^{-1}$), whereas epi-5 β -stigmastanol (666.5 $\mu\text{g g}^{-1}$) shows a predominance in sheep faeces (Fig. 4). For comparison, Prost et al. (2017) did not find epi-5 β -stigmastanol to predominate in sheep faeces. It cannot be excluded, that the predominance of epi-5 β -stigmastanol in our sample is induced by strong epimerization, which is possibly influenced by long dry storage of the sample.

The coprostanol content in modern cattle faeces is 125.7 $\mu\text{g g}^{-1}$, whereas the coprostanol content of 173.4 $\mu\text{g g}^{-1}$ was detected in modern sheep faeces. These low contents of human-related faecal biomarkers in our modern ruminant faeces samples have to be considered for the identification of faeces origin in our soil samples. The plant-derived steroid β -sitosterol shows 347.9 $\mu\text{g g}^{-1}$ in modern cattle faeces and 255.9 $\mu\text{g g}^{-1}$ in modern sheep faeces (Fig. 4).

We calculated ratio 1, 2, 3 and 5 for our modern cattle and sheep faeces regarding to their steroid patterns to get reference ratios for comparing of our analysed soil samples. Bull et al. (1999a) and Prost et al. (2017) introduced ratio 1 for the identification of faeces input. Typical ratio 1 is ~ 0.8 for cattle and sheep, whereas ~ 1 is typical for human faeces. Our results revealed a ratio 1 of 0.8 for both modern faeces samples. These results are in agreement to Prost et al. (2017). Ratio 2 can be used as proxy for the identification of faeces origin. Ratios > 1 are characteristic for human or other omnivore faeces, whereas ratios < 1 represent herbivore/ruminant faeces. Our modern cattle and sheep faeces yielded ratio 2 of 0.1 and 0.2, respectively. These results are also in agreement with data of Prost et al. (2017). Ratio 3 reflects the predominance of plant-derived steroids (β -sitosterol) over livestock-derived steroids (5 β -stigmastanol and epi-5 β -stigmastanol). Ratios ≤ 0.5 indicate dominance of β -sitosterol, whereas ratios ≥ 0.5 show dominance of 5 β -stigmastanol and epi-5 β -stigmastanol. Our modern faeces samples yielded ratio 3 of ~ 0.2 for cattle and sheep, which demonstrates a dominant content of livestock-derived steroids over plant-derived steroids in our ruminant faeces.

The most dominant bile acid in our modern ruminant faeces is DCA (38.7 $\mu\text{g g}^{-1}$ and 55.0 $\mu\text{g g}^{-1}$ in cattle and sheep faeces, respectively), being in agreement with literature data of Kuhajda et al. (2006) and Prost et al. (2017). LCA as marker for the input of human faeces was found in modern ruminant faeces only at low amounts (4.2 $\mu\text{g g}^{-1}$ and 2.9 $\mu\text{g g}^{-1}$ in cattle and sheep faeces, respectively (Fig. 4)). Unexpectedly, our results of modern cattle and sheep faeces show low contents of CDCA (2.6 $\mu\text{g g}^{-1}$ and 2.4 $\mu\text{g g}^{-1}$ in cattle and sheep faeces, respectively). Prost et al. (2017) did not detect CDCA in cattle and sheep faeces, but in goats, horses, geese and human faeces instead.

According to Prost et al. (2017), ratio 5 ranged from 5-21 and 8-12 for cattle and sheep faeces, respectively. Based on our results, ratio 5 showed a ratio of 9.2 for cattle faeces and 18.8 for sheep faeces. For comparison, ratio 5 is typical for human faeces when in a range from 3 to 4.5. Accordingly, ratio 5 is highly promising for distinguishing between ruminant versus human faeces in soils. Unfortunately, steroid patterns of our analyzed ruminant faeces allow no differentiation between cattle and sheep.

All three measurements yielded similar content for sterols, stanols and stanones. By contrast, measurement 1 showed higher contents of the bile acid DCA but lower contents of oxolithocholic acid (OLCA) for both modern faeces samples (Table S4). The latter can be synthesized by oxidation of the hydroxyl groups of DCA (Sakai et al., 1980; Kuhajda et al., 2006; Marion et al., 2019).

Based on the steroid patterns of our modern cattle and sheep faeces from the Ullafelsen and surroundings, it is possible to evaluate the faecal biomarker results of our soils samples. Calculated steroid ratios (Ratio 1, 2, 3 and 5) help us to identify the sources of dominant faeces input in soils.

335 - Figure 4 -

3.2 Ancient faecal marker - contents and patterns in soils

The total sterol, stanol and stanone contents of the 37 soil samples from the Ullafelsen ranged from 1.2 to 198.1 $\mu\text{g g}^{-1}$ (Fig. 5). The OAh1 and OAh3 horizons yielded maxima (12.2 to 198.1 $\mu\text{g g}^{-1}$) coinciding with TOC maxima. The lowest content was measured in the Bh, Bs and BvCv horizons (1.2 to 22.3 $\mu\text{g g}^{-1}$), whereas the content of the E (LL) horizon was intermediate (2.3 to 58.6 $\mu\text{g g}^{-1}$; Fig. 5). In contrast, the sterol, stanol and stanone contents of the 9 soil samples from reference soil profiles from the Fotsch Valley ranged from 3.3 to 106.3 $\mu\text{g g}^{-1}$. Maximum contents were measured in the OAh horizons (24.3 to 106.3 $\mu\text{g g}^{-1}$), whereas Bh horizons yielded minimum contents (3.3 to 6.8 $\mu\text{g g}^{-1}$) of sterols, stanols and stanones (Fig. 5).

The total bile acid content of the 37 soil samples from the Ullafelsen ranged from 0 to 6.8 $\mu\text{g g}^{-1}$ and show their maximum in the topsoil horizons OAh1 and OAh2 (Fig. 5). In comparison to the maximum sterol, stanol and stanone contents in the OAh1 horizon, the maximum bile acid content of 6.8 $\mu\text{g g}^{-1}$ was detected in the OAh2 horizon. Similar to the sterol, stanol and stanone contents, the bile acid contents are much higher in the topsoil horizons OAh1, OAh2, and OAh3 (0.5 to 6.8 $\mu\text{g g}^{-1}$) and lower in the subsoil horizons E (LL), Bh, Bs and BvCv (0 to 0.9 $\mu\text{g g}^{-1}$) (Fig. 5). The total bile acids content of the 9 soil samples from reference soil profiles from the Fotsch Valley ranged from 0.1 to 15.5 $\mu\text{g g}^{-1}$. The OAh horizons yielded maximum contents (1.3 to 15.5 $\mu\text{g g}^{-1}$), whereas minimum contents of bile acids were measured in the Bh horizons (0.1 to 0.2 $\mu\text{g g}^{-1}$, Fig. 5).

Steroids in the subsoils (Bh, Bs and BvCv horizon) can be induced by bioturbation and/or roots of plants (Piironen et al., 2000). The most detected steroid in the subsoils is β -sitosterol, which is a plant-derived Δ^5 -sterol. Thomas and Hale (1983) as well as Verma and Gupta (2013) found that roots or root exudates contain also a significant content of β -sitosterol. We assume that the small contents of steroids in the subsoil horizons of the archaeological soils on the Ullafelsen and the reference soils in the Fotsch Valley are mainly caused by the influence of β -sitosterol due to the strong rooting of grasses and alpine dwarf shrubs in the soil matrix. There is no deformation of soil horizons, thus we exclude the influence of bioturbation.

The most abundant steroid in all soil samples from the Ullafelsen is β -sitosterol with a maximum content of $150.1 \mu\text{g g}^{-1}$ measured in an OAh3 horizon (Fig. 6; Table S1). β -Sitosterol is a typical plant-derived Δ^5 -sterol compound in plant biomass and reflects the predominating vegetation signal (Holtvoeth et al., 2010; Prost et al., 2017; von der Lühе et al., 2018). Due to the plant diet, β -sitosterol is eaten by ruminants and can be detected in their faeces (Haurrault et al., 2019). Cholesterol as the dominating Δ^5 -sterol in animal tissues has a maximum content of $6.2 \mu\text{g g}^{-1}$ detected in an OAh1 horizon (Fig.6; Table S1). Due to microbial degradation in soils, 5α -stigmastanol and 5α -cholestanol (both are 5α -stanols), are partly produced from their Δ^5 -sterol precursors β -sitosterol and cholesterol, respectively (Björkhem and Gustafsson, 1971). These 5α -stanols also occur in small amounts in fresh plant material and animal tissue (Noda et al., 1988; Bull et al., 2002; Prost et al., 2017). The maximum content of 5α -stigmastanol is $32.1 \mu\text{g g}^{-1}$ in an OAh3 horizon, whereas $1.4 \mu\text{g g}^{-1}$ is the maximum content of 5α -cholestanol detected in an OAh1 horizon (Fig. 6; Table S1).

The 5β -stanol compound coprostanol as marker for human faeces was detected at highest content of $0.2 \mu\text{g g}^{-1}$ in an OAh1 and OAh2 horizon. Epicoprostanol as transformation product of coprostanol due to microbial degradation (Bull et al., 1999a; Lauer et al., 2014) was not detectable in our soils (Fig. 6; Table S1). 5β -stigmastanol and epi- 5β -stigmastanol (epimerization product of 5β -stigmastanol) as marker for ruminant faeces have their maximum content of $3.3 \mu\text{g g}^{-1}$ and $3.2 \mu\text{g g}^{-1}$ in an OAh1 horizon, respectively (Fig.6; Table S1).

The most abundant bile acid in the analysed soil samples from the Ullafelsen was DCA with a maximum content of $4.2 \mu\text{g g}^{-1}$ in an OAh1 and OAh2 horizon (Fig. 6; Table S1). It predominates in ruminants such as cattle and sheep (Kuhajda et al., 2006; Prost et al., 2017). LCA as the dominating bile acid in human faeces (Shillito et al., 2020) showed a maximum content of $0.5 \mu\text{g g}^{-1}$ in an OAh2 horizon (Fig.6; Table S1). Wiedner et al. (2015) reported similar DCA and LCA contents for anthropogenic dark earths in northern Germany.

Furthermore, we found CDCA in the topsoil horizons OAh1, OAh2 and OAh3 as well as in the E (LL) horizon with a maximum content of $0.6 \mu\text{g g}^{-1}$, detected in an OAh2 horizon. Results of our modern cattle and sheep faeces showed also low contents of CDCA. Therefore, we explain the detected CDCA in our soils by the input of ruminant faeces.

OLCA was also detected in our soil samples from the Ullafelsen and has a maximum content of $1.7 \mu\text{g g}^{-1}$ in an OAh1 horizon (Fig. 6; Table S1). As discussed in 3.1, ruminant faeces also containing OLCA at low abundance. However, according to Marion et al. (2019), we cannot exclude that OLCA is formed by microorganisms in soils, such as *Clostridium scindens*, due to transformation of DCA to OLCA.

The most abundant sterol in the reference soils was β -sitosterol with the maximum content of $56.0 \mu\text{g g}^{-1}$ (Table S1). Deoxycholic acid is the dominant bile acid in the reference soils with the maximum content of $10.0 \mu\text{g g}^{-1}$ (Table S2). Both steroids were detected in the OAh horizon of reference soil profile 5.6w. Maximum contents of 5β -stigmastanol and deoxycholic acid as livestock-derived steroids were also found in this reference soil profile.

395

Overall, our results indicate that a strong input of faeces from cattle and sheep into the soils on the Ullafelsen occurred. By contrast, low LCA and coprostanol contribution point to a minor influence of human faeces. We suggest that the faeces input by wild animals such as chamois, marmot or fox can be neglected due to the high density of ruminant species at the Ullafelsen and surroundings.

400 According to Lloyd et al. (2012), our steroid results of the archaeological soils on the Ullafelsen and also of the reference soils from the Fotsch Valley show no leaching or transport in deeper soil horizons due to the significant lower steroid contents.

In comparison to the reference soils, the higher contents of total sterols, stanols and stanones in the OAh horizons of the archaeological soils on the Ullafelsen confirm an enrichment at this site. Unexpectedly, the total bile acid contents in the OAh horizons of the reference soils indicate a large range with partly higher contents compared to the archaeological soils and show thus no increased bile acid contents at the Ullafelsen. The highest contents of livestock-derived steroids (5 β -stigmastanol, epi-5 β -stigmastanol, deoxycholic acid) were detected in soil profile 5.6w of all reference soil profiles from the Fotsch Valley. We explain these high contents by a strong input of sheep faeces into the soil of soil profile 5.6w.

3.3 Identification of faeces origin on the Ullafelsen based on specific steroid ratios

410 Ratio 1 ((coprostanol+epicoprostanol)/(coprostanol+epicoprostanol+5 α -cholestanol)) ranged from 0 to 0.3 (Fig. 7). Human faeces normally exhibit ratio 1 > 0.7 (Bull et al., 1999a; Prost et al., 2017). Therefore, our results corroborate a very low input of human faecal markers such as coprostanol and suggest a negligible input of human faeces into soils on the Ullafelsen. Furthermore, we exclude a strong degradation of coprostanol in the topsoil horizons due to the non-detected epicoprostanol as transformation product of coprostanol in terms of epimerization over time. Contents of 5 α -cholestanol (\leq 415 1.4 $\mu\text{g g}^{-1}$) as transformation product of cholesterol (plant-derived) indicate also low degradation in the topsoil horizons.

Ratio 2 ((coprostanol+epicoprostanol)/(5 β -stigmastanol+epi-5 β -stigmastanol)) ranged from 0 to 0.4 (Fig. 7) and showed a maximum in the Bh horizon due to the higher content of epi-5 β -stigmastanol compared with the content of 5 β -stigmastanol (Table S1). Our results confirm a predominance of 5 β -stigmastanol and epi-5 β -stigmastanol, which indicate a strong input of ruminant faeces into soils at the Ullafelsen.

420 Ratio 3 (β -sitosterol/(β -sitosterol+5 β -stigmastanol+epi-5 β -stigmastanol)) ranged from 0.95 to 1 (Fig. 7), demonstrating a strong influence of plant-derived steroids (high β -sitosterol contents) in soils at the Ullafelsen caused by the high contribution of β -sitosterol (Fig. 6). High ratio 3 in the Bh, Bs and BvCv horizons can be explained by decreasing 5 β -stigmastanol and epi-5 β -stigmastanol contents. Roots of plants contribute to an input of β -sitosterol into subsoils (Piironen et al., 2000).

425 Ratio 4 (5 β -stigmastanol/epi-5 β -stigmastanol) ranged from 0 to 1.0 (Fig. 8), decreasing from the topsoils to the subsoils. As discussed in section 3.1, we observed a degradation effect of 5 β -stigmastanol in our soils. Our results corroborate a higher

degradation in the E (LL), Bh, Bs and BvCv horizons. We explain that with slightly higher epi-5 β -stigmastanol contribution in this subsoil horizons (Fig. 8; Table S1) due to epimerization (Bull et al., 1999a; Bull et al., 1999b; Bull et al., 2003; Prost et al., 2017). For a reliable identification of faeces origin, we recommend to consider the degradation effect and to include
430 epi-5 β -stanols (e.g. epi-5 β -stigmastanol and/or epicoprostanol) in steroid ratios (Bull et al., 2001).
Ratio 5 (deoxycholic acid/lithocholic acid) ranged from 4.1 to 23.5 (Fig. 7), decreasing from top to bottom. A ratio < 5 indicates a dominance of human faeces, whereas a ratio > 5 corroborates a dominance of ruminant faeces (Prost et al., 2017). No LCA was detected in the Bs and BvCv horizons. Apart from the outlier of 4.1, our results showed a clear dominance of ruminant faeces input into soils at the Ullafelsen due to the high content of DCA. The outlier of 4.1, detected in an E (LL)
435 horizon, can be allocated to a close-by-trench soil profile (1.9 NE E 1 (LL)), which is beyond the archaeological excavation area on the Ullafelsen. For this E (LL) horizon, we cannot exclude the input of human faeces regarding ratio 5, being based on the content of DCA (0.3 $\mu\text{g g}^{-1}$) and LCA (0.1 $\mu\text{g g}^{-1}$) (Table S2).

- Figure 7 - / - Figure 8 -

440 **4 Conclusions and outlook**

This study presents the first results of faecal biomarker analyses carried out on the prehistorical encampment site of Ullafelsen, Fotsch Valley, Austria. Steroid patterns of contemporary ruminants showed a dominance of 5 β -stigmastanol, whereas epi-5 β -stigmastanol as degradation product of 5 β -stigmastanol has to be considered for sheep. DCA was detected as the dominant bile acid for cattle and sheep. These data together with data of Prost et al. (2017) were used for the
445 interpretation of the faecal biomarker results from the geoarchaeological soil samples.

The highest sterol, stanol, stanone and bile acid contents in archaeological soils on the Ullafelsen were found in the OAh3 and OAh2 horizons, respectively. The dominant steroid in our soils is β -sitosterol as plant-derived Δ^5 -sterol compound. The faecal markers for ruminants 5 β -stigmastanol and DCA occurred in high contents in all topsoils corroborate actual grazing by cattle and sheep, which can be associated to the alpine pastoralism in the Fotsch Valley. Human faeces could be detected
450 only to a minor extent. Calculated ratios (Ratio 1-5) confirmed the negligible input of human faeces and the dominant input of ruminant faeces (cattle and sheep) at the Ullafelsen. Modern vegetation and ruminant faeces, associated with the plant-based diet of cattle and sheep, could have induced the high input of plant-derived steroids in our soils. Reference soils from the Fotsch Valley show lower contents of steroids except for bile acids and confirm the enrichment of sterols, stanols and stanones in the soils on the Ullafelsen. We observed no leaching of steroids into subsoil horizons.

455 Soil profiles on the Ullafelsen represent the modern signal of ruminant faeces and of plant-derived steroids. Given the low contents of human-derived faecal biomarkers combined with our evaluation of faecal biomarker ratios (Fig. 7), we see no evidence for a prehistoric input of human faeces at the Ullafelsen and surroundings. The archaeological site of Ullafelsen was used for fireplaces and social living in the Mesolithic period. However, it cannot be excluded that human faeces markers

will be detected in higher content close-by the archaeological excavation area. Future faecal biomarker analyses on further
460 soil profiles at the Ullafelsen aim at more specific insight in this hypothesis.

A robust age control for faecal biomarkers on the Ullafelsen is challenging because of lacking age chronology. This study
allows thus no reconstruction for the onset of alpine pastoralism. We assume the input of faecal steroids in the Holocene
since the beginning of the alpine pastoralism in the Neolithic and Bronze Age.

In order to chronologically identify the history of land use in the Fotsch Valley, we suggest to investigate faecal biomarkers
465 on mire archives in the Fotsch Valley. Previous studies of two subalpine mire archives in the near surroundings of the
prehistorical encampment site of Ullafelsen demonstrate the high potential for palaeoenvironmental reconstructions.

Supplement

Table S1: Overview over all faecal biomarker soil samples from the prehistorical encampment site of Ullafelsen and
reference sites in the Fotsch Valley, Stubai Alps, Austria. Sterols, stanols and stanones content (in $\mu\text{g g}^{-1}$ dry matter) as
470 well as ratios are presented.

Table S2: Overview over all faecal biomarker soil samples from the prehistorical encampment site of Ullafelsen and
reference sites in the Fotsch Valley, Stubai Alps, Austria. Bile acids content (in $\mu\text{g g}^{-1}$ dry matter) as well as ratios are
presented.

Table S3: Sterols, stanols and stanones content (in $\mu\text{g g}^{-1}$ dry matter) for 3 replication measurements of modern cattle and
475 sheep faeces from the prehistorical encampment site of Ullafelsen and reference sites in the Fotsch Valley, Stubai Alps,
Austria.

Table S4: Bile acids content (in $\mu\text{g g}^{-1}$ dry matter) for 3 replication measurements of modern cattle and sheep faeces from the
prehistorical encampment site of Ullafelsen and reference sites in the Fotsch Valley, Stubai Alps, Austria.

Data availability

480 TOC; TN and TOC/N data are available from Zech et al. (2021). Faecal biomarker data of the archaeological soil profiles on
the Ullafelsen and reference soil profiles from the Fotsch Valley are available in the Supplement.

Authors contributions

The project idea was developed by MZ in cooperation with BG, DS and CG. Fieldwork was done by ML, MZ, JNH, DS and
CG. ML performed most of the laboratory work with contributions made by TB, BG and MZ. The manuscript was prepared
485 by ML. All co-authors contributed to the discussion of the results and read and approved the manuscript.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

We greatly acknowledge Prof. Dr. Sixten Bussemer for support of fieldwork and helpful discussions on pedogenesis of the archaeological soil profiles from the Ullafelsen. Special thanks go to Oliver Kiewert and his team from the Bergheim hostel for excellent food and accommodation during the period of fieldwork. We also thank Corinna Heinrich, Heike Maennicke and Marianne Zech (form. Benesch) for supporting laboratory work and for the pleasant collaboration at any time as well as Eike Hüsing for making a drone video from the Ullafelsen and surroundings. Last but not least, we thank Prof. Dr. Eva Lehndorff and an anonymous reviewer for their constructive reviews that greatly helped to improve our manuscript. The German Research Foundation (GL 327/23-1 and ZE 844/12-1) kindly provided project funding.

References

- Baeten, J., Marinova, E., Laet, V. de, Degryse, P., Vos, D. de and Waelkens, M.: Faecal biomarker and archaeobotanical analyses of sediments from a public latrine shed new light on ruralisation in Sagalassos, Turkey, *Journal of Archaeological Science*, 39, 1143–1159, <https://doi.org/10.1016/j.jas.2011.12.019>, 2012.
- Birk, J. J., Dippold, M., Wiesenberg, G. L. B. and Glaser, B.: Combined quantification of faecal sterols, stanols, stanones and bile acids in soils and terrestrial sediments by gas chromatography–mass spectrometry, *Journal of Chromatography A*, 1242, 1–10, <https://doi.org/10.1016/j.chroma.2012.04.027>, 2012.
- Birk, J. J., Teixeira, W. G., Neves, E. G. and Glaser, B.: Faeces deposition on Amazonian Anthrosols as assessed from 5 β -stanols, *Journal of Archaeological Science*, 38, 1209–1220, <https://doi.org/10.1016/j.jas.2010.12.015>, 2011.
- Björkhem, I. and Gustafsson, J.: Microbial transformation of cholesterol into coprostanol, *European Journal of Biochemistry*, 21, 428–432, <https://doi.org/10.1111/j.1432-1033.1973.tb02968.x>, 1971.
- Bull, I. D., Betancourt, P. P. and Evershed, R. P.: An Organic Geochemical Investigation of the Practice of Manuring at a Minoan site on Pseira Island, Crete, *Geoarchaeology*, 16 (2), 223–242, [https://doi.org/10.1002/1520-6548\(200102\)16:2<223::AID-GEA1002>3.0.CO;2-7](https://doi.org/10.1002/1520-6548(200102)16:2<223::AID-GEA1002>3.0.CO;2-7), 2001.
- Bull, I. D., Elhmmali, M. M., Perret, V., Matthews, W., Roberts, D. J. and Evershed, R. P.: Biomarker evidence of faecal deposition in archaeological sediments at Çatalhöyük, in: *Changing materialities at Çatalhöyük: reports from the 1995-1999 seasons*, edited by: Hodder, I., Cambridge, Great Britain, 415–420, 2005.

- 515 Bull, I. D., Elhmmali, M. M., Roberts, D. J. and Evershed, R. P.: The application of steroidal biomarkers to track the abandonment of a Roman wastewater course at the Agora (Athens, Greece), *Archaeometry*, 45, 149–161, <https://doi.org/10.1111/1475-4754.00101>, 2003.
- Bull, I. D., Lockheart, M. J., Elhmmali, M. M., Roberts, D. J. and Evershed, R. P.: The origin of faeces by means of biomarker detection, *Environmental International*, 27 (8), 647–654, [https://doi.org/10.1016/S0160-4120\(01\)00124-6](https://doi.org/10.1016/S0160-4120(01)00124-6), 2002.
- 520 Bull, I. D., Simpson, I. A., Bergen van, P. F. and Evershed, R. P.: Muck 'n' molecules: organic geochemical methods for detecting ancient manuring, *Antiquity*, 73, 86–96, <https://doi.org/10.1017/S0003598X0008786X>, 1999a.
- Bull, I. D., Simpson, I. A., Dockrill, S. J. and Evershed, R. P.: Organic geochemical evidence for the origin of ancient anthropogenic soil deposits at Tofts Ness, Sanday, Orkney, *Organic Geochemistry*, 30, 535–556, [https://doi.org/10.1016/S0146-6380\(99\)00020-0](https://doi.org/10.1016/S0146-6380(99)00020-0), 1999b.
- 525 Cornelissen, M. and Reitmaier, T.: Filling the gap: Recent Mesolithic discoveries in the central and south-eastern Swiss Alps, *Quaternary International*, 423 (22), 9–22, <https://doi.org/10.1016/j.quaint.2015.10.121>, 2016.
- Egli, M., Mirabella, A. and Sartori, G.: The role of climate and vegetation in weathering and clay mineral formation in late Quaternary soils of the Swiss and Italian Alps, *Geomorphology*, 102, 307–324, <https://doi.org/10.1016/j.geomorph.2008.04.001>, 2008.
- 530 Evershed, R. P., Bethell, P. H., Reynolds, P. J. and Walsh, N. J.: 5 β -Stigmastanol and Related 5 β -Stanols as Biomarkers of Manuring: Analysis of Modern Experimental Material and Assessment of the Archaeological Potential, *Journal of Archaeological Science*, 24, 485–495, <https://doi.org/10.1006/jasc.1996.0132>, 1997.
- Fontana, F., Visentin, D. and Wierer, U.: MesoLife. A Mesolithic perspective on Alpine and neighbouring territories, *Quaternary International*, 423 (22), 1–4, <https://doi.org/10.1016/j.quaint.2016.06.016>, 2016.
- 535 Geitner, C., Bussemer, S., Ehrmann, O., Ikinge, A., Schäfer, D., Traidl, R. and Tscherko, D.: Bodenkundlich-stratigraphische Befunde am Ullafelsen im hinteren Fotschertal sowie ihre landschaftsgeschichtliche Interpretation, in: *Das Mesolithikum-Projekt Ullafelsen (Teil 1). Mensch und Umwelt im Holozän Tirols 1*, edited by: Schäfer, D., Philipp von Zabern, Darmstadt, Germany, 109–151, 2011.
- 540 Geitner, C. and Schäfer, D.: Interdisziplinäre Zusammenarbeit an der Schnittstelle von Archäologie und Bodenkunde im Gebirge – Grundsätzliche Überlegungen und Beispiele des Mesolithfundplatzes Ullafelsen (Tirol), in: *Archäologie in den Alpen: Alltag und Kult*, edited by: Mandl, F. and Stadler, H., ANISA, Verein für alpine Forschung, Haus im Ennstal, 25–42, 2010.

- 545 Geitner, C., Schäfer, D., Bertola, S., Bussemer, S., Heinrich, K. and Waroszewski, J.: Landscape archaeological results and discussion of Mesolithic research in the Fotsch valley (Tyrol), in: From the foreland to the Central Alps – Field trips to selected sites of Quaternary research in the Tyrolean and Bavarian Alps (DEUQUA EXCURSIONS), edited by: Kerschner, H., Krainer, K. and Spötl, C., Geozon Science Media, Berlin, Germany, 106–115, 2014.
- Gilck, F. and Poschlod, P.: The origin of alpine farming: A review of archaeological, linguistic and archaeobotanical studies in the Alps, *The Holocene*, 29 (9), 1503–1511, <https://doi.org/10.1177/0959683619854511>, 2019.
- 550 Glaser, B. and Birk, J. J.: State of the scientific knowledge on properties and genesis of Anthropogenic Dark Earths in Central Amazonia (terra preta de Indio), *Geochimica et Cosmochimica Acta*, 82, 39–51, <https://doi.org/10.1016/j.gca.2010.11.029>, 2012.
- Glaser, B., Haumaier, L., Guggenberger, G. and Zech, W.: The "terra preta" phenomenon: a model for sustainable agriculture in the humid tropics, *Naturwissenschaften*, 88 (1), 37–41, <https://doi.org/10.1007/s001140000193>, 2001.
- Hartmann, M.-A.: Plant sterols and the membrane environment, *Trends in Plant Science*, 3 (5), 170–175, [https://doi.org/10.1016/S1360-1385\(98\)01233-3](https://doi.org/10.1016/S1360-1385(98)01233-3), 1998.
- 555 Haurrault, L., Milek, K., Jarde, E., Jeanneau, L., Derrien, M. and Anderson, D. G.: Faecal biomarkers can distinguish specific mammalian species in modern and past environments, *PLoS ONE*, 14 (2), 1–26, <https://doi.org/10.1371/journal.pone.0211119>, 2019.
- Holtvoeth, J., Vogel, H., Wagner, B. and Wolff, G. A.: Lipid biomarkers in Holocene and glacial sediments from ancient Lake Ohrid (Macedonia, Albania), *Biogeosciences*, 7, 3473–3489, <https://doi.org/10.5194/bg-7-3473-2010>, 2010.
- 560 IUSS Working Group WRB.: World Reference Base for Soil Resources 2014, update 2015. International soil classification system for naming soils and creating legends for soil maps., 106, World Soil Resources Reports FAO, Rome, 2015.
- Kemmer, I.: Die rezente Vegetation im inneren Fotschertal / Nördliche Stubaier Alpen, in: *Das Mesolithikum-Projekt Ullafelsen (Teil 1). Mensch und Umwelt im Holozän Tirols 1*, edited by: Schäfer, D., Philipp von Zabern, Darmstadt, Germany, 155–193, 2011.
- 565 Kuhajda, K., Kandrac, J., Kevresan, S., Mikov, M. and Fawcett, J. P.: Structure and origin of bile acids: An overview, *European Journal of Drug Metabolism and Pharmacokinetics*, 31 (3), 135–143, <https://doi.org/10.1007/BF03190710>, 2006.
- 570 Lauer, F., Prost, K., Gerlach, R., Pätzold, S., Wolf, M., Urmersbach, S., Lehndorff, E., Eckmeier, E. and Amelung, W.: Organic Fertilization and Sufficient Nutrient Status in Prehistoric Agriculture? – Indications from Multi-Proxy Analyses of Archaeological Topsoil Relicts, *PLoS ONE*, 9 (9), <https://doi.org/10.1371/journal.pone.0106244>, 2014.


- Lloyd, C. E. M., Michaelides, K., Chadwick, D. R., Dungait, J. A. J. and Evershed, R. P.: Tracing the flow-driven vertical transport of livestock-derived organic matter through soil using biomarkers, *Organic Geochemistry*, 43, 56–66, <https://doi.org/10.1016/j.orggeochem.2011.11.001>, 2012.
- 575 Marion, S., Studer, N., Desharnais, L., Menin, L., Escrip, S., Meibom, A., Hapfelmeier, S. and Bernier-Latmani, R.: In vitro and in vivo characterization of *Clostridium scindens* bile acid transformations, *Gut Microbes*, 10 (4), 481–503, <https://doi.org/10.1080/19490976.2018.1549420>, 2019.
- Nittel, P.: Geologie, Hydrogeologie und Geomorphologie des Fotschertales - Kartierungsergebnisse Projekt "Sellrain", in: Das Mesolithikum-Projekt Ullafelsen (Teil 1). Mensch und Umwelt im Holozän Tirols 1, edited by: Schäfer, D., Philipp von Zabern, Darmstadt, Germany, 61–92, 2011.
- 580 Noda, M., Tanaka, M., Seto, Y., Aiba, T. and Oku, C.: Occurrence of cholesterol as a major sterol component in leaf surface lipids, *Lipids*, 23, 439–444, <https://doi.org/10.1007/BF02535517>, 1988.
- Ossendorf, G., Groos, A. R., Bromm, T., Tekelemariam, M. G., Glaser, B., Lesur, J., Schmidt, J., Akcar, N., Bekele, T., Beldados, A., Demissew, S., Kahsay, T. H., Nash, B. P., Nauss, T., Negash, A., Nemomissa, S., Veit, H., Vogelsang, R., Woldu, Z., Zech, W., Opgenoorth, L. and Mieke, G.: Middle Stone Age foragers resided in high elevations of the
585 glaciated Bale Mountains, Ethiopia, *Science*, 365, 583–587, <https://doi.org/10.1126/science.aaw8942>, 2019.
- Piironen, V., Lindsay, D. G., Miettinen, T. A., Toiva, J. and Lampi, A.: Plant sterols: biosynthesis, biological function and their importance to human nutrition, *Journal of the Scientific of Food and Agriculture*, 80, 939–966, [https://doi.org/10.1002/\(SICI\)1097-0010\(20000515\)80:7<939::AID-JSFA644>3.0.CO;2-C](https://doi.org/10.1002/(SICI)1097-0010(20000515)80:7<939::AID-JSFA644>3.0.CO;2-C), 2000.
- Prost, K., Birk, J. J., Lehndorff, E., Gerlach, R. and Amelung, W.: Steroid Biomarkers Revisited - Improved Source
590 Identification of Faecal Remains in Archaeological Soil Material, *PLoS ONE*, 12 (1), 1–30, <https://doi.org/10.1371/journal.pone.0164882>, 2017.
- Reitmaier, T., Doppler, T., Pike, A. W. G., Deschler-Erb, S., Hajdas, I., Walser, C. and Gerling, C.: Alpine cattle management during the Bronze Age at Ramosch-Mottata, Switzerland, *Quaternary International*, 484, 19–31, <https://doi.org/10.1016/j.quaint.2017.02.007>, 2018.
- 595 Sakai, K., Makino, T., Kawai, Y. and Mutai, M.: Intestinal Microflora and Bile Acids - Effect of Bile Acids on the Distribution of Microflora and Bile Acid in the Digestive Tract of the Rat, *Microbiology and Immunology*, 24 (3), 187–196, <https://doi.org/10.1111/j.1348-0421.1980.tb00578.x>, 1980.
- Schäfer, D.: Das Mesolithikum-Projekt Ullafelsen - Landschaftlicher Rahmen und archäologische Befunde. Arbeitsstand 2009/2010, in: Das Mesolithikum-Projekt Ullafelsen (Teil 1). Mensch und Umwelt im Holozän Tirols 1, edited by:
600 Schäfer, D., Philipp von Zabern, Darmstadt, Germany, 245–351, 2011a.

- Schäfer, D.: Das Mesolithikum-Projekt Ullafelsen (Teil 1). Mensch und Umwelt im Holozän Tirols 1, Philipp von Zabern, Darmstadt, Germany, 2011b.
- Schäfer, D., Bertola, S., Pawlik, A., Geitner, C., Waroszewski, J. and Bussemer, S.: The landscape-archaeological Ullafelsen Project (Tyrol, Austria), *Preistoria Alpina*, 48, 29–38, 2016.
- 605 Schlosser, E.: Das Fotschertal - regionale Klimatologie und gebirgsmeteorologische Aspekte, in: Das Mesolithikum-Projekt Ullafelsen (Teil 1). Mensch und Umwelt im Holozän Tirols 1, edited by: Schäfer, D., Philipp von Zabern, Darmstadt, Germany, 11–16, 2011.
- Schroeter, N., Lauterbach, S., Stebich, M., Kalanke, J., Mingram, J., Yildiz, C., Schouten, S. and Gleixner, G.: Biomolecular Evidence of Early Human Occupation of a High-Altitude Site in Western Central Asia During the Holocene, *Frontiers in Earth Science*, 8, 1–20, <https://doi.org/10.3389/feart.2020.00020>, 2020.
- 610 Shillito, L.-M., Whelton, H. L., Blong, J. C., Jenkins, D. L., Connolly, T. J. and Bull, I. D.: Pre-Clovis occupation of the Americas identified by human fecal biomarkers in coprolites from Paisley Caves, Oregon, *Science Advances*, 6, 1–8, <https://doi.org/10.1126/sciadv.aba6404>, 2020.
- Thomas, L. K. and Hale, M. G.: Effects of kinetin in the rooting medium on root exudation of free fatty acids and sterols from roots of *Arachis hypogaea* L. 'argentine' under axenic conditions, *Soil Biology and Biochemistry*, 15, 125–126, [https://doi.org/10.1016/0038-0717\(83\)90131-1](https://doi.org/10.1016/0038-0717(83)90131-1), 1983.
- 615 Verma, S. and Gupta, R.: Comparative estimation of β -sitosterol in roots, leaves and flowers of *Clerodendrum infortunatum* L., *International Journal of Green Pharmacy*, 7, 131–135, <https://doi.org/10.4103/0973-8258.116394>, 2013.
- von der Lüche, B., Birk, J. J., Dawson, L., Mayes, R. W. and Fiedler, S.: Steroid fingerprints: efficient biomarkers of human decomposition fluids in soil, *Organic Geochemistry*, 124, 228–237, <https://doi.org/10.1016/j.orggeochem.2018.07.016>, 2018.
- 620 von der Lüche, B., Dawson, L., Mayes, R. W., Forbes, S. L. and Fiedler, S.: Investigation of sterols as potential biomarkers for the detection of pig (*S. s. domesticus*) decomposition fluid in soils, *Forensic Science International*, 230, 68–73, <https://doi.org/10.1016/j.forsciint.2013.03.030>, 2013.
- 625 von der Lüche, B., Prost, K., Birk, J. J. and Fiedler, S.: Steroids aid in human decomposition fluid identification in soils of temporary mass graves from World War II, *Journal of Archaeological Science: Reports*, 32, 102431, <https://doi.org/10.1016/j.jasrep.2020.102431>, 2020.
- Wiedner, K., Schneeweiß, J., Dippold, M. and Glaser, B.: Anthropogenic Dark Earth in Northern Germany — The Nordic Analogue to terra preta de Índio in Amazonia, *Catena*, 132, 114–125, <https://doi.org/10.1016/j.catena.2014.10.024>, 2015.

- 630 Zech, M., Lerch, M., Bliedtner, M., Bromm, B., Seemann, F., Szidat, S., Salazar, G., Zech, R., Glaser, B., Haas, J. N., Schäfer, D. and Geitner, C.: Revisiting the subalpine Mesolithic site Ullafelsen in the Fotsch Valley, Stubai Alps, Austria – new insights into pedogenesis and landscape evolution from leaf wax-derived n-alkane biomarkers, black carbon and radiocarbon dating, *E&G Quaternary Science Journal*, 70, 171–186, <https://doi.org/10.5194/egqsj-70-171-2021>, 2021.
- 635 Zech, W., Schad, P. and Hintermaier-Erhard, G.: *Böden der Welt*, 2, Springer Spektrum Verlag, Berlin Heidelberg, Germany, 2014.
- Zech, W. and Wilke, B.: Vorläufige Ergebnisse einer Bodenchronosequenzstudie im Zillertal, *Mitteilungen der Deutschen Bodenkundlichen Gesellschaft*, 25 (1), 571–586, 1977.

Table 1: Overview over all soil profiles, soil and modern faeces samples from the prehistorical encampment site of Ullafelsen and reference sites in the Fotsch Valley, Stubai Alps, Austria. Analytical results of TOC, TN and TOC/N are presented.

	Soil profile	Altitude [m a.s.l.]	Soil profile coordinates	Sample no.	Soil horizon	TOC [%]	TN [%]	TOC/N
Archaeological soil profiles [Ullafelsen]	1.1 C4w	1869	N 47.14702° E 11.21475°	1	OAh1	15.1	0.8	20.0
				2	OAh2	8.5	0.3	24.6
				3	OAh3	4.3	0.1	33.8
				4	E (LL)	3.2	0.1	25.0
				5	Bh	8.3	0.3	28.1
				6	Bs	1.6	0.1	24.4
				7	BvCv	0.8	0.0	17.9
	1.1 B5s	1869	N 47.14704° E 11.21474°	8	OAh1	10.3	0.6	17.9
				9	E (LL)	3.3	0.1	22.5
				10	Bh	6.4	0.2	26.8
				11	Bs	2.2	0.1	24.5
				12	BvCv	0.3	0.0	12.4
	1.1 B5w	1869	N 47.14703° E 11.21474°	13	OAh1	18.2	1.0	18.0
				14	E (LL)	4.5	0.1	37.2
				15	Bh	6.1	0.2	28.3
				16	Bs	2.3	0.1	27.2
				17	BvCv	0.7	0.0	16.2
	1.1 G5n	1869	N 47.14704° E 11.21482°	18	OAh1	14.2	0.7	18.9
				19	OAh2	8.4	0.3	26.5
				20	OAh3	25.0	0.9	28.4
				21	E (LL)	3.3	0.1	25.2
				22	Bh	7.1	0.3	26.8
	1.9 NW	1867	N 47.14698° E 11.21492°	23	OAh1	19.3	1.2	16.7
				24	OAh2	12.5	0.7	18.6
				25	OAh3	28.8	1.0	27.7
				26	E (LL)	2.5	0.1	19.5
				27	Bh	4.8	0.2	24.9
				28	Bs	2.5	0.1	27.1
				29	BvCv	1.6	0.1	26.8
1.9 NE	1867	N 47.14699° E 11.21494°	30	OAh1	13.6	0.7	18.5	
			31	OAh2	11.9	0.5	21.7	
			32	E 1 (LL)	4.3	0.2	28.3	
			33	Bh	5.5	0.2	23.1	
			34	Bs	3.6	0.1	27.8	
			35	E 2 (LL)	1.3	0.1	23.6	
			36	BvCv	2.1	0.1	27.1	
			37	E (LL)	1.3	0.1	16.9	
Reference soil profiles [Fotsch Valley]	4.4	2171	N 47.15060° E 11.20075°	38	OAh	21.9	1.2	18.4
				39	Bh	6.0	0.3	23.3
	4.11	2548	N 47.14373° E 11.17502°	40	OAh	11.9	0.8	15.2
				41	Bh	12.9	0.7	18.6
	4.12	2455	N 47.14584° E 11.18304°	42	OAh	6.6	0.5	13.7
				43	Bh	7.9	0.5	16.0
	5.5	2186	N 47.15025° E 11.19981°	44	OAh	16.0	0.9	17.1
				45	Bh	4.9	0.3	18.5
5.6w	2186	N 47.14583° E 11.20402°	46	OAh	29.4	2.2	13.1	
Faeces I	Cattle		47		43.5	3.0	14.5	
Faeces II	Sheep		48		42.6	2.3	18.4	




645 **Figure 1: The prehistorical encampment site of Ullafelsen in the Fotsch Valley, Stubai Alps, southwest of Innsbruck, Tyrol, Austria. Northward view from the inner Fotsch Valley over the Ullafelsen (1869 m a.s.l.) to the Karwendel mountain range in the Northern Limestone Alps (Photo: E. Hüsing, 2018).**




650 **Figure 2: Left: NNW view over the Ullafelsen in the Fotsch Valley, Stubai Alps, Austria at the upper timberline. The geoarchaeological excavation area on the Ullafelsen plateau with the reopened and sampled soil profiles 1.1 B5, 1.1 C4 and 1.1 G5 and the newly opened and sampled soil profile 1.9 several meters towards the southeast are shown. Right: Reopened archaeological excavation area. View to the southwestern part of the sampled soil profiles (Zech et al., 2021).**



655 **Figure 3:** Left: Soil profile 1.9 NW, which represents a typical soil profile for the Ullafelsen in the Fotsch Valley, Stubai Alps, Austria. Right: Schematic horizons of the soil profiles on the Ullafelsen. Note that the soil profiles reveal a high heterogeneity. Nevertheless, the soil horizons OAh3 and Bh are characterized by a humus-enrichment. The E (LL) horizon ("light layer") reveals the highest relative artifact abundance (~41%) and is overlain by several fireplaces on the Ullafelsen. This horizon is considered as living floor of the Mesolithic hunter-gatherers (Geitner et al., 2011; Geitner et al., 2014). Due to TOC content partly $\geq 15\%$, we adopted the soil horizon classification by Zech et al. (2021).



660 **Figure 4:** Biomarker patterns of modern faeces from predominating livestock at the prehistorical encampment site of Ullafelsen in the Fotsch Valley, Stubai Alps, Austria and surroundings. Steroid (sterols, stanols, stanones and bile acids) contents of cattle and sheep faeces are given in $\mu\text{g g}^{-1}$ dry matter.



665 **Figure 5:** Box plots illustrating the total steroid (sterols, stanols, stanones and bile acids) contents (in $\mu\text{g g}^{-1}$ dry matter) of the investigated soil profiles from the prehistorical encampment site of Ullafelsen and reference sites in the Fotsch Valley, Stubai Alps, Austria, categorized by the soil horizons OAh1 (n=6), OAh2 (n=4), OAh3 (n=3), E (LL) (n=8), Bh (n=6), Bs (n=5) and BvCv (n=5).

670

675






Figure 6: Box plots visualizing the content (in $\mu\text{g g}^{-1}$ dry matter) of steroid (sterols, stanols, stanones and bile acids) patterns for all soil samples from the prehistorical encampment site of Ullafelsen in the Fotsch Valley, Stubai Alps, Austria. For a better overview, the steroids stigmastanol, β -sitosterol and 5α -stigmastanol were plotted to a separate y-axis because of their high contents.



705 **Figure 7: Box plots indicating steroid ratios for estimating the origin of faecal matter. Ratio 1 describes the input of human faeces in soils, whereas ratio 2 and ratio 5 determine the input of human vs. ruminant faeces. Thresholds ratio 1: ratios < 0.7 assume low human faeces input, ratios > 0.7 show high human faeces input. Thresholds ratio 5: ratios < 5 point to a dominant human faeces input, ratios > 5 represent a dominant ruminant faeces input. Ratio 3 considers the input of β -sitosterol as plant-derived steroid.**



710 **Figure 8: Microbial transformation of 5β-stigmastanol to epi-5β-stigmastanol (Ratio 4). Epimerization of 5β-stanols ("Degradation effect") takes place over time due to microbial processes in soils (Bull et al., 2001; Prost et al., 2017).**