

# Growth rate rather than temperature affects the B/Ca ratio in the calcareous red alga *Lithothamnion coralliooides*

Giulia Piazza<sup>1</sup>, Valentina A. Bracchi<sup>1</sup>, Antonio Langone<sup>2</sup>, Agostino N. Meroni<sup>3</sup>, Daniela Basso<sup>1</sup>

<sup>1</sup> Department of Earth and Environmental Sciences, University of Milano-Bicocca, Piazza della Scienza 4, 20126 Milano, Italy

<sup>2</sup> CNR – Institute of Geosciences and Earth Resources, Via Ferrata 1, 27100 Pavia, Italy

<sup>3</sup> Department of Civil and Environmental Engineering, Politecnico of Milan, Piazza Leonardo da Vinci 32, 20133 Milano, Italy

Correspondence to: Giulia Piazza (g.piazza15@campus.unimib.it)

**Abstract.** The B/Ca ratio in calcareous marine species is informative of past seawater  $\text{CO}_3^{2-}$  concentrations, but scarce data exist on B/Ca in coralline algae. Recent studies suggest influences of temperature and growth rates on B/Ca, the effect of which could be critical for the reconstructions of surface ocean pH and atmospheric  $p\text{CO}_2$ . In this paper, we present the first LA-ICP-MS analyses of Mg, Sr, Li and B in the coralline alga *Lithothamnion coralliooides* collected from different geographic settings and depths across the Mediterranean Sea and in the Atlantic Ocean. We produced the first data on putative temperature proxies (Mg/Ca, Li/Ca, Sr/Ca, Mg/Li) and B/Ca in a coralline algal species grown in different Basins ([the Mediterranean Sea and the Atlantic Ocean](#)), from across the photic zone (12 m, 40 m, 45 m, and 66 m depth). We tested the B/Ca correlation with temperature proxies and growth rates, in order to evaluate their possible effect on B incorporation. Our results suggested a growth rate influence on B/Ca, especially in the deepest sample (Pontian Isl., Italy; 66 m) and in the shallowest sample (Morlaix, Atlantic coast of France; 12 m), where the growth rates were respectively 0.11 mm/yr and 0.13 mm/yr and the B/Ca was respectively  $462.8 \pm 49.2 \mu\text{mol/mol}$  and  $726.9 \pm 102.8 \mu\text{mol/mol}$ . A positive correlation between B/Ca and [the temperature proxies](#) was found only in Morlaix, where the seasonal temperature variation ( $\Delta T$ ) was the highest (8.9 °C). [This evidence suggests](#) [We concluded](#) that growth rates, triggered by the different  $\Delta T$  and light availability across depth, affect the B incorporation in *L. coralliooides*.

## 1 Introduction

Warming and acidification are major anthropogenic perturbations of present-day oceans ([Callendar, 1938; Fairhall, 1973; Alverson et al., 2001; Brewer, 1997; Gattuso, 1999; Caldeira, 2005; Feely et al., 2009; Le Quéré et al., 2009; Hansen et al., 2010; Hönisch et al., 2012; Ciais et al., 2013; Masson-Delmotte et al., 2021](#)

[\). Ocean acidification reduces the saturation state of calcite and aragonite, lowering the dissolution threshold of biominerals and threatening habitat-forming species of critical ecological importance such as coralline red algae and corals \(Morse et al., 2006; Hoegh-Guldberg et al., 2007; Andersson et al., 2008, 2011; Basso, 2012; Ragazzola et al., 2012; Ries et al., 2016](#)

[\). Coralline algae, which precipitate high Mg-calcite \(>8-12 mol%  \$\text{MgCO}\_3\$ \) \(Morse et al., 2006](#)

their worldwide distribution and longevity. Importantly, they show indeterminate growth, with no ontogenetic trend (Halfar et al., 2008), that is the growth trend of coralline algae does not slow down asymptotically with age, as in bivalves, thus preserving the resolution of the geochemical signals in all stages of growth (Adey, 1965; Frantz et al., 2005; Halfar et al., 2008). Moreover, coralline algae thin sections under optical microscopy reveal bands that reflect the growth pattern (Cabioch, 1966; Basso, 35 1995a, b; Foster, 2001), similarly to tree rings (Ragazzola et al., 2016) that can be targeted for high-resolution geochemical analyses. Seasonal growth bands, indeed, consist of the perithallial alternation of dark and light bands that together constitute the annual growth patterns (Freiwald and Henrich, 1994; Basso, 1995a, b; Kamenos et al., 2009). Dark bands correspond to slow-growing cells produced in the cold season, which are shorter, thick-walled and with lower Mg contents; while light bands are fast-growing cells produced in the warm season, which are longer, less calcified and with higher Mg concentrations 40 (Kamenos et al., 2009; Ragazzola et al., 2016).

Trace element variations in marine calcareous species inform the reconstruction of changes in the environmental parameters, which characterized the seawater during their growth (Hetzinger et al., 2011; [Montagna and Douville, 2017](#)). Boron is incorporated into the mineral lattice of calcareous marine species during calcite precipitation. In the ocean, B occurs in two molecular species: boric acid  $\text{B}(\text{OH})_3$  and borate ion  $\text{B}(\text{OH})_4^-$  (Dickson, 1990), which are related by the following acid-base 45 equilibrium reaction:



that shows the dependence of the two species concentration on pH. ~~In typical surface seawater conditions, the equilibrium constant of the reaction  $K_B$  is  $10^{-8.6}$  which is equal to a  $pK_B$  of 8.6 (Dickson, 1990), very similar to seawater pH (~8) (Zeebe and Wolf-Gladrow, 2001). Hence, the reaction described in Eq. (1) is at equilibrium. As the pH lowers, based on Eq. (1), the  $[\text{H}^+]$  in seawater increases together with the  $[\text{B}(\text{OH})_3]$ , while the  $[\text{B}(\text{OH})_4^-]$  decreases.~~ Boron has two stable isotopes: the heavier  $^{11}\text{B}$  (80.1% of the total abundance) and the lighter  $^{10}\text{B}$  (19.9%) with an enrichment factor  $\varepsilon^{11}\text{B}$  ( $\delta^{11}\text{B}(\text{OH})_3 - \delta^{11}\text{B}(\text{OH})_4^-$ ) equal to  $1.0272 \pm 0.0006$  (Klochko et al., 2006). ~~The current seawater isotopic composition  $\delta^{11}\text{B}_{\text{sw}}$  is 39.61‰ (Foster et al., 2010).~~ The isotopic composition of  $\text{B}(\text{OH})_3$  and  $\text{B}(\text{OH})_4^-$  changes with pH, being both enriched in  $\delta^{11}\text{B}$  with increasing pH (Dickson, 1990; [Foster et al., 2010](#)). The first analyses of the isotopic signal of marine carbonates evidenced a strong similarity 55 with the isotopic composition of  $\text{B}(\text{OH})_4^-$  in solution, suggesting that borate would preferentially be incorporated into marine carbonates (Vengosh et al., 1991; Hemming and Hanson, 1992; [DeCarlo et al., 2018](#)). Therefore, the  $\delta^{11}\text{B}$  of marine carbonates should increase with pH according to the  $\delta^{11}\text{B}$  of  $\text{B}(\text{OH})_4^-$ . The  $\delta^{11}\text{B}$  of borate is closely related to  $p\text{CO}_2$ , seawater  $[\text{CO}_3^{2-}]$ , pH and  $[\text{B}(\text{OH})_4^-]$ , which in turn are mainly controlled by the balance between the total alkalinity (TA) and the dissolved inorganic carbon (DIC) (Zeebe and Wolf-Gladrow, 2001; Dickson et al., 2007; [Foster and Rae, 2016](#)). ~~Recent magic angle spinning nuclear magnetic resonance (MAS-NMR) techniques, though, revealed that 30–48% of  $\text{B}(\text{OH})_3$  is incorporated in corals, foraminifera, and calcareous red algae (Klochko et al., 2009; Rollion Bard et al., 2011; Cusack et al., 2015). The amount of incorporated boric acid seems to be unrelated to the boron species concentrations in seawater (Cusack et al., 2015) or pH (Mavromatis et al., 2015) and could be the consequence of a coordination change during biomineralization (Klochko et al., 2009; Mavromatis et al., 2015).~~ The above-described theoretical overview frames the role of the B content and its isotopic 60

65 signature ( $\delta^{11}\text{B}$ ) in calcareous marine species, as recorders of information about the seawater carbonate system. The  $\delta^{11}\text{B}$  is used to reconstruct past seawater pH (Hönisch and Hemming, 2005; Foster, 2008; Douville et al., 2010; Paris et al., 2010; Rae et al., 2011). The boron-to-calcium ratio (B/Ca) proved to be informative about past seawater  $\text{CO}_3^{2-}$  concentrations in different empirical studies on benthic foraminifera (Yu and Elderfield, 2007; Yu et al., 2007; Rae et al., 2011) and in synthetic aragonite (Holcomb et al., 2016). Most of the literature on boron studies is focused on its isotopic composition (Hemming and Hönisch, 70 2007; Klochko et al., 2009; Henehan et al., 2013; Fietzke et al., 2015; Cornwall et al., 2017; Ragazzola et al., 2020), whereas less attention has been given to B/Ca records, especially in coralline algae. Recent studies suggest that B/Ca is function of seawater pH, as well as of other environmental variables such as temperature, whose effect should be considered in the attempt to reconstruct surface ocean pH and atmospheric  $p\text{CO}_2$  (Wara et al., 2003; Allen et al., 2012; Kaczmarek et al., 2016).

To achieve the best reliability of geochemical proxies for climate reconstructions, a growing interest has arisen on multiple 75 approaches, by considering multi-proxies for a single environmental factor (D'Olivo et al., 2018; Cuny-Guirrec et al., 2019; Zinke et al., 2019). Moreover, it is important to recognize the influence of multiple factors on a single proxy (Kaczmarek et al., 2016; Donald et al., 2017). For instance, more recently the effects of temperature and growth rate on B incorporation have been investigated through experiments on both synthetic and biogenic carbonates (Wara et al., 2003; Yu et al., 2007; Gabitov et al., 2014; Mavromatis et al., 2015; Uchikawa et al., 2015; Kaczmarek et al., 2016; Donald et al., 2017). In particular, culture 80 experiment on the coralline alga *Neogoniolithon* sp. (Donald et al., 2017) showed a positive correlation of B/Ca with growth rate, and a negative correlation with Sr/Ca, which was proposed by the authors as proxy of DIC. Moreover, a culture experiment on the high latitude species *Clathromorphum compactum* (Kjellman) Foslie 1898 revealed non-significant temperature influences on B/Ca and a significant inverse relationship with growth rate (Anagnostou et al., 2019). The factors which influence the B incorporation in calcareous red algae are therefore still debated. Nevertheless, no studies have been conducted 85 so far on the correlation between temperature proxies (Mg, Sr, Li /Ca) and B/Ca. The Mg/Ca ratio is extensively used as a temperature proxy in calcifying species (Barker et al., 2005), including coralline algae (Halfar et al., 2008; Kamenos et al., 2008; Fietzke et al., 2015; Ragazzola et al., 2020), since the substitution of  $\text{Mg}^{2+}$  with  $\text{Ca}^{2+}$  ions in the calcite lattice is an endothermic reaction. Accordingly, Mg incorporation increases with temperature (Moberly, 1968; Berner, 1975; Ries, 2006; Caragnano et al., 2017). Sr/Ca and Li/Ca ratios in calcareous red algae have also been investigated as climate proxies, showing 90 significant positive correlations with temperature in different species, e.g. *Lithothamnion* spp. (Kamenos et al., 2008; Hetzinger et al., 2011; Caragnano et al., 2014; Darrenougue et al., 2014). The Mg/Li ratio showed a strong correlation with seawater temperature in cultured *Clathromorphum compactum* (Anagnostou et al., 2019), and in empirical studies on high-Mg calcites, including coralline algae (Stewart et al. 2020). Conversely, the Mg/Li calibration did not reveal improvements in the Mg/Ca or Li/Ca proxies in *Lithophyllum* spp. (Caragnano et al., 2014, 2017).

95 Here, we present a laser ablation inductively coupled plasma mass spectrometry (LA-ICP-MS) conducted on a wild grown coralline alga with a wide geographic scope. This technique, which allows high-resolution analysis of a broad range of trace elements in solid-state samples, has been widely used in biogenic carbonates to extract records of seawater temperature, salinity, and water chemistry (Schöne et al., 2005; Corrège, 2006; Hetzinger et al., 2009, 2011; Fietzke et al., 2015; Ragazzola


et al., 2020). Measurements were made on the non-geniculate coralline alga *Lithothamnion corallioides* (P. Crouan & H. 100 Crouan) P. Crouan & H. Crouan 1867, which is widely distributed in the Mediterranean Sea and in the north-eastern Atlantic Ocean, from Scotland to Canary Islands (Irvine and Chamberlain, 1999; Wilson et al., 2004; Carro et al., 2014), usually constituting maerl beds (Potin et al., 1990; Foster, 2001; Martin et al., 2006; Savini et al., 2012; Basso et al., 2017). It forms rhodoliths as unattached branches (Basso et al., 2016) with obvious banding in longitudinal sections (Basso, 1995b). These 105 characteristics combine to make this species a suitable model for the measurement of geochemical proxies, comparing different environmental settings.

In this paper, we provide the first LA-ICP-MS data of putative temperature proxies (Mg/Ca, Sr/Ca, Li/Ca, Mg/Li) and B/Ca measured on *L. corallioides* collected from different geographic settings and depths across the Mediterranean Sea and in the Atlantic Ocean. We test the influence of temperature and growth rate on the B/Ca ratio, which could be crucial in assessing the reliability of B/Ca as a proxy of the seawater carbonate system.

## 110 2 Materials and methods

### 2.1 Sampling sites and collection of *Lithothamnion corallioides*

Samples of the coralline alga *L. corallioides* were collected in the Western Mediterranean Sea and in the Atlantic Ocean (Fig. 1). In the Mediterranean Sea, the samples collected in Pontian Islands (Italy) at 66 m depth were gathered by grab during the 115 cruises of the R/V Minerva Uno, in the framework of the Marine Strategy Campaigns 2017 (Table 1). The last two Mediterranean samples were collected by one of the authors (DB) by SCUBA diving during local surveys at 45 m off the coasts of Pomonte (Elba Island, Italy) (Basso and Brusoni, 2004) and at 40 m depth in the Aegadian Islands (Marettimo, Italy). The Atlantic sample was collected by grab at 12 m depth in Morlaix Bay (Brittany, France) (Table 1).



120 **Figure 1: Map showing the distribution of sampling sites where *Lithothamnion coralliooides* samples were collected: Morlaix Bay (48°34'42"N, 3°49'36"W); Aegadian Islands (37°97'36"N, 12°14'12"E); Elba (42°44'56.4"N, 10°07'08.4"E) and Pontian Islands (40°54'N, 12°45'E). Service layer credits: Source Esri, GEBCO, NOAA, National Geographic, Garmin, geonames.org and other contributors.**

The identification of the algal samples was achieved by morphological analyses of epithallial and perithallial cells using a Field Emission Gun Scanning Electron Microscope (SEM-FEG) Geomini 500 Zeiss. Samples were prepared for SEM according to Basso (1995a). Morphological identification was based on Adey and McKibbin (1970) and Irvine and Chamberlain (1994). Other information about maerl species distribution in Morlaix was provided by Carro et al. (2014) and Melbourne et al. (2017). *L. coralliooides* was selected as the target species because of its presence in both Mediterranean and Atlantic waters. The Atlantic sample (Morlaix) was used as voucher specimen for the subsequent identification of the Mediterranean samples, since *Phymatolithon* spp. and *L. coralliooides* are the only components of the Morlaix maerl (Carro et al., 2014; Melbourne et al., 2017). Hence, once its inclusion under the genus *Phymatolithon* was excluded, the Morlaix sample identified as *L. coralliooides* was used as a reference for the most reliable identification of the other Mediterranean samples.

## 2.2 Sample preparation


The selected algal branches were embedded in Epo-Fix resin, which was stirred for 2 minutes with a hardener (13% w/w); they were then left to dry at room temperature for 24 hours. Afterwards, the treated branches were cut by an IsoMet diamond wafering blade 15HC along the direction of growth. In the laboratory of the Institute of Geosciences and Earth Resources (IGG) of the National Research Council (CNR) in Pavia (Italy), the sections were polished with a MetaServ Grinder-Polisher (400 RPM) using a diamond paste solution, finally cleaned ultrasonically in distilled water for 10 minutes and dried at 30°C for 24 hours.

## 2.3 Trace elements analyses and environmental data

140 LA-ICP-MS analyses were carried out at the IGG-CNR laboratory of Pavia (Italy), on one algal branch per sampling site. <sup>43</sup>Ca, <sup>7</sup>Li, <sup>25</sup>Mg, <sup>88</sup>Sr and <sup>11</sup>B contents were measured using an Agilent ICP-QQQ 8900 quadrupole ICP-MS coupled to an Excimer laser ablation system (193 nm wavelength, [MicroLas with GeoLas optics](#)). Element/Ca ratios were calculated for these isotopes, as well as Mg/Li ratio. Measurements were performed with laser energy densities of 4 J/cm<sup>2</sup> and helium as carrier gas.

145 The laser transects crossed the algal growth direction with a spot size of 50 µm, attempting to target each growth band change which marked the transition between the cells usually produced in the warm season and those usually produced in the cold season, hereinafter referred to as long and short cells (Fig. 2, S1, S2, S3). NIST 612 was used as an external standard (e.g. Fietzke et al., 2010; Jochum et al., 2012), whereas Ca was adopted as an internal standard. Accuracy and precision were ~~better~~higher than 4% [for NIST 612](#) and 8% [for Ca standard respectively](#). [Minimum detection limits \(99% confidence\) of measured elements were: Ca = 16.91 ppm, Li = 0.07 ppm, Mg = 0.11 ppm, Sr = 0.004 ppm, B = 2.64 ppm](#). Each analysis was carried out in MS/MS mode for 3 minutes by acquiring 60 seconds of background before and after the sampling period by the

laser on the polished surface. The first part of the signal was not used for the integration to avoid surface contamination. The Glitter software (v. 4.4.4) was used for data reduction.



155 **Figure 2: (a) Thalli of *Lithothamnion coralliooides* collected in Morlaix (scale bar = 5 mm). (b) Longitudinal section through the *L. coralliooides* branch sampled in Morlaix showing the LA-ICP-MS transects targeting each growth bands (scale bar = 200  $\mu$ m).**

In the absence of in-situ environmental data, the seawater temperature has been extracted by 11 years of monthly reanalysis spanning 1979-2017 from ORAS5 (Ocean ReAnalysis System 5), at 0.25-degree horizontal resolution (Zuo et al., 2019). The time interval of extraction in each site corresponded to 11 years before sample collection. Minimum, maximum, and mean values, as reported in Table 1, refer to the temperature at sampling depth and have been measured on the entire time interval of extraction.

160 For the purpose of this work, we considered temperature data in terms of a) the amplitude of temperature variation ( $\Delta T$ ), and b) the temperature maxima and minima.  $\Delta T$  represents the temperature fluctuations during the algal growth and has been measured as the difference between the maximum and minimum temperature registered in the site over 11 years. We used  $\Delta T$  when comparing the sampling sites, given their differences in depth and geographical regions. The temperature peaks (maximum and minimum) have rather been used when considering data corresponding to long and short cells, since they are related to warm and cold periods of growth, respectively. We used the temperature peaks over the entire time interval of extraction (11 years) when comparing the mean elemental ratios of long and short cells per sampling sites. The maximum and minimum temperature within each year have been used instead for the reconstruction of the algal age model. In the sample 165 from Morlaix Bay, indeed, the good visibility of the growth bands allowed us to relate the temperature record with the algal growth at annual resolution. We therefore plotted all the element ratios against the average seawater temperature values of the coldest and warmest months of the year to reconstruct the temperature variations during the algal growth (Moberly, 1970; Corrège, 2006; Williams et al., 2014; Ragazzola et al., 2020; Caragnano et al., 2014). Missing element ratios possibly due to 170 non-targeted consecutive bands were calculated as the means of known values.

175 Carbon system parameters for each site have also been estimated. Even if they were not available in the same time interval of temperature data, the seasonal variations occurring in the extracted period have allowed the characterization of the sampling sites. Monthly mean seawater pH has been derived by the CMEMS (E.U. Copernicus Marine Service Information) global biogeochemical hindcast spanning 1999-2017, at 0.25-degree horizontal resolution. Monthly means of DIC in the time interval 1999-2017 have been extracted by CMEMS biogeochemical reanalysis for the Mediterranean Sea, at 0.042-degree horizontal  
180 resolution (Teruzzi et al., 2021). In the Atlantic site, monthly means of DIC in 1999-2017 were derived from CMEMS IBI biogeochemical model, at 0.083-degree horizontal resolution. The pH and DIC data showed consistent variations among sites, despite being derived from different biogeochemical models. The mean values of DIC and pH, as reported in Table 1, refer to sampling depth and have been measured on the entire time interval of extraction. The complete timeline of temperature and carbon data used in this paper is shown in Supplement Fig. S5, S6, S7, S8.

## 185 2.4 Growth rate estimation

Growth rates were estimated under light microscope by measuring the length of the LA-ICP-MS transect and dividing it by the number of annual growth bands crossed by the transect. The obtained values have been expressed in linear extension over year (mm/yr). In the samples where the growth bands were not easily detectable under microscope, i.e. the Elba sample, we also used the Mg/Ca results in order to check for the correspondence of Mg peaks with growth bands.

## 190 2.5 Statistical analysis

Statistics were performed on both the dataset with all the spot analyses and the dataset with the records from long and short cells separately. Short cells were referred to slow-growing cells in dark bands, usually produced in the cold season; long cells corresponded to fast-growing cells in light bands, usually produced in the warm season (Kamenos et al., 2009; Ragazzola et al., 2016). For each spot, a distinction between the cells was thus made by image analyses, except for the Elba sample, given  
195 the poor resolution of the growth bands. The Spearman's correlation was tested to provide the statistical comparisons between Mg/Ca, Li/Ca, Sr/Ca and B/Ca records from the LA-ICP-MS analyses in *L. corallioides*. The Kruskal-Wallis test followed by the Dunn's test for comparisons and the One-way ANOVA followed by the Tukey's test for post-hoc analysis were used to compare the geochemical signals among sampling sites and to evidence the differences between group medians and means. All statistical analyses were run in R 3.6.3 software.

## 200 3 Results

### 3.1 Environmental data

The temperature data obtained by ORAS5 reanalysis revealed a lower amplitude of the seasonal temperature change in the Mediterranean samples with respect to the Atlantic site, as shown by the standard deviation and  $\Delta T$  values in Table 1. This difference is explained in terms of the different sampling depths, with the seasonal variations decreasing with increasing depth.

205 Temperature variations in Morlaix Bay (Atlantic Ocean) were higher, registering overall a mean seawater temperature of 12.4 °C (Table 1). Among Mediterranean samples, mean seawater temperatures were highest in the Aegadian Isl., followed by the Elba and the Pontian Isl. (Table 1). Aegadian Isl. also registered the highest temperature variations among the Mediterranean sites (Table 1). Moderate temperature variations characterized the site in Elba, which registered the lowest monthly mean temperature among Mediterranean sites (Table 1). At the Pontian Isl., consistently with the fact that it is the deepest sampling  
210 site at 66 m depth, the lowest seawater temperature variations were found (Table 1).

The pH estimates at the Mediterranean sites were all similarly high ~8.13 and less variable than the Atlantic site (8.05). The mean pH had slightly higher values in Pontian Isl. and Elba, than Aegadian Isl. (Table 1). Similarly, DIC was higher in the Mediterranean sites and decreased in Morlaix, as this is largely dictating the pH (Table 1).

215 **Table 1: Environmental data in each sampling site. The minimum and maximum monthly means of temperature are indicated, as well as the highest temperature variation (ΔT), the mean and the standard deviation of the time series. Data from monthly means extracted by 11 years of ORAS5 reanalysis. pH and DIC in each sampling site are also indicated. The minimum, maximum, mean, and standard deviation values have been measured on the time interval 1999-2017. Carbonate system parameters have been extracted from monthly means biogeochemical data provided by CMEMS.**

Sampling site	Depth (m)	Collection date	Temperature (°C)				pH		DIC (µmol/kg)		Growth rates (mm/yr)
			min	max	ΔT	mean	st. dev.	mean	st. dev.	mean	
Aegadian Isl.	40	25/08/1993	13.7	18.8	5.1	15.9	1.3	8.11	0.02	2303.72	11.58
Elba	45	01/12/1990	12.9	17.5	4.6	15.0	1.2	8.13	0.02	2322.24	7.08
Pontian Isl.	66	14/07/2017	13.9	16.7	2.8	15.0	0.6	8.13	0.01	2321.29	6.71
Morlaix	12	02/05/1991	8.3	17.2	8.9	12.4	2.5	8.05	0.04	2155.60	15.27

### 3.2 Mg/Ca, Li/Ca, Sr/Ca and Mg/Li

220 Both Li/Ca and Sr/Ca records had positive correlations with Mg/Ca in our samples of *L. corallinaoides* (Fig. 3 and Supplement Fig. S4). The overall mean Mg/Ca was  $225.3 \pm 30.4$  mmol/mol, registering the minimum value in the sample from Aegadian Isl. (171.7 mmol/mol) and the maximum value in Morlaix (311.2 mmol/mol) (Fig. 4; Table 2). The Kruskal-Wallis test did not show significant differences in Mg/Ca among samples (Table A1; Fig. 4). Among Mediterranean sites, the algal sample from Aegadian Isl. had the highest Mg/Ca mean value, followed by Elba and Pontian Isl., which had the lowest Mg/Ca mean  
225 value of all sampling sites (Fig. 4). The highest mean Mg/Ca was registered in the sample from Morlaix bay, which also showed a large dispersion of data above the median Mg/Ca value (Fig. 4).





Figure 3: Correlation plots of Mg/Ca with Li/Ca and Sr/Ca. For each analyses the Spearman's coefficient  $r$ , the  $p$ -value and the line equation are given.




230

Figure 4: Box plot of the statistical tests performed to evaluate the differences of Mg/Ca in *L. corallinaoides* collected in different sampling sites. The horizontal black lines indicate the median values. The black filled circles and the numbers inside the plot indicate the mean values.

Long cells had high Mg/Ca values; conversely, short cells corresponded to areas with low Mg/Ca ratio.

235 The ANOVA test followed by the Tukey's test for multiple comparisons evidenced a significant variability of algal Mg/Ca among 3 sites in long cells (Table A2; Fig. 5). In the long cells of *L. corallinaoides* collected from Aegadian Isl. and Pontian Isl.,

the Mg/Ca data showed quite similar distribution (Table A2; Fig. 5). The Mg/Ca of the alga from Pontian Isl. was the lowest (Fig. 5). In Morlaix a higher Mg/Ca mean value was registered, significantly different compared to Aegadian and Pontian Isl. (Table A2; Fig. 5). In short cells, the differences in Mg/Ca among samples were not statistically significant (Table A1). The 240 magnesium incorporation was slightly higher in Morlaix and very similar between Aegadian and Pontian Isl. samples (Fig. 5).



**Figure 5: Box plots of the statistical tests performed to evaluate the differences of Mg/Ca in the long and short cells of *L. coralliooides* collected in different sampling sites. The horizontal black lines indicate the median values. The black filled circles and the numbers inside the plot indicate the mean values.**

245 Mg/Li values in long and short cells fell in the range found by Anagnostou et al. (2019) for cultured *Clathromorphum compactum* (Fig. 8). When plotted against the extracted seawater temperature in Morlaix (Fig. 9), Mg/Li results did not reflect the seasonal oscillations in temperature.




Figure 6: Correlation plot between Mg/Li and seawater temperature. Data are shown for cultured *C. compactum* (Anagnostou et al., 250 2019) and *L. corallioides* (this paper). *L. corallioides* results are shown separately in long and short cells, per sampling site.





Figure 7: Mg/Li ratio of *L. corallioides* collected in Morlaix Bay. Note the lack of cyclic variations in Mg/Li results. In the timeline, the coldest and the warmest months have been reported. Mg/Li in the missing bands (asterisks) have been calculated as the means of the values measured in warm and cold periods. Monthly means of seawater temperature have been extracted by ORASS 255 reanalysis.

### 3.3 B/Ca


The B/Ca ratio in the sample collected from Morlaix showed a moderate positive correlation with all the examined temperature proxies (Mg/Ca, Li/Ca, Sr/Ca), with a more defined trend when plotted against Li/Ca ( $r=0.68$ ) and slightly less defined against Mg/Ca ( $r=0.58$ ) and Sr/Ca ( $r=0.57$ ) (Fig. 8). On the contrary, the Spearman's analyses did not evidence significant correlations 260 between B/Ca and the temperature signals in the algae collected elsewhere ( $p>0.05$ ).

Overall, the B/Ca ratio in *L. corallioides* was  $661.9 \pm 138.9 \mu\text{mol/mol}$ , registering the minimum value in the long cells of the sample from Pontian Isl. ( $356.0 \mu\text{mol/mol}$ ) and the maximum value in Elba ( $954.1 \mu\text{mol/mol}$ ) (Fig. 9; Table 2).

The Kruskal-Wallis coefficient showed a highly significant difference in the B/Ca value among sites, particularly in the *L. coralliooides* from the Pontian Isl., which had the lowest boron incorporation (Table A3; Fig. 9). The algae collected in Aegadian Isl. had still significantly lower B/Ca compared to those collected in Elba and Morlaix (Table A3; Fig. 9). The highest B/Ca mean value was registered in Elba, with medians comparable to Morlaix (Table A3; Fig. 9).



**Figure 8: Correlation plots of B/Ca with Mg/Ca, Li/Ca and Sr/Ca in *L. coralliooides* collected in Morlaix bay. For each analyses the Spearman's coefficient  $r$ , the  $p$ -value and the line equation are given.**




270

**Figure 9: Box plot of the statistical tests performed to evaluate the differences of B/Ca in *L. coralliooides* collected in different sampling sites. The horizontal black lines indicate the median values. The black filled circles and the numbers inside the plot indicate the mean values.**

The ANOVA test followed by the Tukey's test for multiple comparisons by site, for long (Table A4) and short cells (Table 285 A5) separately, showed lower values in the Mediterranean sites and higher values in the Atlantic site (Fig. 10).

The algal cells from Pontian Isl. had the lowest mean B/Ca in both seasons, being significantly different from both the samples from Morlaix and Aegadian Isl. (Table A4, A5; Fig. 10). Morlaix had the highest mean B/Ca in both long and short cells (Table A4, A5; Fig. 10). *L. coralliooides* from Aegadian Isl. had intermediate B/Ca mean value in long cells, differing significantly from both the Morlaix and Pontian Isl. samples (Table A4; Fig. 10). In short cells, the sample from Aegadian Isl. slightly 280 differed from the one in Morlaix (Table A5; Fig. 10).

Interestingly, the long cells of all samples had higher median B/Ca values compared to short cells (Fig. 10), although only in Morlaix, the differences between B/Ca measured in long and short cells were statistically significant ( $\chi^2=8.4899$ ,  $p<0.01$ ).



285 **Figure 10: Box plots of the statistical tests performed to evaluate the differences of B/Ca in the long and short cells of *L. coralliooides* collected in different sampling sites. The horizontal black lines indicate the median values. The black filled circles and the numbers inside the plot indicate the mean values.**

**Table 2: Element/Ca ratio measurements in *L. coralliooides*.**

Sampling site	B/Ca (μmol/mol)		Mg/Ca (mmol/mol)		Li/Ca (μmol/mol)		Sr/Ca (mmol/mol)		Mg/Li (mol/mmol)	
	mean	st. dev.	mean	st. dev.	mean	st. dev.	mean	st. dev.	mean	st. dev.
Aegadian Isl.	610.8	63.9	224.9	30.3	81.9	18.4	3.7	0.4	2.8	0.5
Elba	757.7	75.5	223.4	26.4	85.2	14.3	3.6	0.4	2.7	0.3
Pontian Isl.	462.8	49.2	216.1	21.9	79.6	14.6	3.3	0.3	2.8	0.4
Morlaix	726.9	102.8	239.5	41.2	113.1	32.9	4.1	0.5	2.2	0.4
total	<b>661.9</b>	<b>138.9</b>	<b>225.3</b>	<b>30.4</b>	<b>89.0</b>	<b>23.3</b>	<b>3.7</b>	<b>0.5</b>	<b>2.6</b>	<b>0.4</b>

### 3.4 Growth rates

290 In the sample from Aegadian Isl., the LA-ICP-MS transect was 1.31 mm long and 10 years of growth have been detected by coupling microscopical imaging and Mg/Ca peaks, resulting in 0.13 mm/yr of growth rate. In the Elba sample the laser transect was 1.15 mm long, crossing 8 years of growth, with a resulting growth rate of 0.14 mm/yr. The Pontian Isl. sample had 1.08 mm of transect including 11 years of growth, hence a growth rate of 0.10 mm/yr. Finally, the transect from Morlaix sample was 1.38 mm long, counting 11 years and resulting in 0.13 mm/yr of growth rate.

295 Growth rates did not show any linear relationship with Mg, Li and Sr/Ca, but they were positively correlated with the samples mean B/Ca values (Fig. 11).




Figure 11: Correlation plots of growth rates and seawater temperature with B/Ca in *L. coralliooides* samples analysed in this study. Spearman's coefficient  $r$ , the  $p$ -value and the line equation are given. Temperature variations ( $\Delta T$ ) correspond to the differences between the maximum and minimum temperature registered over 11 years of monthly reanalysis (ORAS5). The B/Ca means measured in long and short cells correspond respectively to the maximum and minimum temperature.

### 4 Discussion

In general, temperature variations affect many physiological processes involved in the biomineralization, and the rate of calcification, along with the preservation state of mineral structures, influences the content of trace elements in carbonates 305 (Lorens, 1981; Rimstidt et al., 1998; Gussone et al., 2005; Noireaux et al., 2015; Kaczmarek et al., 2016). Trace elements concentrations recorded from the four *L. coralliooides* branches analysed in this study were consistent with previously published values for other calcareous red algae (Chave, 1954; Hemming and Hanson, 1992; Hetzinger et al., 2011; Darrenougue et al., 2014). Particularly, the Mg/Ca ratios recorded in this study ranged from 172 to 311 mmol/mol, comparable to previous studies on rhodoliths of *Lithothamnion glaciale* Kjellman 1883 grown at 6-15 °C (148-326 mmol/mol) (Kamenos et al., 2008). The 310 B/Ca ratios in *L. coralliooides* from our results range from 356 to 954 μmol/mol, wider than the range measured by solution ICP-MS on bulk samples of *Neogoniolithon* sp. (352-670 μmol/mol) (Donald et al., 2017) and *C. compactum* (320-430 μmol/mol) (Anagnostou et al., 2019), both cultured with controlled  $p\text{CO}_2$  and a pH ranging from 7.2 to 8.2. The high resolution

given by laser ablation should be more effective in measuring the heterogeneity of B/Ca across the thallus, thus explaining the wider range of our data.


315 In the sample from Elba the growth bands were not clearly visible, preventing the analyses of trace elements in long and short cells separately (Fig. S1, S2, S3). To measure the growth rate in this sample, we considered the results from the spots where the light/dark bands resulting from the image analyses corresponded to obvious Mg/Ca peaks.

The Mg/Ca results evidenced a strong relationship with the seawater temperatures extracted from ORAS5 (Table 1; Fig. 12), as expected. *L. coralliooides* from Aegadian Isl. had slightly higher Mg/Ca values, followed by Elba and Pontian Isl. (Fig. 4).

320 This was consistent with local temperature values in the Mediterranean (Table 1), since Pontian Isl. registered the lowest mean value and the lowest  $\Delta T$ , while Aegadian Isl. showed the highest mean temperature and  $\Delta T$ . On the contrary, the sample from Morlaix, collected at 12 m depth, showed high Mg/Ca values in both long and short cells (Table A2; Fig. 5). The monthly mean temperatures had the highest variations during the year ( $\Delta T$  in Table 1), due to the shallow depth (12 m) and the geographical location. Temperature correlates to seasons, influencing primary production, respiration, and calcification in *L.*

325 *coralliooides* (Payri, 2000; Martin et al., 2006) as well as other calcareous red algae (Roberts et al., 2002). The high seasonality that characterized the sample from Morlaix, represented by the high  $\Delta T$  (Table 1), was probably responsible for the highest variation of Mg/Ca values and undoubtedly accounted for most of the differences with Mediterranean samples. For the first time, we confirmed here the reliability of the temperature proxies Li/Ca and Sr/Ca on a wild-grown coralline alga collected at different depths and locations. Li/Ca and Sr/Ca records were positively correlated with Mg/Ca in *L. coralliooides* (Fig. 3), 330 which, in turn, showed a strong relationship with seawater temperature. Moreover, both Li and Sr/Ca showed periodical oscillations in correspondence to long and short cells, related to seasonal temperature variations (Fig. 12). Therefore, Li/Ca and Sr/Ca could be regarded as temperature proxy in *L. coralliooides*, as for Mg/Ca. The coupling of the Mg/Ca ratio with Li/Ca and Sr/Ca represents a multi-proxy approach that minimize the possible species effect and can be considered a useful tool to gather information about past temperature for paleoclimate reconstructions (Halfar et al., 2011; Caragnano et al., 2014; 335 Williams et al., 2014; Fowell et al., 2016; Cuny-Guirriec et al., 2019).

The B/Ca ratio in coralline algae has been rarely measured and it is not clear how the environmental factors control its incorporation. The carbonate system primarily drives the changes in B incorporation (Hemming and Hanson, 1992; Yu and Elderfield, 2007). In benthic foraminifera, B/Ca increases with  $[CO_3^{2-}]$  (Yu and Elderfield, 2007), whereas there is no consensus on the effect of  $[CO_3^{2-}]$  on Mg/Ca and Sr/Ca (Rosenthal et al., 2006; Dueñas-Bohórquez et al., 2011). Experiments 340 with inorganic calcite showed a positive correlation between B/Ca and [DIC] (Uchikawa et al., 2015). Nevertheless, in culture experiments of the coralline algae *Neogoniolithon* sp. (Donald et al., 2017) and corals (Gagnon et al., 2021), [DIC] had a negative effect on B/Ca. The occurrence of low DIC concentrations in Morlaix (Table 1), compared to the Mediterranean sites, could therefore contribute to the high B/Ca. Nevertheless, based on the extracted pH and DIC data, significantly higher values of B/Ca should be expected in the sample from Pontian Isl., similarly to the other Mediterranean sites (Table 1).



**Figure 12: Elemental ratios in *L. corallioides* collected in Morlaix Bay (scale bar = 200  $\mu$ m). Mg, Li and Sr/Ca show cyclic variations mirroring the local seawater temperature. In the timeline, the coldest and the warmest months have been reported, which correspond to dark and light bands of growth. Elements/Ca in the missing bands (asterisks) have been calculated as the means of the values measured in warm or cold periods. Monthly means of temperature have been extracted by ORAS5 reanalysis.**

350 The estimated growth rate of *L. corallioides* was  $0.13 \pm 0.02$  mm/yr and it was supposed to decrease with increasing depth as a direct consequence of lower light availability (Halfar et al., 2011); indeed, the growth rate of the sample from Pontian Isl. was the lowest (0.10 mm/yr). As already suggested by previous studies on both synthetic and biogenic calcite, B incorporation is likely affected by growth rate (Gabitov et al., 2014; Mavromatis et al., 2015; Noireaux et al. 2015; Uchikawa et al., 2015; Kaczmarek et al., 2016). Indeed, in the cultured calcareous red alga *Neogoniolithon* sp. the B/Ca increases with increasing 355 growth rate (Donald et al., 2017). In Pontian Isl., the slow growth rate could have contributed to the low B/Ca values. The mean annual growth rate of the shallowest sample (Morlaix) was equal to the one in Aegadian Isl., at 40 m depth (0.13 mm/yr). In Morlaix, the alga probably significantly slowed down the growth in cold months, when the monthly mean seawater temperature was the lowest of all the sampling sites (Table 1). Nevertheless, its growth rate likely speeded up in the warm season due to the abundant light availability at shallow depth and the warming of seawater (Table 1), contributing to the 360 significantly higher B/Ca values in long cells (Fig. 10). According to this interpretation, the effect of the growth rate on B/Ca might be significant across depth and geographical regions (Fig. 11). In Morlaix, B/Ca showed higher values in comparison with Aegadian Isl. and Pontian Isl., and a positive correlation with temperature proxies (Mg/Ca, Li/Ca and Sr/Ca; Fig. 8). A positive correlation between B/Ca and Mg/Ca was already observed in planktonic foraminifera (Wara et al., 2003; Yu et al., 2007). Therefore, we reconstructed the elemental variations during the algal growth in the Morlaix sample at annual resolution, 365 in order to highlight the influence of temperature (Fig. 12). Contrary to Mg, Li, and Sr/Ca, the B/Ca did not mirror the seasonal temperature variations. In the sample from the Pontian Isl., the seasonal  $\Delta T$ , Mg/Ca and B/Ca values were the lowest among sites. In particular, B/Ca was significantly low (Fig. 9), differing more from the other samples than the results on Mg/Ca (Fig. 4). This suggests that in this sample the B incorporation should be influenced by factors other than those affecting Mg. In general, the poor correlation with seawater temperature (Fig. 11), and most of all the lack of distinct seasonal oscillations in 370 B/Ca across the algal thallus (Fig. 12), excludes the suitability of B/Ca as a temperature proxy and suggests a closer relationship with growth rate rather than temperature.

Knowing the biogeochemistry and the variation of the environmental parameters of seawater is crucial for a more comprehensive picture of the reliability of geochemical proxies, like the ones we investigated in this paper (Mg, Li, Sr/Ca and B/Ca). Boron incorporation in marine carbonates is still debated, rising questions about the boron isotopic fractionation, the 375 mechanisms of boron incorporation into marine carbonates, the so-called “vital effects” (i.e. the metabolic activities that can bias the isotopic signal), and the seawater isotopic composition. Moreover, the paucity of B/Ca measurements from coralline algae and, most of all, the complete absence of these data on specimens grown in nature makes it difficult to compare our B/Ca data with the literature. This observation takes stock of the significance of our results and emphasizes the importance of collecting more representative B/Ca data in coralline algae. Further studies on *L. corallioides* and other calcareous red algae

380 should be carried out to clarify the environmental factors influencing the B/Ca in these organisms, and to ensure the reliability  
of this proxy for paleoclimate reconstructions.

## 5 Conclusion

This paper presents the first measures of trace elements (Mg, Sr, Li and B) from the coralline alga *L. coralliooides* collected  
across the Mediterranean Sea and in the Atlantic Ocean, at different oceanographic settings and depths (12 m, 40 m, 45 m, and  
385 66 m depth), providing the first geochemical data of a wild-grown coralline algal species, sampled at different depths and  
geographical locations. LA-ICP-MS records of Mg/Ca, Sr/Ca and Li/Ca have shown a similar trend, primarily controlled by  
seawater temperatures in the algal habitat. On the contrary, Mg/Li did not provide a valuable temperature proxy in this species.  
In order to evaluate the control exerted by temperature over B incorporation, we also tested the correlation between B/Ca with  
390 Mg/Ca, Li/Ca and Sr/Ca. This led us to provide the first B/Ca data on wild grown coralline algae from across the photic zone  
in different Basins. The correlation between B/Ca and Mg/Ca in *L. coralliooides* was statistically significant only in the shallow  
Atlantic waters of Morlaix, where seasonality, hence the seasonal temperature variations, during the algal growth was the  
strongest among sites. Accordingly, B incorporation differences between long and short cells of *L. coralliooides* strongly depend  
on seasonality, being statistically significant just in Morlaix. Nevertheless, in contrast to Mg, Li and Sr/Ca, B/Ca oscillations  
across the algal growth showed a poor relationship with seasonal variations in seawater temperature. We found high B/Ca  
395 values in the Atlantic sample, where pH and DIC were the lowest. Carbon data did not explain the low B concentration in the  
Pontian Isl. sample (66 m depth), though, where pH and DIC were similar to the other Mediterranean sites. The estimation of  
growth rate, that was low in the Pontian Isl. sample (0.11 mm/yr) and got higher in the other Mediterranean samples and in  
Morlaix (~0.13 mm/yr), led us to conclude that B/Ca relates to growth rate rather than seawater temperature. B incorporation  
is therefore subject to the specific algal growth patterns and rates, whose knowledge is essential in order to assess the reliability  
400 of B/Ca in tracing seawater carbon variations.

## Appendix A

**Table A1: (a) Statistically non-significant results of tests performed to evaluate (a) the differences of Mg/Ca in *L. coralliooides* and  
(b) the differences of Mg/Ca in the short cells of *L. coralliooides* collected in different sampling sites. Test significance at  $\alpha = 0.05$ .**

(a) Kruskal-Wallis test (Mg/Ca)			
	Df	$\chi^2$	P
SITE	3	3.799	0.284

(b) One-way ANOVA test (Mg/Ca)					
Short cells					
	Df	Sum sq.	Mean sq.	F value	Pr(>F)

SITE	2	788.1	394.0	1.4647	0.2496
Residuals	26	6994.5	269.0		
Shapiro-Wilk normality test					P=0.6442
Bartlett's K-squared					P=0.5856

405

**Table A2:** Results of statistical tests performed to evaluate the differences of Mg/Ca in the long cells of *L. coralliooides* collected in different sampling sites. Statistically significant p-values are given in bold. ANOVA test significance at  $\alpha = 0.05$ ; Tukey's test significant at  $p \leq \alpha$ .

One-way ANOVA test (Mg/Ca)					
Long cells					
	Df	Sum sq.	Mean sq.	F value	Pr(>F)
SITE	2	10897.7	5448.9	16.413	<b>0.0001</b>
Residuals	20	6639.8	332.0		
Shapiro-Wilk normality test					P=0.1440
Bartlett's K-squared					P=0.5826
Tukey's test					
Multiple comparisons of means					
SITE	Mean difference		95% confidence interval		P. adjusted
	SITE		lower bound	upper bound	
Morlaix-Aegadian Isl.	38.32918	15.09816	61.56019		<b>0.00130</b>
Pontian Isl.-Aegadian Isl.	-10.84361	-35.48382	13.79661		0.51716
Pontian Isl.-Morlaix	-49.17278	-72.40380	-25.94177		<b>0.00009</b>

410

**Table A3:** Results of statistical tests performed to evaluate the differences of B/Ca in *L. coralliooides* collected in different sampling sites. Statistically significant p-values are given in bold. Kruskal-Wallis test significance at  $\alpha = 0.05$ ; Dunn's test significant at  $p \leq \alpha/2$ .

Kruskal-Wallis test (B/Ca)			
	Df	$\chi^2$	P
SITE	3	79.816	<b>&lt;2.2e-16</b>
Dunn's test			
Comparisons by SITE (Bonferroni)			
Z P. adjusted	Aegadian Isl.	Elba	Morlaix
Elba	-4.64580 <b>0.00000</b>		
Morlaix	-3.07755 <b>0.00630</b>	1.17249 0.72300	2.80564 8.38673
		6.15663	

Pontian Isl.	<b>0.01510</b>	<b>0.00000</b>	<b>0.00000</b>
--------------	----------------	----------------	----------------

Table A4: Results of statistical tests performed to evaluate the differences of B/Ca in the long cells of *L. coralliooides* collected in different sampling sites. Statistically significant p-values are given in bold. ANOVA test significance at  $\alpha = 0.05$ ; Tukey's test significant at  $p \leq \alpha$ .

One-way ANOVA test (B/Ca)					
Long cells					
	Df	Sum sq.	Mean sq.	F value	Pr(>F)
SITE	2	428364	214182	33.066	<b>0.0000</b>
Residuals	20	129546	6477		
Shapiro-Wilk normality test				P=0.5527	
Bartlett's K-squared				P=0.5470	
Tukey's test					
Multiple comparisons of means					
SITE	Mean difference	95% confidence interval		P. adjusted	
	SITE	lower bound	upper bound		
Morlaix-Aegadian Isl.	190.11730	87.50374	292.73094	<b>0.00040</b>	
Pontian Isl.-Aegadian Isl.	-135.42490	-244.26303	-26.58672	<b>0.01342</b>	
Pontian Isl.-Morlaix	-325.54220	-428.15581	-222.92862	<b>0.00000</b>	

Table A5: Results of statistical tests performed to evaluate the differences of B/Ca in the short cells of *L. coralliooides* collected in different sampling sites. Statistically significant p-values are given in bold. ANOVA test significance at  $\alpha = 0.05$ ; Tukey's test significant at  $p \leq \alpha$ .

One-way ANOVA test (B/Ca)					
Short cells					
	Df	Sum sq.	Mean sq.	F value	Pr(>F)
SITE	2	216232	108116	35.360	<b>0.0000</b>
Residuals	26	79497	3058		
Shapiro-Wilk normality test				P=0.1699	
Bartlett's K-squared				P=0.0576	
Tukey's test					
Multiple comparisons of means					
SITE	Mean difference	95% confidence interval		P. adjusted	
	SITE	lower bound	upper bound		
Morlaix-Aegadian Isl.	43.09640	-19.61932	105.81212	0.22146	
Pontian Isl.-Aegadian Isl.	-156.90170	-223.66771	-90.13574	<b>0.00001</b>	
Pontian Isl.-Morlaix	-199.99810	-260.58727	-139.40898	<b>0.00000</b>	

## 420 Data availability

Data resulting from this study are available at <https://doi.org/10.1594/PANGAEA.932201>.

Environmental data were provided by E.U. Copernicus Marine Service Information.

DIC data in the Mediterranean: [https://doi.org/10.25423/CMCC/MEDSEA\\_MULTIYEAR\\_BGC\\_006\\_008\\_MEDBFM3](https://doi.org/10.25423/CMCC/MEDSEA_MULTIYEAR_BGC_006_008_MEDBFM3)

DIC data in the Atlantic Ocean:

425 [https://resources.marine.copernicus.eu/?option=com\\_csw&view=details&product\\_id=IBI\\_MULTIYEAR\\_BGC\\_005\\_003](https://resources.marine.copernicus.eu/?option=com_csw&view=details&product_id=IBI_MULTIYEAR_BGC_005_003)  
pH data:

[https://resources.marine.copernicus.eu/?option=com\\_csw&view=details&product\\_id=GLOBAL\\_REANALYSIS\\_BIO\\_001\\_029](https://resources.marine.copernicus.eu/?option=com_csw&view=details&product_id=GLOBAL_REANALYSIS_BIO_001_029)

Temperature data: <https://icdc.cen.uni-hamburg.de/daten/reanalysis-ocean/easy-init-ocean/ecmwf-oras5.html>

## 430 Author contributions

DB, VB, and GP conceptualized the research question and study design. AL, DB, and VB conducted the experimental work; AM and GP the environmental data extraction. GP performed the data analysis and prepared the draft of the paper. All authors contributed to the editing and reviewing of the paper.

## Competing interests

435 The authors declare that they have no conflict of interest.

## Acknowledgements

This paper is a contribution to the Project MIUR-Dipartimenti di Eccellenza 2018-2022 DISAT-UNIMIB.

The Pontian Isl. sample has been collected in the framework of "Convenzione MATTM-CNR per i Programmi di Monitoraggio per la Direttiva sulla Strategia Marina (MSFD, Art. 11, Dir. 2008/56/CE)". Captain, crew and scientific staff of RV Minerva

440 Uno cruise STRATEGIA MARINA LIGURE-TIRRENO are acknowledged for their efficient and skilful cooperation at sea.  
Financial support for GP was provided by Italian MIUR as a PhD fellowship.

## References

Adey, W. H.: The genus *Clathromorphum* (Corallinaceae) in the Gulf of Maine, *Hydrobiologia*, 26, 539–573, doi:10.1007/BF00045545, 1965.

445 Adey, W. H., and McKibbin, D.: Studies on the maerl species *Phymatolithon calcareum* (Pallas) nov. comb. and *Lithothamnion coralloides* Crouan in the Ria de Vigo, Botanica marina, 13, 100–106, doi:10.1515/botm.1970.13.2.100, 1970.

Allen, K. A., Hönisch, B., Eggins, S. M., and Rosenthal, Y.: Environmental controls on B/Ca in calcite tests of the tropical planktic foraminifer species *Globigerinoides ruber* and *Globigerinoides sacculifer*, Earth. Planet. Sci. Lett., 351–352, 270–280, <https://doi.org/10.1016/j.epsl.2012.07.004>, 2012.

450 ~~Alverson, K., Bradley, R., Briffa, K., Cole, J., Hughes, M., Larocque, I., Pedersen, T., Thompson, L., and Tudhope, S.: A global paleoclimate observing system, Science, 293, 47–48, doi:10.1126/science.293.5527.47e, 2001.~~

Anagnostou, E., Williams, B., Westfield, I., Foster, G. L., and Ries, J. B.: Calibration of the pH- $\delta^{11}\text{B}$  and temperature-Mg/Li proxies in the long-lived high-latitude crustose coralline red alga *Clathromorphum compactum* via controlled laboratory experiments, Geochim. Cosmochim. Acta, 254, 142–155, doi:10.1016/j.gca.2019.03.015, 2019.

455 Andersson, A. J., Mackenzie, F. T., and Bates, N. R.: Life on the margin: implications on Mgcalcite, high latitude and cool-water marine calcifiers, Mar. Ecol. Prog. Ser., 373, 265–273, doi:10.3354/meps07639, 2008.

Andersson, A. J. and Mackenzie, F. T.: Technical comment on Kroeker et al. (2010) Meta-analysis reveals negative yet variable effects of ocean acidification on marine organisms, Ecol. Lett., 13, 1419–1434, doi:10.1111/j.1461-0248.2011.01646.x, 2011.

Barker, S., Cacho, I., Benway, H. and Tachikawa, K.: Planktonic foraminiferal Mg/Ca as a proxy for past oceanic temperatures: 460 a methodological overview and data compilation for the Last Glacial Maximum, Quat. Sci. Rev., 24, 821–834, doi:10.1016/j.quascirev.2004.07.016, 2005.

Basso, D.: Study of living calcareous algae by a paleontological approach: the non-geniculate Corallinaceae (Rhodophyta) of the soft bottoms of the Tyrrhenian Sea (western Mediterranean). The genera *Phymatolithon* Foslie and *Mesophyllum* Lemoine, Riv. Ital. Paleontol. S., 100, 575–596, doi:10.13130/2039-4942/8602, 1995a.

465 Basso, D.: Living calcareous algae by a paleontological approach: the genus *Lithothamnion* Heydrich nom. cons. from the soft bottoms of the Tyrrhenian Sea (Mediterranean), Riv. Ital. Paleontol. S., 101, 349–366, doi:10.13130/2039-4942/8592, 1995b.

Basso, D.: Carbonate production by calcareous red algae and global change, Geodiversitas, 34 (1), 13–33, doi:10.5252/g2012n1a2, 2012.

Basso, D., Babbini, L., Kaleb, S., Bracchi, V.A. and Falace, A.: Monitoring deep Mediterranean rhodolith beds, Aquat. 470 Conserv., 26, 549–561, doi:10.1002/aqc.2586, 2016.

Basso, D., Babbini, L., Ramos-Esplá, A.A. and Salomidi, M.: Mediterranean rhodolith beds, in: Rhodolith/maërl beds: A global perspective, edited by: Riosmena-Rodriguez, R., Wendy Nelson, W. and Aguirre, J., Springer, Cambridge, 281–298, doi:10.1007/978-3-319-29315-8\_11, 2017.

Basso, D. and Brusoni, F.: The molluscan assemblage of a transitional environment: the Mediterranean maerl from off the 475 Elba Island (Tuscan Archipelago, Tyrrhenian Sea), Boll. Malacol., 40 (1–4), 37–45, 2004.

Berner, R. A.: The role of magnesium in the crystal growth of calcite and aragonite from sea water, Geochim. Cosmochim. Acta, 39, 489–504, doi:10.1016/0016-7037(75)90102-7, 1975.

Brewer, P. G.: Ocean chemistry of the fossil fuel signal: The haline signature of “Business as Usual”, Geophys. Res. Lett., 24, 1367–1369, 1997.

480 Cabioch, J.: Contribution à l'étude morphologique, anatomique et systématique de deux Mélobésées: *Lithothamnium calcareum* (Pallas) Areschoug et *Lithothamnium coralliooides* Crouan, Bot. Mar., 9, 33–53, 1966.

Caldeira, K., Akai, M., Brewer, P., Chen, B., Haugan, P., Iwama, T., Johnston, P., Kheshgi, H., Li, Q., Ohsumi, T., Poertner, H., Sabine, C., Shirayama, Y., and Thomson, J.: Ocean Storage, in: Carbon Dioxide Capture and Storage: A Special Report of IPCC Working Group III, Cambridge University Press, Cambridge UK, 2005.

485 Callendar, G. S.: The artificial production of carbon dioxide and its influence on temperature. Quarterly Journal Roy. Met. Soc., 64, 223–240, 1938.

Caragnano, A., Basso, D., Jacob, D. E., Storz, D., Rodondi, G., Benzoni, F. and Dutrieux, E.: Coralline red alga *Lithophyllum kotschyanum* f. affine as proxy of climate variability in the Yemen coast, Gulf of Aden (NW Indian Ocean), Geochim. Cosmochim. Acta, 124, 1–17, doi:10.1016/j.gca.2013.09.021, 2014.

490 Caragnano, A., Basso, D., Storz, D., Jacob, D. E., Ragazzola, F., Benzoni, F. and Dutrieux, E.: Elemental variability in the coralline alga *Lithophyllum yemenense* as an archive of past climate in the Gulf of Aden (NW Indian Ocean), J. Phycol., 53, 381–395, doi:10.1111/jpy.12509, 2017.

Carro, B., Lopez, L., Peña, V., Bárbara, I. and Barreiro, R.: DNA barcoding allows the accurate assessment of European maelr diversity: a proof-of-concept study, Phytotaxa, 190, 176–189, doi:10.11646/phytotaxa.190.1.12, 2014.

495 Chave, K.E.: Aspects of the biogeochemistry of magnesium 1. Calcareous marine organisms, J. Geol., 62 (3), 266–283., 1954.

Ciais, P., Sabine, C., Bala, G., Bopp, L., Brovkin, V., Canadell, J., Chhabra, A., DeFries, R., Galloway, J., Heimann, M., Jones, C., Le Quéré, C., Myneni, R.B., Piao, S. and Thornton, P.: Carbon and Other Biogeochemical Cycles, in: Climate Change 2013: The Physical Science Basis, Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, edited by: Stocker, T. F., Qin, D., Plattner, G. K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., Cambridge University Press, Cambridge, 465–570, 2013.

500 Cornwall, C. E., Comeau, S. and Mcculloch, M. T.: Coralline algae elevate pH at the site of calcification under ocean acidification, Glob. Change Biol., 23 (10), 1–12, doi:10.1111/gcb.13673, 2017.

Corrège, T.: Sea surface temperature and salinity reconstruction from coral geochemical tracers, Palaeogeogr., Palaeoclimatol., Palaeoecol., 232, 408–428, doi:10.1016/j.palaeo.2005.10.014, 2006.

505 Cuny-Guiriec, K., Douville, E., Reynaud, S., Allemand, D., Bordier, L., Canesi, M., Mazzoli, C., Taviani, M., Canese, S., McCulloch, M., Trotter, J., Rico-Esenaro, S. D., Sanchez-Cabeza, J.-A., Ruiz-Fernàndez, A. C., Carricart-Ganivet, J. P., Scott, P. M., Sadekov, A. and Montagna, P.: Coral Li/Mg thermometry: Caveats and constraints, Chem. Geol., 523, 162–178, doi:10.1016/j.chemgeo.2019.03.038, 2019.

Cusack, M., Kamenos, N. A., Rollion Bard, C. and Tricot, G.: Red coralline algae assessed as marine pH proxies using <sup>11</sup>B MAS NMR, Sci. Rep., 5, 8175, doi:10.1038/srep08175, 2015.

Darrenougue, N., De Deckker, P., Eggins, S. and Payri, C.: Sea-surface temperature reconstruction from trace elements variations of tropical coralline red algae, *Quat. Sci. Rev.*, 93, 34–46, doi:10.1016/j.quascirev.2014.03.005, 2014.

DeCarlo, T. M., Holcomb, M., and McCulloch, M. T.: Reviews and syntheses: Revisiting the boron systematics of aragonite and their application to coral calcification, *Biogeosciences*, 15, 2819–2834, doi:10.5194/bg-15-2819-2018, 2018.

515 Dickson, A.G.: Thermodynamics of the dissociation of boric acid in synthetic seawater from 273.15 to 318.15 K, *Deep-Sea Res.*, 37, 755–766, doi:10.1016/0198-0149(90)90004-F, 1990.

Dickson, A.G., Sabine, C.L. and Christian, J.R.: Guide to Best Practices for Ocean CO<sub>2</sub> measurements, in: PICES Special Publication 3, North Pacific Marine Science Organization (PICES), Sidney, 2007.

D’Olivo, J. P., Sinclair, D. J., Rankenburg, K. and McCulloch, M. T.: A universal multi-trace element calibration for 520 reconstructing sea surface temperatures from long-lived *Porites* corals: removing ‘vital-effects’, *Geochim. Cosmochim. Acta* 239, 109–135, doi:10.1016/j.gca.2018.07.035, 2018.

Donald, H. K., Ries, J. B., Stewart, J. A., Fowell, S. E. and Foster, G. L.: Boron isotope sensitivity to seawater pH change in a species of *Neogoniolithon* coralline red alga, *Geochim. Cosmochim. Acta*, 217, 240–253, doi:10.1016/j.gca.2017.08.021, 2017.

525 Douville, É., Paterne, M., Cabioch, G., Louvat, P., Gaillardet, J., Juillet-Leclerc, A. and Ayliffe, L.: Abrupt sea surface pH change at the end of the Younger Dryas in the central sub-equatorial Pacific inferred from boron isotope abundance in corals (*Porites*), *Biogeosciences*, 7(8), 2445–2459, doi:10.5194/bg-7-2445-2010, 2010.

Dueñas-Bohrquez, A., Raitzsch, M., Nooijer, L. J. and Reichart, G.-J.: Independent impacts of calcium and carbonate ion concentration on Mg and Sr incorporation in cultured benthic foraminifera, *Mar. Micropaleontol.*, 81 (3–4), 122–130, 530 doi:10.1016/j.marmicro.2011.08.002, 2011.

Fairhall, A. W.: Accumulation of fossil CO<sub>2</sub> in the atmosphere and the sea, *Nature*, 245, 20–23 (Erratum *Nature* v. 246, 106), 1973.

Feely, R. A., Doney, S. C. and Cooley, S. R.: Ocean acidification: present conditions and future changes in a high CO<sub>2</sub> World, *Oceanography*, 22(4), 36–47, doi:10.5670/oceanog.2009.95, 2009.

535 Fietzke, J., Heinemann, A., Taubner, I., Böhm, F., Erez, J., Eisenhauer, A.: Boron isotope ratio determination in carbonates via LA-MC-ICP-MS using soda-lime glass standards as reference material, *J. Anal. At. Spectrom.*, 12, doi:10.1039/C0JA00036A, 2010.

Fietzke, J., Ragazzola, F., Halfar, J., Dietze, H., Foster, L. C., Hansteen, T. H., Eisenhauer, A. and Steneck, R. S.: Century-scale trends and seasonality in pH and temperature for shallow zones of the Bering Sea, *Proc. Natl. Acad. Sci.*, 112, 2960–540 2965, doi:10.1073/pnas.1419216112, 2015.

Foster, G. L.: Seawater pH, pCO<sub>2</sub> and [CO<sub>3</sub><sup>2-</sup>] variations in the Caribbean Sea over the last 130 kyr: a boron isotope and B/Ca study of planktic foraminifera, *Earth and Planet. Sci. Lett.*, 271, 254–266, doi:10.1016/j.epsl.2008.04.015, 2008.

Foster, G. L., Pogge von Strandmann, P. A. E. and Rae, J. W. B.: Boron and magnesium isotopic composition of seawater, *Geochem. Geophys.,* 11 (8), Q08015, doi:10.1029/2010GC003201, 2010.

545 [Foster, G. L. and Rae, J. W. B.: Reconstructing Ocean pH with boron isotopes in foraminifera, Annu. Rev. Earth Planet. Sci., 44 \(1\), 207–237, 2016.](#)  
Foster, M.S.: Rhodoliths: between rocks and soft places, *J. Phycology*, 37, 659–667, doi:10.1046/j.1529-8817.2001.00195.x, 2001.

550 Fowell, S.E., Sandford, K., Stewart, J.A., Castillo, K.D., Ries, J.B. and Foster, G.L.: Intrareef variations in Li/Mg and Sr/Ca sea surface temperature proxies in the Caribbean reef-building coral *Siderastrea sidereal*, *Paleoceanography*, 31, 1315–1329, doi:10.1002/2016PA002968, 2016.

Frantz, B.R., Foster, M.S. and Riosmena-Rodríguez, R.: *Clathromorphum nereostratum* (Corallinales, Rhodophyta): The oldest alga?, *J. Phycology*, 41, 770–773, doi:10.1111/j.1529-8817.2005.00107.x, 2005.

555 Freiwald, A. and Henrich, R.: Reefal coralline algal build-ups within the Arctic Circle: morphology and sedimentary dynamics under extreme environmental seasonality, *Sedimentology*, 41, 963–984, doi:10.1111/j.1365-3091.1994.tb01435.x, 1994.

Gabitov, R. I., Rollion-Bard, C., Tripati, A. and Sadekov, A.: In situ study of boron partitioning between calcite and fluid at different crystal growth rates, *Geochim. Cosmochim. Acta*, 137, 81–92, doi:10.1016/j.gca.2014.04.014, 2014.

560 Gagnon, A. C., Gothmann, A. M., Branson, O., Rae, J. W. B. and Stewart, J. A.: Controls on boron isotopes in a cold-water coral and the cost of resilience to ocean acidification, *Earth Planet. Sci. Lett.*, 554, 116662, doi:10.1016/j.epsl.2020.116662, 2021.

[Gattuso, J.-P., Frankignoulle, M., and Smith, S. V.: Measurement of community metabolism and significance in the coral reef CO<sub>2</sub> source-sink debate, Proc. Natl. Acad. Sci. USA, 96, 13017–13022, 1999.](#)

565 Gussone, N., Böhm, F., Eisenhauer, A., Dietzel, M., Heuser, A., Teichert, B. M. A., Reitner, J., Wörheide, G. and Dullo, W.-C.: Calcium isotope fractionation in calcite and aragonite, *Geochim. Cosmochim. Acta*, 69, 4485–4494, doi:10.1016/j.gca.2005.06.003, 2005.

Halfar, J., Steneck, R. S., Joachimski, M., Kronz, A. and Wanamaker, A. D. Jr.: Coralline red algae as high-resolution climate recorders, *Geology*, 36, 463–466, doi:10.1130/G24635A.1, 2008.

570 Halfar, J., Williams, B., Hetzinger, S., Steneck, R.S., Lebednik, P., Winsborough, C., Omar, A., Chan, P. and Wanamaker, A.D.: 225 years of Bering Sea climate and ecosystem dynamics revealed by coralline algal growth-increment widths, *Geology*, 39 (6), 579–582, doi:10.1130/g31996.1, 2011.

Hall-Spencer, J.M., Kelly, J. and Maggs, C.A.: Background document for maërl beds, in: OSPAR Commission Biodiversity Series, OSPAR Commission, London, 2010.

[Hansen, J., Ruedy, R., Sato, M. and Lo, K.: Global surface temperature change, Rev. Geophys., 48, RG4004, doi:10.1029/2010RG000345, 2010.](#)

575 Hemming, N. G. and Hanson, G. N.: Boron isotopic composition and concentration in modern marine carbonates, *Geochim. Cosmochim. Acta*, 56 (1), 537–543, doi:10.1016/0016-7037(92)90151-8, 1992.

Hemming, N. G. and Hönisch, B.: Boron isotopes in marine carbonate sediments and the pH of the ocean, in: *Proxies in Late Cenozoic Paleoceanography*, edited by: Hillaire-Marcel, C. and de Vernal, A., Elsevier, 717–734, 2007.

Henehan, M. J., Rae, J. W. B., Foster, G. L., Erez, J., Prentice, K. C., Kucera, M., Bostock, H. C., Martinez-Botí, M. A.,  
580 Milton, J. A., Wilson, P. A., Marshall, B. J. and Elliott, T.: Calibration of the boron isotope proxy in the planktonic foraminifera  
*Globigerinoides ruber* for use in palaeo-CO<sub>2</sub> reconstruction, *Earth Planet. Sci. Lett.*, 364, 111–122,  
doi:10.1016/j.epsl.2012.12.029, 2013.

Henehan, M.J., Foster, G.L., Rae, J.W.B., Prentice, K.C., Erez, J., Bostock, H.C., Mar-shall, B.J. and Wilson, P.A.: Evaluating  
the utility of B/Ca ratios in planktic foraminifera as a proxy for the carbonate system: a case study of *Globigerinoides ruber*:  
585 investigating controls on *G. ruber* B/Ca, *Geochem. Geophys.*, 16, 1052–1069, doi:10.1002/2014GC005514, 2015.

Hetzinger, S., Halfar, J., Kronz, A., Steneck, R., Adey, W. H., Philipp, A. L. and Schöne, B.: High-resolution Mg/Ca ratios in  
a coralline red alga as a proxy for Bering Sea temperature variations from 1902–1967, *Palaois*, 24, 406–412,  
doi:10.2110/palo.2008.p08-116r, 2009.

Hetzinger, S., Halfar, J., Zack, T., Gamboa, G., Jacob, D.E., Kunz, B.E., Kronz, A., Adey, W., Lebednik, P.A. and Steneck,  
590 R.S.: High-resolution analysis of trace elements in crustose coralline algae from the North Atlantic and North Pacific by laser  
ablation ICP-MS, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 302 (1-2), 81–94, doi:10.1016/j.palaeo.2010.06.004, 2011.

Hoegh-Guldberg, O., Mumby, P. J., Hooten, A. J., Steneck, R. S., Greenfield, P., Gomez, E., Harvell, C. D., Sale, P. F.,  
Edwards, A. J., Caldeira, K., Knowlton, N., Eakin, C. M., Iglesias-Prieto, R., Muthiga, N., Bradbury, R. H., Dubi, A. and  
Hatzios, M. E.: Coral reefs under rapid climate change and ocean acidification, *Science*, 318 (5857), 1737–1742,  
595 doi:10.1126/science.1152509, 2007.

Holcomb, M., DeCarlo, T. M., Gaetani, G. A., and McCulloch, M.: Factors affecting B/Ca ratios in synthetic aragonite, *Chem. Geol.*, 437, 67–76, doi:10.1016/j.chemgeo.2016.05.007, 2016.

Hönisch, B. and Hemming, N. G.: Surface ocean pH response to variations in pCO<sub>2</sub> through two full glacial cycles, *Earth Planet. Sci. Lett.*, 236, 305–314, doi:10.1016/j.epsl.2005.04.027, 2005.

600 Hönisch, B., Ridgwell, A., Schmidt, D. N., Thomas, E., Gibbs, S. J., Sluijs, A., Zeebe, R., Kump, L., Martindale, R. C., Greene,  
S. E. et al.: The geological record of ocean acidification, *Science*, 335 (6072), 1058–1063, doi:10.1126/science.1208277, 2012.

Irvine, L. M. and Chamberlain, Y. M. (Eds.): Volume 1 Rhodophyta Part 2B. Corallinales, Hildenbrandiales, Natural History  
Museum, London, 1994.

Jochum, K. P., Scholz, D., Stoll, B., Weis, U., Wilson, S. A., Yang, Q., Schwalb, A., Börner, N., Jacob, D. E., Andreae, M.  
605 O.: Accurate trace element analysis of speleothems and biogenic calcium carbonates by LA-ICP-MS, *Chem. Geol.*, 318-319,  
31-44, doi:10.1016/j.chemgeo.2012.05.009, 2012.

Kaczmarek, K., Nehrke, G., Misra, S., Bijma, J. and Elderfield, H.: Investigating the effects of growth rate and temperature on  
the B/Ca ratio and  $\delta^{11}\text{B}$  during inorganic calcite formation, *Chem. Geol.*, 421, 81–92, doi:10.1016/j.chemgeo.2015.12.002,  
2016.

610 Kamenos, N. A., Cusack, M. and Moore, P. G.: Coralline algae are global paleothermometers with bi-weekly resolution,  
*Geochim. Cosmochim. Acta*, 72, 771–779, doi:10.1016/j.gca.2007.11.019, 2008.

Kamenos, N. A., Cusack, M., Huthwelker, T., Lagarde, P. and Scheibling, R. E.: Mg-lattice associations in red coralline algae, *Geochim. Cosmochim. Acta*, 73, 1901–1907, doi:10.1016/j.gca.2009.01.010, 2009.

Keul, N., Langer, G., Thoms, S., de Nooijer, L. J., Reichart, G. J. and Bijma, J.: Exploring foraminiferal Sr/Ca as a new  
615 carbonate system proxy, *Geochim. Cosmochim. Acta*, 202, 374–386, doi:10.1016/j.gca.2016.11.022, 2017.

Klochko, K., Kaufman, A. J., Yao, W., Byrne, R. H. and Tossell, J. A.: Experimental measurement of boron isotope fractionation in seawater, *Earth Planet. Sci. Lett.*, 248, 276–285, doi:10.1016/j.epsl.2006.05.034, 2006.

Klochko, K., Cody, G. D., Tossell, J. A., Dera, P. and Kaufman, A. J.: Re-evaluating boron speciation in biogenic calcite and aragonite using  $^{11}\text{B}$  MAS NMR, *Geochim. Cosmochim. Acta*, 73, 1890–1900, doi:10.1016/j.gca.2009.01.002, 2009.

620 ~~Le Quéré, C., Raupach, M. R., Canadell, J. G., Marland, G., et al.: Trends in the sources and sinks of carbon dioxide, *Nat. Geosci.*, 2, 831–836, doi:10.1038/ngeo689, 2009.~~

Lorens, R. B.: Sr, Cd, Mn and Co distribution coefficients in calcite as a function of calcite precipitation rate, *Geochim. Cosmochim. Acta*, 45, 553–561, doi:10.1016/0016-7037(81)90188-5, 1981.

Martin, S., Castets, M.-D. and Clavier, J.: Primary production, respiration and calcification of the temperate free-living  
625 coralline alga *Lithothamnion coralliooides*, *Aquat. Bot.*, 85, 121–128, doi:10.1016/j.aquabot.2006.02.005, 2006.

Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, Zhou, B. et al.: IPCC: Climate Change 2021: The Physical Science Basis, in: Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge University Press, Cambridge UK, 2021.

Mavromatis, V., Montouillout, V., Noireaux, J., Gaillardet, J. and Schott, J.: Characterization of boron incorporation and  
630 speciation in calcite and aragonite from co-precipitation experiments under controlled pH, temperature and precipitation rate, *Geochim. Cosmochim. Acta*, 150, 299–313, doi:10.1016/j.gca.2014.10.024, 2015.

Melbourne, L.A., Hernández-Kantún, J., Russell, S. and Brodie, J.: There is more to maerl than meets the eye: DNA barcoding reveals a new species in Britain, *Lithothamnion erinaceum* sp. nov. (Hapalidiales, Rhodophyta), *Eur. J. Phycol.*, 52(2), 166–178, doi:10.1080/09670262.2016.1269953, 2017.

635 Moberly, R.: Composition of magnesian calcites of algae and pelecypods by electron microprobe analysis, *Sedimentology*, 11: 61–82, doi:10.1111/j.1365-3091.1968.tb00841.x, 1968.

Moberly, R.: Microprobe study of diagenesis in calcareous algae, *Sedimentology*, 14: 113–123, doi: 10.1111/j.1365-3091.1970.tb00185.x, 1970.

640 Montagna, P. and Douville, E.: Geochemical proxies in marine biogenic carbonates: New developments and applications to global change, *Chem. Geol.*, 533: 119411, doi:10.1016/j.chemgeo.2019.119411, 2020.

Morse, J. W., Andersson, A. J. and Mackenzie, F. T.: Initial responses of carbonate-rich shelf sediments to rising atmospheric  $\text{pCO}_2$  and “ocean acidification”: role of high Mg-calcites, *Geochim. Cosmochim. Acta*, 70, 5814–5830, doi:10.1016/j.gca.2006.08.017, 2006.

Noireaux, J., Mavromatis, V., Gaillardet, J., Schott, J., Montouillout, V., Louvat, P., Rollion-Bard, C. and Neuville, D. R.:  
645 Crystallographic control on the boron isotope paleo-pH proxy, *Earth Planet. Sci. Lett.*, 430, 398–407, doi:10.1016/j.epsl.2015.07.063, 2015.

Paris, G., Bartolini, A., Donnadieu, Y., Beaumont, V. and Gaillardet, J.: Investigating boron isotopes in a middle Jurassic micritic sequence: primary vs. diagenetic signal, *Chem. Geol.*, 275, 117–126, doi:10.1016/j.chemgeo.2010.03.013, 2010.

Payri, C. E.: Production primaire et calcification des algues benthiques en milieu corallien, *Oceanis*, 26, 427–463, 2000.

650 Potin, P., Floc'h, J.-Y., Augris, C. and Cabioch, J.: Annual growth rate of the calcareous red algae *Lithothamnion coralliooides* (Corallinales, Rhodophyta) in the Bay of Brest, France, *Hydrobiologia*, 204–205, 263–267, 1990.

Rae, J. B. R., Foster, G. L., Schmidt, D. N. and Elliott, T.: Boron isotopes and B/Ca in benthic foraminifera: proxies for the deep ocean carbonate system, *Earth Planet. Sci. Lett.*, 302, 403–413, doi:10.1016/j.epsl.2010.12.034, 2011.

Ragazzola, F., Foster, L. C., Form, A., Anderson, P. S. L., Hansteen, T. H. and Fietzke, J.: Ocean acidification weakens the 655 structural integrity of coralline algae, *Glob. Change Biol.*, 18, 2804–2812, doi:10.1111/j.1365-2486.2012.02756.x, 2012.

Ragazzola, F., Foster, L. C., Jones, C. J., Scott, T. B., Fietzke, J., Kilburn, M. R. and Schmidt, D. N.: Impact of high CO<sub>2</sub> on the geochemistry of the coralline algae *Lithothamnion glaciale*, *Sci. Rep.*, 6, 20572, doi:10.1038/srep20572, 2016.

Ragazzola, F., Caragnano, A., Basso, D., Schmidt, D. N. and Fietzke, J.: Establishing temperate crustose Early Holocene 660 coralline algae as archived for paleoenvironmental reconstructions of the shallow water habitats of the Mediterranean Sea, *Paleontology*, 63, 155–170, doi:10.1111/pala.12447, 2020.

Ries, J. B.: Mg fractionation in crustose coralline algae: geochemical, biological and sedimentological implications of secular variation in Mg/Ca ratio of seawater, *Geochim. Cosmochim. Acta*, 70, 891–900, doi:10.1016/j.gca.2005.10.025, 2006.

Ries, J. B., Ghazaleh, M. N., Connolly, B., Westfield, I. and Castillo, K. D.: Impacts of seawater saturation state ( $\Omega_A = 0.4$ – 665 4.6) and temperature (10, 25 °C) on the dissolution kinetics of whole-shell biogenic carbonates, *Geochim. Cosmochim. Acta*, 192, 318–337, doi:10.1016/j.gca.2016.07.001, 2016.

Rimstidt, J. D., Balog, A. and Webb, J.: Distribution of trace elements between carbonate minerals and aqueous solutions, *Geochim. Cosmochim. Acta*, 62 (11), 1851–1863, doi:10.1016/S0016-7037(98)00125-2, 1998.

Roberts, R. D., Kühl, M., Glud, R. N. and Rysgaard, S.: Primary production of crustose coralline algae in a high Arctic Fjord, *J. Phycol.*, 38 (2), 273–283, doi:10.1046/j.1529-8817.2002.01104.x, 2002.

670 ~~Rollion Bard, C., Blamart, D., Trebose, J., Tricot, G., Mussi, A. and Cuif, J. P.: Boron isotopes as pH proxy: A new look at boron speciation in deep sea corals using <sup>11</sup>B MAS NMR and EELS, *Geochim. Cosmochim. Acta*, 75, 1003–1012, doi:10.1016/j.gca.2010.11.023, 2011.~~

Rosenthal, Y., Lear, C. H., Oppo, D. W. and Braddock, K. L.: Temperature and carbonate ion effects on Mg/Ca and Sr/Ca ratios in benthic foraminifera: Aragonitic species *Hoeglundina elegans*, *Paleoceanography*, 21, PA 1007, 675 doi:10.1029/2005PA001158, 2006.

Savini, A., Basso, D., Bracchi, V. A., Corselli, C. and Pennetta, M.: Maerl-bed mapping and carbonate quantification on submerged terraces offshore the Cilento peninsula (Tyrrhenian Sea, Italy), *Geodiversitas*, 34 (1), 77–98, doi:10.5252/g2012n1a5, 2012.

Schöne, B.R., Fiebig, J., Pfeiffer, M., Gleß, R., Hickson, J., L. A. Johnson, A., Dreyer, W. and Oschmann, W.: Climate records from a bivalved Methuselah (*Arctica islandica*, Mollusca; Iceland), *Palaeogeogr., Palaeoclimatol., Palaeoecol.*, 228, 130–148, doi:10.1016/j.palaeo.2005.03.049, 2005.

Teruzzi, A., Di Biagio, V., Feudale, L., Bolzon, G., Lazzari, P., Salon, S., Di Biagio, V., Coidessa, G. and Cossarini, G.: Mediterranean Sea Biogeochemical Reanalysis (CMEMS MED-Biogeochemistry, MedBFM3 system) (Version 1) [Data set], Copernicus Monitoring Environment Marine Service (CMEMS), 2021.

Uchikawa, J., Penman, D. E., Zachos, J. C. and Zeebe, R. E.: Experimental evidence for kinetic effects on B/Ca in synthetic calcite: Implications for potential  $B(OH)_4^-$  and  $B(OH)_3$  incorporation, *Geochim. Cosmochim. Acta*, 150, 171–191, doi:10.1016/j.gca.2014.11.022, 2015.

Vengosh, A., Kolodny, Y., Starinsky, A., Chivas, A.R. and McCulloch, M.T.: Coprecipitation and isotopic fractionation of boron in modern biogenic carbonates, *Geochim. Cosmochim. Acta*, 55, 2901–2910, doi:10.1016/0016-7037(91)90455-E, 1991.

Wara, M. W., Delaney, M. L., Bullen, T. D. and Ravelo, A. C.: Possible roles of pH, temperature, and partial dissolution in determining boron concentration and isotopic composition in planktonic foraminifera, *Paleoceanography*, 18 (4), 1100, doi:10.1029/2002PA000797, 2003.

Williams, B., Halfar, J., Delong, K. L., Hetzinger, S., Steneck, R. S. and Jacob, D. E.: Multi-specimen and multi-site calibration of Aleutian coralline algal Mg/Ca to sea surface temperature, *Geochim. Cosmochim. Acta*, 139, 190–204, doi:10.1016/j.gca.2014.04.006, 2014.

Wilson, S., Blake, C., Berges, J. A. and Maggs, C. A.: Environmental tolerances of free-living coralline algae (maerl): implications for European marine conservation, *Biol. Conserv.*, 120, 279–289, doi:10.1016/j.biocon.2004.03.001, 2004.

Yu, J. M. and Elderfield, H.: Benthic foraminiferal B/Ca ratios reflect deep water carbonate saturation state, *Earth Planet. Sci. Lett.*, 258, 73–86, doi:10.1016/j.epsl.2007.03.025, 2007.

Yu, J. M., Elderfield, H. and Hönnisch, B.: B/Ca in planktonic foraminifera as a proxy for surface water pH, *Paleoceanography*, 22, 2202, doi:10.1029/2006PA001347, 2007.

Yu, J., Day, J., Greaves, M. and Elderfield, H.: Determination of multiple element/calcium ratios in foraminiferal calcite by quadrupole ICP-MS, *Geochem. Geophys.,* 6, Q08P01, doi:10.1029/2005GC000964, 2015.

Zeebe, R. E. and Wolf-Gladrow, D. (Eds.):  $CO_2$  in seawater: equilibrium, kinetics, isotopes, Elsevier Oceanography Book Series, 65, Amsterdam, 2001.

Zinke, J., D'Olivo, J. P., Gey, C. J., McCulloch, M. T., Bruggemann, J. H., Lough, J. M. and Guillaume, M. M. M.: Multi-trace-element sea surface temperature coral re-construction for the southern Mozambique Channel reveals teleconnections with the tropical Atlantic, *Biogeosciences*, 16, 695–712, doi:10.5194/bg-16-695-2019, 2019.

710 Zuo, H., Balmaseda, M. A., Tietsche, S., Mogensen, K. and Mayer, M.: The ECMWF operational ensemble reanalysis–analysis  
system for ocean and sea ice: a description of the system and assessment, *Ocean Sci.*, 15, 779–808, doi:10.5194/os-15-779-  
2019, 2019.