



## Contrasting patterns of carbon cycling and DOM processing in two phytoplankton-bacteria communities

Samu Elovaara<sup>1,2</sup>, Eeva Eronen-Rasimus<sup>2</sup>, Eero Asmala<sup>1</sup>, Tobias Tamelander<sup>1</sup>, Hermanni Kaartokallio<sup>2</sup>


<sup>1</sup>Tvärminne Zoological Station, University of Helsinki, Helsinki, 00014, Finland

5 <sup>2</sup>Finnish Environment Institute, Helsinki, 00790, Finland

Correspondence to: Samu Elovaara ([samu.elovaara@syke.fi](mailto:samu.elovaara@syke.fi))



**Abstract.** Microbial consumption of phytoplankton-derived organic carbon in the pelagic food web is an important component of the global C cycle. We studied C cycling in two phytoplankton-bacteria systems (non-axenic cultures of a dinoflagellate *Apocalathium malmogiense* and a cryptophyte *Rhodomonas marina*) in two experiments. In the first experiment we grew phytoplankton and bacteria in nutrient replete conditions and followed C processing at early exponential growth phase and at two later phases. Primary production and total community respiration were up to 4 and 7 times higher, respectively, in the *A. malmogiense* treatments. Based on the optical signals, accumulating dissolved organic C (DOC) was degraded more in the *R. marina* treatments and the rate of bacterial production to primary production was higher. Thus, the flow of C from phytoplankton to bacteria was relatively higher in *R. marina* treatments than in *A. malmogiense* treatments which was further supported by faster  $^{14}\text{C}$  transfer from phytoplankton to bacterial biomass. In the second experiment we investigated consumption of the phytoplankton-derived DOC by bacteria. DOC consumption and transformation, bacterial production and bacterial respiration were all higher in *R. marina* treatments. In both experiments *A. malmogiense* supported a bacterial community predominated by bacteria specialized in the utilization of less labile DOC (class Bacteroidia) whereas *R. marina* supported a community predominated by copiotrophic Alpha- and Gammaproteobacteria. Our findings suggest that large dinoflagellates cycle relatively more C between phytoplankton biomass and the inorganic C pool whereas small cryptophytes direct relatively more C to the microbial loop.



Created with Biorender.com

**Graphical abstract: Differences in C cycling between *A. malmogiense* and *R. marina*.**



## 1 Introduction

Dissolved organic carbon (DOC) forms the largest aquatic organic C pool (~660 Pg C, (Hansell et al., 2009)), comparable in magnitude to atmospheric CO<sub>2</sub> (~780 Pg C, (Emerson and Hedges, 2008)). Phytoplankton are the most important source of autochthonous DOC in marine systems (Thornton, 2014). DOC is the main energy source for pelagic heterotrophic bacteria (Ducklow and Carlson, 1992), which quickly consume the most bioavailable organic molecules. As a result, bulk of the marine DOC pool consists of refractory DOC (Jiao et al., 2010). Depending on the composition of DOC and surrounding conditions, DOC may accumulate in the water column (Hedges, 1992; Jiao et al., 2010; Mari et al., 2017), aggregate and sink (Engel et al., 2004), or be consumed (Azam et al., 1983; Kujawinski, 2011). The rates of these processes determine the prevalent fate of DOC and thus greatly determine total C cycling pathways.

35

The proportion of a phytoplankton species in a mixed community may affect the release of dissolved organic matter (DOM) within that community. The composition of phytoplankton-derived DOM is generally affected by growth phase (Urbani et al., 2005), environmental conditions (e.g. increased C:nutrient ratios of released DOM under nutrient limitation (Saad et al., 2016)) and physiological state of the phytoplankton community (e.g. release of specific compounds as a result of cell death (Orellana et al., 2013)). Different phytoplankton species produce different kinds of DOM (Becker et al., 2014; Mühlenbruch et al., 2018; Romera-Castillo et al., 2010; Sarmiento et al., 2013), which shape the composition of bacterial community depending on the composition of the released DOM (Romera-Castillo et al., 2011; Sarmiento et al., 2013; Sarmiento and Gasol, 2012; Teeling et al., 2012). Bacteria remineralize and transform organic matter and as a result produce different types of DOC (Kawasaki and Benner, 2006), nutrients (Amin et al., 2009; Christie-Oleza et al., 2017) and other substances (Croft et al., 2005) that become available to phytoplankton. Interactions among phytoplankton and bacteria may affect the composition of DOM released by phytoplankton (reviewed by Mühlenbruch et al. 2018).

45

Metabolic capability to utilize rapid pulses of phytoplankton-derived DOM varies among bacteria and thus phytoplankton blooms are followed by distinct succession patterns of various bacterial genera commonly from classes Gammaproteobacteria, Alphaproteobacteria and Bacteroidia (Mühlenbruch et al., 2018; Teeling et al., 2012). Marine bacteria are often functionally divided in copiotrophs and oligotrophs based on their C uptake strategies. Oligotrophs are specialized in low nutrient concentrations, whereas copiotrophs thrive in high nutrient and DOM concentrations. Labile DOM attracts copiotrophic bacteria capable of quickly draining the DOM pool of its most bioavailable labile components (Pedler et al., 2014). Copiotrophic bacteria can become abundant e.g. after phytoplankton blooms, but because oceans mostly are a low-nutrient environment copiotrophic bacteria are not as ubiquitous as oligotrophic bacteria, e.g. alphaproteobacterial SAR11 clade (Morris et al., 2002).

55



Optical properties of colored and fluorescent DOM (CDOM and FDOM respectively) can be used as proxies of DOM bioavailability and source (Coble, 1996). Proxies for properties such as molecule size and amino acid content can be used to make predictions of the ecological function of the DOM pool by e.g. identifying DOM produced by phytoplankton blooms (Suksomjit et al., 2009), DOM degraded by bacteria (Kinsey et al., 2018) or DOM of freshwater origin (Coble, 1996). CDOM produced by phytoplankton differ in composition depending on phytoplankton species (Fukuzaki et al., 2014; Romera-Castillo et al., 2010). The composition is further altered by bacterial DOM utilization (Guillemette and del Giorgio, 2012; Romera-Castillo et al., 2011).

Mixed species communities mask C cycling differences that stem from the traits of individual phytoplankton species. Even during single species blooms previous environmental conditions may affect C cycling and DOM processing. Knowledge on the full cascade of C cycling through manipulated phytoplankton-bacterial communities aids to understand the contribution of individual phytoplankton species to C cycling in mixed communities. This is especially important because the composition of natural mixed phytoplankton communities seems to have little effect on the chemical composition of the accumulated autochthonous DOM, apparently due to rapid bacterial DOM processing (Haraguchi et al., 2019). In mixed phytoplankton communities it is also difficult to detect how the age and physiological state of individual phytoplankton species affects bacterial community composition (Grossart et al., 2005).

Environmental change has affected the composition of phytoplankton communities (Li et al., 2009). In the Baltic Sea spring blooms have shifted towards dinoflagellate predominance (Klais et al., 2011) and the ecological consequences of this shift are currently being investigated (Spilling et al., 2018). In this study we investigated how ecophysiology of two phytoplankton species affects microbial C cycling. We compared two common coastal phytoplankton species; a larger dinoflagellate *Apocalathium malmogiense* and a smaller, fast growing, cryptophyte *Rhodomonas marina*. In broad sense they could be considered a K-strategist and an R-strategist, respectively (*A. malmogiense* can produce cysts (Kremp and Heiskanen, 1999), can use allelopathy to inhibit growth of competitors (Suikkanen et al., 2011) and has slower growth rate than *R. marina*). *A. malmogiense* is a common, bloom-forming species in the Baltic Sea during spring, and *R. marina* was chosen as a general model organism representing a smaller, faster growing phytoplankton. We hypothesized that these two phylogenetically and physiologically different phytoplankton species show differences in C cycling. The focus of the experiment was to investigate whether these differences can be detected consistently on all levels of C cycling, from dissolved inorganic C (DIC) fixation via DOM release to bacterial DOM uptake and processing. Using phytoplankton cultures inoculated with natural bacterial community from the Baltic Sea, we experimentally investigated how species-specific differences in primary production (PP) and DOM production affect C flow from phytoplankton to bacteria, and bacterial DOM consumption, production and community composition. Our results on the effects of individual species on C cycling will increase understanding on how community shifts driven by environmental change will affect C cycling in aquatic environments.




## 2 Methods

### 2.1 Experimental setup

The ecophysiology of two different phytoplankton species and its effect on microbial C cycling from DIC uptake to bacterial DOC processing was investigated in an experimental study design. A larger (cell volume 3391-12764  $\mu\text{m}^3$ , (Olenina et al., 2006)) dinoflagellate *Apocalathium malmogiense* (G.Sjöstedt) Craveiro, Daugbjerg, Moestrup & Calado 2016 was compared to a smaller (mean cell volume 217  $\mu\text{m}^3$ , (Olenina et al., 2006)), fast growing, cryptophyte *Rhodomonas marina* (P.A.Dangeard) Lemmermann 1899. Both species are common in the Baltic Sea during spring season. Phytoplankton cultures were acquired from the FINMARI culture collection/SYKE Marine Research Centre (*A. malmogiense* (syn. *Scrippsiella hangoei*), culture id: SHTV-2, isolated in Storfjärden, Tvärminne by Anke Kremp in 2002; *R. marina*, culture id: Crypto08-A2, isolated in Storfjärden, Tvärminne by Anke Kremp in 2008). Phytoplankton cultures were grown in artificial sea water to minimize the effect of growth medium on optical DOM properties. Cultures were inoculated with the natural bacterial community from the Baltic Sea (hereafter called *A. malmogiense* treatment and *R. marina* treatment) and then investigated experimentally for the effect of species-specific differences in PP and DOM production on C flow from phytoplankton to bacteria, and bacterial DOM consumption, production and community composition.

The experiment was conducted at Tvärminne Zoological Station (59.844966, 23.249642) during the winter 2017-2018 in two parts; the *DOM release experiment* and the *DOM consumption experiment* (Fig. 1). The purpose of the DOM release experiment was to study the long-term net accumulation and alterations of DOM in conditions where phytoplankton produce new DOM and bacteria consume it. The purpose of the DOM consumption experiment was to study the effect of bacteria on DOM processing when the phytoplankton are removed. The experiment was timed to winter and spring months because the phytoplankton used were spring bloom species and we wanted the natural bacterial inoculum to represent winter and spring bloom bacteria. Non-axenic unialgal batch cultures were grown in two triplicate series, one for each part of the experiment with identical growth conditions. Cultures were grown in F/2 growth medium in 5 L Erlenmeyer flasks in 4 °C in approximately 60  $\mu\text{mol photons s}^{-1} \text{m}^{-2}$  under light-dark regime of 14 h and 10 h. The growth medium was prepared in artificial sea water (autoclaved MQ water adjusted to salinity 6 using Tropic Marin Classic Sea Salt). Vials were stirred manually every 1-2 days and prior to any sampling.



120 **Figure 1.** A schematic description of the DOM release experiment (a) and the DOM consumption experiment (b). Both experiments  
 were conducted separately for *A. malmogiense* and *R. marina*. Black timeline arrow at the far left starts from the inoculation of  
 phytoplankton into 5 L growth vials. Grey arrows depict the flow of water through different filtration, mixing and measurement  
 steps. (a, bracketed area) Procedure for an individual key point incubation, which were conducted thrice for each species. On day 1  
 bacterial inoculum was prepared, on day 2 the incubation was initiated (i.e. phytoplankton treatments were established) and  
 125 measurements (production line: primary production, bacterial production, <sup>14</sup>C flow; DOM line: DOC, CDOM, bacterial abundance)  
 were taken at 4 h intervals and on day 3 the final primary production measurement was taken (for net primary production). (b,  
 bracketed area) Incubation of the DOM consumption experiment, which was conducted once for each species. On preparation day  
 1 bacterial inoculum was prepared, on preparation day 2 the cultures were filtered, on day 1 the incubation was initiated (i.e.  
 phytoplankton treatments were established) and measurements (DOC, CDOM, bacterial abundance, production, respiration,  
 community composition and growth efficiency) were taken daily until day 7. The variables measured at each phase of the two



130 experiments are listed in Table A1. KPI = Key Point Incubation (see text), BR = optical O<sub>2</sub> consumption (bacterial respiration) measurement, F/2 = F/2 growth medium (control), TFF = concentration of bacteria by tangential flow filtration

In the first part of the experiment, the DOM release experiment (Fig. 1a), the phytoplankton and bacteria present in the cultures were grown together for over 4 months and phytoplankton and bacterial abundance and optical properties of DOM were monitored 1-3 times a week. At the beginning of the exponential growth phase and at two later stages the C flow from phytoplankton to bacteria, DOM alterations and bacterial activity were measured using day-long incubations. During these  
135 three measurement occasions (hereafter referred to as 1<sup>st</sup>, 2<sup>nd</sup> and 3<sup>rd</sup> KPI after Key Point Incubation) subsamples of phytoplankton cultures were incubated with an inoculum of natural bacterial community, for 24 h and sampled at 0, 4, 8 and 12 h (+ extra sampling at 24 h for net PP). This experiment addressed the C flow in a combined phytoplankton-bacterial community system.

140 In the DOM release experiment concentrations of Chlorophyll *a* (Chl *a*) and particulate organic C and N (POC and PON, respectively) were measured, and bacterial community composition was determined, before each KPI. Two separate sample sets were incubated at each KPI. In the first set, hereafter referred to as *production line* (Fig. 1a), phytoplankton cultures and bacteria (90% vol phytoplankton culture + 10% vol ml bacterial inoculum in 10 ml aliquots) were incubated in light and PP, bacterial production (BP) and <sup>14</sup>C transfer from <sup>14</sup>C-NaHCO<sub>3</sub> via phytoplankton to DOC pool and bacterial biomass were  
145 measured. Transfer of <sup>14</sup>C to DOC was investigated by filtering PP samples through 0.45 μm GD/X (Whatman) syringe filters and by measuring the radioactivity in the filtrate. Transfer of <sup>14</sup>C from DOC to bacterial biomass was investigated by incubating the previously mentioned filtrate for 4 h in dark after which the incubation was stopped by addition of 50% trichloroacetic acid and the particulate biomass in the samples was centrifuged for analysis of radioactivity. The protocol for the production line is depicted in detail in Fig. A1.

150 In the second set, hereafter referred to as *DOM line* (Fig. 1a), phytoplankton were removed by 0.8 μm filtration and bacteria (225 ml filtered phytoplankton culture + 25 ml bacterial inoculum) were incubated in dark and DOC concentration, optical properties of DOM, and bacterial abundance were measured. This incubation was conducted in the dark to see if the bacteria influence the DOM pool already during the 12 h KPI when photosynthesis is stopped. Inorganic nutrients (NO<sup>3-</sup>, including  
155 NO<sup>2-</sup>, and PO<sub>4</sub><sup>3-</sup>) were measured in untreated culture and DOM line samples at 0 h and 12 h.

Except for PP measurements, there were no trends in any other production line or DOM line variables from 0 to 12 h measurements. Therefore the measurements at all time points within a KPI were pooled for statistical analysis and presentation.

160 In the second part of the experiment, the DOM consumption experiment (Fig. 1b), the phytoplankton were grown to high density (*A. malmogiense*: ~1 × 10<sup>4</sup> cells ml<sup>-1</sup>, *R. marina* ~9 × 10<sup>4</sup> cells ml<sup>-1</sup>) after which the phytoplankton and most of the bacteria were removed by filtering (GF/F filters pre-combusted 450 °C for 4 h; Whatman). The filtrate, inoculated with natural



bacterial community (1480 ml filtered phytoplankton culture + 120 ml bacterial inoculum), was incubated for 7 days to study the DOM processing and C flow in the bacterial compartment without phytoplankton present. To ensure nutrient replete conditions, 18  $\mu\text{mol NH}_4\text{Cl}$  and 11  $\mu\text{mol NaH}_2\text{PO}_4$  was added in the experimental mixtures at the start of the incubation. Temperature during the incubation was increased to 10 °C to enhance the bacterial processes for easier detection. A control treatment containing only F/2 medium and natural bacterial community inoculum was used to investigate how the natural bacterial community develops and how their DOC processing differs in the growth medium in the absence of DOM derived from the cultured phytoplankton and competition from cultured bacteria. During the 7-day incubation of the DOM consumption experiment DOC concentration, optical properties of DOM and bacterial abundance, production, respiration, community composition and growth efficiency were measured daily. The variables measured at each phase of the DOM release experiment and the DOM consumption experiment are listed in Table A1.

The natural bacterial community inoculum was prepared the same way for both parts. Sea water was collected at the pier of the station and bacterial abundance and community composition were measured. Seawater was vacuum filtered using 10 mmHg pressure through 0.8  $\mu\text{m}$  pore size polycarbonate membrane filter ( $\text{\O}$  47mm; Whatman) to remove grazers including heterotrophic nanoflagellates and 3 L of the filtrate was concentrated to about 30 mL using tangential filtration (Pall Minimate 100 kDa TFF Capsule), and then diluted to 300 mL with artificial seawater, after which bacterial abundance was measured again. All handling of seawater and bacterial concentrate was done in 4 °C. Purpose of this treatment was to concentrate the seawater bacterial concentration 10-fold and to remove most of the marine DOM. Phytoplankton culture and bacterial concentrate were mixed in volume ratio of 90:10% in an attempt to recreate the natural concentration of sea water bacteria. Time limitations of filtrations forced the use of only 92.5:7.5% in the DOM consumption experiment. However, the concentration of sea water bacteria proved to be inefficient and the final ratio of sea water bacteria to bacteria present in the culture was small (DOM release experiment: *A. malmogiense*: 7.53%, 0.02% and 0.03%, *R. marina*: 18.11%, 0.03% and 0.02%, at 1<sup>st</sup>, 2<sup>nd</sup> and 3<sup>rd</sup> KPI, respectively; DOM consumption experiment: *A. malmogiense*: 3.38%, *R. marina*: 1.70%).

## 2.2 Laboratory analyses

### 2.2.1 Cell abundance

Phytoplankton and bacterial abundance were analyzed using flow cytometry (BD Accuri C6 Plus). Phytoplankton abundance was analyzed in untreated samples by plotting red fluorescence (670 nm long pass filter, 488 nm excitation) against forward scatter. Samples for bacterial abundance were fixed with paraformaldehyde (final concentration: 0.9%) and glutaraldehyde (final concentration: 0.045%), incubated in room temperature for 30 min, frozen in liquid N and stored in -80 °C until analysis. After thawing samples were diluted 10-100 fold with pH 8 TE-buffer, stained with Sybr Green I nucleic acid stain (final concentration 1:10000 vol.) and incubated in dark at room temperature for 10 min. Heterotrophic bacteria were detected and counted by plotting green (530/30 bandpass filter, 488 nm excitation) fluorescence against red (670 nm long pass filter, 488





195 nm excitation) fluorescence so that they could be differentiated from cells containing Chl *a*. Cytometer data were analyzed with FCS Express 5 software (De Novo software).

### 2.2.2 Primary and bacterial production and <sup>14</sup>C transfer

DIC was analyzed with Elektro-Dynamo URAS-3E C analyzer against NaHCO<sub>3</sub> standards. PP was measured from mean of light sample <sup>14</sup>C-activity corrected with dark sample <sup>14</sup>C-activity according to Gargas (1975) with the modifications described in Fig. A1. PP measured at each KPI represents cumulative gross PP divided by time (GPP). PP at 24 h is the net PP including the dark period (NPP). PP was used to calculate community respiration according to Spilling et al. (2019) using equation 1. Actual respiration measurements were not available for either phytoplankton species so uniform respiration rates for light and dark periods were assumed.

205 eq 1: respiration = (GPP × 14) – (NPP × 24)

GPP in *A. malmogiense* treatments decreased from 4 to 12 h at the 2<sup>nd</sup> and 3<sup>rd</sup> KPI. This might have been caused by high respiration (see results) but also by insufficient addition of <sup>14</sup>C-NaHCO<sub>3</sub> which might have resulted in underestimation of GPP which we did not want to carry over to other results. Therefore GPP at 4 h was used in calculation of community respiration and <sup>14</sup>C flow percentages.

The transfer of <sup>14</sup>C originating from <sup>14</sup>C-NaHCO<sub>3</sub> measured in percent from phytoplankton via DOC to bacteria was quantified by dividing the accumulation rate of <sup>14</sup>C in each compartment with the accumulation rate in the previous compartment (i.e. GPP:DIC, DOC:GPP and bacterial biomass:DOC). Accumulation of <sup>14</sup>C-activity in DOC and in bacterial biomass was calculated by dividing the time normalized activity in samples after 12 h incubation with specific activity of <sup>14</sup>C-NaHCO<sub>3</sub>. Before calculations, <sup>14</sup>C accumulation in DOC pool and in bacterial biomass were corrected for the ratio of <sup>14</sup>C-DIC to ambient DIC concentration. This was done by multiplying them with the ratio of PP to bulk <sup>14</sup>C accumulation rate in phytoplankton biomass.

### 2.2.3 Bacterial production and respiration

220 Thymidine and leucine-based BP were measured with the centrifugation method (Smith and Azam, 1992) with modifications for the DOM release experiment described in Fig. A1. <sup>3</sup>H-thymidine incorporation was converted to bacterial biomass increase by using conversion factors  $1.1 \times 10^{18}$  cells mol<sup>-1</sup> (Riemann et al., 1987) and  $0.12 \text{ pg C} \times (\mu\text{m}^3 \text{ cell}^{-1})^{0.7}$  (Norland, 1993) using theoretical bacterial cell volume of  $0.063 \mu\text{m}^3 \text{ cell}^{-1}$  (Kuparinen, 1988). <sup>14</sup>C-leucine incorporation was converted to bacterial biomass increase with a conversion factor of  $1.55 \text{ kg C mol}^{-1}$ .

225



From each experimental treatment in the DOM release experiment 100 mL was enclosed in air-tight septum sealed Duran bottles at the start of the incubation for bacterial respiration (BR) measurements (Fig. 1b). Needle sheathed oxygen optode (PreSens NTH-PSt1-L5-TF-NS120/0.8-YOP) was pierced through the septum to monitor bacterial oxygen consumption with an OXY-4 micro oxygen meter (PreSens). Prior to measurements oxygen optodes were calibrated to 0% and 100% air saturation by exposing the optode to Na<sub>2</sub>SO<sub>3</sub> solution and water vapor-saturated air respectively. Relative oxygen concentration was recorded every 10 minutes through the incubation. BR for each day was calculated by dividing the difference in relative O<sub>2</sub> saturation between the start of the experiment and each day with time. Oxygen solubility of 678.8 mmol l<sup>-1</sup> (at 10 °C, 1 atm and salinity 6) was used to convert relative O<sub>2</sub> saturation to molar concentration, which was then converted to units of mol C L<sup>-1</sup> h<sup>-1</sup>.

Bacterial growth efficiency (BGE) at each day during DOM consumption experiment was calculated separately for thymidine and leucine-based BP using the equation 2

$$\text{eq 2: } \text{BGE} = \text{BP} / (\text{BP} + \text{BR})$$

where BP is bacterial production and BR is bacterial respiration calculated as the change in C during the previous day and converted to h<sup>-1</sup>. Therefore BGE is reported starting from day 2.

#### 2.2.4 Bacterial community composition

500 mL (sea water) or 100 mL (DOM release experiment water (DOM line mix) and DOM consumption experiment water) was vacuum filtered onto sterile 0.22 µm pore size membrane filters (Ø 47mm; Whatman), frozen in liquid N and stored in -80 °C until analysis. DNA was extracted from filters with DNeasy Power Soil kit (Qiagen) 6 months after the experiments and stored in -80°C for further processing. In addition, negative controls without a sample were extracted. Only one replicate from seawater (both experiments) and cultures (DOM release experiment) was sequenced (Fig. 1). For sequencing, 16S ribosomal RNA gene region V4 was amplified with a polymerase chain reaction, using the universal bacterial primers 341F and 785R (Klindworth et al., 2013). A two-step polymerase chain reaction and Illumina MiSeq (Illumina Inc, San Diego, CA, USA) paired-end multiplex sequencing were performed at the Institute of Biotechnology, University of Helsinki, Finland. In total 16 × 10<sup>6</sup> paired raw reads were obtained with the Illumina MiSeq platform. Primer removal was done with Cutadapt (settings - m 1 -O 15 -e 0.2, V 2.1 with Python 3.5.3, (Martin, 2011)). Reads were merged and processed according to DADA2 pipeline (DADA2 V 2.1.10 Rcpp V 1.0.0, (Callahan et al., 2016)) with filterAndTrim maxEE = 3. After filtering and trimming, a total of 11.2 × 10<sup>6</sup> sequences remained from which 10.7 × 10<sup>6</sup> were merged and 9.1 × 10<sup>6</sup> were non chimeric and used for further analyses. Taxonomic classification of the Amplicon Sequence Variants (ASVs) was done with DADA2 default parameters (minBoot = 50) using Silva for DADA2 (v. 132, (Quast et al., 2013), <https://zenodo.org/record/1172783#.Xila11MzZgg>). Before the statistical analyses, chloroplasts and mitochondria were removed ending up with 4545 ASVs. Raw reads are



deposited in the Sequence Read Archive of National Center for Biotechnology Information under BioProject accession number  
260 PRJNA647035.

### 2.2.5 Dissolved C and N and optical properties of DOM

DOC and CDOM samples were prepared by filtering 20 mL of water through acid washed and pre-combusted GF/F filters (450 °C, 4 h) into acid washed and pre-combusted glass vials which were then sealed with a septum cap. DOC samples were acidified to pH 2 with 2 M HCl and stored in -20 °C until analysis of DOC with Shimadzu TOC-V CPH total organic carbon  
265 analyzer. Filtered CDOM samples were analyzed within 24 h. CDOM absorption was measured using a Shimadzu 2401PC spectrophotometer with a 4 cm quartz cuvette over the spectral range from 200 to 800 nm with 1 nm resolution. Ultrapure water (MQ) was used as the blank for all samples. Excitation-emission matrices (EEMs) of FDOM were measured with a Varian Cary Eclipse fluorometer (Agilent). Processing of the EEMs was done using the eemR package for R software (Massicotte, 2016). A blank sample of ultrapure water was subtracted from the EEMs, and the Rayleigh and Raman scattering  
270 bands were removed from the spectra after calibration. EEMs were calibrated by normalizing to the area under the Raman water scatter peak 11 (excitation wavelength of 350 nm) of an MQ water sample run on the same session as the samples, and were corrected for inner filter effects with absorbance spectra (Murphy et al., 2010). For assessing the characteristics and the quality of the DOM pool, fluorescence peaks (Coble, 1996) were extracted from the EEMs. In this study the following optical variables were used as proxies for DOM characteristics: absorbance coefficient at 254 nm ( $a_{CDOM(254)}$ ) as a general indicator  
275 of optically active molecules and light attenuation, absorption spectral slope between 275 and 295 nm as a proxy of molecular size ( $S_{275-295}$ , (Helms et al., 2008)), fluorescence peaks T and C (Coble, 1996) as proxies of protein-like and humic-like DOM, respectively, and humification index (HIX, (Zsolnay et al., 1999)) as an indicator of relative humification of DOM. Additional optical variables were collected, but these were not included in the detailed analysis and are only presented in Appendix B (Fig. B1, Fig. B2 and Table B1).

### 280 2.2.6 Particulate organic C and N, Chl *a* and nutrients

For POC/N and Chl *a* measurements 20 mL of sample water was filtered through GF/F filters (for POC/N they were pre-combusted in 450 °C for 4 h). The POC/N filters were wrapped in a foil and stored in -20 °C until analysis with Europa Scientific ANCA-MS 20-20 <sup>15</sup>N/<sup>13</sup>C mass spectrometer. Chl *a* filters were stored in EtOH in dark at room temperature overnight, after which they were stored in -20 °C until fluorometric analysis with a Varian Cary Eclipse spectrofluorometer.  
285 300 µL of sample was added into a well plate and the fluorescence was measured (excitation/emission: 430/670 nm). Fluorescence intensity was converted to Chl *a* concentration using Chl *a* standards (Sigma).

Nutrient samples were frozen immediately after sampling in -20 °C and stored frozen until measurement according to Grasshoff et al. (1999) using Thermo Scientific Aquakem 250 photometric analyzer.



## 290 2.2.7 Statistical analyses

All statistical analyses were done using R version 3.6.1 (R Core Team, 2019) and figures using package ggplot2 (Wickham, 2016). Differences in variables between treatments (species) and among KPIs were analyzed using Welch-ANOVA, which allows for some difference in variance among treatments. If there was no trend in measurements of all the time points within a KPI all the measurements were pooled for the statistical analyses. For cumulative variables (<sup>14</sup>C accumulation rate in DOC  
295 and in bacterial biomass) measurements at 12 h were used in the statistical analyses. For GPP measurements at 4 h were chosen for statistical analyses (justified in results). Significant differences among KPIs were investigated using Games-Howell post hoc test (Peters, 2018). Differences in ANOVA were considered significant at a  $p < 0.05$ . Results of all Welch-ANOVA tests are given in Appendix C.

300 All multivariate analyses for bacterial community analysis were performed on the Bray–Curtis dissimilarity matrix derived from square-root transformed values. The bacterial community dynamics in the experiments was visualized with Principal Coordinates Analysis (PCoA). To determine whether the bacterial communities differed significantly between different phytoplankton species a PERMANOVA (permutational multivariate analysis of variance (Anderson, 2001)) was performed using the function adonis (9999 permutations) in the R package vegan (Oksanen et al., 2019). Due to the lack of replicates,  
305 differences between phytoplankton treatments with seawater and control treatments could not be tested. The homogeneity of dispersion was tested using the function betadisper (9999 permutations) in the R package vegan (Oksanen et al., 2019). To determine the association between the environmental parameters and bacterial community composition Distancebased Redundancy Analysis (dbRDA) with 9999 permutations (capscale, (Oksanen et al., 2019)) was done. Significance of the model and the explanatory variables were tested with analysis of variance (anova, (Oksanen et al., 2019)), using 9999 permutations.  
310 Biologically most relevant factors with Variance inflation factor less than 10 that were statistically significant ( $p < 0.05$ ) were chosen for the analysis (vif.cca, (Oksanen et al., 2019)). Bar plots and PCoA (Bray-Curtis dissimilarity) were made with phyloseq v. 1.26.1 (McMurdie and Holmes, 2013).


## 3 Results

### 3.1 Phytoplankton growth and primary production

315 In the DOM release experiment *R. marina* grew faster to maximum density and ended the growth phase sooner than *A. malmogiense* (Fig. 2a). The average cell size of both phytoplankton species remained unchanged throughout the experiment, as indicated by the forward scatter results from flow cytometry (data not shown). There was no indication of N limitation during the experiment as nitrate concentrations remained high (Table B1). P was depleted in filtered *A. malmogiense* treatments but not in unfiltered treatments (Table B1), suggesting intracellular phosphate storing. PP (gross, net and cell specific) was  
320 higher in *A. malmogiense* treatments (Fig. 2b, Table C1).



The timing of KPIs aimed to capture comparable growth phases for these short incubations but *A. malmogiense* cultures were still growing when spring bloom was closing in, so we had to initiate the measurements at earlier stages while the natural bacterial communities still resembled winter and spring communities (Fig. 2a).



325

**Figure 2. (a) Abundance of phytoplankton (mean of three replicate treatments, error bars indicate one standard deviation) during the DOM release experiment. Vertical lines from left to right in each panel mark the times of 1<sup>st</sup>, 2<sup>nd</sup> and 3<sup>rd</sup> KPI. (b) Cell specific gross primary production in *A. malmogiense* (white) and *R. marina* (grey) treatments at each KPI. Number above the boxplots shows the number of measurements. Lower and upper hinges of boxes mark the first and third quartiles, respectively, while the whiskers extend to the lowest and the highest value within 1.5 times the interquartile range.**

330


335

In *A. malmogiense* treatments also a second population distinguished by flow cytometry based on lower red fluorescence (Chl *a* fluorescence) started to grow at the same time with the main population (Table B1). This population grew linearly to about 15% of maximum density of the main population. These were considered to possibly be cysts and were not included in *A. malmogiense* abundance in further analyses, although *A. malmogiense* does not usually produce cysts in the low temperature used in this study (Kremp et al., 2009). This decision brings a certain bias to the interpretation of the results, but we did relevant calculations with and without these cells and the results did not change significantly. We chose to exclude the group because we could not be certain of what the group consists of and we wanted to avoid including e.g. cysts in the population of active *A. malmogiense* cells. We considered that doing the opposite would have introduced unknown distortion to our interpretation.



### 3.2 Bacterial production and $^{14}\text{C}$ transport

340 In the DOM release experiment differences in bacterial abundance between the phytoplankton were modest at each KPI (Fig. 3a, Table C1). Ratio of bacteria to phytoplankton was most of the time between  $1 \times 10^4$  and  $3 \times 10^4$  except at the 1<sup>st</sup> KPI for *R. marina*, when it was much lower (Table B1, C1). Thymidine incorporation was slightly higher in *A. malmogiense* treatments at each KPI (Fig. 3b, Table C1), while leucine incorporation was of equal magnitude between the species, except at the 1<sup>st</sup> KPI (Fig. 3c, Table C1). As a result the ratio of leucine to thymidine incorporation was higher in *R. marina* treatments at the 2<sup>nd</sup> and 3<sup>rd</sup> KPI (Table B1, C1). In the DOM consumption experiment bacterial abundance increased considerably faster and thymidine and leucine incorporation was higher in *R. marina* treatments than in *A. malmogiense* treatments (Fig. 3d-f). These observations from both experiments suggest that *R. marina* community can support a more productive bacterial community in proportion to PP.




350 **Figure 3.** Bacterial abundance (a, d), cell specific bacterial thymidine (b, e) and leucine (c, f) incorporation at each KPI (0-12 h measurements pooled) in the DOM release experiment (boxplots) and in the DOM consumption experiment (line graphs). Lines A, B and C mark different replicates (culture filtrate + sea water bacteria) and ctrl marks the control (F/2 + sea water bacteria). Number above the boxplots shows the number of measurements. Lower and upper hinges of boxes mark the first and third quartiles, respectively, while the whiskers extend to the lowest and the highest value within 1.5 times the interquartile range. White boxes = A.



355 *malmogiense*, grey boxes = *R. marina*. Due to a measurement error only 12 h measurements of bacterial abundance are available in  
*A. malmogiense* treatments at 3<sup>rd</sup> KPI.

*A. malmogiense* cells were more efficient in incorporating DIC (i.e. higher PP, Fig. 2b) but they also respired more than *R.*  
*marina* cells, as shown by higher cell-specific community respiration in the DOM release experiment (Fig. 4a, Table C1). In  
the DOM consumption experiment cell-specific BR was higher in *R. marina* treatments (Fig. 4b). Although BP was higher in  
360 *R. marina* treatments, the lower BR in *A. malmogiense* treatments led to comparable BGE in both treatments (Fig. 4c). Since  
BR was consistently higher in *R. marina* treatments (DOM consumption experiment) and the ratio of bacteria to phytoplankton  
cells was higher in the *A. malmogiense* treatments only at the 1<sup>st</sup> KPI (DOM release experiment, Table B1), the higher cell-  
specific community respiration in *A. malmogiense* treatments is likely mainly caused by respiration of phytoplankton.



365 **Figure 4. (a) Community respiration divided by phytoplankton abundance in *A. malmogiense* treatments (white) and *R. marina***  
**treatments (grey) at each KPI in the DOM release experiment. Cell specific BR (b) and BGE calculated from leucine incorporation-**  
**based BP (c) in the DOM consumption experiment. Lines A, B and C mark different replicates (culture filtrate + sea water bacteria)**  
**and ctrl marks the control (F/2 + sea water bacteria). Number above the boxplots shows the number of measurements. Lower and**  
370 **upper hinges of boxes mark the first and third quartiles, respectively, while the whiskers extend to the lowest and the highest value**  
**within 1.5 times the interquartile range.**

<sup>14</sup>C-DOC, originating from <sup>14</sup>C-NaHCO<sub>3</sub>, was produced by both species at each KPI (Fig. B3). <sup>14</sup>C originating from <sup>14</sup>C-  
NaHCO<sub>3</sub> was incorporated in bacterial biomass at 2<sup>nd</sup> and 3<sup>rd</sup> KPI. There were considerable uncertainties with this  
measurement; <sup>14</sup>C-DOC was produced also in dark controls and surprisingly much at the 1<sup>st</sup> KPI compared to the latter KPIs  
when phytoplankton biomass was much higher (Fig. B3). Regardless, these uncertainties concern both phytoplankton species  
375 so we consider the analysis to be suitable for comparing the species, despite uncertainties in the absolute quantities of <sup>14</sup>C in  
different compartments. Higher PP of *A. malmogiense* led to higher fraction of <sup>14</sup>C-NaHCO<sub>3</sub> pool being incorporated into  
phytoplankton biomass compared to *R. marina* (Table 1). However, a larger fraction of PP ended in filtrate in *R. marina*  
treatments (Table 1). In *R. marina* treatments also a larger fraction of <sup>14</sup>C-organic matter was incorporated into bacterial  
biomass at 2<sup>nd</sup> and 3<sup>rd</sup> KPI (Table 1), although the difference was very small at 3<sup>rd</sup> KPI (at the 1<sup>st</sup> KPI no activity was detected  
380 in *R. marina* treatments so comparisons could not be made). Of all the <sup>14</sup>C that was fixed by PP about 5 and 4 times more  
ended up in bacterial biomass in *R. marina* treatments at 2<sup>nd</sup> and 3<sup>rd</sup> KPI respectively (Table 1).



385 **Table 1. Flow of  $^{14}\text{C}$  between different phases of C cycle. Numbers are percentages of  $^{14}\text{C}$  accumulation rates between the phases indicated in the left column (PP:DIC is an exception as PP is a rate but DIC is a concentration). Stars indicate the significance ( $p$ -values) at the side of the significantly higher percentage between species at the corresponding KPI, compared with Welch-ANOVA (\*\*\*)  $< 0.001 < ** < 0.01 < * < 0.05$ ). Welch-ANOVA results are presented in Table C2.**

Species KPI	<i>A. malmogiense</i>			<i>R. marina</i>		
	1 <sup>st</sup>	2 <sup>nd</sup>	3 <sup>rd</sup>	1 <sup>st</sup>	2 <sup>nd</sup>	3 <sup>rd</sup>
PP:DIC	3.02*	5.24***	4.88***	0.68	0.67	0.64
DOC:PP	0.402	0.119	0.128	1.095**	0.420***	0.419***
bacteria:DOC	1.39	24.07	18.15	NA	33.87*	22.84*
bacteria:PP	0.0049	0.0283	0.0233	NA	0.1429***	0.0965**


390 Given that comparable amounts of DOC accumulated during the DOM release experiment it can be assumed that, given higher PP, relatively larger share of fixed C is directed to respiration by *A. malmogiense* and to DOC release by *R. marina*. These results also suggest that DOM released by *R. marina* is more bioavailable to bacteria than DOM released by *A. malmogiense*, and that the microbial loop is favored more strongly when DOM originates from *R. marina*.

### 3.3 Bacterial community

395 In both experiments, bacterial communities resembled those in the cultures and were distinct from the seawater community (bacterial inoculum), suggesting that the addition of seawater bacteria had a negligible contribution to the composition of the total bacterial community (Fig. 5).


400 In the DOM release experiment classes Alphaproteobacteria, Gammaproteobacteria and Bacteroidia predominated the bacterial communities (Fig. 5). Alphaproteobacteria increased and Gammaproteobacteria decreased from the 1<sup>st</sup> KPI to the 2<sup>nd</sup> KPI, whereas Bacteroidia had its peak at the 2<sup>nd</sup> KPI. The relative share of different classes differed between the treatments: Bacteroidia (average: 45.6%, genera *Algoriphagus* and *Polaribacter*) and Alphaproteobacteria (average: 44.7%, genera *Pseudorhodobacter* and *Sphingorhabdus*) were the most abundant classes in *A. malmogiense* treatments while Alpha- (average: 49.9%, genus *Pseudorhodobacter*) and Gammaproteobacteria (average: 38.8%, genera *Rheinheimera* and RS62 marine group) predominated in the *R. marina* treatments. Interestingly, class Actinobacteria (average: 2%, genus *Candidatus Aquiluna*) appeared in *A. malmogiense* treatments and slightly increased along the experiment. Bacterial communities in both  
 405 phytoplankton treatments changed between the KPIs (PCoA, Fig. 6a, c). There were also differences in bacterial communities between the phytoplankton species in relation to selected environmental variables: thymidine-based BP correlated with *A. malmogiense* at the 2<sup>nd</sup> and 3<sup>rd</sup> KPI whereas  $a_{\text{CDOM}(254)}$  and  $S_{275-295}$  correlated with *R. marina* at the 2<sup>nd</sup> and 3<sup>rd</sup> KPI (Fig. 6c). In total, dbRDA axis 1 and 2 explained 51.33% of the variation in bacterial community analysis.





410

**Figure 5.** Class level (upper panels) and genus level (lower panels) bacterial diversity of 16S ribosomal RNA (rRNA) gene sequences representing >1% of all amplicon sequence variants (ASVs) in DOM release experiment (a, c) and in DOM consumption experiment (b, d). (a, c) The bars labeled control refer to cultures before addition of bacterial inoculum. (b, d) The bars labeled control refer to the control treatments (F/2 + sea water bacteria).



415

**Figure 6.** (a, b) Principal coordinate analysis plots showing bacterial community dynamics in different experiments and (c, d) dbRDA biplots showing relationship between bacterial communities and selected, significant environmental variables (anova,  $p < 0.05$ ) in DOM release experiment (a, c) and DOM consumption experiment (b, d). THY = thymidine based bacterial production, a254 =  $a_{CDOM(254)}$ , S275-295 =  $S_{275-295}$ .

420 In the DOM consumption experiment class Bacteroidia (average: 70%, genera *Algoriphagus* and *Polaribacter*) and Alphaproteobacteria (average: 19.9%, genera *Pseudorhodobacter*, *Sphingorhabdus* and *Seohaecicola*) predominated bacterial communities in *A. malmogiense* treatments (Fig. 5). Congruently with DOM release experiment, class Actinobacteria (average: 5.7%, genus *Candidatus Aquiluna*) was present throughout the experiment. In the small share of class Gammaproteobacteria (average: 1.7%), the most abundant were the order Betaproteobacteriales (genera *Hydrogenophaga*, *Kerstersia*, *Limnobacter*,  
 425 *Methylotenera*). In *R. marina* Alphaproteobacteria (day 1-3 average 82%, genus *Pseudorhodobacter*) predominated the bacterial communities until the day 3, after which they began to decrease (day 4-7 average: 36.1%) and Bacteroidia (day 4-7 average: 49.9%), genus *Flavobacterium*) increased. Also Gammaproteobacteria increased slightly towards the end of the experiment (day 4-7 average: 7.7%, genera *Shewanella*, *Marinomonas* and *Polynucleobacter*). In *R. marina*, bacterial community composition changed along time (PCoA, Fig. 6b). Bacterial communities differed significantly between the different phytoplankton treatments (adonis:  $R^2 = 0.67$ ,  $p < 0.001$ ), but due to the tight grouping in *A. malmogiense*, the homogeneity of variance was violated (betadisp:  $p > 0.05$ , Fig. 6d). However, because groups were not overlapping, it can be assumed that the observed differences are true. The shift in bacterial community was observed also in relation to selected environmental parameters: Bacterial communities on days 1-3 correlated with peak T and  $a_{CDOM(254)}$  whereas on days 4-7 they  
 430




435 correlated with thymidine-based BP (Fig. 6d). In total, dbRDA axis 1 and 2 explained 68.89% of the variation in bacterial community analysis.

In the control treatments of the DOM consumption experiment (seawater inoculum + growth media) bacterial communities were comparable with seawater community in the beginning of the experiment in both experimental treatments but later developed into cultures which were different from communities in both sea water and experimental treatments (Fig. 5).  
440 Interestingly, class Campylobacteria (genus *Argobacter*), which was not abundant in either of the phytoplankton treatments, began to increase in both control treatments on day 4 (Fig. 5 b, d).

### 3.4. DOM transformations

During the DOM release experiment DOC concentrations increased in both phytoplankton treatments, however, only a very small increase from 2<sup>nd</sup> to 3<sup>rd</sup> KPI was observed in *R. marina* treatments (Fig. 7a). DOM absorbance and fluorescence generally  
445 started to increase when the phytoplankton started to grow (Fig. 7, Fig. B1). The general trend was the accumulation of lower molecular weight and potentially more refractory molecules, as seen by e.g. increase in  $a_{CDOM(254)}$ ,  $S_{275-295}$ , humic-like DOM peak C, and HIX.  $S_{275-295}$ , peak C, and HIX increased faster in *R. marina* treatments (Table C1). While  $a_{CDOM(254)}$  increased in both species during the experiment, DOC-normalized absorbance at 254 nm (Weishaar et al., 2003) increased slightly in *R. marina* treatments, whereas it decreased in *A. malmogiense* treatments (Table B1). This suggests that the increase in  $a_{CDOM(254)}$   
450 in *R. marina* treatments was caused mainly by the increase in DOC absorbing in UV region with relatively higher intensity than the bulk material whereas in *A. malmogiense* treatments the increase in  $a_{CDOM(254)}$  was caused by increased absorbance due to higher bulk DOC concentration. This was also supported by the three times higher  $a_{CDOM(254)}$  production by *R. marina* when the  $a_{CDOM(254)}$  consumption by bacteria during the DOM consumption experiment is subtracted from the net accumulation of  $a_{CDOM(254)}$  in the DOM release experiment.





460 **Figure 7. DOC concentration (a, d), absorbance coefficient at 254 nm (b, e), spectral slope between 275 and 295 nm (c, f) fluorescence peaks T (g, j) and C (h, k) and humification index (i, l) at each KPI (0-12 h measurements pooled) in the DOM release experiment (boxplots) and in the DOM consumption experiment (line graphs). Lines A, B and C mark different replicates (culture filtrate + sea water bacteria) and ctrl marks the control (F/2 + sea water bacteria). Number above the boxplots shows the number of measurements. Lower and upper hinges of boxes mark the first and third quartiles, respectively, while the whiskers extend to the lowest and the highest value within 1.5 times the interquartile range. White boxes = *A. malmogiense*, grey boxes = *R. marina*. Optical variables not included in the detailed analysis are presented in Fig. B1 and Table B1 (DOM release experiment), and Fig. B2 (DOM consumption experiment).**

465 At the beginning of the DOM consumption experiment DOC concentrations were comparable and higher than in the control treatments (Fig. 7d), indicating that considerable DOC production by phytoplankton had occurred in both treatments despite the difference in the phytoplankton abundance before the start of the incubation (*R. marina*:  $\sim 9 \times 10^4$  cells mL<sup>-1</sup>, *A. malmogiense*:  $\sim 1 \times 10^4$  cells mL<sup>-1</sup>). During the incubation the DOC concentration did not change much in *A. malmogiense* treatments but decreased in *R. marina* treatments, especially during the first four days.

470 Contrary to the DOM release experiment, in DOM consumption experiment DOM absorbance decreased during the incubation, although often no clear change could be detected in *A. malmogiense* treatments (Fig. 7, Fig. B2). Peak C and HIX increased at first, as in the DOM release experiment, but started to decline at day 4 (Fig. 7k, l). Likely in the DOM release experiment the continuous production of fresh DOM by phytoplankton supplied the bacteria with bioavailable DOM, which was consumed and transformed to more refractory, UV-absorbing material. In the DOM consumption experiment the phytoplankton were no  
475 longer present as a fresh DOM source, so the bacteria started to use the more refractory material. This would also explain the bell-shaped curves (increase until day 4 and then decrease) of peak C and HIX in the DOM consumption experiment. Until day 4 the bacteria still used more bioavailable material which was left from the phytoplankton and converted it to optically active molecules, but on day 4 this material ran out and the bacteria switched to consuming more refractory, optically active material.

## 480 **4 Discussion**

### **4.1 DOC production, transformation and consumption**

In the first, DOM production experiment, the trends in DOC concentration and optical DOM characteristics were similar through KPIs 1-3, suggesting that there was no qualitative shift from production to consumption of any DOM fraction detected by the optical methods. Also the trends were similar between the species which further suggest that the observed changes in  
485 the optical DOM properties were more related to the age of the culture than to growth phases. Of course, the optical method does not detect changes in the concentrations of optically inert molecules, such as simple carbohydrates, and there may have been growth phase dependent changes in their production (Chen and Wangersky, 1996; Urbani et al., 2005).



490 The decline in the abundance of *R. marina* was not fast nor linear and occasionally abundance increased again, suggesting that  
conditions were still quite favorable for *R. marina* during all KPIs. The resumption of growth might have been due to the cells  
turning to heterotrophy, as some *Rhodomonas* species are known to be mixotrophs (Ballen-Segura et al., 2017). As nutrient  
limitation was most likely not significant, C limitation could be another possible cause for population decline and a switch to  
support growth with heterotrophy. Total dissolved C was high in both *R. marina* and *A. malmogiense* treatments even at the  
3<sup>rd</sup> KPI, but since pH was not measured the relative fractions of different forms of inorganic C are not known. To our  
495 knowledge, the capacity of *A. malmogiense* to use different forms of inorganic C is not known, but many dinoflagellates are  
be able to use bicarbonate (Nimer et al., 1997) suggesting that *A. malmogiense* was likely not C limited. The potential for C  
limitation of *R. marina* is not clear, since the use of different forms of inorganic C by *R. marina* is not known. Some  
*Rhodomonas* species use only free CO<sub>2</sub> (Elzenga et al., 2000) while some also seem to use bicarbonate (Camiro-vargas et al.,  
2005).

500

Optical characteristics of DOM revealed potential sources and consumption patterns in the experiments. Usually fluorescence  
peak T is interpreted as a proxy for bioavailable DOM (Nieto-Cid et al., 2006), but it increased in both treatments together  
with the signals for less labile DOM throughout the DOM release experiment. Increase in protein-like DOM fluorescence has  
been connected to phytoplankton growth during simulated (Stedmon and Markager, 2005) and natural (Suksomjit et al., 2009)  
505 phytoplankton blooms, but bacterial processing can decrease protein-like fluorescence while increasing humic-like  
fluorescence (Romera-Castillo et al., 2011; Yamashita and Tanoue, 2004b). Therefore, simultaneous increase in peaks T and  
C likely occurred because of (1) excess production of protein-like DOM by phytoplankton, (2) production of less labile protein-  
like DOM by phytoplankton, (3) production of protein-like DOM by bacteria or by combinations of these. Not all protein-like  
DOM fractions are equally degradable (Yamashita and Tanoue, 2004a) and some protein-like FDOM can accumulate in the  
510 pelagic environment (Asmala et al., 2018; Yamashita et al., 2017). Production of peak T by bacteria might be due to bacterial  
reworking of initially labile (non-colored) autochthonous DOM into small, UV-absorbing molecules (Asmala et al., 2018;  
Berggren et al., 2009). In the case of *R. marina*, this could possibly result from the bacterial consumption of monosaccharides,  
which *R. marina* can produce in high amounts (Fernandes et al., 2017), as several bacterial species have been shown to produce  
peak T when grown on glucose (Fox et al., 2017).

515

Just like the simultaneous increase of most optical DOM variables in the DOM release experiment, the decrease of most of the  
FDOM variables towards the end of the DOM consumption experiment is surprising, given that bacterial processing of  
phytoplankton-derived DOM is usually connected to increase of FDOM (Romera-Castillo et al., 2011). The high abundance  
of *Pseudorhodobacter* might explain part of this as Rhodobacteraceae have been connected to reduced FDOM intensities when  
520 using dinoflagellate-derived DOM (Tada et al., 2017). Bacteria may also change from net source of protein-like FDOM to a  
net sink as bacterial activity increases (Guillemette and del Giorgio, 2012). This is in agreement with the decreasing peak T  
during the DOM consumption experiment, as the higher temperature used in the DOM consumption experiment may have



525 directly enhanced bacterial activity. Guillemette and del Giorgio (2012) also showed that production of humic-like FDOM increases with increasing BGE, which is in line with the increase of humic-like peak C and HIX concurrently with BGE until day 4, although after that the FDOM signals decreased while BGE did not. The change in DOM processing patterns on day 4, which was suspected to have been caused by the depletion of fresh labile DOM originating from phytoplankton, was interesting also because the production of humic-like DOM should increase when bacteria shift from processing labile DOM to semi-labile DOM (Jørgensen et al., 2015).

530 The overall differences between *A. malmogiense* and *R. marina* are similar to those in a previous study with dinoflagellates *Heterocapsa circularisquama* and *Alexandrium catenella* and a cryptophyte *Rhodomonas ovalis* (Fukuzaki et al., 2014). They observed higher biomass production for the dinoflagellates and higher apparent percentage of net photosynthetic extracellular release for *R. ovalis*. In addition to the inherent species-specific physiological differences between *A. malmogiense* and *R. marina*, some fraction of the different DOM release might be caused by more general traits, such as the size difference between  
535 the species. Higher release of bioavailable DOM from *R. marina* might simply be caused by the smaller size of *R. marina* cells and, therefore, higher passive release of DOC (Bjørnsen, 1988).

Even though both of the phytoplankton species can be assumed to be mixotrophic (Ballen-Segura et al., 2017; Rintala et al., 2007) and phytoplankton can take up DOM in mixed communities (Bronk and Glibert, 1993; Moneta et al., 2014), significant  
540 DOM consumption by phytoplankton during this experiment was unlikely. Uptake of organic N or P would be energetically unlikely at the presence of light and available inorganic N and P. Towards the end of the experiment, if the decline of *R. marina* was caused by C limitation, DOM consumption would have been more likely and an unknown fraction of changes in the properties of DOM could maybe be attributed to reuptake by phytoplankton. However, because there was a shift from increase to decrease of some optical DOM properties between the DOM release experiment and the DOM consumption experiment,  
545 the principal role of *R. marina* was likely still the production of DOM rather than its consumption throughout the DOM release experiment.

Because the observed changes in optical DOM properties seem to be independent of the ratio of bacterial to phytoplankton abundance, the observed changes in DOM characteristics have to arise primarily from the traits of individual phytoplankton  
550 species (rate and type of produced DOM (Fukuzaki et al., 2014)) or bacterial species (rate and type of consumed and produced DOM (Fox et al., 2017; Romera-Castillo et al., 2011)) instead of only from the ratio of producers to consumers. A general conclusion from DOM quality indicators is that *R. marina* produce comparatively more DOM, when normalized to PP, than *A. malmogiense* and that this DOM seems to be more efficiently consumed and altered by bacteria. However, DOC release and the rate of DOC production to PP do not necessarily reflect natural conditions precisely for either phytoplankton species  
555 since the fraction of PP released as DOC from phytoplankton is generally higher in situ than in cultures (Thornton, 2014).



#### 4.2 Response of bacteria to DOC

The higher leucine:thymidine incorporation ratio in *R. marina* treatments indicates that bacteria struggled to get enough C and/or energy from DOM for balanced growth. A likely explanation for this is that the bacterial community in *R. marina* treatments efficiently depleted the readily available labile DOM pool and the stable DOM release from *R. marina* could not  
560 keep up with the demands of the bacterial growth. This idea was supported by much higher BP:PP ratio in *R. marina* treatments during the 2<sup>nd</sup> and the 3<sup>rd</sup> KPI. In the DOM consumption experiment bacteria growing on *R. marina* filtrate invested more in thymidine incorporation (lower leucine:thymidine ratio) than in the DOM release experiment. This was most likely caused by the relaxed resource competition due to dilution of bacterial abundance during the filtration and further suggests that the higher leucine:thymidine ratio in DOM release experiment was caused by intense competition for DOM among bacteria.

565

In general, Bacteroidia, Alphaproteobacteria and Gammaproteobacteria predominated bacterial communities in both DOM release and consumption experiments and the communities reflected those in the phytoplankton cultures indicating that the bacterial communities emerged from the phytoplankton cultures. In the beginning of the DOM consumption experiment in *R. marina*, class Alphaproteobacteria (mostly genus *Pseudorhodobacter*) comprised 82% of the bacterial community, which was  
570 related to the high peak T and high BP. This kind of ‘feast and famine’ growth mode is typical for copiotrophic bacteria (Lauro et al., 2009). Alphaproteobacteria benefit from phytoplankton blooms when there is high concentration of labile DOM available (Allers et al., 2007) and they are efficient in using amino acids (Cottrell and Kirchman, 2000; Gasol et al., 2008). The predicted high production of monosaccharides by *R. marina* (Fernandes et al., 2017) may explain the higher proportions of Alpha- and Gammaproteobacteria in *R. marina* treatments. *Pseudorhodobacter* has been detected also in a previous  
575 mesocosm study with Baltic Sea water (Camarena-Gómez et al., 2018) as well as in Baltic Sea bacterioplankton (Herlemann et al., 2011).

In general, the share of class Bacteroidia (genera *Algoriphagus* and *Polaribacter*) was higher in *A. malmogiense* treatments in both experiments, likely reflecting the more stable and less optically active DOM pool. In the *R. marina* treatments class  
580 Bacteroidia (genus *Flavobacter*) became abundant only after the day 3 congruently with the drop in peak T, implicating that the ratio of labile to semi-labile DOM dropped on day 4 and caused the shift in bacterial community composition. Phylum Bacteroidetes is well known of its capability to degrade high-molecular weight DOM (Cottrell and Kirchman, 2000; Romera-Castillo et al., 2011) with their polysaccharide utilizing enzymes (Grondin et al., 2017). Both *Polaribacter* and *Flavobacterium* are common moderate copiotrophs and detected from phytoplankton blooms (Mühlenbruch et al., 2018; Teeling et al., 2012).  
585 In addition, *Polaribacter* and *Algoriphagus* have been detected in previous Baltic Sea mesocosm studies (Camarena-Gómez et al., 2018; Herlemann et al., 2017). The observed pattern in the bacterial community composition support the interpretation that DOM was more labile in *R. marina* treatments than in *A. malmogiense*. The difference between control treatments and





experimental replicates suggests that phytoplankton-derived DOM, not the growth medium, is the main driver for bacterial community and DOM processing dynamics.

590

Actinobacteria, which were present in *A. malmogiense* treatments, are members of autochthonous bacterioplankton in the Baltic Sea (Riemann et al., 2008) occupying several different niches and thus have likely various different functions in the Baltic Sea food web (Holmfeldt et al., 2009). They have also occurred with dinoflagellates in a previous mesocosm experiment with Baltic Sea water (Camarena-Gómez et al., 2018). In the Baltic Sea, some Actinobacteria are linked to high DOC concentrations and terrestrial DOM close to the land (Holmfeldt et al., 2009) and others are outcompeted by fast-growing copiotrophs when phytoplankton-derived DOM is available (Pérez and Sommaruga, 2006). Possibly, in *A. malmogiense* treatments the presumably less labile DOM allowed them to compete better with the copiotrophic Alpha and Gammaproteobacteria. However, it is also possible that the 0.8 µm filtration in the DOM consumption experiment caused a bias and favored them due to their small size (Hahn et al., 2003).

600

In the DOM consumption experiment, a shift in the preferred substrate for bacterial consumption and a concurrent shift in the bacterial community was obvious in *R. marina* treatments even though DOC concentration was still high after the incubation. This highlights the strong connection between phytoplankton DOM release and bacterial processes. The existing DOM pool explains only part of the mechanisms which structure the bacterial community. The fast flow of <sup>14</sup>C from DIC pool through phytoplankton to DOC pool and bacterial biomass in DOM release experiment supports this statement.

605

In both experiments the final bacterial communities were similar and seemingly unaffected by the addition of seawater bacteria. Most likely the low number of bacteria in the bacterial inoculum could not compete with the high number of pre-existing bacteria. This suggests that the phytoplankton-bacteria communities in the cultures were somewhat stable and resistant to minor introductions of foreign bacteria. This is in line with other studies which have shown stable and predictable bacterial communities associated with certain phytoplankton species (e.g. Schäfer et al. 2002, Sapp et al. 2007, Goecke et al. 2013, Buchan et al. 2014, Krohn-Molt et al. 2017, Mönnich et al. 2020). A variety of mutualistic or algicidal interactions between bacteria and phytoplankton are known (Seymour et al., 2017). Phytoplankton might e.g. affect bacterial community composition by producing certain amino acids (Tada et al., 2017) which may in part explain why the development of the bacterial community in the DOM consumption experiment was connected to peak T. The minor differences in the thymidine and leucine incorporation between the species in the DOM release experiment despite the major differences in PP and DOC processing, and the comparable BGE between species in the DOM consumption experiment, suggest that the bacterial communities, while different in composition, are functionally optimized to grow using the DOC produced by the host phytoplankton.

615



#### 620 4.3 Ecological implications of species-specific DOC dynamics

Recent study connected a dinoflagellate community consisting of *A. malmogiense* and related species to lower BP and distinct bacterial community, compared to communities with common spring bloom diatom species (Camarena-Gómez et al., 2018). The results with *A. malmogiense* support their view that DOC released from some dinoflagellate species may lead to lower efficiency of the microbial loop.

625

When dinoflagellate blooms are not terminated in mass encystment they are expected to lyse in the water column and contribute to pelagic DOC pool (Spilling et al., 2018). Our results indicate that blooms predominated by *A. malmogiense* indeed release high amounts of DOC, but this DOC may not be readily bioavailable for bacteria coinciding with phytoplankton blooms and may, therefore, stay in the pelagic system for longer. High biomass production combined with release of less bioavailable

630

DOC could lead to direct grazing being favored over microbial loop. Thus, a probable long-term effect of *A. malmogiense* predominance in natural communities on C cycling is the accumulation of less bioavailable DOC at the expense of sedimentation and microbial loop. Also, pronounced cycling of C between phytoplankton biomass and DIC pool can be expected, as community respiration was high and dinoflagellates are generally considered to have high respiration rates (Taylor and Pollinger, 1987).

635

Compared to *A. malmogiense*, *R. marina* produces less phytoplankton biomass and the DOC it releases is more bioavailable. Thus, blooms predominated by *R. marina* may favor microbial loop and DOC processing over grazing. Since Alpha and Gammaproteobacteria, which were common in *R. marina* treatments, are heavily grazed by heterotrophic nanoflagellates (Alonso-Sáez et al., 2009) *R. marina* predominance may increase C transfer through microbial loop. In addition, the higher

640 BR in DOM release experiment may indicate that total C fixation is lower during such blooms. *Rhodomonas* species have not traditionally been connected to periods of high DOC release from phytoplankton (Storch and Saunders, 1978) and, according to our results, this might be the result of fast bacterial consumption of DOC released by *Rhodomonas* species. The very fast consumption and transformation of DOC in *R. marina* treatments in the DOM consumption experiment support the assumption of fast DOC depletion in natural *R. marina* predominated blooms.

645

Strong extrapolations of these results to related phytoplankton species or to phytoplankton of similar size should be made with caution, as even much more closely related phytoplankton species may support differing bacterial communities (Grossart et al., 2005) and, consequently, different C cycling dynamics. Instead, we want to highlight the importance of studying C cycling between individual phytoplankton species and related bacterial communities, in order to understand the mixed phytoplankton

650 communities of the natural environments. These community manipulation experiments should also include protozoan grazers as their impact on DOM composition and processing can be significant (Kujawinski et al., 2016). Grazing may, for example, alter bacterial community composition by removing groups which are less resistant to grazing (Alonso-Sáez et al., 2009),



655 enhance DOM production (Strom et al., 1997) and affect the lability of the produced DOM (Fouilland et al., 2014), all of which can be assumed to affect C cycling. Better knowledge on C cycling on species level will help in predicting how the large-scale change in phytoplankton community composition will affect C cycling on ecosystem level.


## Conclusions

Two common phytoplankton species in the Baltic Sea, *A. malmogiense* and *R. marina*, produce DOM with different bioavailability and support distinct bacterial communities specialized in utilizing this specific DOM source. This results in different C cycling patterns: *A. malmogiense* cells circulate more C between DIC and phytoplankton biomass, while producing less labile DOC. *R. marina* releases more labile DOC and relatively more C is thus directed towards the bacterial community. DOC released by *R. marina* is taken up, incorporated, and respired faster than DOC released by *A. malmogiense*. Differences were clear at every level of C cycling: PP, flow of  $^{14}\text{C}$  from DIC to bacterial biomass, optical properties of DOM and the response in the composition and activity of the bacterial community. This experiment supports the view that phytoplankton and bacteria are intimately connected through the rapid bacterial consumption of DOM released by phytoplankton, and that this connection explains bacterioplankton dynamics better than the composition of the ambient DOM pool. An experimental approach based on monocultures was necessary to quantify these differences in C pathways. To better understand C cycling in a natural environment, it may be beneficial to see also natural pelagic microbial communities as collections of various linked phytoplankton-bacteria associations with distinctive C cycling patterns.



## Appendices

### 670 Appendix A: Additional details of the experimental setup



675 **Figure A1. Schematic description of the production line of the DOM release experiment.** The schematic starts at the top from the  
 “Production line mix” which consisted of phytoplankton culture (90% vol.) and inoculum of sea water bacteria (explained in the  
 methods and in Fig. 1a). This was divided into five 10 ml aliquots in 20 ml scintillation vials, which were spiked with 125 µL of 23.43  
 680 µCi mL<sup>-1</sup> 14C-NaHCO<sub>3</sub> and incubated in light for 0 to 24 h. At 0, 4, 8 and 12 h one set of incubation vials was divided in two. One  
 half was terminated with 0.1% formaldehyde for 14C-NaHCO<sub>3</sub>-incorporation measurement (i.e. primary production). 24 h  
 acidification with HCl was used to remove remaining 14C-NaHCO<sub>3</sub> prior to scintillation counting. The other half was filtered to  
 remove majority of 14C-labeled phytoplankton and bacterial cells and mixed (50:50 vol.) with non-spiked production line mix  
 (termed “mix” in the lower left part of the schematic). This new mixture was then spiked with either only 3H-thymidine or both 3H-  
 685 thymidine and 14C-leucine and incubated for 4 h. The mixture spiked with 3H-thymidine was used to measure incorporation of  
 14C-labeled DOC originating from phytoplankton into bacterial biomass (3H-thymidine incorporation was measured as a control  
 for bacterial activity). The mixture spiked with 3H-thymidine and 14C-leucine was used to measure incorporation rate of both  
 radioisotope tracers and, subsequently, bacterial production based on both tracers. These mixtures were terminated with  
 trichloroacetic acid (TCA, final concentration: 5%) and measured according the centrifugation method (Smith and Azam, 1992).  
 Instagel Plus (PerkinElmer) was used as the scintillation cocktail and liquid scintillation counting was done with Wallac 1414 LSC.

690 **Table A1. Measured variables at each stage of both parts of the experiment (DOM release experiment and DOM line experiment).**  
 The stages of the experiments are explained in the methods and in the Fig. 1. In short, in the DOM release experiment monitoring  
 of culture growth refers to the total time the cultures were grown. During this time three shorter key point incubations (KPI)  
 were conducted. In the DOM consumption experiment monitoring of culture growth refers to the period before the experimental  
 incubation.

DOM release experiment


DOM consumption experiment



Measurement	Monitoring of culture growth	KPI: Phytoplankton culture only	KPI: Production line mix	KPI: DOM line mix	Monitoring of culture growth	Experimental incubation (7 days)
Phytoplankton abundance	1-3 times per week				1-3 times per week	
Bacterial abundance				0, 4, 8 and 12 h		Daily
Bacterial community composition		At the start of each KPI (also from sea water)	At the start of each KPI			Daily (also from sea water at the start of the incubation)
<sup>14</sup> C-incorporation			0, 4, 8, 12 and 24 h			
<sup>3</sup> H and <sup>14</sup> C-incorporation			0, 4, 8 and 12 h			Daily
Bacterial respiration						Daily
Nutrients (NO <sup>3-</sup> (including NO <sup>2-</sup> ) and PO <sub>4</sub> <sup>3-</sup> )		At the start and end of each KPI (same time as from DOM line samples)		0 and 12 h		
Particulate organic C and N, and Chl <i>a</i>		At the start of each KPI				
Dissolved organic C and total dissolved N				0, 4, 8 and 12 h		Daily
Optical properties of dissolved organic matter	1-3 times per week			0, 4, 8 and 12 h		Daily




## Appendix B: Variables which were measured but not included in the detailed analysis



695

**Figure B1.** CDOM and FDOM variables in phytoplankton cultures during the DOM release experiment. (a-h) Absorption coefficients at different wavelengths. (i, j, l) Spectral slopes between different wavelength ranges. (k) Ratio of spectral slopes between 275-295 nm and 350-400 nm (Helms et al., 2008). (n) Peak M, marine humic-like DOM (Coble, 1996). (p) FI, fluorescence index (Mcknight et al., 2001). (q) HIX, humification index (Zsolnay et al., 1999). (r) BIX, biological index (Huguet et al., 2009). Vertical lines from left to right in each panel mark the times of 1st, 2nd and 3rd KPI. Error bars indicate one standard deviation



700 Figure B2. Those CDOM and FDOM variables in the DOM consumption experiment which were measured but not included in the  
 more detailed analysis. (a-g) Absorption coefficients at different wavelengths. (h, j) Spectral slopes between different wavelength  
 ranges. (i) Ratio of spectral slopes between 275-295 nm and 350-400 nm (Helms et al., 2008). (k) Peak M, marine humic-like DOM  
 (Coble, 1996). (l) FI, fluorescence index (Mcknight et al., 2001). (m) BIX, biological index (Huguet et al., 2009). (n) SUVA<sub>254</sub>, DOC-  
 705 normalized absorbance at 254 nm (Weishaar et al., 2003). Lines A, B and C mark different replicates (culture filtrate + sea water  
 bacteria) and ctrl marks the control (F/2 + sea water bacteria).




Figure B3.  $^{14}\text{C}$ -activity originating from  $^{14}\text{C}\text{-NaHCO}_3$  in DOC pool in the filtrates from light (a) and dark (b) primary production incubation samples.  $^{14}\text{C}$ -activity originating from  $^{14}\text{C}\text{-NaHCO}_3$  in bacterial biomass after 4 h dark incubation of filtrate from light (c) and dark (d) primary production incubation samples amended with non-spiked cultures. Number above the boxplots shows the number of measurements. Lower and upper hinges of boxes mark the first and third quartiles, respectively, while the whiskers extend to the lowest and the highest value within 1.5 times the interquartile range. White boxes = *A. malmogiense*, grey boxes = *R. marina*.

710





715

**Table B1. Mean±1SD of additional variables at each KPI in the DOM release experiment. S ratio = Ratio of spectral slopes between 275-295 nm and 350-400 nm (Helms et al., 2008). Peak M = marine humic-like DOM (Coble, 1996). FI = fluorescence index (Mcknight et al., 2001). BIX = biological index (Huguet et al., 2009).**

Species	<i>A. malmogiense</i>			<i>R. marina</i>		
	1 <sup>st</sup>	2 <sup>nd</sup>	3 <sup>rd</sup>	1 <sup>st</sup>	2 <sup>nd</sup>	3 <sup>rd</sup>
Low chlorophyll <i>a</i> fluorescence cells (cells mL <sup>-1</sup> )	1813 ± 296	4770 ± 1288	6868 ± 609	0	0	0
Bacteria:phytoplankton	569.311 ± 5.259	329.050 ± 50.469	181.504 ± 59.284	637.657 ± 11.579	428.095 ± 17.734	399.943 ± 13.467
chlorophyll <i>a</i> (µmol L <sup>-1</sup> )	0.149 ± 0.046	0.746 ± 0.016	1.245 ± 0.154	0.027 ± 0.008	0.174 ± 0.025	0.149 ± 0.013
POC (µmol L <sup>-1</sup> )	67.735 ± 9.371	406.580 ± 129.780	723.624 ± 167.338	38.865 ± 8.774	247.752 ± 13.753	306.471 ± 19.531
PON (µmol L <sup>-1</sup> )	10.430 ± 1.613	67.271 ± 18.906	117.577 ± 15.612	5.384 ± 1.547	46.558 ± 2.055	58.427 ± 4.350
DOC:phytoplankton (pmol cell <sup>-1</sup> )	44.060 ± 7.097	17.842 ± 0.978	12.474 ± 1.335	15.930 ± 4.367	9.492 ± 1.651	14.976 ± 0.618
NO <sub>3</sub> -N, unfiltered (µmol L-1)	549.003 ± 165.250	422.896 ± 104.020	158.170 ± 95.621	658.442 ± 183.104	424.477 ± 43.252	401.662 ± 36.279
NO <sub>3</sub> <sup>-</sup> -N, 0.8 µm filtered (µmol L-1)	743.355 ± 17.682	383.821 ± 91.189	95.527 ± 99.425	765.352 ± 11.973	406.647 ± 49.715	362.545 ± 29.791
PO <sub>4</sub> <sup>3-</sup> -P, unfiltered (µmol L-1)	18.695 ± 1.107	10.738 ± 1.676	12.453 ± 1.269	21.559 ± 4.673	16.462 ± 0.912	15.983 ± 1.020
PO <sub>4</sub> <sup>3-</sup> -P, 0.8 µm filtered (µmol L-1)	18.643 ± 1.415	0.224 ± 0.156	0.335 ± 0.221	23.752 ± 0.591	8.690 ± 0.642	7.861 ± 1.169



DIC:NO <sub>3</sub> <sup>-</sup> -N, unfiltered	0.404 ± 1.631 ± 21.066 ± 19.435	0.761 ± 1.355 ± 1.804 ± 0.200
	0.024 0.639	0.035 0.218
DIC:PO <sub>4</sub> <sup>3-</sup> -P, unfiltered	16.194 ± 5239.512 ± 5225.060 ± 24.503 ± 62.920 ± 83.907 ±	
	1.788 5870.309 5813.222	0.832 7.155 11.968
NO <sub>3</sub> <sup>-</sup> -N:PO <sub>4</sub> <sup>3-</sup> -P, unfiltered	39.978 ± 2662.517 ± 404.746 ± 32.229 ± 46.684 ± 46.693 ±	
	2.086 2107.333 330.627	0.535 2.393 6.597
Gross primary production:net primary production	3.309 ± 4.061 ± 3.742 ± 0.059	1.904 ± 2.786 ± 2.776 ± 0.316
	0.497 0.046	0.224 0.505
Community respiration:gross primary production	0.277 ± 0.337 ± 0.316 ± 0.004	0.053 ± 0.217 ± 0.220 ± 0.043
	0.045 0.003	0.062 0.061
leucine incorporation:thymidine incorporation	43.833 ± 16.341 ±	57.234 ± 25.009 ± 31.403 ±
	7.075 1.090 19.264 ± 1.347	21.333 0.332 0.936
Absorption coefficient at 275 nm (m <sup>-1</sup> )	3.344 ± 5.690 ± 7.600 ± 0.786	3.920 ± 6.640 ± 7.830 ± 0.188
	0.096 0.388	0.226 0.284
Absorption coefficient at 295 nm (m <sup>-1</sup> )	2.720 ± 4.428 ± 5.618 ± 0.582	2.980 ± 4.620 ± 5.283 ± 0.137
	0.088 0.254	0.088 0.166
Absorption coefficient at 300 nm (m <sup>-1</sup> )	2.572 ± 4.025 ± 5.023 ± 0.561	2.754 ± 4.217 ± 4.784 ± 0.124
	0.071 0.230	0.094 0.166
Absorption coefficient at 350 nm (m <sup>-1</sup> )	0.557 ± 1.367 ± 2.001 ± 0.463	0.336 ± 1.612 ± 1.833 ± 0.091
	0.030 0.101	0.051 0.109
Absorption coefficient at 375 nm (m <sup>-1</sup> )	0.437 ± 0.888 ± 1.267 ± 0.225	0.187 ± 1.089 ± 1.243 ± 0.082
	0.046 0.082	0.025 0.087
Absorption coefficient at 400 nm (m <sup>-1</sup> )	0.360 ± 0.629 ± 0.825 ± 0.109	0.120 ± 0.715 ± 0.830 ± 0.088
	0.052 0.072	0.022 0.068



Absorption coefficient at 440 nm (m <sup>-1</sup> )	0.278 ± 0.374 ± 0.046 0.080	± 0.465 ± 0.071	0.014 ± 0.437 ± 0.025 0.058	± 0.499 ± 0.087
Spectral slope between 300-650 nm	23.229 ± 18.158 ± 4.315 1.370	± 17.010 ± 0.349	41.557 ± 15.955 ± 1.370 1.155	± 15.766 ± 1.448
Spectral slope between 350-400 nm	0.009 ± 0.016 ± 0.001 0.001	± 0.018 ± 0.002	0.026 ± 0.017 ± 0.002 0.001	± 0.017 ± 0.001
S ratio	1.074 ± 0.824 ± 0.084 0.064	± 0.871 ± 0.087	0.509 ± 1.032 ± 0.020 0.081	± 1.141 ± 0.054
DOC-normalized absorbance at 254 nm (m <sup>2</sup> g <sup>-1</sup> )	0.969 ± 0.638 ± 0.141 0.074	± 0.563 ± 0.030	1.058 ± 1.091 ± 0.057 0.031	± 1.079 ± 0.127
Peak M (R.U.)	0.011 ± 0.078 ± 0.001 0.010	± 0.101 ± 0.013	0.033 ± 0.097 ± 0.011 0.007	± 0.124 ± 0.008
FI	1.837 ± 1.713 ± 0.367 0.104	± 1.853 ± 0.061	1.801 ± 2.062 ± 0.555 0.062	± 2.095 ± 0.034
BIX	0.913 ± 0.772 ± 0.121 0.024	± 0.894 ± 0.024	0.836 ± 0.718 ± 0.086 0.042	± 0.680 ± 0.022

### Appendix C: Results of statistical tests

**Table C1. Welch-ANOVA results of the DOM release experiment. Each line presents the test of difference between the phytoplankton species for the given variable at the given KPI.**

Variable	KPI	<i>F</i>	df1	df2	<i>p</i>	ω <sup>2</sup>
GPP:phytop.	1	28.549	1	1.0071	0.117	0.85
	2	552.26	1	2.1329	0.001	0.99
	3	244.08	1	2.7835	<0.001	0.98
Bacterial abundance	1	45.304	1	13.272	<0.001	0.65



	2	<0.001	1	21.818	0.978	<0
	3	21.407	1	2.7879	0.022	0.58
Bacteria:phytop.	1	239.57	1	8.2775	<0.001	0.92
	2	12.756	1	20.362	0.002	0.33
	3	321.87	1	12.884	<0.001	0.96
Thymidine incorporation	1	62.749	1	19.94	<0.001	0.72
	2	39.464	1	17.64	<0.001	0.62
	3	31.385	1	2.1297	0.026	0.67
Leucine incorporation	1	24.025	1	21.648	<0.001	0.49
	2	1.0029	1	17.246	0.330	<0.001
	3	9.7471	1	2.2395	0.077	0.35
Leucine:thymidine incorporation ratio	1	1.6961	1	11.854	0.218	0.02
	2	77.527	1	14.929	<0.001	0.76
	3	45.127	1	12.946	<0.001	0.65
Community respiration	1	16.681	1	1.0094	0.151	0.78
	2	483.87	1	2.2395	0.001	0.99
	3	207.59	1	2.9805	<0.001	0.97
S <sub>275-295</sub>	1	22.482	1	14.652	<0.001	0.47
	2	40.022	1	12.081	<0.001	0.62
	3	345.94	1	15.087	<0.001	0.93
Peak C	1	7.0792	1	15.902	0.017	0.20



	2	394.08	1	17.46	<0.001	0.94
	3	347.17	1	18.884	<0.001	0.94
HIX	1	6.7755	1	21.858	0.016	0.19
	2	279.75	1	14.952	<0.001	0.92
	3	242.71	1	13.075	<0.001	0.91

720

**Table C2. Welch-ANOVA results of  $^{14}\text{C}$  flow (Table 1). Each line presents the test of difference between the phytoplankton species for the given ratio at the given KPI.**

Ratio	KPI	<i>F</i>	df1	df2	<i>p</i>	$\omega^2$
PP:DIC	1	18.907	1	2.1995	0.041	0.75
	2	2283.2	1	2.1064	<0.001	>0.99
	3	867.2	1	3.7377	<0.001	>0.99
DOC:PP	1	17.073	1	9.407	0.002	0.57
	2	396.06	1	7.5657	<0.001	0.97
	3	99.106	1	5.2151	<0.001	0.89
Bacteria:DOC	1	NA				
	2	9.396	1	8.1184	0.015	0.41
	3	9.6009	1	5.0832	0.026	0.42
Bacteria:PP	1	NA				
	2	72.008	1	5.0338	<0.001	0.86
	3	43.611	1	5.0487	0.001	0.78



### Data availability

725 Raw reads are deposited in the Sequence Read Archive of National Center for Biotechnology Information under BioProject  
accession number PRJNA647035. Other data have been submitted to PANGAEA on 2021-08-13, but no DOI is available yet.  
For review purposes the data is available on request from the corresponding author.

### Author contribution

730 SE, TT and HK designed the experiment. SE conducted the experiment. EER processed bacterial community composition  
data. EA processed DOM optics data. SE processed remaining data. SE wrote the manuscript with contributions from the other  
authors.

### Competing interests

The authors declare that they have no conflict of interest.

### Acknowledgements

735 The study was funded by Walter and Andrée de Nottbeck Foundation (WANS), Onni Talas Foundation, Maa- ja vesiteknii-  
tuki ry and Societas Pro Fauna et Flora Fennica. EA was supported by the Academy of Finland (Grant No. 309748). TT was  
funded by WANS and by the Swedish Cultural Heritage Foundation. We thank Timo Tikka (funded by Waldemar von  
Frenckell Foundation) for help in establishing the experiments, Tvärminne Zoological Station for producing facilities, practical  
aid and analyses, and Heidi Hällfors for aid and expertise related to phytoplankton cultures. We thank anonymous reviewers  
740 for considerably improving the manuscript. The study utilized the Finnish Environment Institute Marine Research Centre  
marine research infrastructure as a part of the national FINMARI RI consortium.

### References

- Allers, E., Gómez-Consarnau, L., Pinhassi, J., Gasol, J. M., Šimek, K. and Pernthaler, J.: Response of Alteromonadaceae and  
Rhodobacteriaceae to glucose and phosphorus manipulation in marine mesocosms, *Environ. Microbiol.*, 9(10), 2417–2429,  
745 doi:10.1111/j.1462-2920.2007.01360.x, 2007.
- Alonso-Sáez, L., Unanue, M., Latatu, A., Azua, I., Ayo, B., Artolozaga, I. and Iriberry, J.: Changes in marine prokaryotic  
community induced by varying types of dissolved organic matter and subsequent grazing pressure, *J. Plankton Res.*, 31(11),  
1373–1383, doi:10.1093/plankt/fbp081, 2009.
- Amin, S. A., Green, D. H., Hart, M. C., Küpper, F. C., Sunda, W. G. and Carrano, C. J.: Photolysis of iron – siderophore



- 750 chelates promotes bacterial – algal mutualism, *Proc. Natl. Acad. Sci.*, 106(40), 17071–17076, doi:10.1073/pnas.0905512106, 2009.
- Anderson, M. J.: A new method for non-parametric multivariate analysis of variance, *Austral Ecol.*, 26, 32–46, doi:10.1111/j.1442-9993.2001.01070.pp.x, 2001.
- Asmala, E., Haraguchi, L., Markager, S., Massicotte, P., Riemann, B., Staehr, P. A. and Carstensen, J.: Eutrophication Leads  
755 to Accumulation of Recalcitrant Autochthonous Organic Matter in Coastal Environment, *Global Biogeochem. Cycles*, 32(11), 1673–1687, doi:10.1029/2017GB005848, 2018.
- Azam, F., Fenchel, T. M., Field, J. G., Gray, J. S., Meyer-Reil, L. A. and Thingstad, F.: The ecological role of water-column microbes in the sea, *Mar. Ecol. Prog. Ser.*, 10(January), 257–263, doi:10.3354/meps010257, 1983.
- Ballen-Segura, M., Felip, M. and Catalan, J.: Some mixotrophic flagellate species selectively graze on archaea, *Appl. Environ. Microbiol.*, 83(2), 1–11, doi:10.1128/AEM.02317-16, 2017.
- 760 Becker, J. W., Berube, P. M., Follett, C. L., Waterbury, J. B., Chisholm, S. W., DeLong, E. F. and Repeta, D. J.: Closely related phytoplankton species produce similar suites of dissolved organic matter, *Front. Microbiol.*, 5, 1–14, doi:10.3389/fmicb.2014.00111, 2014.
- Berggren, M., Laudon, H. and Jansson, M.: Aging of allochthonous organic carbon regulates bacterial production in  
765 unproductive boreal lakes, *Limnol. Oceanogr.*, 54(4), 1333–1342, doi:10.4319/lo.2009.54.4.1333, 2009.
- Bjørnsen, P. K.: Phytoplankton exudation of organic matter: Why do healthy cells do it?, *Limnol. Oceanogr.*, 33(1), 151–154, doi:10.4319/lo.1988.33.1.0151, 1988.
- Bronk, D. A. and Glibert, P. M.: Application of a <sup>15</sup>N tracer method to the study of dissolved organic nitrogen uptake during spring and summer in Chesapeake Bay, *Mar. Biol.*, 115, 501–508, 1993.
- 770 Buchan, A., LeClerc, G. R., Gulvik, C. A. and González, J. M.: Master recyclers : features and functions of bacteria associated with phytoplankton blooms, *Nat. Rev. Microbiol.*, 12, 686–698, doi:10.1038/nrmicro3326, 2014.
- Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A. and Holmes, S. P.: DADA2 : High-resolution sample inference from Illumina amplicon data, *Nat. Methods*, 13(7), 581–583, doi:10.1038/nmeth.3869, 2016.
- Camarena-Gómez, M. T., Lipsewers, T., Piiparinen, J., Eronen-Rasimus, E., Perez-Quemaliños, D., Hoikkala, L., Sobrino, C.  
775 and Spilling, K.: Shifts in phytoplankton community structure modify bacterial production , abundance and community composition, *Aquat. Microb. Ecol.*, 81(March), 149–170, doi:10.3354/ame01868, 2018.
- Camiro-vargas, T. K., Hernández-Ayón, J. M., Valenzuela-espinoza, E., Delgadillo-hinojosa, F. and Ramón, C.-M.: Dissolved inorganic carbon uptake by *Rhodomonas* sp . and *Isochrysis* aff . *galbana* determined by a potentiometric technique, *Aquac. Eng.*, 33, 83–95, doi:10.1016/j.aquaeng.2004.10.001, 2005.
- 780 Chen, W. and Wangersky, P. J.: Production of dissolved organic carbon in phytoplankton cultures as measured by high-temperature catalytic oxidation and ultraviolet photo-oxidation methods, *J. Plankton Res.*, 18(7), 1201–1211, 1996.
- Christie-Oleza, J. A., Sousoni, D., Lloyd, M., Armengaud, J. and Scanlan, D. J.: Nutrient recycling facilitates long-term stability of marine microbial phototroph-heterotroph interactions, *Nat. Microbiol.*, 2, 1–10, doi:10.1038/nmicrobiol.2017.100,



- 2017.
- 785 Coble, P. G.: Characterization of marine and terrestrial DOM in the seawater using exciting-emission matrix spectroscopy, *Mar. Chem.*, 51, 325–346, doi:10.1016/0304-4203(95)00062-3, 1996.
- Cottrell, M. T. and Kirchman, D. L.: Natural Assemblages of Marine Proteobacteria and Members of the Cytophaga-Flavobacter Cluster Consuming Low- and High- Molecular-Weight Dissolved Organic Matter, *Appl. Environ. Microbiol.*, 66(4), 1692–1697, 2000.
- 790 Croft, M. T., Lawrence, A. D., Raux-Deery, E., Warren, M. J. and Smith, A. G.: Algae acquire vitamin B12 through a symbiotic relationship with bacteria, *Nature*, 438(7064), 90–93, doi:10.1038/nature04056, 2005.
- Ducklow, H. W. and Carlson, C. A.: *Oceanic Bacterial Productivity*, in *Advances in Microbial Ecology*, Plenum Press, New York, NY., 1992.
- Elzenga, J. T. M., Prins, H. B. A. and Stefels, J.: The role of extracellular carbonic anhydrase activity in inorganic carbon  
795 utilization of *Phaeocystis globosa* (Prymnesiophyceae): A comparison with other marine algae using the isotopic disequilibrium technique, *Limnol. Oceanogr.*, 45(2), 372–380, 2000.
- Emerson, S. R. and Hedges, J. I.: *Chemical Oceanography and the Marine Carbon Cycle*, Cambridge University Press, New York, NY., 2008.
- Engel, A., Thoms, S., Riebesell, U., Rochelle-Newall, E. and Zondervan, I.: Polysaccharide aggregation as a potential sink of  
800 marine dissolved organic carbon, *Nature*, 428(6986), 929–932, doi:10.1038/nature02453, 2004.
- Fernandes, T., Fernandes, I., Andrade, C. A. P., Ferreira, A. and Cordeiro, N.: Marine microalgae monosaccharide fluctuations as a stress response to nutrients inputs, *Algal Res.*, 24(February), 340–346, doi:10.1016/j.algal.2017.04.023, 2017.
- Fouilland, E., Tolosa, I., Bonnet, D., Bouvier, C., Bouvier, T., Bouvy, M., Got, P., Le Floc’h, E., Mostajir, B., Roques, C., Sempéré, R., Sime-Ngando, T. and Vidussi, F.: Bacterial carbon dependence on freshly produced phytoplankton exudates  
805 under different nutrient availability and grazing pressure conditions in coastal marine waters, *FEMS Microbiol. Ecol.*, 87(3), 757–769, doi:10.1111/1574-6941.12262, 2014.
- Fox, B. G., Thorn, R. M. S., Anesio, A. M. and Reynolds, D. M.: The in situ bacterial production of fluorescent organic matter; an investigation at a species level, *Water Res.*, 125, 350–359, doi:10.1016/j.watres.2017.08.040, 2017.
- Fukuzaki, K., Imai, I., Fukushima, K., Ishii, K. I., Sawayama, S. and Yoshioka, T.: Fluorescent characteristics of dissolved  
810 organic matter produced by bloom-forming coastal phytoplankton, *J. Plankton Res.*, 36(3), 685–694, doi:10.1093/plankt/fbu015, 2014.
- Gasol, J. M., Pinhassi, J., Alonso-Sáez, L., Ducklow, H., Herndl, G. J., Koblížek, M., Labrenz, M., Luo, Y., Morán, X. A. G., Reinthaler, T. and Simon, M.: Towards a better understanding of microbial carbon flux in the sea, *Aquat. Microb. Ecol.*, 53(September), 21–38, doi:10.3354/ame01230, 2008.
- 815 Goecke, F., Thiel, V., Wiese, J., Labes, A. and Imhoff, J. F.: Algae as an important environment for bacteria – phylogenetic relationships among new bacterial species isolated from algae, *Phycologia*, 52(1), 14–24, doi:10.2216/12, 2013.
- Grasshoff, K., Kremling, K. and Ehrhardt, M.: *Methods of Seawater Analysis*, 3rd ed., Wiley-VCH, Weinheim., 1999.





- Grondin, J. M., Tamura, K., Déjean, G., Abbott, D. W. and Brumer, H.: Polysaccharide Utilization Loci : Fueling Microbial Communities, *J. Bacteriol.*, 199(15), 1–15, 2017.
- 820 Grossart, H., Levold, F., Allgaier, M., Simon, M. and Brinkhoff, T.: Marine diatom species harbour distinct bacterial communities, *Environ. Microbiol.*, 7(6), 860–873, doi:10.1111/j.1462-2920.2005.00759.x, 2005.
- Guillemette, F. and del Giorgio, P. A.: Simultaneous consumption and production of fluorescent dissolved organic matter by lake bacterioplankton, *Environ. Microbiol.*, 14(6), 1432–1443, doi:10.1111/j.1462-2920.2012.02728.x, 2012.
- Hahn, M. W., Lünsdorf, H., Wu, Q., Schauer, M., Höfle, M. G., Boenigk, J. and Stadler, P.: Isolation of Novel  
825 Ultramicrobacteria Classified as Actinobacteria from Five Freshwater Habitats in Europe and Asia, *Appl. Environ. Microbiol.*, 69(3), 1442–1451, doi:10.1128/AEM.69.3.1442, 2003.
- Hansell, D. A., Carlson, C. A., Repeta, D. J. and Schlitzer, R.: Dissolved organic matter in the ocean: a controversy stimulates new insights, *Oceanography*, 22(4), 202–211, 2009.
- Haraguchi, L., Asmala, E., Jakobsen, H. H. and Carstensen, J.: Composition of natural phytoplankton community has minor  
830 effects on autochthonous dissolved organic matter characteristics, *Mar. Biol. Res.*, doi:10.1080/17451000.2019.1662449, 2019.
- Hedges, J. I.: Global biogeochemical cycles: progress and problems, *Mar. Chem.*, 39, 67–93, 1992.
- Helms, J. R., Stubbins, A., Ritchie, J. D., Minor, E. C., Kieber, D. J. and Mopper, K.: Absorption spectral slopes and slope ratios as indicators of molecular weight, source, and photobleaching of chromophoric dissolved organic matter, *Limnol.*  
835 *Oceanogr.*, 53(3), 955–969, doi:10.4319/lo.2009.54.3.1023, 2008.
- Herlemann, D. P. R., Labrenz, M., Jürgens, K., Bertilsson, S., Waniek, J. J. and Andersson, A. F.: Transitions in bacterial communities along the 2000 km salinity gradient of the Baltic Sea, *ISME J.*, 5, 1571–1579, doi:10.1038/ismej.2011.41, 2011.
- Herlemann, D. P. R., Manecki, M., Dittmar, T. and Jürgens, K.: Differential responses of marine, mesohaline and oligohaline bacterial communities to the addition of terrigenous carbon, *Environ. Microbiol.*, 19(8), 3098–3117, doi:10.1111/1462-  
840 2920.13784, 2017.
- Holmfeldt, K., Dziallas, C., Titelman, J., Pohlmann, K., Grossart, H. P. and Riemann, L.: Diversity and abundance of freshwater Actinobacteria along environmental gradients in the brackish northern Baltic Sea, *Environ. Microbiol.*, 11(8), 2042–2054, doi:10.1111/j.1462-2920.2009.01925.x, 2009.
- Huguet, A., Vacher, L., Relexans, S., Saubusse, S., Froidefond, J. M. and Parlanti, E.: Properties of fluorescent dissolved  
845 organic matter in the Gironde Estuary, *Org. Geochem.*, 40(6), 706–719, doi:10.1016/j.orggeochem.2009.03.002, 2009.
- Jiao, N., Herndl, G. J., Hansell, D. A., Benner, R., Kattner, G., Wilhelm, S. W., Kirchman, D. L., Weinbauer, M. G., Luo, T., Chen, F. and Azam, F.: Microbial production of recalcitrant dissolved organic matter: Long-term carbon storage in the global ocean, *Nat. Rev. Microbiol.*, 8(8), 593–599, doi:10.1038/nrmicro2386, 2010.
- Jørgensen, L., Stedmon, C. A., Kaartokallio, H., Middelboe, M. and Thomas, D. N.: Changes in the composition and  
850 bioavailability of dissolved organic matter during sea ice formation, *Limnol. Oceanogr.*, 60, 817–830, doi:10.1002/lno.10058, 2015.



- Kawasaki, N. and Benner, R.: Bacterial release of dissolved organic matter during cell growth and decline: Molecular origin and composition, *Limnol. Oceanogr.*, 51(5), 2170–2180, doi:10.4319/lo.2006.51.5.2170, 2006.
- Kinsey, J. D., Corradino, G., Ziervogel, K., Schnetzer, A. and Osburn, C. L.: Formation of Chromophoric Dissolved Organic Matter by Bacterial Degradation of Phytoplankton-Derived Aggregates, *Front. Mar. Sci.*, 4(January), 1–16, doi:10.3389/fmars.2017.00430, 2018.
- Klais, R., Tamminen, T., Kremp, A., Spilling, K. and Olli, K.: Decadal-scale changes of Dinoflagellates and Diatoms in the Anomalous Baltic Sea spring bloom, *PLoS One*, 6(6), e21567, doi:10.1371/journal.pone.0021567, 2011.
- Klindworth, A., Pruesse, E., Schweer, T., Peplies, J., Quast, C., Horn, M. and Glöckner, F. O.: Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-generation sequencing-based diversity studies, *Nucleic Acids Res.*, 41(1), 1–11, doi:10.1093/nar/gks808, 2013.
- Kremp, A. and Heiskanen, A. S.: Sexuality and cyst formation of the spring-bloom dinoflagellate *Scrippsiella hangoei* in the coastal northern Baltic Sea, *Mar. Biol.*, 134(4), 771–777, doi:10.1007/s002270050594, 1999.
- Kremp, A., Rengefors, K. and Montresor, M.: Species-specific encystment patterns in three Baltic cold-water dinoflagellates: The role of multiple cues in resting cyst formation, *Limnol. Oceanogr.*, 54(4), 1125–1138, doi:10.4319/lo.2009.54.4.1125, 2009.
- Krohn-Molt, I., Alawi, M., Förstner, K. U., Wiegandt, A., Burkhardt, L., Indenbirken, D., Thieß, M., Grundhoff, A., Kehr, J., Tholey, A. and Streit, W. R.: Insights into Microalga and bacteria interactions of selected phycosphere biofilms using metagenomic, transcriptomic, and proteomic approaches, *Front. Microbiol.*, 8(OCT), 1–14, doi:10.3389/fmicb.2017.01941, 2017.
- Kujawinski, E. B.: The Impact of Microbial Metabolism on Marine Dissolved Organic Matter, *Ann. Rev. Mar. Sci.*, 3(1), 567–599, doi:10.1146/annurev-marine-120308-081003, 2011.
- Kujawinski, E. B., Longnecker, K., Barott, K. L., Weber, R. J. M. and Kido Soule, M. C.: Microbial community structure affects marine dissolved organic matter composition, *Front. Mar. Sci.*, 3(APR), 1–15, doi:10.3389/fmars.2016.00045, 2016.
- Kuparinen, J.: Development of bacterioplankton during winter and early spring at the entrance to the Gulf of Finland, Baltic Sea, *Int. Vereinigung für Theor. und Angew. Limnol. Verhandlungen*, 23, 1869–1878, 1988.
- Lauro, F. M., McDougald, D., Thomas, T., Williams, T. J., Egan, S., Rice, S., Demaree, M. Z., Ting, L., Ertan, H., Johnson, J., Ferreria, S., Lapidus, A., Anderson, I., Kyrpides, N., Munk, A. C., Detter, C., Han, C. S., Brown, M. V., Robb, F. T., Kjelleberg, S. and Cavicchioli, R.: The genomic basis of trophic strategy in marine bacteria, *Proc. Natl. Acad. Sci.*, 106(37), 15527–15533, doi:10.1073/pnas.0903507106, 2009.
- Li, W. K. W., Mclaughlin, F. A., Lovejoy, C. and Carmack, E. C.: Smallest Algae Thrive As the Arctic Ocean Freshens, *Science* (80-. ), 326, 539, doi:10.1126/science.1179798, 2009.
- Mari, X., Passow, U., Migon, C., Burd, A. B. and Legendre, L.: Transparent exopolymer particles: Effects on carbon cycling in the ocean, *Prog. Oceanogr.*, 151, 13–37, doi:10.1016/j.pocean.2016.11.002, 2017.
- Martin, M.: Cutadapt removes adapter sequences from high-throughput sequencing reads, *EMBnet.journal*, 17.1, 10–12, 2011.



- Massicotte, P.: eemR: Tools for Pre-Processing Emission-Excitation-Matrix (EEM) Fluorescence Data. R package version 0.1.5.9000. Retrieved from <https://github.com/PMassicotte/eemR>, 2016.
- Mcknight, D. M., Boyer, E. W., Westerhoff, P. K., Doran, P. T., Kulbe, T. and Andersen, D. T.: Spectrofluorometric characterization of dissolved organic matter for indication of precursor organic material and aromaticity, , 46(1), 38–48, 2001.
- 890 McMurdie, P. J. and Holmes, S.: phyloseq : An R Package for Reproducible Interactive Analysis and Graphics of Microbiome Census Data, PLoS One, 8(4), e61217, doi:10.1371/journal.pone.0061217, 2013.
- Moneta, A., Veuger, B., van Rijswijk, P., Meysman, F., Soetaert, K. and Middelburg, J. J.: Dissolved inorganic and organic nitrogen uptake in the coastal North Sea : A seasonal study, Estuar. Coast. Shelf Sci., 147, 78–86, doi:10.1016/j.ecss.2014.05.022, 2014.
- 895 Mönnich, J., Tebben, J., Bergemann, J., Case, R., Wolhrab, S. and Harder, T.: Niche-based assembly of bacterial consortia on the diatom *Thalassiosira rotula* is stable and reproducible, ISME J., 14, 1614–1625, doi:10.1038/s41396-020-0631-5, 2020.
- Morris, R. M., Rappé, M. S., Connon, S. A., Vergin, K. L., Siebold, W. A., Carlson, C. A. and Giovannoni, S. J.: SAR11 clade dominates ocean surface bacterioplankton communities, Nature, 420(December), 806–810, doi:10.1038/nature01281.1., 2002.
- Mühlenbruch, M., Grossart, H. P., Eigemann, F. and Voss, M.: Mini-review: Phytoplankton-derived polysaccharides in the  
900 marine environment and their interactions with heterotrophic bacteria, Environ. Microbiol., 20(8), 2671–2685, doi:10.1111/1462-2920.14302, 2018.
- Murphy, K. R., Butler, K. D., Spencer, R. G. M., Stedmon, C. A., Boehme, J. R. and Aiken, G. R.: Measurement of dissolved organic matter fluorescence in aquatic environments: An interlaboratory comparison, Environ. Sci. Technol., 44, 9405–9412, doi:10.1021/es102362t, 2010.
- 905 Nieto-Cid, M., Álvarez-Salgado, X. A. and Pérez, F. F.: Microbial and photochemical reactivity of fluorescent dissolved organic matter in a coastal upwelling system, Limnol. Oceanogr., 51(3), 1391–1400, 2006.
- Nimer, N. A., Iglesias-rodriguez, M. D. and Merrett, M. J.: Bicarbonate utilization by marine phytoplankton species, J. Phycol., 631, 625–631, 1997.
- Norland, S.: The Relationship Between Biomass and Volume of Bacteria, in Handbook of Methods in Aquatic Microbial  
910 Ecology, pp. 303–306, Routledge., 1993.
- Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlenn, D., Minchin, P. R., O’Hara, R. B., Simpson, G. L., Solymos, P., Stevens, M. H. H., Szoecs, E. and Wagner, H.: vegan: Community Ecology Package. R package version 2.5-5. <https://CRAN.R-project.org/package=vegan>, 2019.
- Olenina, I., Hajdu, S., Edler, L., Andersson, A., Wasmund, N., Busch, S., Göbel, J., Gromisz, S., Huseby, S., Huttunen, M.,  
915 Jaanus, A., Kokkonen, P., Ledaine, I. and Niemkiewicz, E.: Biovolumes and Size-Classes of Phytoplankton in the Baltic Sea, HELCOM Balt. Sea Environ. Proc., 106, 2006.
- Orellana, M. V, Pang, W. L., Durand, P. M., Whitehead, K. and Baliga, N. S.: A Role for Programmed Cell Death in the Microbial Loop, PLoS One, 8(5), e62595, doi:10.1371/journal.pone.0062595, 2013.
- Pedler, B. E., Aluwihare, L. I. and Azam, F.: Single bacterial strain capable of significant contribution to carbon cycling in the



- 920 surface ocean, *Proc. Natl. Acad. Sci.*, 111(20), 7202–7207, doi:10.1073/pnas.1401887111, 2014.
- Pérez, M. T. and Sommaruga, R.: Differential effect of algal- and soil-derived dissolved organic matter on alpine lake bacterial community composition and activity, *Limnol. Oceanogr.*, 51(6), 2527–2537, doi:10.4319/lo.2006.51.6.2527, 2006.
- Peters, G.: *\_userfriendlyscience: Quantitative analysis made accessible\_*. doi: 10.17605/osf.io/txequ (URL: <https://doi.org/10.17605/osf.io/txequ>), R package version 0.7.2, <URL: <https://userfriendlyscience.com>>, 2018.
- 925 Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J. and Glöckner, F. O.: The SILVA ribosomal RNA gene database project: improved data processing and web-based tools, *Nucleic Acids Res.*, 41, 590–596, doi:10.1093/nar/gks1219, 2013.
- R Core Team: R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>, 2019.
- 930 Riemann, B., Bjørnsen, P. K., Newell, S. and Fallon, R.: Calculation of cell production of coastal marine bacteria based on measured incorporation of [3H]thymidine, *Limnol. Oceanogr.*, 32(2), 471–476, doi:10.4319/lo.1987.32.2.0471, 1987.
- Riemann, L., Leitet, C., Pommier, T., Simu, K., Holmfeldt, K., Larsson, U. and Hagstro, Å.: The Native Bacterioplankton Community in the Central Baltic Sea Is Influenced by Freshwater Bacterial Species □ †, , 74(2), 503–515, doi:10.1128/AEM.01983-07, 2008.
- 935 Rintala, J. M., Spilling, K. and Blomster, J.: Temporary cyst enables long-term dark survival of *Scrippsiella hangoei* (Dinophyceae), *Mar. Biol.*, 152(1), 57–62, doi:10.1007/s00227-007-0652-x, 2007.
- Romera-Castillo, C., Sarmiento, H., Álvarez-Salgado, X. A., Gasol, J. M. and Marrase, C.: Production of chromophoric dissolved organic matter by marine phytoplankton, *Limnol. Oceanogr.*, 55(1), 446–454, 2010.
- Romera-Castillo, C., Sarmiento, H., Álvarez-Salgado, X. A., Gasol, J. M. and Marrase, C.: Net Production and Consumption of Fluorescent Colored Dissolved Organic Matter by Natural Bacterial Assemblages Growing on Marine Phytoplankton Exudates, *Appl. Environ. Microbiol.*, 77(21), 7490–7498, doi:10.1128/AEM.00200-11, 2011.
- 940 Saad, E. M., Longo, A. F., Chambers, L. R., Huang, R., Benitez-Nelson, C., Dyhrman, S. T., Diaz, J. M., Tang, Y. and Ingall, E. D.: Understanding marine dissolved organic matter production: Compositional insights from axenic cultures of *Thalassiosira pseudonana*, *Limnol. Oceanogr.*, 61, 2222–2233, doi:10.1002/lno.10367, 2016.
- 945 Sapp, M., Schwaderer, A. S., Wiltshire, K. H., Hoppe, H.-G., Gerdt, G. and Wichels, A.: Species-specific bacterial communities in the phycosphere of microalgae?, *Microb. Ecol.*, 53(4), 683–699, doi:10.1007/s00248-006-9162-5, 2007.
- Sarmiento, H. and Gasol, J. M.: Use of phytoplankton-derived dissolved organic carbon by different types of bacterioplankton, *Environ. Microbiol.*, 14(9), 2348–2360, doi:10.1111/j.1462-2920.2012.02787.x, 2012.
- Sarmiento, H., Romera-Castillo, C., Lindh, M., Pinhassi, J., Sala, M. M., Gasol, J. M., Marrasé, C. and Taylor, G. T.: 950 Phytoplankton species-specific release of dissolved free amino acids and their selective consumption by bacteria, *Limnol. Oceanogr.*, 58(3), 1123–1135, doi:10.4319/lo.2013.58.3.1123, 2013.
- Schäfer, H., Abbas, B., Witte, H. and Muyzer, G.: Genetic diversity of “satellite” bacteria present in cultures of marine diatoms, *FEMS Microbiol. Ecol.*, 42(1), 25–35, doi:10.1016/S0168-6496(02)00298-2, 2002.



- Seymour, J. R., Amin, S. A., Raina, J. B. and Stocker, R.: Zooming in on the phycosphere: The ecological interface for  
955 phytoplankton-bacteria relationships, *Nat. Microbiol.*, 2(May), 1–12, doi:10.1038/nmicrobiol.2017.65, 2017.
- Smith, D. C. and Azam, F.: A simple, economical method for measuring bacterial protein synthesis rates in seawater using  
3H-leucine, *Mar. Microb. Food Webs*, 6(2), 107–114, 1992.
- Spilling, K., Olli, K., Lehtoranta, J., Kremp, A., Tedesco, L., Tamelander, T., Klais, R., Peltonen, H. and Tamminen, T.:  
960 Shifting Diatom—Dinoflagellate Dominance During Spring Bloom in the Baltic Sea and its Potential Effects on  
Biogeochemical Cycling, *Front. Mar. Sci.*, 5, 327, doi:10.3389/fmars.2018.00327, 2018.
- Spilling, K., Fuentes-Lema, A., Quemaliños, D., Klais, R. and Sobrino, C.: Primary production, carbon release, and respiration  
during spring bloom in the Baltic Sea, *Limnol. Oceanogr.*, 64(4), 1779–1789, doi:10.1002/lno.11150, 2019.
- Stedmon, C. A. and Markager, S.: Tracing the production and degradation of autochthonous fractions of dissolved organic  
matter by fluorescence analysis, *Limnol. Oceanogr.*, 50(5), 1415–1426, 2005.
- 965 Storch, T. A. and Saunders, G. W.: Phytoplankton extracellular release and its relation to the seasonal cycle of dissolved  
organic carbon in a eutrophic lake, *Limnol. Oceanogr.*, 23(1), 112–119, doi:10.4319/lo.1978.23.1.0112, 1978.
- Strom, S. L., Benner, R., Ziegler, S. and Dagg, M. J.: Planktonic grazers are a potentially important source of marine dissolved  
organic carbon, *Limnol. Oceanogr.*, 42(6), 1364–1374, doi:10.4319/lo.1997.42.6.1364, 1997.
- Suikkanen, S., Hakanen, P., Spilling, K. and Kremp, A.: Allelopathic effects of Baltic Sea spring bloom dinoflagellates on co-  
970 occurring phytoplankton, *Mar. Ecol. Prog. Ser.*, 439, 45–55, doi:10.3354/meps09356, 2011.
- Suksomjit, M., Nagao, S., Ichimi, K., Yamada, T. and Tada, K.: Variation of dissolved organic matter and fluorescence  
characteristics before, during and after phytoplankton bloom, *J. Oceanogr.*, 65(6), 835–846, doi:10.1007/s10872-009-0069-x,  
2009.
- Tada, Y., Nakaya, R., Goto, S., Yamashita, Y. and Suzuki, K.: Distinct bacterial community and diversity shifts after  
975 phytoplankton-derived dissolved organic matter addition in a coastal environment, *J. Exp. Mar. Bio. Ecol.*, 495(August 2016),  
119–128, doi:10.1016/j.jembe.2017.06.006, 2017.
- Taylor, F. J. R. and Pollinger, U.: Ecology of Dinoflagellates, in *Biology of Dinoflagellates*, Blackwell Scientific  
Publications, Oxford., 1987.
- Teeling, H., Fuchs, B. M., Becher, D., Klockow, C., Gardebrecht, A., Bennke, C. M., Kassabgy, M., Huang, S., Mann, A. J.,  
980 Waldmann, J., Weber, M., Klindworth, A., Otto, A., Lange, J., Bernhardt, J., Reinsch, C., Hecker, M., Peplies, J., Bockelmann,  
F. D., Callies, U., Gerdt, G., Wichels, A., Wiltshire, K. H., Glöckner, F. O., Schweder, T. and Amann, R.: Substrate-controlled  
succession of marine bacterioplankton populations induced by a phytoplankton bloom, *Science* (80-. ), 336(6081), 608–611,  
doi:10.1126/science.1218344, 2012.
- Thornton, D. C. O.: Dissolved organic matter (DOM) release by phytoplankton in the contemporary and future ocean, *Eur. J.*  
985 *Phycol.*, 49(1), 20–46, doi:10.1080/09670262.2013.875596, 2014.
- Urbani, R., Magaletti, E., Sist, P. and Cicero, A. M.: Extracellular carbohydrates released by the marine diatoms *Cylindrotheca*  
*closterium*, *Thalassiosira pseudonana* and *Skeletonema costatum*: Effect of P-depletion and growth status, *Sci. Total Environ.*,



- 353, 300–306, doi:10.1016/j.scitotenv.2005.09.026, 2005.
- 990 Weishaar, J. L., Aiken, G. R., Bergamaschi, B. A., Fram, M. S., Fujii, R. and Mopper, K.: Evaluation of specific ultraviolet absorbance as an indicator of the chemical composition and reactivity of dissolved organic carbon, *Environ. Sci. Technol.*, 37, 4702–4708, doi:10.1021/es030360x, 2003.
- Wickham, H.: *ggplot2: Elegant Graphics for Data Analysis*, Springer-Verlag, New York., 2016.
- Yamashita, Y. and Tanoue, E.: Chemical characteristics of amino acid-containing dissolved organic matter in seawater, *Org. Geochem.*, 35(6), 679–692, doi:10.1016/j.orggeochem.2004.02.007, 2004a.
- 995 Yamashita, Y. and Tanoue, E.: In situ production of chromophoric dissolved organic matter in coastal environments, *Geophys. Res. Lett.*, 31(14), 1–5, doi:10.1029/2004GL019734, 2004b.
- Yamashita, Y., Hashihama, F., Saito, H., Fukuda, H. and Ogawa, H.: Factors controlling the geographical distribution of fluorescent dissolved organic matter in the surface waters of the Pacific Ocean, *Limnol. Oceanogr.*, 62(6), 2360–2374, doi:10.1002/lno.10570, 2017.
- 1000 Zsolnay, A., Baigar, E., Jimenez, M., Steinweg, B. and Saccomandi, F.: Differentiating with fluorescence spectroscopy the sources of dissolved organic matter in soils subjected to drying, *Chemosphere*, 38(1), 45–50, doi:10.1016/S0045-6535(98)00166-0, 1999.