

1 **What can we learn from amino acids about oceanic organic matter cycling**
2 **and degradation?**

3

4 **Birgit Gaye¹, Niko Lahajnar¹, Natalie Harms¹, Sophie Anna Luise Paul^{2, 3}, Tim Rixen^{1,4}**
5 **and Kay-Christian Emeis¹**

6 *¹Institute for Geology, Universität Hamburg, 20146 Hamburg, Germany*

7 *²Department of Physics and Earth Sciences, Jacobs University Bremen, 28759 Bremen,*
8 *Germany*

9 *³GEOMAR, Helmholtz-Zentrum für Ozeanforschung, 24148 Kiel, Germany*

10 *⁴Leibniz Centre for Tropical Marine Research (ZMT), 28359 Bremen, Germany*

11

12 **Correspondence:** Birgit Gaye (birgit.gaye@uni-hamburg.de)

13

14 **Abstract**

15 Amino acids (AA) mainly bound in proteins are major constituents of living biomass and non-
16 living organic material in the oceanic particulate and dissolved organic matter pool. Uptake and
17 cycling by heterotrophic organisms lead to characteristic changes in AA composition so that
18 AA based biogeochemical indicators are often used to elucidate processes of organic matter
19 cycling and degradation. We analyzed particulate AA in a large sample set collected in various
20 oceanic regions covering sinking and suspended particles in the water column, sediment
21 samples as well as dissolved AA from water column and pore water samples. The aim of this
22 study was to test and improve the use of AA derived biogeochemical indicators as proxies for
23 organic matter sources and degradation, and to better understand particle dynamics and
24 interaction between the dissolved and particulate organic matter pools. A principal component
25 analysis (PCA) of all data delineates diverging AA compositions of sinking and suspended
26 particles with increasing water depth. A new sinking particle and sediment degradation
27 indicator (SDI) allows a fine-tuned classification of sinking particles and sediments with respect
28 to the intensity of degradation, which is associated with changes of bulk $\delta^{15}\text{N}$ ratios. This new
29 indicator furthermore is sensitive to sedimentary redox conditions and can be used to detect
30 past anoxic early diagenesis. A second indicator emerges from the AA spectra of suspended

31 particulate matter (SPM) in the epipelagic and that of the meso- and bathypelagic ocean and is
32 a residence time indicator (RTI). The characteristic changes in AA patterns from shallow to
33 deep SPM are recapitulated in the AA spectra of the dissolved organic matter (DOM) pool, so
34 that deep SPM is more similar to DOM than to any of the other organic matter pools. This
35 implies that there is equilibration between finely dispersed SPM and DOM in the deep sea,
36 which may be driven by microbial activity combined with annealing and fragmentation of gels.
37 As these processes strongly depend on physico-chemical conditions in the deep ocean, changes
38 in quality and degradability of DOM may strongly affect the relatively large pool of suspended
39 and dissolved AA in the ocean that amounts to 15 Pg amino acid carbon (AAC) and 89 ± 29 Pg
40 AAC, respectively.

41 **1 Introduction**

42 Amino acids are ubiquitous in living organisms and comprise a major share of characterized
43 organic matter in the particulate and dissolved pool in the ocean (Lee, 1988; Wakeham et al.,
44 1984; Zhang et al., 2016; Davis et al., 2009; Lee et al., 2004). AA comprise more than 80 % of
45 total organic carbon in fresh autochthonous plankton while allochthonous organic matter from
46 continental runoff and atmospheric deposition has lower AA contents (Degens and Ittekkot,
47 1983; Degens and Ittekkot, 1985). Most of the primary productivity occurs in the sunlit surface
48 layer of the ocean and most of the allochthonous material is also transported into surface waters.
49 Thus, organic matter concentrations including their major biogenic constituents generally, have
50 a surface maximum and decrease with depth (Peters et al., 2018; Gaye et al., 2013b; Wakeham
51 and Lee, 1993). The main mechanism behind this depth dependent distribution is that most of
52 the organic matter is recycled in surface waters while only a small proportion of surface
53 particles leaves the surface waters by gravitational settling in the form of macroaggregates or
54 fecal pellets. Particles containing organic matter, shells, frustules of organisms and mineral
55 matter sink at speeds of 200 m day^{-1} on average and constitute the export from the surface mixed
56 layer or euphotic zone into the deep ocean, where part of it can ultimately reach the sediments
57 (Alldredge and Silver, 1988; Alldredge, 1998; Pilskaln and Honjo, 1987; Fowler and Knauer,
58 1986; Karl et al., 1988; Rixen et al., 2019b). Sinking particles are caught by moored or floating
59 sediment traps while SPM is sampled by filtration or ultrafiltration of water from water
60 samplers or by pump systems (Yamaguchi and McCarthy, 2018). SPM is too small to sink and
61 therefore – like DOM – predominantly enters deep water by subduction of surface waters
62 (Resplandy et al., 2019; Boyd et al., 2019) and is transported passively following the route of
63 ocean water along the ocean conveyor belt (Silver et al., 1998; McCave, 1984). It has thus been
64 surmised that the long residence time of SPM in the water column should result in a more
65 degraded state compared with organic matter of sinking particles (McCave, 1984; Degens and
66 Ittekkot, 1984). Studies of pigments, AA and fatty acids, however, do not find such a systematic
67 difference between the two types of particles and even indicate that SPM can be less degraded
68 than sinking particles (Abramson et al., 2011; Rontani et al., 2011; Wakeham and Canuel,
69 1988). In two studies of AA composition in the Benguela Upwelling System and in the Arabian
70 Sea it was shown that the degradation pathways of SPM and sinking particles differ as their AA
71 compositions diverge with depth (Gaye et al., 2013b; Nagel et al., 2009). These studies
72 suggested that there is only little interaction between suspended and sinking particle pools
73 below the euphotic zone. Due to its long residence time in the ocean, SPM appears to interact

74 with DOM (Gaye et al., 2013b) and therefore carries different AA signatures related to genesis
75 and history of organic matter cycling in its specific water mass (Nagel et al., 2016). Whereas
76 information on the composition of sediment trap samples has been compiled in comprehensive
77 studies (Honjo et al., 2008; Wilson et al., 2012; Rixen et al., 2019a, b), similar compilations of
78 the profuse literature on suspended matter are yet missing.

79 On the way to the deep sea the flux of sinking particles is reduced by disaggregation and organic
80 matter degradation. Suess (1980) empirically derived the first power function for organic
81 carbon decay based on sediment trap data. Subsequently, a large number of similar functions
82 were calculated for various oceanic areas based on trap experiments (Rixen et al., 2019b;
83 Rixen et al., 2002; Armstrong et al., 2002; Martin et al., 1987). Early work on AA had produced
84 similar decay functions combining data from Atlantic and Pacific trap experiments (Lee and
85 Cronin, 1982, 1984). As AA decay faster than bulk organic carbon (Haake et al., 1993b; Haake
86 et al., 1992; Haake et al., 1996; Lee et al., 2004; Wakeham and Lee, 1989; Whelan and Emeis,
87 1992), they are often considered as “labile” constituents of bulk organic matter. This is
88 supposedly due to their preferential uptake as a nitrogen (N) source for further synthesis of AA
89 or as a source of essential AA for heterotrophs (Ittekkot and Arain, 1986; Ittekkot et al., 1986).
90 This has been questioned, as a large proportion of the oceanic organic N pool is comprised of
91 AA that are not bioavailable (Aluwihare et al., 2005). In addition to the quantification of AA
92 decay, degradation state of organic matter (proteins) can be assessed by characteristic changes
93 in AA monomer composition which, furthermore, have the potential to elucidate sources of
94 organic matter and degradation processes (Ittekkot et al., 1984a; Ittekkot et al., 1984b; Dauwe
95 and Middelburg, 1998; Dauwe et al., 1999; Jennerjahn and Ittekkot, 1997).

96 Ratios of individual amino acids such as the Reactivity Index (RI) (Jennerjahn and Ittekkot,
97 1997) or the Degradation Index (DI) normalizing AA data to the results of a principal
98 component analyses (PCA) (Dauwe et al., 1999; Dauwe and Middelburg, 1998) have often been
99 used to scale organic matter degradation (Niggemann et al., 2018; Unger et al., 2005; Ingalls et
100 al., 2006; Ingalls et al., 2004; Pantoja et al., 2004; Möbius et al., 2010). These biogeochemical
101 indicators of organic matter quality were essentially developed for marine sinking particles and
102 sediments. Although based on marine sediments only (Dauwe et al., 1999) the DI was applied
103 for example to SPM samples from the brackish environment (Unger et al. 2005) or even to trace
104 dissolved AA degradation (Davis and Benner, 2005; Guo et al., 2018). Other work used
105 individual and adapted indices to differentiate the states of degradation in SPM or DOM

106 samples and samples from lakes and rivers (Abramson et al., 2011; Gaye et al., 2007; Goutx et
107 al., 2007; Menzel et al., 2013; Sheridan et al., 2002; Kaiser and Benner, 2009).

108 Understanding and quantifying AA degradation is required to estimate the diagenetic imprint
109 on $\delta^{15}\text{N}$ ratios of particulate matter. This is important as $\delta^{15}\text{N}$ ratios track major shifts between
110 N pools and are commonly used to reconstruct the N cycle from sedimentary archives
111 (Galbraith et al., 2013). Amino acid nitrogen (AAN) comprises 80-100 % of N in fresh organic
112 matter and is the precursor of most of the N buried in sediments and ultimately stored in the
113 form of ammonium, adsorbed to clay minerals (Boyd, 2001; Waples and Sloan, 1980; Müller,
114 1977). Considerable AA degradation already occurs in the water column and progresses during
115 organic matter burial in the sediments so that the impact of diagenetic processes on $\delta^{15}\text{N}$ has to
116 be accounted for (Möbius et al., 2010; Möbius et al., 2011; Niggemann et al., 2018; Carr et al.,
117 2016). Ammonification leads to a diagenetic increase of $\delta^{15}\text{N}$ values by up to 6.5 ‰ in deep
118 sea sediments while there is little effect during organic matter burial in shelf and slope
119 sediments due to the higher sedimentation rates and sub- to anoxic diagenetic conditions
120 (Tesdal et al., 2013; Robinson et al., 2012; Möbius, 2013; Gaye-Haake et al., 2005). Such $\delta^{15}\text{N}$
121 increases were shown to correlate with AA derived degradation indicators so that the primary
122 $\delta^{15}\text{N}$ signal from the water column can be reconstructed (Gaye-Haake et al., 2005; Gaye et al.,
123 2009; Möbius et al., 2011).

124 DOM comprising the largest oceanic organic matter pool is defined by the pore size of the
125 filters it passes through which is 0.2-0.7 μm (Carlson and Hansell, 2015) and thus includes
126 some picoplankton cells and all viruses (Aristegui et al., 2009). DOM in surface water is partly
127 labile and can originate from the exudates and lysis of organisms, passive diffusion, or
128 “overflow” out of phytoplankton and bacteria; grazers can excrete or egest DOM, it can
129 furthermore be leached from their fecal pellets or released by sloppy zooplankton feeding and
130 is thus primarily released and also taken up in the surface ocean (Carlson and Hansell, 2015).
131 Moreover, terrestrially derived DOM is transported into surface waters by rivers and via the
132 atmosphere (Benner et al., 2005). Deep DOM has a different source than simply transport of
133 surface DOM by intermediate and deep water formation and mixing, as deep DOM is refractory
134 in nature and has been heterotrophically altered by cycling and degradation processes
135 (Yamaguchi and McCarthy, 2018) discernible e.g. from their composition of dissolved AA
136 (Kaiser and Benner, 2009; McCarthy et al., 2004). The possible source of deep DOM may be
137 the release from sinking or suspended particles associated with microbial degradation on

138 particles and in the ambiance of particles by processes such as solubilizing organic matter by
139 ectohydrolase (Cho and Azam, 1988; Ciais et al., 2014; Aristegui et al., 2009). DOM can also
140 be released from sediment pore water into overlying waters (Lahajnar et al., 2005b). Stable
141 isotope ratios of nitrogen ($\delta^{15}\text{N}$) in ultrafiltered DOM (UDOM) showed no systematic change
142 with depth and suggested a common microbial source or viral lysis (McCarthy et al., 2007).

143 In the following synoptic compilation of AA data, we will examine the differences in AA
144 spectra of a large data set that combines dissolved and particulate AA from plankton, suspended
145 and sinking material, and sediments from different oceanic regions, as well as from riverine to
146 brackish-marine conditions. Focusing on processes in the water column the data serve to (i) test
147 existing AA based biogeochemical indicators of organic matter sources and degradation, (ii)
148 better understand transformation and degradation processes of organic matter in aquatic
149 environments reflected by AA composition in sinking and suspended particles and total
150 dissolved AA (TDAA), (iii) investigate the impact of such processes on the $\delta^{15}\text{N}$ values and
151 (iv) identify open questions which may be pursued with the help of AA analyses in the future.

152


153 **2. Materials and Methods**

154 **2.1 Sampling**

155 A total of 1425 samples were taken for AA analyses in different oceanic areas and water depths
156 between 1993 and 2017 and include 218 sediment trap samples, 489 sediment samples, 608
157 SPM samples and 110 water and pore water samples (Fig. 1a-d). Five additional plankton
158 samples were taken in the Arabian Sea and from the Namibian upwelling area by plankton tows
159 between 0-100 m and between 100-700 m water depths. In the Kara Sea - a shallow shelf sea
160 strongly impacted by water and suspended matter discharge from the rivers Ob and Yenisei -
161 sediment traps, surface sediments, and suspended matter were sampled (Gaye et al., 2007;
162 Gaye-Haake et al., 2003; Nagel et al., 2009; Unger et al., 2005). In the deep Mediterranean Sea
163 sediment traps and surface sediments were sampled (Möbius et al., 2010). SPM from the
164 Mediterranean Sea was only analyzed for $\delta^{15}\text{N}$ values (Emeis et al., 2010). Sediment trap as
165 well as surface sediment, SPM and water samples were taken along cross shelf transects off
166 Namibia (Nagel et al., 2013; Nagel et al., 2016). Sediment trap samples and short sediment
167 cores were taken at two stations in the northeastern Atlantic (Lahajnar et al., 2005a;
168 Turnewitsch et al., 2017; Turnewitsch et al., 2015). In the Arabian Sea sediment trap, SPM, and

169 surface sediment samples were taken in the deep ocean and on the continental slope including
170 a core within the oxygen minimum zone at water depths of 775m (Gaye et al., 2013b; Gaye et
171 al., 2013a; Rixen et al., 2014; Gaye-Haake et al., 2005; Suthhof et al., 2001; Suthhof et al.,
172 2000). In the Indian Ocean Subtropical Gyre sediment trap, SPM and water samples were taken
173 (Harms et al., 2019; Harms et al., 2021) and samples from the equatorial North Pacific and
174 eastern South Pacific comprise bottom water, pore water and sediment core samples (Paul et
175 al., 2018).

176 Sea water was filtered through glass fiber filters (Whatman GF/F) with a nominal pore size of
177 0.7 μm and filters were dried at 40°C in order to obtain SPM samples. At some stations water
178 samples were taken by deep freezing an aliquot of the filtrate for TDAA analyses. In addition,
179 18 water samples taken off Namibia were separated into two size classes by ultrafiltration
180 (Brockmeyer and Spitzky, 2013). The size classes 50 kDa-0.7 μm and 1 kDa-0.7 μm were used
181 for TDAA analyses. Sediment trap samples were wet sieved on board and comprise the <1mm
182 fraction, filtered with polycarbonate nuclepore filters of 0.45 μm pore size and dried at 40°C.
183 Sediment samples from multicores, box grabs, box cores, or gravity cores were taken by spatula
184 or syringes from cold stored cores and were freeze dried before analyses. Surface samples
185 represent either the upper 0.5 cm or 1 cm of a sediment core. Pore-water samples were taken
186 by rhizons with a mean pore size of 0.15 μm and stored frozen before analyses (see methods in
187 Paul et al., 2018).



188

189 Figure 1: Stations of sediment trap deployments (a), sediment (b), SPM (c) water and pore water
 190 sampling (d).

191

192 2.2 Analytical methods

193 Total carbon and N were measured with a Carlo Erba Nitrogen Analyser 1500 (Milan, Italy) or
 194 a EURO EA3000 elemental analyzer. Particulate organic carbon (POC) was measured after
 195 treatment of weighed samples with 1N HCl to remove carbonate. The precision of this method
 196 is 0.05% for carbon and 0.005% for N. Carbonate carbon was calculated by subtracting organic
 197 carbon from total carbon. Ratios of $^{15}\text{N}/^{14}\text{N}$ of particulate N were determined using a Thermo
 198 Finnigan MAT 252 isotope ratio mass spectrometer connected with a ConFlo-III interface after
 199 high-temperature flash combustion in a Thermo Finnigan Flash EA 1112 at 1050°C. Part of the
 200 samples were measured with an Elementar IsoPrime 100 isotope ratio mass spectrometer after
 201 high temperature combustion in an Elementar CHNOS Vario isotope elemental analyzer at 950
 202 °C. Pure tank N_2 calibrated against the reference standards IAEA-N1 (ammonium sulfate,
 203 $\delta^{15}\text{N} = +0.4 \text{ ‰}$ versus air N_2) and IAEAN2 (ammonium sulfate, $\delta^{15}\text{N} = +20.3 \text{ ‰}$) of the

204 International Atomic Energy Agency was used as a working standard. Duplicate measurements
205 of samples differ by less than 0.15 ‰. The laboratory's long-term standard deviation for IAEA-
206 N1 standard is 0.09 ‰. N-isotope ratios are reported in ‰ using the delta notation and the
207 $^{15}\text{N}/^{14}\text{N}$ of air N_2 as the reference standard:

$$208 \quad \delta^{15}\text{N}_{\text{sample}} = \left(\frac{(^{15}\text{N}/^{14}\text{N})_{\text{sample}}}{(^{15}\text{N}/^{14}\text{N})_{\text{reference standard}}} - 1 \right) * 1000 \quad (1)$$

209 Dissolved organic carbon (DOC) concentrations [mg/L] were determined of water and pore
210 water samples via a high temperature combustion method (POC-V_{CSH} Analyzer, Shimadzu).
211 Inorganic carbon was removed by 2 M HCl prior to injection into the combustion tube where
212 organic carbon is oxidized to CO_2 at 680 °C with a platinum catalyst. A 5-point calibration from
213 0.5 to 5 mg DOC/L was used. The error of measurement is less than 2 % (Brockmeyer and
214 Spitzzy, 2013).

215 TDAA, particulate AA and hexosamines (HA) were analyzed with a Biochrom 30 Amino Acid
216 Analyzer. Acid hydrolysis with 6N HCl for 22 h at 110°C under a pure argon atmosphere was
217 carried out on ca. 3 ml of filtrate of water and pore water samples, on 1-2 mg of suspended
218 matter collected on Whatman GF/F filters, on 1-2 mg of sediment trap samples, or on 1-50 mg
219 of freeze dried surface sediments. A particle free aliquot was evaporated three times to dryness
220 in order to remove the unreacted HCl; the residue was taken up in an acidic buffer (pH 2.2).
221 After injection and subsequent separation with a cation exchange resin, the individual AA
222 monomers were post-column derivatized with o-phthaldialdehyde in the presence of 2-
223 mercaptoethanol and detected with a Merck Hitachi L-2480 fluorescence detector. Duplicate
224 analysis of a standard solution according to this method results in a relative error of 0.1 to 1.3%
225 for the concentrations of individual AA monomers and 0.2 to 3.0% for individual AA
226 monomers of water or particulate matter samples. Due to acid hydrolysis, aspartic acid (ASP)
227 and asparagine (Asn) are both measured as Asp and glutamic acid (Glu) and glutamine (Gln)
228 are both measured as Glu. The other AA measured are threonine (Thr), serine (Ser), glycine
229 (Gly), alanine (Ala), valine (Val), methionine (Met), isoleucine (Ile), leucine (Leu), tyrosine
230 (Tyr), phenylalanine (Phe), β -Alanine (β -Ala), γ -aminobutyric acid (γ -Aba), histidine (His),
231 ornithine (Orn), lysine (Lys) and arginine (Arg). The HA together with AA are glucosamine
232 (Gluam) and galactosamine (Galam) and their total contents were corrected with a factor of 1.4
233 for loss during hydrolysis (Muller et al., 1986).

234

235 **2.3 Amino acid derived biogeochemical indicators of organic matter origin and** 236 **degradation**

237 Amino acid concentrations and the contribution of AAC and AA nitrogen (AAN) as
238 percentages of total organic carbon (AAC/C %) or total N (AAN/N %) are used to determine
239 the degradation state of organic matter in the marine realm as both decrease with increasing
240 organic matter degradation (Wakeham and Lee, 1993; Cowie and Hedges, 1994). AAN/N %
241 >50 % are characteristic of fresh organic matter in the freshwater and marine realm (Menzel et
242 al., 2015; Haake et al., 1992; Haake et al., 1993b). AA contribute >60 % to total organic carbon
243 (AAC/C %) in fresh plankton and suspended matter in surface waters whereas AAC/C % drop
244 to values <20 % in sinking particles and suspended matter from subsurface water (Wakeham
245 and Lee, 1993). AAC/C % values are often below 10 % in freshwater environments and indicate
246 the enhanced input of land plants enriched in carbohydrates and lignin rather than enhanced
247 organic matter degradation (Menzel et al., 2015).

248 Asn, Gln and Glu are the primary products of N assimilation and all other AA are synthesized
249 from them (Loick-Wilde et al., 2018; Riccardi et al., 1989; Hildebrandt et al., 2015). Asp and
250 Glu are enriched in bacteria, vascular plant tissue, phytoplankton, zooplankton and fungi and
251 their high relative contents of Asp and Glu therefore, indicate fresh organic matter (Cowie and
252 Hedges, 1992). The ratios of Asp/ β -Ala and Glu/ γ -Aba are high in fresh organic matter and
253 drop with organic matter degradation as β -Ala and γ -Aba are degradation products of Asp and
254 Glu, respectively (Lee and Cronin, 1984). β -Ala and γ -Aba also become relatively enriched
255 during organic matter degradation as these non-protein AA are not taken up by heterotrophic
256 organisms (Ittekkot et al., 1984b).

257 The relative accumulation of the non-protein AA is also expressed by the RI which is the ratio
258 of the very labile aromatic AA Tyr and Phe and the non-protein AA β -Ala and γ -Aba. The RI
259 is, generally, between 0 (very degraded) and 15 (very fresh) (Jennerjahn and Ittekkot, 1997). It
260 is applicable not only in studies of sinking and suspended matter in marine and brackish
261 environments (Unger et al., 2005; Gaye et al., 2007) but also as a proxy for degradation state in
262 the sediment column (Möbius et al., 2011). The enrichment of Asp and Glu in sediments is
263 related to their enrichment in carbonate shells (Ittekkot et al., 1984a) and to adsorption of
264 primarily acidic AA onto carbonate minerals (King and Hare, 1972), whereas basic AA
265 primarily adsorb onto silicate minerals (Hedges and Hare, 1987; Keil et al., 1994; King, 1975).

266 The DI, the integral of 14 protein AA, assesses the diagenetic alteration of a sample by
 267 comparing it to a set of 28 sediment samples of different degradation states and environments.
 268 Molar percentages of individual AA are standardised by the mean and standard deviations of
 269 the 28-sample data set. The DI then integrates the result of these standardized values weighed
 270 by the factor coefficients for the first axis of the PCA of Dauwe et al. (1999) according to the
 271 formula:

$$272 \quad DI = \sum_i \left[\frac{\text{var}_i - \text{AVGvar}_i}{\text{STDvar}_i} \right] \cdot \text{fac.coef}_i \quad (2)$$

273 where var_i is the original mole percentage of each AA_i , AVGvar_i and STDvar_i are the mean and
 274 standard deviations, respectively, and fac.coef_i is the factor coefficient of the first axis of the
 275 PCA of Dauwe et al. (1999). The DI thus represents the cumulative deviation of AA with
 276 respect to an assumed average molar composition. The DI ranges approximately from -2 to +3
 277 where negative values indicate more and positive values less degradation than the average.

278 An indicator of oxic vs. anoxic organic matter degradation in the water column and in sediments
 279 was proposed by Menzel et al. (2015) for lake samples. Based on work by Cowie et al. (1995)
 280 on marine sediments the ox/anox indicator is the quotient of AA preserved under oxic
 281 diagenetic conditions to those preserved in anoxic water or sediments and is thus higher in oxic
 282 than in anoxic sediments:

$$283 \quad \text{ox/anox} = \frac{\text{Asp} + \text{Glu} + \beta\text{-Ala} + \gamma\text{-Aba} + \text{Lys}}{\text{Ser} + \text{Met} + \text{Ile} + \text{Leu} + \text{Tyr} + \text{Phe}} \quad (3)$$

284 ox/anox ratios <1.0 indicate anoxic and ratios >1.5 oxic diagenesis (Menzel et al., 2015).

285 The stability of AA vs. HA has been discussed since the early research on AA and HA in natural
 286 material. Fresh plankton was observed to have AA/HA ratios of 13-25 (Degens and Mopper,
 287 1975) which is a mixed signal of phytoplankton with an AA/HA ratio of >80 and zooplankton
 288 with a ratio of ~9 due to chitinous skeletons of many zooplankters (Mayzaud and Martin,
 289 1975). Low AA/HA are also observed in cell walls of fungi and bacteria. As the building blocks
 290 of chitin, HA were assumed to be more resistant to degradation than bulk AA (Muller et al.,
 291 1986). This is, however, challenged by studies of enzyme activities which were observed to
 292 respond to substrate availability so that the activity of chitobiase and chitinase is as high as that
 293 of glucosidase (Boetius et al., 2000a; Boetius and Lochte, 1994; Boetius et al., 2000b; Smith et
 294 al., 1992) suggesting intense degradation also of chitin. Gluam is the main constituent of chitin

295 and while Galam is relatively enriched in bacterial cell walls (Walla et al., 1984; Kandler,
296 1979). The Gluam/Galam ratio has, therefore, been used to distinguish bacterial material from
297 zooplankton rich material (Haake et al., 1993b; Benner and Kaiser, 2003; Niggemann and
298 Schubert, 2006). Gluam/Galam ratios > 4 were found in sinking particles (Haake et al., 1993b;
299 Haake et al., 1992; Lahajnar et al., 2007), ratios of < 3 usually indicate relatively high
300 contribution of microbial OM and values between 1 and 2 are characteristic of sediments and
301 indicate an enrichment of microbial biomass (Benner and Kaiser, 2003).

302


303 **2.4 Statistical analyses**

304 To investigate the differences of AA composition and to recognize the interaction and pathways
305 of degradation between the different pools we carried out a PCA of AA monomer contributions
306 in Mol %. Met was excluded as it is below detection limit in many samples. The PCA was
307 carried out using the program SPSS Statistics 22. PCAs have often been used to analyze large
308 databases (Xue et al., 2011) in order to trace organic matter degradation, group and categorize
309 samples and develop indices such as the DI of Dauwe et al. (1999) using summary statistics
310 (see equation 2). A PCA is an orthogonal transformation of a set of variables into a new set of
311 uncorrelated variables called principal components. New axes are chosen in order to explain as
312 much as possible of the variance within the data set on a few main axes of highest correlation.
313 The first component explains most of the variance within the data set, consecutively followed
314 by the remaining components in the order of their decreasing capacity to explain the variance
315 within the data set. The selection of the most relevant components can be done by selecting
316 those with eigenvalues (the variances of the principal components) > 1 . Alternatively, the kink
317 method can be applied selecting those components from a plot of eigenvalues (scree plot),
318 which describe a steep slope of declining variance followed by a “kink” after which the
319 principal components add only small amounts to the variance. The factor loadings of the
320 variables (in this case the individual amino acids) are their projections on the new axis. The
321 factor score of each data set from a sampling location is obtained by multiplying the
322 standardized data with the factor loadings (also called factor coefficients). A high (low) factor
323 score shows that a sample has high (low) concentrations of the variables with high factor
324 loadings. A plot of factor loadings of the variables compared with a plot of the factor scores of
325 samples helps to visualize the relation of the samples to the variables and thus to identify the
326 processes behind the results of the PCA.

327 2 Results

328 3.1 Organic carbon, nitrogen and amino acid content

329 The POC (N) content is 35.9 % (5.9%) in plankton and 1.65-46.4 % (0.21-10.14 %) in sediment
330 trap samples. In sediments, POC (N) contents drop to 0.10-13.5 % (0.02-1.72 %). SPM has
331 POC (N) contents of 0.94-45.4 % (0.09-12.08 %). DOC content in water samples is between
332 0.5-1.1 mg L⁻¹ and DOC in pore water samples is between 3.9-29 mg L⁻¹.



333

334 Figure 2: Box and Whisker plots of AA content in nmol g⁻¹ or nmol L⁻¹ (a) and of AAC/C %
335 (b) in SPM from water depths <200m (SPM<) and >200m (SPM>), in sediment traps at water
336 depth <200m (Trap<) and >200 m (Trap>), in sediments (Sed), in water samples (Water) and
337 in pore water samples (Pore W). Boxes comprise the upper and lower quartile and lines indicate
338 median; whiskers delineate the 10 and 90 percentile; outliers are marked by dots; some outliers
339 above the 90 percentile are cut off in the figure for better perceptibility of trends.


340

341 AA contents are grouped into SPM and trap samples taken at water depths <200m (shallow)
342 and >200m (deep) (Fig. 2, Table 1). AA contents are highest in SPM samples and shallow
343 sediment traps (<200m water depth) with values between 40-4307 $\mu\text{mol g}^{-1}$ (Fig. 2) and
344 averages of 662-908 $\mu\text{mol g}^{-1}$ (Table 1). AA contents are lower in traps from water depth >200m
345 with an average of 164 $\mu\text{mol g}^{-1}$. Sediments have lowest AA contents of all particulate matter
346 samples with an average of 50 $\mu\text{mol g}^{-1}$ (Table 1). TDAA concentrations are between 0.6-44
347 $\mu\text{mol L}^{-1}$ and AA contents are lower in water than in pore water samples with averages of 3.2
348 and 8.8 $\mu\text{mol L}^{-1}$, respectively.

349 The AAC/C is between 5.4-66 % in SPM and traps samples and the AAN/N (not shown) is
350 between 3.7-100 %. The overall pattern found for AAC/C (Fig. 2b) is similar to the pattern of

351 AA contents (Fig. 2a) but there is more overlap of AAC/C between the different groups.
 352 Sediments have AAC/C between 2.7-50 % and AAN/N between 3-78 % (not shown). The
 353 contribution of AAC to DOC (AAC/C) in water samples is between 4-40 % and in pore water
 354 samples between 0.5-9 %.

355 AA contents of sinking and suspended particles decrease with water depth and the most
 356 significant decrease occurs in the upper ocean (Fig. 3a). The decay constant of AA of sinking
 357 particles is twice as high as the decay constant of AA of SPM (Fig. 3 a, b). Kara Sea samples
 358 were excluded from these calculations as their AA contents are low due to the strong dilution
 359 by material from rivers and resuspended sediments in this near-shore environment (see 3.2 and
 360 4.2). It is also notable that AAC/C and AAN/N (not shown) significantly decrease between
 361 shallow and deep traps and from deep traps to sediments while AAC/C of SPM show little
 362 decrease between shallow and deep samples (Fig. 2b).




363
 364 Figure 3: AA contents in nmol g^{-1} in sediment traps (a) and SPM (b). Red triangles mark
 365 samples from the Kara Sea, black dots are samples from the other trap and SPM locations shown
 366 in Figure 1. The decay functions are calculated from samples excluding Kara Sea samples.

367
 368
 369

370 **3.2 Amino acid composition**

371 While the AA contents of sinking particles from traps show a distinct decrease with water
 372 depths the changes in AA composition are lesser so that we averaged all AA spectra irrespective
 373 of water depths and area of study to compare them with plankton and sediment samples (Figure
 374 4a). Dominant AA in plankton samples are Glu, Gly, Ala and Asp. Sinking particles and
 375 sediments are also dominated by these AA but in the order of Gly, Asp, Glu and Ala (Fig. 4a).
 376 Mol% Asp, Gly, β -Ala, γ -Aba and Orn increase from plankton via sinking particles to sediments
 377 while Mol% of Glu, Ala, Val, Met, Ile, Leu, Tyr and Phe decrease (Fig. 4a). These trends are
 378 further continued with depths in sediment cores (not shown).
 379




380

381 Figure 4: Average concentrations of individual AA (Mol%) and 1 σ standard deviation (vertical
 382 bars) in plankton (green), sediment trap (red) and sediment (black) samples (a), in SPM from
 383 water depths <200m (green), >200m (red) and TDAA in water samples (black) (b). Asterisks
 384 mark the AA with increasing Mol% from plankton via sediment trap samples to sediments (a)
 385 and from shallow SPM via deep SPM to TDAA of water samples.
 386

387 AA composition of SPM shows a clear trend with water depth with enrichments of Mol% Ser,
 388 Glu, Gly, Orn and His and decreases of almost all other AA from shallow waters (<200 m) to
 389 deep waters (>200m) and decreases are most pronounced for Asp, Thr and Lys (Fig. 4b). These
 390 trends are partly resumed by TDAA in water samples with a further enrichment of Mol% Ser,
 391 Orn and His, while Mol% Glu and Gly slightly decrease in water samples compared with deep
 392 SPM (Fig. 4b). AA spectra of pore waters (not shown) are very similar to water samples.

393



394
 395 Figure 5: Box and Whisker plot of AA/HA ratios (a) and Gluam/Galam ratios (b), RI (c) and
 396 DI (d) in SPM from water depths <200m (SPM<) and >200m (SPM>), in sediment traps at
 397 water depth <200m (Trap<) and >200 m (Trap>), in sediments (Sed), in water samples (Water)
 398 and in pore water samples (Pore W). Outliers are marked by dots; some outliers above the 90
 399 percentile are cut off for better perceptibility of trends. Note logarithmic scales of AA/HA (a)
 400 and Gluam/Galam (b).

401 Biogeochemical indicators reveal the subtle depth dependent trends in sediment traps and
402 therefore the biogeochemical indicators were averaged separately for shallow and deep
403 sediment trap and SPM samples (Fig. 5, Table 1). The AA/HA in SPM and water samples -
404 with averages between 80.2 and 204.6 - are higher than in traps and sediments (Fig. 5a, b; Table
405 1). The AA/HA decrease from shallow via deep traps to sediments with averages of 25.4, 14.9
406 and 9.1, respectively. The Gluam/Galam is highest in SPM samples, slightly lower in shallow
407 and deep traps and lower in sediments, water and pore water (Fig. 5a, b; Table 1). The RI (Fig.
408 5c; Table 1) shows the same pattern as the ratios of Asp/ β -Ala and the Glu/ γ -Aba (Table 1),
409 with no clear trend between shallow and deep SPM samples and decreases from shallow to deep
410 traps and further to the sediments. Water samples have similar values as sediment samples with
411 average RI of 1.8 and 1.6, respectively, and pore waters have an even lower average RI of 0.9.
412 Similar to the RI the DI is within the same range in shallow and deep SPM samples and the
413 mean values are very close (Fig. 5d; Table 1) while the DI decreases from shallow sediment
414 traps via deep traps to sediments. In contrast to the RI where water samples have lowest values
415 the highest DI values are found in water and pore water samples.

416 In summary, common biogeochemical indicators of organic matter degradation (RI, Asp/ β -Ala,
417 Glu/ γ -Aba) and bacterial OM accumulation (AA/HA, Gluam/Galam) drop and thus imply
418 increasing degradation between shallow and deep sediment traps and between deep traps and
419 sediments, while these indicators reveal little or no degradation with depth in SPM as the
420 patterns of relative enrichment vs. decreases found in SPM and partly also in TDAA of water
421 and pore water samples differ from the degradation pathway of common biogeochemical
422 indicators (Fig. 4, 5; Table 1). The enhanced DI values, furthermore, imply that water and SPM
423 samples are less degraded than deep trap and sediment samples and that TDAA in water and
424 pore water samples are least degraded.

425

426

427

428

429

430

431 Table 1: Mean values and standard deviation (Stdev.) of POC [%], DOC [mg/L], amino acid
 432 (AA) content [$\mu\text{mol/g}$ or $\mu\text{mol/L}$], AAC/C%, AAN/N%, ratios of Asp/ β -Ala, Glu/ γ -Aba,
 433 AA/HA and Gluam/Galam, the RI, DI, SDI*, RTI* and ox/anox ratio summarized in traps at
 434 <200m and >200m water depth, sediments, SPM <200m and >200m water depth, water samples
 435 and pore water samples. *definition of these indicators in part 4.2 below.

		Trap <200m	Trap >200m	Sediment	SPM <200m	SPM >200m	Water	Pore Water
POC; DOC [%; mg/L]	Mean	13.6	5.3	1.8	14.9	10.6	0.8	13.0
	Stdev.	± 11.4	± 1.9	± 2.2	± 8.3	± 4.8	± 0.2	± 8.3
Amino Acids [$\mu\text{mol/g}$; $\mu\text{mol/L}$]	Mean	631.3	164.1	49.8	907.7	661.9	3.2	8.8
	Stdev.	± 602.0	± 93.8	± 82.3	± 637.6	± 434.7	± 3.2	± 7.9
AAC/C [%]	Mean	26.5	15.5	10.8	33.9	32.7	10.1	2.8
	Stdev.	± 8.6	± 4.2	± 5.9	± 12.6	± 10.5	± 6.5	± 2.6
AAN/N [%]	Mean	57.0	38.3	24.2	65.6	61.6		
	Stdev.	± 14.8	± 8.8	± 12.8	± 18.4	± 26.4		
Asp/ β -Ala	Mean	71.8	19.2	10.5	57.9	47.3	10.2	2.6
	Stdev.	± 63.4	± 19.5	± 6.6	± 64.7	± 44.9	± 14.7	± 2.4
Glu/ γ -Aba	Mean	45.7	12.9	8.5	103.6	105.3	8.5	8.8
	Stdev.	± 38.8	± 7.2	± 6.9	± 122.7	± 69.6	± 7.0	± 10.2
AA/HA	Mean	25.4	14.9	9.1	84.6	204.6	80.2	106.8
	Stdev.	± 14.7	± 6.6	± 4.6	± 42.7	± 179.1	± 65.2	± 142.7
Gluam/Galam	Mean	3.2	3.6	1.2	3.0	2.1	1.6	0.3
	Stdev.	± 3.3	± 1.1	± 0.3	± 1.8	± 1.7	± 2.4	± 0.2
RI	Mean	15.1	3.9	1.8	18.8	15.7	1.6	0.9
	Stdev.	± 16.4	± 1.5	± 2.2	± 10.6	± 8.3	± 1.8	± 0.7
DI	Mean	0.9	0.3	-0.5	1.0	1.1	2.1	1.5
	Stdev.	± 0.8	± 0.3	± 0.8	± 0.1	± 0.3	± 1.1	± 0.9
SDI	Mean	1.1	0.0	-0.9	0.8	0.8	-0.7	-0.8
	Stdev.	± 0.2	± 0.2	± 1.0	± 0.2	± 0.3	± 0.4	± 0.3
RTI	Mean	0.7	0.1	0.6	0.2	-1.0	-2.8	-2.7
	Stdev.	± 0.3	± 0.3	± 0.4	± 0.5	± 0.4	± 0.6	± 0.4
ox/anox	Mean	1.2	1.3	2.2	1.1	1.0	0.8	0.7
	Stdev.	± 0.3	± 0.2	± 1.1	± 0.1	± 0.1	± 0.3	± 0.3

436

437

438 4 Discussion

439 4.1 Changes during organic matter degradation

440 Our summary of AA data from various locations in the world ocean corroborates earlier
 441 findings that degradation of organic matter by zooplankton and microbes imparts characteristic
 442 changes to AA spectra so that the differences in AA composition are much larger between types
 443 of organic matter and from different water and sediment depths than between different oceanic
 444 areas (Lee, 1988). The AA spectra track the successive degradation of organic matter during
 445 sedimentation from the plankton source via sinking particles, their incorporation into sediments
 446 and their further degradation after burial. The most characteristic changes along this
 447 sedimentation pathway are the relative enrichments (in Mol%) of Gly, Asp and the non-protein
 448 AA β -Ala, γ -Aba and Orn and the relative decrease of AA produced by fresh plankton such as
 449 Glu, Ala, Val, Met, Ile, Leu, Tyr and Phe (Fig. 4a). These changes are depicted by the common

450 biogeochemical indicators: the ratios of proteinaceous AA vs. non-protein AA (RI and Glu/ γ -
451 Aba) decrease along this pathway. Asp/ β -Ala ratios also decrease because β -Ala becomes
452 relatively more enriched than Asp. The DI, originally derived from sediment samples of
453 different degradation states (Dauwe et al., 1999; Dauwe and Middelburg, 1998), decreases from
454 positive values in fresh plankton and most sinking particles to negative values in sediments as
455 it integrates the products of Asp and Gly multiplied with negative factors, and the products of
456 Glu, Met, Ile, Leu, Tyr and Phe multiplied with positive factors (Dauwe et al., 1999).

457 In contrast, the AA in SPM evolve along a different path than the sedimentation pathway (Gaye
458 et al., 2013b). The increases in Mol% Ser, His and Orn and the decreases in Mol% Asp from
459 shallow (<200 m) to deep SPM (>200m) and even further in TCAA of sea water and pore water
460 are either not depicted or even contrary to the trends depicted by the common biogeochemical
461 indicators (Fig. 4b). The striking difference in AA distribution of SPM (Fig. 4) on the one hand
462 and sinking particles and sediments on the other hand as well as the different depth dependent
463 trends (Fig. 4, 5) suggest that there is little exchange between the two types of particles in the
464 ocean. Sinking particles build up sediments and the degradation pathways evident in the water
465 column - namely the accumulation of degradation products and acidic AA often absorbed to
466 carbonates - continue in the sediments. Novel biogeochemical indicators are required for SPM
467 and possibly also DOM to characterize their AA changes.

468

469 **4.2 Results of a PCA: two new biogeochemical indicators**

470 A PCA of individual AA (Mol %, Fig. 6a) of all samples compiled in this study results in two
471 factors which explain 59 % of the total variance within the data set. The first factor delineates
472 the well-known changes along the degradation pathway from plankton via sinking particles to
473 sediments. Phe, Ile, Leu, Glu and Tyr (enriched in fresh plankton) have the highest F1 loadings
474 while Asp, β -Ala and γ -Aba (accumulating during degradation) have the lowest negative F1
475 loadings. Highest F2 loadings are found for Asp, Thr, Lys and Val while Gly, Orn, His and Ser
476 have the most negative F2 loadings. Factor scores of the individual samples (Fig. 6b) plot in a
477 triangular shape with plankton and fresh organic matter from surface waters at the apex with
478 highest F1 and F2 scores. The diverging sides of the triangle mark sinking particles and
479 sediments decreasing in F1 scores on one side and SPM with decreasing F2 scores on the other
480 side (Fig. 6b). Similar trends were observed in earlier studies based on local data sets (Nagel et
481 al., 2016; Gaye et al., 2013b). That samples from greatly different environments reveal the same


482 divergence between sinking particles and SPM with only little overlap (Fig. 4) suggest a general
483 mechanism operating globally. Most of the overlap encompasses SPM samples from the Kara
484 Sea and sediment traps all sampled at water depths below 100 m. The Kara Sea is characterized
485 by sediment resuspension related to strong riverine input in combination with sea ice dynamics
486 so that many of the Kara Sea SPM and trap samples are mixed with resuspended sediments
487 (Gaye et al., 2007; Unger et al., 2005). TDAA analyzed in water and pore water form a cluster
488 with significantly different AA composition from particulate matter, but instead recapitulating
489 the enrichments of Mol% Ser, His and Orn observed in SPM (Fig. 6b).

490 The precise separation of the degradation pathway of sinking particles and sediments from SPM
491 and DOM by the PCA suggests that we can use the first factor (F1) to calculate a new sinking
492 particle and sediment degradation index (SDI)

$$493 \quad SDI = \sum_i \left(\frac{var_i - AVG.var_i}{STD.var_i} \right) \times Loadings . F1_i \quad (4)$$

494 where var_i is the original mole percentage of each AA_i , $AVG.var_i$ and $STD.var_i$ are the mean and
495 standard deviations, respectively, and $Loading.F1_i$ is the factor loading of the first axis (F1) of
496 the PCA of the individual amino acid $_i$ shown in Table 2. Most of the F1 loadings resemble those
497 of the DI of Dauwe et al. (1999) (Table 2) and the SDI and DI thus are significantly correlated
498 (Table 3).

499 The second factor (F2) - normalized in the same way with the averages and standard deviations
500 of the same PCA - can be used as an indicator of changes in the AA composition of SPM
501 possibly related to the residence time or renewal time of the water mass they are transported
502 with (see discussion in 4.2.2). With longer residence time in the ocean the organic matter in
503 SPM is likely to become more recalcitrant and the indicator and is therefore named residence
504 time index (RTI)



505

506 Figure 6: Results of a PCA of AA (Mol%) of all samples of this study with factor loadings of
 507 amino acids for the first and second factor (a) and factor scores of samples (b). Small arrows in
 508 (a) point to the positions of Val and Ile, respectively. Arrows in (b) indicate progressive
 509 deviation in composition from the plankton source, essentially with increasing water and
 510 sediment depths.

511

$$RTI = \sum_i \left(\frac{var_i - AVG.var_i}{STD.var_i} \right) \times Loadings.F2_i \quad (5)$$

and is calculated in the same way as the SDI but the factor loadings of the second axis (F2) of the PCA of the individual amino acid_i (Table 2) is inserted for the term *Loadings.F2_i*.

The SDI allows a separation of trap samples from shallow water depth from those of greater depths (Figure S1). All samples from deep sediment traps have SDI values below 0.5. Likewise, SPM from >200 m depths have lower RTI than most of the samples from shallower depths. Deep trap samples and deep SPM samples form two clearly separated clusters with different SDI and RTI (see Figure S1 for further details).

Table 2: Factor loadings of F1 and F2 for calculating the SDI and RTI, respectively, average [Mol%] and standard deviations (Std. Dev.) of AA of samples used for the PCA shown in Figure 5 in comparison with the factor loadings (named factor coefficients) of the DI published by Dauwe et al (1999) and their averages [Mol%] and standard deviations used for the DI based on 28 sediment samples.

Amino Acid	Loadings F1 SDI	Average [Mol%]	Std. Dev.	Loadings F2 RTI	DI	DI Average [Mol%]	DI Std. Dev.
Ser	0.067	8.7	3.6	-0.210	0.015	7.2	1.9
His	0.014	2.4	2.3	-0.166	0.158	1.0	0.8
Orn	-0.038	1.2	1.9	-0.152	-	-	-
Tyr	0.162	1.9	1.1	-0.075	0.178	2.1	1.2
Gly	-0.023	16.2	2.8	-0.068	-0.099	17.6	3.8
Glu	0.132	10.5	2.5	-0.001	0.065	10.0	2.3
Leu	0.157	5.9	1.9	0.027	0.169	6.6	1.5
Ile	0.148	3.8	1.0	0.028	0.139	4.5	0.8
β-Ala	-0.161	1.3	2.0	0.036	-	-	-
Phe	0.140	2.9	1.0	0.047	0.134	3.2	1.2
γ-Aba	-0.164	1.0	1.6	0.064	-	-	-
Ala	-0.030	9.1	1.5	0.080	-0.043	11.8	0.8
Arg	0.012	4.2	1.1	0.117	-0.115	6.1	2.3
Val	0.020	5.7	1.1	0.132	-0.044	7.6	1.1
Lys	0.026	5.1	1.2	0.135	-	-	-
Thr	-0.074	5.7	1.2	0.174	-0.129	7.1	1.5
Asp	-0.154	12.2	4.3	0.179	-0.102	13.4	2.7

526

4.2.1 The SDI as an indicator of degradation and oxic vs. anoxic diagenetic conditions of sinking particles and sediments

In order to test the performance of our new degradation indices, we separated SPM samples from sinking particles and sediments and correlated the common biogeochemical indicators and

531 individual AA (Mol %) of SPM with the RTI of individual samples while we correlated the
532 same variables of sinking particles and sediments with the SDI (Table 3). We assume that
533 correlations with Pearson correlation coefficients $R > 0.50$ can be considered as “strong
534 correlations” (Cohen, 1988). The SDI correlates moderately to strongly with the common
535 degradation indicators and the best positive correlation is found between SDI and the DI (Table
536 3). The strong correlation among the degradation indicators with POC contents indicates that
537 this common and often measured variable is a good indicator of relative organic matter quality
538 in sinking particles and sediments and all other degradation indices do not perform better than
539 POC concentrations (see correlation coefficients in Table 3). The DI and the SDI, which are to
540 some extent interchangeable, allow a fine tuning of degradation intensities. The most significant
541 negative correlation of the ox/anox ratio with the SDI is preconditioned, as the ox/anox is the
542 quotient of AA enriched by degradation to those enriched in fresh plankton. It should be noted
543 that this negative correlation is even better than the positive correlation of the DI and the SDI.
544 A close look at the SDI and ox/anox in sediment samples suggests that the SDI can be used to
545 distinguish between oxic and anoxic diagenetic conditions (Fig. 7). The sediment samples
546 deposited in regions of bottom water anoxia (surface samples from Namibian shelf at < 200 m
547 depths; a core from the Arabian Sea slope at 775 m) have lower ox/anox ratios and distinctly
548 higher SDI values compared with the samples from similar depths and oxygenated bottom water
549 (e.g. Mediterranean Sea, Kara Sea) (Fig 7a). SDI performs better than the DI to determine
550 diagenetic conditions as the DI less significantly correlates with the ox/anox indicator (Figure
551 S2). The best fit between the SDI and the ox/anox in sediment samples is an exponential
552 correlation with a correlation coefficient of $R = -0.95$ while the DI and the ox/anox correlate
553 exponentially with an $R = -0.79$ (Fig. 7a, Figure S2). The SDI better depicts the spectral changes
554 in samples deposited under anoxic diagenetic conditions such as those from the Namibian shelf
555 (Nagel et al., 2016) and the Arabian Sea mid-water oxygen minimum zone (Suthhof et al., 2001)
556 while the ox/anox ratio better resolves variations in samples of strong oxic degradation so that
557 the SDI is in fact better suited to determine the threshold of anoxic vs. oxic diagenesis. Another
558 indication of this quality of the SDI is that the anoxic sediments have SDI values in the range
559 of sediment trap samples from the water column while they decrease under oxic diagenetic
560 conditions (Fig. 7b). Further, the SDI also correlates with other indicators of oxic vs. anoxic
561 conditions. The core SO90-111 KL from within the mid-water oxygen minimum impinging on
562 the Pakistan margin, was used to reconstruct changes in oxygenation during the last 60 ka BP
563 based e.g. on $\delta^{15}\text{N}$ values of total N (Suthhof et al., 2001). The $\delta^{15}\text{N}$ values fluctuated between

564 enhanced values in warm phases due to denitrification in the mid-water oxygen minimum and
 565 lower values in cold phases when the oxygen minimum zone was weaker or absent (Suthhof et
 566 al., 2001). The SDI very precisely tracks these changes (Fig. 7c) and in accordance with the
 567 threshold discernable in Fig. 7a and b we propose that the divide between oxic and anoxic
 568 diagenetic conditions is at SDI values between 0 and -0,2 with SDI<-0.2 indicating oxic and
 569 SDI >0 indicating anoxic diagenetic conditions (Fig. 7a, b) and we propose to use this indicator
 570 to reconstruct redox conditions from sediment cores. The work of (Carr et al., 2016) - relying
 571 on the DI – even suggests that signals of changes in redox conditions can be preserved even
 572 down to 200 m core depth.

573

574 Table 3: Pearson correlation coefficients of the SDI, RI, DI, Asp/β-Ala and Glu/γ-Aba with
 575 selected AA*, the RTI, AAC/C, AAN/N, AA ratios and degradation indices, water depth
 576 (Depth), POC and TN contents (%) and AA content (nmol/g) in sediment trap and sediment
 577 samples (column 2-6). Pearson correlation coefficients of the RTI, RI, DI, Asp/β-Ala and Glu/γ-
 578 Aba with selected AA, the SDI, AAC/C, AAN/N, AA ratios and degradation indices, water
 579 depth (Depth), POC and TN contents (%) and AA content (nmol/g) in SPM samples (column
 580 8-12).


581 * Only AA with a correlation coefficient $R \geq 0.50$ with at least one of the indicators are shown.

1	2	3	4	5	6	7	8	9	10	11	12
Traps and Sediments	SDI	RI	DI	Asp/β-Ala	Glu/γ-Aba	SPM	RTI	RI	DI	Asp/β-Ala	Glu/γ-Aba
Asp	-0.68	-0.30	-0.83	-0.25	-0.32	Asp	0.81	0.00	-0.61	0.13	-0.17
Thr	0.12	-0.25	-0.25	0.03	0.07	Thr	0.86	-0.02	-0.57	0.06	-0.16
Ser	0.32	0.03	0.16	0.09	0.16	Ser	-0.91	-0.15	0.08	-0.11	-0.01
Gly	-0.21	-0.32	-0.45	-0.31	-0.31	Gly	-0.90	-0.17	0.12	-0.11	0.04
Ala	0.46	0.07	0.17	0.03	0.11	Ala	0.91	0.09	-0.43	0.11	-0.09
Val	0.67	0.31	0.48	0.29	0.31	Val	0.88	0.07	-0.26	0.08	-0.09
Ile	0.87	0.48	0.76	0.52	0.55	Ile	0.68	0.34	0.26	0.26	0.02
Leu	0.88	0.49	0.76	0.52	0.56	Leu	0.65	0.47	0.28	0.35	0.11
Tyr	0.72	0.39	0.76	0.34	0.32	Tyr	-0.14	0.27	0.66	0.02	0.23
Phe	0.89	0.38	0.79	0.40	0.44	Phe	0.78	0.37	0.04	0.29	0.06
β-Ala	-0.85	-0.27	-0.44	-0.32	-0.35	β-Ala	-0.08	-0.66	-0.19	-0.52	-0.12
γ-Aba	-0.79	-0.25	-0.37	-0.28	-0.35	γ-Aba	0.24	-0.53	-0.65	-0.20	-0.40
Lys	0.56	0.14	0.43	0.15	0.15	Lys	0.71	0.05	-0.20	0.03	-0.06
SDI		0.45	0.82	0.46	0.51	SDI	0.07	0.49	0.83	0.23	0.28
RTI	0.04	-0.01	-0.26	0.03	-0.05	RTI		0.10	-0.34	0.12	-0.08
AAC/C%	0.56	0.51	0.53	0.56	0.56	AAC/C%	-0.08	0.14	0.22	0.06	0.09
AAN/N%	0.55	0.40	0.49	0.47	0.47	AAN/N%	-0.05	0.06	0.03	-0.06	0.08
AA/HA	0.54	0.58	0.54	0.73	0.75	AA/HA	-0.57	0.11	0.32	-0.01	0.16
Gluam/Galam	0.36	0.43	0.44	0.40	0.35	Gluam/Galam	0.27	0.43	0.16	0.28	0.08
Asp/β-Ala	0.46	0.93	0.57		0.85	Asp/β-Ala	0.12	0.74	0.10		0.07
Glu/γ-Aba	0.51	0.93	0.56	0.85		Glu/γ-Aba	-0.08	0.37	0.24	0.07	
RI	0.45		0.49	0.93	0.93	RI	0.10		0.33	0.74	0.37
DI	0.82	0.49		0.51	0.56	DI	-0.34	0.33		0.10	0.24
ox/anox	-0.91	-0.30	-0.71	-0.31	-0.37	ox/anox	0.67	-0.19	-0.50	-0.10	-0.09
Depth	-0.67	-0.28	-0.54	-0.28	-0.37	Depth	-0.55	-0.12	0.23	-0.11	0.04
POC	0.57	0.79	0.67	0.77	0.76	POC	0.27	0.30	0.13	0.25	-0.03
N	0.45	0.46	0.50	0.39	0.39	N	0.29	0.30	0.13	0.24	-0.01
AA	0.51	0.84	0.60	0.79	0.76	AA	0.17	0.28	0.15	0.20	0.01

582

583

584



585

586 Figure 7: SDI indicator plotted against the ox/anox ratio with dots indicating oxic sediments
 587 (black) and suboxic to anoxic sediments from Namibia (green) and the Arabian Sea (blue), the
 588 red line marks the exponential fit to all sediment samples with an $R=0.95$ (a). SDI plotted with
 589 water depth (in m) of sediment trap deployment (red) and of sediment sampling (see color code
 590 of a) (b). SDI and the $\delta^{15}\text{N}$ of total N with sediment depth in sediment core SO90-111 KL
 591 correlated with an $R=0.71$ (c); the blue bar marks the threshold of the SDI delimiting oxic and
 592 anoxic diagenetic conditions at an SDI value of about -0.1.

593

594 4.2.2 The RTI as an indicator of suspended matter residence time

595 Changes in SPM composition between shallow and deep waters (Fig. 4b) are depicted by a
 596 decrease of the RTI which is due to the relative depletion of Asp, Thr, Lys and Val with highest
 597 positive factor loadings and the enrichment of Ser, His and Orn with the most negative factor
 598 loadings (Table 2). Both, Ser and Gly (Mol %) are strongly linearly anticorrelated with the RTI
 599 ($R=-0.91$ and -0.90) showing that they can be used instead of the RTI to characterize SPM if
 600 not all AA used for the RTI can be measured. The anticorrelation of the RTI with water depths
 601 ($R=-0.55$; Table 3) is due to the RTI decrease in the upper 200 m only. Below this depth there
 602 is no further trend in the RTI and values scatter between -0.5 and -1.5 (Fig. 8).

603 The results of our PCA also show that below 200 m SPM becomes distinctly decoupled from
604 sinking aggregates (Fig. 7, Figure S1). Both, sinking particles and SPM sampled in the upper
605 ocean mixed layer and euphotic zone resemble fresh plankton whereas below the surface mixed
606 layer they follow different pathways and the chance that SPM and sinking particles interact
607 obviously decreases with water depth which could be due to the scarcity of both types of
608 particles in the deep ocean (McCave, 1984). Our AA results support previous studies on SPM
609 (using e.g. thorium isotopes, radiocarbon and biomarkers) which found that the interaction
610 between sinking particles and SPM by aggregation and disaggregation strongly decreases from
611 the euphotic zone to the meso- and bathypelagic zone and that SPM rather interacts with DOM
612 due to the long residence time of both in the deep ocean (Lam and Marchal, 2015). The observed
613 constant AA composition of SPM below 200 m water depth could be explained by a recalcitrant
614 nature making the AA barely accessible to further microbial degradation. Alternatively, and in
615 analogy to observations of DOC, it could be due low concentrations of SPM which rather than
616 their recalcitrance limits prokaryotic growth and thus organic matter degradation in the deep
617 ocean (Arrieta et al., 2015). The age of the water masses in the upper ocean mixed layer is less
618 than 100 years while deeper waters have ages of several 100 years to maxima of 1600 years in
619 the deep Indian and Pacific Oceans (England, 1995; Gebbie and Huybers, 2012). The constantly
620 low RTI below the mixed layer is thus related to the long residence time of deep SPM in the
621 deep ocean. An earlier study in fact suggested a relationship between AA degradation and their
622 residence time using detailed water mass ages available from the Atlantic and Pacific
623 subtropical gyres at stations BATS and HOT, respectively (Kaiser and Benner, 2009).

624 Solubilization of particulate matter by exoenzymes and the subsequent uptake in dissolved form
625 (Carlson and Hansell, 2015; Aristegui et al., 2009) leads to an almost complete turnover of
626 originally diverse surface derived organic matter. It is thus feasible that bacterial biomass
627 comprises a large amount of organic matter in compartments of long residence times. However,
628 fresh bacteria and fungi have quite similar AA composition as plankton (Cowie and Hedges,
629 1992) while SPM AA composition is fundamentally different. The high AA/HA ratios not
630 having a clear trend with water depth also suggest that the contribution of bacterial biomass to
631 SPM is small and does not increase with water depth (Table 1; Fig. 5a, b). The observed changes
632 in SPM are thus more likely related to adsorption processes and macromolecule formation of
633 material not digestible to deep sea organisms and resistant to their enzymes. DOM was shown
634 to become adsorbed to mineral surfaces (Keil and Kirchman, 1993; Keil and Kirchman, 1994;
635 Keil et al., 1994; Arnarson and Keil, 2005, 2007). However, degradation of adsorbed AA

636 proceeds on particles (Satterberg et al., 2003; Taylor, 1995). Thus, the constant AA composition
637 in SPM at depths >200 m may indicate that SPM is in equilibrium with TDAA which likewise
638 show no depth dependent changes in AA composition (Figure S3). Feasible candidate processes
639 to explain the homogeneity are AA scavenging by SPM or formation of gels (3D networks =
640 biopolymers) which can anneal to larger sizes so that part of the dissolved AA can be passed
641 from the dissolved to the particulate organic carbon pool (Druffel and Williams, 1990; Orellana
642 and Leck, 2015). This process is, however, reversible so that there is probably an exchange
643 between the gel and particulate matter phase as well as between gels of different sizes and
644 complexities depending on pH, temperature, the presence of ligands, pollutants or UV radiation
645 (Orellana and Leck, 2015). Generally, hydrophobic AA (Ala, Val, Met, Ile, Leu, Phe, Pro, Trp)
646 and aromatic AA (His, Tyr) are more likely to form gels and aggregates (Orellana and Leck,
647 2015). Our results indicate that an equilibrium may be attained between the dissolved phase
648 and SPM after a relatively short time so that the AA composition of SPM is constant below 200
649 m water depth. If there is no further significant scavenging of SPM by sinking particles and no
650 degradation of AA on SPM, their abundance could increase due to further adsorption of DOM
651 with increasing age on the ocean conveyor belt. However, large zooplankters may be able to
652 utilize the SPM pool (Koppelman et al., 2009; Gloeckler et al., 2018; Hannides et al., 2013)
653 and further studies are required to elucidate the fate of SPM in the ocean.

654

655 **4.3 Contents and composition of total dissolved amino acids in sea water and pore water**

656 The spectra of TDAA sampled in the oligotrophic Indian Ocean Subtropical Gyre are uniform
657 with water depth which is also reflected by the RTI. Further, the difference between water and
658 pore water samples is small and Ser, Gly and His are uniformly the major TDAA in sea water
659 and pore water. It is possible that the selective accumulation of these AA in the dissolved phase
660 is due to their excretion or their association with exoenzymes. Ser is present in N-acyl
661 homoserine lactone (AHLs) which is a class of bacterially produced signaling molecules
662 involved in bacterial quorum sensing; these compounds serve to regulate growth by changing
663 gene expressions, for example, in order to influence population density or phenotype (Parsek et
664 al., 1999; Klein et al., 2009). His changes from its protonated to deprotonated form at a pH of
665 6 and is therefore often present at the active sites of enzymes. Ser and Gly may simply remain
666 dissolved in sea water as they are hydrophilic. Once mixed into the deeper ocean the scarcity
667 of bacteria or the incorporation of AA into gels could be the reason for their recalcitrance.

668 However, we do not assume that a considerable part of the TDAA belong to dissolved free AA.
669 Because the differences between samples from different regions are much smaller than the
670 difference between the molecular weight fractions and sea water vs. pore-water (Figure S3), we
671 surmise that the formation and transformation processes of DOC are very uniform in the ocean.
672 This assumption is based on limited data so that these results are rather preliminary. We also
673 do not have enough spatial coverage of SPM and TDAA data in the deep ocean to detect AA
674 utilization by organisms or sorption and desorption processes. Both these organic matter pools
675 are large (see below), so that such investigations are important to estimate the possible role of
676 these pools in oceanic carbon sequestration and the reactions to global change (Ridgwell and
677 Arndt, 2015)(Lonborg et al., 2018).


678 Our AA yield with AA-C/C of $10.1 \pm 6.5\%$ are in the high range of studies from the literature.
679 In some of the previous studies AAC/C was between 0.4 and 4 % with a reduction from 1-4 %
680 AAC/C in surface waters to 0.4-0.8 % in waters >1.000 m. This reduction was moreover,
681 associated with a progressive AA degradation reflected in the DI and a DOM specific
682 degradation indicator at some of the sampling stations (Kim et al., 2017; Davis and Benner,
683 2005; Kaiser and Benner, 2009). These lower yields may however, be due to different
684 hydrolysis conditions as these studies used water vapor hydrolysis at higher temperatures
685 (150°C) but for a much shorter duration (32.5 minutes). Studies using the same hydrolysis
686 conditions as this work reported AAC/C of 5-10 % (Ittekkot, 1981; Keil and Kirchman, 1999;
687 McCarthy et al., 1997).

688

689 **4.4 $\delta^{15}\text{N}$ values in sinking and suspended matter and evidence for nitrogen sources and** 690 **transformation processes**

691 The $\delta^{15}\text{N}$ values in sediments can preserve information on N sources throughout the geological
692 history (Sun et al., 2019; Gaye et al., 2018; Kienast et al., 2008). However, $\delta^{15}\text{N}$ values may be
693 modulated by organic matter cycling and diagenetic processes which are replicated and thus
694 traceable in the AA composition not least because AA are the main identifiable contributors to
695 N in particulate organic matter. The increase of $\delta^{15}\text{N}$ values by about 2 ‰ on average during
696 organic matter burial and early diagenesis in the upper sediments (Robinson et al., 2012; Tesdal
697 et al., 2013) is corroborated by a parallel shift in AA based degradation indicators (Gaye-Haake
698 et al., 2005; Möbius et al., 2010). In contrast to sediments, there are no clear depth related trends
699 in $\delta^{15}\text{N}$ values of sinking particles in the water column of the epi- to mesopelagic ocean (Gaye-

700 Haake et al., 2005; Yang et al., 2017; Altabet, 2006). AA based biogeochemical indicators
 701 revealed degradation with depth at specific trap sites (Haake et al., 1993a) and $\delta^{15}\text{N}$ analyses of
 702 individual amino acids showed that degradation is proceeding on sinking particles with $\delta^{15}\text{N}$
 703 changes of “trophic” AA while $\delta^{15}\text{N}$ of “source” AA remained constant (McCarthy et al., 2007).
 704 However, degradation of sinking particles is much smaller than degradation at the sediment
 705 water interface and in our large data set that integrates many different areas of study the small
 706 to moderate changes in AA degradation are obviously obliterated, as neither AA contents (Fig.
 707 3), nor the SDI (Fig. 7), the AAC/C % (Figure S4) nor AAN/N % (not shown) reveal any
 708 significant trends in sinking particles in the deep ocean.



709
 710 Figure 8: The RTI (black dots) and the $\delta^{15}\text{N}$ values of SPM with water depths [m] (red circles).

711 AA composition of SPM as expressed in the RTI is constant and SPM is rather recalcitrant at
 712 water depths >200 m. Paralleling this, the $\delta^{15}\text{N}$ values of SPM are about 6-8 ‰ on average in
 713 all our studies carried out (Fig. 8). In previous studies $\delta^{15}\text{N}$ values of SPM were reported to

714 increase from ≤ 5 ‰ in surface waters to values between 6-8 ‰ below 200 m water depth which
715 was attributed to organic matter degradation on SPM (Yang et al., 2017; Altabet et al., 1991;
716 Hannides et al., 2013; Emeis et al., 2010). However, SPM samples from the Arabian Sea
717 upwelling area show decreasing $\delta^{15}\text{N}$ values from an average of 8.6 ‰ at water depth above
718 200 m to 7.4 ‰ at depths below 200 m (Gaye et al., 2013b). It is thus reasonable that SPM has
719 a constant $\delta^{15}\text{N}$ value in the mesopelagic and bathypelagic ocean. This is an additional indicator
720 of a common process determining the AA composition and their $\delta^{15}\text{N}$ values of SPM and
721 probably also of DOM sampled below water depths of 200 m (equivalent to an age of ≥ 100
722 years; (England, 1995; Gebbie and Huybers, 2012).

723

724 **4.5 Abundance of amino acids in the ocean**

725 Based on POC, TN and AA fluxes and the area of the open ocean and shallow seas (Costello et
726 al., 2010) we can estimate annual downward fluxes (Table S5). Average POC flux of
727 compilations of trap fluxes were between $1.65 \text{ g m}^{-2} \text{ a}^{-1}$ (Wilson et al., 2012) and $2.74 \text{ g m}^{-2} \text{ a}^{-1}$
728 (Rixen et al., 2019a) while our subset of trap samples from the open ocean (>2000 m water
729 depth) averages to $3.06 \text{ g m}^{-2} \text{ a}^{-1}$. For open ocean traps this results in total fluxes of 0.51-0.94
730 PgC a^{-1} . Our average flux estimates for TN are 0.13 PgN a^{-1} and for AAC are 0.15 PgAA a^{-1} .
731 The flux rates over the shelves and slopes bear, however, large uncertainty because productivity
732 is by several orders of magnitude higher than in offshore areas and spatially variable. Our first
733 estimate, simply based on an average of our fluxes caught in traps deployed in areas of water
734 depth < 2000 m arrives at POC fluxes of 5.4 PgC a^{-1} , TN fluxes of 0.9 PgN a^{-1} and AAC fluxes
735 of $1.36 \text{ Pg AAC a}^{-1}$. Thus 85-90 % of fluxes occur in near shore environments corroborating
736 that 95 % or the total marine organic carbon is buried in these environments (Hedges and Keil,
737 1995). The total sinking fluxes in the proximal plus distal ocean add up to 6.3 PgC a^{-1} , 1.0 PgN
738 a^{-1} and, respectively, $1.51 \text{ Pg AAC a}^{-1}$ (see Table S5 for further details).

739 The largest organic carbon pool in the ocean is DOC with an inventory of $632 \pm 32 \text{ PgC}$ (Carlson
740 and Hansell, 2015; Hansell et al., 2009) and the largest N pool is DON with $77 \pm 23 \text{ PgN}$ (Gruber,
741 2008; Bronk, 2002). Dissolved AA are thus the largest AA pool in the ocean even if AA
742 comprise only a minor amount of DOC. We have only few measurements of AA concentrations,
743 which range between 0.1-0.2 mg/L with an average of 0.16 mg/L in all water samples excluding
744 bottom water. Based on these data we can estimate that AA comprise about $200 \pm 70 \text{ Pg}$ which
745 would contribute about $35 \pm 11 \text{ Pg AAN}$ and about $89 \pm 29 \text{ Pg AAC}$ to the oceanic DON and,

746 respectively, DOC pools. Accordingly, AAC contributes about 14 % to DOC while AAN
747 contributes 45 % to total oceanic DON. This is in the low range of an estimate of 45-86 % AAN
748 based on NMR spectroscopy with acid hydrolysis suggested to recover about half of this AAN
749 pool (Aluwihare et al., 2005).

750 The constant contents and composition of TDAA throughout the ocean indicates that it belongs
751 to the recalcitrant or refractory pool of DOC; this pool is hardly removed in the deep sea and
752 may only be degraded by photochemical reactions as it is returned into surface waters in the
753 course of ocean circulation (Legendre et al., 2015). Our TDAA data reveal no depth dependent
754 trend but our data coverage is not sufficient to detect any spatial variation. The distribution of
755 DOC is, however, well known with its maximum in surface water with 40-80 $\mu\text{mol C kg}^{-1}$ and
756 depletion in deep water with DOC concentrations from $>50 \mu\text{mol C kg}^{-1}$ in the North Atlantic
757 to 39 $\mu\text{mol C kg}^{-1}$ in the North Pacific deep water (Carlson and Hansell, 2015; Hansell et al.,
758 2009). Due to our limited number of measurements we may have missed spatial variations
759 which could elucidate TDAA sources and cycling processes in the ocean as is the case for DOC.
760 Respiration of DOC may be an important removal process in shallower waters (Reinthal et
761 al., 2006) while a large proportion of the DOC reduction on its way to the Pacific on the deep
762 conveyor belt could be related to adsorption to POC, partly via gel formation (Druffel and
763 Williams, 1990).

764 TDAA may be among the constituents of DOC, which interact with SPM as both are transported
765 with their specific water masses by the ocean conveyor belt. Interaction with SPM is suggested
766 by the relative similarity in AA composition of TDAA and SPM. Moreover, SPM carries the
767 second largest pool of POC and AA in the ocean which has not been accounted for in carbon
768 budgets and which role in oceanic biogeochemical cycling has received little attention. The
769 total abundance of POC, TN and AA in SPM can be calculated using average concentrations
770 (Table 1) in the ocean volume between 0-200 m and between 200 m and the sea floor (Costello
771 et al., 2010). These calculations show that there are 443 Pg of total suspended matter in the
772 ocean of which organic carbon comprises 48 PgC, amino acids 35 PgAA and, total nitrogen 6
773 PgN. The relative similarity of AA spectra in SPM and TDAA suggests interaction between the
774 two pools at shallower depths and the build-up of an equilibrium, so that both pools remain
775 constant in concentrations and composition with depths. Like DOC, which was suggested to be
776 recalcitrant in the deep sea (Hansell and Carlson, 2013), SPM may only be affected by
777 degradation and repackaging into aggregates as it is reintroduced into surface water by ocean

778 circulation. Several studies, however, suggest that SPM may be an important food source for
779 deep living zooplankton (Koppelman et al., 2009; Hannides et al., 2013; Gloeckler et al.,
780 2018). If there are no removal processes in the deep ocean, we would expect SPM and their
781 organic constituents to be exported from the Atlantic via the deep ocean circulation and to
782 accumulate in the Pacific.

783

784 **5 Conclusions**

785 The PCA of a set of 1425 samples consisting of sinking particle, SPM, sediment and water
786 samples produced two factors which separate AA in sinking particles and sediments on the one
787 hand from SPM and DOM on the other hand. As the PCA produced two branches diverging
788 with water and, respectively, sediment depth, strong interactions between the sinking and
789 suspended particles pools can be excluded.

790 The relative degradation of sinking particles and sediments, dominated by Gly, Asp, Glu and
791 Ala, can be tracked by a new degradation indicator named Sediment Degradation Index (SDI)
792 derived from the first factor of the PCA and correlated with the often-used degradation index
793 DI. Except the SDI and the DI the other biogeochemical indicators tested here (Asp/-Ala, Glu/-
794 Aba, RI) are not better than POC concentrations for a relative classification of organic matter
795 degradation. The SDI is, moreover, capable to separate oxic and anoxic diagenetic conditions
796 at an SDI between 0 to -0.2 (with values <-0.2 indicating oxic and values >0 indicating anoxic
797 diagenetic conditions). Application of the SDI furthermore, shows that the diagenetic signal
798 from the water column is preserved in sediments deposited under anoxic conditions. The
799 correlation of the SDI with POC shows that anoxic diagenesis enhances POC accumulation in
800 sediments compared to oxic diagenesis.

801 A novel biogeochemical indicator derived from the second factor of the PCA named Residence
802 Time Indicator (RTI) depicts the transformation of SPM enriched in plankton derived AA in
803 the epipelagic ocean to a constant composition in the meso- and bathypelagic ocean. The deep
804 SPM is probably the residue of microbial processing and is not utilizable by enzymes under the
805 present oceanic conditions. This constant composition of SPM is corroborated by a constant
806 $\delta^{15}\text{N}$ value below 200 m irrespective of the area of study.

807 DOM has constant AA composition throughout the water column, dominated by Ser, Gly, His,
808 Ala and Orn, pursuing the same accumulation AA pathway as found in deep SPM. Comparison

809 with literature data shows that the amount of AA released, depends on the intensity of
810 hydrolysis and that about 50 % of the amide linkages detectable by NMR spectroscopy cannot
811 be hydrolyzed. Similar to SPM the proteins are not utilizable by microorganisms. Protein-like
812 dissolved material was determined to be on average 2670 years old (Loh et al., 2004), showing
813 that these refractory molecules are cycled for several times before they can be removed by as
814 yet unknown processes.

815 Based on our AA data we have calculated the total oceanic AA inventory and found that TDAA
816 are the largest oceanic AA pool with a total amount of 200 ± 70 PgAA and AA comprise 14 %
817 of the oceanic DOC and 45 % of oceanic DON.

818 The pool transported with SPM is 35 PgAA. SPM, furthermore, carries 48 PgC and 6 PgN not
819 accounted for in global carbon and nitrogen budgets. At present it is not known how the oceanic
820 DOM and SPM-particulate organic matter pool is formed and how this rather recalcitrant
821 organic matter can be removed from its ambient water mass. It is feasible that these organic matter
822 pools have fluctuated in the past due to change in oceanic physicochemical conditions
823 (Ridgwell and Arndt, 2015). It is intriguing to understand how the accumulation or reduction
824 of this carbon and nitrogen pools has interacted with climate and environmental changes in the
825 geological history but it is vital to understand the response to ongoing and future climate
826 change.

827

828 **Data Availability**

829 Excerpts of the data were used in previous publications (i) from the Kara Sea in Gaye et al.
830 (2007) Nagel et al. (2005) and Unger et al. (2009), (ii) from the northern Indian Ocean in Gaye
831 et al. (2013), Gaye-Haake et al. (2005), Möbius et al. (2011) and Suthhof et al. (2001), (iii) from
832 the Mediterranean Sea in Möbius (2013) and Möbius et al. (2010), (iv) from the Namibian
833 upwelling in Nagel et al. (2016) and (v) from the Pacific in Paul et al. (2018). The entire set of
834 amino acids data was submitted to PANGAEA. Data from the Pacific are available at:
835 <https://doi.pangaea.de/10.1594/PANGAEA.885391>, <https://doi.pangaea.de/10.1594/PANGAEA.881804>,
836 <https://doi.pangaea.de/10.1594/PANGAEA.881813>.

837

838 **Acknowledgements**

839 This work is based on samples taken during cruises of research vessels R/V SONNE, METEOR,
840 MARIA S. MERIAN, PELAGIA, ORV SAGAR KANYA and R/V AKADEMIK PETROV.
841 We are grateful to all officers of crew of these research vessels. We are indebted to Venugopalan
842 Ittekkot who initiated and inspired this work and led many of the research projects. We thank
843 Desmond Gracias, Areef Sardar and Fernando Vijayan from NIO, Goa, India for technical
844 support on board. We thank Inken Preuss, Annika Moje, Tim Jesper Suhrhoff and Seinab
845 Bohsung for help with pore water sampling during the cruises SO239 and SO242. We are
846 indebted to Frauke Langenberg and Marc Metzke for their high-quality analyses and to Dilek
847 Topcu for help with collecting and archiving the large data sets. We are very thankful to Cindy
848 Lee and two anonymous reviewers for their critical comments on the manuscript. We thank the
849 German Federal Ministry of Education and Research for funding (grant no.: 03F0707G) in the
850 framework JPI Oceans EcoMining-DEU - Ecological Aspects of Deep-Sea Mining to Jacobs
851 University Bremen. Sampling in the southern Indian Ocean was conducted within the
852 framework of the INDEX program of the Federal Institute for Geosciences and Natural
853 Resources (BGR). The DFG and BMBF financed the finalized or terminated projects in the
854 northern Indian Ocean and Atlantic (Indian-German Program, JGOFS, BIGSET), the
855 Mediterranean Sea (MEDNIT), the Namibian upwelling (GENUS) and the Kara Sea (SIRRO).

856

857 **Author Contribution**

858 BG, NL, TR and KE designed the study and led the projects in which samples were taken and
859 analyzed. NL developed and refined the AA analyses. NL, NH and SP contributed and analyzed
860 samples from the southern Indian Ocean and the Pacific. BG wrote the manuscript with
861 contributions of all co-authors.

862

863 **Competing interests**

864 The authors declare that they have no conflict of interest.

865

866 **References**

867 Abramson, L., Lee, C., Liu, Z., Wakeham, S. G., and Szlosek, J.: Exchange between suspended
868 and sinking particles in the northwest Mediterranean as inferred from the organic composition
869 of in situ pump and sediment trap samples, *Limnology and Oceanography*, 55, 725-739, 2011.

870 Alldredge, A.: The carbon, nitrogen and mass content of marine snow as a function of aggregate
871 size, *Deep-Sea Research I*, 45, 529-541, 1998.

872 Alldredge, A. L. and Silver, M. W.: Characteristics, dynamics and significance of marine snow,
873 *Progress in Oceanography*, 20, 41-82, 1988.

874 Altabet, M. A.: Isotopic Tracers of the Marine Nitrogen Cycle: Present and Past, in: *Marine*
875 *Organic Matter: Biomarkers, Isotopes and DNA. The Handbook of Environmental Chemistry*,
876 edited by: Volkman, J. K., Springer, Berlin, Heidelberg, 251-293, doi: 10.1007/698_2_008,
877 2006.

878 Altabet, M. A., Deuser, W. G., Honjo, S., and Stienen, C.: Seasonal and depth-related changes
879 in the source of sinking particles in the North Atlantic, *Nature*, 354, 136-139, 1991.

880 Aluwihare, L. I., Repeta, D. J., Pantoja, S., and Johnson, C. G.: Two chemically distinct pools
881 of organic nitrogen accumulate in the ocean, *Science*, 308, 1007-1010, 2005.

882 Aristegui, J., Gasol, J. M., Duarte, C. M., and Herndl, G. J.: Microbial oceanography of the dark
883 ocean's pelagic realm, *Limnology and Oceanography*, 54, 1501-1529,
884 10.4319/lo.2009.54.5.1501, 2009.

885 Armstrong, R. A., Lee, C., Hedges, J. I., Honjo, S., and Wakeham, S. G.: A new, mechanistic
886 model for organic carbon fluxes in the ocean based on the quantitative association of POC with
887 ballast minerals, *Deep-Sea Research Part II*, 49, 219-236, 2002.

888 Arnarson, T. S. and Keil, R. G.: Influence of organic-mineral aggregates on microbial
889 degradation of the dinoflagellate *Scrippsiella trochoidea*, *Geochimica Et Cosmochimica Acta*,
890 69, 2111-2117, 10.1016/j.gca.2004.11.004, 2005.

891 Arnarson, T. S. and Keil, R. G.: Changes in organic matter-mineral interactions for marine
892 sediments with varying oxygen exposure times, *Geochimica et Cosmochimica Acta*, 71, 3545-
893 3556, 10.1016/j.gca.2007.04.027, 2007.

894 Arrieta, J. M., Mayol, E., Hansman, R. L., Herndl, G. J., Dittmar, T., and Duarte, C. M.: Dilution
895 limits dissolved organic carbon utilization in the deep ocean, *Science*, 348, 331-333,
896 10.1126/science.1258955, 2015.

897 Benner, R. and Kaiser, J.: Abundance of amino sugars and peptidoglycan in marine particulate
898 and dissolved organic matter, *Limnology and Oceanography*, 48, 118-128, 2003.

899 Benner, R., Louchouart, P., and Amon, R. M. W.: Terrigenous dissolved organic matter in the
900 Arctic Ocean and its transport to surface and deep waters of the North Atlantic, *Global*
901 *Biogeochemical Cycles*, 19, GB2025, 10.1029/2004GB002398, 2005.

902 Boetius, A. and Lochte, K.: Regulation of microbial enzymatic degradation of organic matter
903 in deep-sea sediments *Mar. Ecol.-Prog. Ser.*, 104, 299-307, 10.3354/meps104299, 1994.

904 Boetius, A., Ferdelman, T., and Lochte, K.: Bacterial activity in sediments of the deep Arabian
905 Sea in relation to vertical flux, *Deep-Sea Res. Part II-Top. Stud. Oceanogr.*, 47, 2835-2875,
906 10.1016/s0967-0645(00)00051-5, 2000a.

907 Boetius, A., Springer, B., and Petry, C.: Microbial activity and particulate matter in the benthic
908 nepheloid layer (BNL) of the deep Arabian Sea, *Deep-Sea Res. Part II-Top. Stud. Oceanogr.*,
909 47, 2687-2706, 10.1016/s0967-0645(00)00045-x, 2000b.

910 Boyd, P., Claustre, H., Levy, M., Siegel, D., and Weber, T.: Multi-faceted particle pumps drive
911 carbon sequestration in the ocean, *Nature*, 568, 327-335, 10.1038/s41586-019-1098-2, 2019.

- 912 Boyd, S. R.: Nitrogen in future biosphere studies, *Chemical Geology*, 176, 1-30,
913 [https://doi.org/10.1016/S0009-2541\(00\)00405-8](https://doi.org/10.1016/S0009-2541(00)00405-8), 2001.
- 914 Brockmeyer, B. and Spitzzy, A.: Evaluation of a Disc Tube Methodology for Nano- and
915 Ultrafiltration of Natural Dissolved Organic Matter, *International Journal of Organic*
916 *Chemistry*, 3, 17-25, 2013.
- 917 Bronk, D. A.: Biogeochemistry of marine dissolved organic matter, in, edited by: Hansell, D.
918 A., and Carlson, C. A., San Diego, 2002.
- 919 Carlson, C. A. and Hansell, D. A.: DOM sources, sinks, reactivity, and budgets,
920 *Biogeochemistry of Marine Dissolved Organic Matter*, 2nd Edition, Academic Press Ltd-
921 Elsevier Science Ltd, London, 1-693 pp.2015.
- 922 Carr, S. A., Mills, C. T., and Mandernack, K. W.: The use of amino acid indices for assessing
923 organic matter quality and microbial abundance in deep-sea Antarctic sediments of IODP
924 Expedition 318, *Marine Chemistry*, 186, 72-82,
925 <http://dx.doi.org/10.1016/j.marchem.2016.08.002>, 2016.
- 926 Cho, B. C. and Azam, F.: Major role of bacteria in biogeochemical fluxes in the ocean's interior,
927 *Nature*, 332, 441-443, 1988.
- 928 Ciais, P., Sabine, C., Bala, G., Bopp, L., Brovkin, V., Canadell, J., Chhabra, A., DeFries, R.,
929 Galloway, J., Heimann, M., Jones, C., Le Quere, C., Myneni, R. B., Piao, S. L., Thornton, P.,
930 Ahlstrom, A., Anav, A., Andrews, O., Archer, D., Arora, V., Bonan, G., Borges, A. V.,
931 Bousquet, P., Bouwman, L., Bruhwiler, L. M., Caldeira, K., Cao, L., Chappellaz, J., Chevallier,
932 F., Cleveland, C., Cox, P., Dentener, F. J., Doney, S. C., Erisman, J. W., Euskirchen, E. S.,
933 Friedlingstein, P., Gruber, N., Gurney, K., Holland, E. A., Hopwood, B., Houghton, R. A.,
934 House, J. I., Houweling, S., Hunter, S., Hurtt, G., Jacobson, A. D., Jain, A., Joos, F., Jungclaus,
935 J., Kaplan, J. O., Kato, E., Keeling, R., Khatiwala, S., Kirschke, S., Goldewijk, K. K., Kloster,
936 S., Koven, C., Kroeze, C., Lamarque, J. F., Lassey, K., Law, R. M., Lenton, A., Lomas, M. R.,
937 Luo, Y. Q., Maki, T., Marland, G., Matthews, H. D., Mayorga, E., Melton, J. R., Metzl, N.,
938 Munhoven, G., Niwa, Y., Norby, R. J., O'Connor, F., Orr, J., Park, G. H., Patra, P., Peregón,
939 A., Peters, W., Peylin, P., Piper, S., Pongratz, J., Poulter, B., Raymond, P. A., Rayner, P.,
940 Ridgwell, A., Ringeval, B., Rodenbeck, C., Saunoy, M., Schmittner, A., Schuur, E., Sitch, S.,
941 Spahni, R., Stocker, B., Takahashi, T., Thompson, R. L., Tjiputra, J., van der Werf, G., van
942 Vuuren, D., Voulgarakis, A., Wania, R., Zaehle, S., and Zeng, N.: Carbon and Other
943 Biogeochemical Cycles, *Climate Change 2013: The Physical Science Basis*, Cambridge Univ
944 Press, Cambridge, 465-570 pp.2014.
- 945 Cohen, J.: *Statistical power analysis for the behavioral sciences* (2nd Ed.) L. Erlbaum
946 Associates, Hillsdale, N.J., 567 pp.1988.
- 947 Costello, M. J., Cheung, A., and De Hauwere, N.: Surface Area and the Seabed Area, Volume,
948 Depth, Slope, and Topographic Variation for the World's Seas, Oceans, and Countries,
949 *Environ. Sci. Technol.*, 44, 8821-8828, [10.1021/es1012752](https://doi.org/10.1021/es1012752), 2010.
- 950 Cowie, G. L. and Hedges, J. I.: Sources and reactivities of amino acids in a coastal marine
951 environment, *Limnology and Oceanography*, 37, 703-724, 1992.
- 952 Cowie, G. L. and Hedges, J. I.: Biogeochemical indicators of diagenetic alteration in natural
953 organic-matter mixtures, *Nature*, 369, 304-307, 1994.
- 954 Cowie, G. L., Hedges, J. I., Prahl, F. G., and De Lange, G. J.: Elemental and major biochemical
955 changes across an oxidation front in a relict turbidite: An oxygen effect, *Geochimica et*
956 *Cosmochimica Acta*, 59, 33-46, 1995.

- 957 Dauwe, B. and Middelburg, J. J.: Amino acids and hexosamines as indicators of organic matter
958 degradation state in North Sea sediments, *Limnology and Oceanography*, 43, 782-798, 1998.
- 959 Dauwe, B., Middelburg, J. J., Hermann, P. M. J., and Heip, C. H. R.: Linking diagenetic
960 alteration of amino acids and bulk organic matter reactivity, *Limnology and Oceanography*, 44,
961 1809-1814, 1999.
- 962 Davis, J. and Benner, R.: Seasonal trends in the abundance, composition and bioavailability of
963 particulate and dissolved organic matter in the Chukchi/Beaufort Seas and western Canada
964 Basin, *Deep Sea Research Part II: Topical Studies in Oceanography*, 52, 3396-3410,
965 <http://dx.doi.org/10.1016/j.dsr2.2005.09.006>, 2005.
- 966 Davis, J., Kaiser, K., and Benner, R.: Amino acid and amino sugar yields and compositions as
967 indicators of dissolved organic matter diagenesis, *Org. Geochem.*, 40, 343-352,
968 10.1016/j.orggeochem.2008.12.003, 2009.
- 969 Degens, E. T. and Ittekkot, V.: Dissolved organic carbon - An overview, *Mitt. Geol.-Paläont.*
970 *Inst. Univ. Hamburg*, 55, 21-38, 1983.
- 971 Degens, E. T. and Ittekkot, V.: A new look at clay-organic interactions, in: *Ein Nord-Süd Profil:*
972 *Zentraleuropa-Mittelmeerraum-Afrika*, edited by: Degens, E. T., Krumbein, W. E., and
973 Prashnowsky, A. A., *Mitteilungen aus dem Geologisch-Paläontologischen Institut der*
974 *Universität Hamburg*, Krause-Druck, Stade, 229-248, 1984.
- 975 Degens, E. T. and Ittekkot, V.: Particulate organic carbon - An overview, *Mitt. Geol.-Paläont.*
976 *Inst. Univ. Hamburg*, SCOPE/UNEP Sonderband, Heft 58, 7-27, 1985.
- 977 Degens, E. T. and Mopper, K.: Early diagenesis of organic matter in marine soils, *Soil Sci.*,
978 119, 65-72, 10.1097/00010694-197501000-00010, 1975.
- 979 Druffel, E. R. M. and Williams, P. M.: Identification of a deep marine source of particulate
980 organic carbon using bomb ¹⁴C *Nature*, 347, 172-174, 10.1038/347172a0, 1990.
- 981 Emeis, K.-C., Mara, P., Schlarbaum, T., Möbius, J., Dähnke, K., Struck, U., Mihalopoulos, N.,
982 and Krom, M.: External inputs and internal N cycling traced by isotope ratios of nitrate,
983 dissolved reduced nitrogen, and particulate nitrogen in the eastern Mediterranean Sea, *Journal*
984 *of Geophysical Research*, 115, 10.1029/2009JG001214, 2010.
- 985 England, M. E.: The Age of Water and Ventilation Timescales in a Global Ocean Model, *J.*
986 *Phys. Oceanogr.*, 25, 2756-2777, 10.1175/1520-0485(1995)025<2756:taowav>2.0.co;2, 1995.
- 987 Fowler, S. W. and Knauer, G. A.: Role of large particles in the transport of elements and organic
988 compounds through the oceanic water column *Progress in Oceanography*, 16, 147-194,
989 10.1016/0079-6611(86)90032-7, 1986.
- 990 Galbraith, E. D., Kienast, M., Albuquerque, A. L., Altabet, M. A., Batista, F., Bianchi, D.,
991 Calvert, S. E., Contreras, S., Crosta, X., De Pol-Holz, R., Dubois, N., Etourneau, J., Francois,
992 R., Hsu, T. C., Ivanochko, T., Jaccard, S. L., Kao, S. J., Kiefer, T., Kienast, S., Lehmann, M.
993 F., Martinez, P., McCarthy, M., Meckler, A. N., Mix, A., Mobius, J., Pedersen, T. F., Pichevin,
994 L., Quan, T. M., Robinson, R. S., Ryabenko, E., Schmittner, A., Schneider, R., Schneider-Mor,
995 A., Shigemitsu, M., Sinclair, D., Somes, C., Studer, A. S., Tesdal, J. E., Thunell, R., Yang, J.
996 Y. T., and Members, N. W. G.: The acceleration of oceanic denitrification during deglacial
997 warming, *Nat. Geosci.*, 6, 579-584, 10.1038/ngeo1832, 2013.
- 998 Gaye-Haake, B., Unger, D., Nöthig, E.-M., Okolodkov, Y., Fahl, K., and Ittekkot, V.: Particle
999 fluxes from short term sediment trap deployments in late summer in the southern Kara Sea, In:
1000 R. Stein, K. Fahl, D.K. Fütterer, E. Galimov and O. Stepanets (Eds.). *Siberian River Run-off in*

- 1001 the Kara Sea: Characterisation, Quantification, Variability, and Environmental Significance,
1002 Proceedings in Marine Science, Elsevier, Amsterdam, 309-328, 2003.
- 1003 Gaye-Haake, B., Lahajnar, N., Emeis, K.-C., Unger, D., Rixen, T., Suthhof, A., Ramaswamy,
1004 V., Schulz, H., Paropkari, A. L., Guptha, M. V. S., and Ittekkot, V.: Stable nitrogen isotopic
1005 ratios of sinking particles and sediments from the northern Indian Ocean, *Marine Chemistry*,
1006 96, 243-255, 2005.
- 1007 Gaye, B., Wiesner, M. G., and Lahajnar, N.: Nitrogen sources in the South China Sea, as
1008 discerned from stable nitrogen isotopic ratios in rivers, sinking particles, and sediments, *Marine*
1009 *Chemistry*, 114, 72-85, 2009.
- 1010 Gaye, B., Nagel, B., Daehnke, K., Rixen, T., and Emeis, K.-C.: Evidence of parallel
1011 denitrification and nitrite oxidation in the ODZ of the Arabian Sea from paired stable isotopes
1012 of nitrate and nitrite, *Global biogeochemical cycles*, 27, 1059-1071, 10.1002/2011GB004115,
1013 2013a.
- 1014 Gaye, B., Nagel, B., Daehnke, K., Rixen, T., Lahajnar, N., and Emeis, K. C.: Amino acid
1015 composition and delta N-15 of suspended matter in the Arabian Sea: implications for organic
1016 matter sources and degradation, *Biogeosciences*, 10, 7689-7702, 10.5194/bg-10-7689-2013,
1017 2013b.
- 1018 Gaye, B., Fahl, K., Kodina, L. A., Lahajnar, N., Nagel, B., Unger, D., and Gebhardt, A. C.:
1019 Particulate matter fluxes in the southern and central Kara Sea compared to sediments: Bulk
1020 fluxes, amino acids, stable carbon and nitrogen isotopes, sterols and fatty acids, *Cont. Shelf*
1021 *Res.*, 27, 2570-2594, 10.1016/j.csr.2007.07.003, 2007.
- 1022 Gaye, B., Boell, A., Segschneider, J., Burdanowitz, N., Emeis, K.-C., Ramaswamy, V.,
1023 Lahajnar, N., Lueckge, A., and Rixen, T.: Glacial-interglacial changes and Holocene variations
1024 in Arabian Sea denitrification, *Biogeosciences*, 15, 507-527, 10.5194/bg-15-507-2018, 2018.
- 1025 Gebbie, G. and Huybers, P.: The Mean Age of Ocean Waters Inferred from Radiocarbon
1026 Observations: Sensitivity to Surface Sources and Accounting for Mixing Histories, *J. Phys.*
1027 *Oceanogr.*, 42, 291-305, 10.1175/jpo-d-11-043.1, 2012.
- 1028 Gloeckler, K., Choy, C. A., Hannides, C. C. S., Close, H. G., Goetze, E., Popp, B. N., and
1029 Drazen, J. C.: Stable isotope analysis of micronekton around Hawaii reveals suspended particles
1030 are an important nutritional source in the lower mesopelagic and upper bathypelagic zones,
1031 *Limnology and Oceanography*, 63, 1168-1180, 10.1002/lno.10762, 2018.
- 1032 Goutx, M., Wakeham, S. G., Lee, C., Duflos, M., Guigue, C., Liu, Z., Moriceau, B., Sempère,
1033 R., Tedetti, M., and Xue, J.: Composition and degradation of marine particles with different
1034 settling velocities in the northwestern Mediterranean Sea, *Limnology and Oceanography*, 52,
1035 1645-1664, 2007.
- 1036 Gruber, N.: The marine nitrogen cycle: Overview and challenges, in: *Nitrogen in the Marine*
1037 *Environment*, 2nd Edition, edited by: Capone, D. G., Bronk, D. A., Mulholland, M. R., and
1038 Carpenter, E., Academic Press, San Diego, 51, 2008.
- 1039 Guo, J., Liang, S.-k., Li, X.-j., Li, W., Wang, Y., and Su, R.-g.: Composition and bioavailability
1040 of dissolved organic matter in different water masses of the East China Sea, *Estuarine, Coastal*
1041 *and Shelf Science*, 212, 189-202, 10.1016/j.ecss.2018.07.009, 2018.
- 1042 Haake, B., Rixen, T., and Ittekkot, V.: Variability of monsoonal upwelling signals in the deep
1043 western Arabian Sea, *Mitt. Geol.-Paläont. Inst. Univ. Hamburg, Scope/UNEP Sonderband Heft*
1044 76, 85-96, 1993a.

- 1045 Haake, B., Ittekkot, V., Honjo, S., and Manganini, S.: Amino acids, hexosamines and
 1046 carbohydrate fluxes to the deep Subarctic Pacific (Station P), *Deep-Sea Research I*, 40, 547-
 1047 560, 1993b.
- 1048 Haake, B., Ittekkot, V., Ramaswamy, V., Nair, R. R., and Honjo, S.: Fluxes of amino acids and
 1049 hexosamines of the deep Arabian Sea, *Marine Chemistry*, 40, 291-314, 1992.
- 1050 Haake, B., Rixen, T., Reemtsma, T., Ramaswamy, V., and Ittekkot, V.: Processes determining
 1051 seasonality and interannual variability of settling particle fluxes to the deep Arabian Sea, in:
 1052 *Particle Flux in the Ocean*, edited by: Ittekkot, V., Schäfer, P., Honjo, S., and Depetris, P. J.,
 1053 John Wiley&Sons Ltd., 251-270, 1996.
- 1054 Hannides, C. C. S., Popp, B. N., Choy, C. A., and Drazen, J. C.: Midwater zooplankton and
 1055 suspended particle dynamics in the North Pacific Subtropical Gyre: A stable isotope
 1056 perspective, *Limnology and Oceanography*, 58, 1931-1946, 10.4319/lo.2013.58.6.1931, 2013.
- 1057 Hansell, D., Carlson, C. A., Repeta, D. J., and Schlitzer, R.: Dissolved organic matter in the
 1058 ocean, *Oceanography*, 22, 202-211, 2009.
- 1059 Hansell, D. A. and Carlson, C. A.: Localized refractory dissolved organic carbon sinks in the
 1060 deep ocean, *Global Biogeochemical Cycles*, 27, 705-710, 2013.
- 1061 Harms, N. C., Lahajnar, N., Gaye, B., Rixen, T., Schwarz-Schampera, U., and Emeis, K.-C.:
 1062 Sediment trap-derived particulate matter fluxes in the oligotrophic subtropical gyre of the South
 1063 Indian Ocean, *Deep Sea Research Part II: Topical Studies in Oceanography*, 183, 104924,
 1064 <https://doi.org/10.1016/j.dsr2.2020.104924>, 2021.
- 1065 Harms, N. C., Lahajnar, N., Gaye, B., Rixen, T., Dähnke, K., Ankele, M., Schwarz-Schampera,
 1066 U., and Emeis, K. C.: Nutrient distribution and nitrogen and oxygen isotopic composition of
 1067 nitrate in water masses of the subtropical southern Indian Ocean, *Biogeosciences*, 16, 2715-
 1068 2732, 10.5194/bg-16-2715-2019, 2019.
- 1069 Hedges, J. I. and Hare, P. E.: Amino acid adsorption by clay minerals in distilled water,
 1070 *Geochimica et Cosmochimica Acta*, 51, 255-259, 1987.
- 1071 Hedges, J. I. and Keil, R. G.: Sedimentary organic matter preservation an assessment and
 1072 speculative synthesis, *Marine Chemistry*, 49, 137-139, 10.1016/0304-4203(95)00013-h, 1995.
- 1073 Hildebrandt, Tatjana M., Nunes Nesi, A., Araújo, Wagner L., and Braun, H.-P.: Amino Acid
 1074 Catabolism in Plants, *Molecular Plant*, 8, 1563-1579,
 1075 <https://doi.org/10.1016/j.molp.2015.09.005>, 2015.
- 1076 Honjo, S., Manganini, S. J., Krishfield, R. A., and Francois, R.: Particulate organic carbon
 1077 fluxes to the ocean interior and factors controlling the biological pump: A synthesis of global
 1078 sediment trap programs since 1983, *Progress in Oceanography*, 76, 217-285,
 1079 10.1016/j.pcean.2007.11.003, 2008.
- 1080 Ingalls, A. E., Liu, Z., and Lee, C.: Seasonal trends in the pigment and amino acid compositions
 1081 of sinking particles in biogenic CaCO₃ and SiO₂ dominated regions of the Pacific sector of the
 1082 Southern Ocean along 170°W, *Deep-Sea Research I*, 53, 836-859, 2006.
- 1083 Ingalls, A. E., Aller, R. C., Lee, C., and Wakeham, S. G.: Organic matter diagenesis in shallow
 1084 water carbonate sediments, *Geochimica et Cosmochimica Acta*, 68, 4263-4379, 2004.
- 1085 Ittekkot, V.: Verteilung von gelöstem organischen Kohlenstoff, gelösten Zuckern und
 1086 Aminosäuren in Fladengrund, nördliche Nordsee (FLEX 1976), *Mitt. Geol.-Paläont. Inst. Univ.*
 1087 *Hamburg*, 51, 115-187, 1981.

- 1088 Ittekkot, V. and Arain, R.: Nature of Particulate Organic-Matter in the River Indus, Pakistan,
1089 *Geochimica Et Cosmochimica Acta*, 50, 1643-1653, 10.1016/0016-7037(86)90127-4, 1986.
- 1090 Ittekkot, V., Degens, E. T., and Honjo, S.: Seasonality in the fluxes of sugars, amino acids, and
1091 amino sugars to the deep ocean: Panama Basin, *Deep-Sea Research*, 31, 1071-1083, 1984a.
- 1092 Ittekkot, V., Deuser, W. G., and Degens, E. T.: Seasonality in the fluxes of sugars, amino acids,
1093 and amino sugars to the deep ocean: Sargasso Sea, *Deep-Sea Research*, 31, 1057-1069, 1984b.
- 1094 Ittekkot, V., Safiullah, S., and Arain, R.: Nature of Organic-Matter in Rivers with Deep-Sea
1095 Connections - the Ganges-Brahmaputra and Indus, *Science of the Total Environment*, 58, 93-
1096 107, 10.1016/0048-9697(86)90080-x, 1986.
- 1097 Jennerjahn, T. and Ittekkot, V.: Organic matter in sediments in the mangrove areas and adjacent
1098 continental margins of Brazil: I. Amino acids and hexosamines, *Oceanol. Acta*, 20, 359-369,
1099 1997.
- 1100 Kaiser, K. and Benner, R.: Biochemical composition and size distribution of organic matter at
1101 the Pacific and Atlantic time-series stations, *Marine Chemistry*, 113, 63-77,
1102 <http://dx.doi.org/10.1016/j.marchem.2008.12.004>, 2009.
- 1103 Kandler, O.: Cell-wall structures in methane bacteria - evolution of procaryotes
1104 *Naturwissenschaften*, 66, 95-105, 10.1007/bf00373500, 1979.
- 1105 Karl, D., Knauer, G. A., and Martin, A. P.: Downward flux of particulate organic matter in the
1106 ocean: a particle composition paradox, *Nature*, 332, 438-441, 1988.
- 1107 Keil, R. G. and Kirchman, D. L.: Dissolved Combined Amino Acids: Chemical Form and
1108 Utilization by Marine Bacteria, *Limnology and Oceanography*, 38, 1256-1270, 1993.
- 1109 Keil, R. G. and Kirchman, D. L.: Abiotic Transformation of Labile Protein to Refractory Protein
1110 in Sea-Water, *Marine Chemistry*, 45, 187-196, 10.1016/0304-4203(94)90002-7, 1994.
- 1111 Keil, R. G. and Kirchman, D. L.: Utilization of dissolved protein and amino acids in the northern
1112 Sargasso Sea, *Aquat. Microb. Ecol.*, 18, 293-300, 1999.
- 1113 Keil, R. G., Montlucon, D. B., Prahl, F. G., and Hedges, J. I.: Sorptive Preservation of Labile
1114 Organic-Matter in Marine-Sediments, *Nature*, 370, 549-552, 10.1038/370549a0, 1994.
- 1115 Kienast, M., Lehmann, M. F., Timmermann, A., Galbraith, E., Bolliet, T., Holbourn, A.,
1116 Normandeau, C., and Laj, C.: A mid-Holocene transition in the nitrogen dynamics of the
1117 western equatorial Pacific: Evidence of a deepening thermocline?, *Geophysical Research
1118 Letters*, 35, 5, 10.1029/2008gl035464, 2008.
- 1119 Kim, T. H., Kim, G., Shen, Y., and Benner, R.: Strong linkages between surface and deep-water
1120 dissolved organic matter in the East/Japan Sea, *Biogeosciences*, 14, 2561-2570, 10.5194/bg-
1121 14-2561-2017, 2017.
- 1122 King, K. J.: Amino acid composition of the silicified matrix in fossil polycystine Radiolaria,
1123 *Micropaleontology*, 21, 215-226, 1975.
- 1124 King, K. J. and Hare, P. E.: Amino acid composition of the test as a taxonomic character for
1125 living and fossil planktonic foraminifera, *Micropaleontology*, 18, 285-293, 1972.
- 1126 Klein, I., von Rad, U., and Durner, J.: Homoserine lactones: do plants really listen to bacterial
1127 talk?, *Plant signaling & behavior*, 4, 50-51, 2009.

- 1128 Koppelman, R., Bottger-Schnack, R., Mobius, J., and Weikert, H.: Trophic relationships of
1129 zooplankton in the eastern Mediterranean based on stable isotope measurements, *J. Plankton*
1130 *Res.*, 31, 669-686, 10.1093/plankt/fbp013, 2009.
- 1131 Lahajnar, N., Wiesner, M. G., and Gaye, B.: Fluxes of amino acids and hexosamines to the deep
1132 South China Sea, *Deep-Sea Res. Part I-Oceanogr. Res. Pap.*, 54, 2120-2144,
1133 10.1016/j.dsr.2007.08.009, 2007.
- 1134 Lahajnar, N., Rixen, T., Gaye-Haake, B., and Ittekkot, V.: Dissolved organic carbon (DOC)
1135 fluxes of deep-sea sediments from the Arabian Sea and NE Atlantic, *Deep-Sea Research II*, 52,
1136 1947-1964, 2005a.
- 1137 Lahajnar, N., Rixen, T., Gaye-Haake, B., Schafer, P., and Ittekkot, V.: Dissolved organic carbon
1138 (DOC) fluxes of deep-sea sediments from the Arabian Sea and NE Atlantic, *Deep-Sea Res. Part*
1139 *II-Top. Stud. Oceanogr.*, 52, 1947-1964, 10.1016/j.dsr2.2005.05.006, 2005b.
- 1140 Lam, P. J. and Marchal, O.: Insights into Particle Cycling from Thorium and Particle Data, in:
1141 *Annual Review of Marine Science*, Vol 7, edited by: Carlson, C. A., and Giovannoni, S. J.,
1142 *Annual Review of Marine Science*, Annual Reviews, Palo Alto, 159-184, 10.1146/annurev-
1143 marine-010814-015623, 2015.
- 1144 Lee, C.: Amino acids and amine biogeochemistry in marine particulate material and sediments,
1145 In: *Nitrogen cycling in coastal marine environments* (eds. T.H. Blackburn, and J. Sørensen),
1146 *SCOPE*, Wiley and Sons, 125-141, 1988.
- 1147 Lee, C. and Cronin, C.: The vertical flux of particulate organic nitrogen in the sea:
1148 decomposition of amino acids in the Peru upwelling area and the equatorial Atlantic, *J. Mar.*
1149 *Res.*, 40, 227-251, 1982.
- 1150 Lee, C. and Cronin, C.: Particulate amino acids in the sea: Effects of primary productivity and
1151 biological decomposition, *J. Mar. Res.*, 42, 1075-1097, 1984.
- 1152 Lee, C., Wakeham, S., and Arnosti, C.: Particulate organic matter in the sea: The composition
1153 conundrum, *Ambio*, 33, 565-575, 10.1639/0044-7447(2004)033[0565:pomits]2.0.co;2, 2004.
- 1154 Legendre, L., Rivkin, R. B., Weinbauer, M. G., Guidi, L., and Uitz, J.: The microbial carbon
1155 pump concept: Potential biogeochemical significance in the globally changing ocean, *Progress*
1156 *in Oceanography*, 134, 432-450, 10.1016/j.pocean.2015.01.008, 2015.
- 1157 Loh, A. N., Bauer, J. E., and Druffel, E. R. M.: Variable ageing and storage of dissolved organic
1158 components in the open ocean, *Nature*, 430, 877-881, 10.1038/nature02780, 2004.
- 1159 Loick-Wilde, N., Weber, S. C., Eglite, E., Liskow, I., Schulz-Bull, D., Wasmund, N., Wodarg,
1160 D., and Montoya, J. P.: De novo amino acid synthesis and turnover during N₂ fixation,
1161 *Limnology and Oceanography*, 63, 1076-1092, 10.1002/lno.10755, 2018.
- 1162 Lonborg, C., Alvarez-Salgado, X. A., Letscher, R. T., and Hansell, D. A.: Large Stimulation of
1163 Recalcitrant Dissolved Organic Carbon Degradation by Increasing Ocean Temperatures,
1164 *Frontiers in Marine Science*, 4, 11, 10.3389/fmars.2017.00436, 2018.
- 1165 Martin, J. H., Knauer, G. A., Karl, D. M., and Broenkow, W. W.: Vertex: carbon cycling in the
1166 northeast Pacific, *Deep-Sea Research*, 34, 267-285, 1987.
- 1167 Mayzaud, P. and Martin, J. L. M.: Some aspects of biochemical and mineral composition of
1168 marine plankton *J. Exp. Mar. Biol. Ecol.*, 17, 297-310, 10.1016/0022-0981(75)90005-2, 1975.
- 1169 McCarthy, M., Pratum, T., Hedges, J., and Benner, R.: Chemical composition of dissolved
1170 organic nitrogen in the ocean, *Nature*, 390, 150-154, 10.1038/36535, 1997.

- 1171 McCarthy, M. D., Benner, R., Lee, C., and Fogel, M. L.: Amino acid nitrogen isotopic
1172 fractionation patterns as indicators of heterotrophy in plankton, particulate, and dissolved
1173 organic matter, *Geochimica et Cosmochimica Acta*, 71, 4727-4744,
1174 <https://doi.org/10.1016/j.gca.2007.06.061>, 2007.
- 1175 McCarthy, M. D., Benner, R., Lee, C., Hedges, J. I., and Fogel, M. L.: Amino acid carbon
1176 isotopic fractionation patterns in oceanic dissolved organic matter: an unaltered
1177 photoautotrophic source for dissolved organic nitrogen in the ocean?, *Marine Chemistry*, 92,
1178 123-134, 2004.
- 1179 McCave, I. N.: Size spectra and aggregation of suspended particles in the ocean, *Deep-Sea*
1180 *Research*, 31, 329-352, 1984.
- 1181 Menzel, P., Anupama, K., Basavaiah, N., Das, B. K., Gaye, B., Herrmann, N., and Prasad, S.:
1182 The use of amino acid analyses in (palaeo-) limnological investigations: A comparative study
1183 of four Indian lakes in different climate regimes, *Geochimica et Cosmochimica Acta*, 160, 25-
1184 37, <http://dx.doi.org/10.1016/j.gca.2015.03.028>, 2015.
- 1185 Menzel, P., Gaye, B., Wiesner, M. G., Prasad, S., Stebich, M., Das, B. K., Anoop, A., Riedel,
1186 N., and Basavaiah, N.: Influence of bottom water anoxia on nitrogen isotopic ratios and amino
1187 acid contributions of recent sediments from small eutrophic Lonar Lake, Central India,
1188 *Limnology and Oceanography*, 58, 1061-1074, 2013.
- 1189 Möbius, J.: Isotope fractionation during nitrogen remineralization (ammonification):
1190 Implications for nitrogen isotope biogeochemistry, *Geochimica et Cosmochimica Acta*, 105,
1191 422-432, 2013.
- 1192 Möbius, J., Lahajnar, N., and Emeis, K.-C.: Diagenetic control on nitrogen isotope ratios in
1193 Holocene sapropels and recent sediments from the Eastern Mediterranean Sea, *Biogeosciences*
1194 7, 3901-3914, 10.5194/bg-7-3901-2010, 2010.
- 1195 Möbius, J., Gaye, B., Lahajnar, N., Bahlmann, E., and Emeis, K.-C.: Influence of diagenesis on
1196 sedimentary $\delta^{15}\text{N}$ in the Arabian Sea over the last 130 kyr, *Mar. Geol.*, 284, 127-138; doi:
1197 110.1016/j.margeo.2011.1003.1013, 2011.
- 1198 Muller, P. J., Suess, E., and Ungerer, C. A.: Amino acids and amino sugars of surface particulate
1199 and sediment trap material from waters of the Scotia Sea Deep-Sea Research Part a-
1200 *Oceanographic Research Papers*, 33, 819-838, 10.1016/0198-0149(86)90090-7, 1986.
- 1201 Müller, P. J.: C/N ratios in Pacific deep sea sediments: Effect of inorganic ammonium and
1202 organic nitrogen compounds sorbed by clays, *Geochimica et Cosmochimica Acta*, 11, 765-776,
1203 1977.
- 1204 Nagel, B., Gaye, B., Kodina, L. A., and Lahajnar, N.: Stable carbon and nitrogen isotopes as
1205 indicators for organic matter sources in the Kara Sea, *Mar. Geol.*, 266, 42-51,
1206 10.1016/j.margeo.2009.07.010, 2009.
- 1207 Nagel, B., Gaye, B., Lahajnar, N., Struck, U., and Emeis, K.-C.: Effects of current regimes and
1208 oxygenation on particulate matter preservation on the Namibian shelf: Insights from amino acid
1209 biogeochemistry, *Marine Chemistry*, 186, 121-132,
1210 <http://dx.doi.org/10.1016/j.marchem.2016.09.001>, 2016.
- 1211 Nagel, B., Emeis, K.-C., Flohr, A., Rixen, T., Schlarbaum, T., Mohrholz, V., and van der Plas,
1212 A.: N-cycling and balancing of the N-deficit generated in the oxygen minimum zone over the
1213 Namibian shelf—An isotope-based approach, *Journal of Geophysical Research:*
1214 *Biogeosciences*, 118, 361-371, 2013.

- 1215 Niggemann, J. and Schubert, C. J.: Sources and fate of amino sugars in coastal Peruvian
1216 sediments, *Geochimica et Cosmochimica Acta*, 70, 2229-2237, 2006.
- 1217 Niggemann, J., Lomstein, B. A., and Schubert, C. J.: Diagenesis of amino compounds in water
1218 column and sediment of Lake Baikal, *Org. Geochem.*, 115, 67-77,
1219 [10.1016/j.orggeochem.2017.10.008](https://doi.org/10.1016/j.orggeochem.2017.10.008), 2018.
- 1220 Orellana, M. V. and Leck, C.: *Marine Microgels, Biogeochemistry of Marine Dissolved
1221 Organic Matter*, 2nd Edition, Academic Press Ltd-Elsevier Science Ltd, London, 451-480 pp.,
1222 [10.1016/b978-0-12-405940-5.00009-1](https://doi.org/10.1016/b978-0-12-405940-5.00009-1), 2015.
- 1223 Pantoja, S., Sepúlveda, J., and González, H. E.: Decomposition of sinking proteinaceous
1224 material during fall in the oxygen minimum zone off northern Chile, *Deep-Sea Research I*, 51,
1225 55-70, 2004.
- 1226 Parsek, M. R., Val, D. L., Hanzelka, B. L., Cronan, J. E., Jr., and Greenberg, E. P.: Acyl
1227 homoserine-lactone quorum-sensing signal generation, *Proc. Natl. Acad. Sci. U. S. A.*, 96,
1228 4360-4365, 1999.
- 1229 Paul, S. A. L., Gaye, B., Haeckel, M., Kasten, S., and Koschinsky, A.: Biogeochemical
1230 Regeneration of a Nodule Mining Disturbance Site: Trace Metals, DOC and Amino Acids in
1231 Deep-Sea Sediments and Pore Waters, *Frontiers in Marine Science*, 5,
1232 [10.3389/fmars.2018.00117](https://doi.org/10.3389/fmars.2018.00117), 2018.
- 1233 Peters, B. D., Lam, P. J., and Casciotti, K. L.: Nitrogen and oxygen isotope measurements of
1234 nitrate along the US GEOTRACES Eastern Pacific Zonal Transect (GP16) yield insights into
1235 nitrate supply, remineralization, and water mass transport, *Marine Chemistry*, 201, 137-150,
1236 <https://doi.org/10.1016/j.marchem.2017.09.009>, 2018.
- 1237 Pilskałn, C. H. and Honjo, S.: The fecal pellet fraction of biogeochemical particle fluxes to the
1238 deep sea, *Global Biogeochemical Cycles*, 1, 31-48, 1987.
- 1239 Reinthaler, T., van Aken, H., Veth, C., Aristegui, J., Robinson, C., Williams, P., Lebaron, P.,
1240 and Herndl, G. J.: Prokaryotic respiration and production in the meso- and bathypelagic realm
1241 of the eastern and western North Atlantic basin, *Limnology and Oceanography*, 51, 1262-1273,
1242 [10.4319/lo.2006.51.3.1262](https://doi.org/10.4319/lo.2006.51.3.1262), 2006.
- 1243 Resplandy, L., Lévy, M., and McGillicuddy Jr., D. J.: Effects of Eddy-Driven Subduction on
1244 Ocean Biological Carbon Pump, *Global Biogeochemical Cycles*, 33, 1071-1084,
1245 <https://doi.org/10.1029/2018GB006125>, 2019.
- 1246 Riccardi, G., Derossi, E., and Milano, A.: Amino acid biosynthesis and its regulation in
1247 cyanobacteria *Plant Sci.*, 64, 135-151, [10.1016/0168-9452\(89\)90018-6](https://doi.org/10.1016/0168-9452(89)90018-6), 1989.
- 1248 Ridgwell, A. and Arndt, S.: *Why Dissolved Organics Matter: DOC in Ancient Oceans and Past
1249 Climate Change*, *Biogeochemistry of Marine Dissolved Organic Matter*, 2nd Edition, Academic
1250 Press Ltd-Elsevier Science Ltd, London, 1-20 pp., [10.1016/b978-0-12-405940-5.00001-7](https://doi.org/10.1016/b978-0-12-405940-5.00001-7),
1251 2015.
- 1252 Rixen, T., Gaye, B., and Emeis, K.-C.: The monsoon, carbon fluxes, and the organic carbon
1253 pump in the northern Indian Ocean, *Progress in Oceanography*, 175, 24-39,
1254 <https://doi.org/10.1016/j.pocean.2019.03.001>, 2019a.
- 1255 Rixen, T., Guptha, M. V. S., and Ittekkot, V.: Sedimentation, in: *Report of the Indian Ocean
1256 Synthesis Group on the Arabian Sea Process Study*, edited by: L. Watts, e. a., JGOFS
1257 International Project Office, Bergen, 65-73, 2002.

- 1258 Rixen, T., Baum, A., Gaye, B., and Nagel, B.: Seasonal and interannual variations in the
1259 nitrogen cycle in the Arabian Sea, *Biogeosciences*, 11, 5733-5747, 10.5194/bg-11-5733-2014,
1260 2014.
- 1261 Rixen, T., Gaye, B., Emeis, K. C., and Ramaswamy, V.: The ballast effect of lithogenic matter
1262 and its influences on the carbon fluxes in the Indian Ocean, *Biogeosciences*, 16, 485-503,
1263 10.5194/bg-16-485-2019, 2019b.
- 1264 Robinson, R. S., Kienast, M., Albuquerque, A. L. S., Altabet, M., Contreras, S., and al, e.: A
1265 review of nitrogen isotopic alteration in marine sediments, *Paleoceanography*, 27,
1266 10.1029/2012PA002321, 2012.
- 1267 Rontani, J.-F., Zabeti, N., and Wakeham, S. G.: Degradation of particulate organic matter in
1268 the equatorial Pacific Ocean: Biotic or abiotic?, *Limnology and Oceanography*, 56, 333-349,
1269 2011.
- 1270 Satterberg, J., Arnarson, T. S., Lessard, E. J., and Keil, R. G.: Sorption of organic matter from
1271 four phytoplankton species to montmorillonite, chlorite and kaolinite in seawater, *Marine*
1272 *Chemistry*, 81, 11-18, 10.1016/s0304-4203(02)00136-6, 2003.
- 1273 Sheridan, C. C., Lee, C., Wakeham, S. G., and Bishop, J. K. B.: Suspended particle organic
1274 composition and cycling in surface and midwaters of the equatorial Pacific Ocean, *Deep-Sea*
1275 *Research I*, 49, 1983-2008, 2002.
- 1276 Silver, M. W., Coale, S. L., Pilskaln, C. H., and Steinberg, D. R.: Giant aggregates: Importance
1277 as microbial centers and agents of material flux in the mesopelagic zone, *Limnology and*
1278 *Oceanography*, 43, 498-507, 1998.
- 1279 Smith, D. C., Simon, M., Alldredge, A. L., and Azam, F.: Intense hydrolytic enzyme activity
1280 on marine aggregates and implications for rapid particle dissolution, *Nature*, 359, 139-142,
1281 1992.
- 1282 Suess, E.: Particulate organic carbon flux in the oceans - surface productivity and oxygen
1283 utilization, *Nature*, 288, 260-263, 1980.
- 1284 Sun, Y., Zulla, M., Joachimski, M., Bond, D., Wignall, P., Zhang, Z., and Zhang, M.:
1285 Ammonium ocean following the end-Permian mass extinction, *Earth Planet. Sci. Lett.*, 518,
1286 211-222, 10.1016/j.epsl.2019.04.036, 2019.
- 1287 Suthhof, A., Ittekkot, V., and Gaye-Haake, B.: Millennial-scale oscillation of denitrification
1288 intensity in the Arabian Sea during the late Quaternary and its potential influence on
1289 atmospheric N₂O and global climate, *Global Biogeochemical Cycles*, 15, 637-650, 2001.
- 1290 Suthhof, A., Jennerjahn, T. C., Schäfer, P., and Ittekkot, V.: Nature of organic matter in surface
1291 sediments from the Pakistan continental margin and the deep Arabian Sea: amino acids, *Deep-*
1292 *Sea Research II*, 47, 329-351, 2000.
- 1293 Taylor, G. T.: Microbial degradation of sorbed and dissolved protein in seawater, *Limnology*
1294 *and Oceanography*, 40, 875-885, 1995.
- 1295 Tesdal, J. E., Galbraith, E. D., and Kienast, M.: Nitrogen isotopes in bulk marine sediment:
1296 linking seafloor observations with subseafloor records, *Biogeosciences*, 10, 101-118,
1297 10.5194/bg-10-101-2013, 2013.
- 1298 Turnewitsch, R., Lahajnar, N., Haeckel, M., and Christiansen, B.: An abyssal hill fractionates
1299 organic and inorganic matter in deep-sea surface sediments, *Geophysical Research Letters*, 42,
1300 7663-7672, 10.1002/2015gl065658, 2015.

- 1301 Turnewitsch, R., Dale, A., Lahajnar, N., Lampitt, R. S., and Sakamoto, K.: Can neap-spring
1302 tidal cycles modulate biogeochemical fluxes in the abyssal near-seafloor water column?,
1303 *Progress in Oceanography*, 154, 1-24, 10.1016/j.pocean.2017.04.006, 2017.
- 1304 Unger, D., Gaye-Haake, B., Gebhardt, A. C., and Ittekkot, V.: Biogeochemistry of suspended
1305 and sedimentary material from the Ob and Yenisei rivers and the adjacent Kara Sea: Amino
1306 acids and amino sugars, *Cont. Shelf Res.*, 25, 437-460, 2005.
- 1307 Wakeham, S. G. and Canuel, E. A.: Organic geochemistry of particulate matter in the eastern
1308 tropical North Pacific; Implications for particle dynamics, *J. Mar. Res.*, 46, 183-213, 1988.
- 1309 Wakeham, S. G. and Lee, C.: Organic geochemistry of particulate matter in the ocean - the role
1310 of particles in oceanic sedimentary cycles *Org. Geochem.*, 14, 83-96, 10.1016/0146-
1311 6380(89)90022-3, 1989.
- 1312 Wakeham, S. G. and Lee, C.: Production, Transport, and Alteration of Particulate Organic
1313 Matter in the Marine Water Column, in: *Organic Geochemistry: Principles and Applications*,
1314 edited by: Engel, M. H., and Macko, S. A., Springer US, Boston, MA, 145-169, 10.1007/978-
1315 1-4615-2890-6_6, 1993.
- 1316 Wakeham, S. G., Lee, C., Farrington, J. W., and Gagosian, R. B.: Biogeochemistry of particulate
1317 organic matter in the oceans: results from sediment trap experiments, *Deep-Sea Research*, 31,
1318 509-528, 1984.
- 1319 Walla, M. D., Lau, P. Y., Morgan, S. L., Fox, A., and Brown, A.: Capillary gas
1320 chromatography-mass spectrometry of carbohydrate components of legionellae and other
1321 bacteria, *Journal of Chromatography A*, 288, 399-413, [https://doi.org/10.1016/S0021-
1322 9673\(01\)93716-1](https://doi.org/10.1016/S0021-9673(01)93716-1), 1984.
- 1323 Waples, D. W. and Sloan, J. R.: Carbon and nitrogen diagenesis in deep-sea sediments
1324 *Geochimica et Cosmochimica Acta*, 44, 1463-1470, 10.1016/0016-7037(80)90111-8, 1980.
- 1325 Whelan, J. K. and Emeis, K.-C.: Preservation of amino acids and carbohydrates in marine
1326 sediments, in: *Organic Matter: Productivity, Accumulation, and Preservation in Recent and
1327 Ancient Sediments*, edited by: Whelan, J. K., and Farrington, J., Columbia University Press,
1328 Palisades, N. Y., 176-200, 1992.
- 1329 Wilson, J. D., Barker, S., and Ridgwell, A.: Assessment of the spatial variability in particulate
1330 organic matter and mineral sinking fluxes in the ocean interior: Implications for the ballast
1331 hypothesis, *Global Biogeochemical Cycles*, 26, 15, 10.1029/2012gb004398, 2012.
- 1332 Xue, J. H., Lee, C., Wakeham, S. G., and Armstrong, R. A.: Using principal components
1333 analysis (PCA) with cluster analysis to study the organic geochemistry of sinking particles in
1334 the ocean, *Org. Geochem.*, 42, 356-367, 10.1016/j.orggeochem.2011.01.012, 2011.
- 1335 Yamaguchi, Y. T. and McCarthy, M. D.: Sources and transformation of dissolved and
1336 particulate organic nitrogen in the North Pacific Subtropical Gyre indicated by compound-
1337 specific $\delta^{15}\text{N}$ analysis of amino acids, *Geochimica et Cosmochimica Acta*, 220, 329-347,
1338 <https://doi.org/10.1016/j.gca.2017.07.036>, 2018.
- 1339 Yang, J.-Y. T., Kao, S.-J., Dai, M., Yan, X., and Lin, H.-L.: Examining N cycling in the northern
1340 South China Sea from N isotopic signals in nitrate and particulate phases, *Journal of
1341 Geophysical Research: Biogeosciences*, 122, 2118-2136, 10.1002/2016JG003618, 2017.
- 1342 Zhang, P. Y., Yang, G. P., Chen, Y., Leng, W. S., and Ji, C. X.: Temporal and spatial variations
1343 of particulate and dissolved amino acids in the East China Sea, *Marine Chemistry*, 186, 133-
1344 144, 10.1016/j.marchem.2016.09.004, 2016.

1345

i