

1 **What can we learn from amino acids about oceanic organic matter cycling**
2 **and degradation?**

3

4 **Birgit Gaye¹, Niko Lahajnar¹, Natalie Harms¹, Sophie Anna Luise Paul^{2, 3}, Tim Rixen^{1,4}**
5 **and Kay-Christian Emeis¹**

6 *¹Institute for Geology, Universität Hamburg, 20146 Hamburg, Germany*

7 *²Department of Physics and Earth Sciences, Jacobs University Bremen, 28759 Bremen,*
8 *Germany*

9 *³GEOMAR, Helmholtz-Zentrum für Ozeanforschung, 24148 Kiel, Germany*

10 *⁴Leibniz Centre for Tropical Marine Research (ZMT), 28359 Bremen, Germany*

11

12 **Correspondence:** Birgit Gaye (birgit.gaye@uni-hamburg.de)

13

14 **Abstract**

15 Amino acids (AA) mainly bound in proteins are major constituents of living biomass and non-
16 living organic material in the oceanic particulate and dissolved organic matter pool. Uptake and
17 cycling by heterotrophic organisms lead to characteristic changes in AA composition so that
18 AA based biogeochemical indicators are often used to elucidate processes of organic matter
19 cycling and degradation. We analyzed particulate AA in a large sample set collected in various
20 oceanic regions covering sinking and suspended particles in the water column, sediment
21 samples as well as dissolved AA from water column and pore water samples. The aim of this
22 study was to test and improve the use of AA derived biogeochemical indicators as proxies for
23 organic matter sources and degradation, and to better understand particle dynamics and
24 interaction between the dissolved and particulate organic matter pools. A principal component
25 analysis (PCA) of all data delineates diverging AA compositions of sinking and suspended
26 particles with increasing water depth. A new sinking particle and sediment degradation
27 indicator (SDI) allows a fine-tuned classification of sinking particles and sediments with respect
28 to the intensity of degradation, which is associated with changes of stable isotopic ratios of
29 nitrogen ($\delta^{15}\text{N}$). This new indicator furthermore is sensitive to sedimentary redox conditions
30 and can be used to detect past anoxic early diagenesis. A second indicator emerges from the

31 AA spectra of suspended particulate matter (SPM) in the epipelagic and that of the meso- and
32 bathypelagic ocean and is a residence time indicator (RTI). The characteristic changes in AA
33 patterns from shallow to deep SPM are recapitulated in the AA spectra of the dissolved organic
34 matter (DOM) pool, so that deep SPM is more similar to DOM than to any of the other organic
35 matter pools. This implies that there is equilibration between finely dispersed SPM and DOM
36 in the deep sea, which may be driven by microbial activity combined with annealing and
37 fragmentation of gels. As these processes strongly depend on physico-chemical conditions in
38 the deep ocean, changes in quality and degradability of DOM may strongly affect the relatively
39 large pool of suspended and dissolved AA in the ocean that amounts to 15 Pg amino acid carbon
40 (AAC) and 89 ± 29 Pg AAC, respectively.

41 **1 Introduction**

42 Amino acids (AA) are ubiquitous in living organisms and comprise a major share of
43 characterized organic matter in the particulate and dissolved pool in the ocean (Lee, 1988;
44 Wakeham et al., 1984; Zhang et al., 2016; Davis et al., 2009; Lee et al., 2004). AA contribute
45 more than 80 % of total organic carbon in fresh autochthonous plankton while allochthonous
46 organic matter from continental runoff and atmospheric deposition has lower AA contents
47 (Degens and Ittekkot, 1983; Degens and Ittekkot, 1985). Most of the primary productivity
48 occurs in the sunlit surface layer of the ocean and most of the allochthonous material is also
49 transported into surface waters. Thus, organic matter concentrations including their major
50 biogenic constituents generally, have a surface maximum and decrease with depth (Peters et
51 al., 2018; Gaye et al., 2013b; Wakeham and Lee, 1993). The main mechanism behind this depth
52 dependent distribution is that most of the organic matter is recycled in surface waters while
53 only a small proportion of surface particles leaves the surface waters by gravitational settling
54 in the form of macroaggregates or fecal pellets. Particles containing organic matter, shells,
55 frustules of organisms and mineral matter sink at speeds of 200 m day⁻¹ on average and
56 constitute the export from the surface mixed layer or euphotic zone into the deep ocean, where
57 part of it can ultimately reach the sediments (Alldredge and Silver, 1988; Alldredge, 1998;
58 Pilskaln and Honjo, 1987; Fowler and Knauer, 1986; Karl et al., 1988; Rixen et al., 2019b).
59 Sinking particles are caught by moored or floating sediment traps while suspended matter
60 (SPM) is sampled by filtration or ultrafiltration of water from water samplers or by pump
61 systems (Yamaguchi and McCarthy, 2018). SPM is too small to sink and therefore – like
62 dissolved organic matter (DOM) – predominantly enters deep water by subduction of surface
63 waters (Resplandy et al., 2019; Boyd et al., 2019) and is transported passively following the
64 route of ocean water along the ocean conveyor belt (Silver et al., 1998; McCave, 1984). It has
65 thus been surmised that the long residence time of SPM in the water column should result in a
66 more degraded state compared with organic matter of sinking particles (McCave, 1984; Degens
67 and Ittekkot, 1984). Studies of pigments, AA and fatty acids, however, do not find such a
68 systematic difference between the two types of particles and even indicate that SPM can be less
69 degraded than sinking particles (Abramson et al., 2011; Rontani et al., 2011; Wakeham and
70 Canuel, 1988). In two studies of AA composition in the Benguela Upwelling System and in
71 the Arabian Sea it was shown that the degradation pathways of SPM and sinking particles differ
72 as their AA compositions diverge with depth (Gaye et al., 2013b; Nagel et al., 2009). These
73 studies suggested that there is only little interaction between suspended and sinking particle

74 pools below the euphotic zone. Due to its long residence time in the ocean, SPM appears to
75 interact with DOM (Gaye et al., 2013b) and therefore carries different AA signatures related to
76 genesis and history of organic matter cycling in its specific water mass (Nagel et al., 2016).
77 Whereas information on the composition of sediment trap samples has been compiled in
78 comprehensive studies (Honjo et al., 2008; Wilson et al., 2012; Rixen et al., 2019a, b), similar
79 compilations of the profuse literature on suspended matter are yet missing.

80 On the way to the deep sea the flux of sinking particles is reduced by disaggregation and organic
81 matter degradation. Suess (1980) empirically derived the first power function for organic
82 carbon decay based on sediment trap data. Subsequently, a large number of similar functions
83 were calculated for various oceanic areas based on trap experiments (Rixen et al., 2019b;
84 Rixen et al., 2002; Armstrong et al., 2002; Martin et al., 1987). Early work on AA had produced
85 similar decay functions combining data from Atlantic and Pacific trap experiments (Lee and
86 Cronin, 1982, 1984). As AA decay faster than bulk organic carbon (Haake et al., 1993b; Haake
87 et al., 1992; Haake et al., 1996; Lee et al., 2004; Wakeham and Lee, 1989; Whelan and Emeis,
88 1992), they are often considered as “labile” constituents of bulk organic matter. This is
89 supposedly due to their preferential uptake as a nitrogen (N) source for further synthesis of AA
90 or as a source of essential AA for heterotrophs (Ittekkot and Arain, 1986; Ittekkot et al., 1986).
91 This has been questioned, as a large proportion of the oceanic organic N pool is comprised of
92 AA that are not bioavailable (Aluwihare et al., 2005). In addition to the quantification of AA
93 decay, degradation state of organic matter (proteins) can be assessed by characteristic changes
94 in AA monomer composition which, furthermore, have the potential to elucidate sources of
95 organic matter and degradation processes (Ittekkot et al., 1984a; Ittekkot et al., 1984b; Dauwe
96 and Middelburg, 1998; Dauwe et al., 1999; Jennerjahn and Ittekkot, 1997).

97 Ratios of individual amino acids such as the Reactivity Index (RI) (Jennerjahn and Ittekkot,
98 1997) or the Degradation Index (DI) normalizing AA data to the results of a principal
99 component analyses (PCA) (Dauwe et al., 1999; Dauwe and Middelburg, 1998) have often been
100 used to scale organic matter degradation (Niggemann et al., 2018; Unger et al., 2005; Ingalls et
101 al., 2006; Ingalls et al., 2004; Pantoja et al., 2004; Möbius et al., 2010). These biogeochemical
102 indicators of organic matter quality were essentially developed for marine sinking particles and
103 sediments. Although based on marine sediments only (Dauwe et al., 1999) the DI was applied
104 for example to SPM samples from the brackish environment (Unger et al. 2005) or even to trace
105 dissolved AA degradation (Davis and Benner, 2005; Guo et al., 2018). Other work used
106 individual and adapted indices to differentiate the states of degradation in SPM or DOM

107 samples and samples from lakes, groundwater and rivers (Abramson et al., 2011; Gaye et al.,
108 2007; Goutx et al., 2007; Kaiser and Benner, 2009; Menzel et al., 2013; Peter et al. 2012;
109 Sheridan et al., 2002).

110 Understanding and quantifying AA degradation is required to estimate the diagenetic imprint
111 on $\delta^{15}\text{N}$ ratios of particulate matter. This is important as $\delta^{15}\text{N}$ ratios track major shifts between
112 N pools and are commonly used to reconstruct the N cycle from sedimentary archives
113 (Galbraith et al., 2013). Amino acid nitrogen (AAN) comprises 80-100 % of N in fresh organic
114 matter and is the precursor of most of the N buried in sediments and ultimately stored in the
115 form of ammonium, adsorbed to clay minerals (Boyd, 2001; Waples and Sloan, 1980; Müller,
116 1977). Considerable AA degradation already occurs in the water column and progresses during
117 organic matter burial in the sediments so that the impact of diagenetic processes on $\delta^{15}\text{N}$ has to
118 be accounted for (Möbius et al., 2010; Möbius et al., 2011; Niggemann et al., 2018; Carr et al.,
119 2016). Ammonification leads to a diagenetic increase of $\delta^{15}\text{N}$ values by up to 6.5 ‰ in deep
120 sea sediments while there is little effect during organic matter burial in shelf and slope
121 sediments due to the higher sedimentation rates and sub- to anoxic diagenetic conditions
122 (Tesdal et al., 2013; Robinson et al., 2012; Möbius, 2013; Gaye-Haake et al., 2005). Such $\delta^{15}\text{N}$
123 increases were shown to correlate with AA derived degradation indicators so that the primary
124 $\delta^{15}\text{N}$ signal from the water column can be reconstructed (Gaye-Haake et al., 2005; Gaye et al.,
125 2009; Möbius et al., 2011).

126 DOM comprising the largest oceanic organic matter pool is defined by the pore size of the
127 filters it passes through which is 0.2-0.7 μm (Carlson and Hansell, 2015) and thus includes
128 some picoplankton cells and all viruses (Aristegui et al., 2009). DOM in surface water is partly
129 labile and can originate from the exudates and lysis of organisms, passive diffusion, or
130 “overflow” out of phytoplankton and bacteria; grazers can excrete or egest DOM, it can
131 furthermore be leached from their fecal pellets or released by sloppy zooplankton feeding and
132 is thus primarily released and also taken up in the surface ocean (Carlson and Hansell, 2015).
133 Moreover, terrestrially derived DOM is transported into surface waters by rivers and via the
134 atmosphere (Benner et al., 2005). Deep DOM has a different source than simply transport of
135 surface DOM by intermediate and deep water formation and mixing, as deep DOM is refractory
136 in nature and has been heterotrophically altered by cycling and degradation processes
137 (Yamaguchi and McCarthy, 2018) discernible e.g. from their composition of dissolved AA
138 (Kaiser and Benner, 2009; McCarthy et al., 2004). The possible source of deep DOM may be

139 the release from sinking or suspended particles associated with microbial degradation on
140 particles and in the ambience of particles by processes such as solubilizing organic matter by
141 ectohydrolase (Cho and Azam, 1988; Ciais et al., 2014; Aristegui et al., 2009). DOM can also
142 be released from sediment pore water into overlying waters (Lahajnar et al., 2005). Ratios of
143 $\delta^{15}\text{N}$ in ultrafiltered DOM (UDOM) showed no systematic change with depth and suggested a
144 common microbial source or viral lysis (McCarthy et al., 2007).

145 In the following synoptic compilation of AA data, we will examine the differences in AA
146 spectra of a large data set that combines dissolved and particulate AA from plankton, suspended
147 and sinking material, and sediments from different oceanic regions, as well as from riverine to
148 brackish-marine conditions. Focusing on processes in the water column the data serve to (i) test
149 existing AA based biogeochemical indicators of organic matter sources and degradation, (ii)
150 better understand transformation and degradation processes of organic matter in aquatic
151 environments reflected by AA composition in sinking and suspended particles and total
152 dissolved AA (TDAA), (iii) investigate the impact of such processes on the $\delta^{15}\text{N}$ values and
153 (iv) identify open questions which may be pursued with the help of AA analyses in the future.


154

155 **2. Materials and Methods**

156 **2.1 Sampling**

157 A total of 1425 samples were taken for AA analyses in different oceanic areas and water depths
158 between 1993 and 2017 and include 218 sediment trap samples, 489 sediment samples, 608
159 SPM samples and 110 water and pore water samples (Fig. 1a-d). Five additional plankton
160 samples were taken in the Arabian Sea and from the Namibian upwelling area by plankton tows
161 between 0-100 m and between 100-700 m water depths. In the Kara Sea - a shallow shelf sea
162 strongly impacted by water and suspended matter discharge from the rivers Ob and Yenisei -
163 sediment traps, surface sediments, and suspended matter were sampled (Gaye et al., 2007;
164 Gaye-Haake et al., 2003; Nagel et al., 2009; Unger et al., 2005). In the deep Mediterranean Sea
165 sediment traps and surface sediments were sampled (Möbius et al., 2010). SPM from the
166 Mediterranean Sea was only analyzed for $\delta^{15}\text{N}$ values (Emeis et al., 2010). Sediment trap as
167 well as surface sediment, SPM and water samples were taken along cross shelf transects off
168 Namibia (Nagel et al., 2013; Nagel et al., 2016). Sediment trap samples and short sediment
169 cores were taken at two stations in the northeastern Atlantic (Lahajnar et al., 2005; Turnewitsch

170 et al., 2017; Turnewitsch et al., 2015). In the Arabian Sea sediment trap, SPM, and surface
 171 sediment samples were taken in the deep ocean and on the continental slope including a core
 172 within the oxygen minimum zone at water depths of 775m (Gaye et al., 2013b; Gaye et al.,
 173 2013a; Rixen et al., 2014; Gaye-Haake et al., 2005; Suthhof et al., 2001; Suthhof et al., 2000).
 174 In the Indian Ocean Subtropical Gyre sediment trap, SPM and water samples were taken
 175 (Harms et al., 2019; Harms et al., 2021) and samples from the equatorial North Pacific and
 176 eastern South Pacific comprise bottom water, pore water and sediment core samples (Paul et
 177 al., 2018).



178
 179 Figure 1: Stations of sediment trap deployments (a), sediment (b), SPM (c) water and pore water
 180 sampling (d).

181
 182 Sea water was filtered through glass fiber filters (Whatman GF/F) with a nominal pore size of
 183 0.7 μm and filters were dried at 40°C in order to obtain SPM samples. At some stations water
 184 samples were taken by deep freezing an aliquot of the filtrate for TDAA analyses. In addition,
 185 18 water samples taken off Namibia were separated into two size classes by ultrafiltration
 186 (Brockmeyer and Spitzzy, 2013). The size classes 50 kDa-0.7 μm and 1 kDa-0.7 μm were used

187 for TDAA analyses. Sediment trap samples were wet sieved on board and comprise the <1mm
188 fraction, filtered with polycarbonate nuclepore filters of 0.45 µm pore size and dried at 40°C.
189 Sediment samples from multicores, box grabs, box cores, or gravity cores were taken by spatula
190 or syringes from cold stored cores and were freeze dried before analyses. Surface samples
191 represent either the upper 0.5 cm or 1 cm of a sediment core. Pore-water samples were taken
192 by rhizons with a mean pore size of 0.15 µm and stored frozen before analyses (see methods in
193 Paul et al., 2018).

194

195 **2.2 Analytical methods**

196 Total carbon and N were measured with a Carlo Erba Nitrogen Analyser 1500 (Milan, Italy) or
197 a EURO EA3000 elemental analyzer. Particulate organic carbon (POC) was measured after
198 treatment of weighed samples with 1N HCl to remove carbonate. The precision of this method
199 is 0.05% for carbon and 0.005% for N. Carbonate carbon was calculated by subtracting organic
200 carbon from total carbon. Ratios of ¹⁵N/¹⁴N of particulate N were determined using a Thermo
201 Finnigan MAT 252 isotope ratio mass spectrometer connected with a ConFlo-III interface after
202 high-temperature flash combustion in a Thermo Finnigan Flash EA 1112 at 1050°C. Part of the
203 samples were measured with an Elementar IsoPrime 100 isotope ratio mass spectrometer after
204 high temperature combustion in an Elementar CHNOS Vario isotope elemental analyzer at 950
205 °C. Pure tank N₂ calibrated against the reference standards IAEA-N1 (ammonium sulfate,
206 δ¹⁵N= + 0.4 ‰ versus air N₂) and IAEAN2 (ammonium sulfate, δ¹⁵N= + 20.3 ‰) of the
207 International Atomic Energy Agency was used as a working standard. Duplicate measurements
208 of samples differ by less than 0.15 ‰. The laboratory's long-term standard deviation for IAEA-
209 N1 standard is 0.09 ‰. N-isotope ratios are reported in ‰ using the delta notation and the
210 ¹⁵N/¹⁴N of air N₂ as the reference standard:

$$211 \delta^{15}\text{N}_{\text{sample}} = ((^{15}\text{N}/^{14}\text{N})_{\text{sample}} / (^{15}\text{N}/^{14}\text{N})_{\text{reference standard}} - 1) * 1000 \quad (1)$$

212 Dissolved organic carbon (DOC) concentrations [mg/L] were determined of water and pore
213 water samples via a high temperature combustion method (POC-V_{CSH} Analyzer, Shimadzu).
214 Inorganic carbon was removed by 2 M HCl prior to injection into the combustion tube where
215 organic carbon is oxidized to CO₂ at 680 °C with a platinum catalyst. A 5-point calibration from
216 0.5 to 5 mg DOC/L was used. The error of measurement is less than 2 % (Brockmeyer and
217 Spitzzy, 2013).

218 TDAA, particulate AA and hexosamines (HA) were analyzed with a Biochrom 30 Amino Acid
219 Analyzer. Acid hydrolysis with 6N HCl for 22 h at 110°C under a pure argon atmosphere was
220 carried out on ca. 3 ml of filtrate of water and pore water samples, on 1-2 mg of suspended
221 matter collected on Whatman GF/F filters, on 1-2 mg of sediment trap samples, or on 1-50 mg
222 of freeze dried surface sediments. A particle free aliquot was evaporated three times to dryness
223 in order to remove the unreacted HCl; the residue was taken up in an acidic buffer (pH 2.2).
224 After injection and subsequent separation with a cation exchange resin, the individual AA
225 monomers were post-column derivatized with o-phthaldialdehyde in the presence of 2-
226 mercaptoethanol and detected with a Merck Hitachi L-2480 fluorescence detector. Duplicate
227 analysis of a standard solution according to this method results in a relative error of 0.1 to 1.3%
228 for the concentrations of individual AA monomers and 0.2 to 3.0% for individual AA
229 monomers of water or particulate matter samples. Due to acid hydrolysis, aspartic acid (ASP)
230 and asparagine (Asn) are both measured as Asp and glutamic acid (Glu) and glutamine (Gln)
231 are both measured as Glu. The other AA measured are threonine (Thr), serine (Ser), glycine
232 (Gly), alanine (Ala), valine (Val), methionine (Met), isoleucine (Ile), leucine (Leu), tyrosine
233 (Tyr), phenylalanine (Phe), β -Alanine (β -Ala), γ -aminobutyric acid (γ -Aba), histidine (His),
234 ornithine (Orn), lysine (Lys) and arginine (Arg). The HA measured together with AA are
235 glucosamine (Gluam) and galactosamine (Galam) and their total contents were corrected with
236 a factor of 1.4 for loss during hydrolysis (Muller et al., 1986).

237

238 **2.3 Amino acid derived biogeochemical indicators of organic matter origin and** 239 **degradation**

240 Amino acid concentrations and the contribution of AA carbon (AAC) and AA nitrogen (AAN)
241 as percentages of total organic carbon (AAC/C %) or total N (AAN/N %) are used to determine
242 the degradation state of organic matter in the marine realm as both decrease with increasing
243 organic matter degradation (Wakeham and Lee, 1993; Cowie and Hedges, 1994). AAN/N %
244 >50 % are characteristic of fresh organic matter in the freshwater and marine realm (Menzel et
245 al., 2015; Haake et al., 1992; Haake et al., 1993b). AA contribute >60 % to total organic carbon
246 (AAC/C %) in fresh plankton and suspended matter in surface waters whereas AAC/C % drop
247 to values <20 % in sinking particles and suspended matter from subsurface water (Wakeham
248 and Lee, 1993). AAC/C % values are often below 10 % in freshwater environments and indicate

249 the enhanced input of land plants enriched in carbohydrates and lignin rather than enhanced
250 organic matter degradation (Menzel et al., 2015).

251 Asn, Gln and Glu are the primary products of N assimilation and all other AA are synthesized
252 from them (Loick-Wilde et al., 2018; Riccardi et al., 1989; Hildebrandt et al., 2015). Asp and
253 Glu are enriched in bacteria, vascular plant tissue, phytoplankton, zooplankton and fungi and
254 their high relative contents of Asp and Glu therefore, indicate fresh organic matter (Cowie and
255 Hedges, 1992). The ratios of Asp/ β -Ala and Glu/ γ -Aba are high in fresh organic matter and
256 drop with organic matter degradation as β -Ala and γ -Aba are degradation products of Asp and
257 Glu, respectively (Lee and Cronin, 1984). β -Ala and γ -Aba also become relatively enriched
258 during organic matter degradation as these non-protein AA are not taken up by heterotrophic
259 organisms (Ittekkot et al., 1984b).

260 The relative accumulation of the non-protein AA is also expressed by the RI which is the ratio
261 of the very labile aromatic AA Tyr and Phe and the non-protein AA β -Ala and γ -Aba. The RI
262 is, generally, between 0 (very degraded) and 15 (very fresh) (Jennerjahn and Ittekkot, 1997). It
263 is applicable not only in studies of sinking and suspended matter in marine and brackish
264 environments (Unger et al., 2005; Gaye et al., 2007) but also as a proxy for degradation state in
265 the sediment column (Möbius et al., 2011). The enrichment of Asp and Glu in sediments is
266 related to their enrichment in carbonate shells (Ittekkot et al., 1984a) and to adsorption of
267 primarily acidic AA onto carbonate minerals (King and Hare, 1972), whereas basic AA
268 primarily adsorb onto silicate minerals (Hedges and Hare, 1987; Keil et al., 1994; King, 1975).

269 The DI, the integral of 14 protein AA, assesses the diagenetic alteration of a sample by
270 comparing it to a set of 28 sediment samples of different degradation states and environments.
271 Molar percentages of individual AA are standardised by the mean and standard deviations of
272 the 28-sample data set. The DI then integrates the result of these standardized values weighed
273 by the factor coefficients for the first axis of the PCA of Dauwe et al. (1999) according to the
274 formula:

$$275 \quad DI = \sum_i \left[\frac{\text{var}_i - \text{AVGvar}_i}{\text{STDvar}_i} \right] \cdot \text{fac.coef}_i \quad (2)$$

276 where var_i is the original mole percentage of each AA_i , AVGvar_i and STDvar_i are the mean and
277 standard deviations, respectively, and fac.coef_i is the factor coefficient of the first axis of the
278 PCA of Dauwe et al. (1999). The DI thus represents the cumulative deviation of AA with

279 respect to an assumed average molar composition. The DI ranges approximately from -2 to +3
280 where negative values indicate more and positive values less degradation than the average.

281 A specifically designed index for dissolved substances (DOM-DI) calculated in the same way
282 as the DI was based on a PCA of a set of marine DOM samples and resulted in DOM-DI values
283 from 5 in surface waters to -3 in deep waters (Kaiser and Benner, 2009). The values to calculate
284 the marine DOM-DI (averages, standard deviations and factor coefficients of F1) can be found
285 in Peters et al. (2012).

286 An indicator of oxic vs. anoxic organic matter degradation in the water column and in sediments
287 was proposed by Menzel et al. (2015) for lake samples. Based on work by Cowie et al. (1995)
288 on marine sediments the ox/anox indicator is the quotient of AA preserved under oxic
289 diagenetic conditions to those preserved in anoxic water or sediments and is thus higher in oxic
290 than in anoxic sediments:

$$ox/anox = \frac{Asp + Glu + \beta-Ala + \gamma-Aba + Lys}{Ser + Met + Ile + Leu + Tyr + Phe} \quad (3)$$

292 ox/anox ratios <1.0 indicate anoxic and ratios >1.5 oxic diagenesis (Menzel et al., 2015).

293 The stability of AA vs. HA has been discussed since the early research on AA and HA in natural
294 material. Fresh plankton was observed to have AA/HA ratios of 13-25 (Degens and Mopper,
295 1975) which is a mixed signal of phytoplankton with an AA/HA ratio of >80 and zooplankton
296 with a ratio of ~9 due to chitinous skeletons of many zooplankters (Mayzaud and Martin,
297 1975). Low AA/HA are also observed in cell walls of fungi and bacteria. As the building blocks
298 of chitin, HA were assumed to be more resistant to degradation than bulk AA (Muller et al.,
299 1986). This is, however, challenged by studies of enzyme activities which were observed to
300 respond to substrate availability so that the activity of chitobiase and chitinase is as high as that
301 of glucosidase (Boetius et al., 2000a; Boetius and Lochte, 1994; Boetius et al., 2000b; Smith et
302 al., 1992) suggesting intense degradation also of chitin. Gluam is the main constituent of chitin
303 while Galam is relatively enriched in bacterial cell walls (Walla et al., 1984; Kandler, 1979).
304 The Gluam/Galam ratio has, therefore, been used to distinguish bacterial material from
305 zooplankton rich material (Haake et al., 1993b; Benner and Kaiser, 2003; Niggemann and
306 Schubert, 2006). Gluam/Galam ratios > 4 were found in sinking particles (Haake et al., 1993b;
307 Haake et al., 1992; Lahajnar et al., 2007), ratios of < 3 usually indicate relatively high

308 contribution of microbial OM and values between 1 and 2 are characteristic of sediments and
309 indicate an enrichment of microbial biomass (Benner and Kaiser, 2003).

310 **2.4 Statistical analyses**

311 To investigate the differences of AA composition and to recognize the interaction and pathways
312 of degradation between the different pools we carried out a PCA of AA monomer contributions
313 in Mol %. Met was excluded as it is below detection limit in many samples. The PCA was
314 carried out using the program SPSS Statistics 22. PCAs have often been used to analyze large
315 databases (Xue et al., 2011) in order to trace organic matter degradation, group and categorize
316 samples and develop indices such as the DI of Dauwe et al. (1999) using summary statistics
317 (see equation 2). A PCA is an orthogonal transformation of a set of variables into a new set of
318 uncorrelated variables called principal components. New axes are chosen in order to explain as
319 much as possible of the variance within the data set on a few main axes of highest correlation.
320 The first component explains most of the variance within the data set, consecutively followed
321 by the remaining components in the order of their decreasing capacity to explain the variance
322 within the data set. The selection of the most relevant components can be done by selecting
323 those with eigenvalues (the variances of the principal components) >1. Alternatively, the kink
324 method can be applied selecting those components from a plot of eigenvalues (scree plot),
325 which describe a steep slope of declining variance followed by a “kink” after which the
326 principal components add only small amounts to the variance. The factor loadings of the
327 variables (in this case the individual amino acids) are their projections on the new axis. The
328 factor score of each data set from a sampling location is obtained by multiplying the
329 standardized data with the factor loadings (also called factor coefficients). A high (low) factor
330 score shows that a sample has high (low) concentrations of the variables with high factor
331 loadings. A plot of factor loadings of the variables compared with a plot of the factor scores of
332 samples helps to visualize the relation of the samples to the variables and thus to identify the
333 processes behind the results of the PCA.


334

335 **2 Results**

336 **3.1 Organic carbon, nitrogen and amino acid content**

337 The POC (N) content is 35.9 % (5.9%) in plankton and 1.65-46.4 % (0.21-10.14 %) in sediment
338 trap samples. In sediments, POC (N) contents drop to 0.10-13.5 % (0.02-1.72 %). SPM has

339 POC (N) contents of 0.94-45.4 % (0.09-12.08 %). DOC concentration in water samples is
 340 between 0.5-1.1 mg L⁻¹ and DOC in pore water samples is between 3.9-29 mg L⁻¹.




341
 342 Figure 2: Box and Whisker plots of AA concentrations in nmol g⁻¹ or nmol L⁻¹ (a) and of AAC/C
 343 % (b) in SPM from water depths <200m (SPM<) and >200m (SPM>), in sediment traps at
 344 water depth <200m (Trap<) and >200 m (Trap>), in sediments (Sed), in water samples (Water)
 345 and in pore water samples (Pore W). Boxes comprise the upper and lower quartile and lines
 346 indicate median; whiskers delineate the 10 and 90 percentile; outliers are marked by dots; some
 347 outliers above the 90 percentile are cut off in the figure for better perceptibility of trends.

348
 349 AA concentrations are grouped into SPM and trap samples taken at water depths <200m
 350 (shallow) and >200m (deep) (Fig. 2, Table 1). AA concentrations are highest in SPM samples
 351 and shallow sediment traps (<200m water depth) with values between 40-4307 μmol g⁻¹ (Fig.
 352 2) and averages of 662-908 μmol g⁻¹ (Table 1). AA concentrations are lower in traps from water
 353 depth >200m with an average of 164 μmol g⁻¹. Sediments have lowest AA concentrations of all
 354 particulate matter samples with an average of 50 μmol g⁻¹ (Table 1). TDAA concentrations are
 355 between 0.6-44 μmol L⁻¹ and AA concentrations are lower in water than in pore water samples
 356 with averages of 3.2 and 8.8 μmol L⁻¹, respectively. TDAA concentrations of water samples
 357 decrease from the epipelagic (2.5±2.9 μmol L⁻¹) to the meso- and bathypelagic ocean (1.2±0.5
 358 μmol L⁻¹) whereas bottom waters have enhanced TDAA concentrations even higher than those
 359 in surface waters (6.0±3.4 μmol L⁻¹).

360 The AAC/C is between 5.4-66 % in SPM and trap samples and the AAN/N (not shown) is
 361 between 3.7-100 %. The overall pattern found for AAC/C (Fig. 2b) is similar to the pattern of
 362 AA concentrations (Fig. 2a) but there is more overlap of AAC/C between the different groups.
 363 Sediments have AAC/C between 2.7-50 % and AAN/N between 3-78 % (not shown). The

364 contribution of AAC to DOC (AAC/C) in water samples is between 4-40 % and in pore water
365 samples between 0.5-9 %.




366
367 Figure 3: AA concentrations in nmol g⁻¹ in sediment traps (a) and SPM (b). Red triangles mark
368 samples from the Kara Sea, black dots are samples from the other trap and SPM locations shown
369 in Figure 1. The decay functions are calculated from samples excluding Kara Sea samples.

370
371 AA concentrations of sinking and suspended particles decrease with water depth and the most
372 significant decrease occurs in the upper ocean (Fig. 3a). The decay constant of AA of sinking
373 particles is twice as high as the decay constant of AA of SPM (Fig. 3 a, b). Kara Sea samples
374 were excluded from these calculations as their AA concentrations are low due to the strong
375 dilution by material from rivers and resuspended sediments in this near-shore environment (see
376 4.2). It is also notable that AAC/C and AAN/N (not shown) significantly decrease between
377 shallow and deep traps and from deep traps to sediments while AAC/C of SPM show little
378 decrease between shallow and deep samples (Fig. 2b).

379
380 **3.2 Amino acid composition**
381 While the AA concentrations of sinking particles from traps show a distinct decrease with water
382 depths the changes in AA composition are lesser so that we averaged all AA spectra irrespective

383 of water depths and area of study to compare them with plankton and sediment samples (Figure
 384 4a). Dominant AA in plankton samples are Glu, Gly, Ala and Asp. Sinking particles and
 385 sediments are also dominated by these AA but in the order of Gly, Asp, Glu and Ala (Fig. 4a).
 386 Mol% Asp, Gly, β -Ala, γ -Aba and Orn increase from plankton via sinking particles to sediments
 387 while Mol% of Glu, Ala, Val, Met, Ile, Leu, Tyr and Phe decrease (Fig. 4a). These trends are
 388 further continued with depths in sediment cores (not shown).
 389




390
 391 Figure 4: Average concentrations of individual AA (Mol%) and 1 σ standard deviation (vertical
 392 bars) in plankton (green), sediment trap (red) and sediment (black) samples (a), in SPM from
 393 water depths <200m (green), >200m (red) and TDAA in water samples (black) (b). Asterisks

394 mark the AA with increasing Mol% from plankton via sediment trap samples to sediments (a)
 395 and from shallow SPM via deep SPM to TDAA of water samples.
 396

397 AA composition of SPM shows a clear trend with water depth with enrichments of Mol% Ser,
 398 Glu, Gly, Orn and His and decreases of almost all other AA from shallow waters (<200 m) to
 399 deep waters (>200m) and decreases are most pronounced for Asp, Thr and Lys (Fig. 4b). These
 400 trends are partly resumed by TDAA in water samples with a further enrichment of Mol% Ser,
 401 Orn and His, while Mol% Glu and Gly slightly decrease in water samples compared with deep
 402 SPM (Fig. 4b). AA spectra of pore waters (not shown) are very similar to water samples.

403



404

405 Figure 5: Box and Whisker plot of AA/HA ratios (a) and Gluam/Galam ratios (b), RI (c) and
 406 DI (d) in SPM from water depths <200m (SPM<) and >200m (SPM>), in sediment traps at
 407 water depth <200m (Trap<) and >200 m (Trap>), in sediments (Sed), in water samples (Water)
 408 and in pore water samples (Pore W). Outliers are marked by dots; some outliers above the 90
 409 percentile are cut off for better perceptibility of trends. Note logarithmic scales of AA/HA (a)
 410 and Gluam/Galam (b).

411

412 Biogeochemical indicators reveal the subtle depth dependent trends in sediment traps and
 413 therefore the biogeochemical indicators were averaged separately for shallow and deep

414 sediment trap and SPM samples (Fig. 5, Table 1). The AA/HA in SPM and water samples -
415 with averages between 80.2 and 204.6 - are higher than in traps and sediments (Fig. 5a, b; Table
416 1). The AA/HA decrease from shallow via deep traps to sediments with averages of 25.4, 14.9
417 and 9.1, respectively. The Gluam/Galam is highest in SPM samples, slightly lower in shallow
418 and deep traps and lower in sediments, water and pore water (Fig. 5a, b; Table 1). The RI (Fig.
419 5c; Table 1) shows the same pattern as the ratios of Asp/ β -Ala and the Glu/ γ -Aba (Table 1),
420 with no clear trend between shallow and deep SPM samples and decreases from shallow to deep
421 traps and further to the sediments. Water samples have similar values as sediment samples with
422 average RI of 1.8 and 1.6, respectively, and pore waters have an even lower average RI of 0.9.
423 Similar to the RI the DI is within the same range in shallow and deep SPM samples and the
424 mean values are very close (Fig. 5d; Table 1) while the DI decreases from shallow sediment
425 traps via deep traps to sediments. In contrast to the RI where water samples have lowest values,
426 the highest DI values are found in water and pore water samples. The DOM-DI averages of
427 2.0 ± 0.6 and 2.1 ± 0.7 in water and, respectively, pore water samples are in fact very close to the
428 DI averages (Table 1).

429 In summary, common biogeochemical indicators of organic matter degradation (RI, Asp/ β -Ala,
430 Glu/ γ -Aba) and bacterial OM accumulation (AA/HA, Gluam/Galam) drop and thus imply
431 increasing degradation between shallow and deep sediment traps and between deep traps and
432 sediments, while these indicators reveal little or no degradation with depth in SPM as the
433 patterns of relative enrichment vs. decreases found in SPM and partly also in TDAA of water
434 and pore water samples differ from the degradation pathway depicted by common
435 biogeochemical indicators (Fig. 4, 5; Table 1). The enhanced DI values furthermore, imply that
436 water and SPM samples are less degraded than deep trap and sediment samples and that TDAA
437 in water and pore water samples are least degraded.

438

439

440

441

442

443

444 Table 1: Mean values and standard deviation (Stdev.) of POC [%], DOC [mg/L], amino acid
 445 (AA) concentrations [$\mu\text{mol/g}$ or $\mu\text{mol/L}$], AAC/C%, AAN/N%, ratios of Asp/ β -Ala, Glu/ γ -
 446 Aba, AA/HA and Gluam/Galam, the RI, DI, SDI*, RTI* and ox/anox ratio summarized in traps
 447 at <200m and >200m water depth, sediments, SPM <200m and >200m water depth, water
 448 samples and pore water samples. *definition of these indicators in part 4.2 below.

		Trap <200m	Trap >200m	Sediment	SPM <200m	SPM >200m	Water	Pore Water
POC; DOC [%; mg/L]	Mean	13.6	5.3	1.8	14.9	10.6	0.8	13.0
	Stdev.	± 11.4	± 1.9	± 2.2	± 8.3	± 4.8	± 0.2	± 8.3
Amino Acids [$\mu\text{mol/g}$; $\mu\text{mol/L}$]	Mean	631.3	164.1	49.8	907.7	661.9	3.2	8.8
	Stdev.	± 602.0	± 93.8	± 82.3	± 637.6	± 434.7	± 3.2	± 7.9
AAC/C [%]	Mean	26.5	15.5	10.8	33.9	32.7	10.1	2.8
	Stdev.	± 8.6	± 4.2	± 5.9	± 12.6	± 10.5	± 6.5	± 2.6
AAN/N [%]	Mean	57.0	38.3	24.2	65.6	61.6		
	Stdev.	± 14.8	± 8.8	± 12.8	± 18.4	± 26.4		
Asp/ β -Ala	Mean	71.8	19.2	10.5	57.9	47.3	10.2	2.6
	Stdev.	± 63.4	± 19.5	± 6.6	± 64.7	± 44.9	± 14.7	± 2.4
Glu/ γ -Aba	Mean	45.7	12.9	8.5	103.6	105.3	8.5	8.8
	Stdev.	± 38.8	± 7.2	± 6.9	± 122.7	± 69.6	± 7.0	± 10.2
AA/HA	Mean	25.4	14.9	9.1	84.6	204.6	80.2	106.8
	Stdev.	± 14.7	± 6.6	± 4.6	± 42.7	± 179.1	± 65.2	± 142.7
Gluam/Galam	Mean	3.2	3.6	1.2	3.0	2.1	1.6	0.3
	Stdev.	± 3.3	± 1.1	± 0.3	± 1.8	± 1.7	± 2.4	± 0.2
RI	Mean	15.1	3.9	1.8	18.8	15.7	1.6	0.9
	Stdev.	± 16.4	± 1.5	± 2.2	± 10.6	± 8.3	± 1.8	± 0.7
DI	Mean	0.9	0.3	-0.5	1.0	1.1	2.1	1.5
	Stdev.	± 0.8	± 0.3	± 0.8	± 0.1	± 0.3	± 1.1	± 0.9
SDI	Mean	1.1	0.0	-0.9	0.8	0.8	-0.7	-0.8
	Stdev.	± 0.2	± 0.2	± 1.0	± 0.2	± 0.3	± 0.4	± 0.3
RTI	Mean	0.7	0.1	0.6	0.2	-1.0	-2.8	-2.7
	Stdev.	± 0.3	± 0.3	± 0.4	± 0.5	± 0.4	± 0.6	± 0.4
ox/anox	Mean	1.2	1.3	2.2	1.1	1.0	0.8	0.7
	Stdev.	± 0.3	± 0.2	± 1.1	± 0.1	± 0.1	± 0.3	± 0.3

449

450

451 4 Discussion

452 4.1 Changes during organic matter degradation

453 Our summary of AA data from various locations in the world ocean corroborates earlier
 454 findings that degradation of organic matter by zooplankton and microbes imparts characteristic
 455 changes to AA spectra so that the differences in AA composition are much larger between types
 456 of organic matter and from different water and sediment depths than between different oceanic
 457 areas (Lee, 1988). The AA spectra track the successive degradation of organic matter during
 458 sedimentation from the plankton source via sinking particles, their incorporation into sediments
 459 and their further degradation after burial. The most characteristic changes along this
 460 sedimentation pathway are the relative enrichments (in Mol%) of Gly, Asp and the non-protein
 461 AA β -Ala, γ -Aba and Orn and the relative decrease of AA produced by fresh plankton such as
 462 Glu, Ala, Val, Met, Ile, Leu, Tyr and Phe (Fig. 4a). These changes are depicted by the common

463 biogeochemical indicators: the ratios of proteinaceous AA vs. non-protein AA (RI and Glu/ γ -
464 Aba) decrease along this pathway. Asp/ β -Ala ratios also decrease because β -Ala becomes
465 relatively more enriched than Asp. The DI, originally derived from sediment samples of
466 different degradation states (Dauwe et al., 1999; Dauwe and Middelburg, 1998), decreases from
467 positive values in fresh plankton and most sinking particles to negative values in sediments as
468 it integrates the products of Asp and Gly multiplied with negative factors, and the products of
469 Glu, Met, Ile, Leu, Tyr and Phe multiplied with positive factors (Dauwe et al., 1999).

470 In contrast, the AA in SPM evolve along a different path than the sedimentation pathway (Gaye
471 et al., 2013b). The increases in Mol% Ser, His and Orn and the decreases in Mol% Asp from
472 shallow (<200 m) to deep SPM (>200m) and even further in TCAA of sea water and pore water
473 are either not depicted or even contrary to the trends depicted by the common biogeochemical
474 indicators (Fig. 4b). The striking difference in AA distribution of SPM (Fig. 4) on the one hand
475 and sinking particles and sediments on the other hand as well as the different depth dependent
476 trends (Fig. 4, 5) suggest that there is little exchange between the two types of particles in the
477 ocean. Sinking particles build up sediments and the degradation pathways evident in the water
478 column - namely the accumulation of degradation products and acidic AA often absorbed to
479 carbonates - continue in the sediments. Novel biogeochemical indicators are required for SPM
480 and possibly also DOM to characterize their AA changes.

481

482 **4.2 Results of a PCA: two new biogeochemical indicators**

483 A PCA of individual AA (Mol %, Fig. 6a) of all samples compiled in this study results in two
484 factors which explain 59 % of the total variance within the data set. The first factor delineates
485 the well-known changes along the degradation pathway from plankton via sinking particles to
486 sediments. Phe, Ile, Leu, Glu and Tyr (enriched in fresh plankton) have the highest F1 loadings
487 while Asp, β -Ala and γ -Aba (accumulating during degradation) have the lowest negative F1
488 loadings. Highest F2 loadings are found for Asp, Thr, Lys and Val while Gly, Orn, His and Ser
489 have the most negative F2 loadings. Factor scores of the individual samples (Fig. 6b) plot in a
490 triangular shape with plankton and fresh organic matter from surface waters at the apex with
491 highest F1 and F2 scores. The diverging sides of the triangle mark sinking particles and
492 sediments decreasing in F1 scores on one side and SPM with decreasing F2 scores on the other
493 side (Fig. 6b). Similar trends were observed in earlier studies based on local data sets (Nagel et
494 al., 2016; Gaye et al., 2013b). That samples from greatly different environments reveal the same

495 divergence between sinking particles and SPM with only little overlap (Fig. 4) suggest a general
 496 mechanism operating globally. Most of the overlap encompasses SPM and sediment trap
 497 samples from the Kara Sea all sampled at water depths below 100 m. The Kara Sea is
 498 characterized by sediment resuspension related to strong riverine input in combination with sea
 499 ice dynamics so that many of the Kara Sea SPM and trap samples are mixed with resuspended
 500 sediments (Gaye et al., 2007; Unger et al., 2005). TDAA analyzed in water and pore water form
 501 a cluster with significantly different AA composition from particulate matter, but instead
 502 recapitulating the enrichments of Mol% Ser, His and Orn observed in SPM (Fig. 6b).

503 The precise separation of the degradation pathway of sinking particles and sediments from SPM
 504 and DOM by the PCA suggests that we can use the first factor (F1) to calculate a new sinking
 505 particle and sediment degradation index (SDI)

$$506 \quad SDI = \sum_i \left(\frac{var_i - AVG.var_i}{STD.var_i} \right) \times Loadings . F1_i \quad (4)$$


507 where var_i is the original mole percentage of each AA_i , $AVG.var_i$ and $STD.var_i$ are the mean and
 508 standard deviations, respectively, and $Loading.F1_i$ is the factor loading of the first axis (F1) of
 509 the PCA of the individual amino acid_i shown in Table 2. Most of the F1 loadings resemble those
 510 of the DI of Dauwe et al. (1999) (Table 2) and the SDI and DI thus are significantly correlated
 511 (Table 3).

512 The second factor (F2) - normalized in the same way with the averages and standard deviations
 513 of the same PCA - can be used as an indicator of changes in the AA composition of SPM
 514 possibly related to the residence time or renewal time of the water mass they are transported
 515 with (see discussion in 4.2.2). With longer residence time in the ocean the organic matter in
 516 SPM is likely to become more recalcitrant and the indicator and is therefore named residence
 517 time index (RTI)

$$518 \quad RTI = \sum_i \left(\frac{var_i - AVG.var_i}{STD.var_i} \right) \times Loadings . F2_i \quad (5)$$

519 and is calculated in the same way as the SDI but the factor loadings of the second axis (F2) of
 520 the PCA of the individual amino acid_i (Table 2) is inserted for the term $Loadings.F2_i$.

521



522

523 Figure 6: Results of a PCA of AA (Mol%) of all samples of this study with factor loadings of
 524 amino acids for the first and second factor (a) and factor scores of samples (b). Small arrows in
 525 (a) point to the positions of Val and Ile, respectively. Arrows in (b) indicate progressive
 526 deviation in composition from the plankton source, essentially with increasing water and
 527 sediment depths.

528

529 The SDI allows a separation of trap samples from shallow water depth from those of greater
530 depths (Figure S1). All samples from deep sediment traps have SDI values below 0.5. Likewise,
531 SPM from >200 m depths have lower RTI than most of the samples from shallower depths.
532 Deep trap samples and deep SPM samples form two clearly separated clusters with different
533 SDI and RTI (see Figure S1 for further details).

534 Table 2: Factor loadings of F1 and F2 for calculating the SDI and RTI, respectively, average
535 [Mol%] and standard deviations (Std. Dev.) of AA of samples used for the PCA shown in Figure
536 5 in comparison with the factor loadings (named factor coefficients) of the DI published by
537 Dauwe et al (1999) and their averages [Mol%] and standard deviations used for the DI based
538 on 28 sediment samples.

Amino Acid	Loadings F1 SDI	Average [Mol%]	Std. Dev.	Loadings F2 RTI	DI	DI Average [Mol%]	DI Std. Dev.
Ser	0.067	8.7	3.6	-0.210	0.015	7.2	1.9
His	0.014	2.4	2.3	-0.166	0.158	1.0	0.8
Orn	-0.038	1.2	1.9	-0.152	-	-	-
Tyr	0.162	1.9	1.1	-0.075	0.178	2.1	1.2
Gly	-0.023	16.2	2.8	-0.068	-0.099	17.6	3.8
Glu	0.132	10.5	2.5	-0.001	0.065	10.0	2.3
Leu	0.157	5.9	1.9	0.027	0.169	6.6	1.5
Ile	0.148	3.8	1.0	0.028	0.139	4.5	0.8
β-Ala	-0.161	1.3	2.0	0.036	-	-	-
Phe	0.140	2.9	1.0	0.047	0.134	3.2	1.2
γ-Aba	-0.164	1.0	1.6	0.064	-	-	-
Ala	-0.030	9.1	1.5	0.080	-0.043	11.8	0.8
Arg	0.012	4.2	1.1	0.117	-0.115	6.1	2.3
Val	0.020	5.7	1.1	0.132	-0.044	7.6	1.1
Lys	0.026	5.1	1.2	0.135	-	-	-
Thr	-0.074	5.7	1.2	0.174	-0.129	7.1	1.5
Asp	-0.154	12.2	4.3	0.179	-0.102	13.4	2.7

540

541 **4.2.1 The SDI as an indicator of degradation and oxic vs. anoxic diagenetic conditions of** 542 **sinking particles and sediments**

543 In order to test the performance of our new degradation indices, we separated SPM samples
544 from sinking particles and sediments and correlated the common biogeochemical indicators and
545 individual AA (Mol %) of SPM with the RTI of individual samples while we correlated the
546 same variables of sinking particles and sediments with the SDI (Table 3). We assume that
547 correlations with Pearson correlation coefficients $R > 0.50$ can be considered as “strong
548 correlations” (Cohen, 1988). The SDI correlates moderately to strongly with the common

549 degradation indicators and the best positive correlation is found between SDI and the DI (Table
550 3). The strong correlation among the degradation indicators with POC contents indicates that
551 this common and often measured variable is a good indicator of relative organic matter quality
552 in sinking particles and sediments and all other degradation indices do not perform better than
553 POC concentrations (see correlation coefficients in Table 3). The DI and the SDI, which are to
554 some extent interchangeable, allow a fine tuning of degradation intensities. The most significant
555 negative correlation of the ox/anox ratio with the SDI is preconditioned, as the ox/anox is the
556 quotient of AA enriched by degradation to those enriched in fresh plankton. It should be noted
557 that this negative correlation is even better than the positive correlation of the DI and the SDI.
558 A close look at the SDI and ox/anox in sediment samples suggests that the SDI can be used to
559 distinguish between oxic and anoxic diagenetic conditions (Fig. 7). The sediment samples
560 deposited in regions of bottom water anoxia (surface samples from Namibian shelf at < 200 m
561 depths; a core from the Arabian Sea slope at 775 m) have lower ox/anox ratios and distinctly
562 higher SDI values compared with the samples from similar depths and oxygenated bottom water
563 (e.g. Mediterranean Sea, Kara Sea) (Fig 7a). The SDI performs better than the DI to determine
564 diagenetic conditions as the DI less significantly correlates with the ox/anox indicator (Figure
565 S2). The best fit between the SDI and the ox/anox in sediment samples is an exponential
566 correlation with a correlation coefficient of $R=-0.95$ while the DI and the ox/anox correlate
567 exponentially with an $R=-0.79$ (Fig. 7a, Figure S2). The SDI better depicts the spectral changes
568 in samples deposited under anoxic diagenetic conditions such as those from the Namibian shelf
569 (Nagel et al., 2016) and the Arabian Sea mid-water oxygen minimum zone (Suthhof et al., 2001)
570 while the ox/anox ratio better resolves variations in samples of strong oxic degradation so that
571 the SDI is in fact better suited to determine the threshold of anoxic vs. oxic diagenesis. Another
572 indication of this quality of the SDI is that the anoxic sediments have SDI values in the range
573 of sediment trap samples from the water column while the SDI decreases in sediments under
574 oxic diagenetic conditions (Fig. 7b). Further, the SDI also correlates with other indicators of
575 oxic vs. anoxic conditions. The core SO90-111 KL from within the mid-water oxygen minimum
576 impinging on the Pakistan margin, was used to reconstruct changes in oxygenation during the
577 last 60 ka BP based e.g. on $\delta^{15}\text{N}$ values of total N (Suthhof et al., 2001). The $\delta^{15}\text{N}$ values
578 fluctuated between enhanced values in warm phases due to denitrification in the mid-water
579 oxygen minimum and lower values in cold phases when the oxygen minimum zone was weaker
580 or absent (Suthhof et al., 2001). The SDI very precisely tracks these changes (Fig. 7c) and in
581 accordance with the threshold discernable in Fig. 7a and b we propose that the divide between

582 oxic and anoxic diagenetic conditions is at SDI values between 0 and -0,2 with SDI<-0.2
 583 indicating oxic and SDI >0 indicating anoxic diagenetic conditions (Fig. 7a, b) and we propose
 584 to use this indicator to reconstruct redox conditions from sediment cores. The work of (Carr et
 585 al., 2016) - relying on the DI – suggests that signals of changes in redox conditions can be
 586 preserved even down to 200 m core depth.

587

588 Table 3: Pearson correlation coefficients of the SDI, RI, DI, Asp/β-Ala and Glu/γ-Aba with
 589 selected AA*, the RTI, AAC/C, AAN/N, AA ratios and degradation indices, water depth
 590 (Depth), POC and TN contents (%) and AA concentrations (nmol/g) in sediment trap and
 591 sediment samples (column 2-6). Pearson correlation coefficients of the RTI, RI, DI, Asp/β-Ala
 592 and Glu/γ-Aba with selected AA, the SDI, AAC/C, AAN/N, AA ratios and degradation indices,
 593 water depth (Depth), POC and TN contents (%) and AA concentrations (nmol/g) in SPM
 594 samples (column 8-12).


595 * Only AA with a correlation coefficient $R \geq 0.50$ with at least one of the indicators are shown.

1	2	3	4	5	6	7	8	9	10	11	12
Traps and Sediments	SDI	RI	DI	Asp/β-Ala	Glu/γ-Aba	SPM	RTI	RI	DI	Asp/β-Ala	Glu/γ-Aba
Asp	-0.68	-0.30	-0.83	-0.25	-0.32	Asp	0.81	0.00	-0.61	0.13	-0.17
Thr	0.12	-0.25	-0.25	0.03	0.07	Thr	0.86	-0.02	-0.57	0.06	-0.16
Ser	0.32	0.03	0.16	0.09	0.16	Ser	-0.91	-0.15	0.08	-0.11	-0.01
Gly	-0.21	-0.32	-0.45	-0.31	-0.31	Gly	-0.90	-0.17	0.12	-0.11	0.04
Ala	0.46	0.07	0.17	0.03	0.11	Ala	0.91	0.09	-0.43	0.11	-0.09
Val	0.67	0.31	0.48	0.29	0.31	Val	0.88	0.07	-0.26	0.08	-0.09
Ile	0.87	0.48	0.76	0.52	0.55	Ile	0.68	0.34	0.26	0.26	0.02
Leu	0.88	0.49	0.76	0.52	0.56	Leu	0.65	0.47	0.28	0.35	0.11
Tyr	0.72	0.39	0.76	0.34	0.32	Tyr	-0.14	0.27	0.66	0.02	0.23
Phe	0.89	0.38	0.79	0.40	0.44	Phe	0.78	0.37	0.04	0.29	0.06
β-Ala	-0.85	-0.27	-0.44	-0.32	-0.35	β-Ala	-0.08	-0.66	-0.19	-0.52	-0.12
γ-Aba	-0.79	-0.25	-0.37	-0.28	-0.35	γ-Aba	0.24	-0.53	-0.65	-0.20	-0.40
Lys	0.56	0.14	0.43	0.15	0.15	Lys	0.71	0.05	-0.20	0.03	-0.06
SDI		0.45	0.82	0.46	0.51	SDI	0.07	0.49	0.83	0.23	0.28
RTI	0.04	-0.01	-0.26	0.03	-0.05	RTI		0.10	-0.34	0.12	-0.08
AAC/C%	0.56	0.51	0.53	0.56	0.56	AAC/C%	-0.08	0.14	0.22	0.06	0.09
AAN/N%	0.55	0.40	0.49	0.47	0.47	AAN/N%	-0.05	0.06	0.03	-0.06	0.08
AA/HA	0.54	0.58	0.54	0.73	0.75	AA/HA	-0.57	0.11	0.32	-0.01	0.16
Gluam/Galam	0.36	0.43	0.44	0.40	0.35	Gluam/Galam	0.27	0.43	0.16	0.28	0.08
Asp/β-Ala	0.46	0.93	0.57		0.85	Asp/β-Ala	0.12	0.74	0.10		0.07
Glu/γ-Aba	0.51	0.93	0.56	0.85		Glu/γ-Aba	-0.08	0.37	0.24	0.07	
RI	0.45		0.49	0.93	0.93	RI	0.10		0.33	0.74	0.37
DI	0.82	0.49		0.51	0.56	DI	-0.34	0.33		0.10	0.24
ox/anox	-0.91	-0.30	-0.71	-0.31	-0.37	ox/anox	0.67	-0.19	-0.50	-0.10	-0.09
Depth	-0.67	-0.28	-0.54	-0.28	-0.37	Depth	-0.55	-0.12	0.23	-0.11	0.04
POC	0.57	0.79	0.67	0.77	0.76	POC	0.27	0.30	0.13	0.25	-0.03
N	0.45	0.46	0.50	0.39	0.39	N	0.29	0.30	0.13	0.24	-0.01
AA	0.51	0.84	0.60	0.79	0.76	AA	0.17	0.28	0.15	0.20	0.01

596

597

598



599

600 Figure 7: SDI indicator plotted against the ox/anox ratio with dots indicating oxic sediments
 601 (black) and suboxic to anoxic sediments from Namibia (green) and the Arabian Sea (blue), the
 602 red line marks the exponential fit to all sediment samples with an $R=0.95$ (a). SDI plotted with
 603 water depth (in m) of sediment trap deployment (red) and of sediment sampling (see color code
 604 of a) (b). SDI and the $\delta^{15}\text{N}$ of total N with sediment depth in sediment core SO90-111 KL
 605 correlated with an $R=0.71$ (c); the blue bar marks the threshold of the SDI delimiting oxic and
 606 anoxic diagenetic conditions at an SDI value of about -0.1.

607

608 4.2.2 The RTI as an indicator of suspended matter residence time

609 Changes in SPM composition between shallow and deep waters (Fig. 4b) are depicted by a
 610 decrease of the RTI which is due to the relative depletion of Asp, Thr, Lys and Val with highest
 611 positive factor loadings and the enrichment of Ser, His and Orn with the most negative factor
 612 loadings (Table 2). Both, Ser and Gly (Mol %) are strongly linearly anticorrelated with the RTI
 613 ($R=-0.91$ and -0.90) showing that they can be used instead of the RTI to characterize SPM if
 614 not all AA used for the RTI can be measured. The anticorrelation of the RTI with water depths
 615 ($R=-0.55$; Table 3) is due to the RTI decrease in the upper 200 m only. Below this depth there
 616 is no further trend in the RTI and values scatter between -0.5 and -1.5 (Fig. 8).

617 The results of our PCA also show that below 200 m SPM becomes distinctly decoupled from
618 sinking aggregates (Fig. 7, Figure S1). Both, sinking particles and SPM sampled in the upper
619 ocean mixed layer and euphotic zone resemble fresh plankton whereas below the surface mixed
620 layer they follow different pathways and the chance that SPM and sinking particles interact
621 obviously decreases with water depth which could be due to the scarcity of both types of
622 particles in the deep ocean (McCave, 1984). Our AA results support previous studies on SPM
623 (using e.g. thorium isotopes, radiocarbon and biomarkers) which found that the interaction
624 between sinking particles and SPM by aggregation and disaggregation strongly decreases from
625 the euphotic zone to the meso- and bathypelagic zone and that SPM rather interacts with DOM
626 due to the long residence time of both in the deep ocean (Lam and Marchal, 2015). The observed
627 constant AA composition of SPM below 200 m water depth could be explained by a recalcitrant
628 nature making the AA barely accessible to further microbial degradation. Alternatively, and in
629 analogy to observations of DOC, it could be due low concentrations of SPM which rather than
630 their recalcitrance limits prokaryotic growth and thus organic matter degradation in the deep
631 ocean (Arrieta et al., 2015). The age of the water masses in the upper ocean mixed layer is less
632 than 100 years while deeper waters have ages of several 100 years to maxima of 1600 years in
633 the deep Indian and Pacific Oceans (England, 1995; Gebbie and Huybers, 2012). The constantly
634 low RTI below the mixed layer is thus related to the long residence time of deep SPM in the
635 deep ocean. An earlier study using detailed ventilation ages available from the Atlantic and
636 Pacific subtropical gyres at stations BATS and respectively, HOT showed that changes in AA
637 composition of SPM took place within a few decades (Kaiser and Benner, 2009).

638 Solubilization of particulate matter by exoenzymes and the subsequent uptake in dissolved form
639 (Carlson and Hansell, 2015; Aristegui et al., 2009) leads to an almost complete turnover of
640 originally diverse surface derived organic matter. It is thus feasible that bacterial biomass
641 comprises a large amount of organic matter in compartments of long residence times. However,
642 fresh bacteria and fungi have quite similar AA composition as plankton (Cowie and Hedges,
643 1992) while SPM AA composition is fundamentally different. The high AA/HA ratios not
644 having a clear trend with water depth also suggest that the contribution of bacterial biomass to
645 SPM is small and does not increase with water depth (Table 1; Fig. 5a, b). The observed changes
646 in SPM are thus more likely related to adsorption processes and macromolecule formation of
647 material not digestible to deep sea organisms and resistant to their enzymes. DOM was shown
648 to become adsorbed to mineral surfaces (Keil and Kirchman, 1993; Keil and Kirchman, 1994;
649 Keil et al., 1994; Arnarson and Keil, 2005, 2007). However, degradation of adsorbed AA

650 proceeds on particles (Satterberg et al., 2003; Taylor, 1995). Thus, the constant AA composition
651 in SPM at depths >200 m may indicate that SPM is in equilibrium with TDAA which likewise
652 show no clear depth dependent changes in AA composition (Figure S3, S4). Feasible candidate
653 processes to explain the homogeneity are AA scavenging by SPM or formation of gels (3D
654 networks = biopolymers) which can anneal to larger sizes so that part of the dissolved AA can
655 be passed from the dissolved to the particulate organic carbon pool (Druffel and Williams,
656 1990; Orellana and Leck, 2015). This process is, however, reversible so that there is probably
657 an exchange between the gel and particulate matter phase as well as between gels of different
658 sizes and complexities depending on pH, temperature, the presence of ligands, pollutants or UV
659 radiation (Orellana and Leck, 2015). Generally, hydrophobic AA (Ala, Val, Met, Ile, Leu, Phe,
660 Pro, Trp) and aromatic AA (His, Tyr) are more likely to form gels and aggregates (Orellana and
661 Leck, 2015). Our results indicate that an equilibrium may be attained between the dissolved
662 phase and SPM after a relatively short time so that the AA composition of SPM is constant
663 below 200 m water depth. If there is no further significant scavenging of SPM by sinking
664 particles and no degradation of AA on SPM, their abundance could increase due to further
665 adsorption of DOM with increasing age on the ocean conveyor belt. However, large
666 zooplankters may be able to utilize the SPM pool (Koppelman et al., 2009; Gloeckler et al.,
667 2018; Hannides et al., 2013) and further studies are required to elucidate the fate of SPM in the
668 ocean.

669

670 **4.3 Contents and composition of total dissolved amino acids in sea water and pore water**

671 The TDAA concentrations show a decrease from the epi- to the mesopelagic ocean similar to
672 many earlier findings (Davis et al., 2009; Kaiser and Benner, 2009; Kim et al., 2017) whereas
673 the spectra of TDAA sampled in the oligotrophic Indian Ocean Subtropical Gyre and the deep
674 Pacific are uniform with water depth. This is also reflected in the DOM-DI which does not
675 show any trend neither with depth in the water column nor in sediments (Figure S4). Further,
676 the difference between water and pore water samples is small and Ser, Gly and His are
677 uniformly the major TDAA in sea water and pore water. It is possible that the selective
678 accumulation of these AA in the dissolved phase is due to their excretion or their association
679 with exoenzymes. Ser is present in N-acyl homoserine lactone (AHLs) which is a class of
680 bacterially produced signaling molecules involved in bacterial quorum sensing; these
681 compounds serve to regulate growth by changing gene expressions, for example, in order to

682 influence population density or phenotype (Parsek et al., 1999; Klein et al., 2009). His changes
683 from its protonated to deprotonated form at a pH of 6 and is therefore often present at the active
684 sites of enzymes. Ser and Gly may simply remain dissolved in sea water as they are hydrophilic.
685 Once mixed into the deeper ocean the scarcity of bacteria or the incorporation of AA into gels
686 could be the reason for their recalcitrance. However, we do not assume that a considerable part
687 of the TDAA belong to dissolved free AA. Because the differences between samples from
688 different regions are much smaller than the difference between the molecular weight fractions
689 and sea water vs. pore-water (Figure S3), we surmise that the formation and transformation
690 processes of DOC are very uniform in the ocean. This assumption is based on limited data so
691 that these results are rather preliminary. We also do not have enough spatial coverage of SPM
692 and TDAA data in the deep ocean to detect AA utilization by organisms or sorption and
693 desorption processes. Both these organic matter pools are large (see below), so that such
694 investigations are important to estimate the possible role of these pools in oceanic carbon
695 sequestration and the reactions to global change (Ridgwell and Arndt, 2015; Lonborg et al.,
696 2018).


697 Our AA yield of DOC with AA-C/C of $10.1 \pm 6.5\%$ are in the high range of studies on from the
698 literature. In some of the previous studies AAC/C was between 0.4 and 4 % with a reduction
699 from 1-4 % AAC/C in surface waters to 0.4-0.8 % in waters >1.000 m. This reduction was
700 moreover, associated with a progressive AA degradation reflected in the DI and the DOM-DI
701 at some of the sampling stations (Kim et al., 2017; Davis and Benner, 2005; Kaiser and Benner,
702 2009). These lower yields may however, be due to different hydrolysis conditions as these
703 studies used water vapor hydrolysis at higher temperatures (150°C) but for a much shorter
704 duration (32.5 minutes). Studies using the same hydrolysis conditions as this work reported
705 AAC/C of 5-10 % (Ittekkot, 1981; Keil and Kirchman, 1999; McCarthy et al., 1997).

706

707 **4.4 $\delta^{15}\text{N}$ values in sinking and suspended matter and evidence for nitrogen sources and** 708 **transformation processes**

709 The $\delta^{15}\text{N}$ values in sediments can preserve information on N sources throughout the geological
710 history (Sun et al., 2019; Gaye et al., 2018; Kienast et al., 2008). However, $\delta^{15}\text{N}$ values may be
711 modulated by organic matter cycling and diagenetic processes which are replicated and thus
712 traceable in the AA composition not least because AA are the main identifiable contributors to
713 N in particulate organic matter. The increase of $\delta^{15}\text{N}$ values by about 2 ‰ on average during

714 organic matter burial and early diagenesis in the upper sediments (Robinson et al., 2012; Tesdal
 715 et al., 2013) is corroborated by a parallel shift in AA based degradation indicators (Gaye-Haake
 716 et al., 2005; Möbius et al., 2010). In contrast to sediments, there are no clear depth related trends
 717 in $\delta^{15}\text{N}$ values of sinking particles in the water column of the epi- to mesopelagic ocean (Gaye-
 718 Haake et al., 2005; Yang et al., 2017; Altabet, 2006). AA based biogeochemical indicators
 719 revealed degradation with depth at specific trap sites (Haake et al., 1993a) and $\delta^{15}\text{N}$ analyses of
 720 individual amino acids showed that degradation is proceeding on sinking particles with $\delta^{15}\text{N}$
 721 changes of “trophic” AA while $\delta^{15}\text{N}$ of “source” AA remained constant (McCarthy et al., 2007).
 722 However, degradation of sinking particles is much smaller than degradation at the sediment
 723 water interface and in our large data set that integrates many different areas of study the small
 724 to moderate changes in AA degradation are obviously obliterated, as neither AA concentrations
 725 (Fig. 3), nor the SDI (Fig. 7), the AAC/C % (Figure S5) nor AAN/N % (not shown) reveal any
 726 significant trends in sinking particles in the deep ocean.



727
 728 Figure 8: The RTI (black dots) and the $\delta^{15}\text{N}$ values of SPM with water depths [m] (red circles).

729 AA composition of SPM as expressed in the RTI is constant and SPM is rather recalcitrant at
730 water depths >200 m. Paralleling this, the $\delta^{15}\text{N}$ values of SPM are about 6-8 ‰ on average in
731 all our studies carried out (Fig. 8). In previous studies $\delta^{15}\text{N}$ values of SPM were reported to
732 increase from ≤ 5 ‰ in surface waters to values between 6-8 ‰ below 200 m water depth which
733 was attributed to organic matter degradation on SPM (Yang et al., 2017; Altabet et al., 1991;
734 Hannides et al., 2013; Emeis et al., 2010). However, SPM samples from the Arabian Sea
735 upwelling area show decreasing $\delta^{15}\text{N}$ values from an average of 8.6 ‰ at water depth above
736 200 m to 7.4 ‰ at depths below 200 m (Gaye et al., 2013b). It is thus reasonable that SPM has
737 a constant $\delta^{15}\text{N}$ value in the mesopelagic and bathypelagic ocean. This is an additional indicator
738 of a common process determining the AA composition and their $\delta^{15}\text{N}$ values of SPM and
739 probably also of DOM sampled below water depths of 200 m (equivalent to an age of ≥ 100
740 years; (England, 1995; Gebbie and Huybers, 2012).

741

742 **4.5 Abundance of amino acids in the ocean**

743 Based on POC, TN and AA fluxes and the area of the open ocean and shallow seas (Costello et
744 al., 2010) we can estimate annual downward fluxes (Table S6). Average POC flux of
745 compilations of trap fluxes were between $1.65 \text{ g m}^{-2} \text{ a}^{-1}$ (Wilson et al., 2012) and $2.74 \text{ g m}^{-2} \text{ a}^{-1}$
746 (Rixen et al., 2019a) while our subset of trap samples from the open ocean (>2000 m water
747 depth) averages to $3.06 \text{ g m}^{-2} \text{ a}^{-1}$. For open ocean traps this results in total fluxes of 0.51-0.94
748 PgC a^{-1} . Our average flux estimates for TN are 0.13 PgN a^{-1} and for AAC are 0.15 PgAA a^{-1} .
749 The flux rates over the shelves and slopes bear, however, large uncertainty because productivity
750 is by several orders of magnitude higher than in offshore areas and spatially variable. Our first
751 estimate, simply based on an average of our fluxes caught in traps deployed in areas of water
752 depth < 2000 m arrives at POC fluxes of 5.4 PgC a^{-1} , TN fluxes of 0.9 PgN a^{-1} and AAC fluxes
753 of $1.36 \text{ Pg AAC a}^{-1}$. Thus 85-90 % of fluxes occur in near shore environments corroborating
754 that 95 % or the total marine organic carbon is buried in these environments (Hedges and Keil,
755 1995). The total sinking fluxes in the proximal plus distal ocean add up to 6.3 PgC a^{-1} , 1.0 PgN
756 a^{-1} and, respectively, $1.51 \text{ Pg AAC a}^{-1}$ (see Table S6 for further details).

757 The largest organic carbon pool in the ocean is DOC with an inventory of $632 \pm 32 \text{ PgC}$ (Carlson
758 and Hansell, 2015; Hansell et al., 2009) and the largest N pool is DON with $77 \pm 23 \text{ PgN}$ (Gruber,
759 2008; Bronk, 2002). Dissolved AA are thus the largest AA pool in the ocean even if AA
760 comprise only a minor amount of DOC. We have only few measurements of AA concentrations,

761 which range between 0.1-0.2 mg/L with an average of 0.16 mg/L in all water samples excluding
762 bottom water. Based on these data we can estimate that AA comprise about 200 ± 70 Pg which
763 would contribute about 35 ± 11 Pg AAN and about 89 ± 29 Pg AAC to the oceanic DON and,
764 respectively, DOC pools. Accordingly, AAC contributes about 14 % to DOC while AAN
765 contributes 45 % to total oceanic DON. This is in the low range of an estimate of 45-86 % AAN
766 based on NMR spectroscopy with acid hydrolysis suggested to recover about half of this AAN
767 pool (Aluwihare et al., 2005).

768 The constant composition of TDAA throughout the ocean indicates that it belongs to the
769 recalcitrant or refractory pool of DOC; this pool is hardly removed in the deep sea and may
770 only be degraded by photochemical reactions as it is returned into surface waters in the course
771 of ocean circulation (Legendre et al., 2015). Our TDAA data reveal no depth dependent trend
772 but our data coverage is not sufficient to detect any spatial variation. The distribution of DOC
773 is, however, well known with its maximum in surface water with $40-80 \mu\text{mol C kg}^{-1}$ and
774 depletion in deep water with DOC concentrations from $>50 \mu\text{mol C kg}^{-1}$ in the North Atlantic
775 to $39 \mu\text{mol C kg}^{-1}$ in the North Pacific deep water (Carlson and Hansell, 2015; Hansell et al.,
776 2009). Due to our limited number of measurements we may have missed spatial variations
777 which could elucidate TDAA sources and cycling processes in the ocean as is the case for DOC.
778 Respiration of DOC may be an important removal process in shallower waters (Reinthal et
779 al., 2006) while a large proportion of the DOC reduction on its way to the Pacific on the deep
780 conveyor belt could be related to adsorption to POC, partly via gel formation (Druffel and
781 Williams, 1990).

782 TDAA may be among the constituents of DOC, which interact with SPM as both are transported
783 with their specific water masses by the ocean conveyor belt. Interaction with SPM is suggested
784 by the relative similarity in AA composition of TDAA and SPM. Moreover, SPM carries the
785 second largest pool of POC and AA in the ocean which has not been accounted for in carbon
786 budgets and which role in oceanic biogeochemical cycling has received little attention. The
787 total abundance of POC, TN and AA in SPM can be calculated using average concentrations
788 (Table 1) in the ocean volume between 0-200 m and between 200 m and the sea floor (Costello
789 et al., 2010). These calculations show that there are 443 Pg of total suspended matter in the
790 ocean of which organic carbon comprises 48 PgC, amino acids 35 PgAA and, total nitrogen 6
791 PgN. The relative similarity of AA spectra in SPM and TDAA suggests interaction between the
792 two pools at shallower depths and the build-up of an equilibrium, so that both pools remain

793 constant in concentrations and composition with depths. Like DOC, which was suggested to be
794 recalcitrant in the deep sea (Hansell and Carlson, 2013), SPM may only be affected by
795 degradation and repackaging into aggregates as it is reintroduced into surface water by ocean
796 circulation. Several studies, however, suggest that SPM may be an important food source for
797 deep living zooplankton (Koppelman et al., 2009; Hannides et al., 2013; Gloeckler et al.,
798 2018). If there are no removal processes in the deep ocean, we would expect SPM and their
799 organic constituents to be exported from the Atlantic via the deep ocean circulation and to
800 accumulate in the Pacific.

801

802 **5 Conclusions**

803 The PCA of a set of 1425 samples consisting of sinking particle, SPM, sediment and water
804 samples produced two factors which separate AA in sinking particles and sediments on the one
805 hand from SPM and DOM on the other hand. As the PCA produced two branches diverging
806 with water and, respectively, sediment depth, strong interactions between the sinking and
807 suspended particles pools can be excluded.

808 The relative degradation of sinking particles and sediments, dominated by Gly, Asp, Glu and
809 Ala, can be tracked by a new degradation indicator named Sediment Degradation Index (SDI)
810 derived from the first factor of the PCA and correlated with the often-used degradation index
811 DI. Except the SDI and the DI the other biogeochemical indicators tested here (Asp/ β -Ala,
812 Glu/ γ -Aba, RI) are not better than POC concentrations for a relative classification of organic
813 matter degradation. The SDI is moreover, capable to separate oxic and anoxic diagenetic
814 conditions at an SDI between 0 to -0.2 (with values <-0.2 indicating oxic and values >0
815 indicating anoxic diagenetic conditions). Application of the SDI furthermore, shows that the
816 diagenetic signal from the water column is preserved in sediments deposited under anoxic
817 conditions. The correlation of the SDI with POC shows that anoxic diagenesis enhances POC
818 accumulation in sediments compared to oxic diagenesis.

819 A novel biogeochemical indicator derived from the second factor of the PCA named Residence
820 Time Indicator (RTI) depicts the transformation of SPM enriched in plankton derived AA in
821 the epipelagic ocean to a constant composition in the meso- and bathypelagic ocean. The deep
822 SPM is probably the residue of microbial processing and is not utilizable by enzymes under the

823 present oceanic conditions. This constant composition of SPM is corroborated by a constant
824 $\delta^{15}\text{N}$ value below 200 m irrespective of the area of study.

825 DOM has almost constant AA composition throughout the water column as well as in pore
826 water, dominated by Ser, Gly, His, Ala and Orn, pursuing the same accumulation AA pathway
827 as found in deep SPM. Comparison with literature data shows that the amount of AA released,
828 depends on the intensity of hydrolysis and that about 50 % of the amide linkages detectable by
829 NMR spectroscopy cannot be hydrolyzed. Similar to SPM the proteins are not utilizable by
830 microorganisms. Protein-like dissolved material was determined to be on average 2670 years
831 old (Loh et al., 2004), showing that these refractory molecules are cycled for several times
832 before they can be removed by as yet unknown processes.

833 Based on our AA data we have calculated the total oceanic AA inventory and found that TDAA
834 are the largest oceanic AA pool with a total amount of 200 ± 70 PgAA and AA comprise 14 %
835 of the oceanic DOC and 45 % of oceanic DON.

836 The pool transported with SPM is 35 PgAA. SPM, furthermore, carries 48 PgC and 6 PgN not
837 accounted for in global carbon and nitrogen budgets. At present it is not known how the oceanic
838 DOM and SPM-particulate organic matter pool is formed and how this rather recalcitrant
839 organic matter can be removed from its ambient water mass. It is feasible that these organic
840 matter pools have fluctuated in the past due to change in oceanic physicochemical conditions
841 (Ridgwell and Arndt, 2015). It is intriguing to understand how the accumulation or reduction
842 of this carbon and nitrogen pools has interacted with climate and environmental changes in the
843 geological history but it is vital to understand the response to ongoing and future climate
844 change.

845

846 **Data Availability**

847 Excerpts of the data were used in previous publications (i) from the Kara Sea in Gaye et al.
848 (2007) Nagel et al. (2005) and Unger et al. (2009), (ii) from the northern Indian Ocean in Gaye
849 et al. (2013), Gaye-Haake et al. (2005), Möbius et al. (2011) and Suthhof et al. (2001), (iii) from
850 the Mediterranean Sea in Möbius (2013) and Möbius et al. (2010), (iv) from the Namibian
851 upwelling in Nagel et al. (2016) and (v) from the Pacific in Paul et al. (2018). The entire set of
852 amino acid data was submitted to PANGAEA. Data from the Pacific are available at:
853 <https://doi.pangaea.de/10.1594/PANGAEA.885391>, <https://doi.pangaea.de/10.1594/PANGAEA.881804>,
854 <https://doi.pangaea.de/10.1594/PANGAEA.881813> and for TOC at

855 <https://doi.pangaea.de/10.1594/PANGAEA.884975>, <https://doi.pangaea.de/10.1594/PANGAEA.885023>,
856 <https://doi.pangaea.de/10.1594/PANGAEA.884981>.

857 .

858

859 **Acknowledgements**

860 This work is based on samples taken during cruises of research vessels R/V SONNE, METEOR,
861 MARIA S. MERIAN, PELAGIA, ORV SAGAR KANYA and R/V AKADEMIK PETROV.
862 We are grateful to all officers and crew of these research vessels. We are indebted to
863 Venugopalan Ittekkot who initiated and inspired this work and led many of the research
864 projects. We thank Desmond Gracias, Areef Sardar and Fernando Vijayan from NIO, Goa,
865 India for technical support on board. We thank Inken Preuss, Annika Moje, Tim Jesper Suhrhoff
866 and Seinab Bohsung for help with pore water sampling during the cruises SO239 and SO242.
867 We are indebted to Frauke Langenberg and Marc Metzke for their high-quality analyses and to
868 Dilek Topcu for help with compiling and archiving the large data sets. We are very thankful to
869 Cindy Lee and two anonymous reviewers for their critical comments on the manuscript. We
870 thank the German Federal Ministry of Education and Research for funding (grant no.:
871 03F0707G) in the framework JPI Oceans EcoMining-DEU - Ecological Aspects of Deep-Sea
872 Mining to Jacobs University Bremen. Sampling in the southern Indian Ocean was conducted
873 within the framework of the INDEX program of the Federal Institute for Geosciences and
874 Natural Resources (BGR). The DFG and BMBF financed the finalized projects in the northern
875 Indian Ocean and Atlantic (Indian-German Program, JGOFS, BIGSET), the Mediterranean Sea
876 (MEDNIT), the Namibian upwelling (GENUS) and the Kara Sea (SIRRO).

877

878 **Author Contribution**

879 BG, NL, TR and KE designed the study and led the projects in which samples were taken and
880 analyzed. NL developed and refined the AA analyses. NL, NH and SP contributed and analyzed
881 samples from the southern Indian Ocean and the Pacific. BG wrote the manuscript with
882 contributions of all co-authors.

883

884

885 **Competing interests**

886 The authors declare that they have no conflict of interest.

887

888 **References**

889 Abramson, L., Lee, C., Liu, Z., Wakeham, S. G., and Szlosek, J.: Exchange between suspended
890 and sinking particles in the northwest Mediterranean as inferred from the organic composition
891 of in situ pump and sediment trap samples, *Limnology and Oceanography*, 55, 725-739, 2011.

892 Alldredge, A.: The carbon, nitrogen and mass content of marine snow as a function of aggregate
893 size, *Deep-Sea Research I*, 45, 529-541, 1998.

894 Alldredge, A. L. and Silver, M. W.: Characteristics, dynamics and significance of marine snow,
895 *Progress in Oceanography*, 20, 41-82, 1988.

896 Altabet, M. A.: Isotopic Tracers of the Marine Nitrogen Cycle: Present and Past, in: *Marine*
897 *Organic Matter: Biomarkers, Isotopes and DNA. The Handbook of Environmental Chemistry*,
898 edited by: Volkman, J. K., Springer, Berlin, Heidelberg, 251-293, doi: 10.1007/698_2_008,
899 2006.

900 Altabet, M. A., Deuser, W. G., Honjo, S., and Stienen, C.: Seasonal and depth-related changes
901 in the source of sinking particles in the North Atlantic, *Nature*, 354, 136-139, 1991.

902 Aluwihare, L. I., Repeta, D. J., Pantoja, S., and Johnson, C. G.: Two chemically distinct pools
903 of organic nitrogen accumulate in the ocean, *Science*, 308, 1007-1010, 2005.

904 Aristegui, J., Gasol, J. M., Duarte, C. M., and Herndl, G. J.: Microbial oceanography of the dark
905 ocean's pelagic realm, *Limnology and Oceanography*, 54, 1501-1529,
906 10.4319/lo.2009.54.5.1501, 2009.

907 Armstrong, R. A., Lee, C., Hedges, J. I., Honjo, S., and Wakeham, S. G.: A new, mechanistic
908 model for organic carbon fluxes in the ocean based on the quantitative association of POC with
909 ballast minerals, *Deep-Sea Research Part II*, 49, 219-236, 2002.

910 Arnarson, T. S. and Keil, R. G.: Influence of organic-mineral aggregates on microbial
911 degradation of the dinoflagellate *Scrippsiella trochoidea*, *Geochimica Et Cosmochimica Acta*,
912 69, 2111-2117, 10.1016/j.gca.2004.11.004, 2005.

913 Arnarson, T. S. and Keil, R. G.: Changes in organic matter-mineral interactions for marine
914 sediments with varying oxygen exposure times, *Geochimica et Cosmochimica Acta*, 71, 3545-
915 3556, 10.1016/j.gca.2007.04.027, 2007.

916 Arrieta, J. M., Mayol, E., Hansman, R. L., Herndl, G. J., Dittmar, T., and Duarte, C. M.: Dilution
917 limits dissolved organic carbon utilization in the deep ocean, *Science*, 348, 331-333,
918 10.1126/science.1258955, 2015.

919 Benner, R. and Kaiser, J.: Abundance of amino sugars and peptidoglycan in marine particulate
920 and dissolved organic matter, *Limnology and Oceanography*, 48, 118-128, 2003.

921 Benner, R., Louchouart, P., and Amon, R. M. W.: Terrigenous dissolved organic matter in the
922 Arctic Ocean and its transport to surface and deep waters of the North Atlantic, *Global*
923 *Biogeochemical Cycles*, 19, GB2025, 10.1029/2004GB002398, 2005.

- 924 Boetius, A. and Lochte, K.: Regulation of microbial enzymatic degradation of organic matter
925 in deep-sea sediments *Mar. Ecol.-Prog. Ser.*, 104, 299-307, 10.3354/meps104299, 1994.
- 926 Boetius, A., Ferdelman, T., and Lochte, K.: Bacterial activity in sediments of the deep Arabian
927 Sea in relation to vertical flux, *Deep-Sea Res. Part II-Top. Stud. Oceanogr.*, 47, 2835-2875,
928 10.1016/s0967-0645(00)00051-5, 2000a.
- 929 Boetius, A., Springer, B., and Petry, C.: Microbial activity and particulate matter in the benthic
930 nepheloid layer (BNL) of the deep Arabian Sea, *Deep-Sea Res. Part II-Top. Stud. Oceanogr.*,
931 47, 2687-2706, 10.1016/s0967-0645(00)00045-x, 2000b.
- 932 Boyd, P., Claustre, H., Levy, M., Siegel, D., and Weber, T.: Multi-faceted particle pumps drive
933 carbon sequestration in the ocean, *Nature*, 568, 327-335, 10.1038/s41586-019-1098-2, 2019.
- 934 Boyd, S. R.: Nitrogen in future biosphere studies, *Chemical Geology*, 176, 1-30,
935 [https://doi.org/10.1016/S0009-2541\(00\)00405-8](https://doi.org/10.1016/S0009-2541(00)00405-8), 2001.
- 936 Brockmeyer, B. and Spitzzy, A.: Evaluation of a Disc Tube Methodology for Nano- and
937 Ultrafiltration of Natural Dissolved Organic Matter, *International Journal of Organic*
938 *Chemistry*, 3, 17-25, 2013.
- 939 Bronk, D. A.: Biogeochemistry of marine dissolved organic matter, in, edited by: Hansell, D.
940 A., and Carlson, C. A., San Diego, 2002.
- 941 Carlson, C. A. and Hansell, D. A.: DOM sources, sinks, reactivity, and budgets,
942 *Biogeochemistry of Marine Dissolved Organic Matter*, 2nd Edition, Academic Press Ltd-
943 Elsevier Science Ltd, London, 1-693 pp.2015.
- 944 Carr, S. A., Mills, C. T., and Mandernack, K. W.: The use of amino acid indices for assessing
945 organic matter quality and microbial abundance in deep-sea Antarctic sediments of IODP
946 Expedition 318, *Marine Chemistry*, 186, 72-82,
947 <http://dx.doi.org/10.1016/j.marchem.2016.08.002>, 2016.
- 948 Cho, B. C. and Azam, F.: Major role of bacteria in biogeochemical fluxes in the ocean's interior,
949 *Nature*, 332, 441-443, 1988.
- 950 Ciais, P., Sabine, C., Bala, G., Bopp, L., Brovkin, V., Canadell, J., Chhabra, A., DeFries, R.,
951 Galloway, J., Heimann, M., Jones, C., Le Quere, C., Myneni, R. B., Piao, S. L., Thornton, P.,
952 Ahlstrom, A., Anav, A., Andrews, O., Archer, D., Arora, V., Bonan, G., Borges, A. V.,
953 Bousquet, P., Bouwman, L., Bruhwiler, L. M., Caldeira, K., Cao, L., Chappellaz, J., Chevallier,
954 F., Cleveland, C., Cox, P., Dentener, F. J., Doney, S. C., Erisman, J. W., Euskirchen, E. S.,
955 Friedlingstein, P., Gruber, N., Gurney, K., Holland, E. A., Hopwood, B., Houghton, R. A.,
956 House, J. I., Houweling, S., Hunter, S., Hurtt, G., Jacobson, A. D., Jain, A., Joos, F., Jungclaus,
957 J., Kaplan, J. O., Kato, E., Keeling, R., Khatiwala, S., Kirschke, S., Goldewijk, K. K., Kloster,
958 S., Koven, C., Kroeze, C., Lamarque, J. F., Lassey, K., Law, R. M., Lenton, A., Lomas, M. R.,
959 Luo, Y. Q., Maki, T., Marland, G., Matthews, H. D., Mayorga, E., Melton, J. R., Metzl, N.,
960 Munhoven, G., Niwa, Y., Norby, R. J., O'Connor, F., Orr, J., Park, G. H., Patra, P., Peregón,
961 A., Peters, W., Peylin, P., Piper, S., Pongratz, J., Poulter, B., Raymond, P. A., Rayner, P.,
962 Ridgwell, A., Ringeval, B., Rodenbeck, C., Saunio, M., Schmittner, A., Schuur, E., Sitch, S.,
963 Spahni, R., Stocker, B., Takahashi, T., Thompson, R. L., Tjiputra, J., van der Werf, G., van
964 Vuuren, D., Voulgarakis, A., Wania, R., Zaehle, S., and Zeng, N.: Carbon and Other
965 Biogeochemical Cycles, *Climate Change 2013: The Physical Science Basis*, Cambridge Univ
966 Press, Cambridge, 465-570 pp.2014.
- 967 Cohen, J.: *Statistical power analysis for the behavioral sciences* (2nd Ed.) L. Erlbaum
968 Associates, Hillsdale, N.J., 567 pp.1988.

- 969 Costello, M. J., Cheung, A., and De Hauwere, N.: Surface Area and the Seabed Area, Volume,
 970 Depth, Slope, and Topographic Variation for the World's Seas, Oceans, and Countries,
 971 Environ. Sci. Technol., 44, 8821-8828, 10.1021/es1012752, 2010.
- 972 Cowie, G. L. and Hedges, J. I.: Sources and reactivities of amino acids in a coastal marine
 973 environment, Limnology and Oceanography, 37, 703-724, 1992.
- 974 Cowie, G. L. and Hedges, J. I.: Biogeochemical indicators of diagenetic alteration in natural
 975 organic-matter mixtures, Nature, 369, 304-307, 1994.
- 976 Cowie, G. L., Hedges, J. I., Prah, F. G., and De Lange, G. J.: Elemental and major biochemical
 977 changes across an oxidation front in a relict turbidite: An oxygen effect, Geochimica et
 978 Cosmochimica Acta, 59, 33-46, 1995.
- 979 Dauwe, B. and Middelburg, J. J.: Amino acids and hexosamines as indicators of organic matter
 980 degradation state in North Sea sediments, Limnology and Oceanography, 43, 782-798, 1998.
- 981 Dauwe, B., Middelburg, J. J., Hermann, P. M. J., and Heip, C. H. R.: Linking diagenetic
 982 alteration of amino acids and bulk organic matter reactivity, Limnology and Oceanography, 44,
 983 1809-1814, 1999.
- 984 Davis, J. and Benner, R.: Seasonal trends in the abundance, composition and bioavailability of
 985 particulate and dissolved organic matter in the Chukchi/Beaufort Seas and western Canada
 986 Basin, Deep Sea Research Part II: Topical Studies in Oceanography, 52, 3396-3410,
 987 <http://dx.doi.org/10.1016/j.dsr2.2005.09.006>, 2005.
- 988 Davis, J., Kaiser, K., and Benner, R.: Amino acid and amino sugar yields and compositions as
 989 indicators of dissolved organic matter diagenesis, Org. Geochem., 40, 343-352,
 990 10.1016/j.orggeochem.2008.12.003, 2009.
- 991 Degens, E. T. and Ittekkot, V.: Dissolved organic carbon - An overview, Mitt. Geol.-Paläont.
 992 Inst. Univ. Hamburg, 55, 21-38, 1983.
- 993 Degens, E. T. and Ittekkot, V.: A new look at clay-organic interactions, in: Ein Nord-Süd Profil:
 994 Zentraleuropa-Mittelmeerraum-Afrika, edited by: Degens, E. T., Krumbein, W. E., and
 995 Prashnowsky, A. A., Mitteilungen aus dem Geologisch-Paläontologischen Institut der
 996 Universität Hamburg,, Krause-Druck, Stade, 229-248, 1984.
- 997 Degens, E. T. and Ittekkot, V.: Particulate organic carbon - An overview, Mitt. Geol.-Paläont.
 998 Inst. Univ. Hamburg, SCOPE/UNEP Sonderband, Heft 58, 7-27, 1985.
- 999 Degens, E. T. and Mopper, K.: Early diagenesis of organic matter in marine soils, Soil Sci.,
 1000 119, 65-72, 10.1097/00010694-197501000-00010, 1975.
- 1001 Druffel, E. R. M. and Williams, P. M.: Identification of a deep marine source of particulate
 1002 organic carbon using bomb ¹⁴C Nature, 347, 172-174, 10.1038/347172a0, 1990.
- 1003 Emeis, K.-C., Mara, P., Schlarbaum, T., Möbius, J., Dähnke, K., Struck, U., Mihalopoulos, N.,
 1004 and Krom, M.: External inputs and internal N cycling traced by isotope ratios of nitrate,
 1005 dissolved reduced nitrogen, and particulate nitrogen in the eastern Mediterranean Sea, Journal
 1006 of Geophysical Research, 115, 10.1029/2009JG001214, 2010.
- 1007 England, M. E.: The Age of Water and Ventilation Timescales in a Global Ocean Model, J.
 1008 Phys. Oceanogr., 25, 2756-2777, 10.1175/1520-0485(1995)025<2756:taowav>2.0.co;2, 1995.
- 1009 Fowler, S. W. and Knauer, G. A.: Role of large particles in the transport of elements and organic
 1010 compounds through the oceanic water column Progress in Oceanography, 16, 147-194,
 1011 10.1016/0079-6611(86)90032-7, 1986.

- 1012 Galbraith, E. D., Kienast, M., Albuquerque, A. L., Altabet, M. A., Batista, F., Bianchi, D.,
1013 Calvert, S. E., Contreras, S., Crosta, X., De Pol-Holz, R., Dubois, N., Etourneau, J., Francois,
1014 R., Hsu, T. C., Ivanochko, T., Jaccard, S. L., Kao, S. J., Kiefer, T., Kienast, S., Lehmann, M.
1015 F., Martinez, P., McCarthy, M., Meckler, A. N., Mix, A., Mobius, J., Pedersen, T. F., Pichevin,
1016 L., Quan, T. M., Robinson, R. S., Ryabenko, E., Schmittner, A., Schneider, R., Schneider-Mor,
1017 A., Shigemitsu, M., Sinclair, D., Somes, C., Studer, A. S., Tesdal, J. E., Thunell, R., Yang, J.
1018 Y. T., and Members, N. W. G.: The acceleration of oceanic denitrification during deglacial
1019 warming, *Nat. Geosci.*, 6, 579-584, 10.1038/ngeo1832, 2013.
- 1020 Gaye-Haake, B., Unger, D., Nöthig, E.-M., Okolodkov, Y., Fahl, K., and Ittekkot, V.: Particle
1021 fluxes from short term sediment trap deployments in late summer in the southern Kara Sea, In:
1022 R. Stein, K. Fahl, D.K. Fütterer, E. Galimov and O. Stepanets (Eds.). *Siberian River Run-off in
1023 the Kara Sea: Characterisation, Quantification, Variability, and Environmental Significance,*
1024 *Proceedings in Marine Science*, Elsevier, Amsterdam, 309-328, 2003.
- 1025 Gaye-Haake, B., Lahajnar, N., Emeis, K.-C., Unger, D., Rixen, T., Suthhof, A., Ramaswamy,
1026 V., Schulz, H., Paropkari, A. L., Guptha, M. V. S., and Ittekkot, V.: Stable nitrogen isotopic
1027 ratios of sinking particles and sediments from the northern Indian Ocean, *Marine Chemistry*,
1028 96, 243-255, 2005.
- 1029 Gaye, B., Wiesner, M. G., and Lahajnar, N.: Nitrogen sources in the South China Sea, as
1030 discerned from stable nitrogen isotopic ratios in rivers, sinking particles, and sediments, *Marine
1031 Chemistry*, 114, 72-85, 2009.
- 1032 Gaye, B., Nagel, B., Daehnke, K., Rixen, T., and Emeis, K.-C.: Evidence of parallel
1033 denitrification and nitrite oxidation in the ODZ of the Arabian Sea from paired stable isotopes
1034 of nitrate and nitrite, *Global biogeochemical cycles*, 27, 1059-1071, 10.1002/2011GB004115,
1035 2013a.
- 1036 Gaye, B., Nagel, B., Daehnke, K., Rixen, T., Lahajnar, N., and Emeis, K. C.: Amino acid
1037 composition and delta N-15 of suspended matter in the Arabian Sea: implications for organic
1038 matter sources and degradation, *Biogeosciences*, 10, 7689-7702, 10.5194/bg-10-7689-2013,
1039 2013b.
- 1040 Gaye, B., Fahl, K., Kodina, L. A., Lahajnar, N., Nagel, B., Unger, D., and Gebhardt, A. C.:
1041 Particulate matter fluxes in the southern and central Kara Sea compared to sediments: Bulk
1042 fluxes, amino acids, stable carbon and nitrogen isotopes, sterols and fatty acids, *Cont. Shelf
1043 Res.*, 27, 2570-2594, 10.1016/j.csr.2007.07.003, 2007.
- 1044 Gaye, B., Boell, A., Segschneider, J., Burdanowitz, N., Emeis, K.-C., Ramaswamy, V.,
1045 Lahajnar, N., Lueckge, A., and Rixen, T.: Glacial-interglacial changes and Holocene variations
1046 in Arabian Sea denitrification, *Biogeosciences*, 15, 507-527, 10.5194/bg-15-507-2018, 2018.
- 1047 Gebbie, G. and Huybers, P.: The Mean Age of Ocean Waters Inferred from Radiocarbon
1048 Observations: Sensitivity to Surface Sources and Accounting for Mixing Histories, *J. Phys.
1049 Oceanogr.*, 42, 291-305, 10.1175/jpo-d-11-043.1, 2012.
- 1050 Gloeckler, K., Choy, C. A., Hannides, C. C. S., Close, H. G., Goetze, E., Popp, B. N., and
1051 Drazen, J. C.: Stable isotope analysis of micronekton around Hawaii reveals suspended particles
1052 are an important nutritional source in the lower mesopelagic and upper bathypelagic zones,
1053 *Limnology and Oceanography*, 63, 1168-1180, 10.1002/lno.10762, 2018.
- 1054 Goutx, M., Wakeham, S. G., Lee, C., Duflos, M., Guigue, C., Liu, Z., Moriceau, B., Sempère,
1055 R., Tedetti, M., and Xue, J.: Composition and degradation of marine particles with different

- 1056 settling velocities in the northwestern Mediterranean Sea, *Limnology and Oceanography*, 52,
1057 1645-1664, 2007.
- 1058 Gruber, N.: The marine nitrogen cycle: Overview and challenges, in: *Nitrogen in the Marine*
1059 *Environment*, 2nd Edition, edited by: Capone, D. G., Bronk, D. A., Mulholland, M. R., and
1060 Carpenter, E., Academic Press, San Diego, 51, 2008.
- 1061 Guo, J., Liang, S.-k., Li, X.-j., Li, W., Wang, Y., and Su, R.-g.: Composition and bioavailability
1062 of dissolved organic matter in different water masses of the East China Sea, *Estuarine, Coastal*
1063 *and Shelf Science*, 212, 189-202, 10.1016/j.ecss.2018.07.009, 2018.
- 1064 Haake, B., Rixen, T., and Ittekkot, V.: Variability of monsoonal upwelling signals in the deep
1065 western Arabian Sea, *Mitt. Geol.-Paläont. Inst. Univ. Hamburg, Scope/UNEP Sonderband Heft*
1066 *76*, 85-96, 1993a.
- 1067 Haake, B., Ittekkot, V., Honjo, S., and Manganini, S.: Amino acids, hexosamines and
1068 carbohydrate fluxes to the deep Subarctic Pacific (Station P), *Deep-Sea Research I*, 40, 547-
1069 560, 1993b.
- 1070 Haake, B., Ittekkot, V., Ramaswamy, V., Nair, R. R., and Honjo, S.: Fluxes of amino acids and
1071 hexosamines of the deep Arabian Sea, *Marine Chemistry*, 40, 291-314, 1992.
- 1072 Haake, B., Rixen, T., Reemtsma, T., Ramaswamy, V., and Ittekkot, V.: Processes determining
1073 seasonality and interannual variability of settling particle fluxes to the deep Arabian Sea, in:
1074 *Particle Flux in the Ocean*, edited by: Ittekkot, V., Schäfer, P., Honjo, S., and Depetris, P. J.,
1075 John Wiley&Sons Ltd., 251-270, 1996.
- 1076 Hannides, C. C. S., Popp, B. N., Choy, C. A., and Drazen, J. C.: Midwater zooplankton and
1077 suspended particle dynamics in the North Pacific Subtropical Gyre: A stable isotope
1078 perspective, *Limnology and Oceanography*, 58, 1931-1946, 10.4319/lo.2013.58.6.1931, 2013.
- 1079 Hansell, D., Carlson, C. A., Repeta, D. J., and Schlitzer, R.: Dissolved organic matter in the
1080 ocean, *Oceanography*, 22, 202-211, 2009.
- 1081 Hansell, D. A. and Carlson, C. A.: Localized refractory dissolved organic carbon sinks in the
1082 deep ocean, *Global Biogeochemical Cycles*, 27, 705-710, 2013.
- 1083 Harms, N. C., Lahajnar, N., Gaye, B., Rixen, T., Schwarz-Schampera, U., and Emeis, K.-C.:
1084 Sediment trap-derived particulate matter fluxes in the oligotrophic subtropical gyre of the South
1085 Indian Ocean, *Deep Sea Research Part II: Topical Studies in Oceanography*, 183, 104924,
1086 <https://doi.org/10.1016/j.dsr2.2020.104924>, 2021.
- 1087 Harms, N. C., Lahajnar, N., Gaye, B., Rixen, T., Dähnke, K., Ankele, M., Schwarz-Schampera,
1088 U., and Emeis, K. C.: Nutrient distribution and nitrogen and oxygen isotopic composition of
1089 nitrate in water masses of the subtropical southern Indian Ocean, *Biogeosciences*, 16, 2715-
1090 2732, 10.5194/bg-16-2715-2019, 2019.
- 1091 Hedges, J. I. and Hare, P. E.: Amino acid adsorption by clay minerals in distilled water,
1092 *Geochimica et Cosmochimica Acta*, 51, 255-259, 1987.
- 1093 Hedges, J. I. and Keil, R. G.: Sedimentary organic matter preservation an assessment and
1094 speculative synthesis, *Marine Chemistry*, 49, 137-139, 10.1016/0304-4203(95)00013-h, 1995.
- 1095 Hildebrandt, Tatjana M., Nunes Nesi, A., Araújo, Wagner L., and Braun, H.-P.: Amino Acid
1096 Catabolism in Plants, *Molecular Plant*, 8, 1563-1579,
1097 <https://doi.org/10.1016/j.molp.2015.09.005>, 2015.

- 1098 Honjo, S., Manganini, S. J., Krishfield, R. A., and Francois, R.: Particulate organic carbon
1099 fluxes to the ocean interior and factors controlling the biological pump: A synthesis of global
1100 sediment trap programs since 1983, *Progress in Oceanography*, 76, 217-285,
1101 10.1016/j.pocean.2007.11.003, 2008.
- 1102 Ingalls, A. E., Liu, Z., and Lee, C.: Seasonal trends in the pigment and amino acid compositions
1103 of sinking particles in biogenic CaCO₃ and SiO₂ dominated regions of the Pacific sector of the
1104 Southern Ocean along 170°W, *Deep-Sea Research I*, 53, 836-859, 2006.
- 1105 Ingalls, A. E., Aller, R. C., Lee, C., and Wakeham, S. G.: Organic matter diagenesis in shallow
1106 water carbonate sediments, *Geochimica et Cosmochimica Acta*, 68, 4263-4379, 2004.
- 1107 Ittekkot, V.: Verteilung von gelöstem organischen Kohlenstoff, gelösten Zuckern und
1108 Aminosäuren in Fladengrund, nördliche Nordsee (FLEX 1976), *Mitt. Geol.-Paläont. Inst. Univ.*
1109 *Hamburg*, 51, 115-187, 1981.
- 1110 Ittekkot, V. and Arain, R.: Nature of Particulate Organic-Matter in the River Indus, Pakistan,
1111 *Geochimica Et Cosmochimica Acta*, 50, 1643-1653, 10.1016/0016-7037(86)90127-4, 1986.
- 1112 Ittekkot, V., Degens, E. T., and Honjo, S.: Seasonality in the fluxes of sugars, amino acids, and
1113 amino sugars to the deep ocean: Panama Basin, *Deep-Sea Research*, 31, 1071-1083, 1984a.
- 1114 Ittekkot, V., Deuser, W. G., and Degens, E. T.: Seasonality in the fluxes of sugars, amino acids,
1115 and amino sugars to the deep ocean: Sargasso Sea, *Deep-Sea Research*, 31, 1057-1069, 1984b.
- 1116 Ittekkot, V., Safiullah, S., and Arain, R.: Nature of Organic-Matter in Rivers with Deep-Sea
1117 Connections - the Ganges-Brahmaputra and Indus, *Science of the Total Environment*, 58, 93-
1118 107, 10.1016/0048-9697(86)90080-x, 1986.
- 1119 Jennerjahn, T. and Ittekkot, V.: Organic matter in sediments in the mangrove areas and adjacent
1120 continental margins of Brazil: I. Amino acids and hexosamines, *Oceanol. Acta*, 20, 359-369,
1121 1997.
- 1122 Kaiser, K. and Benner, R.: Biochemical composition and size distribution of organic matter at
1123 the Pacific and Atlantic time-series stations, *Marine Chemistry*, 113, 63-77,
1124 <http://dx.doi.org/10.1016/j.marchem.2008.12.004>, 2009.
- 1125 Kandler, O.: Cell-wall structures in methane bacteria - evolution of procaryotes
1126 *Naturwissenschaften*, 66, 95-105, 10.1007/bf00373500, 1979.
- 1127 Karl, D., Knauer, G. A., and Martin, A. P.: Downward flux of particulate organic matter in the
1128 ocean: a particle composition paradox, *Nature*, 332, 438-441, 1988.
- 1129 Keil, R. G. and Kirchman, D. L.: Dissolved Combined Amino Acids: Chemical Form and
1130 Utilization by Marine Bacteria, *Limnology and Oceanography*, 38, 1256-1270, 1993.
- 1131 Keil, R. G. and Kirchman, D. L.: Abiotic Transformation of Labile Protein to Refractory Protein
1132 in Sea-Water, *Marine Chemistry*, 45, 187-196, 10.1016/0304-4203(94)90002-7, 1994.
- 1133 Keil, R. G. and Kirchman, D. L.: Utilization of dissolved protein and amino acids in the northern
1134 Sargasso Sea, *Aquat. Microb. Ecol.*, 18, 293-300, 1999.
- 1135 Keil, R. G., Montlucon, D. B., Prahl, F. G., and Hedges, J. I.: Sorptive Preservation of Labile
1136 Organic-Matter in Marine-Sediments, *Nature*, 370, 549-552, 10.1038/370549a0, 1994.
- 1137 Kienast, M., Lehmann, M. F., Timmermann, A., Galbraith, E., Bolliet, T., Holbourn, A.,
1138 Normandeau, C., and Laj, C.: A mid-Holocene transition in the nitrogen dynamics of the
1139 western equatorial Pacific: Evidence of a deepening thermocline?, *Geophysical Research*
1140 *Letters*, 35, 5, 10.1029/2008gl035464, 2008.

- 1141 Kim, T. H., Kim, G., Shen, Y., and Benner, R.: Strong linkages between surface and deep-water
 1142 dissolved organic matter in the East/Japan Sea, *Biogeosciences*, 14, 2561-2570, 10.5194/bg-
 1143 14-2561-2017, 2017.
- 1144 King, K. J.: Amino acid composition of the silicified matrix in fossil polycystine Radiolaria,
 1145 *Micropaleontology*, 21, 215-226, 1975.
- 1146 King, K. J. and Hare, P. E.: Amino acid composition of the test as a taxonomic character for
 1147 living and fossil planktonic foraminifera, *Micropaleontology*, 18, 285-293, 1972.
- 1148 Klein, I., von Rad, U., and Durner, J.: Homoserine lactones: do plants really listen to bacterial
 1149 talk?, *Plant signaling & behavior*, 4, 50-51, 2009.
- 1150 Koppelman, R., Bottger-Schnack, R., Mobius, J., and Weikert, H.: Trophic relationships of
 1151 zooplankton in the eastern Mediterranean based on stable isotope measurements, *J. Plankton*
 1152 *Res.*, 31, 669-686, 10.1093/plankt/fbp013, 2009.
- 1153 Lahajnar, N., Wiesner, M. G., and Gaye, B.: Fluxes of amino acids and hexosamines to the deep
 1154 South China Sea, *Deep-Sea Res. Part I-Oceanogr. Res. Pap.*, 54, 2120-2144,
 1155 10.1016/j.dsr.2007.08.009, 2007.
- 1156 Lahajnar, N., Rixen, T., Gaye-Haake, B., Schafer, P., and Ittekkot, V.: Dissolved organic carbon
 1157 (DOC) fluxes of deep-sea sediments from the Arabian Sea and NE Atlantic, *Deep-Sea Res. Part*
 1158 *II-Top. Stud. Oceanogr.*, 52, 1947-1964, 10.1016/j.dsr2.2005.05.006, 2005.
- 1159 Lam, P. J. and Marchal, O.: Insights into Particle Cycling from Thorium and Particle Data, in:
 1160 *Annual Review of Marine Science*, Vol 7, edited by: Carlson, C. A., and Giovannoni, S. J.,
 1161 *Annual Review of Marine Science*, Annual Reviews, Palo Alto, 159-184, 10.1146/annurev-
 1162 marine-010814-015623, 2015.
- 1163 Lee, C.: Amino acids and amine biogeochemistry in marine particulate material and sediments,
 1164 In: *Nitrogen cycling in coastal marine environments* (eds. T.H. Blackburn, and J. Sørensen),
 1165 *SCOPE*, Wiley and Sons, 125-141, 1988.
- 1166 Lee, C. and Cronin, C.: The vertical flux of particulate organic nitrogen in the sea:
 1167 decomposition of amino acids in the Peru upwelling area and the equatorial Atlantic, *J. Mar.*
 1168 *Res.*, 40, 227-251, 1982.
- 1169 Lee, C. and Cronin, C.: Particulate amino acids in the sea: Effects of primary productivity and
 1170 biological decomposition, *J. Mar. Res.*, 42, 1075-1097, 1984.
- 1171 Lee, C., Wakeham, S., and Arnosti, C.: Particulate organic matter in the sea: The composition
 1172 conundrum, *Ambio*, 33, 565-575, 10.1639/0044-7447(2004)033[0565:pomits]2.0.co;2, 2004.
- 1173 Legendre, L., Rivkin, R. B., Weinbauer, M. G., Guidi, L., and Uitz, J.: The microbial carbon
 1174 pump concept: Potential biogeochemical significance in the globally changing ocean, *Progress*
 1175 *in Oceanography*, 134, 432-450, 10.1016/j.pocean.2015.01.008, 2015.
- 1176 Loh, A. N., Bauer, J. E., and Druffel, E. R. M.: Variable ageing and storage of dissolved organic
 1177 components in the open ocean, *Nature*, 430, 877-881, 10.1038/nature02780, 2004.
- 1178 Loick-Wilde, N., Weber, S. C., Eglite, E., Liskow, I., Schulz-Bull, D., Wasmund, N., Wodarg,
 1179 D., and Montoya, J. P.: De novo amino acid synthesis and turnover during N-2 fixation,
 1180 *Limnology and Oceanography*, 63, 1076-1092, 10.1002/lno.10755, 2018.
- 1181 Lonborg, C., Alvarez-Salgado, X. A., Letscher, R. T., and Hansell, D. A.: Large Stimulation of
 1182 Recalcitrant Dissolved Organic Carbon Degradation by Increasing Ocean Temperatures,
 1183 *Frontiers in Marine Science*, 4, 11, 10.3389/fmars.2017.00436, 2018.

- 1184 Martin, J. H., Knauer, G. A., Karl, D. M., and Broenkow, W. W.: Vertex: carbon cycling in the
1185 northeast Pacific, *Deep-Sea Research*, 34, 267-285, 1987.
- 1186 Mayzaud, P. and Martin, J. L. M.: Some aspects of biochemical and mineral composition of
1187 marine plankton *J. Exp. Mar. Biol. Ecol.*, 17, 297-310, 10.1016/0022-0981(75)90005-2, 1975.
- 1188 McCarthy, M., Pratum, T., Hedges, J., and Benner, R.: Chemical composition of dissolved
1189 organic nitrogen in the ocean, *Nature*, 390, 150-154, 10.1038/36535, 1997.
- 1190 McCarthy, M. D., Benner, R., Lee, C., and Fogel, M. L.: Amino acid nitrogen isotopic
1191 fractionation patterns as indicators of heterotrophy in plankton, particulate, and dissolved
1192 organic matter, *Geochimica et Cosmochimica Acta*, 71, 4727-4744,
1193 <https://doi.org/10.1016/j.gca.2007.06.061>, 2007.
- 1194 McCarthy, M. D., Benner, R., Lee, C., Hedges, J. I., and Fogel, M. L.: Amino acid carbon
1195 isotopic fractionation patterns in oceanic dissolved organic matter: an unaltered
1196 photoautotrophic source for dissolved organic nitrogen in the ocean?, *Marine Chemistry*, 92,
1197 123-134, 2004.
- 1198 McCave, I. N.: Size spectra and aggregation of suspended particles in the ocean, *Deep-Sea*
1199 *Research*, 31, 329-352, 1984.
- 1200 Menzel, P., Anupama, K., Basavaiah, N., Das, B. K., Gaye, B., Herrmann, N., and Prasad, S.:
1201 The use of amino acid analyses in (palaeo-) limnological investigations: A comparative study
1202 of four Indian lakes in different climate regimes, *Geochimica et Cosmochimica Acta*, 160, 25-
1203 37, <http://dx.doi.org/10.1016/j.gca.2015.03.028>, 2015.
- 1204 Menzel, P., Gaye, B., Wiesner, M. G., Prasad, S., Stebich, M., Das, B. K., Anoop, A., Riedel,
1205 N., and Basavaiah, N.: Influence of bottom water anoxia on nitrogen isotopic ratios and amino
1206 acid contributions of recent sediments from small eutrophic Lonar Lake, Central India,
1207 *Limnology and Oceanography*, 58, 1061-1074, 2013.
- 1208 Möbius, J.: Isotope fractionation during nitrogen remineralization (ammonification):
1209 Implications for nitrogen isotope biogeochemistry, *Geochimica et Cosmochimica Acta*, 105,
1210 422-432, 2013.
- 1211 Möbius, J., Lahajnar, N., and Emeis, K.-C.: Diagenetic control on nitrogen isotope ratios in
1212 Holocene sapropels and recent sediments from the Eastern Mediterranean Sea, *Biogeosciences*
1213 7, 3901-3914, 10.5194/bg-7-3901-2010, 2010.
- 1214 Möbius, J., Gaye, B., Lahajnar, N., Bahlmann, E., and Emeis, K.-C.: Influence of diagenesis on
1215 sedimentary $\delta^{15}\text{N}$ in the Arabian Sea over the last 130 kyr, *Mar. Geol.*, 284, 127-138; doi:
1216 10.1016/j.margeo.2011.1003.1013, 2011.
- 1217 Muller, P. J., Suess, E., and Ungerer, C. A.: Amino acids and amino sugars of surface particulate
1218 and sediment trap material from waters of the Scotia Sea *Deep-Sea Research Part a-*
1219 *Oceanographic Research Papers*, 33, 819-838, 10.1016/0198-0149(86)90090-7, 1986.
- 1220 Müller, P. J.: C/N ratios in Pacific deep sea sediments: Effect of inorganic ammonium and
1221 organic nitrogen compounds sorbed by clays, *Geochimica et Cosmochimica Acta*, 11, 765-776,
1222 1977.
- 1223 Nagel, B., Gaye, B., Kodina, L. A., and Lahajnar, N.: Stable carbon and nitrogen isotopes as
1224 indicators for organic matter sources in the Kara Sea, *Mar. Geol.*, 266, 42-51,
1225 10.1016/j.margeo.2009.07.010, 2009.
- 1226 Nagel, B., Gaye, B., Lahajnar, N., Struck, U., and Emeis, K.-C.: Effects of current regimes and
1227 oxygenation on particulate matter preservation on the Namibian shelf: Insights from amino acid

- 1228 biogeochemistry, Marine Chemistry, 186, 121-132,
1229 <http://dx.doi.org/10.1016/j.marchem.2016.09.001>, 2016.
- 1230 Nagel, B., Emeis, K.-C., Flohr, A., Rixen, T., Schlarbaum, T., Mohrholz, V., and van der Plas,
1231 A.: N-cycling and balancing of the N-deficit generated in the oxygen minimum zone over the
1232 Namibian shelf—An isotope-based approach, *Journal of Geophysical Research:*
1233 *Biogeosciences*, 118, 361-371, 2013.
- 1234 Niggemann, J. and Schubert, C. J.: Sources and fate of amino sugars in coastal Peruvian
1235 sediments, *Geochimica et Cosmochimica Acta*, 70, 2229-2237, 2006.
- 1236 Niggemann, J., Lomstein, B. A., and Schubert, C. J.: Diagenesis of amino compounds in water
1237 column and sediment of Lake Baikal, *Org. Geochem.*, 115, 67-77,
1238 [10.1016/j.orggeochem.2017.10.008](http://dx.doi.org/10.1016/j.orggeochem.2017.10.008), 2018.
- 1239 Orellana, M. V. and Leck, C.: *Marine Microgels, Biogeochemistry of Marine Dissolved*
1240 *Organic Matter*, 2nd Edition, Academic Press Ltd-Elsevier Science Ltd, London, 451-480 pp.,
1241 [10.1016/b978-0-12-405940-5.00009-1](http://dx.doi.org/10.1016/b978-0-12-405940-5.00009-1), 2015.
- 1242 Pantoja, S., Sepúlveda, J., and González, H. E.: Decomposition of sinking proteinaceous
1243 material during fall in the oxygen minimum zone off northern Chile, *Deep-Sea Research I*, 51,
1244 55-70, 2004.
- 1245 Parsek, M. R., Val, D. L., Hanzelka, B. L., Cronan, J. E., Jr., and Greenberg, E. P.: Acyl
1246 homoserine-lactone quorum-sensing signal generation, *Proc. Natl. Acad. Sci. U. S. A.*, 96,
1247 4360-4365, 1999.
- 1248 Paul, S. A. L., Gaye, B., Haeckel, M., Kasten, S., and Koschinsky, A.: Biogeochemical
1249 Regeneration of a Nodule Mining Disturbance Site: Trace Metals, DOC and Amino Acids in
1250 Deep-Sea Sediments and Pore Waters, *Frontiers in Marine Science*, 5,
1251 [10.3389/fmars.2018.00117](http://dx.doi.org/10.3389/fmars.2018.00117), 2018.
- 1252 Peter, S., Shen Y., Kaiser, K., Benner, R., Durisch-Kaiser, E. : Bioavailability and diagenetic
1253 state of dissolved organic matter in riparian groundwater, *Journal of Geophysical Research OF*
1254 *GEOPHYSICAL RESEARCH*, 117, G04006, doi:10.1029/2012JG002072, 2012.
- 1255 Peters, B. D., Lam, P. J., and Casciotti, K. L.: Nitrogen and oxygen isotope measurements of
1256 nitrate along the US GEOTRACES Eastern Pacific Zonal Transect (GP16) yield insights into
1257 nitrate supply, remineralization, and water mass transport, *Marine Chemistry*, 201, 137-150,
1258 <https://doi.org/10.1016/j.marchem.2017.09.009>, 2018.
- 1259 Pilskaln, C. H. and Honjo, S.: The fecal pellet fraction of biogeochemical particle fluxes to the
1260 deep sea, *Global Biogeochemical Cycles*, 1, 31-48, 1987.
- 1261 Reinthaler, T., van Aken, H., Veth, C., Aristegui, J., Robinson, C., Williams, P., Lebaron, P.,
1262 and Herndl, G. J.: Prokaryotic respiration and production in the meso- and bathypelagic realm
1263 of the eastern and western North Atlantic basin, *Limnology and Oceanography*, 51, 1262-1273,
1264 [10.4319/lo.2006.51.3.1262](http://dx.doi.org/10.4319/lo.2006.51.3.1262), 2006.
- 1265 Resplandy, L., Lévy, M., and McGillicuddy Jr., D. J.: Effects of Eddy-Driven Subduction on
1266 Ocean Biological Carbon Pump, *Global Biogeochemical Cycles*, 33, 1071-1084,
1267 <https://doi.org/10.1029/2018GB006125>, 2019.
- 1268 Riccardi, G., Derossi, E., and Milano, A.: Amino acid biosynthesis and its regulation in
1269 cyanobacteria *Plant Sci.*, 64, 135-151, [10.1016/0168-9452\(89\)90018-6](http://dx.doi.org/10.1016/0168-9452(89)90018-6), 1989.
- 1270 Ridgwell, A. and Arndt, S.: Why Dissolved Organics Matter: DOC in Ancient Oceans and Past
1271 Climate Change, *Biogeochemistry of Marine Dissolved Organic Matter*, 2nd Edition, Academic

- 1272 Press Ltd-Elsevier Science Ltd, London, 1-20 pp., 10.1016/b978-0-12-405940-5.00001-7,
1273 2015.
- 1274 Rixen, T., Gaye, B., and Emeis, K.-C.: The monsoon, carbon fluxes, and the organic carbon
1275 pump in the northern Indian Ocean, *Progress in Oceanography*, 175, 24-39,
1276 <https://doi.org/10.1016/j.pocean.2019.03.001>, 2019a.
- 1277 Rixen, T., Guptha, M. V. S., and Ittekkot, V.: Sedimentation, in: Report of the Indian Ocean
1278 Synthesis Group on the Arabian Sea Process Study, edited by: L. Watts, e. a., JGOFS
1279 International Project Office, Bergen, 65-73, 2002.
- 1280 Rixen, T., Baum, A., Gaye, B., and Nagel, B.: Seasonal and interannual variations in the
1281 nitrogen cycle in the Arabian Sea, *Biogeosciences*, 11, 5733-5747, 10.5194/bg-11-5733-2014,
1282 2014.
- 1283 Rixen, T., Gaye, B., Emeis, K. C., and Ramaswamy, V.: The ballast effect of lithogenic matter
1284 and its influences on the carbon fluxes in the Indian Ocean, *Biogeosciences*, 16, 485-503,
1285 10.5194/bg-16-485-2019, 2019b.
- 1286 Robinson, R. S., Kienast, M., Albuquerque, A. L. S., Altabet, M., Contreras, S., and al, e.: A
1287 review of nitrogen isotopic alteration in marine sediments, *Paleoceanography*, 27,
1288 10.1029/2012PA002321, 2012.
- 1289 Rontani, J.-F., Zabeti, N., and Wakeham, S. G.: Degradation of particulate organic matter in
1290 the equatorial Pacific Ocean: Biotic or abiotic?, *Limnology and Oceanography*, 56, 333-349,
1291 2011.
- 1292 Satterberg, J., Arnarson, T. S., Lessard, E. J., and Keil, R. G.: Sorption of organic matter from
1293 four phytoplankton species to montmorillonite, chlorite and kaolinite in seawater, *Marine
1294 Chemistry*, 81, 11-18, 10.1016/s0304-4203(02)00136-6, 2003.
- 1295 Sheridan, C. C., Lee, C., Wakeham, S. G., and Bishop, J. K. B.: Suspended particle organic
1296 composition and cycling in surface and midwaters of the equatorial Pacific Ocean, *Deep-Sea
1297 Research I*, 49, 1983-2008, 2002.
- 1298 Silver, M. W., Coale, S. L., Pilskaln, C. H., and Steinberg, D. R.: Giant aggregates: Importance
1299 as microbial centers and agents of material flux in the mesopelagic zone, *Limnology and
1300 Oceanography*, 43, 498-507, 1998.
- 1301 Smith, D. C., Simon, M., Alldredge, A. L., and Azam, F.: Intense hydrolytic enzyme activity
1302 on marine aggregates and implications for rapid particle dissolution, *Nature*, 359, 139-142,
1303 1992.
- 1304 Suess, E.: Particulate organic carbon flux in the oceans - surface productivity and oxygen
1305 utilization, *Nature*, 288, 260-263, 1980.
- 1306 Sun, Y., Zulla, M., Joachimski, M., Bond, D., Wignall, P., Zhang, Z., and Zhang, M.:
1307 Ammonium ocean following the end-Permian mass extinction, *Earth Planet. Sci. Lett.*, 518,
1308 211-222, 10.1016/j.epsl.2019.04.036, 2019.
- 1309 Suthhof, A., Ittekkot, V., and Gaye-Haake, B.: Millennial-scale oscillation of denitrification
1310 intensity in the Arabian Sea during the late Quaternary and its potential influence on
1311 atmospheric N₂O and global climate, *Global Biogeochemical Cycles*, 15, 637-650, 2001.
- 1312 Suthhof, A., Jennerjahn, T. C., Schäfer, P., and Ittekkot, V.: Nature of organic matter in surface
1313 sediments from the Pakistan continental margin and the deep Arabian Sea: amino acids, *Deep-
1314 Sea Research II*, 47, 329-351, 2000.

- 1315 Taylor, G. T.: Microbial degradation of sorbed and dissolved protein in seawater, *Limnology*
1316 and *Oceanography*, 40, 875-885, 1995.
- 1317 Tesdal, J. E., Galbraith, E. D., and Kienast, M.: Nitrogen isotopes in bulk marine sediment:
1318 linking seafloor observations with subseafloor records, *Biogeosciences*, 10, 101-118,
1319 10.5194/bg-10-101-2013, 2013.
- 1320 Turnewitsch, R., Lahajnar, N., Haeckel, M., and Christiansen, B.: An abyssal hill fractionates
1321 organic and inorganic matter in deep-sea surface sediments, *Geophysical Research Letters*, 42,
1322 7663-7672, 10.1002/2015gl065658, 2015.
- 1323 Turnewitsch, R., Dale, A., Lahajnar, N., Lampitt, R. S., and Sakamoto, K.: Can neap-spring
1324 tidal cycles modulate biogeochemical fluxes in the abyssal near-seafloor water column?,
1325 *Progress in Oceanography*, 154, 1-24, 10.1016/j.pocean.2017.04.006, 2017.
- 1326 Unger, D., Gaye-Haake, B., Gebhardt, A. C., and Ittekkot, V.: Biogeochemistry of suspended
1327 and sedimentary material from the Ob and Yenisei rivers and the adjacent Kara Sea: Amino
1328 acids and amino sugars, *Cont. Shelf Res.*, 25, 437-460, 2005.
- 1329 Wakeham, S. G. and Canuel, E. A.: Organic geochemistry of particulate matter in the eastern
1330 tropical North Pacific; Implications for particle dynamics, *J. Mar. Res.*, 46, 183-213, 1988.
- 1331 Wakeham, S. G. and Lee, C.: Organic geochemistry of particulate matter in the ocean - the role
1332 of particles in oceanic sedimentary cycles *Org. Geochem.*, 14, 83-96, 10.1016/0146-
1333 6380(89)90022-3, 1989.
- 1334 Wakeham, S. G. and Lee, C.: Production, Transport, and Alteration of Particulate Organic
1335 Matter in the Marine Water Column, in: *Organic Geochemistry: Principles and Applications*,
1336 edited by: Engel, M. H., and Macko, S. A., Springer US, Boston, MA, 145-169, 10.1007/978-
1337 1-4615-2890-6_6, 1993.
- 1338 Wakeham, S. G., Lee, C., Farrington, J. W., and Gagosian, R. B.: Biogeochemistry of particulate
1339 organic matter in the oceans: results from sediment trap experiments, *Deep-Sea Research*, 31,
1340 509-528, 1984.
- 1341 Walla, M. D., Lau, P. Y., Morgan, S. L., Fox, A., and Brown, A.: Capillary gas
1342 chromatography-mass spectrometry of carbohydrate components of legionellae and other
1343 bacteria, *Journal of Chromatography A*, 288, 399-413, [https://doi.org/10.1016/S0021-
1344 9673\(01\)93716-1](https://doi.org/10.1016/S0021-9673(01)93716-1), 1984.
- 1345 Waples, D. W. and Sloan, J. R.: Carbon and nitrogen diagenesis in deep-sea sediments
1346 *Geochimica et Cosmochimica Acta*, 44, 1463-1470, 10.1016/0016-7037(80)90111-8, 1980.
- 1347 Whelan, J. K. and Emeis, K.-C.: Preservation of amino acids and carbohydrates in marine
1348 sediments, in: *Organic Matter: Productivity, Accumulation, and Preservation in Recent and
1349 Ancient Sediments*, edited by: Whelan, J. K., and Farrington, J., Columbia University Press,
1350 Palisades, N. Y., 176-200, 1992.
- 1351 Wilson, J. D., Barker, S., and Ridgwell, A.: Assessment of the spatial variability in particulate
1352 organic matter and mineral sinking fluxes in the ocean interior: Implications for the ballast
1353 hypothesis, *Global Biogeochemical Cycles*, 26, 15, 10.1029/2012gb004398, 2012.
- 1354 Xue, J. H., Lee, C., Wakeham, S. G., and Armstrong, R. A.: Using principal components
1355 analysis (PCA) with cluster analysis to study the organic geochemistry of sinking particles in
1356 the ocean, *Org. Geochem.*, 42, 356-367, 10.1016/j.orggeochem.2011.01.012, 2011.
- 1357 Yamaguchi, Y. T. and McCarthy, M. D.: Sources and transformation of dissolved and
1358 particulate organic nitrogen in the North Pacific Subtropical Gyre indicated by compound-

- 1359 specific $\delta^{15}\text{N}$ analysis of amino acids, *Geochimica et Cosmochimica Acta*, 220, 329-347,
1360 <https://doi.org/10.1016/j.gca.2017.07.036>, 2018.
- 1361 Yang, J.-Y. T., Kao, S.-J., Dai, M., Yan, X., and Lin, H.-L.: Examining N cycling in the northern
1362 South China Sea from N isotopic signals in nitrate and particulate phases, *Journal of*
1363 *Geophysical Research: Biogeosciences*, 122, 2118-2136, 10.1002/2016JG003618, 2017.
- 1364 Zhang, P. Y., Yang, G. P., Chen, Y., Leng, W. S., and Ji, C. X.: Temporal and spatial variations
1365 of particulate and dissolved amino acids in the East China Sea, *Marine Chemistry*, 186, 133-
1366 144, 10.1016/j.marchem.2016.09.004, 2016.
- 1367