



A robust initialization method for accurate soil organic carbon simulations

Eva Kanari^{1,2}, Lauric Cécillon^{1,3}, François Baudin², Hugues Clivot⁴, Fabien Ferchaud⁵, Sabine Houot⁶, Florent Levavasseur⁶, Bruno Mary⁵, Laure Soucémarianadin⁷, Claire Chenu⁶, Pierre Barré¹

5 ¹Laboratoire de Géologie, École normale supérieure, CNRS, Université PSL, IPSL, Paris, France

²ISTeP, UMR 7193 Sorbonne Université, CNRS, Paris, France

³Normandie Univ, UNIROUEN, INRAE, ECODIV, Rouen, France

⁴Université de Reims Champagne Ardenne, INRAE, FARE, UMR A 614, 51097 Reims, France

10 ⁵BioEcoAgro Joint Research Unit, INRAE, Université de Liège, Université de Lille, Université Picardie Jules Verne, F-02000, Barenton-Bugny, France

⁶UMR ECOSYS, INRAE, AgroParisTech, Université Paris-Saclay, Thiverval-Grignon, France

⁷ACTA - les instituts techniques agricoles, Paris, France

15 *Correspondence to:* Eva Kanari (kanari@geologie.ens.fr), Lauric Cécillon (lauric.cecillon@inrae.fr) and Pierre Barré (barre@biotite.ens.fr)

Abstract. Changes in soil organic carbon (SOC) stocks are a major source of uncertainty for the evolution of atmospheric CO₂ concentration during the 21st century. They are usually simulated by models dividing SOC into conceptual pools with contrasted turnover times. The lack of reliable methods to initialize these models, by correctly distributing soil carbon amongst their kinetic pools, strongly limits the accuracy of their simulations. Here, we demonstrate that PARTY_{SOC}, a machine-learning model based on Rock-Eval® thermal analysis optimally partitions the active and stable SOC pools of AMG, a simple and well validated SOC dynamics model, accounting for effects of soil management history. Furthermore, we found that initializing the SOC pool sizes of AMG using machine-learning strongly improves its accuracy when reproducing the observed SOC dynamics in nine independent French long-term agricultural experiments. Our results indicate that multi-compartmental models of SOC dynamics combined with a robust initialization can simulate observed SOC stock changes with excellent precision. We recommend exploring their potential before a new generation of models of greater complexity becomes operational. The approach proposed here can be easily implemented on soil monitoring networks, paving the way towards precise predictions of SOC stock changes over the next decades.



1 Introduction

30 Soil organic carbon (SOC) plays an important role in sustaining soil functions and associated soil ecosystem services worldwide (IPCC, 2019). It is the largest terrestrial organic carbon reservoir, with the upper two meters of soil storing 2400 Pg C, three times more carbon than the atmosphere (Jobbagy and Jackson, 2000). A mere 4 per 1000 annual decrease in SOC stocks (*ca.* 10 Gt C^{year}⁻¹) may double the global annual anthropogenic CO₂ emissions, while an equivalent increase may compensate them (Balesdent and Arrouays, 1999). This is the concept behind the 4 per 1000 initiative (Rumpel et al., 2018)
35 that aims at increasing SOC stocks to fight global warming while ensuring food security, two Sustainable Development Goals of the United Nations (UN General Assembly, 2015). This initiative and other political headway have placed the question of managing SOC stocks and assessing the global SOC sequestration potential at the top of political and scientific agendas (Vermeulen et al., 2019; FAO, 2019; Amelung et al., 2020). Despite this particular attention, the prediction of SOC stock changes remains very uncertain, which makes soils a major source of uncertainty for the evolution of atmospheric CO₂
40 concentration (Todd-Brown et al., 2014; He et al., 2016; Shi et al., 2018).

Models of SOC dynamics can predict future SOC stock evolution by simulating carbon transfer into the soil mostly through plant organic matter inputs, and microbial SOC mineralization resulting in a CO₂ flux from the soil to the atmosphere. They can have structures of various complexities reflecting our mechanistic understanding of SOC dynamics (Shi et al., 2018; Luo et al., 2016). However, most models dedicated to prediction, including those used in Earth System Models, have a simple structure dividing SOC into conceptual pools with contrasted turnover times (Manzoni and Porporato, 2009; He et al., 2016; Todd-Brown et al., 2014). These multi-compartmental models of SOC dynamics are the best option we currently have to foster science-based SOC preservation and sequestration actions, given the strong uncertainty of more complex models (Cécillon, 2021a; Dangal et al., 2021; Lee et al., 2020; Shi et al., 2018; Crowther et al., 2019). Predictions of SOC stocks
50 evolution provided by such simple models are very sensitive to the initial distribution of SOC amongst the different kinetic pools (Luo et al., 2016; Smith and Falloon, 2000; Clivot et al., 2019). This makes the question of SOC kinetic pool partitioning a priority for improving the accuracy of multi-compartmental SOC dynamics models (Luo et al., 2016; Taghizadeh-Toosi et al., 2020).

55 The most commonly used method to initialize the size of SOC kinetic pools is to run spin-up simulations until a steady-state equilibrium for SOC is reached, eventually matching the initial SOC stock measurement (Wutzler and Reichstein, 2007; Taghizadeh-Toosi et al., 2020). However, this method has two well-known limitations. First, climatic, SOC input, and land-use or land-cover data extending over long time periods required by this approach are highly uncertain. Second, assuming steady-state equilibrium for SOC at the onset of model simulations is often unrealistic. This is due to the history of the simulated sites that often includes disturbances (*e.g.*, fire), as well as previous changes in climate, land-use and soil management that prevent SOC pools with slow turnover times from being at equilibrium (Wutzler and Reichstein, 2007;
60



Herbst et al., 2018; Oberholzer et al., 2014; Poeplau et al., 2011; Clivot et al., 2019). Alternative initialization procedures are needed to address these issues (Wutzler and Reichstein, 2007; Bruun and Jensen, 2002; Taghizadeh-Toosi et al., 2020).

65 In some models of SOC dynamics, like the AMG model (Clivot et al., 2019), a default initial SOC pool size distribution is prescribed according to basic information on land-use history (*i.e.* long-term cropland *vs.* long-term grassland) (Clivot et al., 2019). This approach does not take into account the effect of recent changes in land-use or historical soil management practices on SOC pool distribution. To better reflect the effect of the frequent state of non-equilibrium of SOC on its partitioning into conceptual kinetic pools, another approach has been proposed, relating results from SOC fractionation methods with SOC kinetic pool sizes (*e.g.*, Zimmermann et al. (2007a) or Skjemstad et al. (2004) for the RothC model; Dangal et al. (2021) for the DAYCENT model). However, this approach also suffers from important drawbacks. First, SOC fractionation procedures are tedious and cannot be implemented on large-scale studies, though this problem may be solved by using soil infrared spectroscopy or environmental variables and machine-learning (Zimmermann et al., 2007b; Viscarra Rossel et al., 2019; Sanderman et al., 2021; Cotrufo et al., 2019; Lugato et al., 2021; Baldock et al., 2013; Barthès et al., 2008; Lee et al., 2020; Dangal et al., 2021). Second, their reproducibility is questionable (Poeplau et al., 2013, 2018), and third, their use for initializing model SOC pool sizes has never been properly validated. A proper validation would require showing that (1) the size of measured SOC fractions matches the one of model kinetic pools, and that (2) simulations of SOC dynamics are more accurate using this initialization strategy, compared to default simulations (on independent validation sites while other model parameters remain constant). Reasonably good correspondence between measured or soil-spectroscopy-estimated SOC fractions and modelled SOC conceptual pools has been reported in a number of studies, though with some notable discrepancies (Leifeld et al., 2009b; Herbst et al., 2018; Zimmermann et al., 2007a; Dangal et al., 2021). Conversely, the studies that attempted to initialize model SOC pool sizes using a SOC fractionation scheme generally reported no improvement in the accuracy of simulations of SOC dynamics compared to a default or a spin-up initialization approach (Leifeld et al., 2009a; Nemo et al., 2016; Cagnarini et al., 2019). Only two studies showed that an initialization based on a SOC fractionation scheme yielded more accurate simulations of observed SOC dynamics, but at the cost of modifying the decomposition rate of SOC kinetic pools (Skjemstad et al., 2004; Luo et al., 2014).

An alternative approach using Rock-Eval® thermal analysis has recently been proposed — under the name PARTY_{SOC} model — to estimate SOC kinetic pool size (Cécillon et al., 2018, 2021). PARTY_{SOC} is a machine-learning model trained on Rock-Eval® data of soil samples from long-term experiments (LTEs) where the size of the centennially stable SOC fraction can be estimated (*e.g.*, sites including a bare fallow treatment). PARTY_{SOC} incorporates recent key elements of the new understanding of SOC dynamics (Dignac et al., 2017), showing that the centennially stable SOC fraction has specific chemical and energetical characteristics that are measurable quickly (*ca.* 1 h per sample) and at a reasonable cost (less than USD 60) using Rock-Eval®: it is thermally stable (*i.e.* high activation energy) and it is depleted in hydrogen (Barré et al., 2016; Hemingway et al., 2019; Gregorich et al., 2015; Poeplau et al., 2019; Chassé et al., 2021).



In this study, we tested if the PARTY_{SOC} machine-learning model, built on a totally independent data set from North-western Europe, could be used to initialize the distribution of SOC pools of the simple AMG model (Clivot et al., 2019) and improve the accuracy of its simulations. The default version of AMG is currently the most accurate model for reproducing the observed SOC stock dynamics in diverse French agricultural LTEs at the pluri-decadal scale (Martin et al., 2019). The efficient use of this model at sites covering an important pedological and climatic variability (including oceanic, continental, and tropical climate) provides further support to its robustness (Levvasseur et al., 2020; Farina et al., 2021; Saffih-Hdadi and Mary, 2008). In this model, SOC is simply divided into two pools, the “stable C (C_S)” that is considered as inert at the time scale of the simulation and the “active C (C_A)” that has a mean turnover time of a few decades. A recent study (Clivot et al., 2019) determined that the optimal initial proportion of C_S (C_S/C_0) can deviate from the model’s default value (0.65 in croplands), so that a more precise initialization of the C_S/C_0 proportion would significantly improve AMG simulations of SOC dynamics. Here, we hypothesized that the SOC pool partitioning as determined by the PARTY_{SOC} machine-learning (Cécillon et al., 2021, 2018) would be close to the mathematically optimal one for the AMG model, therefore, improving the accuracy of its SOC dynamics simulations compared to default initialization. We tested our hypothesis on 32 treatments from nine independent French agricultural LTEs (experiment duration from 12 to 41 years with a median of 21 years) in which the AMG optimal SOC pool partitioning could be determined by *ex-post* optimization and for which topsoil samples collected at the onset of the experiment were available (Table 1). These LTEs were croplands established in different pedoclimates that have experienced contrasted soil management practices and land-use histories. All available initial topsoil samples were analysed with Rock-Eval® and the results were used in the European version of the PARTY_{SOC} model, PARTY_{SOC}v2.0_{EU} (Cécillon et al., 2021), to compute the centennially stable SOC proportion of each topsoil sample.

2 Materials and methods

2.1 Experimental sites

This work was conducted on nine French agricultural LTEs (Supplementary Material Fig. 1). Seven LTEs including 29 treatments were selected from the dataset presented in Clivot et al. (2019), from sites with availability of initial topsoil samples. Two additional LTEs (Colmar and Feucherolles) including a total of three treatments were obtained from the dataset published in Levvasseur et al. (2020), selecting control treatments without organic amendments and with available initial topsoil samples. Basic site and topsoil characteristics are reported in Table 1 and Supplementary Material Table 1. Information necessary to run AMG simulations on a total of 32 treatments (initial soil physico-chemical properties, detailed information on management practices and observed climatic data during all experiments) were obtained from (Clivot et al., 2019) for the 29 treatments of the seven sites and from (Levvasseur et al., 2020) for the three treatments of the sites of Colmar and Feucherolles.



130

Table 1: Main information on the nine French agricultural long-term experiments used in this study. All sites had been croplands for at least 20 years before the onset of all experiments. Additional site information including climatic variability amongst sites and long-term history of land cover is provided in Supplementary Material Table 1.

	Auzeville	Boigneville	Colmar	Doazit	Feucherolles	Grignon-Folleville	Kerbernez	Mant	Tartas
Soil type (WRB 2014)	Luvisol	Haplic Luvisol	Calcic Cambisol	Luvisol	Gleyic Luvisol	Luvisol	Cambisol	Dystric Luvisol	Luvic Arenosol
LTE onset	1968	1970	2000	1967	1998	1958	1978	1975	1972
Simulated period	1975–2010	1970–2011	2000–2018	1977–1989	1998–2019	1989–2008	1978–2005	1975–1992	1976–1997
Number of treatments	4	12	1	2	2	2	5	2	2
Sampling dates (number of samples)	1975 (4), 2010 (8)	1970 (29), 1998 (10), 2017 (32)	2000 (4), 2018 (6)	1977 (4), 1989 (4)	1998 (8), 2013 (8), 2018(8)	1989 (8), 2008 (8)	1978 (6), 1991 (6), 2005 (12)	1975 (4), 1992 (4)	1976 (4), 1997 (4)
Crop rotation	annual crop rotation	annual crop rotation	annual crop rotation	maize monoculture	annual crop rotation	annual crop rotation	silage maize monoculture (KERB_C incl. raygrass)	maize monoculture	maize monoculture
Considered depth	30	29	28	25	28	30	25	28	28
Initial SOC stock (tC·ha⁻¹)	34.68	42.40	45.20	26.35	43.80	55.85	81.98	38.75	45.25
Reference	(Colomb et al., 2007)	(Dimassi et al., 2014)	(Obriot, 2016)	(Lubet et al., 1993)	(Noirot-Cosson et al., 2016)	(Barré et al., 2008)	(Vertès et al., 2007)	(Messiga et al., 2010)	(Morel et al., 2014)

2.2 Archive soil samples from experimental sites

Our final soil sample set included 181 topsoil samples. At each site the soil was sampled to include the whole ploughing depth (Table 1). At all sites, except Boigneville where the soil was sampled in five sublayers, the ploughing layer was sampled as one homogeneous layer. Of the final samples, 71 were from LTEs starting dates, 24 from intermediate dates and 86 from final dates. All samples were air-dried or dried at 40 °C, sieved to < 2 mm and finely ground to < 250 µm using a ball mill (Retsch, Germany).

135

2.3 Rock-Eval® analysis of archive soil samples

All soil samples were analysed using a Rock-Eval 6® Turbo apparatus (Vinci Technologies). The samples were first pyrolyzed in an inert N₂ atmosphere, then oxidized under ambient air (O₂). The heating routine applied during pyrolysis was as described in (Disnar et al., 2003), starting with a three-minute isotherm at 200 °C followed by a heating ramp of 30 °C·min⁻¹ up to 650 °C. For the oxidation step, a one-minute isotherm was kept at 300 °C and was directly followed by a

140



heating ramp of $20\text{ }^{\circ}\text{C}\cdot\text{min}^{-1}$ until $850\text{ }^{\circ}\text{C}$ was reached, followed by a five-minute isotherm at $850\text{ }^{\circ}\text{C}$ (Baudin et al., 2015)
145 adapted from (Behar et al., 2001)).

Based on five generated Rock-Eval® thermograms, 18 parameters were calculated for each sample, and were then used as predictors by the random forests model. These include Total Organic Carbon (TOC; in $\text{gC}\cdot\text{kg}\text{ soil}^{-1}$) — the amount of organic C released during the analysis as a proportion of sample weight; Pyrolyzed organic Carbon (PC; in $\text{gC}\cdot\text{kg}\text{ soil}^{-1}$) — the sum of C released as CH, CO and CO₂ during pyrolysis step; the ratio of PC to TOC (PC/TOC); S2 peak area ($\text{gC}\cdot\text{kg}\text{ soil}^{-1}$) — the hydrocarbon gas released within the range of the pyrolysis temperature ramp; the ratio of S2 to PC (S2/PC); PseudoS1 peak area ($\text{gC}\cdot\text{kg}\text{ soil}^{-1}$) — the sum of C released as CH, CO and CO₂ during the first 200 seconds of pyrolysis (after Khedim et al., 2021); Hydrogen Index (HI; in $\text{mgHC}\cdot\text{gTOC}^{-1}$) — the amount of hydrocarbons released as a ratio of TOC; the ratio of HI to Oxygen Index (HI/OIRE6) — where OIRE6 is calculated as the amount of oxygen released as CO and CO₂ gases normalized to TOC. Finally, various temperature parameters (T70CH_PYR, T90CH_PYR, T30CO₂_PYR, T50CO₂_PYR, T70CO₂_PYR, T90CO₂_PYR, T70CO₂_OX, T50CO₂_OX, T70CO₂_OX, T90CO₂_OX; in $^{\circ}\text{C}$) are
155 included in the predictors set. They describe evolution steps, namely at which temperature a specific amount (e.g. 30, 50, 70 or 90%) of a given gas was released according to each thermogram (Cécillon et al., 2018). It is important to note that for the calculation of all of the above parameters, only the part of each thermogram corresponding to organic carbon was taken into account. For this purpose, upper temperature integration limits were set at $560\text{ }^{\circ}\text{C}$ for the CO and CO₂ pyrolysis
160 thermograms, and at $611\text{ }^{\circ}\text{C}$ for the CO₂ oxidation thermograms (Cécillon et al., 2018). R scripts used for computing Rock-Eval® parameters are available permanently on the Zenodo platform (Cécillon, 2021b).

2.4 The PARTY_{SOC} machine-learning model

The most up-to-date European version of this model, calibrated on soils from North-western Europe, used in this study, is
165 described in detail in (Cécillon et al., 2021). This model uses the 18 above-mentioned Rock-Eval® thermal analysis parameters as predictors and estimates the centennially stable SOC proportion in a topsoil sample. The model consists of a trained non-parametric machine learning algorithm, using the random forests approach to estimate centennially stable SOC proportions in unknown topsoils from centred and scaled Rock-Eval® parameters. In this study the obtained centennially stable SOC proportion of each topsoil sample, was converted to centennially stable SOC content by multiplying the
170 predicted proportion by the total SOC content. The PARTY_{SOC}V2.0_{EU} model, permanently available on Zenodo (Cécillon, 2021b), was used without any adaptation.



2.5 The AMG model of soil organic carbon dynamics

175 The AMG model (Andriulo et al., 1999) was developed based on the two-compartment SOC model proposed by Hénin and Dupuis (Henin and Dupuis, 1945). It is characterized by a simple structure consisting of three carbon pools: fresh organic matter, and two SOC fractions, an active and a stable pool (Supplementary Material Fig. 2). The model allows transfer of carbon from the fresh organic matter pool either to the atmosphere through microbial mineralization or into the active pool. Organic carbon from the active pool is also subject to mineralization, forming a second direct flux of CO₂ from the soil into
180 the atmosphere. SOM decomposition follows first order kinetics with a rate defined by the coefficient of mineralization k (year⁻¹), controlled by climatic conditions and soil characteristics. The h coefficient controls the yield of crop residues transformation into active carbon and depends on the type of fresh organic matter. No carbon exchange with the stable SOC pool is possible since it is considered inert and remains unchanged over the simulation period (here from 12 to 41 years; see Table 1). Considering the stable SOC pool as mathematically inert at this time scale is in line with consistent observations of
185 a significant pluri-decadal persistent SOC fraction in long-term bare fallows and C3-C4 vegetation change chronosequences (Barré et al., 2010; Balesdent et al., 2018).

The AMG model can be mathematically described by two simple equations (Clivot et al., 2019):

$$QC = QC_S + QC_A, \quad (1)$$

190 $\frac{dQC_A}{dt} = \sum_i m_i h_i - k \cdot QC_A, \quad (2)$

where QC is the total SOC stock (t·ha⁻¹), QC_S is the stable SOC stock (t·ha⁻¹) defined as a fraction of initial SOC stock QC_0 (s. Sect. 2.6) constant for a specific treatment, QC_A is the active SOC stock (t·ha⁻¹), t is the time in years, m_i is the annual C input from organic residue i (t·ha⁻¹·yr⁻¹), h is a coefficient representing the fraction of C inputs which is incorporated in
195 SOM after 1 year related to the type of organic residue, and k is the mineralization rate constant associated with the active C pool (yr⁻¹).

The AMG parameters (h and k) have been determined by experimental results (Clivot et al., 2019). This approach differs from most multi-compartmental SOC dynamics models for which decay rates of slower pools were calibrated indirectly,
200 assuming an equilibrium state for SOC (Wutzler and Reichstein, 2007). The simple structure of the AMG model and the experimental determination of its decomposition rates make it less susceptible to the problem of equifinality compared to other multi-compartmental models of SOC dynamics (Clivot et al., 2019; Luo et al., 2016). Furthermore, AMG has been validated with $\delta^{13}C$ tracer data of long term alternative sequences of C4 and C3 crops (Mary et al., 2020).

205 The version of AMG used in this study was AMGv2, described in detail in (Clivot et al., 2019). Input data necessary to run simulations of SOC stocks with AMG include crop type, annual crop yields and information regarding management of crop



residues. These are used to compute annual aboveground and belowground C inputs from plants, here according to the method proposed by (Bolinder et al., 2007) and adapted by (Clivot et al., 2019). The coefficient of mineralization k (year^{-1}) is calculated according to soil characteristics (clay and carbonate contents, pH and C:N ratio) and climatic conditions (mean annual temperature, precipitation and potential evapotranspiration) (Clivot et al., 2019).

2.6 Soil organic carbon pool partitioning in the AMG model

2.6.1 Default C_s/C_0 initialization

Two default values can be used for initialization of SOC pool distribution in AMG, depending on land-use history before the onset of simulations. The initial proportion of the stable SOC pool equals 0.65 for sites with a long-term arable land-use history. Former long-term grassland sites are expected to have lower C_s/C_0 and the value of 0.40 was used in previous studies (Saffih-Hdadi and Mary, 2008; Clivot et al., 2019). Since all sites used in this study had been under arable land for at least 20 years before the onset of the experiment, a default value of 0.65 was used.

2.6.2 Rock-Eval®-based initialization of C_s/C_0

After obtaining predictions of the centennially stable SOC proportion for individual topsoil samples using the PARTY_{SOC}v2.0_{EU} statistical model, these values were averaged per site. For this step, only initial topsoil samples collected at the LTE's onset were used. The site mean was then used to initialize simulations of SOC stocks for the various treatments of every site (the site standard deviation is reported on Fig. 1 and in Supplementary Material Table 2). Supported by the evident common land-use history shared by the multiple treatments of each site before the onset of simulations and as the SOC stocks and centennially stable SOC content were very homogeneous amongst each site, we also performed simulations of treatments for which soil samples from the onset of the LTE were not available. In these cases, we considered that the centennially stable SOC proportion of the treatment was equal to the mean proportion of the respective site (Supplementary Material Table 1 and 2).

2.6.3 *Ex-post* optimization of C_s/C_0

Following a least squares optimization approach, the best fit on observed SOC time series was obtained and the required initial SOC pool partitioning was estimated accordingly for each site (Clivot et al., 2019). In sites where long-term monitoring data were available, the model was adapted to simultaneously match the observed evolution of C, C3 and C4 stocks (determined using ^{13}C stable isotope measurements; Clivot et al., 2019) for a given treatment.



235 2.7 Statistics

The fit between PARTY_{SOC} predictions of centennially stable SOC proportion and *ex-post* optimized C_s/C₀ was assessed by a linear regression model. The same approach was applied for the evaluation of the agreement between centennially stable SOC content and AMG *ex-post* optimized C_s content of initial samples. The evaluation of the performance of the AMG model, for the different SOC pool partitioning initialization methods, was also based on simple linear regressions between
 240 simulated and observed SOC stock values. Statistical terms used to express the strength and the statistical significance of the relationships were the coefficient of determination (R^2), and the associated probability value (*p-value*). Prediction bias and model error were expressed as mean difference (*BIAS*), and relative mean square error (*RMSE*). The relative root mean square error (*RRMSE*) and the normalized root mean square error (*NRMSE*) were used to compare the error of different data sets (with a different range of predictions) (Smith et al., 1996; Wallach, 2006; Otto et al., 2018).

$$245 \quad R^2 = \left(\frac{\sum_{i=1}^n (O_i - \bar{O}) \cdot (S_i - \bar{S})}{\sqrt{\sum_{i=1}^n (O_i - \bar{O})^2} \cdot \sqrt{\sum_{i=1}^n (S_i - \bar{S})^2}} \right)^2, \quad (3)$$

$$BIAS = \frac{1}{n} \sum_{i=1}^n (S_i - O_i), \quad (4)$$

$$RMSE = \sqrt{\frac{1}{n} \sum_{i=1}^n (S_i - O_i)^2}, \quad (5)$$

$$RRMSE = \frac{RMSE}{\bar{O}}, \quad (6)$$

$$NRMSE = \frac{RMSE}{O_{max} - O_{min}}, \quad (7)$$

250

where: O and S are the observed and simulated values, n is the number of observations, \bar{O} and \bar{S} are the means of observations and simulations, respectively, and O_{max} and O_{min} are the maximum and the minimum value observed.

The observed and simulated total SOC stock change dQC was calculated as follows for each treatment:

$$255 \quad dQC_{obs} = QC_{obs,t_2} - QC_{obs,t_1}, \quad (8)$$

$$dQC_{sim} = QC_{sim,t_2} - QC_{obs,t_1}, \quad (9)$$

where QC_{obs} is the observed SOC stock at time t , QC_{sim} is the SOC stock at time t simulated with AMG, t_1 indicates the start and t_2 the end of simulation period.


260 All data processing and statistical analyses were performed within the “R” programming environment (version 3.4.2) (R Core Team, 2017). For plotting, packages *ggpmisc* (Aphalo, 2016), *reshape2* (Wickham, 2007) and *ggplot2* (Wickham, 2016) were used.



3 Results

3.1 Accurate soil organic carbon pool partitioning

265 Centennially stable SOC proportion values were predicted by the PARTY_{SOC} machine learning model (Cécillon et al., 2021)
using Rock-Eval® data measured on initial topsoil samples. The mean value for each independent site was plotted against
the C_S/C₀ proportion as determined by AMG *ex-post* optimization (Fig. 1). The initial centennially stable SOC proportion
values predicted with PARTY_{SOC} ranged from 0.44 to 0.74, with a mean value of 0.59, whereas optimal *ex-post* estimations
of AMG C_S/C₀ covered almost the same range, from 0.45 to 0.74, with a mean value of 0.61. The two approaches were
270 strongly correlated ($R^2 = 0.63$, significant at the $p < 0.05$ level), with a linear regression slope close to 1 ($a = 0.9$) and
intercept close to 0 ($b = 0.04$), showing an unbiased relationship between Rock-Eval® estimated centennially stable SOC
proportion and the AMG *ex-post* optimized C_S/C₀. Although a slight discrepancy was observed for higher C_S/C₀ values, the
results validate our hypothesis showing that centennially stable SOC proportion determined by Rock-Eval® thermal analysis
and the PARTY_{SOC} machine-learning model built on fully independent data provides a good estimate of the optimal C_S/C₀
275 proportion of the AMG model for unrelated French agricultural soils. When expressed as content (g C·kg soil⁻¹), the fit
between the Rock-Eval®-based predictions of the centennially stable SOC determined on initial topsoil samples and the *ex-*
post optimized C_S values was excellent ($R^2 = 0.95$; Supplementary Material Fig. 3; optimal C_S ranged from 4.37 to 12.75 g
C·kg soil⁻¹ across the nine sites). Furthermore, the method appears to be reliable since additional Rock-Eval® measurements
on topsoil samples from intermediate and final dates of the LTEs showed that the PARTY_{SOC} predictions of the centennially
280 stable SOC content remained remarkably constant during the experimental period at most sites (Supplementary Material Fig.
4).




300 Figure 1: Performance of the PARTY_{SOC} model to predict the centennially stable SOC proportion compared to the AMG *ex-post*
optimized stable SOC proportion. Points represent site-mean values based on initial topsoil samples from nine independent French
long-term experiments. Statistics refer to the linear regression between x and y values (blue solid line). Horizontal error bars show
the uncertainty associated with the optimal C_s/C_0 , calculated as the standard deviation of treatment-wise optimized C_s/C_0 value.
305 Vertical error bars represent the prediction error of the centennially stable SOC proportion values, calculated as the standard
deviation of the PARTY_{SOC} model predictions on initial topsoil samples.



3.2 More accurate soil organic carbon simulations

In a second step, we investigated if a Rock-Eval®-based initialization of the SOC pool partitioning could improve the accuracy of SOC stock simulations of the AMG model. To do so, we compared SOC stock simulations obtained with three different initializations. We first ran AMG using the default initialization method for the SOC pool partitioning ($C_S/C_0 = 0.65$ since all LTEs were under cropland for at least two decades before their onset; Table 1). Then, we ran AMG simulations using the Rock-Eval®-based initialization method by defining C_S/C_0 as the site-mean centennially stable SOC proportion determined by the PARTY_{SOC} model. Finally, we ran AMG using the *ex-post* optimization method to initialize the SOC pool partitioning for each site. For all three initialization procedures, the simulated SOC stock change between the initial and last sampling date for each treatment of each site was plotted against the measured SOC stock change (Fig. 2a–c). Observed SOC stock change ranged from +6 to $-24 \text{ t C}\cdot\text{ha}^{-1}$ for the 32 treatments. In spite of a rather good mean agreement (RMSE = $5.95 \text{ t C}\cdot\text{ha}^{-1}$), the AMG model initialized with the default procedure provided predictions of SOC stock change rather far from what was observed in two out of nine LTEs (Fig. 2a). Using the Rock-Eval®-based initialization method improved AMG simulations compared to the default method, bringing them much closer to the observed SOC stock changes (RMSE = $3.60 \text{ t C}\cdot\text{ha}^{-1}$; Fig. 2a, b). Rock-Eval®-based initialization of AMG resulted to unbiased simulations (BIAS = $0.06 \text{ t C}\cdot\text{ha}^{-1}$) and a strong decrease in the mean error of prediction. Unsurprisingly, AMG initialized using *ex-post* optimized C_S/C_0 proportions predicted SOC stock changes very close to the observed ones (RMSE = $2.12 \text{ t C}\cdot\text{ha}^{-1}$; Fig. 2c). AMG simulations from *ex-post* optimized and Rock-Eval®-based initializations were remarkably comparable (Fig. 2b, c). The SOC stock simulations produced with AMG for each independent treatment are presented in Supplementary Material Fig. 5.

325



330

Figure 2: Observed vs. simulated change in SOC stocks between the initial and final date of 32 treatments from nine French long-term experiments. The three panels show the performance of the AMG model for three different initialization approaches. Initial SOC kinetic pool sizes were defined using a, the default value for cropland ($C_S/C_0 = 0.65$), b, the centennially stable SOC proportion predicted by the PARTY_{soc} model and c, the AMG *ex-post* optimized C_S/C_0 proportion. Statistics refer to the linear regression between x and y values (blue solid line). Points represent the values for the 32 treatments for which AMG simulations were run.



335 It is noteworthy that the Rock-Eval®-based initialization improved the fit between observed and simulated SOC stock
change, compared to AMG default initialization, especially for treatments that experienced the greatest SOC stock loss (Fig.
2a, b). In treatments that experienced no SOC stock change or a slight increase in SOC stock, the Rock-Eval®-based
initialization did not improve the simulations but resulted in highly reliable predictions, similarly to AMG default or
optimized initialization methods (Fig. 2a–c). This is likely explained by the history of land cover and soil management
340 practices of the different sites. Sites presenting treatments with no change or a slight increase in SOC stocks were
predominantly sites with a long cropland history (*e.g.* site of Boigneville; Supplementary Material Table 1), for which the
default AMG C_S/C_0 value of 0.65 is nearly optimal. Conversely, the two sites, Kerbernez and Tartas, where the *ex-post*
optimized C_S/C_0 values were far below the default value (Fig. 1) have a more complex history of land use and soil
management practices. The site of Kerbernez is a former grassland (during the first half of the 20th century; Supplementary
345 Material Table 1) that was converted into a cropland two decades before the implementation of its arable LTE, in 1958. The
site of Tartas was cultivated for a longer time before the LTE onset, however it was turned to grassland for a period in the
19th century (Supplementary Material Table 1) and received applications of poultry manure for several years before the LTE
began. In these two sites, characterized by an optimal AMG C_S/C_0 much lower than the default value, the PARTY_{SOC}
machine-learning model based on Rock-Eval® predicted values very close to the optimal C_S/C_0 values (Fig. 1).

350

4 Discussion

Our study demonstrates that the PARTY_{SOC} method based on Rock-Eval® thermal analysis (Cécillon et al., 2018, 2021) can
estimate the initial SOC pool partitioning of the AMG model of SOC dynamics while improving its accuracy in a series of
diverse and independent French LTEs. Contrary to previous studies (Skjemstad et al., 2004; Luo et al., 2014), no
355 modifications of the decomposition rate of SOC kinetic pools were necessary to improve model predictions. The PARTY_{SOC}
initialization method never severely affected the model simulations while it strongly improved them at sites where SOC
stocks were far from an equilibrium state due to historical changes in soil management or land use. Areas with past changes
in land use and soil management represent a large yet poorly known part of arable land in France and Europe (Fuchs et al.,
2015; Erb et al., 2017) where SOC stocks and slow-cycling SOC pools are far from equilibrium (Wutzler and Reichstein,
360 2007; Herbst et al., 2018; Clivot et al., 2019; Taghizadeh-Toosi et al., 2020). Therefore, by accounting for these legacy
effects of site history on SOC pool partitioning, the Rock-Eval®-based initialization of the AMG model should result in
more accurate simulations of SOC dynamics at a national or continental scale.

Our findings combined with results reported in recent ensemble modelling studies (Martin et al., 2019; Farina et al., 2021),
365 suggest that despite its simple structure and when properly initialized (*e.g.*, using the PARTY_{SOC} model) the AMG model is
unsurpassed for predicting observed SOC stock changes in French agricultural LTEs, and is amongst the best available



370 modelling frameworks of SOC dynamics in European arable land. Our results demonstrate that there is still potential to
increase the accuracy of simple multi-compartmental models of SOC dynamics, bringing their simulations very close to the
observed values of SOC stock changes. Developing other Rock-Eval®-based initialization methods specifically designed to
375 match the carbon pool design of other multi-compartmental SOC dynamics models such as RothC (Coleman et al., 1997) is a
promising research area. More generally, we recommend that the potential of multi-compartmental SOC dynamics models
be fully explored and exploited by soil biogeochemists before a new generation of models of increased complexity becomes
operational (Lehmann et al., 2020; Shi et al., 2018; Crowther et al., 2019; Cécillon, 2021a; Dangal et al., 2021; Lee et al.,
2020). The low prediction error of the AMG model when its SOC pool distribution is initialized with the Rock-Eval®-based
375 method even challenges the ability of more complex modelling approaches to achieve better performance, given the
uncertainty on observed values of SOC stock changes (Schrumpf et al., 2011).

The continental or worldwide implementation of the AMG model with the Rock-Eval®-based initialization of SOC pools
distribution will require additional work. First, the PARTY_{SOC} machine-learning model (Cécillon et al., 2018, 2021) will
380 have to be validated on a wider range of pedoclimates. This method initially built on LTEs coming from North-western
Europe (Cécillon et al., 2018), has now been successfully extended to new soil types and a new climate (tropical) (Cécillon
et al., 2021). The good agreement between optimal AMG C_S/C_0 values and PARTY_{SOC} predictions reported here suggests
that most agricultural LTEs with accurate AMG simulations could be used as reference sites for the PARTY_{SOC} model,
lifting an important technical limitation to its geographical expansion (Cécillon et al., 2021). Second, the improved accuracy
385 of model simulations using a Rock-Eval®-based initialization will also have to be demonstrated for a wider pedoclimatic
range (*i.e.* worldwide LTEs; such as those referenced by the International Soil Carbon Network (Nave et al., 2015)). Third,
Rock-Eval® data from the new application areas will be required. Rock-Eval® is a high-throughput technique that is well
adapted to the analysis of large soil sample sets provided by large-scale soil monitoring programs. We recommend
implementing Rock-Eval® measurements in national and continental soil monitoring networks.

390

5 Conclusions

Combining Rock-Eval® thermal analysis with the PARTY_{SOC} machine-learning model should be considered as an emerging
key approach with demonstrated ability to improve the accuracy of simulations of SOC dynamics, complementary to other
SOC cycling proxies (Bailey et al., 2018; Wiesmeier et al., 2019). The progressive large-scale delivery of these
395 complementary data related to SOC dynamics will strengthen model predictions of SOC stock changes at the national to
global scale, necessary for implementing efficient climate change mitigation policies (FAO, 2020).



Code availability

Code used to produce the results presented in this study, as well as the AMG model can be made available upon reasonable
400 request. The code of the Rock-Eval®-based PARTY_{SOCV2.0EU} random forests model is available on Zenodo at the permanent
link <https://doi.org/10.5281/zenodo.4446138>.

Data availability

Pedoclimatic and land management information used to run the AMG model are published in (Clivot et al., 2019) and can be
obtained from H.C. and F.F. upon request concerning seven sites, and from F.L. and S.H. concerning the remaining two sites
405 (Levvasseur et al., 2020). The Rock-Eval® data obtained and processed during this study are available upon request from
the corresponding authors.

Author contributions

E.K. performed the research and produced the first version of the manuscript. P.B. and L.C. designed the study with
contributions from C.C. and F.B., whereas B.M., F.F., H.C., F.L. and S.H. organized the search for archive soil samples,
410 provided data needed to run AMG simulations as well as the latest version of the AMG model, and helped with the
interpretation of the results. F.L. performed AMG simulations for the sites of Colmar and Feucherolles. F.B. conducted all
Rock-Eval® analyses. All authors participated in multiple scientific discussions during the preparation of this paper and
finally reviewed and provided valuable feedback during the writing process.

415 Competing interests

The authors declare no competing interests.

Acknowledgements

This work would not have been possible without the financial support of ADEME (French ecological transition agency,
under the Ministry of Ecological Transition; PhD grant for E.K, PROTERR project) and the ANR (French national research
420 agency; PhD grant for E.K., StoreSoilC project). The authors would also like to thank Florence Savignac for her help in
performing Rock-Eval® analyses, and all the people involved in maintaining and monitoring the field trials used in this
study over multiple decades. We would also like to thank everyone involved in the conception and creation of the AMG
consortium, the AIAL database and the SOERE PRO network for kindly providing access to their data.



References

- 425 Amelung, W., Bossio, D., de Vries, W., Kögel-Knabner, I., Lehmann, J., Amundson, R., Bol, R., Collins, C., Lal, R., Leifeld, J., Minasny, B., Pan, G., Paustian, K., Rumpel, C., Sanderman, J., van Groenigen, J. W., Mooney, S., van Wesemael, B., Wander, M., and Chabbi, A.: Towards a global-scale soil climate mitigation strategy, 11, 1–10, <https://doi.org/10.1038/s41467-020-18887-7>, 2020.
- Andriulo, A. E., Mary, B., and Guerif, J.: Modelling soil carbon dynamics with various cropping sequences on the rolling
430 pampas, 19, 365–377, <https://doi.org/10.1051/agro>, 1999.
- R Core Team (2017). R: A language and environment for statistical computing.
- Aphalo, P. J.: Learn R ...as you learnt your mother tongue., 2016.
- Bailey, V. L., Bond-Lamberty, B., DeAngelis, K., Grandy, A. S., Hawkes, C. V., Heckman, K., Lajtha, K., Phillips, R. P., Sulman, B. N., Todd-Brown, K. E. O., and Wallenstein, M. D.: Soil carbon cycling proxies: Understanding their critical role
435 in predicting climate change feedbacks, 24, 895–905, <https://doi.org/10.1111/gcb.13926>, 2018.
- Baldock, J. A., Hawke, B., Sanderman, J., and Macdonald, L. M.: Predicting contents of carbon and its component fractions in Australian soils from diffuse reflectance mid-infrared spectra, *Soil Res.*, 51, 577, <https://doi.org/10.1071/SR13077>, 2013.
- Balesdent, J. and Arrouays, D.: Usage des terres et stockage de carbone dans les sols du territoire français. Une estimation des flux nets pour la période 1900-1999, 85, 265–277, 1999.
- 440 Balesdent, J., Basile-Doelsch, I., Chadoeuf, J., Cornu, S., Derrien, D., Fekiacova, Z., and Hatté, C.: Atmosphere–soil carbon transfer as a function of soil depth, *Nature*, 559, 599–602, <https://doi.org/10.1038/s41586-018-0328-3>, 2018.
- Barré, P., Montagnier, C., Chenu, C., Abbadie, L., and Velde, B.: Clay minerals as a soil potassium reservoir: Observation and quantification through X-ray diffraction, 302, 213–220, <https://doi.org/10.1007/s11104-007-9471-6>, 2008.
- Barré, P., Eglin, T., Christensen, B. T., Ciais, P., Houot, S., Kätterer, T., van Oort, F., Peylin, P., Poulton, P. R.,
445 Romanenkov, V., and Chenu, C.: Quantifying and isolating stable soil organic carbon using long-term bare fallow experiments, *Biogeosciences*, 7, 3839–3850, <https://doi.org/10.5194/bg-7-3839-2010>, 2010.
- Barré, P., Plante, A. F., Cécillon, L., Lutfalla, S., Baudin, F., Bernard, S., Christensen, B. T., Eglin, T., Fernandez, J. M., Houot, S., Kätterer, T., Le Guillou, C., Macdonald, A., van Oort, F., and Chenu, C.: The energetic and chemical signatures of persistent soil organic matter, *Biogeochemistry*, 130, 1–12, <https://doi.org/10.1007/s10533-016-0246-0>, 2016.
- 450 Barthès, B. G., Brunet, D., Hien, E., Enjalric, F., Conche, S., Freschet, G. T., d’Annunzio, R., and Toucet-Louri, J.: Determining the distributions of soil carbon and nitrogen in particle size fractions using near-infrared reflectance spectrum of bulk soil samples, *Soil Biology and Biochemistry*, 40, 1533–1537, <https://doi.org/10.1016/j.soilbio.2007.12.023>, 2008.
- Baudin, F., Disnar, J. R., Aboussou, A., and Savignac, F.: Guidelines for Rock-Eval analysis of recent marine sediments, 86, 71–80, <https://doi.org/10.1016/j.orggeochem.2015.06.009>, 2015.
- 455 Behar, F., Beaumont, V., and De B. Penteadó, H. L.: Rock-Eval 6 Technology: Performances and Developments, 56, 111–134, <https://doi.org/10.2516/ogst:2001013>, 2001.



- Bolinder, M. A., Janzen, H. H., Gregorich, E. G., Angers, D. A., and VandenBygaart, A. J.: An approach for estimating net primary productivity and annual carbon inputs to soil for common agricultural crops in Canada, 118, 29–42, <https://doi.org/10.1016/j.agee.2006.05.013>, 2007.
- 460 Bruun, S. and Jensen, L. S.: Initialisation of the soil organic matter pools of the Daisy model, 153, 291–295, 2002.
- Cagnarini, C., Renella, G., Mayer, J., Hirte, J., Schulin, R., Costerousse, B., Della Marta, A., Orlandini, S., and Menichetti, L.: Multi-objective calibration of RothC using measured carbon stocks and auxiliary data of a long-term experiment in Switzerland, 70, 819–832, <https://doi.org/10.1111/ejss.12802>, 2019.
- Cécillon, L.: A dual response, *Nat. Geosci.*, 14, 262–263, <https://doi.org/10.1038/s41561-021-00749-6>, 2021a.
- 465 Cécillon, L.: lauric-cecillon/PARTYsoc: Second version of the PARTYsoc statistical model, <https://doi.org/10.5281/zenodo.4446138>, 2021b.
- Cécillon, L., Baudin, F., Chenu, C., Houot, S., Jolivet, R., Kätterer, T., Luffalla, S., Macdonald, A., Van Oort, F., Plante, A. F., Savignac, F., Soucémariadin, L. N., and Barré, P.: A model based on Rock-Eval thermal analysis to quantify the size of the centennially persistent organic carbon pool in temperate soils, 15, 2835–2849, <https://doi.org/10.5194/bg-15-2835-2018>,
- 470 2018.
- Cécillon, L., Baudin, F., Chenu, C., Christensen, B. T., Franko, U., Houot, S., Kanari, E., Kätterer, T., Merbach, I., van Oort, F., Poeplau, C., Quezada, J. C., Savignac, F., Soucémariadin, L. N., and Barré, P.: Partitioning soil organic carbon into its centennially stable and active fractions with machine-learning models based on Rock-Eval® thermal analysis (PARTYsocv2.0 and PARTYsocv2.0eu), *Geosci. Model Dev.*, 14, 3879–3898, <https://doi.org/10.5194/gmd-14-3879-2021>,
- 475 2021.
- Chassé, M., Luftalla, S., Cécillon, L., Baudin, F., Abiven, S., Chenu, C., and Barré, P.: Long-term bare fallow soil fractions reveal thermo-chemical properties controlling soil organic carbon dynamics, 18, 1703–1718, <https://doi.org/10.5194/bg-18-1703-2021>, 2021.
- Clivot, H., Mouny, J. C., Duparque, A., Dinh, J. L., Denoroy, P., Houot, S., Vertès, F., Trochard, R., Bouthier, A., Sagot, S., and Mary, B.: Modeling soil organic carbon evolution in long-term arable experiments with AMG model, 118, 99–113, <https://doi.org/10.1016/j.envsoft.2019.04.004>, 2019.
- Coleman, K., Jenkinson, D. S., Crocker, G. J., Grace, P. R., Klír, J., Körschens, M., Poulton, P. R., and Richter, D. D.: Simulating trends in soil organic carbon in long-term experiments using RothC-26.3, 81, 29–44, [https://doi.org/10.1016/S0016-7061\(97\)00079-7](https://doi.org/10.1016/S0016-7061(97)00079-7), 1997.
- 485 Colomb, B., Debaeke, P., Jouany, C., and Nolot, J. M.: Phosphorus management in low input stockless cropping systems: Crop and soil responses to contrasting P regimes in a 36-year experiment in southern France, 26, 154–165, <https://doi.org/10.1016/j.eja.2006.09.004>, 2007.
- Cotrufo, M. F., Ranalli, M. G., Haddix, M. L., Six, J., and Lugato, E.: Soil carbon storage informed by particulate and mineral-associated organic matter, 12, 989–994, <https://doi.org/10.1038/s41561-019-0484-6>, 2019.



- 490 Crowther, T. W., van den Hoogen, J., Wan, J., Mayes, M. A., Keiser, A. D., Mo, L., Averill, C., and Maynard, D. S.: The global soil community and its influence on biogeochemistry, *Science*, 365, eaav0550, <https://doi.org/10.1126/science.aav0550>, 2019.
- Dangal, S. R. S., Schwalm, C., Cavigelli, M. A., Gollany, H. T., Jin, V. L., and Sanderman, J.: Improving soil carbon estimates by linking conceptual pools against measurable carbon fractions in the DAYCENT Model Version 4.5, *ESSOAR*,
495 <https://doi.org/10.1002/essoar.10507190.1>, 2021.
- Dignac, M.-F., Derrien, D., Barré, P., Barot, S., Cécillon, L., Chenu, C., Chevallier, T., Freschet, G. T., Garnier, P., Guenet, B., Hedde, M., Klumpp, K., Lashermes, G., Maron, P.-A., Nunan, N., Roumet, C., and Basile-Doelsch, I.: Increasing soil carbon storage: mechanisms, effects of agricultural practices and proxies. A review, *Agron. Sustain. Dev.*, 37, 14, <https://doi.org/10.1007/s13593-017-0421-2>, 2017.
- 500 Dimassi, B., Mary, B., Wylleman, R., Labreuche, J. Ô., Couture, D., Piraux, F., and Cohan, J. P.: Long-term effect of contrasted tillage and crop management on soil carbon dynamics during 41 years, 188, 134–146, <https://doi.org/10.1016/j.agee.2014.02.014>, 2014.
- Disnar, J. R., Guillet, B., Keravis, D., Di-Giovanni, C., and Sebag, D.: Soil organic matter (SOM) characterization by Rock-Eval pyrolysis: Scope and limitations, 34, 327–343, [https://doi.org/10.1016/S0146-6380\(02\)00239-5](https://doi.org/10.1016/S0146-6380(02)00239-5), 2003.
- 505 Erb, K. H., Luyssaert, S., Meyfroidt, P., Pongratz, J., Don, A., Kloster, S., Kuemmerle, T., Fetzel, T., Fuchs, R., Herold, M., Haberl, H., Jones, C. D., Marín-Spiotta, E., McCallum, I., Robertson, E., Seufert, V., Fritz, S., Valade, A., Wiltshire, A., and Dolman, A. J.: Land management: data availability and process understanding for global change studies, 23, 512–533, <https://doi.org/10.1111/gcb.13443>, 2017.
- FAO: Recarbonization of global soils: a dynamic response to offset global emissions, 8 pp., 2019.
- 510 FAO: Technical specifications and country guidelines for Global Soil Organic Carbon Sequestration Potential Map (GSOCseq), Rome, 2020.
- Farina, R., Sándor, R., Abdalla, M., Álvaro-Fuentes, J., Bechini, L., Bolinder, M. A., Brillì, L., Chenu, C., Clivot, H., De Antoni Migliorati, M., Di Bene, C., Dorich, C. D., Ehrhardt, F., Ferchaud, F., Fitton, N., Francaviglia, R., Franko, U., Giltrap, D. L., Grant, B. B., Guenet, B., Harrison, M. T., Kirschbaum, M. U. F., Kuka, K., Kulmala, L., Liski, J., McGrath,
515 M. J., Meier, E., Menichetti, L., Moyano, F., Nendel, C., Recous, S., Reibold, N., Shepherd, A., Smith, W. N., Smith, P., Soussana, J. F., Stella, T., Taghizadeh-Toosi, A., Tsutsikh, E., and Bellocchi, G.: Ensemble modelling, uncertainty and robust predictions of organic carbon in long-term bare-fallow soils, 27, 904–928, <https://doi.org/10.1111/gcb.15441>, 2021.
- Fuchs, R., Herold, M., Verburg, P. H., Clevers, J. G. P. W., and Eberle, J.: Gross changes in reconstructions of historic land cover/use for Europe between 1900 and 2010, 21, 299–313, <https://doi.org/10.1111/gcb.12714>, 2015.
- 520 Gregorich, E. G., Gillespie, A. W., Beare, M. H., Curtin, D., Sanei, H., and Yanni, S. F.: Evaluating biodegradability of soil organic matter by its thermal stability and chemical composition, *Soil Biology and Biochemistry*, 91, 182–191, <https://doi.org/10.1016/j.soilbio.2015.08.032>, 2015.



- He, Y., Trumbore, S. E., Torn, M. S., Harden, J. W., Vaughn, L. J. S., Allison, S. D., and Randerson, J. T.: Radiocarbon constraints imply reduced carbon uptake by soils during the 21st century, 353, 1419–1424, 2016.
- 525 Hemingway, J. D., Rothman, D. H., Grant, K. E., Rosengard, S. Z., Eglinton, T. I., Derry, L. A., and Galy, V. V.: Mineral protection regulates long-term global preservation of natural organic carbon, *Nature*, 570, 228–231, <https://doi.org/10.1038/s41586-019-1280-6>, 2019.
- Henin, S. and Dupuis, M.: Essai de bilan de la matière organique du sol, in: *Ann. Agron.* 11, Dunod (impr. de Chaix), Paris, 17–29, 1945.
- 530 Herbst, M., Welp, G., Macdonald, A., Jate, M., Hädicke, A., Scherer, H., Gaiser, T., Herrmann, F., Amelung, W., and Vanderborght, J.: Correspondence of measured soil carbon fractions and RothC pools for equilibrium and non-equilibrium states, 314, 37–46, <https://doi.org/10.1016/j.geoderma.2017.10.047>, 2018.
- IPCC: Climate Change and Land: an IPCC special report on climate change, desertification, land degradation, sustainable land management, food security, and greenhouse gas fluxes in terrestrial ecosystems, 2019.
- 535 Jobbagy, E. G. and Jackson, R. B.: The vertical distribution of Soil Organic Carbon and its relation to climate and vegetation, 10, 423–436, 2000.
- Khedim, N., Cécillon, L., Poulénard, J., Barré, P., Baudin, F., Marta, S., Rabatel, A., Dentant, C., Cauvy-Fraunié, S., Anthelme, F., Gielly, L., Ambrosini, R., Franzetti, A., Azzoni, R. S., Caccianiga, M. S., Compostella, C., Clague, J., Tielidze, L., Messenger, E., Choler, P., and Ficitola, G. F.: Topsoil organic matter build-up in glacier forelands around the world, *Glob. Change Biol.*, 27, 1662–1677, <https://doi.org/10.1111/gcb.15496>, 2021.
- 540 Lee, J., Viscarra Rossel, R. A., Luo, Z., and Wang, Y. P.: Simulation of soil carbon dynamics in Australia under a framework that better connects spatially explicit data with Roth C, *Biogeosciences Discussions*, <https://doi.org/10.5194/bg-2020-150>, 2020.
- Lehmann, J., Hansel, C. M., Kaiser, C., Kleber, M., Maher, K., Manzoni, S., Nunan, N., Reichstein, M., Schimel, J. P., Torn, M. S., Wieder, W. R., and Kögel-Knabner, I.: Persistence of soil organic carbon caused by functional complexity, 13, 529–534, <https://doi.org/10.1038/s41561-020-0612-3>, 2020.
- 545 Leifeld, J., Reiser, R., and Oberholzer, H. R.: Consequences of conventional versus organic farming on soil carbon: Results from a 27-year field experiment, 101, 1204–1218, <https://doi.org/10.2134/agronj2009.0002>, 2009a.
- Leifeld, J., Zimmermann, M., Fuhrer, J., and Conen, F.: Storage and turnover of carbon in grassland soils along an elevation gradient in the Swiss Alps, 15, 668–679, <https://doi.org/10.1111/j.1365-2486.2008.01782.x>, 2009b.
- 550 Levvasseur, F., Mary, B., Christensen, B. T., Duparque, A., Ferchaud, F., Kätterer, T., Lagrange, H., Montenach, D., Resseguier, C., and Houot, S.: The simple AMG model accurately simulates organic carbon storage in soils after repeated application of exogenous organic matter, 117, 215–229, <https://doi.org/10.1007/s10705-020-10065-x>, 2020.
- Lubet, E., Plénet, D., and Juste, C.: Effet à long terme de la monoculture sur le rendement en grain du maïs (*Zea mays* L) en conditions non irriguées., 13, 673–683, <https://doi.org/10.1051/agro:19930801>, 1993.



- Lugato, E., Lavallee, J. M., Haddix, M. L., Panagos, P., and Cotrufo, M. F.: Different climate sensitivity of particulate and mineral-associated soil organic matter, *Nat. Geosci.*, 14, 295–300, <https://doi.org/10.1038/s41561-021-00744-x>, 2021.
- Luo, Y., Ahlström, A., Allison, S. D., Batjes, N. H., Brovkin, V., Carvalhais, N., Chappell, A., Ciais, P., Davidson, E. A., Finzi, A., Georgiou, K., Guenet, B., Hararuk, O., Harden, J. W., He, Y., Hopkins, F., Jiang, L., Koven, C., Jackson, Robert, 560 B., Jones, C. D., Lara, M. J., Liang, J., McGuire, A. D., Parton, W., Peng, C., Randerson, J. T., Salazar, A., Sierra, C. A., Smith, M. J., Tian, H., Todd-Brown, K. E. O., Torn, M., Van Groenigen, K. J., Wang, Y. P., West, T. O., Wei, Y., Wieder, W. R., Xu, X., Xu, X., and Zhou, T.: Towards more realistic projections of soil carbon dynamics by Earth system models, 30, 40–56, <https://doi.org/10.1002/2015GB005239>, 2016.
- Luo, Z., Wang, E., Fillery, I. R. P., Macdonald, L. M., Huth, N., and Baldock, J.: Modelling soil carbon and nitrogen 565 dynamics using measurable and conceptual soil organic matter pools in APSIM, 186, 94–104, <https://doi.org/10.1016/j.agee.2014.01.019>, 2014.
- Manzoni, S. and Porporato, A.: Soil carbon and nitrogen mineralization: Theory and models across scales, 41, 1355–1379, <https://doi.org/10.1016/j.soilbio.2009.02.031>, 2009.
- Martin, M., Dimassi, B., Millet, F., Picaud, C., Bounoua, E.-M., Bardy, M., Bispo, A., Boulonne, L., Bouthier, A., 570 Duparque, A., Eglin, T., Guenet, B., Huard, F., Mary, B., Mathias, E., Mignolet, C., Robert, C., Saby, N., Sagot, S., Schott, C., and Toutain, B. R.: Méthodes de comptabilisation du stockage de carbone organique des sols sous l’effet des pratiques culturales (CSopra), <https://www.ademe.fr/csopra-methodes-comptabilisation-stockage-carbone-organique-sols-sous-leffet-pratiques-culturales>, 2019.
- Mary, B., Clivot, H., Blaszczyk, N., Labreuche, J., and Ferchaud, F.: Soil carbon storage and mineralization rates are 575 affected by carbon inputs rather than physical disturbance: Evidence from a 47-year tillage experiment, 299, 106972, <https://doi.org/10.1016/j.agee.2020.106972>, 2020.
- Messiga, A. J., Ziadi, N., Plénet, D., Parent, L. E., and Morel, C.: Long-term changes in soil phosphorus status related to P budgets under maize monoculture and mineral P fertilization, 26, 354–364, <https://doi.org/10.1111/j.1475-2743.2010.00287.x>, 2010.
- 580 Morel, C., Ziadi, N., Messiga, A., Bélanger, G., Denoroy, P., Jeangros, B., Jouany, C., Fardeau, J. C., Mollier, A., Parent, L. E., Proix, N., Rabeharisoa, L., and Sinaj, S.: Modeling of phosphorus dynamics in contrasting agroecosystems using long-term field experiments, 94, 377–387, <https://doi.org/10.4141/CJSS2013-024>, 2014.
- Nave, L., Johnson, K., Ingen, C. van, Agarwal, D., Humphrey, M., and Beekwilder, N.: International Soil Carbon Network (ISCN) Database V3-1, <https://doi.org/10.17040/ISCN/1305039>, 2015.
- 585 Nemo, Klumpp, K., Coleman, K., Dondini, M., Goulding, K., Hastings, A., Jones, M. B., Leifeld, J., Osborne, B., Saunders, M., Scott, T., Teh, Y. A., and Smith, P.: Soil Organic Carbon (SOC) Equilibrium and Model Initialisation Methods: an Application to the Rothamsted Carbon (RothC) Model, 22, 215–229, <https://doi.org/10.1007/s10666-016-9536-0>, 2016.



- Noirot-Cosson, P. E., Vaudour, E., Gilliot, J. M., Gabrielle, B., and Houot, S.: Modelling the long-term effect of urban waste compost applications on carbon and nitrogen dynamics in temperate cropland, 94, 138–153, <https://doi.org/10.1016/j.soilbio.2015.11.014>, 2016.
- Oberholzer, H. R., Leifeld, J., and Mayer, J.: Changes in soil carbon and crop yield over 60 years in the Zurich Organic Fertilization Experiment, following land-use change from grassland to cropland, 177, 696–704, <https://doi.org/10.1002/jpln.201300385>, 2014.
- Obriot, F.: Epannage de produits résiduels organiques et fonctionnement biologique des sols : De la quantification des impacts sur les cycles carbone et azote à l'évaluation multicritère de la pratique à l'échelle de la parcelle., AgroParisTech, 194–195 pp., 2016.
- Otto, S. A., Kadin, M., Casini, M., Torres, M. A., and Blenckner, T.: A quantitative framework for selecting and validating food web indicators, 84, 619–631, <https://doi.org/10.1016/j.ecolind.2017.05.045>, 2018.
- Poeplau, C., Don, A., Vesterdal, L., Leifeld, J., Van Wesemael, B., Schumacher, J., and Gensior, A.: Temporal dynamics of soil organic carbon after land-use change in the temperate zone - carbon response functions as a model approach, 17, 2415–2427, <https://doi.org/10.1111/j.1365-2486.2011.02408.x>, 2011.
- Poeplau, C., Don, A., Dondini, M., Leifeld, J., Nemo, R., Schumacher, J., Senapati, N., and Wiesmeier, M.: Reproducibility of a soil organic carbon fractionation method to derive RothC carbon pools, 64, 735–746, <https://doi.org/10.1111/ejss.12088>, 2013.
- Poeplau, C., Don, A., Six, J., Kaiser, M., Benbi, D., Chenu, C., Cotrufo, M. F., Derrien, D., Gioacchini, P., Grand, S., Gregorich, E., Griepentrog, M., Gunina, A., Haddix, M., Kuzyakov, Y., Kühnel, A., Macdonald, L. M., Soong, J., Trigalet, S., Vermeire, M. L., Rovira, P., van Wesemael, B., Wiesmeier, M., Yeasmin, S., Yevdokimov, I., and Nieder, R.: Isolating organic carbon fractions with varying turnover rates in temperate agricultural soils – A comprehensive method comparison, 125, 10–26, <https://doi.org/10.1016/j.soilbio.2018.06.025>, 2018.
- Poeplau, C., Barré, P., Cécillon, L., Baudin, F., and Sigurdsson, B. D.: Changes in the Rock-Eval signature of soil organic carbon upon extreme soil warming and chemical oxidation - A comparison, *Geoderma*, 337, 181–190, <https://doi.org/10.1016/j.geoderma.2018.09.025>, 2019.
- Rumpel, C., Amiraslani, F., Koutika, L. S., Smith, P., Whitehead, D., and Wollenberg, E.: Put more carbon in soils to meet Paris climate pledges, 564, 32–34, <https://doi.org/10.1038/d41586-018-07587-4>, 2018.
- Saffih-Hdadi, K. and Mary, B.: Modeling consequences of straw residues export on soil organic carbon, 40, 594–607, <https://doi.org/10.1016/j.soilbio.2007.08.022>, 2008.
- Sanderman, J., Baldock, J. A., Dangal, S. R. S., Ludwig, S., Potter, S., Rivard, C., and Savage, K.: Soil organic carbon fractions in the Great Plains of the United States: an application of mid-infrared spectroscopy, 1, <https://doi.org/10.1007/s10533-021-00755-1>, 2021.
- Schrumpf, M., Schulze, E. D., Kaiser, K., and Schumacher, J.: How accurately can soil organic carbon stocks and stock changes be quantified by soil inventories?, 8, 1193–1212, <https://doi.org/10.5194/bg-8-1193-2011>, 2011.



- Shi, Z., Crowell, S., Luo, Y., and Moore, B.: Model structures amplify uncertainty in predicted soil carbon responses to climate change, *Nat Commun*, 9, 2171, <https://doi.org/10.1038/s41467-018-04526-9>, 2018.
- Skjemstad, J. O., Spouncer, L. R., Cowie, B., and Swift, R. S.: Calibration of the Rothamsted organic carbon turnover model (RothC ver. 26.3), using measurable soil organic carbon pools, 42, 79–88, <https://doi.org/10.1071/SR03013>, 2004.
- 625 Smith, J. U., Smith, P., and Addiscot, T.: Quantitative methods to evaluate and compare soil organic matter (SOM) models, in: *Evaluation of Soil Organic Matter Models Using existing Long-Term Datasets*, Springer Berlin, Heidelberg, 181–199, 1996.
- Smith, P. and Falloon, P. D.: Modelling refractory soil organic matter, 30, 388–398, <https://doi.org/10.1007/s003740050019>,
630 2000.
- Taghizadeh-Toosi, A., Cong, W. F., Eriksen, J., Mayer, J., Olesen, J. E., Keel, S. G., Glendining, M., Kätterer, T., and Christensen, B. T.: Visiting dark sides of model simulation of carbon stocks in European temperate agricultural soils: allometric function and model initialization, 450, 255–272, <https://doi.org/10.1007/s11104-020-04500-9>, 2020.
- Todd-Brown, K. E. O., Randerson, J. T., Hopkins, F., Arora, V., Hajima, T., Jones, C., Shevliakova, E., Tjiputra, J., Volodin, E., Wu, T., Zhang, Q., and Allison, S. D.: Changes in soil organic carbon storage predicted by Earth system models during the 21st century, 11, 2341–2356, <https://doi.org/10.5194/bg-11-2341-2014>, 2014.
- 635 UN General Assembly: UN General Assembly, *Transforming our world : the 2030 Agenda for Sustainable Development*, 21 October 2015, A/RES/70/1, 2015.
- Vermeulen, S., Bossio, D., Lehmann, J., Luu, P., Paustian, K., Webb, C., Augé, F., Bacudo, I., Baedeker, T., Havemann, T., Jones, C., King, R., Reddy, M., Sunga, I., Von Unger, M., and Warnken, M.: A global agenda for collective action on soil carbon, 2, 2–4, <https://doi.org/10.1038/s41893-018-0212-z>, 2019.
- 640 Vertès, F., Simon, J.-C., Laurent, F., and Besnard, A.: Prairies et qualité de l’eau. Evaluation des risques de lixiviation d’azote et optimisation des pratiques, 192, 423–440, 2007.
- Viscarra Rossel, R. A., Lee, J., Behrens, T., Luo, Z., Baldock, J., and Richards, A.: Continental-scale soil carbon composition and vulnerability modulated by regional environmental controls, 12, 547–552, <https://doi.org/10.1038/s41561-019-0373-z>, 2019.
- 645 Wallach, D.: *Evaluating crop models*, 11–53, 2006.
- Wickham, H.: *Reshaping Data with the reshape Package*, 21, 1–20, <https://doi.org/10.18637/jss.v021.i12>, 2007.
- Wickham, H.: *ggplot2: Elegant Graphics for Data Analysis*, Springer-Verlag, New York, 2016.
- 650 Wiesmeier, M., Urbanski, L., Hobbey, E., Lang, B., von Lützow, M., Marin-Spiotta, E., van Wesemael, B., Rabot, E., Ließ, M., Garcia-Franco, N., Wollschläger, U., Vogel, H. J., and Kögel-Knabner, I.: Soil organic carbon storage as a key function of soils - A review of drivers and indicators at various scales, 333, 149–162, <https://doi.org/10.1016/j.geoderma.2018.07.026>, 2019.
- Wutzler, T. and Reichstein, M.: Soils apart from equilibrium - Consequences for soil carbon balance modelling, 4, 125–136,
655 <https://doi.org/10.5194/bg-4-125-2007>, 2007.



Zimmermann, M., Leifeld, J., Schmidt, M. W. I., Smith, P., and Fuhrer, J.: Measured soil organic matter fractions can be related to pools in the RothC model, 58, 658–667, <https://doi.org/10.1111/j.1365-2389.2006.00855.x>, 2007a.

Zimmermann, M., Leifeld, J., and Fuhrer, J.: Quantifying soil organic carbon fractions by infrared-spectroscopy, 39, 224–231, <https://doi.org/10.1016/j.soilbio.2006.07.010>, 2007b.