

To what extent can soil moisture and soil Cu contamination stresses affect nitrous species emissions? Estimation through calibration of a nitrification/denitrification model

5 Laura Sereni^{1*}, Bertrand Guenet², Charlotte Blasi^{1,3}, Olivier Crouzet^{1,4}, Jean-Christophe Lata^{5,6}
Isabelle Lamy¹

¹Université Paris-Saclay, INRAE, AgroParisTech, UMR ECOSYS, Ecotoxicology Team, 78026, Versailles, France

²Laboratoire de Géologie de l'ENS, PSL Research University, CNRS, UMR 8538, IPSL, Paris, France

³Present address: Centre Sève, Département de Chimie, Université de Sherbrooke, Sherbrooke, QC, Canada

10 ⁴Present address: Office national de la chasse et de la faune sauvage, Site d'Auffargis-Saint-Benoist 78612 Le-Perray-en-Yvelines, France

⁵Sorbonne Université, Université de Paris, UPEC, CNRS, INRAE, IRD, UMR 7618, Institute of Ecology and Environmental Sciences – Paris, iEES Paris, 7 quai St Bernard 75252, Paris, France

15 ⁶Department of Geocology and Geochemistry, Institute of Natural Resources, Tomsk Polytechnic University, 30, Lenin Street, Tomsk, 634050, Russia

*Correspondence to Laura Sereni (laura.sereni@inrae.fr)

Abstract

Continental biogeochemical models are commonly used to predict the effect of land use, exogenous organic matter input or climate change on soil greenhouse gas emission. However, they cannot be used for this purpose to investigate the effect of soil contamination, while contamination affects several soil processes and concerns a large fraction of land surface. For that, in this study we implemented a commonly used model estimating soil nitrogen (N) emissions, the DeNitrification DeComposition (DNDC) model, with a function taking into account soil copper (Cu) contamination in nitrate production control. Then, we aimed at using this model to predict N_2O-N , NO_2-N , $NO-N$ and NH_4-N emissions in the presence of contamination and in the context of changes in precipitations. Initial incubations of soils were performed at different soil moistures in order to mimic expected rainfall patterns during the next decades and in particular drought and excess of water. Then, a bioassay was used in the absence or presence of Cu to assess the effect of the single (moisture) or double stress (moisture and Cu) on soil nitrate production. Data of nitrate production obtained through a gradient of Cu under each initial moisture incubation were used to parameterize the DNDC model and to estimate soil N emission considering the various effects of Cu. Whatever the initial moisture incubation, experimental results showed a NO_3-N decreasing production when Cu was added but with different sharpness depending on soil moisture. The DNDC-Cu version we proposed was able to reproduce these observed Cu effects on soil nitrate concentration with $r^2 > 0.99$ and $RMSE < 10\%$ for all treatments in the DNDC-Cu calibration range ($> 40\%$ of the water holding capacity) but showed poor performances for the dry treatments. We modelled a Cu-effect inducing an increase in NH_4-N soil concentration and emissions due to a reduced nitrification activity, and therefore a decrease in NO_3-N , N_2O-N and NO_x-N concentrations and emissions. The effect of added Cu predicted by the model was larger on N_2-N and N_2O-N emissions than on the other N species and larger for the soils incubated under constant than variable moisture. Our work shows that soil contamination can be considered in continental biogeochemical models to better predict soil greenhouse gas emissions.

Keywords: bioassay, DNDC modelling, rainfall pattern, ecotoxicology, soil function

1. Introduction

45 The increase in atmospheric greenhouse gases [GHG] like CO₂, CH₄, or N₂O is expected to induce a global climate change with e.g. higher mean temperature or changes in rainfall patterns with projections of increased precipitations or droughts depending on regions (Knutti and Sedláček, 2012). These modifications in rainfall patterns may impact soil moisture which is one of the main drivers of soil microbial activity (Moyano et al., 2013; Schimel, 2018; Stark and Firestone, 1995). Microbial communities ensure key activities supporting numerous ecosystem functions, such as those involved in nitrogen (N) cycle influencing N₂O emissions (Butterbach-Bahl et al., 2013; Galloway et al., 2008) and are at the origin of more than 80% of N₂O fluxes (IPCC, 2019). In particular, nitrification/denitrification processes are largely controlled by the local (an-)oxic treatments and therefore by soil moisture (Borken and Matzner, 2009; Fierer et al., 2003; Guo et al., 2014; Schimel, 2018), denitrification being the main source of soil N₂O emission for moist soils whereas for dry soils N₂O emissions are mainly due to nitrification (Bateman and Baggs, 2005). N soil fluxes dynamics are thus particularly difficult to predict at a large scale because of this strong dependency to local soil O₂ availability (Khalil et al., 2004) disrupting the realization of nitrification/denitrification reactions and N species diffusion (Conrad 1996; Schurgers et al. 2006). Despite this, some continental biogeochemical models have shown improved predictions when N cycle was explicitly represented (Butterbach-Bahl et al., 2009; Kesik et al., 2005; Vuichard et al., 2019).

60 In addition to climate change, human activities introduce significant quantities of contaminants into the environment, such as trace elements (TE) which are persistent and can be toxic for soil biota (Bech et al., 1997; Giller et al., 2009). Indeed, the contamination of soils by TE has become a major concern at global scale (De Vleeschouwer et al. 2007; Khan et al. 2008) coming from atmospheric sources (Steinnes et al., 1997) or through the use of pesticides (Nicholson et al., 2003). In particular, TE contaminations are known to largely affect soil microorganisms (Bååth, 1989; Giller et al., 2009) and their activities, such as nitrification/denitrification processes (Broos et al., 2007; Mertens et al., 2010). Therefore, the combined effect of climate change and of soil contamination may largely impact the emissions of NO_x and N₂O from soils (Holtan-Hartwig et al., 2002; Vásquez-Murrieta et al. 2006). However, the effect of the interactions between climate change and soil contamination on the GHG emissions is still poorly documented (Rillig et al., 2019; Zandalinas et al., 2021).

70 Despite recent progress, the Earth system models (ESMs) used to predict future climate change still don't take into account soil contamination effect on GHG emissions (Anav et al., 2013) whereas at a large spatial scale many soils are listed as contaminated (FAO, 2008; Lado et al., 2008). Furthermore, soil biogeochemical models are often used to estimate loss or accumulation of N species (ammoniac NH₄ volatilization, nitrate NO₃ leaching) (Giltrap et al., 2010) or they respective concentrations under scenarii of organic fertilizer amendments, but do not take into account the contamination which often occurs simultaneously (Wuana and Okieimen, 2011). Thus, there is a growing need to provide continental models combining ecotoxicological/contamination and climate change concerns. Among the biogeochemical models, DeNitrification DeCompostion (DNDC, Changsheng Li et al., 1992) is a relatively simple model handling both biogeochemistry of denitrification and microbial growth (Li et al., 2000), and on which Land Surface Model-soil N component -a part of ESMs- like ORCHIDEE are built (Vuichard et al., 2019).

In order to improve model outputs, this study combines in an innovative way experimental and modelling approaches to evaluate the impact of soil moisture on the sensitivity of nitrification to copper (Cu) toxicity and consequently on GHG-N emissions. Cu was chosen as a model of soil contamination due to both its relevance in agricultural soils and available data in the literature (Broos et al., 2007; Mertens et al., 2010; Sauvé et al., 1999).

85 It is not straightforward to assess distinct effects between punctual or chronic contamination on microbial structure or soil functions (Brandt et al., 2010; Oorts et al., 2006; Smolders et al., 2009). Here, we designed experiments to assess the conjugated effects of a trace metal contamination and a soil moisture stress on soil N cycle. Soil initial incubations were run during five weeks by applying a given soil moisture from drought to water saturation. Then, a bioassay with a gradient of Cu added by spiking was performed to estimate NO_3^-

90 production. The experimental data were used to calibrate a new model, DNDC-Cu, able to predict NO_x and N_2O emissions with the implementation of new functions considering the effect of Cu concentration ([Cu]) on nitrification/denitrification processes. Our hypothesis is that the building of such a model allows a gain in the understanding of the effect of a soil [Cu] on NO_x and N_2O and NH_4 cycling in a climate change context. Hence, data are also used here to discuss knowledge gaps in such modelling approaches, and to question the matter of

95 soil contamination data in climate change scenarii.

2. Materials and Methods

2.1 Soil sampling

The soil was sampled in January 2017 at the surface layer (0–20 cm) of a control plot at the Qualiagro experimental site (48°87'N, 1° 97'E - https://www6.inrae.fr/valor-pro_eng/Experimental-devices/QualiAgro/QualiAgro-web-site). The soil sample was immediately wet sieved at 5mm and shortly stored at 4°C until microcosm build-up. Aliquots of this sieved soil were used to measure the initial water content in addition to the maximum water holding capacity (WHC) for the further microcosm experiments. This site is located at Feucherolles near Paris, France, and had been designed to evaluate urban compost fertility together with the monitoring of contaminant inputs

105 (Cambier et al., 2019). Soil is a luvisol with 15% clay, 78% silt and 7% sand, a pH of 6.9, organic carbon (Corg) and total N contents at 10.5 ± 0.2 and 1.00 ± 0.03 g kg^{-1} soil, respectively, and with a CEC of 7.9 ± 0.8 $\text{cmol}^+ \text{kg}^{-1}$ soil. This soil is not contaminated with Cu, and geochemical [Cu] background measured by ICP-AES after HF-HClO_4 extraction was of 12 mg Cu.kg^{-1} soil.

2.2 Experimental setup

110 In order to evaluate the impact of soil moisture on the sensitivity of nitrification to Cu toxicity, we carried out a two-step experiment. The first step consisted in initial incubations at 5 different WHC during 5 weeks, and the second step in a 3-day bioassay with spiked Cu gradient (Fig. 1).

For the 5 weeks' initial incubation, five microcosms were built up with about 5g of sampled soil. Three of them were set up with a constant moisture corresponding to 30%, 60% and 90% of their WHC in order to span

115 respectively limiting, optimal, and saturating conditions for the microbial activities. These three samples will be called thereafter "30%, 60% and 90%", respectively. Their water contents were verified by weighting every two

120 days and water added if necessary. The two other microcosms were incubated in order to simulate two kinds of drought and Dry-Rewet cycles. One, thereafter called “Drought” (or DO), started with one week at 60% WHC and then the soil was left for 3 weeks without added water to mimic a dry period until 10% of the WHC before rewetting at the initial 60% WHC. The other, thereafter called “Dry-Rewet” (or DR) encountered 2 cycles of one-week near-saturation period (90% WHC) followed by one-week dry period (10% of the WHC) ending by one week near saturation period. Drying was performed by natural evaporation (gentle air-drying at the laboratory temperature, i.e. 20°C) and controlled by weighting.

125 At the end of the initial incubation period, we performed a nitrification bioassay using 3 replicates originating from soils and following an adaptation of the method proposed by Petersen et al. (2012). Bioassay consisted in nitrate production measurement over a short-term aerobic incubation in soil slurries (ratio soil:solution 1:10) with ammonium in excess and in the presence of gradients of Cu. Briefly, 3.5 g of fresh soil (approximately 3 g of soil equivalent dry weight), were mixed in a 50 mL Falcon® tubes with 29mL of a 10 mM HEPES buffer solution (hydroxyethyl piperazineethanesulfonic acid, Sigma-Aldrich, France) to maintain a constant pH under Cu
130 spiking and nitrification activity, and containing the substrate (NH₄)₂SO₄ (3 mM) (Sigma-Aldrich, France). Soils were first spiked with a gradient of increasing Cu²⁺ in the presence of an excess of NH₄⁺ and the resulting potential nitrification activity (PNA) measured. The microcosms incubated at constant moisture were kept at their moisture level (30, 60 or 90% of WHC) whereas those incubated at variable moisture were set at 60% WHC. The NO₃⁻ production rates were measured in soil slurries over a short-term aerobic incubation, for each Cu added
135 concentration. Briefly, 1mL of Cu solution at different concentrations was added in soil slurries to reach added [Cu] of 50, 100, 250, 500, 750, 1000 and 2000 mg Cu.kg soil⁻¹ (final soil [Cu] of 62, 112, 262, 512, 762, 1012 and 2012 mg Cu.kg soil⁻¹ and control with 12 mgCu.kg soil⁻¹). The pH was adjusted to 7. Then, microcosms were incubated on a rotary shaker (150rpm) under aerobic conditions at 25°C until 72h. After 0, 24 and 72h of incubation, 2 ml aliquots of 3g were transferred in Eppendorf vials and centrifuged. The supernatants were collected and stored
140 in microplates at – 20 °C until analyses of NO₃⁻ and NO₂⁻ by colorimetric determinations, following the reduction of NO₃⁻ in NO₂⁻ by vanadium(III) and then the detection of NO₂⁻ by the acidic Griess reaction (Miranda et al., 2001). Finally, PNA (µg NO₃-N g⁻¹ soil h⁻¹) was calculated on the basis of NO₃⁻-N + NO₂⁻-N concentrations measured at different time steps. In our bioassay, [NO₂⁻] were negligible and PNA was thus calculated following Eq. (1), by checking the linear production rate of NO₃⁻ between 2 h, 24 h and 72h:

$$145 \quad (1) PNA = \frac{[NO_3^-]_{T_{final}} - [NO_3^-]_{T_{initial}}}{T_{final} - T_{initial}} \times V_S \div W$$

with V_S : Volume of solution

W : Weight of fresh soil

T : Time of incubation.

150 Cu in solution was measured by centrifugation of the soil+solution mixture of each bioassay, followed by a determination of Cu in solution by Flame Atomic Absorption Spectroscopy. Cu in solution values are provided in Suppl. Table 1.

2.3 Nitrification/denitrification model

155 Nitrification and denitrification processes are represented following the DNDC model proposed by Changsheng Li et al. (1992) and Li et al. (2000). In this study, we used a simplified version of DNDC adapted by Zaehle and Friend (2010) initially calibrated for soil WHC >40%, that we intended here to test for 30% of WHC. This simplified version needs less boundary data but keeps a mechanistic description of the main processes. Modelled N species are expressed in amount of N, i.e. NH₄-N, NO₃-N, NO_x-N and N₂O-N. To be able to represent both nitrification and denitrification processes occurring in aerobic and anaerobic sites, the soil is split into aerobic and anaerobic fractions based on an empirical relationship linking O₂ consumption to soil respiration. In aerobic microsites, nitrification takes places following Eq. (2):

$$(2) \text{Nitrification} = f(SWC) \times f(temp) \times f(pH) \times k_{Nit} \times (1 - anv) \times NH_4$$

165 with NH₄-N being the stock of ammonium (in gN.m⁻²), (1-anv) the aerobic fraction of the soil described thereafter in Eq. (21), k_{Nit} the nitrification rate (day⁻¹), $f(SWC)$, $f(temp)$ and $f(pH)$ three rate modifiers representing the effect of soil water content (m³ m⁻³), temperature (K) and pH as scalar respectively. They are described by the following Eq. (3), (4) and (5):

$$(3) f(SWC) = 0.0243 + 0.9975 \times SWC + 5.6358 \times SWC^2 + 17.651 \times SWC^3 + 12.904 \times SWC^4$$

$$(4) f(temp) = 0.0233 + 0.3094 \times temp + 0.2234 \times temp^2 + 0.1566 \times temp^3 + 0.0272 \times temp^4$$

$$(5) f(pH) = 1.2314 + 0.7347 \times pH + 0.0604 \times pH^2$$

170

The NH₄-N nitrified is transformed into N₂O-N, NO-N or NO₃-N due to microbial processes and chemonitrification following Eq. (6),(7) and (8):

$$(6) \text{Nitrification}_{N_2O} = f_{tv} \times SWC \times k_{Nitrif_{N_2O}} \times \text{Nitrification}$$

$$175 (7) \text{Nitrification}_{NO} = f_{tv} \times SWC \times k_{Nitrif_{NO}} \times \text{Nitrification} + 496950 \times e^{-1.62 \times pH} \times e^{-31494 / (temp \times R)} \times \text{Nitrification}$$

$$(8) \text{Nitrification}_{NO_3} = \text{Nitrification} - \text{Nitrification}_{NO} - \text{Nitrification}_{N_2O}$$

180 with $k_{Nitrif_{NO}}$ and $k_{Nitrif_{N_2O}}$ two fixed rates (d⁻¹), f_{tv} a rate modifier controlled by temperature and given in Eq. (9) and R the ideal gas constant.

$$(9) f_{tv} = 2.72^{\left(34.6 - \frac{9615}{temp}\right)}$$

Then, the $\text{NO}_3\text{-N}$ produced during the nitrification process enters the denitrification module where it is reduced sequentially into $\text{NO}_x\text{-N}$, $\text{N}_2\text{O-N}$ or $\text{N}_2\text{-N}$ following Eq. (10) to (12):

185

$$(10) \text{Denitrification}_{\text{NO}_x} = \text{anvf} \times \left(\frac{\mu_{\text{NO}_3}}{0.401} + 0.09 \times \frac{\text{NO}_3}{\text{N}_{\text{tot}}} \right) \times B$$

$$(11) \text{Denitrification}_{\text{N}_2\text{O}} = \text{anvf} \times \left(\frac{\mu_{\text{NO}_x}}{0.428} + 0.035 \times \frac{\text{NO}_x}{\text{N}_{\text{tot}}} \right) \times B$$

$$(12) \text{Denitrification}_{\text{N}_2} = \text{anvf} \times \left(\frac{\mu_{\text{N}_2\text{O}}}{0.151} + 0.079 \times \frac{\text{N}_2\text{O}}{\text{N}_{\text{tot}}} \right) \times B$$

The anaerobic fraction anvf is described following Eq. (13):

190

$$(13) \text{anvf} = 0.85 \times \left(1 - \frac{p_{\text{soil O}_2}}{p_{\text{air O}_2}} \right)$$

with $p_{\text{air O}_2}$, $p_{\text{soil O}_2}$ being the partial pressure in the air and in the soil respectively. $p_{\text{soil O}_2}$ is calculated following Eq. (14)

195

$$(14) \frac{\partial p_{\text{soil O}_2}}{\partial t} = p_{\text{soil O}_2} - p_{\text{O}_2\text{resp}} \times k \times \text{SOC} \times f_{\text{Cu}}$$

with SOC being the soil organic carbon stock (gC m^{-2}), k the decomposition rate, $p_{\text{O}_2\text{resp}}$ the O_2 partial pressure related to the respiration, and f_{Cu} the effect of Cu on CO_2 emissions as define in Eq.(15), following (Sereni et al., 2021) Eq. (5):

200

$$(15) f_{\text{CuCO}_2} = \exp(-0.1 - 0.1 \times \log(\text{Cu}) + 0.12 \times \text{pH})$$

The relative growth rate of $\text{NO}_3\text{-N}$, $\text{NO}_x\text{-N}$ and $\text{N}_2\text{O-N}$ denitrifiers are described respectively by μ_{NO_3} , μ_{NO_x} , $\mu_{\text{N}_2\text{O}}$ following Eq. (16), (17) and (18).

205

$$(16) \mu_{\text{NO}_3} = \frac{0.67 \times f_{\text{denit}}(\text{temp}) \times f_{\text{denit}_{\text{NO}_3}}(\text{pH}) \times \text{NO}_3}{\text{NO}_3 + 166}$$

$$(17) \mu_{\text{NO}_x} = \frac{0.34 \times f_{\text{denit}}(\text{temp}) \times f_{\text{denit}_{\text{NO}_x}}(\text{pH}) \times \text{NO}_x}{\text{NO}_x + 166}$$

$$(18) \mu_{\text{N}_2\text{O}} = \frac{0.34 \times f_{\text{denit}}(\text{temp}) \times f_{\text{denit}_{\text{N}_2\text{O}}}(\text{pH}) \times \text{N}_2\text{O}}{\text{N}_2\text{O} + 166}$$

210 with $f_{\text{denit}}(\text{temp})$, $f_{\text{denit}_{\text{NO}_3}}(\text{pH})$, $f_{\text{denit}_{\text{NO}_x}}(\text{pH})$, $f_{\text{denit}_{\text{N}_2\text{O}}}(\text{pH})$ being rates modifiers depending on air temperature and soil pH described in Eq. (19) to (22).

$$(19) f_{\text{denit}}(\text{temp}) = 2^{(\text{temp} - 22.5)/10}$$

$$(20) f_{\text{denit}_{\text{NO}_3}}(\text{pH}) = 1 - \frac{1}{1 + e^{(4.25 \times \text{pH})/0.5}}$$

215

$$(21) f_{\text{denit}_{\text{NO}_x}}(\text{pH}) = 1 - \frac{1}{1 + e^{(5.25 \times \text{pH})}}$$

$$(22) f_{\text{denit}_{\text{N}_2\text{O}}}(\text{pH}) = 1 - \frac{1}{1 + e^{(6.25 \times \text{pH})/1.5}}$$

The denitrifier biomass dynamic B (kg m^{-2}) is described following Eq. (23).

$$(23) \frac{\partial B}{\partial t} = (anvf \times (\mu_{NO_3} + \mu_{NOX} + \mu_{N_2O}) - 3.82 \times 10^{-3}) \times B$$

Finally, all the gaseous forms of mineral N are emitted into the atmosphere. It is important to note that we did not use directly the DNDC model but a simplified version adapted by Zaehle and Friend, (2010). The original code was in fortran and we translated it in R to facilitate its manipulation. The time step of the model was 30 minutes and most of the parameters were kept to the original values of Changsheng Li et al. (1992) and Li et al. (2000) except k_{nit} that was modified to 0.1743 instead of 0.2 to better fit the data from the control. Furthermore, the amounts of $\text{NH}_4\text{-N}$ fixed to the clay were reduced to 0 as the bioassay was performed in excess of $\text{NH}_4\text{-N}$ (see 2.2.0).

We used measures of N species at the end of initial incubation period as initial values of N species for DNDC (Table 1a and Fig. 2). To estimate the anaerobic volume fraction during the 3 days bioassay, we used a C mineralization rate k (Eq. 14) determined on the basis of measurements performed on the same soil (Annabi et al., 2007) and ran DNDC for a 45 days equilibrium period. We then extracted the initial anaerobic volume fraction and partial O_2 pressure.

2.4 Statistical analysis

The dose-response curves of PNA during the bioassay to Cu gradient were plotted and tested with linear, quadratic or cubic functions as fitting models. Our aim was to find, if possible, a similar modelling fit function for all moisture initial incubation treatments. Thus, for each moisture treatment, the two best functions of fit were selected through AIC and R^2 criteria, and compared with ANOVA. After selection of a common type of functions, the permutability of the different functions parameters was tested with the Chow test (gap v.1.2.2 package which tested the regression 1 on the basis of the samples 2 and vice-versa). If the p-value exceeds its critical values, regressions cannot be considered equal (Zhao, 2007).

To estimate the effect of [Cu] and soil moisture on the different variables measured, we used nonparametric Kruskal-Wallis test. The fits between the model and the data of soil nitrate concentration during the bioassays were measured using root mean square error (RMSE, Eq. (24)):

$$(24) RMSE = \sqrt{\frac{1}{N} \sum_{i=1}^N (X_i - Y_i)^2}$$

where i is the number of observations (1 to N), X is the predicted value and Y is the observed value. RMSE was decomposed in standard bias (Eq. (25)), non-unity slope (Eq. (26)) and lack of correlation (Eq. (27)) component following Gauch et al. (2003), with \bar{X} and \bar{Y} the mean modelled and observed values, b the slope of the least square regression of Y on X and r^2 the square of the correlation:

$$(25) : SB = (\bar{X} - \bar{Y})^2$$

$$(26) NU = (1 - b)^2 \times \sum \frac{x_n^2}{N}$$

$$(27) LC = (1 - r^2) \times \sum \frac{y_n^2}{N}$$

All analysis were done with R 3.2.3 (R Core Team, 2015).

255

3 Results

3.1. Effect of Cu on potential nitrification activity (PNA): statistical model selection

260 The soil N species measured at the end of the soil initial incubations in each soil moisture treatment were used to initialise the DNDC model (Table 1). Two anomalous points leading to anomalous calculated NO₂-N values were excluded from the experimental results because of technical problems during measurements (the C replicates in the DR and DO cases).

265 The bioassay experiments performed at the end of the soil initial incubations allowed us to determine the rate of nitrate production as a function of soil [Cu] for each soil moisture (Fig. 1). In all cases, the PNA values were found to decrease with the increase in soil [Cu] but at different rates depending on the moisture treatment. Based on AIC values (suppl. Table 2), we first selected one model per moisture incubation that better fitted the data. For 30 and 60% of WHC, a quadratic model was found to provide the better compromise between the number of parameters and the prediction capacity. For 90% WHC, no significant difference was found between the cubic and the quadratic models (ANOVA, p.v=0.07). For DR, no significant difference was found between linear and quadratic models (supplementary Tables 2a and 2b) whereas for DO the cubic model provided a substantially better fit than the quadratic model (AIC and adj. R2 score, suppl. table 2). Finally, we found that the quadratic model fitted correctly all the sets of data, allowing to be homogeneous across the initial moisture incubation treatments (Fig. 2b). The quadratic function was thus chosen to quantify the Cu effect on PNA including the DO treatment.

275 The parameters of the five quadratic functions (one for each moisture treatment) were found different from each other, except for 60 and 90% WHC (p.v=0.001, Chow test). A single function was thus used to adjust PNA to soil [Cu] curves at 60 and 90% WHC but with different intercepts for these two WHC treatments (suppl. Table 3 and Fig. 2).

The final 4 quadratic equations are as follow: Eq. (28) for 30% WHC, Eq. (29) for 60 and 90% WHC, Eq. (30) for DR, and Eq. (31) for DO. (Fig. 2).

$$280 (28) F_{Cu30} = 0.782 - 0.000451 \times Cu + 9.49 \times 10^{-8} \times Cu^2$$

$$(29) F_{Cu60/90} = b - 0.000342 \times Cu + 4.30 \times 10^{-8} \times Cu^2$$

with b= 0.795 for 60% WHC and b= 0.796for 90% WHC

$$(30) F_{CuDR} = 0.552 - 0.000164 \times Cu + 6.09 \times 10^{-8} \times Cu^2$$

$$(31) F_{CuDO} = 0.625 - 0.000192 \times Cu + 2.82 \times 10^{-8} \times Cu^2$$

285 According to the fitted equations, the decrease in nitrate production rates as a function of soil [Cu] depended on initial incubation treatment. Decreases were found steeper following 30% WHC > 60-90% WHC > DO>DR.

These 4 equations were then added in the DNDC model, allowing to adjust the Eq. (2) which regulates the nitrate production to soil Cu contents.

290 3.2. Modelling soil nitrate concentrations in Cu contaminated treatments using a DNDC-Cu model.

3.2.a. Set up of the DNDC-Cu model

The DNDC model was originally constructed to model both C and N soil cycles. The relative proportion of nitrification and denitrification processes thus depends on soil aerobic fraction determined both by soil C respiration and soil moisture (Eqs. (13) and (14)). Before any addition of Cu function in DNDC, we estimated this soil aerobic fraction using C mineralisation. Previous data from a 366 days incubations made on the same uncontaminated soil (Annabi et al., 2007) were first used to fit a C mineralisation coefficient rate, k. The resulting k coefficient ($k = 1.234 \cdot 10^{-4} \text{ gC}\cdot\text{m}^{-2}\cdot\text{30min}^{-1}$) was introduced in the DNDC model and forced to equilibrium (45 days) without soil Cu contamination effect. This provided a basal aerobic volume fraction for each soil moisture through Eq. (13), corresponding to $3.52 \cdot 10^{-3}$ at 30%, $6.167 \cdot 10^{-3}$ at 60% (and DR/DO to which bio assays were performed at 60% WHC) and $2.705 \cdot 10^{-2}$ at 90% of the WHC. The partial O₂ pressure was calculated as 211.4 hPa at 30% WHC, 210.7 hPa at 60% WHC, DR and DO and 205.4 hPa at 90% WHC. These values were used to initiate the DNDC-Cu version. We then ran the DNDC-Cu version for a 3-day simulation. The constant rate of C mineralisation, k, was adjusted to take into account the Cu contents with the Eq.(14) while the Eqs. (28)-(31) were used to adjust NO₃-N production rate (Fig. 1) to Cu.

305 3.2.b. DNDC-Cu model validation

Our DNDC-Cu model has been evaluated by comparing experimental data of soil nitrate concentration measured after 1 and 3 days of the bioassay incubation with the model outputs. A good fit was provided for 60 and 90% of WHC in the range of the DNDC calibration compared to 30% WHC where the nitrate production is largely underestimated (more than twice after 3 days of incubation, Fig. 3a). The regression slopes between modelled and measured soil [nitrate] for 60 and 90% WHC were respectively 0.94 ± 0.01 and 0.91 ± 0.01 ($R^2=0.99$ in both cases, Fig. 3a.) whereas for 30% WHC the regression slope was 1.21 ± 0.08 ($R^2=0.92$) (Fig. 3a). For DR, the soil nitrate stocks were either overestimated (at $762 \text{ mgCu}\cdot\text{kg soil}^{-1}$ soil) or underestimated (at $2012 \text{ mgCu}\cdot\text{kg soil}^{-1}$, Fig. 3b) but overall modelling adequately fitted the data with a regression slope at 0.95 ± 0.02 and $R^2=0.99$. For DO, the regression slope between modelled and measured soil nitrate stocks was 0.95 ± 0.02 too. The Supplementary Fig. 1 shows the improvement of the DNDC-Cu version to model NO₃-N soil concentration for contaminated soils with the differences between modelled and measured [NO₃-N] using the default DNDC version compared to our DNDC-Cu version for each [Cu].

Considering all the moisture treatments, RMSE was about 57.3 as a mean ($46.4 \text{ gNO}_3\text{-N}\cdot\text{m}^{-2}$ standard error) for a mean soil nitrate measured at $390 \text{ gNO}_3\text{-N}\cdot\text{m}^{-2}$ ($69 \text{ gNO}_3\text{-N}\cdot\text{m}^{-2}$ standard error) after 3 days of incubation.

320 However, for the 30% WHC, RMSE was 139.9 thus 3.7 times more than for the other treatments (Suppl. Fig. 2).
Despite the reduction in nitrate production rate from 0.20 to 0.18 gN.hour⁻¹ (see material and methods), soil nitrate
stock was still slightly overestimated in the 90% WHC as shown by the largest lack of correlation in this case
compared to the 60% WHC treatment (Fig. 3a, Suppl. Fig. 2). Lack of correlation was reduced for all tested
moisture treatments (mean \sqrt{LC} = 23.0, standard error =5.4 which is roughly 1/20 of the produced nitrate in 3
325 days in uncontaminated treatment). Results showed that our DNDC-Cu version was able to reproduce the
variability observed in Cu contaminated soils except for the 30% WHC treatment where soil nitrate stocks were
largely underestimated. The following results thus focused on the use of DNDC-Cu for DR, DO, 60 and 90% of
WHC treatments to predict soil N emissions.

330 3.3 Use of DNDC-Cu to predict N fluxes in contaminated soils.

3.3.a. Effect of soil [Cu] on soil N stocks.

The soil Cu function we included in the DNDC-Cu model modified specifically the default nitrification
equation in complement to pH, soil moisture and O₂ availability (Eq (2.)). In the presence of low [Cu] (12-512
mgCu.kg soil⁻¹), the predicted NO₃-N soil stocks were found equivalent between 60% WHC and DO and, to a less
335 extent, DR treatments (Suppl. Fig. 3). When soil [Cu] increased, soil [NO₃-N] decreased but with different rates
depending on the moisture of initial incubations (Eqs. 28-31). The evolutions of concentrations in soils and
emissions fluxes of each species in response to [Cu] gradient were also found highly different depending on the
species and on the moisture of initial incubations. However, the relative evolution in term of both soils
concentration and emissions fluxes were identical for each species and each initial incubation treatment and are
340 represented in table 2. Largest variations were modelled for N₂O-N decrease (around -63% for the constant
moisture treatments and -54% for the DR at 2012 mgCu.kg soil⁻¹) while smallest variations were modelled for
NH₄-N increase (8-10% for the 60 and 90% WHC against 5-7% for the DR and DO initially incubated soils at
2012 mgCu.kg soil⁻¹). Due to the different evolutions with Cu gradient, concentrations or intensities of fluxes for
a given specie may reversed between two moisture treatment with an increase in soil [Cu].

345 For instance, up to 548 mgCu.kg soil⁻¹, we modelled the lowest NO₃-N stocks in DR incubated soils. Above
it, NO₃-N soil stocks were the smallest for the 60% WHC treatment as a result of the sharpest decrease in NO₃-N
production due to soil [Cu]. NO₃-N soil stock for initial incubation at 90% WHC were the highest for soil [Cu]
below 1432 mgCu.kg soil⁻¹. Between 1432 and 2000 mgCu.kg soil⁻¹, NO₃-N soil stocks were similar for 90%
WHC, DR and DO (Suppl. Fig. 3).

350 In the absence of Cu, NO₃-N/NH₄-N ratios were similar among soil moisture treatments. However, the
variations in NH₄-N and NO₃-N stocks in response to Cu gradient were different across soil moistures. Indeed, the
increase in soil [Cu] resulted in a decrease in nitrification rate, thus in an increase in soil NH₄-N stocks (Suppl. Fig
4). The NO₃-N/NH₄-N stocks ratios decreased faster for 60-90% WHC than for DR and DO with an increase in
soil [Cu] (suppl Fig. 5; Table 2).

355 The decrease in soil $\text{NO}_3\text{-N}$ stocks at high [Cu], induced a decrease in the modelled growth of denitrifying
bacteria that is directly related to $[\text{NO}_3\text{-N}]$ (Eq. (13)). Consequently, the modelled denitrifying bacterial pool was
reduced when soil [Cu] increases (Fig. 4). Whatever the soil [Cu], denitrification was modelled roughly twice
larger in the soils incubated at 90% WHC than in the other treatment as this moist treatment is defined as perfect
condition for denitrifying bacteria in the DNDC model (Changsheng Li et al., 1992). Soils incubated at 60% WHC
360 were modelled with the lowest denitrifying bacterial pool. No difference between the DR and DO soils was found
due to uncertainties in the modelled denitrifying bacterial pool which resulted from the different concentrations in
N species used to initialise DNDC-Cu (Table 1). The soil $\text{N}_2\text{O-N}$ stocks and dissolved $\text{NO}_x\text{-N}$ being directly
related to denitrifying bacteria, they followed similar trends than soil $\text{NO}_3\text{-N}$ stocks with a global decrease in soil
stocks with an increase in soil [Cu] (table 2) and larger stocks at the wetter treatment.

365

3.3.b. Estimation of soil N emissions under various moistures

Large differences are predicted in the $\text{NH}_4\text{-N}$, $\text{NO}_x\text{-N}$ and $\text{N}_2\text{O-N}$ fluxes between the 90% WHC soil and the
3 other soil moisture treatments (Fig. 5). Due to the different evolutions of fluxes in response to Cu, $\text{NH}_4\text{-N}$ fluxes
were modelled smallest for the DR soils than for the 60% WHC incubated for soil Cu below $1774 \text{ mgCu.kg soil}^{-1}$
370 and higher above $1774 \text{ mgCu.kg soil}^{-1}$ (Fig. 5(a)). The emissions of $\text{NH}_4\text{-N}$ in the DO treatment were predicted to
be higher than those of the DR treatment for soil Cu higher than $1290 \text{ mgCu.kg soil}^{-1}$ and smallest below 1290
 mgCu.kg^{-1} (Fig. 5a). In the studied range of added Cu, $\text{NO}_x\text{-N}$ fluxes predicted by the model are largest from 60%
WHC to DO, DR and 90% WHC (Fig. 5(b.)) for a moderate Cu input (~ below $1380 \text{ mgCu.kg soil}^{-1}$). The decrease
in $\text{NO}_x\text{-N}$ emission with the increase in soil [Cu] was however steeper for soils incubated at 60% WHC (Tables 2
375 (a.) and 2(b.)). Hence, at $2012 \text{ mgCu.kg soil}^{-1}$ $\text{NO}_x\text{-N}$ fluxes in soil incubated at 60% WHC were similar to those
in the soils incubated under drought treatment (Fig. 5 (b.)). The smallest fluxes of $\text{N}_2\text{O-N}$ were predicted for the
wetter treatment despite higher modelled $\text{N}_2\text{O-N}$ stocks at 90% WHC whatever the [Cu] (Table 2 (a.) and Fig 5
(c.)). The $\text{N}_2\text{O-N}$ emissions fluxes in the presence of Cu were predicted to be 4 times smallest in the 90% WHC
treatment compared to the others. $\text{N}_2\text{O-N}$ fluxes had similar trends than $\text{NO}_x\text{-N}$ for moderate Cu inputs but fluxes
380 were still largest from 60% WHC to DO, DR and 90% WHC (Fig. 5 (c.)), and $\text{N}_2\text{-N}$ emissions were larger at the
wettest treatment (Fig. 5 (d.)). The ratio of emitted $\text{N}_2\text{O-N}$ per denitrification products (i.e. $\text{N}_2\text{O-N} / \text{N}_2\text{O-N} + \text{N}_2\text{-N}$)
was hence smallest in the moistest soils, which means that the largest soils $\text{N}_2\text{O-N}$ stocks in the case of 90%
WHC had more chance to be transformed rather than emitted (Fig. 6).

385 4 DISCUSSION

4.1. From laboratory experiment to soil N emission modelling

Thanks to our laboratory experiments, we were able to define a function modulating the soil $\text{NO}_3\text{-N}$ production
rates in relation with soil [Cu] and depending on soil moisture. Our results showed that soil nitrate decreases with
an increase in soil [Cu]. Initial incubation treatment significantly affects the response of soil nitrate production rate
390 to subsequent Cu stress with steeper decrease in the order 30% WHC > 60-90% WHC > DO > DR for the Cu
range studied. The lowest sensitivity of Cu in soils initially incubated with dry-rewet events suggest that it might

have selected more resistant communities (Barnard et al., 2013; Gleeson et al., 2008). More complex dose response functions have been used in (Sereni et al., 2022) to assess thresholds and loss of functions after such a double stress. These results are in relatively good agreement with those presented here using the quadratic fit, especially for the highest half of [Cu]. However, they also presented a limited increase in nitrification rate for small Cu input that we weren't able to emphasize in the present study. In the present article we used simple functions of fit to describe the response of soil nitrate production to Cu gradient after the first moisture stress as they further have to be included in the DNDC model. After implementing these quadratics Cu modulating functions into the DNDC-Cu model, we were however able to reproduce the observed soil nitrate stock particularly for the soils incubated at 60 and 90% of WHC. The variability around the mean due to the Cu effect was also reproduced by our DNDC-Cu version at 30% of WHC despite strong underestimation of mean soil nitrate stocks due to model moisture-limit (Changsheng Li et al., 1992). In the case of the DR and DO incubated soils, the so-called "Cu function" also accounted for the effect of drought stress. In fact, our Cu functions were defined on the basis of soil nitrate production against the whole gradient of Cu thus also considering the control without Cu. However, the double stress effect was also well reproduced in nitrate production.

4.2. Expected ecological implications of soil Cu contamination

Based on nitrate production measurements, we modelled a decrease in denitrifying activities with an increase in soil [Cu] as a consequence of the decrease in soil nitrate stocks. However, the experiments performed here did not allow us to determine if the soil Cu contamination rather affects nitrifying bacteria (e.g. decrease in nitrifying activity and in $\text{NO}_3\text{-N}$ production) or denitrifying bacteria (e.g. increase in denitrifying activities and $\text{NO}_3\text{-N}$ consumption). The effect of soil contamination on $\text{N}_2\text{O-N}$ production is debated because i) microbial species involved are not clearly identified (Wrage-Mönning et al., 2018), ii) species richness is not necessarily related to soil functions (Ruyters et al., 2013) and iii) denitrifying communities could be differently sensitive than the nitrifying to soil contamination (Hund-Rinke and Simon, 2008; Vásquez-Murrieta et al., 2006). Also, our approach to model $\text{N}_2\text{O-N}$, $\text{N}_2\text{-N}$ and $\text{NO}_x\text{-N}$ production in the contaminated context could have been more constrained with measurement of denitrification rate to assess the effect of Cu on proportion of production and consumption of $\text{NO}_3\text{-N}$.

Based on our simulations, the soil Cu contamination was expected to substantially modify the proportion of available N in soils with the increase in $\text{NH}_4\text{-N}$ stock at the expense of $\text{NO}_3\text{-N}$. $\text{NH}_4\text{-N}$ accumulation and the large expected decrease in $\text{NO}_3\text{-N}/\text{NH}_4\text{-N}$ ratio in contaminated soils (around 50% for the 60% WHC) may lead to shift in plant community structures with different preferences in N assimilation (Cui and Song, 2007; Peacock et al., 2001). Therefore, Cu stress could not only have implications in microbial community patterns as a stressor, but could also induce further shifts due to N species redistributions in soils.

4.3. From $\text{N}_2\text{O-N}$, $\text{N}_2\text{-N}$ and $\text{NO}_x\text{-N}$ soil stocks to emissions

In the present study, we predicted highest soil $\text{N}_2\text{-N}$, $\text{N}_2\text{O-N}$ and $\text{NO}_x\text{-N}$ stocks in the moistest treatments. Indeed this species are produced by the denitrifying bacteria expected to behave optimally at 90% WHC or after DR cycles (Changsheng Li et al., 1992; Homyak et al., 2017). However, $\text{N}_2\text{O-N}$ and $\text{NO}_x\text{-N}$ emissions were

430 modelled higher in the driest soils, whereas numerous studies (Dobbie and Smith 2003; Xiong et al. 2007; Manzoni
et al. 2012) reported high measured N₂O-N emissions with high moisture. In the present version of DNDC-Cu, the
soil N emissions were directly controlled by their diffusion in soil, calculated on the basis of clay and soil moisture
content. The diffusion of each species would hence be 11 times smaller under 90% WHC ($D_s=0.00357$) than
435 under the 60% WHC treatment ($D_s=0.0306$) because the model described the diffusion as a whole and do not
separated pores with or without water. Diffusion was hence slower in the water than in the air. Thus, the weighted
mean diffusion was lower in the high moisture treatment. Without Cu soil nitrous stocks being roughly 1.6 times
and soils N₂-N stocks 11.1 larger under 90% WHC treatment than the other, the emission of N₂O-N were larger
under driest treatment even if stocks were smaller.

Several studies also reported flushing event with Dry-Rewet cycles which would enhance C mineralization,
440 known as the Birch effect (Birch, 1958; Göransson et al., 2013), hence reducing soil O₂ concentration. Moreover,
soil [O₂] is closely related to the pore size distribution, being of major importance in nitrification/denitrification
control (Khalil et al., 2004) with a dominating nitrification for aggregates up to 0.25cm (Kremen et al., 2005). Pore
size distribution under dry/rewet events is controlled by cracking, (des)aggregation (Cosentino et al., 2006; Deneff
et al., 2001) or gas displacement (Kemper et al., 1985) that we weren't able to take into account in the present
445 study. In DNDC, the calculation of denitrification rate and diffusion was based on a rough description of anaerobic
zone with approximation of soil pore space distribution (Blagodatsky et al., 2011; Li et al., 2000). The soil pore
space distribution approach has been demonstrated to be more generally applicable (Arah and Vinten 1995;
Schurgers et al. 2006) whereas soil aggregates have been shown to control the extend of nitrification and
denitrification (Kremen et al., 2005; Schlüter et al., 2018). However, if models have been proposed to take O₂
450 availability at the aggregate size into account in the nitrous oxide production (Kremen et al., 2005; Leffelaar, 1988),
they also point out the difficulty in parametrization which need a large panel of soil measurements. Moreover, they
are rarely transposable at the meso-and regional scale due to high spatial variations in soil structure (Butterbach-
Bahl et al., 2013). The DNDC-Cu version we used here particularly pointed out the difficulty in dealing on
biogeochemistry model with physical processes, with large discrepancies between modelled soils stocks and
455 emissions. The validation we performed focused on soil nitrates stocks and a second step to go further on would
be the measure of gaseous species to ensure that emissions were also impacted by soils treatment. Moreover, we
assumed here that soil [Cu] affected the C mineralisation with a decrease in soil O₂ production leading to an
increase in denitrification and N₂O-N, NO_x-N. Nevertheless, the present DNDC-Cu version didn't take into
account the retroaction between C and N cycles. Further research would thus be required to include Cu
460 contamination into C and N interacting cycles.

4.4. Climate change could substantially modify contaminated soil N emission

It is well known that climate change and rainfall patterns could substantially modify the soil N balance and
its GHG emissions (Galloway et al. 2003, 2008; Butterbach-Bahl et al. 2013). Despite limitation in DNDC
accuracy for nitrous emissions (Foltz et al., 2019), our results tend to showed that increased Cu contamination as
465 well might affect soil N emissions with smallest emissions of NO_x-N and N₂O-N. These two gases are of major
importance in GHG mitigation with a warming potential per mass 300 and 40 times greater than CO₂, respectively.
Agricultural soils being the dominating source of N₂O-N (Beauchamp, 1997; Signor and Cerri, 2013), even a
limited decrease in their emissions could have major implication for climate. Based on our modelling, the join

effect of soil moisture and [Cu] was particularly important with larger differences in N₂O-N and NO_x-N emissions
470 between rainfall patterns at high [Cu] (3.3.b.). Sereni et al., (2022) also showed that soil Cu contamination
differently affect soils nitrification depending of primary soil moisture stress. Here we showed that the N₂O-N and
NO_x-N emission variations are significantly more sensitive to the combined effect of Cu and precipitation regime
than the nitrate stock. Based on these results, soil Cu inputs on moistest soils would lead to a largest decrease in
475 soil N₂O-N and NO_x-N emission compared to that on driest soils, and even more than on soils submitted to abrupt
and intense shifts in rainfall patterns as the DR and DO soils.

5. CONCLUSION

In the present study, we aimed at combining ecotoxicological experiments and biogeochemical modelling
focusing on the effect of soil Cu contamination on soil N emission under different soil moisture treatments constant
480 moisture (30, 60 or 90% WHC) or to a single long drought period (DO) or several Dry-Rewet cycle (DR). Based
on a 3-day bioassay measuring soil NO₃-N over time, we were able to adjust the DNDC model to take into account
the Cu effect on soil N emission. The DNDC-Cu version we proposed was able to reproduce the observed Cu
effect on soil nitrate stock with R²>0.99 and RMSE<10% for all treatments in the DNDC calibration range (>40%
WHC).

485 We modelled a Cu effect inducing a decrease in denitrifying bacterial pool leading to an increase in NH₄-N
soil stocks at the expense of NO₃-N, N₂O-N and NO_x-N stocks. We showed that the effect of soil Cu contamination
was different among moisture treatment and N species. For instance, we modelled that the largest [Cu] (2012mg
Cu.kg soil⁻¹) provoked a decrease in soil nitrate stocks from -28% in the DR case to -44% in the 60% WHC whereas
N₂O-N emissions were expected to decrease up to 63% in the 90% WHC (-62% in the 60% WHC case, -54% in
490 the DO case). However, our results tended to show that the amount of N₂O-N emitted from denitrification would
decrease with an increase in soil [Cu] and from 60% WHC to DR, DO and 90% WHC, so that less N₂O-N produced
would be converted to N₂-N. This result points out two main difficulties in biogeochemical modelling: i) the
difficulty to take into account hydrological dynamics (produced NO₃-N and NH₄-N could be expected to leach)
and soil structures at different spatial scale (denitrification is estimated based on rough estimation on anaerobic
495 soil volume which also controlled emissions rates through diffusion processes) and ii) linking soil function to
microbial dynamics, in particular in this case were only the NO₃-N stock was measured (without dealing between
production and consumption for instance). Despite these two main points of uncertainty, the combination of
incubations and of modelisation we conducted here emphasize the need to account for soil contamination when
dealing with soil GHG emission modelling and climate change, as both contamination and rainfall patterns affect
500 in a different way the soil NO_x-N and N₂O-N emissions.

Data availability: Data are available at: <https://doi.org/10.15454/ZUKN90>

Authors Contributions: the authors contributed as follows:

Laura Sereni: Methodology, formal analysis, data processing, writing original draft.

- 505 Bertrand Guenet: Methodology, conceptualization, writing review and editing, supervision
Charlotte Blasi: Experimentations and draft initialization
Olivier Crouzet: Methodology, conceptualization, writing review and editing, supervision
Jean-Christophe Lata: writing review and editing, supervision
Isabelle Lamy: Methodology conceptualization, writing review and editing, supervision project administration,
510 funding acquisition.
Competing Interests: authors have no competing interests to declare.

Acknowledgments

- 515 Parts of this study were financially supported by the French National Research Agency ANR CESA-13-0016-01 through the CEMABS project and the Labex BASC through the Connexion project. LS thanks the Ecole Normale Supérieure (ENS) for funding her PhD. The authors thank Amélie Trouvé for her help in soil data analysis, Sébastien Breuil for soil processing and Christelle Marraud for bioassay design.

References

- 520 Anav, a., Friedlingstein, P., Kidston, M., Bopp, L., Ciais, P., Cox, P., Jones, C., Jung, M., Myneni, R. and Zhu, Z.: Evaluating the Land and Ocean Components of the Global Carbon Cycle in the Cmp5 Earth System Models, *J. Clim.*, 130401082723008, doi:10.1175/JCLI-D-12-00417.1, 2013.
- Annabi, M., Houot, S., Francou, C., Poitrenaud, M. and Le Bissonnais, Y.: Soil Aggregate Stability Improvement with Urban Composts of Different Maturities, *Soil Sci. Soc. Am. J.*, 71(2), 413–423,
525 doi:10.2136/sssaj2006.0161, 2007.
- Arah, J. R. M. And Vinten, A. J. A.: Simplified Models Of Anoxia And Denitrification In Aggregated And Simple - Structured Soils, *Eur. J. Soil Sci.*, 46(4), 507–517, doi:10.1111/j.1365-2389.1995.tb01347.x, 1995.
- Bååth, E.: Effects of heavy metals in soil on microbial processes and populations (a review), *Water. Air. Soil Pollut.*, 47(3–4), 335–379, doi:10.1007/BF00279331, 1989.
- 530 Barnard, R. L., Osborne, C. A. and Firestone, M. K.: Responses of soil bacterial and fungal communities to extreme desiccation and rewetting, *ISME J.*, 7(11), 2229–2241, doi:10.1038/ismej.2013.104, 2013.
- Bateman, E. J. and Baggs, E. M.: Contributions of nitrification and denitrification to N₂O emissions from soils at different water-filled pore space, *Biol. Fertil. Soils*, 41(6), 379–388, doi:10.1007/s00374-005-0858-3, 2005.
- Beauchamp, E. G.: Nitrous oxide emission from agricultural soils, *Can. J. Soil Sci.*, 77, 113–123,
535 doi:10.4141/S96-101, 1997.
- Bech, J., Poschenrieder, C., Llugany, M., Barceló, J., Tume, P., Tobias, F. J., Barranzuela, J. L. and Vásquez, E.

- R.: Arsenic and heavy metal contamination of soil and vegetation around a copper mine in Northern Peru, *Sci. Total Environ.*, 203(1), 83–91, doi:10.1016/S0048-9697(97)00136-8, 1997.
- Birch, H. F.: The effect of soil drying on humus decomposition and nitrogen availability, *Plant Soil*,
540 doi:10.1007/BF01343734, 1958.
- Blagodatsky, S., Grote, R., Kiese, R., Werner, C. and Butterbach-Bahl, K.: Modelling of microbial carbon and nitrogen turnover in soil with special emphasis on N-trace gases emission., 2011.
- Borken, W. and Matzner, E.: Reappraisal of drying and wetting effects on C and N mineralization and fluxes in soils, *Glob. Chang. Biol.*, 15(4), 808–824, doi:10.1111/j.1365-2486.2008.01681.x, 2009.
- 545 Brandt, K. K., Frandsen, R. J. N., Holm, P. E. and Nybroe, O.: Development of pollution-induced community tolerance is linked to structural and functional resilience of a soil bacterial community following a five-year field exposure to copper, *Soil Biol. Biochem.*, 42(5), 748–757, doi:10.1016/j.soilbio.2010.01.008, 2010.
- Broos, K., Warne, M. S. J., Heemsbergen, D. A., Stevens, D., Barnes, M. B., Correll, R. L. and McLaughlin, M. J.: Soil factors controlling the toxicity of copper and zinc to microbial processes in Australian soils, *Environ. Toxicol. Chem.*, 26(4), 583–590, doi:10.1897/06-302R.1, 2007.
- 550 Butterbach-Bahl, K., Kahl, M., Mykhayliv, L., Werner, C., Kiese, R. and Li, C.: A European-wide inventory of soil NO emissions using the biogeochemical models DNDC/Forest-DNDC, *Atmos. Environ.*, 43(7), 1392–1402, doi:10.1016/j.atmosenv.2008.02.008, 2009.
- Butterbach-Bahl, K., Baggs, E. M., Dannenmann, M., Kiese, R. and Zechmeister-Boltenstern, S.: Nitrous oxide emissions from soils: How well do we understand the processes and their controls?, *Philos. Trans. R. Soc. B Biol. Sci.*, 368(1621), doi:10.1098/rstb.2013.0122, 2013.
- 555 Cambier, P., Michaud, A., Paradelo, R., Germain, M., Mercier, V., Guérin-Lebourg, A., Revallier, A. and Houot, S.: Trace metal availability in soil horizons amended with various urban waste composts during 17 years – Monitoring and modelling, *Sci. Total Environ.*, 651, 2961–2974, doi:10.1016/j.scitotenv.2018.10.013, 2019.
- 560 Changsheng Li, Frolking, S. and Frolking, T. A.: A model of nitrous oxide evolution from soil driven by rainfall events: 1. Model structure and sensitivity, *J. Geophys. Res.*, 97(D9), 9759–9776, doi:10.1029/92jd00509, 1992.
- Conrad, R.: Soil microorganisms as controllers of atmospheric trace gases (H₂, CO, CH₄, OCS, N₂O, and NO), *Microbiol. Rev.*, 60(4), 609–640, doi:10.1128/membr.60.4.609-640.1996, 1996.
- Cosentino, D., Chenu, C. and Le Bissonnais, Y.: Aggregate stability and microbial community dynamics under drying-wetting cycles in a silt loam soil, *Soil Biol. Biochem.*, 38(8), 2053–2062,
565 doi:10.1016/j.soilbio.2005.12.022, 2006.
- Cui, X. and Song, J.: Soil NH₄⁺/NO₃⁻ nitrogen characteristics in primary forests and the adaptability of some coniferous species, *Front. For. China*, 2, 1–10, doi:10.1007/s11461-007-0001-8, 2007.
- Denef, K., Six, J., Bossuyt, H., Frey, S. D., Elliott, E. T., Merckx, R. and Paustian, K.: Influence of dry-wet

- 570 cycles on the interrelationship between aggregate, particulate organic matter, and microbial community dynamics, *Soil Biol. Biochem.*, 33, 1599–1611, doi:10.1016/S0038-0717(01)00076-1, 2001.
- Dobbie, K. E. and Smith, K. A.: Nitrous oxide emission factors for agricultural soils in Great Britain: The impact of soil water-filled pore space and other controlling variables, *Glob. Chang. Biol.*, 9(2), 204–218, doi:10.1046/j.1365-2486.2003.00563.x, 2003.
- 575 FAO: Soil Pollution a hidden reality., 2008.
- Fierer, N., Schimel, J. P. and Holden, P. A.: Influence of drying-rewetting frequency on soil bacterial community structure, *Microb. Ecol.*, 45(1), 63–71, doi:10.1007/s00248-002-1007-2, 2003.
- Foltz, M. E., Zilles, J. L. and Koloutsou-Vakakis, S.: Prediction of N₂O emissions under different field management practices and climate conditions, *Sci. Total Environ.*, 646, 872–879, doi:10.1016/j.scitotenv.2018.07.364, 2019.
- 580 Galloway, J. N., Townsend, A. R., Erisman, J. W., Bekunda, M., Cai, Z., Freney, J. R., Martinelli, L. A., Seitzinger, S. P. and Sutton, M. A.: Transformation of the nitrogen cycle: Recent trends, questions, and potential solutions, *Science* (80-.), 320(5878), 889–892, doi:10.1126/science.1136674, 2008.
- Galloway, J. N., Aber, J. D., Erisman, J. W., Seitzinger, S. P., Howarth, R. W., Cowling, E. B. And Cosby, B. J.: The Nitrogen Cascade, *Bioscience*, 53(4), 341–356, doi:10.1641/0006-3568(2003)053[0341:tnc]2.0.co;2, 2003.
- 585 Gauch, H. G., Hwang, J. T. G. and Fick, G. W.: Model Evaluation by Comparison of Model-Based Predictions and Measured Values, *Agron. J.*, 95(6), 1442–1446, doi:10.2134/agronj2003.1442, 2003.
- Giller, K. E., Witter, E. and McGrath, S. P.: Heavy metals and soil microbes, *Soil Biol. Biochem.*, 41(10), 2031–2037, doi:10.1016/j.soilbio.2009.04.026, 2009.
- 590 Giltrap, D. L., Li, C. and Sagggar, S.: DNDC: A process-based model of greenhouse gas fluxes from agricultural soils, *Agric. Ecosyst. Environ.*, 136(3–4), 292–300, doi:10.1016/j.agee.2009.06.014, 2010.
- Gleeson, D. B., Herrmann, A. M., Livesley, S. J. and Murphy, D. V.: Influence of water potential on nitrification and structure of nitrifying bacterial communities in semiarid soils, *Appl. Soil Ecol.*, 40(1), 189–194, doi:10.1016/j.apsoil.2008.02.005, 2008.
- 595 Göransson, H., Godbold, D. L., Jones, D. L. and Rousk, J.: Bacterial growth and respiration responses upon rewetting dry forest soils: Impact of drought-legacy, *Soil Biol. Biochem.*, 57, 477–486, doi:10.1016/j.soilbio.2012.08.031, 2013.
- Guo, X., Drury, C. F., Yang, X., Daniel Reynolds, W. and Fan, R.: The Extent of Soil Drying and Rewetting Affects Nitrous Oxide Emissions, Denitrification, and Nitrogen Mineralization, *Soil Sci. Soc. Am. J.*, 78(1), 194–204, doi:10.2136/sssaj2013.06.0219, 2014.
- 600 Homyak, P. M., Allison, S. D., Huxman, T. E., Goulden, M. L. and Treseder, K. K.: Effects of Drought Manipulation on Soil Nitrogen Cycling: A Meta-Analysis, *J. Geophys. Res. Biogeosciences*, 122(12), 3260–

3272, doi:10.1002/2017JG004146, 2017.

605 Hund-Rinke, K. and Simon, M.: Bioavailability assessment of contaminants in soils via respiration and nitrification tests, *Environ. Pollut.*, 153(2), 468–475, doi:10.1016/j.envpol.2007.08.003, 2008.

IPCC: IPCC Special Report on Climate Change and Land: Chapter 4: Land Degradation, , 1–112 [online] Available from: <https://www.ipcc.ch/report/srcccl/>, 2019.

Kemper, W. D., Rosenau, R. and Nelson, S.: Gas Displacement and Aggregate Stability of Soils, *Soil Sci. Soc. Am. J.*, 49, 25–28, doi:10.2136/sssaj1985.03615995004900010004x, 1985.

610 Kesik, M., Ambus, P., Baritz, R., Brüggemann, N., Butterbach-Bahl, K., Damm, M., Duyzer, J., Horváth, L., Kiese, R., Kitzler, B., Leip, A., Li, C., Pihlatie, M., Pilegaard, K., Seufert, G., Simpson, D., Skiba, U., Smiatek, G., Vesala, T. and Zechmeister-Boltenstern, S.: Inventories of N₂O and NO emissions from European forest soils, *Biogeosciences*, 2(4), 353–373, doi:10.5194/bg-2-353-2005, 2005.

615 Khalil, K., Mary, B. and Renault, P.: Nitrous oxide production by nitrification and denitrification in soil aggregates as affected by O₂ concentration, *Soil Biol. Biochem.*, 36, 687–699, doi:10.1016/j.soilbio.2004.01.004, 2004.

Khan, S., Cao, Q., Zheng, Y. M., Huang, Y. Z. and Zhu, Y. G.: Health risks of heavy metals in contaminated soils and food crops irrigated with wastewater in Beijing, China, *Environ. Pollut.*, 152(3), 686–692, doi:10.1016/j.envpol.2007.06.056, 2008.

620 Knutti, R. and Sedláček, J.: Robustness and uncertainties in the new CMIP5 climate model projections, *Nat. Clim. Chang.*, 3(4), 369–373, doi:10.1038/nclimate1716, 2012.

Kremen, A., Bear, J., Shavit, U. and Shaviv, A.: Model demonstrating the potential for coupled nitrification denitrification in soil aggregates, *Environ. Sci. Technol.*, 39(11), 4180–4188, doi:10.1021/es048304z, 2005.

625 Lado, L. R., Hengl, T. and Reuter, H. I.: Heavy metals in European soils: A geostatistical analysis of the FOREGS Geochemical database, *Geoderma*, 148(2), 189–199, doi:10.1016/j.geoderma.2008.09.020, 2008.

Leffelaar, P. A.: Dynamics of partial anaerobiosis, denitrification, and water in a soil aggregate: Simulation, *Soil Sci.*, 146(6), 427–444, doi:10.1097/00010694-198812000-00004, 1988.

630 Li, C., Aber, J., Stange, F., Butterbach-Bahl, K. and Papen, H.: A process-oriented model of N₂O and NO emissions from forest soils: 1. Model development, *J. Geophys. Res. Atmos.*, 105(D4), 4369–4384, doi:10.1029/1999JD900949, 2000.

Manzoni, S., Schimel, J. P. and Porporato, A.: Responses of soil microbial communities to water stress: Results from a meta-analysis, *Ecology*, 93(4), 930–938, doi:10.1890/11-0026.1, 2012.

635 Mertens, J., Wakelin, S. A., Broos, K., McLaughlin, M. J. and Smolders, E.: Extent of copper tolerance and consequences for functional stability of the ammonia-oxidizing community in long-term copper-contaminated Soils, *Environ. Toxicol. Chem.*, 29(1), 27–37, doi:10.1002/etc.16, 2010.

- Miranda, K. M., Espey, M. G. and Wink, D. A.: A rapid, simple spectrophotometric method for simultaneous detection of nitrate and nitrite, *Nitric Oxide - Biol. Chem.*, 5(1), 62–71, doi:10.1006/niox.2000.0319, 2001.
- Moyano, F. E., Manzoni, S. and Chenu, C.: Responses of soil heterotrophic respiration to moisture availability: An exploration of processes and models, *Soil Biol. Biochem.*, 59, 72–85, doi:10.1016/j.soilbio.2013.01.002, 640 2013.
- Nicholson, F. A., Smith, S. R., Alloway, B. J., Carlton-Smith, C. and Chambers, B. J.: An inventory of heavy metals inputs to agricultural soils in England and Wales, *Sci. Total Environ.*, 311(1–3), 205–219, doi:10.1016/S0048-9697(03)00139-6, 2003.
- Oorts, K., Bronckaers, H. and Smolders, E.: Discrepancy of the microbial response to elevated copper between freshly spiked and long-term contaminated soils, in *Environmental Toxicology and Chemistry*, vol. 25, pp. 845–853., 2006. 645
- Peacock, A. D., Mullen, M. D., Ringelberg, D. B., Tyler, D. D., Hedrick, D. B., Gale, P. M. and White, D. C.: Soil microbial community responses to dairy manure or ammonium nitrate applications, *Soil Biol. Biochem.*, 33, 1011–1019, doi:10.1016/S0038-0717(01)00004-9, 2001.
- Petersen, D. G., Blazewicz, S. J., Firestone, M., Herman, D. J., Turetsky, M. and Waldrop, M.: Abundance of microbial genes associated with nitrogen cycling as indices of biogeochemical process rates across a vegetation gradient in Alaska, *Environ. Microbiol.*, 14(4), 993–1008, doi:10.1111/j.1462-2920.2011.02679.x, 2012. 650
- Rillig, M. C., Ryo, M., Lehmann, A., Aguilar-Trigueros, C. A., Buchert, S., Wulf, A., Iwasaki, A., Roy, J. and Yang, G.: The role of multiple global change factors in driving soil functions and microbial biodiversity, *Science* (80-.), 366(6467), 886–890, doi:10.1126/science.aay2832, 2019. 655
- Ruyters, S., Springael, D. and Smolders, E.: Recovery of Soil Ammonia Oxidation After Long-Term Zinc Exposure Is Not Related to the Richness of the Bacterial Nitrifying Community, *Microb. Ecol.*, 66(2), 312–321, doi:10.1007/s00248-013-0210-7, 2013.
- Sauvé, S., Dumestre, A., McBride, M., Gillett, J. W., Berthelin, J. and Hendershot, W.: Nitrification potential in field-collected soils contaminated with Pb or Cu, *Appl. Soil Ecol.*, 12(1), 29–39, doi:10.1016/S0929-1393(98)00166-8, 1999. 660
- Schimel, J. P.: Life in Dry Soils: Effects of Drought on Soil Microbial Communities and Processes, *Annu. Rev. Ecol. Evol. Syst.*, 49(1), 409–432, doi:10.1146/annurev-ecolsys-110617-062614, 2018.
- Schlüter, S., Henjes, S., Zawallich, J., Bergaust, L., Horn, M., Ippisch, O., Vogel, H. J. and Dörsch, P.: Denitrification in soil aggregate analogues-effect of aggregate size and oxygen diffusion, *Front. Environ. Sci.*, 6(17), doi:10.3389/fenvs.2018.00017, 2018. 665
- Schurgers, G., Dörsch, P., Bakken, L., Leffelaar, P. and Haugen, L. E.: Modelling soil anaerobiosis from water retention characteristics and soil respiration, *Soil Biol. Biochem.*, 38(9), 2637–2644, doi:10.1016/j.soilbio.2006.04.016, 2006.

- 670 Sereni, L., Guenet, B. and Lamy, I.: Does Copper Contamination Affect Soil CO₂ Emissions? A Literature Review, *Front. Environ. Sci.*, 9, 29, doi:10.3389/fenvs.2021.585677, 2021.
- Sereni, L., Guenet, B., Crouzet, O. and Lamy, I.: Responses of Soil Nitrification Activities To Copper After A Moisture Stress, *Environ. Sci. Pollut. Res.*, 1–18, 2022.
- Signor, D. and Cerri, C. E. P.: Nitrous oxide emissions in agricultural soils: a review, *Pesqui. Agropecuária Trop.*, 43(3), 322–338, doi:10.1590/s1983-40632013000300014, 2013.
- 675 Smolders, E., Oorts, K., Van Sprang, P., Schoeters, I., Janssen, C. R., McGrath, S. P. and McLaughlin, M. J.: Toxicity of trace metals in soil as affected by soil type and aging after contamination: Using calibrated bioavailability models to set ecological soil standards, *Environ. Toxicol. Chem.*, 28(8), 1633–1642, doi:10.1897/08-592.1, 2009.
- 680 Stark, J. M. and Firestone, M. K.: Mechanisms for soil moisture effects on activity of nitrifying bacteria, *Appl. Environ. Microbiol.*, 61(1), 218–221, doi:10.1128/aem.61.1.218-221.1995, 1995.
- Steinnes, E., Allen, R. O., Petersen, H. M., Rambøll, J. P. and Varskog, P.: Evidence of large scale heavy-metal contamination of natural surface soils in Norway from long-range atmospheric transport, *Sci. Total Environ.*, 205(2–3), 255–266, doi:10.1016/S0048-9697(97)00209-X, 1997.
- 685 Vásquez-Murrieta, M. S., Cruz-Mondragón, C., Trujillo-Tapia, N., Herrera-Arreola, G., Govaerts, B., Van Cleemput, O. and Dendooven, L.: Nitrous oxide production of heavy metal contaminated soil, *Soil Biol. Biochem.*, 38(5), 931–940, doi:10.1016/j.soilbio.2005.08.007, 2006.
- De Vleeschouwer, F., Gérard, L., Goormaghtigh, C., Mattielli, N., Le Roux, G. and Fagel, N.: Atmospheric lead and heavy metal pollution records from a Belgian peat bog spanning the last two millenia: Human impact on a regional to global scale, *Sci. Total Environ.*, 377(2–3), 282–295, doi:10.1016/j.scitotenv.2007.02.017, 2007.
- 690 Vuichard, N., Messina, P., Luyssaert, S., Guenet, B., Zaehle, S., Ghattas, J., Bastrikov, V. and Peylin, P.: Accounting for Carbon and Nitrogen interactions in the Global Terrestrial Ecosystem Model ORCHIDEE (trunk version, rev 4999): multi-scale evaluation of gross primary production, *Geosci. Model Dev. Discuss.*, 12(11), 4751–4779, doi:10.5194/gmd-2018-261, 2019.
- 695 Wrage-Mönnig, N., Horn, M. A., Well, R., Müller, C., Velthof, G. and Oenema, O.: The role of nitrifier denitrification in the production of nitrous oxide revisited, *Soil Biol. Biochem.*, 123(April), A3–A16, doi:10.1016/j.soilbio.2018.03.020, 2018.
- Wuana, R. A. and Okieimen, F. E.: Heavy Metals in Contaminated Soils: A Review of Sources, Chemistry, Risks and Best Available Strategies for Remediation, *ISRN Ecol.*, 2011(February 2015), 1–20, doi:10.5402/2011/402647, 2011.
- 700 Xiong, Z.-Q., Xing, G.-X. And Zhu, Z.-L.: Nitrous Oxide And Methane Emissions As Affected By Water, Soil And Nitrogen, *Pedosphere*, 17(2), 146–155, Doi:10.1016/s1002-0160(07)60020-4, 2007.

705 Zaehle, S. and Friend, A.: Carbon and nitrogen cycle dynamics in the O-CN land surface model: 1. Model description, site-scale evaluation, and sensitivity to parameter estimates, *Global Biogeochem. Cycles*, 24(1), doi:10.1029/2009GB003521, 2010.

Zandalinas, S. I., Fritschi, F. B. and Mittler, R.: Global Warming, Climate Change, and Environmental Pollution: Recipe for a Multifactorial Stress Combination Disaster, *Trends Plant Sci.*, xx(xx), 1–12, doi:10.1016/j.tplants.2021.02.011, 2021.

710 Zechmeister-Boltenstern, S., Schaufler, G. and Kitzler, B.: NO, NO₂, N₂O, CO₂ and CH₄ fluxes from soils under different land use: temperature sensitivity and effects of soil moisture, *Geophys. Res. Abstr.*, 2007.

Zhao, J. H.: Gap: Genetic analysis package, *J. Stat. Softw.*, 23(8), 1–18, doi:10.18637/jss.v023.i08, 2007.

715 Table 1: N species measured in the soils at the end of initial incubation period further used to initialise the DNDC model, mean modelled NO₃-N stocks and mean emissions of NH₄-N, N₂-N, N₂O-N, NO_x-N modelled without Cu. 90 = 90% WHC; 60= 60% WHC; DO =Dry-Only; DR =Dry Rewet treatment during initial incubation. A, B and C are replicates.

| Ech | Measured (µg.g soil ⁻¹) | | | Modelled (gN.m ⁻² h ⁻¹ for emissions, gN.m ⁻² for stocks) | | | | | |
|------|-------------------------------------|--------------------|--------------------|--|-----------------------------|------------------------------|------------------------------|---------------------------|---|
| | NH ₄ -N | NO ₂ -N | NO ₃ -N | NH ₄ -N emissions | N ₂ -N emissions | N ₂ O-N emissions | NO _x -N emissions | NO ₃ -N stocks | NO ₃ -N /NH ₄ -N stocks |
| 30_A | 4.3 | 0.1 | 15.3 | NA | NA | NA | NA | NA | NA |
| 30_B | 4.0 | 0.2 | 14.4 | | | | | | |
| 30_C | 4.5 | 0.2 | 14.3 | | | | | | |
| 60_A | 6.9 | 0.1 | 18.8 | 2.28 .10 ⁻¹⁰ | 2.26 .10 ⁻⁷ | 1.3 .10 ⁻⁴ | 1.3 .10 ⁻³ | 456.3 | 0.21 |
| 60_B | 6.9 | 0.2 | 18.8 | | | | | | |
| 60_C | 6.7 | 0.2 | 18.7 | | | | | | |
| 90_A | 8.2 | 0.2 | 23.6 | 2.64 .10 ⁻¹¹ | 6.21 .10 ⁻⁷ | 2.7 .10 ⁻⁵ | 2.7 .10 ⁻⁴ | 509.8 | 0.24 |
| 90_B | 12.6 | 0.9 | 24.0 | | | | | | |
| 90_C | 8.8 | 0.2 | 24.2 | | | | | | |
| DO_A | 5.4 | 0.2 | 26.1 | 2.35 .10 ⁻¹⁰ | 4.3 .10 ⁻⁷ | 1.1 .10 ⁻⁴ | 1.1 .10 ⁻³ | 432.0 | 0.19 |
| DO_B | 5.9 | 0.3 | 29.8 | | | | | | |
| DO_C | 7.4 | 0.9 | 26.4 | | | | | | |
| DR_A | 3.7 | 0.2 | 28.4 | 2.36.10 ⁻¹⁰ | 3.72 .10 ⁻⁷ | 9.4 .10 ⁻⁵ | 1.1 .10 ⁻³ | 454.5 | 0.21 |
| DR_B | 3.4 | 0.2 | 29.8 | | | | | | |
| DR_C | 5.0 | 0.3 | 29.9 | | | | | | |


720 Table 2: Percentage of variation in soil NO₃-N stocks, soil NO₃-N /NH₄-N stocks, NH₄-N, N₂-N, NO_x-N and N₂O-N emissions in response to soil [Cu] in the various initial incubation treatments for a 3-day modelisation.

a.


| Moisture treatment | Added Cu (mgCu.kg soil ⁻¹) | NO ₃ -N soils stocks | Emission NH ₄ -N | Emission N ₂ -N | Emission NO _x -N | Emission N ₂ O-N | Soil stocks NO ₃ -N /NH ₄ -N |
|--------------------|--|---------------------------------|-----------------------------|----------------------------|-----------------------------|-----------------------------|--|
| 60 | 0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| 60 | 50 | -1.3 | 0.3 | -17.9 | -3.5 | -2.1 | -1.5 |
| 60 | 100 | -2.6 | 0.6 | -24.4 | -5.5 | -4.1 | -3.2 |
| 60 | 250 | -6.7 | 1.5 | -35.0 | -10.5 | -9.8 | -8.0 |
| 60 | 500 | -13.3 | 2.9 | -45.6 | -17.8 | -19.0 | -15.7 |
| 60 | 750 | -19.5 | 4.3 | -53.4 | -24.5 | -27.7 | -22.8 |
| 60 | 1000 | -25.4 | 5.5 | -59.8 | -30.6 | -35.8 | -29.3 |
| 60 | 2000 | -44.5 | 9.7 | -78.0 | -50.5 | -62.3 | -49.4 |
| 90 | 0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| 90 | 50 | -1.0 | 0.3 | -16.4 | -6.7 | -3.1 | -1.2 |
| 90 | 100 | -2.2 | 0.6 | -22.4 | -9.4 | -5.3 | -2.7 |
| 90 | 250 | -6.0 | 1.5 | -32.3 | -14.5 | -11.1 | -7.3 |
| 90 | 500 | -12.1 | 3.0 | -42.7 | -20.8 | -20.1 | -14.7 |
| 90 | 750 | -18.0 | 4.5 | -50.7 | -26.1 | -28.4 | -21.5 |
| 90 | 1000 | -23.6 | 5.8 | -57.4 | -30.8 | -36.2 | -27.8 |
| 90 | 2000 | -41.8 | 10.3 | -76.4 | -46.0 | -61.6 | -47.2 |

b.

| Moisture treatment | Added Cu (mgCu.kg soil ⁻¹) | NO ₃ -N soils stocks | Emission NH ₄ -N | Emission N ₂ -N | Emission NO _x -N | Emission N ₂ O-N | Soil stocks NO ₃ -N /NH ₄ -N |
|--------------------|--|---------------------------------|-----------------------------|----------------------------|-----------------------------|-----------------------------|--|
| DO | 0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| DO | 50 | -0.7 | 0.2 | -17.7 | -3.2 | -1.7 | -0.8 |
| DO | 100 | -1.5 | 0.3 | -23.9 | -4.8 | -3.2 | -1.8 |
| DO | 250 | -3.9 | 0.8 | -33.5 | -8.4 | -7.6 | -4.7 |
| DO | 500 | -8.1 | 1.7 | -42.8 | -13.6 | -14.8 | -9.6 |
| DO | 750 | -12.3 | 2.6 | -49.8 | -18.4 | -22.1 | -14.5 |
| DO | 1000 | -16.5 | 3.5 | -55.8 | -23.1 | -29.3 | -19.3 |
| DO | 2000 | -33.3 | 7.0 | -75.7 | -41.6 | -58.3 | -37.7 |
| DR | 0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| DR | 50 | -0.6 | 0.1 | -17.6 | -3.6 | -1.6 | -0.7 |
| DR | 100 | -1.3 | 0.3 | -23.8 | -5.3 | -3.1 | -1.6 |
| DR | 250 | -3.5 | 0.7 | -33.3 | -9.1 | -7.3 | -4.2 |
| DR | 500 | -7.2 | 1.4 | -42.4 | -14.2 | -14.3 | -8.6 |
| DR | 750 | -10.9 | 2.2 | -49.1 | -19.0 | -21.2 | -12.8 |
| DR | 1000 | -14.5 | 2.9 | -54.8 | -23.5 | -27.9 | -16.9 |
| DR | 2000 | -28.6 | 5.7 | -73.2 | -40.7 | -54.1 | -32.5 |



730 **Fig. 1:** Schematic representation of the experimental and modelling procedures. Left refers to the experimental part and centre to right to the modelling part. Soils were first incubated 5 weeks at different constant percentage of the water holding capacity (WHC) or at two variable moistures, Dry-Only (DO) and Dry-Rewet (DR). Then NO₂-N, NO₃-N and NH₄-N soil concentrations were measured after this initial incubation, and values were used to initialise DNDC, while a bioassay was also applied on soil aliquots. The 3 days bioassay included NH₄⁺ in
735 excess and copper (Cu) spikes at 0, 50, 100, 250, 500, 750, 1000, 2000 mg Cu.kg soil⁻¹ of soil. After 1 and 3 days of bioassay incubation, NO₃-N production was measured in the supernatant. NO₃-N productions against [Cu] gradients were used to define the functions of eq. 28 to 31 in §3.1 (see text). Soil respiration values were extracted from the curve C_i of Fig 1 in Annabi et al. (2007).



740

Fig. 2: Fitted functions of potential nitrifying activities (PNA) against total soil copper concentrations [Cu] for each initial moisture incubation treatment. Points are the measured nitrate production and lines the fitted quadratic function with their 95% confidence interval. (a). Constant moisture treatments: green circle is for 30% WHC, red square for 60% WHC and purple diamond for 90% WHC. The black line is the common fitting function used for 60 and 90% WHC moisture treatments. (b). Variable initial moisture treatments: brown star is for Dry-Rewet (DR) and yellow triangle for Dry-Only (DO).

745





Fig 3: Comparison between modelled and measured soil [nitrate] incubated in different moisture with 1:1 line

(a) = the 3 initial incubations under constant moisture. (b) = the two initial incubations under variable moisture

755 Dry-Rewet (DR) and Dry-Only (DO) treatments. For 30% WHC, Model=1.84 * Measure and R²=0.93; for 60% WHC Model=0.93 *measure. R²=0.99; for 90% WHC Model=0.90 * measure. R²=0.99 ; for Dry -rewetting (DR) model = 0.96*measure. R²=0.98; for Dry-Only (DO)Model=0.95*measure. R²=0.99



760

Fig 4. Modelled soil denitrifying bacterial pool after 3 days (gN.m⁻³ soil) for the 4 moisture treatments. Purple diamond is for 90% WHC, red square for 60% WHC, brown star for Dry-Rewet (DR) and yellow triangle for Dry-Only (DO). Red, brown and yellow curves being superposed. Pools were modelled for 12, 62, 112, 262, 512, 762, 1012 and 2012 mg Cu,kg soil⁻¹ as represented by cross. Quadratic fits were used for representation.

765




Fig 5: Modelled N emission fluxes at 3 days in gN.m⁻².30min⁻¹ under the different moisture treatments. a.) NH₄-N emission fluxes. b.) NOx-N emission fluxes c.) N₂O-N emission fluxes and d.) N₂-N emission fluxes. Purple diamond is for 90% WHC, red square for 60% WHC, brown star for Dry-Rewet (DR) and yellow triangle for Dry-Only (DO). Fluxes were modelled for 12, 62, 112, 262, 512, 762, 1012 and 2012 mg Cu.kg soil⁻¹ as represented by cross. Quadratic fits were used for representation.




Fig. 6: Proportion of N₂O-N emitted arising from the denitrification calculated as N₂O-N / (N₂O-N + N₂-N) modelled fluxes in response to soil Cu concentration for the various moisture treatments. Red square is for 60% WHC, purple diamond is for 90% WHC, yellow circle for Dry-Rewet (DR) and brown star for Dry-Only (DO). Red, yellow and brown curves are superposed. Fluxes were modelled for 12, 62, 112, 262, 512, 762, 1012 and 2012 mg Cu.kg soil⁻¹ as represented by cross. Quadratic fits were used for representation.

785