

Blue Carbon Stocks and Exchanges Along the California Coast

Melissa A. Ward^{1,2}, Tessa M. Hill¹, Chelsey Souza¹, Tessa Filipczyk¹, Aurora M. Ricart^{1,3}, Sarah Merolla¹, Lena R. Capece¹, Brady C. O'Donnell¹, Kristen Elsmore¹, Walter C. Oechel², and Kathryn M. Beheshti⁴.

Correspondence to: Melissa A. Ward (maward@ucdavis.edu)

¹Bodega Marine Laboratory, University of California, Davis, 95616, USA

²San Diego State University, 92182, USA

³Bigelow Laboratory for Ocean Sciences, 04544, USA

⁴University of California, Santa Cruz, 95064, USA

Abstract. Salt marshes and seagrass meadows can sequester and store high quantities of organic carbon (OC) in their sediments relative to other marine and terrestrial habitats. Assessing carbon stocks, carbon sources, and the transfer of carbon between habitats within coastal seascapes are each integral in identifying the role of blue carbon habitats in coastal carbon cycling. Here, we quantified carbon stocks, sources, and exchanges in seagrass meadows, salt marshes, and unvegetated sediments in six bays along the California coast. In the top 20 cm of sediment, the salt marshes studied here contained approximately twice as much OC as did seagrass meadows, 4.92 ± 0.36 kg OC m⁻² compared to 2.20 ± 0.24 kg OC m⁻², respectively. Both salt marsh and seagrass sediment carbon stocks were higher than previous estimates from this region but lower than global and U.S.-wide averages, respectively. Seagrass-derived carbon was deposited annually into adjacent marshes during fall seagrass senescence. However, isotope mixing models estimate that negligible amounts of this seagrass material were ultimately buried in underlying sediment. Rather, the vast majority of OC in sediment across sites was likely derived from planktonic/benthic diatoms and/or C3 salt marsh plants.

Keywords: Carbon cycling, lateral fluxes, seagrass meadow, salt marsh, carbon sequestration

1 Introduction

As carbon dioxide (CO₂) concentrations in the oceans and atmosphere continue to rise, interest in measuring the relative quantities of carbon stored within natural ecosystems has increased. These assessments can help improve global and regional climate models, the prediction of future CO₂ concentrations related to sources and sinks, and our broader understanding of nature-based climate change solutions (Serrano et al., 2019). Coastal habitats including seagrasses, salt marshes, and mangroves have earned the moniker ‘blue carbon’ habitats for their ability to store disproportionately high levels of organic carbon (OC) in their

39 sediments relative to other habitat types, thereby potentially serving in a management context to
40 provide carbon mitigation (Lovelock and Duarte, 2019; McLeod et al., 2011). This ability can be
41 largely attributed to the tendency for these habitats to exhibit high sediment accretion rates and
42 low decomposition rates (Peck et al., 2020; Serrano et al., 2019). This value has led to increasing
43 interest in blue carbon habitats, given their conservation can prevent significant emissions of
44 carbon (Lovelock et al., 2017; Pendleton et al., 2012) and their restoration can lead to increased
45 drawdown of atmospheric CO₂ (Freedman et al., 2009; Greiner et al., 2013).

46

47 **1.1 Carbon stock assessments and their significance**

48 Despite global interest, many regions are still lacking basic information on carbon stocks
49 and burial rates in blue carbon habitats. This information forms the foundation for more
50 advanced scientific research and can be extremely valuable within management contexts to
51 develop informed local and regional climate assessments. Salt marshes and seagrass meadows
52 cover extensive portions of North America's west coast, yet their carbon stocks have been
53 relatively understudied compared to other habitats in North America and other blue carbon
54 habitats in different regions of the world (Ouyang and Lee, 2014; Postlethwaite et al., 2018).

55 Eelgrass (*Zostera marina*) is the dominant seagrass species in North America and
56 occupies coastal waterways from Alaska to Mexico (Green & Short, 2003). Until recently,
57 seagrass carbon stock data were almost entirely absent along the west coast of North America
58 (see Capece et al., 2019; Kauffman et al., 2020; O'Donnell, 2017; Poppe & Rybczyk, 2018;
59 Postlethwaite et al., 2018; Röhr et al., 2018). Of these studies, very few data came from Central
60 or Southern California seagrass meadows, despite the fact that this region represents a key
61 temperate to subtropical transition in *Z. marina*'s range (Cabello-Pasini et al., 2003). As a result

62 of the previous paucity of data from the west coast of North America, this region was not
63 represented in previous global syntheses of seagrass carbon stocks (Duarte et al., 2010;
64 Fourqurean et al., 2012; Mazarrasa et al., 2015). Existing work on global seagrass carbon storage
65 identifies that two species of seagrasses in the Mediterranean (*Posidonia oceanica*) and Australia
66 (*Posidonia australis*) store significantly more carbon than other seagrass species, including *Z.*
67 *marina* (Fourqurean et al., 2012; Lavery et al., 2013; Prentice et al., 2020). These *Posidonia*
68 species were overrepresented in some early assessments of total global seagrass carbon storage –
69 making these global estimates unreliable when applied to management decisions or climate
70 models (Johannessen & Macdonald, 2016; Kennedy et al., 2010). In reality, the geographic and
71 interspecies variability in carbon stocks is likely greater than was initially anticipated (Macreadie
72 et al., 2018; Postlethwaite et al., 2018).

73 Salt marsh carbon stocks are similarly understudied in western North America, with
74 published carbon stock data from only four estuaries in the region and very minimal spatial
75 coverage and analyses performed in three of these locations (Brevik and Homburg, 2004;
76 Callaway et al., 2012; Kauffman et al., 2020; Patrick and DeLaune, 1990). Existing analyses of
77 North American freshwater wetlands and salt marshes are typically dominated by studies along
78 the east coast (Nahlik & Fennessy, 2016; Wilkinson et al., 2018; with the exception of Holmquist
79 et al. 2018). Within west coast studies, considerable variation in carbon stocks is observed
80 (Callaway et al., 2012; Chmura et al., 2003). Similar to seagrass meadows, these regions have
81 distinctly different oceanographic and geomorphological regimes, which can drive differences in
82 sediment carbon storage. Thus, seagrass and salt marsh carbon storage data collected from
83 understudied regions and across varying environmental gradients are necessary for understanding
84 carbon stock variability and its drivers.

85

86 **1.2 Blue carbon sources and drivers**

87 Despite an increasing breadth of information on blue carbon habitats in the last decade, a
88 number of questions remain before we can fully understand their role in regional carbon cycling
89 and climate adaptation (Macreadie et al., 2019). We do not yet fully understand what drives
90 variation in carbon stocks – a key aspect of ensuring the protection and enhancement of these
91 carbon services in the future. Previous work demonstrates that numerous factors can control
92 carbon accumulation in coastal sediments including overlying biomass, topography, hydrology,
93 mineralogy, and remineralization rates (Kelleway et al., 2016; Lima et al., 2020; Mazarrasa et
94 al., 2018; Prentice et al., 2019). In particular, sediment grain size has been demonstrated as a
95 significant predictor of carbon stocks (Dahl et al., 2016; O’Donnell, 2017; Serrano et al., 2016),
96 as it affects decomposition rates, which is likely related to the deposition of small particles and
97 low resuspension from the attenuation of water flow by seagrasses (Bos et al., 2007; Conley et
98 al., 2017; Gambi et al., 1990; Hendriks et al., 2008). Through similar pathways, fine, silty
99 sediments trapped in tidal salt marshes can also increase carbon storage (e.g. Zhou et al., 2007).

100 The relative importance of each potential driver of carbon stock variability may be highly
101 dependent on the environmental setting, species under study, and the interactive effects of these
102 drivers. For example, in Australian salt marshes, overlying vegetation was shown to only
103 significantly affect carbon stocks in sandy and mixed grain size sediments, having no effect on
104 stocks in fine sediments (Kelleway et al., 2016). However, in other cases vegetation has been the
105 primary predictor of carbon stocks in salt marshes, irrespective of grain size (Lovelock et al.,
106 2014; Saintilan et al., 2013). Similar region and species-specific complexities between carbon
107 stocks, overlying vegetation, and mineralogy have been described in seagrass meadows (e.g.

108 Lima et al., 2020; Serrano et al., 2016), meriting further investigation to understand these
109 complex interactions.

110 Knowing the relative contributions of locally produced (autochthonous) and imported
111 (allochthonous) carbon also elucidates the underlying mechanisms by which blue carbon habitats
112 store and accumulate carbon. Overlying vegetation can significantly impact sediment carbon
113 stocks; however, it is rarely the dominant source of carbon buried within blue carbon habitats
114 (Kennedy et al., 2010; Lewis et al., 2020; Mazarrasa et al., 2015; Prentice et al., 2019). In many
115 cases, contributions from terrestrial habitats, macroalgae, and suspended particulate organic
116 material contribute as much or more to carbon buried in blue carbon sediments than do
117 autochthonous sources (Drexler et al. 2020; Kennedy et al., 2010; Leorri et al., 2018; Ricart et
118 al., 2020). Thus, local primary production could significantly contribute to net annual carbon
119 drawdown within a given habitat yet play a minor role in carbon burial due to lateral export or
120 remineralization of this carbon. Understanding this phenomenon elucidates the role of blue
121 carbon habitats in broader, system-wide energy flows and carbon cycling (Hyndes et al., 2014;
122 Ricart et al., 2015). For instance, much of the carbon from laterally exported biomass may be
123 remineralized in the water column or as wrack in nearby habitats (Attard et al., 2019; Liu et al.,
124 2019), serving an important ecological role and altering the production and biogeochemical
125 cycles of recipient systems (Hyndes et al., 2014; Ince et al., 2007; Valiela and Cole, 2002).
126 Given that blue carbon habitats are highly productive (Duarte & Cebrián, 1996), support high
127 carbon burial (e.g. McLeod et al., 2011), and can co-occur within small geographic ranges
128 (Alongi, 2018; Bouillon & Connolly, 2009), laterally exported carbon from one blue carbon
129 habitat may be entrapped and buried in a neighboring habitat. From this landscape-scale
130 perspective, exchanges between blue carbon habitats could increase their capacity for carbon

131 burial or alter the sources of buried carbon. While some coastal studies have estimated lateral
132 carbon fluxes (Jiménez et al., 2017; Liu et al., 2019), very few studies place these lateral fluxes
133 into landscape-scale contexts that also address carbon burial in recipient habitats (Bouillon &
134 Connolly, 2009; Duarte & Krause-Jensen, 2017; Ricart et al., 2017).

135

136 The study presented here addresses this globally relevant topic of research in a relatively
137 understudied region by answering the following questions:


- 138 1) How much organic carbon is stored in seagrass and salt marsh sediments across a
139 California latitudinal gradient?
- 140 2) What are the sources of carbon buried within these sediments?
- 141 3) Is carbon exchanged between blue carbon habitats within the coastal landscape?

142

143 **2 Methods**

144 **2.1 Study Sites**

145 Sediment cores from salt marshes and seagrass meadows, along with neighboring
146 unvegetated sediments (hereafter ‘bare sediment’ near seagrass meadows, and ‘pan’ near salt
147 marshes) were collected from six bays across a latitudinal gradient in California (Fig. 1a). A total
148 of 82 sediment cores were collected, 30 of which have been discussed previously (O’Donnell,
149 2017) and are included here for comparison. The number of cores collected in each site and
150 general site characteristics are described in Table 1. While all sampled seagrass meadows were



151

152 **Figure 1:** Sediment cores were collected from
 153 six estuaries across a latitudinal gradient in
 154 California, where N = number of cores (A).
 155 Seagrass wrack deposition into Walker salt
 156 marsh from surrounding seagrass meadows
 157 (Merkel & Associates, 2017) was estimated,
 158 while sediment cores were taken from all four
 159 labeled habitats (B). Three sediment cores
 160 were collected beneath persistent wrack lines
 161 in Walker salt marsh, while three were
 162 collected from areas that do not receive
 163 regular, annual tidal deposition of wrack (C).
 164

165

166 dominated by a single seagrass species (*Zostera marina*), salt marshes contained a mixed
 167 community of halophytes, predominantly composed of pickleweed (*Sarcocornia pacifica*), and
 168 to a lesser extent salt grass (*Distichlis spicata*) and marsh jaumea (*Jaumea carnosa*). Bare
 169 sediment cores were collected in unvegetated areas nearby each seagrass meadow at a minimum
 170 of 20 meters away from the meadows and at similar depths. Pan cores

171 were collected from patches of unvegetated sediment (2-4m diameter) found within the salt
 172 marsh interior, a natural and semi-permanent feature of salt marsh habitats formed by elevational
 173 depressions (Escapa et al., 2015). None of our sampling sites were actively restored and, to our
 174 knowledge, respective vegetation has persisted through time.

175 We estimated OC stocks in seagrass, salt marsh, pan, and bare sediments (see details below).
 176 In addition, we analyzed sediment OC sources for two individual salt marshes in Elkhorn Slough
 177 ('Elkhorn salt marsh') and Tomales Bay ('Walker salt marsh'), and we examined carbon
 178 exchange between a single seagrass meadow and the neighboring Walker salt marsh. Walker salt
 179 marsh is located approximately 5.5 km from the mouth of Tomales Bay and is in close proximity
 180 to extensive meadows of *Z. marina* – the dominant seagrass in each of the meadows sampled
 181 (Fig. 1b). This marsh lies where the mouth of Walker Creek meets bay waters, and thus can
 182 receive terrestrial and riverine inputs, while simultaneously receiving marine inputs from tidal
 183 exchange.

Bay	Coordinates	Number of cores	Site description
Humboldt Bay	40°48'12 N, 124°10'52 W	Seagrass: 4 Bare sediment: 3	A large semi-enclosed bay with a narrow mouth, stabilized by jetties. Primary freshwater sources come from the Elk River and its tributaries, with Freshwater Creek inputs entering the bay near the sampling site. Freshwater input is small relative to tidal flushing but can locally effect sedimentation rates within the bay (Schlosser and Eicher, 2012)
Bodega Harbor	38°19'60 N, 123°02'53 W	Seagrass: 15 Bare sediment: 3	There are no significant rivers or creeks that provide freshwater or sediment inputs. Sediment input in Bodega Bay is largely controlled by jetties at the mouth of the harbor and is influenced by its history of routine dredging to maintain shipping channels and control wind-blown sand entering from nearby dunes

Tomales Bay	38°10'17 N, 122°54'46 W	Seagrass: 15 Bare sediment: 9 Salt Marsh: 6	A long, narrow bay that receives periodic freshwater inputs from two primary tributaries, Walker Creek and Lagunitas Creek, which peaks seasonally during winter storms.
Elkhorn Slough	36°49'15 N, 121°45'06 W	Salt Marsh: 11 Pan: 3	Receives periodic freshwater inputs from the Old Salinas River and Bennett and Moro Coho Sloughs. After a long history of increasing sediment deposition and slowing tidal flow to the bay, levee breaches and removals in 1953 rapidly converted it to a higher tidal energy, erosional system, as it remains today (Van Dyke and Wasson, 2005). As a result, the extent of 'high-quality' salt marsh in 2000 was 23% of what it had been a century earlier (Van Dyke and Wasson, 2005)
Newport Bay	33°38'07 N, 117°53'17 W	Seagrass: 4 Bare sediment: 4	An urban bay, that receives terrestrial and urban inputs from the San Diego Creek Watershed. After a long history of high sedimentation and water quality nutrient impairments from urban and agricultural sources, a program was implemented in 1998 to meet water quality targets. Since its commencement, both sedimentation and nutrients in the bay have significantly decreased (Trimble, 2003).
Mission Bay	32°47'24 N, 117°13'40 W	Seagrass: 3 Bare sediment: 2	An urban, enclosed bay, that receives terrestrial and urban input from the San Diego River watershed and the Rose Creek tributary. Experiences long residence times and is primarily controlled by tidal flushing processes rather than the minimal freshwater inputs (Cyronak et al., 2018; Largier et al., 1997)

184
185
186
187

Table 1: Location and brief description of each bay sampled in this study, along with the total number of cores collected from each habitat type in each of these bays.

188 **2.2 Carbon stocks**

189 All sediment cores were sampled by manually inserting transparent, open-barrel PVC pipes
190 (20 cm length, 5.08 cm diameter). Compaction occurred in 19% of cores, and a compaction

191 factor was applied when calculating carbon stocks according to Howard et al. (2014). Once
192 extracted, cores were capped and transported to the laboratory upright to prevent mixing of
193 sediment layers. Cores were then immediately extruded into sections at 2 cm intervals. Coarse
194 living plant material (> 1cm) was manually removed. Each section was dried at 60°C and
195 weighed, and dry bulk density (DBD) was determined by dividing dry bulk mass by the volume
196 of sampling interval.

197 Each section was then homogenized and divided into three subsamples of 10 g each, and the
198 remaining sample was archived. One of the three subsamples was acidified using 1.12 M HCl to
199 remove and measure total inorganic carbon (Milliman, 1974). The second subsample was
200 analyzed for total organic material (TOM) by loss-on-ignition method for 4 hours at 550° C
201 (Dean, 1974). Total C content (%) and $\delta^{13}\text{C}$ were additionally determined on a set of 44 of the
202 acidified subsamples randomly selected across each habitat type using an elemental analyzer
203 (PDZ Europa ANCA-GSL, SD \pm 0.25 %) integrated with a continuous flow isotope ratio mass
204 spectrometer (PDZ Europa 20-20, SD \pm 0.2 ‰) at the UC Davis Stable Isotope Facility. $\delta^{13}\text{C}$
205 ratios are expressed in parts per thousand (‰) relative to VPDB (Vienna Pee Dee Belemnite)
206 according to standard notation ($\delta^{13}\text{C} = [(R_{\text{sample}} / R_{\text{standard}}) - 1] \times 1000$, where R is the ratio
207 $^{13}\text{C}/^{12}\text{C}$). Organic carbon in each core section was then estimated using a power model developed
208 between measured TOM and measured OC in this set of subsamples (as in Craft et al., 1991).
209 Specifically, the equation $y = 0.22x^{1.1}$ was applied ($r^2 = 0.84$; Fig. S1). A power model was
210 selected over a linear model for these data to avoid negative estimates of carbon stocks at low
211 levels of TOM (Fig. S1). Carbon stocks were determined by multiplying DBD (g cm^{-3}) by %
212 OC.

213 In each core section, the proportion of fine sediments – the silt and clay fraction (<63 μm),
214 hereafter “mud” – was quantified. The grain size analysis was conducted by mass loss in the
215 third subsample from each section in 70 of the 82 total cores. The subsample was rinsed through
216 a 63 μm sieve with deionized water, and the remaining sediments were dried at 60°C and re-
217 weighed.

218 Comparable to previous studies, a core depth of 20 cm was selected (Fourqurean et al. 2012;
219 Prentice et al., 2020; Röhr et al., 2018). As such, stock estimates are presented in kg OC m^{-2} in
220 the top 20 cm of sediment. However, given the selected depths for stock estimates vary
221 throughout the literature, we extrapolate all data to 1 m when comparing across studies. Some
222 research suggests that OC content remains relatively constant below 10 cm to depths up to 1 m,
223 making this extrapolation appropriate (Callaway et al., 2012; Prentice et al., 2020; Fig. S3),
224 while evidence of downcore variability in other studies makes extrapolation less appropriate
225 (e.g., St. Laurent et al., 2020; Serrano et al., 2012). When comparing the carbon stocks estimated
226 here to those in previous studies, we include each studies’ sampled core depths for clarity.

227

228 **2.3 Carbon sources and exchange**

229 We applied mixing models to assess salt marsh sediment carbon sources to understand
230 within-estuary exchange of carbon among the sampled habitats. Two separate mixing models
231 were produced for 1) Elkhorn salt marsh in Elkhorn Slough and Walker salt marsh in Tomales
232 Bay, and 2) salt marsh sediments under seagrass wrack, versus not under wrack in Walker salt
233 marsh. At Walker salt marsh, a total of six sediment cores were collected. Three of these
234 sediment cores were collected from the interior marsh, while three were collected from
235 underneath a seagrass wrack line along the tidal edge of the marsh (Fig. 1c). *Z. marina* in this

236 region is known to undergo a period of senescence as photoperiod shortens in the fall, as is
237 common in all temperate seagrasses (Fourqurean et al., 1997). Historical imagery of the site
238 shows persistent seagrass wrack concentrated along these tide lines that consistently appears in
239 early fall as seagrass senesces (Fig. S2; Google Earth, 2020). The biomass of seagrass wrack
240 along this tide line at the time of core collection was quantified along an 80 m transect within the
241 marsh by collecting all seagrass present in a 1 m² quadrat every 10 meters. This material was
242 taken back to the lab, sorted into above ground biomass (AGB) and below ground biomass
243 (BGB), rinsed, dried (60°C), and weighed. Biomass data from seagrass in nearby meadows was
244 previously published (see O'Donnell, 2017) and is used here for reference. Sediment cores
245 collected from beneath wrack lines were sectioned at 2 cm intervals, analyzed for total OC and
246 $\delta^{13}\text{C}$ content according to the methods described above. Total nitrogen content was also
247 determined on an unacidified portion of the same 44 sediment subsamples used in carbon
248 analyses (ThermoFinnigan Flash 1112 Series elemental analyzer, SD \pm 0.02%). Data from
249 Walker salt marsh are also displayed as 'shallow' (surface sediment to 10 cm deep) and 'deep'
250 (10 cm and deeper), to facilitate interpretation of changes with depth.

251 The contributions of carbon sources to each core section were then estimated with a mixing
252 model using $\delta^{13}\text{C}$ and N:C ratios as tracers. Given $\delta^{15}\text{N}$ can be altered during early diagenesis
253 (e.g. Benner et al., 1991), we selected $\delta^{13}\text{C}$ and C:N ratios as the primary geochemical tracers,
254 and therefore did not include fractionation factors into the model (Craven et al., 2017). N:C
255 ratios are utilized rather than C:N ratios because mixed fractions returned by the model are based
256 on the denominator, and thus $^{13}\text{C} / ^{12}\text{C}$ and N:C both estimate the fractional contribution of ^{12}C
257 (Craven et al., 2017; Perdue and Koprivnjak, 2007). However, C:N ratios are used in text and
258 figures given this format is more typically presented in the literature. We used a Bayesian mixing

259 model SIAR 4.2 (Parnell and Jackson, 2013) to estimate the contributions of several source
260 groups selected according to the dominant plant and algal species observed within the selected
261 salt marshes. The sources included C3 salt marsh plants, C4 salt marsh plants (*Distichlis*
262 *spicata*), seagrass (*Z. marina*), and plankton/benthic diatoms. Seagrass geochemistry was
263 estimated from seagrass leaves collected from meadows near the selected salt marshes in
264 Newport Bay, Tomales Bay, and Bodega Bay (Fig. 1a; Capece et al., 2019). Representative
265 geochemistry for all other sources has been published previously and was used here, from
266 samples collected in San Francisco Bay, approximately 60 km south of Walker salt marsh (see
267 Cloern et al., 2002 for full methods). All salt marsh sediment cores were collected from areas of
268 the marsh that were dominated by the aforementioned species, and as such, other marsh species
269 were not considered (e.g., *Spartina sp.*). While lower densities of other C3 marsh plants may
270 have been present in some cored sites, the variability demonstrated by the selected source
271 samples likely encompasses much of this variability given the utilization of the same
272 photosynthetic pathway (Cloern et al., 2002). Plankton samples were collected from estuarine
273 water samples while benthic diatom samples were collected from both salt marsh surface
274 sediments and neighboring mudflats (Cloern et al., 2002). These sources have similar isotopic
275 values and are pooled here and referred to as “diatoms” for simplicity, acknowledging that this
276 encompasses contributions from multiple planktonic and benthic sources. Given the overlapping
277 isotopic values of C3 plants and diatoms, these two sources were pooled in mixing models,
278 allowing an estimate of diatoms and/or C3 plants contributions to marsh sediments, in addition to
279 contributions from seagrass and C4 plants.

280

281 **2.4 Statistical Analyses**

282 To evaluate decay of carbon through time, we tested for significant changes in OC in each
283 habitat type with core section depth (a proxy for time) by fitting data to a generalized linear
284 mixed model (GLMM) using maximum likelihood with ‘depth’ and ‘site’ as fixed effects and
285 ‘core’ as a random effect, using a gamma distribution and log link function to account for non-
286 normality. In sites and habitats that demonstrated significant OC changes with depth, the rate of
287 decay was estimated from the slope of its associated model. In all analyses to follow, OC across
288 all sections in each core were averaged, and statistics were performed on these core averages.
289 After inspecting data for normality and homogeneity of variance, differences in OC and grain
290 size between habitat types and between sites were analyzed using simple linear models (SLM;
291 significance defined by $\alpha = 0.05$). Specifically, these data were fit to a linear model using
292 ordinary least squares with ‘site’ and ‘habitat’ (i.e., seagrass, salt marsh, bare sediment, pan) as
293 fixed effects, including their interaction. The relationships between TOM (%) and grain size
294 were analyzed using simple linear regressions, whereby a grain size filter was selectively applied
295 to determine the point at which the relationship between the two was no longer significant ($p <$
296 0.05). Differences in $\delta^{13}\text{C}$ (‰) or C:N ratios between sediment depth (‘surface’ versus ‘deep’),
297 and between sediments collected under wrack versus not under wrack, were also tested with
298 SLMs using ordinary least squares with ‘depth’ or ‘under wrack’ as fixed effects. When
299 necessary, data were log transformed. Tukey’s post-hoc analyses were conducted for multiple
300 comparisons. All statistical analyses were performed in R software (R Core Team, 2018).

301

302 **3 Results**

303 **3.1 Carbon Stock Assessments**




Figure 2 (left): Down core trends (surface = 0 cm) in average (\pm SE) OC (g cm^{-3}) for each habitat type and site. Data show an insignificant decline in OC through time (downcore) in all sites except Newport Bay salt marshes.

326

327

328

329

330

331

332


333

334


335

Down core OC demonstrates high variability, resulting in few significant differences in OC with depth (Fig. 2). Specifically, only Newport Bay salt marsh sediments exhibited significant loss of OC down core, which declined at a rate of $0.001 \text{ g C cm}^{-3}$ per cm depth (Fig 2e; GLMM, T-value = -4.7, SE = 0.01, $p < 0.05$). Significant differences between OC stocks emerged when cores were compared between habitats, with salt marshes containing significantly more carbon than both bare sediment and seagrass meadows (Fig. 3b; SLM, $p < 0.01$, F-statistic = 13.3, DF = 3). Specifically, within the top 20 cm of sediment, salt marsh sediments contained $4.92 \pm 0. \text{ kg OC m}^{-2}$, while seagrass meadows contained $2.20 \pm 0.24 \text{ kg OC m}^{-2}$ (Fig. 3b; mean \pm SE). Seagrass meadow carbon stocks were not significantly different than those of nearby bare

336 sediments (Tukey's post-hoc analysis, $p > 0.05$), which contained an average of 2.47 ± 0.32 kg
337 OC m⁻². Pan sediments also contained higher carbon than did seagrass meadows (Tukey's post-
338 hoc analysis, $p > 0.05$). A full list of carbon stocks is displayed in Table S1. Tukey's post-hoc
339 analysis indicates that in Tomales Bay, salt marshes contained significantly more carbon than
340 seagrass meadows (Fig. 3a; $p < 0.05$) while bare sediment and seagrass carbon stocks did not
341 significantly differ from one another ($p > 0.05$). We did not detect any other significant
342 differences when comparing carbon stocks across habitat types within each individual site
343 (Tukey's post-hoc analysis, $p > 0.05$).



344




345

346 **Figure 3:** Average (\pm SE) total organic carbon (OC) stocks (kg m^{-2}) across all cores collected
 347 from each habitat type within each site in this study (A). Average (\pm SE) total organic carbon
 348 stocks (kg m^{-2}) across all cores collected within each habitat type in this study (B). Stocks are
 349 representative of the top 20 cm of sediment.
 350

351 We observed a very strong relationship between grain size and storage of organic material,
 352 especially at lower TOM (%) values (Fig. 4). Specifically, the observed linear relationship
 353 between grain size and TOM is strongest when sediment is 8.5% TOM or less (linear model, $r^2 =$

354 0.95, $p < 0.05$, F-statistic = 4957, DF = 269). However, as the proportion of fine sediments in
355 each sample increases, the relationship between grain size and TOM weakens rapidly (from $r^2 =$
356 0.73 to 0.53) after sediments consisting of more than 82% mud are included (linear model, p
357 < 0.05 , $r^2 = 0.53$, F-statistic = 233.1, DF = 201). Our results indicate that grain size was similar
358 between sites (SLM, $p > 0.05$, F-statistic = 2.7, DF = 3) but differed between habitat types (Fig.
359 5a; SLM, $p < 0.05$, F-statistic = 5.7, DF = 3), with salt marsh sediments demonstrating
360 significantly greater % mud than both seagrass and bare sediment (Tukey's post-hoc analysis, p
361 < 0.05). Recognizing that the sample size of pan cores was low, pan sediments displayed
362 comparable % mud to salt marsh sediments but there were no significant differences between
363 pan grain size and the other habitat types' grain sizes (Tukey's post-hoc analysis, $p > 0.05$).
364 Tukey's post-hoc analyses did not indicate any significant differences in grain size between
365 habitat types within each site (Fig. 5a; $p > 0.05$).


366




367
368 **Figure 4:** Biplot showing the relationship among total organic material (TOM) (%) of each 2-cm
369 core section from each habitat type plotted against sediment grain size (% Mud).

370

371



372



373

374 **Figure 5:** Average (\pm SE) % mud of all sediments quantified in each site within each habitat
375 type (A), average (\pm SE) % mud in each habitat type across all sites measured (B).

376

377 3.2 Carbon Sources and Exchange

378 3.2.1 Wrack Deposition

379 While we conducted only one ground survey of seagrass wrack lines within Walker salt
 380 marsh in October 2019, historical aerial imagery was utilized to verify that these wrack lines
 381 appear reliably in this season in similar locations each year (Fig. S2). This phenomenon can be
 382 viewed as far back as 2002, before which images are of low quality or unavailable (Google
 383 Earth, 2020). These wrack lines typically persist throughout the winter, becoming indiscernible
 384 via aerial imagery by spring. Previous data collected by the authors (O'Donnell, 2017) quantified
 385 seagrass senescence within the nearby Tom's Point seagrass meadow (Fig. 1b), where average
 386 summer seagrass biomass was $440 \pm 59.4 \text{ g m}^{-2}$ and winter seagrass biomass was $115 \pm 16.5 \text{ g m}^{-2}$
 387 ² (AGB and BGB, mean \pm SE; Table 2). Within the neighboring Walker salt marsh, fall estimates
 388 of wrack demonstrated that $106 \pm 24.6 \text{ g m}^{-2}$ of seagrass (dry weight) was deposited along tide
 389 lines (Table 2). While both seagrass AGB and BGB are included in this value, seagrass BGB
 390 only accounted for 3.5% of total seagrass biomass measured (Table 2). Analysis of seagrass
 391 leaves collected from Tomales Bay demonstrated that seagrass material was composed of 31.6%
 392 of OC. Thus, we estimate Walker salt marsh receives $33.4 \pm 7.6 \text{ g OC m}^{-2}$ in the form of seagrass
 393 along wrack lines each year.


	Seagrass (winter) (g/m ²)	Seagrass (summer) (g/m ²)	Wrack (g/m ²)	Wrack carbon (gC/m ²)
AGB	58.6 ± 14	361.3 ± 47.8	102.8 ± 24	32.5 ± 7.6
BGB	57.1 ± 4.6	79 ± 15.5	3.7 ± 2.6	1.2 ± 0.8
Total	115.7 ± 16.5	440.3 ± 59.4	105.7 ± 24.6	33.4 ± 7.8

394
 395 **Table 2:** Living seagrass above ground biomass (AGB) and below ground biomass (BGB) was
 396 collected in Tom's Point seagrass meadow (Fig. 1b) in both summer and winter (g m^{-2}). Total
 397 seagrass wrack delivered to Walker salt marsh was quantified as AGB and BGB (g m^{-2}) and
 398 converted to carbon (g C m^{-2}) using a 31.6% carbon conversion rate.
 399

400 3.2.2 Sediment Carbon Sources

401 In Walker salt marsh, surface (< 10 cm) wrack sediments had higher $\delta^{13}\text{C}$ values than
402 sediments collected from the interior of the marsh (non-wrack sediments) and thus were more
403 similar to the $\delta^{13}\text{C}$ values of seagrass (Fig. 6; SLM, $p < 0.05$; F-statistic =27.3; DF = 18).
404 Shallow wrack sediments had an average $\delta^{13}\text{C}$ of -22.5 ± 0.38 ‰, while non-wrack sediments
405 had an average $\delta^{13}\text{C}$ of -24.9 ± 0.26 ‰ (mean \pm SE). However, when both shallow and deep
406 (>10cm) sediments were included, wrack sediment $\delta^{13}\text{C}$ did not significantly differ from non-
407 wrack sediments (SLM, $p > 0.05$, F-statistic = 0.34, DF = 31). C:N sediment ratios did not
408 significantly differ from one another regardless of collection depth or location (SLMs, $p > 0.05$).
409 When data were applied to a mixing model, apparent discrepancies in seagrass-derived carbon
410 contributions between shallow and deep, and wrack and non-wrack sediments were insignificant
411 (SIAR mixing model, Table S2). Similarly, no significant quantity of seagrass-derived carbon
412 was detected in Elkhorn salt marsh sediments (SIAR mixing model, Table S2). Rather, models
413 estimate that sediments were derived almost entirely from either diatoms or C3 plants – 83% in
414 Elkhorn Slough and 88% in Walker salt marsh (Fig. 7; Table S2). Thus, regardless of site, depth,
415 or the presence of wrack, model results estimate no significant storage of seagrass-derived
416 carbon in sediment (Table S2). Instead, C3 plants and/or diatoms are the primary significant
417 contributors to underlying carbon storage across salt marshes in all locations.

418



419
 420
 421
 422
 423
 424
 425
 426
 427

Figure 6: Points represent the $\delta^{13}\text{C}$ and C:N ratios of Walker Salt Marsh sediment data, collected from underneath persistent wrack lines (blue) and from areas free of wrack (red) (see Fig. 1c). Sediment subsamples collected from the surface to 10cm deep are labelled as surface samples (square) and below 10cm are labelled as deep samples (circle). Colored boxes represent sources (means \pm SD) from diatoms (red), seagrass (green), C4 salt marsh plants (blue), and C3 salt marsh plants (red).




Figure 7 (left): Mixing model results displaying the average source contributions to salt marsh sediment organic carbon (OC) for two salt marshes. Exact values and significance are shown in Table S2.

441 4 Discussion

442 4.1 Carbon Stocks

443 The lack of clear downcore trends (Fig. 2) could suggest relative OC stability with depth and
444 through time or that there have been minimal changes to factors such as vegetation and grain size
445 through time. The persistence of this pattern is supported by the longer cores sampled in seagrass
446 and bare sediment sites (Fig. S3). One possible explanation for significant downcore OC losses
447 in Newport Bay salt marsh is change in historic hydrography. Specifically, previous sediment
448 profiles from this region also found finer surface sediments above coarser materials – associated
449 with the conversion from an erosional to a depositional system in the early 20th century due to
450 urbanization of the surrounding area (Trimble, 2003). This change in grain size down core may
451 drive the associated OC loss down core, discussed further below. Although we did not quantify
452 sediment carbon accumulation rates here, previous work in Tomales Bay seagrass meadows
453 indicate rates of 11.37 - 15.16 g C m⁻² yr⁻¹ via ²¹⁰Pb dating (O'Donnell, 2017). By these
454 accumulation rate estimates, we estimate the top 20 cm of sediment sampled here accumulated
455 over approximately 100-130 years. Given the relatively slow accumulation rates and lack of
456 down core trends, we interpret that our data represent realistic stock assessments for each
457 respective habitat and location, rather than being reflective of shallow surface sediment carbon
458 stocks.

459 Determining drivers of OC storage variation across habitats is notably complex, where a mix
460 of factors such as grain size, elevation, hydrodynamic energy, and vegetation type may influence
461 underlying sediment OC storage (Kelleway et al., 2016; Lima et al., 2020; Miyajima et al.,
462 2015). Here, sediment grain size and its associated drivers allow us to further examine variation
463 in sediment OC storage variation. Specifically, grain size likely contributed to the observed inter-

464 habitat differences in OC storage, given its strong correlation with TOM. Despite no significant
465 differences in grain size between habitat types within each site (likely due to high grain size
466 variability and low relative sample sizes), apparent qualitative differences elude to possible
467 explanations of stock differences. For example, seagrass sediments had lower % mud than salt
468 marsh sediments in Tomales Bay, a trend that was reversed in Newport Bay (Fig. 5a; Table S1).
469 Similarly, seagrass sediments did not always contain a higher % mud than neighboring bare
470 sediments, contrary to what is typically reported (Bos et al., 2007; Conley et al., 2017; Mazarassa
471 et al., 2015), which may explain their apparent lack of sediment OC differences. Overall, salt
472 marshes had greater OC storage and smaller average grain size than seagrass meadows. The
473 periodic, lower energy inundation of the sampled marsh and pan habitats may facilitate the
474 observed smaller grain sizes – by preventing the resuspension of small particles that might occur
475 in higher hydrodynamic energy or submerged systems (Christiansen et al., 2000; Yang et al.,
476 2008).

477 In marsh and pan habitats, biological factors such as primary production and decomposition
478 rates may become more important drivers of carbon burial than would be true in dominantly
479 sandy sediments (Miyajima et al., 2017). It is possible that root systems of overlying vegetation
480 add bulk organic material into available sediment space, contributing to increased carbon
481 deposition as well as decreased space for mineral accumulation (e.g., Rogers et al., 2019). Yet
482 despite the absence of vegetation in pan sites, we did not observe significant differences in
483 carbon stocks between pan sediments and surrounding salt marsh sediments. This could be
484 attributed to 1) significant carbon contributions from surrounding salt marsh biomass to pan sites
485 overtime, 2) historical marsh recovery along pan edges and subsequent burial of vegetation
486 captured in pan cores, 3) variable decomposition rates in both pan and salt marsh sediments, or

487 4) because the expected effect of canopy-vegetation on salt marsh sediment is too small relative
 488 to other drivers, producing statistically insignificant results. In sum, although seagrasses and salt
 489 marshes can facilitate carbon storage by altering grain size distributions, hydrodynamics and
 490 geomorphology play a critical role in grain size distributions and therefore, carbon storage. The
 491 hydrographic changes in Newport Bay salt marshes associated with decreased grain sizes further
 492 emphasize this point – drawing attention to the importance of watershed and sediment
 493 management to regional carbon storage. Furthermore, although grain size is clearly a key driver
 494 in carbon storage, our data demonstrate that it becomes of limited use as a predictor of carbon
 495 stocks after the proportion of mud exceeds 82% (Fig. 4). This demonstrates that using grain size
 496 as a cost-effective way to estimate carbon stocks (as has been suggested, e.g. Serrano et al.,
 497 2016) only appears possible in sandy and mixed grain size sediments within the sites studied
 498 here. Carbon stocks in fine sediment sites above this threshold cannot be estimated using grain
 499 size alone – a management-relevant finding for efforts to incorporate habitat-specific carbon
 500 storage into regional climate plans.

501

502 **4.2 Global and Regional Comparisons**

503 With recent reviews on coastal carbon stocks in salt marshes and seagrass meadows, we can
 504 compare our data to global and regional averages (no sufficient syntheses of pan or bare
 505 sediment carbons stocks were identified). We find that carbon storage in California seagrass
 506 sediments studied here is lower than global estimates yet is higher than regional estimates of Z.

507

Study	Habitat Type/Species	Location(s)	Value (Mg C_{org} ha⁻¹)	Methods (core length and extrapolation depth)
Seagrass				

This study	Seagrass (<i>Z. marina</i>)	California	110 ± 11.8	20 cm cores extrapolated to 1 m
Kauffman et al. (2020)	Seagrass (<i>Z. marina</i>)	*U.S. west coast	80 ± 12	1 m cores collected (no extrapolation)
Prentice et al. (2020)	Seagrass (<i>Z. marina</i>)	U.S. west coast	65.12	25 cm cores extrapolated to 1 m
Röhr et al. (2018)	Seagrass (<i>Z. marina</i>)	*U.S. west coast *Temperate Northern Hemisphere	69.4 ± 8.4 108.9 ± 39.56	25 cm cores extrapolated to 1 m
Fourqurean et al. (2012)	Seagrass (many species)	Global	¹ 139.7	A combined estimate from short cores (≥ 20cm & < 1 m) extrapolated to 1 m and full cores (1 m)
Salt Marsh				
This study	Salt marsh dominated by <i>Sarcocornia pacifica</i> , <i>Distichlis spicata</i> and <i>Jaumea carnosa</i>	California	235 ± 17.7	20 cm cores extrapolated to 1m
Kauffman et al. (2020)	Salt marsh dominated by <i>Distichlis spicata</i> and <i>Sarcocornia perennis</i>	*U.S. west coast	190 ± 16	1 m cores collected (no extrapolation)
Holmquist et al. (2018)	Tidal wetlands (mixed, dominated by estuarine emergent wetlands)	*Conterminous U.S. tidal wetlands	270 ± 1.4	A combined estimate from short cores (≥ 10cm & < 1 m) extrapolated to 1 m and full cores (1 m)

508
509
510
511
512
513

Table 3: Summary of previously reported carbon stocks in seagrass meadows and tidal wetlands, all normalized to the top 1m of sediment. Studies that included any data from California are denoted by an asterisk. Values are reported as mean ± SE, unless otherwise noted. ¹Fourqurean et al. (2012) value represents median OC storage.

514
515

marina from the U.S. west coast (Table 3). Early estimates from the U.S. west coast were markedly low, with one study of *Z. marina* from British Columbia measuring average stocks of

516 1.34 kg C m⁻³ with a maximum OC of 1.3%, compared to the 11.01 ± 1.18 kg C m⁻³ and a
517 maximum OC of 7.0 % estimated here (Postlethwaite et al., 2018). While estimates from studies
518 averaging over broader U.S. west coast regions are higher than this Canadian case study (Table
519 3), the data presented here are still higher than regional estimates, suggesting possible latitudinal
520 difference in carbon storage between the northern and southern regions of the North American
521 west coast. This is not surprising given the variation in environmental and geomorphic
522 conditions over this large stretch of coastline, which likely influence the carbon storage capacity
523 along with the factors discussed here such as hydrodynamics or grain size conditions.

524 Recent studies of the U.S. west coast and of conterminous U.S. tidal wetlands allow for
525 comparison with salt marsh data collected here. Much like with seagrass meadows, California
526 salt marsh carbon storage was also higher than previous estimates from the U.S. west coast
527 (Kauffman et al., 2020; Table 3). Both the data presented here as well as those from Kauffman et
528 al. (2020) (Table 3) were collected from similar marsh elevation zones and species compositions,
529 reducing some potential for these factors to result in the observed differences in carbon stocks.
530 Previous work has found that high-marsh zone sediments are likely to contain greater OC than
531 mid- and lower-marsh zones, and that less inundation can facilitate increased root productivity
532 and increased OC (the sites presented here would be considered ‘lower-marsh zones’) (Blum,
533 1993; Connor et al., 2001; Zhou et al., 2007). Moreover, sediments beneath other common marsh
534 species in this region (e.g. *Spartina sp.*) were not sampled, yet may have differing productivity
535 levels, resulting in OC stock differences. This may contribute to the variation between our
536 estimated salt marsh carbon stock and that of conterminous U.S. tidal wetlands (Holmquist et al.,
537 2018), which includes a variety of habitat types including forested and shrub-dominated tidal
538 wetlands.

539

540 **4.3 Carbon Sources and Exchanges**

541 Across both salt marshes assessed in this study, diatoms and/or C3 plants were the
542 dominant carbon sources in sediment. Although the isotopic values resulting from C4 and
543 seagrass sources are poorly resolved, lack of a significant values from either source allows for
544 interpretation. Specifically, seagrasses do not significantly contribute to any OC sources in
545 sediment, while C4 plants may contribute minimally to OC in sediment in Elkhorn Slough (1.1-
546 11.9%; Table S2). Thus, despite similar isotopic values between seagrass and C4 plants, if
547 seagrass-derived organic matter was buried under wrack sediments, we would expect to see
548 higher $\delta^{13}\text{C}$ values in wrack sediments with depth. The insignificant differences in $\delta^{13}\text{C}$ between
549 wrack and non-wrack sediments suggest that seagrass is not ultimately being buried, but rather,
550 carbon derived from C3 plants and diatoms are ultimately retained in sediment.

551 Several underlying causes may lead to this high proportion of diatom and/or C3-derived
552 carbon (Fig. 7). The recalcitrance of both autochthonous and allochthonous material in sediment
553 can vary depending on the material's composition (e.g. Burdige, 2007), and thus knowing the
554 sources and composition of deposited material aids in understanding its chance of long-term
555 burial. In seagrass meadows, previous work demonstrated that seagrass BGB likely contributes
556 most to autochthonous carbon burial when compared to AGB, due to higher proportions of
557 refractory compounds and decreased grazing pressure (Trevathan-Tackett et al., 2017). This may
558 serve to explain our lack of seagrass signal in salt marsh sediments under wrack. Seagrass wrack
559 material deposited on top of the salt marsh is predominantly AGB, likely breaking down or being
560 transported elsewhere before any appreciable portion reaches long-term sediment carbon pools.
561 Instead, the dominant C3 signal is likely driven by the presence of pickleweed species

562 (*Sarcocornia sp.*), which have a considerable portion of below ground biomass. Despite the
563 presence of salt grass (*Distichlis spicata*), the relative lack of C4-derived carbon in sediment
564 may be due to a few potential factors: 1) salt grass is less common relative to pickleweed within
565 our sites, 2) based on our field observations, root systems of salt grass do not seem to penetrate
566 as deeply or contain as much BGB as do those of pickleweed, and 3) salt grass contains less
567 woody tissue than pickleweed, making it less refractory (Jepson Flora Project, 2020). In the case
568 of diatom and planktonic sources, their significant presence in sediment may be due to a greater
569 abundance overall, or due to complex preservation pathways such as the facilitation of carbon
570 burial by binding sediments through extracellular polymeric substances (Drexler et al., 2020;
571 Macreadie et al., 2019; Oakes and Eyre, 2013).

572 From the seasonal senescence indicated by seagrass meadow biomass data (Table 2), we
573 infer that a significant amount of seagrass AGB is either degraded within the meadow or
574 exported for remineralization or deposition elsewhere. Given that strong tidal flows can occur
575 within meadows and that high densities of seagrass wrack were observed in a neighboring salt
576 marsh, lateral export of AGB from the meadow is likely. Nonetheless, the majority of OC
577 deposited into the marsh in the form of seagrass wrack is likely remineralized over the course of
578 the year – with wrack decomposition outpacing a sediment accumulation rate that might support
579 carbon preservation. While wrack remineralization can support local estuarine food web
580 metabolism, it can also produce emissions (Jiménez et al., 2017; Liu et al., 2019). However,
581 these emissions are small when compared to the levels of carbon sequestration within the marsh
582 – if all of the seagrass wrack along the wrack lines was remineralized annually, this degradation
583 would only contribute $33 \pm 7.8 \text{ g C m}^{-2}\text{yr}^{-1}$ to the atmosphere (Table 2). For comparison, dating
584 (primarily ^{137}Cs) from salt marshes nearby estimate carbon accumulation as $174 \pm 45 \text{ g C m}^{-2}\text{yr}^{-1}$

585 (Ouyang & Lee, 2014). Thus, even along wrack lines, carbon accumulation far outpaces carbon
586 release from breakdown of allochthonous material. Although estimates of decomposition and
587 accumulation rates in this site could further inform these concepts, the slow annual cycle of
588 seagrass wrack deposition and disappearance observed in the aerial imagery (Fig. S2), and the
589 lack of seagrass-derived carbon in underlying sediment make a compelling case that little of this
590 material is ultimately buried.

591

592 **5 Conclusions**

593 We find that California salt marshes can store approximately twice as much carbon than do
594 seagrass meadows within this region. Grain size – an easier metric to quantify than carbon stock
595 – can be used to estimate regional carbon storage in sandy and mixed-grain size sediments. This
596 information can serve to inform local and regional management plans in efforts to prioritize and
597 quantify carbon storage across these habitat types. While seagrass meadows may act as local
598 sinks for carbon, they also export substantial amounts of AGB annually, which can be
599 remineralized and converted to CO₂, rather than buried in neighboring blue carbon habitats. C3
600 salt marsh plants and/or diatoms contributed to the OC pools in all three salt marshes under study
601 – likely due to their prevalence within our sites and their resistance to degradation. This
602 comprehensive study assesses several key research needs in blue carbon science informing
603 current efforts to prioritize and quantify carbon storage across these habitat types.

604

605 **6 Data Availability**

606 All data included in this manuscript will be made publicly available on the Dryad data
607 repository.

608

609 **7 Competing Interests**

610 All data included in this manuscript will be made publicly available on the Dryad data
611 repository.

612

613 **8 Author Contributions**

614 Ward – conceptualization, investigation, methodology, writing (original draft preparation),
615 formal analysis; Hill – conceptualization, funding support, writing (review & editing); Souza –
616 investigation (field and lab support), writing (review & editing); Filipczyk – investigation (field
617 and lab support), writing (review & editing); Ricart – methodology, investigation, writing
618 (review & editing); Merolla – investigation (field and lab support), writing (review & editing);
619 Capece – writing (review & editing); O’Donnell – investigation (field and lab support),
620 methodology; writing (review & editing);); Elsmore – investigation (field and lab support),
621 writing (review & editing); Oechel – writing (review & editing); Beheshti – investigation (field
622 and lab support), writing (review & editing).

623

624 **9 Acknowledgements**

625 We acknowledge Al Carranza, Grant Susner, Amanda Nordstrom, Siena Watson, Camille Frias,
626 Daphne Bradley, Anya Morrill, Natalie Rossi, and Jezebel Powers for laboratory and field
627 assistance. This work was supported by California Sea Grant (R/HCME-03, support to Tessa
628 Hill, Brian Gaylord, Eric Sanford, and Kristy Kroeker), the California Ocean Protection Council
629 (support to Tessa Hill, Brian Gaylord, Eric Sanford, and Kristy Kroeker), the CSU Council on
630 Ocean Affairs, Science & Technology (support to Melissa Ward), and the San Diego ARCS

631 foundation (support to Melissa Ward). Dr. Jim Cloern also generously provided data to support
632 isotopic mixing models. We also thank Audubon Canyon Ranch for site access from their
633 Cypress Grove Research Center. Lastly, we would like to thank Dr. Peter Bowler, whose
634 mentorship and legacy in wetland conservation has sparked lifelong collaborations and scientific
635 careers.

636
637 **10 References**

- 638
639 Alongi, D. M. Blue carbon coastal sequestration for climate change mitigation.
640 Springer International Publishing, Briefs in Climate Studies, doi: 10.1007/978-3-319-91698-9,
641 2018.
- 642
643 Attard, K. M., Rodil, I. F., Berg, P., Norkko, J., Norkko, A. and Glud, R. N.: Seasonal
644 metabolism and carbon export potential of a key coastal habitat: The perennial canopy-forming
645 macroalga *Fucus vesiculosus*, *Limnology and Oceanography*, 64(1), 149–164,
646 doi:10.1002/lno.11026, 2019.
- 647
648 Benner, R., Fogel, M. L. and Sprague, E. K.: Diagenesis of belowground biomass of *Spartina*
649 *alterniflora* in salt-marsh sediments, *Limnology and Oceanography*, 36(7), 1358–1374,
649 doi:10.4319/lo.1991.36.7.1358, 1991.
- 650
651 Blum, L. K.: *Spartina alterniflora* root dynamics in a Virginia marsh, *Marine Ecology Progress*
651 *Series*, 102, 169–178, 1993.
- 652
653 Bos, A. R., Bouma, T. J., de Kort, G. L. J. and van Katwijk, M. M.: Ecosystem engineering by
654 annual intertidal seagrass beds: Sediment accretion and modification, *Estuarine, Coastal and*
654 *Shelf Science*, 74(1–2), 344–348, doi:10.1016/j.ecss.2007.04.006, 2007.
- 655
656 Bouillon, S. and Connolly, R. M.: Carbon Exchange Among Tropical Coastal Ecosystems, in
657 *Ecological Connectivity among Tropical Coastal Ecosystems*, edited by I. Nagelkerken, pp. 45–
657 70, Springer Netherlands, Dordrecht., 2009.
- 658
659 Brevik, E. C. and Homburg, J. A.: A 5000 year record of carbon sequestration from a coastal
660 lagoon and wetland complex, Southern California, USA, *CATENA*, 57(3), 221–232,
660 doi:10.1016/j.catena.2003.12.001, 2004.
- 661
662 Burdige, D. J.: Preservation of Organic Matter in Marine Sediments: Controls, Mechanisms, and
663 an Imbalance in Sediment Organic Carbon Budgets?, *Chem. Rev.*, 107(2), 467–485,
663 doi:10.1021/cr050347q, 2007.

- 664 Cabello-Pasini, A., Muñiz-Salazar, R. and Ward, D. H.: Annual variations of biomass and
665 photosynthesis in *Zostera marina* at its southern end of distribution in the North Pacific, *Aquatic*
666 *Botany*, 76(1), 31–47, doi:10.1016/S0304-3770(03)00012-3, 2003.
- 667 Callaway, J. C., Borgnis, E. L., Turner, R. E. and Milan, C. S.: Carbon Sequestration and
668 Sediment Accretion in San Francisco Bay Tidal Wetlands, *Estuaries and Coasts*, 35(5), 1163–
669 1181, doi:10.1007/s12237-012-9508-9, 2012.
- 670 Capece, L. The origin of sedimentary organic carbon in temperate seagrass meadows in
671 California estuaries. Thesis 22619435, University of California, Davis, ProQuest Dissertations
672 Publishing, 2019.
- 673
674 Chmura, G. L., Anisfeld, S. C., Cahoon, D. R. and Lynch, J. C.: Global carbon sequestration in
675 tidal, saline wetland soils, *Global Biogeochemical Cycles*, 17(4), doi:10.1029/2002GB001917,
676 2003.
- 677 Christiansen, T., Wiberg, P. L. and Milligan, T. G.: Flow and sediment transport on a tidal salt
678 marsh surface, *Estuarine, Coastal and Shelf science*, 50(3), 315–331,
679 <https://doi.org/10.1006/ecss.2000.0548>, 2000.
- 680 Cloern, J. E., Canuel, E. A. and Harris, D.: Stable carbon and nitrogen isotope composition of
681 aquatic and terrestrial plants of the San Francisco Bay estuarine system, *Limnology and*
682 *Oceanography*, 47(3), 713–729, doi:10.4319/lo.2002.47.3.0713, 2002.
- 683 Conley, D. C., Austin, M., Davidson, I., Buscombe, D. and Masselink, G.: Grain size selection in
684 seagrass beds, *coastal dynamics*, (200), 11, 2017.
- 685 Connor, R. F., Chmura, G. L. and Beecher, C. B.: Carbon accumulation in bay of fundy salt
686 marshes: Implications for restoration of reclaimed marshes, *Global Biogeochemical Cycles*,
687 15(4), 943–954, doi:10.1029/2000GB001346, 2001.
- 688 Craft, C. B., Seneca, E. D. and Broome, S. W.: Loss on Ignition and Kjeldahl Digestion for
689 Estimating Organic Carbon and Total Nitrogen in Estuarine Marsh Soils: Calibration with Dry
690 Combustion, *Estuaries*, 14(2), 175, doi:10.2307/1351691, 1991.
- 691 Craven, K. F., Edwards, R. J. and Flood, R. P.: Source organic matter analysis of saltmarsh
692 sediments using SIAR and its application in relative sea-level studies in regions of C₄ plant
693 invasion, *Boreas*, 46(4), 642–654, doi:10.1111/bor.12245, 2017.
- 694 Cyronak, T., Andersson, A. J., D’Angelo, S., Bresnahan, P., Davidson, C., Griffin, A.,
695 Kindeberg, T., Pennise, J., Takeshita, Y. and White, M.: Short-Term Spatial and Temporal
696 Carbonate Chemistry Variability in Two Contrasting Seagrass Meadows: Implications for pH
697 Buffering Capacities, *Estuaries and Coasts*, 41(5), 1282–1296, doi:10.1007/s12237-017-0356-5,
698 2018.
- 699 Dahl, M., Deyanova, D., Gütschow, S., Asplund, M. E., Lyimo, L. D., Karamfilov, V., Santos,
700 R., Björk, M. and Gullström, M.: Sediment properties as important predictors of carbon storage

- 701 in *Zostera marina* meadows: A Comparison of four European areas, PLoS ONE, 11(12),
702 e0167493, <https://doi.org/10.1371/journal.pone.0167493>, 2016.
- 703 Dean, W. E.: Determination of carbonate and organic matter in calcareous sediments and
704 sedimentary rocks by loss on ignition; comparison with other methods, *Journal of Sedimentary*
705 *Research*, 44, 242–248, <https://doi.org/10.1306/74D729D2-2B21-11D7-8648000102C1865D>,
706 1974.
- 707
- 708 Drexler, J. Z., Davis, M. J., Woo, I. and De La Cruz, S.: Carbon sources in the sediments of a
709 restoring vs. historically unaltered salt marsh, *Estuaries and Coasts*, 43(6), 1345–1360,
710 <https://doi.org/10.1007/s12237-020-00748-7>, 2020.
- 711 Duarte, C., Marbà, N., Gacia, E., Fourqurean, J. W., Beggins, J., Barrón, C. and Apostolaki, E.
712 T.: Seagrass community metabolism: Assessing the carbon sink capacity of seagrass meadows,
713 *Global Biogeochemical Cycles*, 24(4), doi:10.1029/2010GB003793, 2010.
- 714 Duarte, C. M. and Cebrián, J.: The fate of marine autotrophic production, *Limnology and*
715 *Oceanography*, 41(8), 1758–1766, doi:10.4319/lo.1996.41.8.1758, 1996.
- 716 Duarte, C. M. and Krause-Jensen, D.: Export from seagrass meadows contributes to marine
717 carbon sequestration, *Front. Mar. Sci.*, 4, doi:10.3389/fmars.2017.00013, 2017.
- 718 Escapa, M., Perillo, G. M. E. and Iribarne, O.: Biogeomorphically driven salt pan formation in
719 *Sarcocornia*-dominated salt-marshes, *Geomorphology*, 228, 147–157,
720 doi:10.1016/j.geomorph.2014.08.032, 2015.
- 721 Fourqurean, J. W., Duarte, C. M., Kennedy, H., Marbà, N., Holmer, M., Mateo, M. A.,
722 Apostolaki, E. T., Kendrick, G. A., Krause-Jensen, D., McGlathery, K. J. and Serrano, O.:
723 Seagrass ecosystems as a globally significant carbon stock, *Nature Geoscience*, 5(7), 505–509,
724 doi:10.1038/ngeo1477, 2012.
- 725 Fourqurean, J. W., Moore, T. O., Fry, B., and Hollibaugh, J. T.: Spatial and temporal variation in
726 C:N:P ratios, $\delta^{15}\text{N}$, and $\delta^{13}\text{C}$ of eelgrass *Zostera marina* as indicators of ecosystem processes,
727 Tomales Bay, California, USA, 157, 147–157, <https://doi.org/10.3354/meps157147>, 1997.
- 728 Freedman, B., Stinson, G. and Lacoul, P.: Carbon credits and the conservation of natural areas,
729 *Environ. Rev.*, 17(NA), 1–19, doi:10.1139/A08-007, 2009.
- 730 Gambi, M., Nowell, A. and Jumars, P.: Flume observations on flow dynamics in *Zostera marina*
731 (eelgrass) beds, *Mar. Ecol. Prog. Ser.*, 61, 159–169, doi:10.3354/meps061159, 1990.
- 732 Google Earth. Tomales Bay, California, USA. 38°12'42.4"N 122°55'39.7"W. 2020.
- 733
- 734 Greiner, J. T., McGlathery, K. J., Gunnell, J. and McKee, B. A.: Seagrass restoration enhances
735 “Blue Carbon” sequestration in coastal waters, PLoS ONE, 8(8), e72469,
736 doi:10.1371/journal.pone.0072469, 2013.
- 737 Green, E.P. and Short, F.T. (Eds.) (2003). *World Atlas of Seagrasses. University of California*

738 *Press, Berkeley, USA*, 324 pp.
739

740 Hendriks, I., Sintes, T., Bouma, T. and Duarte, C.: Experimental assessment and modeling
741 evaluation of the effects of the seagrass *Posidonia oceanica* on flow and particle trapping,
742 *Marine Ecology Progress Series*, 356, 163–173, doi:10.3354/meps07316, 2008.

743 Holmquist, J. R., Windham-Myers, L., Bliss, N., Crooks, S., Morris, J. T., Megonigal, J. P.,
744 Troxler, T., Weller, D., Callaway, J., Drexler, J., Ferner, M. C., Gonneea, M. E., Kroeger, K. D.,
745 Schile-Beers, L., Woo, I., Buffington, K., Breithaupt, J., Boyd, B. M., Brown, L. N., Dix, N.,
746 Hice, L., Horton, B. P., MacDonald, G. M., Moyer, R. P., Reay, W., Shaw, T., Smith, E., Smoak,
747 J. M., Sommerfield, C., Thorne, K., Velinsky, D., Watson, E., Grimes, K. W. and Woodrey, M.:
748 accuracy and precision of tidal wetland soil carbon mapping in the conterminous United States,
749 *Scientific Reports*, 8(1), 9478, doi:10.1038/s41598-018-26948-7, 2018.

750 Howard, J., Hoyt, S., Isensee, K., Pidgeon, E., Telszewski, M. (eds.) (2014). Coastal Blue
751 Carbon: Methods for assessing carbon stocks and emissions factors in mangroves, tidal salt
752 marshes, and seagrass meadows. Conservation International, Intergovernmental Oceanographic
753 Commission of UNESCO, International Union for Conservation of Nature. Arlington, Virginia,
754 USA.

755 Hyndes, G. A., Nagelkerken, I., McLeod, R. J., Connolly, R. M., Lavery, P. S. and Vanderklift,
756 M. A.: Mechanisms and ecological role of carbon transfer within coastal seascapes, *Biological*
757 *Reviews*, 89(1), 232–254, doi:10.1111/brv.12055, 2014.

758 Ince, R., Hyndes, G. A., Lavery, P. S. and Vanderklift, M. A.: Marine macrophytes directly
759 enhance abundances of sandy beach fauna through provision of food and habitat, *Estuarine,*
760 *Coastal and Shelf Science*, 74(1–2), 77–86, doi:10.1016/j.ecss.2007.03.029, 2007.

761 Jepson Flora Project (eds.) 2020, Jepson eFlora, <https://ucjeps.berkeley.edu/eflora/>, accessed on
762 February 5th, 2020.

763 Jiménez, M. A., Beltran, R., Traveset, A., Calleja, M. L., Delgado-Huertas, A. and Marbà, N.:
764 Aeolian transport of seagrass (*Posidonia oceanica*) beach-cast to terrestrial systems, *Estuarine,*
765 *Coastal and Shelf Science*, 196, 31–44, doi:10.1016/j.ecss.2017.06.035, 2017.

766 Johannessen, S. C. and Macdonald, R. W.: Geoengineering with seagrasses: is credit due where
767 credit is given?, *Environmental Research Letters*, 11, 113001, [https://doi.org/10.1088/1748-](https://doi.org/10.1088/1748-9326/11/11/113001)
768 [9326/11/11/113001](https://doi.org/10.1088/1748-9326/11/11/113001), 2016.

769 Kauffman, J. B., Giovanonni, L., Kelly, J., Dunstan, N., Borde, A., Diefenderfer, H., Cornu, C.,
770 Janousek, C., Apple, J. and Brophy, L.: Total ecosystem carbon stocks at the marine-terrestrial
771 interface: Blue carbon of the Pacific Northwest Coast, United States, *Glob. Change Biol.*, 26(10),
772 5679–5692, doi:10.1111/gcb.15248, 2020.

773 Kelleway, J. J., Saintilan, N., Macreadie, P. I. and Ralph, P. J.: Sedimentary factors are key
774 predictors of carbon storage in SE Australian saltmarshes, *Ecosystems*, 19(5), 865–880,
775 doi:10.1007/s10021-016-9972-3, 2016.

776 Kennedy, H., Beggins, J., Duarte, C. M., Fourqurean, J. W., Holmer, M., Marbà, N. and
777 Middelburg, J. J.: Seagrass sediments as a global carbon sink: Isotopic constraints, *Global*
778 *Biogeochem. Cycles*, 24(4), GB4026, doi:10.1029/2010GB003848, 2010.

779 Largier, J. L., Hollibaugh, J. T. and Smith, S. V.: Seasonally hypersaline estuaries in
780 mediterranean-climate regions, *Estuarine, Coastal and Shelf Science*, 45(6), 789–797,
781 doi:10.1006/ecss.1997.0279, 1997.

782 Lavery, P. S., Mateo, M.-Á., Serrano, O. and Rozaimi, M.: Variability in the carbon storage of
783 seagrass habitats and its implications for global estimates of Blue Carbon ecosystem service,
784 *PLoS ONE*, 8(9), e73748, doi:10.1371/journal.pone.0073748, 2013.

785 Leorri, E., Zimmerman, A. R., Mitra, S., Christian, R. R., Fatela, F. and Mallinson, D. J.:
786 Refractory organic matter in coastal salt marshes-effect on C sequestration calculations, *Science*
787 *of the Total Environment*, 633, 391–398, doi:10.1016/j.scitotenv.2018.03.120, 2018.

788 Lewis, C. J., Young, M. A., Ierodiaconou, D., Baldock, J. A., Hawke, B., Sanderman, J., Carnell,
789 P. E. and Macreadie, P. I.: Drivers and modelling of blue carbon stock variability in sediments of
790 southeastern Australia, *Biogeosciences*, 17(7), 2041–2059, doi:10.5194/bg-17-2041-2020, 2020.

791 Lima, M. do A. C., Ward, R. D. and Joyce, C. B.: Environmental drivers of sediment carbon
792 storage in temperate seagrass meadows, *Hydrobiologia*, doi:10.1007/s10750-019-04153-5, 2020.

793 Liu, S., Trevathan-Tackett, S. M., Ewers Lewis, C. J., Ollivier, Q. R., Jiang, Z., Huang, X. and
794 Macreadie, P. I.: Beach-cast seagrass wrack contributes substantially to global greenhouse gas
795 emissions, *Journal of Environmental Management*, 231, 329–335,
796 doi:10.1016/j.jenvman.2018.10.047, 2019.

797 Lovelock, C. E., Adame, M. F., Bennion, V., Hayes, M., O’Mara, J., Reef, R. and Santini, N. S.:
798 Contemporary rates of carbon sequestration through vertical accretion of sediments in mangrove
799 forests and saltmarshes of South East Queensland, Australia, *Estuaries and Coasts*, 37(3), 763–
800 771, doi:10.1007/s12237-013-9702-4, 2014.

801 Lovelock, C. E., Atwood, T., Baldock, J., Duarte, C. M., Hickey, S., Lavery, P. S., Masque, P.,
802 Macreadie, P. I., Ricart, A. M., Serrano, O. and Steven, A.: Assessing the risk of carbon dioxide
803 emissions from blue carbon ecosystems, *Frontiers in Ecology and the Environment*, 15(5), 257–
804 265, doi:10.1002/fee.1491, 2017.

805 Lovelock, C. E. and Duarte, C. M.: Dimensions of Blue Carbon and emerging perspectives,
806 *Biology Letters*, 15, 20180781, <https://doi.org/10.1098/rsbl.2018.0781>, 2019.

807 Macreadie, P. I., Ewers-Lewis, C. J., Whitt, A. A., Ollivier, Q., Trevathan-Tackett, S. M.,
808 Carnell, P., and Serrano, O.: Comment on ‘Geoengineering with seagrasses: is credit due where
809 credit is given?’, *Environ. Res. Lett.*, 13, 028002, <https://doi.org/10.1088/1748-9326/aaa7ad>,
810 2018.

811 Macreadie, P. I., Anton, A., Raven, J. A., Beaumont, N., Connolly, R. M., Friess, D. A.,
812 Kelleway, J. J., Kennedy, H., Kuwae, T., Lavery, P. S., Lovelock, C. E., Smale, D. A.,

813 Apostolaki, E. T., Atwood, T. B., Baldock, J., Bianchi, T. S., Chmura, G. L., Eyre, B. D.,
814 Fourqurean, J. W., Hall-Spencer, J. M., Huxham, M., Hendriks, I. E., Krause-Jensen, D.,
815 Laffoley, D., Luisetti, T., Marbà, N., Masque, P., McGlathery, K. J., Megonigal, J. P.,
816 Murdiyarso, D., Russell, B. D., Santos, R., Serrano, O., Silliman, B. R., Watanabe, K. and
817 Duarte, C. M.: The future of Blue Carbon science, *Nature Communications*, 10(1), 1–13,
818 doi:10.1038/s41467-019-11693-w, 2019.

819 Mazarrasa, I., Lovelock, C. E., Serrano, O., Lavery, P. S., Fourqurean, J. W., Kennedy, H.,
820 Mateo, M. A., Krause-Jensen, D., Steven, A. D. L. and Duarte, C. M.: Seagrass meadows as a
821 globally significant carbonate reservoir, *Biogeosciences*, 12(16), 4993–5003,
822 doi:https://doi.org/10.5194/bg-12-4993-2015, 2015.

823 Mazarrasa, I., Samper-Villarreal, J., Serrano, O., Lavery, P. S., Lovelock, C. E., Marbà, N.,
824 Duarte, C. M. and Cortés, J.: Habitat characteristics provide insights of carbon storage in
825 seagrass meadows, *Marine Pollution Bulletin*, 134, 106–117,
826 doi:10.1016/j.marpolbul.2018.01.059, 2018.

827 Mcleod, E., Chmura, G. L., Bouillon, S., Salm, R., Björk, M., Duarte, C. M., Lovelock, C. E.,
828 Schlesinger, W. H., and Silliman, B. R.: A blueprint for blue carbon: toward an improved
829 understanding of the role of vegetated coastal habitats in sequestering CO₂, *Frontiers in Ecology
830 and the Environment*, 9, 552–560, https://doi.org/10.1890/110004, 2011.

831 Merkel & Associates, Inc. 2017 Tomales Bay Eelgrass Inventory, prepared the National Oceanic
832 Atmospheric Administration (NOAA) Greater Farallones National Marine Sanctuary. December
833 2017.
834

835 Milliman, J. D.: Precipitation and Cementation of Deep-Sea Carbonate Sediments, in *Deep-Sea
836 Sediments: Physical and Mechanical Properties*, edited by A. L. Inderbitzen, pp. 463–476,
837 Springer US, Boston, MA., 1974.

838 Miyajima, T., Hori, M., Hamaguchi, M., Shimabukuro, H., Adachi, H., Yamano, H. and
839 Nakaoka, M.: Geographic variability in organic carbon stock and accumulation rate in sediments
840 of East and Southeast Asian seagrass meadows, *Global Biogeochemical Cycles*, 29(4), 397–415,
841 doi:10.1002/2014GB004979, 2015.

842 Miyajima, T., Hori, M., Hamaguchi, M., Shimabukuro, H., and Yoshida, G.: Geophysical
843 constraints for organic carbon sequestration capacity of *Zostera marina* seagrass meadows and
844 surrounding habitats, 62, 954–972, https://doi.org/10.1002/lno.10478, 2017.

845 Nahlik, A. M. and Fennessy, M. S.: Carbon storage in US wetlands, *Nat Commun*, 7(1), 13835,
846 doi:10.1038/ncomms13835, 2016.

847 O'Donnell, B. C. Carbon sequestration within Northeastern Pacific seagrass meadows, Thesis
848 10607483, University of California, Davis, ProQuest Dissertations Publishing, 2017.
849

850 Oakes, J. M. and Eyre, B. D.: Transformation and fate of microphytobenthos carbon in
851 subtropical, intertidal sediments: potential for long-term carbon retention revealed by ¹³C-
852 labeling, *Biogeosciences*, 11(7), 1927–1940, <https://doi.org/10.5194/bg-11-1927-2014>, 2014.

853 Ouyang, X. and Lee, S. Y.: Updated estimates of carbon accumulation rates in coastal marsh
854 sediments, *Biogeosciences*, 11(18), 5057–5071, doi:10.5194/bg-11-5057-2014, 2014.

855 Parnell, A. C. and Jackson, A.L. SIAR: stable isotope analysis in R. R package version 4.2. <http://CRAN.R-project.org/package=siar>, 2013.
856
857

858 Patrick, Wm. H. and DeLaune, R. D.: Subsidence accretion and sea level rise in south San
859 Francisco Bay marshes, *Limnology and Oceanography*, 35(6), 1389–1395,
860 doi:10.4319/lo.1990.35.6.1389, 1990.

861 Peck, E. K., Wheatcroft, R. A. and Brophy, L. S.: Controls on Sediment Accretion and Blue
862 Carbon Burial in Tidal Saline Wetlands: Insights From the Oregon Coast, USA, *J. Geophys. Res.*
863 *Biogeosci.*, 125(2), doi:10.1029/2019JG005464, 2020.

864 Pendleton, L., Donato, D. C., Murray, B. C., Crooks, S., Jenkins, W. A., Sifleet, S., Craft, C.,
865 Fourqurean, J. W., Kauffman, J. B., Marbà, N., Megonigal, P., Pidgeon, E., Herr, D., Gordon, D.
866 and Baldera, A.: Estimating global “Blue Carbon” emissions from conversion and degradation of
867 vegetated coastal ecosystems, edited by S. Thrush, *PLoS ONE*, 7(9), e43542,
868 doi:10.1371/journal.pone.0043542, 2012.

869 Perdue, E. M. and Koprivnjak, J. F.: Using the C/N ratio to estimate terrigenous inputs of organic
870 matter to aquatic environments, *Estuarine, Coastal and Shelf Science*, 73(1–2), 65–72,
871 doi:10.1016/j.ecss.2006.12.021, 2007.

872 Poppe, K. L. and Rybczyk, J. M.: Carbon Sequestration in a Pacific Northwest Eelgrass (*Zostera*
873 *marina*) Meadow, *BioOne*, 92(2), 80–91, doi:10.3955/046.092.0202, 2018.

874 Postlethwaite, V. R., McGowan, A. E., Kohfeld, K. E., Robinson, C. L. K. and Pellatt, M. G.:
875 Low blue carbon storage in eelgrass (*Zostera marina*) meadows on the Pacific Coast of Canada,
876 *PLOS ONE*, 13(6), e0198348, doi:10.1371/journal.pone.0198348, 2018.

877 Prentice, C., Hessing-Lewis, M., Sanders-Smith, R. and Salomon, A. K.: Reduced water motion
878 enhances organic carbon stocks in temperate eelgrass meadows, *Limnology and Oceanography*,
879 64(6), 2389–2404, doi:10.1002/lno.11191, 2019.

880 Prentice, C., Poppe, K. L., Lutz, M., Murray, E., Stephens, T. A., Spooner, A., Hessing-Lewis,
881 M., Sanders-Smith, R., Rybczyk, J. M., Apple, J., Short, F. T., Gaeckle, J., Helms, A., Mattson,
882 C., Raymond, W. W., and Klinger, T.: A Synthesis of Blue Carbon Stocks, Sources, and
883 Accumulation Rates in Eelgrass (*Zostera marina*) Meadows in the Northeast Pacific, *Global*
884 *Biogeochem. Cycles*, 34, <https://doi.org/10.1029/2019GB006345>, 2020.
885

886 R Core Team (2018). R: A language and environment for statistical computing. R Foundation for
887 Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.
888

- 889 Ricart, A., Dalmau, A., Pérez, M. and Romero, J.: Effects of landscape configuration on the
890 exchange of materials in seagrass ecosystems, *Mar. Ecol. Prog. Ser.*, 532, 89–100,
891 doi:10.3354/meps11384, 2015.
- 892 Ricart, A. M., Pérez, M. and Romero, J.: Landscape configuration modulates carbon storage in
893 seagrass sediments, *Estuarine, Coastal and Shelf Science*, 185, 69–76,
894 doi:10.1016/j.ecss.2016.12.011, 2017.
- 895 Ricart, A. M., York, P. H., Bryant, C. V., Rasheed, M. A., Ierodiaconou, D. and Macreadie, P. I.:
896 High variability of Blue Carbon storage in seagrass meadows at the estuary scale, *Scientific*
897 *Reports*, 10(1), 5865, doi:10.1038/s41598-020-62639-y, 2020.
- 898 Rogers, K., Kelleway, J. J., Saintilan, N., Megonigal, J. P., Adams, J. B., Holmquist, J. R., Lu,
899 M., Schile-Beers, L., Zawadzki, A., Mazumder, D., and Woodroffe, C. D.: Wetland carbon
900 storage controlled by millennial-scale variation in relative sea-level rise, 567, 91–95,
901 <https://doi.org/10.1038/s41586-019-0951-7>, 2019.
- 902 Röhr, M. E., Holmer, M., Baum, J. K., Björk, M., Boyer, K., Chin, D., Chalifour, L., Cimon, S.,
903 Cusson, M., Dahl, M., Deyanova, D., Duffy, J. E., Eklöf, J. S., Geyer, J. K., Griffin, J. N.,
904 Gullström, M., Hereu, C. M., Hori, M., Hovel, K. A., Hughes, A. R., Jorgensen, P.,
905 Kiriakopolos, S., Moksnes, P.-O., Nakaoka, M., O’Connor, M. I., Peterson, B., Reiss, K.,
906 Reynolds, P. L., Rossi, F., Ruesink, J., Santos, R., Stachowicz, J. J., Tomas, F., Lee, K.-S.,
907 Unsworth, R. K. F., and Boström, C.: Blue carbon storage capacity of temperate Eelgrass
908 (*Zostera marina*) meadows, *Global Biogeochemical Cycles*, 32, 1457–1475,
909 <https://doi.org/10.1029/2018GB005941>, 2018.
- 910 Saintilan, N., Rogers, K., Mazumder, D. and Woodroffe, C. Allochthonous and autochthonous
911 contributions to carbon accumulation and carbon store in southeastern Australian coastal
912 wetlands, *Estuarine, Coastal and Shelf Science*, 128, 84–92, doi:10.1016/j.ecss.2013.05.010,
913 2013.
- 914 Schlosser, S., and A. Eicher. The Humboldt Bay and Eel River Estuary Benthic Habitat Project.
915 California Sea Grant Publication T-075. 246 p, 2012.
- 916 Serrano, O., Mateo, M. A., Renom, P., and Julià, R.: Characterization of soils beneath a
917 *Posidonia oceanica* meadow, *Geoderma*, 185–186, 26–36,
918 <https://doi.org/10.1016/j.geoderma.2012.03.020>, 2012.
- 919 Serrano, O., Lavery, P. S., Duarte, C. M., Kendrick, G. A., Calafat, A., York, P. H., Steven, A.
920 and Macreadie, P. I.: Can mud (silt and clay) concentration be used to predict soil organic carbon
921 content within seagrass ecosystems?, *Biogeosciences*, 13(17), 4915–4926, doi:10.5194/bg-13-
922 4915-2016, 2016.
- 923 Serrano, O., Kelleway, J. J., Lovelock, C. and Lavery, P. S.: Conservation of Blue Carbon
924 Ecosystems for Climate Change Mitigation and Adaptation, *Coastal Wetlands*, pp. 965–996,
925 Elsevier., 2019.

- 926 St. Laurent, K. A., Hribar, D. J., Carlson, A. J., Crawford, C. M., and Siok, D.: Assessing coastal
927 carbon variability in two Delaware tidal marshes, *Journal of Coast Conservation*, 24, 65,
928 <https://doi.org/10.1007/s11852-020-00783-3>, 2020.
- 929 Trevathan-Tackett, S. M., Macreadie, P. I., Sanderman, J., Baldock, J., Howes, J. M. and Ralph,
930 P. J.: A Global Assessment of the Chemical Recalcitrance of Seagrass Tissues: Implications for
931 Long-Term Carbon Sequestration, *Front. Plant Sci.*, 8, 925, doi:10.3389/fpls.2017.00925, 2017.
- 932 Trimble, S. W.: Historical hydrographic and hydrologic changes in the San Diego creek
933 watershed, Newport Bay, California, *Journal of Historical Geography*, 29(3), 422–444,
934 doi:10.1006/jhge.2002.0485, 2003.
- 935 Valiela, I. and Cole, M. L.: Comparative evidence that salt marshes and mangroves may protect
936 seagrass meadows from land-derived nitrogen loads, *Ecosystems*, 5(1), 92–102,
937 doi:10.1007/s10021-001-0058-4, 2002.
- 938 Van Dyke, E. and Wasson, K.: Historical ecology of a central California estuary: 150 years of
939 habitat change, *Estuaries*, 28(2), 173–189, doi:10.1007/BF02732853, 2005.
- 940 Wilkinson, G. M., Besterman, A., Buelo, C., Gephart, J. and Pace, M. L.: A synthesis of modern
941 organic carbon accumulation rates in coastal and aquatic inland ecosystems, *Scientific Reports*,
942 8(1), 15736, doi:10.1038/s41598-018-34126-y, 2018.
- 943 Yang, S. L., Li, H., Ysebaert, T., Bouma, T. J., Zhang, W. X., Wang, Y. Y., Li, P., Li, M. and
944 Ding, P. X.: Spatial and temporal variations in sediment grain size in tidal wetlands, Yangtze
945 Delta: On the role of physical and biotic controls, *Estuarine, Coastal and Shelf Science*, 77(4),
946 657–671, <https://doi.org/10.1016/j.ecss.2007.10.024>, 2008.
- 947 Zhou, J., Wu, Y., Kang, Q. and Zhang, J.: Spatial variations of carbon, nitrogen, phosphorous
948 and sulfur in the salt marsh sediments of the Yangtze Estuary in China, *Estuarine, Coastal and
949 Shelf Science*, 71(1–2), 47–59, doi:10.1016/j.ecss.2006.08.012, 2007