

# Fire in lichen-rich subarctic tundra changes carbon and nitrogen cycling between ecosystem compartments but has minor effects on stocks

Ramona J. Heim<sup>1</sup>, Andrey Yurtaev<sup>2</sup>, Anna Bucharova<sup>1,3</sup>, Wieland Heim<sup>1,4</sup>, Valeriya Kutskir<sup>2</sup>, Klaus-Holger Knorr<sup>5</sup>, Christian Lampei<sup>1</sup>, Alexandr Pechkin<sup>6</sup>, Dora Schilling<sup>1</sup>, Farid Sulkarnaev<sup>2</sup>, and Norbert Hölzel<sup>1</sup>

<sup>1</sup>Institute of Landscape Ecology, Biodiversity and Ecosystem Research Group, University of Muenster, Heisenbergstraße 2, 48149 Münster, Germany

<sup>2</sup>Research Institute of Ecology and Natural Resources Management, Tyumen State University, 6 Volodarskogo Street, Tyumen, Russia

<sup>3</sup>Department of Biology, Conservation Biology Group, University of Marburg, Karl-von-Frisch-Straße 8, 35043 Marburg, Germany

<sup>4</sup>Department of Biology, Animal Ecology, University of Turku, Vesilinnantie 520500, Turku, Finland

<sup>5</sup>Institute of Landscape Ecology, Ecohydrology and Biogeochemistry Group, University of Muenster, Heisenbergstraße 2, 48149 Münster, Germany

<sup>6</sup>Research Center of the Yamal-Nenets Autonomous District, Salekhard 629008, Russia

**Correspondence:** Ramona Heim (ramona.heim@uni-muenster.de)

**Abstract.** Fires are predicted to increase in Arctic regions due to ongoing climate change. Tundra fires can alter carbon and nutrient cycling and release a substantial amount of greenhouse gases with global consequences. Yet, the long-term effects of tundra fires on carbon (C) and nitrogen (N) stocks and cycling are still unclear. Here we used a space-for-time approach to investigate the long-term fire effects on C and N stocks and cycling in soil and aboveground living biomass. We collected data from three large fire scars (>44, 28 and 12 years old) and corresponding control areas and used linear mixed-effects models in a Bayesian framework to analyse long-term development of C and N stocks and cycling after fire.

We found that tundra fires had no long-term effect on total vegetation C and N stocks because a major part of the stocks was located belowground in soils, which were largely unaltered by fire. However, fire had a strong long-term effect on stocks in the aboveground vegetation, mainly due to the reduction of the lichen layer. Fire reduced N concentrations in graminoids and herbs on the younger fire scars, which affected respective C/N ratios and may indicate an increased post-fire competition between vascular plants. Aboveground plant biomass was depleted in <sup>13</sup>C in all three fire scars. In soil, the relative abundance of <sup>13</sup>C changed with time after fire.

Our results indicate that in lichen-rich subarctic tundra ecosystems, the contribution of fires to the release of additional carbon to the atmosphere might be relatively small as soil stocks appear to be resilient within the observed time frame.

## 15 1 Introduction

Arctic regions are warming faster than the worldwide average, with strong impacts on the global carbon cycle (Post et al., 2019; Schuur et al., 2015). Warming is expected to expose ecosystems at high latitudes to increased fire frequency and extent (Chen et al., 2021; Hu et al., 2015; Moritz et al., 2012; Young et al., 2016). Wildfires in the Arctic can feed back on climate change because they release carbon stored in the tundra ecosystems (Lasslop et al., 2020; Veraverbeke et al., 2021). This effect may be substantial because Arctic and sub-arctic regions store huge amounts of the global carbon (C) (ca. 1,700 billion tons in terrestrial soils, Schuur et al., 2015).

Wildfires enhance the release of stored carbon from active ecosystems in multiple ways. Combustion of living and dead biomass rapidly transfers large amounts of stored C to the atmosphere (Mack et al., 2011). Wildfires also burn the insulating and reflecting layers of soil organic material and vegetation, which increases soil temperatures through irradiation (Chambers et al., 2005; Jiang et al., 2015). Warmer soil temperatures raise microbial activity, which in turn promotes faster decomposition processes (Jansson and Hofmockel, 2020). Soil warming induces permafrost thaw and active layer deepening, which exposes organic matter in deeper layers to decomposition, resulting in a massive release of carbon to the atmosphere (Estop-Aragonés et al., 2020; Schuur et al., 2009). In summary, the rate of C loss through fires is substantial and faster than other climate-driven processes influencing C cycling in northern latitudes (Mack et al., 2011).

Fires also affect the stored N, which has consequences for vegetation and ecosystem functioning, because N limits productivity in tundra ecosystems (Oulehle et al., 2016). Fires release substantial parts of the accumulated nitrogen (N) pool to the atmosphere. However, soil warming and related deeper thaw may increase plant-available inorganic N through higher mineralisation rates (Aerts, 2006; Salmon et al., 2016). This increases the productivity of shrubs and graminoids because they can reach the nutrients mineralised in deeper soil layers than cryptogams (Dormann and Woodin, 2002; Oulehle et al., 2016; Salmon et al., 2016). As a consequence, the cover of vascular plants increases after fires at the expense of previously dominating cryptogams, which cannot profit from the enhanced nitrogen availability as they do not reach deep soil layers or cannot take up nitrogen from the soil (Bret-Harte et al., 2013; Jandt et al., 2008; Narita et al., 2015; Turetsky et al., 2012). Such profound changes in vegetation cover, in turn, influence ecosystem functioning and processes, such as carbon and nitrogen cycling and C and N stocks (Longton, 1997; Sancho et al., 2016; Turetsky, 2003).

Studies including nutrient cycles in soil and vegetation related to long-term effects of tundra fires are rare and, therefore, the effect of fires on the carbon cycle of tundra ecosystems is relatively unknown (Mack et al., 2011). This is unfortunate because the reactions of the tundra ecosystem to altered temperature have a time lag and are thus relatively slow (Rinnan et al., 2007). The effects of fire on the functioning of the tundra ecosystem may thus be visible only after more than a decade (Blok et al., 2018). Therefore, long-term studies are essential (Chapin Iii et al., 2000).

One widely used tool to unravel C and nutrient cycling processes is the use of stable isotopes (Peterson and Fry, 1987). The relative abundance of the heavy carbon isotope ( $^{13}\text{C}$ ) in the living aboveground biomass and in the soil can provide insight into post-fire environmental site conditions. In soil, increased activity of microbes after fire is reflected in the residual enrichment of  $^{13}\text{C}$  in the remaining soil organic matter, as decomposers prefer isotopically light material ( $^{12}\text{C}$ ) (Ehleringer et al., 2000). In

the aboveground biomass, the isotope ratio is related to the isotopic composition of the source CO<sub>2</sub>, and thus the isotope ratio  
50 in leaves may reflect the lower levels of <sup>13</sup>C in CO<sub>2</sub> which is available for photosynthesis through respiration from soils during  
decomposition (Dawson et al., 2002; Farquhar et al., 1989).

The isotope ratios of N can be used for a better understanding of post-fire N cycling processes by providing information about  
the association with mycorrhizal fungi. During N transfer from mycorrhizal fungi to the plant, the heavy <sup>15</sup>N gets enriched in  
the mycorrhizas and depleted in the plants. During succession, the associated mycorrhizal fungi have to colonize the plants  
55 first, and therefore <sup>15</sup>N would be expected to be enriched in early successional stages and depleted in later successional stages  
(Hobbie et al., 2005).

Here, we studied the long-term effect of fire on tundra nutrient stocks and cycling using a space-for-time approach. We focused  
on three large fire scars (>44, 28, and 12 years old) that represent a gradient of post-fire succession. We analysed C and N  
stocks, concentrations, and isotope ratios in vegetation and soil. We hypothesized that:

- 60 1) Fire leads to decreased ecosystem C and N stocks.  
2) Fire affects long-term C and N cycling, evident through  
a) an increased N concentration and a decreased C/N ratio in vascular plants after fire.  
b) a depletion of <sup>13</sup>C in aboveground biomass and a relative enrichment of <sup>13</sup>C in soil after fire.  
c) increased relative abundance of <sup>15</sup>N in leaves of vascular plants on the younger fire scars.


65

## 2 Material and Methods

### 2.1 Study area

The study area was located in Western Siberia within the Yamalo-Nenets Autonomous Okrug (Region) between the rivers Pur  
and Taz, north of the Arctic Circle (centre at 67° 1'19.59"N, 79° 1'53.53"E, total study area size ca. 70 km<sup>2</sup>) (Heim et al.,  
70 2021). The region has a subarctic climate with a growing season from mid-June to early September, and with a mean annual  
temperature of -8.1 °C, mean January temperature of -26.2 °C, mean July temperature of 14.4 °C, and annual precipitation of  
482 mm (Kazakov, 2019). The vegetation in the region represents the transition zone between the forest-tundra to the south and  
the shrub-tundra to the north (Yurkovskaya, 2011). Compared to other previously investigated tundra areas, e.g. moist acidic  
tundra in the Anaktuvuk River region in Alaska (Mack et al., 2011), the region is a relatively dry and well-drained upland area.  
75 The vegetation is dominated by reindeer lichens (mostly *Cladonia* spp., around 70% cover), with abundant shrubs and dwarf  
shrubs, such as *Betula nana* (around 25% cover) and *Vaccinium uliginosum* (around 10% cover), and occasional graminoids  
and bryophytes with low cover. The landscape is sparsely dotted with larch trees (*Larix sibirica*).

Soils are Cryosols (IUSS Working Group WRB, 2015), which developed in silty, loess-like parent material and soil organic  
layer thickness was generally low at all sites. Bulk density ranged from 0.6 g/cm<sup>3</sup> to 1.6 g/cm<sup>3</sup> and pH from 3.8 to 6.3 (Fig.  
80 S1). The mineral soil was covered by living lichen and moss in unburned sites, lying directly on the mineral layer. The 0-5  
cm layer in our study is the upper layer of the mineral soil, which was in unburned sites found directly under the lichen layer.



**Figure 1.** Study area with fire scars in the north of Western Siberia. Edited figure from (Heim et al., 2021). Satellite image source: Copernicus Sentinel-2 data (2019).

Soil temperature in 12 cm depth was higher on burnt sites but regenerated to control levels after > 44 years (predicted mean difference to control plots after 12, 28, and < 44 years: 6.29 °C, 6.29 °C, 0.24 °C) (Heim et al., 2021). This was also the case in active layer depth, which was lower on burnt sites but regenerated to control levels after > 44 years (predicted mean difference to control plots after 12, 28, and < 44 years: -33.02 cm, -32.20 cm, 6.62 cm) (Heim et al., 2021).



The main cause of fires in the study region is lightning (Kornienko, 2018), but the number of human-induced fires increases because of expanding transport and settlement infrastructure due to oil and gas exploitation (Mollicone et al., 2006; Vilchek and Bykova, 1992; Yu et al., 2015).

## 90 **2.2 Sampling design and data collection**

To examine the long-term impact of fire on C and N cycling in soil and aboveground biomass, we studied three large fire scars (aged 12, 28, and >44 years) and adjacent unburnt control sites (Fig. 1). All three fire scars were located close to each other (< 10 km distance between scars) and therefore had similar environmental conditions. We detected the three fire scars visually with the help of satellite images. We used annual Landsat images back to the year 1985 in Google Earth Timelapse (Gorelick et al., 2017) and older Landsat images back to 1973, which we downloaded via the USGS earth explorer (U.S. Geological Survey, 2018) (for details see Heim et al., 2021). The oldest fire scar (3,500 ha) was already visible on the first satellite image from 1973 (Landsat1). The medium-aged scar (ca. 12,500 ha) burnt in 1990, and the youngest scar (542 ha) burnt in 2005. We collected samples in July 2017 at fire scars burned before 1973 and in 2005 (that is >44 and 12 years after fire, respectively) and in July 2018 at the area burnt in 1990 (28 years after fire). Environmental conditions of the tundra ecosystem are stable with low inter-annual variability (Dahl, 1975) and climatic conditions in the two sampling years were similar (Kazakov, 2019). It is thus unlikely that collecting in two subsequent years biased the results.

We placed 10 sampling locations along the fire border of each fire scar (at least 300 m apart from each other). At each location, we placed one sampling plot in the burnt and one sampling plot in the unburnt site, resulting in a total of 60 plots (30 pairs). Each site included a 10 m by 10 m plot and a soil profile, located 2-5 m next to the plot. The plots at each location were placed as close to each other as possible, but at least 100 m apart (50 m minimum plot distance from the fire border, to avoid edge effects).

We sampled aboveground biomass in five subplots of the 10 by 10 m plot. The subplots were situated in the four corners and in the centre of the plot. While the subplots for shrubs, herbs, and graminoids were squares of 0.3 m x 0.3 m, the subplots for lichens and bryophytes were 0.1 m x 0.1 m and were randomly placed within the 0.3 m x 0.3 m squares. Biomass of each type (shrubs, graminoids, herbs, bryophytes, lichens) was separately sampled and all subplots were pooled in one bag per type. Soil was sampled in layers ranging from 0-5, 5-30, and 30-60 cm depth. In each layer, we placed three cylinders (200 cm<sup>3</sup>) horizontally and distributed them over the whole range and sampled the soil of one layer in one plastic bag.

## **2.3 Laboratory analyses and stock calculations**

In the field, aboveground biomass was stored well ventilated, while soil was stored in airtight plastic bags. We dried aboveground biomass and soil in the laboratory at 60 °C to a constant weight. For determination of C and N concentrations and stable isotopic composition, samples were ground in a ball mill (tungsten carbide cups, MM400, Retsch, Haan, Germany) and weighed into tin capsules for analyses on an elemental analyser (EA 3000, Eurovector, Padua, Italy) coupled to an isotope ratio mass spectrometer (EA-IRMS, NuHorizon, Nu Instruments, Wrexham, UK). Calibrations were done using certified working standards (IVA Analysentechnik, Meerbusch, Germany) and reference materials (IAEA 600 (Coplen et al., 2006), USGS 35

120 (Böhlke et al., 2003)). We could not analyse nitrogen isotopic composition in the soil samples, as the N concentrations were generally too low.

We express the stable isotope composition using the common delta notation as a ratio relative to an internationally accepted reference standard:  $\delta\text{XXE} = 1000 \cdot (R_{\text{sample}}/R_{\text{standard}} - 1)$ , ‰. Where E is the element, xx the mass of the heavier isotope, and R is the abundance ratio of the isotopes (e.g.,  $^{15}\text{N}:^{14}\text{N}$ ) (Dawson et al., 2002). Higher  $\delta$  values indicate a higher abundance of the heavier isotope (Dawson et al., 2002). We furthermore analysed bulk density. We dried a subsample of soil samples at 105°C and calculated bulk density using the following formula: bulk density ( $\text{g}/\text{cm}^3$ ) = dry soil weight (g) / soil volume ( $\text{cm}^3$ ). Based on the bulk density, we calculated soil stocks for each range with following formula: stock<sub>soil</sub> ( $\text{kg}/\text{m}^2$ ) = bulk density ( $\text{g}/\text{cm}^3$ ) \* concentration/100 \* range size (cm) (e.g. 25 cm for 5-30 cm depth range) / 1000 ( $\text{g}/\text{cm}^3$  in  $\text{kg}/\text{m}^2$ ). C and N stocks for vegetation were calculated as follows: stock<sub>vegetation</sub> ( $\text{kg}/\text{m}^2$ ) = dry mass (kg) \* sampling area in  $\text{m}^2$  \* concentration/100.

130 We use the term ‘total stocks’ for combined soil and vegetation stocks.

## 2.4 Statistical analysis

We carried out all statistics in R Version 4.0.3. (R Core Team, 2020) and fitted all models in a Bayesian framework using the function brm() from the package brms (Bürkner, 2017, 2018). For testing anticipation 1 (Fire leads to decreased ecosystem C and N stocks), we summarised aboveground biomass and soil stocks per sampling site. We fitted two linear models with log-transformed stocks (C stocks, N stocks) as dependent variables. Fire scar (12, 28, >44 years), burn status (burnt, unburnt), and their interaction were included as independent variables and location (paired plots) as a random factor. To obtain the posterior distribution, we used improper priors and ran 6000 iterations (warmup = 3000) with four chains. The posterior distribution is a probability distribution that summarises updated beliefs about the parameter after observing the data and is thus a result of the prior distribution and the likelihood function (Korner-Nievergelt et al., 2015).

135

We present mean values together with the 95% credible interval (CrI) of the simulated posterior distribution. The 95% CrI is the range in which the true value is expected with a probability of 0.95. We calculated the posterior probabilities by using the proportion of simulated values of the posterior distribution of C and N stocks on burnt sites being smaller than the proportion of simulated values of the posterior distribution of stocks on unburnt control sites. Therefore, a posterior probability of 1 would indicate that the stocks on the burnt sites were significantly lower than on unburnt control sites. A posterior probability of 0 would mean the opposite and a posterior probability of 0.5 indicates that stocks at burnt and unburnt sites did not differ.

140

145

To analyse the impacts of fire on aboveground biomass and soil stocks separately, we fitted four linear models with log-transformed stocks (aboveground biomass C stocks, aboveground biomass N stocks, soil C stocks, soil N stocks) as dependent variable. Fire scar (12, 28, >44 years), burn status (burnt, unburnt), and vegetation type for biomass (shrub leaves, herbs, graminoids, bryophytes, lichens) or depth for soil (0-5 cm, 5-30 cm, 30-60 cm) were included as independent variables with all possible interactions. Sites were nested in location as random factor. To obtain the posterior distribution, we used improper priors and ran 4000 iterations (warmup = 2000) with four chains. For the models of C/N ratio in biomass and C concentration in soil, we used adapt\_delta=0.95 and run 6000 iterations (warmup=3000) for better convergence. Bayesian  $R^2$  was calculated with the function bayes\_ $R^2$  from the package brms (Bürkner, 2017, 2018).

150

For testing the anticipation 2 (Fire affected long-term C and N cycling), we first calculated the differences between burnt and unburnt control plots (burnt - unburnt) for concentrations, ratios, and isotope ratios as this enabled us a more straightforward and clearer interpretation (see raw data in Fig. S2, Fig. S3). A negative difference of the C concentration, therefore, indicates that fire decreased the C concentration. A negative difference in the C/N ratio indicates that fire decreased the C/N ratio – either through decreased C concentrations or increased N concentrations. A negative difference of  $\delta^{13}\text{C}$  ( $\Delta^{13}\text{C}$ ) indicates that fire decreased  $\delta^{13}\text{C}$  (and thus caused a depletion of  $^{13}\text{C}$ ).

We fitted 9 linear models with concentrations, C/N-ratio, and isotope ratios (C in biomass, C in soil, N in biomass, N in soil, C/N in biomass, C/N in soil,  $\delta^{13}\text{C}$  in biomass,  $\delta^{13}\text{C}$  in soil,  $\delta^{15}\text{N}$  in biomass) as dependent variable. Fire scar age (12, 28, >44 years), burn status (burnt, unburnt), and vegetation type for biomass (shrub leaves, herbs, graminoids, bryophytes, lichens) or depth for soil (0-5 cm, 5-30 cm, 30-60 cm) were included as independent variables with all possible interactions. Sites were nested in location as random factor. We controlled for non-constant variances in type and depth adapting sigma (e.g. sigma 0 + type). To obtain the posterior distribution, we used improper priors and ran 4000 iterations (warmup = 2000) with four chains.


### 3 Results

#### 3.1 Fire influence on C and N stocks in aboveground biomass and soil

Fire had no long-term effects on combined soil and vegetation stocks (hereafter total stocks) (C:  $P_{\text{burnt}_{12}<\text{control}_{12}} = 0.68$ ,  $P_{\text{burnt}_{28}<\text{control}_{28}} = 0.94$ ,  $P_{\text{burnt}_{44}<\text{control}_{44}} = 0.16$ , N:  $P_{\text{burnt}_{12}<\text{control}_{12}} = 0.40$ ,  $P_{\text{burnt}_{28}<\text{control}_{28}} = 0.81$ ,  $P_{\text{burnt}_{44}<\text{control}_{44}} = 0.18$ ) (Fig. 2, Table S6, Table S7). The net losses/gains for the youngest fire scar were  $-2.17 \text{ kgC/m}^2$  and  $0.09 \text{ kgN/m}^2$ , for the intermediate  $-9.85 \text{ kgC/m}^2$  and  $-0.33 \text{ kgN/m}^2$  and for the oldest  $5.41 \text{ kgC/m}^2$  and  $0.30 \text{ kgN/m}^2$ .

Soils at burnt plots had less C at the intermediate fire scar in 5-30 cm depth. In the oldest fire scar, soils stored more N and C in the upper soil layer in comparison to the unburnt control (Table 1, Table S7, Table S8) pointing towards a replenishment from the belowground organs of vascular plants in the course of succession.

In contrast to soil, C and N stocks in vegetation were strongly reduced through fire (Fig. 2). In the two younger stages vegetation C-stocks had only about 15 % of the unburnt reference and even after more than 44 years they reached on average only 39 % of the unburnt reference. Vegetation N-stocks recovered significantly fast and reached in the oldest stage already 74 % of the unburnt reference (Fig. 2). This was mainly because fire substantially reduced lichens and consequently the C and N stored in them. The depletion of C and N in lichens lasted for more than four decades (Table 1, Table S7, Table S8). In contrast, fire increased C and N stocks in herbs in the intermediate fire scar and in bryophytes in the oldest one (Tab. 1). In the oldest fire scars the proportion of C and N stored in bryophytes clearly dominated the stocks of C and N in aboveground living vegetation. For both elements, absolute values were ca. three times higher than in unburned controls (Fig. 2). Thus, after 44 years the dominant role for storing C and N in aboveground vegetation shifted from lichens to bryophytes and lichens were still far from gaining back their former absolutely dominant importance as C and N sink.



**Figure 2.** Predicted mean values for C and N stocks in aboveground biomass and soil in regard to time after fire on burnt and control plots. Vertical lines are 95% credible intervals (CrI). Model structure and parameters as well as corresponding CrI's and  $R^2$  are shown in Table S6, Table S7 and Table S8. C and N stocks for shrub stems are calculated as described in Figure S4. Compare A and B with aboveground dry weight in Figure S5.

**Table 1.** Probabilities of stocks on burnt plots to be lower than stocks on unburnt control plots. If probabilities are higher than 0.975 (stocks on burnt plots are lower compared to unburnt plots) or smaller 0.025 (stocks on burnt plots are higher) they are written in bold. If probabilities are 0.5 there is no difference between stocks on burnt plots and stocks on control plots.

Vegetation Type	$C_{burnt} < C_{control}$			$N_{burnt} < N_{control}$		
	12	28	>44	12	28	>44
Lichens	<b>1.00</b>	<b>1.00</b>	<b>1.00</b>	<b>1.00</b>	<b>1.00</b>	<b>1.00</b>
Bryophytes	0.22	0.76	<b>0.01</b>	0.29	0.79	<b>0.01</b>
Graminoids	0.08	0.90	0.76	0.16	0.95	0.81
Herbs	0.18	<b>0.00</b>	0.67	0.32	<b>0.00</b>	0.69
Shrub leaves	0.76	0.53	0.29	0.85	0.59	0.27
Shrub stems	0.86	0.67	0.34	0.91	0.70	0.32
Soil	$C_{burnt} < C_{control}$			$N_{burnt} < N_{control}$		
Depth	12	28	>44	12	28	>44
0-5 cm	0.75	0.84	<b>0.02</b>	0.43	0.84	0.03
5-30cm	0.69	<b>0.98</b>	0.94	0.51	0.78	0.94
30-60 cm	0.17	0.61	0.25	0.09	0.57	0.38

### 3.2 Fire influence on C and N cycling

Fire affected C and N concentrations and isotope ratios in aboveground biomass and soil (Fig. 3 A-H, Table S9, Table S10). While fire had generally strong long-term effects on C and N concentrations as well as on isotope ratios in vegetation, only the upper soil layer showed differences in concentrations and isotope ratios. Overall, long-term fire effects on C and N concentrations and isotope ratios were most pronounced in lichens.


Fire had a positive effect on N concentrations in lichens and a negative on N concentrations in graminoids and herbs (Fig. 3 C). Those effects disappeared in all groups after >44 years post fire. The N concentrations were reflected in the C/N ratios, which increased in graminoids and herbs but decreased in lichens (Fig. 3 E). While fire significantly increased C and N concentrations in the upper soil layer on the oldest fire scar, we could not detect a fire effect on the deeper soil layers (Fig. 3 D, F).

195 Lichens on all three fire scars were relatively depleted in  $^{13}\text{C}$  in comparison to unburnt plots (Fig. 3 G). This was also the case in graminoids on the two younger fire scars (Fig. 3 G).

In soil, the abundance of  $^{13}\text{C}$  changed with time after fire. The upper soil layer of the youngest fire scar was enriched in  $^{13}\text{C}$  in comparison to unburnt plots (Fig. 3 H). This relationship turned with time since fire, and we observed a relative depletion of  $^{13}\text{C}$  in the soils of the oldest fire scar in comparison to unburnt plots.

200 Fire increased the abundance of  $^{15}\text{N}$  in lichens (Fig. 3 I). This effect was still visible four decades after fire. We found no significant long-term fire effects on  $\delta^{15}\text{N}$  in vascular plants.





**Figure 3.** Difference (burnt - unburnt) between posterior means of C and N concentrations and isotope ratios in regard to time after fire in aboveground biomass and soil. Colour intensity indicates the probability  $P$  of the value from an unburnt control plot to be higher than the value from the burnt plot. If  $P$  is very high, there is a high probability of a negative fire impact. If  $P$  is very low, there is a high probability of a positive fire impact.

## 4 Discussion

Fire had no long-term effects on total C and N stocks in subarctic dry tundra. This is probably because most of it was stored in soils (97.4% of C and 99.7% of N), which were largely unaffected by fire. In contrast, stocks in aboveground vegetation were strongly reduced through fire, and the effect persisted even after more than 44 years. Striking results reporting high losses of carbon through tundra fires (Mack et al., 2011) received a lot of attention and were covered by media (e.g. Rosen, 2017). While (Mack et al., 2011) investigated mainly the combustion loss, we investigated the long-term fire consequences on stocks. In our study, recovery of vegetation took place and hence a regeneration of the carbon stocks. The effect of fire on tundra carbon and nitrogen stocks is variable and dependent on many factors. First, losses are strongly mediated by fire severity through the consumption of organic matter (Schuur et al., 2003). Fire severity was relatively high for the fire investigated in Alaska (Mack et al., 2011) but intermediate in our study (Heim et al., 2021), which may partly explain the lack of a strong effect in our study (Maslov et al., 2020).

Our results support recent findings of (He et al., 2021), who claim that soil organic carbon loss through tundra fires may be overestimated because the soil organic layer in many tundra areas is much lower (see also our findings) than in the area of the extreme Anaktuvuk River Fire with predominant wet organic soils (Histosols).

In addition to the thick organic layer which usually is combusted by fire (He et al., 2021) moist tundra ecosystems typically store much higher amounts of carbon per unit area because of a higher soil organic matter content in comparison to dry tundra ecosystems (Marion et al., 1997). Through increased temperature sensitive decomposition after fire, (De Baets et al., 2016) burning can thus indirectly lead to higher decomposition rates and ecosystem carbon loss in wet tundra ecosystems. Depending on tundra type, fire impacts can thus have varying severity of fire impact on biogeochemical cycling. Second, the thickness of the soil organic matter plays an important role. Because the decomposition of plant material is limited in tundra ecosystems by low temperatures and the often wet and anaerobic conditions, organic matter accumulates (Jonasson et al., 2001). A reduction of this soil organic layer leads to decreases in the amount of stored C (Mack et al., 2004). Thus, exceptionally large losses of C and N stocks were reported from wildfires in rather wet tundra ecosystems with a thick soil organic layer or from peatlands (Mack et al., 2011; Turetsky et al., 2015). Our study was located in an upland area with well-drained mineral soils and a thin organic layer, and in such conditions, fires seem to cause only minor losses (Loranty et al., 2014). Thick organic soil layers, as e.g. reported for a site in Alaska (21.5 cm, Mack et al., 2011) may store particularly large amounts of C and N that are potentially lost if burnt. However, much shallower organic layers have been reported for North-Eastern Siberia (6.3 cm, Loranty et al., 2014) and were also typical of our study area (average thickness 4.3 cm). Therefore, potential losses upon burning can be expected to be much lower, and, moreover, the importance of the more stable mineral soil carbon stock less susceptible to losses from burning relatively increases. However, a direct comparison of these studies is not possible, as we investigated the system during regeneration and not directly after combustion. Further studies are needed to draw more general conclusions about the role of vegetation type and environmental settings (for example well drained upland soils) for wildfire-induced losses of C and N in tundra ecosystems. In the long term, fire even increased C and N stocks in the upper soil layer, which was probably due to plant community turnover. The vegetation on burnt plots did not return to control levels within >44 years and

was dominated by bryophytes, herbaceous plants, and shrubs, especially *Betula nana* (Heim et al., 2021). Vascular plants, and especially woody species, accumulate more soil C in roots and litter (Cahoon et al., 2016), and due to root exudates and their ability of symbiotic N fixation burnt plots can also contain more N than unburnt lichen dominated plots (Maslov et al., 2018). In contrast to soil, fire substantially reduced C and N stocks in vegetation. Nevertheless, as vegetation stored much less C and  
240 N in comparison to soil, those effects were negligible for the estimation of total ecosystem stocks. In unburnt tundra, a large proportion of the aboveground C and N stocks was stored in reindeer lichens which dominated the vegetation before fire (Heim et al., 2021). As reindeer lichens recover very slowly, they were rare even several decades after fire (Frost et al., 2020; Heim et al., 2021) and did not reach the C and N stocks of unburnt sites. Instead, C and N were stored in shrubs and bryophytes, a pattern that reflects the vegetation recovery (Frost et al., 2020). However, shrubs and bryophytes did not compensate for C and  
245 N loss in the burnt lichen biomass.

#### 4.1 Fire affected long-term C and N cycling

Fire had a long-term effect on C and N concentrations and stable isotope ratios in aboveground vegetation, especially lichens, which suggests a strong, long-lasting effect of fire on C and N cycling in the ecosystem. The long-term fire effects on nutrient  
250 cycling in soil were minor and most apparent in the topsoil.

Fire did not increase N concentrations in vascular plants. Originally, we expected that fire enhances mineralization rates, which releases previously unavailable N that can be taken up by vascular plants (Aerts, 2006; Salmon et al., 2016). However, our results do not support this hypothesis. On the contrary, we even found signs of nitrogen limitation in herbs and graminoids at the younger fire scars, as indicated by lower N concentration and higher C/N ratios in the biomass. A possible explanation  
255 for this pattern may be linked to enhanced competition for nitrogen among vascular plants, which increased during post-fire succession (Bret-Harte et al., 2013; Heim et al., 2021). Unburnt plots were dominated by lichens, which obtain large parts of their nutrients from the atmosphere (Asplund and Wardle, 2017) and thus did not compete for available soil N. Therefore, vascular plants at unburnt plots may have relatively more available soil-N. Lichens reflected long-term impacts of fire on N cycling. We found high N concentrations on burnt plots of the youngest fire scar. The disappearance of this effect with time  
260 since fire might be related to the fact that younger lichens generally contain more N (Kytöviita and Crittenden, 2007).

Soil N concentrations were only increased in the upper soil layer of the oldest fire scar. This pattern is less likely linked to the temperature-mediated increased microbial activity, as the soil temperature in the oldest fire scar recovered to control levels (Heim et al., 2021). Rather, this might be explained by the increased cover of vascular plants, which produce more root exudates, have symbiotic nitrogen fixation, and easily decomposable falloff litter (Maslov et al., 2018; McLaren et al., 2017).  
265 The lower soil  $\delta^{13}\text{C}$  on burnt plots of the oldest fire scar (where soil temperatures returned to control levels) is probably linked to the altered vegetation composition with more shrubs and compound-specific variations of  $\delta^{13}\text{C}$  in litter. In plants, easily decomposing substances are comparatively enriched in  $^{13}\text{C}$ , while more refractory substances, such as lignin, are relatively depleted in  $^{13}\text{C}$  (Ågren et al., 1996). Increased shrub litter and a higher proportion of lignin at the oldest fire scar may thus result in a depletion of heavy  $^{13}\text{C}$  in the remaining organic matter of the soil (Ågren et al., 1996).

270 Why  $\delta^{13}\text{C}$  in vascular and lower plants on fire scars is decreased in our study remains relatively unclear as variations in  $\delta^{13}\text{C}$  are usually complex and not straightforward to interpret (Dawson et al., 2002). Therefore, our findings could not be related with certainty to a process described in literature. One reason for the decreased  $\delta^{13}\text{C}$  might be the lower  $^{13}\text{C}$  content of the  $\text{CO}_2$  in the ambient air of fire scars. A lower  $^{13}\text{C}$  content of the  $\text{CO}_2$  can be explained by increased decomposition rates (Dawson et al., 2002; Lakatos et al., 2007). However, we could not detect a decrease in C stocks in our data that would allow the assumption  
275 of increased mineralisation.

In our case, water availability may be a better explanation for the observed shifts towards lower  $^{13}\text{C}$  values in vegetation, as increased water availability is common in post-fire permafrost landscapes (Holloway et al., 2020). This is in line with the frequently observed negative correlations of indicators for humidity and tree ring  $^{13}\text{C}$  (e.g. Holzkämper et al., 2012), as  $^{13}\text{C}$  reflects stomatal conductance as affected by moisture availability or drought stress. Dawson 2002 states that “But unlike in  
280 vascular plants, delta tends to increase with water limitation in nonvascular plant taxa (Williams and Flanagan 1996, 1998) “. Our soil moisture data, however, does not support this, which might be due to the timing of sampling (soil moisture is also probably the most variable parameter in space and time). We did not find an effect of fire on the abundance of  $^{15}\text{N}$  in the biomass of vascular plants. This was surprising because we expected that  $^{15}\text{N}$  in leaves of vascular plants on younger fire scars would be increased, while it would decrease in later successional stages, through the progressing colonization with mycorrhizas  
285 (Hobbie et al., 2005).

The lack of an effect of past fires on  $\delta^{15}\text{N}$  may indicate that the belowground mycorrhizas remained relatively unaffected by fire, and the resprouting strategy of tundra shrubs facilitates the resilience of dominant fungi against tundra fires (Hewitt et al., 2013). However, on closer inspection of the shrub species *Betula nana* on the intermediate fire scar, we detected an enrichment of  $^{15}\text{N}$  in leaves of the burnt plots in comparison to plants on unburnt plots (Fig. S11). This indicates that the shrubs on the fire  
290 scar may not have been re-colonised again by mycorrhizas (Hobbie et al., 2005). We probably did not find an overall effect of fire on shrubs because we included too many species in the analysis that interact with different mycorrhizas or no mycorrhiza at all - which makes a pattern unrecognizable. While fire had no significant effect on  $\delta^{15}\text{N}$  in vascular plants, the positive effect of fire on  $\delta^{15}\text{N}$  in lichens probably reflects the age of the thallus. Ellis et al. (2003) showed that there is a consistent pattern of  $\delta^{15}\text{N}$  distribution in the lichen thallus, with the highest  $\delta^{15}\text{N}$  content in the apices and a minimum located at 2-4 cm below the  
295 apex. The high  $\delta^{15}\text{N}$  can thus be related to the fact that lichen thalli on burnt areas are younger and thus smaller in comparison to unburnt stands (Heim et al., 2021).

## 5 Conclusions

While our results demonstrate that tundra fires do not generally reduce C and N stocks, we show that fire disturbance is an important long-term driver of C and N cycling in the subarctic tundra ecosystem of northern Siberia. We found that lichens  
300 play an important role in the storage of C and N in the aboveground biomass.

We did not find recovery to pre-fire conditions in terms of C and N cycling, even after more than 44 years.

Overall, our results strongly suggest that in lichen-dominated subarctic tundra ecosystems, the contribution of wildfires to the release of additional carbon to the atmosphere might be relatively small as soil stocks appear to be resilient.

## 6 Funding

305 RJH was supported by a scholarship from the Studienstiftung des deutschen Volkes. We acknowledge support from the Open Access Publication Fund of the University of Münster.

*Code and data availability.* The data and R-scripts that support the findings of this study are openly available in “Zenodo” at <https://doi.org/10.5281/zenodo.5582683> and <https://doi.org/10.5281/zenodo.5583511>

*Author contributions.* RJH: Conceptualization, Data curation, Formal analysis, Funding acquisition, Investigation, Methodology, Project administration, Visualization, Writing – original draft preparation, AY: Data curation, Investigation, Methodology, Project administration, Resources, Validation, Writing – review and editing, AB: Supervision, Validation, Writing – review and editing, WH: Investigation, Validation, Writing – review and editing, VK: Investigation, KHK: Methodology, Resources, Validation, Writing – review and editing, CL: Formal analysis, Investigation, Validation, Writing – review and editing, AP: Investigation, DS: Investigation, FS: Investigation, NH: Conceptualization, Funding acquisition, Methodology, Project administration, Resources, Supervision, Validation, Writing – review and editing

315 *Competing interests.* The authors declare that they have no conflict of interest.

*Acknowledgements.* We would like to thank Julia Aminova, Denis Bazyk, Leya Brodt, Marvin Diek, Bettina Haas, Liv Jessen, Olga Konkova, Ksenia Maryasicheva, Michael Moskowchenko and Daniel Rieker for help with data collection. We are grateful to the Department for Science and Innovation of the Yamalo-Nenets Autonomous Okrug (Alexei Titovsky), the Interregional Expedition Center “Arctic” (Elena Nyukina), as well as to the Arctic Research Center of the Yamalo-Nenets Autonomous Okrug (Andrey Lobanov) for the great financial and logistic support and fruitful cooperation. For the trustful collaboration, we thank also Andrei Tolstikov (University of Tyumen). The C and N analyses of this study were carried out in the laboratory of the Institute of Landscape Ecology of the University of Muenster. The assistance of Ulrike Berning-Mader is greatly acknowledged.

## References

- Aerts, R.: The freezer defrosting: global warming and litter decomposition rates in cold biomes, *Journal of Ecology*, 94, 713–724, 2006.
- 325 Ågren, G. I., Bosatta, E., and Balesdent, J.: Isotope discrimination during decomposition of organic matter: a theoretical analysis, *Soil Science Society of America Journal*, 60, 1121–1126, 1996.
- Asplund, J. and Wardle, D. A.: How lichens impact on terrestrial community and ecosystem properties, *Biological Reviews*, 92, 1720–1738, <https://doi.org/10.1111/brv.12305>, 2017.
- Blok, D., Faucherre, S., Banyasz, I., Rinnan, R., Michelsen, A., and Elberling, B.: Contrasting above-and belowground organic matter  
330 decomposition and carbon and nitrogen dynamics in response to warming in High Arctic tundra, *Global change biology*, 24, 2660–2672, 2018.
- Böhlke, J. K., Mroczkowski, S. J., and Coplen, T. B.: Oxygen isotopes in nitrate: New reference materials for 18O: 17O: 16O measurements and observations on nitrate-water equilibration, *Rapid Communications in Mass Spectrometry*, 17, 1835–1846, 2003.
- Bret-Harte, M. S., Mack, M. C., Shaver, G. R., Huebner, D. C., Johnston, M., Mojica, C. A., Pizano, C., and Reiskind, J. A.: The response of  
335 Arctic vegetation and soils following an unusually severe tundra fire, *Philosophical transactions of the Royal Society of London. Series B, Biological sciences*, 368, 20120490, <https://doi.org/10.1098/rstb.2012.0490>, 2013.
- Bürkner, P.-C.: brms: An R Package for Bayesian Multilevel Models Using Stan, *Journal of Statistical Software*, 80, 1–28, <https://doi.org/10.18637/jss.v080.i01>, 2017.
- Bürkner, P.-C.: Advanced Bayesian Multilevel Modeling with the R Package brms, *The R Journal*, 10, 395–411, <https://doi.org/10.32614/RJ-2018-017>, 2018.  
340
- Cahoon, S. M. P., Sullivan, P. F., and Post, E.: Greater abundance of *Betula nana* and early onset of the growing season increase ecosystem CO<sub>2</sub> uptake in West Greenland, *Ecosystems*, 19, 1149–1163, 2016.
- Chambers, S. D., Beringer, J., Randerson, J. T., and Chapin, I. S.: Fire effects on net radiation and energy partitioning: Contrasting responses of tundra and boreal forest ecosystems, *Journal of Geophysical Research D: Atmospheres*, 110, 1–9,  
345 <https://doi.org/10.1029/2004JD005299>, 2005.
- Chapin III, F. S., McGuire, A. D., Randerson, J., Pielke, R., Baldocchi, D., Hobbie, S. E., Roulet, N., Eugster, W., Kasischke, E., and Rastetter, E. B.: Arctic and boreal ecosystems of western North America as components of the climate system, *Global Change Biology*, 6, 211–223, 2000.
- Chen, Y., Hu, F. S., and Lara, M. J.: Divergent shrub-cover responses driven by climate, wildfire, and permafrost interactions in Arctic tundra  
350 ecosystems, *Global change biology*, 27, 652–663, 2021.
- Coplen, T. B., Brand, W. A., Gehre, M., Gröning, M., Meijer, H. A. J., Toman, B., and Verkouteren, R. M.: New guidelines for  $\delta^{13}\text{C}$  measurements, *Analytical Chemistry*, 78, 2439–2441, 2006.
- Dahl, E.: Stability of tundra ecosystems in Fennoscandia, in: *Fennoscandian tundra ecosystems*, pp. 231–236, Springer, 1975.
- Dawson, T. E., Mambelli, S., Plamboeck, A. H., Templer, P. H., and Tu, K. P.: Stable isotopes in plant ecology, *Annual review of ecology and systematics*, 33, 507–559, 2002.  
355
- De Baets, S., Van de Weg, M. J., Lewis, R., Steinberg, N., Meersmans, J., Quine, T. A., Shaver, G. R., and Hartley, I. P.: Investigating the controls on soil organic matter decomposition in tussock tundra soil and permafrost after fire, *Soil Biology and Biochemistry*, 99, 108–116, 2016.



- Dormann, C. F. and Woodin, S. J.: Climate change in the Arctic: using plant functional types in a meta-analysis of field experiments, *Functional Ecology*, 16, 4–17, 2002.
- 360 Ehleringer, J. R., Buchmann, N., and Flanagan, L. B.: Carbon isotope ratios in belowground carbon cycle processes, *Ecological Applications*, 10, 412–422, 2000.
- Ellis, C. J., Crittenden, P. D., Scrimgeour, C. M., and Ashcroft, C.: The natural abundance of  $^{15}\text{N}$  in mat-forming lichens, *Oecologia*, 136, 115–123, 2003.
- 365 Estop-Aragonés, C., Olefeldt, D., Abbott, B. W., Chanton, J. P., Czimeczik, C. I., Dean, J. F., Egan, J. E., Gandois, L., Garnett, M. H., and Hartley, I. P.: Assessing the Potential for Mobilization of Old Soil Carbon After Permafrost Thaw: A Synthesis of  $^{14}\text{C}$  Measurements From the Northern Permafrost Region, *Global Biogeochemical Cycles*, 34, e2020GB006672, 2020.
- Farquhar, G. D., Ehleringer, J. R., and Hubick, K. T.: Carbon isotope discrimination and photosynthesis, *Annual review of plant biology*, 40, 503–537, 1989.
- 370 Frost, G. V., Loehman, R. A., Saperstein, L. B., Macander, M. J., Nelson, P. R., Paradis, D. P., and Natali, S. M.: Multi-decadal patterns of vegetation succession after tundra fire on the Yukon-Kuskokwim Delta, Alaska, *Environmental Research Letters*, 15, 25003, 2020.
- Gorelick, N., Hancher, M., Dixon, M., Ilyushchenko, S., Thau, D., and Moore, R.: Google Earth Engine: Planetary-scale geospatial analysis for everyone, *Remote Sensing of Environment*, 202, 18–27, 2017.
- He, J., Chen, D., Jenkins, L., and Loboda, T. V.: Impacts of wildfire and landscape factors on organic soil properties in Arctic tussock tundra, *Environmental Research Letters*, 16, 085004, 2021.
- 375 Heim, R. J., Bucharova, A., Brodt, L., Kamp, J., Rieker, D., Soromotin, A. V., Yurtaev, A., and Hoelzel, N.: Post-fire vegetation succession in the Siberian subarctic tundra over 45 years, *Science of The Total Environment*, 760, 143425, <https://doi.org/https://doi.org/10.1016/j.scitotenv.2020.143425>, 2021.
- Hewitt, R. E., Bent, E., Hollingsworth, T. N., Chapin III, F. S., and Taylor, D. L.: Resilience of Arctic mycorrhizal fungal communities after wildfire facilitated by resprouting shrubs, *Ecoscience*, 20, 296–310, 2013.
- 380 Hobbie, E. A., Jumpponen, A., and Trappe, J.: Foliar and fungal  $^{15}\text{N}$ :  $^{14}\text{N}$  ratios reflect development of mycorrhizae and nitrogen supply during primary succession: testing analytical models, *Oecologia*, 146, 258–268, 2005.
- Holloway, J. E., Lewkowicz, A. G., Douglas, T. A., Li, X., Turetsky, M. R., Baltzer, J. L., and Jin, H.: Impact of wildfire on permafrost landscapes: A review of recent advances and future prospects, *Permafrost and Periglacial Processes*, 31, 371–382, 2020.
- 385 Holzkämper, S., Tillman, P. K., Kuhry, P., and Esper, J.: Comparison of stable carbon and oxygen isotopes in *Picea glauca* tree rings and *Sphagnum fuscum* moss remains from subarctic Canada, *Quaternary Research*, 78, 295–302, 2012.
- Hu, F. S., Higuera, P. E., Duffy, P., Chipman, M. L., Rocha, A. V., Young, A. M., Kelly, R., and Dietze, M. C.: Arctic tundra fires: Natural variability and responses to climate change, *Frontiers in Ecology and the Environment*, 13, 369–377, <https://doi.org/10.1890/150063>, 2015.
- 390 IUSS Working Group WRB: World Reference Base for Soil Resources 2014, update 2015 International soil classification system for naming soils and creating legends for soil maps. World Soil Resources Reports No. 106., FAO, Rome, 2015.
- Jandt, R., Joly, K., Meyers, C. R., and Racine, C.: Slow Recovery of Lichen on Burned Caribou Winter Range in Alaska Tundra: Potential Influences of Climate Warming and Other Disturbance Factors, *Arctic, Antarctic, and Alpine Research*, 40, 89–95, <https://doi.org/10.1657/1523-0430>, 2008.
- 395 Jansson, J. K. and Hofmockel, K. S.: Soil microbiomes and climate change, *Nature Reviews Microbiology*, 18, 35–46, <https://doi.org/10.1038/s41579-019-0265-7>, 2020.

- Jiang, Y., Rocha, A. V., O'Donnell, J. A., Drysdale, J. A., Rastetter, E. B., Shaver, G. R., and Zhuang, Q.: Contrasting soil thermal responses to fire in Alaskan tundra and boreal forest, *Journal of Geophysical Research: Earth Surface*, 120, 363–378, <https://doi.org/10.1002/2014jfg003180>, 2015.
- 400 Jonasson, S., Chapin Iii, F. S., and Shaver, G. R.: Biogeochemistry in the Arctic: patterns, processes, and controls, in: *Global biogeochemical cycles in the climate system*, pp. 139–150, Elsevier, 2001.
- Kazakov, K.: Spravochno-informatsionnyy portal "Pogoda i klimat" [Reference information portal "Weather and Climate"], [WWW document] [accessed 2 May 2019], <http://www.pogodaiklimat.ru/msummary.php?m=all{&}y=all{&}id=23256>, 2019.
- Korner-Nievergelt, F., Roth, T., Felten, S. V., Guelat, J., Almasi, B., and Korner-Nievergelt, P.: *Bayesian Data Analysis in Ecology using Linear Models with R, BUGS and Stan*, Elsevier, London, 2015.
- 405 Kornienko, S. G.: Cartography of pyrogenic violations of the vegetation cover on the Tazovsky Peninsula with satellite data, *Actual problems of oil and gas*, 1, 1–11, 2018.
- Kytöviita, M.-M. and Crittenden, P. D.: Growth and nitrogen relations in the mat-forming lichens *Stereocaulon paschale* and *Cladonia stellaris*, *Annals of botany*, 100, 1537–1545, 2007.
- 410 Lakatos, M., Hartard, B., and Máguas, C.: The stable isotopes  $\delta^{13}\text{C}$  and  $\delta^{18}\text{O}$  of lichens can be used as tracers of microenvironmental carbon and water sources, *Terrestrial Ecology*, 1, 77–92, 2007.
- Lasslop, G., Hantson, S., Harrison, S. P., Bachelet, D., Burton, C., Forkel, M., Forrest, M., Li, F., Melton, J. R., and Yue, C.: Global ecosystems and fire: multi-model assessment of fire-induced tree cover and carbon storage reduction, *Global Change Biology*, 2020.
- Longton, R. E.: The role of bryophytes and lichens in polar ecosystems, *Special Publication-British Ecological Society*, 13, 69–96, 1997.
- 415 Loranty, M. M., Natali, S. M., Berner, L. T., Goetz, S. J., Holmes, R. M., Davydov, S. P., Zimov, N. S., and Zimov, S. A.: Siberian tundra ecosystem vegetation and carbon stocks four decades after wildfire, *Journal of Geophysical Research: Biogeosciences*, 119, 2144–2154, <https://doi.org/10.1002/2014jg002730>, 2014.
- Mack, M. C., Schuur, E. A. G., Bret-Harte, M. S., Shaver, G. R., and Chapin, F. S.: Ecosystem carbon storage in arctic tundra reduced by long-term nutrient fertilization, *Nature*, 431, 440–443, 2004.
- 420 Mack, M. C., Bret-Harte, M. S., Hollingsworth, T. N., Jandt, R., Schuur, E. A., Shaver, G. R., and Verbyla, D. L.: Carbon loss from an unprecedented Arctic tundra wildfire, *Nature*, 475, 489–492, <https://doi.org/10.1038/nature10283>, 2011.
- Marion, G. M., Bockheim, J. G., and Brown, J.: Arctic soils and the ITEX experiment, *Global Change Biology*, 3, 33–43, 1997.
- Maslov, M. N., Maslova, O. A., Pozdnyakov, L. A., and Kopeina, E. I.: Biological activity of soils in mountain tundra ecosystems under postpyrogenic restoration, *Eurasian Soil Science*, 51, 692–700, 2018.
- 425 Maslov, M. N., Maslova, O. A., and Kopeina, E. I.: Changes in the pools of total and labile soil organic carbon during post-fire succession in the Khibiny Mountain tundra ecosystems, *Eurasian Soil Science*, 53, 330–338, 2020.
- McLaren, J. R., Buckeridge, K. M., van de Weg, M. J., Shaver, G. R., Schimel, J. P., and Gough, L.: Shrub encroachment in Arctic tundra: *Betula nana* effects on above-and belowground litter decomposition, *Ecology*, 98, 1361–1376, 2017.
- Mollicone, D., Eva, H. D., and Achard, F.: Ecology: Human role in Russian wild fires, *Nature*, 440, 436, 2006.
- 430 Moritz, M. A., Parisien, M.-A., Batllori, E., Krawchuk, M. A., Van Dorn, J., Ganz, D. J., and Hayhoe, K.: Climate change and disruptions to global fire activity, *Ecosphere*, 3, 1–22, 2012.
- Narita, K., Harada, K., Saito, K., Sawada, Y., Fukuda, M., and Tsuyuzaki, S.: Vegetation and Permafrost Thaw Depth 10 Years after a Tundra Fire in 2002, Seward Peninsula, Alaska, Arctic, Antarctic, and Alpine Research, 47, 547–559, <https://doi.org/10.1657/AAAR0013-031>, 2015.

- 435 Oulehle, F., Rowe, E. C., Myška, O., Chuman, T., and Evans, C. D.: Plant functional type affects nitrogen use efficiency in high-Arctic tundra, *Soil Biology and Biochemistry*, 94, 19–28, 2016.
- Peterson, B. J. and Fry, B.: Stable isotopes in ecosystem studies, *Annual review of ecology and systematics*, 18, 293–320, 1987.
- Post, E., Alley, R. B., Christensen, T. R., Macias-Fauria, M., Forbes, B. C., Gooseff, M. N., Iler, A., Kerby, J. T., Laidre, K. L., and Mann, M. E.: The polar regions in a 2° C warmer world, *Science Advances*, 5, eaaw9883, 2019.
- 440 R Core Team: R: A Language and Environment for Statistical Computing, Vienna, Austria, 2020.
- Rinnan, R., Michelsen, A., Bååth, E., and Jonasson, S.: Fifteen years of climate change manipulations alter soil microbial communities in a subarctic heath ecosystem, *Global Change Biology*, 13, 28–39, 2007.
- Rosen, Y.: Warming, fires, warming, fires: How tundra wildfires could create an unstoppable cycle, <https://www.adn.com/alaska-news/science/2016/05/30/warming-fires-warming-fires-how-tundra-wildfires-could-create-an-unstoppable-cycle/>, 2017.
- 445 Salmon, V. G., Soucy, P., Mauritz, M., Celis, G., Natali, S. M., Mack, M. C., and Schuur, E. A. G.: Nitrogen availability increases in a tundra ecosystem during five years of experimental permafrost thaw, *Global Change Biology*, 22, 1927–1941, 2016.
- Sancho, L. G., Belnap, J., Colesie, C., Raggio, J., and Weber, B.: Carbon budgets of biological soil crusts at micro-, meso-, and global scales, in: *Biological soil crusts: an organizing principle in drylands*, pp. 287–304, Springer, 2016.
- Schuur, E. A. G., Trumbore, S. E., Mack, M. C., and Harden, J. W.: Isotopic composition of carbon dioxide from a boreal forest fire: Inferring
- 450 carbon loss from measurements and modeling, *Global Biogeochemical Cycles*, 17, 1, 2003.
- Schuur, E. A. G., Vogel, J. G., Crummer, K. G., Lee, H., Sickman, J. O., and Osterkamp, T. E.: The effect of permafrost thaw on old carbon release and net carbon exchange from tundra, *Nature*, 459, 556–559, 2009.
- Schuur, E. A. G., McGuire, A. D., Schädel, C., Grosse, G., Harden, J. W., Hayes, D. J., Hugelius, G., Koven, C. D., Kuhry, P., and Lawrence, D. M.: Climate change and the permafrost carbon feedback, *Nature*, 520, 171–179, 2015.
- 455 Turetsky, M. R.: The role of bryophytes in carbon and nitrogen cycling, *The bryologist*, 106, 395–409, 2003.
- Turetsky, M. R., Bond-Lamberty, B., Euskirchen, E., Talbot, J., Frohling, S., McGuire, A. D., and Tuittila, E.: The resilience and functional role of moss in boreal and arctic ecosystems, *New Phytologist*, 196, 49–67, 2012.
- Turetsky, M. R., Benscoter, B., Page, S., Rein, G., Van Der Werf, G. R., and Watts, A.: Global vulnerability of peatlands to fire and carbon loss, *Nature Geoscience*, 8, 11–14, <https://doi.org/10.1038/ngeo2325>, 2015.
- 460 U.S. Geological Survey: Earth Explorer, <https://earthexplorer.usgs.gov/>, 2018.
- Veraverbeke, S., Delcourt, C. J. F., Kukavskaya, E., Mack, M., Walker, X., Hessilt, T., Rogers, B., and Scholten, R. C.: Direct and longer-term carbon emissions from arctic-boreal fires: a short review of recent advances, *Current Opinion in Environmental Science & Health*, p. 100277, 2021.
- Vilchek, G. E. and Bykova, O. Y.: The origin of regional ecological problems within the northern Tyumen Oblast, Russia, *Arctic and Alpine*
- 465 *Research*, 24, 99–107, 1992.
- Young, A. M., Higuera, P. E., Duffy, P. A., and Hu, F. S.: Climatic thresholds shape northern high-latitude fire regimes and imply vulnerability to future climate change, *Ecography*, 39, 1–12, 2016.
- Yu, Q., Epstein, H. E., Engstrom, R., Shiklomanov, N., and Streletskiy, D.: Land cover and land use changes in the oil and gas regions of Northwestern Siberia under changing climatic conditions, *Environmental Research Letters*, 10, 124 020, 2015.
- 470 Yurkovskaya, T. K.: Vegetation map [Karta rastitel'nosti], in: *National Atlas of Soils of the Russian Federation [Natsional'nyy atlas pochv Rossiyskoy Federatsii]*, edited by Shoba, S. A., Alyabina, I. O., Urusevskaya, I. S., and Chernova, O. V., pp. 46–48, Moscow State University, Moscow, 2011.