

Soil greenhouse gas fluxes from tropical coastal wetlands and alternative agricultural land uses

10 Naima Iram¹, Emad Kavehei¹, Damien. T. Maher², Stuart. E. Bunn¹, Mehran Rezaei Rashti¹, Bahareh
5 Shahrabi Farahani¹, Maria Fernanda Adame¹

¹Australian Rivers Institute, Griffith University, Nathan, QLD, 4111, Australia

²School of Environment, Science and Engineering, Southern Cross University, Lismore, NSW, 2480, Australia

10 *Correspondence to;* Naima Iram (naima.iram@griffithuni.edu.au)

Abstract. Coastal wetlands are essential for regulating the global carbon budget through soil carbon sequestration and greenhouse gas fluxes (GHG: CO₂, CH₄ and N₂O). The conversion of coastal wetlands to agricultural land alters the magnitude and direction (uptake/release) of these fluxes. However, the extent and drivers of change of GHG fluxes is still unknown for many tropical regions. We measured soil GHG fluxes from three natural coastal wetlands: mangroves, saltmarsh, and freshwater tidal forests, and two alternative agricultural land use, sugarcane farming and pastures for cattle grazing (ponded and dry conditions). We assessed variations throughout different climatic conditions (dry-cool, dry-hot and wet-hot) within two years of measurements (2018-2020) in tropical Australia. The wet pasture had by far the highest CH₄ emissions with 1,231 ± 386 mg m⁻² d⁻¹, which were 200-fold higher than any other site. Dry pastures and sugarcane were the highest emitters of N₂O with 55 ± 9 mg m⁻² d⁻¹ (wet-hot period) and 11 ± 3 mg m⁻² d⁻¹ (hot-dry period, coinciding with fertilisation), respectively. Dry pastures were also the highest emitters of CO₂ with 20 ± 1 g m⁻² d⁻¹ (wet-hot period). The three coastal wetlands measured had lower emission, with saltmarsh taking -0.55 ± 0.23 of N₂O and -1.19 ± 0.08 g m⁻² d⁻¹ of CO₂ during the dry-hot period. During the sampled period, sugarcane and pastures had higher total cumulative soil GHG emissions (CH₄ + N₂O) of 7,142 and 56,124 CO₂eq kg ha⁻¹ y⁻¹ compared to coastal wetlands with 144 to 884 CO₂eq kg ha⁻¹ y⁻¹. Restoring unproductive sugarcane land or pastures (especially ponded ones) to coastal wetlands could provide significant GHG mitigation.

1 Introduction

Coastal wetlands are found at the interface of terrestrial and marine ecosystems and account for 10% of the global wetland area (Lehner and Döll 2004). They are highly productive and provide various ecosystem services such as water quality improvement, biodiversity, and carbon sequestration (Duarte et al, 2013). For instance, mangroves can accumulate five times 30 more soil carbon than terrestrial forests (Kauffman et al, 2020). However, the high productivity and anoxic soil conditions that promote carbon sequestration can also favour potent greenhouse gas emissions (GHGs), including CO₂, CH₄ and N₂O (Whalen, 2005; Conrad, 2009).

The GHG emissions in coastal wetlands primarily result from microbial processes in the soil-water-atmosphere 35 interface (Bauza et al, 2002; Whalen, 2005). The emission of CO₂ is a result of respiration, where fixed carbon by photosynthesis is partially released back into the atmosphere (Oertel et al., 2016). Emissions of CH₄ result from anaerobic and aerobic respiration by methanogenic bacteria, mostly in waterlogged conditions (Angle et al., 2017; Saunois et al, 2020). Finally, N₂O emissions are caused by denitrification in anoxic conditions and nitrification in aerobic soils, both driven by 40 nitrogen content and soil moisture (Ussiri and Lal 2013). Thus, the total GHG emissions from a wetland are driven by environmental conditions that favour these microbial processes, all of which result in highly variable emissions from wetlands worldwide (Kirschke et al., 2013; Oertel et al. 2016).

Despite potential high GHG emissions from coastal wetlands, these are likely to be lower than those from alternative agricultural land uses (Knox et al., 2015), which emit GHGs from their construction throughout their productive lives. Firstly, 45 when wetlands are converted to agricultural land, the oxidation of sequestered carbon in the organic-rich soils release significant amounts of CO₂ (Drexler, de Fontaine, & Deverel, 2009; Hooijer et al, 2012). Secondly, removing tidal flow and converting coastal wetlands to freshwater systems, such as during the creation of ponded pastures, dams or agricultural ditches, can result in very high CH₄ emissions (Deemer et al., 2016; Grinham et al, 2018; Ollivier et al, 2019). For instance, agricultural 50 ditches contribute up to 3% of the total anthropogenic CH₄ emissions globally (Peacock et al., 2021). Finally, the use of fertilisers significantly increases N₂O emissions (Rashti et al, 2015). Thus, emissions of GHG from land-use change can be mitigated through the reversal of these activities, for instance, reduction of fertiliser use and the reinstallation of tidal flow on unused agricultural land (Rashti et al, 2016; Kroeger et al. 2017).

This study measured the annual GHG fluxes (CO₂, CH₄ and N₂O) from three natural coastal wetlands (mangroves, 55 saltmarsh and freshwater tidal forests) and two agricultural land use sites (sugarcane plantation and pasture) in tropical Australia. The objectives were to assess the difference in GHG fluxes throughout different seasons that characterise tropical climates (dry-cool, dry-hot and wet-hot) and to identify environmental factors associated with these GHG fluxes. These data


will inform emission factors for converting wetlands to agricultural land uses and vice versa, filling in a knowledge gap identified in Australia (Baldock et al., 2012) and tropical regions worldwide (IPCC, 2013).


60 2 Materials and Methods

2.1 Study sites


The study area is located within the Herbert River catchment in Queensland, northeast Australia (Fig 1a). The region has a tropical climate with a mean monthly minimum temperature from 14 to 23°C and mean monthly maximum temperature from 25 to 33°C (Australian Bureau of Meteorology, ABM, 2020; 1968-2020; Table S3). The average rainfall is 2,158 mm y^{-1} , with 65 the highest values of 476 mm during February (ABM 2020; 1968-2020; Table S3). The Herbert basin covers 9,842 km 2 , from which 56% is grazing, 31% is conserved wetlands and forestry, 8% is sugarcane, and 4% is other land uses (Department of Science and Environment, QLD, DES, WetlandInfo, 2020). Wetlands in this region were heavily deforested in the past century (1943- 1996) due to rapid agricultural development, primarily for sugarcane farming (Griggs. 2018). Before clearing, the land was mostly covered by rainforest and coastal wetlands, mainly *Melaleuca* forest, grass and sedge swamps (Johnson, Ebert, & 70 Murray, 1999).

We selected five sites, including three natural coastal wetlands (Fig. 1): a mangrove forest (18° 53' 42" S, 146° 15' 51" E), a saltmarsh (18° 53' 43" S, 146° 15' 52" E) and a freshwater tidal forest (18° 53' 45" S, 146° 15' 52" E), and two common agricultural land use types of the region, a sugarcane farm (18° 53' 44.6" S, 146° 15' 53.2" E) and a pasture for fodder 75 grazing. The pasture had different levels of inundation; some areas were covered with shallow ponds (50-100 cm depth), wet grassy areas (hereafter “wet pasture”; 18° 43' 8" S, 146° 15' 50" E) and drier areas (hereafter, “dry pasture”; 18° 43' 7" S, 146° 15' 50" E). The natural coastal wetlands and the sugarcane site were located within the same property at Insulator Creek and were located < 200 m apart (Fig. 1B), while the ponded pasture was 20 km north at Mungalla Station (Fig. 1A). The mangroves were dominated by *Avicennia marina* with few plants of *Rhizophora stylosa*, and the saltmarsh was dominated by 80 *Sueda salsa* and *Sporobolus* spp. Landwards, the freshwater tidal freshwater forest, a wetland commonly known as “tea tree swamp”, was dominated by *Melaleuca quinquenervia* trees. While the mangroves and saltmarsh are directly submerged by tides (5- 30 cm), the tidal freshwater forest is indirectly affected by tidal fluctuations, such as during large spring tides, when tidal water can push groundwater above the forest. The coastal wetlands were adjacent to a sugarcane farm with an area of ~110 ha (Fig. 1b). The sugarcane is fertilised once a year with urea at a rate of 135 kg N ha $^{-1}$ and harvested during May-June, 85 while the foliage is left on the soil surface (trash blanket) after harvest. The ponded pastures in Mungalla Station extend over 2,500 ha and support ~900 cattle throughout the year by providing fodder to cattle during dry periods. The selected ponded pastures were covered by *Eichhornia crassipes* (water hyacinth) and *Hymenachne amplexicaulis* (Fig. 1g-h). Each of the five sites was sampled during three periods dry-cool (May-September), dry-hot (October-December) and wet-hot (January-April; Table 1). During each time, soil physicochemical properties and GHG fluxes were measured as detailed below.






(c) Mangroves




(d) Saltmarsh




(e) Freshwater tidal forest



(f) Sugarcane



(g) Dry pasture



(h) Wet pasture

Figure 1: a) Location of sampling sites (Insulator Creek and Mungalla) in the Herbert River catchment, northeast Australia, (b) natural wetlands adjacent to sugarcane farm in Insulator Ck and sampling locations, and (c) mangroves, (d) saltmarsh, (e) freshwater tidal forest, (f) sugarcane, (g) dry pasture and (h) wet pasture. Pictures by N. Iram and MF Adame.

Table 1. Mean daily air temperature and rainfall (Ingham, Qld. weather station) during sampling.

Season	Study period	Daily min temperature (°C)	Daily max temperature (°C)	Rainfall (mm d ⁻¹)
Dry-cool	17/06/2018	13.4 - 14.6	27.7 - 28.2	0
Dry-hot	23-29/10/2018	15.7 - 21.1	32.2 - 36.2	0
Dry-cool	31/05 to 6/06/2019	10.9 - 17.5	21.6 - 28.2	0-25
Wet-hot	17-22/02/2020	23.9 - 25.3	33.6 - 34.5	0-86

100

2.2 Soil sampling and analysis of physicochemical properties

Soil physicochemical characteristics were measured by taking composite soil samples next to each gas chamber location (n = 5; 0-30 cm) for all study sites during the dry-hot season. The samples were obtained by inserting an open steel corer to a depth of 30 cm; the core was divided into three depths: 0-10, 10-20 and 20-30 cm. Soil samples were oven-dried at 105 °C for 48 h to determine volumetric water content through gravimetric analysis. The volumetric water content was divided by total soil porosity to determine water-filled pore space (WFPS). Total soil porosity was calculated through the following equation from Rashti et al. (2015, Eq. 1).

110

$$\text{Total soil porosity} = 1 - [(\text{soil bulk density (mg cm}^{-3}) / 2.65)]$$

Equation 1.

115

Soil texture analysis (% sand, silt, clay) was done with a simplified method for particle size determination (Kettler et al, 2001). Soil electrical conductivity (EC) and pH were measured using a conductivity meter (WP-84 TPS, Australia) in soil/water slurry at 1:5. Soil subsamples were air-dried, sieved (2mm), ground (Retch™ mill) and analysed for %N and %C using an elemental analyser connected to a gas-isotope ratio mass spectrometer (EA-Delta V Advantage IRMS, Griffith University). Additionally, soil samples from the top 10 cm were collected during each sampling to measure gravimetric soil moisture content and bulk density.

2.3 Greenhouse gas fluxes

120 We measured GHG fluxes (CO₂, CH₄ and N₂O) at each site for three consecutive days during each sampling period except for the dry-cool period of 2018, when mangroves, saltmarsh and sugarcane were surveyed for one day. The sampling was done between 9:00 to 11:00 am, representing the mean daily temperatures, thus, minimising variability of cumulative seasonal fluxes

based on intermittent manual flux measurements (Reeves et al, 2016). Additionally, we assessed the variability of our measurements with tidal inundation in mangroves and saltmarsh, which were regularly inundated (~10-30 cm) during hot-dry
125 season. For this, we measured GHG emissions during a low (0.7m) and a high tide (2.8m; Lucinda, 18° 31' S; 146° 23' E) in the dry cool period of 2019. We found that CH₄ fluxes did not significantly vary between the low and high tide within all coastal wetlands. Contrarily, for saltmarsh, CO₂ was taken during the high tide ($1.12 \pm 0.24 \text{ g m}^{-2} \text{ d}^{-1}$) but emitted ($0.69 \pm 0.4 \text{ g m}^{-2} \text{ d}^{-1}$) during the low tide ($F_{1,28} = 20.06, p < 0.001$). Finally, for N₂O, fluxes differed in all coastal wetlands, with higher uptakes in the high tide for mangroves ($F_{1,28} = 38.28, p < 0.001$; $F_{1,28} = 13.53, p = 0.001$) and higher release for saltmarsh ($F_{1,28} = 38.31, p < 0.001$) during low tide (Table S4). These results suggested that for CO₂ and N₂O fluxes, there was a probability of variation depending on the time of sampling. Thus, fHowever, due to logistic constraint, further sampling was conducted only during low tides.

135 We used static, manual gas chambers made of high-density, round polyvinyl chloride pipe, which consisted of two units: a base ($r = 12 \text{ cm}$, $h = 18 \text{ cm}$) and a detachable collar ($h = 12 \text{ cm}$; Hutchinson and Mosier, 1981; Kavehei et al, 2020). The chambers had lateral holes that could be left covered with rubber bungs at low water levels and left open at high water levels to allow for water movement between sampling events. When the wetlands were inundated for the experiments, we used PVC extensions ($h = 18 \text{ cm}$). Five chambers were set ~ 5cm deep in the soil separated one to two meters from each other selectively located on soil with minimal vegetation, roots, and crab burrows.

140 Five chambers were set ~ 5cm deep in the soil at random locations one day before sampling to minimise the disturbance of installation during the experiment (Rashti et al, 2015). The chambers were selectively located on soil with minimal vegetation, roots, and crab burrows. We were careful not to tramp around the chambers during installation and sampling. The fact that emissions were not significantly different among days ($p > 0.05$) provided us with confidence that disturbance due to installation was not problematic.

145 At the start of the experiment, gas chambers were closed. A sample was taken at time zero and then after one hour with a 20 ml syringe and transferred to a 12 mL-vacuumed exetainer (Exetainer, Labco Ltd., High Wycombe, UK). During the dry-hot season, linearity tests of GHG fluxes with time were conducted by sampling at 0, 20, 40 and 60 min (Rashti et al, 2016). For the rest of the experiments, linearity tests were performed in one of the five chambers at each site; R^2 values were
150 consistently above 0.70. During each experiment, soil temperature was measured next to each chamber. At the end of the experiment, the depth of the base was recorded from five points within each chamber to calculate the headspace volume. The obtained volumetric unit concentrations were converted to mass-based units using the Ideal Gas Law (Hutchinson and Mosier, 1981).

155 The GHG concentrations of all samples were analysed within two weeks of sampling with a gas chromatograph (Shimadzu GC-2010 Plus). For N₂O analysis, an electron capture detector was used with helium as the carrier gas, while CH₄

was analysed on a flame ionisation detector with nitrogen as the carrier gas. For CO₂ determination, the gas chromatograph was equipped with a thermal conductivity detector. Peak areas of the samples were compared against standard curves to determine concentrations (Chen et al, 2012). Seasonal cumulative GHG fluxes were calculated by modifying the equation 160 described by Shaaban et al. (2015; Eq. 2):

$$\text{Seasonal cumulative GHG fluxes (mg or } \mu\text{g m}^{-2} \text{ yr}^{-1}) = \sum_{i=1}^n (R_i \times 24 \times D_i \times 17.38)$$

Equation 2

Where;

R_i = Gas emission rate (mg m⁻² hr⁻¹ for CO₂ and μ g m⁻² hr⁻¹ for CH₄ and N₂O),

165 D_i = Mean daily GHG emission rate in a season, (mg m⁻² d⁻¹ for CO₂ and μ g m⁻² d⁻¹ for CH₄ and N₂O) during low tide

17.38 = number of weeks in each season , assuming these conditions were representative of the annual cycle (see Table 1).

Annual cumulative soil GHG fluxes (CH₄ + N₂O) were calculated by integrating cumulative seasonal fluxes. These estimations did not account for soil CO₂ values as our methodology with dark chambers only accounted for emissions from respiration and excluded uptake from primary productivity. The CO₂-equivalent (CO₂-eq) values were estimated by multiplying

170 CH₄ and N₂O emissions by 25 and 298, respectively (Solomon, 2007), which represent the radiative balance of these gases (Neubauer, 2021). For annual cumulative soil GHG flux calculations from coastal wetlands, we used GHG fluxes measured during low tide therefore our values did not incorporate effect of tidal fluctuations. It's worth noting that spatial and temporal replication of our studies was limited due to safety, technical and financial constraints therfore we assumed that our sampling design represented the GHG flux variation between sites. Therefore, results of the study should be carefully used considering
175 the study limitations regarding sampling replicates. For annual cumulative soil GHG flux calculations from coastal wetlands, we used GHG fluxes measured during low tide; therefore, our values did not incorporate the effect of tidal fluctuations. The spatial and temporal replication of this study targeted spatial variation within soil type (< 50 cm, five chambers), days (three days per sampling) and seasons (three seasons per year). However, our replication within land use and wetland type was limited; thus, generalisations for all wetlands and land uses should be done acknowledging this limitation.

180

2.4 Statistical analyses

GHG flux data were tested for normality through Kolmogorov-Smirnov and Shapiro-Wilk tests. The data was then analysed for spatial and temporal differences with one-way Analyses of Variance (ANOVA), where site and season were the predictive 185 factors and the replicate (chamber) was the random factor of the model. When data were not normal, they were transformed (log10 or 1/x) to comply with the assumptions of normality and homogeneity of variances. Some variables were not normally distributed despite transformations and were analysed with the non-parametric Kruskal-Wallis test and Mann-Whitney U Test.

A Pearson correlation test was run to evaluate the correlation of GHG with measured environmental factors. Analyses were done with SPSS (v25, IBM, New York, USA), and values are presented as mean \pm standard error (SE).

190 **3 Results**

3.1 Soil physicochemical properties

Soil physical and chemical parameters (mean values 0-30 cm) varied among sites (Table 2, see full results of statistical analyses in Supplementary Material). As expected, gravimetric moisture content was highest in the coastal wetlands and wet pasture ($> 26\%$) and lowest in the sugarcane and the dry pasture ($< 14\%$). All soils were acidic, especially the freshwater tidal forest and 195 the wet pastures with values < 5 throughout the sediment column; mangroves had the highest pH with 6.0 ± 0.1 . The lowest EC was recorded in the pastures (247 ± 38 and $190 \pm 39 \mu\text{S cm}^{-1}$ for the dry and wet pasture, respectively), and highest in the three natural coastal wetlands with $1,418 \pm 104$, $8,049 \pm 276$ and $8,930 \pm 790 \mu\text{S cm}^{-1}$ for tidal freshwater wetland, saltmarsh and mangroves, respectively.

Soil bulk density was highest in sugarcane ($1.5 \pm 0.1 \text{ g cm}^{-3}$) and lowest in the freshwater tidal wetland ($0.6 \pm 0.1 \text{ g cm}^{-3}$). For all sites, %C was highest in the top 10 cm of the soil and decreased with depth, with highest values in the freshwater tidal wetland ($5.1 \pm 0.6\%$) and lowest in the saltmarsh ($1.2 \pm 0.1\%$). Soil %N ranged from 0.1 ± 0.0 to $0.4 \pm 0.1\%$ at all sites, except in the freshwater tidal wetland, where it reached values of $0.6 \pm 0.0\%$ in the top 10 cm (Table 2).

3.2 Greenhouse gas fluxes

Soil emissions for CO₂ were significantly different among sites and times of the year ($t = 155.09$, $n = 237$, $p < 0.001$; Fig. 2a). 205 The highest CO₂ emissions were measured during the wet-hot period in the dry pasture, where values reached $20.31 \pm 1.95 \text{ g m}^{-2} \text{ d}^{-1}$ while the lowest values were measured in the saltmarsh, the only site that acted as a sink of CO₂ with an uptake rate of $-0.59 \pm 0.15 \text{ g m}^{-2} \text{ d}^{-1}$. In the pastures, CO₂ emissions were twice as high when dry with cumulative annual emissions of $5,748 \pm 303 \text{ g m}^{-2} \text{ y}^{-1}$ compared to when wet, with $2,163 \pm 465 \text{ g m}^{-2} \text{ y}^{-1}$. For the coastal wetlands, cumulative annual CO₂ emissions were highest in freshwater tidal forests with $2,213 \pm 284 \text{ g m}^{-2} \text{ y}^{-1}$, followed by mangroves with $1,493 \pm 111 \text{ g m}^{-2} \text{ y}^{-1}$ and 210 lowest at the saltmarsh with uptake rates of $-264 \pm 29 \text{ g m}^{-2} \text{ y}^{-1}$.

For CH₄ fluxes, significant differences were observed among sites and seasons ($t = 182.33$, $n = 237$, $p < 0.001$). The differences between different sites were substantial, with wet pasture having significantly higher CH₄ emissions than any other site, with rates ~ 200 times higher throughout the measured period (Fig. 2b). For tidal coastal wetlands, emissions of CH₄ were 215 highest during the wet-hot season in all the sites except for the mangroves, which had similar emissions throughout the year (Fig. 2b). Overall, cumulative annual CH₄ emissions were $209 \pm 36 \text{ g m}^{-2} \text{ y}^{-1}$ for the wet pasture followed by mangroves (0.73

$\pm 0.13 \text{ g m}^{-2} \text{ y}^{-1}$), dry pasture ($0.15 \pm 0.03 \text{ g m}^{-2} \text{ y}^{-1}$), freshwater tidal forest ($0.14 \pm 0.03 \text{ g m}^{-2} \text{ y}^{-1}$), saltmarsh ($0.04 \pm 0.01 \text{ g m}^{-2} \text{ y}^{-1}$), and sugarcane ($-0.04 \pm 0.02 \text{ g m}^{-2} \text{ y}^{-1}$).

220 For N_2O fluxes, the highest emissions ($55 \pm 9 \text{ mg m}^{-2} \text{ d}^{-1}$) were measured in the dry pasture in the hot-wet season, followed by sugarcane ($20 \pm 3 \text{ mg m}^{-2} \text{ d}^{-1}$) during the hot-dry period, which coincides with the post-fertilisation period (Fig. 2c). Overall, dry pastures had the highest cumulative annual N_2O emissions ($7.99 \pm 2.26 \text{ mg m}^{-2} \text{ d}^{-1}$), followed by sugarcane ($2.37 \pm 0.68 \text{ mg m}^{-2} \text{ d}^{-1}$), wet pasture ($1.32 \pm 0.33 \text{ mg m}^{-2} \text{ d}^{-1}$), saltmarsh ($0.33 \pm 0.11 \text{ mg m}^{-2} \text{ d}^{-1}$), freshwater tidal forests ($0.04 \pm 0.0 \text{ mg m}^{-2} \text{ d}^{-1}$) and finally, mangroves ($0.02 \pm 0.04 \text{ mg m}^{-2} \text{ d}^{-1}$). However, these differences were only statistically significant

225 when considering the interaction between time of the year and site ($t=100.21, n=237, p < 0.001$).

We found that CH_4 fluxes did not significantly vary between the low and high tide within all coastal wetlands.

Contrarily, for saltmarsh, CO_2 was taken during the high tide ($1.12 \pm 0.24 \text{ g m}^{-2} \text{ d}^{-1}$) but emitted ($0.69 \pm 0.4 \text{ g m}^{-2} \text{ d}^{-1}$) during the low tide ($F_{1,28}=20.06, p < 0.001$). Finally, for N_2O , fluxes differed in all coastal wetlands, with higher uptakes in the high tide for mangroves ($F_{1,28}=38.28, p < 0.001; F_{1,28}=13.53, p = 0.001$) and higher release for saltmarsh ($F_{1,28}=38.31, p < 0.001$)

230 during low tide (Table S4). These results suggested that for CO_2 and N_2O fluxes, there was a probability of variation depending on the time of sampling.

The wet pasture had the highest total cumulative soil GHG emissions ($\text{CH}_4 + \text{N}_2\text{O}$) with $56,124 \text{ CO}_2\text{eq kg ha}^{-1} \text{ y}^{-1}$ followed by dry pasture $23,890 \text{ CO}_2\text{eq kg ha}^{-1} \text{ y}^{-1}$ and sugarcane $7,142 \text{ CO}_2\text{eq kg ha}^{-1} \text{ y}^{-1}$. While coastal wetlands had comparatively very lower total cumulative soil GHG emissions with $884, 235$ and $144 \text{ CO}_2\text{eq kg ha}^{-1} \text{ y}^{-1}$ for saltmarsh, 235 mangroves and freshwater tidal forests, respectively. Overall, the three coastal wetlands measured in this study had lower total cumulative GHG emissions at $1,263 \text{ CO}_2\text{eq kg ha}^{-1} \text{ yr}^{-1}$ compared to the alternate agricultural land uses, which emitted $87,156 \text{ CO}_2\text{eq kg ha}^{-1} \text{ yr}^{-1}$.


3.3 Greenhouse gas emissions and environmental factors

Overall, we found that not one single parameter measured in this could explain GHG fluxes for all sites except land-use. The 240 CO_2 emissions were not significantly correlated to bulk density ($R^2 = 0.026, p = 0.918, n = 18$), % WFPS ($R^2 = -0.003, p = 0.99, n = 18$), or soil temperature ($R^2 = 0.296, p = 0.233, n = 18$). Soil CH_4 emissions were neither correlated with bulk density ($R^2 = -0.096, p = 0.706, n = 18$), % WFPS ($R^2 = -0.224, p = 0.372, n = 18$) or soil temperature ($R^2 = 0.286, p = 0.25, n = 18$). Finally, no correlation was found between N_2O emissions and bulk density ($R^2 = -0.349, p = 0.156, n = 18$), % WFPS ($R^2 = -0.34, p = 0.168, n = 18$), or soil temperature ($R^2 = -0.241, p = 0.335, n = 18$). Full raw dataset of GHG fluxes are provided in 245 Table S1 and S4,

Table 2. Physicochemical characteristics for the soil of natural coastal wetlands, sugarcane and pastures (dry and ponded) for the top 30 cm of soil in tropical Australia. C= carbon, N = Nitrogen, EC = Electrical Conductivity. Values are mean \pm standard error ($n = 5$).

250

Site	Depth (cm)	Gravimetric moisture content (%)	pH	EC ($\mu\text{s cm}^{-1}$)	Bulk density (g cm^{-3})	%C	%N
Mangroves	0-10	41.7 \pm 1.1	5.9 \pm 0.1	12,550 \pm 524	1.14 \pm 0.05	2.3 \pm 0.1	0.18 \pm 0.01
	10-20	34.6 \pm 0.7	5.9 \pm 0.3	12,164 \pm 5,560	1.34 \pm 0.03	1.7 \pm 0.2	0.12 \pm 0.01
	20-30	31.3 \pm 0.6	6.2 \pm 0.1	5,560 \pm 365	1.95 \pm 0.12	0.9 \pm 0.1	0.07 \pm 0.01
	Mean	35.9 \pm 1.2	6.0 \pm 0.1	8,930 \pm 790	1.48 \pm 0.10	1.6 \pm 0.2	0.12 \pm 0.01
Saltmarsh	0-10	25.6 \pm 1.2	5.8 \pm 0.2	8,442 \pm 435	1.12 \pm 0.04	1.4 \pm 0.1	0.11 \pm 0.01
	10-20	26.6 \pm 0.3	5.8 \pm 0.1	8,666 \pm 437	1.47 \pm 0.05	1.3 \pm 0.1	0.12 \pm 0.01
	20-30	26.4 \pm 0.2	5.9 \pm 0.3	7,040 \pm 316	1.56 \pm 0.03	1.0 \pm 0.3	0.10 \pm 0.02
	Mean	26.2 \pm 0.4	5.8 \pm 0.1	8,049 \pm 276	1.38 \pm 0.06	1.2 \pm 0.1	0.11 \pm 0.01
Freshwater tidal forest	0-10	33.4 \pm 0.5	4.4 \pm 0.2	1,099 \pm 17	0.46 \pm 0.05	7.8 \pm 0.1	0.62 \pm 0.03
	10-20	24.9 \pm 0.6	4.2 \pm 0.0	1,272 \pm 164	0.71 \pm 0.02	5.4 \pm 0.0	0.46 \pm 0.04
	20-30	22.4 \pm 0.7	4.2 \pm 0.1	1,882 \pm 47	0.83 \pm 0.03	2.2 \pm 0.1	0.10 \pm 0.00
	Mean	26.9 \pm 1.3	4.3 \pm 0.1	1,418 \pm 104	0.59 \pm 0.05	5.1 \pm 0.6	0.39 \pm 0.06
Sugarcane	0-10	9.1 \pm 0.4	5.7 \pm 0.1	429 \pm 12	1.35 \pm 0.08	1.5 \pm 0.1	0.10 \pm 0.00
	10-20	12.1 \pm 0.6	5.3 \pm 0.3	365 \pm 11	1.46 \pm 0.06	1.5 \pm 0.1	0.12 \pm 0.01
	20-30	13.7 \pm 0.2	4.7 \pm 0.2	351 \pm 2	1.64 \pm 0.10	1.3 \pm 0.1	0.10 \pm 0.00
	Mean	11.7 \pm 0.6	5.2 \pm 0.2	382 \pm 11	1.48 \pm 0.05	1.4 \pm 0.1	0.11 \pm 0.00
Pasture	Dry	0-10	12.4 \pm 0.3	4.1 \pm 0.0	378 \pm 21	0.78 \pm 0.06	3.1 \pm 0.3
		10-20	13.6 \pm 0.1	4.4 \pm 0.1	279 \pm 60	1.21 \pm 0.14	1.6 \pm 0.4
		20-30	14.5 \pm 0.7	4.4 \pm 0.3	84 \pm 4	1.32 \pm 0.19	1.6 \pm 0.2
		Mean	13.5 \pm 0.3	4.3 \pm 0.1	247 \pm 38	1.10 \pm 0.10	2.1 \pm 0.3
	Wet	0-10	52.1 \pm 0.4	4.8 \pm 0.0	358 \pm 71	0.62 \pm 0.06	3.6 \pm 0.3
		10-20	47.7 \pm 0.4	4.9 \pm 0.1	117 \pm 11	1.30 \pm 0.02	1.7 \pm 0.1
		20-30	46.4 \pm 0.2	5.1 \pm 0.1	95 \pm 6	1.31 \pm 0.02	1.5 \pm 0.1
		Mean	48.7 \pm 0.7	4.9 \pm 0.0	190 \pm 39	1.07 \pm 0.09	2.3 \pm 0.3






Figure 2: Greenhouse gas fluxes ($\text{mg m}^{-2} \text{d}^{-1}$) of (a) CO₂ ($\text{mg m}^{-2} \text{d}^{-1}$), (b) CH₄ ($\text{mg m}^{-2} \text{d}^{-1}$) and (c) N₂O ($\text{mg m}^{-2} \text{d}^{-1}$) from soils of tropical coastal wetlands: mangroves, saltmarsh, freshwater (Fw) tidal forest, and two alternative land uses: sugarcane and pastures (wet and dry) during three periods of the year: dry-cold, dry-hot and wet-hot

4 Discussion

In this study, we found that the three coastal tropical wetlands measured in this study (mangroves, saltmarshes and freshwater tidal forests) had significantly lower GHG emissions compared to two alternative land uses common in tropical Australia (sugar cane and grazing pastures). Notably, we found that coastal wetlands had 200 times lower CH₄ emissions and seven times lower N₂O compared to wet pastures and sugarcane soils, respectively. While future studies should measure GHG from other wetlands, land uses, and within other tropical regions, these results support the idea that the management or conversion of unused agricultural land could be converted to coastal wetlands could result in significant GHG mitigation.

265

The variability of GHG fluxes was best explained by land use and wetland type; however, some trends with seasons were evident. For instance, CO₂ and N₂O emissions were lowest during the dry-cool periods. Reduced emissions at low temperatures are expected as the temperature is a main driver of any metabolic process, including respiration and nitrification-denitrification. Mangroves tend to have higher CO₂ emissions as temperature increases (Liu and Lai 2019), and terrestrial forests have significantly higher N₂O emissions during warm seasons (Schindlbacher et al, 2004). Emissions of CH₄ also tend to increase with temperature as the activity of soil methane-producing microbes (Ding et al, 2004) and the availability of carbon is higher in warmer conditions (Yvon-Durocher et al, 2011). However, as most of the studies on GHG fluxes, were conducted in temperate and subtropical locations where differences in temperature throughout the year are much larger than those in tropical regions. For tropical regions, increased GHG emissions are likely to be strongly affected by the “Birch effect”, which refers to short-term but a substantial increase of respiration from soils under the effect of precipitation during the early wet season (Fernandez-Bou et al, 2020).

270
275
280
285
290

The main factor associated with GHG fluxes was land use and type of wetland. Notably, coastal wetlands, even the freshwater tidal forests, had much lower emissions compared to the wet pastures. This large difference could be attributed to the presence of terminal electron acceptors in the soils (e.g. iron, sulphate, manganese) of the coastal wetlands, which could inhibit methanogenesis (Kögel-Knabner et al, 2010; Sahrawat, 2004). Sulphate reducing bacteria are also likely to outcompete methane-producing bacteria (methanogens) in the presence of high sulphate concentrations in tidal wetlands, resulting in low CH₄ production. Competition between methanogens and methanotrophs may result in a net balance of low CH₄ production despite freshwater conditions (Maietta et al. 2020). Additionally, microorganism living within the bark of *Melaleuca* trees can consume CH₄ (Jeffrey et al, 2021), so it is possible that similar bacteria within the soil could reduce CH₄ emissions. Interestingly, variability within CH₄ fluxes among sites was very high, despite them being very close to each other (Fig. 1b). These differences highlight the importance of land use in GHG fluxes, which are likely to significantly alter the microbial community composition and abundance, which can change rapidly over small spatial scales (Martiny et al, 2006; Drenovsky et al, 2009).

Our results are consistent with other studies, which have shown the importance of land use in GHG emissions. For instance, in a Mediterranean climate, drained agricultural land use types, pasture and corn, were larger CO₂ emitters compared to restored wetlands (Knox et al. 2015). Clearing of wetlands for agricultural development, such as the drainage of peatlands, results in very high CO₂ emissions (Nieveen et al, 2005; Veenendaal et al, 2007; Hirano et al, 2012), and restoration of these wetlands could decrease these emissions (Cameron et al, 2020). Additionally, some of the wetland types, such as marshes, were occasional sinks of CO₂ and CH₄, consistent with previous studies where intertidal wetlands sink of GHG at least under some conditions or during some times of the year (Knox et al, 2015; Maher et al, 2016).

300 The fluxes measured in the coastal wetlands of this study (1,191 to 10,970 mg m⁻² d⁻¹ for CO₂; 0.2 to 3.9 mg m⁻² d⁻¹ for CH₄; and -0.2 to 2.8 mg m⁻² d⁻¹ for N₂O) are within the range of those measured in other wetlands, worldwide. For CO₂, fluxes can range between -139 and 22,000 mg m⁻² d⁻¹ (Stadmark and Leonardson 2005; Morse et al. 2012), for CH₄, from -1 to 418 mg m⁻² d⁻¹ (Allen et al. 2007; Mitsch et al 2013; Cabezas et al. 2018), and for N₂O, from -0.3 to 3.9 mg m⁻² d⁻¹ (Hernandez and Mitsch 2006; Morse et al. 2012). Despite being in tropical regions, the fluxes from this study were not notably higher compared to wetlands in other climates. The fluxes measured in the coastal wetlands of this study, -1191 to 10,970 mg m⁻² d⁻¹ for CO₂, -0.3 to 3.9 mg m⁻² d⁻¹ for CH₄, and -0.2 to 2.8 mg m⁻² d⁻¹ for N₂O, were within the range of those measured in other subtropical/tropical wetlands, worldwide (Table 3). For CO₂, fluxes can range between 44 and 11,328 mg m⁻² d⁻¹, for CH₄, from 0.03 to 1255 mg m⁻² d⁻¹ and for N₂O, from 0.1 to 279 mg m⁻² d⁻¹ (Table 3). Despite being in tropical regions, GHG fluxes from this study were lower compared to other climates (Table 3). The general lower nitrogen pollution in Australia's soils and waterways compared to other countries may partially explain the lower emissions. However, the GHG flux measurements from this study did not account for the effects of vegetation, which can alter fluxes. For instance, some plant species of rice paddies (Timilsina et al., 2020) and *Miscanthus sinensis* (Lenhart et al., 2019) can increase N₂O emissions, and some tree species can facilitate CH₄ efflux from the soil (Pangala et al. 2013). Finally, changes in emissions between low and high tides were detected for CO₂ and N₂O. Thus, future studies that include vegetation and changes within tidal cycles will improve GHG flux estimates for coastal wetlands.

Formatted: Indent: Left: 0 cm, Space After: 6 pt, Adjust space between Latin and Asian text, Adjust space between Asian text and numbers

315

Table: 3. Comparison of GHG fluxes (mg m⁻² d⁻¹) from literature review with the rates reported in this study

Reference	Climate	Country	Ecosystem	CO ₂ fluxes	CH ₄ fluxes	N ₂ O fluxes
Allen et al., 2011	Subtropical	Australia	Mangroves estuary	-	1.5-51	-
Cabezas et al., 2018	Subtropical	USA	Mangroves estuary	-	0.3-2.2	-

Li and Mitsch, 2016	Subtropical	USA	Flooded brackish marsh	-	212 ± 51	-
Morse, Ardón and Bernhardt, 2012	Subtropical	USA	Forested wetlands	7224-11328	118-1255	46-279
Musenze et al., 2014	Subtropical	Australia	Mangroves estuary	-	5-448	0.1-3.4
Whiting and Chanton, 2001	Subtropical	USA	Typha marsh	409-477	189-264	-
Mitsch et al., 201	Tropical	South Africa	Seasonally flooded wetland	-	264±29	-
Krithika et al., 2008	Tropical	India	Mangroves	-	25-50	-
Kristensen et al., 2008	Tropical	Tanzania	Mangroves	44-3521	1.9-6.5	-
Biswas et al., 2007	Tropical	India	Mangroves estuary	-	0.03-2.16	-
Purvaja et al., 2004	Tropical	India	Mangroves estuary	-	10-85	-
Kreuzwieser et al., 2003	Tropical	Australia	Mangroves	-	0.6-11	-
Kiese and Butterbach-Bahl, 2002	Tropical	Australia	Tropical rain forest	2208-3288		1.9-3.2
Purvaja and Ramesh, 2000	Tropical	India	Mangroves	-	63-434	-
Sotomayor et al. 1994	Tropical	Puerto Rico	Mangroves	-	5-110	-

Barnes et al., 2006	Tropical	India	Mangroves	-	9-15	-
Melling et al. 2012	Tropical	Malaysia	Peat swamp forest	3384	21-29	
This study	Tropical	Australia	Freshwater tidal forest	1640-10970	0.16-0.59	-0.19-0.7
This study	Tropical	Australia	Saltmarsh	-594-(-1191)	-0.25-0.12	-0.22-2.76
This study	Tropical	Australia	Mangroves	2852-5669	0.44-3.95	0.04-0.16
Oertel et al., 2016	(Sub) Tropical	Global	Wetlands	-	-1.08-1169	-
Oertel et al., 2016	Temperate	Global	Wetlands	-	-1.49-1510	-
Oertel et al., 2016	Mediterranean	Global	Wetlands	-	-	-2.6-9.4
Al-Haj and Fulweiler, 2020	-	Global	Mangroves	-	-1.1-1169	-0.2-6.3
Al-Haj and Fulweiler, 2020	-	Global	Saltmarshes	6844-34983	0.38-3002	-7.39-28.52
Rosentreter et al., 2021	-	Global	Mangroves	4563-30800	-0.69-10.78	-1.69-4.65
Rosentreter et al., 2021	-	Global	Saltmarshes	3802-20914	107-168	4.96
IPCC 2013	Tropical	Global	Swamp forests		30.76-2149	

Note: Hyphen means no data was available; GHG fluxes as CO₂-C, CH₄-C and N₂O-N were multiplied by 3.66, 1.34 and 1.57 respectively to calculate CO₂, CH₄ and N₂O fluxes (National Greenhouse Accounts Factors, Australian Government Department of Industry, Science, Energy and Resources, 2020).

4.1 Management implications

Under the Paris Agreement, Australia has committed to reducing GHG emissions 26 - 28% below its 2005 levels by 2030.

With annual emissions of 153 million tonnes of carbon dioxide equivalent (Mt CO₂-eq y⁻¹), Queensland is a major GHG emitter

325 in Australia (~ 28.7% of the total in 2016; DES, 2016). Of these emissions, about 18.3 Mt CO₂-eq y⁻¹ (14%) are attributed to agriculture, while land-use change and forestry emit 12.1 Mt CO₂-eq y⁻¹ (DES, 2016). Production of CH₄ from ruminant animals, primarily cattle, contribute 82% of agriculture-related emissions (DES, 2016). Therefore, any GHG mitigation strategy from land-use change could be important for Australia to achieve its national goals.

330 This study supports the application of three management actions that could reduce GHG emissions. First, the conversion of ponded pastures to coastal wetlands is likely to reduce soil GHG emissions. Our results showed that wet pastures emit 56 ton CO₂-eq ha⁻¹ y⁻¹ of total GHG (CH₄ + N₂O) compared with 0.2 ton CO₂-eq ha⁻¹ y⁻¹, 0.1 ton CO₂-eq ha⁻¹ y⁻¹ and 0.9 ton CO₂-eq ha⁻¹ y⁻¹ from mangroves, freshwater tidal forest, and saltmarshes, respectively. This implies that about 55 ton CO₂-eq ha⁻¹ y⁻¹ emissions from the soils could be potentially avoided by converting wet pastures to coastal wetlands. The carbon 335 mitigation for GHG emissions from soil solely could provide ~ AUD 860 ha⁻¹ yr⁻¹ assuming a carbon value of AUD 15.37 per ton of CO₂-eq (Australian Government Clean Energy Regulator, 2018). This mitigation could be added up to the carbon sequestration through sediment accumulation and tree growth that results from wetland restoration. Legal enablers in Queensland are in place to manage unproductive agricultural land this way (Bell-James and Lovelock 2019), and could provide an alternative income source for farmers.

340

A second management option would be to reduce the time pastures are kept under water. Dry pastures produced significantly less CH₄ with ~0.005 kg ha⁻¹ d⁻¹ than wet pastures with 6 kg ha⁻¹ d⁻¹. For comparison, an average cow produces 141 g CH₄ d⁻¹ (McGinn et al, 2004), and our study area supported around 900 cattle over 2,500 ha throughout the year, 345 equivalent to 18519 kg ha⁻¹ y⁻¹ compared to 2 kg ha⁻¹ y⁻¹ and 2090 kg ha⁻¹ y⁻¹ CH₄ from dry and wet pasture respectively. This implies that nearly 99% of the CH₄ emissions came from wet pastures, while dry pasture and grazing cattle had a low share in total CH₄ emissions. Therefore, land use management of wet pastures which are used to feed grazing cattle in Queensland may be a significant opportunity to reduce agriculture-related CH₄ emissions. Future studies should increase the number of sites of 350 ponded pastures to account for variability in hydrology, fertilisation, and on-farm cattle density cattle use. However, the very high difference (2-3 orders of magnitude) between dry and ponded pastures provides confidence that pasture management could provide significant GHG mitigation throughout the year.

Finally, fertiliser management in sugarcane could reduce N_2O emissions. Higher N_2O emissions of $17.63 \text{ mg m}^{-2} \text{ d}^{-1}$ were measured in sugarcane following fertilisation during the dry-hot season. Comparatively, natural wetlands had low N_2O emissions (0.16 to $2.79 \text{ mg m}^{-2} \text{ d}^{-1}$); even the saltmarsh was an occasional sink. Thus, improved management of fertiliser applications could result in GHG emission mitigation. Some activities include split application of nitrogen fertiliser in combination with low irrigation, reduction in fertiliser application rates, the substitution of nitrate-based fertiliser for urea (Rashti et al, 2015), removing mulch layer before fertiliser application (Pinheiro et al, 2019; Xu et al, 2019 Zaehle and Dalmonech, 2011) or conversion of unproductive sugarcane to coastal wetlands.

360 5 Conclusion

The GHG emissions from three coastal wetlands in tropical Australia (mangroves, saltmarsh and freshwater tidal forests) were consistently lower than those from two common agricultural land use of the region (sugarcane and pastures) throughout three climatic conditions (dry-cool, dry-hot and wet-hot). Ponded pastures, which emitted 200 times more CH_4 , and sugarcane emitted seven times more than any natural coastal wetland. If these high emissions are persistent in other locations and within 365 other tropical regions, conversion of pastures and sugarcane to similar coastal wetlands could provide significant GHG mitigation. As nations try to reach their emission reduction targets, projects aimed at converting or restoring coastal wetland can financially benefit farmers and provide additional co-benefits derived from coastal wetland restoration.

370

Author contribution

Iram, N. and M.F. Adame designed the project, Iram, N, B. Shahrabi Farahani and E. Kavehei carried out experiments, Iram, N., E. Kavehei and M.F. Adame analysed the data. Iram, N and M.F. Adame prepared the manuscript with contributions from 375 D.T. Maher, S.E. Bunn, and M. Rezaei Rashti.

Competing interests

The authors declare that they have no conflict of interest.

380 Acknowledgements

We acknowledge the Traditional Owners of the land in which the field study was conducted, especially the Nywaigi people from Mungalla Station, where this study was conducted. We are also thankful to Sam and Santo Lamari for allowing us to work on their property and for sharing their local knowledge. We are thankful to Charles Cadier and Julieta Gamboa for their contribution to the fieldwork. This project was financially supported by an Advance Queensland Industry Research Fellowship

385 to MF Adame. We are very thankful to anonymous referees for their valuable feedback which helped to improve the quality
of the article.

References

Al-Haj, A. N., and Fulweiler, R. W.: A synthesis of methane emissions from shallow vegetated coastal ecosystems. *Global Change Biology*, 26(5), 2988– 3005. <https://doi.org/10.1111/gcb.15046>, 2020.

Allen, D., Dalal, R. C., Rennenberg, H. and Schmidt, S.: Seasonal variation in nitrous oxide and methane emissions from subtropical estuary and coastal mangrove sediments, Australia, *Plant Biol.*, 13(1), 126–133, doi:10.1111/j.1438-8677.2010.00331.x, 2011.

Angle, J. C., Morin, T. H., Soden, L. M., Narrowe, A. B., Smith, G. J., Borton, M. A., Rey-Sanchez, C., Daly, R. A., Mirfenderesgi, G., Hoyt, D. W., Riley, W. J., Miller, C. S., Bohrer, G. and Wrighton, K. C.: Methanogenesis in oxygenated soils is a substantial fraction of wetland methane emissions, , doi:10.1038/s41467-017-01753-4, 2017.

Barnes, J., Ramesh, R., Purvaja, R., Rajkumar, A. N., Kumar, B. S., Krishika, K., Ravichandran, K., Uher, G. and Upstill-Goddard, R.: Tidal dynamics and rainfall control N2O and CH4 emissions from a pristine mangrove creek, *Geophys. Res. Lett.*, 33(15), doi:10.1029/2006GL026829, 2006.

Bauza, J. F., Morell, J. M. and Corredor, J. E.: Biogeochemistry of nitrous oxide production in the red mangrove (Rhizophora mangle) forest sediments, *Estuar. Coast. Shelf Sci.*, 55(5), 697–704, doi:10.1006/ecss.2001.0913, 2002.

Bell-James, J. and Lovelock, C. E.: Legal barriers and enablers for reintroducing tides: An Australian case study in reconvert ing ponded pasture for climate change mitigation, *Land use policy*, 88, 104192, doi:10.1016/J.LANDUSEPOL.2019.104192, 2019.

Biswas, H., Mukhopadhyay, S. K., Sen, S. and Jana, T. K.: Spatial and temporal patterns of methane dynamics in the tropical mangrove dominated estuary, NE coast of Bay of Bengal, India, *J. Mar. Syst.*, 68(1–2), 55–64, doi:10.1016/j.jmarsys.2006.11.001, 2007.

Cabezas, A., Mitsch, W. J., MacDonnell, C., Zhang, L., Bydałek, F. and Lasso, A.: Methane emissions from mangrove soils in hydrologically disturbed and reference mangrove tidal creeks in southwest Florida, *Ecol. Eng.*, 114, 57–65, doi:10.1016/j.ecoleng.2017.08.041, 2018.

Cameron, C., Hutley, L. B., Munksgaard, N. C., Phan, S., Aung, T., Thinn, T., Aye, W. M. and Lovelock, C. E.: Impact of an extreme monsoon on CO2 and CH4 fluxes from mangrove soils of the Ayeyarwady Delta, Myanmar, *Sci. Total Environ.*, 760, 143422, doi:10.1016/J.SCITOTENV.2020.143422, 2021.

Chen, G. C., Tam, N. F. Y. and Ye, Y.: Spatial and seasonal variations of atmospheric N2O and CO2 fluxes from a subtropical mangrove swamp and their relationships with soil characteristics, *Soil Biol. Biochem.*, 48, 175–181, doi:10.1016/j.soilbio.2012.01.029, 2012.

Conrad, R.: The global methane cycle: recent advances in understanding the microbial processes involved, *Env. Microbiol. Rep.*, 1, 285–292, <https://doi.org/10.1111/j.1758-2229.2009.00038.x>, 2009.

Deemer, B. R., Harrison, J. A., Li, S., Beaulieu, J. J., Delsontro, T., Barros, N., Bezerra-Neto, J. F., Powers, S. M., Santos, M. A. Dos and Vonk, J. A.: Greenhouse Gas Emissions from Reservoir Water Surfaces: A New Global Synthesis, *Bioscience*,

66(11), doi:10.1093/biosci/biw117, 2016.

Department of Environment and Science, Queensland, 2016. Total greenhouse gas emissions. <https://www.stateoftheenvironment.des.qld.gov.au/pollution/greenhouse-gas-emissions/total-annual-greenhouse-gas-emissions>. Last assessed, 25/02/2020.

425 Department of Science and Environment. 2020. WetlandInfo, 2020. Herbert River drainage sub-basin — facts and maps, <https://wetlandinfo.des.qld.gov.au/wetlands/facts-maps/sub-basin-herbert-river/> viewed 20 March 2020.

Ding, W. X., Cai, Z. C. and Tsuruta, H.: Cultivation, nitrogen fertilization, and set-aside effects on methane uptake in a drained marsh soil in Northeast China, *Glob. Chang. Biol.*, 10(10), 1801–1809, doi:10.1111/j.1365-2486.2004.00843.x, 2004.

430 Drenovsky, R. E., Steenwerth, K. L., Jackson, L. E. and Scow, K. M.: Land use and climatic factors structure regional patterns in soil microbial communities, *Glob. Ecol. Biogeogr.*, 19(1), 27–39, doi:10.1111/j.1466-8238.2009.00486.x, 2010.

Drexler, J. Z., De Fontaine, C. S. and Deverel, S. J.: The legacy of wetland drainage on the remaining peat in the Sacramento-San Joaquin delta, California, USA., 2009.

Duarte, C. M., Losada, I. J., Hendriks, I. E., Mazarrasa, I. and Marbà, N.: The role of coastal plant communities for climate change mitigation and adaptation, *Nat. Clim. Chang.*, 3(11), 961–968, doi:10.1038/nclimate1970, 2013.

435 Fernandez-Bou, A. S., Dierick, D., Allen, M. F. and Harmon, T. C.: Precipitation-drainage cycles lead to hot moments in soil carbon dioxide dynamics in a Neotropical wet forest, *Glob. Chang. Biol.*, 26(9), 5303–5319, doi:10.1111/gcb.15194, 2020.

Griggs, P. D.: Too much water: drainage schemes and landscape change in the sugar-producing areas of Queensland, 1920–90, *Aust. Geogr.*, 49(1), 81–105, doi:10.1080/00049182.2017.1336965, 2018.

440 Grinham, A., Albert, S., Deering, N., Dunbabin, M., Bastviken, D., Sherman, B., Lovelock, C. E. and Evans, C. D.: The importance of small artificial water bodies as sources of methane emissions in Queensland, Australia, *Hydrol. Earth Syst. Sci.*, 22(10), 5281–5298, doi:10.5194/hess-22-5281-2018, 2018.

Hirano, T., Segah, H., Kusin, K., Limin, S., Takahashi, H. and Osaki, M.: Effects of disturbances on the carbon balance of tropical peat swamp forests, *Glob. Chang. Biol.*, 18(11), 3410–3422, doi:10.1111/j.1365-2486.2012.02793.x, 2012.

445 Hooijer, A., Page, S., Jauhainen, J., Lee, W. A., Lu, X. X., Idris, A. and Anshari, G.: Subsidence and carbon loss in drained tropical peatlands, *Biogeosciences*, 9(3), 1053–1071, doi:10.5194/bg-9-1053-2012, 2012.

Hutchinson, G. L. and Mosier, A. R.: Improved Soil Cover Method for Field Measurement of Nitrous Oxide Fluxes, *Soil Sci. Soc. Am. J.*, 45(2), 311–316, doi:10.2136/sssaj1981.03615995004500020017x, 1981.

450 IPCC 2013. Supplement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories: Wetlands ed T Hiraishi, T Krug, K Tanabe, N Srivastava, J Baasansuren, M Fukuda and T G Troxler (Geneva: Intergovernmental Panel on Climate Change).

Jeffrey, L. C., Maher, D. T., Chiri, E., Leung, P. M., Nauer, P. A., Arndt, S. K., Tait, D. R., Greening, C. and Johnston, S. G.: Bark-dwelling methanotrophic bacteria decrease methane emissions from trees, , doi:10.1038/s41467-021-22333-7, 2021.

Johnson, A. K. L., Ebert, S. P. and Murray, A. E.: Distribution of coastal freshwater wetlands and riparian forests in the Herbert River catchment and implications for management of catchments adjacent the Great Barrier Reef Marine Park, *Environ.*

455 Conserv., 26(3), 229–235, doi:10.1017/S037689299000314, 1999.

Kauffman, J. B., Adame, M. F., Arifanti, V. B., Schile-Beers, L. M., Bernardino, A. F., Bhomia, R. K., Donato, D. C., Feller, I. C., Ferreira, T. O., Jesus Garcia, M. del C., MacKenzie, R. A., Megonigal, J. P., Murdiyarso, D., Simpson, L. and Hernández Trejo, H.: Total ecosystem carbon stocks of mangroves across broad global environmental and physical gradients, *Ecol. Monogr.*, 90(2), doi:10.1002/ecm.1405, 2020.

460 Kavehei, E., Iram, N., Rezaei Rashti, M., Jenkins, G. A., Lemckert, C. and Adame, M. F.: Greenhouse gas emissions from stormwater bioretention basins, *Ecol. Eng.*, 159, doi:10.1016/j.ecoleng.2020.106120, 2021.

Kettler, T. A., Doran, J. W. and Gilbert, T. L.: Simplified Method for Soil Particle-Size Determination to Accompany Soil-Quality Analyses, *Soil Sci. Soc. Am. J.*, 65(3), 849–852, doi:10.2136/sssaj2001.653849x, 2001.

Kiese, R. and Butterbach-Bahl, K.: N 2 O and CO 2 emissions from three different tropical forest sites in the wet tropics of 465 Queensland, Australia. [online] Available from: www.elsevier.com/locate/soilbio, 2002.

Kirschke, S., Bousquet, P., Ciais, P., Saunois, M., Canadell, J. G., Dlugokencky, E. J., Bergamaschi, P., Bergmann, D., Blake, D. R., Bruhwiler, L., Cameron-Smith, P., Castaldi, S., Chevallier, F., Feng, L., Fraser, A., Heimann, M., Hodson, E. L., Houweling, S., Josse, B., Fraser, P. J., Krummel, P. B., Lamarque, J. F., Langenfelds, R. L., Le Quéré, C., Naik, V., O'doherty, S., Palmer, P. I., Pison, I., Plummer, D., Poulter, B., Prinn, R. G., Rigby, M., Ringeval, B., Santini, M., Schmidt, M., Shindell, 470 D. T., Simpson, I. J., Spahni, R., Steele, L. P., Strode, S. A., Sudo, K., Szopa, S., Van Der Werf, G. R., Voulgarakis, A., Van Weele, M., Weiss, R. F., Williams, J. E. and Zeng, G.: Three decades of global methane sources and sinks, *Nat. Geosci.*, 6(10), 813–823, doi:10.1038/ngeo1955, 2013.

Knox, S. H., Sturtevant, C., Matthes, J. H., Koteen, L., Verfaillie, J. and Baldocchi, D.: Agricultural peatland restoration: Effects of land-use change on greenhouse gas (CO2 and CH4) fluxes in the Sacramento-San Joaquin Delta, *Glob. Chang. Biol.*, 475 21(2), 750–765, doi:10.1111/gcb.12745, 2015.

Kögel-Knabner, I., Amelung, W., Cao, Z., Fiedler, S., Frenzel, P., Jahn, R., Kalbitz, K., Kölbl, A. and Schloter, M.: Biogeochemistry of paddy soils, *Geoderma*, 157(1–2), 1–14, doi:10.1016/J.GEODERMA.2010.03.009, 2010.

Kreuzwieser, J., Buchholz, J. and Rennenberg, H.: Emission of Methane and Nitrous Oxide by Australian Mangrove Ecosystems, *Plant Biol.*, 5(4), 423–431, doi:10.1055/s-2003-42712, 2003.

480 Kristensen, E., Flindt, M. R., Uломи, S., Borges, A. V., Abril, G. and Bouillon, S.: Emission of CO2 and CH4 to the atmosphere by sediments and open waters in two Tanzanian mangrove forests, *Mar. Ecol. Prog. Ser.*, 370, 53–67, doi:10.3354/meps07642, 2008.

Krithika, K., Purvaja, R. and Ramesh, R.: Current Science Association Fluxes of methane and nitrous oxide from an Indian mangrove., 2008.

485 Kroeger, K. D., Crooks, S., Moseman-Valtierra, S. and Tang, J.: Restoring tides to reduce methane emissions in impounded wetlands: A new and potent Blue Carbon climate change intervention, *Sci. Rep.*, 7(1), doi:10.1038/s41598-017-12138-4, 2017.

Lehner, B. and Döll, P.: Development and validation of a global database of lakes, reservoirs and wetlands, *J. Hydrol.*, 296(1–4), 1–22, doi:10.1016/j.jhydrol.2004.03.028, 2004.

Lenhart, K., Behrendt, T., Greiner, S., Steinkamp, J., Well, R., Giesemann, A. and Keppler, F.: Nitrous oxide effluxes from
490 plants as a potentially important source to the atmosphere, *New Phytol.*, 221(3), 1398–1408, doi:10.1111/nph.15455, 2019.

Li, X. and Mitsch, W. J.: Methane emissions from created and restored freshwater and brackish marshes in southwest Florida, USA, *Ecol. Eng.*, 91, 529–536, doi:10.1016/j.ecoleng.2016.01.001, 2016.

Liu, J. and Lai, D. Y. F.: Subtropical mangrove wetland is a stronger carbon dioxide sink in the dry than wet seasons, *Agric. For. Meteorol.*, 278, 107644, doi:10.1016/J.AGRFORMAT.2019.107644, 2019.

495 Maher, D. T., Sippo, J. Z., Tait, D. R., Holloway, C. and Santos, I. R.: Pristine mangrove creek waters are a sink of nitrous oxide OPEN, *Nat. Publ. Gr.*, doi:10.1038/srep25701, 2016.

Maietta, C. E., Hondula, K. L., Jones, C. N. and Palmer, M. A.: Hydrological Conditions Influence Soil and Methane-Cycling Microbial Populations in Seasonally Saturated Wetlands, *Front. Environ. Sci.*, 8, doi:10.3389/fenvs.2020.593942, 2020.

Martiny, J. B. H., Bohannan, B. J. M., Brown, J. H., Colwell, R. K., Fuhrman, J. A., Green, J. L., Horner-Devine, M. C., Kane,
500 M., Krumins, J. A., Kuske, C. R., Morin, P. J., Naeem, S., Øvreås, L., Reysenbach, A. L., Smith, V. H. and Staley, J. T.: Microbial biogeography: Putting microorganisms on the map, *Nat. Rev. Microbiol.*, 4(2), 102–112, doi:10.1038/nrmicro1341, 2006.

McGinn, S. M., Turner, D., Tomkins, N., Charmley, E., Bishop-Hurley, G. and Chen, D.: Methane Emissions from Grazing Cattle Using Point-Source Dispersion, *J. Environ. Qual.*, 40(1), 22–27, doi:10.2134/jeq2010.0239, 2011.

505 Melling, L., Goh, K. J., Kloni, A., & Hatano, R. (2012). Is water table the most important factor influencing soil C flux in tropical peatlands. In *Peatlands in Balance, Proceedings of the 14th International Peat Congress*.

Mitsch, W. J., Bernal, B., Nahlik, A. M., Mander, Ü., Zhang, L., Anderson, C. J., Jørgensen, S. E. and Brix, H.: Wetlands, carbon, and climate change, *Landsc. Ecol.*, 28(4), 583–597, doi:10.1007/s10980-012-9758-8, 2013.

Morse, J. L., Ardón, M., and Bernhardt, E. S.: Greenhouse gas fluxes in southeastern U . S . coastal plain wetlands under
510 contrasting land uses, *Ecol. Appl.*, 22(1), 264–280, 2012.

Musenze, R. S., Werner, U., Grinham, A., Udy, J. and Yuan, Z.: Methane and nitrous oxide emissions from a subtropical estuary (the Brisbane River estuary, Australia), *Sci. Total Environ.*, 472, 719–729, doi:10.1016/j.scitotenv.2013.11.085, 2014.

National Greenhouse Accounts Factors, Australian Government Department of Industry, Science, Energy and Resources. 2020. Retrieved from (<https://www.industry.gov.au/sites/default/files/2020-10/national-greenhouse-accounts-factors-2020.pdf>) last access on 09/08/2021.

515 Neubauer, S. C.: Global Warming Potential Is Not an Ecosystem Property, *Ecosystems*, doi:10.1007/s10021-021-0063, n.d.

Nieveen, J. P., Campell, D. I., Schipper, L. A. and Blair, I. J.: Carbon exchange of grazed pasture on drained peat soil, *Glob. Chang. Biol.*, 11(4), 607–618, doi:10.1111/j.1365-2486.2005.00929.x, 2005.

Oertel, C., Matschullat, J., Zurba, K., Zimmermann, F. and Erasmi, S.: Greenhouse gas emissions from soils—A review,
520 *Chemie der Erde*, 76(3), 327–352, doi:10.1016/j.chemer.2016.04.002, 2016.

Ollivier, Q. R., Maher, D. T., Pitfield, C. and Macreadie, P. I.: Punching above their weight: Large release of greenhouse gases from small agricultural dams, *Glob. Chang. Biol.*, 25(2), 721–732, doi:10.1111/gcb.14477, 2019.

Pangala, S. R., Moore, S., Hornibrook, E. R. C. and Gauci, V.: Trees are major conduits for methane egress from tropical

forested wetlands, *New Phytol.*, 197(2), 524–531, doi:10.1111/nph.12031, 2013.

525 Pinheiro, P. L., Recous, S., Dietrich, G., Weiler, D. A., Schu, A. L., Bazzo, H. L. S. and Giacomini, S. J.: N₂O emission increases with mulch mass in a fertilized sugarcane cropping system, *Biol. Fertil. Soils*, 55(5), 511–523, doi:10.1007/s00374-019-01366-7, 2019.

Purvaja, R. and Ramesh, R.: Human impacts on methane emission from mangrove ecosystems in India., n.d.

Purvaja, R., Ramesh, R. and Frenzel, P.: Plant-mediated methane emission from an Indian mangrove, *Glob. Chang. Biol.*, 530 10(11), 1825–1834, doi:10.1111/j.1365-2486.2004.00834.x, 2004.

Reeves, S., Wang, W., Salter, B. and Halpin, N.: Quantifying nitrous oxide emissions from sugarcane cropping systems: Optimum sampling time and frequency, *Atmos. Environ.*, 136, 123–133, doi:10.1016/j.atmosenv.2016.04.008, 2016.

Rezaei Rashti, M., Wang, W. J., Harper, S. M., Moody, P. W., Chen, C. R., Ghadiri, H. and Reeves, S. H.: Strategies to mitigate greenhouse gas emissions in intensively managed vegetable cropping systems in subtropical Australia, *Soil Res.*, 53(5), 475–535 484, doi:10.1071/SR14355, 2015.

Rezaei Rashti, M., Wang, W. J., Reeves, S. H., Harper, S. M., Moody, P. W. and Chen, C. R.: Linking chemical and biochemical composition of plant materials to their effects on N₂O emissions from a vegetable soil, *Soil Biol. Biochem.*, 103, 502–511, doi:10.1016/j.soilbio.2016.09.019, 2016.

Rosentreter, J. A., Al-Haj, A. N., Fulweiler, R. W. and Williamson, P.: Methane and Nitrous Oxide Emissions Complicate 540 Coastal Blue Carbon Assessments, *Global Biogeochem. Cycles*, 35(2), doi:10.1029/2020GB006858, 2021.

Sahrawat, K. L.: Terminal electron acceptors for controlling methane emissions from submerged rice soils, *Commun. Soil Sci. Plant Anal.*, 35(9–10), 1401–1413, doi:10.1081/CSS-120037554, 2004.

Saunois, M., Stavert, A. R., Poulter, B., Bousquet, P., Canadell, J. G., Jackson, R. B., Raymond, P. A., Dlugokencky, E. J., Houweling, S., Patra, P. K., Ciais, P., Arora, V. K., Bastviken, D., Bergamaschi, P., Blake, D. R., Brailsford, G., Bruhwiler, 545 Carlson, K. M., Carroll, M., Castaldi, S., Chandra, N., Crevoisier, C., Crill, P. M., Covey, K., Curry, C. L., Etiope, G., Frankenberg, C., Gedney, N., Hegglin, M. I., Höglund-Isaksson, L., Hugelius, G., Ishizawa, M., Ito, A., Janssens-Maenhout, G., Jensen, K. M., Joos, F., Kleinen, T., Krummel, P. B., Langenfelds, R. L., Laruelle, G. G., Liu, L., Machida, T., Maksyutov, S., McDonald, K. C., McNorton, J., Miller, P. A., Melton, J. R., Morino, I., Müller, J., Murguia-Flores, F., Naik, V., Niwa, Y., Noce, S., O'Doherty, S., Parker, R. J., Peng, C., Peng, S., Peters, G. P., Prigent, C., Prinn, R., Ramonet, M., Regnier, P., Riley, 550 W. J., Rosentreter, J. A., Segers, A., Simpson, I. J., Shi, H., Smith, S. J., Steele, L. P., Thornton, B. F., Tian, H., Tohjima, Y., Tubiello, F. N., Tsuruta, A., Viovy, N., Voulgarakis, A., Weber, T. S., van Weele, M., van der Werf, G. R., Weiss, R. F., Worthy, D., Wunch, D., Yin, Y., Yoshida, Y., Zhang, W., Zhang, Z., Zhao, Y., Zheng, B., Zhu, Q., Zhu, Q. and Zhuang, Q.: The Global Methane Budget 2000–2017, *Earth Syst. Sci. Data*, 12(3), 1561–1623, doi:10.5194/essd-12-1561-2020, 2020.

Schindlbacher, A., Zechmeister-Boltenstern, S. and Butterbach-Bahl, K.: Effects of soil moisture and temperature on NO, 555 NO₂ and N₂O emissions from European forest soils, *J. Geophys. Res.*, 109, 17302, doi:10.1029/2004JD004590, 2004.

Shaaban, M., Peng, Q. an, Hu, R., Wu, Y., Lin, S. and Zhao, J.: Dolomite application to acidic soils: a promising option for

mitigating N₂O emissions, *Environ. Sci. Pollut. Res.*, 22(24), 19961–19970, doi:10.1007/s11356-015-5238-4, 2015.

Solomon S.: Climate Change: The Physical science Basis, *Clim. Chang. Phys. Sci. Basis* [online] Available from: <http://ci.nii.ac.jp/naid/20001628955/en/> (Accessed 13 August 2021), 2007.

560 Sotomayor, D., Corredor, J.E., Morell, J.M., 1994. Methane flux from mangrove sediments along the southwestern coast of Puerto Rico. *Estuaries* 17, 140–147.

Timilsina, A., Bizimana, F., Pandey, B., Kailash, R., Yadav, P., Dong, W. and Hu, C.: plants Nitrous Oxide Emissions from Paddies: Understanding the Role of Rice Plants, , doi:10.3390/plants9020180, 2018.

565 Ussiri, D., & Lal, R. Soil emission of nitrous oxide and its mitigation. Springer Science & Business Media.Germany, 378, ISBN 978-94-007-5364-8, 2012.

Whalen, S.: Biogeochemistry of methane exchange between natural wetlands and the atmosphere, *Environ. Eng. Sci.*, 22, 73–94, <https://doi.org/10.1089/ees.2005.22.73>, 2005.

570 Veenendaal, E. M., Kolle, O., Leffelaar, P. A., Schrier-Uijl, A. P., Van Huissteden, J., Walssem, J. Van, Möller, F. and Berendse, F.: CO₂ exchange and carbon balance in two grassland sites on eutrophic drained peat soils. [online] Available from: www.biogeosciences.net/4/1027/2007/, 2007.

Whiting, G. J. and Chanton, J. P.: Greenhouse carbon balance of wetlands: Methane emission versus carbon sequestration, *Tellus, Ser. B Chem. Phys. Meteorol.*, 53(5), 521–528, doi:10.3402/tellusb.v53i5.16628, 2001.

Xu, C., Han, X., Ru, S., Cardenas, L., Rees, R. M., Wu, D., Wu, W. and Meng, F.: Crop straw incorporation interacts with N 575 fertilizer on N₂O emissions in an intensively cropped farmland, *Geoderma*, 341, 129–137, doi:10.1016/J.GEODERMA.2019.01.014, 2019.

Yvon-Durocher, G., Montoya, J. M., Woodward, G., Jones, J. I. and Trimmer, M.: Warming increases the proportion of primary production emitted as methane from freshwater mesocosms, *Glob. Chang. Biol.*, 17(2), 1225–1234, doi:10.1111/j.1365-2486.2010.02289.x, 2011.

580 Zaehle, S. and Dalmonech, D.: Carbon-nitrogen interactions on land at global scales: Current understanding in modelling climate biosphere feedbacks, *Curr. Opin. Environ. Sustain.*, 3(5), 311–320, doi:10.1016/j.cosust.2011.08.008, 2011.