

1 Supporting information for

2 **Global modelling of soil carbonyl sulfide exchanges**

3 Camille Abadie¹, Fabienne Maignan¹, Marine Remaud¹, Jérôme Ogée², J. Elliott Campbell³,
4 Mary E. Whelan⁴, Florian Kitz⁵, Felix M. Spielmann⁵, Georg Wohlfahrt⁵, Richard Wehr⁶, Wu
5 Sun⁷, Nina Raoult¹, Ulli Seibt⁸, Didier Hauglustaine¹, Sinikka T. Lennartz^{10,11}, Sauveur
6 Belviso¹, David Montagne¹² and Philippe Peylin¹.

7 ¹Laboratoire des Sciences du Climat et de l'Environnement, LSCE/IPSL, CEA-CNRS-UVSQ, Université Paris-
8 Saclay, Gif-sur-Yvette, France

9 ²INRA, UMR 1391 ISPA, 33140 Villenave d'Ornon, France

10 ³Sierra Nevada Research Institute, University of California, Merced, California 95343, USA

11 ⁴Department of Environmental Sciences, Rutgers University, New Brunswick, NJ 08901, USA

12 ⁵Department of Ecology, University of Innsbruck, Innsbruck, 6020, Austria

13 ⁶Center for Atmospheric and Environmental Chemistry, Aerodyne Research, Inc., Billerica, Massachusetts, 01821,
14 USA.

15 ⁷Department of Global Ecology, Carnegie Institution for Science, Stanford, CA 94305, USA

16 ⁸Department of Atmospheric & Oceanic Sciences, University of California Los Angeles, California 90095, USA

17 ¹⁰Institute of Chemistry and Biology of the Marine Environment, University of Oldenburg, 26129 Oldenburg,
18 Germany

19 ¹¹Department of Earth, Atmospheric and Planetary Sciences, Massachusetts Institute of Technology, Cambridge,
20 02139, MA, USA

21 ¹²AgroParisTech, INRAE, Université Paris-Saclay, UMR ECOSYS, 78850 Thiverval-Grignon, France

22

23 **Table S1. Soil textures for the USDA texture classification.**

USDA texture classes
1- Sand
2- Loamy sand
3- Sandy loam
4- Silt loam
5- Silt
6- Loam
7- Sandy clay loam
8- Silty clay loam
9- Clay loam
10- Sandy clay
11- Silty clay
12- Clay

24

25

26 **Table S2. USDA textures initially assigned in ORCHIDEE and the substituted textures from the observations at the**
27 **studied sites. Textures are in bold when the imposed texture differs from the one initially assigned in ORCHIDEE.**

	Initial	Substituted
ES-LMA	3- Sandy loam	3- Sandy loam
DK-SOR	3- Sandy loam	3- Sandy loam
IT-CRO	3- Sandy loam	4- Silt loam

AT-NEU	3- Sandy loam	3- Sandy loam
ET-JA	3- Sandy loam	3- Sandy loam
FI-HYY	2- Loamy sand	3- Sandy loam
US-HA	1- Sand	3- Sandy loam

28

29

30 **Table S3: First-order parameters.**

Parameter name in ORCHIDEE	Parameter name in the model description	Description (unit)	Specificity	Range
Empirical model				
ksoil	k_{soil}	Proportionality factor for soil COS fluxes (pmol COS μmol^{-1} CO ₂)	(-)	± 1.08 pmol COS μmol^{-1} CO ₂
Mechanistic model				
FCA	f_{CA}	CA enhancement factor (unitless)	PFT-dependent	See Meredith et al., (2019) Table 1
α	α	COS production parameter (unitless)	PFT-dependent	See Text S1 and Table S5
β	β	COS production parameter ($^{\circ}\text{C}^{-1}$)	PFT-dependent	See Text S1 and Table S5

31

32

33

34 **Table S4: Second-order parameters.**

Parameter name in ORCHIDEE	Description (unit)	Specificity	Range
Photosynthesis parameters			
Vcmax25	Maximum rate of Rubisco activity-limited carboxylation at 25°C ($\mu\text{mol m}^{-2} \text{s}^{-1}$)	PFT-dependent	$\pm 45\%$ (Mahmud et al., 2021)
Zroot	Root profile in empirical plant water stress function calculation (m)	PFT-dependent	See Mahmud et al. (2021)
Tmin	Minimum photosynthesis temperature ($^{\circ}\text{C}$)	PFT-dependent	Vegetated PFTs: -9, 1 (Mahmud et al., 2021)
Tmax	Maximum photosynthesis temperature ($^{\circ}\text{C}$)	PFT-dependent	Vegetated PFTs: 50, 60 (Mahmud et al., 2021)
Conductance parameters			
g0	Residual stomatal conductance when irradiance approaches zero ($\text{mol m}^{-2} \text{s}^{-1} \text{bar}^{-1}$)	PFT-dependent (C ₃ or C ₄ plant types)	C ₃ plants: 0.00565, 0.00685 C ₄ plants: 0.01675, 0.02075 (Mahmud et al., 2021)
Phenology parameters			
SLA	Specific leaf area ($\text{m}^2 \text{gC}^{-1}$)	PFT-dependent	See Mahmud et al. (2021)
Soil hydrology parameters			

n	Van Genuchten water retention curve coefficient n (unitless)	Soil texture-dependent	±40% (Dantec-Nédélec et al., 2016)
a	Van Genuchten water retention curve coefficient a (unitless)	Soil texture-dependent	±50% (Dantec-Nédélec et al., 2016)
Ks	Hydraulic conductivity at saturation	Soil texture-dependent	±40%
θWP	Volumetric water content at wilting point (%)	Soil texture-dependent	±20% (Dantec-Nédélec et al., 2016)
θFC	Volumetric water content at field capacity (%)	Soil texture-dependent	±20% (Dantec-Nédélec et al., 2016)
θR	Residual volumetric water content (m ³ m ⁻³)	Soil texture-dependent	±20% (Dantec-Nédélec et al., 2016)
θSAT	Saturated volumetric water content (m ³ m ⁻³)	Soil texture-dependent	±20% (Dantec-Nédélec et al., 2016)
θTransp_max	Fraction of saturated volumetric soil moisture above which transpiration is maximum (unitless)	Soil texture-dependent	±20% (Dantec-Nédélec et al., 2016)
C_dry	Dry soil heat capacity (J m ⁻³ K ⁻¹)	Soil texture-dependent	±20% (Dantec-Nédélec et al., 2016)
Post carbon uptake and allocation parameters			
soilC	Scalar on the active soil C pool content to account for uncertainty in spinup (unitless)	(-)	0.5, 2 (Mahmud et al., 2021)

soil_Q10	Temperature dependency factor for heterotrophic respiration ($Q_{10} = \exp^{SOIL_Q10}$) (unitless)	(-)	FI-HYY: 0.53, 1.36 (Barba et al., 2018) US-HA: 0.88, 1.37 (Giasson et al., 2013)
Min_SWC_resp	Minimum soil wetness to limit the heterotrophic respiration (unitless)	(-)	0.1, 0.6 (Mahmud et al., 2021)

35

36

37

38

39 **Text S1: Determination of the variation range for α and β parameters.**

40

41 The values of the α and β parameters are found in Whelan et al. (2016), but no range of variation is given. A
42 similar expression of the production term is defined in Meredith et al. (2018),

$$43 \quad P_{Meredith} = P_{ref} * Q_{10}^{\frac{(T - T_{ref})}{10}} \quad (S1)$$

44 with P_{ref} ($\text{mol m}^{-3} \text{s}^{-1}$) the COS flux at T_{ref} ($^{\circ}\text{C}$).

45 Using the correspondence between the production term describe in Whelan et al. (2016) and Meredith et al. (2018),

$$46 \quad e^{\alpha + \beta * T} = P_{ref} * Q_{10}^{\frac{(T - T_{ref})}{10}} \quad (S2)$$

$$47 \quad \left[\begin{array}{l} \alpha = \log \left(P_{ref} * Q_{10}^{\frac{T_{ref}}{10}} \right) \end{array} \right. \quad (S3)$$

$$48 \quad \left[\begin{array}{l} \beta = \frac{1}{10} \log (Q_{10}) \end{array} \right. \quad (S4)$$

49 Using the identity $\log(a * b^x) = \log(a) + x * \log(b)$ and taking the derivatives, we obtain the following error
50 propagation:

$$51 \quad \left[\begin{array}{l} \Delta\alpha = \frac{\Delta P_{ref}}{P_{ref}} + \frac{\Delta Q_{10}}{Q_{10}} \end{array} \right. \quad (S5)$$

$$52 \quad \left[\begin{array}{l} \Delta\beta = \frac{1}{10} * \frac{\Delta Q_{10}}{Q_{10}} \end{array} \right. \quad (S6)$$

53 Meredith et al. (2018) indicate a ± 1 uncertainty on Q_{10} and a 50% uncertainty on P_{ref} .

$$54 \quad \left[\begin{array}{l} \Delta\alpha = \frac{0.5 * P_{ref}}{P_{ref}} + \frac{1}{e^{10 * \beta}} \end{array} \right. \quad (S7)$$

$$55 \quad \left[\begin{array}{l} \Delta\beta = \frac{1}{10} * \frac{1}{e^{10 * \beta}} \end{array} \right. \quad (S8)$$

56

57

58 **Table S5: Ranges of variation for α and β parameters.**

PFT	$\Delta\alpha$	$\Delta\beta$
1 - Bare soil	NA	NA
2 - Tropical broad-leaved evergreen	1.41	0.0364
3 - Tropical broad-leaved raingreen	1.41	0.0364
4 - Temperate needleleaf evergreen	1.26	0.0304
5 - Temperate broad-leaved evergreen	1.26	0.0304
6 - Temperate broad-leaved summergreen	1.26	0.0304
7 - Boreal needleleaf evergreen	1.26	0.0304
8 - Boreal broad-leaved summergreen	1.26	0.0304
9 - Boreal needleleaf summergreen	1.26	0.0304
10 - C ₃ grass	1.35	0.0340
11 - C ₄ grass	1.35	0.0340
12 - C ₃ agriculture	1.46	0.0383
13 - C ₄ agriculture	1.46	0.0383
14 - Tropical C ₃ grass	1.35	0.0340
15 - Boreal C ₃ grass	1.35	0.0340


59

60

61

62

LONGITUDE : 178.1E(-181.9) to 178.1W (XY ave)
LATITUDE : 90.9S to 90.9N (XY ave) DATA SET: atmcoe_seasonal_2010_2019
FERRET Ver. 8.06
NOVA/FMRL TRAP
21-NOV-2021 15:30:12



ATMCOS (ppt)


63

64

Figure S1: Mean seasonal cycle of monthly atmospheric COS concentrations over 2010-2019.

65

Atmospheric COS concentration (2010-2019)



66


67

68

Figure S2: Mean spatial distribution of atmospheric COS concentrations over 2010-2019.

69

70



71

72 **Figure S3: Spatial distribution of dominant plant functional types (PFTs) in ORCHIDEE over 2010-2019. The map**
 73 **resolution is 0.5°x0.5°.**

74

75

76