

# Observed and ~~project~~forecasted global warming pressure on coastal hypoxia

Michael M. Whitney<sup>1</sup>

<sup>1</sup>Department of Marine Sciences, University of Connecticut, 1080 Shennecossett Road, Groton, CT, USA

5 Correspondence to: Michael M. Whitney ([michael.whitney@uconn.edu](mailto:michael.whitney@uconn.edu))

**Abstract.** Coastal hypoxia is a major environmental problem of increasing severity. A global 40-year observational gridded climate data record and 21<sup>st</sup> century ~~forecasts~~projections from the Community Earth System Model (CESM) under RCP-8.5 forcing are analyzed for long-term linear trends with a focus on warming-related pressures on coastal oxygen ~~conditions~~levels. Projected surface temperature and oxygen conditions are compared to global observations over the 16-year overlapping period (2006-2021). The ~~forecasts~~projected median linear trends for 2006-2100 along the global coast are 0.32 °C, -1.6 mmol m<sup>-3</sup>, and -1.2 mmol m<sup>-3</sup> per decade for sea-surface temperature (SST), oxygen saturation concentration at the surface (surface oxygen capacity), and vertical-minimum oxygen concentration, respectively. These trends point to more rapid deterioration in coastal conditions than experienced over the last four decades; the ~~forecasts~~projected median coastal trends for SST and oxygen capacity are 148% and 118% offaster than the corresponding observed rates. Companion analysis of other models and  
15 climate scenarios indicates projected coastal oxygen trends for the more moderate RCP4.5 and updated SSP5-8.5 scenarios respectively are 37-77% and 103-196% of the CESM RCP8.5 projections. Median rates for the coast and documented hypoxic areas are higher than in the global ocean. Warming and oxygen declines tend to be fastest at high latitudes, one region where new hypoxic areas may emerge as oxygen conditions deteriorate. ~~Over 19% of the coast has extremely rapid forecasts change upwards of 0.60 °C per decade warming and 3.0 mmol m<sup>-3</sup> per decade oxygen change.~~ There is considerable pressure on  
20 current hypoxic areas since future oxygen declines of any magnitude will make hypoxia more severe. The ~~coastal forecasts~~projections can inform coastal environmental management strategies to protect future water quality and ecosystem services.

## 1 Introduction

Hypoxia in coastal waters is a major environmental problem of increasing severity confronted around the world (Hoegh-  
25 Guldberg et al., 2018). “Dead zones in the coastal oceans have spread exponentially since the 1960s and have serious consequences for ecosystem functioning” (Diaz and Rosenberg, 2008). To date, over 500 hypoxic areas in estuaries, coastal seas, and on continental shelves have been documented globally (Breitburg et al., 2018). Furthermore, the severity of hypoxia has increased in many areas (Rabalais et al., 2010). Coastal o<sub>2</sub> concentrations (within 30 km from global coast) have been decreasing an order of magnitude faster than surface-layer concentrations in the open ocean (Gilbert et al., 2010).

30 Worsening coastal oxygen conditions are attributable to the dual human pressures of nutrient overloading fueling eutrophication and anthropogenic climate change (e.g. Rabalais and Turner, 2001; Paerl, 2006; Diaz and Rosenberg, 2011). Coastal oxygen conditions are influenced by many aspects of climate controls including warming waters, altered storm patterns, changing precipitation and river flow, sea-level rise, ~~ocean acidification~~, and shifting ocean circulation (Altieri and Gedan, 2015). This paper focuses on warming-related pressures on coastal hypoxia. Oxygen ~~saturation concentration capacity~~ (oxygen ~~capacity-saturation concentration~~) decreases with increased water temperatures (Weiss, 1970; Garcia and Gordon, 1992), which can exacerbate hypoxia. Oxygen capacity is a succinct synonym for oxygen saturation concentration which is borrowed from physiological research on blood oxygen levels (e.g. Haldane and Smith, 1900; Black, 1940; Maio and Neville, 1965; Bernal et al., 2018) and has been applied to dissolved oxygen in the coastal and open ocean (Helm et al., 2011; Deignan-Schmidt and Whitney, 2017). ~~Furthermore,~~ metabolic rates and related oxygen demands also rise with temperature (Brown et al., 2004). Warming also can intensify and extend the duration of summertime thermal stratification that inhibits ventilation of near-bottom hypoxic waters (Cloern, 2001).

Earth system models provide ~~forecasts~~projections of 21<sup>st</sup> century conditions relevant to coastal hypoxia. Altieri and Gedan (2015) analyzed sea surface temperature (SST) ~~forecasts~~projections from the Community Climate System Model (Collins et al., 2006) for the greenhouse gas emissions A1B scenario (Nakicenovic and Swart, 2000). Under this emission scenario, almost 45 all ~~documented~~known coastal hypoxic areas (based on the Diaz and Rosenberg (2008) dataset) would experience 2 °C or greater water temperature rise by the century end (Altieri and Gedan, 2015). There is a need to update this type of analysis with more recent climate modeling and analyze oxygen trends for Earth system models that incorporate ocean ecosystem dynamics. The present study analyzes Community Earth System Model (CESM) results that include ocean biogeochemistry (Kay et al., 2015a) for the Representative Concentration Pathway (RCP) 8.5 (Moss et al., 2010); which is part of the Climate Model Intercomparison Project phase 5 (CMIP5). The RCP-8.5 exhibits more global warming than the earlier A1B emission scenario (Melillo et al., 2014). The study also compares CESM RCP8.5 results to CMIP5 and CMIP6 for different models and climate scenarios.

Earth system models offer great value in ~~project~~forecasting future conditions, but they do have limitations that affect coastal applications. The relatively coarse horizontal and bathymetric resolution of CESM and most Earth system models, particularly 55 those including ocean biogeochemistry, limits the representation of coastal processes. Earth system models do not currently resolve estuary ecosystems, but they do have the resolution to represent coastal conditions at regional scales. Regional patterns strongly influence rates of change for coastal SST (e.g. Pershing et al., 2015) and temperature-related changes in oxygen capacity. Coastal oxygen concentrations should vary with the regional variations in temperature and oxygen capacity resolved by Earth system models, but oxygen levels also depend on other abiotic and biotic factors. Oxygen concentrations tend to be 60 highly variable on small spatial scales that are not resolved by CESM, particularly within smaller estuarine waters. For these reasons, ~~it is likely that forecasted~~projected changes for coastal oxygen concentrations may be less reliable than SST and oxygen capacity. Open-ocean results at 300 m (Oschlies et al., 2017) and 100-600 m (Cocco et al., 2013) point to model limitations in representing the observed distribution of dissolved oxygen trends. It is reasonable to expect model-observation

oxygen mismatches also occur ~~in open-ocean surface waters and~~ along coasts. ~~The forecasts~~projections for temperature and  
65 oxygen changes, nevertheless, are worth considering and evaluating given the importance of anticipating potentially worsening  
conditions in coastal hypoxic areas. ~~Forecasts~~Projections from Earth system models should be considered in the context of  
observed trends in coastal conditions and compared to available global coastal observations. Gridded SST climate data records  
can provide sufficient global coastal data coverage and allow for computation of oxygen capacities to evaluate whether Earth  
system models provide reasonable representations of coastal conditions affecting hypoxia. Comparison to estuarine oxygen  
70 conditions can be made using long-term observations within some estuaries, but is not practical for all estuaries globally.  
Frankly discussing the limitations and appropriate application of Earth system model projections for hypoxic estuaries is  
important and should motivate future modeling improvements.

The main objective of this paper is to quantify~~studying~~ global patterns exacerbating coastal hypoxia by analyzing linear trends  
in SST, surface oxygen capacity, and ~~(vertical-minimum)~~ oxygen concentrations (the minimum dissolved oxygen in the water  
75 column at each location). Observations from a satellite-derived SST global climate data record (Merchant et al., 2019; Embury  
and Good, 2021) are analyzed for coastal SST and oxygen-capacity trends over the last four decades, which provide context  
for the forecasts~~projections~~. New analysis of 21<sup>st</sup> century RCP8.5 forecasts~~projections~~ from the CESM Large Ensemble Project  
(Kay et al., 2015a,b) is completed for coastal areas and compared to open-ocean rates. Observed and project~~forecasted~~ coastal  
SST and oxygen capacities are compared for the first 16 years of the forecast-projection period that already have occurred. The  
80 study not only investigates forecasts~~projections~~ for documented coastal hypoxic locations, but also considers the entire global  
coast to include unknown and potentially emerging hypoxic areas. Companion analysis of other CMIP5 RCP8.5 models, the  
more moderate RCP4.5 scenario, and the corresponding updated CMIP6 Shared Socioeconomic Pathways (SSP) 2-4.5 and 5-  
8.5 provides context for the CESM RCP8.5 coastal results. Comparisons to prior studies are included and applications to some  
well-studied hypoxic estuaries are discussed. Results are placed in the context provided by other studies and the~~Implications~~  
85 for future coastal hypoxia and management ~~–are discussed,~~ limitations of current Earth system model projections are  
considered, and modeling techniques for better projections of coastal oxygen conditions are recommended.

## 2 Methods

### 2.1 Observations

The global observational dataset analyzed is the satellite-based SST time-series described in Merchant et al. (2019) and  
90 available with updates at the Climate Data Store of the Copernicus Climate Change Service (Embury and Good, 2021). The  
Level-4 (version 2.0) product combines SST data from several satellite platforms to construct a high-quality climate data  
record that has been validated with *in situ* observations. The dataset is a daily product on a regular grid with 0.05° (latitude and  
longitude) resolution. The Level-4 product is gap-filled so that each grid point has an SST value for every day from September  
1981 up to within a month of present time. The SST data variable used in this study has been corrected from skin temperature  
95 to a nominal depth of 0.2 m in order to pair well with *in situ* surface observations (Merchant et al., 2019). This study analyzes

40 years of data spanning 1982-2021. Daily data during each August and February are used to represent summer conditions in the northern and southern hemispheres, respectively. The rationale for analyzing these months is they are the summer months in each hemisphere when water temperatures tend to be highest and oxygen levels tend to be lower. Daily data are averaged to create August-averaged and February-averaged SST time series for the northern and southern hemispheres, respectively.

100 Oxygen capacities ~~are the oxygen saturation concentrations~~ is calculated with the Garcia and Gordon (1992) ~~oxygen saturation concentration~~ equations using the monthly averaged SST data and a constant 35 salinity. The constant salinity is used because the Merchant et al. (2019) product does not include salinity and because this straightforward approach is sufficient to provide observational context for ~~the forecasts~~ projections. The sensitivity of oxygen capacity trends to the choice of salinity is assessed with different salinities specified in the calculations. Coastal points are defined as grid points with at least on  
105 neighboring land cell (directly to the east, west, north, or south). Techniques for comparing observations to ~~forecasts~~ projections for overlapping years, for calculating observed linear trends, and extracting information at documented hypoxic areas are described in subsequent sections.

## **2.2 Forecasts** **CESM RCP8.5 projections**

Forecasts Projections of 21<sup>st</sup> century water temperatures and oxygen conditions are derived from the CESM Large Ensemble  
110 Project that includes ocean biogeochemistry (Kay et al., 2015a), which is a contributor to Climate Model Intercomparison Project phase CMIP-5 (CMIP5, Taylor et al., 2012). Monthly-averaged results for 2006-2100 are accessed via the Earth System Grid (Kay et al., 2015b). Multiple ensemble members for RCP-8.5 forcing (following CMIP5 protocols) are used: 35 and ~~297~~  
ensemble members for ~~temperature~~ SST and oxygen ~~variable~~ capacity, respectively. Runs 1-35 are included for SST (the  
115 “SST” variable) and temperature at fixed vertical levels (the “TEMP” variable). ~~Runs~~ Runs 1, 2, and 9-35 are analyzed for  
surface oxygen ~~saturation concentration~~ capacity (saturation concentration at the surface, the “O2SAT” variable), ~~and~~ vertical-  
minimum oxygen concentration (minimum dissolved oxygen concentration in each water column, the “O2\_ZMIN” variable),  
oxygen concentrations at fixed vertical levels (the “O2” variable), and the apparent oxygen utilization at the same levels (the  
“AOU” variable). Runs 3-8 are omitted for ~~O2SAT and O2\_ZMIN~~ oxygen variables because these results are not available on  
120 the Earth System Grid. Additional runs (101 and higher) were avoided because of a documented unexplained positive bias in  
global temperature relative to other ensemble members. The surface level (representing 0-10 m depth) is subsampled from the  
O2 variable to obtain surface oxygen concentrations. All included runs are averaged together to produce ensemble-mean  
monthly time series of SST, surface oxygen ~~saturation concentration (oxygen capacity)~~, surface oxygen concentration, and  
vertical-minimum oxygen concentrations at each ocean grid cell. Results from individual runs are analyzed to assess  
uncertainty associated with differences among ensemble members. Supplementary analysis of temperatures, oxygen  
125 concentrations, and AOU at 10-m intervals down to 100 m deep is included to describe transitions from surface conditions to  
deeper levels in the coastal water column.

For each year at each CESM ocean grid cell (at 1° latitude nominal resolution), the month with minimum surface oxygen saturation concentration is selected to construct annual time series of minimum oxygen capacity and coincident SST and

~~vertical-minimum~~ oxygen concentrations. August and February are the median months associated with minimum surface oxygen capacity in the northern and southern hemispheres, respectively. The global set of all coastal points includes all CESM ocean grid cells with at least one neighboring land cell (there are 4,899 coastal cells). ~~Coastal points are defined as grid points with at least one neighboring land cell. Globally, there are 4,899 cells coastal cells.~~ Techniques for comparing ~~forecastsprojections~~ and observations for overlapping years, for calculating ~~forecastedprojected~~ linear trends, and extracting information at documented hypoxic areas are described in the following sections.

### 2.3 Comparison of observations and ~~forecastsprojections~~

To characterize the reliability of CESM in coastal areas, observations and ~~forecastsprojections~~ are compared for global coastal points. There are 16 ~~forecast- projection-period~~ years spanning 2006-2021 that now overlap with observed conditions. Every CESM coastal point is matched with the closest coastal point on the observational grid, which has higher spatial resolution than the ~~CESMforecast~~ grid. The summer month values (calculated as described in previous section) for the overlapping period are averaged together to determine mean observed and ~~forecastedprojected~~ values at each coastal point. The resulting ~~mean~~ SST and oxygen capacity values are ~~quantitatively compared to each other used to assess local and global biases relative to observations.~~ Linear regression is completed to indicate the global relationship between observed and ~~forecastedprojected~~ SST and oxygen capacity for coastal waters. Regression results are included in the supporting dataset (Whitney, 2021) and are reported with the slope, offset, p-value (p) of the F-statistic, correlation coefficient squared ( $r^2$ ), ~~and~~ root mean square error (RMSE), ~~and standard error of the regression slope ( $\sigma_s$ ).~~ The level of agreement between observed and ~~forecastedprojected~~ mean values for the overlapping period assesses the quality of CESM performance in coastal waters. Observed and ~~forecastedprojected~~ temporal trends are not compared for the overlapping period because observations have high interannual variability that can obscure longer-term trends over shorter periods. Individual ensemble members have comparably large interannual vari~~ations~~bility, but the ensemble-mean ~~projectionsforecast~~ results have much less interannual variability.

### 2.4 Trend analysis

Linear regression analysis is applied to characterize long-term temporal trends. ~~It is worth noting that nonlinear trends are possible, but the strategy is to focus on constant long-term rates of change and the results indicate linear trends fit the data well in most locations globally.~~ For ~~the~~ observations, the entire 40-year observational period is regressed with time for SST and surface oxygen capacity. For ~~the CESM RCP8.5 forecastsprojections~~, the entire 94-year ~~projectionforecast~~ period is regressed for ~~ensemble-mean~~ SST, surface oxygen capacity, and ~~surface and vertical-minimum~~ oxygen concentrations. The regression slopes are reported as rates of change associated with each variable and are included in the supporting dataset (Whitney, 2021). ~~These rates are graphed spatially and the spatial median rates are reported for the global ocean, coastal grid cells, and documented hypoxic areas.~~ Regression statistics are calculated. The p-values of the F-statistic are used to categorize results depending on whether or not they are statistically significant at the 90% confidence level ( $p=0.10$ ). Correlation coefficients are calculated but not emphasized because ensemble-mean ~~forecastsprojections~~ average out interannual variability present in

ensemble members and therefore tend to have high  $r^2$  values; while the observational record has higher interannual variability and correspondingly lower  $r^2$  values. Uncertainty associated with the regression fit is quantified with  $\sigma_s$ , the standard error of the regression slope. Based on the  $\sigma_s$  values, the corresponding standard error of the spatial median values ( $\sigma_m$ ) are reported for global, coastal, and documented hypoxic areas. For CESM RCP8.5 projections, variability among ensemble members is characterized by calculating regression slopes for each member and then calculating the associated standard error of the ensemble-mean rates of change ( $\sigma_e$ ) at each grid cell. The spatial median rates for each ensemble member also are calculated and the corresponding standard deviation of these median rates ( $\sigma_{me}$ ) are reported as another way to characterize intra-ensemble variability. Rates of change are shown for the entire global ocean and emphasis is placed on change in coastal waters. The documented coastal hypoxic areas are further highlighted and are identified as described in the next section.

## 170 **2.5 Documented coastal hypoxic areas**

The 532 documented coastal hypoxic areas analyzed are from the Diaz et al. (2011) database. The dataset is an expanded version of the Diaz and Rosenberg (2008) database, which is similar to the coastal dataset in Breitburg et al. (2018). The oxygen concentration threshold used to identify hypoxia in the dataset is  $63 \text{ mmol m}^{-3}$  ( $2 \text{ mg L}^{-1}$ ) (Diaz and Rosenberg, 2008). The Diaz et al. (2011) database also includes 244 additional locations classified as eutrophic (but not hypoxic); these documented eutrophic points that are not isolated as a group in this analysis, but are included among the global coastal points analyzed included in this analysis. Rates of change (for ~~SST, oxygen capacity, and oxygen concentration~~ temperature and oxygen variables) for the documented hypoxic areas are extracted from rates at the nearest observational coastal grid point or CESM coastal ocean grid cell. The observational grid resolution is fine enough to represent each individual documented coastal hypoxic area with a separate observational grid point. The coarser (nominally  $1^\circ$  latitude) CESM resolution causes some documented locations to be represented by the same CESM coastal cell. The 532 documented locations map to only 240 unique CESM ocean grid cells.

## 180 **2.6 Other projections**


Projections from other models and climate scenarios provide context for the CESM RCP8.5 ensemble results. Data are accessed online via the Earth System Grid Federation. Only models with ocean biogeochemistry are included. Monthly-averaged results for vertical-minimum oxygen concentrations (the “o2min” variable) are analyzed. The CMIP5 RCP4.5 and RCP8.5 sets include five other models: HadGEM2-ES, IPSL-CM5A-LR, IPSL-CM5A-MR, MPI-ESM-LR, and MPI-ESM-MR (Collins et al., 2011; Dufresne et al., 2013; Giorgetta et al., 2013). The CESM2 (CMIP6) SSP2-4.5 and SSP5-8.5 sets include the three available ensemble members (r4i1p1f1, r10i1p1f1, and r11i1p1f1) with oxygen data (Danabasoglu et al., 2020). The CMIP6 SSP5-8.5 set of other models includes CanESM5 (r1i1p2f1 run), IPSL-CM6A-LR, MPI-ESM1-2-LR, and MPI-ESM1-2-HR (Swart et al., 2019; Boucher et al., 2020; Mauritsen et al., 2019). The r1i1p1f1 runs are selected, unless already noted. Oxygen trends are calculated following the exact same procedures as described above. Analysis years are 2006-2100 and 2015-2100 for CMIP5 and CMIP6 models, respectively.


### 3 Results

#### 195 3.1 Observed trends

The 40-year observational SST record (updated from Merchant et al., 2019) indicates warming has occurred throughout the world ocean (Fig. 1a). SST rates are stronger in the Northern Hemisphere, with the most rapid warming occurring in Arctic areas near coasts. Just over half (55%) of the ocean grid points have linear trends with  $p \leq 0.10$  (Table 1). P-values are lower where calculated rates are lower in parts of the Atlantic, much of the South Pacific, and most of the Southern Ocean. Where  $p \leq 0.10$ , the mean  $r^2 = 0.20$ ; indicating that linear trends describe an appreciable part of the observed variance, but interannual variability is larger. Since the long-term linear trends describe only part of the variance, the uncertainty of the regression slopes is relatively large. Among the locations where  $p \leq 0.10$ , the average  $\sigma_s$  is 30% of the local rate of change. The observed global median SST trend (including only points with  $p \leq 0.10$ ) is  $0.22 \text{ }^\circ\text{C per decade}$  (Table 1). The  $\sigma_m$ , the standard error of the global median trend, is exceedingly small ( $2 \times 10^{-4} \text{ }^\circ\text{C per decade}$ ) because of the large number of data points (Table 1). The median

205





**Figure 1: Observed SST trends: (a) spatial distribution for all ocean observational points, (b) latitudinal dependence for global coastal points and documented coastal hypoxic areas, and (c) histograms of global coastal points and documented coastal hypoxic areas (having linear trends with  $p \leq 0.10$ ). Locations that have linear trends with lower confidence ( $p > 0.10$ ) are marked with grey points. Global coastal data are shown as red points and bars. Documented coastal hypoxic areas are marked in green.**

-trend with all points included (regardless of p-value) is  $0.13^\circ\text{C}$  per decade. This rate is consistent with the global mean SST rate of  $0.1^\circ\text{C}$  per decade from 1982-2013 (Pershing et al., 2015).

Observed rates along the global coast indicate conditions relevant to coastal hypoxia. The median SST trend for global coastal points ( $0.27^\circ\text{C}$  per decade, for points with  $p \leq 0.10$ ) is 22% faster than the observed ocean median rate (Table 1). SST rates tend to be near the median coastal value from  $60^\circ\text{S}$  to  $30^\circ\text{N}$  (Fig. 1b) and increase towards higher latitudes. Similar to the global points, the average  $\sigma_s$  is 34% of the local rate of change for coastal points with  $p \leq 0.10$ . The  $\sigma_m$  is much smaller than the median coastal value (Table 1). The observed median trend for documented hypoxic areas ( $0.24^\circ\text{C}$  per decade) is 6% faster than the ocean median rate (Table 1). The rate in documented hypoxic areas is somewhat (11%), but is lower than the median global coastal rate because there are few documented hypoxic areas at high latitudes where warming is fastest. The histograms of SST rates (Fig. 1c) indicate the documented hypoxic areas have a narrower spectrum than the global coastal points.

The global distribution of oxygen capacity (~~saturation concentration~~) trends (Fig. 2a) at the surface is tied to observed SST rates (Fig. 1a). Since oxygen ~~saturation concentrations~~capacity decreases nonlinearly with temperature with a greater response at lower

temperatures, oxygen capacity decreases are amplified at high latitudes where waters are colder and ~~forecasted~~projected warming rates are faster (Weiss, 1970; Garcia and Gordon, 1992; Altieri and Gedan, 2015). Oxygen capacity linear trends with  $p \leq 0.10$  occur in the same locations as for SST trends and  $r^2$  values are similar to those for SST. The average  $\sigma_s$  is 38% of the local rate of change. The observed global median oxygen capacity trend at the surface (including only points with  $p \leq 0.10$ ) is -



0.89 mmol m<sup>-3</sup> per decade and  $\sigma_m$  is very small (Table 1). The calculated global median rate is several times faster than the median rate of -0.2 mmol m<sup>-3</sup> per decade observed in offshore (>100 km from coast) upper-ocean (0-300 m) waters for 1976-2000 (Gilbert et al., 2010). The mismatch with the prior study is likely due to the more recent time period and the reliance on surface, rather than upper-ocean, observations. Due to these methodological differences, matching rates between studies is not expected, but the earlier study does provide context.

**Table 1: Observed (1982-2021) and forecasted/projected (2006-2100) median linear trends in SST, surface oxygen capacity, surface oxygen concentrations (projected only), and vertical-minimum oxygen concentrations (forecasted/projected only) for the ocean, global coastal cells, and documented coastal hypoxic areas. All points included in median calculations have linear trends with  $p \leq 0.10$ , the corresponding percent coverage of  $p \leq 0.10$  is given in square brackets. Median rates are listed with  $\pm \sigma_m$ , the standard errors of the median values.**

Rate	Ocean	Coastal	Hypoxic
Observed SST (°C per decade)	0.218 $\pm 2 \times 10^{-4}$ [55.12%]	0.265 $\pm 10^{-3}$ [70.64%]	0.236 $\pm 0.01$ [76.13%]
Observed oxygen capacity (mmol m <sup>-3</sup> per decade)	-0.82 $\pm 10^{-3}$ [55.06%]	-1.34 $\pm 0.01$ [70.64%]	-0.78 $\pm 0.04$ [76.13%]
Forecasted/Projected SST (°C per decade)	0.352 $\pm 3 \times 10^{-5}$ [100%]	0.392 $\pm 2 \times 10^{-4}$ [100%]	0.402 $\pm 2 \times 10^{-4}$ [100%]
Forecasted/Projected oxygen capacity (mmol m <sup>-3</sup> per decade)	-1.23 $\pm 2 \times 10^{-4}$ [100%]	-1.58 $\pm 2 \times 10^{-3}$ [99.98%]	-1.39 $\pm 10^{-3}$ [100%]
Projected surf. oxygen conc. (mmol m <sup>-3</sup> per decade)	-1.33 $\pm 6 \times 10^{-4}$ [99.19%]	-1.51 $\pm 4 \times 10^{-3}$ [98.12%]	-1.41 $\pm 2 \times 10^{-3}$ [100%]
Forecasted/Projected min. oxygen conc. (mmol m <sup>-3</sup> per decade)	-0.65 $\pm 4 \times 10^{-4}$ [97.24%]	-1.15 $\pm 3 \times 10^{-3}$ [97.39%]	-1.38 $\pm 5 \times 10^{-3}$ [100%]


The observed median oxygen capacity trend for global coastal points (-1.34 mmol m<sup>-3</sup> per decade, for points with  $p \leq 0.10$ ) is 62% faster than the surface ocean median rate (Table 1). Similar to the global points, the average  $\sigma_s$  is 34% of the local rate of change for coastal points with  $p \leq 0.10$  and  $\sigma_m$  is much smaller than the median coastal value (Table 1). The observed median coastal rate is half of the rate calculated for a global coastal band (within 30 km of the coast) for 1976-2000 (Gilbert et al., 2010). For the reasons mentioned above, a match between the studies is not expected. It is interesting that the current study has faster global rates and slower coastal rates; the underlying reasons are not explored here. Rates tend to be near the median coastal value from 60 °S to 30 °N (Fig. 2b). At higher latitudes, rates tend to increase with latitude. The median oxygen capacity trend for documented hypoxic areas is -0.8 mmol m<sup>-3</sup> per decade (Table 1). This rate is lower magnitude than for all global coastal points because of little high-latitude coverage, where oxygen capacity rates are largest. Similar to SST rates, the histograms for oxygen capacity (Fig. 2c) indicate the documented hypoxic areas have a narrower spectrum than the global

coastal points. Overall, the observational SST and oxygen capacity analysis provides context for the ~~forecasts~~projections and new information about global coastal conditions influencing coastal hypoxia.

255 ~~As described in the methods, oxygen capacity is calculated from SST and a constant 35 salinity. This approach neglects changes in oxygen capacity due to long-term salinity variability and the constant salinity choice may bias the calculate oxygen capacity trends. Sensitivity calculations with constant salinity values of 32 and 34 had RMSE (relative to the original calculations) of at most  $10^{-3}$  mmol m<sup>-3</sup> per decade. Introducing a long-term linear salinity trend of 34.9 to 35.1 (or vice versa) over the observation period created similarly small RMSE of  $3 \times 10^{-3}$  mmol m<sup>-3</sup> per decade. The sensitivity calculations indicate that~~  
260 ~~assuming a constant salinity does not introduce much error in the oxygen capacity trends.~~

### 3.2 Comparison of observations and ~~forecasts~~projections


The observational SST record is compared to CESM ~~RCP8.5 forecasted~~projected coastal conditions for the overlapping 16 years spanning 2006-2021 (Fig. 3a). ~~As described in the methods, the comparison involves mean summer-month values for the overlapping period rather than comparing time series with relatively short-term interannual variability.~~ There is a ~~nearly~~essentially one-to-one relationship between observed and ~~forecasted~~projected coastal SST with a low p-value ( $p < 0.001$ ) and high correlation ( $r^2 = 0.97$ ). The ~~projected temperatures have a small positive bias (0.3 °C) and a moderate~~is approximate 5 °C scatter around the linear relationship, but the RMSE is small (0.031.9 °C) relative to observations. This level of agreement indicates that CESM results are broadly representative of global coastal SST conditions. The comparison of observed and  
270 ~~forecasted~~projected coastal oxygen capacities averaged over the overlapping period (Fig. 3b) show a near one-to-one relationship with a low p-value ( $p < 0.001$ ) and high correlation ( $r^2 = 0.94$ ). Projected oxygen capacities have a bias and RMSE of -5 and 17 mmol m<sup>-3</sup>, respectively. The scatter away from the regression line is larger at higher oxygen capacities (in colder waters). The nonlinear relationship between temperature and oxygen capacity (~~noted above~~) means that temperature scatter in colder waters translates into more oxygen capacity scatter. This analysis points to the degree of reliability of CESM results for  
275 representing coastal oxygen capacities. Overall, the comparison of observations and CESM results suggest that CESM can provide reliable ~~forecasts~~projections of coastal conditions relevant to coastal hypoxia.



280 **Figure 2: Observed surface oxygen capacity (saturation concentration) trends: (a) spatial distribution for all ocean observational points, (b) latitudinal dependence for global coastal points and documented coastal hypoxic areas, and (c) histograms of global coastal points and documented coastal hypoxic areas. The format and color coding follow Fig. 1.**

### 3.3 Projected trends


285 The CESM Large Ensemble Project ensemble-mean projection for the RCP8.5 scenario indicates SST will appreciably increase throughout the world ocean over the 21<sup>st</sup> century (Fig. 4a). All of the ocean cells have linear trends with  $p \leq 0.10$  (Table 1). SST rates account for more than 90% of the variance in the ensemble-mean projection ( $r^2 > 0.90$ ) for most of the ocean; the only




**Figure 3: Comparison of ~~forecasted~~ projected to observed coastal conditions: (a) SST and (b) surface oxygen capacity. Mean values for the overlapping 2006-2021 period are compared. Global coastal points and documented coastal hypoxic areas are marked in red and green, respectively. The one-to-one line (dashed) and linear regression (solid) are included along with associated regression statistics ( $p < 0.001$  for both regressions).**

exceptions are some areas to the north and south of Greenland and near parts of Antarctica. The average  $\sigma_s$  is only 2% of the local rate of change for points with  $p < 0.10$ . The  $\sigma_s$  values for the projections are much smaller than for observations. The ~~forecasted~~ projected global median SST trend is 0.35 °C per decade and the associated  $\sigma_m$  is negligible (Table 1). Global distributions of SST warming have been studied in detail for multiple models and RCP scenarios (e.g. Bopp et al., 2013). Bopp et al. (2013) includes CESM simulations in an analysis of ten models running the RCP-8.5 scenario and finds the global average SST increase is 0.27 °C per decade (from the 1990s to 2090s) when averaged across all included models. The smaller warming rate for the Bopp et al. (2013) results is connected to including multiple models and due to calculating rates relative to the 1990-1999 historical period instead of limiting analysis to the 2006-2100 CMIP5 projection period.—The CESM ~~forecasted~~ projected global median SST trend is 61% higher than the observed global warming rate (0.22 °C per decade) calculated in the previous section (Table 1). Under the RCP-8.5 scenario, the ocean SST will increase considerably faster than the observed linear trend over the last four decades.

The global distribution of SST warming indicates variations among oceans, with the Arctic Ocean ~~forecasted~~ projected to experience at least twice the median warming rates (Fig. 4a). The influence of ocean circulation patterns on warming rates also is evident.—Observations (Fig. 1a) also indicate rapid SST increases near Arctic coasts and influences of ocean circulation are evident. There are, however, clear differences in the spatial structure of ~~forecasted~~ projected and observed rates. Differences away from the coast in the Arctic Ocean are immediately apparent: the projected SST rates are much stronger than observed rates away from the coasts. These offshore differences are not explored further here, as the focus is on coastal conditions. The high p-values of



310



**Figure 4: ~~Forecasted~~Projected SST trends: (a) spatial distribution for all CESM ocean grid cells, (b) latitudinal dependence for global coastal points and documented coastal hypoxic areas, and (c) histograms of global coastal points and documented coastal hypoxic areas. The format and color coding follow Fig. 1.**

315 observed SST trends in much of the Southern Ocean ( $p > 0.10$ ) preclude comparisons of projected and observed spatial structure in this region.


Focusing on warming in global coastal areas reveals new information directly relevant to coastal hypoxia. The ~~forecasted~~projected median SST trend for global coastal points ( $0.39 \text{ }^\circ\text{C per decade}$ ) is 11% faster than the ~~forecasted~~projected ocean median rate and 48% higher than the observed median coastal rate (Table 1). Local coastal SST rates (with  $\sigma_3$  averaging

320 2% of the coastal rates) tend to be near the median coastal value from 60 °S to 30 °N (Fig. 4b). Above 30 °N, warming rates tend to increase with latitude and variability among coasts increases. The latitudinal patterns are broadly similar for forecastedprojected and observed coastal SST rates (Fig. 4b and Fig. 1b). The forecastedprojected median SST trend for documented hypoxic areas (0.40 °C per decade) is 14% faster than the forecastedprojected ocean median rate (Table 1). For documented hypoxic areas, the forecastedprojected median trend is 70% higher than the median observed rate (Table 1) and  
325 75% faster than the median trend under moderate A1B emissions scenario (Altieri and Gedan, 2015). The documented hypoxic areas sample much of the variability in coastal SST rates from 45 °S to 60 °N. The hypoxic area database, however, has little coverage at higher latitudes, where the most rapid warming is forecastedprojected. Histograms (Fig. 4c) indicate most coastal locations (92%) and documented hypoxic areas (96%) are forecastedprojected to warm more than 0.3 °C per decade; 0.1 °C per decade faster than the A1B results in Altieri and Gedan (2015). A significant portion of the global coast (18%) are  
330 forecastedprojected to warm faster than 0.60 °C per decade, but few of the documented hypoxic sites (2%) are in this range because most are not at high latitudes. Supplementary trend analysis of temperatures at 10-m intervals down to 100 m deep also indicates robust warming trends (not shown), but the coastal warming rates at the 20-30 m, 40-50 m, and 90-100 m levels decrease to 92%, 80%, and 70% the SST trend magnitude, respectively.

The global distribution of forecastedprojected oxygen capacity (saturation concentration) trends (Fig. 5a) at the surface is  
335 tightly linked to SST rates (Fig. 4a). All of the ocean cells have linear trends with  $p \leq 0.10$  (Table 1). The forecasted oxygen capacity trends have  $r^2 > 0.90$  over most of the ocean with exceptions in the areas where SST rates have  $r^2 \leq 0.90$ . The  $\sigma_s$  values on average are 2% of local rates. The forecastedprojected global median oxygen capacity trend at the surface is  $-1.2 \text{ mmol m}^{-3}$  per decade and  $\sigma_m$  is diminishingly small (Table 1). The median rate is 50% higher than the observed global median trend (Table 1). The forecastedprojected median rate for ocean waters above 60 °N ( $-5.3 \text{ mmol m}^{-3}$  per decade) is several times  
340 higher than the total ocean median rate. Projected oxygen capacity trends in the Arctic Ocean, particularly offshore, are much stronger than observed rates; this Arctic pattern echoes the differences in SST rates.

The forecastedprojected median oxygen capacity trend for global coastal points ( $-1.6 \text{ mmol m}^{-3}$  per decade) is 28% faster than the surface ocean median rate (Table 1). The forecastedprojected median coastal rate is 18% faster than observed. Rates tend to be near the median value from 30 °S to 30 °N (Fig. 5b). Outside that latitude range, rates tend to increase with latitude and  
345 variability among coasts increases, particularly in the northern hemisphere where rates can exceed  $-10.0 \text{ mmol m}^{-3}$  per decade. The latitudinal pattern in coastal oxygen capacity trends (Fig. 5b) is similar to the observed coastal pattern (Fig. 2b). The median oxygen capacity trend for documented hypoxic areas is  $-1.4 \text{ mmol m}^{-3}$  per decade (Table 1). This rate is lower magnitude than for all global coastal points because of little high-latitude coverage, where oxygen capacity rates are largest. The forecastedprojected median trend for documented hypoxic areas is 78% higher than observed (Table 1). Histograms (Fig.  
350 5c) indicate for most coastal locations (95%) and all documented hypoxic areas the oxygen capacity trend is forecastedprojected to be faster than  $-0.9 \text{ mmol m}^{-3}$  per decade. A significant portion of the global coast (28%) has forecastedprojected rates faster than  $-3.0 \text{ mmol m}^{-3}$  per decade, but few of the documented hypoxic sites (4%) are in this range because most are not at high latitudes.

The distribution of surface oxygen concentration trends (not shown) is very similar in terms of magnitudes and spatial patterns to the distribution for oxygen capacity trends (Figure 5). The median surface oxygen concentration trends for global, coastal,




**Figure 5: ~~Forecasted~~Projected surface oxygen capacity (saturation concentration) trends:** (a) spatial distribution for all CESM ocean grid cells, (b) latitudinal dependence for global coastal points and documented coastal hypoxic areas, and (c) histograms of global coastal points and documented coastal hypoxic areas. The format and color coding follow Fig. 1.

and documented hypoxic areas are within 8% of the corresponding median oxygen capacity trends. The link between surface oxygen capacity and concentrations diminishes with depth.

The projected global distribution of vertical-minimum oxygen concentration trends indicates decreases in most areas (Fig. 6a). In CESM and in nature, the depth of the vertical-minimum oxygen concentration tends to be near-bottom in coastal areas and

365



**Figure 6: ~~Forecasted~~Projected** vertical-minimum oxygen concentration trends: (a) spatial distribution for all CESM ocean grid cells, (b) latitudinal dependence for global coastal points and documented coastal hypoxic areas, and (c) histograms of global coastal points and documented coastal hypoxic areas. The format and color coding follow Fig. 1 except for the blue shading indicating increasing oxygen concentrations.

370



is bathymetrically constrained to be relatively close to the surface (within ~100 m), while the minimum oxygen concentration can occur much deeper in the open ocean and consequently have a more remote connection to surface oxygen capacity. The projections for most ocean cells have linear trends with  $p \leq 0.10$  (Table 1). On average, the  $\sigma_s$  is 6% of the local trends. The global median trend in oxygen is  $-0.7 \text{ mmol m}^{-3}$  per decade and  $\sigma_m$  is small (Table 1). The global distribution of vertical-minimum oxygen concentration rates (Fig. 6a) is broadly consistent with the global distribution of the RCP8.5 model-average projected changes in oxygen concentrations (at 200-600 m depth) shown in Bopp et al. (2013). The total ocean oxygen content decrease (from the 1990s to 2090s) calculated in Bopp et al. (2013) translates to a  $-0.6 \text{ mmol m}^{-3}$  per decade trend, which is close to the current results. It is noteworthy that some areas in the tropics and Arctic with projected vertical-minimum oxygen increases in spite of surface oxygen capacity decreases. The interplay between oxygen capacities and concentrations is described below in a coastal context.

The median trend in vertical-minimum oxygen concentrations for global coastal points ( $-1.2 \text{ mmol m}^{-3}$  per decade) is 77% faster than the ocean median rate (Table 1). Similar to oxygen capacity and surface oxygen concentrations, vertical-minimum oxygen concentrations rates tend to be near the median value from  $30^\circ\text{S}$  to  $30^\circ\text{N}$  and increase at higher latitudes (exceeding  $-10.0 \text{ mmol m}^{-3}$  per decade), but there is a lot of scatter (Fig. 6b). Similar to the global points,  $\sigma_s$  averages only 5% of the trends at coastal points. The median trend for documented hypoxic areas ( $-1.4 \text{ mmol m}^{-3}$  per decade) is somewhat stronger than for all coastal points ~~because of little high latitude coverage and approximately the same as the corresponding oxygen capacity decrease~~ (Table 1). Histograms (Fig. 6c) indicate most coastal locations (72%) and documented hypoxic areas (89%) have ~~foreeastedprojected~~ vertical-minimum oxygen concentration declines. Some coastal (19%) and documented hypoxic areas (8%) have ~~foreeastedprojected~~ trends faster than  $-3.0 \text{ mmol m}^{-3}$  per decade.

The median trends for vertical-minimum oxygen concentrations in the global ocean and for coastal points are respectively only 49% and 76% of the corresponding trends for surface oxygen capacity. The trends in oxygen concentration weaken with depth (for the upper 100-m range analyzed). The weakening of warming trends with depth (mentioned above) accounts for part of this difference due to the temperature-dependence of oxygen capacity. Notwithstanding, the median trends for oxygen concentrations are smaller than the oxygen capacity trends at the corresponding depths. This situation is consistent with decreasing AOU, as indicated by supplementary analysis of AOU trends within the upper 100 m. It should be noted that some coastal areas having stronger, equal, or weaker oxygen rates than oxygen capacity trends and some areas even have oxygen increases despite decreasing oxygen capacity. The ecosystem dynamics for the variety of coastal oxygen situations occurring within the model warrant further investigation beyond this study.

#### 400 3.4 Intra-ensemble variability and context from other projections

-Linear trends calculated for each individual member of the CESM RCP8.5 ensemble differ from the ensemble-mean trends already described. The intra-ensemble variability is characterized by  $\sigma_e$ , the standard error of the ensemble-mean rates of change, which is calculated at each grid cell. Median  $\sigma_e$  values for ocean, coastal, and documented hypoxic area are included in Table 2. For SST, and oxygen capacity, and surface oxygen concentration trends,  $\sigma_e$  is small locally and the median  $\sigma_e$

405 values are only 5-7% of the median ensemble-mean trends (Table 2). The  $\sigma_e$  values are larger for vertical-minimum oxygen  
concentration trends. The median  $\sigma_e$  is 10-24% of the median ensemble-mean trends; this source of uncertainty is highest in  
the global ocean and lowest for documented hypoxic areas (Table 2). Overall, the relatively small  $\sigma_e$  values indicate that each  
ensemble member projects long-term trends that are similar to each other and the ensemble-mean projections. Another way to  
410 **Table 2: Variability among CESM RCP8.5 ensemble members for ocean, global coastal cells, and documented coastal hypoxic areas**  
**where  $p \leq 0.10$ . Median  $\sigma_e$  values (over all included grid cells) are reported along with  $\sigma_{me}$  values. Corresponding percentages in**  
**brackets express the ratio of  $\sigma_e$  and/or  $\sigma_{me}$  to the median ensemble-mean trends (Table 1).**

<u>Rate</u>	<u>Ocean</u>	<u>Ocean</u>	<u>Coastal</u>	<u>Coastal</u>	<u>Hypoxic</u>	<u>Hypoxic</u>
	<u><math>\sigma_e</math></u>	<u><math>\sigma_{me}</math></u>	<u><math>\sigma_e</math></u>	<u><math>\sigma_{me}</math></u>	<u><math>\sigma_e</math></u>	<u><math>\sigma_{me}</math></u>
<u>Projected SST</u> <u>(°C per decade)</u>	<u>0.02</u> <u>{6%}</u>	<u><math>5 \times 10^{-3}</math></u> <u>{1%}</u>	<u>0.02</u> <u>{6%}</u>	<u><math>6 \times 10^{-3}</math></u> <u>{2%}</u>	<u>0.02</u> <u>{5%}</u>	<u>0.01</u> <u>{2%}</u>
<u>Projected oxygen capacity</u> <u>(mmol m<sup>-3</sup> per decade)</u>	<u>0.08</u> <u>{7%}</u>	<u>0.02</u> <u>{2%}</u>	<u>0.09</u> <u>{6%}</u>	<u>0.05</u> <u>{3%}</u>	<u>0.08</u> <u>{5%}</u>	<u>0.05</u> <u>{4%}</u>
<u>Projected surf. oxygen conc.</u> <u>(mmol m<sup>-3</sup> per decade)</u>	<u>0.08</u> <u>{6%}</u>	<u>0.02</u> <u>{2%}</u>	<u>0.10</u> <u>{7%}</u>	<u>0.03</u> <u>{2%}</u>	<u>0.13</u> <u>{6%}</u>	<u>0.04</u> <u>{3%}</u>
<u>Projected min. oxygen conc.</u> <u>(mmol m<sup>-3</sup> per decade)</u>	<u>0.15</u> <u>{24%}</u>	<u>0.03</u> <u>{5%}</u>	<u>0.23</u> <u>{20%}</u>	<u>0.03</u> <u>{3%}</u>	<u>0.13</u> <u>{10%}</u>	<u>0.04</u> <u>{3%}</u>

415 look at the intra-ensemble variability is considering  $\sigma_{me}$ , the standard deviation of the spatial median rates from each ensemble  
member (analogous the median values in Table 1). The  $\sigma_{me}$  values are only 1-5% of the corresponding median ensemble-mean  
rates.- Thus, the reported median trends are robust regardless of whether an individual run or ensemble-mean is analyzed.

420 The CESM RCP8.5 ensemble results can be placed within a context of projections from different models with various climate  
scenarios. The focus is on projections from Earth system models that include biogeochemistry and the ranges of median rates  
for vertical-minimum oxygen concentrations are compared (Table 3). Means and standard deviations of the median rates within  
each model set are not calculated due to the small number of runs in most sets. The more moderate CMIP5 RCP4.5 scenario  
and its updated companion CMIP6 SSP2-4.5 have the smallest rates, particularly in coastal and documented hypoxic areas,  
but still project appreciable declines in oxygen concentrations. For instance, the CMIP5 RCP4.5 coastal projections are 37-

425 **Table 3: Projected median linear trends in vertical-minimum oxygen concentrations (mmol m<sup>-3</sup> per decade) for the ocean, global**  
**coastal cells, and documented coastal hypoxic areas. All points included in median calculations have linear trends with  $p \leq 0.10$ . The**  
**reported numbers are the range of median rates from individual runs included in each model set. The number of models in each set**  
**are listed in the count column. The CESM (CMIP5) RCP8.5 set includes the ensemble mean data and all individual ensemble**  
430 **members with oxygen data.**

<u>Model set</u>	<u>Count</u>	<u>Ocean</u>	<u>Coastal</u>	<u>Hypoxic</u>
<u>Other CMIP5 RCP4.5</u>	<u>5</u>	<u>-0.77 to -0.26</u>	<u>-0.88 to -0.43</u>	<u>-0.74 to -0.34</u>
<u>CESM2 (CMIP6) SSP2-4.5</u>	<u>3</u>	<u>-0.73 to -0.68</u>	<u>-0.81 to -0.76</u>	<u>-0.82 to -0.81</u>
<u>Other CMIP5 RCP8.5</u>	<u>5</u>	<u>-1.61 to -0.34</u>	<u>-1.72 to -0.66</u>	<u>-1.33 to -0.63</u>
<b><u>CESM (CMIP5) RCP8.5</u></b>	<b><u>30</u></b>	<b><u>-0.80 to -0.65</u></b>	<b><u>-1.30 to -1.15</u></b>	<b><u>-1.46 to -1.32</u></b>
<u>CESM2 (CMIP6) SSP5-8.5</u>	<u>3</u>	<u>-1.02 to -0.94</u>	<u>-1.52 to -1.50</u>	<u>-1.65 to -1.63</u>
<u>Other CMIP6 SSP5-8.5</u>	<u>4</u>	<u>-1.74 to -0.71</u>	<u>-2.25 to -1.18</u>	<u>-2.00 to -1.13</u>

77% the CESM RCP8.5 median coastal trend. The CESM RCP8.5 ensemble members are in the middle range of other CMIP5 RCP8.5 projections for ocean and coastal areas, but are on the high end of rates in documented hypoxic areas. The CESM2 (CMIP6) SSP5-8.5 projected oxygen trends are 13-40% stronger than the CESM RCP8.5 projections. The CESM RCP8.5 and CESM2 SSP5-8.5 projections mostly fall within the low to mid ranges of the CMIP6 SSP5-8.5 projections from other models. The SSP5-8.5 coastal projections from all analyzed models are 103-196% of the CESM RCP8.5 median coastal trend. It is notable that median oxygen trends are not larger in documented hypoxic areas than coastal areas for all projections, though this is the pattern in CESM RCP8.5 and CESM2 SSP5-8.5 runs. Larger median trends in coastal areas than the ocean is a robust pattern for all projections. The projections all point to long-term oxygen declines of order 1 mmol m<sup>-3</sup> per decade through the 21<sup>st</sup> century.

#### 4 Discussion

The results of this study provide a global perspective on ~~some of the~~ climate warming pressures confronting existing and emerging coastal hypoxic areas. The ~~forecasted~~ projected warming and declining oxygen conditions will exert considerable pressure on much of the global coast and on current coastal hypoxic areas. Observed and projected rates along coasts are considerably higher than the open ocean. These differences point to the increased climate vulnerability of coastal regions and the need to focus on coastal conditions separately from open-ocean conditions. Observations indicate the warming and reduced oxygen capacities that coastal waters have been experiencing and the CESM RCP8.5 projection points to even more rapid warming and oxygen declines throughout the 21<sup>st</sup> century. The new analysis indicates the median warming trend in documented hypoxic areas is much (70-75%) stronger than the observed rate and the median trend reported in Altieri and Gedan (2015) for the moderate A1B emissions scenario. The new results suggest that warming-related pressures likely will accelerate the severity of coastal hypoxia around the world. The documented hypoxic areas already experience oxygen concentrations at or below the 63 mmol m<sup>-3</sup> threshold (applied in Diaz and Rosenberg, 2008). If concentrations were at this threshold in 2000, the ~~forecasted~~ projected median trend in oxygen capacity and concentration represents a 20% reduction by 2100. Furthermore, future oxygen declines of any magnitude will have deleterious ecosystem effects as hypoxia worsens (in intensity and duration) in existing hypoxic areas. Such decreases may bottom out at anoxic conditions in some areas; particularly those locations already experiencing oxygen levels considerably below the hypoxic threshold. ~~The forecasted oxygen declines can erode~~

~~oxygen gains achieved in systems improved by wastewater treatment and nutrient management. For example, oxygen conditions in Long Island Sound have improved with reduced nutrient loading, but current and forecasted future warming favor deteriorating oxygen conditions (Whitney and Vlahos, 2021).~~

460 Declines in oxygen conditions along the global coast will create emerging hypoxic areas. ~~Forecasted~~Projected warming and oxygen declines are particularly severe at high latitudes in the northern hemisphere. Two adjacent fjords in Norway (Trysfjord and Ofotfjord) are the only sites above the Arctic circle within the documented hypoxic area database (Dommasnes et al., 1994; Diaz et al., 2011), but it is likely new Arctic coastal hypoxic areas will emerge. For instance, recent observations suggest Jago Lagoon, Alaska may be on the brink of hypoxia (Smith, 2012; Beaufort Lagoon Ecosystems LTER, 2020). High-latitude  
465 waters are colder and therefore tend to have higher ~~oxygen capacities~~saturation concentrations farther from the hypoxic threshold, but oxygen conditions are ~~forecasted~~projected to decline most rapidly in these coastal waters. Emphasis also should be placed on rapidly growing coastal megacities at low and mid-latitudes, which tend to struggle with wastewater treatment infrastructure as populations increase and are likely to experience emerging or worsening hypoxia due to such pressures and warming (von Glasow et al., 2013; Varis et al., 2006).

470 It is important to note that ecosystem problems can arise before conditions deteriorate down to the canonical hypoxic threshold ( $63 \text{ mmol m}^{-3}$ ), as many organisms experience physiological stresses above this threshold and have differing tolerances for low oxygen conditions (Vaquer-Sunyer and Duarte, 2008). In addition to directly reducing oxygen capacities, warming also increases metabolic rates, related biological oxygen demand, and thermal stratification (Brown et al., 2004; Cloern, 2001; Breitburg et al., 2018). Thus, attention should be paid to all coastal areas with lowering oxygen conditions, not just the areas  
475 already experiencing seasonal hypoxia.

The projected oxygen declines can erode oxygen gains achieved in systems improved by wastewater treatment and nutrient management. For example, oxygen concentrations in Long Island Sound have risen with reduced nutrient loading after decades of nitrogen management, but hypoxia still occurs and oxygen conditions would have been better if not for warming-related oxygen capacity decreases (Whitney and Vlahos, 2021). Projections made following essentially the same methods as this study point to warming and deteriorating oxygen conditions that will erode gains made by management (Whitney and Vlahos, 2021). It is noteworthy that the projected rate of oxygen capacity decreases for Long Island Sound is smaller than the trend observed in recent decades. This mismatch may be partially due to the different time periods for the observations and projections and also due to limits in the resolution and dynamics of the CESM RCP8.5 results (Whitney and Vlahos, 2021; discussed further  
480 below). In other systems, hypoxia has not decreased in spite of major nutrient management efforts. The Baltic Sea is a well-studied hypoxic system with large managed reductions in nutrient loads entering from its watersheds. Hypoxic areas, however, have grown in recent decades (Conley et al., 2007; Meier et al., 2019). This study projects warming and decreases in oxygen capacity and concentrations in the Baltic. Projections from a Baltic Sea model point to increasing hypoxia due to warming, increased nutrient loads, and intensified nutrient cycling (Meier et al., 2011). Nitrogen loads have been reduced in the  
485 Mississippi watershed, which feeds eutrophication in the Gulf of Mexico coastal hypoxic zone (Scavia et al. 2019; Giudice et  
490

al., 2020; USGS, 2022). Despite the nitrogen management, hypoxic extent has not consistently decreased and remains well above the management goal (Rabalais and Turner, 2019). Water temperatures in the area have been rising and projections indicate warming will continue to exert a pressure on oxygen conditions (Turner et al., 2016; Guidice et al., 2020; this study). The projections from this study are consistent with a Gulf of Mexico hypoxia study that points to more severe, prolonged, and extensive hypoxia by the end of the century; primarily due to warming-related oxygen solubility reductions (Laurent et al., 2018). Hypoxia in the Chesapeake Bay has not been reduced despite extensive nitrogen management and somewhat decreased nitrogen loads (Murphy et al., 2011; Maryland Department of Natural Resources, 2021; Chesapeake Bay Program, 2022). Moderate oxygen increases tied to load reductions have been overwhelmed by long-term oxygen declines mainly associated with warming (Ni et al., 2020). Projections based on climate downscaling suggest hypoxic volume will increase substantially by mid-century (Ni et al., 2019). In other areas such as the Bohai Sea and Pearl River estuary, nutrient loads have not been reined in by management and hypoxia is worsening in response to increased anthropogenic loads and warming (Qian et al., 2018; Zhai et al., 2019). In general, ameliorating coastal hypoxia through nutrient management has proved challenging. The ongoing and projected warming pressure make efforts to improve coastal oxygen conditions more daunting. Future management efforts should incorporate projected warming-driven oxygen decreases. It would be wise to consider progressively decreasing maximum loads to contend with decreasing oxygen capacities.

Projections from CMIP5 and CMIP6 are widely used and provide valuable information for potential climate scenarios. Earth system models, however, have differences in representing oxygen dynamics. This study has a coastal focus, but it is worth noting that some features of global ocean oxygen patterns in models warrant further investigation. For instance, the Arctic is an area with projected decreases in oxygen capacity and offshore increases in vertical-minimum oxygen concentrations for the CESM RCP8.5 projections. This tendency appears in some of the other CMIP5 models, but not in the CMIP6 SSP5-8.5 projections. The locations and extent of other areas with projected vertical-minimum oxygen increases (mostly in the tropics) vary among models. In most cases, the projected offshore oxygen increases do not reach the coasts. In coastal regions, the vertical-minimum oxygen level is bathymetrically constrained to be closer to the surface and therefore more closely tied to projected oxygen capacity declines. This factor favors closer agreement in coastal areas among models that have similar warming rates. On the other hand, the relatively coarse resolution and global application of Earth system models make it challenging to represent physical and biogeochemical dynamics in coastal waters. ~~The results of this study provide a global perspective on some of the climate pressures confronting existing and emerging coastal hypoxic areas. Observations indicate the warming and reduced oxygen capacities that coastal waters have been experiencing and the CESM forecast for the RCP 8.5 scenario points to even more rapid warming and oxygen declines throughout the 21<sup>st</sup> Century.~~ It is encouraging that the new analysis indicates CESM RCP8.5 coastal performance is broadly consistent with observed SST and oxygen capacities during overlapping years. The ~~forecasted~~ projected and observed coastal oxygen capacities have a similar latitudinal pattern. These coastal results indicate more latitudinal pattern agreement between observations and model results than found for open-ocean oxygen concentrations at the thermocline (Oschlies et al., 2017). The nominal 1° latitude CESM resolution, however, offers only a limited representation of coastal processes. Regional scales of variability are resolved, but smaller scales along

525 continental shelves and within estuaries are not. The analysis pairs each documented hypoxic area with the nearest CESM coastal grid point, but most estuaries are not resolved in the CESM grid. Consequently, the projections may not represent oxygen conditions near the heads of estuaries where hypoxia often occurs. The nearest coastal points reflect conditions in the vicinity of the estuaries. In nature, the influence of surrounding coastal waters on estuarine hypoxic areas is exerted via estuarine exchange (e.g. Kuo et al., 1991; Roegner et al., 2011; Coogan et al., 2021).

530 Research on future coastal oxygen conditions can be advanced with ~~forecasts~~projections from local high-resolution models (Fennel and Testa, 2019), as for systems such as the Gulf of Mexico (Justic et al., 2007), Chesapeake Bay (Ni et al., 2019), and Baltic Sea (Meier et al., 2019). Such regional or estuary-specific models better resolve processes on continental shelves and within estuaries and often are compared to local datasets for oxygen and biogeochemical variables. Further steps forward will come as Earth system models ~~increase~~improve resolution and improve the representation of physical and ecosystem

535 processes in global coastal areas (e.g. Holt et al., 2009; Holt et al., 2017). One way forward is extending the box model approach for estuarine mixing applied in CESM and CESM2 to estuarine biogeochemical cycling (Sun et al., 2017). Other ways forward are nested coastal grids, resolution refinements approaching the coast, or hybrid approaches with higher resolution for some coastal systems and analytical or box model representations in others (Ward et al., 2020). For existing and future Earth system models, the coastal biogeochemical results and representation should be compared to coastal observations and coastal model

540 results where available. Such analyses of biogeochemical performance have been completed for southwest Pacific and northwest Atlantic continental shelf; though oxygen is not included in the analyses (Rickard et al., 2016; Laurent et al., 2021). This approach can be extended to other areas and other biogeochemical variables including oxygen.

This study has focused on warming-related pressures on hypoxia, but it is important to note that the development of hypoxia and long-term changes in its prevalence depend on many factors. As described above, anthropogenic increases in nutrient

545 loading and related management efforts play important roles in hypoxic extent and intensity. Interannual and longer-term variability in river flow directly influence terrestrial nutrient loads entering estuaries and other coastal waters. Such changes can be related to climate controls on storm tracks and precipitation (Altieri and Gedan, 2015). In addition, long-term increases in estuary depths due to sea-level rise and stratification due to warming and intensified freshwater inputs can inhibit ventilation of near-bottom waters and increase hypoxia (Cloern, 2001). Overall, the observed and projected deterioration in coastal oxygen

550 conditions are attributable to nutrient overloading fueling eutrophication and anthropogenic climate change (e.g. Rabalais and Turner, 2001; Paerl, 2006; Diaz and Rosenberg, 2011). The present study contributes by describing global coastal distributions of trends in temperature, oxygen capacity, and vertical-minimum oxygen concentration. Future studies should assess the relative importance of the multitude of stressors exacerbating coastal hypoxia, both regionally and globally. Such studies will be facilitated by advances in Earth system models with biogeochemistry.

## 555 5 Conclusions

Linear trend analyses of a global 40-year observational gridded climate data record (updated from Merchant et al., 2019) and CESM 21<sup>st</sup> ~~c~~Century ~~forecasts~~projections for the RCP-8.5 scenario indicate warming-related pressures on oxygen conditions are increasing. The median trends ~~foreeasted~~projected along the global coast are 0.39 °C, -1.6 mmol m<sup>-3</sup>, and -1.2 mmol m<sup>-3</sup> per decade for SST, surface oxygen capacity (saturation concentration), and vertical-minimum oxygen concentration, respectively. These trends are considerably faster than the median ~~foreeasted~~projected for the entire surface ocean. The ~~foreeasted~~projected median coastal trends for SST and oxygen capacity are 48% and 18% faster than the corresponding observed rates. Significant portions of the global coast (upwards of 189%) are ~~foreeasted~~projected to change even more rapidly, with SST warming more than 0.60 °C per decade and oxygen capacity and concentrations rates faster than -3.0 mmol m<sup>-3</sup> per decade. Observed and ~~foreeasted~~projected rates tend to increase with latitude. Most (89%) of the documented hypoxic areas (Diaz et al., 2011) are in the mid-latitude (20-60 °N) northern hemisphere where trends tend to be larger, but not as extreme as coastal rates in the Arctic. The database documents only a few coastal hypoxic areas at high latitudes, where it is likely that new hypoxic areas will emerge due to warming and rapidly deteriorating oxygen conditions. Coastal megacities (at low and mid-latitudes) are likely to experience emerging or worsening hypoxia with the dual pressures of warming and increasing populations. Projections from other models and other climate scenarios (including CIMP5 RCP4.5 and CMIP6 SSP2-4.5 and SSP5-8.5) point to long-term oxygen decreases of order 1 mmol m<sup>-3</sup> per decade through the 21<sup>st</sup> century and have larger median trends for coastal waters than for the open ocean. The ~~foreeasted~~projected warming and declining oxygen conditions will exert considerable pressure on current hypoxic areas, since future oxygen declines of any magnitude will have deleterious ecosystem effects as hypoxic intensity and duration worsens. The ~~foreeasted~~projected oxygen declines can erode oxygen gains achieved in systems improved by nutrient/wastewater management and should be incorporated into coastal environmental management strategies to protect future water quality and ecosystem services.

**Author contribution:** M.W. completed the research and wrote the manuscript.

**Competing interests:** The author declares that he has no conflict of interest.

**Acknowledgements:** This research was supported by the U.S. Environmental Protection Agency Long Island Sound Study (LISS01719).

## 580 References

Altieri, A. H. and Gedan, K. B.: Climate change and dead zones. *Global Change Biol.*, 21, 1395–1406, <https://doi.org/10.1111/gcb.12754>, 2015.

- 585 Bopp, L., Resplandy, L., Orr, J. C., Doney, S. C., Dunne, J. P., Gehlen, M., Halloran, P., Heinze, C., Ilyina, T., Séférian, R., Tjiputra, J., and Vichi, M.: Multiple stressors of ocean ecosystems in the 21st century: Projections with CMIP5 models. *Biogeosciences*, 10, 6225–6245, <https://doi.org/10.5194/bg-10-6225-2013>, 2013.
- [Bernal, D., J. P. Reid, J. M. Roessig, S. Matsumoto, C. A. Sepulveda, J. J. Cech, and J. B. Graham: Temperature effects on the blood oxygen affinity in sharks. \*Fish Physiol. Biochem.\*, 44, 949–967. <https://doi.org/10.1007/s10695-018-0484-2>, 2018.](#)
- [Black, E. C.: The transport of oxygen by the blood of freshwater fish. \*Biological Bull.\*, 79\(1\), 215-229. <https://doi.org/10.2307/1537841>, 1940.](#)
- 590 [Boucher, O., Servonnat, J., Albright, A. L., Aumont, O., Balkanski, Y., Bastrikov, V., Bekki, S., Bonnet, R., Bony, S., Bopp, L., Braconnot, P., Brockmann, P., Cadule, P., Caubel, A., Cheruy, F., Codron, F., Cozic, A., Cugnet, D., D’Andrea, F., Davini, P., Lavergne, C. de, Denvil, S., Deshayes, J., Devilliers, M., Ducharne, A., Dufresne, J.-L., Dupont, E., Éthé, C., Fairhead, L., Falletti, L., Flavoni, S., Foujols, M.-A., Gardoll, S., Gastineau, G., Ghattas, J., Grandpeix, J.-Y., Guenet, B., Guez, L., Guilyardi, É., Guimberteau, M., Hauglustaine, D., Hourdin, F., Idelkadi, A., Joussaume, S., Kageyama, M.,](#)
- 595 [Khodri, M., Krinner, G., Lebas, N., Levvasseur, G., Lévy, C., Li, L., Lott, F., Lurton, T., Luyssaert, S., Madec, G., Madeleine, J.-B., Maignan, F., Marchand, M., Marti, O., Mellul, L., Meurdesoif, Y., Mignot, J., Musat, I., Ottlé, C., Peylin, P., Planton, Y., Polcher, J., Rio, C., Rochetin, N., Rousset, C., Sepulchre, P., Sima, A., Swingedouw, D., Thiéblemont, R., Traore, A. K., Vancoppenolle, M., Vial, J., Vialard, J., Viovy, N. and Vuichard, N.: Presentation and evaluation of the IPSL-CM6A-LR climate model, \*J. Adv. Model. Earth Syst.\*, 12, e2019MS002010, <https://doi.org/10.1029/2019MS002010>, 2020.](#)
- 600 Beaufort Lagoon Ecosystems LTER: *Physiochemical water column parameters and hydrographic time series from river, lagoon, and open ocean sites along the Alaska Beaufort Sea coast*, 2018-ongoing ver 1. Environmental Data Initiative, <https://doi.org/10.6073/pasta/e0e71c2d59bf7b08928061f546be6a9a>, 2020.
- Breitburg, D., Levin, L. A., Oschlies, A., Grégoire, M., Chavez, F. P., Conley, D. J., Garçon, V., Gilbert, D., Gutiérrez, D., Isensee, K., Jacinto, G. S., Limburg, K. E., Montes, I., Naqvi, S. W. A., Pitcher, G. C., Rabalais, N. N., Roman, M. R., Rose,
- 605 K. A., Seibel, B. A., Telszewski, M., Yasuhara, M., and Zhang, J.: Declining oxygen in the global ocean and coastal waters. *Science*, 359(6371), p.eaam7240, <https://doi.org/10.1126/science.aam7240>, 2018.
- Brown, J. H., Gillooly, J. F., Allen, A. P., Savage, V. M., and West, G. B.: Toward a metabolic theory of ecology. *Ecology*, 85, 1771–1789, <https://doi.org/10.1890/03-9000>, 2004.
- 610 [Chesapeake Bay Program: State of the Chesapeake: Pollution, <https://www.chesapeakebay.net/state/pollution>, accessed 2022.](#)
- Cloern, J. E.: Our evolving conceptual model of the coastal eutrophication problem. *Mar. Ecol. Prog. Ser.*, 210, 223–253, <https://doi.org/10.3354/meps210223>, 2001.



- 615 Cocco, V., Joos, F., Steinacher, M., Frölicher, T. L., Bopp, L., Dunne, J., Gehlen, M., Heinze, C., Orr, J., Oschlies, A., Schneider, B., Segschneider, J., and Tjiputra, J.: Oxygen and indicators of stress for marine life in multi-model global warming projections, *Biogeosciences*, 10, 1849-1868, <https://doi.org/10.5194/bg-10-1849-2013>, 2013.
- Collins, W. D., Bitz, C. M., Blackmon, M. L., Bonan, G. B., Bretherton, C. S., Carton, J. A., Chang, P., Doney, S. C., Hack, J. J., Henderson, T. B., Kiehl, J. T., Large, W. G., McKenna, D. S., Santer, B. D., and Smith, R. D.: The Community Climate System Model Version 3 (CCSM3). *J. Climate*, 19, 2122-2143, <https://doi.org/10.1175/JCLI3761.1>, 2006.
- 620 [Collins, W. J., Bellouin, N., Doutriaux-Boucher, M., Gedney, N., Halloran, P., Hinton, T., Hughes, J., Jones, C. D., Joshi, M., Liddicoat, S., Martin, G., O'Connor, F., Rae, J., Senior, C., Sitch, S., Totterdell, I., Wiltshire, A., and Woodward, S.: Development and evaluation of an Earth-System model – HadGEM2. \*Geosci. Model Dev.\*, 4, 1051–1075, <https://doi.org/10.5194/gmd-4-1051-2011>, 2011.](#)
- [Conley, D. J., Carstensen, J., Ærtebjerg, G., Christensen, P. B., Dalsgaard, T., Hansen, J. L. S., Josefson, A. B.: Long-term changes and impacts of hypoxia in Danish coastal waters. \*Ecol. Appl.\*, 17 \(5\) Supplement, S165–S184, <https://doi.org/10.1890/05-0766.1>, 2007.](#)
- 625 [Coogan, J., Dzwonkowski, B., Lehrter, J., Park, K., and Collini, R. C.: Observations of dissolved oxygen variability and physical drivers in a shallow highly stratified estuary. \*Estuar. Coast. Shelf Sci.\*, 259, 107482, <https://doi.org/10.1016/j.ecss.2021.107482>, 2021.](#)
- [Danabasoglu, G., Lamarque, J. F., Bacmeister, J., Bailey, D. A., DuVivier, A. K., Edwards, J., Emmons, L. K., Fasullo, J., Garcia, R., Gettelman, A., Hannay, C., Holland, M. M., Large, W. G., Lauritzen, P. H., Lawrence, D. M., Lenaerts, J. T. M., Lindsay, K., Lipscomb, W. H., Mills, M. J., Neale, R., Oleson, K. W., Otto-Bliesner, B., Phillips, A. S., Sacks, W., Tilmes, S., van Kampenhout, L., Vertenstein, M., Bertini, A., Dennis, J., Deser, C., Fischer, C., Fox-Kemper, B., Kay, J. E., Kinnison, D., Kushner, P. J., Larson, V. E., Long, M. C., Mickelson, S., Moore, J. K., Nienhouse, E., Polvani, L., Rasch, P. J., Strand, W. G.: The Community Earth System Model version 2 \(CESM2\). \*J. Adv. Model. Earth Syst.\*, 12\(2\), \[e2019MS001916. https://doi.org/10.1029/2019MS001916\]\(https://doi.org/10.1029/2019MS001916\), 2020.](#)
- 630 [Deignan-Schmidt, S. R. and Whitney, M. M.: A model study on the summertime distribution of river waters in Long Island Sound. \*Estuaries and Coasts\*, 41\(4\), <https://doi.org/10.1007/s12237-017-0348-5>, 1002-1020.](#)
- Diaz, R., Selman, M. and Chique, C.: *Global Eutrophic and Hypoxic Coastal Systems*. World Resources Institute. Eutrophication and Hypoxia: Nutrient Pollution in Coastal Waters. Available online at:
- 640 <https://datasets.wri.org/dataset/eutrophication-hypoxia-map-data-set>, 2011.
- Diaz R. J., and Rosenberg, R.: Spreading dead zones and consequences for marine ecosystems. *Science*, 321, 926–929, <https://doi.org/10.1126/science.1156401>, 2008.

- Diaz, R.J. and Rosenberg, R.: Introduction to environmental and economic consequences of hypoxia. *Int. J. Water Resour. D.*, 27, 71-82, <https://doi.org/10.1080/07900627.2010.531379>, 2011.
- 645 Dommasnes, A., Rey, F., and Røttingen, I.: Reduced oxygen concentrations in herring wintering areas. *ICES J. Mar. Sci.*, 51, 63-69, <https://doi.org/10.1006/jmsc.1994.1006>, 1994.
- [Dufresne, J.-L., Foujols, M.-A., Denvil, S., Caubel, A., Marti, O., Aumont, O., Balkanski, Y., Bekki, S., Bellenger, H., Benschila, R., Bony, S., Bopp, L., Braconnot, P., Brockmann, P., Cadule, P., Cheruy, F., Codron, F., Cozic, A., Cugnet, D., Noblet, N., Duvel, J.-P., Ethé, C., Fairhead, L., Fichet, T., Flavoni, S., Friedlingstein, P., Grandpeix, J.-Y., Guez, L., Guilyardi, E., Hauglustaine, D., Hourdin, F., Idelkadi, A., Ghattas, J., Joussaume, S., Kageyama, M., Krinner, G., Labetoulle, S., Lahellec, A., Lefebvre, M.-P., Lefevre, F., Levy, C., Li, Z. X., Lloyd, J., Lott, F., Madec, G., Mancip, M., Marchand, M., Masson, S., Meurdesoif, Y., Mignot, J., Musat, I., Parouty, S., Polcher, J., Rio, C., Schulz, M., Swingedouw, D., Szopa, S., Talandier, C., Terray, P., Viovy, N., and Vuichard, N.: Climate change projections using the IPSL-CM5 Earth System Model: from CMIP3 to CMIP5, \*Clim. Dynam.\*, 40, 2123–2165, <https://doi.org/10.1007/s00382-012-1636-1>, 2013.](#)
- 655 Embury, O. and Good, S.: Sea surface temperature daily data from 1981 to present derived from satellite observations, Level-4 Product. *Climate Data Store*, <https://cds.climate.copernicus.eu/cdsapp#!/dataset/satellite-sea-surface-temperature>, 2021.
- Fennel, K. and Testa, J. M.: Biogeochemical controls on coastal hypoxia. *Ann. Rev. Mar. Sci.*, 11, 105-130, <https://doi.org/10.1146/annurev-marine-010318-095138>, 2019.
- 660 Garcia, H. E., and Gordon, L. I.: Oxygen solubility in seawater: better fitting equations. *Limnol. Oceanogr.*, 37, 1307-1312, <https://doi.org/10.4319/lo.1992.37.6.1307>, 1992.
- [Giorgetta, M. A., Jungclaus, J., Reick, C. H., Legutke, S., Bader, J., Böttinger, M., Brovkin, V., Crueger, T., Esch, M., Fieg, K., Glushak, K., Gayler, V., Haak, H., Hollweg, H.-D., Ilyina, T., Kinne, S., Kornblueh, L., Matei, D., Mauritsen, T., Mikolajewicz, U., Mueller, W., Notz, D., Pithan, F., Raddatz, T., Rast, S., Redler, R., Roeckner, E., Schmidt, H., Schnur, R., Segschneider, J., Six, K. D., Stockhause, M., Timmreck, C., Wegner, J., Widmann, H., Wieners, K.-H., Claussen, M., Marotzke, J., and Stevens, B.: Climate and carbon cycle changes from 1850 to 2100 in MPI-ESM simulations for the Coupled Model Intercomparison Project phase 5, \*J. Adv. Model. Earth Syst.\*, 5, 572–597, <https://doi.org/10.1002/jame.20038>, 2013.](#)
- 665 Gilbert, D., Rabalais, N. N., Diaz, R. J., Zhang, J.: Evidence for greater oxygen decline rates in the coastal ocean than in the open ocean. *Biogeosciences Discuss.*, 7, 2283-2296, <https://doi.org/10.5194/bgd-6-9127-2009>, 2010.
- [Giudice, D. D., Matli, V. R. R., Obenour, D. R.: Bayesian mechanistic modeling characterizes Gulf of Mexico hypoxia: 1968-2016 and future scenarios, \*Ecol. Appl.\*, 30 \(2\), e02032, <https://doi.org/10.1002/eap.2032>, 2020.](#)

- Haldane, J. and Smith, J. L.: The mass and oxygen capacity of the blood in man. *J. Physiol.*, 25(5), 331, <https://doi.org/10.1113/jphysiol.1900.sp000800>, 1900.
- 675 Helm, K. P., Bindoff, N. L., and Church, J. A.: Observed decreases in oxygen content of the global ocean. *Geophys. Res. Lett.*, 38(23), <https://doi.org/10.1029/2011GL049513>, 2011.
- Hoegh-Guldberg, O., Jacob, D., Taylor, M., Bindi, M., Brown, S., Camilloni, I., Diedhiou, A., Djalante, R., Ebi, K. L., Engelbrecht, F., Guiot, J., Hijoka, Y., Mehrotra, S., Payne, A., Seneviratne, S. I., Thomas, A., Warren, R., Zhou, G., and Tschakert, P.: Impacts of 1.5°C Global Warming on Natural and Human Systems. In: *Global Warming of 1.5°C*, [Masson-Delmotte, V., Zhai, P., Pörtner, H. -O., Roberts, D., Skea, J., Shukla, P. R., Pirani, A., Moufouma-Okia, W., Péan, C., Pidcock, R., Connors, S., Matthews, J. B. R., Chen, Y., Zhou, X., Gomis, M. I., Lonnoy, E., Maycock, T., Tignor, M., and Waterfield, T. (eds.)], Geneva, Switzerland: World Meteorological Organization, <https://www.ipcc.ch/sr15/chapter/chapter-3/>, 2018.
- 680 Holt, J., Harle, J., Proctor, R., Michel, S., Ashworth, M., Batstone, C., Allen, I., Holmes, R., Smyth, T., Haines, K., Bretherton, D., and Smith, G.: Modelling the global coastal ocean. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 367(1890), 939-951, <https://doi.org/10.1098/rsta.2008.0210>, 2009.
- Holt, J., Hyder, P., Ashworth, M., Harle, J., Hewitt, H. T., Liu, H., New, A. L., Pickles, S., Porter, A., Popova, E., Allen, J. I., Siddorn, J., and Wood, R.: Prospects for improving the representation of coastal and shelf seas in global ocean models. *Geosci. Model Dev.*, 10, 499-523, <https://doi.org/10.5194/gmd-10-499-2017>, 2017.
- 690 Justic, D., Bierman, V. J., Scavia, D., and R. D. Hetland, R. D.: Forecasting Gulf's Hypoxia: The Next 50 Years? *Estuar. Coasts*, 30, 791-801, <https://doi.org/10.1007/BF02841334>, 2007.
- Kay, J. E., Deser, C., Phillips, A., Mai, A., Hannay, C., Strand, G., Arblaster, J., Bates, S., Danabasoglu, G., Edwards, J., Holland, M., Kushner, P., Lamarque, J. -F., Lawrence, D., Lindsay, K., Middleton, A., Munoz, E., Neale, R., Oleson, K., Polvani, L., and Vertenstein, M.: The Community Earth System Model (CESM) Large Ensemble Project: A Community Resource for Studying Climate Change in the Presence of Internal Climate Variability, *B. Am. Meteorol. Soc.*, 96, 1333-1349. <https://doi.org/10.1175/BAMS-D-13-00255.1>, 2015a.
- 695 Kay, J. E., Deser, C., Phillips, A., Mai, A., Hannay, C., Strand, G., Arblaster, J., Bates, S., Danabasoglu, G., Edwards, J., Holland, M., Kushner, P., Lamarque, J. -F., Lawrence, D., Lindsay, K., Middleton, A., Munoz, E., Neale, R., Oleson, K., Polvani, L., and Vertenstein, M.: CESM1 CAM5 BGC 20C + RCP8.5 Large Ensemble, *Earth System Grid*, <https://www.earthsystemgrid.org/dataset/ucar.cgd.cesm4.cesmLE.html>, 2015b.
- 700 Kuo, A. Y., Park, K. and Moustafa, M. Z.: Spatial and temporal variabilities of hypoxia in the Rappahannock River, Virginia. *Estuaries*, 14, 113–121. <https://doi.org/10.2307/1351684>, 1991.

- Laurent, A.; Fennel, K., Ko, D. S., Lehrter, J.: Climate Change Projected to Exacerbate Impacts of Coastal Eutrophication in the Northern Gulf of Mexico. *J. Geophys. Res.: Oceans*, 123, 3408-3426. <https://doi.org/10.1002/2017JC013583>, 2018.
- 705 Laurent, A., Fennel, K. and Kuhn, A.: An observation-based evaluation and ranking of historical Earth system model simulations in the northwest North Atlantic Ocean. *Biogeosciences*, 18(5), 1803-1822. <https://doi.org/10.5194/bg-18-1803-2021>, 2021.
- Maryland Department of Natural Resources, Chesapeake Bay Hypoxia Report – Year-End 2021. <https://news.maryland.gov/dnr/2021/11/30/chesapeake-bay-hypoxia-report-year-end-2021/>, 2021.
- 710 Maio, D. A. and J. R. Neville: Polarographic determination of oxygen content and capacity in a single blood sample. *J. Appl. Physiol.*, 20(4), 774-778. <https://doi.org/10.1152/jappl.1965.20.4.774>, 1965.
- Mauritsen, T., Bader, J., Becker, T., Behrens, J., Bittner, M., Brokopf, R., Brovkin, V., Claussen, M., Crueger, T., Esch, M., and Fast, I.: Developments in the MPI-M Earth System Model version 1.2 (MPI-ESM1.2) and its response to increasing CO<sub>2</sub>. *J. Adv. Model. Earth Syst.*, 11, 998–1038. <https://doi.org/10.1029/2018MS001400>, 2019.
- 715 Meier, H. E. M., Eilola, K., Almroth-Rosell, E., Schimanke, S., Kniebusch, M., Höglund, A., Pemberton, P., Liu, Y., Väli, G., Saraiva, S.: Disentangling the impact of nutrient load and climate changes on Baltic Sea hypoxia and eutrophication since 1850. *Clim. Dynam.*, 53, 1145–1166. <https://doi.org/10.1007/s00382-018-4296-y>, 2019.
- Meier, H. E. M., Andersson, H. C., Eilola, K., Gustafsson, B. G., Kuznetsov, I., Müller-Karulis, B., Neumann, T., Savchuk, O. P.: Hypoxia in future climates: a model ensemble study for the Baltic Sea. *Geophys. Res. Lett.*, 38, L24608. <https://doi.org/10.1029/2011GL049929>, 2011.
- 720 Melillo, J. M., Richmond, T. C., and Yohe, G. W. (eds.): *Climate Change Impacts in the United States: The Third National Climate Assessment*. U.S. Global Change Research Program, 841 pp. <https://doi.org/10.7930/J0Z31WJ2>, 2014.
- Merchant, C. J., O. Embury, C. E. Bulgin, T. Block, G. K. Corlett, E. Fiedler, S. A. Good, J. Mittaz, N. A. Rayner, D. Berry, S. Eastwood, M. Taylor, Y. Tsushima, A. Waterfall, R. Wilson, and C. Donlon: Satellite-based time-series of sea surface temperature since 1981 for climate applications. *Nature, Scientific Data*, 6:223, 18 pp., <https://doi.org/10.1038/s41597-019-0236-x>, 2019.
- 725 Moss, R. H., Edmonds, J. A., Hibbard, K. A., Manning, M. R., Rose, S. K., van Vuuren, D. P., Carter, T. R., Emori, S., Kainuma, M., Kram, T., Meehl, G. A., Mitchell, J. F. B., Nakicenovic, N., Riahi, K., Smith, S. J., Stouffer, R. J., Thomson, A. M., Weyant, J. P., and Wilbanks, T. J.: The next generation of scenarios for climate change research and assessment. *Nature*, 463(7282), 747-756. <https://doi.org/10.1038/nature08823>, 2010.
- Murphy, R. R., Kemp, W. M., and Ball, W. P.: Long-Term Trends in Chesapeake Bay Seasonal Hypoxia, Stratification, and Nutrient Loading. *Estuaries Coast.*, 34, 1293–1309. <https://doi.org/10.1007/s12237-011-9413-7>, 2011.

- Nakicenovic, N. and Swart, R. (eds.): *Emission Scenarios: Intergovernmental Panel on Climate Change Special Report on Emission Scenarios*, Cambridge University Press, 570 pp.,  
735 [https://www.ipcc.ch/site/assets/uploads/2018/03/emissions\\_scenarios-1.pdf](https://www.ipcc.ch/site/assets/uploads/2018/03/emissions_scenarios-1.pdf), 2000.
- [Ni, W., Li, M., Testa, J. M.: Discerning effects of warming, sea level rise and nutrient management on long-term hypoxia trends in Chesapeake Bay. \*Sci. Total Environ.\*, 737, 139717, <https://doi.org/10.1016/j.scitotenv.2020.139717>, 2020.](#)
- Ni, W., M. Li, M., Ross, A. C., and Najjar, R. G.: Large Projected Decline in Dissolved Oxygen in a Eutrophic Estuary Due to Climate Change. *J. Geophys. Res.: Oceans*, 124, 8271-8289, <https://doi.org/10.1029/2019JC015274>, 2019.
- 740 Oeschies, A., Duteil, O., Getzlaff, J., Koeve, W., Landolfi, A., Schmidtko, S.: Patterns of deoxygenation: sensitivity to natural and anthropogenic drivers. *Phil. Trans. R. Soc. A*, 375, 20160325. <https://doi.org/10.1098/rsta.2016.0325>, 2017.
- Paerl, H. W.: Assessing and managing nutrient-enhanced eutrophication in estuarine and coastal waters: Interactive effects of human and climatic perturbations. *Ecol. Eng.*, 26, 40-54, <https://doi.org/10.1016/j.ecoleng.2005.09.006>, 2006.
- Pershing, A.J., Alexander, M. A., Hernandez, C. M., Kerr, L. A., Le Bris, A., Mills, K. E., Nye, J. A., Record, N. R.,  
745 Scannell, H. A., Scott, J. D., Sherwood, G. D., and Thomas, A. C.: Slow adaptation in the face of rapid warming leads to collapse of the Gulf of Maine cod fishery. *Science*, 350(6262), 809-812, <https://doi.org/10.1126/science.aac9819>, 2015.
- [Qian, W., Gan, J., Liu, J., He, B., Lu, Z., Guo, X., Wang, D., Guo, L., Huang, T., Dai, M.: Current status of emerging hypoxia in a eutrophic estuary: The lower reach of the Pearl River Estuary, China. \*Estuar. Coast. Shelf Sci.\*, 205, 58-67, <https://doi.org/10.1016/j.ecss.2018.03.004>, 2018.](#)
- 750 Rabalais, N.N. and Turner, R. E. (Eds.): *Coastal Hypoxia: Consequences for Living resources and Ecosystems*. Coastal and Estuarine Studies 58. American Geophysical Union, Washington, DC, 454 pp., <https://doi.org/10.1029/CE058>, 2001.
- Rabalais, N. N., Diaz, R. J., Levin, L. A., Turner, R. E., Gilbert, D., and Zhang, J.: Dynamics and distribution of natural and human-caused hypoxia. *Biogeosciences*, 7, 585–619, <https://doi.org/10.5194/bg-7-585-2010>, 2010.
- [Rabalais, N. N. and Turner, R. E.: Gulf of Mexico Hypoxia: Past, Present, and Future. \*Limnol. Oceanogr. Bull.\*, 28\(4\), 117-124, <https://doi.org/10.1002/lob.10351>, 2019.](#)
- [Rickard, G. J., Behrens, E., and Chiswell, S. M.: CMIP5 earth system models with biogeochemistry: An assessment for the southwest Pacific Ocean, \*J. Geophys. Res.: Oceans\*, 121\(10\), 7857–7879, <https://doi.org/10.1002/2016JC011736>, 2016.](#)
- [Roegner, G. C., Needoba, J. A., Baptista, A. M.: Coastal upwelling supplies oxygen-depleted water to the Columbia River estuary. \*PloS one\*, 6\(4\), e18672, <https://doi.org/10.1371/journal.pone.0018672>, 2011.](#)
- 760 [Scavia, D.; Justić, D.; Obenour, D.R.; Craig, J.K.; Wang, L. Hypoxic volume is more responsive than hypoxic area to nutrient load reductions in the northern Gulf of Mexico—and it matters to fish and fisheries. \*Environ. Res. Lett.\*, 14 \(2\), p.024012, <https://doi.org/10.1088/1748-9326/aaf938>, 2019.](#)

- Smith, S. D.: *Chemical Parameters of Northern Alaska Coastal Lagoons*. Unpublished term paper for CE 394K: GIS in Water Resources (D. Maidment), University of Texas, 23 pp,  
765 <https://www.caee.utexas.edu/prof/maidment/giswr2012/TermPaper/Smith.pdf>, 2012.
- <https://doi.org/10.1016/j.ocemod.2017.03.004>, 139-153.
- <https://doi.org/10.5194/gmd-12-4823-2019>, 2019.
- 770 Taylor, K. E., Stouffer, R. J., and Meehl, G. A.: An overview of CMIP5 and the experiment design. *B. Am. Meteorol. Soc.*, 93, 485-498, <https://doi.org/10.1175/BAMS-D-11-00094.1>, 2012.
- 775 Turner, R. E., Rabalais, N. N., Justic, D.: Summer bottom-water temperature trends on the northern Gulf of Mexico continental shelf, 1985 to 2015. *PLoS One*, 12, e0184350, <https://doi.org/10.1371/journal.pone.0184350>, 2017.
- [https://nrtwq.usgs.gov/mississippi\\_loads/#/GULF](https://nrtwq.usgs.gov/mississippi_loads/#/GULF), accessed 2022.
- Vaquer-Sunyer, R. and Duarte, C. M.: Thresholds of hypoxia for marine biodiversity, *PNAS*, 105, 15452–15457, <https://doi.org/10.1073/pnas.0803833105>, 2008.
- 780 Varis, O., Biswas, A. K., Tortajada, C., and Lundqvist, J.: Megacities and Water Management, *Water Res. Dev.*, 22:2, 377-394, <https://doi.org/10.1080/07900620600684550>, 2006.
- von Glasow, R., Jickells, T. D., Baklanov, A., Carmichael, G. R., Church, T. M., Gallardo, L., Hughes, C., Kanakidou, M., Liss, P. S., Mee, L. and Raine, R.: Megacities and large urban agglomerations in the coastal zone: interactions between atmosphere, land, and marine ecosystems. *Ambio*, 42(1), 13-28, <https://doi.org/10.1007/s13280-012-0343-9>, 2013.
- 785 Ward, N. D., Megonigal, J. P., Bond-Lamberty, B., Bailey, V. L., Butman, D., Canuel, E.A., Diefenderfer, H., Ganju, N. K., Goñi, M. A., Graham, E. B. and Hopkinson, C. S., Khangaonkar, T., Langley, J. A., McDowell N. G., Myers-Pigg, A. N., Neumann, R. B., Osburn, C. L., Price, R. M., Rowland, J., Sengupta, A., Simard, M., Thornton, P. E., Tzortziou, M., Vargas, R., Weisenhorn, P. B., Windham-Myers, L.: Representing the function and sensitivity of coastal interfaces in Earth system models. *Nat. Commun.*, 11, 2458, <https://doi.org/10.1038/s41467-020-16236-2>, 2020.
- 790 Weiss R. F.: Solubility of nitrogen, oxygen and argon in water and seawater. *Deep-Sea Res.*, 17, 721–735, [https://doi.org/10.1016/0011-7471\(70\)90037-9](https://doi.org/10.1016/0011-7471(70)90037-9), 1970.

Whitney, M. M.: Supporting dataset for observed and ~~foreeasted~~projected global warming pressure on coastal hypoxia. Department of Marine Sciences, *OpenCommons@UConn*, [https://opencommons.uconn.edu/marine\\_sci/13](https://opencommons.uconn.edu/marine_sci/13), 2021.

795

Zhai, W. D., Zhao, H. D., Su, J. L., Liu, P. F., Li, Y. W., Zheng, N.: Emergence of summertime hypoxia and concurrent carbonate mineral suppression in the central Bohai Sea, China. *J. Geophys. Res. Biogeosciences*, 124 (9), 2768-2785, <https://doi.org/10.1029/2019JG005120>, 2019.