SMOS L-VOD shows that post-fire recovery of dense forests is slower than what is depicted with X- and C-VOD and optical indices

Emma Bousquet¹, Arnaud Mialon¹, Nemesio Rodriguez-Fernandez¹, Stéphane Mermoz², and Yann Kerr¹

 ¹Centre d'Etudes Spatiales de la Biosphère (CESBIO), Université de Toulouse (CNES/CNRS/INRAE/IRD/UPS), 18 av. Edouard Belin, bpi 2801, 31401 Toulouse CEDEX 9, France
 ²GlobEO, 31400 Toulouse, France

Correspondence to: Emma Bousquet (emma.bousquet@cesbio.cnes.fr)

Abstract. Anthropogenic climate change is now considered to be one of the main factors causing an increase in both frequency and severity of wildfires. These fires are prone to release substantial quantities of CO_2 in the atmosphere and to endanger natural ecosystems and biodiversity. Depending on the ecosystem and climate regime, fires have distinct triggering factors and impacts. To better analyse and describe fire impact on different biomes, we investigated pre and post fire vegetation anomalies at global scale. The study was performed using several remotely sensed quantities ranging from visible-infrared vegetation indices (the enhanced vegetation index (EVI)) to vegetation opacities obtained at several passive

15 microwave wavelengths (X-band, C-band, and L-band vegetation optical depth (X-VOD, C-VOD, and L-VOD)), ranging from 2 to 20 cm. It was found that C- and X-VOD are mostly sensitive to fire impact on low vegetation areas (grass and small bushes) or on tree leaves; while L-VOD depicts better the fire impact on tree trunks and branches. As a consequence, L-VOD is probably a better way of assessing fire impact on biomass. The study shows that L-VOD can be used to monitor fire affected areas as well as post-fire recovery, especially over densely vegetated areas.

20 1 Introduction

Fires are a natural part of many ecosystems, being historically triggered by lightning strikes (de Groot et al., 2013). Nevertheless, most wildfires are now ignited by human activities (95% in the Mediterranean basin, 85% in Asia and South America; FAO, 2006). In recent years, and in spite of various efforts, wildfires were proven to increase both in frequency and in severity worldwide, largely due to anthropogenic climate change and human pressure (Weber and Stocks, 1998; Jin et

- 25 al., 2012). The 2020 fire season became historically significant in southern Australia and in western US, linked with extreme vegetation dryness (Higuera and Abatzoglou, 2020). Summer 2021 saw an unprecedented number of fires around the Mediterranean Sea, in Siberia and in North America (CAMS, 2021). In tropical rainforests, the Amazon in particular, wildfires have become increasingly prevalent over the past decades due to more frequent droughts and periodic El Niño events (Aragão et al., 2018; Chen et al., 2013; Cochrane, 2003), but also to selective logging and deforestation, that lead to
- 30 forest desiccation and reduce rainfall (Asner et al., 2010).

Wildfire likelihood factors were categorized into climatic (e.g. precipitation, temperature, air humidity, wind speed), topographic, in situ, historical, and anthropogenic factors (Mhawej et al., 2015). Drought, i.e. the concomitant increase of air dryness and decrease of soil moisture, was identified as the most significant fire likelihood factor (Ray et al., 2005). Indirectly, drought also causes vegetation drying, leaf shedding, and branch losses, which increases forest flammability

- 35 (Nepstad et al., 2001; Chuvieco et al., 2012). Surveying the soil moisture (SM) and the vegetation water content (VWC) could then be a good indicator for fire risk detection, and passive microwave remote sensing is a useful tool for that. Indeed, the SM deficit monitored with AMSR-E was previously proven to be a major driving factor for the evolution of extreme fire events in Siberia (Forkel et al., 2012). GRACE-assimilated SM was also exploited for fire risk assessment in the United States (Jensen et al., 2018; Farahmand et al., 2020). SMOS SM anomalies have been found to explain singular fire episodes
- 40 in the northwestern Iberian Peninsula (Chaparro et al., 2016) and in Canada (Ambadan et al., 2020). SMOS SM has been used as an alternative source of moisture information in the McArthur Forest Fire Danger Index (FFDI; Holgate et al., 2017). Finally, AMSR-E vegetation optical depth (VOD) was successfully used in data-driven fire models (Forkel et al., 2017; Kuhn-Régnier et al., 2021).
- In addition to endangering populations, wildlife, ecosystems, and to releasing overwhelming quantities of CO2 in the 45 atmosphere (CAMS, 2021), wildfires have several negative effects on soil and vegetation properties. They cause deterioration of soil structure and porosity, ash entrapment, removal of organic matter and nutrients, decreasing of microbial and invertebrate communities, etc. (Certini, 2005). Plant cover removal also increases soil water repellency and runoff, which can lead to floods and erosion (Shakesby and Doerr, 2005). Post-fire vegetation regeneration highly depends on the ecosystem and on the fire severity (Chu and Guo, 2013). In humid tropical forests, the Amazon in particular, wildfires can
- 50 significantly reduce above-ground biomass (AGB) for decades by amplifying tree mortality (Barlow et al., 2003; Silva et al., 2018; de Faria et al., 2021). Conversely, some ecosystems can recover much faster. For instance, some coniferous trees (e.g. jack pine, black spruce) evolved to become fire resistant and to use the flames as a means for spreading their seeds, as the heat causes the opening of cones (Weber and Stocks, 1998). Some eucalyptus communities of South-East Australia are also able to survive fire by activating dormant vegetative buds to produce regrowth (Heath et al., 2016). In savannas, recurrent
- 55 seasonal fires help maintaining the structure, species composition, and biological diversity (Menaut et al., 1990). In forests, prescribed burning enables to reduce hazardous accumulations of fuel, and thus to mitigate the severity of wildfires (Sackett, 1975). Fires can even be necessary for canopy regeneration: a decline of sequoias population was observed when fires were suppressed in California (Parsons and DeBenedetti, 1979). Vegetation can thus recover from fire and if plants succeed in promptly recolonising the burnt area, the pre-fire level of most properties can be recovered and even enhanced (Certini, 60 2005).

Post-fire vegetation conditions are then essential to monitor, and satellite remote sensing proved its abilities for that in addition to field campaigns (Chu and Guo, 2013). Indicators and metrics based on multispectral satellite imagery (visible and infrared) are the most frequently used, such as the normalized difference vegetation index (NDVI), the enhanced vegetation index (EVI), and the normalized burned ratio (NBR) (Pérez-Cabello et al., 2021). Despite their fast saturation over dense

- 65 forests, they still provide a good proxy for green vegetation regrowth. Microwave data also shown a good potential to monitor post-fire recovery. L-band SAR was used to assess forest regrowth in South-East Asia (Mermoz and Le Toan, 2016); and to estimate the tree survival in eucalyptus forests of Western Australia (Fernandez-Carrillo et al., 2019). C-band VOD was used to analyse the Amazon canopy dynamics during the 2019 fire season (Zhang et al., 2021). Authors found a lower magnitude of canopy damage and a longer recovery period for C-VOD than for optical-based indices (NDVI, EVI, NBR).
- 70 Indeed, the optical-based indices only represent the canopy greenness, whereas microwave measurements are more sensitive to woody components (Guglielmetti et al., 2007; Frappart et al., 2020). Microwave VODs are also sensitive to VWC and can help to monitor the biomass status (Fan et al., 2018; Konings et al., 2019).

With the arrival of L-band radiometers such as the Soil Moisture and Ocean Salinity (SMOS) satellite, it is now possible to infer surface soil moisture, biomass (i.e. fuel) and its water content at deeper sensing depth. The rationale of this study is to

- 75 investigate how L-band radiometry responds to fire events in various ecosystems and climates. The SMOS satellite has been operating for over 12 years now and we have access to a large catalogue of major fires. This study also presents for the first time L-VOD used in conjunction with other sensors, from visible-infrared (EVI) to microwave X- and C-VOD, for the study of post-fire vegetation recovery. The complementarity of these vegetation variables along with climate variables (air temperature (T), precipitation (P), soil moisture (SM), and terrestrial water storage (TWS)) was used to identify the fire
- 80 likelihood factors and the immediate and long-term fire impacts on vegetation. To do this, we first observed three particular cases of large fires in various environments and then extended the analysis to the global scale.

2 Data

2.1 Fires

Fires were obtained from the National Aeronautics and Space Administration (NASA) MODerate resolution Imaging Spectroradiometer (MODIS) Active Fire product (MOD14A1_M). The product is a quantification of the number of fires observed within a 1000 km² area over a month. A fire must cover at least ~ 1000 m² to be detected, and must not be covered with clouds, heavy smoke, or tree canopy (Giglio et al., 2020). The Active Fire product is based on the 1 km fire channels at 3.9 and 11 µm of MODIS Terra and Aqua satellites (Justice et al., 2006). It is distributed at 0.1 deg resolution and at a monthly time scale by NASA Earth Observations (NEO) portal.

90 2.2 Precipitation

Precipitation (P) data come from the Precipitation Estimation from Remotely Sensed Information using Artificial Neural Networks- Climate Data Record (PERSIANN-CDR). The precipitation estimate uses the PERSIANN algorithm on GridSat-B1 infrared satellite data, and training of the artificial neural network on the National Centers for Environmental Prediction (NCEP) hourly precipitation data (Ashouri et al., 2015). The dataset is distributed by National Oceanic and Atmospheric

95 Administration (NOAA) at a daily time scale, and at 0.25 deg resolution in the latitude band $60^{\circ}S - 60^{\circ}N$.

2.3 Soil Moisture

The soil moisture (SM) dataset comes from the SMOS satellite, launched by the European Space Agency (ESA) in 2009 (Kerr et al., 2001). It performs passive measurements of the thermal emission of the Earth at L-band (1.4 GHz, 21 cm). L-band VOD and SM are derived from SMOS brightness temperatures using the L-band Microwave Emission of the Biosphere

100 (L-MEB) radiative transfer model (Wigneron et al., 2007; Kerr et al., 2012). L-band SM is the volume of water per volume of soil (m³ m⁻³) in the top surface soil layer (~ 5 cm). The footprint size is ~ 43 km in average (Kerr et al., 2010). We considered the ESA level 2 SM dataset in version 7.2 (L2 v720) resampled to the global cylindrical Equal-Area Scalable Earth (EASE) Grid version 2.0 (Brodzik et al., 2012) at 625 km² spatial sampling (25 km × 25 km at 30 deg of latitude). Ascending (6 am) and descending (6 pm) overpasses were averaged, from June 2010 to December 2020.

105 2.4 Terrestrial Water Storage

Terrestrial water storage (TWS) anomalies from the Gravity Recovery and Climate Experiment (GRACE) satellite were also considered. We used monthly GRACE/GRACE-Follow On (FO) Level-3 product provided through the Gravity Information Service (GravIS) web portal of the German Research Centre for Geosciences (GFZ) at 1 deg latitude-longitude grids (Boergens et al., 2019). TWS anomalies represent the water mass anomalies from snow, surface water, soil moisture, and

110 deep groundwater. They are derived from measurements of temporal changes in the Earth's gravity field. Data were lacking for 35 dates of the ten-year dataset. One-time gaps were filled by linear interpolation; consecutive missing months were not considered (Sep.-Nov. 2016, Jul. 2017-May 2018, and Aug.-Oct. 2018, 17 months in total).

2.5 Temperature

Temperature (T) data come from the Land Surface Temperature (LST) dataset from MODIS Terra satellite (NASA).
115 Daytime and night time measurements were averaged (MOD11C3 Version 6 product in a Climate Modeling Grid (CMG), LST_Day_CMG and LST_Night_CMG, Wan et al., 2015). These datasets are obtained using MODIS thermal infrared bands from 3 to 15 µm, and distributed by NASA Land Processes Distributed Active Archive Center (LP DAAC) at a monthly time scale and at 0.05 deg resolution.

2.6 Vegetation Optical Depth

120 Vegetation optical depth (VOD) is a remotely sensed indicator related to AGB and to VWC (Kerr and Njoku, 1990; Jackson and Schmugge, 1991; Jones et al., 2011; Rahmoune et al., 2014; Vittucci et al., 2016; Rodriguez-Fernandez et al., 2018; Mialon et al., 2020). No clear approach exists for disentangling the contributions of AGB and VWC in the VOD because of

the co-sensitivity of microwave observables to both quantities (Konings et al., 2019). The lower frequencies have better capabilities to penetrate deeper within the canopy (Ulaby et al., 1981). At L-band, VOD is sensitive to coarse woody

- 125 elements, such as trunks, stems, and branches. At C- and X-band, VOD is more sensitive to thin stems and leaves (Guglielmetti et al., 2007). L-VOD is then more sensitive to high AGB values, and is a good proxy for dense vegetation (Rodriguez-Fernandez et al., 2018). In this paper, L-VOD comes from SMOS level 2 dataset in version 7.2 (L2 v720) measured at 1.4 GHz ($\lambda = 21$ cm), resampled to EASE-Grid 2.0 at 625 km² resolution (25 km × 25 km at 30 deg of latitude). In the SMOS retrieval algorithm, the vegetation attenuation is taken into account by the τ parameter of the $\tau - \omega$ model (Mo
- et al., 1982) which corresponds to the L-VOD. Data from June 2010 to December 2020 were considered, and ascending (6 am) and descending (6 pm) overpasses were averaged. C- and X-VOD from the Japan Aerospace Exploration Agency (JAXA) Global Change Observation Mission (GCOM) Advanced Microwave Scanning Radiometer (AMSR)-2 dataset were also considered (Imaoka et al., 2010). C- and X-VOD are measured at 6.9 GHz (λ = 4.3 cm) and 10.7 GHz (λ = 2.8 cm) respectively. C2-band (7.3 GHz, λ = 4.1 cm) was not discussed in this paper as the data were mostly redundant with C1-band (6.9 GHz). We used the daily L3 V001 VOD products, from July 2012 to December 2020, processed with the Land
- Parameter Retrieval Model (LPRM) algorithm (Owe et al., 2008) and distributed by NASA on a regular grid at 25 km × 25 km resolution. Ascending (1:30 pm) and descending (1:30 am) overpasses (LPRM_AMSR2_A_SOILM3 and LPRM_AMSR2_D_SOILM3) were averaged.

2.7 Enhanced Vegetation Index

140 VODs values were compared with the visible-near infrared based enhanced vegetation index (EVI) from MODIS (NASA) MOD13C2 and MYD13C2 Version 6 for Aqua and Terra Satellites respectively, distributed at 5600 m resolution (Didan, 2015). EVI represents canopy greenness, with an improved sensitivity over high AGB regions compared to NDVI. It is obtained by combining measurements at red ($\lambda = 0.6-0.7 \mu m$, f ~ 460 THz) and near infrared wavelengths ($\lambda = 0.7-1.1 \mu m$, f ~ 330 THz).

145 2.8 Auxiliary data

2.8.1 Year of gross forest cover loss event

The year of gross forest cover loss event map (the so-called lossyear product) from Hansen et al. (2013) was used to observe the forest loss rate and year within a SMOS pixel, for the three major fires studied (Fig. 2). This map represents the first year of detected tree loss during the period 2000–2020, defined as a stand-replacement disturbance, or a change from a forest to

150 non-forest state. This dataset is based on Landsat images and is distributed at ~ 30 m resolution with 10x10 square degree tiles at https://glad.earthengine.app/view/global-forest-change. Each year of the period 2010-2020 was extracted from the forest loss product and averaged into SMOS EASE-Grid 2.0, so as to obtain a yearly percentage of forest loss.

2.8.2 Land cover

A land surface climatology map based on 10 years (2001–2010) of the MODIS MCD12Q1 product at 500 m resolution (Broxton et al., 2014) was used to filter the data and to distinguish four different vegetation types (see Sect. 3). This land cover map allows to identify 17 ecosystems based on the IGBP (International Geosphere-Biosphere Programme) class labels.

2.8.3 Above-Ground Biomass

The global map of AGB (Mg ha⁻¹) from Santoro et al. (2021) was used to distinguish sparse from dense forests (see Sect. 3.3). This map is distributed through the ESA Climate Change Initiative (CCI) Biomass at 100 m resolution. It combines a large pool of spaceborne remote sensing observations from two synthetic aperture radar (SAR) missions (Envisat and ALOS), and uses optical (Landsat) and LiDAR (Icesat GLAS) data to support the model calibration procedure. The ESA CCI Biomass map representative of 2010 was used here because it provides an AGB information prior to the studied fire events (2011–2020).

2.8.4 Snow and Ice

165 The Interactive Multisensor Snow and Ice Mapping System (IMS) database was used to mask areas covered with snow or ice (see Sect. 3.1). We used the IMS Daily Northern Hemisphere Snow and Ice Analysis at 4 km resolution, version 1 (Helfrich et al., 2007), provided by the National Snow and Ice Data Center (NSIDC).

2.8.5 Flooding

Flooded areas were filtered out (see Sect. 3.1) based on the Global Inundation Estimate from Multiple Satellites (GIEMS-2) dataset (Prigent et al., 2019). It provides long-term monthly estimates of surface water extent, including open water, wetlands, and rice paddies. The methodology combines passive and active microwaves, visible and near-infrared observations (SSM/I, ERS, AVHRR). The water fraction is delivered globally from 1992 to 2015, on an equal area grid of 0.25 deg × 0.25 deg at the equator (~ 28 km × 28 km). Flooded areas were detected with a climatology over the 1992–2015 period.

175 **2.8.6 Topography**

Strong topographies were also filtered out for this study (see Sect. 3.1). They were flagged using a mask created for SMOS retrieval (Mialon et al., 2008) based on a digital elevation model (DEM) provided by the Shuttle Radar Topography Mission (SRTM), a joint project between the National Aeronautics and Space Administration (NASA) and the National Geospatial-Intelligence Agency (NGA), conducted in 2000 (Jarvis et al., 2006).

180 **3 Methods**

First, we investigated three various regions which recently experienced a severe fire. These areas consist in: i) a eucalyptus open forest in a human-affected environment, under dry El Niño conditions in Australia; ii) a mixed area of needleleaf forests, woody savannas, and human activities under a Mediterranean climate in California; and iii) a remote primary rainforest in a tropical wet climate in Amazonia (see Sect. 3.2). Secondly, the study was extended to the global scale, for five vegetation types, by selecting the major fires of the last decade (see Sect. 3.3). The rationale was to capture significant events occurring over an area large enough to be observed with the SMOS satellite without any ambiguity. Four climate variables related to the fire risk were considered: precipitation, SM, TWS, and temperature. Wind is another predominant fire likelihood factor (Albini, 1993), but was not studied here due to the lack of reliable data at the required spatio-temporal scale. Vegetation status before, during, and after fire was monitored with four vegetation variables: EVI, X-VOD, C-VOD, and L-VOD.

3.1 Data preprocessing

Data from June 2010 to December 2020 were considered (10.5 years), except for C- and X-VOD from AMSR2 which were only available from July 2012. Only the latitude band 60° S – 60° N was kept, in order to be consistent with the precipitation dataset extent. Monthly averages of all datasets were computed and resampled to SMOS EASE-Grid 2.0 (~ 25 km

- 195 resolution) with a weighted average interpolation, using GDAL (GDAL/OGR contributors, 2020). SMOS data (SM and L-VOD) were filtered from RFI impacts by using a 20% maximum threshold on the RFI probability, provided by SMOS Level 2 product. Only the centre part of the swath was considered (less than 450 km away from the sub-satellite track) so as to only use optimal retrievals. Microwave measurements were also proven to be disturbed by strong topography (Mialon et al., 2008), snow (Schwank et al., 2014), and standing water (Ye et al., 2015; Jones et al., 2011; Bousquet et al., 2021). Hence, for
- 200 all datasets, we removed strong topography areas based on SMOS topography mask; snow-covered months based on IMS database (20% maximum snow coverage); water contaminated areas based on the land cover map (50% maximum water fraction); and flooded ones based on GIEMS-2 climatology (20% maximum water fraction).

3.2 Case study : analysis of three major fires

3.2.1 Wildfires in the South Coast of New South Wales in Australia

205 The first studied area is located in the South Coast of New South Wales in Australia, between [33.53°S – 37.72°S] and [149.40°E – 150.17°E] (Fig. 1) and covers thirteen SMOS pixels. The dominant vegetation type is eucalyptus open forest (McColl, 1969; DEWR 2007). The climate is warm temperate with dry summer (Kottek et al., 2006). The mean rainfall is ~ 1000 mm year⁻¹, and the mean temperature is ~ 15°C (McColl, 1969). The topography varies between 0 to 600 m above sea level. The 2019–2020 wildfires in Australia were influenced by El Niño Southern Oscillation (Dowdy, 2018). They became

210 historically significant as they were widespread and extremely severe, in particular in New South Wales (Ehsani et al., 2020). The tree cover loss map (Hansen et al., 2013) indicates a 25% forest loss in 2020 in the studied area (Fig. 2).

3.2.2 Mendocino Complex fire in California

The second studied area is located in California, near Lakeport, between [38.96°N – 39.46°N] and [122.68°W – 123.20°W] (Fig. 1). It corresponds to four SMOS pixels. The area is covered with evergreen needleleaf forest and woody savannas
(Broxton et al., 2014), and is much urbanised. The climate is warm temperate (Kottek et al., 2006), with dry, windy, and often hot weather conditions from spring through late autumn that can produce severe wildfires (Crockett and Westerling, 2018). The 2018 fire season was the most extreme on record in Northern California (now second to the 2020 fire season) in terms of number of fatalities, destroyed structures, and burned areas (Brown et al., 2020). The Mendocino complex is the largest fire complex in state history, and burned nearly 1860 km² of vegetation between July and September 2018. It included two wildfires: the Ranch fire at the North, which was the largest single fire in state history and burned 1660 km², and the River fire at the West, which burned 198 km² (BLM, 2018). The Mendocino complex caused a 34% vegetation loss in this region (26% in 2018 and 8% in 2019, Fig. 2), and was predominantly classified as moderate severity (62%; BLM, 2018).



225 Figure 1 - Global maps of SMOS L-VOD (left) and SM (right), in average for 2011–2020. The red dots show the locations of the three areas of interest: the Mendocino complex in California, Santarem in Amazonia, and the South Coast of New South Wales in Australia.



Figure 2 – Yearly forest loss (%) attributed to the three burnt areas under study, from Hansen et al. "lossyear" product.

3.2.3 Santarem wildfire in the Amazon

The third studied area is located in the Amazon rainforest near Santarem city (Brazil), between $[3.14^{\circ}S - 2.75^{\circ}S]$ and $[53.95^{\circ}W - 54.13^{\circ}W]$ (Fig. 1) and covers two SMOS pixels. The evergreen broadleaf forest is dense (L-VOD = 1.02; AGB = 280 Mg ha⁻¹ in average over the area). The climate is hot and humid, with annual mean temperature of 25°C and mean precipitation of 1920 mm year⁻¹ (Berenguer et al., 2018). During the strong El Niño event in December 2015, a severe drought caused large fires in this area, with no link with anthropic deforestation (Berenguer et al., 2018). They induced a 20% forest loss in 2016 in the studied area (Fig. 2).

3.2.4 Anomaly time series computation

Anomaly time series of EVI, X-, C-, L-VOD, P, SM, TWS, and T were plotted over the three studied sites. The anomaly time 240 series of a variable x is the difference between the original time series and the mean climatology, which is the mean seasonal cycle of this variable. They are defined as:

$$anom(x(t)) = x(t) - climatology(x(m))$$
⁽¹⁾

and

$$climatology(x(m)) = \frac{1}{y_n} \sum_{y=1}^{y_n} x(m + (y - 1) + 12), \forall m = 1:12$$
(2)

- where t is the month number from January 2010 (6 to 132 in this study); m is the month of the year, between 1 and 12, with $m = (t-1 \mod 12) + 1$; y is the year number, from 1 to y_n , with $y_n = 11$ here as the climatology was computed on the period 2010–2020. Plotting the anomaly time series enables to remove the natural seasonal cycle so as to observe only the variations due to specific events. The average pre-fire variable value was subtracted from the anomaly time series, only if at least twelve months of data were available before the fire event. It enables to observe the anomalies with respect to the pre-250 disturbance value. The time series of the number of fires were plotted in absolute values.

3.3 Extension to the global scale

Fires were then studied at the global scale to assess the general factors and impacts of fire according to the ecosystem. Only the latitude band 60°S – 60°N was retained in order to match with the precipitation dataset extent. Australia was also excluded because numerous fires were detected in 2012 in the Outback (shrublands) and in 2019/2020 in the South-East (broadleaf forests), which prevailed over the global dataset (~55% of the points) and prevented to perform a robust pre- and post-fire study. The global scale was divided into five land cover classes: needleleaf forests (IGBP labels 1 and 3), sparse broadleaf forests (IGBP labels 2 and 4, AGB ≤ 150 Mg ha⁻¹), dense broadleaf forests (IGBP labels 2 and 4, AGB > 150 Mg

235

ha⁻¹), savannas and shrublands (IGBP labels 6, 7, 8, and 9), and grasslands and croplands (IGBP labels 10, 12, and 14). Only

the range July 2012-December 2020 was conserved here for all datasets so as to match with AMSR2 time-period. To

- properly observe the factors and impacts of a fire event over a long time period without any other disturbance, only 25 km regions showing a unique and intense fire over the nine-year period were considered. This excluded areas with regular seasonal fires, such as the Sahel region. For that, a minimum threshold of 5 was applied on the maximum number of fires; and a maximum threshold of 2 was applied outside the main fire event period (i.e. outside the period -6/+6 months around the fire event). These thresholds were defined empirically. Anomalies were computed with Eq. (1) and (2), with a
- 265 climatology over all dates excepted the period -5/+6 months from the fire date, in order to remove these exceptional values. The anomaly time series were then shifted to collocate in time all fire events, and averaged by ecosystem. To ensure the spatial representativeness of each ecosystem, only the months with available data for at least one quarter of the sites were kept.

Pre-fire climatic anomalies and post-fire vegetation anomalies were also aggregated at different time frames and plotted, in order to compare their temporal behaviour in different ecosystems. The standard error of the mean of the measurements σ was also computed with Eq. (3):

$$\sigma(p) = \frac{std(p)}{\sqrt{n}} \tag{3}$$

where std is the standard deviation of the population p and n is the number of samples.

4 Results

270

275 4.1 Case study: analysis of three major fires

In evergreen forests of the South Coast of New South Wales in Australia (Fig. 3a), fires reached a maximum in January 2020 (mean number of fires = 8). They are associated with high temperature and low precipitation $(anom(T) = +3^{\circ}C, anom(P) = -80 \text{ mm})$. The drought started 3 years before fire (decrease in precipitation, SM, and TWS). All vegetation data exhibit the same pattern, which is i) a constant and mild decrease since 2012; ii) a strong decrease just before and during the fire event

280 (~-0.15); and iii) a rapid post-fire recovery (~ 1 year). C-VOD is the most affected vegetation variable.

In California, no major pre-fire drought is visible in summer 2018 (Fig. 3b). The Mendocino Complex was the strongest of the three case studies, with 20 fires observed on average in August 2018. It provoked a decrease in all vegetation variables, particularly in L-VOD (anom(L-VOD) = -0.08) and in EVI (anom(EVI) = -0.10). C- and X-VOD regained rapidly their pre-fire values (~ 1 year), but EVI and L-VOD did not.

In the dense rainforest near Santarem (Brazilian Amazon), the detected number of fires in December 2015 is quite low (~ 4.5) (Fig. 3c), but this value may be underestimated due to cloud coverage (Giglio et al., 2020). Vegetation variables are stable before the fire event, even if L-VOD signal is quite noisy because only two SMOS pixels were considered here. Strong positive temperature anomalies (+3°C), negative precipitation anomalies (-160 mm) and TWS anomalies (-60) are





Figure 3 – Time series of the number of fires, and anomaly time series of EVI, X-, C-, L-VOD, P, SM, TWS, and T on (a) South-East Australia (13 SMOS pixels), (b) the Mendocino Complex, California (4 SMOS pixels), and (c) Santarem (2 SMOS pixels).

4.2 Extension to the global scale

290

In this section, the major fires from July 2012 to December 2020 were analysed at the global scale, by shifting the anomaly time series of all variables on the fire date t_{fire} . The considered fires are well spread spatially and temporally over the nine-

year period (Fig. 4). In savannas and grasslands (Fig. 5a and 5b), pre-fire hydrologic variables are stable (positive for

- 300 grasslands) and temperature anomalies are negative during 2 years before fire. Concurrently, vegetation variables start to increase and reach a maximum a few months before the fire event (particularly C- and X-VOD over grasslands). Vegetation variables anomalies also show a light surplus over needleleaf forests just before the fire event (Fig. 5c). Over forests (Fig. 5c, 5d and 5e), a one-year pre-fire drought is visible through the temperature increase and the precipitation, SM, and TWS decrease. For all ecosystems, these drought conditions intensify just before and during fire, and end a few months after fire.
- 305 During fire, all vegetation variables abruptly decrease in all ecosystems, EVI being the most impacted one, excepted over dense forests where L-VOD heavily decreases (Fig. 5e). For all ecosystems, EVI recovers more rapidly than VODs. L-VOD is particularly long to recover over dense broadleaf forests (more than 4 years, Fig. 5e). Needleleaf forests (Fig. 5c) exhibit a slow recovery time for all vegetation variables, with ~ 3 years for EVI and ~ 4 years for VODs. VODs even continue to decrease during 1 year post-fire in this ecosystem. In low vegetation ecosystems (Fig. 5a and 5b), C- and X-VOD never
- 310 regain their immediately pre-fire values, which were particularly high.

Climate variables anomalies were also averaged in space and in time, within time frames of 6 months, from 24 to 1 month pre-fire, in order to observe their general trends (Fig. 6). The error bars were computed with Eq. (3). Precipitation anomalies (Fig. 6a) are negative from 6 months pre-fire for all classes, and reach $-10 \text{ mm month}^{-1}$ in average before the fire event. The precipitation deficit is more intense in dense broadleaf forests, where it starts two years pre-fire and reaches $-60 \text{ mm month}^{-1}$

- 315 before the fire event. SM anomalies (Fig. 6b) are similar for the three forest ecosystems. The SM deficit starts one year prefire and reaches -0.04 m³ m⁻³ before the fire event. Savannas and grasslands are affected later (6 months pre-fire) and to a lesser extent (~ -0.015 m³ m⁻³), as previously observed in Fig. 5. TWS anomalies (Fig. 6c) are negative from 12 months prefire, excepted for grasslands. Again, dense broadleaf forests are the most impacted ecosystem, with a minimum TWS anomaly of -7 before fire. Temperature anomalies (Fig. 6d) show significant negative anomalies in grasslands, savannas, and
- 320 needleleaf forests from two to one year pre-fire. From 6 months pre-fire, temperature anomalies show a surplus in nearly all ecosystems, and reach +0.95°C in needleleaf forests and +0.80°C in dense broadleaf forests before the fire event. In summary, pre-fire drought is mainly observed in forests, with particularly low hydrological values in dense forests (rainforests), and particularly high temperatures in needleleaf forests (boreal ecosystems). Savannas and grasslands barely suffer from pre-fire drought; temperatures are even mild one year pre-fire.
- 325 Vegetation variables anomalies were averaged within time frames of 6 months, from 1 to 36 months post-fire, in order to observe the global impacts and recovery time (Fig. 7). We considered that a variable has totally recovered when its anomaly is between -0.005 and +0.005. Immediately after fire, EVI is the most impacted variable, with average anomalies of -0.030 over grasslands, -0.033 over savannas, -0.036 over needleleaf forests, -0.052 over sparse broadleaf forests, and -0.059 over dense broadleaf forests (Fig. 7a). EVI recovers rapidly, in about 25 to 30 months over needleleaf forests, and 19 to 24
- 330 months over other ecosystems. X-VOD is less affected over forests (-0.015) than over low vegetation (-0.025) (Fig. 7b). X-VOD gets back to normal within three years, needleleaf forests being the longest to recover. C-band is mainly impacted over forests and savannas (~ -0.03, Fig. 7c). C-VOD recovers slower than X-VOD, in particular over forests. L-VOD is mainly

affected over broadleaf forests, and particularly over the densest ones (Fig. 7d). There, negative anomalies decrease up to - 0.05 one year post-fire, then slowly increase. L-VOD is less affected than C-VOD elsewhere. It also shows a delayed impact by one year over needleleaf forests, as for C- and X-VOD.



Figure 4 – Location of the selected fires and histograms of the fire dates, for grasslands and croplands (IGBP label 10, 12, and 14), savannas and shrublands (IGBP labels 6, 7, 8, and 9), needleleaf forests (IGBP labels 1 and 3), sparse broadleaf forests (IGBP labels 2 and 4, AGB ≤ 150 Mg ha⁻¹), and dense broadleaf forests (IGBP labels 2 and 4, AGB > 150 Mg ha⁻¹). Australia was excluded as well as areas affected by water, snow, or strong topography (see Sect. 3.1).



Figure 5 – Time series of the number of fires, and anomaly time series of EVI, X-, C-, L-VOD, P, SM, TWS, and T, shifted on the fire date, for (a) 669 points in the grasslands and croplands biome; (b) 591 points in the savannas and shrublands biome; (c) 387
points in the needleleaf forest biome; (d) 79 points in the sparse broadleaf forest biome; and (e) 66 points in the dense broadleaf forest biome. The missing values are mainly due to snow filtering.



Figure 6 – Anomalies of (a) precipitation, (b) SM, (c) TWS, and (d) temperature, for each land cover class, at several pre-fire timescales. The error bars were computed with Eq. (3).

Sparse broadleaf forests Dense broadleaf forests



Figure 7 – Anomalies of (a) EVI, (b) X-VOD, (c) C-VOD, and (d) L-VOD, for each land cover class, at several post-fire time-scales. The error bars were computed with Eq. (3).

5 Discussion

5.1 Case study : analysis of three major fires

- 355 In South-East Australia, a strong pre-fire drought is visible in the climate variables but also in the mild decrease of vegetation variables (Fig. 3a), linked with VWC deficit. Ehsani et al. (2020) stated that the air temperature from December 2019 to February 2020 was about 1°C higher than usual, which increased evapotranspiration, while the lack of precipitation prevented the soil from satisfying the moisture demand, and led to a significant vegetation drying (fuel) that facilitated the propagation of fires. After the fire event, L-VOD regained its pre-fire values within a year, meaning that the woody biomass 360 was not entirely destroyed. Indeed, these eucalyptus forests are known to be somewhat fire resistant (Wilkinson and Jennings)
- 1993; Caccamo et al., 2015). They can regenerate branches and leaves by resprouting from heat-resistant buds (Burrows, 2002). The rapid recovery of vegetation data can also be explained by the recovery of VWC, linked with the post-fire increase in precipitation and SM (Konings et al., 2021). Indeed, in 2020, SM values exceeded those of the previous decade (anom(SM) = +0.15 m³ m⁻³), corresponding to the end of the severe drought affecting South-East Australia associated with the 2020/2021 La Niña event (BoM, 2021). The increase of SM and precipitation may also have expedited the extinction of fires (Ehsani et al., 2020).

In California, the study by Brown et al. (2020) provides a comprehensive analysis of the climate and fuel conditions leading to the 2018 Mendocino complex, and reports several events that are also noticeable in our analysis. Among these, positive rainfall and SM anomalies in winter 2016/2017 are depicted in Fig. 3b, which led to the second consecutive spring with above average accumulation of fine fuel (grasses). Positive temperature anomalies in winter 2017/2018 are also visible, when a lack of storm enabled the survival of grasses. In April 2018, precipitation and warm temperatures led to above normal spring brush and grass growth. No major drought is visible in summer 2018, but low rainfall and warm temperatures led to a rapid drying of fuels, and induced a poor overnight humidity recovery. All these similarities with the findings by Brown et al. support our observations. The dramatic fire impacted EVI and L-VOD in the long term. Eucalyptus, pine trees and chapparal were burnt. Even if this type of vegetation is fire-adapted, the strength of the fire seemed to have destroyed most of it (34% vegetation loss, Hansen et al., 2013). Increased forest fire activity in recent decades in California has likely been enabled by the legacy of fire suppression, human settlement, and anthropogenic climate change (Abatzoglou and

- Williams, 2016). Stephens et al. (2018) stated that the massive current tree mortality in California will undoubtedly provoke severe "mass fires" in the coming decades, driven by the amount of dry and combustible wood.
- 380 In the Santarem region (Amazon), the winter 2015 wildfire was attributed to high temperature and low precipitation linked with El Niño event (Berenguer et al., 2018), which clearly emerges from Fig. 3c. These extreme drought conditions worsened during the fire, and may explain its strength. Contrary to TWS and precipitation, SM stayed stable during the fire, maybe because of the reduced accuracy of SM measurements under very dense forest. The three-year recovery time of EVI after the severe fire indicates a moderate regrowth of leaves and grasses. In contrast, L-VOD never regained its pre-fire values, meaning that trunks were impacted on the long term.

5.2 Extension to the global scale

At global scale, grasslands, croplands, shrublands and savannas do not show signs of pre-fire drought (Fig. 5a, 5b, 6). A substantial increase in vegetation variables occurs 1–2 years before fire, which implies an increase in vegetation density, e.g. available fuel. Immediately before fire, both VOD and SM values drop down, suggesting a decrease in VWC, especially over

- 390 grasslands (Fig. 5a). The increase of vegetation material combined with the decrease of VWC may contribute to trigger large wildfires (Forkel et al., 2017; Kuhn-Régnier et al., 2021). Indeed, the fire risk in savannas is highest for a dry vegetation with enough fuel to enable a drastic burning (Mbow et al., 2004). This vegetation growth might be enabled by negative pre-fire temperature anomalies, and light positive pre-fire hydrological variables anomalies (Fig. 6). Vegetation variables are less impacted by fires (Fig. 7) because they are rapid and burn through the grass layer, resulting in less destruction than in forests
- 395 (Menaut et al., 1990; Gignoux et al., 1997). L-VOD in particular is slightly impacted because the burnt vegetation is mainly leaf biomass. EVI quickly recovers after fire, probably because fire burns most of the AGB of grass species, but spares their large underground root systems, resulting in a rapid establishment of new shoots (Hochberg et al., 1994). The exceptionally high pre-fire vegetation variables values are never regained.
- In needleleaf forests anomaly time series (Fig. 5c), the numerous missing values correspond to the filtering of snow in 400 winter. These wildfires are located in the Northern hemisphere temperate and boreal forests, and mostly occur in late spring and summer (Fig. 4). De Groot et al. (2013) explained that most fires in Canada occur during summer, due to lightning strikes; whereas most fires in Russia occur during spring and are human-caused. We found a strong pre-fire drought in this ecosystem (low SM and high temperature one year pre-fire, Fig. 6), which is well documented for previous fire episodes (Weber and Stocks, 1998). Our results are in line with those of Forkel et al. (2012), who found that previous-summer SM
- 405 was a good predictor for burned area in Siberian larch forests. Indeed, negative summer anomalies led to low frozen water the following winter, and to less water released during the following spring-summer season, which in turn eased the outbreak of large wildfires. VODs also showed a light surplus before fire, possibly linked with litter thickening (e.g. dead needles, cured grass, leaf litter), which also facilitate fire propagation (de Groot et al., 2013). We found a delayed impact of fire on vegetation variables, and a longer recovery time than in other ecosystems, of about 3–4 years (Fig. 5c and 7). This duration is
- 410 slightly less than what was found in other studies (5 years in Canada, Goetz et al., 2006; 5 to 8 years in North America, Jin et al., 2012), but our findings still confirm previous results from Yang et al. (2017), who showed with NDVI analyses over North America that the fire effect on needleleaf trees was stronger and longer than on other vegetation types. Fires in North America are predominantly stand-replacing and high-intensity crown fires (Stocks et al., 2004; Jin et al., 2012), whereas fires in Eurasia are generally lower intensity surface fires and less destructive for the vegetation (de Groot et al., 2013). These
- 415 different fire regimes are influenced by tree species (Rogers et al., 2015). Time series were plotted separately over each continent (Fig. S1). L-VOD and EVI recover slower in North-America than in Eurasia (~ 4 years vs ~ 2 years for L-VOD, ~ 3 years vs ~ 2 years for EVI), confirming these different boreal fire regimes. Moreover, we found that L-VOD is moderately

impacted during fire. This can be attributed to the dominant destruction of needles and branches by boreal fires (Alexander and Cruz, 2011).

- 420 Sparse broadleaf forests (AGB ≤ 150 Mg ha⁻¹) subject to wildfires are mostly located in subtropical and temperate areas of South America, North America, West Africa, and South-East Asia (Fig. 4). A drying trend is visible one year pre-fire (Fig. 5d and 6). The link between drought and wildfires was previously observed by de Marzo et al. (2021) in the Argentine Gran Chaco; by Cheng et al. (2013) in the Mexican Yucatan forest; and by Vadrevu et al. (2019) in South-East Asian forests, with a prominent influence of precipitation variations over temperature variations. L-VOD and EVI are particularly impacted by
- 425 fire, but they recover quickly (1 year for EVI, 2 years for L-VOD). Yang et al. (2017) also found a rapid recovery time over North American broadleaf trees due to their fire-adaptive resprouting regeneration mode. Same observations were made in the fire-prone Argentine Chaco forest by Torres et al. (2014).

Dense broadleaf forests are mostly located in the tropics (Fig. 4). We can notice few fires in the densest rainforests (Congo basin, central Amazon) because i) they are usually too humid to burn (Cochrane, 2003; Forkel et al., 2017); ii) MODIS active fire detections are underestimated under thick cloud coverage or for understory fires (Giglio et al., 2020); and iii) seasonally flooded areas were excluded in order to use only robust VOD estimations (Bousquet et al., 2021). A consistent drought is visible 8 months before fire events (Fig. 5e), with high negative SM, TWS, and precipitation anomalies (Fig. 6). Chen et al. (2013) also found TWS deficits before severe fire seasons across the southern Amazon. Indeed, rainfall shortage generates high water deficits (i.e. high negative TWS and SM anomalies), which cause tree mortality, leaf shedding (visible in pre-fire

- 435 EVI decrease) and thus increase fuel availability (Aragão et al., 2018). Nevertheless, no pre-fire VOD decrease is observed here, showing that tree species of dense forests can maintain their VWC. Drought-related fires were suggested to prevail over deforestation fires in the Amazon, and are predicted to increase in the near future (Aragão et al., 2018). The opening of forest canopies also boosts incident radiation levels which leads to temperature rise (Ray et al., 2005). The combination of fuel increase in a drier and hotter environment converts forests into fire-prone ecosystems (Aragão et al., 2018). We also
- 440 found that the dense broadleaf forest biome was the one most impacted by fire (Fig. 7), because the absolute values of vegetation variables before fires are higher in this biome, and because it is not a fire-adapted ecosystem (Cochrane, 2003). L-VOD in particular decreases strongly and recovers very slowly (Fig. 7d), as previously observed over Santarem fire (Fig. 3c). The strong post-fire decrease in L-VOD is due to biomass destruction but also to water stress in the remaining vegetation (Konings et al., 2021). This finding confirms the significant and damaging impact of fires in the dense broadleaf ecosystem previously observed by Silva et al. (2018) and de Faria et al. (2021). L-VOD was previously proven to be more
- 445 ecosystem previously observed by Silva et al. (2018) and de Faria et al. (2021). L-VOD was previously proven to be more sensitive to high AGB values than C- and X-VOD (Rodriguez-Fernandez et al., 2018). Here, we suggest that L-VOD depicts better the fire impact on high AGB areas than the other vegetation variables.

For all biomes, EVI is the most rapid index to recover, because leaves rapidly resprout. EVI and X-VOD seem better adapted for grasslands fire monitoring, C-VOD for savanna fire monitoring, and L-VOD for forest fire monitoring.

450 5.3 The potential of L-VOD for vegetation recovery monitoring over dense forests

Vegetation variables anomalies were also plotted with respect to the number of fires in the dense broadleaf ecosystem, immediately after fire (1–3 months post-fire, Fig. 8a) and over a longer period (1–2 years post-fire, Fig. 8b). A quasi-linear relationship is visible for all vegetation estimates. As previously observed in Sect. 4.2, EVI and L-VOD are the most impacted variables immediately after fire (Fig. 8a), while L-VOD is still significantly affected 1 to 2 years after fire (up to - 0.05, Fig. 8b). L-VOD then shows a clear response to fire events over high AGB areas, immediately but also in the long term, and proportionally to the number of fires within a SMOS pixel. Thanks to its sensitivity to coarse woody elements and to its deep penetration through the vegetation layer, L-VOD is better correlated to high AGB than other vegetation variables (Rodriguez-Fernandez et al., 2018), and could be used for post-fire recovery monitoring over dense forests. One must keep

in mind that not only the biomass volume (AGB) but also the biomass status (VWC) is depicted in the VOD.

460

455



Figure 8 – Anomalies of vegetation variables (V) averaged (a) from 1 to 3 months post-fire, and (b) from 24 to 35 months post-fire, with respect to the number of fires by pixel (MODIS), for dense broadleaf forests only. The anomalies were normalized with the 99th quantile of each variable V_{max} (EVI_{max} = 0.60, X-VOD_{max} = 1.02, C-VOD_{max} = 1.20, and L-VOD_{max} = 1.21).

465 6 Conclusion

In this paper, we analysed the pre-fire behaviour of four fire likelihood factors, including SMOS SM which provides access to top surface soil moisture at the global scale. In forests, which generally maintain a steady humidity, we found that fires are linked with intense and prolonged drought. Pre-fire temperature anomalies are particularly high in boreal needleleaf forests. In savannas and grasslands, in agreement with previous studies (Mbow et al., 2004), we found evidences of an increase in

470 available fuel prior to fire events, enabled by humid and cold conditions a few years before. We also found that vegetation variables recover rapidly in these ecosystems, as wildfires are often rapid and mildly destructive for trees. In contrast, over forests, fires can damage the vegetation in the long term. Zhang et al. (2021) demonstrated the potential of C-band vegetation optical depth to detect the vegetation change patterns caused by fire in the southern Amazon. Our study confirms these

findings and extends it to the global scale, and to two extra wavelengths. Dense broadleaf forest fires particularly impact the

- 475 L-band emission, which represents coarse woody elements (trunks and stems); whereas sparse vegetation fires affect more C- and X-bands, which are more sensitive to small branches and leaves. For all biomes, the visible-infrared index (EVI) drops down after fire but recovers quickly, as it represents only herbage and canopy foliage. The long term impact on L-VOD in dense broadleaf forests shows that fires in this ecosystem are severely destructive for trunks, while smaller woody elements and leaves resprout faster. Thus, L-VOD seems the best adapted vegetation variable for the monitoring of dense
- 480 vegetation recovery after large fires.

The increasing number of wildfires threatens the stability of several ecosystems. It is then particularly important to monitor the vegetation health and L-band proved to be complementary to existing measurements, especially over dense forests.

Author contributions

E.B., A.M., N.J.R.F., and Y.H.K. planned the research discussed in this manuscript. E.B. performed most of the
 computations. S.M. provided the AGB dataset and expertise on AGB and on forest loss estimation. All authors participated in
 the writing and provided comments and suggestions.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

490 E.B., A.M., N.J.R.F., Y.H.K. acknowledge support by CNES (Centre National d'Etudes Spatiales) TOSCA program. SMOS L2 was obtained from ESA's DPGS (Data Processing Ground Segment). The authors would like to thank the European Space Agency (ESA). AMSR2 data were provided by Vrije Universiteit Amsterdam (Richard de Jeu) and NASA GSFC (Manfred Owe) (2014), AMSR2/GCOM-W1 surface soil moisture (LPRM) L3 1 day 25 km × 25 km ascending V001, Edited by Goddard Earth Sciences Data and Information Services Center (GES DISC) (Bill Teng), Greenbelt, MD, USA, 495 Goddard Earth Sciences Data and Information Services Center (GES DISC), Accessed: 29/07/2020, doi:10.5067/M5DTR2QUYLS2. TWS data were obtained from GravIS web portal, the Gravity Information Service of the German Research Centre for Geosciences (GFZ). Fires, temperature and EVI data came from NASA Earth Observations (NEO) portal and Land Processes Distributed Active Archive Center (LP DAAC). IMS Daily Northern Hemisphere Snow and Ice Analysis at 4 km resolution, Version 1, came from the United States National Ice Center (USNIC), Boulder, 500 Colorado USA, delivered by the National Snow and Ice Data Center (NSIDC), doi: https://doi.org/10.7265/N52R3P,

Accessed: 01/08/2020. The authors wish to thank David Chaparro and Matthias Forkel for their relevant comments and reviews which considerably improved the manuscript.

References

Abatzoglou, J. T., and Williams, A. P.: Impact of anthropogenic climate change on wildfire across western US forests, P. 505 Natl. Acad. Sci. USA, 113, 11770–11775, https://doi.org/10.1073/pnas.1607171113, 2016.

Albini, F. A.: Dynamics and modeling of vegetation fires: observations, in: Fire in the environment: the ecological, atmospheric, and climatic importance of vegetation fires, edited by: Crutzen, P. J. and Goldammer, J. G., John Wiley & Sons, Chichester, England, 39–52, 1993.

Alexander M. E. and Cruz M. G.: Crown fire dynamics in conifer forests, in: Synthesis of knowledge of extreme fire
behavior, USDA Forest Service, Pacific Northwest Research Station, General technical report PNW-GTR-854, 1, 107–142, 2011.

Ambadan, J. T., Oja, M., Gedalof, Z. E., and Berg, A. A.: Satellite-Observed Soil Moisture as an Indicator of Wildfire Risk, Remote Sensing, 12, 1543, https://doi.org/10.3390/rs12101543, 2020.

Aragão, L. E., Anderson, L. O., Fonseca, M. G., Rosan, T. M., Vedovato, L. B., Wagner, F. H., Silva, C. V. J., Silva Junior, C.

515 H. L., Arai, E., Aguiar, A. P., Barlow, J., Berenguer, E., Deeter, M. N., Domingues, L. G., Gatti, L., Gloor, M., Malhi, G., Marengo, J. A., Miller, J. B., Phillips, O. L., and Saatchi, S.: 21st Century drought-related fires counteract the decline of Amazon deforestation carbon emissions, Nat. commun., 9, 1–12, https://doi.org/10.1038/s41467-017-02771-y, 2018.

Asner, G. P., Loarie, S. R., and Heyder, U.: Combined effects of climate and land-use change on the future of humid tropical forests, Conserv. Lett., 3, 395–403, https://doi.org/10.1111/j.1755-263X.2010.00133.x, 2010.

520 Ashouri, H., Hsu, K. L., Sorooshian, S., Braithwaite, D. K., Knapp, K. R., Cecil, L. D., Nelson, B. R., and Prat, O. P.: PERSIANN-CDR: Daily precipitation climate data record from multisatellite observations for hydrological and climate studies, B. Am. Meteorol. Soc., 96, 69–83, https://doi.org/10.1175/BAMS-D-13-00068.1, 2015.

Barlow, J., Peres, C. A., Lagan, B. O., and Haugaasen, T.: Large tree mortality and the decline of forest biomass following Amazonian wildfires, Ecol. Lett., 6, 6–8, https://doi.org/10.1046/j.1461-0248.2003.00394.x, 2003.

525 Berenguer, E., Malhi, Y., Brando, P., Cardoso Nunes Cordeiro, A., Ferreira, J., França, F., Chesini Rossi, L., Moraes de Seixas, M. M., and Barlow, J.: Tree growth and stem carbon accumulation in human-modified Amazonian forests following drought and fire, Phil. Trans. R. Soc. B., 373, 20170308, https://doi.org/10.1098/rstb.2017.0308, 2018.

BLM: Mendocino Complex Fire, Watershed Modeling Report, available at: https://www.blm.gov/sites/blm.gov/files/uploads/ Programs Fire California Watershed Modeling Report-Mendocino-Complex-Fire.pdf, accessed 14 December 2021, 2018.

530 Boergens, E., Dobslaw, H., and Dill, R.: GFZ GravIS RL06 Continental Water Storage Anomalies, V. 0002, GFZ Data Services, http://doi.org/10.5880/GFZ.GRAVIS 06 L3 TWS, 2019.

BoM: Annual climate statement 2020, available at: http://www.bom.gov.au/climate/current/annual/aus/2020/, accessed 14 December 2021, 2021.

Bousquet, E., Mialon, A., Rodriguez-Fernandez, N., Prigent, C., Wagner, F. H., and Kerr, Y. H.: Influence of surface water
variations on VOD and biomass estimates from passive microwave sensors, Remote Sens. Environ., 257, 112345, https://doi.org/10.1016/j.rse.2021.112345, 2021.

Brando, P. M., Soares-Filho, B., Rodrigues, L., Assunção, A., Morton, D., Tuchschneider, D., Fernandes, E.C.M, Macedo, M.N., Oliveira, U., and Coe, M. T.: The gathering firestorm in southern Amazonia, Sci. Adv., 6, eaay1632, https://doi.org/10.1126/sciadv.aay1632, 2020.

540 Brodzik, M. J., Billingsley, B., Haran, T., Raup, B., and Savoie, M. H.: EASE-Grid 2.0: Incremental but Significant Improvements for Earth-Gridded Data Sets, ISPRS Int. J. Geo-Inf., 1, 32–45, https://doi.org/10.3390/ijgi1010032, 2012.

Broxton, P. D., Zeng, X., Sulla-Menashe, D., and Troch, P. A.: A global land cover climatology using MODIS data, J. Appl. Meteorol. Clim., 53, 1593–1605, https://doi.org/10.1175/JAMC-D-13-0270.1, 2014.

Brown, T., Leach, S., Wachter, B., and Gardunio, B.: The Northern California 2018 Extreme Fire Season, in: Explaining
545 Extremes of 2018 from a Climate Perspective, B. Am. Meteorol. Soc., 101, S1–S4, https://doi.org/10.1175/BAMS-D-19-0275.1, 2020.

Burrows, G. E.: Epicormic strand structure in Angophora, Eucalyptus and Lophostemon (Myrtaceae): implications for fire resistance and recovery, New Phytol., 111–131, 2002.

Caccamo, G., Bradstock, R., Collins, L., Penman, T., and Watson, P.: Using MODIS data to analyse post-fire vegetation recovery in Australian eucalypt forests, J. Spat. Sci., 60, 341–352, https://doi.org/10.1080/14498596.2015.974227, 2015. CAMS, Copernicus Atmosphere Monitoring Service: Copernicus: A summer of wildfires saw devastation and record emissions around the Northern Hemisphere, https://atmosphere.copernicus.eu/copernicus-summer-wildfires-saw-devastation-and-record-emissions-around-northern-hemisphere, accessed 30 September 2021.

Certini, G.: Effects of fire on properties of forest soils: a review, Oecologia, 143, 1–10, https://doi.org/10.1007/s00442-004-555 1788-8, 2005.

Chaparro, D., Piles, M., Vall-Llossera, M., and Camps, A.: Surface moisture and temperature trends anticipate drought conditions linked to wildfire activity in the Iberian Peninsula, Eur. J. Remote Sens., 49, 955–971, 2016.

Chen, Y., Velicogna, I., Famiglietti, J. S., and Randerson, J. T.: Satellite observations of terrestrial water storage provide early warning information about drought and fire season severity in the Amazon, J. Geophys. Res.-Biogeo., 118, 495–504, https://doi.org/10.1002/jgrg.20046, 2013.

Cheng, D., Rogan, J., Schneider, L., and Cochrane, M.: Evaluating MODIS active fire products in subtropical Yucatán forest, Remote Sens. Lett., 4, 455–464, https://doi.org/10.1080/2150704X.2012.749360, 2013.

Chu, T. and Guo, X.: Remote sensing techniques in monitoring post-fire effects and patterns of forest recovery in boreal forest regions: A review, Remote Sensing, 6, 470–520, https://doi.org/10.3390/rs6010470, 2013.

565 Chuvieco, E., Aguado, I., Jurdao, S., Pettinari, M. L., Yebra, M., Salas, J., Hantson, S., de la Riva, J., Ibarra, P., Rodrigues, M., Echeverría, M., Azqueta, D., Román, M. V., Bastarrika, A., Martínez, S., Recondo, C., Zapico, E., and Vega, J. M.: Integrating geospatial information into fire risk assessment. Int. J. wildland fire. 23. 606-619. https://doi.org/10.1071/WF12052, 2012.

Cochrane, M. A.: Fire science for rainforests, Nature, 421, 913–919, https://doi.org/10.1038/nature01437, 2003.

570 Crockett, J. L. and Westerling, A. L.: Greater temperature and precipitation extremes intensify Western US droughts, wildfire severity, and Sierra Nevada tree mortality, J. Climate, 31, 341–354, https://doi.org/10.1175/JCLI-D-17-0254.1, 2018.

de Faria, B. L., Brando, P. M., Macedo, M. N., Panday, P. K., Soares-Filho, B. S., and Coe, M. T.: Current and future patterns of fire-induced forest degradation in Amazonia, Environ. Res. Lett., 12, 095005, https://doi.org/10.1088/1748-9326/aa69ce, 2017.

575 de Faria, B. L., Marano, G., Piponiot, C., Silva, C. A., Dantas, V. D. L., Rattis, L., Rech, A.R., and Collalti, A.: Model-based estimation of Amazonian forests recovery time after drought and fire events, Forests, 12, 8, https://doi.org/10.3390/f12010008, 2021.

de Groot, W. J., Cantin, A. S., Flannigan, M. D., Soja, A. J., Gowman, L. M., and Newbery, A.: A comparison of Canadian and Russian boreal forest fire regimes, Forest Ecol. Manag., 294, 23–34, https://doi.org/10.1016/j.foreco.2012.07.033, 2013.

580 de Marzo, T., Pflugmacher, D., Baumann, M., Lambin, E. F., Gasparri, I., and Kuemmerle, T.: Characterizing forest disturbances across the Argentine Dry Chaco based on Landsat time series, Int. J. Appl. Earth Obs., 98, 102310, https://doi.org/10.1016/j.jag.2021.102310, 2021.

DEWR, Department of the Environment and Water Resources: Australia's native vegetation: a summary of Australia's major vegetation groups, Australian Government, Canberra, http://www.environment.gov.au/resource/australias-native-vegetation-summary-australias-major-vegetation-groups, accessed 20 July 2021, 2007.

585 sum

Didan, K.: MOD13C2 MODIS/Terra Vegetation Indices Monthly L3 Global 0.05Deg CMG V006 EVI. NASA EOSDIS Land Processes DAAC, https://doi.org/10.5067/MODIS/MOD13C2.006, accessed 01 July 2021, 2015.

Dowdy, A. J.: Climatological variability of fire weather in Australia, J. Appl. Meteorol. Clim., 57, 221–234, https://doi.org/10.1175/JAMC-D-17-0167.1, 2018.

Ehsani, M. R., Arevalo, J., Risanto, C. B., Javadian, M., Devine, C. J., Arabzadeh, A., Venegas-Quiñones, H. L., Dell'Oro, A. P., and Behrangi, A.: 2019–2020 Australia fire and its relationship to hydroclimatological and vegetation variabilities, Water, 12, 3067, https://doi.org/10.3390/w12113067, 2020.

Fan, L., Wigneron, J. P., Xiao, Q., Al-Yaari, A., Wen, J., Martin-StPaul, N., Dupuy, J. L., Pimont, F., Al Bitar, A., Fernandez-Moran, R., and Kerr, Y. H.: Evaluation of microwave remote sensing for monitoring live fuel moisture content in the
Mediterranean region, Remote Sensing of Environment, 205, 210–223, https://doi.org/10.1016/j.rse.2017.11.020, 2018.

FAO, Food and Agriculture Organization: Fire Management – Global Assessment 2006. A Thematic Study Prepared in the Framework of the Global Forest Resources Assessment 2005, Rome, 2006.

Farahmand, A., Stavros, E. N., Reager, J. T., Behrangi, A., Randerson, J. T., and Quayle, B.: Satellite hydrology observations as operational indicators of forecasted fire danger across the contiguous United States, Nat. Hazard Earth Sys., 20, 1097-1106, https://doi.org/10.5194/nhess-20-1097-2020, 2020.

600 1106, https://doi.org/10.5194/nhess-20-1097-2020, 202

Fernandez-Carrillo, A., McCaw, L., and Tanase, M. A.: Estimating prescribed fire impacts and post-fire tree survival in eucalyptus forests of Western Australia with L-band SAR data, Remote Sens. Environ., 224, 133–144, https://doi.org/10.1016/j.rse.2019.02.005, 2019.

Forkel, M., Thonicke, K., Beer, C., Cramer, W., Bartalev, S., and Schmullius, C.: Extreme fire events are related to previous-605 year surface moisture conditions in permafrost-underlain larch forests of Siberia, Environ. Res. Lett., 7, 044021, https://doi.org/10.1088/1748-9326/7/4/044021, 2012.

Forkel, M., Dorigo, W., Lasslop, G., Teubner, I., Chuvieco, E., and Thonicke, K.: A data-driven approach to identify controls on global fire activity from satellite and climate observations (SOFIA V1), Geosci. Model Dev., 10, 4443–4476, https://doi.org/10.5194/gmd-10-4443-2017, 2017.

610 Frappart, F., Wigneron, J. P., Li, X., Liu, X., Al-Yaari, A., Fan, L., Wang, M., Moisy, C., Le Masson, E., Aoulad Lafkih, Z., Vallé, C., Ygorra, B., and Baghdadi, N.: Global monitoring of the vegetation dynamics from the Vegetation Optical Depth (VOD): A review, Remote Sensing, 12, 2915, https://doi.org/10.3390/rs12182915, 2020.

GDAL/OGR contributors: GDAL/OGR Geospatial Data Abstraction software Library, Open Source Geospatial Foundation, https://gdal.org, accessed in 2020.

615 Giglio, L., Schroeder, W., Hall, J. V., and Justice, C. O.: MODIS Collection 6 Active Fire Product User's Guide Revision C, Department of Geographical Sciences, University of Maryland, 2020.

Gignoux, J., Clobert, J., and Menaut, J. C.: Alternative fire resistance strategies in savanna trees, Oecologia, 110, 576–583, https://doi.org/10.1007/s004420050198, 1997.

Goetz, S. J., Fiske, G. J., and Bunn, A. G.: Using satellite time-series data sets to analyze fire disturbance and forest recovery
across Canada, Remote Sens. Environ., 101, 352–365, https://doi.org/10.1016/j.rse.2006.01.011, 2006.

Guglielmetti, M., Schwank, M., Mätzler, C., Oberdörster, C., Vanderborght, J., and Flühler, H.: Measured microwave radiative transfer properties of a deciduous forest canopy, Remote Sens. Environ., 109, 523–532, https://doi.org/10.1016/j.rse.2007.02.003, 2007.

Hansen, M. C., Potapov, P. V., Moore, R., Hancher, M., Turubanova, S. A., Tyukavina, A., Thau, D., Stehman, S. V., Goetz,
S. J., Loveland, T. R., Kommareddy, A., Egorov, A., Chini, L., Justice, C. O., and Townshend, J. R. G.: High-resolution global maps of 21st-century forest cover change, Science, 342, 850–853, https://doi.org/10.1126/science.1244693, 2013.

Heath, J. T., Chafer, C. J., Bishop, T. F., and Van Ogtrop, F. F.: Post-fire recovery of eucalypt-dominated vegetation communities in the Sydney Basin, Australia, Fire Ecol., 12, 53–79, https://doi.org/10.4996/fireecology.1203053, 2016.

Helfrich, S., McNamara, D., Ramsay, B., Baldwin, T., and Kasheta, T.: Enhancements to, and forthcoming developments in

630 the Interactive Multisensor Snow and Ice Mapping System (IMS), Hydrol. Process., 21, 1576–1586, https://doi.org/10.1002/ hyp.6720, 2007.

Higuera, P. E. and Abatzoglou, J. T.: Record-setting climate enabled the extraordinary 2020 fire season in the western United States, Glob. change biol., https://doi.org/10.1111/gcb.15388, 2020.

Hochberg, M. E., Menaut, J. C., and Gignoux, J.: The influences of tree biology and fire in the spatial structure of the West 635 African savannah, J. Ecol., 82, 217–226, 1994.

Holgate, C. M., van Dijk, A. I., Cary, G. J., and Yebra, M.: Using alternative soil moisture estimates in the McArthur Forest Fire Danger Index, Int. J. Wildland Fire, 26, 806-819, https://doi.org/10.1071/WF16217, 2017.

Imaoka, K., Kachi, M., Kasahara, M., Ito, N., Nakagawa, K., and Oki, T.: Instrument performance and calibration of AMSR-E and AMSR2, Int. Arch. Photogramm., 38, 13–18, 2010.

640 Jackson, T. J. and Schmugge, T. J.: Vegetation effects on the microwave emission of soils, Remote Sens. Environ., 36, 203– 212, 1991.

Jarvis, A., Reuter, H., Nelson, A., and Guevara, E.: Hole-filled seamless SRTM data V3, International Centre for Tropical Agriculture (CIAT), available from CGIAR-CSI SRTM 90 m database, http://srtm.csi.cgiar.org, accessed 30 September 2021, 2006.

645 Jensen, D., Reager, J. T., Zajic, B., Rousseau, N., Rodell, M., and Hinkley, E.: The sensitivity of US wildfire occurrence to pre-season soil moisture conditions across ecosystems, Environ. Res. Lett., 13, 014021, https://doi.org/10.1088/1748-9326/aa9853, 2018.

Jin, Y., Randerson, J. T., Goetz, S. J., Beck, P. S., Loranty, M. M., and Goulden, M. L.: The influence of burn severity on postfire vegetation recovery and albedo change during early succession in North American boreal forests, J. Geophys. Res.-Biogeo., 117, G01036, https://doi.org/10.1029/2011JG001886, 2012.

650

655

Jones, M. O., Jones, L. A., Kimball, J. S., and McDonald, K. C.: Satellite passive microwave remote sensing for monitoring global land surface phenology, Remote Sens. Environ., 115, 1102–1114, https://doi.org/10.1016/j.rse.2010.12.015, 2011.

Justice, C., Giglio, L., Boschetti, L., Roy, D., Csiszar, I., Morisette, J., and Kaufman, Y.: Algorithm technical background document MODIS fire products, MODIS Science Team, Washington DC, USA, https://eospso.nasa.gov/sites/default/files/atbd/atbd mod14.pdf, accessed 01 January 2020, 2006.

Kerr, Y. H. and Njoku, E.G.: A semiempirical model for interpreting microwave emission from semiarid land surfaces as seen from space, IEEE T. Geosci. Remote, 28, 384–393, https://doi.org/10.1109/36.54364, 1990.

Kerr, Y. H., Waldteufel, P., Wigneron, J. P., Martinuzzi, J. A. M. J., Font, J., and Berger, M.: Soil moisture retrieval from space: The Soil Moisture and Ocean Salinity (SMOS) mission, IEEE T. Geosci. Remote, 39, 1729–1735, https://doi.org/10.1109/36.942551, 2001.

660

Kerr, Y. H., Waldteufel, P., Wigneron, J. P., Delwart, S., Cabot, F., Boutin, J., Escorihuela, M. J., Font, J., Reul, N., Gruhier, C., Juglea, S. E., Drinkwater, M. R., Hahne, A., Martin-Neira, M., and Mecklenburg, S.: The SMOS mission: New tool for monitoring key elements of the global water cycle, P. IEEE, 98, 666–687, https://doi.org/10.1109/JPROC.2010.2043032, 2010.

665 Kerr, Y. H., Waldteufel, P., Richaume, P., Wigneron, J. P., Ferrazzoli, P., Mahmoodi, A., Al Bitar, A., Cabot, F., Gruhier, C., Juglea, S. E., Leroux, D., Mialon, A., and Delwart, S.: The SMOS Soil Moisture Retrieval Algorithm, Geosci. Remote Sens., 50, 1384–1403, https://doi.org/10.1109/TGRS.2012.2184548, 2012.

Konings, A. G., Rao, K., and Steele-Dunne, S. C.: Macro to micro: microwave remote sensing of plant water content for physiology and ecology, New Phytol., 223, 1166–1172, https://doi.org/10.1111/nph.15808, 2019.

670 Konings, A. G., Holtzman, N., Rao, K., Xu, L., and Saatchi, S. S.: Interannual Variations of Vegetation Optical Depth Are Due to Both Water Stress and Biomass Changes, Geophys. Res. Lett., 48, e2021GL095267, https://doi.org/10.1029/2021GL095267, 2021.

Kottek, M., Grieser, J., Beck, C., Rudolf, B., and Rubel, F.: World map of the Köppen-Geiger climate classification updated, Meteorol. Z., 15, 259–263, https://doi.org/10.1127/0941-2948/2006/0130, 2006.

675 Kuhn-Régnier, A., Voulgarakis, A., Nowack, P., Forkel, M., Prentice, I. C., and Harrison, S. P.: The importance of antecedent vegetation and drought conditions as global drivers of burnt area, Biogeosciences, 18, 3861–3879, https://doi.org/10.5194/bg-18-3861-2021, 2021.

Mbow, C., Goïta, K., and Bénié, G. B.: Spectral indices and fire behavior simulation for fire risk assessment in savanna ecosystems, Remote Sens. Environ., 91, 1–13, https://doi.org/10.1016/j.rse.2003.10.019, 2004.

680 McColl, J. G.: Soil-Plant Relationships in a Eucalyptus Forest on the South Coast of New South Wales, Ecology, 50, 354– 362, https://doi.org/10.2307/1933883, 1969. Menaut, J. C., Gignoux, J., Prado, C., and Clobert, J.: Tree community dynamics in a humid savanna of the Cote-d'Ivoire: modelling the effects of fire and competition with grass and neighbours, J. Biogeogr., 471–481, 1990.

Mermoz, S. and Le Toan, T.: Forest disturbances and regrowth assessment using ALOS PALSAR data from 2007 to 2010 in Vietnam, Cambodia and Lao PDR, Remote Sensing, 8, 217, https://doi.org/10.3390/rs8030217, 2016.

Mhawej, M., Faour, G., and Adjizian-Gerard, J.: Wildfire likelihood's elements: a literature review, Challenges, 6, 282–293, https://doi.org/10.3390/challe6020282, 2015.

Mialon, A., Coret, L., Kerr, Y. H., Sécherre, F., and Wigneron, J.-P.: Flagging the topographic impact on the SMOS signal, IEEE T. Geosci. Remote, 46, 689–694, https://doi.org/10.1109/TGRS.2007.914788, 2008.

690 Mialon, A., Rodriguez-Fernandez, N. J., Santoro, M., Saatchi, S., Mermoz, S., Bousquet, E., Kerr, Y. H.: Evaluation of the Sensitivity of SMOS L-VOD to Forest Above-Ground Biomass at Global Scale, Remote Sensing, 12, 1450, https://doi.org/10.3390/rs12091450, 2020.

Mo, T., Choudhury, B. J., Schmugge, T. J., Wang, J. R., and Jackson, T. J.: A model for microwave emission from vegetationcovered fields, J. Geophys. Res., 87, 11229–11237, https://doi.org/10.1029/JC087iC13p11229, 1982.

Nepstad, D., Carvalho, G., Barros, A. C., Alencar, A., Capobianco, J. P., Bishop, J., Moutinho, P., Lefebvre, P., Silva Junior, U. L., and Prins, E.: Road paving, fire regime feedbacks, and the future of Amazon forests, Forest Ecol. Manag., 154(3), 395–407, https://doi.org/10.1016/S0378-1127(01)00511-4, 2001.

Owe, M., de Jeu, R., and Holmes, T.: Multisensor historical climatology of satellite-derived global land surface moisture, J. Geophys. Res., 113, F01002, https://doi.org/10.1029/2007JF000769, 2008.

700 Pérez-Cabello, F., Montorio, R., and Alves, D. B.: Remote Sensing Techniques to assess Post-Fire Vegetation Recovery, Curr. Opin. Env. Sci. Health, 21, 100251, https://doi.org/10.1016/j.coesh.2021.100251, 2021.

Parsons, D. J. and DeBenedetti, S. H.: Impact of fire suppression on a mixed-conifer forest, Forest Ecol. Manag., 2, 21-33, 1979.

Prigent, C., Jimenez, C., and Bousquet, P.: Satellite-derived global surface water extent and dynamics over the last 25 years (GIEMS-2), J. Geophys. Res.-Atmos., 125, e2019JD030711, https://doi.org/10.1029/2019JD030711, 2019.

Rahmoune, R., Ferrazzoli, P., Singh, Y., Kerr, Y. H., Richaume, P., and Al Bitar, A.: SMOS Retrieval Results Over Forests: Comparisons With Independent Measurements, IEEE J. Sel. Top. Appl., 7, 3858–3866, https://doi.org/10.1109/JSTARS.2014.2321027, 2014.

Ray, D., Nepstad, D., and Moutinho, P.: Micrometeorological and canopy controls of fire susceptibility in a forested Amazon
landscape, Ecol. Appl., 15, 1664–1678, https://doi.org/10.1890/05-0404, 2005.

Rodriguez-Fernandez, N. J., Mialon, A., Mermoz, S., Bouvet, A., Richaume, P., Al Bitar, A., Al-Yaari, A., Brandt, M., Kaminski, T., Le Toan, T., Kerr, Y. H., and Wigneron, J. P.: An evaluation of SMOS L-band vegetation optical depth (L-VOD) data sets: high sensitivity of L-VOD to above-ground biomass in Africa, Biogeosciences, 15, 4627–4645, https://doi.org/10.5194/bg-15-4627-2018, 2018.

715 Rogers, B. M., Soja, A. J., Goulden, M. L., and Randerson, J. T.: Influence of tree species on continental differences in boreal fires and climate feedbacks, Nat. Geosci., 8, 228–234, https://doi.org/10.1038/ngeo2352, 2015.

Sackett, S. S.: Scheduling prescribed burns for hazard reduction in the southeast, J. Forest., 73, 143–147, https://doi.org/10.1093/jof/73.3.143, 1975.

Santoro, M., Cartus, O., Carvalhais, N., Rozendaal, D., Avitabile, V., Araza, A., de Bruin, S., Herold, M., Quegan, S.,
Rodríguez-Veiga, P., Balzter, H., Carreiras, J., Schepaschenko, D., Korets, M., Shimada, M., Itoh, T., Moreno Martínez, A.,
Cavlovic, J., Cazzolla Gatti, R., da Conceição Bispo, P., Dewnath, N., Labrière, N., Liang, J., Lindsell, J., Mitchard, E. T. A.,
Morel, A., Pacheco Pascagaza, A. M., Ryan, C. M., Slik, F., Laurin, G. V., Verbeeck, H., Wijaya, A., and Willcock, S.: The
global forest above-ground biomass pool for 2010 estimated from high-resolution satellite observations, Earth Syst. Sci.
Data, 13, 3927–3950, https://doi.org/10.5194/essd-13-3927-2021, 2021.

725 Schwank, M., Rautiainen, K., Mätzler, C., Stähli, M., Lemmetyinen, J., Pulliainen, J., Vehviläinen, J., Kontu, A., Ikonen, J., Ménard, C. B., Drusch, M., Wiesmann, A., and Wegmüller, U.: Model for microwave emission of a snow-covered ground with focus on L band, Remote Sens. Environ., 154, 180–191, https://doi.org/10.1016/j.rse.2014.08.029, 2014.

Shakesby, R. A. and Doerr, S. H.: Wildfire as a hydrological and geomorphological agent, Earth-Sci. Rev., 74, 269–307, https://doi.org/10.1016/j.earscirev.2005.10.006, 2005.

730 Silva, C. V., Aragão, L. E., Barlow, J., Espirito-Santo, F., Young, P. J., Anderson, L. O., Berenguer, E., Brasil, I., Brown, I. F., Castro, B., Farias, R., Ferreira, J., França, F., Graça, P. M. L. A., Kirsten, L., Lopes, A. P., Salimon, C., Scaranello, M. A., Seixas, M., Souza, F. C., and Xaud, H. A. M.: Drought-induced Amazonian wildfires instigate a decadal-scale disruption of forest carbon dynamics, Philos. T. Roy. Soc. B., 373, https://doi.org/10.1098/rstb.2018.0043, 2018. Stephens, S. L., Collins, B. M., Fettig, C. J., Finney, M. A., Hoffman, C. M., Knapp, E. E., North, M. P., Safford, H., and
Wayman, R. B.: Drought, tree mortality, and wildfire in forests adapted to frequent fire, BioScience, 68, 77–88, https://doi.org/10.1093/biosci/bix146, 2018.

Stocks, B. J., Alexander, M. E., and Lanoville, R. A.: Overview of the International Crown Fire Modelling Experiment (ICFME), Can. J. Forest Res., 34, 1543–1547, https://doi.org/10.1139/x04-905, 2004.

Torres, R. C., Giorgis, M. A., Trillo, C., Volkmann, L., Demaio, P., Heredia, J., and Renison, D.: Post-fire recovery occurs
overwhelmingly by resprouting in the Chaco Serrano forest of Central Argentina, Austral Ecol., 39, 346–354, https://doi.org/ 10.1111/aec.12084, 2014.

Vadrevu, K. P., Lasko, K., Giglio, L., Schroeder, W., Biswas, S., and Justice, C.: Trends in vegetation fires in south and southeast Asian countries, Sci. Rep.-UK, 9, 1–13, https://doi.org/10.1038/s41598-019-43940-x, 2019.

 Vittucci, C., Ferrazzoli, P., Kerr, Y., Richaume, P., Guerriero, L., Rahmoune, R., and Laurin, G. V.: SMOS retrieval over
 forests: Exploitation of optical depth and tests of soil moisture estimates, Remote Sens. Environ., 180, 115–127, https://doi.org/10.1016/j.rse.2016.03.004, 2016.

Wan, Z., Hook, S., and Hulley, G.: MOD11C3 MODIS/Terra Land Surface Temperature/Emissivity Monthly L3 Global 0.05Deg CMG V006, LST_Day_CMG and LST_Night_CMG, NASA EOSDIS Land Processes DAAC, https://doi.org/10.5067/MODIS/MOD11C3.006, accessed 25 August 2021, 2015.

750 Weber, M. G. and Stocks, B. J.: Forest fires and sustainability in the boreal forests of Canada, Ambio, 27(7), 545–550, 1998.

Wigneron, J. P., Kerr, Y. H., Waldteufel, P., Saleh, K., Escorihuela, M. J., Richaume, P., Ferrazzoli, P., de Rosnay, P., Gurney, R., Calvet, J. C., Grant, J., Guglielmetti, M., Hornbuckle, B., Mätzler, C., Pellarin, T., and Schwank, M.: L-band Microwave Emission of the Biosphere (L-MEB) Model: Description and calibration against experimental data sets over crop fields, Remote Sens. Environ., 107, 639–655, https://doi.org/10.1016/j.rse.2006.10.014, 2007.

Wilkinson, G. and Jennings, S.: Survival and recovery of Eucalyptus obliqua regeneration following wildfire, Tasforests, 5, 1–11, 1993.

Yang, J., Pan, S., Dangal, S., Zhang, B., Wang, S., and Tian, H.: Continental-scale quantification of post-fire vegetation greenness recovery in temperate and boreal North America, Remote Sens. Environ., 199, 277–290, https://doi.org/10.1016/j.rse.2017.07.022, 2017.

760 Ye, N., Walker, J. P., Guerschman, J., Ryu, D., and Gurney, R. J.: Standing water effect on soil moisture retrieval from Lband passive microwave observations, Remote Sens. Environ., 169, 232–242, 2015.

Zhang, H., Hagan, D. F. T., Dalagnol, R., and Liu, Y.: Forest Canopy Changes in the Southern Amazon during the 2019 Fire Season Based on Passive Microwave and Optical Satellite Observations, Remote Sensing, 13, 2238, https://doi.org/10.3390/rs13122238, 2021.