Riverine impact on future projections of marine primary 1 production and carbon uptake 2 Shuang Gao^{1,2}, Jörg Schwinger³, Jerry Tjiputra³, Ingo Bethke¹, Jens Hartmann⁴, Emilio 3 4 Mayorga⁵, Christoph Heinze¹ 5 ¹Geophysical Institute, University of Bergen, Bjerknes Centre for Climate Research, Bergen, Norway 6 ²Institute of Marine Research, Bergen, Norway ³NORCE Norwegian Research Centre, Bjerknes Centre for Climate Research, Bergen, Norway 89 ⁴Institute of Geology, Center for Earth System Research and Sustainability (CEN), Universität Hamburg, Hamburg, Germany 10 5Applied Physics Laboratory, University of Washington, Seattle, WA, USA Deleted: 5 Applied 11 Correspondence to: Shuang Gao (shuang.gao@hi.no) 12 Abstract. Riverine transport of nutrients and carbon from inland waters to the coastal and finally the open ocean 13 alters marine primary production (PP) and carbon (C) uptake regionally and globally. So far, this process has not Deleted: contribution is 14 been fully represented and evaluated in the state-of-the-art Earth system models. Here we assess changes in marine Deleted: with limited effort 15 PP and C uptake projected under the Representative Concentration Pathway 4.5 climate scenario using the 16 Norwegian Earth system model, with four riverine transport configurations for nutrients (nitrogen, phosphorus, 17 silicon and iron), carbon and total alkalinity: deactivated, fixed at a recent-past level, coupled to simulated 18 freshwater runoff, and following four plausible future scenarios. The inclusion of riverine nutrients and carbon at 19 1970's level improves the simulated contemporary spatial distribution of annual mean PP and air-sea CO2 fluxes 20 relative to observations, especially on the continental margins (5.4% reduction in root mean square error [RMSE] 21 for PP) and in the North Atlantic region (7.4% reduction in RMSE for C uptake). While the riverine nutrients and 22 C input is kept constant, its impact on projected PP and C uptake expresses differently in future period from the 23 historical period. Riverine nutrient inputs lessen nutrient limitation under future warmer conditions as 24 stratification increases, and thus lessen the projected decline in PP by up to 0.66 ± 0.02 Pg C yr⁻¹ (29.5%) globally, Deleted: future 25 when comparing 1950-1999 with 2050-2099 period. The riverine impact on projected C uptake depends on the Deleted: 7 26 balance between the net effect of riverine nutrient induced C uptake and riverine C induced CO₂ outgassing. In Deleted: These two opposite impacts are comparable in magnitudes when they are globally integrated. Therefore, in 27 the two idealized riverine configurations the <u>riverine</u> inputs result in a weak net C sink of $0.03-0.04 \pm 0.01$ Pg C Deleted: river 28 yr¹, while in the more plausible riverine configurations the <u>riverine</u> inputs cause a net C source of 0.11 ± 0.03 Pg Deleted: river 29 C yr⁻¹. It implies that the effect of increased riverine C may be larger than the effect of nutrient inputs in the future Deleted: ~ 30 on the projections of ocean C uptake, while in historical period increased nutrient inputs are considered as the Deleted: 1 31 largest driver. The results are subject to model limitations related to resolution and process representations that 32 potentially cause underestimation of impacts. High-resolution global or regional models with an adequate 33 representation of physical and biogeochemical shelf processes should be used to assess the impact of future 34 riverine scenarios more accurately.

(Formatted: Right: 0.63 cm

46 1 Introduction

47 At global scale, the major sources of both dissolved and particulate materials to the oceans are river runoff, 48 atmospheric deposition and hydrothermal inputs; of these three, river runoff plays an essential role in transporting 49 nutrients into the ocean which stimulate biological primary production (PP) in the ocean (Meybeck, 1982; Smith 50 et al., 2003; Chester, 2012). For some substances riverine transport even acts as the absolutely dominant source, 51 such as total phosphorus (~90%) and total silicon (>70%) (Chester, 2012). River transport of carbon into the ocean 52 influences the air-sea CO2 exchange, local oxygen balance and acidification level, thus further affecting marine 53 ecosystem health (Meybeck and Vörösmarty, 1999; Liu et al., 2021). Despite our limited understanding on 54 the riverine carbon fluxes, they could play an important role in closing the global carbon budget (Friedlingstein 55 et al., 2021). A recent study on global carbon cycle has emphasized the importance of the carbon transport through 56 the land-to-ocean aquatic continuum (Regnier et al., 2022). 57 With an increasing world population and a perturbed hydrological cycle under climate change, riverine transport 58 of nutrients and carbon from land to oceans has a potentially growing impact on the marine biogeochemistry and

59 ecosystem (Seitzinger et al., 2010; van der Struijk and Kroeze, 2010). Furthermore, the impacts of anthropogenic 60 activity, particularly agriculture (Bouwman et al., 2009; Garnier et al., 2021), wastewater discharges (Van Drecht 61 et al., 2009) and extensive damming (Eiriksdottir et al., 2016; Zhang et al., 2022), have greatly perturbed the 62 riverine transport of nitrogen (N), phosphorus (P) and silicon (Si) to the oceans. Seitzinger et al. (2010) estimated 63 that there was an increase in global riverine fluxes of dissolved inorganic nitrogen (DIN) and phosphorus (DIP) 64 by 35% and 29%, respectively, between 1970 and 2000, and a further possible change of -2% to +29% in DIN 65 and +37% to +57% in DIP between 2000 and 2050, depending on the future scenarios used in their study. Beusen 66 et al. (2016) estimated that river nutrient transport to the ocean increased from 19 to 37 Tg N yr¹ and from 2 to 4 67 Tg P yr⁻¹ over the 20th century, taking into account of both increased nutrient input to rivers and intensified 68 retention/removal of nutrients in freshwater systems. The riverine carbon input is highly influenced by the 69 magnitude of continental runoff (Liu et al., 2020; Frigstad et al., 2020), permafrost melting and leaching of post-70 glacial peat deposits (Wild et al., 2019; Pokrovsky et al., 2020; Mann et al., 2022), all of which are sensitive to 71 climate change. In addition, anthropogenic change, such as land-use and land-cover changes, lake and reservoir 72 eutrophication and sewage emissions of organic material into rivers may become an important factor in the future 73 (Meybeck and Vörösmarty, 1999).

74 Some regions such as the Arctic Ocean and large river estuaries may receive a higher impact from changes in 75 riverine inputs than other regions. The Arctic Ocean accounts for only 4% of the global ocean area (Jakobsson, 76 2002), but takes 11% of the global river discharge (McClelland et al., 2012), and it is estimated that about one 77 third of its net PP is sustained by nutrients originated from rivers and coastal erosion (Terhaar et al., 2021). 78 Therefore, one can expect that Arctic PP will be affected by altered riverine transport of nutrients and carbon 79 under future climate changes. Previous studies have shown that enhanced riverine nutrient input increases PP in 80 the Arctic Ocean (Letscher et al., 2013; Le Fouest et al., 2013, 2015, 2018; Terhaar et al., 2019), while large 81 riverine dissolved organic carbon (DOC) delivery reduces CO2 uptake in Siberian shelf seas (Anderson et al., 82 2009; Manizza et al., 2011). Considering large river estuaries, van der Struijk and Kroeze (2010) have 83 demonstrated potentially higher eutrophication or hypoxia risk in the coastal waters of South America by 2050,

Forn

2

Formatted: Right: 0.63 cm

.

85 enhanced N inputs will continue to dominate river DIN yields in the future and impose a challenge of N 86 eutrophication in Changijang river basin. 87 The latest generation of Earth system models (ESMs) have implemented some forms of riverine inputs in their 88 ocean biogeochemistry modules (Séférian et al., 2020). The models that include riverine inputs use different 89 implementations, from constant contemporary fluxes (e.g., IPSL-SM6A-LR and NorESM2; Aumont et al., 2015; 90 Tjiputra et al., 2020), to temporally varying fluxes (CESM2; Danabasoglu et al., 2020), and to interactive with 91 terrestrial nutrient leaching transported by dynamical river routing (e.g., CNRM-ESM2-1 and MIROC-ES2L; 92 Séférian et al., 2019; Hajima et al., 2020), and they typically use the Redfield ratio to convert from one chemical 93 compound to the others. For instance, in the latest version of IPSL model (IPSL-SM6A-LR: Aumont et al., 2015) 94 riverine nutrients (DIN, DIP, Si), dissolved organic nitrogen (DON), dissolved organic phosphorus (DOP), 95 dissolved inorganic carbon (DIC) and total alkalinity (TA) are implemented as constant contemporary fluxes 96 based on data sets from Global NEWS 2 (NEWS 2; Mayorga et al., 2010) and the Global Erosion Model of 97 Ludwig et al. (1996). Further, in the CESM2 (Danabasoglu et al., 2020) DIN and DIP are taken from the Integrated 98 Model to Assess the Global Environment-Global Nutrient Model (IMAGE-GNM; Beusen et al., 2015, 2016) and 99 vary from 1900 to 2005, which is more sophisticated than using constant fluxes. The other riverine nutrients, DIC 100 and TA are held constant using data from NEWS 2 (Mayorga et al., 2010). Some ESMs have implemented 101 interactive riverine nutrients input from terrestrial processes, e.g., in the CNRM-ESM2-1 the riverine DOC is 102 calculated actively from litter and soil carbon leaching in the land model, and the supply of the other nutrients, 103 DIC and TA have been parameterized using the global average ratios to DOC from Mayorga et al. (2010) and 104 Ludwig et al. (1996). In MIROC-ES2L model (Hajima et al., 2020), N cycle is coupled between the ocean and 105 land ecosystems, therefore, the inorganic N leached from the soil is transported by rivers and subsequently as an 106 input to the ocean ecosystem. The riverine P is calculated from N using the Redfield ratio, but riverine carbon 107 input is not implemented. Existing models with interactive riverine inputs typically do not consider 108 biogeochemical processes in the freshwater system such as sedimentation. 109 A few modelling studies have assessed the impact of riverine nutrients and carbon on marine biogeochemistry. 110 For example, Bernard et al. (2011) and Aumont et al. (2001) evaluated riverine impact on marine Si and carbon 111 cycle, respectively. Lacroix et al. (2020) estimated and implemented pre-industrial riverine loads of nutrients and 112 carbon in a global ocean biogeochemistry model, and concluded that the riverine (mainly inorganic and organic) 113 carbon inputs lead to a net global oceanic CO2 outgassing of 231 Tg C yr⁻¹ and an opposing response of an uptake 114 of 80 Tg C yr⁻¹ due to riverine nutrient inputs. Additionally, the riverine inputs at pre-industrial level lead to a 115 strong PP increase in some regions, e.g., +377%, +166% and +71% in Bay of Bengal, tropical west Atlantic and 116 the East China Sea, respectively (Lacroix et al., 2020). Tivig et al. (2021), on the other hand, found that riverine 117 N supply alone has limited impact on global marine PP (\leq +2%) due to the negative feedback of reduced N₂ 118 fixation and increased denitrification. This negative feedback could also overcompensate the N addition by river 119 supply locally, e.g., in Bay of Bengal where PP decreased due to riverine N input (Tivig et al., 2021). A couple 120 of modelling studies have also assessed the impact of changing riverine inputs on marine PP and CO₂ fluxes.

where increasing trends in DIN and DIP are detected. Yan et al. (2010) have reported that anthropogenically

121 Cotrim da Cunha et al. (2007) assessed riverine impact, using a coarse resolution ocean biogeochemistry model,

Deleted: a model (

Formatted: Right: 0.63 cm

3

.

	4		Formatted: Header
123	with single or combined nutrients from zero input to a high input corresponding to a world population of 12 billion		
124	people, and reported changes in PP from -5% to $+5\%$ for the open ocean, and from -16% to $+5\%$ for the coastal		
125	ocean, compared to the present-day simulation. Liu et al. (2021) demonstrated an increase in global coastal net		
126	PP of +4.6% response to a half-century (1961-2010) increase in river N loads. In a recent study by Lacroix et al.		
127	(2021a) the impact of changing riverine N and P in a historical period (1905–2010) on marine net PP and air-sea		Deleted: 2021
128	CO ₂ fluxes was investigated by applying an eddy-permitting fine resolution (~0.4°) ocean biogeochemistry model.		
129	Their result revealed an enhancement of 2.15 Pg C yr ⁻¹ of the global marine PP, corresponding to a relative		
130	increase of +5% over the studied period, induced by increased terrigenous nutrient inputs. The PP increase in		
131	coastal ocean averaged to 14% with regional increase exceeding 100% and the global coastal ocean CO2 uptake		
132	increased by 0.02 Pg C yr ⁻¹ due to the increased riverine nutrient inputs (Lacroix et al., <u>2021a</u>). In the Arctic,		Deleted: 2021
133	doubling riverine nutrient delivery increased PP by 11% on average and by up to 35% locally, while the riverine		
134	DOC input induced CO ₂ outgassing resulted in 25% reduction in C uptake in the Arctic Ocean (Terhaar et al.,		
135	2019).		
136	Although the historical and contemporary impacts of riverine nutrients and carbon have been considered		
137	increasingly, their impacts on future projections of marine biogeochemistry have not been sufficiently addressed.		
138	Taking advantage of the latest improvement of global river nutrient/carbon export datasets, e.g., NEWS 2		
139	(https://marine.rutgers.edu/globalnews/datasets.htm) and GLORICH		
140	(https://doi.pangaea.de/10.1594/PANGAEA.902360), and responding to the demand of development of ESMs		
141	with increasing model resolution, the assessment of the impact of riverine nutrients and carbon on projections of		Deleted: future
142	marine biogeochemistry becomes feasible and desired.		
143	In this study, we aim to assess the impact of riverine nutrients and carbon on the projected changes in regional		
144	and global marine PP and air-sea CO2 exchange by addressing the following questions:		
145	1) How does the presence of riverine fluxes of nutrient and carbon affect the contemporary representation of		
146	marine PP and C uptake in our model?		
147	2) How does the presence of riverine fluxes of nutrient and carbon affect the projections of marine PP and C		Deleted: future
148	uptake?		
149	3) How important is the consideration of transient changes in riverine fluxes of nutrient and carbon on the		
150	projections?		Deleted: future
151	We explore these questions by performing a series of transient historical and 21 st century climate simulations		
152	under the RCP 4.5 (middle-of-the-road) scenario with the fully coupled Norwegian Earth system model (NorESM)		
153	under four different riverine input configurations. Another objective of the study is to explore the best practical		
154	way of implementing riverine inputs into future versions of NorESM. Because of the coarse resolution of the		
155	version used here, a series of processes in the coastal zone cannot be represented in our study such as the high		
156	accumulation of organic sediment in shallow waters and respective remineralization rates of previously deposited		
157	material (Arndt et al., 2013; Regnier et al., 2013). These processes can only be presented in models of much higher		
158	spatial resolution, which are at present too costly to be integrated long enough to simulate the large-scale water		
159	masses adequately and project long-term scale climatic change. Given missing contributions from unresolved		
160	processes, our results are to be interpreted as lower bound estimates.		
	• · · •		Formatted: Right: 0.63 cm
	4-	and the second se	

166 2 Methods

167 2.1 Model description

168 All simulations in this study have been performed with the Norwegian Earth System Model version 1 (NorESM1-169 ME, hereafter NorESM) (Bentsen et al., 2013), a climate model that provided input to the Fifth Coupled Model 170 Intercomparison Project (CMIP5) (Taylor et al., 2011). The model is based on the Community Earth System 171 Model version 1 (CESM1) (Hurrell et al., 2013). The atmospheric, land and sea ice components are the 172 Community Atmosphere Model (CAM4) (Neale et al., 2013), the Community Land Model (CLM4) (Oleson et 173 al., 2010; Lawrence et al., 2011) and the Los Alamos National Laboratory sea ice model (CICE4) (Holland et al., 174 2011), respectively. An interactive aerosol-cloud-chemistry module has been added to the atmospheric component 175 (Kirkevåg et al., 2013). The physical ocean component-the Bergen Layered Ocean Model (BLOM, formerly 176 called NorESM-O) (Bentsen et al., 2013)-is an updated version of the Miami Isopycnic Coordinate Ocean Model 177 (MICOM) (Bleck and Smith, 1990; Bleck et al., 1992) and features a stack of 51 isopycnic layers (potential 178 densities ranging from 1028.2 to 1037.8 kg m⁻³ referenced to 2000 dbar) with a two-layer bulk mixed layer on 179 top. The depth of the bulk mixed layer varies in time and the thickness of the topmost layer is limited to 10 m in 180 order to allow for a faster air-sea flux exchange. The ocean and sea ice components are implemented on a dipolar 181 curvilinear horizontal grid with a 1° nominal resolution that is enhanced at the Equator and towards the poles, and 182 its northern grid pole singularity is rotated over Greenland. The atmosphere and land components are configured 183 on a regular 1.9° x 2.5° horizontal grid. 184 The ocean biogeochemistry component of NorESM is based on the Hamburg Ocean Carbon Cycle Model 185 (HAMOCC5) (Maier-Reimer et al., 2005). The component has been tightly coupled to NorESM-O such that both 186 components share the same horizontal grid as well as vertical layers and that all tracers are transported by the 187 physical component at model time step (Assmann et al., 2010). Tuning choices and further improvements to the 188 biogeochemistry component are detailed in Tjiputra et al. (2013). Here we only summarise features of particular 189 importance to this study, and refer to the HAMOCC version used here as HAMOCCNorESMI. The partial pressure Deleted: 190 of CO2 (pCO2) in seawater is calculated as a function of surface temperature, salinity, pressure, dissolved 191 inorganic carbon (DIC) and total alkalinity (TA). Dissolved iron is released to the surface ocean with a constant 192 fraction (3.5%) of the climatology monthly aerial dust deposition (Mahowald et al., 2005), but only 1% of this is 193 assumed to be bio-available. Nitrogen fixation by cyanobacteria occurs when nitrate in the surface water is 194 depleted relative to phosphate according to the Redfield ratio (Redfield et al., 1934). Phytoplankton growth in the 195 model depends on temperature, availability of light and on the most limiting nutrient among phosphate, nitrate 196 and iron. Constant stoichiometric ratios for the biological fixation of C, N, P and ΔO_2 (122 : 16 : 1 : -172) are 197 prescribed in HAMOCC_{NorFSM1}, and are extended by fixed Si : P (25 : 1) and Fe : P (3.66 x 10⁻⁴ : 1) stoichiometric 198 ratios. HAMOCCNorESMIA prognostically simulates export production of particulate organic carbon (POC). It is 199 assumed that a fraction of POC production is associated with diatom silica production, and the remaining fraction 200 is associated with calcium carbonate production by coccolithophorides. The fraction of diatom-associated 201 production is calculated from silicate availability, effectively assuming that diatoms are able to out-compete other 202 phytoplankton growth under favorable (high surface silicate concentration) growth conditions. Particles, including

Deleted: state-of-the-art

Deleted: HAMOCC5
Deleted: HAMOCC5

Formatted: Subscript

Formatted: Right: 0.63 cm

207 POC, biogenic silica, calcium carbonate and dust are advected by ocean circulation in the model. Those particles 208 sink through the water column with constant sinking speeds and are remineralized at constant rates. 209 HAMOCCNorESMI includes an interactive sediment module with 12 biogeochemically active vertical layers. 210 Permanent burial of particles out of the deepest sediment layer represents a net loss of POC, calcium carbonate 211 and silica from the ocean/sediment system and is compensated by atmospheric and riverine inputs on a time scale 212 of several thousand model-years. More detailed model description and parameters are documented in previous 213 publications (Bentsen et al., 2013; Tjiputra et al., 2013). 214 2.2 Model evaluation

215 The overall performance of the physical and biogeochemistry ocean components has been evaluated elsewhere

216 (Bentsen et al., 2013; Tjiputra et al., 2013). For example, simulated alkalinity, phosphate, nitrate and silicic acid

- 217 have been evaluated in previous works (Tjiputra et al., 2013; Tjiputra et al., 2020). Here we only briefly review 218
- the model performance of the mostly relevant variables for this study, namely PP and air-sea CO₂ fluxes.
- 219 The simulated global annual mean PP is 40.1 Pg C yr⁻¹ during 2003–2012, which is lower than the satellite-based 220 model estimates, ranging from 55 to 61 Pg C yr⁻¹ (Behrenfeld and Falkowski, 1997; Westberry et al., 2008).
- 221 However, the distribution of annual mean surface PP is generally consistent with the remote sensing-based 222 estimates from Behrenfeld and Falkowski (1997), with the largest model-data deviation in the eastern equatorial
- 223 Pacific and parts of the Southern Ocean (known as High-Nutrient-Low-Chlorophyll regions), where the model 224
- overestimates PP (the Arctic Ocean was not assessed in that study; Tjiputra et al., 2013). Along the continental 225
- margins, the simulated PP is generally underestimated compared to the remote sensing-based estimates (Tjiputra 226 et al., 2013), which may relate to the lack of riverine inputs and/or unresolved shelf processes due to coarse model
- 227 resolution. Additionally, our model simulates a comparable magnitude of projected decrease in PP, by the end of
- 228 the 21th century compared to historical period, with other global models (see detailed discussion in section 4.1).
- 229 In the Arctic Ocean, the simulated PP in our model is biased towards lower values. In the study by Skogen et al. 230 (2018), the NorESM model is compared with a regional model that comprises part of the Arctic region, and it
- 231 shows that the NorESM simulates too late and too short bloom period than the regional model, hence the annual 232 integrated PP is too low. In a multi-model study (Lee et al., 2016) that assesses the relative skills of 21 regional 233 and global biogeochemical models in reproducing the observed contemporary Arctic PP, the NorESM is shown
- 234 to have a negative bias of -0.49, but is well within the multi-model mean bias of -0.31±0.39. Many
- 235 coarse/intermediate resolution global models also show considerably lower net PP in the Arctic (Terhaar et al., 236 2019). Such common shortcomings in global scale marine biogeochemical models can partly be attributed by the
- 237 simplified, not regionally adapted ecosystem parameterization, which can be improved through data assimilation
- 238 (Tjiputra et al., 2007; Gharamti et al., 2017). Additionally, lack of adequate representation of riverine input in
- 239 some ESMs can also lead to underestimate of PP, since around one third of current Arctic marine PP is sustained
- 240 by terrigenous nutrient input (Terhaar el al., 2021). Despite the biased low PP under the contemporary climate,
- 241 the projected absolute change of 70 Tg C yr⁻¹ by the end of the 21th century is well within the range estimated
- 242 from other ESMs (Vancoppenolle et al., 2013).

Deleted: HAMOCC5

Formatted: Header

Deleted: in good agreement

6

Tjiputra et al. (2013) also evaluated the simulated mean annual sea-air CO₂ fluxes for the 1996–2005 period against observational-based estimates by Takahashi et al. (2009) and concluded that the model broadly agrees with the observations in term of spatial variation, although in the equatorial Indian Ocean and in the polar Southern

248 Ocean (South of 60° S) the model underestimates outgassing and overestimates C uptake, respectively.

249 2.3 Riverine data

250 The influx of carbon and nutrients from over 6000 rivers to the coastal oceans has been implemented in 251 <u>HAMOCC_{NorESML}</u> based on previous work of Bernard et al. (2011) but with modifications that are outlined in the

following paragraphs.The riverine influx includes carbon, nitrogen and phosphorus, each in dissolved inorganic, dissolved organic, and

254 particulate forms, as well as TA, dissolved silicon and iron (Fe). Except for DIC, TA and Fe, all data are provided 255 by the NEWS 2 model (Mayorga et al., 2010), which is a hybrid of empirical, statistical and mechanistic model 256 components that simulate steady-state annual riverine fluxes as a function of natural processes and anthropogenic 257 influences. The NEWS 2 data product contains historical (year 1970 and 2000) and future (year 2030 and 2050) 258 estimates of riverine fluxes of carbon and nutrients. The future products are developed based on four Millennium 259 Ecosystem Assessment scenarios (Alcamo et al., 2006): Global Orchestration (GNg), Order from Strength (GNo), 260 Technogarden (GNt) and Adapting Mosaic (GNa). These scenarios represent different focuses of future society 261 on e.g., globalization or regionalization, reactive or proactive environmental management and their respective 262 influences on efficiency of nutrient use in agriculture, nutrient release from sewage, total crop and livestock 263 production along with others (see Table 1 for a brief summary; Seitzinger et al., 2010). The NEWS 2 riverine 264 dataset has been calibrated and assessed against measured yields (Mayorga et al., 2010) and has been widely used 265 and evaluated for different river estuaries (van der Struijk and Kroeze, 2010; Terhaar et al., 2019; Tivig et al., 266 2021). For example, van der Struijk and Kroeze (2010) compared the NEWS 2 nutrient yields to observed values 267 for South American rivers and indicated that the NEWS 2 models in general perform reasonably well for South 268 American rivers with the variations in yields among rivers described well, although the model performs better for 269 some rivers such as the Amazon than for others. We have compared DIN and dissolved organic nitrogen (DON) 270 from NEWS 2 with measured data from PARTNERS Project (Holmes et al., 2012) for the six largest Arctic rivers 271 around year 2000 (Table C1). The NEWS 2 dataset compares fairly well with the measured data, especially for 272 the Eurasian Arctic rivers with 3.5-28.6% deviation in DIN and 7.3-34.8% in DON, while the discrepancy is larger 273 in the Canadian-Alaska Arctic rivers (i.e., Yukon and Mackenzie rivers) with up to 80.8% and 100% deviation in 274 DIN and DON, respectively. 275 The DIC and TA fluxes, provided by Hartmann (2009), are produced from a high-resolution model for global 276 CO2 consumption by chemical weathering and are aggregated within catchment basins defined by the NEWS 2

277 study for each river, Riverine Fe flux is calculated as a proportion of a global total input of 1.45 Tg yr⁻¹ (Chester,

1278 1990), weighted by water runoff of each river. Only 1% of the riverine Fe is added to the oceanic dissolved Fe,

- 279 under the assumption that upto 99% of the fluvial gross dissolved Fe is removed during estuarine mixing (Boyle
- et al., 1977; Figuères et al., 1978; Sholkovitz and Copland, 1981; Shiller and Boyle, 1991).

Deleted: HAMOCC5

Deleted: upto

-(Deleted: implemented to
•(Formatted
(Formatted
1	Deleted: basin map

285 At the river mouths, all fluxes are interpolated to the ocean grid in the same way as the freshwater runoff, which 286 is distributed as a function of river mouth distance with an e-folding length scale of 1000 km and cutoff of 300 287 km. 288 In HAMOCC_{NorESM1}, there is one dissolved organic pool (DOM) and one particulate organic pool (DET, detritus). 289 First, we calculate the riverine organic P-N-C ratios for both dissolved and particulate forms, then add the least 290 abundant species (scaled by the Redfield ratio) to the DOM and DET pools, respectively, (see equations below). $DOM_{riv} = \min(DOP, \frac{DON}{16}, \frac{DOC}{122})$ 291 _____(1) $DET_{riv} = \min\left(POP, \frac{PON}{16}, \frac{POC}{122}\right)$ 292 (2) 293 POP and PON denote particulate organic phosphorus and particulate organic nitrogen, respectively, The excess 294 budget from the remaining two species both in dissolved and in particulate forms are assumed to be directly

remineralized into inorganic form and added to the corresponding dissolved inorganic pools (i.e., DIP, DIN, and

Formatted: English (US)

Deleted: HAMOCC5

Deleted:

297 2.4 Experimental design

fluxes are deactivated.

DIC) in the ocean.

298 The fully coupled NorESM model is spun up for 900 years with external forcings fixed at preindustrial year-1850 299 levels prior to our experiments (Tjiputra et al., 2013). The atmospheric CO2 mixing ratio is set to 284.7 ppm 300 during the spin-up. Nutrients and oxygen concentrations in the ocean are initialised with the World Ocean Atlas 301 dataset (Garcia et al., 2013a, b). Initial DIC and TA fields are taken from the Global Data Analysis Project (Key 302 et al., 2004). After 900 years, the ocean physical- and biogeochemical tracer distributions reach quasi-equilibrium 303 states. We extended the spin-up for another 200 years with riverine input for each experiment (except for the 304 reference run) and then performed a set of transient climate simulations for the industrial era and the 21st century 305 (1850-2100). The simulations use external climate forcings that follow the CMIP5 protocol (Taylor et al., 2011). 306 For the historical period (1850-2005), observed time-varying solar radiation, atmospheric greenhouse gas 307 concentrations (including CO₂), natural and anthropogenic aerosols are prescribed. For the future period (2006-308 2100), the Representative Concentration Pathway (RCP) 4.5 (van Vuuren et al., 2011) is applied. Here, we 309 consider RCP4.5 as the representative future scenario following the CO2 emission rate based on the submitted 310 Intended Nationally Determined Contributions, which projects a median warming of 2.6-3.1°C by 2100 (Rogelj 311 et al., 2016). The riverine input configurations employed in this study are summarized in Figure 1. The evolution 312 of global total fluxes of each nutrient/carbon species are shown in Figure 2. The experiment configurations are 313 described as follows: 314 REF: Reference run. Riverine nutrient and carbon supply is deactivated. 315 FIX and FIXnoc: Fixed at recent-past level. FIX: A constant riverine nutrient and carbon supply, 316 representative for the year 1970 as provided by NEWS 2, is applied to the model throughout the whole 317 experiment duration. FIXnoc: As FIX but only with nutrients supply, all carbon (DIC, DOC, POC) and TA 318

319 RUN: Coupled to simulated freshwater runoff. Riverine nutrient and carbon supply representative for the 320 year 1970 is linearly scaled with the on-line simulated freshwater runoff divided by the climatological mean 321 runoff over 1960-1979 of the model. Thus, the inputs follow the seasonality and long-term trend of the

Formatted: Right: 0.63 cm

8

295

simulated runoff. We assume that the nutrient and carbon concentrations in the rivers are constant at the
level of 1970, but the fluxes fluctuate with freshwater runoff.
GNS: Four different transient inputs following future projections of NEWS 2. A constant riverine nutrient

and carbon supply representative for year 1970 has been applied from year 1850 to 1970. Between year 1970,
2000, 2030 and 2050 the annual riverine supply is linearly interpolated. From year 2050 to 2100 the annual
riverine supply is linearly extrapolated. From year 2000, riverine supplies of the four NEWS 2 future
scenarios (GNa, GNg, GNo and GNt) are applied.

By comparing FIX versus REF, we assess how the presence of riverine inputs affect the contemporary marine PP and C uptake representation and also the projected changes. By comparing RUN versus FIX we assess the potential effects of riverine nutrient and carbon long-term trends associated with an intensifying global hydrological cycle on marine PP and C uptake. RUN represents a first step towards coupling riverine nutrient and carbon fluxes to the simulated hydrological cycle. By comparing the GNS configurations versus FIX we assess how plausible, realistic future evolutions in riverine nutrient and carbon fluxes may impact marine PP and C uptake projections. We span the uncertainty in future riverine nutrient and carbon fluxes by considering multiple

338 NEWS 2 scenarios.

339 3 Results

340 3.1 Effect of including riverine inputs on contemporary marine PP and C uptake

We start with assessing how the inclusion of riverine nutrients and carbon affects the contemporary representation of the global marine PP and C uptake in our model by comparing the annual mean output over the years 2003– 2012 between the REF and FIX experiments. We also compare with satellite and observational based estimates to see if the inclusion of riverine nutrients and carbon improves the marine PP and C uptake representation in our model. The spatially integrated values presented in this and following sections are summarized and supplemented with statistical robustness information in Tables B1 and B2 in Appendix B.

The annual net primary production (PP) is 40.1 and 43.0 Pg C yr⁻¹ in the REF and FIX experiments, respectively. 347 348 The increase of PP in FIX occurs along continental margins (where seafloor is shallower than 300 m) and also in 349 the North Atlantic region (0°N-65°N, 0°W-90°W), accounting for 15.4% and 24.9% of the global total increase, 350 respectively (Figure 3c). The simulated global total PP in both REF and FIX are lower than the satellite-based 351 model estimates, including Vertically Generalized Production Model (VGPM), Eppley-VGPM and Carbon-based 352 Production Model (CbPM) over the same time period (data source: 353 http://www.science.oregonstate.edu/ocean.productivity), ranging from 55 to 61 Pg C yr⁻¹ (Behrenfeld and 354 Falkowski, 1997; Westberry et al., 2008). Although the total PP in FIX is still considerably lower than the satellite-355 based estimates, the inclusion of riverine nutrients and carbon does slightly improve the distribution of PP 356 especially on continental margins (Figure 3), according to our area-weighted root mean square error (RMSE) 357 analysis. The RMSE of REF relative to mean observational estimates (mentioned above) averages 10.7 mol C m⁻ 358 ² yr⁻¹ globally, while the value of FIX is 10.3 mol C m⁻² yr⁻¹, which is reduced by 3.7%. For the continental margins, 359 the RMSE is reduced by 5.5% from 29.0 mol C m^{-2} yr⁻¹ in REF to 27.4 mol C m^{-2} yr⁻¹ in FIX.

0

360	The ocean annual net uptake of CO2 is 2.8 and 2.9 Pg C yr ⁻¹ in REF and FIX, respectively, with a FIX-REF	
361	difference of 0.1 Pg C yr ⁻¹ equivalent to 3.1% relative change, which is statistically significant (see Table B2). In	
362	FIX the ocean carbon uptake is generally enhanced everywhere except for the upwelling regions of the Southern	
363	Ocean and in the subpolar North Atlantic between approximately 50°N-65°N and 60°W-10°W (Figure 4c). To	
364	isolate the impact of riverine nutrients input from carbon input, an additional experiment (FIXnoc) was conducted,	
365	where the nutrient fluxes are implemented the same as in FIX, while all carbon (DIC, DOC, POC) and TA fluxes	
366	are eliminated. As shown in Figure 4d, the nutrients input results in more CO2 uptake not only at large river	
367	estuaries but also in the subtropical gyres due to enhanced primary production. In the subpolar North Atlantic and	
368	in the Southern Ocean upwelling region, the addition of riverine nutrients leads to enhanced outgassing. The	
369	riverine carbon input, on the other hand, leads to CO2 outgassing mainly at river estuaries (Figure 4e), but also in	
370	a band along the gulf stream extending into the North Atlantic, where it accounts for 18.1% of the CO2 outgassing	
371	in the subpolar region (50°N-65°N, 60°W-10°W). Along the continental margins the nutrients input increases the	
372	CO2 uptake, while the carbon input has an opposite effect which induces more outgassing. The net effect of both	
373	nutrient and carbon inputs shows that the uptake of CO2 dominates over the outgassing, along the continental	
374	margins and in subtropical gyres (Figure 4c). Compared to the observational based estimates of Landschützer et	
375	al. (2017) (Figure 4a) and according to our RMSE analysis, the inclusion of riverine nutrients and carbon does	
376	not improve the simulated air-sea CO2 fluxes globally. The RMSE of FIX relative to observational estimates	
377	averages to 0.83 mol C m ⁻² yr ⁻¹ globally, which does not differ much from the value of REF (0.84 mol C m ⁻² yr ⁻¹	
378	¹). However, there is a distinguishable improvement of the distribution of air-sea CO ₂ fluxes in the subpolar North	
379	Atlantic (RMSE is reduced by 8.2%, from 0.73 mol C m ⁻² yr ⁻¹ in REF to 0.67 mol C m ⁻² yr ⁻¹ in FIX), with slight	
380	degradations in some other regions (Figure 4c).	
381	3.2 Effect of including contemporary riverine inputs on projections of marine PP and C uptake	Deleted: future
382	We now address how the inclusion of riverine nutrient and carbon fluxes affects projections of marine PP and C	Deleted: future
383	uptake by comparing the average output between a future period (2050-2099) and a historical period (1950-1999)	
384	of FIX versus REF.	
385	In both experiments the projections of global PP averaged over the years 2050-2099 are lower than their	Deleted: future
386	corresponding 1950-1999 averages (Figure 5a). However, when riverine input of nutrient and carbon is included,	
387	the projected decrease of global PP is mitigated from -2.2 Pg C yr ⁻¹ in REF to -1.9 Pg C yr ⁻¹ in FIX (by 13.6%).	
388	Spatially, the decrease of PP in REF occurs largely in upwelling regions such as the tropical eastern Pacific and	
389	tropical Atlantic, as well as along a latitude band around 40°S (Figure 6a). The riverine inputs alleviate the	
390	projected PP decrease in those regions (see further discussion in Section 4.2) and reinforce the projected PP	
391	increase in high latitudes (Figure 6b, c). The projections of PP in the Arctic Ocean show significant increases in	Deleted: future
392	both REF and FIX. Climate change alone (REF, without riverine inputs) almost doubles the simulated PP in the	
393	Arctic from 0.08 Pg C yr ⁻¹ during 1950–1999 to 0.15 Pg C yr ⁻¹ in 2050–2099 (Figure 5b), likely as a consequence	
394	of sea ice retreat. FIX, which includes riverine inputs, exhibits a slightly larger (but significant, see Table B1)	
395	absolute Arctic PP increase (from 0.10 to 0.18 Pg C yr ⁻¹) in its future projection than REF.	
		Formatted: Right: 0.63 cm

10

rmatted: Right: 0.63

For global net uptake rate of CO₂, both experiments (REF and FIX) project a significant increase under the RCP4.5 (Figure 7a). The inclusion of riverine inputs leads to a slightly higher (but significant, see Table B2) (2.4%) projected increase of 1.28 Pg C yr⁻¹ in FIX compared with 1.25 Pg C yr⁻¹ in REF. The increase rate of CO₂ uptake in the Arctic closely follows the global trend (Figure 7b). Spatially, there is a widespread simulated increase in ocean uptake of CO₂ under future climate change except in the subtropical gyres (Figure 8a). Riverine nutrients input slightly increases the projected carbon uptake at large river estuaries, while decreases the projected uptake in subpolar North Atlantic (Figure 8d).

407 **3.3** Effect of future changes in riverine inputs on marine PP and C uptake projections

Finally, we address how future changes in riverine fluxes of nutrients and carbon affect marine PP and C uptake
by comparing the projected changes for the time period 2050–2099 relative to 1950–1999 among FIX, RUN and
the four GNS experiments.
The future projected decrease of PP in the four GNS averages to -1.6 Pg C yr⁻¹, which is less in magnitude

412 compared to FIX (-1.9 Pg C yr⁻¹) and RUN (-1.8 Pg C yr⁻¹) (Figure 5a). Spatial distributions of projected PP 413 changes in GNS and their respective differences relative to FIX are shown in Figure 9. The latter occur 414 predominantly on the continental shelf in Southeast Asia, where the future projected increase in riverine nutrient 415 load is the largest in the world in GNS (Seitzinger et al., 2010). Interestingly, the projected increase in PP in 416 Southeast Asia, induced by riverine nutrient inputs in GNS, is of the same order of magnitude as the projected 417 decrease in PP due to future climate change in REF. Thus, in GNS the PP are projected to slightly increase on the

418 continental shelf of Southeast Asia (Figure 9a-d). The riverine nutrient induced PP increase in FIX or RUN is not

419 large enough to compensate the PP decline due to climate change, since the projected changes in riverine nutrient

420 inputs are not taken into account in FIX or locally underestimated in RUN.

421 On the other hand, the future projected global uptake of CO_2 in GNS (1.13 Pg C yr⁻¹ in average) is reduced

 $422 \qquad \text{compared to REF (1.25 Pg C yr^{-1}), which shows an opposite change than FIX (1.28 Pg C yr^{-1}) and RUN (1.29 Pg C yr^{-1})}$

423 C yr⁻¹). The changes in riverine inputs in GNS emerge along continental margins, especially around large river

424 estuaries (Figure 10e-h), where the dissolved organic matter (DOM), that is projected to increase in GNS, enters

 $425 \qquad \text{the ocean and releases CO}_2 \text{ to the atmosphere (Seitzinger et al., 2010)}.$

426 Despite the regional differences, there is no significant difference in the projected changes in either globally 427 integrated PP or CO₂ uptake among the four GNS in our model (Figures 5 and 7, see further discussion in Section

428 4.3).

429 4 Discussion

430	4.1 Projected marine PP and C uptake under climate change		Deleted: changes
431 432	Jn our model, PP is roughly linearly related to the concentrations of the most limiting nutrient (Nut), light intensity (I), temperature (T) and the available phytoplankton concentration (Phy), i.e., $PP \sim Nut \cdot I \cdot f(T) \cdot Phy$. It is shown		Deleted: The projected global total PP shows up to 29.5% less decrease, if riverine inputs are present in the model. This
433	in Figure 6a that under climate change the projected decrease in PP occurs mainly in low- and mid-latitudes.		is mainly because the riverine nutrient inputs into the surface ocean alleviates the increasing nutrient limitation caused by stronger stratification under future climate warming.
434	Nitrate is the limiting nutrient (in REF) in almost everywhere except in the Central indo-Pacific region, in the	(Formatted: Right: 0.63 cm

I

••• Formatted: Header

441 South Pacific subtropical gyre, in the Bering Sea and part of the Arctic, where Fe is limiting (Figure A1). Projected 442 reduction in surface nitrate concentrations (Figure A2b), which is tightly linked to the upper ocean warming and 443 increased vertical stratification (Bopp et al., 2001; Behrenfeld et al., 2006; Steinacher et al., 2010; Cabré et al., 444 2015), contributes to the projected decrease in PP in our model. The simulated global mean PP over 2050-2099 445 is 38.9 Pg C yr⁻¹ in REF, which is 2.24 Pg C yr⁻¹ lower than the value over 1950–1999. This -5.4% projected 446 change in PP is comparable with the multi-model mean estimate of projected change of $-3.6 \pm 5.7\%$ in the 2090s 447 relative to the 1990s for RCP4.5 (Bopp et al., 2013) and sits in the range of 2-13% decrease projected by four 448 ESMs over the 21st century under the SRES A2 scenario (Steinacher et al., 2010). It is also still within the range of the 13 multi-model mean projected PP change of -1.13 \pm 5.81% under the CMIP6 Shared Socioeconomic 449 450 Pathways SSP2-4.5 when comparing mean values in 2080-2099 relative to 1870-1899 (Kwiatkowski et al., 2020), 451 given that the inter-model uncertainties in projected PP have increased in CMIP6 compared to CMIP5 (Tagliabue 452 et al., 2021). In contrast to the global PP, there are considerable increases in the future projected PP in the Arctic 453 in REF (Figure 5b). In polar regions light and temperature are the primary limiting factors for phytoplankton 454 growth, therefore PP increases when light and temperature become more favourable owing to sea-ice melting 455 under warmer conditions (Sarmiento et al., 2004; Bopp et al., 2005; Doney, 2006; Steinacher et al., 2010). On the 456 other hand, the fresher and warmer surface water increases stratification, prohibiting nutrients upwelling 457 (Vancoppenolle et al., 2013; Figure A2b), which counteracts the increase in PP. 458 The ocean annual net uptake of CO2 increases significantly during 2050-2099 compared with the uptake during 459 1950-1999 in REF (Figure 7a), which is mainly driven by increasing difference in air-sea partial pressure of CO2. 460 4.2 Changes in projected marine PP and C uptake due to riverine input 461 When riverine nutrient fluxes are added into coastal surface waters in FIX, the PP is higher in both historical and 462 future periods compared to REF (Figure 5a), due to alleviated nutrient limitation. Interestingly, the effect of 463 riverine inputs on PP for the historical and future time periods is not the same, suggesting a different nutrient 464 depletion level (Figure A2b). The projected decrease in PP is lessened from -5.4% in REF to -4.4% in FIX. It

465 implies that during 1950-1999 the riverine nutrients are not depleted by primary producers, while during 2050-466 2099 the riverine nutrients are utilized to a greater extent due to the exacerbated nutrient limitation (Figure A2b) 467 and potentially to higher phytoplankton growth rate in warmer climate. Figure 12 illustrates this in a schematic 468 diagram that shows the impact of riverine nutrients on projected PP in low- and mid-latitudes. Moreover, the 469 inclusion of constant riverine inputs (FIX) can potentially explain one tenth of the ~10% (2-13%, Steinacher et 470 al., 2010) inter-model spread, In RUN and GNS, the projected decline in PP is further alleviated to -4.1% and -471 3.6% (averaged over four GN scenarios), respectively, compared to -4.4% in FIX, owing to the varying (mostly 472 increase) nutrients input. In the Arctic, when riverine nutrients input is present in the model, it helps to sustain the

- projected PP increase against the stronger stratification under future climate warming, although this effect is only
- 474 minor (Figure 5b)
- The riverine inputs have a two-fold effect on the ocean C uptake. It is the competition between the riverine
 (inorganic and organic) nutrients input induced CO₂ uptake and the riverine carbon input induced CO₂ outgassing,
 which determines whether the shelf is a C sink or a C source. However, the composition of the riverine organic

(Moved (insertion) [1]

Deleted:

In contrast to the global PP, there are considerable increases in the future projected PP in the Arctic in REF (Figure 5b). In polar regions light and temperature are the primary limiting factors for phytoplankton growth, therefore PP increases when light and temperature become more favourable owing to sea-ice melting under warmer conditions (Sarmiento et al., 2004; Bopp et al., 2005; Doney, 2006; Steinacher et al., 2010). On the other hand, the fresher and warmer surface water increases stratification, prohibiting nutrients upwelling (Figure A2b), which counteracts the increase in PP. Therefore

Deleted: in the Arctic

Moved up [1]: The ocean annual net uptake of CO₂ increases significantly during 2050–2099 compared with the uptake during 1950–1999 in REF (Figure 7a), which is mainly driven by increasing difference in air-sea partial pressure of CO₂.

Deleted:

12-

Formatted: Right: 0.63 cm

ļ

• (Formatted: Header

496 matter (i.e., carbon to nutrient ratio) and the degradation timescales which are the key factors, have been debated 497 over the last three decades (Ittekkot, 1988; Hedges et al., 1997; Cai, 2010; Bianchi, 2011; Blair and Aller, 2011; 498 Lalonde et al., 2014; Galy et al., 2015). It is generally agreed that the riverine organic carbon to nutrient ratio is 499 high (e.g., C:P weight ratio larger than 700, Seitzinger et al., 2010) and the degradation and resuspension rates in 500 shallow shelf seas/sediment are higher than the open ocean (Krumins et al., 2013). It suggests that at shallow and 501 near-shore areas the riverine carbon input usually results in a CO2 source to the atmosphere, while at deeper outer 502 shelf areas the riverine nutrient input causes PP increase and a CO2 sink, and the magnitudes of the C source and 503 sink on the continental shelves almost compensate each other. This phenomenon has been discussed by both 504 measurement-based studies (Borges and Frankignoulle, 2005; Chen and Borges, 2009) and modelling studies (e.g., 505 Lacroix et al., 2020). However, the spatial resolution in our model is not fine enough to differentiate the near-506 shore and outer shelf processes. This partly contributes to comparable CO2 outgassing near shore (due to riverine 507 C) and CO2 ingassing on outer shelves (due to riverine inorganic and organic nutrients input), leading to a globally 508 weak integrated C sink on the continental margins in FIX and RUN experiments for both historical and future 509 time periods. Although the riverine input of nutrients and C are constant for both time periods in FIX, the riverine 510 induced C uptake is slightly (but significantly) bigger (0.03 Pg C yr⁻¹) during 2055–2099 compared to 1950–1999, 511 which indicates that the riverine nutrients input is slightly dominant over riverine C input in FIX, and the riverine 512 nutrients are utilized more in the future period. A recent modelling study (Lacroix et al., 2021a) with improved Deleted: On the other hand, in GNS 513 shelf processes, has also reported a 0.03 Pg C yr⁻¹ increase in global C uptake induced by temporally varying 514 terrestrial nutrients input during 1905-2010. They conclude that due to large historical perturbation, the increased 515 nutrient inputs are the largest driver of change for the CO₂ uptake at the regional scale. In GNS, on the other hand, 516 the riverine inputs reduce globally integrated C uptake for both historical and future time periods, but not equally, Deleted: 517 (Figure 7a). It reduces more in the future period (2050-2099) than the historical period (1950-1999), which 518 implies that the effect of riverine C input in the future scenarios are more dominant over nutrients input. 519 Simulations with high-resolution global or regional models with more realistic representation of shelf processes Deleted: A recent modelling study (Lacroix et al., 2021), which uses a finer resolution (~0.4°) global model with 520 are required to accurately assess the impact of riverine inputs on carbon cycling in the coastal ocean. improved shelf processes, has also reported a 0.03 Pg C yr⁻¹ increase in global C uptake induced by terrestrial nutrients input during 1905-2010, although they have applied 521 4.3 Sensitivity of projected marine PP and C uptake to riverine configuration temporally varying (increasing) nutrients and no riverine C input. ... 522 By exploring different riverine configurations (FIX, RUN, GNS) we investigate how uncertainties in future Deleted: 2 Different 523 riverine fluxes translate into uncertainties in projected PP and C uptake changes. In RUN we assume constant **Deleted:** configurations 524 concentrations (at 1970's level) of riverine nutrient and carbon over time and couple them to the simulated 525 freshwater runoff. Thus, the annual global total fluxes of nutrient and carbon vary with time following the 526 variability of runoff (Figure 2), in contrast to the constant fluxes in FIX. The global total simulated runoff, under RCP4.5 in our model, is on average higher during 2050-2099 than the runoff during 1950-1999, indicating an 527 528 intensified hydrological cycle under future climate change. Hence, the global riverine fluxes of nutrient and carbon 529 during 2050-2099 are higher than those during 1950-1999 in RUN. However, the temporal changes in global 530 riverine fluxes in RUN are relatively small compared with the absolute flux values in FIX, which explains the

slightly larger projected changes in global PP and ocean carbon uptake in RUN compared to FIX. It is noteworthy

that the large inter-annual variability in the riverine fluxes of nutrient and carbon in RUN does not increase the

531

532

13

----- Formatted: Header

544 inter-annual variability in simulated PP and ocean carbon uptake either globally or on the continental margins 545 (Figure 11), something that warrants further investigation. The approach of RUN serves as a trial to introduce 546 seasonal and inter-annual variability in riverine nutrient and C inputs that is linked to hydrological variability. It 547 should be explored in future works if RUN and GNS can be integrated to produce more realistic long-term trends 548 in riverine nutrient and C inputs as well as short-term variability. Although the RUN approach is more 549 sophisticated when compared to FIX, it employs a linear relationship between the future riverine nutrient and C 550 fluxes and the simulated hydrological cycle, which is a highly simplified assumption (see discussion in section 551 44). 552 Figure 2 shows that the inputs of DIN and DIP are considerably lower, while the dissolved silicon (DSi) and 553 particulate organic matter (POM) are higher in the future period in RUN compared to GNS. This is because many 554 anthropogenic processes that are important for determining the future riverine fluxes are not considered in RUN, 555 but are considered in NEWS 2 model system, from which the GNS' future scenarios are simulated. For example, 556 the nutrient management in agriculture, the sewage treatment and phosphorus detergent use, and the increased 557 reservoirs from global dam construction in river system (Seitzinger et al., 2010; Beusen et al., 2009) are the key

558 factors affecting future riverine fluxes of DIN, DIP, and DSi/POM, respectively. Therefore, it is worth exploring 559 the merits of using GNS in future projections of marine biogeochemistry. The four future scenarios provide a 560 range of potential outcomes resulting from different choices tending toward either globalization or regional 561 orientation, either reactive or proactive approach to environmental threats (see Table 1). A large range of the 562 riverine inputs in GNS, e.g., temporal changes in DIN fluxes across scenarios ranging 24.8-63.0% of the annual 563 flux in FIX, do not transfer to large uncertainties in future projections of global marine PP in our model, which 564 can primarily be attributed to unresolved shelf processes due to coarse model resolution. However, the scenario 565 differences might be of importance in regional projections, such as in seas surrounded by highly populated nations 566 and near river estuaries. Simulations with high-resolution global or regional models with a good representation of

shelf processes are required to accurately assess the local impact of riverine inputs.

568 4.4 Limitations and uncertainties

567

569 We acknowledge several limitations of our study, particularly related to the resolution and complexity of our 570 model. Firstly, coarse-resolution models tend to underestimate PP along the coast. Such well-known model issues 571 may offset the impact induced by riverine inputs. Secondly, shelf processes, which are not well represented in our 572 model due to coarse resolution, modify a large fraction of some riverine species, e.g., conversion of organic carbon 573 to CO2 occurs rapidly via remineralization in estuaries before they are transported to the open ocean. Further, 574 some simplified processes of the model may introduce bias in the results, e.g., how the model deals with the 575 riverine dissolved organic and particulate matter. In our model, there is only one dissolved organic pool (DOM) 576 and one particulate organic pool (DET), and the Redfield ratio (P-N-C) needs to be kept. Therefore, the P-N-C 577 ratios of riverine input for both dissolved organic matter (including DON, DOP and DOC) and particulate 578 (inorganic and organic) matter (including particulate nitrogen, particulate phosphorus and POC) are calculated, 579 then the least abundant species (scaled by the Redfield ratio) are added to the DOM and DET pools, respectively. 580 The excess budget from the remaining two species (of P, N or C) are assumed to be directly remineralized into

Deleted: 3

Deleted: 3

Deleted: Given that the riverine nutrient and carbon inputs account for only a small proportion of the total amount of nutrients and carbon in the euphotic zone of the ocean, we

Deleted: and resolution

Deleted: ESM

14

588 inorganic form and added to the corresponding dissolved inorganic pools (i.e., DIP, DIN, or DIC) in the ocean. 589 This simplification may result in overestimation of riverine dissolved inorganic nutrients and thereby riverine 590 induced PP enhancement. Especially, in NEW 2 dataset particulate P is typically dominated by inorganic forms 591 (Mayorga et al., 2010), which means that it is likely not directly bio-available. Therefore, we have assessed the 592 bias due to the direct remineralization of the riverine dissolved organic and particulate matter. We calculated 593 firstly the proportion of directly remineralized matter from the total riverine dissolved organic matter (DOM) and 594 particulate (inorganic and organic) matter (PM) by using the following equation, i.e., [X/(DOM_{riv}+PM_{riv})*100%] 595 (X is the directly remineralized dissolved organic and particulate matter). The directly remineralized part on 596 average accounts for 64.8%, 27.8% and 62.8% of the total riverine organic and particulate matter of P, N and C, 597 respectively. In a recent study by Lacroix et al. (2021b) who used an enhanced version of HAMOCC (horizontal 598 resolution of $\sim 0.4^{\circ}$) with improved representation of riverine inputs and organic matter dynamics in the coastal 599 ocean, they quantified that around 50% of the riverine DOM and 75% of the POM are mineralized in global shelf 600 waters. Therefore, our model assumption is on track with the finer-resolution-model estimates and this direct 601 remineralization compensates to some extent the under-represented organic matter degradation rate on the ocean shelf. This bias in riverine dissolved nutrient input may further lead to bias in the enhanced PP. We calculated the 602 603 contribution of the directly remineralized part on the enhanced PP, by comparing X with the corresponding total 604 riverine dissolved nutrient additions as [X/(X+DIXriv)*100%] (DIXriv denotes the corresponding riverine 605 dissolved nutrient additions), which accounts for 80.5%, 33.3%, and 41.1% for P, N, and C, respectively. 606 Assuming that all coastal regions are nutrient limited, this direct remineralization could be theoretically 607 responsible for 33.3%-80.5% of the enhanced PP, depending on which nutrient species is limiting the PP. In our 608 model, phosphate is rarely limiting (Figure A1), therefore, the impact of this direct remineralization on PP is likely 609 on the lower end of this range (33.3%-80.5%). Given that the proportion of the direct remineralized organic N 610 (27.8%, see the calculation above) in our model is comparable to or lower than the reported values by field studies 611 (~38.8% of DON decomposed during transition from Arctic rivers to coastal ocean; Kattner et al., 1999; Lobbes 612 et al., 2000; Dittmar et al., 2001), which indicates that the bias on enhanced PP is likely less than 33.3%. 613 Some approximation and assumption in the experimental setup may also induce uncertainties in our results. Our 614 spin-up experiment uses riverine nutrient and carbon inputs fixed at 1970 levels, as provided by NEWS 2. As a 615 caveat, our post-1970 simulated changes in marine PP and CO2 fluxes miss out any legacy effects from riverine 616 input changes that occurred before 1970. The fixed inputs likely overestimate the accumulated inputs prior 1970, 617 causing potential underestimation of the projected change impacts. However, Beusen et al. (2016) found that 618 changes in riverine N and P are relatively small before 1970 compared to changes after 1970. Therefore, we expect 619 the impact due to missing legacy effects to be minor. Moreover, in FIX we applied riverine inputs at 1970 level 620 over available inputs at 2000 level, because the former are more representative for the 1950-1999 baseline period. 621 However, the use of 1970 level input is suboptimal when evaluating simulated PP and CO₂ fluxes against 622 observations obtained after 2000. Beusen et al. (2016) have shown that the riverine N and P has increased by 623 ~40.0% and 28.6%, respectively, from 1970 to 2000. Therefore, the riverine impact may be underestimated when 624 comparing with the observations during 2003-2012. In RUN, we assume constant concentrations of riverine 625 nutrient and carbon over time and the fluxes vary with freshwater runoff. This may be applicable for some

Deleted: 2021

Deleted: matters
Deleted: those
Deleted: Lacroix
Deleted: . (2021
Deleted: there is a considerable fraction of organic matters that remineralize in shelf waters,

15

633 nutrients such as DIN or within a certain limit of runoff change such as for dissolved Si (Figure A3). However, 634 this may not be appropriate for all nutrient/carbon species. Furthermore, the variability of runoff is subject to 635 inter-annual to decadal climate variability, which partially masks the centennial trend. This caveat can be 636 overcome through performing multi-realization ensemble simulations.

637 Lastly, riverine Fe flux is weighted by water runoff of each river and integrated globally as a total input of 1.45

638 Tg yr⁻¹ (Chester, 1990). To the best of our knowledge, the available global riverine iron dataset is rare. Previous

639 studies have used various approximation approaches, e.g., constant Fe to dissolved inorganic carbon (DIC) ratio

640 (Aumont et al., 2015), Fe to phosphorus ratio (Lacroix et al., 2020). In the study by Aumont et al. (2015), the Fe:

641 DIC ratio is determined so that the total Fe supply also equals 1.45 Tg Fe yr⁻¹ as estimated by Chester (1990). We

642 are aware that our approximation likely has bias in regional scales, especially in Fe limiting regions like the Arctic.

643 However, it has likely a minor impact on the projected PP, since light rather than riverine nutrients input is the

644 primary control of the projected Arctic PP in our model. Also, we have conducted all simulations only under one

645 IPCC representative concentration pathway scenario (the intermediate RCP 4.5), which may lead to a narrower

646 possible range of the riverine fluxes induced impact on the projected marine PP and C uptake.

647 5 Conclusions

649

648 In this study, we apply a fully coupled Earth system model to assess the impact of riverine nutrients and carbon

of uncertainty in future riverine fluxes on the projected changes, using several riverine input configurations.

delivery to the ocean on the contemporary and future marine PP and carbon uptake. We also quantify the effects

651 Compared to satellite- and observation-based estimates, the inclusion of riverine nutrients and carbon improves

652 the contemporary spatial distribution only slightly for PP (3.6% reduction in RMSE) and insignificantly for ocean

653 carbon uptake (0.1% reduction in RMSE) on a global scale, with larger improvements on the continental margins

654 (5.4% reduction in RMSE for PP) and the North Atlantic region (7.4% reduction in RMSE for carbon uptake).

655 Concerning future projected changes, decline in nutrients supply in tropical and subtropical surface waters, due

656 to upper ocean warming and increased vertical stratification, is projected by our model to reduce PP over the 21st 657 century. Riverine nutrient inputs into surface coastal waters alleviate the nutrient limitation and considerably

658 lessen the projected future decline in PP from -5.4% without riverine inputs to -4.4%, -4.1% and -3.6% in FIX,

659 RUN and GNS (averaged over GNa, GNg, GNo and GNt), respectively. Different from the global value, the

660 projected PP in the Arctic increases considerably, because light and temperature—the primary limiting factors for

661 phytoplankton growth in polar regions—become more <u>favorable</u> due to sea-ice melting under warmer future

conditions. When riverine nutrient inputs are presented in the model, they further enhance the projected increase

in PP in the Arctic, counteracting the nutrient decline effect due to stronger stratification in the fresher and warmersurface water.

```
\begin{array}{ll} 665 & \text{Depending on the riverine scenarios, where the riverine nutrients input dominates over the C input, the projected} \\ 666 & \text{net uptake of } CO_2 \text{ further enhances along continental margins via photosynthesis process. Conversely, where the} \end{array}
```

667 riverine C input is dominant over the nutrients input, the projected net uptake of CO_2 is reduced, especially at

668 large river estuaries, due to higher CO₂ outgassing.

Deleted: favourable

16•

(Formatted: Right: 0.63 cm

l

Deleted: idealised

(Formatted: Right: 0.63 cm

17•

670 We have explored a range of riverine input configurations from temporally constant fluxes (FIX), to idealized 671 time-varying fluxes following variations in simulated hydrological cycle (RUN), to plausible future scenarios 672 (GNS) from a set of global assumptions. The large range of the uncertainty of the riverine input does not transfer 673 to large uncertainty of the projected global PP and ocean C uptake in our simulations likely due to model 674 limitations related to resolution and shelf process representations. Our study suggests that applying transient 675 riverine inputs in the ESMs with coarse or intermediate model resolution (~1°) does not significantly reduce the 676 uncertainty in global marine PP and C uptake projections, but it may be of importance for regional studies such 677 as in the North Atlantic and along the continental margins. 678 Future modelling studies that include riverine input to the ocean can benefit from using high or at least adequate 679 model resolution, so that shelf processes, such as realistic remineralization rate for riverine organic matter in the 680 coastal water and shelf sediment as well as lateral transport, can be better resolved. Better constraints on riverine 681 C to nutrient ratios are needed to accurately assess the net riverine impact on ocean C uptake. Further exploration

682 of various future scenarios of riverine input is clearly warranted in order to better assess projected changes in

683 <u>ocean PP and C uptake, especially in regional scales.</u>

684 Appendix A



685 686

l





18

696 We assessed statistical significance of time-averaged differences using Student's t-test. We performed the test on 697 annual data with α set to 0.05 and N set to the number of years in the respective average period, assuming the 698 internal climate variability exhibits most power on interannual and shorter timescales. We removed the main part 699 of the externally forced signal by subtracting the linear trend of the annual timeseries prior to performing the t-691 test if the timeseries contained more than 20 years. For shorter time series, we therefore did not remove the linear 691 trend as it potentially has a large internal variability component. 702 All differences presented in the main text, summarized in Tables B1 and B2, were found to be statistically

703 significant and the plots feature only differences for which the t-test locally rejected the null-hypothesis. We found 704 even small inter-simulation differences statistically significant because these differences were less affected by 705 internal variability. In our model setup, the marine biogeochemistry does not feedback on the physical climate. 706 Consequently, the climate variability and climate trends are the same in all experiments and the interannual 707 variability in the biogeochemical parameters-which is predominantly driven by the physical climate 708 variability-is also virtually the same. As illustrated in Figure B1, any uncertainty related to internal climate 709 variability is effectively removed in the computation of the inter-experiment differences. In this manner, we were 710 able to obtain statistically robust results for short time-slices without having to perform multi-member simulation 711 ensembles for each experiment.

712 Detectability of inter-simulation differences does, however, not guarantee that the differences are large enough to

713 be competitive with real-world internal variability to have real-world implications. Therefore, we additionally

714 compared the inter-simulation differences against the internal variability of the absolute field (i.e., not the

715 difference field). We estimated the joint internal variability of the absolute field for N-year time averages as

716

$$\sigma_{\mu_{AB}} = \frac{\sqrt{\sigma_A^2 + \sigma_B^2}}{\sqrt{2N}}$$

717 where σ_A and σ_B are the interannual standard deviations for experiment A and B, respectively. As for the t-test,

718 we removed the externally forced signal by subtracting the linear trend of the annual timeseries prior to computing

719 standard-deviations if N>20. On all difference plots we marked the areas where inter-simulation differences

720 exceed $\sigma_{\mu_{AB}}$ and thus are large enough to have real-world implications.

721 722

723 724

725

Table B1: Global and regional statistics of simulated primary production. Shown are the time-mean μ and twice its standard-deviation σ_{μ} (rounded up to two decimals) derived from annual values. The t_{his} and t_{fut} denote the time periods 1950–1999 and 2050–2099, respectively.

Variable	Experiment	Period	Region	$\mu \pm 2 \sigma_{\mu}$
RMSE of PP	REF	2003-2012	Global	10.70 ± 0.18
(mol C m ⁻² yr ⁻¹)			Continental margins	28.96 ± 0.18
	FIX		Global	10.31 ± 0.21
			Continental margins	27.43 ± 0.19
	FIX-REF		Global	$\textbf{-0.39}\pm0.04$
			Continental margins	-1.52 ± 0.04
PP	REF	2003-2012	Global	40.06 ± 0.50
(Pg C yr ⁻¹)	FIX			42.99 ± 0.51
	FIX-REF			2.93 ± 0.02
PP	REF	t _{his}	Arctic	0.08 ± 0.01
projection		t _{fut}		0.15 ± 0.01
(Pg C yr ⁻¹)		t _{fut} -t _{his}		0.07 ± 0.01

Deleted: Values in brackets denote relative changes in percentage.

10

	FIX	t _{his}		0.10 ± 0.01
		t _{fut}		0.18 ± 0.01
		t _{fut} -t _{his}		0.08 ± 0.01
	FIX-REF	t _{fut} -t _{his}		0.01 ± 0.01
PP projection	REF	t _{his}	Global	41.14 ± 0.26
(Pg C yr ⁻¹)		t _{fut}		38.90 ± 0.23
		t _{fut} -t _{his}		$\textbf{-2.24}\pm0.37$
	FIX	t _{his}		43.99 ± 0.26
		t _{fut}		42.06 ± 0.24
		t _{fut} -t _{his}		$\textbf{-1.93}\pm0.38$
	RUN	t _{fut} -t _{his}		$\textbf{-1.82}\pm0.38$
	GNS	-		-1.57 ± 0.38
	FIX-REF			0.31 ± 0.01
	GNS-REF			0.66 ± 0.02

Table B2: Global and regional statistics of simulated ocean carbon uptake. Shown are the time-mean μ and twice its standard-deviation σ_{μ} (rounded up to two decimals) derived from annual values. The t_{tils} and t_{ful} denote the time periods 1950–1999 and 2050–2099, respectively. Values in brackets denote relative changes in percentage.

		r	L	
Variable	Experiment	Period	Region	$\mu \pm 2 \sigma_{\mu}$
RMSE of C	REF	2003-2012	Global	0.84 ± 0.05
uptake			Subpolar North Atlantic	0.73 ± 0.09
(mol C m ⁻² yr ⁻¹)	FIX		Global	0.83 ± 0.05
			Subpolar North Atlantic	0.67 ± 0.08
	FIX-REF		Global	-0.01 ± 0.01
			Subpolar North Atlantic	-0.06 ± 0.01
			-	(8.2±0.1%)
C uptake	REF	2003-2012	Global	2.77 ± 0.06
(Pg C yr ⁻¹)	FIX			2.86 ± 0.07
	FIX-REF			0.09 ± 0.01
				(3.1±0.1%)
C uptake	REF	t _{fut} -t _{his}	Global	1.25 ± 0.03
projection	FIX			1.28 ± 0.04
(Pg C yr ⁻¹)	RUN			1.29 ± 0.04
	GNS			1.13 ± 0.04
	FIX-REF			0.03 ± 0.01
	GNS-REF	7		-0.11 ± 0.03

733



Figure B1. Global integrated primary production (PP) time-series from single experiments (top) versus difference between two experiments (bottom). The PP variability of REF and FIX closely follow each other because the simulations feature the exact same physical variability. As a result, the interannual variability largely cancels out in the computation of FIX-REF differences and the FIX-REF difference times-series exhibits a standard-deviation that is an order of magnitude smaller than the standard-deviations of REF and FIX.

20-

Formatted: Font: 9 pt

734 Appendix C – Comparison between NEWS 2 dataset and measurement-based riverine data

735 Table C1: Comparison between NEWS 2 dataset (Mayorga et al., 2010) and measurement-based (provided by 736 PARTNERS Project; Holmes et al., 2012) riverine dissolved inorganic nitrogen (DIN) and dissolved organic nitrogen 737 (DON) in the 6 largest Arctic rivers around year 2000.

River	D	IN	DC	DN		
	(Pg N	(Pg N yr ⁻¹)		(Pg N yr ⁻¹)		
	NEWS 2	Measurement	NEWS 2	Measurement		
Ob	89	86	102	110		
Yenisei	47	51	132	111		
Lena	30	33	88	135		
Kolyma	9	7	21	17		
Yukon	5	26	14	47		
Mackenzie	22	27	62	31		

Note that the data from NEWS 2 are for the year 2000, while measured data from PARTNERS Project are calculated over
 1999–2008 (missing discharge data restricted the Yukon estimates to 2001–2008).

740 Code and data availability

741 The model code and riverine data used can be provided by the corresponding author upon request.

742 Author contribution

- 743 SG and IB designed the model experiments and SG developed the model code and performed the simulations with
- 744 the help from IB. JS and JT contributed to the interpretation and analyzation of the results. JS, JT, IB and CH
- 745 contributed to editing the manuscript. CH supervised the project work. JH and EM provided riverine data and
- $746 \qquad \text{consultation. SG prepared the manuscript with contributions from all co-authors.}$

747 Competing interests

748 The authors declare that they have no conflict of interest.

749 Disclaimer

- 750 This article reflects only the authors' view the funding agencies as well as their executive agencies are not
- responsible for any use that may be made of the information that the article contains.

752 Acknowledgement

- 753 This work was supported through project CRESCENDO (Coordinated Research in Earth Systems and
- 754 Climate: Experiments, kNowledge, Dissemination and Outreach; Horizon 2020 European Union's Framework
- 755 Programme for Research and Innovation, grant no. 641816, European Commission). Computing and storage
- 756 resources have been provided by UNINETT Sigma2 (nn2345k, nn2980k, ns2345k, ns2980k). JT acknowledge
- the Research Council Funded project Downscaling Climate and Ocean Change to Services (CE2COAST; 318477).

21

(Formatted: Right: 0.63 cm

ļ

758 IB received funding from the Trond Mohn Foundation through the Bjerknes Climate Prediction Unit

- 759 (BFS2018TMT01) and NFR Climate Futures (309562). JH benefited from financial support from the Deutsche
- 760 Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy EXC
- 761 2037 'Climate, Climatic Change, and Society' project number 390683824, contribution to the Center for Earth
- 762 System Research and Sustainability (CEN) of Universität Hamburg. We gratefully acknowledge that the
- 763 schematic diagram of Figure 12 has been prepared by Jadelynn Fong. We thank Fabrice Lacroix and one
- 764 <u>anonymous reviewer for their constructive comments which improved the manuscript greatly.</u>

765 References

- Alcamo, J., van Vuuren, D., Rosegrant, M., Alder, J., Bennett, E., Lodge, D., Masui, T., Morita, T., Ringler,
 C., Sala, O., Schulze, K., Zurek, M., Eickhout, B., Maerker, M., and Kok, K.: Changes in ecosystem
 services and their drivers across the scenarios, in: Ecosystems and Human Well-being: Scenarios, edited by:
 Carpenter, S. R., Pingali, P. L., Bennett, E. M., and Zurek, M. B., Island Press, Washington, 279-354, 2006.
- Anderson, L. G., Jutterström, S., Hjalmarsson, S., Wåhlström, I., and Semiletov, I. P.: Out-gassing of CO2
 from Siberian Shelf seas by terrestrial organic matter decomposition, Geophysical Research Letters, 36, https://doi.org/10.1029/2009GL040046, 2009.
- Arndt, S., Jørgensen, B. B., LaRowe, D. E., Middelburg, J. J., Pancost, R. D., and Regnier, P.: Quantifying
 the degradation of organic matter in marine sediments: A review and synthesis, Earth-Science Reviews,
 123, 53-86, https://doi.org/10.1016/j.earscirev.2013.02.008, 2013.
- Assmann, K. M., Bentsen, M., Segschneider, J., and Heinze, C.: An isopycnic ocean carbon cycle model,
 Geosci. Model Dev., 3, 143-167, 10.5194/gmd-3-143-2010, 2010.
- Aumont, O., Orr, J. C., Monfray, P., Ludwig, W., Amiotte-Suchet, P., and Probst, J.-L.: Riverine-driven
 interhemispheric transport of carbon, Global Biogeochemical Cycles, 15, 393-405,
 https://doi.org/10.1029/1999GB001238, 2001.
- Aumont, O., Ethé, C., Tagliabue, A., Bopp, L., and Gehlen, M.: PISCES-v2: an ocean biogeochemical model for carbon and ecosystem studies, Geosci. Model Dev., 8, 2465-2513, 10.5194/gmd-8-2465-2015, 2015.
- Behrenfeld, M. J. and Falkowski, P. G.: Photosynthetic rates derived from satellite-based chlorophyll
 concentration, Limnology and Oceanography, 42, 1-20, https://doi.org/10.4319/lo.1997.42.1.0001, 1997.
- Behrenfeld, M. J., O'Malley, R. T., Siegel, D. A., McClain, C. R., Sarmiento, J. L., Feldman, G. C.,
 Milligan, A. J., Falkowski, P. G., Letelier, R. M., and Boss, E. S.: Climate-driven trends in contemporary
 ocean productivity, Nature, 444, 752-755, 10.1038/nature05317, 2006.
- Bentsen, M., Bethke, I., Debernard, J. B., Iversen, T., Kirkevåg, A., Seland, Ø., Drange, H., Roelandt, C.,
 Seierstad, I. A., Hoose, C., and Kristjánsson, J. E.: The Norwegian Earth System Model, NorESM1-M –
 Part 1: Description and basic evaluation of the physical climate, Geosci. Model Dev., 6, 687-720,
 10.5194/gmd-6-687-2013, 2013.
- Bernard, C. Y., Dürr, H. H., Heinze, C., Segschneider, J., and Maier-Reimer, E.: Contribution of riverine nutrients to the silicon biogeochemistry of the global ocean – a model study, Biogeosciences, 8, 551-564, 10.5194/bg-8-551-2011, 2011.
- Beusen, A. H. W., Bouwman, A. F., Dürr, H. H., Dekkers, A. L. M., and Hartmann, J.: Global patterns of
 dissolved silica export to the coastal zone: Results from a spatially explicit global model, Global
 Biogeochemical Cycles, 23, https://doi.org/10.1029/2008GB003281, 2009.
- Beusen, A. H. W., Van Beek, L. P. H., Bouwman, A. F., Mogollón, J. M., and Middelburg, J. J.: Coupling
 global models for hydrology and nutrient loading to simulate nitrogen and phosphorus retention in surface
 water & ndash; description of IMAGE–GNM and analysis of performance, Geosci. Model Dev., 8, 40454067, 10.5194/gmd-8-4045-2015, 2015.

22

Formatted: Right: 0.63 cm

l

- Beusen, A. H. W., Bouwman, A. F., Van Beek, L. P. H., Mogollón, J. M., and Middelburg, J. J.: Global
 riverine N and P transport to ocean increased during the 20th century despite increased retention along the
 aquatic continuum, Biogeosciences, 13, 2441-2451, 10.5194/bg-13-2441-2016, 2016.
- Bianchi, T. S.: The role of terrestrially derived organic carbon in the coastal ocean: A changing paradigm
 and the priming effect, Proceedings of the National Academy of Sciences, 108, 19473-19481,
 10.1073/pnas.1017982108, 2011.
- Blair, N. E. and Aller, R. C.: The Fate of Terrestrial Organic Carbon in the Marine Environment, Annual
 Review of Marine Science, 4, 401-423, 10.1146/annurev-marine-120709-142717, 2011.
- 811Bleck, R. and Smith, L. T.: A wind-driven isopycnic coordinate model of the north and equatorial Atlantic812Ocean: 1. Model development and supporting experiments, Journal of Geophysical Research: Oceans, 95,8133273-3285, 10.1029/JC095iC03p03273, 1990.
- Bleck, R., Rooth, C., Hu, D., and Smith, L. T.: Salinity-driven Thermocline Transients in a Wind- and
 Thermohaline-forced Isopycnic Coordinate Model of the North Atlantic, Journal of Physical Oceanography,
 22, 1486-1505, 10.1175/1520-0485(1992)022<1486:SDTTIA>2.0.CO;2, 1992.
- 817 Bopp, L., Monfray, P., Aumont, O., Dufresne, J.-L., Le Treut, H., Madec, G., Terray, L., and Orr, J. C.:
 818 Potential impact of climate change on marine export production, Global Biogeochemical Cycles, 15, 81-99,
 819 https://doi.org/10.1029/1999GB001256, 2001.
- Bopp, L., Aumont, O., Cadule, P., Alvain, S., and Gehlen, M.: Response of diatoms distribution to global
 warming and potential implications: A global model study, Geophysical Research Letters, 32,
 https://doi.org/10.1029/2005GL023653, 2005.
- Bopp, L., Resplandy, L., Orr, J. C., Doney, S. C., Dunne, J. P., Gehlen, M., Halloran, P., Heinze, C., Ilyina,
 T., Séférian, R., Tjiputra, J., and Vichi, M.: Multiple stressors of ocean ecosystems in the 21st century:
 projections with CMIP5 models, Biogeosciences, 10, 6225-6245, 10.5194/bg-10-6225-2013, 2013.
- Borges, A. V., Delille, B., and Frankignoulle, M.: Budgeting sinks and sources of CO2 in the coastal ocean:
 Diversity of ecosystems counts, Geophysical Research Letters, 32, https://doi.org/10.1029/2005GL023053,
 2005.
- Bourgeois, T., Orr, J. C., Resplandy, L., Terhaar, J., Ethé, C., Gehlen, M., and Bopp, L.: Coastal-ocean uptake of anthropogenic carbon, Biogeosciences, 13, 4167-4185, 10.5194/bg-13-4167-2016, 2016.
- Bouwman, A. F., Beusen, A. H. W., and Billen, G.: Human alteration of the global nitrogen and phosphorus
 soil balances for the period 1970–2050, Global Biogeochemical Cycles, 23,
 https://doi.org/10.1029/2009GB003576, 2009.
- Boyle, E. A., Edmond, J. M., and Sholkovitz, E. R.: The mechanism of iron removal in estuaries, Geochim
 Cosmochim Ac, 41, 1313-1324, https://doi.org/10.1016/0016-7037(77)90075-8, 1977.
- Cabré, A., Marinov, I., and Leung, S.: Consistent global responses of marine ecosystems to future climate
 change across the IPCC AR5 earth system models, Climate Dynamics, 45, 1253-1280, 10.1007/s00382 014-2374-3, 2015.
- Cai, W.-J.: Estuarine and Coastal Ocean Carbon Paradox: CO2 Sinks or Sites of Terrestrial Carbon
 Incineration?, Annual Review of Marine Science, 3, 123-145, 10.1146/annurev-marine-120709-142723,
 2010.
- Chen, C.-T. A. and Borges, A. V.: Reconciling opposing views on carbon cycling in the coastal ocean:
 Continental shelves as sinks and near-shore ecosystems as sources of atmospheric CO2, Deep Sea Research
- 844
 Part II: Topical Studies in Oceanography, 56, 578-590, https://doi.org/10.1016/j.dsr2.2009.01.001, 2009.
- 845
 Chester, R.: The transport of material to the oceans: the river pathway, in: Marine Geochemistry, Springer,

 846
 Dordrecht, https://doi.org/10.1007/978-94-010-9488-7_3, 1990.
- Chester, R.: The Input of Material to the Ocean Reservoir, in: Marine Geochemistry, Wiley Online Books,
 7-10, https://doi.org/10.1002/9781118349083.ch2, 2012.

23

849 Cotrim da Cunha, L., Buitenhuis Erik, T., Le Quéré, C., Giraud, X., and Ludwig, W.: Potential impact of 850 changes in river nutrient supply on global ocean biogeochemistry, Global Biogeochemical Cycles, 21, 851 10.1029/2006GB002718, 2007. 852 853 854 855 856 857 858 859 Danabasoglu, G., Lamarque, J. F., Bacmeister, J., Bailey, D. A., DuVivier, A. K., Edwards, J., Emmons, L. K., Fasullo, J., Garcia, R., Gettelman, A., Hannay, C., Holland, M. M., Large, W. G., Lauritzen, P. H., Lawrence, D. M., Lenaerts, J. T. M., Lindsay, K., Lipscomb, W. H., Mills, M. J., Neale, R., Oleson, K. W., Otto-Bliesner, B., Phillips, A. S., Sacks, W., Tilmes, S., van Kampenhout, L., Vertenstein, M., Bertini, A., Dennis, J., Deser, C., Fischer, C., Fox-Kemper, B., Kay, J. E., Kinnison, D., Kushner, P. J., Larson, V. E., Long, M. C., Mickelson, S., Moore, J. K., Nienhouse, E., Polvani, L., Rasch, P. J., and Strand, W. G.: The Community Earth System Model Version 2 (CESM2), Journal of Advances in Modeling Earth Systems, 12, e2019MS001916, https://doi.org/10.1029/2019MS001916, 2020. 860 Dittmar, T., Fitznar, H. P., and Kattner, G.: Origin and biogeochemical cycling of organic nitrogen in the 861 eastern Arctic Ocean as evident from D- and L-amino acids, Geochim Cosmochim Ac, 65, 4103-4114, 862 https://doi.org/10.1016/S0016-7037(01)00688-3, 2001. 863 Doney, S. C.: Plankton in a warmer world, Nature, 444, 695-696, 10.1038/444695a, 2006. 864 Figuères, G., Martin, J. M., and Meybeck, M.: Iron behaviour in the Zaire estuary, Netherlands Journal of 865 Sea Research, 12, 329-337, https://doi.org/10.1016/0077-7579(78)90035-2, 1978. 866 Eiriksdottir, E., Oelkers, E., Hardardóttir, J., and Gislason, S.: The impact of damming on riverine fluxes to 867 the ocean: A case study from Eastern Iceland, Water Research, 113, 10.1016/j.watres.2016.12.029, 2016. 868 Friedlingstein, P., Jones, M. W., O'Sullivan, M., Andrew, R. M., Bakker, D. C. E., Hauck, J., Le Quéré, C., 869 Peters, G. P., Peters, W., Pongratz, J., Sitch, S., Canadell, J. G., Ciais, P., Jackson, R. B., Alin, S. R., 870 Anthoni, P., Bates, N. R., Becker, M., Bellouin, N., Bopp, L., Chau, T. T. T., Chevallier, F., Chini, L. P., 871 872 Cronin, M., Currie, K. I., Decharme, B., Djeutchouang, L., Dou, X., Evans, W., Feely, R. A., Feng, L., Gasser, T., Gilfillan, D., Gkritzalis, T., Grassi, G., Gregor, L., Gruber, N., Gürses, Ö., Harris, I., Houghton, 873 R. A., Hurtt, G. C., Iida, Y., Ilyina, T., Luijkx, I. T., Jain, A. K., Jones, S. D., Kato, E., Kennedy, D., Klein 874 Goldewijk, K., Knauer, J., Korsbakken, J. I., Körtzinger, A., Landschützer, P., Lauvset, S. K., Lefèvre, N., 875 Lienert, S., Liu, J., Marland, G., McGuire, P. C., Melton, J. R., Munro, D. R., Nabel, J. E. M. S., Nakaoka, 876 S. I., Niwa, Y., Ono, T., Pierrot, D., Poulter, B., Rehder, G., Resplandy, L., Robertson, E., Rödenbeck, C., 877 Rosan, T. M., Schwinger, J., Schwingshackl, C., Séférian, R., Sutton, A. J., Sweeney, C., Tanhua, T., Tans, 878 P. P., Tian, H., Tilbrook, B., Tubiello, F., van der Werf, G., Vuichard, N., Wada, C., Wanninkhof, R., 879 Watson, A., Willis, D., Wiltshire, A. J., Yuan, W., Yue, C., Yue, X., Zaehle, S., and Zeng, J.: Global 880 Carbon Budget 2021, Earth Syst. Sci. Data Discuss., 2021, 1-191, 10.5194/essd-2021-386, 2021. 881 Frigstad, H., Kaste, Ø., Deininger, A., Kvalsund, K., Christensen, G., Bellerby, R. G. J., Sørensen, K., 882 Norli, M., and King, A. L.: Influence of Riverine Input on Norwegian Coastal Systems, Frontiers in Marine 883 Science, 7, 2020. 884 Galy, V., Peucker-Ehrenbrink, B., and Eglinton, T.: Global carbon export from the terrestrial biosphere controlled by erosion, Nature, 521, 204-207, 10.1038/nature14400, 2015. 885 Garcia, H. E., Locarnini, R. A., Boyer, T. P., Antonov, J. I., Baranova, O. K., Zweng, M. M., Reagan, J. R., and Johnson, D. R.: World Ocean Atlas 2013. Vol. 4: Dissolved Inorganic Nutrients 886 887 888 (phosphate,nitrate,silicate), NOAA Atlas NESDIS 76, 25 pp, 2013a. 889 Garcia, H. E., Locarnini, R. A., Boyer, T. P., Antonov, J. I., Mishonov, A. V., Baranova, O. K., Zweng, M. 890 M., Reagan, J. R., and Johnson, D. R.: World Ocean Atlas 2013. Vol. 3: Dissolved Oxygen, Apparent 891 Oxygen Utilization, and Oxygen Saturation, NOAA Atlas NESDIS 75, 27 pp, 2013b. 892 Garnier, J., Billen, G., Lassaletta, L., Vigiak, O., Nikolaidis, N. P., and Grizzetti, B.: Hydromorphology of 893 coastal zone and structure of watershed agro-food system are main determinants of coastal eutrophication, Environmental Research Letters, 16, 023005, 10.1088/1748-9326/abc777, 2021. 894 895 Gharamti, M. E., Tjiputra, J., Bethke, I., Samuelsen, A., Skjelvan, I., Bentsen, M., and Bertino, L.: 896 Ensemble data assimilation for ocean biogeochemical state and parameter estimation at different sites, 897 Ocean Modelling, 112, 65-89, https://doi.org/10.1016/j.ocemod.2017.02.006, 2017. 898 Hajima, T., Watanabe, M., Yamamoto, A., Tatebe, H., Noguchi, M. A., Abe, M., Ohgaito, R., Ito, A., 899 Yamazaki, D., Okajima, H., Ito, A., Takata, K., Ogochi, K., Watanabe, S., and Kawamiya, M.: Formatted: Right: 0.63 cm 24



I

I		Formatted: Header
950	Autotrophic and a CO2 Sink?, Global Biogeochemical Cycles, 35, e2020GB006603,	
951	https://doi.org/10.1029/2020GB006603, 2021b.	
952 953	Lalonde, K., Vähätalo, A. V., and Gélinas, Y.: Revisiting the disappearance of terrestrial dissolved organic matter in the ocean: a δ 13C study, Biogeosciences, 11, 3707-3719, 10.5194/bg-11-3707-2014, 2014.	
954 955 956	Landschützer, P., Gruber, N., and Bakker, D.: An updated observation-based global monthly gridded sea surface pCO 2 and air-sea CO 2 flux product from 1982 through 2015 and its monthly climatology. (website: https://www.nodc.noaa.gov/ocads/oceans/SPCO2 1982 2015 ETH SOM FFN.html), 2017.	Deleted:
957 958 959 960	Lawrence, D. M., Oleson, K. W., Flanner, M. G., Thornton, P. E., Swenson, S. C., Lawrence, P. J., Zeng, X., Yang, ZL., Levis, S., Sakaguchi, K., Bonan, G. B., and Slater, A. G.: Parameterization improvements and functional and structural advances in Version 4 of the Community Land Model, Journal of Advances in Modeling Earth Systems, 3, 10.1029/2011MS00045, 2011.	
961 962 963 964 965 966 967	Lee, Y. J., Matrai, P. A., Friedrichs, M. A. M., Saba, V. S., Aumont, O., Babin, M., Buitenhuis, E. T., Chevallier, M., de Mora, L., Dessert, M., Dunne, J. P., Ellingsen, I. H., Feldman, D., Frouin, R., Gehlen, M., Gorgues, T., Ilyina, T., Jin, M., John, J. G., Lawrence, J., Manizza, M., Menkes, C. E., Perruche, C., Le Fouest, V., Popova, E. E., Romanou, A., Samuelsen, A., Schwinger, J., Séférian, R., Stock, C. A., Tjiputra, J., Tremblay, L. B., Ueyoshi, K., Vichi, M., Yool, A., and Zhang, J.: Net primary productivity estimates and environmental variables in the Arctic Ocean: An assessment of coupled physical-biogeochemical models, Journal of Geophysical Research: Oceans, 121, 8635-8669, https://doi.org/10.1002/2016JC011993, 2016.	
968 969	Le Fouest, V., Babin, M., and Tremblay, J. É.: The fate of riverine nutrients on Arctic shelves, Biogeosciences, 10, 3661-3677, 10.5194/bg-10-3661-2013, 2013.	
970 971 972	Le Fouest, V., Manizza, M., Tremblay, B., and Babin, M.: Modelling the impact of riverine DON removal by marine bacterioplankton on primary production in the Arctic Ocean, Biogeosciences, 12, 3385-3402, 10.5194/bg-12-3385-2015, 2015.	
973 974 975 976	Le Fouest, V., Matsuoka, A., Manizza, M., Shernetsky, M., Tremblay, B., and Babin, M.: Towards an assessment of riverine dissolved organic carbon in surface waters of the western Arctic Ocean based on remote sensing and biogeochemical modeling, Biogeosciences, 15, 1335-1346, 10.5194/bg-15-1335-2018, 2018.	
977 978	Letscher, R. T., Hansell, D. A., Kadko, D., and Bates, N. R.: Dissolved organic nitrogen dynamics in the Arctic Ocean, Marine Chemistry, 148, 1-9, https://doi.org/10.1016/j.marchem.2012.10.002, 2013.	
979 980 981	Liu, D., Bai, Y., He, X., Chen, CT. A., Huang, TH., Pan, D., Chen, X., Wang, D., and Zhang, L.: Changes in riverine organic carbon input to the ocean from mainland China over the past 60 years, Environment International, 134, 105258, https://doi.org/10.1016/j.envint.2019.105258, 2020.	
982 983 984	Liu, X., Stock, C. A., Dunne, J. P., Lee, M., Shevliakova, E., Malyshev, S., and Milly, P. C. D.: Simulated Global Coastal Ecosystem Responses to a Half-Century Increase in River Nitrogen Loads, Geophysical Research Letters, 48, e2021GL094367, https://doi.org/10.1029/2021GL094367, 2021.	
985 986	Lobbes, J. M., Fitznar, H. P., and Kattner, G.: Biogeochemical characteristics of dissolved and particulate organic matter in Russian rivers entering the Arctic Ocean, Geochim Cosmochim Ac, 64, 2973-2983, 2000.	
987 988	Ludwig, W., Probst, JL., and Kempe, S.: Predicting the oceanic input of organic carbon by continental erosion, Global Biogeochemical Cycles, 10, 23-41, https://doi.org/10.1029/95GB02925, 1996.	
989 990 991	Mahowald, N. M., Baker, A. R., Bergametti, G., Brooks, N., Duce, R. A., Jickells, T. D., Kubilay, N., Prospero, J. M., and Tegen, I.: Atmospheric global dust cycle and iron inputs to the ocean, Global Biogeochemical Cycles, 19, 10.1029/2004GB002402, 2005.	
992 993 994	Maier-Reimer, E., Kriest, I., Segschneider, J., and Wetzel, P.: The Hamburg oceanic carbon cycle circulation model HAMOCC5.1—Technical Description Release 1.1, Tech. Rep. 14, Reports on Earth System Science, Max Planck Institute for Meteorology, Hamburg, Germany, 2005.	
995 996 997	Manizza, M., Follows, M. J., Dutkiewicz, S., Menemenlis, D., McClelland, J. W., Hill, C. N., Peterson, B. J., and Key, R. M.: A model of the Arctic Ocean carbon cycle, Journal of Geophysical Research: Oceans, 116, https://doi.org/10.1029/2011JC006998, 2011.	
		Enermatted: Dight: 0.62 cm
1	76.	Tormatted. Right. 0.03 Cm
	20*	

I

1000 Polimene, L., Overduin, P., Mollenhauer, G., Grosse, G., Rachold, V., Sobczak, W. V., Spencer, R. G. M., 1001 and Juhls, B.: Degrading permafrost river catchments and their impact on Arctic Ocean nearshore processes, Ambio, 51, 439-455, 10.1007/s13280-021-01666-z, 2022. 1002 1003 Mayorga, E., Seitzinger, S. P., Harrison, J. A., Dumont, E., Beusen, A. H. W., Bouwman, A. F., Fekete, B. 1004M., Kroeze, C., and Van Drecht, G.: Global Nutrient Export from WaterSheds 2 (NEWS 2): Model 1005 development and implementation, Environmental Modelling & Software, 25, 837-853, 1006 https://doi.org/10.1016/j.envsoft.2010.01.007, 2010. 1007 McClelland, J. W., Holmes, R. M., Dunton, K. H., and Macdonald, R. W.: The Arctic Ocean Estuary, 1008 Estuaries and Coasts, 35, 353-368, 10.1007/s12237-010-9357-3, 2012. 1009 Meybeck, M.: Carbon, nitrogen, and phosphorus transport by world rivers, American Journal of Science, 1010 282, 401, 10.2475/ajs.282.4.401, 1982.

Meybeck, M. and Vörösmarty, C.: Global transfer of carbon by rivers, Global Change Newsletter, 37, 18 19, 1999.

Mann, P. J., Strauss, J., Palmtag, J., Dowdy, K., Ogneva, O., Fuchs, M., Bedington, M., Torres, R.,

- Neale, R. B., Richter, J., Park, S., Lauritzen, P. H., Vavrus, S. J., Rasch, P. J., and Zhang, M.: The Mean
 Climate of the Community Atmosphere Model (CAM4) in Forced SST and Fully Coupled Experiments,
 Journal of Climate, 26, 5150-5168, 10.1175/JCLI-D-12-00236.1, 2013.
- 1016 Oleson, K. W., Lawrence, D. M., Bonan, G. B., Flanner, M. G., Kluzek, E., Lawrence, P. J., Levis, S.,
 1017 Swenson, S. C., Thornton, P. E., Dai, A., Decker, M., Dickinson, R., Feddema, J., Heald, C. L., Hoffman,
 1018 F., Lamarque, J.-F., Mahowald, N., Niu, G.-Y., Qian, T., Randerson, J., Running, S., Sakaguchi, K., Slater,
 1019 A., Stockli, R., Wang, A., Yang, Z.-L., Zeng, X., and Zeng, X.: Technical Description of version 4.0 of the
 1020 Community Land Model (CLM) (No. NCAR/TN-478+STR), University Corporation for Atmospheric
 1021 Research, doi:10.5065/D6FB50WZ, 2010.
- Pokrovsky, O. S., Manasypov, R. M., Kopysov, S. G., Krickov, I. V., Shirokova, L. S., Loiko, S. V., Lim,
 A. G., Kolesnichenko, L. G., Vorobyev, S. N., and Kirpotin, S. N.: Impact of Permafrost Thaw and Climate
 Warming on Riverine Export Fluxes of Carbon, Nutrients and Metals in Western Siberia, Water, 12,
 1025 10.3390/w12061817, 2020.
- 1026Redfield, A. and Daniel, R. J.: On the proportions of organic derivations in sea water and their relation to
the composition of plankton, in: James Johnstone Memorial Volume, University Press of Liverpool, 177-
192, 1934.
- Regnier, P., Friedlingstein, P., Ciais, P., Mackenzie, F. T., Gruber, N., Janssens, I. A., Laruelle, G. G.,
 Lauerwald, R., Luyssaert, S., Andersson, A. J., Arndt, S., Arnosti, C., Borges, A. V., Dale, A. W., GallegoSala, A., Goddéris, Y., Goossens, N., Hartmann, J., Heinze, C., Ilyina, T., Joos, F., LaRowe, D. E., Leifeld,
 J., Meysman, F. J. R., Munhoven, G., Raymond, P. A., Spahni, R., Suntharalingam, P., and Thullner, M.:
 Anthropogenic perturbation of the carbon fluxes from land to ocean, Nature Geoscience, 6, 597-607,
 1038/ngco1830, 2013.
- 1035Regnier, P., Resplandy, L., Najjar, R. G., and Ciais, P.: The land-to-ocean loops of the global carbon cycle,
Nature, 603, 401-410, 10.1038/s41586-021-04339-9, 2022.
- 1037
 Rogelj, J., den Elzen, M., Höhne, N., Fransen, T., Fekete, H., Winkler, H., Schaeffer, R., Sha, F., Riahi, K.,

 1038
 and Meinshausen, M.: Paris Agreement climate proposals need a boost to keep warming well below 2 °C,

 1039
 Nature, 534, 631-639, 10.1038/nature18307, 2016.
- 1040Sarmiento, J. L., Gruber, N., Brzezinski, M. A., and Dunne, J. P.: High-latitude controls of thermocline1041nutrients and low latitude biological productivity, Nature, 427, 56-60, 2004.
- Seitzinger, S. P., Mayorga, E., Bouwman, A. F., Kroeze, C., Beusen, A. H. W., Billen, G., Van Drecht, G.,
 Dumont, E., Fekete, B. M., Garnier, J., and Harrison, J. A.: Global river nutrient export: A scenario analysis
 of past and future trends, Global Biogeochemical Cycles, 24, https://doi.org/10.1029/2009GB003587, 2010.
- Séférian, R., Nabat, P., Michou, M., Saint-Martin, D., Voldoire, A., Colin, J., Decharme, B., Delire, C.,
 Berthet, S., Chevallier, M., Sénési, S., Franchisteguy, L., Vial, J., Mallet, M., Joetzjer, E., Geoffroy, O.,
- Bothe, B., Choune, M., Schumargey, L., Valt, J., Marte, M., Soch, E., Schuber, S., Soch, S.,

27

Formatted: Right: 0.63 cm

I

1049Evaluation of CNRM Earth System Model, CNRM-ESM2-1: Role of Earth System Processes in Present-1050Day and Future Climate, Journal of Advances in Modeling Earth Systems, 11, 4182-4227,1051https://doi.org/10.1029/2019MS001791, 2019.

- 1052 Séférian, R., Berthet, S., Yool, A., Palmiéri, J., Bopp, L., Tagliabue, A., Kwiatkowski, L., Aumont, O.,
- Christian, J., Dunne, J., Gehlen, M., Ilyina, T., John, J. G., Li, H., Long, M. C., Luo, J. Y., Nakano, H.,
 Romanou, A., Schwinger, J., Stock, C., Santana-Falcón, Y., Takano, Y., Tjiputra, J., Tsujino, H., Watanab
- Romanou, A., Schwinger, J., Stock, C., Santana-Falcón, Y., Takano, Y., Tjiputra, J., Tsujino, H., Watanabe,
 M., Wu, T., Wu, F., and Yamamoto, A.: Tracking Improvement in Simulated Marine Biogeochemistry
 Between CMIP5 and CMIP6, Current Climate Change Reports, 6, 95-119, 10.1007/s40641-020-00160-0,
- 1057 2020.1058 Shiller, A. M. and Boyle, E. A.: Trace elements in the Mississippi River Delta outflow region: B
- 1058Shiller, A. M. and Boyle, E. A.: Trace elements in the Mississippi River Delta outflow region: Behavior at
high discharge, Geochim Cosmochim Ac, 55, 3241-3251, https://doi.org/10.1016/0016-7037(91)90486-0,
1991.
- Sholkovitz, E. R. and Copland, D.: The coagulation, solubility and adsorption properties of Fe, Mn, Cu, Ni,
 Cd, Co and humic acids in a river water, Geochim Cosmochim Ac, 45, 181-189,
 https://doi.org/10.1016/0016-7037(81)90161-7, 1981.
- Skogen, M. D., Hjøllo, S. S., Sandø, A. B., and Tjiputra, J.: Future ecosystem changes in the Northeast
 Atlantic: a comparison between a global and a regional model system, ICES Journal of Marine Science, 75,
 2355-2369, 10.1093/icesjms/fsy088, 2018.
- Smith, S. V., Swaney, D. P., Talaue-Memanus, L., Bartley, J. D., Sandhei, P. T., McLaughlin, C. J., Dupra,
 V. C., Crossland, C. J., Buddemeier, R. W., Maxwell, B. A., and Wulff, F.: Humans, Hydrology, and the
 Distribution of Inorganic Nutrient Loading to the Ocean, BioScience, 53, 235-245, 10.1641/00063568(2003)053[0235:HHATDO]2.0.CO;2, 2003.
- Steinacher, M., Joos, F., Frolicher, T. L., Bopp, L., Cadule, P., Cocco, V., Doney, S. C., Gehlen, M.,
 Lindsay, K., Moore, J. K., Schneider, B., and Segschneider, J.: Projected 21st century decrease in marine
 productivity: a multi-model analysis, Biogeosciences, 7, 979-1005, 2010.
- 1074 Tagliabue, A., Kwiatkowski, L., Bopp, L., Butenschön, M., Cheung, W., Lengaigne, M., and Vialard, J.:
 1075 Persistent Uncertainties in Ocean Net Primary Production Climate Change Projections at Regional Scales
 1076 Raise Challenges for Assessing Impacts on Ecosystem Services, Frontiers in Climate, 3, 2021.
- Takahashi, T., Sutherland, S. C., Wanninkhof, R., Sweeney, C., Feely, R. A., Chipman, D. W., Hales, B.,
 Friederich, G., Chavez, F., Sabine, C., Watson, A., Bakker, D. C. E., Schuster, U., Metzl, N., YoshikawaInoue, H., Ishii, M., Midorikawa, T., Nojiri, Y., Körtzinger, A., Steinhoff, T., Hoppema, M., Olafsson, J.,
 Arnarson, T. S., Tilbrook, B., Johannessen, T., Olsen, A., Bellerby, R., Wong, C. S., Delille, B., Bates, N.
 R., and de Baar, H. J. W.: Climatological mean and decadal change in surface ocean pCO2, and net sea-air
 CO2 flux over the global oceans, Deep Sea Research Part II: Topical Studies in Oceanography, 56, 554577, https://doi.org/10.1016/j.dsr2.2008.12.009, 2009.
- 1084Taylor, K. E., Stouffer, R. J., and Meehl, G. A.: An Overview of CMIP5 and the Experiment Design,1085Bulletin of the American Meteorological Society, 93, 485-498, 10.1175/BAMS-D-11-00094.1, 2011.
- Terhaar, J., Orr, J. C., Ethé, C., Regnier, P., and Bopp, L.: Simulated Arctic Ocean Response to Doubling of
 Riverine Carbon and Nutrient Delivery, Global Biogeochemical Cycles, 33, 1048-1070,
 https://doi.org/10.1029/2019GB006200, 2019.
- Terhaar, J., Lauerwald, R., Regnier, P., Gruber, N., and Bopp, L.: Around one third of current Arctic Ocean
 primary production sustained by rivers and coastal erosion, Nature Communications, 12, 169,
 10.1038/s41467-020-20470-z, 2021.
- Tivig, M., Keller, D. P., and Oschlies, A.: Riverine nitrogen supply to the global ocean and its limited
 impact on global marine primary production: a feedback study using an Earth system model,
 Biogeosciences, 18, 5327-5350, 10.5194/bg-18-5327-2021, 2021.
- Tjiputra, J. F., Polzin, D., and Winguth, A. M. E.: Assimilation of seasonal chlorophyll and nutrient data into an adjoint three-dimensional ocean carbon cycle model: Sensitivity analysis and ecosystem parameter optimization, Global Biogeochemical Cycles, 21, https://doi.org/10.1029/2006GB002745, 2007.

284

Formatted: Right: 0.63 cm

ļ

- Tjiputra, J. F., Roelandt, C., Bentsen, M., Lawrence, D. M., Lorentzen, T., Schwinger, J., Seland, Ø., and
 Heinze, C.: Evaluation of the carbon cycle components in the Norwegian Earth System Model (NorESM),
 Geosci. Model Dev., 6, 301-325, 10.5194/gmd-6-301-2013, 2013.
- 1101
 Tjiputra, J. F., Schwinger, J., Bentsen, M., Morée, A. L., Gao, S., Bethke, I., Heinze, C., Goris, N., Gupta,

 1102
 A., He, Y. C., Olivié, D., Seland, Ø., and Schulz, M.: Ocean biogeochemistry in the Norwegian Earth
- 1103
 System Model version 2 (NorESM2), Geosci. Model Dev., 13, 2393-2431, 10.5194/gmd-13-2393-2020,

 1104
 2020.
- Vancoppenolle, M., Bopp, L., Madec, G., Dunne, J., Ilyina, T., Halloran, P. R., and Steiner, N.: Future
 Arctic Ocean primary productivity from CMIP5 simulations: Uncertain outcome, but consistent
 mechanisms, Global Biogeochemical Cycles, 27, 605-619, https://doi.org/10.1002/gbc.20055, 2013.
- 1108
 van der Struijk, L. F. and Kroeze, C.: Future trends in nutrient export to the coastal waters of South

 1109
 America: Implications for occurrence of eutrophication, Global Biogeochemical Cycles, 24,

 1110
 https://doi.org/10.1029/2009GB003572, 2010.
- van Vuuren, D. P., Edmonds, J., Kainuma, M., Riahi, K., Thomson, A., Hibbard, K., Hurtt, G. C., Kram, T.,
 Krey, V., Lamarque, J.-F., Masui, T., Meinshausen, M., Nakicenovic, N., Smith, S. J., and Rose, S. K.: The
 representative concentration pathways: an overview, Climatic Change, 109, 5, 10.1007/s10584-011-0148-z,
 2011.
- 1115Van Drecht, G., Bouwman, A. F., Harrison, J., and Knoop, J. M.: Global nitrogen and phosphate in urban1116wastewater for the period 1970 to 2050, Global Biogeochemical Cycles, 23,
- 1117 https://doi.org/10.1029/2009GB003458, 2009.
- 1118
 Westberry, T., Behrenfeld, M. J., Siegel, D. A., and Boss, E.: Carbon-based primary productivity modeling

 1119
 with vertically resolved photoacclimation, Global Biogeochemical Cycles, 22,

 1120
 https://doi.org/10.1029/2007GB003078, 2008.
- Wild, B., Andersson, A., Bröder, L., Vonk, J., Hugelius, G., McClelland, J. W., Song, W., Raymond, P. A., and Gustafsson, Ö.: Rivers across the Siberian Arctic unearth the patterns of carbon release from thawing permafrost, Proceedings of the National Academy of Sciences of the United States of America, 116, 10280-10285, 2019.
- Yan, W., Mayorga, E., Li, X., Seitzinger, S. P., and Bouwman, A. F.: Increasing anthropogenic nitrogen inputs and riverine DIN exports from the Changjiang River basin under changing human pressures, Global Biogeochemical Cycles, 24, https://doi.org/10.1029/2009GB003575, 2010.
- 1128Zhang, X., Fang, C., Wang, Y., Lou, X., Su, Y., and Huang, D.: Review of Effects of Dam Construction on1129the Ecosystems of River Estuary and Nearby Marine Areas, Sustainability, 14, 10.3390/su14105974, 2022.
- 1130

1131

1132

1133

1134

1135

1136

1137

1138

29





I

1141

I





32-

Formatted: Header

l















1158

I







1165 1166 Table 1. Brief introduction to future scenarios for river nutrient export used in Global NEWS 2 (Seitzinger et al., 2010)

Scenario	Agricultural trends	Sewage
Adapting Mosaic (GNa) a world with a focus on regional and local socio-ecological management	-medium productivity increase -2% of cropland area for energy crops -fertilizer efficiency: moderate increase in N and P fertilizer use in all countries; better integration of animal manure and recycling of human N and P from households with improved sanitation but lacking a sewage connection	-constant fraction of population with access to sanitation and sewage connection -moderate increase in N and P removal by wastewater treatment
Global Orchestration (GNg) a globalized world with an economic development focus and rapid economic growth	-high productivity increase -4% of cropland area for energy crops -fertilizer efficiency: no change in countries with a soil nutrient surplus; rapid increase in N and P fertilizer use in countries with soil nutrient depletion	-towards full access to improved sanitation and sewage connection -rapid increase in N and P removal by wastewater treatment
Order from Strength (GNo) a regionalized world with a focus on security	 -low productivity increase -1% of cropland area for energy crops -fertilizer efficiency: no change in countries with a soil nutrient surplus; moderate increase in N and P fertilizer use in countries with soil nutrient depletion 	-same as GNa
Technogarden (GNt) a globalized world with a focus on environmental technology	-medium-high productivity increase -28% of cropland area for energy crops -fertilizer efficiency: rapid increase in N and P fertilizer use in countries with a soil nutrient surplus; rapid increase in countries with soil nutrient depletion	-same as GNg

1167

41•