

Geodiversity influences limnological characteristics and freshwater ostracode species distributions across broad spatial scales in the northern Neotropics

5 Laura Macario-González^{1,2}, Sergio Cohuo^{2,3}, Philipp Hoelzmann⁴, Liseth Pérez², Manuel Elías-Gutiérrez⁵, Margarita Caballero⁶, Alexis Oliva⁷, Margarita Palmieri⁸, María Renée Álvarez⁸ and Antje Schwab²

¹Tecnológico Nacional de México/I. T. de la Zona Maya. Carretera Chetumal-Escárcega km 21.5, ejido Juan Sarabia, 77965 Quintana Roo, México.

10 ²Institut für Geosysteme und Bioindikation, Technische Universität Braunschweig, Langer Kamp 19c, 38106 Braunschweig, Germany.

³Tecnológico Nacional de México/I. T. de Chetumal. Av. Insurgentes 330, Chetumal, 77013 Quintana Roo, México.

⁴Institut für Geographische Wissenschaften, Freie Universität Berlin, Malteserstrasse 74-100, 12249 Berlin, Germany.

⁵El Colegio de la Frontera Sur (ECOSUR), Unidad Chetumal, Av. Centenario km 5.5, Chetumal, 77014 Quintana Roo, México.

15 ⁶Instituto de Geofísica, Universidad Nacional Autónoma de México, Ciudad Universitaria, 04510 Ciudad de México, México

⁷Asociación de municipios del lago de Yojoa y su Área de influencia (Amuprolago), Aldea Monte Verde, Santa Cruz de Yojoa, Cortés, 21000 Honduras.

⁸Departamento de Biología. Universidad del Valle de Guatemala, 01015 Guatemala, Guatemala.

20 *Correspondence to:* Sergio Cohuo (sergio.cd@chetumal.tecnm.mx)

Abstract. Geodiversity is recognized as one of the most important drivers of ecosystem characteristics and biodiversity globally. However, in the northern Neotropics, the contribution of highly diverse landscapes and geological history in structuring large-scale patterns of aquatic environments and aquatic species associations remains poorly understood. We evaluated the relationships among geodiversity, limnology and freshwater ostracodes from southern Mexico to Nicaragua.

25 Cluster analysis, based on geological, water-column physical and chemical variables, and geochemical and mineralogical characteristics of 76 aquatic ecosystems (karst, volcanic, tectonic) revealed two main limnological regions: (1) karst plateaus of the Yucatán Peninsula and northern Guatemala, and (2) volcanic terrains of the Guatemalan highlands, mid-elevation sites in El Salvador and Honduras, and the Nicaraguan lowlands. In addition, seven subregions were recognized, demonstrating a high heterogeneity of aquatic environments. Principal Component Analysis identified water chemistry (ionic composition) and mineralogy as most influential for aquatic ecosystem classification. Multi-parametric analyses, based on biological data, revealed that ostracode species associations represent disjunct faunas. Five species associations, distributed according to limnological regions, were recognized. Structural equation modelling (SEM) revealed that geodiversity explains the limnological regions and subregions of the study area. Limnology further explained species composition, but not species richness. The influence of conductivity and elevation were statistically significant for ostracode species composition, though not for species richness. We conclude that geodiversity has a strong influence on the limnological characteristics of aquatic systems, which, in turn, influence ostracode species composition in lakes of the northern Neotropical region.

1 Introduction

Geodiversity is defined as the natural variety of geological ([bedrock](#)), geomorphological ([elevation](#)), [soil features and assemblages \(mineralogy, sedimentology\)](#), [fossil and](#) hydrological features of a landscape (Gray, [2004](#); 2019; [Schrodt et al., 2018](#)). Geodiversity reflects aspects of regional or local climate, hydrology, and soil properties (Vartanyan, 2006a; Hu et al., 2020). By interacting with the biosphere and the atmosphere, geodiversity can also contribute, through sediment delivery, to the input of nutrients into ecosystems, and determine the chemical composition of environments (Vartanyan, 2006b; Bravo-Cuevas et al., 2021). Geodiversity has been fundamental for ecosystem development and biological evolution at scales ranging from local to global (Mittelbach et al., 2007; Etienne and Apol, 2009; Smith et al., 2010; Bryson et al., 2013; Gillespie and Roderick, 2014; Steinbauer et al., 2016).

Biological diversity, defined as the variety of life forms in a place on Earth (Huston, 1995), is strongly related to geodiversity, as species are distributed in response to landscape features. Biological diversity is dynamic. Species evolve and distribution patterns change at rates different from rates of change of geodiversity. This suggests that other intrinsic (e.g., ecological) and extrinsic (e.g., climate) factors, either ancient or recent, have interacted to shape regional species pools. In areas with highly dynamic and complex biological systems, such as tropical regions (Rull, 2011; Antonelli et al., 2018; Matzke-Karasz et al., 2019; Moguel et al., 2021), it is difficult to discern the contributions of individual versus joint factors that shaped the distribution of past and modern biota (Rossetti and Toledo, 2016). Understanding the role of geodiversity in the structure of biological systems is key to identifying evolutionary traits and mechanisms that control species distributions, and can be applied to conservation, ecosystem management, and prediction of ecosystem responses to future climate scenarios (Martiny et al., 2006; Hulsey et al., 2010; Jiménez-Alfaro et al., 2018).

The northern Neotropical region extends from central-southern Mexico to Central America and includes the Caribbean. It is characterized by a dynamic geological history, caused by the interplay of the North American, Cocos and Caribbean tectonic plates (Molnar and Sykes, 1969; Marshall, 2007). The region is characterized by broad ranges of elevation and soil types, displays frequent volcanic and seismic activity, and has been subjected to repeated marine regressions and transgressions (Brezonik and Fox, 1974; Horn and Haberyan, 1993; Umaña et al., 1999; Haberyan et al., 2003; Obrist-Farner et al., 2021). At least 16 biogeographic provinces are recognized, based on terrestrial plant and animal taxa (Morrone, 2014).

Numerous studies have attempted to elucidate the relationship between geodiversity and biological diversity and identify the factors that account for the current biogeographic patterns in the northern Neotropics (Wallace, 1853; Patton et al., 1994; Gillespie and Roderick, 2014). Most evidence from terrestrial taxa suggests *in situ* diversification, resulting from repeated colonization events by North and South American taxa, before and after the closure of the Isthmus of Panama, estimated to have occurred between 15 and 4 Mya (Bacon et al., 2015; Montes et al., 2015). Molecular evidence suggests that extant Mesoamerican terrestrial taxa (i.e., angiosperms, ferns, birds, reptiles and mammals) originated primarily in the Amazon Basin, with ancestors arriving by dispersal during the last 10 Mya (Antonelli et al., 2018).

70 These large-scale species movements between the American continents were mainly associated with large-amplitude
Pleistocene climate fluctuations such as glacial and interglacial cycles and episodes of shorter, centennial to millennial
fluctuations, such as Late Glacial Heinrich and Dansgaard-Oeschger stadials (Behling et al., 2000; Carnaval and Moritz, 2008;
Bouimetarhan et al., 2018; Baker et al., 2001, 2020). It is therefore assumed that both geodiversity and climate change have
been fundamental in shaping the current patterns of species diversity and distribution of terrestrial taxa (Haffer, 1969; Wiens
and Donoghue, 2004; Peterson and Nyári, 2008; Smith et al., 2012; Leite et al., 2016).

75 In aquatic environments of the northern Neotropics, diversification and drivers of species distribution are less well known than
those that operated in terrestrial environments. For instance, most aquatic species have dispersal limitations generated by
system morphologies and environmental variables, such as sediment texture, geochemistry, and water physico-chemical
properties, and how they have been modified through time (Albert and Reis, 2011; Bagley and Johnson 2014; Cabassi et al.,
2019). The limnology of a region, defined as the set of physical, chemical, and biological components of inland waters, and
80 its interactions with terrestrial, atmospheric, anthropogenic, and geological elements, in a spatial and temporal framework
(Last, 2002; Azim, 2009; McCullough et al., 2021), is generally accepted as a fundamental driver of the diversification and
distribution of aquatic species (Matamoros et al., 2015). The relevance of limnology for recent or past biological community
composition and species richness in the northern Neotropics, however, has been poorly evaluated.

85 During the last 50 years, anthropogenic influences on aquatic environments have altered biological diversity and species
distributions because of the modification of natural conditions (Albert and Reis, 2011; Wehrtmann et al., 2016; Franco-Gaviria
et al., 2018). Currently, most aquatic environments in the northern Neotropics are used as potable water sources and for
agriculture; in large lakes, fishing and aquaculture have caused eutrophication and introduced invasive species (e.g.,
Oreochromis niloticus). In addition, some lakes located close to urban centers and farmlands are used as disposal sites for
waste waters, agrochemicals, and mine residues (McKaye et al., 1995; Soto et al., 2020). The high heterogeneity of geodiversity
90 in the northern Neotropics, and the poor knowledge of the limnological attributes of aquatic ecosystems in the region make it
difficult to: (1) understand the complex relations among geology, water-column physico-chemical variables and biological
characteristics (community composition and species richness) and (2) distinguish different limnological regions and determine
their geographic coverage. There is great uncertainty about how geodiversity and limnological variables influenced aquatic
biota over time, and what factors have determined current patterns of diversity and distribution.

95 Freshwater ostracodes are a well-suited group to evaluate past and present drivers of species distribution in the northern
Neotropics. Ostracodes are bivalved microcrustaceans that are abundant, diverse, and widely distributed in aquatic ecosystems
(Pérez et al., 2011b, 2013a; Cohuo et al., 2016, 2020; Macario-González et al., 2018; Echeverria-Galindo et al., 2019). This
taxonomic group shows levels of endemism (restricted distribution) as high as 74%, sometimes confined to a single lake, or
100 found in surface waters throughout the region (Cohuo et al., 2016). In sediment sequences from lakes of the northern
Neotropics, ostracode remains are abundant, particularly in late Pleistocene deposits (Pérez et al., 2011b, 2013a; Cohuo et al.,
2020). The greatest limitation for using freshwater ostracodes to identify drivers of species distribution in the northern

Neotropics is the scarcity of integrated and comparable regional studies, for which detailed spatial and temporal limnological and biological data were collected.

105 We conducted a limnological and biological survey of aquatic systems across the northern Neotropics, from southeast Mexico to Nicaragua, to obtain environmental data across broad spatial scales to answer two main questions: (1) To what extent does geodiversity control limnological variables and thus define limnological regions?; (2) how does geodiversity (water and sediment geochemistry characteristics, as influenced by geology) affect diversity and distributions of ostracode species?

110 **2. Materials and Methods**

2.1. Study area

Our study area covers the northernmost Neotropics, ranging from southern Mexico (Yucatán Peninsula) to Nicaragua (Fig. 1). This region is considered a biodiversity hotspot (Mesoamerican hotspot; Myers et al., 2000), with more than 5000 endemic vascular plants (De Albuquerque et al., 2015), and about 1120 bird (~200 endemic), 440 mammal (65 endemic), 690 reptile
115 (~240 endemic), 550 amphibians (~350 endemic), and 500 fish species (~350 endemic) (CEPF, 2021). Species from North and South America converge in this region (Myers et al., 2000; Ojeda et al., 2003; DeClerck et al., 2010; Rull, 2011). The orography is highly irregular, and elevations range from sea level to more than 4500 m a.s.l. (Molnar and Sykes, 1969; Marshall et al., 2003, 2007). Plate tectonic interactions are responsible for active volcanism along the Central American Volcanic Arc and high seismic activity (Marshall et al., 2003, 2007). The climate is typically tropical (Köppen-type group A-climate
120 “tropical/megathermal climate”, Peel et al., 2007) and predominantly warm at low elevations (26°C mean annual temperature) (Waliser et al., 1999). Because of the irregular orography, at least a dozen climate sub-zones are distinguished (Taylor and Alfaro, 2005). Mean annual precipitation in the study area ranges from <500 to >3000 mm and is highly seasonal, governed by the seasonal migration of the Intertropical Convergence Zone (ITCZ). The northern position of the ITCZ during summer results in the so called “rainy season,” during which precipitation increases from ~240 mm in April to more than 1600 mm in
125 September-October (Hastenrath, 1967; Magaña et al., 1999). The hurricane season extends from July to December and is an important contributor to the humidity budget because on average, 300 mm day⁻¹ of rain falls during tropical storms and hurricanes (Jury, 2011). The region is rich in aquatic systems that are of different origins, shapes, and hydrological dynamics, as well as water chemistry and sediment composition. The karst Yucatán Peninsula, for example, with about 8000 *cenotes* (sinkholes), many lakes, and both surface and subterranean rivers, is considered a unique hydrological region (Schmitter-Soto
130 et al., 2002a, Alcocer and Bernal-Brooks, 2010). In Central America, lakes and wetlands cover more than 8% of the total land area (Ellison, 2004). Most important aquatic ecosystems in the northern Neotropics include coastal, tectonic and volcanic lakes, such as crater lakes and maars, karst waterbodies including lakes, *cenotes* and *aguadas* (water accumulated in topographic depressions under canopy cover), flooded caves, subterranean rivers, and both permanent and ephemeral ponds (Brezonik and Fox 1974; Pérez et al., 2011a; Delgado-Martínez et al., 2018; Echeverría-Galindo et al. 2019; Obrist-Farner and
135 Rice, 2019).




Figure 1: Simplified geological map of the northern Neotropical region showing the locations of the 76 studied aquatic ecosystems. Colors indicate geological units based on bedrock type and age of the sediments. Geological data were obtained from Garrity and Soller (2009). Black dots and numbers represent sampling localities. Detailed information on sampling sites can be found in Table 1. Legend: K_ Cretaceous sedimentary rocks; Kg_ Cretaceous Plutonic rocks; PZ, PZvf, PZx_ Paleozoic sedimentary rocks; Q_ Quaternary sedimentary rocks; Qvf, Qvm, TQv_ Quaternary volcanic rocks; T_ Tertiary sedimentary rocks of undetermined age; TRJ_ Jurassic sedimentary rocks; eT_ Eocene sedimentary rocks; mT_ Miocene sedimentary rocks; mTvfi_ Miocene volcanic rocks; nT_ Eocene sedimentary rocks; oT_ Oligocene sedimentary rocks; paT, pgT_ Paleocene sedimentary rocks.

145 2.2. Sampling aquatic environments in the northern Neotropics

A total of 76 aquatic ecosystems located in five countries across the northern Neotropical region (Fig. 1) were sampled during July-October 2013, coinciding with the rainy season in the region. These systems are situated on the Yucatán Peninsula, Mexico (n=28), Guatemala (n=26), El Salvador (n=14), Honduras (n=5) and Nicaragua (n=3) (Fig. 1). For all water bodies,

150 physical and chemical variables (temperature, dissolved oxygen, pH, conductivity) were measured *in situ* with a WTW Multi Set 350i multi-parameter probe at a water depth of 0.5 m. Maximum water depth at each site was measured with an echosounder Fishfinder GPSMAP 178C. Location of sites, including elevation, latitude, and longitude, was determined with a navigator Garmin GPSmap 60c.

155 Water samples for analysis of major anions (Cl^- , SO_4^{2-} , CO_3^{2-} , HCO_3^-) and cations (Ca^{2+} , K^+ , Mg^{2+} , Na^+) were collected at water depths of 0.5 m below surface using a Ruttner-type water sampling bottle. All water samples were filtered *in situ* using a 0.45- μm pore size Whatman glass microfiber filter. For cation analysis, filtered samples were acidified with HNO_3 to pH 2. Waters were stored under refrigeration until analysis.

160 Biological samples were collected from the littoral zone and deepest area of the profundal zone. [In most lakes](#), we collected five samples [in the littoral areas](#), using a 250- μm -mesh hand net [and five surface sediment samples collected from the littoral to the deepest area with an Ekman grab](#). Collection sites were distributed at regularly spaced intervals across the systems and were mostly characterized by submersed vegetation. [For sediment samples, to ensure we collect only extant specimens, we only analyzed the uppermost 3 cm of each grab. In six large lakes \(~75 km²\), such as Chichancanab \(Mexico\), Petén, Atilán \(Guatemala\), Ilopango \(El Salvador\), Yojoa \(Honduras\), Nicaragua, and Managua \(Nicaragua\), seven water and seven sediment samples were taken.](#)

165 **2.3 Environmental variables: water chemistry, sediment geochemistry, mineralogy, and geology**

170 Water chemistry analysis: Ionic composition was analyzed following Armienta et al. (2008). Bicarbonate was measured by acid titration to pH 4.6, using a mixed indicator of methyl red and bromocresol green. Concentrations of calcium and magnesium were obtained by complexometric titration with EDTA, whereas sodium and potassium were measured by atomic emission spectroscopy. Chloride was potentiometrically determined using an ion selective electrode, adding a 5 M solution of NaNO_3 as an ionic strength adjuster. Sulfate was determined by turbidimetry. Analytical quality was checked by ionic charge balance and most samples were balanced, with < 5% error. Major ions concentrations [were](#) expressed in mg L^{-1} , but the data were transformed to meq L^{-1} and percentages to determine anion and cation dominance and water type and test the charge balance. Sodium and potassium were summed. Ternary plots were constructed using the PAST 4.03 software (Hammer et al., 2001).

175 Geochemical analysis: Total carbon (TC) and total nitrogen (TN) contents in sediments were determined by combustion under oxygen saturation with a LECO TruSpec Macro CHN analyzer. Total inorganic carbon (TIC) was quantified with a Woesthoff Carmhograph C-16 after dissolution with phosphoric acid (45% H_3PO_4) and detection of the CO_2 -induced conductivity change in NaOH. Total organic carbon (TOC) was calculated by subtracting TIC from TC.

180 Mineralogical analysis: Qualitative and semi-quantitative mineralogical compounds in sediments were examined by x-ray diffraction with a RIGAKU Miniflex600. For the identification and semi-quantification of the minerogenic components, the software Philips Highscore was used. All sediment analyses used are described in detail in Vogel et al. (2016).

Geological analysis: The geological map of North and Central America generated by the Geological Society of America (GSA) (Reed et al., 2005) and adapted and converted to a geographic information system (GIS) by Garrity and Soller (2009), was used to identify geological regions in our study area. ArcGIS software was used to identify geological attributes of sampling sites such as bedrock type and age of sediments. Three major types of bedrock were distinguished: sedimentary, volcanic and plutonic rocks. Ten geologic periods and epochs, respectively, were defined: Jurassic, Cretaceous, Tertiary of undetermined age, Paleogene of undetermined age, Paleocene, Eocene, Oligocene, Neogene of undetermined age, Miocene, and Quaternary.

2.4 Limnological regionalization using cluster analysis and PCA

The final data set included 23 variables, of which 21 were numerical and the remaining two were categorical (Table S1). Numerical variables included elevation, water physical and chemical properties (temperature, dissolved oxygen, pH, conductivity, CO_3^{2-} , HCO_3^- , SO_4^{2-} , Cl^- , Na^+ , K^+ , Ca^{2+} and Mg^{2+}), sediment geochemistry (total carbon (TC), total inorganic carbon (TIC), total organic carbon (TOC), total nitrogen (TN)) and mineralogy (quartz, carbonate, phyllosilicates, and feldspars). Categorical data were geological properties represented by bedrock type and age of sediments.

Prior to statistical analysis, numerical data were log-transformed, except for pH, which is already a log-transformed value, to achieve an approximate normal distribution of variables. Normality was verified for all variables using the Shapiro-Wilk test. Missing data were substituted with the mean value of the respective variable, as missing values represented < 8% of the data set (Jakobsen et al., 2017).

We performed a cluster analysis (CA) to define groups of lakes based on the similarity of their measured attributes. For this analysis, we included all numerical variables. We used the unweighted pair group method with arithmetic mean (UPGMA) for the CA, and Euclidean distance to investigate the grouping similarity of sampling points. Calculations were conducted in R software (R Development Core Team, 2015), using the Vegan package (Oksanen et al., 2017).

We then used a Principal Component Analysis (PCA) for each of the main groups discriminated by the cluster, to identify correlated and explanatory variables of the data sets. For each group, the first PCA run included all 23 variables measured (numerical and categorical) and those represented by superimposed arrows in the graphs were considered correlated and excluded from further statistical analysis. A second PCA run, using uncorrelated variables, was used to identify explanatory variables of the data sets. The PCA-mixdata package implemented in R software (Chavent et al., 2014) was used because of its ability to handle quantitative and categorical data simultaneously. The loading values for all parameters were obtained using normalized rotation.

To provide a graphical representation of the most meaningful variables of the data sets detected in the PCA, and further evaluate latitudinal and/or altitudinal environmental gradients, we created an environmental variable-specific map using kriging interpolation. We used an empirical semi-variogram to quantify the spatial composition and structure of the data (Wagner, 2003; Bivand et al., 2008). We then fitted a theoretical variogram using the nugget effect (0.5), and a partial sill (0.5) in a linear model. We ultimately obtained a map representing measured data and estimates from unmeasured locations. The software Surfer® from Golden Software, LLC (www.goldensoftware.com) was used for calculations.

2.5. Biological analysis: ostracode abundances, identification, and statistics

Ostracode extraction and counting were carried out using 15 cm³ of wet sediment. Specimens were picked using a [Leica MZ75](#) stereomicroscope [and sorted according to external morphology](#). Only adult specimens, represented by single valves, empty carapaces or complete organisms (carapace and soft body) were counted. [For morphological identification down to species level, at least](#) three adult specimens with complete soft parts [were dissected](#). Identification using more than a single organism is appropriate in ostracodes, given morphological plasticity and mutations that can modify morphology. Specimens were dissected using distilled water and glycerin (3%) under a stereomicroscope. Selected shells from identified species were photographed with a Zeiss Axio Imager 2 microscope. Shells were stored in micropaleontology slides. Dissected soft parts were mounted on individual slides with Hydromatrix®. Species-level identification was conducted using species keys provided by Karanovic (2012). When keys did not fully resolve taxonomic identities, original descriptions were consulted. Taxonomic classification follows Cohuo et al. (2016). Undissected material was preserved in Eppendorf plastic vials with 70% ethanol and is currently available at the ostracod collection of the Instituto Tecnológico de Chetumal, Mexico. Shannon diversity index, used for species diversity metrics, was calculated with PAST 4.03 software (Hammer et al., 2001).

Ostracode species associations: Species associations were examined by means of non-metric multidimensional scaling (NMDS) (Legendre and Legendre, 1998). This procedure generates an ordination in a two-dimensional space, representing the pairwise dissimilarity between species according to their occurrences. We used the Bray-Curtis dissimilarity coefficient on a presence-absence database (Sørensen Coefficient), as count data were highly heterogenous (with species presence in subsets of samples and absences in most of the database), to which NMDS is sensitive. Only species with at least two occurrences were included in this analysis. In NMDS graph, species associations were determined with a hierarchical cluster analysis based on Ward distances. To test significance between species groups discriminated in NMDS, a permutational multivariate analysis of variance (PERMANOVA) was performed. We used a permutation with 9999 replicates and applied the Bonferroni correction. Calculations were done with R software, using the Vegan package (Oksanen et al., 2017).

Relating environmental variables and ostracode abundances: The relative importance of geochemical, mineralogy, and water physical and chemical conditions for ostracode species composition and distribution was assessed by multivariate constrained ordination techniques. We used ostracode relative abundance and a set of uncorrelated variables for calculations. First, we performed a Detrended Correspondence Analysis (DCA) with detrending by segments and non-linear rescaling to estimate the extent of the environmental gradient (Hill and Gauch, 1980) and to decide which type of ordination would be more appropriate for our data set. Species–environmental relationships were then analyzed using Canonical Correspondence Analysis (CCA) (Ter Braak, 1986). Environmental variables were standardized and added by forward selection using the Monte Carlo permutation test with 999 permutations ($\alpha = 0.05$). Calculations and final ordination graphs of the DCA and CCA were performed using [the Canoco software](#) version 5 (Šmilauer and Lepš, 2014).

2.6. Structural equation modeling


250 To disentangle the relationships between geodiversity, limnology and species composition (as a function of distribution) and
richness we used structural equation modeling (SEM), [the R software and the package Lavaan \(Rosseel, 2012\)](#). This is a
multivariate statistical technique that enables one to model pre-defined causal relationships between observed and non-
observed (latent) variables and test their statistical significance (Fan et al., 2016; Sarstedt and Ringle, 2020). Our conceptual
255 model for SEM was based on the assumption that geodiversity in the northern Neotropics is heterogenous, relatively constant
in the Yucatán Peninsula and Guatemalan lowlands, but highly variable in the rest of our study area (Guatemalan highlands-
Nicaragua). Aquatic systems [were](#), consequently, partially, or entirely influenced by underlying geodiversity, and thus
environmental variables such as sediment geochemistry and mineralogy, water ionic composition, temperature, and
conductivity of lake waters (limnology) [were](#) modified in response to geodiversity changes. At the regional scale, geodiversity
and limnology [were](#) expected to exert a direct or indirect influence on ostracode species richness and community composition.
260 The individual influence of variables forward selected in the CCA as explanatory for species-environment relationships, and
those that may display environmental gradients such as elevation, conductivity, and TOC, were also tested in the models
evaluated.

For this conceptual framework, ‘geodiversity’ (latent and exogenous variable) was constructed with all or a subset of
geological (bedrock type and age and elevation) and mineralogical variables. ‘Limnology’ (latent and endogenous variable)
265 was constructed using geochemistry (TOC), and water physical and chemical (major anion and cation, temperature, pH, and
conductivity) variables. Species richness was treated as an observed variable, and the latent variable ‘species composition’
was constructed using NMDS associations. Using a covariance matrix with a set of uncorrelated variables, we fitted five
models using this conceptual framework. For all models, statistical significance was tested with Root Mean Square Error of
Approximation (RMSEA), comparative fit index (CFI) and standardized root mean squared residuals (SRMR). The predictive
270 power of the model (R-square) was measured based on the amount of variation of the biological data. The most parsimonious
model fitting our data set was selected as the explanatory model (Section S1 in the Supplement).

3 Results

275 3.1 Limnological regionalization in the northern Neotropics

Cluster analysis identified two main groups that represent limnological regions (Fig. 2). The first group (YG: Yucatán and
Guatemala) consists of lowland lakes from the Yucatán Peninsula (Mexico), the Petén district (northern Guatemala) and the
Pacific lowlands of southern Guatemala. The second group (GSHN: Guatemala, Salvador, Honduras, Nicaragua) consists of
Guatemalan highland lakes, El Salvador and Honduras mid-elevation lakes and Nicaraguan lowland lakes.



280

Figure 2 (a) Cluster analysis dendrogram of the 76 studied aquatic ecosystems using 23 water physico-chemical, sediment geochemistry, mineralogical, and geological variables. Blue line indicates cut-off criterion for cluster partition. Two major groups and seven subgroups were detected and named according to their position in the study area: Yucatán Peninsula/Northern Guatemala (YG) and Guatemala highlands, El Salvador, Honduras, Nicaragua (GSHN). In order to provide a graphic representation of cluster grouping, a color bar was assigned to each group and lakes within these groups were plotted on (b) YG map and (c) GSHN map using the same color. Lake full names of codes presented in cluster and maps are given in Table 1. Numbers of lake name codes correspond to that in Figure 1.

285

290 Table 1 shows a list of all the studied aquatic ecosystems located in the YG and GSHN limnological regions and subregions, as well as information about selected attributes determined in this study. Detailed results of water physico-chemical, geochemical, mineralogical, and geological analyses for all studied water ecosystems can be founded in Table S1.

Table 1. Limnological regions in the northern Neotropical region, with lakes main water physico-chemical, mineralogical, and geological properties.

295

Limnology		Country	Site		ID	Coordinates		Origin	Elevation (m.a.s.l)	Surface area (km ²)	Depth ^a (m)	Water type (ionic dominance)	Dominant minerals in sediment
group	Sub group		number	Site		N	W						
YG	YG1	GUA	1	Lachuá	LAC	15.92	90.67 32	tectonic	170	4	378	SO ₄ <<H CO ₃ --Ca	Calcite
		MEX	2	Candelaria River	CAN	18.18 1	91.04 93	lectono- karstic	32	river	1.8	HCO ₃ << SO ₄ --Ca	Calcite
		MEX	3	Encantada	ENC	18.49 9	88.38 95	karst	4	0.07	3	SO ₄ -- Ca<<Mg	Calcite
		MEX	4	Bacalar	BAC	18.70 1	88.38 19	karst	4	51	16	SO ₄ -- Ca<<Mg	Calcite
		MEX	5	Sabana de Chetumal	SAB	18.51 7	88.34 88	karst	7	0.88	1.6	SO ₄ --Ca- -Mg	Nd
		GUA	6	El Rosario	ROS	16.52 6	90.16 01	karst	126	0.02	3	HCO ₃ << SO ₄ -- Ca<<Mg	Dolomite
		GUA	7	Petexbatún	PET	16.42 1	90.18 62	tectono- karstic	120	5.6	40	HCO ₃ << SO ₄ -- Ca<<Mg	Calcite
		GUA	8	Salpetén	SAL	16.98 1	89.67 55	tectono- karstic	105	2.77	10	SO ₄ --Ca- -Mg	Calcite
		MEX	9	Chichancanab	CHI	19.88 4	88.76 82	tectonic	1	5.1	15	SO ₄ -- Ca<<Mg	Calcite
	YG2	GUA	10	La Gloria ^b	GLO	16.95 5	90.37 89	tectono- karstic	138	3.6	65	SO ₄ <<Cl --Ca	Nd
		MEX	11	Camp	CAMP	18.03 7	90.98 87	Flooded depression	43	2.59	3.5	SO ₄ <<Cl --Ca	Nd
		GUA	12	Las Pozas	PO2	16.34 3	90.16 60	karst	152	2.16	35	HCO ₃ -- Mg<<Ca	Nd
		GUA	13	Petenchel	PE1	16.91 6	89.82 88	tectono- karstic	118	0.73	Nd	HCO ₃ -- Ca	Phyllosilic ates
		GUA	14	Poza azul	PO1	16.95 4	90.78 84	karst	160	0.08	78.1	HCO ₃ -- Mg<<Ca	Nd
		GUA	15	San Diego ^b	DIE	16.92 2	90.42 31	tectono- karstic	156	3.8	8.1	SO ₄ <<Cl --Ca	Nd
		GUA	16	Quexil	QUE	16.92 3	89.80 99	tectono- karstic	120	2.2	25.5	HCO ₃ -- Ca	Nd
		GUA	17	Oquevix	OQU	16.65 1	89.74 27	karst	150	1.6	10	HCO ₃ -- Ca	Quartz
		GUA	18	Yaxhá	YAX	17.05 8	89.38 97	tectono- karstic	164	7	22	HCO ₃ -- Ca	Quartz and

												phyllosilicates
	GUA	19	Peten Itzá ^b	ITZ	16.98 6	89.69 38	tectono- karstic	112	100	165	SO4<<H CO3-- Ca<<Mg	Nd
	GUA	20	El muchacho	MU1	13.88 9	90.19 18	Flooded depression	3	0.36	2	HCO3<< Cl-- Na<<Ca	Feldspars
	GUA	21	Grande	GRA	13.89 0	90.17 03	Flooded depression	5	0.95	2	HCO3-- Mg<<Ca	Phyllosilicates
	GUA	22	Comandador	COM	13.96 0	90.25 44	Flooded depression	20	0.65	1.7	HCO3-- Ca--Mg	Phyllosilicates
	GUA	23	Sepalau	SEP	15.78 6	90.21 67	karst	266	0.03	10.3	HCO3-- Ca	Quartz and phyllosilicates
	GUA	24	Sacnab	SAC	17.05 8	89.37 25	tectono- karstic	170	4.28	9	HCO3-- Ca	Quartz and phyllosilicates
YG3	MEX	25	Señor	SEN	19.87 6	88.07 75	karst	3	1.06	2	Cl<<SO4 -- Na<<Mg	Calcite
	MEX	26	Muyil	MUY	20.07 5	87.60 37	karst	-1	2.52	16	Cl<<HCO 3--Na-- Mg	Calcite
	MEX	27	Colac	COL	20.90 9	88.86 69	karst	11	0.0002	120	Cl<<HCO 3--Na-- Mg--Ca	Calcite
	MEX	28	Kaná	KAN	19.50 1	88.39 54	Flooded depression	5	1.01	2.5	HCO3<< SO4--Na- -Ca	Nd
	MEX	29	Sabak-ha	CSA B	20.58 0	89.58 81	karst	18	0.0002	90	HCO3<< Cl-- Na<<Ca	Calcite
	MEX	30	Mucuyche	MU2	20.62 4	89.60 65	karst	17	Cavern	18	Cl<<HCO 3--Na-- Ca	Ankerite
	MEX	31	Oxolá	OXO	20.67 8	89.24 17	karst	18	Cavern	16	HCO3<< Cl--Ca-- Mg	Calcite
	MEX	32	San Miguel	GMI	19.93 5	88.99 83	karst	32	Cavern	1.2	HCO3<< Cl--Ca-- Mg-Na	Calcite
	MEX	33	Caobas	CAO	18.44 5	89.10 06	Flooded depression	126	0.13	4	HCO3<< SO4--Ca	Calcite
	MEX	34	Yumku	YUM	20.57 8	89.60 52	karst	16	Cavern	15	HCO3<< Cl--Ca-- Mg--Na	Calcite
	MEX	35	Yalahau	YAL	20.65 7	89.21 70	karst	2	0.25	12	Cl<<HCO 3--Na-- Mg	Nd
YG4	MEX	36	Silvituc	SIL	18.64 3	90.27 27	unknown	69	7.9	3	HCO3-- Ca<<Na	Quartz and phyllosilicates
	MEX	37	San José de la Montaña	JOS	18.36 9	89.01 20	karst	118	2	3	HCO<<Cl --Na<<Ca	Nd

		MEX	38	Chacanbacab	LCH2	18.47 8	89.08 69	Flooded depression	109	257.29	3	HCO3-- Ca<<Na	Phyllosilic ates
		MEX	39	Sijil Noh ha	SIJ	19.47 3	88.05 54	karst	0	0.25	8	SO4<<Cl --Na--Mg	Nd
		MEX	40	Chacchoben	LCH1	19.03 7	88.18 11	Flooded depression	6	0.57	4	HCO3<< Cl-- Ca<<Na	Nd
		MEX	41	Vallehermoso	VAL	19.17 8	88.52 16	Flooded depression	18	0.22	3	HCO3-- Ca<<Na	Quartz
		MEX	42	Emiliano Zapata	EMI	19.19 7	88.46 91	Flooded depression	23	2.52	5	HCO3-- Ca<<Na	Quartz and phyllosilic ates
		MEX	43	Noh-bec	NOH	19.14 6	88.17 62	Flooded depression	1	8.5	2.5	Cl<<SO4 --Ca--Mg	Nd
		MEX	44	Miguel Hidalgo	MIG	18.78 6	88.36 74	Flooded depression	31	20.22	4	HCO3<< SO4-- Ca<<Na	Calcite
		MEX	45	Misteriosa	MIS	18.04 2	90.49 80	unknown	53	5	5.8	SO4<<Cl --Ca	Quartz
GSHN	GSHN1	SAL	46	Olomega	OLO	13.30 7	88.05 5	volcanic	66	25.2	2.5	HCO3-- Na<<Mg	Phyllosilic ates
		SAL	47	Apastepeque	APA	13.69 2	88.74 5	volcanic, crater lake	509	0.38	47	HCO3-- Mg<<Ca	Feldspars
		SAL	48	Gūujija ^b	GUI	14.26 1	89.50 1	volcanic, lava flows	431	45	22	SO4<<Cl --Ca<<Na	Nd
		SAL	49	Jocotal	JOC	13.33 7	88.25 2	volcanic, lava flows	26	8.7	3	HCO3-- Na--Mg	Feldspars and phyllosilic ates
		SAL	50	Nagualapa	NAG	13.47 0	89.00 2	volcanic	43	0.12	1	HCO3-- Na--Mg	Nd
		SAL	51	El espino	ESP	13.95 3	89.86 5	volcanic	689	0.99	5.5	HCO3<< SO4-- Mg<<Na	Phyllosilic ates
		SAL	52	Los negritos	NEG	13.28 3	87.93 7	volcanic	102	0.69	2	HCO3<< Cl-- Na<<Mg	Phyllosilic ates
		NIC	53	Masaya	MAS	11.99 6	86.11 6	volcano- tectonic	222	8.33	ND	HCO3-- Na<<Mg	Feldspars
		NIC	54	Managua	MAN	12.27 0	86.47 7	volcano- tectonic	41	1061.1 7	ND	HCO3<< Cl-- Na<<Mg	Feldspars
		SAL	55	Chanmico	CHA	13.77 9	89.35 4	volcanic, crater lake	477	0.78	51	HCO3<< SO4-- Mg<<Na	Feldspars
		SAL	56	Aramuaca	ARA	13.42 9	88.10 7	volcanic, maar lake	76	0.4	107	SO4<<H CO3-- Na<<Mg	Feldspars
		MEX	57	Tekom	TEK	20.73	89.46 60	karst	24	<0.01	3	HCO3<< SO4-- Ca<<Na	Calcite
		HON	58	Jucutuma	JUC	15.51 2	87.90 3	Flooded depression	27	4.34	2	HCO3-- Ca<<Na	Quartz and phyllosilic ates

	HON	59	Ticamaya	TIC	15.55 1	87.89 0	Flooded depression	17	2.91	2	HCO ₃ << Cl--Ca-- Na	Nd
	HON	60	Yojoa	YOJ	14.86 1	87.98 5	volcanic	639	79.7	22	HCO ₃ -Ca	Quartz
	SAL	61	Metapan	MET	14.30 9	89.46 6	volcanic	450	16	6	HCO ₃ -Ca	Quartz
GSHN2	SAL	62	Ilopango	ILO	13.68 2	89.07 1	volcanic, caldera lake	446	70.28	177	Cl<<HCO 3--Na	Nd
	SAL	63	Coatepeque	COA	13.86 2	89.55 3	volcanic, caldera lake	743	26	119	Cl<<HCO 3-- Na<<Mg	Feldspars
	NIC	64	Nicaragua	NIC	11.46 0	85.77 4	volcano- tectonic	37	8264	ND	HCO ₃ << Cl-- Na<<Ca	Nd
GSHN3	GUA	65	Chicabal	CH1	14.78 8	91.65 6	volcanic, crater lake	2726	0.21	10.3	SO ₄ --Mg	feldspars
	GUA	66	Calderas	CAL	14.41 2	90.59 1	volcanic, crater lake	1790	0.35	26	HCO ₃ -- Mg<<Ca	feldspars
	SAL	67	Verde	VER	13.89 1	89.78 7	volcanic, crater lake	1609	0.1	12	HCO ₃ -- Mg<<Ca	Feldspars
	GUA	68	San Juan Bautista	JUA	14.04 2	90.07 2	Flooded depression	1285	0.07	2	HCO ₃ -- Mg<<Ca	feldspars
	SAL	69	Alegría	ALE	13.49 3	88.49 3	volcanic, crater lake	1272	0.09	10	SO ₄ -- Ca<<Mg	Feldspars
	GUA	70	Atitlán	ATI	14.68 4	91.22 4	volcanic, caldera lake	1556	125	340	HCO ₃ -- Mg--Na	Nd
	GUA	71	Amatitlán	AMA	14.43 6	90.54 8	Volcanic	1193	15.2	23	HCO ₃ -- Cl-- Na<<Mg	Calcite
	GUA	72	Magdalena	MAG	15.54 3	91.39 6	tectonic	2863	0.01	2.8	HCO ₃ -Ca	Calcite
	HON	73	Madre vieja	MAD	14.35 7	88.13 8	Flooded depression	1866	0.1	3.4	HCO ₃ -- Mg<<Ca	phyllosilicates
	HON	74	Chiligatoro	CH2	14.37 6	88.18 3	volcanic	1925	0.04	5.5	SO ₄ --Mg	Quartz
	GUA	75	El pino	PIN	14.34 5	90.39 4	volcanic	1038	0.64	6	HCO ₃ -- Mg<<Ca	phyllosilicates
	GUA	76	Ipala	IPA	14.55 7	89.63 9	volcanic, crater lake	1495	0.56	25	HCO ₃ -- Mg<<Na	feldspars

a= Maximum sampled water depth


b= Pérez et al., 2011

Nd=not determined

300 For the YG region, the first PCA run identified thirteen uncorrelated variables. The second PCA run clearly explained the variation of the data set (Fig. 3a). The first (PC1) and second (PC2) components explained 38% of the total variance of the data set (Fig. 3a, Table S2.1). The PC1 accounted for 23.4%, and the PC2 for 14.6% of the total variance, respectively. The biplot based on component 1 and 2 indicated that conductivity (ranging from 175 to 3479 $\mu\text{S cm}^{-1}$) and related ions sodium

305


(Na⁺), chloride (Cl) and magnesium (Mg²⁺) were the variables that exhibit the highest correlations (<0.64) with the first principal component Fig. 3a-c). Thus, they represented the most influential variables differentiating aquatic ecosystems in the YG region. pH, ranging from 6.9 to 9.9, was highly correlated (>0.73) with the second component (PC2), suggesting that it is the second most influential variable of the YG aquatic environments (Fig. 3a, d, Table S2.1). Figures 3 b, c and d show regional distribution of the most meaningful variables for the YG region.



310 Figure 3 (a) PCA biplot based on 13 variables from 45 aquatic ecosystems of the Yucatán Peninsula and northern Guatemala (YG
 limnological region). Conductivity and related ions were the most meaningful variables characterizing aquatic ecosystems as they
 explain 23.4% of the total variance. The second most important variable is pH, accounting for 14.6% of the total variance. Arrows
 represent variables. Spatial representations of meaningful variables are presented in (b-d) interpolated maps of conductivity (b);
 315 chloride (c); pH (d). Abbreviations are as follows: temperature (temp) (°C), conductivity (cond) (μS cm⁻¹), magnesium (Mg²⁺),
 chloride (Cl⁻), bicarbonate (HCO₃⁻), potassium (K⁺), calcium (Ca²⁺), sodium (Na⁺), sulfates (SO₄²⁻) dissolved oxygen (DO) (μmol L⁻¹)

and total organic carbon (TOC). Major ions concentrations are expressed in mg L⁻¹. Blue numbers in maps represent measured values, black numbers are interpolated values.

320 For the GSHN region, the PCA based on 13 uncorrelated variables, explained 49.6% of the total variance of the data set, within the first (PC1) and second (PC2) components (Fig. 4a, Table S2.2). PC1 accounted for 26.8% and PC2 for the 22.8% of the total variance. The PCA biplot based on components 1 and 2, respectively, showed that water ionic composition, particularly content of bicarbonates (HCO₃⁻, ranging from 4 to 373 mg L⁻¹), sodium (Na⁺, ranging from 2 to 180 mg L⁻¹) and chloride, was the most influential in discriminating aquatic systems in the GSHN region, as it correlated >0.62 with the first component (Fig. 4a-c, Table S2.2). Geochemical and geological variables such as TOC (ranging from 5 to 22.9%), age and bedrock were the second most influential variables as they are strongly correlated with the second component (> 0.75) (Fig. 4a, d). Figure 4 b, c, d show the spatial distribution of meaningful variables for the GSHN region.



330 Figure 4 (a) PCA biplot based on 13 variables from 31 aquatic ecosystems of central-southern Guatemala, El Salvador, Honduras
and Nicaragua (GSHN limnological region). Ionic composition HCO_3^- , Na^+ , Cl^- were the most meaningful variables characterizing
aquatic ecosystems, as they explain 26.8% of the total variance. Sediment geochemistry and geology were the second most important
variables, accounting for 22.8% of the total variance. Arrows represent variables. Spatial representations of meaningful variables
are presented in (b-d) interpolated maps of bicarbonates (b); sodium (c); TOC (d). Abbreviations are as follows: conductivity (cond)
335 ($\mu\text{S cm}^{-1}$), magnesium (Mg^{2+}), sodium (Na^+), chloride (Cl^-), bicarbonate (HCO_3^-), carbonates (CO_3^{2-}), calcium (Ca^{2+}), total nitrogen
(TN) and total organic carbon (TOC). Major ion concentrations are expressed in mg L^{-1} . Blue numbers in maps represent measured
values, black numbers are interpolated values.

340

Water ionic dominance was graphically evaluated with ternary plots (Fig. S1 and Fig. S2) and information on lake water types
is shown in Table 1. Here, we highlight the most relevant characteristics for YG and GSHN limnological subregions:

3.2 The YG region composed by karst aquatic systems

In the YG limnological region, four subregions were identified in the cluster analysis. The first subregion (YG1; n=9) within
345 YG included systems located in central-southern Yucatán and in the northern Petén district in Guatemala (Fig. 2). Lakes are
located in the lowlands (<170 m a.s.l.) and most of them are relatively shallow (<16 m depth), such as Bacalar, Encantada,
Sabana Chetumal, Rosario, Salpetén, Chichancanab, except for Lake Petexbatún (40 m depth) and Lachuá (378 m depth). The
latter constitutes the deepest lake in the study area. Waters of these systems are dominated primarily by sulfates, followed by
calcium and magnesium. The YG limnological subregion YG2 (n=15) contained lakes that are mostly restricted to the
350 Guatemalan lowlands (Fig. 2). Most of these systems are large lakes, including Lake Petén Itzá, one of the largest (100 km^2),
deepest (165 m) and oldest (~400 ka) lakes of the northern Neotropics. Lake waters are dominated by carbonates, and therefore
calcium prevails in these lakes. Aquatic systems located in the central and northern portion of the Yucatán Peninsula were
grouped in YG3 (n=11) (Fig. 2). Colac, Sabak ha, Mucuyche, Oxolá, Gruta Miguel Hidalgo and Yumku are *cenotes* in northern
Yucatán, and lakes Yalahau, Caobas, Kaná, Señor and Muyil are located in the central-northern Peninsula. Carbonates
355 dominate lake waters and chloride shows high values. Limnological subregion YG4 (n=10) was constituted of lakes located
in the central-southern Yucatán Peninsula (Fig. 2). Lakes are relatively large, shallow and far from the Caribbean and Gulf of
Mexico coasts, at least >60 km, e.g., Silvituc and Chacanbacab. Waters are dominated by carbonates and calcium.
Mineralogical analysis reveals that most lakes are dominated by carbonates in the YG limnological region. In subregions YG1
and YG3 (central and northern Yucatán Peninsula), most lake sediments have calcite as the dominant mineral (Table 1, Fig.
360 S2). Lakes Chichancanab and Salpetén both belong to YG, but show carbonates with a co-dominance of phyllosilicates and
gypsum. Lakes from YG2 are mainly dominated by phyllosilicates and feldspars. The mineralogical composition of lake
sediments of the YG4 subgroup varies. Sediments of the lakes such as Vallehermoso, Emiliano Zapata and Chacanbacab are
dominated by phyllosilicates with or without feldspars. Sediments of lakes such as Miguel Hidalgo, on the other hand, are
dominated by carbonates, with calcite as main mineral, whereas Lake Silvituc is characterized by exotic minerals such as silver
365 and gold (Table S1).

3.3 The GSHN limnological region is composed by volcanic aquatic systems

In the GSHN region, three subregions were identified by cluster analysis. Lakes of GSHN1 (n=16) are located in Central America at low and middle elevations, ranging from 17 to 689 m a.s.l. Lake waters show a clear dominance of carbonates, followed by sodium, potassium and magnesium (Table 1, Fig. S3). The GSHN2 is composed of three of the largest lakes in Central America: Ilopango, Coatepeque and Nicaragua. These three lakes originated from volcanic activity. Lakes Ilopango and Coatepeque are caldera lakes, whereas Lake Nicaragua surrounds Volcanoes Concepción and Maderas. There is no clear pattern for ions in the water column, but sodium and potassium dominated, followed by magnesium. The GSHN3 (n= 12) is formed mainly by crater lakes located in the highlands of Guatemala, Honduras and El Salvador. Most lake waters are dominated by bicarbonates, followed by magnesium, whereas Chiligatoro, Chicabal and Alegría are dominated by sulfates (Table 1, Fig. S3).

Sediment mineralogy of Central American lakes (GSHN region) shows that most subgroups are dominated by feldspars. Co-dominance with other minerals such as phyllosilicates and carbonates occurs. The GSHN1 combined lakes dominated by phyllosilicates and feldspars, whereas quartz is the dominant mineral in Lakes Yojoa and Ticamaya (Honduras). GSHN2 included large lakes dominated by feldspars. In Lake Nicaragua, this dominance is also shared with phyllosilicates. For the GSHN3, we detected two main mineral assemblages. The first is dominated by phyllosilicates and feldspars, with clay minerals and feldspars as main minerals, and the second is dominated by feldspars and phyllosilicates.

3.4 Ostracode species associations and their relationship with limnological regions

We found ostracode species in 74 of the 76 of the aquatic systems we studied in the northern Neotropics. In the volcanic Lake Alegría (El Salvador) and the karstic cave San Miguel (Yucatán, Mexico), ostracodes were not observed. Living adult specimens were encountered in samples from all systems, except those from Lakes Chicabal, Tekoh, Yaxhá, Verde and Cenote Mucuyche, where only empty shells or single valves were recovered. Taxonomic analysis of species enabled us to identify 70 species (Table S3), out of which 31 were recorded at single sites, whereas the remaining 39 were observed in at least two systems. Species richness ranged between 1 to 9 with an average of 4 species per site, whereas the Shannon Diversity Index (H) maximum value was 2.1, corresponding to Lakes Bacalar and Petén Itzá and the Candelaria River. For all other lakes the index averaged 1.1. The list of ostracode species found in our study is presented in Table S3.

NMDS ordination, based on species occurrence data, revealed five major species associations (OST 1-5) with a reliable stress value of 0.08 (Fig. 5a) (Clarke, 1993). The PERMANOVA test showed statistically significant differences between group centroids ($F = 1.19$, $p = 0.0001$), thus supporting NMDS ordinations. Ostracode groups 1 and 2 are located in the YG limnological region (karst terraces), and groups 3-5 in the limnological region GSHN (volcanic Guatemalan highlands and Central American mid-elevations and lowlands). The first species group (OST1) consisted of twelve ostracode species, recorded from lakes and *cenotes* from the eastern Yucatán Peninsula (Fig. 5b). Most of these species are tolerant of high conductivity (particularly related to Na^+ and Cl^-), such as *Heterocypris punctata* and *Limnocythere floridensis* (Fig. 5c)

400 | *Cyprideis* cf. *salebrosa* and *Perissocytheridea* cf. *cribrosa*. The second species group (OST 2) included 11 species, distributed
in lakes and ponds of the southern Yucatán Peninsula and northern Guatemala (Fig. 5b). Some of these species (Fig. 5b), such
as *Cypria petenensis*, *Cypretta campechensis* and *Paracythereis opesta*, are considered endemic (Cohuo et al., 2016). Some
others, such as *Alicenula serricaudata*, *Pseudocandona antilliana* and *Cytheridella ilosvayi*, have a wide Neotropical
distribution (Cohuo et al., 2016). The third species group (OST 3) was integrated of seven species distributed mainly in mid-
405 | elevation lakes from Guatemala, El Salvador, and Honduras (Fig. 5b). Several of these species have very restricted distributions
and correspond to the *Strandesia*, *Keysercypria* and *Cypridopsis* genera (Fig. 5c). The fourth species group (OST 4) is
composed of six species: *Heterocypris nicaraguensis*, *Potamocypris islagrandensis*, *Cypria granadae*, *Limnocytherina royi*,
Perissocythere marginata and *Cyprideis* sp., distributed exclusively in lakes in Nicaragua (Figs. 5a, b). The fifth group (OST
5) included only three species from highland lakes of Guatemala: *Chlamydotheca* cf. *colombiensis*, *Cypria* sp. 4 and
Cypridopsis sp. 7 (Figs. 5b, c).

and Lepš, 2014). Using the forward selection technique, we found a set of eight variables that best explain the variance of data (Table S4.1). The first two axes of the CCA accounted for 47.4% of the variance of the species-environment relationship and 13.1% of the variance of species data. The summary of performance and ordination of this analysis are presented in Table S4.2. The first axis of the CCA plot (Fig. S3) explained 25.8% of the correlation between species and explanatory variables, with a canonical coefficient of 0.95 and an eigenvalue of 0.81. Sedimentary bedrock, TOC, temperature, and feldspars were the most important factors that influenced species composition on axis 1 (Fig. S3). Axis 2 was strongly correlated with plutonic bedrock, elevation, and phyllosilicates. This axis explained 21.6% of the species-environment relationship, with a canonical coefficient of 0.90 and an eigenvalue of 0.68. In the CCA ordination, OST1 was positively correlated with conductivity and bedrock of sedimentary origin. OST 3 and OST 4 were positively correlated to volcanic and plutonic bedrock. OST 5 was associated with sediments dominated by feldspars. OST2 was positioned close to the center of the axes, suggesting that species are broadly tolerant of the variables analyzed.


3.5 Structural equation modelling

Five models of the relationships of geodiversity, limnology and species composition and richness were tested with structural equation modelling (SEM). Four models used geodiversity only as exogenous variables (models 1-2, 4-5), and in one model (model 3) limnology was also considered as an exogenous variable (two exogenous variables, explaining an endogenous variable). Description of the rationale behind variable selection and relationships tested in each model are found in section S1 of the Supplementary Material. Model 1 evaluated the direct influence of geodiversity on limnology and the influence of limnology on species composition and richness, resulting in the following metrics of global fit: CFI 0.63 (values close to 1.00 indicate better fit of the model), RMSEA 0.19 (values < 0.05 are considered reliable; Fabrigar et al., 1999), and SRMR 0.12 (values < 0.08 are generally considered a good fit; Hu and Bentler, 1999). Model 2, tested relationships similar to model 1, but additionally the relevance of TOC on species composition and richness, with the following statistics: CFI 0.79, RMSEA 0.13, and SRMR 0.14. Models 1 and 2 received relatively low values of global fit, all below the threshold of statistical significance. Values of a Chi-square test were < 0.05 in these two models, and they were therefore rejected as explanatory models. Model 3 evaluated the influence of two exogenous variables, geodiversity and limnology on species composition. Geodiversity and limnology were considered independent and without any influence on one another. Metrics of global fit of this model were better than in models 1 and 2, (CFI 0.94, RMSEA 0.06, and SRMR 0.14). However, the data were too small to calculate the statistical significance of both exogenous variables, and then, interpretation of the results is not reliable. This model was also rejected.

Model 4 and model 5 evaluated the same relationships between geodiversity, limnology, and species composition and richness as in model 1, but additionally, the individual influence of elevation and conductivity (model 4) and TOC and latitude (model 5) were analyzed. Model 5 received the following metrics of global fit CFI 0.62, RMSEA 0.20, and SRMR 0.17, all below the

threshold of significance, and the model was therefore rejected. The optimal model was model 4 (Fig. 6). This is supported by the following metrics of global fit, CFI = 0.94), RMSEA = 0.01, and SRMR=0.03. The optimal model suggests that geodiversity strongly influenced limnology. Limnology explained species composition (distribution), but not species richness. Elevation and conductivity did not explain species richness but demonstrated significant influences on species composition. The direct influence of limnology on species composition was statistically significant, whereas it did not exert an influence on species richness (p-value > 0.05).

All paths in figure 6 are statistically significant (p-value <0.05), except for the direct effect of limnology, conductivity, and elevation on species richness.



465

Figure 6. Structural equation modelling of the influence of geodiversity and related variables with freshwater ostracode species composition and distribution. Metrics of global fit of the optimal model CFI = 0.941, RMSEA = 0.01, and SRMR=0.035. Arrows indicate the direction of influence. Black arrows are paths statistically significant at p-value <0.05, and gray paths are not significant. Numbers next to arrows are the path standardized coefficients.

470

4. Discussion

4.1 Geodiversity defines two main limnological regions in the northern Neotropics

Our cluster analysis based on 76 lakes and 23 lake attributes shows limnological regionalization in the northern Neotropics. Two main regions, corresponding to Yucatán Peninsula-northern Guatemala (Group YG) and northern Central America (Group GSHN), were identified (Fig. 2). The group YG is located in karstic plateaus of sedimentary origin, dominated by limestone, dolomite, evaporites and carbonate-rich impact breccia (Hildebrand et al., 1995; Schmitter-Soto et al., 2002a, b; Vázquez-Domínguez and Arita, 2010). The Group GSHN is located in volcanic bedrock terrains of Guatemala, El Salvador, Honduras and Nicaragua, where pyroclastic and volcanic epiclastic materials, usually reworked, are abundant, reflecting active or past volcanic activity (Dengo et al., 1970; Stoiber and Carr, 1973; Carr, 1984). The YG and the GSHN groups were further subdivided into four (YG1-4) and three subgroups (GSHN1-3), representing limnological subregions. This proposed regionalization, therefore, reveals high heterogeneity of aquatic systems in the northern Neotropics. Multivariate statistics (PCA) shows that regions and subregions can be distinguished by ionic composition of waters. Geochemical variables related to sediments, such as TOC and mineral composition, are recognized as the second most important characteristics (Figs. 3, 4). In the YG karst region, lakes are characterized by carbonate-, calcium- and calcite-signatures, which is expected because waters interact with limestone and dolomite-rich bedrock on the Peninsula (Schmitter-Soto et al., 2002a, b; Perry et al., 2009). This is also responsible for the dominance of calcium, sodium, and magnesium ions in waters, which in turn are related to the generally alkaline surface waters in most aquatic systems of the region (Alcocer et al., 1998; Schmitter-Soto et al., 2002a, b) (Figs. 3a, d). In specific areas, such as YG3, dominance of chloride is also relevant. This can be explained by two main processes, 1) marine intrusion and 2) input of subterranean waters that have interacted with evaporites. The spatial distribution map of chloride contents in lake waters (Fig. 3c) shows a clear tendency to higher values on the northern Yucatán Peninsula where marine intrusion is probably the most important source of chloride (Sánchez-Sánchez et al., 2015; Saint-Loup et al., 2018). Marine intrusions in northern Yucatán have been mapped as far as 100 km inland (Steinich and Marín, 1996). Pérez-Ceballos et al. (2012) found that several water systems, mainly *cenotes*, in this same region are characterized by marine waters below freshwater lenses, with water intermixing. Sulfate is an interesting component of some lakes of the YG1 systems (Socki et al., 2002; Pérez-Ceballos et al., 2012). The presence of sulfates in lake waters may be attributed to the K/T anhydrite/gypsum-bearing impact breccia and dissolution of CaSO_4 of evaporites (Rosencrantz, 1990). This suggests that lakes with high sulfate contents receive high ground water input, as evaporites are only present at depths greater than 170 m below surface, or that these lakes developed along sites with past tectonic activity such as faulting and uplift (Perry et al., 2002). Some lakes from the YG1 with high content of sulfates did, in fact, develop along fault zones. For example, Lake Chichancanab is associated with the Sierrita de Ticul fault (Hodell et al., 2005) and Lachuá to the Polochic fault (Erdlac and Anderson, 1982), whereas other lakes are related to high ground water input, such as the Bacalar hydrological system (Perry et al., 2009). High TOC values in sediments of most lakes of YG2 probably reflect the trophic state of lake waters. Our data confirm results by Pérez et al. (2011a), who recorded a TOC increase from north to south on the Yucatán Peninsula. This may be attributed to

505 the combined effect of soil, precipitation and vegetation type, which changes from north to south. The northern part of the Yucatán Peninsula is characterized by leptosols, which are shallow soils with high amounts of exposed hard rock and calcareous material (Bautista et al., 2011; Estrada-Medina et al., 2013). There, precipitation of about 450 mm yr⁻¹ (Pérez et al., 2011a), gives rise to low-stature deciduous forests with low biological productivity. In the south of the peninsula, where precipitation increases to >3200 mm yr⁻¹ (Pérez et al., 2011a), luvisols and vertisols, clayey and fertile soils, support the growth
510 of tropical evergreen forests that generate high amounts of organic matter runoff, particularly during the rainy season.

[In](#) Central American volcanism [is the most common mode of lake formation](#). These lakes are classified as caldera lakes, crater lakes in (partially) active or inactive volcanoes, maar lakes, or are located in volcanic bedrock basins (Golombek and Carr, 1978; Newhall and Dzurisin, 1988; Dull et al., 2001; Vallance and Calvert, 2003) (Table 1). The existence of at least three
515 limnological subregions highlights that these lakes are additionally influenced by regional factors related to orography (elevation), climate and the level of volcanic activity, including magmatic heat and gas input. Ionic dominance of Central American lakes is highly variable, but anions Cl⁻, SO₄²⁻, and cations K⁺, Mg²⁺, Na⁺ were dominant (Fig. 4a). This ionic composition reveals that two main processes control water chemistry: 1) active volcanic activity with strong interaction with lakes, and 2) precipitation-evaporation rates, especially at high-elevation sites. Dominance of magnesium, chloride and
520 sulfates, such as in GSHN3, can be attributed to volcanic activity and hydrothermal systems. Chloride and sulfates are strongly influenced by volcanic gas input, by incorporation of HCl and SO₂, and interaction with igneous rocks. Lakes with such ionic dominance also display high contents of phyllosilicates and feldspars, which may have formed through dissolution by hot and likely acid groundwater, heated by hydrothermal activity. Lakes located in the Transmexican Volcanic Belt of central Mexico (TMVB; Armienta et al., 2008; Sigala, et al., 2017) have similar ionic and sediment chemical composition as lakes in Central
525 America. The ionic dominance of the currently extinct crater lakes of Volcano Popocatepetl, for example, changed from sulfate to calcium-magnesium dominance and finally magnesium dominance, resulting from heating of andesite rocks after a period of increased volcanic activity (Armienta et al., 2000, 2008). This suggests that the influence of active volcanism on lake water chemistry may exert similar influences along the northern Neotropical region and American transition zone. High rates of evaporation are anticipated to represent an important driver for water chemistry in Central America as well, because of
530 dominance of carbonates, bicarbonates and sodium (GSHN1 subregion). This ionic composition can be attributed to interaction and weathering of volcanic rocks, capture and dissolution of CO₂, and high evaporation rates. Although Central America can be considered a tropical and humid region, high temperatures and solar radiation, especially at high elevations, may produce high evaporation rates leading to ion- specific signatures such as dominance of carbonates-bicarbonates.

Given the origin of discriminating variables in PCA for both YG and GSHN, we found three main sources controlling
535 limnology in the northern Neotropics: (i) bedrock type, which determines specific mineral and ionic composition of lake sediments and host waters; (ii) volcanic and marine influence, which determines the presence of dominant and conservative ions such as Mg²⁺ and Cl⁻, and (iii) precipitation-evaporation balance across altitudinal and latitudinal gradients, which determines the concentration of solutes and therefore conductivity. This is consistent with results from limnological studies in

540 other regions of Central America and southern Mexico, which suggested that geology, through bedrock types and volcanic
input to lakes, and marine interactions, are the most influential factors for aquatic environments (Löffler, 1972; Haberyan and
Horn, 1999; Haberyan et al., 2003; Cervantes-Martínez et al., 2002; Perry et al., 2002; Schmitter-Soto et al., 2002a; Socki et
al., 2002; Pérez et al., 2011a). Several authors, however, consider additional features relevant for lake classification, such as
545 temperature, pH and altitude (orography) (Brezonik and Fox, 1974; Horn and Haberyan, 1993; Umaña et al., 1999; Haberyan
et al., 2003). In our study, water temperature, pH and elevation scored relatively low on the PCA, suggesting that their influence
is relevant only on a local scale. Considering the determinants of limnological variability of aquatic systems at both regional
and local scales, the interaction with marine environments and the high variability of aquatic system morphology and origin,
it is evident that the current geological configuration is the main factor driving aquatic ecosystem properties and defining
limnological regions in the northern Neotropics.

550 **4.2 Geodiversity as a determinant of distribution of freshwater ostracodes in the northern Neotropics**

We found ostracode species in almost all the aquatic systems we studied. Seventy species were recognized, demonstrating that
this group is abundant and diverse in the northern Neotropics. The number of species per lake (species richness), however,
was relatively low. In most sites we found between two and six species, and in large lakes, such as Petén Itzá and Nicaragua,
we found a maximum of nine species. Diversity metrics substantiated this tendency, with values of the Shannon Index always
555 < 2, which reflects low diversity in aquatic environments (Margalef, 1957; Chao and Shen, 2003). Low species richness was
also observed in previous studies in the region. In Lake Petén Itzá, for example, a maximum of 11 species was reported by
Pérez et al., (2010a). In Lake Nicaragua, the largest lake in Central America, seven species were found (Hartmann, 1959). In
contrast, in regions such as the Palearctic, a single lake may host 32 species, such as ancient Lake Ohrid (Macedonia/Albania),
although there, the Shannon Diversity Index revealed low diversity ($H=1.2$) (Lorenschat et al., 2014), whereas small lakes in
560 Europe, such as Lake Maarsseveen may harbor 9 to 15 species (Sluys, 1981). In Japan, 40 species were recorded from Lake
Biwa (Karanovic, 2015). This global species richness pattern contrasts with the Latitudinal Diversity Gradient theory, that
postulates an increase in species richness from the poles to the Equator (Mannion et al., 2014; Schumm et al., 2019).
Nevertheless, a similar pattern, i.e. a decline in diversity at lower latitudes is observed for other aquatic microorganisms in the
northern Neotropics, such as testate amoebae (Sigala et al., 2016; Sigala Regalado et al., 2018; Charqueño-Celis et al., 2019),
565 chironomids (Hamerlík et al., 2018), and cladocerans (Vázquez-Molina et al., 2016; Wojewódka et al., 2016). Low species
richness in lakes of the northern Neotropics may be related to local environmental and evolutionary processes that occurred in
recent geologic time, namely during the Pleistocene and Holocene. For instance, in southern Mexico and northern Guatemala,
severe droughts were inferred from study of sediment cores from Lake Petén Itzá (Mueller et al., 2010; Cohuo et al., 2018).
Some of the droughts, such as those that occurred during Heinrich Stadials, probably caused most shallow lakes to desiccate
570 (Cohuo et al., 2018). Repetitive droughts in the Pleistocene, with their consequent effect on surficial freshwater availability,
must have impacted the biological evolutionary processes of ostracodes, which occupy benthic habitats, thus limiting species
diversification and adaptation. An alternative explanation for the low ostracode species richness in the northern Neotropics is

575 related to intensification of human activities in lakes and their watersheds. In Central America, water extraction, fisheries and fish farms, eutrophication and wastewater disposal, are among the most frequent alterations in freshwater ecosystem (Cuadra et al., 2006; Arcega-Cabrera et al., 2014; Campuzano et al., 2014). These conditions are known to strongly affect aquatic taxa, and particularly endemic species, sometimes causing loss of biodiversity (Moyle and Leidy, 1992). In the Neotropics, loss of both genetic and phenotypic biodiversity of aquatic taxa has been poorly evaluated (Johann et al., 2019), but is certainly an important issue that must be considered for lake protection and management (Mercado-Salas et al., 2013).

580 By comparison, in the flood plains of the Upper Parana River, South America, species richness can be as high as 44 (Higuti et al., 2017), which is higher than in lakes of Central America, and comparable to richness of lakes in temperate regions. In Colombian aquatic systems, however, such as La Fé reservoir (Saldarriaga and Martínez, 2010) and the Magdalena River basin (Roessler, 1990a, b), the number of ostracode species is similar to that in Central America (~6 species per lake). Therefore, to clarify structural patterns of ostracodes in the Neotropical region, more intensive sampling in lakes and rivers is needed. Evidence from other tropical regions around the world will be valuable to understand patterns of ostracode diversity (tropical vs temperate) and how latitudinal diversity gradients act in the Ostracoda group.

590 The NMDS analysis shows the existence of at least five species associations in the region (OST1-OST5), emphasizing that ostracodes do not conform to a faunal unit, but rather display disjunct faunas (Fig. 5a), similar to what is observed in freshwater fishes of the region (Miller, 1976; Matamoros et al., 2015). Ostracode associations are geographically delimited, and no overlap was observed (Fig. 5b). Ostracode groups OST1 and OST2, belong to the YG limnological region, whereas the OST3, OST4 and OST5 associations belong to the GHSN limnological region (Figs. 5a, b). Few species were present in more than three limnological subregions, and these can be considered of wide Neotropical distribution, e.g., *C. ilosvayi* and *C. vidua*.

600 Correspondence between species associations and limnological subregions suggests a major influence of physical and chemical properties of lake environments on biological systems. The structural equation modelling (SEM) analysis exposed the significant influence of geodiversity on limnology, and of limnology on species associations, identified in the NMDS. This illustrates that limnology, particularly geochemistry, mineralogy, and water chemistry, is the primary factor responsible for species distributions in the study area. We were, however, unable to find statistical significance to explain the relationships between geodiversity and limnology with ostracode species richness. This suggest that the number of species per lake may not be fully governed by our predictors. For instance, conductivity and elevation in model 4 and TOC and latitude in model 5, also failed to explain species richness. This reveal that the main environmental gradients in the northern Neotropics, and the theorized increase in the number of species at lower latitudes (Latitudinal Diversity Gradient theory), do not apply in the northern Neotropics and, therefore, other intrinsic or extrinsic factors such historical water level fluctuations and precipitation/evaporation balance, might instead control species richness. In our SEM models, we also tested the direct influence of geodiversity and its indirect influence through limnology on species distribution and richness. The optimal SEM model demonstrated the significance of paths describing the direct influence of geodiversity on species richness and composition, but significance of standardized coefficients was of [minor importance](#) (0.1 and 0.04, respectively). Conversely, strong ties were discovered when we analyzed for the indirect effect of geodiversity (via limnology) on species composition

(>2.0). This implies that ostracode species composition is more predictable from limnological variables such as water physical and chemical and geochemistry than from geodiversity variables such as bedrock and mineralogy. Our SEM model also revealed that elevation is an important predictor of species composition (Fig. 6). The negative correlation suggested that an increase in elevation causes a decrease in the number of species per site. Most highland lakes were characterized by up to three species and, more commonly, by a single species, except for large lakes Amatitlán and Atitlán in Guatemala.

The CCA further demonstrates that geological components of the landscape such as bedrock, elevation (temperature), mineralogy and ionic composition are important for species distribution (Fig. S3). For instance, the OST1 group is represented by lakes of the YG3 subregion, associated with the Caribbean Sea coast (Fig. 5b). Similarly, the OST 2 group corresponds to lakes of the YG1 and YG2 subregions. These lakes are characterized by high TOC and abundant submerged vegetation. In Central America, patterns of coincidence between ostracodes and lake regions are evident, as well (Fig. 5b). The OST 4 and OST 5 groups are associated with volcanic-influenced lakes in the lowlands (GSHN1) and highlands (GSHN3), respectively. The CCA and SEM analyses show that the influence of geodiversity over species composition and distribution is only relevant when limnological attributes are taken into account. Future biogeographic studies should focus on combining different zooplankton and zoobenthos groups and explore the role of geodiversity in driving patterns of composition and distribution.

5 Conclusions

The northern Neotropics is a region characterized by [diverse environmental conditions](#), abundant aquatic systems, and high biodiversity. Our limnological survey of 76 aquatic environments identified two main limnological regions in the northern Neotropics. The YG region is associated with karst plateaus in southern Mexico and northern Guatemala, whereas the GSHN is associated with landscapes formed by volcanic activity in southern Guatemala, El Salvador, Honduras and Nicaragua. At least seven limnological subregions were identified, illustrating the high heterogeneity of aquatic systems in the northern Neotropics. Analysis of ostracode species assemblages shows that water chemistry and mineralogical attributes of the host environments drives species composition (distribution). Low ostracode species richness in the northern Neotropics also seems to be strongly related to the geological history of the region. The low number of species per lake contrasts with the number of species per lake in temperate regions, which is at least five times higher. Questions about the applicability of the Latitudinal Diversity Gradient theory for Ostracoda and other zooplankton and zoobenthos groups remain unresolved, and sampling in the Neotropics and other tropical regions of the world is necessary. SEM analysis highlights that geodiversity has a direct influence on limnological regions, and an indirect but relevant influence on freshwater ostracodes. This is the first study to integrate data on watershed geology, physical and chemical characteristics of the water column, sediment mineralogy and geochemistry, and biota in aquatic ecosystems of southern Mexico and Central America. Further studies should focus on establishment of a more detailed regionalization, by including a greater number of lakes, more environmental variables, and samples collected at different times throughout the seasonal cycle.

640 **Data availability.** Water physico-chemical, sediment geochemistry, mineralogical and geological data from the 76 aquatic ecosystems sampled in this study are available in Table S1 in the Supplementary Material and in the Pangaea repository: <https://doi.org/10.1594/PANGAEA.940538>. Ostracode relative abundances at each sampling site can be found at the Pangaea repository: <https://doi.org/10.1594/PANGAEA.940254>.

645 **Financial support.** Funding was provided by the Deutsche Forschungsgemeinschaft (DFG, SCHW 671/16-1), CONACYT (Mexico; project number 319857) and Tecnológico Nacional de México (projects number 14502.21-P and 14698.22-P).

650 **Author contributions.** LM and SC conducted the fieldwork, performed data analysis, and wrote the manuscript. AS, LP and ME developed, managed and coordinated the project and contributed to data interpretation and manuscript writing. PH contributed and interpreted mineralogical data and gave scientific input to the manuscript. MC developed water chemistry analyses and gave scientific input to the manuscript. AO, MP and MA provided support for sampling in their respective countries and organized sampling permits.

Competing interests. The authors declare that they have no conflict of interest.

655 **Acknowledgments.** We thank all colleagues and institutions involved in this work, including the student team: Christian Vera, León E. Ibarra, Miguel A. Valadéz and Cuauhtémoc Ruiz (Instituto Tecnológico de Chetumal, Mexico); Ramón Beltran (Centro Interdisciplinario de Ciencias Marinas, Mexico) and Lisa Heise (Universidad Autónoma de San Luis Potosí, Mexico) for their excellent contributions during field work. We also thank the following colleagues and institutions that made field sampling and water chemistry analysis possible: The team from the Asociación de Municipios del Lago de Yojoa y su área de influencia (Amuprolago, Honduras); Margaret Dix, Eleonor de Tott, Roberto Moreno (Universidad del Valle de Guatemala, Guatemala); Consejo Nacional de Áreas Protegidas (CONAP, Guatemala); Néstor Herrera (Ministerio de Medio Ambiente, San Salvador), Teresa Álvarez (El Colegio de la Frontera sur, Chetumal Unit, Mexico); María Aurora Armienta (Laboratorio de Química Analítica, Instituto de Geofísica, Universidad Nacional Autónoma de México) and Adriana Zavala (El Colegio de la Frontera Sur, Mexico). We also gratefully acknowledge anonymous reviewers and Mark Brenner for their constructive feedback and language editing.

References

665 Albert, J. S., and Reis, R. E. (Eds.): Historical Biogeography of Neotropical Freshwater Fishes, University of California Press, Berkeley, CA, 2011.

Alcocer, J. and Bernal-Brooks, F. W.: Limnology in Mexico, *Hydrobiologia*, 644, 15–68, <https://doi.org/10.1007/s10750-010-0211-1>, 2010.

- Alcocer, J., Lugo, A., Marín, L., and Escobar, E.: Hydrochemistry of waters from five cenotes and evaluation of their suitability for drinking-water supplies, northeastern Yucatan, Mexico, *Hydrogeol. J.*, 6, 293–301, <https://doi.org/10.1007/s100400050152>, 1998.
- Antonelli, A., Zizka, A., Carvalho, F. A., Scharn, R., Bacon, C.D., Silvestro, D., and Condamine, F. L.: Amazonia is the primary source of Neotropical biodiversity, *Proc. Natl. Acad. Sci. U.S.A.*, 115, 6034–6039, <https://doi.org/10.1073/pnas.1713819115>, 2018.
- Arcega-Cabrera, F., Velázquez-Tavera, N., Fargher, L., Derrien, M., and Noreña-Barroso, E.: Fecal sterols, seasonal variability, and probable sources along the ring of cenotes, Yucatan, Mexico, *J. Contam. Hydrol.*, 168, 41–49, <https://doi.org/10.1016/j.jconhyd.2014.08.007>, 2014.
- Armienta, M. A., De la Cruz-Reyna, S., and Macías, J. L.: Chemical characteristics of the crater lakes of Popocatepetl, El Chichon, and Nevado de Toluca volcanoes, Mexico, *J. Volcanol. Geotherm. Res.*, 97, 105–125, [https://doi.org/10.1016/S0377-0273\(99\)00157-2](https://doi.org/10.1016/S0377-0273(99)00157-2), 2000.
- Armienta, M., Vilaclara, G., De la Cruz-Reyna, S., Ramos, S., Ceniceros, N., Cruz, O., Aguayo, A., and Arcega-Cabrera, F.: Water chemistry of lakes related to active and inactive Mexican volcanoes. *J. Volcanol. Geotherm. Res.*, 178, 249–258, <https://doi.org/10.1016/j.jvolgeores.2008.06.019>, 2008.
- Azim, M. E.: Photosynthetic Periphyton and Surfaces, in: *Encyclopedia of Inland Waters*, edited by Gene, E. L., Academic Press, 184–191, 2009.
- Bacon, C. D., Silvestro, D., Jaramillo, C., Smith, B.T., Chakrabarty, P., and Antonelli, A.: Biological evidence supports an early and complex emergence of the Isthmus of Panama. *Proc. Natl. Acad. Sci. U.S.A.*, 112, 6110–6115, <https://doi.org/10.1073/pnas.1423853112>, 2015.
- Bagley, J., and Johnson, J.: Phylogeography and biogeography of the lower Central American Neotropics: Diversification between two continents and between two seas. *Biol. Rev.*, 89, 767–790, <https://doi.org/10.1111/brv.12076>, 2014.
- Baker P. A., Fritz S. C., Dick C. W., Battisti D. S., Vargas O. M., Asner G. P., Martin R. E., Wheatley A., and Prates, I.: Beyond Refugia: New Insights on Quaternary Climate Variation and the Evolution of Biotic Diversity in Tropical South America, in: *Neotropical Diversification: Patterns and Processes*, Fascinating Life Sciences, edited by: Rull V., and Carnaval, A., Springer, Cham, 51–70, https://doi.org/10.1007/978-3-030-31167-4_3, 2020.

- 695 Baker, P. A., Seltzer, G. O., Fritz, S. C., Dunbar, R. B., Grove, M., Tapia, P., Cross, S., Rowe, H., and Broda, J.: The history of South American tropical climate for the past 25,000 years, *Science*, 291, 640–643, <https://doi.org/10.1126/science.291.5504.640>, 2001.
- Bautista F., Palacio, G., Quintana, P., and Zinck, A.J.: Spatial distribution and development of soils in tropical karst areas from the Peninsula of Yucatán, Mexico, *Geomorphology*, 135, 308–321, <https://doi.org/10.1016/j.geomorph.2011.02.014>, 2011.
- 700 Behling, H., Arz, H. W., Patzold, J., and Wefer, G.: Late Quaternary vegetational and climate dynamics in northeastern Brazil, inferences from marine core GeoB 3104-1, *Quat. Sci. Rev.*, 19, 981–994, [https://doi.org/10.1016/S0277-3791\(99\)00046-3](https://doi.org/10.1016/S0277-3791(99)00046-3), 2000.
- Bivand, R. S., Pebesma, E. J., Gomez-Rubio, V., and Pebesma, E. J. (Eds.): *Applied spatial data analysis with R*, Springer, New York, 2008.
- 705 Bouimetarhan, I., Chiessi, C., Gonzalez-Arango, C. G., Dupont, L., Voigt, I., Prange, M., and Zonneveld, K.: Intermittent development of forest corridors in northeastern Brazil during the last deglaciation: climatic and ecologic evidence, *Quat. Sci. Rev.*, 192, 86–96, <https://doi.org/10.1016/j.quascirev.2018.05.026>, 2018.
- Bravo-Cuevas, V., González-Rodríguez, K., Cabral-Perdomo, M., Cuevas-Cardona, C., and Pulido-Silva, M.: Geodiversity and its implications in the conservation of biodiversity: Some case studies in central Mexico, *CIENCIA Ergo-Sum*, 28(3), 1–15, <https://doi.org/10.30878/ces.v28n3a8>, 2021.
- 710 Brezonik, P., and Fox, J.: The limnology of selected Guatemalan lakes, *Hydrobiologia*, 45, 467–487, <https://doi.org/10.1007/BF00012032>, 1974.
- Bryson, R., Riddle, B., Graham, M., Smith, B., and Prendini, L.: As Old as the hills: montane scorpions in Southwestern North America reveal ancient associations between biotic diversification and landscape history, *PLoSOne*, 8:e52822, <https://doi.org/10.1371/journal.pone.0052822>, 2013.
- 715 Cabassi, J., Capecchiacci, F., Magi, F., Vaselli, O., Tassi, F., Montalvo, F., Esquivel, I., Grassa, F., and Caprai, A.: Water and dissolved gas geochemistry at Coatepeque, Ilopango and Chanmico volcanic lakes (El Salvador, Central America), *J. Volcanol. Geotherm. Res.*, 378, 1–15, <https://doi.org/10.1016/j.jvolgeores.2019.04.009>, 2019.
- Campuzano, C., Hansen, A. M., De Stefano, L., Martínez-Santos, P., Torrente, D., and Willaarts, B. A.: Water resources assessment, in: *Water for Food and Wellbeing in Latin America and the Caribbean. Social and Environmental Implications*

for a Globalized Economy, edited by: Willaarts, B. A., Garrido, A., Llamas, M. R., Routledge, Oxon and New York, 27–53, 2014.

Carnaval, A. C., and Moritz, C.: Historical climate modeling predicts patterns of current biodiversity in the Brazilian Atlantic forest, *J. Biogeogr.*, 35, 1187–1201, <https://doi.org/10.1111/j.1365-2699.2007.01870.x>, 2008.

725 Carr, M.: Symmetrical and segmented variation of physical and geochemical characteristics of the Central American volcanic front, *J. Volcanol. Geoth. Res.*, 20, 231–252, [https://doi.org/10.1016/0377-0273\(84\)90041-6](https://doi.org/10.1016/0377-0273(84)90041-6), 1984.

Cervantes-Martínez, A., Elías-Gutiérrez, M., and Suárez-Morales, E.: Limnological and morphometrical data of eight karstic systems 'cenotes' of the Yucatán Peninsula, Mexico, during the dry season (February–May, 2001), *Hydrobiologia*, 482, 167–177, <https://doi.org/10.1023/A:1021260131757>, 2002.

730 Chao, A., Shen, T. J.: Nonparametric estimation of Shannon's index of diversity when there are unseen species in sample. *Environ. Ecol. Stat.*, 10, 429–443, <https://doi.org/10.1023/A:1026096204727>, 2003.

Charqueño-Celis, N. F., Garibay, M., Sigala, I., Brenner, M., Echeverría Galindo, P., Lozano García, S., Massaferró, J., and Pérez, L.: Testate amoebae (Amoebozoa: Arcellinidae) as indicators of dissolved oxygen concentration and water depth in lakes of the Lacandón Forest, Southern Mexico, *J. Limnol.*, 79, 82–91, <https://doi.org/10.4081/jlimnol.2019.1936>, 2019.

735 Chavent, M., Kuentz-Simonet, V., Labenne, A., and Saracco, J.: Multivariate analysis of mixed data: The PCAmixdata R package. arXiv preprint arXiv:1411.4911v1, 2014.

Clarke, K.: Non-parametric multivariate analyses of changes in community structure. *Aust. J. Ecol.*, 18, 117–143, <https://doi.org/10.1111/j.1442-9993.1993.tb00438.x>, 1993.

740 Cohuo, S., Macario-González, L., Pérez, L., and Schwalb, L.: Overview of Neotropical-Caribbean freshwater ostracode fauna (Crustacea, Ostracoda): identifying areas of endemism and assessing biogeographical affinities, *Hydrobiologia*, 786, 5–21, <https://doi.org/10.1007/s10750-016-2747-1>, 2016.

Cohuo, S., Macario-González, L., Pérez, L., Sylvestre, F., Paillès, C., Curtis, J., Kutterolf, S., Wojewódka, M., Zawisza, E., Sze-roczyńska, K., and Schwalb, A.: Ultrastructure and aquatic community response to Heinrich Stadial (HS5a-HS1) in the continental northern Neotropics, *Quat. Sci. Rev.*, 19, 75–91, <https://doi.org/10.1016/j.quascirev.2018.07.015>,

745 2018.

Cohuo, S., Macario-González, L., Wagner, S., Naumann, K., Echeverría-Galindo, P., Pérez, L., Curtis, J., Brenner, M., and Schwalb, A.: Influence of late Quaternary climate on the biogeography of Neotropical aquatic species as reflected by non-marine ostracodes, *Biogeosciences*, 17, 145–161, <https://doi.org/10.5194/bg-17-145-2020>, 2020.

CEPF (Critical Ecosystem Partnership Fund) - The Biodiversity hotspots (2021, November).

750 Cuadra, S. N., Linderholm, L., Athanasiadou, M., and Jakobsson, K.: Persistent organochlorine pollutants in children working at a waste-disposal site and in young females with high fish consumption in Managua, Nicaragua, *Ambio*, 35, 109–116, [https://doi.org/10.1579/0044-7447\(2006\)35\[109:popicw\]2.0.co;2](https://doi.org/10.1579/0044-7447(2006)35[109:popicw]2.0.co;2), 2006.

De Albuquerque, F. S., Benito, B., Beier, P., Assunção-Albuquerque, M. J., and Cayuela, L.: Supporting underrepresented forests in Mesoamerica, *Natureza e Conservacao*, 13(2), 152–158, <https://doi.org/10.1016/j.ncon.2015.02.001>, 2015.

755 DeClerck, F., Chazdon, R., Robin, H., Karen, D., Milder, J., Finegan, B., Martínez-Salinas, A., Imbach, P., Canet, L., and Ramos, Z.: Biodiversity conservation in human-modified landscapes of Mesoamerica: Past, present and future. *Biol. Conserv.*, 143, 2301–2313, <https://doi.org/10.1016/j.biocon.2010.03.026>, 2010.

760 Delgado-Martínez, C. M., Alvarado, F., Mendoza, E., Flores-Hernández, S., Navarrete, A., Navarrete, E., and Botello, F.: An ignored role of sartenejas to mitigate water shortage hazards for tropical forest vertebrates, *Ecology*, 99(3), 758–760, <https://doi.org/10.1002/ecy.2078>, 2018.

Dengo, G., Bohenberger, O., and Bonis, S.: Tectonics and volcanism along the Pacific Marginal Zone of Central America, *Geol. Rundsch.*, 59, 1215–1235, <https://doi.org/10.1007/BF02042290>, 1970.

Dull, R. A., Southon, J. R. and Sheets, P.: Volcanism, Ecology and Culture: A Reassessment of the Volcan Ilopango Tbj eruption in the Southern Maya Realm, *Lat. Am. Antiq.*, 12, 25–44, <https://doi.org/10.2307/971755>, 2001.

765 Echeverría-Galindo P. G., Pérez, L., Correa-Metrio, A., Avendaño, C., Moguel, B., Brenner, M., Cohuo, S., Macario, L., and Schwalb, A.: Tropical freshwater ostracodes as environmental indicators across an altitude gradient in Guatemala and Mexico *Rev. Biol. Trop.*, 67, 1037–1058, <http://dx.doi.org/10.15517/rbt.v67i4.33278>, 2019.

Ellison, A.M.: Wetlands of Central America, *Wetl. Ecol. Manag.*, 12, 3–55, <https://doi.org/10.1023/B:WETL.0000016809.95746.b1>, 2004.

770 Erdlac, T., and Anderson, H.: The Chixoy-Polochic fault and its associated fractures in western Guatemala. *GSA Bull.*, 93, 57–67, [https://doi.org/10.1130/0016-7606\(1982\)93<57:TCFAIA>2.0.CO;2](https://doi.org/10.1130/0016-7606(1982)93<57:TCFAIA>2.0.CO;2), 1982.

- Estrada-Medina, H., Bautista, F., Jiménez-Osornio, J. J., González-Iturbe, J. A., and Aguilar-Cordero, W.: Maya and WRB Soil Classification in Yucatan, Mexico: Differences and Similarities. *Int. Sch. Res. Notices*, 634260, 1–10, <https://doi.org/10.1155/2013/634260>, 2013.
- 775 Etienne, R., and Apol, M.: Estimating speciation and extinction rates from diversity data and the fossil record, *Evolution*, 63, 244–255, <https://doi.org/10.1111/j.1558-5646.2008.00537.x>, 2009.
- Fabrigar L. R., MacCallum R. C., Wegener D. T., and Strahan E. J.: Evaluating the use of exploratory factor analysis in psychological research, *Psychol. Methods*, 4, 272–299, <https://doi.org/10.1037/1082-989X.4.3.272>, 1999.
- Fan, Y., Chen, J., Shirkey, G., John, R., Wu, S., Park, H., and Shao, C.: Applications of structural equation modeling (SEM) in ecological studies: an updated review. *Ecol. Process.*, 5, 19, <https://doi.org/10.1186/s13717-016-0063-3>, 2016.
- 780 Franco-Gaviria, F., Caballero-Rodríguez, D., Correa-Metrio, A., Pérez, L., Schwalb, A., Cohuo, S., and Macario-González, L.: The human impact imprint on modern pollen spectra of the Maya lands, *Bol. Soc. Geol. Mex.*, 70, 61–78, 2018.
- Garrity, C., and Soller, C.: Database of the Geologic Map of North America; adapted from the map by: Reed Jr., J. C. et al. (2005), US Geological Survey Data Series 424, 2009.
- 785 Gillespie, R., and Roderick, G.: Geology and climate drive diversification, *Nature*, 509, 297–298, <https://doi.org/10.1038/509297a>, 2014.
- Golombek, M. P., and Carr, M. J.: Tidal triggering of seismic and volcanic phenomena during the 1879–1880 eruption of Islas Quemadas volcano in El Salvador, Central America, *J. Volcanol. Geoth. Res.*, 3, 299–307, [https://doi.org/10.1016/0377-0273\(78\)90040-9](https://doi.org/10.1016/0377-0273(78)90040-9), 1978.
- 790 Gray, M.: Geodiversity, geoheritage and geoconservation for society. *International Journal of Geoheritage and Parks*, 7, 226–236, <https://doi.org/10.1016/j.ijgeop.2019.11.001>, 2019.
- [Gray, M. \(Ed.\): Geodiversity: Valuing and Conserving Abiotic Nature, John Wiley and Sons Ltd., Chichester, 2004.](#)
- Haberyan, K., and Horn, S.: Chemical and Physical Characteristics of Seven Lakes in Costa Rica, *Brenesia*, 51, 85–95, 1999.
- Haberyan, K., Horn, S., and Umaña, G.: Basic limnology of fifty-one lakes in Costa Rica, *Rev. Biol. Trop.*, 51, 107–122, 2003.
- 795 Haffer, J.: Speciation in Amazonian forest birds, *Science*, 165, 131–137, <https://doi.org/10.1126/science.165.3889.131>, 1969.

- Hamerlík, L., Wojewodka, M., Zawisza, E., Cohuo Duran, S., Macario-Gonzalez, L., Pérez, L., and Szeroczynska, K.: Subfossil Chironomidae (Diptera) in surface sediments of the sinkholes (cenotes) of the Yucatan Peninsula: Diversity and distribution, *J. Limnol.*, 77 (s1), 213–219, <https://doi.org/10.4081/jlimnol.2018.1769>, 2018.
- 800 Hammer, Ø., Harper, D.A.T., and Ryan, P.D.: PAST: Paleontological Statistics Software Package for Education and Data Analysis. *Palaeontologia Electronica*, 4(1), 1–9, 2001.
- Hartmann, G.: Beitrag zur Kenntnis des Nicaragua-Sees unter besonderer Berücksichtigung seiner Ostracoden (mit Beschreibung von 5 neuen Arten), *Zool. Anz.*, 162, 269–294, 1959.
- Hastenrath, S.: Rainfall distribution and regime in Central America, *Arch. Met. Geoph. Biokl. B.*, 15, 201–241, <https://doi.org/10.1007/BF02243853>, 1967.
- 805 Higuti, J., Conceição, E. O., Campos, R., Ferreira, V. G., Rosa, J., Pinto, M. B. O., and Martens, K.: Periphytic community structure of Ostracoda (Crustacea) in the river-floodplain system of the Upper Paraná River. *Acta Limnol. Bras.*, 29(e.120), <https://doi.org/10.1590/S2179-975X12217>, 2017.
- Hildebrand, A. R., Pilkington, M., Connors, M., Ortiz-Aleman, C., and Chavez, R. E.: Size and structure of the Chicxulub crater revealed by horizontal gravity gradients and cenotes, *Nature*, 376, 415–417, <https://doi.org/10.1038/376415a0>, 1995.
- 810 Hill, M. O., and Gauch, H. G.: Detrended correspondence analysis: An improved ordination technique, *Vegetatio*, 42, 47–58, <https://doi.org/10.1007/BF00048870>, 1980.
- Hodell, D. A., Brenner, M., and Curtis, J.: Terminal Classic drought in the northern Maya lowlands inferred from multiple sediment cores in Lake Chichancanab (Mexico), *Quat. Sci. Rev.*, 24, 1413–1427, <https://doi.org/10.1016/j.quascirev.2004.10.013>, 2005.
- 815 Horn, S., and Haberyan, K.: Costa Rican lakes: physical and chemical properties, *Nat. Geog. Res. Explor.*, 9, 86–103, 1993.
- Hu, A., Wang, J., Sun, H., Niu, B., Si, G., Wang, J., Yeh, C., Zhu, X., Lu, X., Zhou, J., Yang, Y., Ren, M., Hu, Y., Dong, H. D., and Zhang, G.: Mountain biodiversity and ecosystem functions: interplay between geology and contemporary environments, *ISME J.*, 14, 931–944, <https://doi.org/10.1038/s41396-019-0574-x>, 2020.
- Hu, L., and Bentler, P. M.: Cutoff criteria for fit indexes in covariance structure analysis: Conventional criteria versus new 820 alternatives. *Struct. Equ. Model.*, 6, 1–55, <https://doi.org/10.1080/10705519909540118>, 1999.

- Hulsey, C. D., Hollingsworth, P. R., and Fordyce, J. A.: Temporal diversification of Central American cichlids. *BMC Evol. Biol.*, 10, 279, <https://doi.org/10.1186/1471-2148-10-279>, 2010.
- Huston, M. A.: *Biological Diversity: the coexistence of species on changing landscapes*, xix, 681, Cambridge University Press, 1995.
- 825 Jakobsen, J. C., Gluud, C., Wetterslev, J., and Winkel, P.: When and how should multiple imputation be used for handling missing data in randomised clinical trials - a practical guide with flowcharts, *BMC Med. Res. Methodol.* 17 (1), 162, <https://doi.org/10.1186/s12874-017-0442-1>, 2017.
- Jiménez-Alfaro, B., Girardello, M., Chytry, M., Svenning, J. C., Willner, W., Gegou, J. C., Agrillo, E., Campos, J. A., Jandt, U., Kački, Z., Šilc, U., Slezák, M., Tichý, L., Tsiripidis, L., Turtureanu, P. D., Ujházyová, M., and Wohlgemuth, T.: History and environment shape species pools and community diversity in European beech forests, *Nat. Ecol. Evol.*, 2, 483–490, <https://doi.org/10.1038/s41559-017-0462-6>, 2018.
- Johann, A., Mangolin, L., Sanches, P., Sebastián, N., Topan, D., Piana, P., and Gomes, B.: Urbanized Tributary Causes Loss of Biodiversity in a Neotropical River Segment. *Water Air Soil Poll.*, 230, 118, <https://doi.org/10.1007/s11270-019-4164-3>, 2019.
- 835 Jury, M.: Long-term variability and trends in the Caribbean Sea. *Int. J. Oceanogr.*, ID 465810, 1–9, <https://doi.org/10.1155/2011/465810>, 2011.
- Karanovic, I. (Ed.): *Recent Freshwater Ostracods of the World, Crustacea, Ostracoda, Podocopida*, Springer, Berlin, 2012.
- Karanovic, I.: Barcoding of ancient lake ostracods (Crustacea) reveals cryptic speciation with extremely low distances, *PLoS ONE*, 10, e0121133, <https://doi.org/10.1371/journal.pone.0121133>, 2015.
- 840 Legendre, P., and Legendre, L. (Eds.): *Numerical Ecology*, Elsevier Scientific, Oxford, 1998.
- Leite, Y. L., Costa, L. P., Loss, A. C., Rocha, R. G., Batalha-Filho, H., Bastos, A. C., Quaresma, V. S., Fagundes, V., Paresque, R., Passamani, M., and Pardini, R.: Neotropical forest expansion during the last glacial period challenges refuge hypothesis, *Proc. Natl. Acad. Sci. U.S.A.*, 113, 1008–1013, <https://doi.org/10.1073/pnas.1513062113>, 2016.
- Löffler, H.: Contribution to the limnology of high mountain lakes in Central America, *Int. Rev. Hydrobiol.*, 57, 397–408, <https://doi.org/10.1002/iroh.19720570304>, 1972.
- 845

- Lorenschat, J., Pérez, L., Correa-Metrio, A., Brenner, M., Von Bramann, U., and Schwalb, A.: Diversity and spatial distribution of extant freshwater ostracodes (Crustacea) in ancient Lake Ohrid (Macedonia/Albania), *Diversity*, 6, 524–550, <https://doi.org/10.3390/d6030524>, 2014.
- 850 Macario-González, L., Cohuo, S., Elías, M., Vences, M., Perez, L., and Schwalb, A.: Integrative taxonomy of freshwater ostracodes (Crustacea: Ostracoda) of the Yucatán Peninsula, implications for paleoenvironmental reconstructions in the northern Neotropical region. *Zool. Anz.*, 275, 20–36, <https://doi.org/10.1016/j.jcz.2018.04.002>, 2018.
- Magaña, V., Amador J., and Medina, S.: The midsummer drought over Mexico and Central America, *J. Climate*, 12, 1577–1588, [https://doi.org/10.1175/1520-0442\(1999\)012<1577:TMDOMA>2.0.CO;2](https://doi.org/10.1175/1520-0442(1999)012<1577:TMDOMA>2.0.CO;2), 1999.
- 855 Mannion, P.D., Upchurch, P., Benson, R.B., Goswami, A.: The latitudinal biodiversity gradient through deep time, *Trends Ecol. Evol.*, 29, 42–50, <https://doi.org/10.1016/j.tree.2013.09.012>, 2014.
- Margalef, R.: La teoría de la información en Ecología. *Mems R. Acad. Cienc. Artes de Barcelona*, 32, 373–436, 1957.
- Marshall, J. S., Idleman, B. D., Gardner, T. W., and Fisher, D. M.: Landscape evolution within a retreating volcanic arc, Costa Rica, Central America, *Geology*, 31, 419–422, [https://doi.org/10.1130/0091-7613\(2003\)031<0419:LEWARV>2.0.CO;2](https://doi.org/10.1130/0091-7613(2003)031<0419:LEWARV>2.0.CO;2), 2003.
- 860 Marshall, J.: Geomorphology and physiographic provinces of Central America. Central America, in: *Central America: Geology, Resources and Hazards*, edited by: Bundschuh, J., and Alvarado, G., Taylor and Francis Group, London, 75–122, 2007.
- 865 Martiny, J. B., Bohannon, B. J., Brown, J. H., Colwell, R. K., Fuhrman, J. A., Green, J. L., Horner-Devine, M. C., Kane, M., Krumins, J. A., Kuske, C. R., Morin, P. J., Naeem, S., Ovreås, L., Reysenbach, A. L., Smith, V. H., and Staley, J. T.: Microbial biogeography: putting microorganisms on the map, *Nat. Rev. Microbiol.*, 4, 102–112, <https://doi.org/10.1038/nrmicro1341>, 2006.
- Matamoros, W. A., McMahan, C. D., Chakrabarty, P., Albert, J. S., and Schaefer, J. F.: Derivation of the freshwater fish fauna of Central America revisited: Myers's hypothesis in the twenty-first century, *Cladistics*, 31, 177–188, <https://doi.org/10.1111/cla.12081>, 2015.

- 870 Matzke-Karasz, R., Serrano-Sánchez, M. d. L., Pérez, L., Keyser, D., Pipík, R., and Vega, F. J.: Abundant assemblage of Ostracoda (Crustacea) in Mexican Miocene amber sheds light on the evolution of the brackish-water tribe Thalassocypridini, *Hist. Biol.*, 31, 65–101, <https://doi.org/10.1080/08912963.2017.1340471>, 2019.
- McCullough, I., Fergus, E., Scott, C.: *Macrosystems Limnology and Beyond: Re-Envisioning the Scale of Limnology*, in: *Encyclopedia of Inland Waters*, edited by: Cheruvilil, K., Elsevier, U.S.A., 2021.
- 875 McKaye, K. R., Ryan, J. D., Stauffer, J. R., Lopez-Perez, J. L., Vega G. I., van den Berghe. E. P.: African Tilapia in Lake Nicaragua. *BioScience*, 45, 406-411, <https://doi.org/10.2307/1312721>, 1995.
- Mercado-Salas, N. F., Morales-Vela, B., Suárez-Morales, E., and Iliffe, T. M.: Conservation status of the inland aquatic crustaceans in the Yucatan Peninsula, Mexico: shortcomings of a protection strategy, *Aquat. Conserv.*, 23, 939–951, <https://doi.org/10.1002/aqc.2350>, 2013.
- 880 Miller, R. R.: Geographical distribution of Central American freshwater fishes, *Copeia*, 773– 802, <https://doi.org/10.2307/1441406>, 1966.
- Mittelbach, G. G., Schemske, D. W., Cornell, H. V., Allen, A. P., Brown, J. M., Bush, M. B., Harrison, S. P., Hurlbert, S. P., Knowlton, N., Lessios, H. A., McCain, C. M., McCune, A.R., McDade, L. A., McPeck, M. A., Near, T. I., Price, T. D., Ricklefs, R. E., Roy, K., Sax, D. F., Schluter, D., Sobel, A. M., and Turelli, M.: Evolution and the latitudinal diversity gradient: speciation, extinction and biogeography, *Ecol. Lett.*, 10, 315–331, <https://doi.org/10.1111/j.1461-0248.2007.01020.x>, 2007.
- 885 Moguel, B., Pérez, L., Alcaraz, L. D., Blaz, J., Caballero, M., Muñoz-Velasco, I., Becerra, A., Laclette, J. P., Ortega-Guerrero, B., Romero-Oliva, C. S., Herrera-Estrella, L., and Lozano-García, S.: Holocene life and microbiome profiling in ancient tropical Lake Chalco, Mexico, *Sci. Rep.*, 11, 13848, <https://doi.org/10.1038/s41598-021-92981-8>, 2021.
- 890 Molnar, P., and Sykes. L.: Tectonics of the Caribbean and Middle American regions from focal mechanisms and seismicity, *GSA Bulletin*, 80, 1639–1684, [https://doi.org/10.1130/0016-7606\(1969\)80\[1639:TOTCAM\]2.0.CO;2](https://doi.org/10.1130/0016-7606(1969)80[1639:TOTCAM]2.0.CO;2), 1969.
- Montes, C., Cardona, A., Jaramillo, C., Pardo, A., Silva, J.C., Valencia, V., Ayala, C., Pérez-Angel, L.C., Rodríguez-Parra, L.A., Ramírez, V., and Niño, H.: Middle Miocene closure of the Central American Seaway, *Science*, 348, 226–229, <https://doi.org/10.1126/science.aaa2815>, 2015.

- 895 Morrone, J. J.: Biogeographical regionalization of the Neotropical region, *Zootaxa*, 3782, 1–110, <https://doi.org/10.11646/zootaxa.3782.1.1>, 2014.
- Moyle, P. B., and Leidy, R. A.: Loss of Biodiversity in Aquatic Ecosystems: Evidence from Fish Faunas, in: *Conservation Biology*, edited by: Fiedler, P.L., and Jain, S. K., Springer, Boston, MA, https://doi.org/10.1007/978-1-4684-6426-9_6, 127–169, 1992.
- 900 Mueller, A., Anselmetti, F., Ariztegui, D., Brenner, D., Hodell, D., Curtis, J., Escobar, J., Gilli, A., Grzesik, D., Guilderson, T., Kutterolf, S., and Plötze, M.: Late Quaternary palaeoenvironment of northern Guatemala: evidence from deep drill cores and seismic stratigraphy of Lake Petén Itzá, *Sedimentology*, 57,1220–1245, <https://doi.org/10.1111/j.1365-3091.2009.01144.x>, 2010.
- Myers, N., Mittermeier, R. A., Mittermeier, C. G., da Fonseca, G. A. B., and Kent, J.: Biodiversity hotspots for conservation priorities, *Nature*, 403, 853–858, <https://doi.org/10.1038/35002501>, 2000.
- 905 Newhall, C.G., and Dzurisin, D.: Historical unrest at large calderas of the world, *US Geol. Surv. Bull.*, 2, 210–213, <https://doi.org/10.3133/b1855>, 1988.
- Obrist-Farner, J., and Rice, P. M.: Nixtun-Ch'ich' and its environmental impact: Sedimentological and archaeological correlates in a core from Lake Petén Itzá in the southern Maya lowlands, Guatemala, *J. Archaeol. Sci. Rep.*, 26, 101868, <https://doi.org/10.1016/j.jasrep.2019.05.033>, 2019.
- 910 Obrist-Farner, J., Brenner, M., Stone, J., Wojewódka, M., Bauersachs, T., Eckert, A., Locmelis, M., Curtis, J., Zimmerman, S., Correa-Metrio, A., Schwark, L., Duarte, E., Schwalb, A., Niewerth, E., Echeverría Galindo, P., and Pérez, L.: New estimates of the magnitude of the sea-level jump during the 8.2 ka event, *Geology*, XX, p., <https://doi.org/10.1130/G49296.1>, 2021.
- 915 Ojeda, R. A., Stadler, J., and Brandl, R.: Diversity of mammals in the tropical-temperate neotropics: Hotspots on a regional scale, *Biodivers. Conser.*, 12, 1431–1444, <https://doi.org/10.1023/A:1023625125032>, 2003.
- Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P. R., O'Hara, R. B., Simpson, G. L., Solymos, Stevens, M. H. H., Szoecs, E., and Wagner, H.: *vegan: Community Ecology Package*, R package version 2.4-5, <https://CRAN.R-project.org/package=vegan>, 2017.

- 920 Patton, J. L., da Silva, M. N. F., and Malcolm, J. R.: Gene genealogy and differentiation among arboreal spiny rats (Rodentia: Echimyidae) of the Amazon Basin: a test of the riverine barrier hypothesis, *Evolution*, 48, 1314–1323, <https://doi.org/10.1111/j.1558-5646.1994.tb05315.x>, 1994.
- Peel, M. C., Finlayson, B. L., and McMahon, T. A.: Updated world map of the Köppen–Geiger climate classification, *Hydrol. Earth Syst. Sci.*, 11, 1633–1644, <https://doi.org/10.5194/hess-11-1633-2007>, 2007.
- 925 Pérez-Ceballos, R., Pacheco-Ávila, J., Euán-Ávila J. I., and Hernández-Arana, H.: Regionalization based on water chemistry and physicochemical traits in the ring of cenotes, Yucatan, Mexico. *J. Caves Karst Stud.*, 74, 90–102, <https://doi.org/10.4311/2011es0222>, 2012.
- Pérez, L., Bugja, R., Lorenschat, J., Brenner, M., Curtis, J., Hoelzmann, P., Islebe, G., Scharf, B., and Schwalb, A.: Aquatic ecosystems of the Yucatán Peninsula (Mexico), Belize, and Guatemala. *Hydrobiologia*, 661, 407–433, <https://doi.org/10.1007/s10750-010-0552-9>, 2011a.
- 930 Pérez, L., Frenzel, P., Brenner, M., Escobar, J., Hoelzmann, P., Scharf, B., and Schwalb, A.: Late Quaternary (24–10 ka BP) environmental history of the Neotropical lowlands inferred from ostracodes in sediments of Lago Petén Itzá, Guatemala, *J. Paleolimnol.*, 46, 59–74, <https://doi.org/10.1007/s10933-011-9514-0>, 2011b.
- Pérez, L., Lorenschat, J., Brenner, M., Scharf, B., and Schwalb, A.: Extant freshwater ostracodes (Crustacea: Ostracoda) from Lago Petén Itzá, Guatemala, *Rev. Biol. Trop.*, 58, 871–895, <https://doi.org/10.15517/rbt.v58i2.5252>, 2010a.
- 935 Pérez, L., Lorenschat, J., Massaferró, J., Paillès, C., Sylvestre, F., Hollwedel, W., Brandorff, G.O., Brenner, M., Islebe, G., Lozano, M.S., Scharf, B., and Schwalb, A.: Bioindicators of climate and trophic state in lowland and highland aquatic ecosystems of the Northern Neotropics, *Rev. Biol. Trop.*, 61, 603–644, 2013a.
- Perry, E., Paytan, A., Pedersen, B., and Velazquez-Oliman, G.: Groundwater geochemistry of the Yucatan Peninsula, Mexico: Constraints on stratigraphy and hydrogeology. *J. Hydrol.*, 367, 27–40, <https://doi.org/10.1016/j.jhydrol.2008.12.026>, 2009.
- 940 Perry, E., Velazquez-Oliman, G., Marin, L.: The Hydrogeochemistry of the Karst Aquifer System of the Northern Yucatan Peninsula, Mexico. *Int. Geol. Review*, 44, 191–221, <https://doi.org/10.2747/0020-6814.44.3.191>, 2002.
- Peterson, A. T., and Nyári, A. S.: Ecological niche conservatism and Pleistocene refugia in the Thrush-like Mourner, *Schiffornis* sp., in the neotropics, *Evolution*, 62, 173–183, <https://doi.org/10.1111/j.1558-5646.2007.00258.x>, 2008.

- 945 R Core Team: R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria, <https://www.R-project.org/>, 2017.
- Reed, J. C., Wheeler, J. O., and Tucholke, B. E.: Decade of North American Geology, Geologic Map of North America, 1:5000000 continent scale map 001, Geological Society of America, Inc. Boulder, CO., 2005.
- Roessler, E. W.: Estudios sobre los ostracodos de agua dulce en Colombia – 6, Parte 4: Estudio taxonómico del grupo
950 *Strandesia psittacea psittacea* (Sars, 1901)', *Caldasia*, 16, 215–230, 1990a.
- Roessler, E.W.: Estudios taxonómicos, ontogenéticos, ecológicos y etológicos sobre los ostrácodos de agua dulce en Colombia (Ostracoda, Podocopida, Cyprididae) – 6: Estudio taxonómico del género *Strandesia* Stuhlmann, 1888 – Parte 3: El grupo *Strandesia elliptica* (Sars, 1901). *Revista Acad. Colomb. Ci. Exact.*, 17, 795–804, 1990b.
- Rosencrantz, E.: Structure and tectonics of the Yucatan Basin, caribbean Sea, as determined from seismic reflection
955 studies, *Tectonics*, 9, 1037–1059, <https://doi.org/10.1029/TC009i005p01037>, 1990.
- Rosseel, Y.: Lavaan: An R package for structural equation modeling and more. Version 0.5–12 (BETA). *Journal of Statistical Software*, 48(2), 1–36, <https://cutt.ly/xgw1RZJ>, 2012.
- Rossetti, D. P., and Toledo, P. M.: Biodiversity from a Historical Geology Perspective: A Case Study from Marajó Island, Lower Amazon, *Geobiology*, 4, 215–223, <http://dx.doi.org/10.1111/j.1472-4669.2006.00080.x>, 2006.
- 960 Rull, V.: Neotropical biodiversity: timing and potential drivers. *Trends Ecol.*, 26, 508–513
<https://doi.org/10.1016/j.tree.2011.05.011>, 2011.
- Saint-Loup, R., Felix, T., Maqueda, A., Schiller, A., and Renard, P.: A survey of groundwater quality in Tulum region, Yucatan Peninsula, Mexico, *Environ. Earth Sci.*, 77, 1–20 <https://doi.org/10.1007/s12665-018-7747-1>, 2018.
- Saldarriaga, A. T., and Martínez J.I.: Ecology of non – marine ostracoda from La Fe
965 reservoir (El Retiro, Antioquia) and their potential application in paleoenvironmental studies, *Rev. Acad. Col. Cienc.*, 34, 397–409, 2010.
- Sánchez-Sánchez, J. A., Álvarez-Legorreta, T., Pacheco-Ávila, J. G., González-Herrera, R. A., and Carrillo-Briebiezca, L.: Caracterización hidrogeoquímica de las aguas subterráneas del sur del Estado de Quintana Roo, México. *Rev. Mex. Cienc. Geol.*, 32, 62-76, 2015.

- 970 Sarstedt, M., Ringle, C. M.: *Structural Equation Models: From Paths to Networks* (Westland 2019), *Psychometrika*, 85, 841–844, <https://doi.org/10.1007/s11336-020-09719-0>, 2020.
- Schmitter-Soto, J., Comín, F., Escobar-Briones, E., Herrera, J., Alcocer, J., Suarez-Morales, E., Elías-Gutiérrez, M., Díaz, V., Marin, L., and Steinich, B.: Hydrogeochemical and biological characteristics of cenotes in the Yucatan Peninsula (SE Mexico), *Hydrobiologia*, 467, 215–228, <https://doi.org/10.1023/A:1014923217206>, 2002a.
- 975 Schmitter-Soto, J., Escobar-Briones, E., Alcocer, J., Suarez-Morales, E., Elías-Gutiérrez, M., and Marín, L.: Los cenotes de la Península de Yucatán, in *Lagos y Presas de México*, edited by: De La Lanza, G., and García-Calderón, J. L., AGT, México, 338–376, 2002b.
- [Schrodt, F., Santos, M. J., Bailey, J. J., and Field, R.: Challenges and opportunities for biogeography—What can we still learn from von Humboldt?, *J. Biogeogr.*, 46, 1631–1642. <https://doi.org/10.1111/jbi.13616>, 2019.](https://doi.org/10.1111/jbi.13616)
- 980 Schumm M., Edie S. M., Collins K. S., Gómez-Bahamón V., Supriya K., White A. E., Price T. D., and Jablonski D.: Common latitudinal gradients in functional richness and functional evenness across marine and terrestrial systems, *Proc. R. Soc. B.*, 286, 20190745, <http://doi.org/10.1098/rspb.2019.0745>, 2019.
- Sigala, I., Caballero, M., Correa-Metrio, A., Lozano-García, S., Vázquez, G., Pérez-Alvarado, L., and Zawisza, E.: Basic limnology of 30 continental waterbodies of the Transmexican Volcanic Belt across climatic and environmental gradients. *B. Soc. Geol. Mex.*, 69, 313–370, <https://doi.org/10.18268/bsgm2017v69n2a3>, 2017.
- 985 Sigala, I., Lozano-García, S., Escobar, J., Pérez, L., and Gallegos-Neyra, E.: Testate Amoebae (Amebozoa: Arcellinida) in Tropical Lakes of Central Mexico, *Rev. Biol. Trop.*, 64, 377–397, <https://doi.org/10.15517/rbt.v64i1.18004>, 2016.
- Sigala Regalado, I., Lozano García, S., Pérez Alvarado, L., Caballero, M., and Lugo Vázquez, A.: Ecological drivers of testate amoeba diversity in tropical water bodies of central Mexico, *J. Limnol.*, 77 (3), 385–399, <https://doi.org/10.4081/jlimnol.2018.1699>, 2018.
- 990 Sluys, R.: Ostracods (Ostracoda, Crustacea) from two freshwater lakes near Maarssen, The Netherlands, *Hydrobiol. Bull.*, 15, 92–93, <https://doi.org/10.1007/BF02260263>, 1981.
- Šmilauer, P., and Lepš, J.: *Multivariate Analysis of Ecological Data using Canoco 5*, Cambridge University Press, Cambridge, 1–362, 2014.

- 995 Smith, B. T., Amei, A., and Klicka, J.: Evaluating the role of contracting and expanding rainforest in initiating cycles of speciation across the Isthmus of Panama, *Proc. Biol. Sci.*, 279, 3520–3526, <https://doi.org/10.1098/rspb.2012.0706>, 2012.
- Smith, G. R., Badgley, C., Eiting, T. P., and Larson, P. S.: Species diversity gradients in relation to geological history in North American freshwater fishes, *Evol. Ecol. Res.*, 12, 693–726, 2010.
- 1000 Socki, R., Perry, E., and Romanek, C.: Stable isotope systematics of two cenotes from the northern Yucatan Peninsula, Mexico, *Limnol. Oceanogr.*, 47, 1808–1818, <https://doi.org/10.4319/lo.2002.47.6.1808>, 2002.
- Soto, D. X., Sánchez-Murillo, R., Ortega, L., Quiroz-Londoño, O. M., Araguás-Araguás, L. J. and Martins, V.: Environmental isotope applications in Latin America and the Caribbean region. *Isot. Environ. Health Stud.*, 56, 387–390, <https://doi.org/10.1080/10256016.2020.1839448>, 2020.
- 1005 Steinbauer, M. J., Field, R., Grytnes, J., Trigas, P., Ah-Peng, C., Attorre, F., Birks, H. J. B., Borges, P. A. V., Cardoso, P., Chou, C., Sanctis, M. D., de Sequeira, M. M., Duarte, M. C., Elias, R. B., Fernández-Palacios, J. M., Gabriel, R., Gereau, R. E., Gillespie, R. G., Greimler, J., Harter, D. E. V., Huang, T., Irl, S. D. H., Jeanmonod, D., Jentsch, A., Jump, A. S., Kueffer, C., Nogué, S., Otto, R., Price, J., Romeiras, M. M., Strasberg, D., Stuessy, T., Svenning, J., Vetaas, O. R., Beierkuhnlein, C., and Gillespie, T.: Topography-driven isolation, speciation and a global increase of endemism with elevation, *Global Ecol. Biogeogr.*, 25, 1097–1107, <https://doi.org/10.1111/geb.12469>, 2016.
- 1010 Steinich, B., and Marín, L. E.: Hydrogeological investigations in northwestern Yucatán, México, using resistivity surveys, *Groundwater*, 34, 640–646, <https://doi.org/10.1111/j.1745-6584.1996.tb02051.x>, 1996.
- Stoiber, R. E., and Carr, M. J.: Quaternary volcanic and tectonic segmentation of Central America. *Bull. Volcanol.*, 37, 304–325, <https://doi.org/10.1007/BF02597631>, 1973.
- 1015 Taylor, M., and Alfaro, E.: Climate of Central America and the Caribbean, in: *The Encyclopedia of World Climatology*, edited by: Oliver J., Springer Press, Dordrecht, The Netherlands, 183–189, https://doi.org/10.1007/1-4020-3266-8_37, 2005.
- Ter Braak, C. J. F.: Canonical correspondence analysis: a new eigenvector technique for multivariate direct gradient analysis, *Ecology*, 67, 1167–1179, <https://doi.org/10.2307/1938672>, 1986.
- Umaña, G., Haberyan, K., and Horn, S.: Limnology in Costa Rica, in *Limnology in Developing Countries*, edited by: Wetzel, R., and Gopal, B., International Scientific Publications, New Delhi, 33–62, 1999.

- 1020 Vallance, J. W., and Calvert, A. T.: Volcanism during the past 84 ka at Atitlan caldera, Guatemala. AGU, Fall Meeting 2003, 2003.
- Vartanyan, G.: Influence of Modern Geological Processes on Evolution of Ecosystems, in: *Geology and Ecosystems*, edited by: Zektser, I. S., Marker, B., Ridgway, J., Rogachevskaya, L., and Vartanyan, G., Springer, Boston, MA, https://doi.org/10.1007/0-387-29293-4_3, 2006b.
- 1025 Vartanyan, G.: The Geological Environment and Ecosystems, in: *Geology and Ecosystems*, edited by: Zektser, I. S., Marker, B., Ridgway, J., Rogachevskaya, L., and Vartanyan, G., Springer, Boston, MA, https://doi.org/10.1007/0-387-29293-4_2, 2006a.
- Vázquez-Domínguez, E., and Arita, H.: The Yucatán peninsula: biogeographical history 65 million years in the making, *Ecology*, 33, 212–219, <https://doi.org/10.1111/j.1600-0587.2009.06293.x>, 2010.
- 1030 Vázquez-Molina, Y., Correa-Metrio, A., Zawisza, E., Franco-Gaviria, J. F., Pérez, L., Romero, F., Prado, B., Charqueño-Celis, F., and Esperón-Rodríguez, M.: Decoupled lake history and regional climates in the middle elevations of tropical Mexico, *Rev. Mex. Cienc. Geol.*, 33, 355–364, <https://doi.org/10.22201/cgeo.20072902e.2016.3.450>, 2016.
- Vogel, S., Maerker, M., Rellini, I., Hoelzmann, P., Wulf, S., Robinson, M., Steinhübel, L., Maio, G. D., Imperatore, C., Kastenmeier, P., Liebmann, L., Esposito, D., and Seiler, F. M.: From a stratigraphic sequence to a landscape evolution model: Late Pleistocene and Holocene volcanism, soil formation and land use in the shade of Mount Vesuvius (Italy), *Quater. Int.*, 394, 155–179, <https://doi.org/10.1016/j.quaint.2015.02.033>, 2016.
- 1035 Wagner, H. H.: Spatial covariance in plant communities: integrating ordination, geostatistics, and variance testing, *Ecology*, 84, 1045–1057, <http://www.jstor.org/stable/3108047>, 2003.
- Waliser, D. E., Shi, Z., Lanzante, J. R., and Oort, A. H.: The Hadley circulation: assessing NCEP/NCAR reanalysis and sparse in-situ estimates, *Clim. Dyn.*, 15, 719–735, <https://doi.org/10.1007/s003820050312>, 1999.
- 1040 Wallace, A.: On the insects used for food by the Indians of the Amazon, *Trans. R. Entomol. Soc. Lond.*, 7, 241–244, <https://doi.org/10.1111/j.1365-2311.1854.tb02224.x>, 1853.
- Wehrtmann, I., Ramirez, A., and Perez-Reyes, O.: Freshwater Decapod Diversity and Conservation in Central America and the Caribbean, in: *A Global Overview of the Conservation of Freshwater Decapod Crustaceans*, edited by Kawai T., and
1045 Cumberlidge, N., Springer, Cham., https://doi.org/10.1007/978-3-319-42527-6_9, 2016.

Wiens, J. J., and Donoghue, M. J.: Historical biogeography, ecology and species richness, *Trends Ecol. Evol.*, 19, 639–644, <https://doi.org/10.1016/j.tree.2004.09.011>, 2004.

1050 Wojewódka, M., Zawisza, E., Cohuo, S., Macario-González, L., Schwalb, A., Zawiska, I., and Pérez, L.: Ecology of Cladocera species from Central America based on subfossil assemblages, *Adv. Oceanogr. Limnol.*, 7 (2), 151–162, <https://doi.org/10.4081/aiol.2016.6266>, 2016.