

1 **Effect of plateau pikas' presence on the ecosystem services of alpine meadows**

2

3 **Yingying Chen<sup>1</sup>, Huan Yang<sup>1</sup>, Gensheng Bao<sup>2</sup>, Xiaopan Pang<sup>1</sup>, Zhenggang Guo<sup>1</sup>**

4 <sup>1</sup>Engineering Research Center of Grassland Industry, Ministry of Education; Key Laboratory

5 of Grassland Livestock Industry Innovation, Ministry of Agriculture and Rural Affairs;

6 College of Pastoral Agriculture Science and Technology, Lanzhou University, Lanzhou,

7 730020, P. R. China

8 <sup>2</sup>Academy of Animal and Veterinary Sciences, Qinghai University (Qinghai Academy of

9 Animal and Veterinary Sciences), Xining, China

10

11 **Correspondence:** Zhenggang Guo (guozhg@lzu.edu.cn)

12

13 **Abstract**

14 The activity of small mammalian herbivores influences grassland ecosystem services in arid  
15 and semi-arid regions. Plateau pika (*Ochotona curzoniae*) was considered as a focal organism  
16 to investigate the effect of small mammalian herbivores on meadow ecosystem services in  
17 alpine regions. In this study, a home-range scale was used to measure the ecological service of  
18 forage available to livestock, water conservation, carbon sequestration, and soil nutrient  
19 maintenance (total nitrogen, phosphorus, and potassium) in the topsoil layer; and a quadrat  
20 scale was used to assess the ecological service of biodiversity conservation of alpine  
21 meadows. This study showed that the presence of plateau pikas led to lower ecological  
22 services of forage available to livestock and water conservation, and led to higher ecological  
23 services of biodiversity conservation, carbon sequestration, soil nitrogen, and phosphorus  
24 maintenance of meadow ecosystems. In contrast, it had no impact on ecological service of  
25 soil potassium maintenance of meadow ecosystems in alpine regions. With the increase of  
26 disturbance intensity of plateau pikas, the forage available to livestock, biodiversity  
27 conservation, and soil nutrient maintenance of meadow ecosystems in alpine regions first  
28 increased and then decreased; the water conservation tended to decrease linearly with the  
29 increasing disturbance intensity of plateau pikas. These results present a possible pattern of  
30 plateau pikas influencing the ecosystem services of meadow ecosystems in alpine regions,  
31 richening the small mammalian herbivores in relation to grassland ecosystem services.

32 **1 Introduction**

33 Grasslands provide multiple ecosystem services, mainly including provisioning services  
34 of food and water, regulating services of carbon sequestration and water conservation,  
35 supporting services of soil nutrient maintenance and biodiversity conservation, and cultural

36 services of landscapes and recreation tourism (Millennium Ecosystem Assessment, 2005).  
37 These ecosystem services sustain animal production, flora and fauna, and other human  
38 welfare (Costanza et al., 1997; Zhang et al., 2018; Dong et al., 2020); however, they are  
39 affected by multiple biotic factors, such as soil microbial communities (Van Eekeren et al.,  
40 2010), grazing by large herbivores (Lu et al., 2017), and the presence of small herbivores  
41 (Delibes-Mateos et al., 2011; Martínez-Estévez et al., 2013).

42 Small mammalian herbivores are common biotic factors (Davidson et al., 2012). These  
43 herbivores usually create extensive disturbances on grassland vegetation and soil (Pang et al.,  
44 2020a, 2020b) by developing burrow systems (Delibes-Mateos et al., 2008; Sun et al., 2015),  
45 excreting feces and urine (Zhang et al., 2016), consuming plants (Eldridge and Myers, 2001;  
46 Liu et al., 2017), clipping tall plants (Zhang et al., 2020), and producing bare soil patches  
47 (Guo et al., 2012a, 2012b; Yu et al., 2017a, 2017b) or mounds (Yang et al., 2021). Previous  
48 studies have shown that the presence of prairie dogs (*Cynomys ludovicianus*) can increase the  
49 ecological services of forage available to livestock, water conservation, carbon sequestration,  
50 and biodiversity conservation of grassland ecosystems in arid regions (Ceballos et al., 1999,  
51 Martínez-Estévez et al., 2013), whereas the presence of European rabbit (*Oryctolagus*  
52 *cuniculus*) can decrease the ecological service of forage available to livestock  
53 (Delibes-Mateos et al., 2008; Eldridge and Myers, 2001), and increase ecological services of  
54 biodiversity conservation (Delibes-Mateos et al., 2008) and nitrogen maintenance (Willott et  
55 al., 2000) of grassland ecosystems in semi-arid regions. In addition to grasslands in arid and  
56 semi-arid regions, vast alpine meadows exist in high latitude and altitude regions throughout  
57 the world (Zhang et al., 2018; Dong et al., 2020). However, how small mammalian herbivores

58 influence the ecosystem services in alpine meadows as much as they do in arid and semi-arid  
59 regions has not been well documented.

60 The plateau pika (*Ochotona curzoniae*) is a common, small mammalian herbivore that  
61 mainly lives in alpine meadows of the Qinghai-Tibetan Plateau (Smith and Foggin, 1999).  
62 This small mammalian herbivore with averaging 150 g are diurnally active and  
63 non-hibernating (Smith and Wang, 1991; Fan et al., 1999), and preferentially consume  
64 dicotyledons (Zhao et al., 2013; Pang and Guo, 2017). Plateau pikas, a sexual monomorphism  
65 (Dobson et al., 1998), often construct a family warren with numerous burrow entrances and  
66 develop a complex burrow system with average 13 m length and 30 cm depth (Fan et al.,  
67 1999). This mammalian herbivore is social and philopatric (Dobson et al., 1998) and its  
68 young offspring stay with its family during its birth year (Wang et al., 2020). Plateau pikas are  
69 generally considered a pest in China (Harris, 2010; Pang and Guo, 2017) as they often  
70 exacerbate the degradation of alpine meadows (Liu et al., 2013; Zhang et al., 2016). However,  
71 some studies have argued that plateau pika is a key species in alpine meadow ecosystems  
72 (Smith and Foggin, 1999; Delibes-Mateos et al., 2011). This disagreement has encouraged  
73 professionals to re-evaluate the role of plateau pikas in alpine meadow ecosystems. Thus, the  
74 effects of plateau pikas' presence on ecosystem services of alpine meadows allow insight into  
75 the role of plateau pikas in alpine meadow ecosystems. Previous studies have demonstrated  
76 that the presence of plateau pikas decreases (Liu et al., 2013) or has no significant effect on  
77 plant biomass (Pang and Guo, 2017), increases (Liu et al., 2017; Pang and Guo, 2017) or  
78 decreases (Sun et al., 2015) plant-species richness, and increases (Yu et al., 2017a; Pang et al.,  
79 2020a, 2020b) or decreases (Sun et al., 2015) soil carbon and nutrients. In addition, previous

80 studies have shown that the disturbance intensity of plateau pikas affects plant-species  
81 richness, and soil nutrient stocks of alpine meadows (Yu et al., 2017a; Pang and Guo, 2018).  
82 These findings imply that plateau pikas may have an impact on the ecosystem services of  
83 alpine meadows. Thus, further studies are needed to test whether the presence of plateau pikas  
84 and its disturbance intensity influence the ecosystem services of alpine meadows, which can  
85 richen the presence of small mammalian herbivores in relation to grassland ecosystem  
86 services.

87 Since soil carbon and nutrients differ between vegetated and bare soil patches in the  
88 home range (Yu et al., 2017b), Pang et al. (2020a; 2020b) proposed that the home-range scale  
89 is a better proxy than the quadrat scale to estimate the complete effects of the presence of  
90 plateau pikas on soil carbon and nutrient stocks. Although the provisioning, regulation,  
91 support, and cultural services of alpine meadows can be estimated by multiple indicators  
92 (Egoh et al., 2012; Brown et al., 2014), one or two can be used to verify whether the presence  
93 and intensity of plateau pikas influence each ecosystem service. In previous studies, palatable  
94 plant biomass for livestock has been used to evaluate the provisioning services  
95 (Martínez-Estévez et al., 2013; Wen et al., 2013); soil-water storage and soil organic carbon  
96 stock have been used to evaluate the regulating services (Wen et al., 2013; Li and Xie, 2015;  
97 Tang et al., 2019;); and plant-species richness and soil total nutrient stocks can be used to  
98 evaluate the supporting services (Wen et al., 2013). Notably, cultural services are particularly  
99 related to the spatial scale, as many are perceived visually over distant views (Norton et al.,  
100 2012). The plateau pika is territorial and its habitat use is patchy within a given area (Pang et  
101 al., 2020a), which leads to mismatches between the spatial scale and cultural services (De

102 Groot et al., 2010). Therefore, the present study used ecological services of forage available to  
103 livestock, water conservation, carbon sequestration and soil nutrient maintenance, and  
104 biodiversity conservation to test how the presence of plateau pikas influences the ecosystem  
105 services of alpine meadows across five sites. In this study, we hypothesized that (1) the  
106 presence of plateau pikas leads to lower ecological service of forage available to livestock  
107 because of lower palatable plant biomass in the presence of small mammalian herbivores; (2)  
108 the presence of plateau pikas leads to higher ecological services of water conservation and  
109 carbon sequestration because small mammalian herbivores can increase soil-water storage  
110 and carbon stocks; and (3) the presence of plateau pikas leads to higher ecological services of  
111 biodiversity conservation and soil nutrient maintenance because small mammalian herbivores  
112 can increase plant-species richness and soil nutrient stocks.

## 113 **2 Materials and methods**

### 114 **2.1 Study site descriptions**

115 Plateau pikas can live in various habitats with different soil types, topographies, and  
116 microclimates on the Qinghai-Tibetan Plateau. To determine how the presence of plateau  
117 pikas generally influences the ecosystem services of alpine meadows, five survey sites were  
118 selected in Luqu (102°22'12"E, 34°15'51"N), Gangcha (100°26'26"E, 37°36'12"N), Haiyan  
119 (100°54'33"E, 36°57'50"N), Qilian (100°34'48"E, 37°43'26"N), and Gonghe (99°47'11"E,  
120 36°43'48"N) counties on the Qinghai-Tibetan Plateau. These five survey sites have a similar  
121 typical plateau continental climate, with elevations ranging from 3194 m at the Gonghe  
122 survey site to 3550 m at the Luqu survey site. Based on 5-year weather data, the mean annual  
123 temperatures are 3.1, 0.9, 1.9, 2.2, 3.3°C at Luqu, Gangcha, Haiyan, Qilian, and Gonghe,

124 respectively, of which the average temperature in warm season are 9.3, 8.3, 9.6, 10.3, 9.9°C  
125 and in cold season are -3.1, -6.5, -5.8, -5.9, -3.4°C. The mean annual precipitation is 439.5,  
126 258.9, 257.4, 257.0, and 239.8 mm, of which the warm season accounts for 83.4%, 92.8%,  
127 89.3%, 91.5%, 91.4% at Luqu, Gangcha, Haiyan, Qilian, and Gonghe, respectively.  
128 According to the Chinese soil classification system (Gong, 2001), the soil type at each site is  
129 alpine meadow soil, similar to Cambisol in the WRB soil classification system.

130 Animal husbandry is the dominant use of alpine meadows on the Qinghai-Tibetan  
131 Plateau, and herders traditionally graze their livestock seasonally on cold and warm  
132 grasslands. The survey sites in this study were all situated in cold grasslands, in which alpine  
133 meadows were fenced from mid-April to September, and fences were opened to grazing yaks  
134 from mid-October to early April (Zhang et al., 2020). All field data were collected in August  
135 when the annual population of plateau pikas was the highest and reproduction had largely  
136 ceased (Qu et al., 2013; Pang et al., 2020a, 2020b). In addition, the growing season for plants  
137 is short on the Qinghai-Tibetan Plateau, and some plants can turn green until July. Therefore,  
138 sampling in August is optimal because August is good time to identify all plants and ensure  
139 an accurate survey of the plant species. Notably, the small burrowing herbivore at each survey  
140 site was only plateau pikas.

## 141 **2.2 Field survey design**

142 The adult dispersal of plateau pikas is a gradual process (Pang et al., 2020a), it is easy to  
143 identify reference sites without plateau pikas, even though these sites might be potential as  
144 suitable habitats. In this study, a home-range scale was used to calculate the ecological  
145 services of forage available to livestock, water conservation, carbon sequestration, and soil

146 nutrient maintenance, and a quadrat scale was used to calculate the ecological service of  
147 biodiversity conservation.

148 A stratified random and paired design was used to select plots. The home range of the  
149 plateau pika was approximately 1262.5 m<sup>2</sup> (Fan et al., 1999), and the plot size was 35 × 35 m,  
150 which was similar to the average area of the plateau pika's home range. At each of the five  
151 sites, this study first selected 10 plots where plateau pikas were present, or where active  
152 burrow entrances were observed. The second plot was identified along the road when the first  
153 plot with plateau pikas was selected. The distance between the two plots with plateau pikas  
154 was more than 3 km, which ensured that plateau pikas of the same family would not appear in  
155 two plots at the same time. Second, a paired adjacent plot without plateau pikas and active  
156 burrow entrances was selected for each plot with plateau pikas. The plots without plateau  
157 pikas were in any direction of plots with plateau pikas. The distance between each plot with  
158 plateau pikas and its paired plot without plateau pikas ranged from 500 to 1000 m. If the  
159 distance between each paired plot was too close, the plateau pikas could move between plots  
160 with and without plateau pikas. To ensure that each plot with plateau pikas was paired with a  
161 plot without plateau pikas, each paired plot shared the alpine meadow with same dominant  
162 plant, with no obvious differences in soil type, topography, or microclimate. In total, there  
163 were 10 paired plots at each site and 100 plots across five sites, including 50 with and 50  
164 without plateau pikas. Each paired plot shared the same grazing intensity during the cold  
165 season; however, 50 paired plots consisted of different yak grazing intensity, and this can  
166 permit the general pattern relating to the effect of plateau pika disturbance on alpine meadow  
167 ecosystem services.



### 168 **2.3 Field sampling**

169 Field surveys and sampling were conducted in early August 2020. First, the active  
170 burrow entrance at each plot with plateau pikas was estimated by the “plugging tunnels  
171 method,” in which the burrow entrances were plugged with hay for 3 days (Sun et al., 2015),  
172 and the number of plugs cleared by the plateau pikas to allow access to the meadow surface  
173 was recorded (Guo et al., 2012a). The average number of burrow entrances with cleared plugs  
174 after 3 days was taken as the density of active burrow entrances. For plots with plateau pikas,  
175 the density of active burrow entrances was used as a proxy for the intensity of the disturbance  
176 (Guo et al., 2012a; Sun et al., 2015). Second, this study was restricted to plateau pikas in  
177 relation to the ecosystem services of alpine meadows. However, bare soil patches caused by  
178 other factors (no plateau pikas) is simultaneously existed on the vegetated surface in the  
179 presence/absence of plateau pikas. To actual quantify the effect of plateau pikas on ecosystem  
180 services of alpine meadows, this study only measured the area of bare soil patches caused by  
181 plateau pikas, although there exist multiple types of bare soil patches in alpine meadows. The  
182 area of each bare soil patch (created by plateau pikas) in the plot with plateau pikas was  
183 measured. Each bare soil patch was identified as regular shape or irregular shape. If one bare  
184 soil patch was identified as regular shape, such as rectangle, circle, trapezoid, etc; a ruler was  
185 used to measure its length, width, height, diameter, upper and lower bottom, and these data  
186 was used to calculate the area of that bare soil patch. If one bare soil patch was identified as  
187 irregular shape, this bare soil patch was divided into several regular shapes; the areas of these  
188 regular shapes were calculated, respectively; the area sum of these regular shapes form  
189 irregular bare soil patch was considered as the area of that irregular bare soil patch (Han et al.,

2011). Then, the sum of all bare soil patch areas in each plot with plateau pikas was calculated and defined as the bare soil area for that plot. Third, five vegetated quadrats (1 × 1 m) were placed on the vegetated surface approximately 8 m apart along a W pattern in all plots (with or without plateau pikas). In plot with plateau pikas, if quadrat was justly covered with the bare patches caused by plateau pikas, the quadrat was slightly moved to avoid it; if quadrat was justly covered with the bare patches caused by other factors, the quadrat was not moved. Fourth, alpine meadows in plot with plateau pikas consisted of bare and vegetated surface, and a bare soil patch was selected as a paired bare soil quadrat for each vegetated quadrat in the plot with plateau pikas, and the distance between each paired bare soil quadrat and vegetated quadrat was as short as possible (less than 1 m). Bare soil patch quadrat and each vegetated quadrat were beneficial to accurately measure the soil nutrient, carbon concentrations and plant biomass, which reflected the effect of plateau pikas' presence on ecosystem services in alpine meadows by comparing the parameters between plots with and without plateau pikas at home range scale. Thus, there were 5 paired quadrats, consisting of 5 vegetated and 5 bare soil quadrats in each plot with plateau pikas. Additionally, there were 5 vegetated quadrats in each plot without plateau pikas, since this study focused on bare soil patches induced by plateau pikas.

In each vegetated quadrat of the plot with or without plateau pikas, all vascular plant species were identified, and the number of plant species were recorded as plant-species richness. Then, all plants rooted in a quadrat were harvested into palatable and unpalatable plants, in which palatable plants or unpalatable plants were for yak and Tibetan sheep (Pang and Guo, 2017). Finally, all palatable plant samples were placed in envelopes and transported

212 to the laboratory.

213 Generally, most burrows derived from the activities of plateau pikas are less than 20 cm  
214 in depth (Yu et al., 2017b), although a few burrows extend to depths of 60 cm (Fan et al.,  
215 1999). In addition, the majority of plant roots in alpine meadows of the Qinghai-Tibetan  
216 Plateau are in the top 20 cm of the soil. The soil samples were collected at a depth of 20 cm.  
217 Soil samples were collected from vegetated and bare soil quadrats for each plot with plateau  
218 pikas and vegetated quadrats for each plot without plateau pikas. Before collecting the soil  
219 samples, plants and litter were removed from the soil surface. First, a 5 cm diameter soil  
220 auger was used to collect soil samples, which were used to measure soil organic carbon and  
221 soil nutrient concentrations (total nitrogen, phosphorus, and potassium). Second, soil profiles  
222 in each quadrat were obtained using a spade, and a stainless-steel cutting ring (with a volume  
223 of 100 cm<sup>3</sup>) was used to collect soil cores to determine soil bulk density and soil water  
224 content. Soil samples used to determine soil bulk density were packed into aluminum boxes  
225 with recorded weights, and each aluminum box was numbered. The aluminum boxes  
226 containing fresh soil were immediately weighed, recorded, stored at 4°C, and then transported  
227 to the laboratory. Thus, in this study, 10 soil samples were collected to analyze the soil carbon,  
228 nitrogen, phosphorus, and potassium concentrations, and 10 soil samples were obtained to  
229 determine the soil bulk density in each plot with plateau pikas. Because this study is  
230 constricted with bare soil patch caused by plateau pikas, bare soil quadrats was not considered  
231 in plot without plateau pikas; therefore, 5 soil samples were used to determine the soil carbon,  
232 nitrogen, phosphorus, and potassium concentrations, and 5 samples were obtained for the  
233 analysis of soil bulk density in each plot without plateau pikas. 5 soil samples in each plot

234 were individually measured. The average value of five soil samples in one plot was  
235 considered as the representative data of that plot.

#### 236 **2.4 Analysis of samples**

237 In the laboratory, palatable plant samples were dried in an oven at 80°C for 48 h and  
238 weighed. The soil samples used to measure soil bulk density and soil-water content were  
239 dried to a constant weight at 105±2°C, and the aluminum boxes with dry soil were weighed  
240 and the values were recorded. The soil samples used to measure soil organic carbon, total  
241 nitrogen, phosphorus, and potassium concentrations were air-dried, gravel and roots were  
242 manually picked out, and the proportion of gravel larger than 2.0 mm in the soil sample was  
243 determined by passing through a 2.0 mm sieve. Finally, soil samples were sieved at 0.15 mm  
244 to analyze soil organic carbon, nitrogen, phosphorus, and potassium concentrations in the  
245 laboratory. Soil organic carbon was measured using the dichromate heating-oxidation  
246 (Naelson and Sommers, 1982). Soil total nitrogen concentration was measured using the  
247 Kjeldahl procedure. Soil total phosphorus concentration was measured using the  
248 Molybdenum blue colorimetric method. Soil total potassium concentration was measured  
249 using flame photometry.

250 Soil bulk density, soil organic carbon, and nutrient concentrations (total nitrogen,  
251 phosphorus, and potassium) were used to calculate the soil organic carbon, total nitrogen,  
252 phosphorus, and potassium stocks. Soil bulk density and soil-water content were used to  
253 calculate the soil-water storage (Jia et al., 2020).

#### 254 **2.5 Calculations**

255 The bare soil area consisted of all bare soil patches, and the vegetated surface area was

256 estimated from the plot areas minus the bare soil areas. This study only measured the area of  
257 bare soil patches caused by plateau pikas, although there were other kinds of bare soil patches  
258 in alpine meadows. Therefore, bare soil areas caused by plateau pikas were considered to be  
259 zero in each plot without plateau pikas, and the vegetated surface area was considered to be  
260 100%.

261 The palatable plant biomass was calculated using the following equation:

$$262 \quad GB = B_q \times \delta_{va} \quad (1)$$

263 where  $GB$ ,  $B_q$ , and  $\delta_{va}$  are the palatable plant biomass of the plot, palatable plant biomass  
264 on the quadrat scale ( $\text{g m}^{-2}$ ), and vegetated surface area, respectively.

265 The plant-species richness in a quadrat ( $1 \times 1 \text{ m}$ ) was measured using the species number  
266 of each quadrat.

267 Soil-water storage was determined using the method recommended by Jia et al. (2020),  
268 and it was calculated by the following equation:

$$269 \quad SWS_{pika} = (SWC_{BA} \times BD_{BA} \times T \times (1 - \theta_{BA}) \times 0.01 \times BA) + (SWC_{VA} \times BD_{VA} \times T \times (1 - \theta_{VA}) \times 0.01 \times VA) \quad (2)$$

270 Where  $SWS_{pika}$ ,  $SWC_{BA}$ ,  $BD_{BA}$ , and  $\theta_{BA}$  were soil-water storage in a plot with plateau  
271 pikas, water content ( $\text{g kg}^{-1}$ ), soil bulk density ( $\text{g cm}^{-3}$ ) and soil fraction of gravel larger than 2  
272 mm in bare soil areas of plots with plateau pikas, respectively;  $BA$  was the percentage of  
273 bare soil areas in plots with plateau pikas;  $SWC_{VA}$ ,  $BD_{VA}$ , and  $\theta_{VA}$  were water content ( $\text{g kg}^{-1}$ ),  
274 soil bulk density ( $\text{g cm}^{-3}$ ) and soil fraction of gravel larger than 2 mm in vegetated areas of a  
275 plot with plateau pikas, respectively; and  $T$  was soil thickness (20 cm);  $VA$  was the percentage  
276 of vegetated surface area in plots with plateau pikas;  $SWC_{BA}$  and  $SWC_{VA}$  was measured by  
277 oven drying method.

278 
$$SWS_{no\ pika} = SWC_{no\ pika} \times BD_{no\ pika} \times T \times (1 - \theta_{no\ pika}) \times 0.01 \times 100\% \quad (3)$$

279 Where  $SWS_{no\ pika}$ ,  $SWC_{no\ pika}$ ,  $BD_{no\ pika}$  and  $\theta_{no\ pika}$  were soil-water storage in a plot without  
 280 plateau pikas, soil water content ( $g\ kg^{-1}$ ), soil bulk density ( $g\ cm^{-3}$ ) and soil fraction of gravel  
 281 larger than 2 mm in plots without plateau pikas, respectively; and  $T$  is soil thickness (20 cm).

282 The soil organic carbon stock per plot was estimated using the method described by Pang  
 283 et al. (2020b), and it was calculated by following equation:

284 
$$SOCS_{pika} = (SOC_{BA} \times BD_{BA} \times T \times (1 - \theta_{BA}) \times 0.01 \times BA) + (SOC_{VA} \times BD_{VA} \times T \times (1 - \theta_{VA}) \times 0.01 \times VA) \quad (4)$$

285 Where  $SOCS_{pika}$  was soil organic carbon stock in a plot with plateau pikas ( $kg\ m^{-2}$ );  
 286  $SOC_{BA}$ ,  $BD_{BA}$ , and  $\theta_{BA}$  were soil organic carbon concentration ( $g\ kg^{-1}$ ), soil bulk density ( $g\ cm^{-3}$ )  
 287 and soil fraction of gravel larger than 2 mm in bare soil areas of plots with plateau pikas,  
 288 respectively;  $BA$  was the percentage of bare soil areas in plots with plateau pikas;  $SOC_{VA}$ ,  
 289  $BD_{VA}$ , and  $\theta_{VA}$  were organic carbon concentration ( $g\ kg^{-1}$ ), soil bulk density ( $g\ cm^{-3}$ ) and soil  
 290 fraction of gravel larger than 2 mm in vegetated areas of a plot with plateau pikas,  
 291 respectively; and  $T$  was soil thickness (20 cm);  $VA$  was the percentage of vegetated surface  
 292 area in plots with plateau pikas.

293 
$$SOCS_{no\ pika} = SOC_{no\ pika} \times BD_{no\ pika} \times T \times (1 - \theta_{no\ pika}) \times 0.01 \times 100\% \quad (5)$$

294 Where  $SOCS_{no\ pika}$  was soil organic carbon stock in the plot without plateau pikas ( $kg\ m^{-2}$ );  
 295 and  $SOC_{no\ pika}$ ,  $BD_{no\ pika}$  and  $\theta_{no\ pika}$  were soil organic carbon concentration ( $g\ kg^{-1}$ ), soil  
 296 bulk density ( $g\ cm^{-3}$ ) and soil fraction of gravel larger than 2 mm in plots without plateau  
 297 pikas, respectively; and  $T$  was soil thickness (20 cm).

298 The soil total nitrogen, phosphorus, and potassium stocks per plot were quantified using  
 299 the method described by Pang et al. (2020a), and it was calculated by the following  
 300 equation:

$$301 \quad SNSi_{pika} = (SNi_{BA} \times BD_{BA} \times T \times (1 - \theta_{BA}) \times 0.01 \times BA) + (SNi_{VA} \times BD_{VA} \times T \times (1 - \theta_{VA}) \times 0.01 \times VA) \quad (6)$$

302 Where  $SNSi_{pika}$  was soil total nitrogen, phosphorus, potassium stock in plot with plateau  
 303 pikas ( $\text{kg m}^{-2}$ ), and  $SNi_{BA}$ ,  $BD_{BA}$ , and  $\theta_{BA}$  were soil nutrient concentration ( $\text{g kg}^{-1}$ ), soil bulk  
 304 density ( $\text{g cm}^{-3}$ ) and soil fraction of gravel larger than 2 mm in bare soil area of plots with  
 305 plateau pikas, respectively;  $BA$  was the percentage of bare soil areas in plots with plateau  
 306 pikas;  $SNi_{VA}$ ,  $BD_{VA}$ , and  $\theta_{VA}$  were soil nutrient concentration ( $\text{g kg}^{-1}$ ), soil bulk density ( $\text{g cm}^{-3}$ )  
 307 and soil fraction of gravel larger than 2 mm in vegetated areas of a plot with plateau pikas,  
 308 respectively; and  $T$  was soil thickness (20 cm);  $VA$  was the percentage of vegetated surface  
 309 area in plots with plateau pikas.

$$310 \quad SNSi_{no\ pika} = SNi_{no\ pika} \times BD_{no\ pika} \times T \times (1 - \theta_{no\ pika}) \times 0.01 \times 100\% \quad (7)$$

311 Where  $SNSi_{no\ pika}$  was soil total nitrogen, phosphorus, potassium stock in the plot without  
 312 plateau pikas ( $\text{kg m}^{-2}$ ),  $SNi_{no\ pika}$ ,  $BD_{no\ pika}$  and  $\theta_{no\ pika}$  were soil nutrient concentration ( $\text{g kg}^{-1}$ ),  
 313 soil bulk density ( $\text{g cm}^{-3}$ ) and soil fraction of gravel larger than 2 mm in plots without plateau  
 314 pikas, respectively; and  $T$  was soil thickness (20 cm).

## 315 2.6 Data analysis

316 Data from 50 disturbed plots and 50 undisturbed plots were used to examine the  
 317 difference in ecosystem services of alpine meadows between the presence of plateau pikas  
 318 and the absence of plateau pikas; and then data from 50 disturbed plots were used to examine  
 319 the responses of each ecosystem service of alpine meadows to the disturbance intensity of  
 320 plateau pikas.

321 All data variables (palatable plant biomass, plant-species richness, soil-water storage,  
 322 soil organic carbon stock, soil total nitrogen stock, soil total phosphorus stock, and soil total  
 323 potassium stock) were assessed for the normality and homogeneity by using the Shapiro-Wilk

324 test. If necessary, the data were base-10 log-transformed to fit the assumption of normality  
325 and homogeneity for further variance analysis.

326 A Linear Mixed Model (LMM) with the function “lmer” from the lme4 package was  
327 used to examine differences in palatable plant biomass, plant-species richness, soil-water  
328 storage, soil organic carbon stock, soil total nitrogen stock, soil total phosphorus stock, and  
329 soil total potassium stock between the presence and absence of plateau pikas across the five  
330 sites. In linear mixed models, the abovementioned parameters acted as response variables, the  
331 absence/presence were introduced as fixed factor, and the paired plots nested within each site  
332 as a random factor.

333 To clarify the responses of palatable plant biomass, plant-species richness, soil-water  
334 storage, soil organic carbon stock, soil total nitrogen stock, soil total phosphorus stock, and  
335 soil total potassium stock to the disturbance caused by plateau pikas, a linear model (LM) was  
336 used to examine the relationships between these variables and active burrow entrance  
337 densities in all plots with plateau pikas. The densities of active burrow entrances by plateau  
338 pikas were considered to be the fixed factor, and were used to construct the regression  
339 analysis between palatable plant biomass, plant-species richness, soil-water storage, soil  
340 organic carbon stock, soil total nitrogen stock, soil total phosphorus stock, and active burrow  
341 entrances densities. To select the final regression models, likelihood ratio tests were used to  
342 compare simple linear regression and polynomial regression models. After likelihood ratio  
343 tests, the models with  $p < 0.05$  and the smaller Akaike Information Criterion (AIC) were used  
344 as the final regression models.

345 The Bonferroni’s test used to adjust  $P$  values and made to correct for experiment-wise




346 error rates. All statistical analyses were performed with R 4.0.2 (R Foundation for Statistical  
 347 Computing, Vienna, Austria).

### 348 3 Results

#### 349 3.1 Effects of plateau pikas' presence on the ecosystem services of alpine meadows

350 The palatable plant biomass (Fig. 1A) and soil-water storage (Fig. 1B) were lower in the  
 351 plots with plateau pikas than in the plots without plateau pikas, whereas soil organic carbon  
 352 stock (Fig. 1C), plant-species richness (Fig. 1D), soil total nitrogen (Fig. 1E) and total  
 353 phosphorus stocks (Fig. 1F) in the plots with plateau pikas was higher than those in the plots  
 354 without plateau pikas. In addition, there was no difference in the soil total potassium stock  
 355 between the plots with and without plateau pikas (Fig. 1G).




356  
 357 **Figure 1.** Palatable plant biomass (A,  $F = 46.254$ ,  $p < 0.001$ ), soil-water storage (B,  $F =$   
 358  $35.189$ ,  $p < 0.001$ ), soil organic carbon stock (C,  $F = 87.628$ ,  $p < 0.001$ ), plant-species  
 359 richness (D,  $F = 63.569$ ,  $p = 0.003$ ), soil total nitrogen stock (E,  $F = 22.477$ ,  $p < 0.001$ ), soil

360 total phosphorus stock (F,  $F = 11.724$ ,  $p = 0.004$ ), and soil total potassium stock (G,  $F = 0.026$ ,  
 361  $p = 0.88$ ) of plots with and without plateau pikas (mean  $\pm$  standard error). Lower case  
 362 represents a significant difference between the absence and presence of pika based on an  
 363 LMM.

### 364 3.2 Effects of plateau pikas' disturbance intensity on the ecosystem services of alpine 365 meadows

366 The palatable plant biomass (Fig. 2A), soil organic carbon stock (Fig. 2C), plant-species  
 367 richness (Fig. 2D), soil total nitrogen (Fig. 2E), and phosphorus (Fig. 2F) stocks significantly  
 368 increased at first and then decreased gradually as the disturbance intensity of plateau pikas  
 369 increased. While the soil-water storage of the topsoil layer (Fig. 2B) decreased linearly with  
 370 increasing disturbance intensity of plateau pikas. In addition, the disturbance intensity of  
 371 plateau pikas had no obvious relationship with soil total potassium (Fig. 2G).



372  
 373 **Figure 2.** The palatable plant biomass (A,  $F = 68.534$ ), soil-water storage (B,  $F = 69.102$ ),

374 soil organic carbon stock (C,  $F = 12.642$ ), plant-species richness (D,  $F = 3.292$ ), soil total  
375 nitrogen stock (E,  $F = 22.901$ ), soil total phosphorus stock (F,  $F = 23.652$ ), soil total  
376 potassium stock (G) for different disturbance intensity of plateau pikas based on linear models  
377 (LMs). An adjusted local smoothed regression line (black) with its 95% confident interval  
378 (gray) was used to determine the relationship between the disturbance intensity of plateau  
379 pikas and the above indicators. ABED: active burrow entrance densities

#### 380 **4 Discussion**

381 Prairie dogs and European rabbits have been shown to affect grassland ecosystem  
382 services in arid and semi-arid regions (Delibes-Mateos et al., 2011; Martínez-Estévez et al.,  
383 2013). This study combined the home-range scale and a quadrat scales to test how the  
384 presence of plateau pikas and its disturbance intensity influence the ecosystem services of  
385 alpine meadows, and found that the presence of plateau pikas and its disturbance intensity  
386 indeed impacts the ecosystem services of alpine meadows, similar to prairie dogs and  
387 European rabbits in grassland ecosystem services in arid and semi-arid regions.

388 Lower palatable plant biomass in the presence of plateau pikas indicates that the  
389 presence of plateau pikas reduces the ecological service of forage available to livestock,  
390 which is consistent with the results of European rabbits in semi-arid regions (Eldridge and  
391 Myers, 2001; Delibes-Mateos et al., 2008), and is not consistent with results from prairie dogs  
392 in arid regions (Martínez-Estévez et al., 2013). Prairie dogs benefit perennial plants in arid  
393 grasslands, in which blue gramma (*Bouteloua gracilis*) and vine mesquite (*Panicum obtusum*)  
394 are palatable perennials for livestock (Sierra-Corona et al., 2015), whereas European rabbits  
395 increase unpalatable plants (*Marrubium vulgare* and *Colchicum melitensis*) because they

396 prefer grasses (Leigh et al., 1989; Eldridge and Myers, 2001). Plateau pikas enable more  
397 unpalatable broad-leaved plants to grow in alpine meadows (Pang and Guo, 2018) and can  
398 bury many plants (Pang and Guo, 2017). However, their consumption patterns can benefit the  
399 growth of palatable plants (Pang and Guo, 2017), because plateau pikas preferentially  
400 consume unpalatable dicotyledons (Zhao et al., 2013; Pang and Guo, 2017). The tradeoff  
401 between the decrease and increase in palatable plant biomass contributes to a negative effect  
402 on palatable plant biomass on a home-range scale, resulting in a decrease in forage available  
403 to livestock. These results demonstrate that the presence of small mammalian herbivores  
404 affects the ecological service of forage available to livestock of grassland ecosystems may be  
405 related to environmental conditions. Specific performance is that the presence of small  
406 mammalian herbivores is disadvantageous to the ecological service of forage available to  
407 livestock in semi-arid and alpine regions, but it is beneficial to the ecological service of  
408 forage available to livestock in arid regions.

409       The presence of plateau pikas has different impacts on regulating services of alpine  
410 meadows, when assessed by different indicators. The presence of plateau pikas leads to lower  
411 soil-water storage, resulting in a decrease in the ecological services of water conservation,  
412 whereas the presence of plateau pikas can lead to higher soil organic carbon stock, implying  
413 that the presence of plateau pikas can increase the ecological services of carbon sequestration.  
414 Lower ecological services of water conservation of alpine meadows in relation to the presence  
415 of plateau pikas is consistent with the effect of European rabbits' presence on ecological  
416 services of water conservation of grasslands in semi-arid regions (Eldridge et al., 2010),  
417 whereas it is inconsistent with the presence of prairie dogs in relation to ecological services of

418 water conservation in arid regions (Martínez-Estévez et al., 2013). This difference is ascribed  
419 to evaluation indicators for ecological services of water conservation. The water infiltration  
420 rate is considered as an index to evaluate the effect of prairie dogs on ecological services of  
421 water conservation of grasslands in arid regions (Martínez-Estévez et al., 2013). In contrast,  
422 the water storage of topsoil is used to evaluate the effects of European rabbits and plateau  
423 pikas on ecological services of water conservation of grasslands in semi-arid grassland and  
424 alpine meadow (Eldridge et al., 2010). The activities of European rabbits and plateau pikas  
425 can reduce the crust cover of grasslands and increase water infiltration from top soil to deep  
426 soil in semi-arid regions (Eldridge et al., 2010; Li et al., 2015), contributing to a negative  
427 effect on ecological services of water conservation in the topsoil layer. This study shows that  
428 the presence of plateau pikas leads to higher ecological services of carbon sequestration in  
429 alpine meadows, similar to the effect of the presence of prairie dogs in arid regions  
430 (Martínez-Estévez et al., 2013) and European rabbits in semi-arid regions (Delibes-Mateos et  
431 al., 2011). Plateau pikas can input extra organic matter through the deposition of uneaten food  
432 (Liu et al., 2009; Zhang et al., 2016; Yu et al., 2017a) and the excretion of urine and feces  
433 (James et al., 2009; Yu et al., 2017b), which increases the soil organic carbon stock and  
434 contributes to an increase in ecological services of carbon sequestration of alpine meadows.  
435 These results indicate that the presence of small mammalian herbivores can increase the  
436 ecological services of carbon sequestration of grasslands.

437 Higher plant-species richness in the presence of plateau pikas shows that the presence of  
438 plateau pikas can lead to higher biodiversity conservation, similar to the effect of European  
439 rabbits in semi-arid regions (Delibes-Mateos et al., 2008) and prairie dogs in arid regions

440 (Davidson et al., 2012). The mechanisms by which small mammalian herbivores lead to  
441 higher plant-species richness have been discussed in many previous studies (Zhang et al.,  
442 2020; Pang et al., 2021). The presence of plateau pikas can lead to higher soil total nitrogen  
443 and total phosphorus stocks, demonstrating that plateau pikas can increase the ecological  
444 services of soil nitrogen and phosphorus maintenance. In addition, there was no difference in  
445 the soil total potassium stock between the areas with and without plateau pikas, indicating that  
446 the presence of plateau pikas had no effect on ecological services of soil potassium  
447 maintenance. This effect was also observed with prairie dogs and European rabbits in arid  
448 (Delibes-Mateos et al., 2011) and semi-arid regions (Delibes-Mateos et al., 2008; Willott,  
449 2001). Some of the following factors explain the higher soil nitrogen and phosphorus stocks  
450 caused by plateau pikas. The presence of plateau pikas can increase the input of soil organic  
451 material (Liu et al., 2013; Zhang et al., 2016; Pang et al., 2020a). Secondly, the presence of  
452 plateau pikas can result in higher organic nitrogen and phosphorus stocks (Yu et al., 2017b),  
453 which contributes to higher ecological services of soil nitrogen and phosphorus maintenance.  
454 These results suggest that a general pattern can be identified regarding the effect of the  
455 presence of small mammalian herbivores on the supporting services of biodiversity  
456 conservation, soil nitrogen, and phosphorus maintenance.

457 In addition to the presence of plateau pikas, this study found that the disturbance  
458 intensity of plateau pikas also affects the palatable plant biomass, plant-species richness,  
459 soil-water storage, soil organic carbon stock, and soil total nitrogen and phosphorus stocks.  
460 These imply that the disturbance intensity of plateau pikas also affects the meadow ecosystem  
461 in alpine regions. With the increasing disturbance intensity of plateau pikas, the palatable

462 plant biomass, plant-species richness, soil organic carbon stock, soil total nitrogen and  
463 phosphorus stocks first increase and then decrease, demonstrating that the intermediate  
464 disturbance intensity of plateau pikas to maximize ecological services of forage available to  
465 livestock, biodiversity conservation, carbon sequestration, and soil nitrogen and phosphorus  
466 maintenance. At intermediate disturbance intensity, the presence of plateau pikas can enhance  
467 soil total nitrogen (Li et al., 2014), organic carbon accumulation (Yu et al., 2017b), palatable  
468 plant biomass (Pang and Guo, 2018) by improving the growth potential of grass plants (Wang  
469 et al., 2012), and encourage more hygrophytes and mesophytes, annual and perennial,  
470 common and rare plants to coexist (Guo et al., 2012b), contributing to higher ecological  
471 services of forage available to livestock, biodiversity conservation, carbon sequestration, soil  
472 total nitrogen and phosphorus maintenance services. When the disturbance intensity of  
473 plateau pikas is below the intermediate disturbance intensity, stronger competition of  
474 dominant sedges often restrains the grass to grow well (Pang and Guo, 2018) and the rare  
475 plants to coexist (Wang et al., 2012), which leads the ecological services of forage available to  
476 livestock and biodiversity conservation of alpine meadows to be maintained at a low level.  
477 The increase in soil organic matter input caused by plateau pikas at low disturbance intensity  
478 is less than the intermediate disturbance intensity (Pang and Guo, 2017; Pang et al., 2020b),  
479 which enables ecological services of soil organic carbon sequestration and soil nitrogen and  
480 phosphorus maintenance of alpine meadows at low disturbance intensity of plateau pikas to  
481 maintain a relatively low level. Once the disturbance intensity of plateau pikas surpasses its  
482 intermediate disturbance intensity, low soil water content in alpine meadows (Liu et al., 2013)  
483 only sustains the xerophytes and mesophytes, most of which are unpalatable (Pang and Guo,

484 2018). This contributes to relatively low ecological services of forage available to livestock  
485 and biodiversity conservation. Low vegetation biomass at high disturbance intensity of  
486 plateau pikas decreases the input resources of soil organic matter (Sun et al., 2015; Pang and  
487 Guo, 2017), contributing to a decrease in ecological services of soil organic carbon  
488 sequestration and soil nitrogen and phosphorus maintenance of alpine meadows. Additionally,  
489 the linearly negative relationship between ecological services of water conservation of alpine  
490 meadow and disturbance intensity of plateau pikas is ascribed to evaporation and more water  
491 infiltration on bare soil patches, as the amount of water evaporation and infiltration tends to  
492 increase as the area of bare soil increases (Liu et al., 2013).

493 Together with previous studies (Delibes-Mateos et al., 2011; Martínez-Estévez et al.,  
494 2013), this study demonstrates that the presence of small mammalian herbivores has similar  
495 impacts on ecological services of biodiversity conservation, soil nutrient maintenance, and  
496 carbon sequestration of grasslands throughout the arid, semi-arid, and alpine regions, whereas  
497 the effects of the presence of small mammalian herbivores on ecological services of forage  
498 available to livestock and water conservation are dependent on environmental conditions.  
499 This study further verifies that the disturbance intensity of plateau pikas also has a significant  
500 impact on the ecosystem services of alpine ecosystems. These results concur with the findings  
501 in research fields of small mammalian herbivores in relation to grassland ecosystem services.

## 502 **5 Conclusions**

503 This study focused on plateau pikas to investigate the responses of ecological services of  
504 forage available to livestock, water conservation, carbon sequestration, soil nutrient  
505 maintenance, and biodiversity conservation of meadow ecosystems to the presence of a small



506 mammalian herbivore and its disturbance intensity across five sites. This will provide insight  
507 into the relationship between small mammalian herbivores and ecosystem services of  
508 grasslands. The results of this study showed that the presence of plateau pikas led to higher  
509 ecological services of biodiversity conservation, soil nitrogen and phosphorus maintenance,  
510 and carbon sequestration of alpine meadows, whereas it led to lower ecological services of  
511 forage available to livestock and water conservation of alpine meadows. Furthermore, this  
512 study found that the effect of plateau pikas disturbance intensity on ecological services of  
513 forage available to livestock, biodiversity conservation, soil maintenance of nitrogen and  
514 phosphorus, and carbon sequestration also conformed to the moderate disturbance hypothesis.  
515 These results verified that plateau pikas could affect the ecosystem services of meadow  
516 ecosystems in alpine regions and present a relatively complete pattern of small mammalian  
517 herbivores influencing grassland ecosystem services.  
518

519 *Author contributions.* YC and ZG conceived the ideas and designed the methodology; YC, XP,  
520 GB and HY collected the data; YC analysed the data; YC and ZG wrote the manuscript. All of  
521 the authors contributed critically to the drafts and gave their final approval for publication.

522

523 *Competing interests.* The authors declare that they have no known competing financial  
524 interests or personal relationships that could have influenced to influence the work reported in  
525 this paper.

526

527 *Acknowledgments.* The authors would like to thank Jing Zhang, Qian Wang, Haipeng Xu,  
528 Wenna Zhang, Juan Wang, Ding Yang, Jie Li, Fuyun Qiao, Digang Zhi, Haohao Qi, Ganlin  
529 Feng and Yuanyuan Duan from Lanzhou University for the contributions made to this study  
530 through their field assistance and laboratory analysis.

531

532 *Financial support.* This study was funded by the National Natural Science Foundation of  
533 China (32171675), the Key Laboratory of Superior Forage Germplasm in the Qinghai-Tibetan  
534 Plateau (2020-ZJ-Y03) and the Open Project of State Key Laboratory of Plateau Ecology and  
535 Agriculture, Qinghai University.

536

537

538

539

540

541 **References**

- 542 Brown, C., Reyers, B., Ingwall-King, L., Mapendembe, A., Nel, J., O'Farrell, P., Dixon, M.,  
543 and Bowles-Newark, N.J.: Measuring Ecosystem Services: Guidance on developing  
544 ecosystem service indicators, *Unep-Wcmc*, 72, [https://doi.org/10.13140/RG.2.2.11321.](https://doi.org/10.13140/RG.2.2.11321.83043)  
545 83043, 2014.
- 546 Ceballos, G., Pacheco, J., and List, R.: Influence of prairie dogs (*Cynomys ludovicianus*) on  
547 habitat heterogeneity and mammalian diversity in Mexico, *J. Arid. Environ.*, 41, 161–172,  
548 <https://doi.org/10.1006/jare.1998.0479>, 1999.
- 549 Costanza, R., D'Arge, R., De Groot, R., Farber, S., Grasso, M., Hannon, B., Limburg, K.,  
550 Naeem, S., O'Neill, R., Paruelo, J., Raskin, R.G., Sutton, P., and Van Den Belt, M.: The  
551 value of the world's ecosystem services and natural capital, *Nature*, 387, 253–260,  
552 <https://doi.org/10.1038/387253a0>, 1997.
- 553 Davidson, A. D., Detling, J. K., and Brown, J. H.: Ecological roles and conservation  
554 challenges of social, burrowing, herbivorous mammals in the world's grasslands. *Front.*  
555 *Ecol. Environ.*, 10, 477-486. <https://doi.org/10.1890/110054>, 2012.
- 556 Delibes-Mateos, M., Delibes, M., Ferreras, P., and Villafuerte, R.: Key role of European  
557 rabbits in the conservation of the western Mediterranean Basin hotspot. *Conserv. Biol.*, 22,  
558 1106–1117, <https://doi.org/10.1111/j.1523-1739.2008.00993.x>, 2008.
- 559 Delibes-Mateos, M., Smith, A. T., Slobodchikoff, C. N., and Swenson, J. E.: The paradox of  
560 keystone species persecuted as pests: A call for the conservation of abundant small  
561 mammals in their native range, *Biol. Conserv.*, 144, 1335–1346, [https://doi.org/10.1](https://doi.org/10.1016/j.biocon.2011.02.012)  
562 016/j.biocon.2011.02.012, 2011.

563 De Groot, R. S., Alkemade, R., Braat, L., Hein, L., and Willemsen, L.: Challenges in  
564 integrating the concept of ecosystem services and values in landscape planning,  
565 management and decision making, *Ecol. Complex.*, 7, 260–272, <https://doi.org/10.1016/j.ecocom.2009.10.006>, 2010.

567 Dobson, F. S., Smith, A. T., and Gao, W. X.: Social and ecological influences on dispersal and  
568 philopatry in the plateau pika (*Ochotona curzoniae*), *Behav. Ecol.*, 9, 622–635, <https://doi.org/10.1093/beheco/9.6.622>, 1998.

570 Dong, S. K., Shang, Z. H., Gao, J. X., and Boone, R. B.: Enhancing sustainability of grassland  
571 ecosystems through ecological restoration and grazing management in an era of climate  
572 change on Qinghai-Tibetan Plateau, *Agr. Ecosyst. Environ.*, 287, <https://doi.org/10.1016/j.agee.2019.106684>, 2020.

574 Egoh, B., Drakou, E. G., Dunbar, M. B., Maes, J., and Willemsen, L.: Indicators for mapping  
575 ecosystem services: a review, European Commission, Joint Research Centre (JRC) (p. 111).  
576 <https://doi.org/10.2788/41823>, 2012.

577 Eldridge, D. J., and Myers, C. A.: The impact of warrens of the European rabbit (*Oryctolagus*  
578 *cuniculus* L.) on soil and ecological processes in a semi-arid Australian woodland, *J. Arid*  
579 *Environ.*, 47, 325–337. <https://doi.org/10.1006/jar e.2000.0685>, 2001.

580 Eldridge, D. J., Bowker, M. A., Maestre, F. T., Alonso, P., Mau, R. L., Papadopoulos, J., and  
581 Escudero, A.: Interactive effects of three ecosystem engineers on infiltration in a semi-arid  
582 Mediterranean grassland. *Ecosystems*, 13, 499-510, DOI: 10.1007/s10021-010-9335-4,  
583 2010.

584 Fan, N., Zhou, W., Wei, W., Wang, Q., and Jiang, Y.: Rodent pest management in the

585 Qinghai-Tibet alpine meadow ecosystem, in *EcologicallyBased Rodent Management*, eds  
586 G. R. Singleton, L. A. Hinds, H. Leirs, and Z. Zhang Canberra, ACT: Australian Centre  
587 International Agricultural Research, 285-304, 1999.

588 Gong, Z.: *Chinese soil taxonomy*. Science Press, China (in Chinese), 2001.

589 Guo, Z. G., Zhou, X. R., and Hou, Y.: Effect of available burrow densities of plateau pika  
590 (*Ochotona curzoniae*) on soil physicochemical property of the bare land and vegetation  
591 land in the Qinghai-Tibetan Plateau, *Acta Ecologica Sinica*, 32, 104-110, [https://doi.or](https://doi.org/10.1016/j.chnaes.2012.02.002)  
592 [g/10.1016/j.chnaes.2012.02.002](https://doi.org/10.1016/j.chnaes.2012.02.002), 2012a.

593 Guo, Z. G., Li, X. F., Liu, X. Y., and Zhou, X. R.: Response of alpine meadow communities to  
594 burrow density changes of plateau pika (*Ochotona curzoniae*) in the Qinghai-Tibet  
595 Plateau. *Acta Ecologica Sinica*, 32, 44-49, <https://doi.org/10.1016/j.chnaes.2011.12.002>,  
596 2012b.

597 Han, L. H., Shang, Z. H., Ren, G. H., Wang, Y. L., Ma, Y. S., Li, X. L., and Long, R. J.: The  
598 response of plants and soil on black soil patch of the Qinghai-Tibetan Plateau to variation  
599 of bare-patch areas, *Acta Prataculturae Sinica*, 20, 1-6, DOI:1004-5759(2011)01-001-06,  
600 2011.

601 Harris, R. B.: Rangeland degradation on the Qinghai-Tibetan plateau: A review of the  
602 evidence of its magnitude and causes. *J. Arid. Environ.*, 74, 1-12. [https://doi.org/10.101](https://doi.org/10.1016/j.jaridenv.2009.06.014)  
603 [6/j.jaridenv.2009.06.014](https://doi.org/10.1016/j.jaridenv.2009.06.014), 2010.

604 James, A. I., Eldridge, D. J., and Hill, B. M.: Foraging animals create fertile patches in an  
605 Australian desert shrubland, *Ecography*, 32, 723-732, [https://doi.org/10.1111/j.1600-0](https://doi.org/10.1111/j.1600-0587.2009.05450.x)  
606 [587.2009.05450.x](https://doi.org/10.1111/j.1600-0587.2009.05450.x), 2009.

607 Jia, Q. M., Xu, R. R., Chang, S. H., Zhang, C., Liu, Y. J., Shi, W., Peng, Z. C., and Hou, F. J.:  
608 Planting practices with nutrient strategies to improves productivity of rain-fed corn and  
609 resource use efficiency in semi-arid regions, *Agr. Water Manage.*, 228, 105879, [https://](https://doi.org/10.1016/j.agwat.2019.105879)  
610 [doi.org/10.1016/j.agwat.2019.105879](https://doi.org/10.1016/j.agwat.2019.105879), 2020.

611 Leigh, J. H., Wood, D. H., Slee, A. V, and Stanger, M. G.: Effects of rabbit and kangaroo gra  
612 zing on two semi-arid grassland communities in central-western new south wales, *Aust. J.*  
613 *Bot.*, 37, 375-396, <https://doi.org/10.1071/BT9890375>, 1989.

614 Li, J., Zhang, F. W., Lin, L., Li, H. Q., Du, Y. G., Li, Y. K., and Cao, G. M.: Response of the  
615 plant community and soil water status to alpine Kobresia meadow degradation gradients on  
616 the Qinghai-Tibetan Plateau, China, *Ecol. Res.*, 30, 589-596, <https://doi.org/10.1007/s1>  
617 [1284-015-1258-2](https://doi.org/10.1007/s1284-015-1258-2), 2015.

618 Li, S. M., and Xie, G. D.: Spatial and temporal heterogeneity of water conservation service  
619 for meadow ecosystem, *Chinese Journal of Grassland*, 37, 88-93, DOI: CNKI: SUN:ZG  
620 CD.0.2015-02-015, 2015.

621 Liu, W., Zhang, Y., Wang, X., Zhao, J. Z., Xu, Q. M., and Zhou, L.: The relationship of the  
622 harvesting behavior of plateau pikas with the plant community, *Acta Theriologica Sinica*,  
623 29, 40-49, <http://doi.org/10.16829/j.slxb.2009.01.007>, 2009.

624 Liu, Y. S., Fan, J. W., Harris, W., Shao, Q. Q., Zhou, Y. C., Wang, N., and Li, Y. Z.: Effects of  
625 plateau pika (*Ochotona curzoniae*) on net ecosystem carbon exchange of grassland in the  
626 Three Rivers Headwaters region, Qinghai-Tibet, China, *Plant. Soil.*, 366, 491-504, [https://](https://doi.org/10.1007/s11104-012-1442-x)  
627 [doi.org/10.1007/s11104-012-1442-x](https://doi.org/10.1007/s11104-012-1442-x), 2013.

628 Liu, Y. S., Fan, J. W., Shi, Z. J., Yang, X. H., and Harris, W.: Relationships between plateau

629 pika (*Ochotona curzoniae*) densities and biomass and biodiversity indices of alpine  
630 meadow steppe on the Qinghai-Tibet Plateau China, *Ecol. Eng.*, 102, 509-518, <https://doi.org/10.1016/j.ecoleng.2017.02.026>, 2017.

632 Lu, X. Y., Kelsey, K. C., Yan, Y., Sun, J., Wang, X. D., Cheng, G. W., and Neff, J. C.: Effects  
633 of grazing on ecosystem structure and function of alpine grasslands in Qinghai-Tibetan  
634 Plateau: A synthesis, *Ecosphere*, 8, e01656, <https://doi.org/10.1002/ecs2.1656>, 2017.

635 Martínez-Estévez, L., Balvanera, P., Pacheco, J., and Ceballos, G.: Prairie dog decline reduces  
636 the supply of ecosystem services and leads to desertification of semiarid grasslands, *PLoS*  
637 *One*, 8, e75229, <https://doi.org/10.1371/journal.pone.0075229>, 2013.

638 Millennium Ecosystem Assessment.: Ecosystems and human well-being: Synthesis, Island  
639 Press, Washington, DC, 2005.

640 Nelson, D. W., and Sommers, L. E.: Total carbon, organic carbon, and organic matter.  
641 Methods of soil analysis. In: Part 3-Chemical and Microbiological Properties, pp. 539-579,  
642 1982.

643 Norton, L. R., Inwood, H., Crowe, A., and Baker, A.: Trialling a method to quantify the  
644 “cultural services” of the English landscape using Countryside Survey data, *Land Use*  
645 *Policy*, 29, 449–455, <https://doi.org/10.1016/j.landusepol.2011.09.002>, 2012.

646 Pang, X. P., and Guo, Z. G.: Plateau pika disturbances alter plant productivity and soil  
647 nutrients in alpine meadows of the Qinghai-Tibetan Plateau, China, *Rangeland J.*, 39,  
648 133-144, <https://doi.org/10.1071/RJ16093>, 2017.

649 Pang, X. P., and Guo, Z. G.: Effects of plateau pika disturbance levels on the plant diversity  
650 and biomass of an alpine meadow. *Grassland Science*, 64, 159-166, <https://doi.org/10.11>

651 11/grs.12199, 2018.

652 Pang, X. P., Yu, C. Q., Zhang, J., Wang, Q., Guo, Z. G., and Tian, Y.: Effect of disturbance by  
653 plateau pika on soil nitrogen stocks in alpine meadows, *Geoderma*, 372, 114392, <https://doi.org/10.1016/j.geoderma.2020.114392>, 2020a.

654

655 Pang, X. P., Wang, Q., Zhang, J., Xu, H. P., Zhang, W. N., Wang, J., and Guo, Z. G.:  
656 Responses of soil inorganic and organic carbon stocks of alpine meadows to the  
657 disturbance by plateau pikas, *Eur. J. of Soil Sci.*, 71, 706-715, [https://doi.org/10.1111/ejss.](https://doi.org/10.1111/ejss.12895)  
658 12895, 2020b.

659 Pang, X. P., Wang, Q., Guo, Z. G.: The impact of the plateau pika on the relationship between  
660 plant aboveground biomass and plant species richness, *Land. Degrad. Dev.*, 32, 1205-1212,  
661 <https://doi.org/10.1002/ldr.3790>, 2021.

662 Qu, J. P., Li, W. J., Yang, M., Ji, W. H., and Zhang, Y. M.: Life history of the plateau pika  
663 (*Ochotona curzoniae*) in alpine meadows of the Tibetan Plateau, *Mamm. Biol.*, 78, 68-72,  
664 <https://doi.org/10.1016/j.mambio.2012.09.005>, 2013.

665 Sierra-Corona, R., Davidson, A., Fredrickson, E. L., Luna-Soria, H., Suzan-Azpiri, H.,  
666 Ponce-Guevara, E., and Ceballos, G.: Black-tailed prairie dogs, cattle, and the conservation  
667 of North America's Arid Grasslands, *PLoS One*, 10, e0118602, [https://doi.org/10.1371/](https://doi.org/10.1371/journal.pone.0118602)  
668 [journal.pone.0118602](https://doi.org/10.1371/journal.pone.0118602), 2015.

669 Smith, A. T., and Wang, X. G.: Social relationships of adult black-lipped pikas (*Ochotona*  
670 *curzoniae*), *J. Mammal.*, 72, 231–247, <https://doi.org/10.2307/1382094>, 1991.

671 Smith, A. T., and Foggin, J. M.: The plateau pika (*Ochotona curzoniae*) is a keystone species  
672 for biodiversity on the Tibetan plateau. *Anim. Conserv.*, 2, 235–240, <https://doi.org/10.1017>



673 /S1367943099000566, 1999.

674 Sun, F. D., Chen, W. Y., Liu, L., Liu, W., Cai, Y. M., and Smith, P.: Effects of plateau pika  
675 activities on seasonal plant biomass and soil properties in the alpine meadow ecosystems of  
676 the Tibetan Plateau. *Grassland Science*, 61, 195-203, <https://doi.org/10.1111/grs.12101>,  
677 2015.

678 Tang, Y. K., Wu, Y. T., Wu, K., Guo, Z. W., Liang, C. Z., Wang, M. J., and Chang, P. J.:  
679 Changes in trade-offs of grassland ecosystem services and functions under different grazing  
680 intensities, *Chinese Journal of Plant Ecology*, 43, 408-417, DOI: 10.17521/cjpe.2018.0289,  
681 2019.

682 Van Eekeren, N., de Boer, H., Hanegraaf, M., Bokhorst, J., Nierop, D., Bloem, J., Schouten,  
683 T., de Goede, R., and Brussaard, L.: Ecosystem services in grassland associated with biotic  
684 and abiotic soil parameters, *Soil. Biol. Biochem.*, 42, 1491-1504, <https://doi.org/10.1016/j.soilbio.2010.05.016>, 2010.

686 Wang, C. T., Wang, G. X., Liu, W., Wang, Q. L., and Xiang, Z. Y.: Vegetation roots and soil  
687 physical and chemical characteristics in degeneration succession of the *Kobresia pygmaea*  
688 meadow, *Ecology and Environmental Sciences*, 21, 409-416, DOI: 10.16258/j.cnki.  
689 1674-5906.2012.03.002, 2012.

690 Wang, Q., Guo, Z. G., Pang, X. P., Zhang, J., and Yang, H.: Effects of small-herbivore  
691 disturbance on the clonal growth of two perennial graminoids in alpine meadows, *Alpine.*  
692 *Bot.*, 130, 115-127, <https://doi.org/10.1007/s00035-020-00240-9>, 2020.

693 Wen, L., Dong, S. K., Li, Y. Y., Li, X. Y., Shi, J. J., Wang, Y. L., Liu, D. M., and Ma, Y. S.:  
694 Effect of degradation intensity on grassland ecosystem services in the alpine region of

695 Qinghai-Tibetan Plateau, China, PLoS One, 8, e58432, <https://doi.org/10.1371/journal.pone.0058432>, 2013.

696

697 Willott, S. J., Miller, A. J., Incoll, L. D., and Compton, S. G.: The contribution of rabbits  
698 (*Oryctolagus cuniculus*) to soil fertility semi-arid. Biol. Fert. Soils., 31, 379–384, <https://doi.org/10.1007/s003749900183>, 2000.

699

700 Yang, D., Pang, X. P., Jia, Z. F., and Guo, Z. G.: Effect of plateau zokor on soil carbon and  
701 nitrogen concentrations of alpine meadows. CATENA, 207: 105625. <https://doi.org/10.1016/j.catena.2021.105625>, 2021.

702

703 Yu, C., Pang, X. P., Wang, Q., Jin, S. H., Shu, C. C., and Guo, Z. G.: Soil nutrient changes  
704 induced by the presence and intensity of plateau pika (*Ochotona curzoniae*) disturbances in  
705 the Qinghai-Tibet Plateau, China, Ecol. Eng., 106, 1-9, <https://doi.org/10.1016/j.ecoleng.2017.05.029>, 2017a.

706

707 Yu, C., Zhang, J., Pang, X. P., Wang, Q., Zhou, Y. P., and Guo, Z. G.: Soil disturbance and  
708 disturbance intensity: Response of soil nutrient concentrations of alpine meadow to plateau  
709 pika bioturbation in the Qinghai-Tibetan Plateau, China, Geoderma, 307, 98-106,  
710 <https://doi.org/10.1016/j.geoderma.2017.07.041>, 2017b.

711

712 Zhang, Y., Dong, S. K., Gao, Q. Z., Liu, S. L., Liang, Y., and Cao, X. J.: Responses of alpine  
713 vegetation and soils to the disturbance of plateau pika (*Ochotona curzoniae*) at burrow  
714 level on the Qinghai-Tibetan Plateau of China, Ecol. Eng., 88, 232-236, <https://doi.org/10.1016/j.ecoleng.2015.12.034>, 2016.

715

716 Zhang, H. Y., Fan, J. W., Cao, W., Zhong, H. P., Harris, W., Gong, G. L., and Zhang, Y. X.:  
Changes in multiple ecosystem services between 2000 and 2013 and their driving factors in

717 the Grazing Withdrawal Program, China, *Ecol. Eng.*, 116, 67-79, <https://doi.org/10.1016/j.ecoleng.2018.02.028>, 2018.

718

719 Zhang, W. N., Wang, Q., Zhang, J., Pang, X. P., Xu, H. P., Wang, J., and Guo, Z. G.: Clipping  
720 by plateau pikas and impacts to plant community. *Rangeland. Ecol. Manag.*, 73(3), 368-374.  
721 <https://doi.org/10.1016/j.rama.2020.01.010>, 2020.

722 Zhao, G. Q., Li, G. Y., Ma, W. H., Zhao, D. Z., and Li, X. Y.: Impacts of *Ochotona pallasii*  
723 disturbance on alpine grassland community characteristics, *Chinese Journal of Applied*  
724 *Ecology*, 24, 2122-2128, <http://doi.org/10.13287/j.1001-9332.2013.0398>, 2013.