

Supplementary Material

Shuang Ma^{1,2,3}, Lifen Jiang¹, Rachel M. Wilson⁴, Jeff P. Chanton⁴, Scott Brigham⁵, Shuli Niu⁶, Colleen M. Iversen⁷, Avni Malhotra^{7,8}, Jiang Jiang⁹, Xingjie Lu¹⁰, Yuanyuan Huang¹¹, Jason Keller¹², Xiaofeng Xu¹³, Daniel M. Ricciuto⁷, Paul J. Hanson⁷, Yiqi Luo^{1,2}

¹ Center for Ecosystem Science and Society, Northern Arizona University, Flagstaff, Arizona, USA

² Department of Biological Sciences, Northern Arizona University, Flagstaff, Arizona, USA

³ Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA, USA

⁴ Earth, Ocean and Atmospheric Sciences, Florida State University, Tallahassee, Florida, USA

⁵ Institute of Ecology & Evolution, University of Oregon, Eugene, Oregon, USA

⁶ Key Laboratory of Ecosystem Network Observation and Modeling, Institute of Geographic Sciences and Natural Resources Research, Chinese Academy of Sciences, Beijing, China

⁷ Environmental Sciences Division and Climate Change Science Institute, Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA

⁸ Department of Earth System Science, Stanford University, Stanford, California, USA

⁹ Department of Soil and Water Conservation, Nanjing Forestry University, Nanjing, Jiangsu, China

¹⁰ School of Atmospheric Sciences, Sun Yat-sen University, Guangzhou, Guangdong, China

¹¹ CSIRO Oceans and Atmosphere, Aspendale, Victoria, Australia

¹² Schmid College of Science and Technology, Chapman University, Orange, USA

¹³ Biology Department, San Diego State University, San Diego, California, USA

Correspondence to: Yiqi Luo (yiqi.luo@nau.edu)

© 2021. All rights reserved.




Figure S1: Small bubbles at 20cm depth belowground, captured by manual minirhizotron at treed area from an ambient plot, August, 2016. Photo credit: John Latimer and Colleen M Iversen.