

Response to the comments of Referee 2

We are thankful for the constructive remarks of Anne Morée. In the following we reply to the comments point by point.

General comments

Comment 1 In the Introduction, a stronger argument could be made for why you decided to compare the different parameterizations, and why specifically these two. Some earlier studies state that model $\delta^{13}\text{C}$ distributions are not very sensitive to the chosen parameterization (e.g., Schmittner 2013), especially not in the surface ocean (e.g. Jahn et al., 2015). It should be discussed in Sect. 3 why your model shows something else. An interesting study on biological fractionation is e.g. that of Young et al. (2013) – such results should be discussed in light of your own.

Response: We compare different parameterisations to choose one that is more suitable for our model. We choose the parameterisations of Popp and Laws because 1) they are of different complexities and 2) their input variables are explicitly computed in our model. Furthermore, there is, to our knowledge, only a recent study by Dentith et al. (2020) that systematically addressed the impact of Popp and Laws parameterisations on both d^{13}POC and d^{13}DIC distributions. The above text is incorporated in the revised Introduction.

We don't agree with the Referee #2 that our model shows contrasting conclusions from Schmittner et al. (2013) and Jahn et al. (2015). Jahn et al. (2015) show similar surface d^{13}DIC when using different parameterisations as seen in our simulations (comparing our Fig. 7 to their Fig. 5). The similar surface d^{13}DIC for different parameterisation is an expected result because the same atmospheric CO_2 and $^{13}\text{CO}_2$ are used. In the ocean interior, Jahn et al. (2015) show lower d^{13}DIC for stronger biological fractionation, which is again consistent with this study.

Thank you for bringing the study of Young et al. (2013) to our attention. Our simulated rate of change in ϵ_p for 1960-2009 has a global mean value of -0.026‰ yr^{-1} in Hist_Popp, which it is close to Young et al. (2013)'s estimate of -0.022‰ yr^{-1} . Hist_Laws shows changes in ϵ_p with a global-mean rate of -0.005‰ yr^{-1} because Laws parameterisation is less sensitive to the change of $\text{CO}_2(\text{aq})$. Spatially Hist_Popp simulates relatively small ϵ_p change rates in eastern tropical Pacific and south of 60°S , in agreement with Young et al. (2013). The discussion on the change rate of ϵ_p is added in Section 3.1 (Isotopic signature of particular organic carbon in the surface ocean) in the revised manuscript. The spatial distributions of ϵ_p change rate are added in the Appendix.

Comment 2 As you in Sect.3 on model results mostly discuss d^{13}C , I think it is important to add a section under Sect. 3 where you introduce the reader to the model performance for the other tracers: Plus, even though this study is on the biogeochemistry, I think it is important for the reader

to get an introduction to the physical ocean model performance and specifics as well, and how it performs compared to obs (fe AMOC / Drake / SST / SSS / T / S / sea ice) - which probably has been described in a separate study but would be good to repeat here. I mainly stress this because d13C is governed by both circulation and biological processes, so any results obtained with your model setup also depend on the simulated circulation. Please summarize the biogeochemical performance shortly as well (i.e. how does e.g. PO4, O2, DIC and NO3 distributions compare to obs).

Response: We fully agree that d13C is strongly affected by both circulation and biological processes in the model. When discussing the performance of 13C tracers, we have already included the performance of several physical and biogeochemical variables. Examples can be found for the upwelling and mixed layer depth, AMOC geometry and ventilation in the North Atlantic, the Equatorial Intermediate Current System and Equatorial Deep Jets, PO4 and AOU.

In the revised manuscript, we further provide the volume transport across the Drake Passage and AMOC stream function. The distributions of sea water temperature and salinity, O2, DIC and NO3 are shown for the surface and for the zonal mean in the Atlantic, Pacific and Indian Ocean.

Comment 3 Section 4.2 is difficult to follow. Help the reader by clarifying why you are recalculating the E17 approach beyond that you have more data available as you use a model. Extra subsections, a thorough shortening and more focus on the results would help. I have to note that I am not an expert on the Suess effect or the E17 approach, but I feel I should be able to follow it based on my experience with d13C modelling. It seems quite some text is used to describe the figures: one could instead refer to the figures and only highlight the most important features of the figures.

Response: We test the E17 approach because our model simulation provides an opportunity to gain more insights into the source of the E17 approach's uncertainty because of its satisfactory performance in simulating the oceanic $\delta^{13}\text{C}$ in the late 20th century, the oceanic anthropogenic CO_2 sink, as well as the invasion of CFC-12 into the ocean.

To improve the readability of this section, we restructure it and divide it into subsections. We also move the detailed description of the E17 approach and calculations procedures to the Appendix to make this section more focused on the results and discussion.

Specific comments

Abstract

Comment 4 p1, l2-3: 'Direct comparison between paleo oceanic $\delta^{13}\text{C}$ records and model results facilitates assessing simulated distributions and properties of water masses in the past.'

This is true, but your study (mostly) focuses on the use of d13C in understanding the contemporary ocean, as you are able to use observations of POC-d13C and explore the Suess effect. This first sentences of your abstract sounds as if the goal of implementing d13C in HAMOCC was motivated by paleoceanographic questions only. Please rephrase.

Response: The above-mentioned sentence is changed to “The stable carbon isotopic composition $\delta^{13}\text{C}$ is an important variable to study ocean carbon cycle across different time scales.”

Comment 5 P1, l11: ‘because the latter results in a too strong preference for 12C’

I think here the reader could get confused (how can d13C_DIC be OK for both parametrizations but d13C_POC be better for Popp?). Maybe add something like ‘.. during C fixation, resulting in too low d13C_POC.’

Response: The suggested phrase is added.

Comment 6 p1, l15: It is not entirely clear from this sentence where the ‘that’ refers to: Has your model ample spatial and temporal data coverage? I think you want to stress here that you can repeat the Eide et al (2017) procedure with the advantage that your model has higher temporal and spatial resolution. Please clarify/rephrase.

Response: We delete this half sentence starting with ‘that’ because it is confusing. When applying the E17 approach we also used spatially-sparse model data that were sub-sampled at the geographic locations of E17. When evaluating the underestimation we compare the estimated Suess effect to the “true” simulated Suess effect.

Comment 7 P1, l14-20: This part about the Eide et al. approach makes up for almost one third of your abstract and does not connect so well to the first part. Why did you focus on the Eide et al approach, and how did you apply the findings from the first part of the study to the Eide et al part?

Response: We test Eide’s approach because “The satisfactory model performance using ϵ_p^{Popp} , regarding the present-day oceanic $\delta^{13}\text{C}$ distribution and the anthropogenic CO_2 uptake, allows us to further investigate the potential uncertainties of Eide et al. (2017a)’s approach for estimating the oceanic ^{13}C Suess effect.” This sentence is incorporated in the revised text.

Comment 8 It would also be good to finish the abstract with an outlook or overarching conclusion/summary.

Response: Thank you for the suggestion. We add a summary statement: “The new ^{13}C module in the ocean biogeochemical component of MPI-ESM shows satisfying performance. It is a useful tool to study the ocean carbon sink under the anthropogenic influences and it will be applied to investigating variations of ocean carbon cycle in the past. ”

Sect. 1 Introduction

Comment 9 p2, l36: Here one could add additional studies such as referring to HAMOCC2s by Heinze and Maier-Reimer (1999).

Response: The suggested reference is added.

Comment 10 p2 l43-45: Note that the biogeochemical model in NorESM is also called HAMOCC – one could confuse the reader here by making general statements suggesting there is only one ‘HAMOCC’. See Tjiputra et al., 2020 for the description of the implementation of d13C in NorESM-OC.

Response: We now specify the model version HAMOCC3 here.

Comment 11 p3, l55: After this sentence I would expect a paragraph on both d13C_DIC (which you provide) and d13C_POC (which is missing). Let the reader know already here what data you’ve used of d13C_POC like you do for DIC.

Response: We remove the specification of $\delta^{13}\text{C}_{\text{POC}}$ and $\delta^{13}\text{C}_{\text{DIC}}$ here because in fact we not only evaluate these two variables but also the simulated ocean physical and other ocean biogeochemical variables. In the next paragraph we focus on the oceanic 13C Suess effect, i.e. the $\delta^{13}\text{C}_{\text{DIC}}$ decrease due to anthropogenic CO2 emission.

Sect 2. Model description

Comment 12 p 4, l106: It is not only small, but also very uncertain (e.g., Zeebe and Wolf-Gladrow 2001). one could also mention here that this is commonly omitted in modeling studies. To name a few:

Schmittner, A. et al. Biology and air-sea gas exchange controls on the distribution of carbon isotope ratios ($\delta^{13}\text{C}$) in the ocean. *Biogeosciences* 10, 5793-5816, doi:10.5194/bg-10-5793-2013 (2013).

Lynch-Stieglitz, J., Stocker, T. F., Broecker, W. S. & Fairbanks, R. G. The influence of air-sea exchange on the isotopic composition of oceanic carbon: Observations and modeling. *Global Biogeochemical Cycles* 9, 653-665, doi:10.1029/95GB02574 (1995)

Tjiputra, J. F., Schwinger, J., Bentsen, M., Morée, A. L., Gao, S., Bethke, I., Heinze, C., Goris, N., Gupta, A., He, Y. C., Olivíe, D., Seland, Ø., and Schulz, M.: Ocean biogeochemistry in the Norwegian Earth System Model version 2 (NorESM2), *Geosci. Model Dev.*, 13, 2393-2431, 10.5194/gmd-13-2393-2020, 2020.

Response: Thank you, we add the suggested argument and references in the revised manuscript.

Comment 13 p4, l108-109: I suggest to move this sentence down. This would make it clearer that you first discuss total C exchange and then go into the isotope exchange.

Response: The suggested change is implemented.

Comment 14 P5, l124: You actually deviate here from the OMIP protocol of Orr et al 2017, who recommend taking 0.88 permil

Comment 15 p5, l126: Similarly here, you deviate from the OMIP protocol / the original formula who use 0.0144 and 0.107.

Comment 16 p5, l130: Why did you decide to simplify the equations here, when it computationally is not a large burden to include the whole equation? Also, I would argue again that you are not following the CMIP protocol here if you decide to simplify the air-sea gas exchange equations.

Response: This is the reply to Comments 14-16.

Thank you for catching these errors. Indeed equations 5 and 7 deviate from the OMIP protocol (Orr et al., 2017). We actually adopted these equations, as well as the simplification of Eq. (7) from Schmittner et al. (2013). We ran a short simulation and proved the small differences in Eqs. 5 and 7 lead to negligible changes in model results. Thus we didn't rerun and expensive simulations in this study.

In the revised manuscript we remove the statement about following the OMIP protocol. And we make a note to the differences in Eqs. 5 and 7 between this study and OMIP protocol.

Comment 17 P5, l132: Use 'is preferred over' instead of 'is preferentially utilised than' or rephrase in another way.

Response: We rephrase it as "is preferentially utilised over".

Comment 18 P7, l177-178: so do I understand it correctly that you base your ^{13}C model field on ^{12}C (i.e. total C) of the model and the PO_4 of the model, after the initial spinup without the ^{13}C ? Please clarify. The initialization process of an isotope model is important for spinup duration, so detailed information on this can be valuable to the readers of your work.

Response: Yes, here DIC (total C) and PO_4 are from the quasi-equilibrium state of the spin-up run without ^{13}C . This is now clarified in the revised manuscript.

Comment 19 P8, l182-184: This input rate is the input to compensate for the loss to the sediments, right? Is it equally distributed over the surface ocean?

Response: Yes, it is uniformly distributed over the surface ocean. This information is added.

Comment 20 P8, 1186-187: the inventory adjusts to be consistent with the simulated processes – what does that mean? Is the result agreeing with observational fields? This relates maybe also to my general comment that I miss an overview/summary of the non-13C performance of the model regarding both circulation and biogeochemistry.

Response: This sentence is inaccurate and redundant, so it is removed in the revised manuscript. A summary of the performance of ocean physical and non-13C biogeochemical variables is added in the Appendix.

Comment 21 P8, 1195: Is the sediment 13C also already equilibrated after the 2500y spinup?

Response: The sediment 13C is not in equilibrium. We note this in the revised manuscript.

Sect. 3 Model results and observations in the late 20th century

Comment 22 p8, 1199: One could mention here that you are not using the Eide et al. estimate of pre-industrial d13C because that is based on her estimate of the Suess effect, which you have re-evaluated. That said, you could (like Eide et al. have done) share your Suess effect estimate as a gridded dataset. On p21, 1419-420 you also explain that the E17 dataset and the Schmittner dataset are not so different, which you could mention earlier.

Response: Thank you for the suggestion. We add a paragraph to note the reasons for not using the Eide et al. estimate of pre-industrial d13C.

Our Suess effect estimate will be archived in the Max Planck Society Publication Repository as primary data and available to the public.

Please note the “E17 dataset” mentioned in the above comment is the field measurements of d13C, DIC, PO4, etc., which Eide et al. (2017a) used to estimate 13C Suess effect, not the gridded data of Eide et al. (2017b). The E17 dataset is not mentioned in Section 3 because it does not provide additional useful information.

Comment 23 P8, 1200-202: Why not regrid the obs data to the model grid, instead of regridding to 1x1 and then doing the same for the model? Also you mean the model-obs comparison for d13C_DIC here, because for POC in Fig. 4 for example you are comparing model and obs without regridding – did you then take the nearest model value?

Response: By re-gridding the observational data to a 1°x1° grid rather than to model grid (nominal resolution 1.5°) we keep slightly more detailed spatial features. The different means of re-gridding observations only has a marginal impact on model evaluation (compare for instance

the gridded d13DIC observation in our Figs. 7 and 9 to Jahn et al. (2015)'s non-gridded d13DIC observation in Figs. 5 and 7).

For Figure 4 we also re-gridded d13POC observations. In the revised text we specify: "For the model-observation comparison, we first grid the observed $\delta^{13}\text{C}_{\text{POC}}$ and $\delta^{13}\text{C}_{\text{DIC}}$ horizontally onto a $1^\circ \times 1^\circ$ grid ..."

Comment 24 P9, l207: Why do you start with POC here instead of DIC? I expect more readers will be familiar with the d13C.DIC distributions. Also, this Section is about simulated isotopic signature, maybe add this to the title?

Response: The vertical gradient of d13DIC mainly results from the biological fractionation. A number of discussions on d13DIC depend on the results and discussion about d13POC. Thus, it seems to us more logical to begin with the discussion on d13POC.

This section includes both simulation results and comparison to observations. Thus, we keep the original section title.

Comment 25 P9, l221-224: I think you should not only refer to Appendix B when it comes to what the model simulates, but also refer to Appendix B when it comes to how that compares to observational estimates. You could summarize the most important results of Appendix B here. Also, Fig B1 is for the Popp results? Note that for clarity if you refer to model results (like also in e.g. Fig D3) it is good to say which model run you mean.

Response: In the revised manuscript, Appendix B is now referred to at the beginning of Section 3 to note the performance of the model.

Phytoplankton growth rate in Fig B1 is identical for Popp and Laws because it is for the total carbon and is not affected by ^{13}C . This is now noted in the revised Appendix.

Comment 26 P10, l242: I understand you want to make an evaluation of the model performance here of $\text{CO}_2(\text{aq})$ around 45S in Fig 4g, but why choose this particular dataset - aren't there more recent data with better coverage available?

Response: Thank you for the question. We chose this particular dataset because it provides contemporaneous measurements of both d13POC and $\text{CO}_2(\text{aq})$. This is clarified in the revised manuscript.

Comment 27 P10, l245: Hist_Laws captures more than Hist_Popp, but not all and more importantly offset by a few permil. If this offset is really very constant, wouldn't it be possible to adjust the Laws et al parameterization. One could evaluate the offset needed and see if after that offset Popp is still superior?

Response: As the optimisation of the Laws parameterisation is beyond the scope of this study, we use a back-of-the-envelope calculation to refit the slope and intercept of ϵ_p^{Laws} (Eq. 7).

Because $\alpha_{\text{Phy} \leftarrow \text{DIC}}$, $\alpha_{\text{aq} \leftarrow \text{g}}$ and $\alpha_{\text{DIC} \leftarrow \text{g}}$ are close to unity, $\epsilon_{\text{Phy} \leftarrow \text{DIC}} \approx \delta^{13}\text{C}_{\text{Phy}} - \delta^{13}\text{C}_{\text{DIC}}$ and $\epsilon_{\text{Phy} \leftarrow \text{DIC}} \approx \epsilon_p + \epsilon_{\text{aq} \leftarrow \text{DIC}} \approx \epsilon_p + \epsilon_{\text{aq} \leftarrow \text{g}} - \epsilon_{\text{DIC} \leftarrow \text{g}}$. As $\delta^{13}\text{C}_{\text{Phy}} \approx \delta^{13}\text{C}_{\text{POC}}$, $\delta^{13}\text{C}_{\text{POC}}$ can be approximated as

$$\delta^{13}\text{C}_{\text{POC}} \approx \epsilon_p + \epsilon_{\text{aq} \leftarrow \text{g}} - \epsilon_{\text{DIC} \leftarrow \text{g}} + \delta^{13}\text{C}_{\text{DIC}}. \quad (\text{R1})$$

As sea water temperature (determining $\epsilon_{\text{DIC} \leftarrow \text{g}}$), $\text{CO}_2(\text{aq})$ and growth rate are independent of the choice of ϵ_p , and the surface $\text{d}13\text{DIC}$ is only marginally affected by the choice of ϵ_p (see Section 3.2 and Figs. 5a and 5b), we can approximate ϵ_p^{Laws} using existing monthly output of the above model variables. The surface $\text{d}13\text{POC}$ obtained with Eq. (R1) ($r=0.74$, $\text{NRMSE}=2.8$, Figs. R1a-R1c) is indeed very close to the simulated $\text{d}13\text{POC}$ ($r=0.71$, $\text{NRMSR}=2.5$, Figs. 4b, 4d and 4f).

$$\epsilon_p^{\text{Laws}} = 68.3 \frac{\mu}{\text{CO}_2(\text{aq})} - 24.7, \quad (7)$$

$$\epsilon_p^{\text{Laws.V1}} = 68.3 \frac{\mu}{\text{CO}_2(\text{aq})} - 20, \quad (\text{R2})$$

$$\epsilon_p^{\text{Laws.V2}} = 68.3 \frac{\mu}{\text{CO}_2(\text{aq})} - 16.7, \quad (\text{R3})$$

$$\epsilon_p^{\text{Laws.V3}} = 100 \frac{\mu}{\text{CO}_2(\text{aq})} - 20. \quad (\text{R4})$$

As the intercept increases (comparing Eq. 7, R2 and R3), $\text{d}13\text{POC}$ generally increases (Figs. R1a-R1i) and accordingly NRMSE decreases from 2.8 for Eq. (7) to 1.4 for Eq. (R2) and 0.75 for Eq. (R3), while the spatial relation coefficient remains unchanged.

As the slope increases (comparing Eq. R2 and R4), the $\text{d}13\text{POC}$ difference between low and high latitudes increases (Figs. R1d-R1f, R1j-R1l) and NRMSE slightly decrease from 1.4 to 1.3. However, $\text{d}13\text{POC}$ in the low latitude of the Atlantic shows too high variability compared to the observation. Accordingly, the spatial correlation coefficient decreases from 0.74 to 0.66.

In summary, a systematic refitting of the Laws parameterisation could be an interesting step to improve the model performance and might be considered in our future work.

Comment 28 P11, Fig. 4: Are there uncertainty estimates available for the obs data, or would it be possible to at least shade an estimate of the uncertainty?

Response: The uncertainties for observations are added.

Comment 29 P12, l281: this seems a bit repetitive, wouldn't a too steep vertical gradient always lead too too low deep $\text{d}13\text{C}_{\text{DIC}}$ if surface $\text{d}13\text{C}_{\text{DIC}}$ is reasonable? This could be rephrased.




Figure R1: As Figure 4b, 4d and 4f in the manuscript, but for $\delta^{13}\text{C}_{\text{POC}}$ estimated for ϵ_p^{Laws} (Eq. 7) using Eq. (R1) (a-c). (d-f), (g-i) and (j-l): As (a-c), but for $\epsilon_p^{\text{Laws-V1}}$ (Eq. R2), $\epsilon_p^{\text{Laws-V2}}$ (Eq. R3) and $\epsilon_p^{\text{Laws-V3}}$ (Eq. R4), respectively.

Response: This sentence is rephrased: “Hist_Laws generally shows too strong vertical gradients of $\delta^{13}\text{C}_{\text{DIC}}$ and therefore too low $\delta^{13}\text{C}_{\text{DIC}}$ values in the ocean interior”.

Comment 30 P13, l284-285: this is a very important transition point for the reader, where you decide to mainly focus on Popp from now on. This could be mentioned earlier (or even in the abstract), or denoted by a new section here. For example, make a 3.2.1 and a 3.2.2 section.

Response: This transition does not affect the statements in the earlier text because before this transition the performance of both Hist_Popp and Hist_Laws is discussed.

We now divide this section into subsections: Section 3.2.1 on the comparison between Hist_Popp and Hist_Laws and to observed $\delta^{13}\text{C}_{\text{DIC}}$, Section 3.2.2 on the source of surface $\delta^{13}\text{C}_{\text{DIC}}$ biases in Hist_Popp and Section 3.2.3 on the source of $\delta^{13}\text{C}_{\text{DIC}}$ biases in the interior ocean of Hist_Popp.

Comment 31 P13, l286: I got slightly lost here. Maybe explain the reader how this $\delta^{13}\text{C}_{\text{DIC}}$ comparison is different from the one in Fig. 7. It could help to make Fig 8a,b,c one figure, and present the $\delta^{13}\text{C}_{\text{bio}}$ and $\delta^{13}\text{C}_{\text{resi}}$ and the net air-sea CO_2 flux separately. Regarding $\delta^{13}\text{C}_{\text{bio}}$ and $\delta^{13}\text{C}_{\text{resi}}$ I think it helps with comparison to earlier studies if you show the absolute values and not only the model-obs difference.

Response: We rephrase “interior-ocean $\delta^{13}\text{C}_{\text{DIC}}$ ” to “zonal-mean $\delta^{13}\text{C}_{\text{DIC}}$ in the Atlantic, Pacific and Indian Ocean” to make a distinction from Fig. 7.

Figure 8 is divided into two as suggested. The absolute values of $\delta^{13}\text{C}_{\text{bio}}$ and $\delta^{13}\text{C}_{\text{resi}}$ are added.

Comment 32 P13, l294: Why do you use Δ_{photo} and not ϵ_{p} like before here?

Response: Because they are different variables: Δ_{photo} equals $(\alpha_{\text{Phy}\leftarrow\text{DIC}} - 1) \times 10^3$ (see Eq. 8), whereas $\epsilon_{\text{p}} = (\alpha_{\text{Phy}\leftarrow\text{aq}} - 1) \times 10^3$.

Comment 33 P13, l295: Do you use the same R_C:P for the components calculations of both obs and model?

Response: Yes, the same R_C:P is used. We specify this in the revised manuscript.

Comment 34 P13, l296: I think it would be appropriate here to remind the reader that ϵ_{p} actually varies, referring e.g. to Fig. 3b,d. Also, what model MO DIC, PO4 and d13C were used? Is the R_C:P of 122 the one used in the model for consistency?

Response: Sentences are added regarding the spatial variation of Δ_{photo} , the mean values of DIC, PO4 and d13DIC.

Yes, R_C:P of 122 is used in the model for consistency, we specify this in the revised manuscript.

Comment 35 P15, l302-304: Their MO values should be included here as done for the obs.

Response: The MO values for the model are added.

Comment 36 P17, l312-314: This is some information on the physical model performance that I think should be introduced earlier and possibly in an own subsection under Sect. 2.

Response: Additional figures on physical model performance are added in the revised Appendix.

Comment 37 p17, l324-326: One could quantify the effect on d13C_DIC by analysing the bio and resi components for the PI run instead of the or in addition to the hist runs.

Response: The purpose of decomposing d13C_DIC into d13_bio and d13_resi components is to better understand the sources of model-observation differences. There is no observational information to evaluate d13_bio and d13_resi components for the pre-industrial periods. Thus we focus on the discussion and model-data comparison for the late 20th century.

Comment 38 P17, l329: Here you continue to Fig. 9- the reader could use a bit more guidance here: what are you going to present and discuss here in this section/paragraph? Why do you go away from showing d13Cbio and d13Cresi?

Response: This transition is now marked by a new subsection on the source of $\delta^{13}\text{C}_{\text{DIC}}$ biases in the interior ocean of Hist_Popp.

Sect. 4 Oceanic ^{13}C Suess effect

Comment 39 p18, l 367-369: Clarify here why this is important for your discussion of the Suess effect. You could also add a comment here that even though the model does well simulating the total anthropogenic C uptake, locally air-sea exchange fluxes deviate from obs (Fig. 8f).

Response: Our motivation to discuss Suess effect is stated in the previous paragraph: “The oceanic ^{13}C Suess effect, could serve as benchmark for ocean models to evaluate the uptake and re-distribution of the anthropogenic CO_2 emissions in the ocean”.

A statement on the local biases in air-sea flux is added.

Comment 40 P19, l385: This section is quite long and heavy, and I have to admit I did not follow all of it. I would suggest to cut it up in different subsections, which can discuss the different aspects which you have investigated. Start early in Sect. 4.2 with why it is relevant to re-evaluate the approach by E17 (e.g., is it often used?). Large parts of the text also feel like a methods section ? could more of this section be moved to a supplement/appendix C, such that more focus on the results can be given here?

Response: See our reply to Comment 3.

Comment 41 P20, l393: You have only discussed total global anthropogenic C uptake, and you have compared to E17 at depth (Sect. 4.1) - now you are going to explore the E17 underestimation after concluding that your model produces similar results to E17? This is somewhat of a confusing step.

Response: This comparison is for the qualitative behaviour of ^{13}C Suess effect, whereas E17's underestimation only has some quantitative impact. It is expected that our model produces similar pattern as E17. This is because E17's Suess effect estimate is proportional to pCFC-12. The oceanic distributions of CFC-12 and ^{13}C Suess effect are both dominated by the uptake of the atmospheric signal and the subsequent transport in the ocean, and these processes are well simulated in our model.

The above discussion is incorporated in the revised manuscript.

Comment 42 p26, l518: How does this compare to observational estimates (e.g. Young et al., 2013)?

Response: See the last paragraph of our reply to Comment 1.

Sect. 5 Summary and conclusions

Comment 43 p27, l565: One should add a short paragraph here summarizing your $\delta^{13}\text{C}_{\text{bio}}$ and $\delta^{13}\text{C}_{\text{resi}}$ component analysis for the hist_Popp run.

Response: We add sentences on findings based on this decomposition: “Our model slightly overestimates surface $\delta^{13}\text{C}_{\text{DIC}}$. By decomposing $\delta^{13}\text{C}_{\text{DIC}}$ into a biological component and a residual component, we find the overestimation in the high latitude ocean is dominated by biases in the biological component. ”.

Comment 44 P28, l582: the Popp et al., 1989 parameterization has a satisfactory performance for the PI and historical times. I would agree this encourages reliability for paleoclimatic simulations, but I think some more critical remarks are in place (which should come before Sect. 5). E.g., in the past ϵ_p was possibly different due to different ecosystem structures or other influences, Redfield ratios could have changed (Ödalen et al., 2020).

Response: We add a text to note the potential limitations of ϵ_p in long-term simulations for the past climate.

Past changes of C/P ratios discussed in Ödalen et al. (2020) affect the production of ^{13}C -depleted organic matter (according to Eq. 8) and its export to the deep ocean. Changes of C/P ratios do not directly affect parameterisation of the biological fractionation, thus no discussion is added on C/P ratios.

Technical corrections

Comment 45 p6, l146: Zeebe

Comment 46 P17, l335: positive biases between 1000 and 3000m

Comment 47 p19, l387: the ^{13}C Suess effect

Comment 48 p20, l391: at 200 m depth

Comment 49 p27, l561: again yields slightly better agreement

Comment 50 p28, l574: Mode Water and explains

Comment 51 p26, l528: Fig. D7h?

Response: The technical corrections in Comments 45 - 51 are implemented in the revised manuscript.

References

- J. E. Dentith, R. F. Ivanovic, L. J. Gregoire, J. C. Tindall, and L. F. Robinson. Simulating stable carbon isotopes in the ocean component of the famous general circulation model with moses1 (xoavi). *Geoscientific Model Development*, 13(8):3529–3552, 2020. doi: 10.5194/gmd-13-3529-2020.
- M. Eide, A. Olsen, U. S. Ninnemann, and T. Eldevik. A global estimate of the full oceanic ^{13}C suess effect since the preindustrial. *Global Biogeochemical Cycles*, 31(3):492–514, 2017a. doi: 10.1002/2016GB005472.
- M. Eide, A. Olsen, U. S. Ninnemann, and T. Johannessen. A global ocean climatology of preindustrial and modern ocean $\delta^{13}\text{C}$. *Global Biogeochemical Cycles*, 31(3):515–534, 2017b. doi: 10.1002/2016GB005473.
- A. Jahn, K. Lindsay, X. Giraud, N. Gruber, B. L. Otto-Bliesner, Z. Liu, and E. C. Brady. Carbon isotopes in the ocean model of the community earth system model (cesm1). *Geoscientific Model Development*, 8(8):2419–2434, 2015. doi: 10.5194/gmd-8-2419-2015.
- M. Ödalen, J. Nycander, A. Ridgwell, K. I. C. Oliver, C. D. Peterson, and J. Nilsson. Variable C/P composition of organic production and its effect on ocean carbon storage in glacial-like model simulations. *Biogeosciences*, 17(8):2219–2244, 2020. doi: 10.5194/bg-17-2219-2020.
- J. C. Orr, R. G. Najjar, O. Aumont, L. Bopp, J. L. Bullister, G. Danabasoglu, S. C. Doney, J. P. Dunne, J.-C. Dutay, H. Graven, S. M. Griffies, J. G. John, F. Joos, I. Levin, K. Lindsay, R. J. Matear, G. A. McKinley, A. Mouchet, A. Oschlies, A. Romanou, R. Schlitzer, A. Tagliabue, T. Tanhua, and A. Yool. Biogeochemical protocols and diagnostics for the cmip6 ocean model intercomparison project (omip). *Geoscientific Model Development*, 10(6):2169–2199, 2017. doi: 10.5194/gmd-10-2169-2017.
- A. Schmittner, N. Gruber, A. C. Mix, R. M. Key, A. Tagliabue, and T. K. Westberry. Biology and air-sea gas exchange controls on the distribution of carbon isotope ratios ($\delta^{13}\text{C}$) in the ocean. *Biogeosciences*, 10(9):5793–5816, 2013. doi: 10.5194/bg-10-5793-2013.
- J. N. Young, J. Bruggeman, R. E. M. Rickaby, J. Erez, and M. Conte. Evidence for changes in carbon isotopic fractionation by phytoplankton between 1960 and 2010. *Global Biogeochemical Cycles*, 27(2):505–515, 2013. doi: 10.1002/gbc.20045.