

# Contrasting vertical distributions of recent planktic foraminifera off Indonesia during the southeast monsoon: implications for paleoceanographic reconstructions

Raúl Tapia<sup>1</sup>, Sze Ling Ho<sup>1</sup>, Hui-Yu Wang<sup>1</sup>, Jeroen Groeneveld<sup>2</sup>, and Mahyar Mohtadi<sup>3</sup>

<sup>1</sup>Institute of Oceanography, National Taiwan University, No. 1, Sec. 4, Roosevelt Road 10617 Taipei, Taiwan.

<sup>2</sup>Department of Geology, Hamburg University, D-20146 Hamburg, Germany

<sup>3</sup>MARUM – Center for Marine Environmental Sciences, University of Bremen, D-28334 Bremen, Germany

**Correspondence:** Raúl Tapia (raultapia@ntu.edu.tw), Sze Ling Ho (slingho@ntu.edu.tw)

**Abstract.** Planktic foraminifera are widely used in palaeoceanographic and paleoclimatic studies. The accuracy of such reconstructions depends on our understanding of the organisms' ecology. Here we report on field observations of planktic foraminiferal abundances ( $>150\ \mu\text{m}$ ) from 5 depth intervals between 0-500 m water depth at 37 sites in the eastern tropical Indian Ocean. The total planktic foraminiferal assemblage here comprises 29 morphospecies; with 11 morphospecies accounting for 90% of the total assemblage. Both species composition and dominance in the net samples are broadly consistent with those the published data from the corresponding surface sediments.

The abundance and vertical distribution of planktic foraminifera are low offshore west Sumatra, and increase towards offshore south Java and the Lesser Sunda Islands (LSI). Average living depth of *Trilobatus trilobus*, *Globigerinoides ruber*, and *Globigerina bulloides* increases eastward, while that of *Neogloboquadrina dutertrei*, *Pulleniatina obliquiloculata*, and *Globorotalia menardii* remains constant. We interpret the overall zonal and vertical distribution patterns in planktic foraminiferal abundances as a response to the contrasting upper water column conditions during the southeast monsoon, i.e., oligotrophic and stratified offshore Sumatra (non-upwelling) vs. eutrophic and well-mixed offshore Java-LSI (upwelling). Overall, the inferred habitat depths of selected planktic foraminifera species shows a good agreement with those from sediment trap samples and from surface sediments off Sumatra, but not with those from surface sediments off Java-LSI. The discrepancy might stem from the different temporal coverage of these sample types. Our findings highlight the need to consider how foraminiferal assemblages and ecology vary on shorter timescales, i.e. from “snapshots” of the water column captured by plankton net to seasonal and interannual variability as recorded in sediment traps and how these changes are transferred and preserved in deep-sea sediments.

## 1 Introduction

Planktic foraminifera's diversity, community composition, population dynamics as well as their shell chemistry are sensitive to hydrographic parameters (e.g., temperature, salinity, food availability) of the upper ocean (Bemis et al., 1998; Fairbanks

et al., 1980; Katz et al., 2010). Consequently, their assemblage composition and the geochemical signature of their shells are routinely used to reconstruct past ocean conditions (e.g. Ding et al., 2006, 2013; Mohtadi et al., 2017; Steinke et al., 2014).

25 Since the early observations from Murray (1897), many studies have evaluated the relationship between seawater temperature and planktic foraminiferal distribution, and found increasing dominance of cold species with increasing latitudes (Bauer, 1976; Bé and Hamlin, 1967; Eguchi et al., 1999). This relationship with seawater temperature allowed the development of the first comprehensive mapping of past sea surface temperature (SST) (CLIMAP Project Members, 1976). Subsequent assemblage studies have attempted to improve the accuracy of past sea surface temperature (SST) reconstruction via the transfer function approach (e.g., Imbrie and Kipp, 1971; Ortiz and Mix, 1997). Although seawater temperature has a large influence on the global distribution of planktic foraminifera (Kucera, 2007, 2009), other parameters such as salinity, oxygen content, food availability, turbidity, and upwelling intensity can also exert a strong control on the abundance, community structure, and vertical distribution of planktic foraminifera at regional scales (Anderson et al., 1979; Davis et al., 2021; Field, 2004; Lessa et al., 2020; Rebotim et al., 2017; Schiebel and Hemleben, 2017; Tolderlund et al., 1971).

35 In the last decades, geochemical proxies using the calcite tests of foraminifera (e.g., isotope  $\delta^{18}\text{O}$  and trace elements such as Mg/Ca ratios ) have become a popular powerful tool for the reconstruction of past ocean conditions (Katz et al., 2010; Lea, 2014; Schiebel et al., 2018). The robustness of paleoclimate reconstructions derived from foraminiferal calcite proxies is as good as our understanding of the multiple fields that planktic foraminiferal biology encompasses (Schiebel et al., 2018). For example, reconstructing past changes in the water column structure using multiple planktic foraminiferal species (e.g., surface vs. deep dweller) requires understanding of the ecology of the selected species, such as seasonality, habitat depth, and food requirements (Kemle-von Mücke and Oberhänsli, 1999; Kucera, 2007; Marchant et al., 2004; Schiebel et al., 2017). One common way of inferring planktic foraminifera habitat depth is by comparing the reconstructed parameters (typically Mg/Ca-SST) from surface sediments with instrumental data or climatological products (e.g., World Ocean Atlas) (Groeneveld and Chiessi, 2011; Hollstein et al., 2017; Mohtadi et al., 2011; Steinke et al., 2014; Tapia et al., 2015). In this approach, habitat depth is defined as the water depth at which the reconstructed Mg/Ca-temperature or seawater  $\delta^{18}\text{O}$  value show the closest match with the instrumental data or climatological product. However, this approach may be associated with uncertainties arising from a myriad of processes during the settling, deposition, and burial that may lead to varying degrees of proxy signal alteration (Regenberg et al., 2014), selected proxy calibrations, and the instrumental database used for ground truthing the proxy. Furthermore, additional uncertainty stems from the selected proxy calibrations and the instrumental database used for comparison with the proxy. “Snapshots” from the water column where planktic foraminifera dwell are thus providing additional information to improve proxy understanding.

50 The marginal seas of the Indonesian Archipelago are of great climatic importance. Here, the Indonesian Throughflow (ITF) connects the upper water masses of the Pacific and Indian oceans, exerting a strong effect on the salinity and heat exchange between these basins (Gordon and Fine, 1996; Gordon, 2005; Tillinger, 2011). This feature has fueled the scientific interest to better understand the role that heat exchange plays in climatic modulation on several time scales (Smith et al., 2020; Sprintall and Révelard, 2014). Consequently, a growing number of oceanographic reconstructions based on foraminiferal calcite have been published over the last decade to shed light on the changes in the regional oceanographic processes and climate (Dang et

al., 2020; Ding et al., 2013; Fan et al., 2018; Karas et al., 2011; Mohtadi et al., 2014, 2017; Pang et al., 2021; Steinke et al., 2014; Wang et al., 2018). However, our understanding of the ecology of planktic foraminifera in this region, especially their vertical distribution, relies largely on indirect inferences based on surface sediments and sediment trap samples (Ding et al., 2006; Mohtadi et al., 2007, 2009, 2011). Thus far there is only one field observation that is focused on the standing stock of planktic foraminifera in surface ocean sampled using plankton net (Ujiié, 1968), but this study did not investigate the vertical distribution of planktic foraminifera in the water column. To fill this gap, here we present depth stratified (0–500 m) plankton net data from the Indonesian marginal seas off Sumatra, Java and the Lesser Sunda Islands (LSI) (Fig. 1). The main goal of this study is to shed light on the spatial distribution of planktic foraminifera during the southeast (SE) monsoon, on the relationship between foraminiferal abundance and environmental parameters, and how these findings compare with sediment trap and core top assemblage data to further improve our understanding of foraminifera-based proxy reconstructions in this region.

## 1.1 Study area

Modern oceanography off Indonesia is strongly modulated by seasonal monsoons. During the SE monsoon from April to October, the southeasterly winds from Australia induce Ekman pumping that generates upwelling along the coast of southern Sumatra, Java and the LSI. The peak of upwelling-favorable winds occurs at the southernmost coast of Sumatra ( $\sim 105^\circ$  E) during July-August, but the upwelling center moves northwestward reaching as far as  $100^\circ$  E and  $2^\circ$  S in October (Susanto et al., 2001). Seasonal upwelling results in increased chlorophyll-*a* concentrations and reduced thickness of the depth of the mixed layer ( $<20$  m) south of  $\sim 4^\circ$  S (Fig. 1; Table A1 and A2). Furthermore, seasonal upwelling leads to a  $\sim 2^\circ$  C decrease in SST in comparison to the non-upwelling season and a contrast in SST of  $\sim 3^\circ$  C between the northern and southern parts of the study area. During the northwest (NW) monsoon from late October to early April, the wind direction is reversed, resulting in downwelling, lower chlorophyll-*a* concentrations, thicker mixed layer, higher SST south of  $\sim 4^\circ$  S and a more uniform SST distribution offshore southern Sumatra, Java and the LSI (Muskananfolia et al., 2021; Qu et al., 2005; Susanto et al., 2001, 2006). At the interannual time-scale, the SST variability offshore southern Sumatra, Java and the LSI can be larger than  $4^\circ$  C, showing the influence of climatic modes such as El Niño-Southern Oscillation (ENSO) or the Indian Ocean Dipole (IOD) (Qu et al., 2005). During years of the El Niño (La Niña) phase of the ENSO and the positive (negative) phase of the IOD, intensified (weakened) southeasterly winds result in stronger (weaker) coastal upwelling, leading to abnormally low (high) SSTs offshore southern Sumatra and Java (Du and Zhang, 2015; Mohtadi et al., 2011; Qu et al., 2005).

## 2 Materials and methods

Multinet samples were collected between August and September 2005 during FS Sonne cruise SO-184 (Hebbeln and cruise participants, 2006). The 47 sampling sites are grouped into seven land-sea transects off Sumatra ( $n = 4$ ), Java ( $n = 2$ ) and the Lesser Sunda Islands (LSI) ( $n = 1$ ) (Fig. 1; details in Table A1 and A2).

## 2.1 Collection of hydrographic data and foraminiferal samples

The characterization of the physicochemical properties of the water column (i.e., temperature, salinity, nutrients, oxygen, and in situ chlorophyll-*a*) was obtained from CTD and water samples collected using a rosette water sampler equipped with 24 Niskin bottles (10-liter volume each) and a Seabird SBE911 probe (Hebbeln and cruise participants, 2006) across 45 water stations (see Table A1 and A2). These hydrographic data are available in the cruise report of SO184 (Hebbeln and cruise participants, 2006). Briefly, the dissolved oxygen was determined using an automated titrator (Titroline alpha), which is controlled by a redox electrode and a color agent (starch solution), on a self-constructed titration board for ship cruises according to the WOCE protocol (Hebbeln and cruise participants, 2006). Chlorophyll-*a* measurement were performed using a SFM25 spectrofluorometer (KONTRON). The measurement were done with 1 cm cuvettes at a 435 nm excitation wavelength and a 667 nm emission wavelength (Hebbeln and cruise participants, 2006).

The plankton samples ( $n = 37$ ; Table A1 and A2) were obtained using a MultiNet sampler (Hydro-Bios, Kiel, Germany). The gear is comprised of five individual 64  $\mu\text{m}$  nets with an opening of 0.25  $\text{m}^2$ . The nets were lowered to a water depth of 500 m and vertically towed with a maximum winch speed of 0.2  $\text{m s}^{-1}$  along five depth intervals of 500 to 200 m, 200 to 100 m, 100 to 50 m, 50 to 25 and 25 to 0 m depth. At the end of each depth interval the MultiNet sampler was stopped to open the next net automatically closing the previous net. Once on board the samples were poisoned with 1 mL saturated HgCl solution and stored at 4 °C (Hebbeln and cruise participants, 2006). The volume of water passed through the net opening was calculated as the product of the height of the towed intervals and the area of the net opening.

## 2.2 Foraminiferal identification and census count

The taxonomic identification of the planktic foraminifera was based on Parker (1962), Kennett and Srinivasan (1983) and Hemleben et al. (1989). We used the same taxonomic approach as in previous studies based on surface sediments (Mohtadi et al., 2007) and sediment trap (Mohtadi et al., 2009; 2011). The only exceptions are for *Globigerinoides elongatus* and *Trilobatus trilobus*, as the name of these species have been updated recently by Aurahs et al. (2009) and Spezzaferi et al. (2015), respectively. Here the species *Globigerinoides ruber* (white) can occur in two morphotypes (Mohtadi et al., 2009, 2011), namely *G. ruber sensu stricto* (*s.s.*) *sensu stricto* (*s.s.*) and *sensu lato* (*s.l.*) *sensu lato* (*s.l.*). As *G. ruber* (*s.l.*) (*s.l.*) is now identified as a separate species, *Globigerinoides elongatus* (Aurahs et al., 2009), we have included this distinction. The distinction was done according to the approach of Wang (2000); specimens with spherical chambers sitting on the previous suture and high arched primary aperture were classified as *G. ruber* (white), meanwhile, more compressed organisms with subspherical chambers and low arched primary aperture were classified as *G. elongatus*. In the case of the genus *Trilobatus*, we integrated the sacculifer-morphotype, i.e., individuals with a sac-like final chamber, into the counts of *T. trilobus* (individuals with a regular, globular terminal chamber). The distinction of *Neogloboquadrina dutertrei* from *Neogloboquadrina incompta* was made according to the approach of Mohtadi et al. (2009), based on the presence of an umbilical tooth, and the occurrence of more than four chambers per whorl (Parker, 1962). The samples were separated into four size classes (>500, 500–355,

355–250, and 250–150  $\mu\text{m}$ ) and stored in foraminiferal microslides. Here we report the foraminiferal abundances  $>150 \mu\text{m}$  as  
120 individuals per volume of seawater passing through the opening of the net ( $\text{ind. m}^{-3}$ ).

Vertical habitat preferences were estimated using the total abundance of each species in each vertical profile. As we did not  
discriminate between living and dead specimens, this approach may have led to an overestimation of the habitat depth, for  
instance when specimens were found in deep nets. Therefore, we refrain from interpreting individual multinet deployments,  
which may be prone to the aforementioned bias. Instead, we base our interpretations on the median value of transects, each  
125 consisting of at least three multinet deployments. The proportion of dead specimens, if any, likely varies across stations, thus  
averaging over stations yields estimates that are less prone to single-site bias. This approach might also mitigate potential biases  
caused by patchy occurrences of planktic foraminifera in the water column, in both lateral and vertical directions (Meillard  
et al., 2019). To facilitate comparison with previous studies and to avoid bias due to different sampling strategies, we used  
Average Living Depth (ALD) (Jorissen et al., 1995) to discuss the vertical habitat of planktic foraminifera. ALD is defined as:

130 
$$ALD = \frac{\sum n_i \cdot D_i}{N_i}$$

Where  $n_i$  is the number of specimens in the interval  $i$  of a particular species;  $D_i$  is the midpoint of the sampled interval  $i$ ;  
and  $N$  is the total number of individuals for all the depth levels of that particular species. For the calculation of the ALD, we  
followed the approach suggested by Rebotim et al. (2017) by considering only stations with at least five individuals of a given  
species.

135 The 95% confidence interval of the species ALD for upwelling vs non-upwelling region (Table 1) was estimated using non-  
parametric randomization tests i.e., bootstrap (Manly, 1997), as the distributions of ALDs of either region do not meet normal  
distributions required for parametric tests due to small sample sizes. The upwelling region consists of transects 5–7, while  
the non-upwelling region consists of transects 1–3. We permuted the ALDs of each species for upwelling and non-upwelling  
region, respectively, simulating the differences in mean ALDs between these permuted sites with 999 replicates. We then sorted  
140 the sequence of 999 differences in mean ALDs, which approximated all possible outcomes of the lack of differences in mean  
ALDs between upwelling vs. non-upwelling sites. We determined the probability of the observed between-region difference  
in mean ALDs as 1-quantile of the observed estimate in the sorted sequence. The probability (P-value)  $<0.05$  represents  
significant differences in mean ALDs between regions. Statistical analysis was performed in R language (R core Team, 2020),  
code available upon request.

## 145 3 Results

### 3.1 Hydrological conditions

Across the sampled depths (0–500 m) of the plankton nets, the water temperature ranges from  $\sim 8$  to  $\sim 30$   $^{\circ}\text{C}$ , the salinity  
from 33.5 to 35.1  $\text{psu}$ , the in situ chl- $a$  from  $\sim 0.2$  to  $\sim 0.7 \text{ mg m}^{-3}$  and the oxygen concentration from  $1.2$ – $\sim 5 \text{ ml L}^{-1}$  (Fig. 2  
a-c). Sea surface temperature shows a strong zonal contrast, with colder conditions off Java-LSI and warmer conditions off

150 Sumatra (Fig. 2a). This zonal pattern remains across all the sampled depths from the surface to a water depth of 500 m (Fig. A1). Similarly, salinity also shows a strong zonal contrast, as fresher conditions dominate the upper 50 m of the water column off Sumatra while more saline conditions dominate the upper ocean off Java (<50 m) (Fig. 2b). This pattern reverses at depths >50 m, with more saline condition off Sumatra than Java and the LSI.

In situ chl-*a* for 0–25 m water depth is on average  $\sim 0.65 \text{ mg m}^{-3}$ , with no zonal patterns across the study area (Table A1 and Fig. 2c). Lower-than-average chl-*a* values can be found at two oceanic stations from transect 3 off Sumatra, that is, sites GeoB10003 ( $0.28 \text{ mg m}^{-3}$ ) and GeoB10007 ( $0.34 \text{ mg m}^{-3}$ ) (Table A1). Despite the absence of a zonal divide in surface chl-*a*, at depths >50 m the vertical distribution differs between Sumatra and further east ( $\sim 105^\circ \text{ E}$ ; Fig. 2c). Off Sumatra (< $105^\circ \text{ E}$ ), chl-*a* is mainly restricted to the upper 100 m of the water column, while high chl-*a* values  $> 0.3 \text{ mg m}^{-3}$  ( $>105^\circ \text{ E}$ ) can reach as deep as 500 m at the easternmost study area, i.e., Transect 7 around Sumba Island (Fig. 2c). The distribution of oxygen content across the water column matches the distribution of in situ chl-*a*, i.e., high dissolved oxygen concentrations ( $>2 \text{ ml L}^{-1}$ ) are restricted to the upper 100 m of the water column off Sumatra, while high dissolved oxygen concentrations off Java and particularly around Sumba (Transect 7) can reach as deep as 500 m (Fig. 2d).

The thermal mixed layer depth ( $\text{MLD}_{\text{TEMP}}$ ) for this region, defined as the depth where the temperature is  $>0.8^\circ \text{ C}$  colder than the SST (Kara et al., 2000; Qu et al., 2005), roughly marks the top of the thermocline depth and ranges from  $\sim 13$  to  $\sim 91$  m (Table A1, A2 and Fig. 2a). The  $\text{MLD}_{\text{TEMP}}$  varies zonally; off Sumatra the  $\text{MLD}_{\text{TEMP}}$  ( $\sim 74$  m) is on average twice the thickness of that off Java-LSI ( $\sim 33$  m) (see Table A1 and A2). The barrier layer separates the well mixed upper ocean from the thermocline. Here, the barrier layer is defined as the  $\text{MLD}_{\text{TEMP}}$  minus the mixed layer depth calculated using density, namely  $\text{MLD}_{\text{DEN}}$  (Qu and Meyers, 2005). It ranges between 0 to  $\sim 72$  meters of thickness and follows a similar spatial distribution as the  $\text{MLD}_{\text{TEMP}}$ , with an average thickness of  $\sim 45$  m off Sumatra and  $\sim 2$  m off Java-LSI (Table A1 and A2). The upper water column stratification,  $\text{SI}_{0-200}$  is defined as the temperature difference between the sea surface and 200 m (Somavilla et al., 2017). The  $\text{SI}_{0-200}$  values are higher off Sumatra than off Java and the LSI, indicating more stratified upper water column off Sumatra (Table A1 and A2). These observations suggest two contrasting hydrological conditions in the study area, with strong (weak) water column stratification, thick (thin) mixed layer and barrier layer, and low (high) subsurface water entrainment towards the surface off Sumatra (off Java-LSI), in agreement with the observed geographical extension of the coastal upwelling (Fig. 1c) during the SE monsoon (Susanto and Marra, 2005; Susanto et al., 2001).

Multivariate analysis (non-metric multidimensional scaling) performed on the in situ hydrographic data obtained during the cruise provides further evidence that the study area consists of three hydrologically distinct regions (Fig. 3), constituting of transects 1–3, 4, and 5–7, respectively. The cluster off Java and the LSI that encompasses transects 5 to 7 is characterized by lower SST ( $<29^\circ \text{ C}$ ), saltier surface ocean, shallow mixed layer, and chl-*a* with a larger vertical dispersion, i.e., conditions typical of seasonal upwelling during the southeast monsoon. Meanwhile, the cluster off Sumatra that encompasses transects 1 to 3 is characterized by higher SST ( $>29^\circ \text{ C}$ ), fresher sea surface ocean, deep mixed layer, and chl-*a* with low vertical dispersion. A third cluster consists solely of Transect 4, which reflects transitional conditions as here we can observe a large dispersion in the values of the parameters analyzed (see Table A1 and A2) and its location coincides with the northernmost extent of upwelling-favorable winds (see section 1.1). Based on the result of multivariate analysis and hydrographic data, we therefore

185 categorize transects. The multivariate analysis performed on hydrographic data separates the sites into two main groups, i.e.,  
transects 1-3 in non-upwelling sites sector and transects 5-7 as upwelling sites. in upwelling sector.

### 3.1.1 Planktic foraminiferal assemblage and absolute abundance

The total composition of the planktic foraminiferal species comprises 29 morphospecies. Eleven morphospecies, namely *Globigerina bulloides*, *Globigerinella calida*, *Globigerinina glutinata*, *Globigerinoides ruber* (white), *Trilobatus trilobus*, *Globorotalia menardii*, *Neogloboquadrina dutertrei*, *Globorotalia hirsuta*, *Globigerina falconensis*, *Pulleniatina obliquiloculata*, and  
190 *Globigerinella siphonifera*, accounted for 90% of the total assemblage (Fig. 4a). The abundance of planktic foraminifera ranges from  $\sim 3$  to  $\sim 80$  ind.  $m^{-3}$ , and shows a strong zonal divide (Fig. 4b, c). The lowest abundances, with median values ranging between  $\sim 4$  and  $\sim 12$  ind.  $m^{-3}$ , occur off Sumatra (transects 1-3) (Fig. 4b, c) while, the highest abundances, with median values ranging between 18 and 35 ind.  $m^{-3}$ , occur off southern Sumatra and Java-LSI (transects 5-7) (Fig. 4b, c).

### 195 3.2 Vertical distribution of palaeoceanographic-relevant species

Similar to the spatial pattern of total abundance in each transect (Fig. 4b), the vertical dispersion of planktic foraminifera abundance across the water column shows a strong zonal pattern (Fig. 5a). Off Sumatra, the vertical distribution of planktic foraminiferal abundance is characterized by a high concentration in the upper 50 m (i.e., >30% of the total abundance) and a rapid decrease toward deeper levels with  $\leq 15\%$  of the total abundance occurring between 50 and 500 m water depth. Mean-  
200 while, off Java and the LSI, planktic foraminifera can be found throughout the upper 500 m of the water column. Unlike for stations off Sumatra, here the first 50 m of water column are relatively poor in planktic foraminifera and the highest concentration (>20 % of the total abundance) occurs between 50 to 100 m water depth. This general pattern is true for all stations off Java-LSI except station GeoB10062-1 (Fig. 4c).

In the following section, we describe the vertical distribution of six species of planktic foraminifera with paleoceanographic  
205 relevance that are typically used in paleoceanographic studies (for example, Caley et al., 2012; Ding et al., 2013; Mohtadi et al., 2017; Steinke et al., 2014; Tapia et al., 2019), namely *T. trilobus*, *G. ruber* (white), *G. bulloides*, *N. dutertrei*, *P. obliquiloculata*, and *G. menardii* (Caley et al., 2012; Ding et al., 2013; Mohtadi et al., 2017; Steinke et al., 2014; Tapia et al., 2019). *Trilobatus trilobus* exhibits a surface distribution with most of the organisms dwelling in the upper 50 m of the water column. Interestingly, some differences in their vertical distribution can be observed between sectors (i.e., Sumatra vs. Java-LSI) (Fig.  
210 5b). In the Sumatra sector, *T. trilobus* is highly concentrated within the upper 30 m of the water column, showing a small vertical dispersion, as its occurrence below 75 m is rare (Fig. 5b). Meanwhile, offshore Java and the LSI, *T. trilobus* shows a larger vertical dispersion with a relatively high concentration as deep as 100–200 m (Fig. 5b). The vertical distribution of *G. ruber* (white) shares some similarities with *T. trilobus*, that is, high concentration of specimens within the first 50 m of the water column off Sumatra and a larger dispersion off Java and the LSI (Fig. 5c). However, the vertical distribution of *G. ruber* (white) off Java suggests an even larger vertical dispersion than that of *T. trilobus*, as *G. ruber* (white) shows that the  
215 isocline lysocline of 20% of presence stretches from 100 to 400 m of water depth (Fig. 5c). *Globigerina bulloides* shows no clear pattern in its vertical preference off Sumatra, occupying both upper and lower depths. In contrast, off Java and the LSI



220 this species seems to prefer water depths below the thermocline between 100 and 350 m (Fig. 5d). *Pulleniatina obliquiloculata* and *N. dutertrei* (Fig. 5e and 5f) show a similar distribution with high occurrence of individuals between ~50 and ~100 m water depth. This feature seems to be constant throughout the study area. Although *G. menardii* is found mostly below 30 m water depth, its vertical distribution deepens along the study area, from dwelling at ~50 m water depth off Sumatra to ~150 m water depth off Java-LSI (Fig. 5fg). Overall, abundance distribution of the selected species across the water column agrees with previous studies that categorize these species as surface-mixed layer dwellers (*T. trilobus*, and *G. ruber* (white)) or as deep-thermocline dwellers (*N. dutertrei*, *P. obliquiloculata*, and *G. menardii*) (Birch et al., 2013; Faul, 2000; Hemleben et al., 225 1989; Lessa et al., 2020; Rebotim et al., 2017; Steph et al., 2009).

The ALD calculated from the total abundance (living + dead specimens) of the selected species ranges between 47 and 113 m water depth (Fig. 6). The ALD for *G. menardii*, *N. dutertrei*, and *P. obliquiloculata* indicates that the habitat depth of these species is located between ~67 to ~87 m. The ALD value for *G. bulloides* suggests that the habitat depth of this species in the study area is ~113 m, much deeper than the mixed layer in the study area. Meanwhile, the ALD for mixed-layer species 230 is ~49 m for *T. trilobus* and ~69 m for *G. ruber* (white), respectively. Interestingly, the ALD of surface-dwelling *G. ruber* (white) is similar to that of some thermocline-dwelling species.

## 4 Discussion

### 4.1 Planktic foraminiferal abundance and assemblages off Indonesia

235 The total number of species ( $n = 29$ ; size fraction  $>150 \mu\text{m}$ ) observed in the plankton net samples collected during August and September 2005 is higher than the number of species observed in the surface sediments off Indonesia ( $n = 18$ ; size fraction  $>150 \mu\text{m}$ ) (Ding et al., 2006) but lower than that observed in a sediment trap off Java ( $n = 37$ ; size fraction  $>150 \mu\text{m}$ ) (Mohtadi et al., 2009). Higher diversity in our data compared to those of Ding et al. (2006) may be due to the loss of fragile and dissolution-prone species (e.g., *Hastigerinella digitata*, *Turborotalita humilis*, *Globigerinita uvula*) in the sedimentary record, since only 4 of the 13 sites collected from the upwelling area off Java by Ding et al. (2006) are above the lysocline 240 (~2400–2800 m water depth) (Ding et al., 2006; Mohtadi et al., 2007). On the other hand, the longer temporal interval spanned by sediment trap samples off Java (3 months) compared to our sampling period (6 weeks) makes it possible to collect species with sporadic occurrence throughout the year. A recent global compilation study which includes the sediment trap data from Indonesia reported a warm bias in the assemblage due to anthropogenic effects and the likelihood that the water column may have changed over the last few decades (Jonkers et al., 2019). We note that their approach is based on the biogeography 245 of planktic foraminifera, i.e., each species occupies a specific thermal niche, which may span a temperature range of  $>10^\circ\text{C}$  for some tropical species. On the other hand, our main findings about the habitat depth and implications for paleoclimate reconstruction are based on individual species. The selected species are not dwelling at the limit of their thermal niche, thus as long as these species do not substantially shift their thermal niche over time, we do not expect any large bias due to the reported anthropogenic changes in foraminiferal assemblage.



250 The abundance of planktic foraminifera in the upwelling sector of Java-LSI is 4 to 8 times higher than that in the non-upwelling-Sumatra sector (Fig. 4b). Higher foraminiferal abundance in the presence of upwelling is consistent with previous studies, and is likely due to higher food availability when upwelling occurs (Kimoto, 2015; Schiebel and Hemleben, 2005; Schiebel et al., 2001). Species such as *G. bulloides*, *G. glutinata* and *G. falconensis* have been classified as species associated with upwelling conditions, and their abundances are positively correlated with the intensity of the upwelling (Brock et al., 1992; 255 Cayre et al., 1999; Conan and Brummer, 2000; Sautter and Sancetta, 1992). Species such as *N. dutertrei*, *P. obliquiloculata*, and *G. menardii* normally live in nutrient-rich waters below the mixed layer in the thermocline (Sautter and Thunell, 1991). Meanwhile, *T. trilobus* and *G. ruber* (white) occur in areas with a thick mixed layer. These mixed-layer dwellers have a similar life span (of two to four weeks), feeding strategies, and reproduction synchronized with the synodic lunar cycle (Schiebel and Hemleben, 2017 and references therein). Compared to *T. trilobus*, which predominantly occurs under warm oligotrophic 260 conditions, *G. ruber* (white) displays a more opportunistic behavior, as this species can be abundant and occurs in a larger range of trophic conditions, from oligotrophic to eutrophic due to its ability to feed on a greater variety of food sources (Schiebel and Hemleben, 2017; Schiebel et al., 2018).

Off Sumatra without upwelling, *G. ruber* (white), *G. elongatus* and *T. trilobus* account for ~44% of the total assemblage, while the species associated with high productivity or upwelling conditions have a minor presence (*G. bulloides* (~8%) and *G. glutinata* (~9%)) or are rare to absent (*N. dutertrei*, <2%) (Fig. 7a). On the other hand, off Java and the LSI where upwelling 265 occurs, the assemblage composition is strongly dominated by *G. bulloides* (21%), *G. glutinata* (14%) and the deep-dwelling species associated with high productivity i.e., *N. dutertrei*, *G. menardii*, and *P. obliquiloculata*, together contribute 18% to the total assemblage. *Trilobatus trilobus* and *G. ruber* (white), contribute only 7% and 8%, respectively, to the total assemblage off Java and the LSI (Fig. 7b).

270 Interestingly, despite methodological differences (sampling season and water depth, size fraction analyzed), our results are broadly consistent with those of a plankton net study carried out here in late autumn-early winter of 1963 at the end of the upwelling season (Ujjié, 1968). Ujjié found that the assemblage of planktic foraminifera off Java was consisted of a mixture of species associated with nutrient-rich and nutrient-poor waters, dominated by *N. dutertrei* (28%), *G. ruber* (22%) and *T. trilobus* (10%). On the other hand, off Sumatra (100° E) oligotrophic species *T. trilobus* and *G. ruber* accounted for 56% of 275 the total assemblage of planktic foraminifera. Thus, the assemblage and dominant species characterizing these two sectors seem to persist until the end of the upwelling season. The spatial contrast in the composition and abundance of planktic foraminiferal species (Fig. 4 and 7) in upwelling vs. non-upwelling conditions suggests that the hydrographic changes related to upwelling govern the distribution of planktic foraminifera. This finding is consistent with previous studies suggesting that off Indonesia, seasonal upwelling plays a critical role in modulating the ecology of planktic foraminifera (Ding et al., 2006; 280 Mohtadi et al., 2007, 2009, 2011; Ujjié, 1968). Consequently, foraminiferal abundances (Fig. 4b) and species composition in plankton net samples collected during the SE monsoon season reflect the transition, in space, from oligotrophic, deep mixed layer and more stratified upper water column conditions (Sumatra) to a more eutrophic, shallow mixed layer and well-mixed upper water column (Java and the LSI). The transition zone between the two hydrographic regimes for the period August-September is located off the Sunda Strait (~103° – ~105° W) (Fig. 1c) (Susanto et al., 2001). Altogether, the aforementioned

285 observations suggest that changes in the temporal extent, intensity, and zonal coverage of the seasonal upwelling might have a profound effect on the ecology of planktic foraminifera in the study area. Interestingly, the zonal divide in planktonic planktic foraminiferal assemblage between non-upwelling and upwelling regions observed in our net data is also reflected to some degree in surface sediments (Fig. 7c and 7d). Similar to what we observed in the plankton net data of the SE monsoon, the assemblage in sediments off Sumatra is dominated by *G. ruber* (white) while that off Java is dominated by *G. bulloides*.  
290 However, the proportion of less dominant species differs for plankton net and sediment samples. For instance, *G. calida* constitutes 12% of Java net samples but only 4% in the sediments. This discrepancy may reflect the different temporal coverage of these two sample types, i.e., surface sediments integrate over tens to hundreds of years whereas net samples provide only a “snap-shot” of the sampling period during the SE monsoon. The test of *G. calida* is relatively fragile, thus post-depositional processes like carbonate dissolution may also bias the assemblage in sediments (Ding et al., 2006). The effect of dissolution  
295 is likely not severe, as most of the stations are above the lysocline (Mohtadi et al., 2007), and some *G. calida* are found in sediments (Ding et al., 2006) albeit at a lower proportion than in our plankton net data.

#### 4.2 Planktic foraminifera habitat depth off Indonesia

Processes such as daily vertical migration and reproduction may also play a role in the vertical distribution of planktic foraminifera. The effect of daily vertical migration cannot be properly assessed by our sampling design, but there is strong  
300 evidence that argues against daily vertical migration in planktic foraminifera (Meilland et al., 2019). However, it cannot be excluded that lateral patchiness of foraminiferal occurrence affects the vertical distribution, as proposed recently by Meilland et al. (2019). To buffer against this potential caveat as well as potential bias due to dead specimens collected at depths (details in section 2.2), we interpret data averaged over several stations within each individual transects.

In the case of a modified vertical distribution due to synchronized reproduction – if some species reproduced consistently in  
305 phase with the full moon (Schiebel and Hemleben, 2017) – changes in the size class distribution should be noticeable before and after the occurrence of full moon during the sampling period. Among all the species selected, *G. ruber* (white), *N. dutertrei* and *G. menardii* show changes in their size classes distribution consistent with this pattern (Fig. A1.2). The other species show equal distribution of the larger and smaller size fractions before and after full moon. Although we cannot completely rule out ontogenic vertical migration, the lack of coherence between the calculated ALDs and the moon phase argues against ontogenic  
310 vertical migration as primary driver of the habitat depth distribution in the study area.

The discussion in the following sub-sections focuses on species that are commonly used in geochemical analyses for paleoceanographic reconstruction, namely *G. ruber* (white), *T. trilobus*, ~~*G. bulloides*~~, *N. dutertrei*, *P. obliquiloculata*, and *G. menardii* and *G. bulloides*. Importantly, the habitat depth of these species was also the focus of several previous studies in the region based on sediment trap and surface sediments (Mohtadi et al., 2007; 2009; 2011).

#### 315 4.3 Dominant species in nutrient-poor waters: *G. ruber* and *T. trilobus*

Many studies have shown that *G. ruber* (white) and *T. trilobus* have mixed layer habitat preferences in oligotrophic conditions (Bé, 1977; Duplessy et al., 1981; Fairbanks et al., 1980; Kuroyanagi and Kawahata, 2004), thus they are considered reliable

recorders of changes in the surface ocean at different timescales. However, the habitat depth of these species may change from area to area depending on the local hydrography, e.g., the depth of the mixed layer (Schiebel and Hemleben, 2017 and references therein).

The calculated habitat depth for *G. ruber* (white) (median = 69 m) (Fig. 6) is deeper than the habitat depth estimated from surface sediments for the study area (20 – 50 m) (Mohtadi et al., 2007, 2011). In contrast, the calculated habitat depth for *T. trilobus* (median = 49 m) (Fig. 7) is similar to the calcification depth estimated using surface sediments off Indonesia (~50 m) (Mohtadi et al., 2011). The relatively great habitat depth shown by *G. ruber* (white) may be related to the lack of living planktic foraminifera specimens and the use of total counts in the calculation of the ALD (see section 2.2).

Previous studies have suggested that a deepening of the habitat depth due to the use of total counts can be considered marginal (Greco et al., 2019). For example, in the area influenced by the upwelling, including both live and dead (total) specimens in the calculation of ALD leads to an increase of only ~4 to 15 m for *G. ruber* (Lessa et al., 2020; Rebotim et al., 2017). Similar overestimation, 2–13 m, in the habitat depth of *T. trilobus* is observed when dead specimens are included by including inclusion of dead specimens in the ALD, suggesting that the bias on ALD calculation caused by the inclusion of dead specimens should be comparable across mixed-layer dwellers. If this is also true for our study area, the agreement in habitat depth of *T. trilobus* inferred from sediments and our ALD calculation indicates that dead specimens likely do not make up a large portion of the samples, thus the relatively deep ALD calculated for *G. ruber* (white) is likely not severely biased by the inclusion of dead specimens in the calculation. Therefore, the inclusion of dead specimens may not necessarily result in a severe bias in the habitat depth estimates. Furthermore, the agreement in the habitat depth of *T. trilobus* inferred from sediments and our ALD calculation also suggests that dead specimens likely do not make up a large portion of the net samples; the same is probably true for *G. ruber* (white) from the same samples. Together, these observations suggest that the relatively deep ALD calculated for *G. ruber* (white) is likely a robust finding and not severely biased by the inclusion of dead specimens in the calculation.

#### 4.4 Dominant species in nutrient-rich waters: *N. dutertrei*, *P. obliquiloculata*, *G. menardii* and *G. bulloides*

Species such as *N. dutertrei*, *P. obliquiloculata*, *G. menardii* and *G. bulloides* are normally associated with high food availability and dwell in the upper water column across the mixed layer and upper part of the thermocline (Schiebel and Hemleben, 2017 and references therein). The habitat depth based on ALD calculation of the deep-dwelling species, *N. dutertrei* (median = 82 m) and *P. obliquiloculata* (median = 87 m) (Fig. 7) shows a good agreement with the habitat depth inferred from both sediment trap time series and surface sediments, i.e., 75–100 m for *N. dutertrei* and, 60–90 m for *P. obliquiloculata*, respectively (Mohtadi et al., 2009, 2011). These habitat depth estimates are close to the lower end of the range for these species in regions influenced by the Benguela and Canary upwelling system in the Atlantic Ocean, that is, ALDs of  $52 \pm 32$  m for *N. dutertrei* and  $45 \pm 31$  m for *P. obliquiloculata*, respectively (Lessa et al., 2020; Rebotim et al., 2017). Similarly, the habitat depth of *G. menardii* off Indonesia is also at the lower end of its habitat depth observed off Africa in waters influenced by the Benguela Upwelling System (ALD of  $39 \pm 22$  m) (Lessa et al., 2020). Notably, of all the species only *G. menardii* shows a shallower habitat depth (median = 67 m) (Fig. 8) than that inferred from sediment trap data (90 to 110 m) (Mohtadi et al., 2009). A habitat depth of

67–87 m water depth, for *G. menardii*, *N. dutertrei*, and *P. obliquiloculata*, places them just below the lower boundary of the mixed layer, which is on average 52 m (Table A1 and A2).

*Globigerina bulloides* has a median habitat depth of 113 m (Fig. 7); this value is almost twice the mean habitat depth estimated from surface sediments off Indonesia (~50 m) (Mohtadi et al., 2011), and close to the low end of the observed  
355 habitat depth, i.e.,  $57 \pm 10$  to  $102 \pm 67$  m, for this species in the Atlantic (areas under influenced of Benguela and Canary upwelling) (Lessa et al., 2020; Rebotim et al., 2017). *Globigerina bulloides* is an opportunistic species whose abundance and habitat depth has been linked to food availability in the water column (Peeters and Brummer, 2002). Therefore, it is possible that its habitat depth follows the depths where food availability is the highest in the water column. The comparison of its median ALD value off Sumatra (non-upwelling) and off Java-LSI (upwelling) shows a deepening from ~64 m to ~152 m,  
360 respectively. This finding suggests that regional differences in the habitat depth of some planktic foraminifera might occur as a result of seasonal upwelling.

#### 4.5 Zonal differences in the habitat depth: non-upwelling vs. upwelling

Previous studies off Indonesia matching geochemical data ( $\delta^{18}\text{O}_c$ ) ( $\delta^{18}\text{O}_{\text{calcite}}$ ) from core tops and water profiles have suggested differences in the habitat depth of some planktic foraminifera species between the sectors of Sumatra, Java and the LSI  
365 (Mohtadi et al., 2007). Mohtadi et al. (2007) reported that off Sumatra, the *G. ruber*- $\delta^{18}\text{O}$  values reflected a habitat depth <50 m water depth, while off Java and the LSI the *G. ruber*- $\delta^{18}\text{O}$  values were “out of range” (<0 m water depth). The authors speculated that the “out of range” geochemical signature off Java most likely reflects a greater contribution during the non-upwelling period when the water is warmer, fresher and more stratified. A similar situation was observed for *N. dutertrei*, as its geochemical signature suggests a deeper habitat depth in the non-upwelling Sumatra sector (50–75 m) relative to the upwelling  
370 Java sector (20–50 m) (Mohtadi et al., 2007). The notion that the seasonal upwelling off Indonesia may trigger changes in the habitat depth of planktic foraminifera species is further supported by sediment trap data, wherein Mg/Ca and  $\delta^{18}\text{O}$  data from planktic foraminifera (i.e., *G. ruber*, *N. dutertrei*, *G. menardii*, and *P. obliquiloculata*) vary with seasonal upwelling (Mohtadi et al., 2009). Therefore, the coherent, strong zonal shifts in the vertical dispersion of the selected planktic foraminifera in our plankton net samples (Fig. 5 a-g) and hydrographic parameters (i.e., SST, MLD, salinity, and chl-*a*) (Fig. 2a-c) in response to  
375 upwelling call for further scrutiny of potential zonal differences in the habitat depth values (Fig. 6).

The zonal disaggregation of the habitat depth (Fig. 8) into non-upwelling (transect 1 to 3) and upwelling (transect 5 to 7) sectors (see section 3.1) shows that *G. ruber* (white), *T. trilobus*, and *G. bulloides* have a greater habitat depth in the upwelling sector than the non-upwelling sector. There is a two-fold increase in their mean habitat depth (Fig. 8a-c) from non-upwelling to upwelling sector, i.e., from ~33 m to ~85 m for *T. trilobus*, ~58 m to ~97 m for *G. ruber* (white) and, ~64 m to ~152 m  
380 for *G. bulloides*. This finding, i.e., differing habitat depths in upwelling vs. non-upwelling regions for *T. trilobus* and *G. ruber* (white), is further corroborated by the estimates of bootstrap 95% confidence interval and randomization test with replacement (Table 1; details of calculation in Method). Compared to other species, the statistical significance of the ALD difference in *G. bulloides* is less strong ( $P = 0.057$ ), likely due to the large spread in the vertical dispersion of this species. Although deeper than usually assumed for paleoceanographic reconstructions, the ALD values for *G. ruber* (white), *T. trilobus*, and *G. bulloides*

385 in the upwelling sector are indeed within the ranges previously reported elsewhere for areas influenced by upwelling (Lessa  
et al., 2020; Rebotim et al., 2017) (see section 4.2). A clear zonal divide can be observed, i.e., shallow ALD values occurring  
off Sumatra (non-upwelling) vs. deep ALD values occurring off Java-LSI (upwelling). Transect 7 in the LSI is in addition to  
monsoonal upwelling also under the influence of the ITF as it allows the passage of cooler and fresher water (Tillinger, 2011).  
The habitat depth of mixed layer dwellers here is not the deepest in the upwelling region, despite the low presence of mixed-  
390 layer dwellers (*G. ruber* (white) and *T. trilobus*) in the upper 50 m during the sampling period (Fig. 5), especially at stations  
GeoB10065 and GeoB10070 (Fig. 4c). The ALD estimates of *G. ruber* (white) and *T. trilobus* at these two stations are the  
deepest along Transect 7 (the two lowest data points in the panel for *G. ruber* and *T. trilobus* in Figure 8). Station GeoB10070  
is the easternmost site of the study, thus might be under a stronger influence of the ITF than the other stations along the transect.  
This is, however, not the case for station GeoB10065, which is flanked by several stations at which mixed layer dwellers are  
395 present in the upper 50 m. Therefore, it is possible that the vertical distribution at these two stations is not representative of the  
transect nor of the influence of the ITF. Multivariate analysis also indicates that the hydrography at transect 7 is similar to that  
at transect 5 and 6, suggesting a negligible influence of ITF here. Although the calculated habitat depths off Sumatra (Fig. 8)  
of the surface dwellers show a relatively good fit with the estimated habitat depths based on geochemical data ( $\delta^{18}\text{O}_{\text{calcite}}$  and,  
Mg/Ca temperatures) (Mohtadi et al., 2009, 2011), this is not the case for transects off Java and the LSI, where the overall lack  
400 of agreement between the sedimentary data and plankton net results is evident (Figure 8a, b and c).

The low abundance of *G. menardii*, *N. dutertrei* and *P. obliquiloculata* off Sumatra precludes ALD calculation, hence also  
the zonal comparison of their habitat depths (Fig. 8d-f) between Sumatra and Java-LSI. Despite their low abundance in net  
samples collected in August-September, these three species are found in relatively high abundances in surface sediments off  
Sumatra, constituting up to 13% of the assemblage (Fig.7c; Mohtadi et al., 2007). Thus, their occurrence offshore Sumatra  
405 might be temporally restricted to only the final part of the SE monsoon (October) when the upwelling center locked off the  
Sunda Strait starts drifting westward (Susanto et al., 2001), triggering higher productivity offshore Sumatra. Alternatively, they  
might represent the positive IOD or El Niño years, when upwelling is generally stronger in the Eastern Indian Ocean and  
reaches further northerly latitudes. Off Java where their abundance is sufficiently high for ALD calculation, the data suggest  
that these three species share a similar niche at thermocline depths centered at ~90 m water depth, in agreement with the  
410 habitat depth inferred from surface sediments (Fig. 8d-f).

#### 4.6 Possible Implications for palaeoceanographic reconstructions

Field observations (plankton net and sediment trap data) provide insights into the modern ecology of planktonic planktic  
foraminifera, as the habitat depth of some species is known to vary in time and at regional scale (Schiebel and Hemleben,  
2017). Our plankton net data show that the habitat depth of mixed-layer dwelling *G. ruber* (white) and *T. trilobus* deepens in  
415 upwelling conditions, while thermocline-dwelling *N. dutertrei* and *P. obliquiloculata* thrive only in the upwelling region off  
Java and the LSI.

As with many zooplankton, the abundance of planktic foraminifera is linked to food availability, thus some species may  
change their habitat depth to maximize food acquisition. The habitat depth of symbiont-bearing species like *T. trilobus* and

*G. ruber* is typically assumed to be restricted to the surface ocean or at least within the photic zone as they rely on their  
420 photosynthetic symbionts for nutrition. In addition, these species also catch prey and feed on a wide variety of food sources  
(Hemleben et al., 1989 and references therein), plausibly because the photosynthates produced by the symbionts are insufficient  
to sustain the growth of the host (Bé et al., 1981; Caron et al., 1982). Indeed, a recent study showed that the nutritional  
contribution of the symbionts to the host is significantly smaller than that obtained by ingesting copepods, implying that the  
photosymbiosis in planktic foraminifera may not be the primary source of energy when preys are abundant (Takagi et al.,  
425 2018).

During the SE monsoon, the photic zone ranges between ~50 m to ~75 m, and the food availability off Java and the LSI  
is high due to the enhanced predator-prey encounter fueled by the high phytoplanktonic biomass across the water column and  
upwelling-induced vertical mixing (Pécseli et al., 2014). Under this circumstance, it is plausible that omnivorous mixed-layer  
dwellers like *T. trilobus* and *G. ruber* (white) might adopt a feeding strategy that includes both photosymbiosis and preying,  
430 or even primarily the latter. Preying on other zooplanktons like copepods which have ontogenic and daily migration through  
water column means that planktic foraminifera are not limited to the photic zone for food. Moreover, copepods have been  
found to shift to a deeper habitat as a strategy to maximize their retention within a coastal upwelling area (Peterson, 1998;  
Peterson et al., 1979; Verheyen et al., 1991). In combination, the aforementioned factors may thus lead to a greater habitat depth  
for mixed-layer dwellers in the upwelling region off Java compared to the non-upwelling region off Sumatra.

435 The thermal gradient of **between** mixed-layer and deep dwelling species ( $\Delta T$ ) is commonly used as a proxy for the thickness  
of the mixed layer and the position of the thermocline in the water column on glacial-interglacial time scales (Farmer et  
al., 2011; Mohtadi et al., 2017; Steinke et al., 2014; Tapia et al., 2015). The  $\Delta T$  calculated from the abundance-weighted  
temperatures of our plankton net data show that during the SE monsoon larger  $\Delta T$  values occur off Sumatra than off Java-  
LSI where upwelling occurs (Fig. 9 and 10). Due to its relatively great habitat depth off Java-LSI, the abundance-weighted  
440 temperature derived from surface-dwelling *G. ruber* (white) and *T. trilobus* is comparable to that of thermocline-dwelling  
species, and substantially lower than that of their counterpart off Sumatra. **The habitat depth change of the mixed-layer species  
is thus the primary reason for the  $\Delta T$  difference between the two sectors.** Interestingly, smaller  $\Delta T$  during upwelling conditions  
are also evident in the flux-weighted data from a sediment trap off Java (Fig. 9 and 10), which show that the  $\Delta T$  reduces  
from 4.3 °C during non-monsoon period to 1.2 °C during seasonal upwelling period (SE monsoon) (Mohtadi et al., 2011).  
445 Although both plankton net and sediment trap data are based on suspended or sinking foraminifera in the water column,  
they reflect a different signal in time and space – our net data reflect spatial difference during the sampling period spanning  
six weeks during the monsoonal upwelling season, while the trap data reflect temporal changes at one location that is under  
the influence of monsoonal upwelling. Despite their inherent differences, both datasets, however, show that off Indonesia ( $\Delta T$ )  
decreases as a function of upwelling dynamics. A more well-mixed upper water column in upwelling conditions, hence smaller  
450 surface-subsurface temperature difference, is also reasonable taking into account the weaker water column stratification during  
upwelling (Fig. 10). The latter is due to a thinner barrier layer and a shallower  $MLD_{TEMP}$  (Fig. 2a and Table A1 and A2). This  
finding indicates that  $\Delta T$  may be a useful proxy for reconstructing past upwelling conditions off Indonesia, if the foraminifera  
produced during upwelling season dominate foraminiferal test abundance in the sediments.



Mohtadi et al (2009) showed that ~50% of the total annual foraminiferal flux off Java occurs during the SE monsoon season, and the fluxes during this season are largely centered around September, suggesting that our “snapshot” may be reasonably representative of the foraminiferal response to the prevailing ocean conditions during the SE monsoon off Indonesia. Geochemical data ( $\delta^{18}\text{O}_{\text{calcite}}$  and Mg/Ca inferred temperatures) in marine sediments off Java show a broad agreement with the habitat depth estimates from our plankton net data for subsurface dwellers (Fig. 8), but not for mixed-layer dwelling species. **Plankton net data suggest a greater habitat depth for the mixed-layer species and hence also lower inferred temperature. As a result, the  $\Delta T$  off Java-LSI calculated from the plankton net data is smaller than that of surface sediment data.** The situation differs off Sumatra, where the habitat depth estimates for both mixed layer and subsurface species derived from marine sediments and plankton net samples are in agreement, hence also the derived  $\Delta T$ . Consequently, the zonal reconstruction of the  $\Delta T$  based on surface sediments indicates a larger  $\Delta T$  off Java-LSI than off Sumatra (Mohtadi et al., 2011), in contrast to that of our net data (Fig. 9 and 10). In other words, surface sediment data suggest larger  $\Delta T$ s in regions influenced by strong seasonal upwelling.

This discrepancy between surface sediment and plankton net data off Java may stem from the different temporal intervals integrated by each sample type. The plankton net data reflect the conditions in the water column sampled during the SE monsoon, thus can be directly linked to the hydrographic processes that occurred during sampling. To some extent this is also true for sediment trap samples – limited temporal interval integrated by the samples at one location means that process attribution can be better constrained. On the other hand, marine sediments integrate foraminiferal flux over tens to hundreds of years depending on the local sedimentation rate, and may be susceptible to post-depositional processes such as dissolution, bioturbation, and reworking which may alter proxy signal. Together, these issues make it challenging to quantify the relative importance of each processes and thus to ground-truth proxies. Nonetheless, paleoceanographic reconstructions are based on sedimentary material, which may have undergone the same post-depositional processes as the surface sediments. If the relative importance of these processes would stay unchanged over time, then surface sediments are arguably the best modern analogue for paleoceanographic reconstruction. In this case, stronger upwelling off Indonesia is characterized by a larger ( $\Delta T$ ) as indicated by the zonal pattern of surface sediments (Fig. 9 and 10). However, were the fluxes of planktic foraminifera and the post-depositional processes to change over time, it is within the realm of possibility that under some circumstances, e.g., strong positive IOD or El Niño years, the proxy signal produced in the water column during one-off events like upwelling can be preserved in the sediments, especially if the resultant flux increases exponentially. Our findings highlight the need to consider multiple sample types to further constrain the analog used for downcore paleoceanographic reconstructions. **To further shed light on the transfer of proxy signal from the water column to the sediment, longer sediment trap time series and repeated plankton net sampling in the same region would be useful to capture the seasonality of the vertical distribution of planktic foraminifera. Importantly, generating geochemical data on plankton net samples may help to verify the habitat depths and allow a direct comparison with the depth inference from the surface sediments. It would also be helpful to constrain the age of surface sediments to ensure that they are comparable to modern data.**



## 5 Conclusions

We examined the zonal and vertical distribution of planktic foraminifera off Indonesia during the boreal summer of 2005 at 37 stations. The stations were grouped into 7 land-sea transects off Sumatra and off Java-LSI. The factors driving the observed distribution of foraminifer species were assessed using vertically resolved environmental data.

490 Twenty-nine species were identified that can be divided in two basic communities, i.e., one dominated by warm-oligotrophic-stratified water column species vs. cooler-eutrophic-well mixed water column species. Similarly, foraminiferal abundance shows contrasting distributions off Sumatra and off Java-LSI. The concurrent zonal shift in abundance and species composition of planktic foraminiferal and environmental parameters (SST, Chl-*a*, MLD and BL) in response to upwelling imply a close link between upwelling and the ecology of planktic foraminifera in this area. Similar to the abundances and species composition  
495 of planktic foraminifera, their vertical distribution across the water column shows a strong zonal differentiation i.e., shallow depths-low dispersion off Sumatra vs. deeper depths-larger dispersion off Java-LSI.

The calculated ALDs of the selected species are in broad agreement with typically assumed habitat depths in paleoceanographic reconstructions, with the exception of *G. ruber* (white) and *G. bulloides*. The possibility of an overestimation due to possible inclusion of dead specimens in the calculation cannot be entirely ruled out at this point, but several lines of evidence  
500 suggest that a severe overestimation is not likely. The species *G. ruber* (white) and *T. trilobus* show a strong deepening in their habitat depth off Java in comparison to the sector off Sumatra. This zonal divide is not evident in the habitat depth distribution of the subsurface dwellers *N. dutertrei*, *G. menardii*, and *P. obliquiloculata*, in part due to their low abundance off Sumatra. Compared to inferred habitat depth estimates based on surface sediments, those inferred from plankton net data show a better agreement for the subsurface species, while the relatively good agreement for the shallow dwellers is restricted to off Sumatra.  
505 Off Java and the LIS LSI, surface-dwelling species in our net data have a much greater habitat depth than that derived from surface sediments. The discrepancy between plankton net and surface sediment data likely stems from the fact that each sample type integrates over different temporal duration, i.e., net samples reflect the conditions during the sampling period in SE monsoon while the marine sediments integrate over tens to hundreds of years of foraminiferal flux. Whilst each sample type has its pros and cons, our findings highlight the need to consider multiple sample types to further constrain the analog adopted  
510 for paleoceanographic reconstruction.

*Data availability.* Data generated in this study are available in the supplementary material and on request to the main author until their online publication on PANGAEA database (<https://doi.pangaea.de/10.1594/PANGAEA.941895>).

*Author contributions.* MM and SLH conceptualized this study. MM was responsible for the collection of planktic foraminifera and oceanographic data. RT analyzed the planktic foraminifera fauna. JG provided support for taxonomic identification. H-Y W contributed to the  
515 statistical analyses. The manuscript was written by RT and SLH with the contribution from all co-authors who approved its final version.

*Competing interests.* The authors declare that they have no conflict of interest.

*Acknowledgements.* This study was funded by DFG grant HE3412/15-1 (MM), NTU Core Research Team fund 109L892604 and 110L890705 (SLH). RT acknowledges funding by MOST grants 108-2116-M-002-008, 109-2116-M-002-014 and 110-2116-M-002-007. We thank the crew and scientists participating in the PABESIA Cruise (SO-184). We are grateful to T. Sagawa (U. Kanazawa) and Y. Kubota (National  
520 Science Museum) for providing a copy of the paper by Ujiie (1968) stored at National Museum of Nature and Science in Tokyo, Japan.

## References

- Anderson, O. R., Spindler, M., Bé, A. W. H. and Hemleben, Ch.: Trophic activity of planktonic foraminifera, *J. Mar. Biol. Ass.*, 59(3), 791–799, <https://doi.org/10.1017/S002531540004577X>, 1979.
- Aurahs, R., Grimm, G. W., Hemleben, V., Hemleben, C. and Kucera, M.: Geographical distribution of cryptic genetic types in the planktonic foraminifer *Globigerinoides ruber*, *Mol. Ecol.*, 18(8), <https://doi.org/10.1111/j.1365-294X.2009.04136.x>, 2009.
- Bauer, M. A.: Ecology And Distribution Of Living Planktonic In Partial Fulfillment Of The Ecology And Distribution Of Living Planktonic, Doctoral dissertation, Rice University, 1976.
- Bé, A. W. H.: An ecological, zoogeographic and taxonomic review of recent planktonic foraminifer, in *Oceanic Micropaleontology*, edited by A. T. S. Ramsay, pp. 1–100, Academic Press, London, , 1977.
- 530 Bé, A. W. H. and Hamlin, W. H.: Ecology of Recent Planktonic Foraminifera: Part 3: Distribution in the North Atlantic during the Summer of 1962, *Micropaleontology*, 13(1), 87, <https://doi.org/10.2307/1484808>, 1967.
- Bé, A. W. H., Caron, D. A. and Anderson, O. R.: Effects of feeding frequency on life processes of the planktonic foraminifer *Globigerinoides sacculifer* in laboratory culture, *J. Mar. Biol. Ass.*, 61(1), 257–277, <https://doi.org/10.1017/S002531540004604X>, 1981.
- Bemis, B. E., Spero, H. J., Bijma, J. and Lea, D. W.: Reevaluation of the oxygen isotopic composition of planktonic foraminifera: Experimental results and revised paleotemperature equations, *Paleoceanography*, 13(2), 150–160, <https://doi.org/10.1029/98PA00070>, 1998.
- 535 Birch, H., Coxall, H. K., Pearson, P. N., Kroon, D. and O'Regan, M.: Planktonic foraminifera stable isotopes and water column structure: Disentangling ecological signals, *Mar. Micropaleontol.*, 101, 127–145, <https://doi.org/10.1016/j.marmicro.2013.02.002>, 2013.
- Brock, J. C., McClain, C. R., Anderson, D. M., Prell, W. L. and Hay, W. W.: Southwest monsoon circulation and environments of recent planktonic foraminifera in the northwestern arabian sea, *Paleoceanography*, 7(6), 799–813, <https://doi.org/10.1029/92PA01267>, 1992.
- 540 Caley, T., Giraudeau, J., Malaizé, B., Rossignol, L. and Pierre, C.: Agulhas leakage as a key process in the modes of Quaternary climate changes., *Proc Natl Acad Sci USA*, 109(18), 6835–6839, <https://doi.org/10.1073/pnas.1115545109>, 2012.
- Caron, D. A., Bé, A. W. H. and Anderson, O. R.: Effects of variations in light intensity on life processes of the planktonic foraminifer *Globigerinoides sacculifer* in laboratory culture, *J. Mar. Biol. Ass.*, 62(2), 435–451, <https://doi.org/10.1017/S0025315400057374>, 1982.
- Cayre, O., Beaufort, L. and Vincent, E.: Paleoproductivity in the Equatorial Indian Ocean for the last 260,000yr: A transfer function based on planktonic foraminifera, *Quat. Sci. Rev.*, 18(6), 839–857, [https://doi.org/10.1016/S0277-3791\(98\)00036-5](https://doi.org/10.1016/S0277-3791(98)00036-5), 1999.
- 545 CLIMAP Project Members: The surface of the ice-age Earth., *Science*, 191(4232), 1131–1137, <https://doi.org/10.1126/science.191.4232.1131>, 1976.
- Conan, S. M. H. and Brummer, G. J. A.: Fluxes of planktic foraminifera in response to monsoonal upwelling on the Somalia Basin margin, *Deep Sea Res. Part II Top. Stud. Oceanogr.*, 47(9–11), 2207–2227, [https://doi.org/10.1016/S0967-0645\(00\)00022-9](https://doi.org/10.1016/S0967-0645(00)00022-9), 2000.
- 550 Dang, H., Jian, Z., Wang, Y., Mohtadi, M., Rosenthal, Y., Ye, L., Bassinot, F. and Kuhnt, W.: Pacific warm pool subsurface heat sequestration modulated Walker circulation and ENSO activity during the Holocene., *Sci. Adv.*, 6(42), <https://doi.org/10.1126/sciadv.abc0402>, 2020.
- Davis, C. V., Wishner, K., Renema, W. and Hull, P. M.: Vertical distribution of planktic foraminifera through an oxygen minimum zone: how assemblages and test morphology reflect oxygen concentrations, *Biogeosciences*, 18(3), 977–992, <https://doi.org/10.5194/bg-18-977-2021>, 2021.
- 555 Ding, X., Bassinot, F., Guichard, F., Li, Q. Y., Fang, N. Q., Labeyrie, L., Xin, R. C., Adisaputra, M. K. and Hardjawidjaksana, K.: Distribution and ecology of planktonic foraminifera from the seas around the Indonesian Archipelago, *Mar. Micropaleontol.*, 58(2), 114–134, <https://doi.org/10.1016/j.marmicro.2005.10.003>, 2006.

- Ding, X., Bassinot, F., Guichard, F. and Fang, N. Q.: Indonesian Throughflow and monsoon activity records in the Timor Sea since the last glacial maximum, *Mar. Micropaleontol.*, 101, 115–126, <https://doi.org/10.1016/j.marmicro.2013.02.003>, 2013.
- 560 Duplessy, J. C., Bé, A. W. H. and Blanc, P. L.: Oxygen and carbon isotopic composition and biogeographic distribution of planktonic foraminifera in the Indian Ocean, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 33(1–3), 9–46, [https://doi.org/10.1016/0031-0182\(81\)90031-6](https://doi.org/10.1016/0031-0182(81)90031-6), 1981.
- Du, Y. and Zhang, Y.: Satellite and Argo Observed Surface Salinity Variations in the Tropical Indian Ocean and Their Association with the Indian Ocean Dipole Mode, *J. Clim.*, 28(2), 695–713, <https://doi.org/10.1175/JCLI-D-14-00435.1>, 2015.
- 565 Eguchi, N. O., Kawahata, H. and Taira, A.: Seasonal response of planktonic foraminifera to surface ocean condition: Sediment trap results from the central North Pacific Ocean, *Journal of Oceanography*, 55(6), 681–691, 1999.
- Fairbanks, R. G., Wiebe, P. H. and Bé, A. W.: Vertical distribution and isotopic composition of living planktonic foraminifera in the Western north Atlantic, *Science*, 207(4426), 61–63, <https://doi.org/10.1126/science.207.4426.61>, 1980.
- Fan, W., Jian, Z., Chu, Z., Dang, H., Wang, Y., Bassinot, F., Han, X. and Bian, Y.: Variability of the Indonesian Throughflow in the Makassar Strait over the last 30 ka., *Sci. Rep.*, 8(1), 5678, <https://doi.org/10.1038/s41598-018-24055-1>, 2018.
- 570 Farmer, E. J., Chapman, M. R. and Andrews, J. E.: Holocene temperature evolution of the subpolar North Atlantic recorded in the Mg/Ca ratios of surface and thermocline dwelling planktonic foraminifera, *Glob. Planet. Change*, 79(3–4), 234–243, <https://doi.org/10.1016/j.gloplacha.2011.02.003>, 2011.
- Faul, K. L.: Reconstructions of upwelling, productivity, and photic zone depth in the eastern equatorial Pacific ocean using planktonic foraminiferal stable isotopes and abundances, *Journal of Foraminiferal Research*, 30(2), 110–125, <https://doi.org/10.2113/0300110>, 2000.
- 575 Field, D. B.: Variability in vertical distributions of planktonic foraminifera in the California Current: Relationships to vertical ocean structure, *Paleoceanography*, 19(2), <https://doi.org/10.1029/2003PA000970>, 2004.
- Gordon, A. L.: Oceanography of the Indonesian seas and their throughflow, *oceanog.*, 18(4), 14–27, <https://doi.org/10.5670/oceanog.2005.01>, 2005.
- 580 Gordon, A. L. and Fine, R. A.: Pathways of water between the Pacific and the Indonesian seas, *Nature*, 379, 146–149, 1996.
- Greco, M., Jonkers, L., Kretschmer, K., Bijma, J. and Kucera, M.: Depth habitat of the planktonic foraminifera *Neoglobobulimina pachyderma* in the northern high latitudes explained by sea-ice and chlorophyll concentrations, *Biogeosciences*, 16(17), 3425–3437, <https://doi.org/10.5194/bg-16-3425-2019>, 2019.
- Groeneveld, J. and Chiessi, C. M.: Mg/Ca of *Globobulimina inflata* as a recorder of permanent thermocline temperatures in the South Atlantic, *Paleoceanography*, 26(2), <https://doi.org/10.1029/2010PA001940>, 2011.
- 585 Hebbeln, D. and cruise participants: Report and preliminary results of the RV SONNE Cruise SO-184, Pabesia, Durban (South Africa)-Cilacap (Indonesia) - Darwin (Australia), July 8th-September 13th, 2005., 2006.
- Hemleben, C., Spindler, M. and Anderson, O. R.: *Modern Planktonic Foraminifera*, Springer New York, New York, NY., 1989.
- Hollstein, M., Mohtadi, M., Rosenthal, Y., Moffa Sanchez, P., Oppo, D., Martínez Méndez, G., Steinke, S. and Hebbeln, D.: Stable oxygen isotopes and mg/ca in planktic foraminifera from modern surface sediments of the western Pacific warm pool: implications for thermocline reconstructions, *Paleoceanography*, 32(11), 1174–1194, <https://doi.org/10.1002/2017PA003122>, 2017.
- 590 Imbrie, J. and Kipp, N. G.: A new micropaleontological method for quantitative paleoclimatology: Application to a Late Pleistocene Caribbean core, in *The Late Cenozoic Glacial Ages*, edited by K. Turekian, pp. 71–81, Yale Univ. Press, New Haven, Conn., , 1971.
- Jonkers, L., Hillebrand, H. and Kucera, M.: Global change drives modern plankton communities away from the pre-industrial state., *Nature*, 570(7761), 372–375, <https://doi.org/10.1038/s41586-019-1230-3>, 2019.
- 595


- Jorissen, F. J., de Stigter, H. C. and Widmark, J. G. V.: A conceptual model explaining benthic foraminiferal microhabitats, *Mar. Micropaleontol.*, 26(1–4), 3–15, [https://doi.org/10.1016/0377-8398\(95\)00047-X](https://doi.org/10.1016/0377-8398(95)00047-X), 1995.
- Karas, C., Nürnberg, D., Tiedemann, R. and Garbe-Schönberg, D.: Pliocene Indonesian Throughflow and Leeuwin Current dynamics: Implications for Indian Ocean polar heat flux, *Paleoceanography*, 26(2), <https://doi.org/10.1029/2010PA001949>, 2011.
- 600 Kara, A. B., Rochford, P. A. and Hurlburt, H. E.: An optimal definition for ocean mixed layer depth, *J. Geophys. Res.*, 105(C7), 16803–16821, <https://doi.org/10.1029/2000JC900072>, 2000.
- Katz, M. E., Cramer, B. S., Franzese, A., Honisch, B., Miller, K. G., Rosenthal, Y. and Wright, J. D.: Traditional and emerging geochemical proxies in foraminifera, *Journal of Foraminiferal Research*, 40(2), 165–192, <https://doi.org/10.2113/gsjfr.40.2.165>, 2010.
- Kemle-von Mücke, S. and Oberhänsli, H.: The distribution of living planktic foraminifera in relation to southeast atlantic oceanography, 605 in *Use of proxies in Paleoceanography*, edited by G. Fischer and G. Wefer, pp. 91–115, Springer Berlin Heidelberg, Berlin, Heidelberg, [https://doi.org/10.1007/978-3-642-58646-0\\_3](https://doi.org/10.1007/978-3-642-58646-0_3), 1999.
- Kennett, J. P. and Srinivasan, M. S.: *Neogene planktonic foraminifera: A phylogenetic atlas*, Hutchinson Ross Publishing Company, Stroudsburg, Pa., 1983.
- Kimoto, K.: Planktic Foraminifera, in *Marine Protists*, edited by S. Ohtsuka, T. Suzuki, T. Horiguchi, N. Suzuki, and F. Not, pp. 129–178, 610 Springer Japan, Tokyo, [https://doi.org/10.1007/978-4-431-55130-0\\_7](https://doi.org/10.1007/978-4-431-55130-0_7), 2015.
- Kucera, M.: Planktonic foraminifera as tracers of past oceanic environments, in *Proxies in late cenozoic paleoceanography*, vol. 1, pp. 213–262, Elsevier, [https://doi.org/10.1016/S1572-5480\(07\)01011-1](https://doi.org/10.1016/S1572-5480(07)01011-1), 2007.
- Kucera, M.: Determination of past sea surface temperatures, in *Encyclopedia of ocean sciences*, pp. 98–113, Elsevier, <https://doi.org/10.1016/B978-012374473-9.00700-1>, 2009.
- 615 Kuroyanagi, A. and Kawahata, H.: Vertical distribution of living planktonic foraminifera in the seas around Japan, *Mar. Micropaleontol.*, 53(1–2), 173–196, <https://doi.org/10.1016/j.marmicro.2004.06.001>, 2004.
- Lea, D. W.: Elemental and isotopic proxies of past ocean temperatures, in *Treatise on Geochemistry*, pp. 373–397, Elsevier, <https://doi.org/10.1016/B978-0-08-095975-7.00614-8>, 2014.
- Lessa, D., Morard, R., Jonkers, L., Venancio, I. M., Reuter, R., Baumeister, A., Albuquerque, A. L. and Kucera, M.: Distribution of 620 planktonic foraminifera in the subtropical South Atlantic: depth hierarchy of controlling factors, *Biogeosciences*, 17(16), 4313–4342, <https://doi.org/10.5194/bg-17-4313-2020>, 2020.
- Locarnini, R. A., Mishonov, A. V., Baranova, O. K., Boyer, T. P., Zweng, M. M., Garcia, H. E., Reagan, J. R., Seidov, D., Weathers, K., Paver, C. R. and Smolyar, I.: *World Ocean Atlas 2018, Volume 1: Temperature*, edited by A. Mishonov, NOAA Atlas NESDIS 81., 2018.
- Manly, B. F. J.: *Randomization, bootstrap and monte carlo methods in biology*, 2nd ed., Chapman and Hall/CRC., 1997.
- 625 Marchant, M., Hebbeln, D., Giglio, S., Coloma, C. and González, H. E.: Seasonal and interannual variability in the flux of planktic foraminifera in the Humboldt Current System off central Chile (30°S), *Deep Sea Res. Part II Top. Stud. Oceanogr.*, 51(20–21), 2441–2455, <https://doi.org/10.1016/j.dsr2.2004.08.013>, 2004.
- Meilland, J., Siccha, M., Weinkauff, M. F. G., Jonkers, L., Morard, R., Baranowski, U., Baumeister, A., Bertlich, J., Brummer, G.-J., Debray, P., Fritz-Endres, T., Groeneveld, J., Magerl, L., Munz, P., Rillo, M. C., Schmidt, C., Takagi, H., Theara, G. and Kucera, M.: Highly replicated 630 sampling reveals no diurnal vertical migration but stable species-specific vertical habitats in planktonic foraminifera, *J. Plankton Res.*, 41(2), 127–141, <https://doi.org/10.1093/plankt/fbz002>, 2019.

- Mohtadi, M., Max, L., Hebbeln, D., Baumgart, A., Krück, N. and Jennerjahn, T.: Modern environmental conditions recorded in surface sediment samples off W and SW Indonesia: Planktonic foraminifera and biogenic compounds analyses, *Mar. Micropaleontol.*, 65(1–2), 96–112, <https://doi.org/10.1016/j.marmicro.2007.06.004>, 2007.
- 635 Mohtadi, M., Steinke, S., Groeneveld, J., Fink, H. G., Rixen, T., Hebbeln, D., Donner, B. and Herunadi, B.: Low-latitude control on seasonal and interannual changes in planktonic foraminiferal flux and shell geochemistry off south Java: A sediment trap study, *Paleoceanography*, 24(1), <https://doi.org/10.1029/2008PA001636>, 2009.
- Mohtadi, M., Oppo, D. W., Lückge, A., DePol-Holz, R., Steinke, S., Groeneveld, J., Hemme, N. and Hebbeln, D.: Reconstructing the thermal structure of the upper ocean: Insights from planktic foraminifera shell chemistry and alkenones in modern sediments of the tropical eastern Indian Ocean, *Paleoceanography*, 26(3), <https://doi.org/10.1029/2011PA002132>, 2011.
- 640 Mohtadi, M., Prange, M., Oppo, D. W., De Pol-Holz, R., Merkel, U., Zhang, X., Steinke, S. and Lückge, A.: North Atlantic forcing of tropical Indian Ocean climate., *Nature*, 509(7498), 76–80, <https://doi.org/10.1038/nature13196>, 2014.
- Mohtadi, M., Prange, M., Schefuß, E. and Jennerjahn, T. C.: Late Holocene slowdown of the Indian Ocean Walker circulation., *Nat. Commun.*, 8(1), 1015, <https://doi.org/10.1038/s41467-017-00855-3>, 2017.
- 645 Murray, J.: On the distribution of the pelagic foraminifera at the surface and on the floor of the ocean, *Sci. Am.*, 44(1128supp), 18026–18028, <https://doi.org/10.1038/scientificamerican08141897-18026supp>, 1897.
- Muskananfolo, M. R., Jumsar and Wirasatriya, A.: Spatio-temporal distribution of chlorophyll-a concentration, sea surface temperature and wind speed using aqua-modis satellite imagery over the Savu Sea, Indonesia, *Remote Sensing Applications: Society and Environment*, 22, 100483, <https://doi.org/10.1016/j.rsase.2021.100483>, 2021.
- 650 Ortiz, J. D. and Mix, A. C.: Comparison of Imbrie-Kipp Transfer Function and modern analog temperature estimates using sediment trap and core top foraminiferal faunas, *Paleoceanography*, 12(2), 175–190, <https://doi.org/10.1029/96PA02878>, 1997.
- Pang, X., Bassinot, F. and Sepulcre, S.: Indonesian Throughflow variability over the last two glacial-interglacial cycles: Evidence from the eastern Indian Ocean, *Quat. Sci. Rev.*, 256, 106839, <https://doi.org/10.1016/j.quascirev.2021.106839>, 2021.
- Parker, F. L.: Planktonic foraminiferal species in pacific sediments, *Micropaleontology*, 8(2), 219, <https://doi.org/10.2307/1484745>, 1962.
- 655 Pécseli, H. L., Trulsen, J. K. and Fiksen, Ø.: Predator-prey encounter and capture rates in turbulent environments, *Limnol. Oceanogr.*, 4(1), 85–105, <https://doi.org/10.1215/21573689-2768717>, 2014.
- Peeters, F. J. C. and Brummer, G.-J. A.: The seasonal and vertical distribution of living planktic foraminifera in the NW Arabian Sea, *Geological Society, London, Special Publications*, 195(1), 463–497, <https://doi.org/10.1144/GSL.SP.2002.195.01.26>, 2002.
- Peterson, W.: Life cycle strategies of copepods in coastal upwelling zones, *Journal of Marine Systems*, 15(1–4), 313–326, [https://doi.org/10.1016/S0924-7963\(97\)00082-1](https://doi.org/10.1016/S0924-7963(97)00082-1), 1998.
- 660 Peterson, W. T., Miller, C. B. and Hutchinson, A.: Zonation and maintenance of copepod populations in the Oregon upwelling zone, *Deep Sea Research Part A. Oceanographic Research Papers*, 26(5), 467–494, [https://doi.org/10.1016/0198-0149\(79\)90091-8](https://doi.org/10.1016/0198-0149(79)90091-8), 1979.
- Qu, T. and Meyers, G.: Seasonal variation of barrier layer in the southeastern tropical Indian Ocean, *J. Geophys. Res.*, 110(C11), <https://doi.org/10.1029/2004JC002816>, 2005.
- 665 Qu, T., Du, Y., Stachan, J., Meyers, G. and Slingo, J.: Sea surface temperature and its variability in the Indonesian region, *oceanog.*, 18(4), 50–61, <https://doi.org/10.5670/oceanog.2005.05>, 2005.
- Rebotim, A., Voelker, A. H. L., Jonkers, L., Waniek, J. J., Meggers, H., Schiebel, R., Fraile, I., Schulz, M. and Kucera, M.: Factors controlling the depth habitat of planktonic foraminifera in the subtropical eastern North Atlantic, *Biogeosciences*, 14(4), 827–859, <https://doi.org/10.5194/bg-14-827-2017>, 2017.


- 670 RegenberG, M., RegenberG, A., Garbe-SchönberG, D. and Lea, D. W.: Global dissolution effects on planktonic foraminiferal Mg/Ca ratios controlled by the calcite-saturation state of bottom waters, *Paleoceanography*, 29(3), 127–142, <https://doi.org/10.1002/2013PA002492>, 2014.
70. R Core Team: R: A language and environment for statistical computing., Computer software, R Foundation for Statistical Computing, Vienna, Austria., 2021.
- 675 Sautter, L. R. and Sancetta, C.: Seasonal associations of phytoplankton and planktic foraminifera in an upwelling region and their contribution to the seafloor, *Mar. Micropaleontol.*, 18(4), 263–278, [https://doi.org/10.1016/0377-8398\(92\)90043-J](https://doi.org/10.1016/0377-8398(92)90043-J), 1992.
- Sautter, L. R. and Thunell, R. C.: Seasonal Variability in the  $\delta^{18}\text{O}$  and  $\delta^{13}\text{C}$  of Planktonic Foraminifera from an Upwelling Environment: Sediment Trap Results from the San Pedro Basin, Southern California Bight, *Paleoceanography*, 6(3), 307–334, <https://doi.org/10.1029/91PA00385>, 1991.
- 680 Schiebel, R. and Hemleben, C.: Modern planktic foraminifera, *Paläontol. Z.*, 79(1), 135–148, <https://doi.org/10.1007/BF03021758>, 2005
- Schiebel, R. and Hemleben, C.: Planktic foraminifera in the modern ocean, 2d ed., Springer Berlin Heidelberg, Berlin, Heidelberg., 2017.
- Schiebel, R., Waniek, J., Bork, M. and Hemleben, C.: Planktic foraminiferal production stimulated by chlorophyll redistribution and entrainment of nutrients, *Deep Sea Research Part I: Oceanographic Research Papers*, 48(3), 721–740, [https://doi.org/10.1016/S0967-0637\(00\)00065-0](https://doi.org/10.1016/S0967-0637(00)00065-0), 2001.
- 685 Schiebel, R., Spielhagen, R. F., Garnier, J., Hagemann, J., Howa, H., Jentzen, A., Martínez-García, A., Meilland, J., Michel, E., Repschläger, J., Salter, I., Yamasaki, M. and Haug, G.: Modern planktic foraminifera in the high-latitude ocean, *Mar. Micropaleontol.*, 136, 1–13, <https://doi.org/10.1016/j.marmicro.2017.08.004>, 2017.
- Schiebel, R., Smart, S. M., Jentzen, A., Jonkers, L., Morard, R., Meilland, J., Michel, E., Coxall, H. K., Hull, P. M., de Garidel-Thoron, T., Aze, T., Quillévéré, F., Ren, H., Sigman, D. M., Vonhof, H. B., Martínez-García, A., Kučera, M., Bijma, J., Spero, H. J. and Haug, G. H.: Advances in planktonic foraminifer research: New perspectives for paleoceanography, *Revue de Micropaléontologie*, 61(3–4), 113–138, <https://doi.org/10.1016/j.revmic.2018.10.001>, 2018.
- 690 78. Schlitzer, R.: Ocean Data View, Computer software, <https://odv.awi.de.>, 2021.
- Smith, R. A., Castañeda, I. S., Groeneveld, J., De Vleeschouwer, D., Henderiks, J., Christensen, B. A., Renema, W., Auer, G., Bogus, K., Gallagher, S. J. and Fulthorpe, C. S.: Plio-pleistocene Indonesian throughflow variability drove eastern Indian ocean sea surface temperatures, *Paleoceanography and Paleoclimatology*, 35(10), <https://doi.org/10.1029/2020PA003872>, 2020.
- 695 Somavilla, R., González-Pola, C. and Fernández-Díaz, J.: The warmer the ocean surface, the shallower the mixed layer. How much of this is true?, *J. Geophys. Res. Oceans*, 122(9), 7698–7716, <https://doi.org/10.1002/2017JC013125>, 2017.
81. Spezzaferri, S., Kucera, M., Pearson, P. N., Wade, B. S., Rappo, S., Poole, C. R., Morard, R. and Stalder, C.: Fossil and genetic evidence for the polyphyletic nature of the planktonic foraminifera “Globigerinoides”, and description of the new genus *Trilobatus*., *PLoS ONE*, 10(5), e0128108, <https://doi.org/10.1371/journal.pone.0128108>, 2015.
- 700 Sprintall, J. and Révelard, A.: The Indonesian Throughflow response to Indo-Pacific climate variability, *J. Geophys. Res. Oceans*, 119(2), 1161–1175, <https://doi.org/10.1002/2013JC009533>, 2014.
- Steinke, S., Prange, M., Feist, C., Groeneveld, J. and Mohtadi, M.: Upwelling variability off southern Indonesia over the past two millennia, *Geophys. Res. Lett.*, 41(21), 7684–7693, <https://doi.org/10.1002/2014GL061450>, 2014.
- 705 Steph, S., RegenberG, M., Tiedemann, R., Mulitza, S. and Nürnberg, D.: Stable isotopes of planktonic foraminifera from tropical Atlantic/Caribbean core-tops: Implications for reconstructing upper ocean stratification, *Mar. Micropaleontol.*, 71(1–2), 1–19, <https://doi.org/10.1016/j.marmicro.2008.12.004>, 2009.




- Susanto, R. D. and Marra, J.: Effect of the 1997/98 el niño on chlorophyll a variability along the southern coasts of java and sumatra, *oceanog.*, 18(4), 124–127, <https://doi.org/10.5670/oceanog.2005.13>, 2005.
- 710 Susanto, R. D., Gordon, A. L. and Zheng, Q.: Upwelling along the coasts of Java and Sumatra and its relation to ENSO, *Geophys. Res. Lett.*, 28(8), 1599–1602, <https://doi.org/10.1029/2000GL011844>, 2001.
- Susanto, R. D., Moore, T. S. and Marra, J.: Ocean color variability in the Indonesian Seas during the SeaWiFS era, *Geochem. Geophys. Geosyst.*, 7(5), <https://doi.org/10.1029/2005GC001009>, 2006.
- Takagi, H., Kimoto, K., Fujiki, T. and Moriya, K.: Effect of nutritional condition on photosymbiotic consortium of cultured Globigerinoides  
 715 sacculifer (Rhizaria, Foraminifera)., *Symbiosis*, 76(1), 25–39, <https://doi.org/10.1007/s13199-017-0530-3>, 2018.
- Tapia, R., Nürnberg, D., Ronge, T. and Tiedemann, R.: Disparities in glacial advection of Southern Ocean Intermediate Water to the South Pacific Gyre, *Earth and Planetary Science Letters*, 410, 152–164, <https://doi.org/10.1016/j.epsl.2014.11.031>, 2015.
- Tapia, R., Nürnberg, D., Ho, S. L., Lamy, F., Ullermann, J., Gersonde, R. and Tiedemann, R.: Glacial differences of Southern Ocean Intermediate Waters in the Central South Pacific, *Quat. Sci. Rev.*, 208, 105–117, <https://doi.org/10.1016/j.quascirev.2019.01.016>, 2019.
- 720 Tillinger, D.: Physical oceanography of the present day Indonesian Throughflow, Geological Society, London, Special Publications, 355(1), 267–281, <https://doi.org/10.1144/SP355.13>, 2011.
- Tolderlund, D. S., Bé, A. W. H. and Be, A. W. H.: Seasonal distribution of planktonic foraminifera in the western north atlantic, *Micropaleontology*, 17(3), 297, <https://doi.org/10.2307/1485143>, 1971.
- Ujjié, H.: Distribution of living planktonic foraminifera in the southeast Indian Ocean, *Bull. Nat. Sci. Mus. Tokyo*, 11(1), 97–125, 1968.
- 725 Verheye, H. M., Hutchings, L. and Peterson, W. T.: Life history and population maintenance strategies of *Calanoides carinatus* (Copepoda: Calanoida) in the southern Benguela ecosystem, *South African Journal of Marine Science*, 11(1), 179–191, <https://doi.org/10.2989/025776191784287600>, 1991.
- Wang, L.: Isotopic signals in two morphotypes of *Globigerinoides ruber* (white) from the South China Sea: implications for monsoon climate change during the last glacial cycle, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 161(3–4), 381–394,  
 730 [https://doi.org/10.1016/S0031-0182\(00\)00094-8](https://doi.org/10.1016/S0031-0182(00)00094-8), 2000.
- Wang, X., Jian, Z., Lückge, A., Wang, Y., Dang, H. and Mohtadi, M.: Precession-paced thermocline water temperature changes in response to upwelling conditions off southern Sumatra over the past 300,000 years, *Quat. Sci. Rev.*, 192, 123–134, <https://doi.org/10.1016/j.quascirev.2018.05.035>, 2018.




**Figure 1.** (a) Distribution of the in-situ sea surface temperature in the study area during PABESIA cruise on board of R/V Sonne in August 2005 (Hebbeln and cruise participants, 2006). The study area was divided into seven land-sea transects (numbered rectangles). The stations are divided into plankton net (white circles) and water (red diamond). The location of the sediment trap (Mohtadi et al., 2009) and surface sediments (Mohtadi et al., 2007) are indicated by triangle and squares, respectively. Upper ocean chlorophyll-*a* (Chl-*a*) (retrieved from <https://oceancolor.gsfc.nasa.gov>) during (b) January (Non-monsoon) and (c) August (SE-monsoon) 2005 i.e., the sampling period. Temperature map was created with Ocean Data View 5.2.1 (Schlitzer, 2021).




**Figure 2.** Cross sections of (a) water temperature, (b) salinity, (c) Chl-*a* and (d) oxygen content of the upper 600 m of the water column during the sampling period. Location of the stations comprising the cross section are depicted in the inset of panel (a). Thermal mixed layer depth (MLD<sub>TEMP</sub>), density mixed layer depth (MLD<sub>DEN</sub>) and barrier layer (BL) are marked in panel (a). Numbers and horizontal lines on top show each transect (1 to 7). Sampled depths depicted by gray squares (Hebbeln and cruise participants, 2006). Visualisation is based on Ocean Data View 5.2.1 (Schlitzer, R., 2021).




**Figure 3.** Non-metric multidimensional scaling of the in-situ hydrographic parameters shows three main clusters. Cluster 1 includes transects 5 to 7 off Java and the LSI influenced by upwelling (upwelling; blue). Cluster 3 includes transects 1 to 3 located off Sumatra which are not influenced by upwelling (non-upwelling; red). Cluster 2 consists of only transect 4 that represents a transitional condition between Sumatra and Java-LSI (green). Location of the stations are shown in the inset at the top right corner of the figure.



**Figure 4.** Summary of planktic foraminifera census counts off Indonesia. a) Species with contribution >2% to the total assemblage, b) abundance distribution of planktic foraminifera in transects 1 to 7 and, c) abundance of planktic foraminifera by water depth interval at each station. Numbers in (b) represent transects 1 to 7 (italic) and median values of transects (bold); colors in (b) depict sectors off Sumatra (red), transitional (green) and off Java-LSI (blue). The five sampled water depths are: 0–25 m (tangerine), 25–50 m (orange), 50–100 m (white), 100–200 m (lilac), and 200–500 m (purple). Stations are grouped by transect (number). [Schematic of the box plots showing the median value \(horizontal line\), the whisker marks the minimum \(min\) and maximum \(max\) values.](#) Note the different scales for y-axis in b (log scale) and c (linear scale).




**Figure 5.** Cross sections showing the vertical distribution across the water column of a) the total community of planktic foraminifera and the species; b) *T. trilobus*, c) *G. ruber* (white), d) *G. bulloides*, e) *N. dutertrei*, f) *P. obliquiloculata*, and g) *G. menardii*. Numbers and horizontal lines on top of (a) show each transect (1 to 7). Visualisation is based on Ocean Data View 5.2.1 (Schlitzer, R., 2021).




**Figure 6.** Average Living Depth (ALD) of the species *T. trilobus*, *G. ruber* (white), *G. bulloides*, *N. dutertrei*, *P. obliquiloculata* and *G. menardii*. The calculation includes only stations with more than 5 specimens per species (n). Inset is a schematic of the box plots the median value is the horizontal line the whisker marks the minimum (min) and maximum (max) values, white circles indicate ALD values. Note log scale for y-axis.







**Figure 7.** Planktic foraminiferal assemblage in (a and b) plankton nets (this study) and (c and d) surface sediments (data from Mohtadi et al., 2007) off Sumatra (Transect 1–3) and off Java-LSI (Transect 5–7) samples. Mohtadi et al. (2007) did not differentiate between morphotypes; \**G. ruber* data comprise both morphotypes (*G. ruber* (white) + *G. elongatus*). Note that *G. ruber* (white) and *G. elongatus* data from plankton nets are plotted side by side to facilitate visual comparison with \**G. ruber* in sediment data.



**Figure 8.** Average Living Depths (ALD) distribution for a) *T. trilobus*, b) *G. ruber* (white), c) *G. bulloides*, d) *N. dutertrei*, e) *P. obliquiloculata*, and f) *G. menardii* per transect off Sumatra (red), transitional (green), and Java-LSI (blue). The calculation only considers stations with more than 5 counts per species. The yellow and light blue boxes depict the 1st and 3rd quartiles of the ALD values off Sumatra (non-upwelling) and off Java-LSI (upwelling); black bars and numbers depict the median value per sector. Boxes depict the inferred vertical distribution from surface sediments (gray) along Indonesia and from the sediment trap (pattern) off Java (Mohtadi et al., 2009; 2011). Colored dots and vertical lines depict calculated mean ALDs and vertical dispersion based on living foraminifera data from areas influenced by the Benguela (green) and Canary (purple) upwelling systems (Lessa et al., 2020; Rebotim et al., 2017). ALD for other species are presented in Appendices Fig. A.3



**Figure 9.** Comparison of the thermal gradient ( $\Delta T$ ) off Indonesia inferred from surface sediments (Mg/Ca data from Mohtadi et al., 2011), plankton net (abundance-weighted temperatures, this study) and from sediment trap (Mg/Ca data from Mohtadi et al., 2009). During the SE monsoon the  $\Delta T$  calculated from the plankton net data shows that larger  $\Delta T$  values occur off Sumatra (non-upwelling) than off Java (upwelling) in agreement with the seasonal  $\Delta T$  off Java calculated from sediment trap (circles) with larger ( $\Delta T$ ) occurring during the non-upwelling season than the upwelling season. The color represents the inferred averaged temperatures for the mixed layer (combining *G. ruber* and *T. trilobus*; red) and thermocline (combining *N. dutertrei* and *P. obliquiloculata*; blue), numbers represent median values. Outliers and extreme outliers,  $>1.5$  and  $>3$  times the Inner Quartile Range, are depicted by filled and open circles, respectively.




**Figure 10.** Schematic comparing the  $\Delta T$  derived from mixed layer and thermocline-dwellers off Sumatra and Java-LSI, from a) surface sediment (inferred habitat depth from Mohtadi et al., 2011) matching the mean annual conditions and b) ALD from plankton net (ALD) matching the water structure during the sampling period (SE-monsoon; August-September, 2005). Gray lines depict the average annual mean water structure offshore Sumatra and Java-LSI (band of 160 km wide; >200 stations from WOA2018; 0.25°) (Locarnini et al., 2018); dashed lines depict the average water column structure in each sector during the SE-monsoon based on in situ data collected during the PABESIA cruise (August-September, 2005); black line shows the average depth of the  $MLD_{TEMP}$  (Table A1 and A2).

**Table 1.** 95% confidence interval of the habitat depth of 6 species of planktic foraminifera in upwelling vs. non-upwelling regions. Upwelling region consists of transects 5–7, while non-upwelling region consists of transects 1–3. Details of the calculation are described in section Method.


| Species                            | Non-Upwelling    |                  |                           | Upwelling        |                  |                           | P-value |
|------------------------------------|------------------|------------------|---------------------------|------------------|------------------|---------------------------|---------|
|                                    | Lower 95% CI (m) | Upper 95% CI (m) | Mean ALD (upw – nupw) (m) | Lower 95% CI (m) | Upper 95% CI (m) | Mean ALD (upw – nupw) (m) |         |
| <i>T. trilobus</i>                 | 30.5             | 48.4             | 49.5                      | 71.7             | 103.1            | 49.5                      | <0.001* |
| <i>G. ruber</i> <sub>(white)</sub> | 56.5             | 87.1             | 29.1                      | 82.4             | 118.4            | 29.1                      | 0.017*  |
| <i>G. bulloides</i>                | 55.6             | 156.1            | 45.2                      | 118.1            | 175.3            | 45.2                      | 0.057   |
| <i>P. obliquiloculata</i>          | NA               | NA               | NA                        | 81.3             | 105.8            | NA                        | -       |
| <i>N. dutertrei</i>                | NA               | NA               | NA                        | 72.9             | 102.1            | NA                        | -       |

\*Difference in ALD for the two regions is statistically significant. Abbreviation: CI for confidence interval; ALD for average living depth. Upwelling minus Non-upwelling (upw – nupw); NA for no data available

## Appendix A




**Figure A1.** Longitudinal distribution of temperature and salinity anomalies across the upper 500 m of water column. Anomalies are calculated by subtracting the mean of all sites for a given depth interval (e.g., 0–25 m, 25–50 m, etc) from the data of a station for the same depth interval. Positive (negative) values are centered off Sumatra (off Java) showing strong zonal trend. The largest temperature change occurs at 50–100 m, roughly corresponding to the depth of the thermocline.



**Figure A2.** Size class distribution of the six selected species. *Globigerinoides ruber*, *N. dutertrei*, and *G. menardii* show a reduction in the proportion of the size class >250 μm after full moon (blue). The large difference in their median values suggest synchronized reproduction; *P. obliquiloculata* shows no differences in the size class distribution before (red) and after (blue) full moon; *Globigerina bulloides* and *T. trilobus* show an increase in the proportion of larger organisms after the full moon the 1st and 3rd quartiles; median values are shown by bar and circles represent mean values.





**Figure A3.** In addition to species with paleoceanographic relevance, we calculated the (a) ALD for the species *G. glutinata*(84 m), *G. falconensis* (139 m), *G. siphonifera* (85 m), *G. calida* (88 m), and *G. hirsuta* (160 m). The zonal comparison of these species (b-f) is only possible for three species; *G. calida* shows the smallest change in ALD (~5 m) while *G. glutinata* and *G. siphonifera* show a change of 30 and 50 m in their habitat depth between sectors. The yellow and light blue boxes depict the 1st and 3rd quartiles of the ALD values off Sumatra (non-upwelling) and, off Java-LSI (upwelling). Black bars and numbers depict the median value per sector.

**Table A1.** Details of the plankton net stations (location, water depth, and sampling date) during cruise SO-184 (Hebbeln and cruise participants, 2006). The stations are divided into seven transects. Relevant environmental parameters during the sampling include the thermal mixed layer depth (MLD<sub>TEMP</sub>), barrier layer thickness (BL), Stratification index (SI<sub>0-200</sub>), observed sea surface temperature (SST), sea surface salinity (SSS), and in-situ chlorophyll-a (Chl-a) at the surface (0-25 m), and thermocline level (100-200 m) (Table 5.1 in Hebbeln and cruise participants, 2006). Sampling gear: Multinet (MN), Rosette Sampler (Rs), No data available (ND).

| Transect | Station | GeoB   | Lat.    | Long.    | Date     | MLD <sub>TEMP</sub> | BL   | SI <sub>0-200</sub> | SSS   | SST   | Oxygen | Chl-a              |          | Gear   | Depth |
|----------|---------|--------|---------|----------|----------|---------------------|------|---------------------|-------|-------|--------|--------------------|----------|--------|-------|
|          |         |        |         |          |          |                     |      |                     |       |       |        | 0-25 m             | 100-200m |        |       |
|          |         |        | N       | E        | dd.mm.yy |                     | m    | m                   | psu   | °C    | mlL-1  | mg m <sup>-3</sup> |          |        | m     |
| <b>1</b> | 10018   |        | 1.570   | 96.512   | 08.08.05 | 83.6                | 51   | 17.6                | 33.7  | 29.9  | 4.9    | 0.646              | 0.220    | Mn, Rs | 2577  |
|          | 10019   |        | 1.632   | 96.885   | 08.08.05 | 83.2                | 55   | 17.4                | 33.83 | 29.9  | 4.7    | 0.641              | 0.228    | Mn, Rs | 1465  |
|          | 10020   |        | 1.678   | 96.980   | 09.08.05 | 77.8                | 52   | 17.5                | 33.83 | 30.0  | 4.7    | 0.641              | 0.241    | Mn, Rs | 1160  |
|          | 10011   |        | -1.191  | 97.986   | 06.08.05 | 82.3                | 70   | 16.8                | 34.0  | 29.6  | 4.4    | 0.634              | 0.253    | Mn, Rs | 3030  |
|          | 10012   |        | -1.070  | 98.058   | 06.08.05 | 89.4                | 40   | 17.1                | 34.0  | 29.6  | 4.4    | 0.629              | 0.248    | Rs     | 2096  |
| <b>2</b> | 10013   |        | -0.958  | 98.266   | 07.08.05 | 76.0                | 30   | 17.0                | 34.0  | 29.6  | 4.3    | 0.619              | 0.212    | Rs     | 927   |
|          | 10022   |        | -0.051  | 98.850   | 10.08.05 | 71.5                | 52   | 17.1                | 33.8  | 29.7  | 4.4    | 0.639              | 0.227    | Mn, Rs | 707   |
|          | 10023   |        | -0.857  | 99.407   | 11.08.05 | 70.4                | 43   | 17.4                | 33.8  | 29.6  | 4.4    | 0.635              | 0.204    | Mn, Rs | 1557  |
|          | 10024   |        | -0.769  | 99.269   | 11.08.05 | 68.4                | 42   | 17.7                | 33.8  | 29.6  | 4.4    | 0.637              | 0.245    | Mn, Rs | 1384  |
|          | 10025   |        | -0.675  | 99.123   | 11.08.05 | 67.9                | 34   | 17.3                | 33.8  | 29.6  | 4.6    | 0.654              | 0.245    | Mn, Rs | 1148  |
| <b>3</b> | 10026   |        | -0.944  | 99.521   | 12.08.05 | ND                  | ND   | ND                  | ND    | ND    | ND     | ND                 | ND       | Mn     | 1636  |
|          | 10027   |        | -0.809  | 99.653   | 12.08.05 | 64.4                | 52   | 17.6                | ND    | ND    | ND     | ND                 | ND       | Mn, Rs | 876   |
|          | 10028   |        | -0.696  | 99.763   | 12.08.05 | 62.0                | 45   | 18.0                | 33.71 | 29.8  | 4.4    | 0.631              | 0.209    | Mn, Rs | 521   |
|          | 10029   |        | -1.505  | 100.131  | 13.08.05 | 76.3                | 48   | 18.1                | 33.74 | 29.6  | 4.5    | 0.644              | 0.211    | Rs     | 962   |
|          | 10030   |        | -1.638  | 99.774   | 13.08.05 | 62.4                | 39   | 17.6                | 33.73 | 29.5  | 4.6    | 0.662              | 0.209    | Rs     | 1757  |
| <b>4</b> | 10031   |        | -1.708  | 99.607   | 13.08.05 | 55.4                | 21   | 17.8                | 33.80 | 29.5  | 4.4    | 0.627              | 0.248    | Rs     | 1661  |
|          | 10003   |        | -4.751  | 100.767  | 03.08.05 | 90.9                | 72   | ND                  | 33.93 | 29.10 | 4.5    | 0.280              | 0.000    | Rs     | 3176  |
|          | 10007   |        | -4.354  | 100.996  | 04.08.05 | 85.8                | 54   | 16.8                | 33.70 | 29.2  | 4.5    | 0.338              | 0.239    | Mn, Rs | 598   |
|          | 10034   |        | -4.165  | 101.499  | 15.08.05 | 62.8                | 21   | 16.6                | 33.83 | 29.3  | 4.6    | 0.654              | 0.205    | Mn, Rs | 992   |
|          | 10035   |        | -4.036  | 101.733  | 15.08.05 | 66.7                | 25   | 16.9                | 33.82 | 29.3  | 4.5    | 0.652              | 0.228    | Mn, Rs | 997   |
| 10036    |         | -5.338 | 103.657 | 16.08.05 | 45.5     | 10                  | 17.7 | 33.91               | 29.2  | 4.5   | 0.650  | 0.241              | Mn, Rs   | 1498   |       |

**Table A2.** continued Table A1

| Transect | Station | Lat.    | Long.    | Date     | MLD <sub>TEMP</sub> | BL   | SI <sub>0-200</sub> | SSS   | SST   | Oxygen             |                   | Chl- <i>a</i> | Gear   | Depth |
|----------|---------|---------|----------|----------|---------------------|------|---------------------|-------|-------|--------------------|-------------------|---------------|--------|-------|
|          |         |         |          |          |                     |      |                     |       |       | mg m <sup>-3</sup> | 0-2.5 m 100-200 m |               |        |       |
|          | GeoB    | N       | E        | dd.mm.yy | m                   | m    | psu                 | °C    | mlL-1 |                    |                   | m             | m      |       |
| <b>4</b> | 10038   | -5.937  | 103.245  | 17.08.05 | 51.2                | 39   | 16.5                | 33.48 | 28.50 | 4.52               | 0.651             | 0.227         | Mn, Rs | 1887  |
|          | 10039   | -5.867  | 103.294  | 17.08.05 | 53.9                | 36   | 16.7                | 33.54 | 28.56 | 4.50               | 0.646             | 0.246         | Mn, Rs | 1797  |
|          | 10040   | -6.475  | 102.857  | 18.08.05 | 62.0                | 25   | 16.5                | 33.49 | 28.00 | 4.57               | 0.657             | 0.236         | Mn, Rs | 2602  |
|          | 10041   | -6.274  | 103.008  | 18.08.05 | 51.9                | 19   | 16.4                | 33.92 | 28.45 | 4.50               | 0.647             | 0.232         | Mn, Rs | 1540  |
|          | 10043   | -7.310  | 105.062  | 19.08.05 | 34.1                | 3    | 16.8                | 34.07 | 27.55 | 4.47               | 0.642             | 0.398         | Rs     | 2161  |
| <b>5</b> | 10044   | -8.055  | 109.015  | 22.08.05 | 33.5                | 2    | 16                  | 33.93 | 28.53 | 5.39               | 0.775             | 0.301         | Mn, Rs | 3358  |
|          | 10045   | -8.743  | 109.020  | 23.08.05 | 40.3                | 6    | 15.5                | 34.25 | 26.52 | 4.72               | 0.677             | 0.326         | Mn, Rs | 3571  |
|          | 10046   | -9.604  | 109.063  | 24.08.05 | 50.2                | 1    | 14.9                | 34.24 | 25.90 | 4.81               | 0.691             | 0.376         | Mn, Rs | 2604  |
|          | 10048   | -8.255  | 108.147  | 25.08.05 | 43.2                | 0    | 15.8                | 34.21 | 26.74 | 4.78               | 0.687             | 0.335         | Mn, Rs | 3060  |
|          | 10049   | -8.785  | 110.496  | 26.08.05 | 37.7                | 1    | 14.5                | 34.45 | 25.24 | 4.70               | 0.676             | 0.335         | Mn, Rs | 1291  |
| <b>6</b> | 10051   | -9.293  | 110.497  | 27.08.05 | 31.7                | 5    | 14                  | 34.15 | 25.89 | 4.78               | 0.687             | 0.364         | Mn, Rs | 2391  |
|          | 10052   | -8.694  | 110.634  | 28.08.05 | 35.9                | 6    | 14.9                | 34.40 | 25.60 | 4.88               | 0.701             | 0.350         | Mn, Rs | 1000  |
|          | 10053   | -8.677  | 112.872  | 29.08.05 | 23.7                | 0    | 14.6                | 34.36 | 24.43 | 4.17               | 0.600             | 0.321         | Mn, Rs | 1378  |
|          | 10054   | -8.681  | 112.668  | 29.08.05 | 19.0                | 1    | 13.7                | 34.34 | 24.59 | 4.16               | 0.599             | 0.314         | Rs     | 1069  |
|          | 10055   | -9.248  | 113.050  | 30.08.05 | 23.8                | 0    | 14.9                | 34.34 | 24.59 | 4.16               | 0.599             | 0.314         | Mn, Rs | 2615  |
| <b>7</b> | 10057   | -9.822  | 113.107  | 31.08.05 | 26.3                | 1    | 14.3                | 34.26 | 25.94 | 4.76               | 0.684             | 0.348         | Mn, Rs | 1615  |
|          | 10061   | -9.729  | 113.024  | 02.09.05 | 12.7                | 2    | 14.3                | 34.35 | 23.65 | 4.29               | 0.617             | 0.354         | Mn, Rs | 2174  |
|          | 10062   | -11.166 | 115.999  | 03.09.05 | 76.4                | 1    | 12.6                | 34.24 | 26.94 | 4.67               | 0.671             | 0.390         | Mn, Rs | 5851  |
|          | 10063   | -9.646  | 118.149  | 04.09.05 | 24.4                | 0    | 13.8                | 34.15 | 25.61 | 4.78               | 0.687             | 0.385         | Mn, Rs | 2498  |
|          | 10064   | -9.539  | 118.304  | 04.09.05 | ND                  | ND   | ND                  | ND    | ND    | ND                 | ND                | ND            | Mn     | 2033  |
| 10065    | -9.223  | 118.894 | 05.09.05 | 56.4     | 3                   | 13.4 | 34.19               | 25.43 | 4.49  | 0.645              | 0.409             | Mn, Rs        | 1286   |       |
| 10066    | -9.394  | 118.575 | 05.09.05 | 29.5     | 4                   | 13.6 | 34.18               | 26.11 | 4.80  | 0.690              | 0.412             | Mn, Rs        | 1630   |       |
| 10067    | -9.149  | 119.290 | 06.09.05 | 38.0     | 9                   | 14.6 | 34.13               | 26.43 | 4.75  | 0.682              | 0.441             | Mn, Rs        | 1136   |       |
| 10068    | -9.595  | 121.152 | 07.09.05 | 16.0     | 2                   | 15.4 | 34.12               | 26.87 | 4.71  | 0.677              | 0.428             | Rs            | 2011   |       |
| 10069    | -9.608  | 120.921 | 07.09.05 | 14.2     | 2                   | 14.2 | 34.12               | 26.62 | 4.77  | 0.686              | 0.422             | Rs            | 1264   |       |
| 10070    | -10.359 | 121.303 | 08.09.05 | 22.2     | 0                   | 13.6 | 34.13               | 26.94 | 4.82  | 0.693              | 0.406             | Mn, Rs        | 1509   |       |