



## Soil carbon loss in warmed subarctic grasslands is rapid and restricted to topsoil

Niel Verbrigghe<sup>1</sup>, Niki I. W. Leblans<sup>1,2</sup>, Bjarni D. Sigurdsson<sup>3</sup>, Sara Vicca<sup>1</sup>, Chao Fang<sup>1,4,5</sup>, Lucia Fuchslueger<sup>1,6</sup>, Jennifer L. Soong<sup>1,7</sup>, James T. Weedon<sup>8</sup>, Christopher Poeplau<sup>9</sup>, Cristina Ariza-Carricondo<sup>1</sup>, Michael Bahn<sup>10</sup>, Bertrand Guenet<sup>11</sup>, Per Gundersen<sup>12</sup>, Gunnhildur E. Gunnarsdóttir<sup>13</sup>, Thomas Kätterer<sup>14</sup>, Zhanfeng Liu<sup>15</sup>, Marja Maljanen<sup>16</sup>, Sara Marañón-Jiménez<sup>17,18</sup>, Kathiravan Meeran<sup>10</sup>, Edda S. Oddsdóttir<sup>19</sup>, Ivika Ostonen<sup>20</sup>, Josep Peñuelas<sup>17,18</sup>, Andreas Richter<sup>6,21</sup>, Jordi Sardans<sup>17,18</sup>, Páll Sigurðsson<sup>3</sup>, Margaret S. Torn<sup>22</sup>, Peter M. Van Bodegom<sup>23</sup>, Erik Verbruggen<sup>1</sup>, Tom W. N. Walker<sup>24</sup>, Håkan Wallander<sup>25</sup>, and Ivan A. Janssens<sup>1</sup>

<sup>1</sup>Research Group Plants and Ecosystems, University of Antwerp, Antwerp, Belgium.

<sup>2</sup>Climate Impacts Research Centre, Umeå University, Umeå, Sweden.

<sup>3</sup>Agricultural University of Iceland, Hvanneyri, Borgarnes, Iceland.

<sup>4</sup>Institute of Ecology, School of Applied Meteorology, Nanjing University of Information Science and Technology, Nanjing, China.

<sup>5</sup>State Key Laboratory of Grassland Agro-ecosystems, Institute of Arid Agroecology, School of Life Sciences, Lanzhou University, Lanzhou, China.

<sup>6</sup>Centre for Microbiology and Environmental Systems Science, University of Vienna, Vienna, Austria.

<sup>7</sup>Soil and Crop Sciences Department, Colorado State University, Fort Collins, Colorado, USA.

<sup>8</sup>Systems Ecology, Department of Ecological Science, Vrije Universiteit Amsterdam, Amsterdam, The Netherlands.

<sup>9</sup>Thünen Institute of Climate-Smart Agriculture, Braunschweig, Germany.

<sup>10</sup>Department of Ecology, University of Innsbruck, Innsbruck, Austria.

<sup>11</sup>Laboratoire de Géologie, École normale supérieure/CNRS, PSL Research University, Paris, France.

<sup>12</sup>Department of Geosciences and Natural Resource Management, University of Copenhagen, Frederiksberg C, Denmark.

<sup>13</sup>Soil Conservation Service of Iceland, Gunnarsholt, Hella, Iceland.

<sup>14</sup>Department of Ecology, Swedish University of Agricultural Sciences, Uppsala, Sweden.

<sup>15</sup>Key Laboratory of Vegetation Restoration and Management of Degraded Ecosystems & CAS Engineering Laboratory for Vegetation Ecosystem Restoration on Islands and Coastal Zones, South China Botanical Garden, Chinese Academy of Sciences, Guangzhou, China.

<sup>16</sup>Department of Environmental and Biological Sciences, University of Eastern Finland, Kuopio, Finland.

<sup>17</sup>CREAF, Cerdanyola del Vallès, Barcelona, Catalonia, Spain.

<sup>18</sup>CSIC, Global Ecology Unit CREAM-CSIC-UAB, Bellaterra, Barcelona, Spain.

<sup>19</sup>Icelandic Forest Research, Mógilsá, Reykjavík, Iceland.

<sup>20</sup>Institute of Ecology and Earth Sciences, University of Tartu, Tartu, Estonia

<sup>21</sup>International Institute for Applied Systems Analysis (IIASA), Laxenburg, Austria.

<sup>22</sup>Climate and Ecosystem Sciences Division, Berkeley Lab, Berkeley, CA, USA.

<sup>23</sup>Environmental Biology Department, Institute of Environmental Sciences, CML, Leiden University, Leiden, The Netherlands.

<sup>24</sup>Department of Environmental Systems Science, ETH Zürich, Zürich, Switzerland.

<sup>25</sup>MEMEG, Department of Biology, Lund University, Lund, Sweden.

**Correspondence:** Niel Verbrigghe (Niel.Verbrigghe@UAntwerpen.be)



**Abstract.** Global warming may lead to carbon transfers from soils to the atmosphere, yet this positive feedback to the climate system remains highly uncertain, especially in subsoils (Ilyina and Friedlingstein, 2016; Shi et al., 2018). Using natural geothermal soil warming gradients of up to +6.4 °C in subarctic grasslands (Sigurdsson et al., 2016), we show that soil organic carbon (SOC) stocks decline strongly and linearly with warming ( $-2.8 \text{ ton ha}^{-1} \text{ °C}^{-1}$ ). Comparison of SOC stock changes following medium-term (5 and 10 years) and long-term (>50 years) warming revealed that all SOC loss occurred within the first five years of warming, after which continued warming no longer reduced SOC stocks. This rapid equilibration of SOC observed in Andosol suggests a critical role for ecosystem adaptations to warming and could imply short-lived soil carbon-climate feedbacks. Our data further revealed that the soil C loss occurred in all aggregate size fractions, and that SOC losses only occurred in topsoil (0-10 cm). SOC stocks in subsoil (10-30 cm), where plant roots were absent, remained unaltered, even after >50 years of warming. The observed depth-dependent warming responses indicate that explicit vertical resolution is a prerequisite for global models to accurately project future SOC stocks for this soil type and should be investigated for soils with other mineralogies.

## 1 Introduction

Soils store more carbon (C) than the atmosphere and vegetation biomass combined (Batjes, 2016; Scharlemann et al., 2014). Global warming has been hypothesised to lead to increased soil CO<sub>2</sub> emissions that may lead to large reductions in soil organic carbon (SOC) stocks, constituting a positive feedback to the climate system (Davidson and Janssens, 2006; Jenkinson et al., 1991). The strength and even sign of this carbon cycle-climate feedback are, however, highly uncertain (Crowther et al., 2016; Todd-Brown et al., 2018; van Gestel et al., 2018). Accordingly, the World Climate Research Programme has acknowledged it as one of the "Grand Challenges" of climate research (Ilyina and Friedlingstein, 2016).

*In situ* soil warming studies provide ideal tools to study the response of soil SOC stocks to warming (Batjes, 2016), yet challenges remain great. First, ecosystem responses to warming may take decades to stabilise (Walker et al., 2020; Melillo et al., 2017), implying that extrapolations of responses from-, or model parametrisation based on short-term experiments may lead to erroneous estimation of the future evolution of SOC stocks. Second, SOC stock changes are rarely studied in subsoils. The high cost and labour requirements of SOC research, combined with the fact that most biological activity and SOC mineralisation occur in topsoils, explains why soil biology and ecology, including SOC cycling, are rarely studied below a depth of 20-30 cm (Yost and Hartemink, 2020). This is a major issue, because more SOC is stored below this threshold than above (Shi et al., 2020) and therefore the carbon cycle-climate feedback does not stop at 20-30 cm depth. Unfortunately, the very few soil warming experiments that also warmed subsoils and quantified SOC stock changes yielded very different warming responses, ranging from declining to increasing subsoil SOC stocks (Soong et al., 2020a; Hanson et al., 2020).

To address both these challenges, we determined SOC stock changes along natural geothermal gradients at the ForHot research site in Iceland (Sigurdsson et al., 2016) encompassing the full warming range projected for Northern regions (up to +6.4 °C), throughout the topsoil (0-10 cm) and the subsoil (10-30 cm). We compared topsoil and subsoil SOC dynamics along replicate warming gradients exposed to medium-term (5 and 10 years) and long-term (>50 years, but possibly centuries)



warming by sampling permanent study plots twice in a six-year period (2013 and 2018). This enabled us to characterise the  
35 magnitude, shape and temporal dynamics of the temperature response of SOC stocks in these northern, non-permafrost, soils.  
Next to measuring SOC stocks, we gathered data about soil aggregates, carbon inputs to the soil by plants and arbuscular  
mycorrhizal fungi and carbon flux from topsoil to subsoil. This allowed us to elaborate about the possible mechanisms behind  
SOC stock changes along the warming gradient.

The recently warmed grassland we investigated at the ForHot site has been warmed since 2008, when a major earthquake  
40 shifted geothermal systems to previously unwarmed soils, causing increased temperature in the soil above by radiative heating  
(Halldórsson and Sigbjörnsson, 2009; O’Gorman et al., 2014). In contrast, the long-term warmed grassland had been warmed  
for at least 45 years at the time of the earthquake in 2008 (Sigurdsson et al., 2016). The soil type on both study sites is Andosol,  
and they are covered by the same grassland type (Sigurdsson et al., 2016).

We hypothesised that because of the slow reaction of protected SOC pools to temperature change, medium-term warmed  
45 soils would still be losing SOC over time, while the long-term warmed soils would have reached a new equilibrium at lower  
SOC content. We further hypothesised similar subsoil and topsoil SOC loss, given that subsoils were exposed to the same  
warming intensity and duration as topsoils.

## 2 Large, linear and fast topsoil SOC loss


Topsoil (0-10 cm; comprising the A horizon and rooting zone) SOC stocks linearly declined by  $2.8 \pm 0.5$  ( $\pm$  SE) ton SOC ha<sup>-1</sup> °C<sup>-1</sup>  
50 soil warming (or  $8.8 \pm 2.1$  °C<sup>-1</sup>; both  $P < 0.001$ ) for mass-corrected SOC stocks (fig. 1a, 2). These topsoil temperature re-  
sponses did not differ between the medium-term and the long-term warmed grassland, i.e., the warming:grassland interaction  
term was not significant ( $P = 0.47$ ). This clearly suggests that warming induced SOC losses only during the initial five years of  
exposure and that SOC stocks did not change thereafter.

However, several sources of variation such as sampling errors and the large heterogeneity inherent to soils, induced quite  
55 broad uncertainty intervals that reduced the potential to detect statistically significant changes in SOC stocks or in their tempera-  
ture response. Hence, to demonstrate that the warming-induced SOC stock loss had indeed stabilised within five years of warm-  
ing and did not further loose SOC, we calculated what SOC stock decline could have remained undetected given the variability  
in our samples using a one-sided 95 % confidence interval on the soil warming regression coefficient. This shows that average  
additional SOC stock losses smaller than  $0.88$  ton C ha<sup>-1</sup> °C<sup>-1</sup> would not have been detected at  $p < 0.05$ , implying that in the five  
60 year period following the initial warming response (i.e., 2013-2018), annual declines of up to  $0.18$  ton C ha<sup>-1</sup> °C<sup>-1</sup> year<sup>-1</sup> would  
have remained undetected. In the subsequent time span of  $>50$  years, only changes smaller than  $0.018$  ton C ha<sup>-1</sup> °C<sup>-1</sup> year<sup>-1</sup>  
would have remained undetected, i.e., a rate 30-fold less than the SOC loss observed in the initial 5 years of soil warming.


Also when not corrected for warming-induced density changes, SOC stocks and soil C concentrations declined with warming  
(fig. B1; fig. B2) and did not further decrease after five years of soil warming. In contrast to our hypothesis, our data thus  
65 revealed that in topsoil, a stepwise increase in temperature caused a fast SOC stock loss that stabilised within five years of



warming, despite the sustained higher temperatures. Even grasslands that had been warmed at least 55 years exhibited no larger SOC loss than that observed after 5 years of soil warming.




**Figure 1. Soil organic carbon (SOC) stocks (ton C ha<sup>-1</sup>) along soil warming gradients, a) in the topsoil (0-10 cm) and b) in the subsoil (10-30 cm) after a soil mass correction.** The regression (solid line; dashed lines represent the 95 % confidence interval; regression details provided in the inset) for medium-term warmed (MTW) and long-term warmed (LTW) grassland (in topsoil) for both 2013 and 2018 were combined, since no soil temperature x warming-duration interaction effect, nor a main effect for warming-duration or sampling year was found. The soil mass correction is visualised in fig. B8. The uncorrected SOC stocks yield qualitatively similar conclusions (fig. B1), as did the C percentage in top- and subsoil (fig. B2). (n = 78 & 40 for topsoil and subsoil respectively)



**Figure 2.** Warming effect on topsoil (0-10 cm) SOC stocks, as observed during repeated sampling campaigns. Stocks after 5 and 10 years of warming are sampled in the medium-term warmed grasslands, stocks after >50 and >55 years in the long-term warmed grasslands. The data at the start of warming is interpolated from the ambient plots in grasslands combined. Soils are divided in four warming categories for representation. The colours on the heatmap and the smoother lines are based on a linear regression equation per sampling event.

To gain insight in the warming-induced soil physical changes and their effects on SOC stocks, soils were fractionated into different size classes that were analysed separately (see Methods). Aggregate fractionation showed that with increasing warming intensity, the mass of >2 mm fraction declined significantly, in favour of the >250  $\mu\text{m}$  and >63  $\mu\text{m}$  fractions. No significant change was detected in the mass of the smallest (<63  $\mu\text{m}$ ) fraction (fig. 3). Opposed to this contrasting response of relative mass, all soil fractions exhibited similar soil C % declines with soil warming (fig. 3). The relative mass increase of the smaller fractions was compensated for by the soil C % decline, resulting in a stable amount of C in the >250  $\mu\text{m}$  and >63  $\mu\text{m}$  fractions. As a result, all of the warming-induced SOC stock decline we observed in the bulk soil, was attributable to C losses in the >2 mm fraction.



**Figure 3.** Relative mass, soil C % and absolute soil C amount of soil aggregate fractions originating from topsoil in the medium-term warmed (MTW) and long-term warmed (LTW) grassland. Darker lines indicate smaller fractions. Fractions significantly affected by soil warming are represented with solid lines, non-significant relations upon soil warming are shown with dashed lines. ( $n = 17$  &  $16$  for each fraction in MTW and LTW grassland respectively)

We suggest that the rapid topsoil SOC loss observed under warming, as well as its attenuation in the medium-term, emerged from the interplay between soil microbial biomass and activity. Warming at the same study site accelerated microbial growth and respiration (Marañón-Jiménez et al., 2018; Walker et al., 2020), which, in the absence of increased plant inputs to soil (fig. B3), caused the initial SOC loss observed here (fig. B1a). In turn, the warming-induced SOC loss caused a decline in micro-



80 bial biomass, creating a negative feedback on microbial activity that we presume prevented further SOC loss (Walker et al.,  
2018, 2020). Alternatively, ephemeral SOC loss under warming may have resulted from physiological adaptations (Allison  
et al., 2010; Bradford et al., 2019) or compositional shifts (Melillo et al., 2017) in the microbial community, but we found  
no evidence this occurred here. Previous research in these grasslands showed that soil microbial carbon use efficiency (CUE)  
remained constant under short- and long-term warming (Walker et al., 2020), and microbial community composition was only  
85 affected by more intense ( $>9^{\circ}\text{C}$ ) long-term warming (Radujković et al., 2018). Aggregate fractionation suggests C % all size  
fractions were impacted similarly by warming (fig. 3). This likely indicates both particulate organic matter, often occluded in  
large-size aggregates, and mineral-associated organic matter, present in all aggregate sizes, decreased with warming.

### 3 Stable subsoil SOC stocks

We hypothesised that the similar warming intensity across the soil profile (fig. B4) would elicit similar declines in subsoil  
90 SOC stocks than those in topsoil. In contrast, SOC remained constant in the subsoil, even under 50 years of soil warming  
in the long-term warmed grasslands ( $P = 0.63$ ; fig. 1b, 2). This lack of SOC loss from the subsoil may be explained by  
three, potentially co-occurring mechanisms. First, limited fresh C inputs from litter and root exudates below the rooting zone  
( $<10$  cm deep; fig. B3) could be a critical factor (Tian et al., 2016) explaining the lack of a positive warming effect on subsoil  
decomposition, as a plausible positive priming effect often elicited by fresh C inputs would have been restricted to the topsoil.  
95 Second, a large fraction of SOC in the topsoil is particulate organic matter protected in aggregates, whereas most subsoil SOC  
is associated with minerals (Fontaine et al., 2007; Rumpel and Kögel-Knabner, 2011). As such, by accelerating mineralisation  
of plant litter that is deposited only in topsoil (Walker et al., 2018), and thereby reducing aggregate stability and breaking  
up macroaggregates (fig. 3) (Poeplau et al., 2020), warming may have had much greater effect on SOC losses in topsoil than  
in subsoil, where mineral protection dominates. Third, although unlikely given the absence of increased dissolved organic C  
100 (DOC) with warming in subsoil (fig. B6), it can also not be excluded that SOC stocks in subsoils only appear stable, because  
increased losses are compensated for by increased inputs from above (Osher et al., 2003).

Only very few studies have assessed subsoil SOC stocks responses to deep soil warming, and observed responses differ  
strongly in magnitude and even direction. Higher subsoil than topsoil SOC losses were reported in two forests (Lin et al., 2018;  
Soong et al., 2020a), while unresponsive (this study) subsoil SOC stocks or even increases in subsoil SOC stocks were observed  
105 in grasslands (Jia et al., 2019). Further research is needed to unravel the drivers of these contrasting subsoil SOC responses to  
warming among experiments, which may be related to differences in soil properties, aggregate dynamics or rooting depths.

### 4 Implications for carbon-climate feedbacks

Earth System Model (ESM) inter-comparison studies (Eyring et al., 2016) have revealed large variability in both contemporary  
global SOC stock estimates and future SOC stock projections, underlining the need for empirical observations to better con-  
110 strain the response of SOC to temperature change (Nishina et al., 2014). Long-term warming experiments like this study are



thus needed to reduce the uncertainty on model projections (Abramoff et al., 2019). Although geothermally active areas offer long-lasting, continuous and large soil temperature gradients and overcome the technical challenges and high costs associated with warming manipulation experiments (Sigurdsson et al., 2016; O’Gorman et al., 2014), their use as a proxy for climate change has some drawbacks of its own, such as limited aboveground warming and a stepwise increase in soil temperature at the initiation of the geothermal gradient (De Boeck et al., 2015). Also the Andosol, covering only  $\pm 0.8\%$  of the earth’s surface (Baillie, 2001), makes that one should be cautious extrapolating the results to the entire sub-arctic region. Nonetheless, this site offers a unique opportunity to study the direct versus long-term response of SOC stocks to temperature change and the results from this study and other deep soil warming experiments clearly indicate that introducing vertically resolved plant- and microbial dynamics in ESMs is a necessity for more accurate projections of the carbon-climate feedback.

115

120 In conclusion, warming caused a large but rapidly equilibrating SOC loss in the topsoil that increased linearly with warming intensity, while no SOC loss was observed in the subsoil in our subarctic grasslands exposed to decades of soil warming. Future work should focus on understanding whether these observed temporal dynamics are consistent throughout the northern non-permafrost region. Improved understanding of the variation in subsoil SOC responses to warming is also critical for constraining Earth System Models and obtaining reliable climate projections.





## 125 Appendix A: Material and methods

This study was conducted at the ForHot research site, located in the Hengill geothermal area, 40 km east of Reykjavík, Iceland (64°00'01" N, 21°11'09" W; 100 – 225 m a.s.l. (Sigurdsson et al., 2016). The mean annual temperature between 2006 and 2016 was  $5.2 \pm 0.1$  ( $\pm$  SE) °C, and mean annual daily minimum and maximum temperatures were  $2.2 \pm 0.2$  (SE) and  $8.6 \pm 0.2$  ( $\pm$  SE) °C. The mean annual precipitation during the same period was  $1413 \pm 57$  ( $\pm$  SE) mm (Icelandic Meteorological Office; 130 Eyrbakki weather station which closed in 2017). The main vegetation type is unmanaged grassland, dominated by *Agrostis capillaris*, *Ranunculus acris* and *Equisetum pratense* and the underlying soil is classified as Brown Andosol (Arnalds, 2015).

In this study, we define the 0-10 cm layer as topsoil, and the 10-30 cm layers as subsoil. This subsoil layer strongly differed from the topsoil in many ways. First in these soils, the A horizon that is enriched with SOC is maximum 10 cm deep (Arnalds, 2015). Second,  $95.7 \pm 0.4$  (SE) % of the fine root biomass sampled in the top 30 cm layer, was found in upper 10 cm (fig. B5). 135 Third, bulk density in subsoil is significantly higher than in topsoil ( $P < 0.001$ ) (fig. B7).

The site comprises two areas that have been subjected to geothermal soil warming for different periods of time (Sigurdsson et al., 2016). One area (hereafter “medium-term warmed grassland”) has been warmed since May 2008, when a large earthquake shifted geothermal systems to previously unwarmed soils. The second area (2.5 km North-east from the first area; hereafter “long-term warmed grassland”) was already mentioned to be warmed in the early 18th century (Magnússon and Vídalín, 1708) and has thus likely been warmed for centuries. For sure, the warming was registered in a census during the 1960s, and no change 140 in the location of the hotspots has been recorded during the past 50 years (Kristján Sæmundsson, personal communication). The soil warming increment at both sites is relatively constant throughout the year and extreme deviations are rare (Sigurdsson et al., 2016). Soil warming is caused by horizontal heat conduction through the soil, causing fairly homogeneous warming with depth and inducing a fairly natural temperature depth profile (fig. B4). This homogeneous soil warming is in line with CMIP5 145 predictions of rapid transfer of the temperature signal from air to shallow and deeper soils (Soong et al., 2020b). The geothermal water is confined within the bedrock and no signs of soil contamination by geothermal byproducts have been found (Shi et al., 2020). Soil pH (mean:  $5.5 \pm 0.1$  (SE)) and soil moisture did not show major changes along the soil warming gradients, with soil moisture very rarely approaching the permanent wilting point and no relation between soil temperature and the frequency of drought events (Leblans et al., 2017). Further, the plant species composition was very similar between the medium-term and 150 the long term warmed grassland and no drastic changes in dominant plant species occurred up to  $+6.4$  °C warming (Leblans et al., 2017) (which is the RCP8.5 projected annual warming level for high northern latitudes for the year 2100) (IPCC, 2013). More detailed information on the site characteristics can be found in Sigurdsson et al. (2016).

We established five replicate transects in each area (the medium-term and the long-term warmed grassland) in 2012, around two and four geothermal heat sources respectively. In the medium-term warmed grassland, all transects were located on south- 155 west facing slopes, three with the geothermal heat source at the bottom of the slope, and two with the geothermal heat source at the top, to eliminate effects of topography and downward transport of groundwater and nutrients from introducing a bias in the SOC stocks. In the long-term warmed grassland, all transects were located on level ground. Within the long-term and medium-term warmed grassland, all measurement plots had similar microtopography, soil depth and grazing history.



Each transect consists of six 2 x 2 m permanent measurement plots distributed along the soil temperature gradient, including  
160 unwarmed soil (MAT:  $5.7 \pm 0.1$  °C), yielding 60 plots in total. Each 2 x 2 m permanent measurement plot was accompanied by  
two adjacent 0.5 x 0.5 m subplots for destructive measurements. Plot-specific soil warming was recorded hourly at 10 cm soil  
depth using HOBO TidbiT v2 Water Temperature Data Loggers (Onset Computer Corporation, USA). Because the permanent  
plots occurred at different warming intensities in the different transects, we adopted a regression approach (see statistics below).  
More detailed information on the experimental design is provided in Sigurdsson et al. (2016).

165 In July 2013 and 2018, two 0-10 cm soil cores (corer  $\phi = 5.12$  cm) were taken within each subplot. In the medium-term  
warmed grassland, soils were too shallow to sample deeper, but additional 10-30 cm cores were taken in the long-term warmed  
grassland. Cores were analysed for: (1) soil C concentrations; (2) pH (topsoil only); (3) soil bulk density (BD); and (4) SOC  
stocks.

From the first core we obtained fine roots (<2 mm) and soil particles (>2 mm) (necessary to calculate BD) by washing the  
170 cores over two sieves with mesh sizes 2 mm and 0.5 mm. Roots and >2 mm particles were dried and weighed to gain fine root  
biomass ( $\text{g m}^{-2}$ ) and the volume of >2 mm particles ( $\text{g cm}^{-3}$ ) was measured by the water displacement method. The second  
soil core was first dried and weighed (as for aboveground vegetation), and soil was then sieved to obtain soil particles <2 mm  
and split into three aliquots. One aliquot of 2 g was milled (Retsch MM301 Mixer Mill, Haan, Germany) and analysed for  
C concentration (%) by dry combustion (Macro Elemental Analyser, model vario MAX CN, Hanau, Germany). Finally, BD  
175 ( $\text{g cm}^{-3}$ ) and SOC stocks ( $\text{ton ha}^{-1}$ ) were calculated according to the approach described in Bárcena et al. (2014).

To measure dissolved organic matter (DOC), teflon suction cup lysimeters (Prenart Super Quartz, Prenart Equipment Aps,  
Frederiksberg, Denmark) were placed at about 30-40 cm depth in the medium-term and long-term warmed grassland, in Octo-  
ber 2014. Samples were taken during summer 2015, 2016 and 2017. The DOC was analysed with a combined Total Organic  
Carbon (Shimadzu, Kyoto, Japan). Further installation details of the lysimeters are described in Edlinger (2016), as well as  
180 the sampling procedure. The C-input data for arbuscular mycorrhizae originates from Zhang et al. (2020), where the sampling  
procedure is described. Aboveground biomass was sampled by placing a 20x40 cm frame on the plot, after which all vegetation  
was clipped. The samples were taken to the lab and sorted by hand in a grass and moss fraction. Both fractions were dried for  
48 h at 70 °C, weighed and milled. The samples were analysed for C concentration (%) by dry combustion (Macro Elemental  
Analyser, model vario MAX CN, Hanau, Germany). Aggregate fractionation was done in 2018 only. Per plot, a 0-10 cm soil  
185 core was taken (corer  $\phi = 5.12$  cm) and dried at room temperature for a some weeks. Stones were removed and aggregates  
larger than 8 mm were broken up by dry-sieving on a 8 mm soil sieve. The dry-sieved soil was then slaked for 5 min with DI  
water, after which it was wet-sieved on a 2 mm, 250  $\mu\text{m}$  and 63  $\mu\text{m}$ , to separate into four size fractions. Each fraction was dried  
at 70 °C for 72 h, after which all fractions were ground with a ball mill to homogenise and analysed for C concentration (%)  
by dry combustion (Macro Elemental Analyser, model vario MAX CN, Hanau, Germany). Relative mass of the fractions was  
190 calculated by dividing the fraction mass by the sum of all fraction masses of initial sample. The absolute soil C-amount of each  
fraction was calculated by multiplying the fraction soil mass per 100 g of dry soil with the soil C % (fig. 3).


A soil mass correction of the SOC stocks as described in Ellert and Bettany (1995) was necessary to compare stock changes  
across the soil warming gradient as soil compaction increased with warming in the upper soil layers (increasing BD; fig. B7),




195 implying that soil depths in unwarmed soil corresponded to shallower soil depths at warmer soils. The calculation method is explained in detail in fig. B8.

200 The soil warming dependence of bulk soil SOC stocks (corrected and uncorrected for soil compaction), BD, DOC and soil C % were tested with a linear mixed effects model (Pinheiro et al., 2021). Soil warming and warming-duration (medium-term vs. long-term) were included as main effects, while sampling year was used as random effect to account for sampling differences and interannual variabilities between the two sampling campaigns. In all cases, criteria for normality and homoscedasticity were met. For all tests, the dataset was reduced to cover only the warming levels captured by the projections for high northern latitudes for the year 2100 (0 – 6.4 °C warming) (IPCC, 2013). All tests were performed using R software (R Development Core Team, 2011).


## Appendix B: Supplementary




**Figure B1.** Reduction of soil organic carbon (SOC) stock with soil warming, a) in the topsoil (0-10 cm); b) in the subsoil (10-30 cm). All soil samples were taken in July 2013 or July 2018. The regression for medium-term warmed (MTW) and long-term warmed (LTW) grassland (in topsoil) for both 2013 and 2018 were combined, since no soil temperature x warming-duration interaction effect, nor a main effect for warming-duration was found. Soil warming is expressed relative to ambient soil temperature (both at 10 cm depth). In topsoil, a linear relation was observed, while no significant effect was present in subsoil. The 95 % confidence bounds are shown around the topsoil regression slope. (n = 78 & 40 for topsoil and subsoil respectively)




**Figure B2.** Percentage of carbon in topsoil and subsoil, for the long-term warmed (LTW) and medium-term warmed (MTW) grassland. The 95 % confidence bounds are shown around the topsoil linear regression slopes. (topsoil  $n = 42$  &  $40$  for MTW and LTW grassland respectively; subsoil  $n = 40$  for LTW grassland)



**Figure B3.** Proxies for annual soil C-inputs from aboveground biomass (AGB), fine root biomass and arbuscular mycorrhizae (AMF) in the medium-term warmed (MTW) and the long-term warmed (LTW) grasslands. Vascular plant aboveground biomass was used as a proxy for aboveground C-inputs; vascular plant fine root biomass for belowground C-inputs and C sequestered by AMF (Cnew; data from Zhang et al. (2020)) for C-inputs by arbuscular mycorrhizae. For the observed soil warming range (0-6.4 °C warming), no change in C-inputs could be found. P-values were obtained by a linear regression analysis. (n = 20 & 17 for MTW and LTW inputs respectively)




**Figure B4.** Soil warming along the vertical soil profile, based on the reference temperature measured at 10 cm which is used throughout the paper. The median temperature is about 1 °C higher at 5 cm depth, while being slightly lower at 20 cm and 30 cm depth. Due to the shallow soil in the medium-term warmed grassland, the warming profile is given for the long-term warmed grassland only.




**Figure B5.** Fine root density in long-term warmed (LTW) grassland in top-and subsoil sampled in July 2018. For the observed soil warming range (0-6.4 °C warming), no change in C-inputs could be found. P-values were obtained by a linear regression analysis. In the top 30 cm soil layer,  $95.7 \pm 0.4$  (SE) % of the fine root biomass was found in upper 10 cm, which is why we define the 0-10 cm layer as topsoil, and the 10-30 cm layer as subsoil. (n = 17 & 15 for topsoil and subsoil respectively)






**Figure B6.** Dissolved organic carbon (DOC) in medium-term warmed (MTW) and long-term warmed (LTW) grassland during summer 2015, 2016 and 2017. No DOC change was observed with soil warming in the SWT ( $P = 0.95$ ), nor in the LTW ( $P = 0.8$ ), meaning that carbon-inputs into deeper soil layers remained constant over the whole soil temperature gradient. Statistical analysis was done with a linear mixed effects model with soil warming as explanatory variable and sampling year as a random factor. Criteria for normality and homoscedasticity were met. Dissolved organic carbon was sampled at a approximate depth of 30 cm with Prenart Super Quartz (Prenart, Fredriksberg, Denmark) soil water samplers, installed around 1 October 2014. The samples was always inserted from ‘downslope,’ where the soil was deep enough. Analyses of the samples was done at the IGN Biochemistry Lab, University of Copenhagen. ( $n = 25$  &  $29$  for MTW and LTW grassland respectively)



**Figure B7.** Changes in bulk density with soil warming in the long-term warmed (LTW; circles) and the medium-term warmed (MTW; triangles) grassland. Soil warming is expressed relative to ambient soil temperature (at 10 cm depth). Bulk density is separated for the topsoil (0-10 cm) and the subsoil (10-30 cm). The regression for medium-term warmed (MTW) and long-term warmed (LTW) grassland (in topsoil) for both 2013 and 2018 were combined, since no soil temperature  $\times$  warming-duration interaction effect, nor a main effect for warming-duration or sampling year was found. The 95 % confidence bounds are shown around the regression slopes. ( $n = 78$  &  $40$  for topsoil and subsoil respectively)



**Figure B8. Soil mass correction.** Due to significant soil compaction (increasing bulk density (BD)) with increasing soil temperature in the upper soil layers (fig. B7), a certain soil depth in unwarmed soil corresponds to ever shallower soil depths at warmer soil temperatures. Therefore, the SOC stocks were corrected for soil compaction, i.e., the corrected SOC stocks were calculated on the same mass of soil.

For topsoil, we calculated a corrected thickness of a warmed soil layer, corresponding to the same core mass as the the  
 205 unwarmed soils. Using the ratio of corrected and uncorrected layer thickness, we calculated a corrected SOC stock for the  
 warmed topsoil. For warmed subsoils, we calculated the corrected thickness in the same way as for topsoil, but subtracted a  
 surplus thickness of the above topsoil. The corrected subsoil SOC stock was then calculated as the sum of the surplus topsoil  
 SOC stock and the SOC stock in the corrected subsoil layer. The detailed calculation method is shown below.

### Corrections 0-10cm soil layer

210 First, the corrected soil layer thickness of the 0-10 cm layer ( $C.Th_{0-10}$ ) was calculated for the warmed soils:

$$C.Th_{0-10} = \frac{U.Th_{0-10} \times U.BD_{0-10}}{C.BD_{0-10}} \quad (B1)$$

Where  $U.Th_{0-10}$  is the uncorrected soil layer thickness of the 0-10 cm layer (10 cm),  $U.BD_{0-10}$  is the uncorrected BD of the 0-10 cm layer (which corresponds to the BD at ambient soil temperature) and  $C.BD_{0-10}$  is the measured BD for the 0-10 cm layer. Then, the corrected SOC stocks of the 0-10 depth layer ( $C.SOC_{0-10}$ ) were calculated:

$$215 \quad C.SOC_{0-10} = \frac{U.SOC_{0-10} \times C.Th_{0-10}}{U.Th_{0-10}} \quad (B2)$$



Where  $U.SOC_{0-10}$  is the uncorrected SOC stock in the 0-10 cm depth layer, and  $C.Th_{0-10}/U.Th_{0-10}$  corresponds to the proportional thickness of the corrected layer compared to the uncorrected layer.

### Corrections 10-30 cm soil layer (only applicable to the long-term warmed grassland)

First, the thickness of the surplus soil layer from the 0-10 cm layer ( $S.Th_{0-10}$ ) was calculated:

$$220 \quad S.Th_{0-10} = \frac{(U.Th_{0-10} - C.Th_{0-10}) \times U.BD_{0-10}}{C.BD_{0-10}} \quad (B3)$$

The second term is a correction factor for the soil compaction of the surplus soil layer. Then, the corrected thickness of the 10-30 cm soil layer, not yet taking the surplus soil sampled from the 0-10 cm layer ( $S.Th_{0-10}$ ) into account, ( $c.Th_{10-30}$ ) was calculated:

$$c.Th_{10-30} = \frac{U.Th_{10-30} \times U.BD_{10-30}}{C.BD_{10-30}} \quad (B4)$$

225 Where  $U.Th_{10-30}$  is the uncorrected soil layer thickness of the 10-30 cm layer (20 cm),  $U.BD_{10-30}$  is the uncorrected BD of the 10-30 cm layer (which corresponds to the BD at ambient soil temperature) and  $C.BD_{10-30}$  is the measured BD for the 10-30 cm layer.

Subsequently, we took into account the thickness of the surplus soil sampled from the 0-10 cm layer to calculate the final corrected soil thickness of the 10-30 cm soil layer ( $C.Th_{10-30}$ ). Hence,  $C.Th_{10-30}$  is the part of the 10-30 cm layer that  
230 remains after (i) correcting for soil compaction and (ii) subtracting the thickness of the surplus soil sampled at the 0-10 cm layer:

$$C.Th_{10-30} = c.Th_{10-30} - S.Th_{0-10} \quad (B5)$$

Subsequently, the corrected SOC stock for the 10-30 cm layer ( $C.SOC_{10-30}$ ) was calculated:

$$C.SOC_{10-30} = (U.SOC_{0-10} - C.SOC_{0-10}) + \frac{U.SOC_{10-30} \times C.Th_{10-30}}{U.Th_{10-30}} \quad (B6)$$

235 Where  $U.SOC_{10-30}$  is the uncorrected SOC stock in the 10-30 cm depth layer, and  $C.Th_{10-30}/U.Th_{10-30}$  corresponds to the proportional thickness of the corrected layer compared to the uncorrected layer.

*Data availability.* All data used in this manuscript has been made available on Zenodo. DOI: 10.5281/zenodo.4745479

*Author contributions.* BDS, NIWL, SV, JLS, LF, JTW, HW, PMVB, NV and IAJ designed the study. PG, HW, CF, NIWL, BDS, IAJ, SV, KVDV, EV, ZFL, SM-J, NV and MM provided the data. All authors contributed substantially to the analysis and the writing of the manuscript.



240 *Competing interests.* The authors declare that they have no conflict of interest.

*Acknowledgements.* This research was supported by a joint Fonds Wetenschappelijk Onderzoek Flanders (FWO) and Fonds zur Förderung der wissenschaftlichen Forschung (FWF) grant with nos. FWO-G0F2217N & FWF-I-3237, awarded to I.A.J and M.B., the European Research Council Synergy grant 610028 (IMBALANCE-P), and the Research Council of the University of Antwerp (FORHOT TOP-BOF project). This work contributes to the FSC-Sink, CAR-ES and the ClimMani COST Action (ES1308). Reykir - the Icelandic state gardening school, Keldnaholt – the Agricultural University of Iceland and Mogilsá – the Icelandic Forest Research, provided logistical support for the present study. Further, we thank the Lorentz Center in Leiden. We thank Iris Janssens, Jochen Janssens, Inge Van De Putte, Maxime Sepelie, Jana Vynckier, Alexander Meire, Lieven Michielsens, Freja Dreesen, Sebastien Leys, Elín Gudmundsdóttir, Annemie Vinck, Paul Leblans, Kwinten Leblans, Sigvatur, Már Gudmundsson, Elías Óskarsson and Simon Arnar Pálsson for their help in the field. We thank Brita Berglund, Baldur Vigfusson, Nadine Calluy, Tom Van Der Spiet, Anne Cools, Marijke Van den Bruel, Els Oosterbos, Saad El-Rawi, Miguel Portillo Estrada and Wannes Kiebooms for their assistance with the lab analyses.



## References

- Abramoff, R. Z., Torn, M. S., Georgiou, K., Tang, J., and Riley, W. J.: Soil Organic Matter Temperature Sensitivity Cannot Be Directly Inferred from Spatial Gradients, *Global Biogeochemical Cycles*, 33, 761–776, 2019.
- Allison, S. D., Wallenstein, M. D., and Bradford, M. A.: Soil-Carbon Response to Warming Dependent on Microbial Physiology, *Nature Geoscience*, 3, 336–340, <https://doi.org/10.1038/ngeo846>, 2010.
- 255 Arnalds, O.: The Soils of Iceland, *World Soils Book Series*, Springer Netherlands, Dordrecht, <https://doi.org/10.1007/978-94-017-9621-7>, 2015.
- Baillie, I.: Soil survey staff 1999, soil taxonomy: a basic system of soil classification for making and interpreting soil surveys, agricultural handbook 436, Natural Resources Conservation Service, USDA, Washington DC, USA, pp. 869, 2001.
- 260 Bárcena, T. G., Gundersen, P., and Vesterdal, L.: Afforestation Effects on SOC in Former Cropland: Oak and Spruce Chronosequences Resampled after 13 Years, *Global Change Biology*, 20, 2938–2952, 2014.
- Batjes, N.: Harmonized Soil Property Values for Broad-Scale Modelling (WISE30sec) with Estimates of Global Soil Carbon Stocks, *Geoderma*, 269, 61–68, <https://doi.org/10.1016/j.geoderma.2016.01.034>, 2016.
- Bradford, M. A., McCulley, R. L., Crowther, T. W., Oldfield, E. E., Wood, S. A., and Fierer, N.: Cross-Biome Patterns in Soil Microbial  
265 Respiration Predictable from Evolutionary Theory on Thermal Adaptation, *Nature ecology & evolution*, 3, 223, 2019.
- Crowther, T. W., Todd-Brown, K. E. O., Rowe, C. W., Wieder, W. R., Carey, J. C., Machmuller, M. B., Snoek, B. L., Fang, S., Zhou, G., Allison, S. D., Blair, J. M., Bridgham, S. D., Burton, A. J., Carrillo, Y., Reich, P. B., Clark, J. S., Classen, A. T., Dijkstra, F. A., Elberling, B., Emmett, B. A., Estiarte, M., Frey, S. D., Guo, J., Harte, J., Jiang, L., Johnson, B. R., Kröel-Dulay, G., Larsen, K. S., Laudon, H., Lavallee, J. M., Luo, Y., Lupascu, M., Ma, L. N., Marhan, S., Michelsen, A., Mohan, J., Niu, S., Pendall, E., Peñuelas, J., Pfeifer-Meister, L., Poll,  
270 C., Reinsch, S., Reynolds, L. L., Schmidt, I. K., Sistla, S., Sokol, N. W., Templer, P. H., Treseder, K. K., Welker, J. M., and Bradford, M. A.: Quantifying Global Soil Carbon Losses in Response to Warming, *Nature*, 540, 104–108, <https://doi.org/10.1038/nature20150>, 2016.
- Davidson, E. A. and Janssens, I. A.: Temperature Sensitivity of Soil Carbon Decomposition and Feedbacks to Climate Change, *Nature*, 440, 165–173, <https://doi.org/10.1038/nature04514>, 2006.
- De Boeck, H. J., Vicca, S., Roy, J., Nijs, I., Milcu, A., Kreyling, J., Jentsch, A., Chabbi, A., Campioli, M., Callaghan,  
275 T., Beierkuhnlein, C., and Beier, C.: Global Change Experiments: Challenges and Opportunities, *BioScience*, 65, 922–931, <https://doi.org/10.1093/biosci/biv099>, 2015.
- Edlinger, A.: Changes of belowground processes of the carbon and nitrogen cycle in response to soil warming. Will global warming increase the turnover of soil C and N above plant demands and lead to increased leaching losses, Master's thesis, M. Sc. thesis. Copenhagen University, Denmark & University of Natural Resources and Applied Sciences (BOKU), <http://forhot.is/wp-content/uploads/2018/01/Edlinger-Anna-Theresa.-2016.-Changes-of-belowground-processes-of-the-carbon-and-nitrogen.xps>, 2016.  
280
- Ellert, B. H. and Bettany, J. R.: Calculation of Organic Matter and Nutrients Stored in Soils under Contrasting Management Regimes, *Canadian Journal of Soil Science*, 75, 529–538, 1995.
- Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) Experimental Design and Organization, *Geoscientific Model Development*, 9, 1937–1958,  
285 <https://doi.org/10.5194/gmd-9-1937-2016>, 2016.
- Fontaine, S., Barot, S., Barré, P., Bdioui, N., Mary, B., and Rumpel, C.: Stability of organic carbon in deep soil layers controlled by fresh carbon supply, *Nature*, 450, 277–280, 2007.



- Halldórsson, B. and Sigbjörnsson, R.: The Mw6.3 Ölfus Earthquake at 15:45 UTC on 29 May 2008 in South Iceland: ICEARRAY Strong-Motion Recordings, Soil Dynamics and Earthquake Engineering, 29, 1073–1083, <https://doi.org/10.1016/j.soildyn.2008.12.006>, 2009.
- 290 Hanson, P. J., Griffiths, N. A., Iversen, C. M., Norby, R. J., Sebestyen, S. D., Phillips, J. R., Chanton, J. P., Kolka, R. K., Malhotra, A., Oleheiser, K. C., et al.: Rapid net carbon loss from a whole-ecosystem warmed Peatland, AGU Advances, 1, e2020AV000163, 2020.
- Ilyina, T. and Friedlingstein, P.: WCRP Grand Challenge-Carbon Feedbacks in the Climate System, Tech. rep., Tech. rep., WCRP, [https://www.wcrp-climate.org/JSC37/Documents ...](https://www.wcrp-climate.org/JSC37/Documents...), 2016.
- IPCC: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2013.
- 295 Jenkinson, D. S., Adams, D. E., and Wild, A.: Model Estimates of CO<sub>2</sub> Emissions from Soil in Response to Global Warming, Nature, 351, 304–306, 1991.
- Jia, J., Cao, Z., Liu, C., Zhang, Z., Lin, L., Wang, Y., Haghpor, N., Wacker, L., Bao, H., Dittmar, T., Simpson, M. J., Yang, H., Crowther, T. W., Eglinton, T. I., He, J.-S., and Feng, X.: Climate Warming Alters Subsoil but Not Topsoil Carbon Dynamics in Alpine Grassland, Global Change Biology, 25, 4383–4393, <https://doi.org/10.1111/gcb.14823>, 2019.
- 300 Leblans, N. I. W., Sigurdsson, B. D., Vicca, S., Fu, Y., Penuelas, J., and Janssens, I. A.: Phenological Responses of Icelandic Subarctic Grasslands to Short-Term and Long-Term Natural Soil Warming, Global Change Biology, 23, 4932–4945, <https://doi.org/10.1111/gcb.13749>, 2017.
- Lin, W., Li, Y., Yang, Z., Giardina, C. P., Xie, J., Chen, S., Lin, C., Kuzyakov, Y., and Yang, Y.: Warming Exerts Greater Impacts on Subsoil than Topsoil CO<sub>2</sub> Efflux in a Subtropical Forest, Agricultural and Forest Meteorology, 263, 137–146, <https://doi.org/10.1016/j.agrformet.2018.08.014>, 2018.
- Magnússon, Á. and Vídalín, P.: Jarðabók. Annað bindi [The farm register of Iceland], 1708.
- Marañón-Jiménez, S., Soong, J. L., Leblans, N. I. W., Sigurdsson, B. D., Peñuelas, J., Richter, A., Asensio, D., Fransen, E., and Janssens, I. A.: Geothermally Warmed Soils Reveal Persistent Increases in the Respiratory Costs of Soil Microbes Contributing to Substantial C Losses, Biogeochemistry, 138, 245–260, <https://doi.org/10.1007/s10533-018-0443-0>, 2018.
- 310 Melillo, J. M., Frey, S. D., DeAngelis, K. M., Werner, W. J., Bernard, M. J., Bowles, F. P., Pold, G., Knorr, M. A., and Grandy, A. S.: Long-Term Pattern and Magnitude of Soil Carbon Feedback to the Climate System in a Warming World, Science, 358, 101–105, <https://doi.org/10.1126/science.aan2874>, 2017.
- Nishina, K., Ito, A., Beerling, D. J., Cadule, P., Ciais, P., Clark, D. B., Falloon, P., Friend, A. D., Kahana, R., Kato, E., Keribin, R., Lucht, W., Lomas, M., Rademacher, T. T., Pavlick, R., Schaphoff, S., Vuichard, N., Warszawski, L., and Yokohata, T.: Quantifying Uncertainties in Soil Carbon Responses to Changes in Global Mean Temperature and Precipitation, Earth System Dynamics, 5, 197–209, <https://doi.org/10.5194/esd-5-197-2014> <<https://doi.org/10.5194/esd-5-197-2014>>, 2014.
- 315 W., Lomas, M., Rademacher, T. T., Pavlick, R., Schaphoff, S., Vuichard, N., Warszawski, L., and Yokohata, T.: 2014 Quantifying uncertainties in soil carbon responses to changes in global mean temperature and precipitation. Earth System Dynamics, 5 (1), 197-209. <https://doi.org/10.5194/esd-5-197-2014> <<https://doi.org/10.5194/esd-5-197-2014>>, 2014.
- O’Gorman, E. J., Benstead, J. P., Cross, W. F., Friberg, N., Hood, J. M., Johnson, P. W., Sigurdsson, B. D., and Woodward, G.: Climate Change and Geothermal Ecosystems: Natural Laboratories, Sentinel Systems, and Future Refugia, Global Change Biology, 20, 3291–3299, <https://doi.org/10.1111/gcb.12602>, 2014.
- 320 Osher, L. J., Matson, P. A., and Amundson, R.: Effect of Land Use Change on Soil Carbon in Hawaii, Biogeochemistry, 65, 20, 2003.



- 325 Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., and R Core Team: nlme: Linear and Nonlinear Mixed Effects Models, <https://CRAN.R-project.org/package=nlme>, r package version 3.1-152, 2021.
- Poeplau, C., Sigurðsson, P., and Sigurdsson, B. D.: Depletion of Soil Carbon and Aggregation after Strong Warming of a Subarctic Andosol under Forest and Grassland Cover, *SOIL*, 6, 115–129, <https://doi.org/10.5194/soil-6-115-2020>, 2020.
- R Development Core Team, R.: R: A Language and Environment for Statistical Computing, R foundation for statistical computing Vienna, Austria, 2011.
- 330 Radujković, D., Verbruggen, E., Sigurdsson, B. D., Leblans, N. I., Janssens, I. A., Vicca, S., and Weedon, J. T.: Prolonged Exposure Does Not Increase Soil Microbial Community Compositional Response to Warming along Geothermal Gradients, *FEMS microbiology ecology*, 94, fix174, 2018.
- Rumpel, C. and Kögel-Knabner, I.: Deep soil organic matter—a key but poorly understood component of terrestrial C cycle, *Plant and soil*, 338, 143–158, 2011.
- 335 Scharlemann, J. P., Tanner, E. V., Hiederer, R., and Kapos, V.: Global Soil Carbon: Understanding and Managing the Largest Terrestrial Carbon Pool, *Carbon Management*, 5, 81–91, <https://doi.org/10.4155/cmt.13.77>, 2014.
- Shi, Z., Crowell, S., Luo, Y., and Moore, B.: Model Structures Amplify Uncertainty in Predicted Soil Carbon Responses to Climate Change, *Nature communications*, 9, 1–11, 2018.
- 340 Shi, Z., Allison, S. D., He, Y., Levine, P. A., Hoyt, A. M., Beem-Miller, J., Zhu, Q., Wieder, W. R., Trumbore, S., and Randerson, J. T.: The Age Distribution of Global Soil Carbon Inferred from Radiocarbon Measurements, *Nature Geoscience*, 4, 1–5, 2020.
- Sigurdsson, B. D., Leblans, N., Dauwe, S., Guðmundsdóttir, E., Gundersen, P., Gunnarsdóttir, G. E., Holmstrup, M., Ilieva-Makulec, K., Kätterer, T., Marteinsdóttir, B., et al.: Geothermal Ecosystems as Natural Climate Change Experiments: The ForHot Research Site in Iceland as a Case Study, *Icelandic Agricultural Sciences*, 29, 53–71, <https://doi.org/10.16886/IAS.2016.05>, 2016.
- 345 Soong, J. L., Castanha, C., Hicks Pries, C. E., Ofiti, Porras, R. C., and Riley: Five Years of Whole-Soil Warming Led to Loss of Subsoil Carbon Stocks and Increased CO<sub>2</sub> Efflux, *Science Advances*, 0, 2020a.
- Soong, J. L., Phillips, C. L., Ledna, C., Koven, C. D., and Torn, M. S.: CMIP5 models predict rapid and deep soil warming over the 21st century, *Journal of Geophysical Research: Biogeosciences*, 125, e2019JG005 266, 2020b.
- Tian, Q., Yang, X., Wang, X., Liao, C., Li, Q., Wang, M., Wu, Y., and Liu, F.: Microbial Community Mediated Response of Organic Carbon Mineralization to Labile Carbon and Nitrogen Addition in Topsoil and Subsoil, *Biogeochemistry*, 128, 125–139, <https://doi.org/10.1007/s10533-016-0198-4>, 2016.
- 350 Todd-Brown, K. E., Zheng, B., and Crowther, T. W.: Field-Warmed Soil Carbon Changes Imply High 21st-Century Modeling Uncertainty, *Biogeosciences*, 15, 3659–3671, 2018.
- van Gestel, N., Shi, Z., van Groenigen, K. J., Osenberg, C. W., Andresen, L. C., Dukes, J. S., Hovenden, M. J., Luo, Y., Michelsen, A., Pendall, E., Reich, P. B., Schuur, E. A. G., and Hungate, B. A.: Predicting Soil Carbon Loss with Warming, *Nature*, 554, E4–E5, <https://doi.org/10.1038/nature25745>, 2018.
- Walker, T. W. N., Kaiser, C., Strasser, F., Herbold, C. W., Leblans, N. I. W., Woebken, D., Janssens, I. A., Sigurdsson, B. D., and Richter, A.: Microbial Temperature Sensitivity and Biomass Change Explain Soil Carbon Loss with Warming, *Nature Climate Change*, 8, 885–889, <https://doi.org/10.1038/s41558-018-0259-x>, 2018.
- 360 Walker, T. W. N., Janssens, I. A., Weedon, J. T., Sigurdsson, B. D., Richter, A., Peñuelas, J., Leblans, N. I. W., Bahn, M., Bartrons, M., De Jonge, C., Fuchslueger, L., Gargallo-Garriga, A., Gunnarsdóttir, G. E., Marañón-Jiménez, S., Oddsdóttir, E. S., Ostonen, I., Poeplau, C., Prommer, J., Radujković, D., Sardans, J., Sigurðsson, P., Soong, J. L., Vicca, S., Wallander, H., Ilieva-Makulec, K., and Verbruggen,





- E.: A Systemic Overreaction to Years versus Decades of Warming in a Subarctic Grassland Ecosystem, *Nature Ecology & Evolution*, 4, 101–108, <https://doi.org/10.1038/s41559-019-1055-3>, 2020.
- 365 Yost, J. L. and Hartemink, A. E.: How deep is the soil studied—an analysis of four soil science journals, *Plant and soil*, 452, 5–18, 2020.
- Zhang, J., Ekblad, A., Sigurdsson, B. D., and Wallander, H.: The Influence of Soil Warming on Organic Carbon Sequestration of Arbuscular Mycorrhizal Fungi in a Sub-Arctic Grassland, *Soil Biology and Biochemistry*, 147, 107 826, <https://doi.org/10.1016/j.soilbio.2020.107826>, 2020.
- DOI = 10.5281/zenodo.4745479