

Controlling factors on the global distribution of a representative marine ~~heterotrophic~~non-cyanobacterial diazotroph phylotype (Gamma A)

Zhibo Shao¹, Ya-Wei Luo¹

5 ¹State Key Laboratory of Marine Environmental Science and College of Ocean and Earth Sciences, Xiamen University, 361102 Xiamen, Fujian, China

*Correspondence to: Ya-Wei Luo (ywluo@xmu.edu.cn)

Abstract. ~~Heterotrophic~~Non-cyanobacterial diazotrophs are presumably heterotrophic bacteria and emerge as a potentially important~~may be~~ contributors to the global marine N₂ fixation, while the factors controlling their distribution are unclear. Here, we explored what controls the distribution of the most sampled ~~heterotrophic~~non-cyanobacterial diazotroph phylotype, Gamma A, in the global ocean. First, we represented Gamma A abundance by its *nifH* qPCR copies reported in the literature and analyzed the-its relationship ~~to~~between *nifH* based Gamma A abundance and climatological biological and environmental conditions. The ~~carrying capacity of~~maximal observed Gamma A abundance increased with net primary production (NPP) and saturated when NPP reached ~400 mg C m⁻² d⁻¹. The reduction in Gamma A abundance from ~~its~~this NPP-supported ~~carrying capacity~~maximal abundance was mostly related to low temperature, which possibly slowed the decomposition of organic matter, and high concentration of dissolved iron, to which the explanation was elusive but could result from the competition with autotrophic diazotrophs. Using a generalized additive model, these climatological factors together explained ~~41~~39% of the variance in the Gamma A abundance. Second, in addition~~al~~ to the climatological background, we found that mesoscale cyclonic eddies can substantially elevate Gamma A abundance, implying that Gamma A can respond to ~~short-term~~mesoscale features and benefit from stimulated primary production by nutrient inputs. Overall, our results suggest that the distribution of Gamma A is ~~most likely determined~~influenced by the supply of organic matters, not by those factors controlling autotrophic diazotrophs, and therefore provide an insight ~~into~~a niche differentiation between the heterotrophic and autotrophic N₂ fixation. More samplings on Gamma A and other ~~heterotrophic~~non-cyanobacterial diazotroph phylotypes are needed to ~~better~~ reveal the controlling mechanisms of heterotrophic N₂ fixation in the ocean.

25 1 Introduction

Dinitrogen (N₂) fixation, mostly conducted by prokaryotic bacteria (termed “diazotrophs”), is an important bioavailable nitrogen (N) source to the ocean (Moore et al., 2018; Karl et al., 2002). Although autotrophic cyanobacteria have been recognized as important diazotrophs in the ocean (Zehr, 2011), ~~heterotrophic~~non-cyanobacteria diazotrophs (NCDs) that are presumably heterotrophic (probably including photoheterotrophic) bacteria (Bombar et al., 2016) have been widely detected

30 (e.g., Moisander et al., 2008; Langlois et al., 2008; Halm et al., 2012; Moisander et al., 2014; Shiozaki et al., 2014)(Moisander
et al., 2017; Riemann et al., 2010) and sometimes even found to dominate the diazotrophic gene pools in surface oceans
(Farnelid et al., 2011). For example, NCD *nifH* (a gene encoding subunit of nitrogenase enzyme) amplicons had higher relative
abundances than were far superior in number to autotrophic diazotrophs at some sampling sites in the South Pacific Ocean
(Halm et al., 2012; Moisander et al., 2014), Indian Ocean (Shiozaki et al., 2014; Wu et al., 2019) and South China Sea (Ding
35 et al., 2021). Metagenomic studies also revealed the dominant abundant presence of diverse N₂-fixing proteobacteria in ocean
genomic databases (Delmont et al., 2018; Delmont et al., 2021). Additionally, *nifH* of NCDs was also detected in subphotic
seawaters (Benavides et al., 2018a) and oxygen deplete-deficient zones (Jayakumar and Ward, 2020; Loescher et al., 2014)
where nitrogen loss was considered heavy-significant (Lam and Kuypers, 2011). Although the N₂ fixed by heterotrophic
NCDs diazotrophs is still has not well-been quantified, substantial N₂ fixation found in aphotic zones (Rahav et al., 2013; Bonnet
40 et al., 2013) and in experiments with photosynthetic inhibitors (Rahav et al., 2015; Geisler et al., 2020), as well as recovered
transcripts of the NCD *nifH* gene (Fernandez et al., 2011; Gradoville et al., 2017), provided a line of indirect evidence of
heterotrophic supports active heterotrophic N₂ fixation in the ocean.

The major known NCD classes include bacteria such as Alphaproteobacteria, Gammaproteobacteria, Epsilonproteobacteria,
Betaproteobacteria, and Firmicutes belonging to Cluster I of *nifH* clusters, and some obligate anaerobic bacteria and archaea
45 belonging to Cluster III of *nifH* clusters (Zehr et al., 2003; Riemann et al., 2010). Among them, Gamma A is the most sampled
and studied phylotype. Gamma A represents a part of uncultured Gammaproteobacterial sequences isolated from the open
ocean, and its cluster is distantly related to cultured NCD (Langlois et al., 2015). Gamma A *nifH* gene expression has been
widely found in the global ocean, suggesting its important role in marine N₂ fixation (Bird et al., 2005; Moisander et al., 2014;
Langlois et al., 2015; Shiozaki et al., 2017).

50 It is unclear what controls the growth and distribution of NCDs, as most of them, including Gamma A, are uncultivated
(Bombar et al., 2016). Apparently, NCDs are different from their autotrophic counterparts in depending organic matter as their
carbon and energy sources, which can be supported by experimental evidence that N₂ fixation is stimulated by adding dissolved
organic matter (DOM) (Rahav et al., 2016; Rahav et al., 2015; Bonnet et al., 2013; Bentzon-Tilia et al., 2015). However, DOM
addition sometimes did not stimulate *nifH* expression of Gamma A even when its DNA copies were ambient (Benavides et al.,
55 2018b), implying that DOM may not always stimulate the activity of Gamma A. In addition, the response of aphotic N₂ fixation
to different DOM compositions could also vary (Benavides et al., 2015). other studies showed contradictory results in which
DOM addition did not always stimulate the activity of NCDs (Benavides et al., 2018b; Benavides et al., 2015). Due to
sensitivity to O₂ and high energy requirements of N₂ fixation (Bombar et al., 2016), abundant NCDs were found to attach
to associate with particles which that supposedly provide the diazotrophs with a microenvironment with depleted oxygen and
60 rich organic matter (Riemann et al., 2010; Farnelid et al., 2010; Scavotto et al., 2015; Pedersen et al., 2018; Geisler et al.,
2019). NCDs were also detected in diatom mats (Martínez et al., 1983), implying another novel habitat for NCDs probably
releasing their fixed N to support diatom growth (Bonnet et al., 2016). An isolated strain of diazotrophic Alphabacteria from

~~that~~ Baltic Sea found ~~to to equip with~~contain photosynthetic genes (Bentzon-Tilia et al., 2015) may complicate this issue, doubting whether NCDs can be mixotrophic and also depend on light.

65 Although dissolved inorganic nitrogen (DIN) is generally considered to inhibit marine N₂ fixation (Karl et al., 2002; Zehr and Kudela, 2011), NCDs are active in DIN-replete environments such as estuaries (Geisler et al., 2020), coastal zones (Li et al., 2020), upwelling regions (Geisler et al., 2020; Moreira-Coello et al., 2017; Dekaezemacker et al., 2013) and other eutrophic seas (Bird and Wyman, 2013). Culture experiments showed that the DIN inhibition effect on NCDs can be strain specific (Bentzon-Tilia et al., 2015; Martínez-Pérez et al., 2018). Temperature could be another factor controlling NCDs which may
70 prefer warm oligotrophic surface oceans (Langlois et al., 2015; Shiozaki et al., 2017), the same region where the majority of autotrophic N₂ fixation ~~supposably~~ occurs (Wang et al., 2019; Luo et al., 2014). Similar to cyanobacterial diazotrophs, phosphate can also limit the growth of NCDs in oligotrophic environments (Rahav et al., 2015). Regarding other important factors that control autotrophic diazotrophs, iron (Fe) may potentially impact NCDs if they also depend on the high Fe-containing nitrogenases to fix N₂ (Bombar et al. 2016), although, as discussed above, the N₂ fixation by NCDs is still not
75 quantified. Strong stratification may also benefit NCDs by accumulating organic matter in the upper water column (Langlois et al. 2015). However, to our knowledge, no studies that have analyzed the effects of Fe or stratification on NCDs. such as iron (Fe) and stratification, there have been, to our knowledge, no studies reporting its relationship with NCDs.

Mesoscale eddies ~~ean may~~ also impact NCD abundance. Although anticyclonic eddies were generally considered to benefit
80 autotrophic diazotrophs by inhibiting vertical DIN input from deep waters (Liu et al., 2020; Fong et al., 2008; Church et al., 2009), a class of NCDs, ~~the~~ Gammaproteobacteria, were found to dominate diazotrophic communities inside cyclonic eddies in the South China Sea (Zhang et al., 2011; Liu et al., 2020). Different types of mesoscale eddies may have discrepancies in impacting the ecophysiology of NCDs and their autotrophic counterparts (Benavides and Robidart, 2020).

~~The major known NCD classes include bacteria such as Alphaproteobacteria, Gammaproteobacteria, Epsilonproteobacteria, Betaproteobacteria, Firmicutes belonging to Cluster I of *nifH* clusters, and some obligate anaerobic bacteria and archaea belonging to Cluster III of *nifH* clusters (Zehr et al., 2003; Riemann et al., 2010). Among them, Gamma A is the most sampled and studied phylotype. Gamma A represents a part of uncultured Gammaproteobacterial sequences isolated from the open ocean, and its cluster is distantly related to cultured NCD (Langlois et al., 2015). Gamma A *nifH* gene expression has been widely found in the global ocean, revealing its important role in marine N₂ fixation (Bird et al., 2005; Moisander et al., 2014; Langlois et al., 2015; Shiozaki et al., 2017).~~
90 Langlois et al., 2015; Shiozaki et al., 2017).

Langlois et al. (2015) have analyzed the distribution of the Gamma A phylotype in the Pacific and Atlantic Oceans and suggested that Gamma A prefers warm and oligotrophic surface oceans. With more data becoming available in recent years, we collected, to the best of our knowledge, all the reported in situ measurements of Gamma A *nifH* copies using qPCR assays, compiling a dataset with 80% more data than those used in Langlois et al. (2015). We then analyzed the relationship between

95 this *nifH*-based Gamma A abundance and the long-term background of ecological and environmental factors by using their climatological monthly averages. In addition to temperature and concentrations of nitrate, phosphate and silicate that have been used in Langlois et al. (2015), we included 5 more variables (primary production, Fe, DOC concentrations, solar radiation and mixed layer depth) to more thoroughly analyze potential controlling factors on Gamma A. We further explored the influence of mesoscale eddies on Gamma A abundance. In this study, we collected, to our best knowledge, all the reported in situ measurements of Gamma A *nifH* copies. We then analysed the relationship between this *nifH* based Gamma A abundance and the long-term background of ecological and environmental factors by using their climatological monthly averages. We further explored the short-term influence of mesoscale eddies on Gamma A abundance. Our analyses revealed-suggested that local primary productivity, temperature, dissolved Fe concentration and the occurrence of cyclonic eddies can be the main factors impacting the distribution of Gamma A in the global ocean.

105 **2 Methods**

2.1 Data summary and quality control of Gamma A abundance

A total of ~~1690-1795~~ in situ measurements of *nifH* copies of Gamma A in the Pacific, Atlantic and Indian Oceans were collected from 18 published papers (Table 1), and are available in a data repository (<https://doi.org/10.6084/m9.figshare.17284517>) (Shao and Luo, 2021). Gamma A was sometimes also named γ -24774A11 in the collected papers (Moisander et al., 2008). All these data were measured using quantitative polymerase chain reaction (qPCR). Note that the primer of Gamma A used by Langlois et al. (2015) in the North Atlantic was slightly different from other studies. Most samples (~~8788~~%) were collected in the surface-upper 100 m seawater of the water column. In the following analyses, we represented Gamma A abundance using its *nifH* copies, although we noted that variations in *nifH* copies in different cyanobacterial diazotrophic cells were have been reported (White et al., 2018; Sargent et al., 2016) and *nifH* copy numbers in Gamma A genome remain unknown.

115 The non-zero *nifH*-based abundance data of Gamma A were approximately log-normally distributed (Fig. S1). There were 682 data points reporting zero *nifH* copies which theoretically could indicate that Gamma A in the samples was either true absent or its abundance was below the detection limit. As the reported detection limit of qPCR usually ranges from 10^1 to 10^2 copies L^{-1} , the number of the Gamma A *nifH* data that could be below detection in our dataset, according to the log-normal distribution of observed non-zero data, was very likely less than 72 even assuming a high detection limit of 10^2 copies L^{-1} (Fig. S1). The fact that there were far more zero-value data (682) in our dataset indicated that a high fraction of the zero-value data could represent true absence of Gamma A.

125 The zero-value abundance data of Gamma A were not included in our further analyses, mainly because of two reasons. First, the fact that Gamma A was absent in many samples, as well as the spatially mixed distribution of the zero-value and non-zero Gamma A abundance data (see Results), indicated the patchy distribution of Gamma A, which was also widely found for other diazotrophs as a consequence of lateral transport and mixing of water masses (Robidart et al., 2014).

The patchiness of Gamma A implicated that it could be either present or absent even when the environmental conditions were suitable to its growth. That is, the presence of Gamma A requires a suitable environment, but a suitable environment does not necessarily guarantee the presence of Gamma A. If the zero-value data were included, similar environmental conditions could possibly be associated with both high abundance and zero abundance of Gamma A (Fig. S2), which would bias the response function of our statistical analyses, particularly as the fraction of the zero-abundance data was large (~1/3). Second, it is difficult to identify whether the zero-value data represented true absence or below-detection abundance of Gamma A, considering that the accuracy of qPCR was highly sensitive to sample preservation, extraction protocol and the reliance of the standard curve (Smith and Osborn, 2009). There were 616 data points reporting zero *nifH* copies which theoretically could indicate that they were below the detection limit. However, by assuming a log-normal distribution of the data, there should be only 72 data below the common detection limit of qPCR (100 copies L⁻¹) (Luo et al., 2012) (Fig. S1). Therefore, these zero-value data were more likely caused by primer specificity or insufficient volume of water samples, instead of representing the low abundance, and therefore were not used in our analyses.

Chauvenet's criterion was used to identify outliers in the non-zero Gamma A abundance data by first log-transforming all the data (Glover et al., 2011). Two outliers (0.22 copies L⁻¹ and 0.33 copies L⁻¹) were removed because their probability of deviation from the mean was smaller than 1/(2n), where n was the number of data. Even though they can be reliable, we excluded them from the analyses to avoid possible biases.

In the following analyses, we represented Gamma A abundance using its *nifH* copies, although we noted that variations in *nifH* copies in different diazotroph cells were reported (White et al., 2018; Sargent et al., 2016).

Table 1. Data source of compiled Gamma A *nifH* samples.

Reference	Location	Latitude	Longitude	Depth (m)
<i>Pacific Ocean</i>				
Moisander et al. (2008)	South China Sea	9°-12°N	107°- 110°W	0-1700

Bombar et al. (2011)	Mekong River Plume	9°-11°N	106°- 107°W	0
Hamersley et al. (2011)	Southern California Bight	33°N	118°W	5-885
Moisander et al. (2014)	South Pacific Ocean	15°- 30°S	177°E- 155°W	4-175
Shiozaki et al. (2015)	Northwest Pacific	38°- 39°N	141°- 143°W	0-119
Berthelot et al. (2017)	Western Pacific Ocean	3°- 12°S	140°- 160°W	0 -70
Shiozaki et al. (2017)	North Pacific Ocean	0°- 68°N	168°- 170°E	0-157
Shiozaki et al. (2018a)	South Pacific Ocean	0°- 40°S	170°- 100°W	0-215
Shiozaki et al. (2018b)	Kuroshio	25°-33°N	124°- 139°W	0-5
Chen et al. (2019a)	Western Pacific Ocean	0° - 21°N	110°- 159°W	0-150
<u>Chen et al. (2019b)</u>	<u>South China Sea</u>	<u>19° - 22°N</u>	<u>116°- 121°W</u>	<u>5-1000</u>
Cheung et al. (2020)	North Pacific Ocean	7°- 54°N	139°E- 80°W	5
<i>Atlantic Ocean</i>				
Langlois et al. (2008)	North Atlantic Ocean	0° - 40°N	10°- 70°W	5-100
Benavides et al. (2016)	North Atlantic Ocean	0°-21°N	15°-75°W	0-150
Martinez-Perez et al. (2016)	Tropical North Atlantic Ocean	11°-15°N	21°-60°E	5-200
Moreira-Coello et al. (2017)	Upwelling Region off NW Iberia	6°-18°N	18°-54°E	0-157
Moore et al. (2018)	Tropical Atlantic Ocean	0°-21°N	15°-55°W	0
<i>Indian Ocean</i>				
Shiozaki et al. (2014)	Arabian Sea	4°S–20°N	65° –70° E	0-86
Wu et al. (2019)	Bay of Bengal	4°S–10°N	84° –96° E	0-200

2.2 Environmental and ecological parameters

Monthly climatological environmental and ecological parameters were used as predictors for Gamma A abundance (Table 2).

155 Temperature and concentrations of nitrate, phosphate and silicate were the products of the World Ocean Atlas (WOA) 2018
(www.nodc.noaa.gov)(Locarnini et al., 2018; Garcia et al., 2019), and excess phosphate (P^*) was derived from concentrations
of phosphate and nitrate based on the Redfield ratio ($P^* = [\text{phosphate}] - [\text{nitrate}]/16$). Dissolved iron (Fe) concentrations were
obtained from the Community Earth System Model – Biogeochemistry module (Misumi et al., 2014). Dissolved organic carbon
concentration used a product estimated by artificial neural network (Roshan and Devries, 2017). Mixed layer depth (MLD)
160 was downloaded from Ifremer (<http://www.ifremer.fr/>) using the criterion that the potential density of water parcels at the

depth was 0.03 kg m^{-3} higher than that at the surface (De Boyer Montégut et al., 2004). Net primary production used a satellite data based on the VGPM algorithm (Behrenfeld and Falkowski, 1997) (<http://sites.science.oregonstate.edu/ocean.productivity/>). Surface photosynthetically active radiation (PAR) was downloaded from MODIS-Aqua program (<http://oceancolor.gsfc.noaa.gov/>). To estimate vertical profile of PAR, we first obtained the estimated euphotic zone depth Z_e (<https://oceancolor.gsfc.nasa.gov/>) at 1% surface PAR based on an inherent optical property (IOP)-centered approach (Lee et al., 2005), and used it to estimate the attenuation coefficient:

$$k_d = \frac{\ln(0.01)}{Z_e}. \quad (1)$$

The PAR at depth z can be calculated while we assumed organisms in the mixed layer were exposed to PAR homogenously:

$$PAR(z) = \begin{cases} PAR_0 e^{-k_d z} & (z > MLD) \\ \frac{1}{MLD} \int_0^{MLD} PAR_0 e^{-k_d z} dz & (z < MLD) \end{cases}, \quad (2)$$

170 where PAR_0 is the surface PAR.

To identify ~~if whether~~ the Gamma A abundance ~~is was~~ sampled in cyclonic or anticyclonic eddies, we extracted from AVISO program (www.aviso.altimetry.fr) the satellites-merged daily sea level anomaly (SLA) for the sampling days of the Gamma A data. The cores of mesoscale eddies were identified by the outermost closed contour lines of the SLA field. Only those sampling points located in cyclonic (negative SLA) and anticyclonic (positive SLA) eddies cores were recorded. Otherwise, data points were recorded as “outside eddy”. ~~we used daily sea level anomaly data (SLA) provided by AVISO program (www.aviso.altimetry.fr). Only those cyclonic (negative SLA) and anticyclonic (positive SLA) eddies with clear shapes were recorded.~~

All the variables used in the analyses are available in a data repository_ (<https://doi.org/10.6084/m9.figshare.17284517>) (Shao and Luo, 2021)

Table 2. Environmental and ecological parameters.

Environmental parameters	Symbol	Source	Spatial resolution
--------------------------	--------	--------	--------------------

Log-normally
distributed and log-
transformed

Temperature (°C)	T		1°	No
Dissolved inorganic n Nitrogenate (μM)	DIN NO ₃	World Ocean Atlas	1°	Yes
Dissolved inorganic p Phosphate (μM)	PO₄ DIP	2018	1°	Yes
Silicate (μM)	Si		<u>1°</u>	<u>Yes</u>
Excess P (μM)	P*	DIP PO ₄ - DIN NO ₃ /16	1°	No
Fe (nM)	Fe	CESM	1°	Yes
Mixed layer depth (m)	MLD	IFREMER	2°	Yes
Net primary production (mg C m ⁻² d ⁻¹)	NPP	VGPM	1/6°	Yes
Photosynthetic active radiation (E m ⁻² d ⁻¹)	PAR	MODIS-Aqua	1/6°	Yes
Dissolved organic carbon (μM)	DOC	Model	1°	No
Sea level anomaly	SLA	AVISO	<u>1/6°</u>	<u>No</u>

2.3 Statistical analyses

190 For Gamma A data points sampled in the same months and the same depth bins (defined in WOA), they were binned to 2° × 2° grids to help eliminate possible biases caused by concentrated samplings in specific regions, resulting in [893-939](#) binned means of log-10 based Gamma A *nifH* abundance. The corresponding environmental and ecological parameters were also averaged to the same bins when necessary. Univariate Pearson correlation was used to evaluate [the](#) linear relationship between Gamma A abundance and each environmental or ecological variable.

195 We also used the generalized additive model (GAM) using R package ‘mgcv’ (Wood, , 2017) to demonstrate non-linear relationships between the multiple predictors and the Gamma A abundance:

$$\mathbf{y} = \boldsymbol{\alpha} + \sum_{i=1}^n s(\mathbf{x}_i) + \boldsymbol{\varepsilon}, \quad (3)$$

where \mathbf{y} is response variable (Gamma A abundance), \mathbf{x}_i is the i th predictor (i.e., the environmental or ecological variable), $\boldsymbol{\alpha}$ is the intercept, $s(\mathbf{x}_i)$ is a linear combination of smooth functions of predictor \mathbf{x}_i , n is [the](#) number of predictors and $\boldsymbol{\varepsilon}$ is [the](#) standard error. To avoid over-fitting to noise, the Restricted Maximum Likelihood (REML) method was selected for the GAM smoothing parameters of every predictor with the basis function number (k) set to 9 (Wood et al., 2016). In the model selection of GAM, a double penalization approach was used to identify and remove those insignificant predictors with large smoothing parameters and set them to zero functions (Marra and Wood, 2011).

200


205 The scientific color maps are used in several figures to prevent visual distortion of the data and exclusion of readers with color-
vision deficiencies (Cramer et al., 2020).

3 Results and Discussion

3.1 Global distribution of Gamma A *nifH* abundance

210 The *nifH* gene abundance ranges from 1 to 10^7 copies L^{-1} in the global ocean and shows an approximately log-normal
distribution (Fig. S1). High abundance of Gamma A *nifH* abundance over 10^5 copies L^{-1} is prevalent in the subpolar North
Pacific, tropical Atlantic and Bay of Bengal (Indian Ocean) (Fig. 1). Most Gamma A abundance data were sampled above 100
m, particularly in the upper 25 m. The deepest datum with detectable Gamma A *nifH* was sampled at 340-885 m in Southern
California Bight (Hamersley et al., 2011). Available data showed that *nifH* abundance decreased with depth in the
Southwestern Pacific Ocean, the Indian Ocean and the South China Sea, but did not have an apparent trend from the surface
down to 200 m in the tropical Atlantic Ocean and the Indian Ocean (Fig. S2S3). More data particularly in deep waters are
215 needed to better and more reliably reveal the vertical pattern of Gamma A abundance.

Although high Gamma A abundance over 10^6 *nifH* copies L^{-1} was observed in the surface North Pacific Ocean, zero-value
data were also massive (215 in a total of 608 data points) and even located closely to those high-abundance data (Cheung et al.,
2020) (Fig.1), indicating the patchy distribution of Gamma A. As discussed already (Section 2.1), zero-abundance data were
not included in the further analyses due to the patchiness of Gamma A and the limitations of qPCR method in detecting the
220 true absence of Gamma A.







Figure 1. Gamma A abundance (*nifH* copies L⁻¹). The panels show data in depth ranges of (a) 0 – 25 m, (b) 25 – 100 m, (c) 100 – 200 m and (d) below 200 m. For clear demonstration, data are binned to 2° × 2° and geometric means in each bin are shown. Zero-value data were denoted as black triangles. The data are binned to 2° × 2°.

3.2 Primary production determines the carrying capacity of maximal Gamma A abundance

230 The logarithm of Gamma A *nifH* abundance positively correlated to the logarithm of net primary production (NPP) (correlation = 0.21, $p < 0.01$) (Fig. 2). More importantly, the upper bound of Gamma A abundance increased with the NPP [$\log_{10}(\text{Gamma A}) = 5.1 \text{ Log}_{10}\text{NPP} - 6.3$] when NPP < 10^{2.6} (~400) mg C m⁻² d⁻¹, above which the upper bound of Gamma A abundance saturated at ~10⁷ *nifH* copies L⁻¹ (Fig. 2).



235

Figure 2. The relationship between Gamma A abundance and net primary production. Both Gamma A abundance and net primary production (NPP) ~~are were~~ log₁₀-transformed. The data with NPP of 10^{2.0}–10^{2.6} mg C m⁻² d⁻¹ (the “low” NPP range) are divided into 6 groups with equal log-NPP intervals (i.e., divided at NPP of 10^{2.1}, 10^{2.2}, 10^{2.3}, 10^{2.4} and 10^{2.5} mg C m⁻² d⁻¹), and the highest Gamma A abundance in each group is identified (red dots). The NPP-supported maximal Gamma A abundance (red line) is estimated by linearly fitting the red dots in the low NPP range and ~~The NPP-determined Gamma A carrying capacity (red line) in the “low” NPP range [$< 10^{2.6}$ (≈ 400) mg C m⁻² d⁻¹] is estimated by linearly fitting the highest Gamma A abundance values (red dots) in NPP intervals of 10^{0.1} mg C m⁻² d⁻¹. The Gamma A carrying capacity saturates at 10^{7.0} *nifH* copies L⁻¹ for NPP > 10^{2.6} mg C m⁻² d⁻¹ (the “high” NPP range).~~

240

245

250

These results indicated that local NPP could largely determine the carrying capacity maximal Gamma A abundance, which was expected because Gamma A ~~was heterotrophic bacteria and neederequired~~ a sufficient supply of organic matter from primary producers, particularly for their energetically intensive N₂ fixation (Bombar et al., 2016). This conclusion can also be partly supported by previous experimental studies in which the addition of organic carbon enhanced heterotrophic nitrogen fixation and NCD abundance in oligotrophic seas (Benavides et al., 2015; Rahav et al., 2016; Moisaner et al., 2012; Dekaezemacker et al., 2013). Our finding contradicted the hypothesis mentioned above that Gamma A preferred oligotrophic waters based on samples mainly in tropical and subtropical Pacific and Atlantic Oceans, in which Gamma A reached 8×10^4 *nifH* copies L⁻¹

(Shiozaki et al., 2018a; Langlois et al., 2015). However, the new dataset (Cheung et al., 2020) included in the present study showed even higher (over 10^5 nifH copies L^{-1}) Gamma A abundance in the subarctic North Pacific (Fig.1) where nutrient concentrations and NPP are generally high.

We then estimated the upper bound of Gamma A abundance as a function of NPP (the red line in Fig. 2), and termed the estimated upper bound as “NPP-supported maximal Gamma A abundance” hereafter.


3.3 Univariate linear relationships between environmental factors and Gamma A abundance

~~We then analyzed what environmental factors may limit Gamma A abundance from reaching the carrying capacity. We first defined this disparity for each data point, $\Delta_{\text{Gamma A}}$, as the observed Gamma A abundance minus corresponding carrying capacity in logarithmic space. That is, $\Delta_{\text{Gamma A}}$ can be treated as the ‘residual’ of data to the carrying capacity line in Fig. 2.~~

~~Therefore, a positive correlation between $\Delta_{\text{Gamma A}}$ and an environmental factor can indicate that Gamma A prefers the increase of this factor, and vice versa.~~

~~As the Gamma A carrying capacity saturated at an NPP of $10^{2.6}$ (≈ 400) $\text{mg C m}^{-2} \text{d}^{-1}$, we then divided our dataset into a low and a high NPP groups at this threshold (Fig. 2) in further analyses to address possibly different controlling factors and mechanisms on Gamma A abundance.~~

~~In the univariate linear analyses (Fig. 3), the most correlated variable to $\Delta_{\text{Gamma A}}$ was the dissolved Fe concentration in both groups, but surprisingly the relationship was negative. Temperature and DOC were positively correlated with $\Delta_{\text{Gamma A}}$ in the low NPP group, while the relationships turned negative when NPP was high. In terms of inorganic nutrients, $\Delta_{\text{Gamma A}}$ correlated negatively to nitrate and correlated positively to the excess P (P^*) in the low NPP group, while both relationships became insignificant in the high NPP group. Phosphate had no significant relationship with $\Delta_{\text{Gamma A}}$ in either group. Silicate was positively correlated to Gamma A, and the correlation became stronger in high NPP area. The positive correlation between PAR and $\Delta_{\text{Gamma A}}$, particularly when NPP is low, can be a result of a decreasing trend of Gamma A abundance with water depth. Lastly, $\Delta_{\text{Gamma A}}$ and the mixed layer depth were negatively correlated only in the high NPP group.~~



275

~~Figure 3. Correlation between environmental factors and $\Delta_{\text{Gamma-A}}$. The values are univariate Pearson correlation coefficients. Correlations with $p > 0.05$ are considered insignificant and left blank.~~

3.43. Multivariate nonlinear relationships between environmental factors and Gamma A abundance using GAM


280

~~We then analyzed what environmental factors may limit Gamma A abundance from reaching the carrying capacity NPP-supported maximal Gamma A abundance. We first defined this disparity for each data point, $\Delta_{\text{Gamma-A}}$, as the observed Gamma A abundance minus the corresponding NPP-supported maximal Gamma A abundance carrying capacity in logarithmic space. That is, $\Delta_{\text{Gamma-A}}$ can be treated as the “residual” of data to the estimated NPP-supported maximal Gamma A abundance carrying capacity line in Fig. 2. Therefore, a positive correlation between $\Delta_{\text{Gamma-A}}$ and an environmental factor can indicate that Gamma A prefers the increase of this factor, and vice versa.~~

285

~~As the NPP-supported maximal Gamma A abundance Gamma A carrying capacity saturated at an NPP of $10^{2.6}$ (≈ 400) $\text{mg C m}^{-2} \text{d}^{-1}$, we then divided our dataset the Gamma A abundance data into a low- and a high-NPP groups at this threshold (Fig. 2) in further analyses to address possibly different controlling factors and mechanisms on Gamma A abundance in further analyses.~~

~~Although the linear univariate correlations can provide basic information in analyzing relationships between each environmental factor and the Gamma A abundance, false relationships can be generated by intercorrelations existing among the environmental variables. Additionally, the relationships can also be nonlinear. We then used a GAM multivariate analysis to partly avoid these possible problems and to obtain a more reliable~~nonlinear relationship between $\Delta_{\text{Gamma-A}}$ and environment parameters (Fig. 4~~Fig. 3)~~. Note that phosphate was not included in the GAM because ~~it was not correlated to $\Delta_{\text{Gamma-A}}$ in both the low and the high NPP groups (Fig. 3) and~~ its variance can ~~nevertheless~~ be partly represented by P^* .



300 **Figure 43. Partial effects of environmental variables on $\Delta\text{Gamma-A}$ using GAM multivariate analysis.** The analyses are
 conducted for the low-NPP group (a–h) and the high-NPP group (i–p), showing the anomaly of $\Delta\text{Gamma-A}$ contributed by the
 smooth function (blue line) and its 95% confidence interval (shadow) of each environmental variable. Data (circles) are shown
 as partial residuals after all other partial effects have been considered. The numbers in the parentheses of y-axis labels are the
 degree of freedom of the smooth functions. A degree of freedom smaller than 1 is equivalent to a linear line, and higher degrees
 305 of freedom represent more wiggly curves. The blue ticks on the x-axis also indicate the location of the data.

3.43.1 DOC

Although DOC ~~was supposedly~~ is presumed to be one of the major carbon sources for Gamma A, it did not impact $\Delta_{\text{Gamma-A}}$ in the low-NPP group (~~Fig. 4~~Fig. 3a) and even showed a negative linear relationship with $\Delta_{\text{Gamma-A}}$ under high NPP (~~Fig. 4~~Fig. 3i). First, $\Delta_{\text{Gamma-A}}$ was the residual to the NPP-supported maximal Gamma A abundance~~NPP-determined carrying capacity~~ and therefore was a collective indicator in which the effects of organic carbon production had been largely removed. Additionally, low DOC concentrations in high-NPP regions may even indicate that the DOC pool is more labile and can be more easily used (Jiao et al., 2014). Lastly, particulate organic matter (POM) can also supply necessary organic carbon and nutrients to fuel Gamma A and can even create favorable oxygen-deplete microenvironments for Gamma A (Riemann et al., 2010) (~~Farnelid et al., 2019~~), but it was not included in this study because of insufficient data.

3.43.2 Temperature

Temperature had a generally positive relationship with $\Delta_{\text{Gamma-A}}$ (Figs. ~~4b~~3b and ~~4j~~3j). This is consistent with several regional studies in which a strong positive correlation between temperature and Gamma A abundance was also found (Shiozaki et al., 2018a; Moisander et al., 2014). The relationship is expected considering the widely recognized increase in heterotrophic bacterial production with temperature in the ocean because of stimulated bacterial metabolism (Kirchman and Rich, 1997; Pomeroy and Wiebe, 2001). In addition, $\Delta_{\text{Gamma-A}}$ started to rise at a lower temperature ($\sim 15^{\circ}\text{C}$) in the high-NPP group (~~Fig. 4~~Fig. 3j) than that ($\sim 20^{\circ}\text{C}$) in the low-NPP group (~~Fig. 4~~Fig. 3b). The contribution of temperature to $\Delta_{\text{Gamma-A}}$ is larger in the low-NPP group (~~Fig. 4~~Fig. 3b) than that in the high-NPP group (~~Fig. 4~~Fig. 3j). A possible reason is that the consumption rate of less labile DOC produced in less productive regions is more sensitive to temperature (Lønborg et al., 2018; Brewer and Peltzer, 2017; Carlson et al., 2004). ~~This result implied that high temperature could be more important for Gamma A to activate accumulated semi-labile DOC in low-productive oligotrophic oceans.~~

3.43.3 Nitrate and P*

Neither nitrate nor P* had a substantial effect on $\Delta_{\text{Gamma-A}}$ (Figs. ~~4e~~3c-d and ~~4k~~3k-l). This is consistent with a previous review showing that nitrate did not show an immediate inhibition of Gamma A (Moisander et al., 2017). How and to what extent NCDs are inhibited by nitrate remains unknown (Bombar et al., 2016). Abundant Gamma A was found in oceans with high nitrate concentrations (Bird and Wyman, 2013) or shallow nitracline (Shiozaki et al., 2014). The hypothesis that low-nitrate and high-P* environments favor autotrophic diazotrophs is based on the competition of inorganic nutrients between ~~the~~ diazotrophs and other phytoplankton (Karl and Letelier, 2008; Deutsch et al., 2007), while our results tentatively indicate that competition may not occur strongly between NCDs and phytoplankton, although it is still unclear whether NCDs use inorganic or organic P sources.~~while this competition should not occur directly between NCDs and phytoplankton because the former is heterotrophic while the latter is autotrophic.~~ Nevertheless, high inorganic nutrients may still play a role in Gamma A

distribution by indirectly impacting the ~~Gamma A abundance carrying capacity of Gamma A~~ via NPP. If this is true, high nitrate ~~is then is~~ a beneficial, instead of an inhibiting, factor on NCDs.

3.43.4 Iron

340 In both the low- and ~~the~~ high-NPP groups, $\Delta_{\text{Gamma-A}}$ generally showed a decreasing trend with the increasing dissolved Fe, except for a slight increase in $\Delta_{\text{Gamma-A}}$ when the dissolved Fe increased in the range of 0.01–0.1 nM (Figs. ~~4e-3e~~ and ~~4m3m~~). Our dataset showed that a high abundance of Gamma A was prevalently observed in the North Pacific Ocean (Fig. 1a), where Fe was considered as the dominant limiting factor for N₂ fixation (Sohm et al., 2011). Other Gammaproteobacterial phylotypes such as Gamma 3 and Gamma ETSP2 were also found to dominate the diazotrophic community in the eastern South Pacific
345 (Turk-Kubo et al., 2014; Halm et al., 2012) where Fe heavily limited primary production (Knapp et al., 2016; Bonnet et al., 2008). It has also been suggested that Gamma A and unicellular cyanobacterial diazotroph UCYN-B share niches in Fe-depleted western and southern Pacific Oceans (Moisander et al., 2014; Chen et al., 2019a), possibly to avoid competing with other Fe-demanding diazotrophs. Gammaproteobacterial diazotrophs may be equipped with siderophore releasing genes, such as ~~that those~~ already reported in another versatile phylotype Gamma 4 (Cheung et al., 2021), and the released siderophores are
350 an efficient tool ~~in for~~ scavenging low-level Fe in the ocean (Boyd and Ellwood, 2010). Although more studies are certainly needed to further explore the ecological and physiological mechanisms and evolutionary reasons, the good survival of Gamma A in a low-Fe environment is an intriguing finding that may expand our recognized space of active N₂ fixation in the ocean.

3.43.5 Silicate


Our GAM results also ~~revealed-suggested~~ a positive relationship between silicate and $\Delta_{\text{Gamma-A}}$ in both the low- and the high-
355 NPP groups (Figs. ~~4f3f~~ and ~~4n3n~~), indicating a possible association between Gamma A and diatoms. NCDs have been found on the surface of diatoms ~~or on the diatom mats~~ (Martínez et al., 1983) ~~or on the diatom mats~~ as discussed above. Diatom-dominant ecosystems tend to produce abundant large particles either from dead diatoms and their aggregates or the fecal pellets generated by zooplankton (Tréguer et al., 2018). The large particles can be a good habitat for NCDs as already discussed. ~~A metagenomic study also indicated the swimming motility gene expression of NCDs and their potential particle-attached lifestyle (Delmont et al., 2018).~~ Our results then provide indirect evidence for ~~this hypothesis~~ the association between Gamma A and diatoms.

3.43.6 Light

PAR did not show a substantial contribution to the variance of $\Delta_{\text{Gamma-A}}$ in our multivariate GAM analysis (Figs. 4g-3g and 4e3o). The decrease in Gamma A abundance with depth found in the Southwestern Pacific (Fig. S32a, c and d) and in other regions by previous studies (Moisander et al., 2008; Langlois et al., 2015; Chen et al., 2019b; Shiozaki et al., 2014; Wu et al., 2019) may was therefore be attributed to higher productivity, more released photosynthetic products and higher temperature in the surface ocean, instead of the direct effects of light such as the hypothesized photoheterotrophy of Gamma A (Moisander et al., 2014). The nearly constant Gamma A abundance with depth in the Tropical Atlantic Ocean and the Indian Ocean and (Fig. 1 and Figs. S2bS3b-e) can be the results of active transport of organic matter from the surface that fuels heterotrophic N_2 fixations supported the growth of Gamma A in the dark deeper ocean.

3.43.7 Predictions based on GAM

Overall, the multivariate GAM model explained 45% and 39% of the variance in $\Delta_{\text{Gamma-A}}$ in the low- and high-NPP groups, respectively (Figs. 5a4a-b). The predicted $\Delta_{\text{Gamma-A}}$ generally followed the observed values, although it tended to underestimate the observed high $\Delta_{\text{Gamma-A}}$ (> -1) or overestimate the low $\Delta_{\text{Gamma-A}}$ (< -5) (Figs. 5a4a-b). The moderate explained variance indicated that although the tested environmental factors can substantially influence Gamma A abundance, there must be other untested factors and unknown mechanisms that can also substantially impact the Gamma A distribution.



380

Figure 54. Predictivity of GAM. Predicted $\Delta_{\text{Gamma-A}}$ versus observed $\Delta_{\text{Gamma-A}}$ are shown in (a) the low-NPP and (b) the high-NPP data groups. (c) Comparison of predicted versus observed Gamma A *nifH* abundance. The red lines are 1:1 ratio of prediction to observation.


385

As described above, $\Delta_{\text{Gamma-A}}$ was defined as the Gamma A abundance minus the corresponding NPP-supported maximal Gamma A abundance. After $\Delta_{\text{Gamma-A}}$ was predicted using GAM (Figs. 4a-b), the NPP-supported maximal Gamma A

abundance (i.e., the red line in Fig. 2) was added back to $\Delta_{\text{Gamma-A}}$ to form a prediction model for the Gamma A abundance (Fig. 4c). The GAM models for $\Delta_{\text{Gamma-A}}$ (Figs. 5a–b) were added to the estimated NPP-based carrying capacity (Fig. 2) to form a prediction model for the Gamma A abundance (Fig. 5c). Although a substantial fraction of variance in Gamma A abundance was still unexplained ($R^2 = 0.4139$), the predicted and the observed Gamma A abundances were generally consistent (Fig. 5e4c). The predicted Gamma A abundance ranged mostly on the order of 10^1 – 10^6 *nifH* copies L^{-1} , slightly narrower than that of the observations (10^0 – 10^7 *nifH* copies L^{-1}), which was mainly attributed to the performance of the GAM models for $\Delta_{\text{Gamma-A}}$ as discussed above.

Although the overall R^2 was at a moderate level of 41.39%, we applied this model to give a first-order estimate of Gamma A abundance in the surface ocean (Fig. 6a5a) from climatological NPP and environmental factors (Fig. S3S4), admitting that this demonstration did not fully cover the observed spatial variance in Gamma A abundance. The results suggested that the Gamma A was most abundant in the upwelling region in the of the Eastern Tropical South Pacific and in the Southern Ocean, the Southern Ocean and some coastal areas where, however, Gamma A was not sampled (Fig. 1). The predicted high abundance in the Southern Ocean was mostly caused by its high nitrate concentration (Figs. S3gS4g–h). However, the largest uncertainties for the predictions also exist in the Southern Ocean (Fig. 6b5b) as there were no Gamma A samples in this high-nitrate area (Fig. 1). Future sampling in the Southern Ocean can then test our predictions and reduce the uncertainties.

It was interesting that although Gamma A was undetected in all the samples in the South Pacific Gyre (Fig. 1) and all these zero-value data were not included in our GAM analyses, the prediction still showed the lowest Gamma A in this region (Fig. 5a), partly supporting the robustness of our prediction on Gamma A. However, another study suggested that NCDs were major players of N_2 fixation in this region (Halm et al., 2012), which could reflect the possibility that Gamma A may not always be the dominant NCD phylotype in the ocean. For example, Gamma 4 was suggested to be more versatile NCD phylotype in the North Pacific Ocean (Cheung et al., 2021). The prediction also showed the lowest Gamma A in the South Pacific Subtropical Gyre (Fig. 6a) where however the uncertainty was among the highest in the tropical and subtropical regions (Fig. 6b) similarly because it was under sampled (Fig. 1).



415

Figure 65. Prediction of Gamma A abundance. (a) Predicted annual mean surface (0–25 m) Gamma A abundance (*nifH* copies L⁻¹) and (b) the standard errors estimated by the GAM. Black dashed contours in (a) represent NPP of 10^{2.6} (≈400) mg C m⁻² d⁻¹.

3.5.4 Impact of mesoscale eddies on Gamma A

420


The root-mean-square error (RMSE) of 0.86 and an R² of 41.39% in the prediction model (Fig. 5e4c) indicated that there was still substantial unexplained variance in Gamma A abundance. One possible reason was that we used the climatological monthly means for the environmental factors, while the *in situ* conditions can differ greatly from the climatological values. For example, oceanic mesoscale eddies can influence biogeochemical processes not only by advective transport but also by variations in the biological and chemical environments (McGillicuddy, 2016). Particularly, as discussed above, some regional studies have suggested that mesoscale eddies may influence the distribution of autotrophic diazotrophs and/or NCDs. We then explored whether the occurrence of mesoscale eddies can impact Gamma A abundance. can be attributed to the climatological

425

~~monthly means we used in the environmental factors. We then explored whether the occurrence of short-term phenomena, such as the mesoseale eddies, can impact Gamma A abundance.~~

430 In the low- and the high-NPP groups, we identified 39 and 20 data points of Gamma A abundance that were sampled in cyclonic eddies, respectively, while more (~~204~~213 and 55, respectively) were sampled in anticyclonic eddies. This is consistent with the fact that eddies are more likely anticyclonic in the Northern Hemisphere, where our most (74%) sampling points were located (Chelton et al., 2011).

435 The results showed that in the high-NPP group, the average Gamma A abundance within cyclonic eddies was one order of magnitude higher than that in anticyclonic eddies or outside eddies (Fig. ~~7b6b~~), while the difference in the low-NPP group was much smaller and statistically insignificant (t-test, $p > 0.05$) (Fig. ~~7a6a~~). The systematically higher Gamma A abundance is unlikely to be caused by the locations of cyclonic eddies because most of the climatological factors were not significantly different across types of eddies, except for a slightly lower dissolved Fe and DOC in cyclonic eddies in the high-NPP group (t-test, $p < 0.05$) (Fig. ~~S4a-S5a~~ and ~~S4eS5c~~). We then further checked the residuals of the predicted Gamma A abundance using climatological factors (i.e., Fig. ~~5e4c~~), still finding that the Gamma A abundance in cyclonic eddies in the high-NPP group 440 was significantly higher (one-tail t-test, $p < 0.05$) than the climatology-based predictions by on average a half order of magnitude, while this was not the case for samples in anticyclonic eddies or outside eddies (Fig. ~~7d6d~~). Note that the residuals of predicted Gamma A abundance in anticyclonic eddies in the low-NPP group were also significantly but only slightly higher than 0 (one-tailed ~~ed~~ t-test, $p < 0.05$) (Fig. ~~7e6c~~).



445 **Figure 76. Influence of mesoscale eddies on observed Gamma A abundance.** (a) Gamma A abundance and (b) residuals of
 predicted Gamma A abundance using climatological NPP and environmental factors in Fig. 5c grouped according to the data
 were sampled in cyclonic eddies (CE), anticyclonic eddies (AE) or outside eddies. The box plots show the median (central
 line), 25th and 75th percentile of data (upper and lower edges of box), 5th and 95th percentile (error lines) and outliers (red
 crosses). The error bars within boxes show the mean value (purple dots) and its standard error. Values above brackets are p-
 450 p-values of two tailed t-test whether the means of observed Gamma A abundance are equal (a–b) or one-tailed t-test whether the
 residuals are greater than zero (c–d).

These results indicated that cyclonic eddies could stimulate Gamma A abundance, but only in the high productivity oceans ($> 400 \text{ mg C m}^{-2} \text{ d}^{-1}$ in this study). This finding is opposite to a previous hypothesis on autotrophic diazotrophs that anticyclonic eddies form a nitrate-depleted and well-lit environment favorable to N_2 fixation (Davis and Mcgillicuddy, 2006; Fong et al., 2008; Church et al., 2009; Liu et al., 2020). However, a sufficient supply of organic matter can play a prominent role in heterotrophic N_2 fixation when the vertical pumping of nutrition-rich water driven by cyclonic eddies (Mcgillicuddy et al., 1998) can stimulate primary production (Falkowski et al., 1991). Nevertheless, the biogeochemical consequences of mesoscale eddies can be complex (Gaube et al., 2014; Mcgillicuddy Jr, 2016). For example, in addition to vertical pumping, the eddy stirring and trapping generated by mesoscale eddies can also have spatial effects on phytoplankton (Abraham, 1998; Wiebe and Joyce, 1992). Further sampling and studies are still needed to improve our mechanistic understanding of the effects of mesoscale eddies on both autotrophic and heterotrophic N_2 fixation.

3.6-5 Reliability of Gamma A *nifH* data

It is questionable whether the *nifH* copies measured using qPCR and collected in this study can reliably represent the abundance of Gamma A or even NCDs in general. When metadata are used, the reliability of comparison among absolute quantifications of *nifH* copies can be affected by methodological factors of qPCR assays. For example, even highly reproducible standard curves may result in significant variations in quantities of the same template in separated qPCR assays due to the log nature of the curve (Smith et al., 2006). The extraction method of nucleic acids, sample preparation, variations in the efficiencies of qPCR, and differences in the qPCR platform can also impact the quantitative results (Smith and Osborn, 2009). In addition, the copy numbers of the *nifH* gene in Gamma A's genome remains unknown. There exists a large uncertainty regarding the extend to which *nifH* gene copies can represent Gamma A abundance, especially in contrast to its autotrophic counterparts. All these problems will need better technology to be resolved in the future. Previous studies found a large discrepancy between the PCR amplicon library and qPCR copy number, and suggested that there existed preferential PCR amplification of Gamma A *nifH* genes (Shiozaki et al., 2017; Turk et al., 2011). As we already pointed out, the zero values of Gamma A *nifH* copies obtained from qPCR were very likely untrustworthy because they occurred more frequently than expected (Fig. S1). Although high Gamma A abundance over 10^6 *nifH* copies L^{-1} was observed in surface North Pacific Ocean, zero value data were also massive (215 in total 608 data points) and even located close to those high abundance data (Cheung et al., 2020) (Fig. S5a). It has also been found that Gamma 4, whose primer was first designed in the Eastern South Pacific (Halm et al., 2012), might be a more versatile NCD phylotype in north Pacific Ocean (Cheung et al., 2021), although it is much less studied in other regions. Thus, whether preferential PCR amplification or qPCR detectability issue causes sampling bias in Gamma A abundance remains unknown and needs further study.

4. Summary and outlook

485 With more measurements becoming available, we explored in this study what factors controlled the distribution of a
representative phylotype of heterotrophienon-cyanobacterial diazotrophs, Gamma A, in the global ocean. The results of our
study did not fully agree with the conclusion of a previous study that Gamma A preferred warm oligotrophic oceans (Langlois
et al. 2015). Instead, Most-most of our findings imply that the supply of organic matter is the major determinant of Gamma
A's abundance, ~~confirming its heterotrophy~~. These findings ~~include~~ suggest that (1) the maximal Gamma A abundance
that carrying capacity of Gamma A abundance can occur in an environment increases with local primary production and
490 saturates at high local primary production; (2) Gamma A benefits from high temperature probably because of the accelerated
degradation rate of organic matter; and (3) cyclonic eddies may stimulate the growth of Gamma A by introducing nutrients
and elevating primary production. In addition, our analyses also ~~revealed~~ suggested that Gamma A was more abundant in Fe-
depleted areas, possibly to avoid competition with autotrophic diazotrophs in high-Fe environments. Overall, our study
suggests that productivity and Fe can be factors differentiating niches between heterotrophienon-cyanobacterial and
495 autotrophie-cyanobacterial diazotrophs in the ocean, with the former favoring a high productivity and low-Fe niche, while the
latter ~~occupying~~ occupies the opposite.

However, the moderate explanatory power of our prediction model indicates that there must be other unknown factors and
mechanisms that also impacting heterotrophienon-cyanobacterial diazotrophs. For instance, heterotrophienon-cyanobacterial
diazotrophs found in the guts of copepods (Scavotto et al., 2015) imply that they are subject to top-down controls, which was
500 also suggested for marine autotrophic diazotrophs (Landolfi et al., 2021; Wang et al., 2019; Wang and Luo, in press). ~~The~~
~~uneven spatial samplings of Gamma A may also introduce biases into our analyses. Lastly, Future studies should consider~~
~~qPCR primer and probe sets targeting a universal primer is still lacking for detecting Gamma A and~~ other NCDs such as
Alphaproteobacteria and Cluster III phylotypes, which can also be important diazotrophs particularly in previously
unrecognized regions for marine N₂ fixation (Wu et al., 2019; Langlois et al., 2008; Martínez-Pérez et al., 2018; Chen et al.,
505 2019b). The combination of PCR amplification and metagenomic data can identify a broader NCD community (Delmont et
al., 2018) and may help us design a better universal primer targeting major NCDs. Lastly, the uneven spatial samplings of
Gamma A, particularly the relatively scarce samples in the Southern Hemisphere, may also introduce biases into our analyses.
More samples and studies are needed in the future to improve our understandings of the controlling factors, niches and
distributions for heterotrophienon-cyanobacterial diazotrophs, so that their contribution to global marine N₂ fixation can be
510 better evaluated.

Data availability

All the data used in this study are available in a data repository (<https://doi.org/10.6084/m9.figshare.17284517>) (Shao and Luo, 2021).

Author contributions

515 Y.-W.L. conceived and supervised the study. Y.-W.L. and Z.S. designed the study. Z.S. collected and analyzed the data and drafted the first version of the manuscript. Y.-W.L. and Z.S. contributed to the discussion of the results and revised the manuscript.

Competing interests

The authors declare that they have no conflict of interest.

520 Acknowledgements

The authors would like to thank the scientists and crew to sample the data that used in this study. The authors also thank Hua Wang, Yuhong Huang and Xiaoli Lu from XMU for their efforts in data collection and computation support. The authors would also thank Wupeng Xiao, Kuanbo Zhou and Xin Lin for the very useful discussions with them and two anonymous referees for their very constructive comments in improving the manuscript. This study was funded by the National Natural Science Foundation of China (42076153 and 41890802).

References

- Abraham, E. R.: The generation of plankton patchiness by turbulent stirring, *Nature*, 391, 577-580, <https://doi.org/10.1038/35361>, 1998.
- Behrenfeld, M. J. and Falkowski, P. G.: A consumer's guide to phytoplankton primary productivity models, *Limnol Oceanogr*, 42, 1479-1491, <https://doi.org/10.4319/lo.1997.42.7.1479>, 1997.
- 530 Benavides, M. and Robidart, J.: Bridging the spatiotemporal gap in diazotroph activity and diversity with high-resolution measurements, *Front. Mar. Sci.*, 7, <https://doi.org/10.3389/fmars.2020.568876>, 2020.
- Benavides, M., Bonnet, S., Berman-Frank, I., and Riemann, L.: Deep into oceanic N₂ fixation, *Front. Mar. Sci.*, 5, 108, <https://doi.org/10.3389/fmars.2018.00108>, 2018a.
- 535 Benavides, M., Moisaner, P. H., Daley, M. C., Bode, A., and Aristegui, J.: Longitudinal variability of diazotroph abundances in the subtropical North Atlantic Ocean, *J. Plankton Res.*, 38, 662-672, <https://doi.org/10.1093/plankt/fbv121>, 2016.
- Benavides, M., Martias, C., Elifantz, H., Berman-Frank, I., Dupouy, C., and Bonnet, S.: Dissolved organic matter influences N₂ fixation in the New Caledonian Lagoon (Western Tropical South Pacific), *Front. Mar. Sci.*, 5, 89, <https://doi.org/10.3389/fmars.2018.00089>, 2018b.

- Benavides, M., Moisaner, P. H., Berthelot, H., Dittmar, T., Grosso, O., and Bonnet, S.: Mesopelagic N₂ fixation related to organic matter composition in the Solomon and Bismarck Seas (Southwest Pacific), PLoS One, 10, 12, <https://doi.org/10.1371/journal.pone.0143775>, 540 2015.
- Bentzon-Tilia, M., Severin, I., Hansen, L. H., and Riemann, L.: Genomics and ecophysiology of heterotrophic nitrogen-fixing bacteria isolated from estuarine surface water, mBio, 6, 4, <https://doi.org/10.1128/mBio.00929-15>, 2015.
- Berthelot, H., Benavides, M., Moisaner, P. H., Grosso, O., and Bonnet, S.: High-nitrogen fixation rates in the particulate and dissolved pools in the Western Tropical Pacific (Solomon and Bismarck Seas), Geophys. Res. Lett., 44, 8414-8423, 545 <https://doi.org/10.1002/2017gl073856>, 2017.
- Bird, C. and Wyman, M.: Transcriptionally active heterotrophic diazotrophs are widespread in the upper water column of the Arabian Sea, FEMS Microbiology Ecology, 84, 189-200, <https://doi.org/10.1111/1574-6941.12049>, 2013.
- Bird, C., Martinez, M. J., O'Donnell, A. G., and Wyman, M.: Spatial distribution and transcriptional activity of an uncultured clade of planktonic diazotrophic γ -proteobacteria in the Arabian Sea, Applied and Environmental Microbiology, 71, 2079-2085, 550 <https://doi.org/10.1128/AEM.71.4.2079-2085.2005>, 2005.
- Bombar, D., Paerl, R. W., and Riemann, L.: Marine non-cyanobacterial diazotrophs: moving beyond molecular detection, Trends Microbiol., 24, 916-927, <https://doi.org/10.1016/j.tim.2016.07.002>, 2016.
- Bombar, D., Moisaner, P. H., Dippner, J. W., Foster, R. A., Voss, M., Karfeld, B., and Zehr, J. P.: Distribution of diazotrophic microorganisms and *nifH* gene expression in the Mekong River plume during intermonsoon, Mar. Ecol. Prog. Ser., 424, 39-55, 555 <https://doi.org/10.3354/meps08976>, 2011.
- Bonnet, S., Dekaezemacker, J., Turk-Kubo, K. A., Moutin, T., Hamersley, R. M., Grosso, O., Zehr, J. P., and Capone, D. G.: Aphotic N₂ fixation in the Eastern Tropical South Pacific Ocean, PLoS One, 8, 14, <https://doi.org/10.1371/journal.pone.0081265>, 2013.
- Bonnet, S., Berthelot, H., Turk-Kubo, K., Cornet-Barthaux, V., Fawcett, S., Berman-Frank, I., Barani, A., Gregori, G., Dekaezemacker, J., Benavides, M., and Capone, D. G.: Diazotroph derived nitrogen supports diatom growth in the South West Pacific: A quantitative study 560 using nanoSIMS, Limnol Oceanogr, 61, 1549-1562, <https://doi.org/10.1002/lno.10300>, 2016.
- Bonnet, S., Guieu, C., Bruyant, F., Prášil, O., Van Wambeke, F., Raimbault, P., Moutin, T., Grob, C., Gorbunov, M. Y., Zehr, J. P., Masquelier, S. M., Garczarek, L., and Claustre, H.: Nutrient limitation of primary productivity in the Southeast Pacific (BIOCOPE cruise), Biogeosciences, 5, 215-225, <https://doi.org/10.5194/bg-5-215-2008>, 2008.
- Boyd, P. W. and Ellwood, M. J.: The biogeochemical cycle of iron in the ocean, Nat Geosci, 3, 675-682, <https://doi.org/10.1038/ngeo964>, 565 2010.

- Brewer, P. G. and Peltzer, E. T.: Depth perception: the need to report ocean biogeochemical rates as functions of temperature, not depth, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 375, 2102, <https://doi.org/10.1098/rsta.2016.0319>, 2017.
- 570 Carlson, C. A., Giovannoni, S. J., Hansell, D. A., Goldberg, S. J., Parsons, R., and Vergin, K.: Interactions among dissolved organic carbon, microbial processes, and community structure in the mesopelagic zone of the northwestern Sargasso Sea, *Limnol Oceanogr*, 49, 1073-1083, <https://doi.org/10.4319/lo.2004.49.4.1073>, 2004.
- Chelton, D. B., Schlax, M. G., and Samelson, R. M.: Global observations of nonlinear mesoscale eddies, *Prog. Oceanogr.*, 91, 167-216, <https://doi.org/10.1016/j.pocean.2011.01.002>, 2011.
- 575 Chen, M. M., Lu, Y. Y., Jiao, N. Z., Tian, J. W., Kao, S. J., and Zhang, Y.: Biogeographic drivers of diazotrophs in the western Pacific Ocean, *Limnol Oceanogr*, 64, 1403-1421, <https://doi.org/10.1002/lno.11123>, 2019a.
- Chen, T. Y., Chen, Y. L. L., Sheu, D. S., Chen, H. Y., Lin, Y. H., and Shiozaki, T.: Community and abundance of heterotrophic diazotrophs in the northern South China Sea: Revealing the potential importance of a new alphaproteobacterium in N₂ fixation, *Deep-Sea Res Pt I*, 143, 104-114, <https://doi.org/10.1016/j.dsr.2018.11.006>, 2019b.
- 580 Cheung, S., Zehr, J. P., Xia, X., Tsurumoto, C., Endo, H., Nakaoka, S.-i., Mak, W., Suzuki, K., and Liu, H.: Gamma4: a genetically versatile Gammaproteobacterial *nifH* phylotype that is widely distributed in the North Pacific Ocean, *Environ. Microbiol.*, 23, 4246-4259, <https://doi.org/10.1111/1462-2920.15604>, 2021.
- Cheung, S. Y., Nitanaï, R., Tsurumoto, C., Endo, H., Nakaoka, S., Cheah, W., Lorda, J. F., Xia, X. M., Liu, H. B., and Suzuki, K.: Physical forcing controls the basin-scale occurrence of nitrogen-fixing organisms in the North Pacific Ocean, *Global Biogeochem Cy*, 34, 9, <https://doi.org/10.1029/2019GB006452>, 2020.
- 585 Church, M. J., Mahaffey, C., Letelier, R. M., Lukas, R., Zehr, J. P., and Karl, D. M.: Physical forcing of nitrogen fixation and diazotroph community structure in the North Pacific subtropical gyre, *Global Biogeochem Cy*, 23, 2, <https://doi.org/10.1029/2008gb003418>, 2009.
- Cramer, F., Shephard, G. E., and Heron, P. J.: The misuse of colour in science communication, *Nat. Commun.*, 11, 5444, <https://doi.org/10.1038/s41467-020-19160-7>, 2020.
- 590 Davis, C. S. and McGillicuddy, D. J., Jr.: Transatlantic abundance of the N₂-fixing colonial cyanobacterium *Trichodesmium*, *Science*, 312, 1517-1520, <https://doi.org/10.1126/science.1123570>, 2006.
- de Boyer Montégut, C., Madec, G., Fischer, A. S., Lazar, A., and Iudicone, D.: Mixed layer depth over the global ocean: An examination of profile data and a profile-based climatology, *J Geophys Res-Oceans*, 109, C12, <https://doi.org/10.1029/2004JC002378>, 2004.

- 595 Dekaezemacker, J., Bonnet, S., Grosso, O., Moutin, T., Bressac, M., and Capone, D. G.: Evidence of active dinitrogen fixation in surface waters of the eastern tropical South Pacific during El Nino and La Nina events and evaluation of its potential nutrient controls, *Global Biogeochem Cy*, 27, 768-779, <https://doi.org/10.1002/gbc.20063>, 2013.
- Delmont, T. O., Pierella Karlusich, J. J., Veseli, I., Fuessel, J., Eren, A. M., Foster, R. A., Bowler, C., Wincker, P., and Pelletier, E.: Heterotrophic bacterial diazotrophs are more abundant than their cyanobacterial counterparts in metagenomes covering most of the sunlit ocean, *The ISME Journal*, <https://doi.org/10.1038/s41396-021-01135-1>, 2021.
- 600 Delmont, T. O., Quince, C., Shaiber, A., Esen, O. C., Lee, S. T. M., Rappe, M. S., MacLellan, S. L., Lucker, S., and Eren, A. M.: Nitrogen-fixing populations of Planctomycetes and Proteobacteria are abundant in surface ocean metagenomes, *Nat. Microbiol*, 3, 804-813, <https://doi.org/10.1038/s41564-018-0176-9>, 2018.
- Deutsch, C., Sarmiento, J. L., Sigman, D. M., Gruber, N., and Dunne, J. P.: Spatial coupling of nitrogen inputs and losses in the ocean, *Nature*, 445, 163-167, <https://doi.org/10.1038/nature05392>, 2007.
- 605 Ding, C., Wu, C., Li, L., Pujari, L., Zhang, G., and Sun, J.: Comparison of diazotrophic composition and distribution in the South China Sea and the Western Pacific Ocean, *Biology*, 10, 555, <https://doi.org/10.3390/biology10060555>, 2021.
- Falkowski, P. G., Ziemann, D., Kolber, Z., and Bienfang, P. K.: Role of eddy pumping in enhancing primary production in the ocean, *Nature*, 352, 55-58, <https://doi.org/10.1038/352055a0>, 1991.
- 610 Farnelid, H., Tarangkoon, W., Hansen, G., Hansen, P. J., and Riemann, L.: Putative N₂-fixing heterotrophic bacteria associated with dinoflagellate–cyanobacteria consortia in the low-nitrogen Indian Ocean, *Aquat Microb Ecol*, 61, 105-117, <https://doi.org/10.3354/ame01440>, 2010.
- Farnelid, H., Turk-Kubo, K., Ploug, H., Ossolinski, J. E., Collins, J. R., van Mooy, B. A. S., and Zehr, J. P.: Diverse diazotrophs are present on sinking particles in the North Pacific Subtropical Gyre, *ISME J*, 13, 170-182, <https://doi.org/10.1038/s41396-018-0259-x>, 2019.
- 615 Farnelid, H., Andersson, A. F., Bertilsson, S., Abu Al-Soud, W., Hansen, L. H., Sorensen, S., Steward, G. F., Hagstrom, A., and Riemann, L.: Nitrogenase gene amplicons from global marine surface waters are dominated by genes of non-cyanobacteria, *PLoS One*, 6, 9, <https://doi.org/10.1371/journal.pone.0019223>, 2011.
- Fernandez, C., Farias, L., and Ulloa, O.: Nitrogen fixation in denitrified marine waters, *PLoS One*, 6, 6, <https://doi.org/10.1371/journal.pone.0020539>, 2011.
- Fong, A. A., Karl, D. M., Lukas, R., Letelier, R. M., Zehr, J. P., and Church, M. J.: Nitrogen fixation in an anticyclonic eddy in the oligotrophic North Pacific Ocean, *ISME J*, 2, 663-676, <https://doi.org/10.1038/ismej.2008.22>, 2008.

- 620 Garcia, H., Weathers, K., Paver, C., Smolyar, I., Boyer, T., Locarnini, R., Zweng, M., Mishonov, A., Baranova, O., Seidov, D., and Reagan, J.: World Ocean Atlas 2018, Volume 4: Dissolved Inorganic Nutrients (phosphate, nitrate and nitrate+nitrite, silicate). A. Mishonov Technical, NOAA Atlas NESDIS 84, 35 pp. [dataset], 2019.
- Gaube, P., McGillicuddy Jr., D. J., Chelton, D. B., Behrenfeld, M. J., and Strutton, P. G.: Regional variations in the influence of mesoscale eddies on near-surface chlorophyll, *J Geophys Res-Oceans*, 119, 8195-8220, <https://doi.org/10.1002/2014JC010111>, 2014.
- 625 Geisler, E., Bogler, A., Bar-Zeev, E., and Rahav, E.: Heterotrophic nitrogen fixation at the hyper-eutrophic qshon river and estuary system, *Front Microbiol*, 11, 1370, <https://doi.org/10.3389/fmicb.2020.01370>, 2020.
- Geisler, E., Bogler, A., Rahav, E., and Bar-Zeev, E.: Direct detection of heterotrophic diazotrophs associated with planktonic aggregates, *Sci Rep-Uk*, 9, 1-9, <https://doi.org/10.1038/s41598-019-45505-4>, 2019.
- Glover, D. M., Jenkins, W. J., and Doney, S. C.: Modeling methods for marine science, Cambridge University Press, Cambridge, UK, 630 2011.
- Gradoville, M. R., Bombar, D., Crump, B. C., Letelier, R. M., Zehr, J. P., and White, A. E.: Diversity and activity of nitrogen-fixing communities across ocean basins, *Limnol Oceanogr*, 62, 1895-1909, <https://doi.org/10.1002/lno.10542>, 2017.
- Halm, H., Lam, P., Ferdelman, T. G., Lavik, G., Dittmar, T., LaRoche, J., D'Hondt, S., and Kuypers, M. M. M.: Heterotrophic organisms dominate nitrogen fixation in the South Pacific Gyre, *ISME J*, 6, 1238-1249, <https://doi.org/10.1038/ismej.2011.182>, 2012.
- 635 Hamersley, M. R., Turk, K. A., Leinweber, A., Gruber, N., Zehr, J. P., Gunderson, T., and Capone, D. G.: Nitrogen fixation within the water column associated with two hypoxic basins in the Southern California Bight, *Aquat Microb Ecol*, 63, 193-205, <https://doi.org/10.3354/ame01494>, 2011.
- Jayakumar, A. and Ward, B. B.: Diversity and distribution of nitrogen fixation genes in the oxygen minimum zones of the world oceans, *Biogeosciences*, 17, 5953-5966, <https://doi.org/10.5194/bg-17-5953-2020>, 2020.
- 640 Jiao, N., Robinson, C., Azam, F., Thomas, H., Baltar, F., Dang, H., Hardman-Mountford, N. J., Johnson, M., Kirchman, D. L., Koch, B. P., Legendre, L., Li, C., Liu, J., Luo, T., Luo, Y. W., Mitra, A., Romanou, A., Tang, K., Wang, X., Zhang, C., and Zhang, R.: Mechanisms of microbial carbon sequestration in the ocean – future research directions, *Biogeosciences*, 11, 5285-5306, <https://doi.org/10.5194/bg-11-5285-2014>, 2014.
- Karl, D., Michaels, A., Bergman, B., Capone, D., Carpenter, E., Letelier, R., Lipschultz, F., Paerl, H., Sigman, D., and Stal, L.: Dinitrogen 645 fixation in the world's oceans, *Biogeochemistry*, 57, 47-98, <https://doi.org/10.1023/a:1015798105851>, 2002.
- Karl, D. M. and Letelier, R. M.: Nitrogen fixation-enhanced carbon sequestration in low nitrate, low chlorophyll seascapes, *Mar. Ecol. Prog. Ser.*, 364, 257-268, <https://doi.org/10.3354/meps07547>, 2008.

- Kirchman, D. L. and Rich, J.: Regulation of bacterial growth rates by dissolved organic carbon and temperature in the equatorial Pacific Ocean, *Microb Ecol*, 33, 11-20, <https://doi.org/10.1007/s002489900003>, 1997.
- 650 Knapp, A. N., Casciotti, K. L., Berelson, W. M., Prokopenko, M. G., and Capone, D. G.: Low rates of nitrogen fixation in eastern tropical South Pacific surface waters, *P Natl Acad Sci USA*, 113, 4398-4403, <https://doi.org/10.1073/pnas.1515641113>, 2016.
- Lam, P. and Kuypers, M. M. M.: Microbial nitrogen cycling processes in oxygen minimum zones, *Ann Rev Mar Sci*, 3, 317-345, <https://doi.org/10.1146/annurev-marine-120709-142814>, 2011.
- 655 Landolfi, A., Prowe, A. E. F., Pahlow, M., Somes, C. J., Chien, C.-T., Schartau, M., Koeve, W., and Oschlies, A.: Can top-down controls expand the ecological niche of marine N₂ fixers?, *Front Microbiol*, 12, 690200, <https://doi.org/10.3389/fmicb.2021.690200>, 2021.
- Langlois, R., Grokopf, T., Mills, M., Takeda, S., and LaRoche, J.: Widespread distribution and expression of Gamma A (UMB), an uncultured, diazotrophic, gamma-proteobacterial *nifH* phylotype, *PLoS One*, 10, 17, <https://doi.org/10.1371/journal.pone.0128912>, 2015.
- Langlois, R. J., Hummer, D., and LaRoche, J.: Abundances and distributions of the dominant *nifH* phylotypes in the Northern Atlantic Ocean, *Appl Environ Microbiol*, 74, 1922-1931, <https://doi.org/10.1128/AEM.01720-07>, 2008.
- 660 Lee, Z. P., Du, K. P., Arnone, R., Liew, S. C., and Penta, B.: Penetration of solar radiation in the upper ocean: A numerical model for oceanic and coastal waters, *J Geophys Res-Oceans*, 110, C09019, <https://doi.org/10.1029/2004jc002780>, 2005.
- Li, D., Jing, H., Zhang, R., Yang, W., Chen, M., Wang, B., Zheng, M., and Qiu, Y.: Heterotrophic diazotrophs in a eutrophic temperate bay (Jiaozhou Bay) broadens the domain of N₂ fixation in China's coastal waters, *Estuar Coast Mar Sci*, 242, 106778, <https://doi.org/10.1016/j.ecss.2020.106778>, 2020.
- 665 Liu, J. X., Zhou, L. B., Li, J. J., Lin, Y. Y., Ke, Z. X., Zhao, C. Y., Liu, H. J., Jiang, X., He, Y. H., and Tan, Y. H.: Effect of mesoscale eddies on diazotroph community structure and nitrogen fixation rates in the South China Sea, *Reg Stud Mar Sci*, 35, 14, <https://doi.org/10.1016/j.rsma.2020.101106>, 2020.
- 670 Locarnini, M., Mishonov, A., Baranova, O., Boyer, T., Zweng, M., Garcia, H., Reagan, J., Seidov, D., Weathers, K., Paver, C., and Smolyar, I.: World Ocean Atlas 2018, Volume 1: Temperature. A. Mishonov Technical Ed., NOAA Atlas NESDIS 81, 52pp. [dataset], 2018.
- Loescher, C. R., Großkopf, T., Desai, F. D., Gill, D., Schunck, H., Croot, P. L., Schlosser, C., Neulinger, S. C., Pinnow, N., Lavik, G., Kuypers, M. M. M., LaRoche, J., and Schmitz, R. A.: Facets of diazotrophy in the oxygen minimum zone waters off Peru, *ISME J*, 8, 2180-2192, <https://doi.org/10.1038/ismej.2014.71>, 2014.
- 675 Lønborg, C., Álvarez-Salgado, X. A., Letscher, R. T., and Hansell, D. A.: Large stimulation of recalcitrant dissolved organic carbon degradation by increasing ocean temperatures, *Front. Mar. Sci.*, 4, 436, <https://doi.org/10.3389/fmars.2017.00436>, 2018.

- Luo, Y. W., Lima, I. D., Karl, D. M., Deutsch, C. A., and Doney, S. C.: Data-based assessment of environmental controls on global marine nitrogen fixation, *Biogeosciences*, 11, 691-708, <https://doi.org/10.5194/bg-11-691-2014>, 2014.
- Luo, Y. W., Doney, S. C., Anderson, L. A., Benavides, M., Berman-Frank, I., Bode, A., Bonnet, S., Bostrom, K. H., Bottjer, D., Capone, D. G., Carpenter, E. J., Chen, Y. L., Church, M. J., Dore, J. E., Falcon, L. I., Fernandez, A., Foster, R. A., Furuya, K., Gomez, F.,
680 Gundersen, K., Hynes, A. M., Karl, D. M., Kitajima, S., Langlois, R. J., LaRoche, J., Letelier, R. M., Maranon, E., McGillicuddy, D. J., Moisaner, P. H., Moore, C. M., Mourino-Carballido, B., Mulholland, M. R., Needoba, J. A., Orcutt, K. M., Poulton, A. J., Rahav, E., Raimbault, P., Rees, A. P., Riemann, L., Shiozaki, T., Subramaniam, A., Tyrrell, T., Turk-Kubo, K. A., Varela, M., Villareal, T. A., Webb, E. A., White, A. E., Wu, J., and Zehr, J. P.: Database of diazotrophs in global ocean: abundance, biomass and nitrogen fixation rates, *Earth Syst. Sci. Data*, 4, 47-73, <https://doi.org/10.5194/essd-4-47-2012>, 2012.
- 685 Marra, G. and Wood, S. N.: Practical variable selection for generalized additive models, *Comput Stat Data An*, 55, 2372-2387, <https://doi.org/10.1016/j.csda.2011.02.004>, 2011.
- Martínez, L., Silver, M. W., King, J. M., and Alldredge, A. L.: Nitrogen fixation by floating diatom mats: A source of new nitrogen to oligotrophic ocean waters, *Science*, 221, 152-154, <https://doi.org/10.1126/science.221.4606.152>, 1983.
- Martínez-Pérez, C., Mohr, W., Schwedt, A., Dürschlag, J., Callbeck, C. M., Schunck, H., Dekaezemacker, J., Buckner, C. R., Lavik, G.,
690 and Fuchs, B. M.: Metabolic versatility of a novel N₂-fixing Alphaproteobacterium isolated from a marine oxygen minimum zone, *Environ. Microbiol.*, 20, 755-768, <https://doi.org/10.1111/1462-2920.14008>, 2018.
- Martínez-Pérez, C., Mohr, W., Loscher, C. R., Dekaezemacker, J., Littmann, S., Yilmaz, P., Lehnen, N., Fuchs, B. M., Lavik, G., Schmitz, R. A., LaRoche, J., and Kuypers, M. M.: The small unicellular diazotrophic symbiont, UCYN-A, is a key player in the marine nitrogen cycle, *Nat Microbiol*, 1, 16163, <https://doi.org/10.1038/nmicrobiol.2016.163>, 2016.
- 695 McGillicuddy, D. J., Robinson, A. R., Siegel, D. A., Jannasch, H. W., Johnson, R., Dickey, T. D., McNeil, J., Michaels, A. F., and Knap, A. H.: Influence of mesoscale eddies on new production in the Sargasso Sea, *Nature*, 394, 263-266, <https://doi.org/10.1038/28367>, 1998.
- McGillicuddy Jr, D. J.: Mechanisms of physical-biological-biogeochemical interaction at the oceanic mesoscale, *Ann Rev Mar Sci*, 8, 125-159, <https://doi.org/10.1146/annurev-marine-010814-015606>, 2016.
- Misumi, K., Lindsay, K., Moore, J. K., Doney, S. C., Bryan, F. O., Tsumune, D., and Yoshida, Y.: The iron budget in ocean surface waters
700 in the 20th and 21st centuries: projections by the Community Earth System Model version 1, *Biogeosciences*, 11, 33-55, <https://doi.org/10.5194/bg-11-33-2014>, 2014.
- Moisaner, P. H., Beinart, R. A., Voss, M., and Zehr, J. P.: Diversity and abundance of diazotrophic microorganisms in the South China Sea during intermonsoon, *ISME J*, 2, 954-967, <https://doi.org/10.1038/ismej.2008.51>, 2008.
- Moisaner, P. H., Serros, T., Paerl, R. W., Beinart, R. A., and Zehr, J. P.: Gammaproteobacterial diazotrophs and *nifH* gene expression in
705 surface waters of the South Pacific Ocean, *ISME J*, 8, 1962-1973, <https://doi.org/10.1038/ismej.2014.49>, 2014.

- Moisander, P. H., Benavides, M., Bonnet, S., Berman-Frank, I., White, A. E., and Riemann, L.: Chasing after non-cyanobacterial nitrogen fixation in marine pelagic environments, *Front Microbiol*, 8, 1736, <https://doi.org/10.3389/fmicb.2017.01736>, 2017.
- Moisander, P. H., Zhang, R., Boyle, E. A., Hewson, I., Montoya, J. P., and Zehr, J. P.: Analogous nutrient limitations in unicellular diazotrophs and *Prochlorococcus* in the South Pacific Ocean, *ISME J*, 6, 733-744, <https://doi.org/10.1038/ismej.2011.152>, 2012.
- 710 Moore, R. M., Grefe, I., Zorz, J., Shan, S., Thompson, K., Ratten, J., and LaRoche, J.: On the relationship between hydrogen saturation in the tropical Atlantic Ocean and nitrogen fixation by the symbiotic diazotroph UCYN-A, *J Geophys Res-Oceans*, 123, 2353-2362, <https://doi.org/10.1002/2017jc013047>, 2018.
- Moreira-Coello, V., Mourino-Carballido, B., Maranon, E., Fernandez-Carrera, A., Bode, A., and Varela, M. M.: Biological N₂ fixation in the upwelling region off NW Iberia: magnitude, relevance, and players, *Front. Mar. Sci.*, 4, 303, <https://doi.org/10.3389/fmars.2017.00303>,
715 2017.
- Pedersen, J. N., Bombar, D., Paerl, R. W., and Riemann, L.: Diazotrophs and N₂-fixation associated with particles in coastal estuarine waters, *Front Microbiol*, 9, 2759, <https://doi.org/10.3389/fmicb.2018.02759>, 2018.
- Pomeroy, L. R. and Wiebe, W. J.: Temperature and substrates as interactive limiting factors for marine heterotrophic bacteria, *Aquat Microb Ecol*, 23, 187-204, <https://doi.org/10.3354/ame023187>, 2001.
- 720 Rahav, E., Giannetto, M. J., and Bar-Zeev, E.: Contribution of mono and polysaccharides to heterotrophic N₂ fixation at the eastern Mediterranean coastline, *Sci Rep-Uk*, 6, 27858, <https://doi.org/10.1038/srep27858>, 2016.
- Rahav, E., Herut, B., Mulholland, M. R., Belkin, N., Elifantz, H., and Berman-Frank, I.: Heterotrophic and autotrophic contribution to dinitrogen fixation in the Gulf of Aqaba, *Mar. Ecol. Prog. Ser.*, 522, 67-77, <https://doi.org/10.3354/meps11143>, 2015.
- Rahav, E., Bar-Zeev, E., Ohayon, S., Elifantz, H., Belkin, N., Herut, B., Mulholland, M. R., and Berman-Frank, I.: Dinitrogen fixation in
725 aphotic oxygenated marine environments, *Front Microbiol*, 4, 11, <https://doi.org/10.3389/fmicb.2013.00227>, 2013.
- Riemann, L., Farnelid, H., and Steward, G. F.: Nitrogenase genes in non-cyanobacterial plankton: prevalence, diversity and regulation in marine waters, *Aquat Microb Ecol*, 61, 225-237, <https://doi.org/10.3354/ame01431>, 2010.
- Roshan, S. and DeVries, T.: Efficient dissolved organic carbon production and export in the oligotrophic ocean, *Nat. Commun.*, 8, 2036, <https://doi.org/10.1038/s41467-017-02227-3>, 2017.
- 730 Sargent, E. C., Hitchcock, A., Johansson, S. A., Langlois, R., Moore, C. M., LaRoche, J., Poulton, A. J., and Bibby, T. S.: Evidence for polyploidy in the globally important diazotroph *Trichodesmium*, *FEMS microbiology letters*, 363, fnw244, <https://doi.org/10.1093/femsle/fnw244>, 2016.

- Scavotto, R. E., Dziallas, C., Bentzon-Tilia, M., Riemann, L., and Moisaner, P. H.: Nitrogen-fixing bacteria associated with copepods in coastal waters of the North Atlantic Ocean, *Environ. Microbiol.*, 17, 3754-3765, <https://doi.org/10.1111/1462-2920.12777>, 2015.
- 735 Shao, Z. and Luo, Y.-W.: Gamma A *nifH* abundance in the global ocean, figshare [data set], <https://doi.org/10.6084/m9.figshare.17284517>, 2021.
- Shiozaki, T., Kondo, Y., Yuasa, D., and Takeda, S.: Distribution of major diazotrophs in the surface water of the Kuroshio from northeastern Taiwan to south of mainland Japan, *J. Plankton Res.*, 40, 407-419, <https://doi.org/10.1093/plankt/fby027>, 2018a.
- Shiozaki, T., Ijichi, M., Kodama, T., Takeda, S., and Furuya, K.: Heterotrophic bacteria as major nitrogen fixers in the euphotic zone of
740 the Indian Ocean, *Global Biogeochem Cy*, 28, 1096-1110, <https://doi.org/10.1002/2014gb004886>, 2014.
- Shiozaki, T., Bombar, D., Riemann, L., Hashihama, F., Takeda, S., Yamaguchi, T., Ehama, M., Hamasaki, K., and Furuya, K.: Basin scale variability of active diazotrophs and nitrogen fixation in the North Pacific, from the tropics to the subarctic Bering Sea, *Global Biogeochem Cy*, 31, 996-1009, <https://doi.org/10.1002/2017gb005681>, 2017.
- Shiozaki, T., Bombar, D., Riemann, L., Sato, M., Hashihama, F., Kodama, T., Tanita, I., Takeda, S., Saito, H., Hamasaki, K., and Furuya,
745 K.: Linkage between dinitrogen fixation and primary production in the oligotrophic South Pacific Ocean, *Global Biogeochem Cy*, 32, 1028-1044, <https://doi.org/10.1029/2017gb005869>, 2018b.
- Smith, C. J. and Osborn, A. M.: Advantages and limitations of quantitative PCR (Q-PCR)-based approaches in microbial ecology, *FEMS Microbiology Ecology*, 67, 6-20, 10.1111/j.1574-6941.2008.00629.x, 2009.
- Smith, C. J., Nedwell, D. B., Dong, L. F., and Osborn, A. M.: Evaluation of quantitative polymerase chain reaction-based approaches for
750 determining gene copy and gene transcript numbers in environmental samples, *Environ. Microbiol.*, 8, 804-815, 2006.
- Sohm, J. A., Webb, E. A., and Capone, D. G.: Emerging patterns of marine nitrogen fixation, *Nat. Rev. Microbiol.*, 9, 499-508, <https://doi.org/10.1038/nrmicro2594>, 2011.
- Tréguer, P., Bowler, C., Moriceau, B., Dutkiewicz, S., Gehlen, M., Aumont, O., Bittner, L., Dugdale, R., Finkel, Z., Iudicone, D., Jahn, O.,
Guidi, L., Lasbleiz, M., Leblanc, K., Levy, M., and Pondaven, P.: Influence of diatom diversity on the ocean biological carbon pump, *Nat*
755 *Geosci*, 11, 27-37, <https://doi.org/10.1038/s41561-017-0028-x>, 2018.
- Turk, K. A., Rees, A. P., Zehr, J. P., Pereira, N., Swift, P., Shelley, R., Lohan, M., Woodward, E. M. S., and Gilbert, J.: Nitrogen fixation and nitrogenase (*nifH*) expression in tropical waters of the eastern North Atlantic, *ISME J*, 5, 1201-1212, <https://doi.org/10.1038/ismej.2010.205>, 2011.
- Turk-Kubo, K. A., Karamchandani, M., Capone, D. G., and Zehr, J. P.: The paradox of marine heterotrophic nitrogen fixation: abundances
760 of heterotrophic diazotrophs do not account for nitrogen fixation rates in the Eastern Tropical South Pacific, *Environ Microbiol*, 16, 3095-3114, <https://doi.org/10.1111/1462-2920.12346>, 2014.

- Wang, H. and Luo, Y.-W.: Top-down control on major groups of global marine diazotrophs, *Acta Oceanol. Sin.*, in press.
- Wang, W. L., Moore, J. K., Martiny, A. C., and Primeau, F. W.: Convergent estimates of marine nitrogen fixation, *Nature*, 566, 205-213, <https://doi.org/10.1038/s41586-019-0911-2>, 2019.
- 765 White, A. E., Watkins-Brandt, K. S., and Church, M. J.: Temporal variability of *Trichodesmium* spp. and diatom-diazotroph assemblages in the North Pacific subtropical gyre, *Front. Mar. Sci.*, 5, 27, <https://doi.org/10.3389/fmars.2018.00027>, 2018.
- Wiebe, P. H. and Joyce, T. M.: Introduction to interdisciplinary studies of Kuroshio and Gulf Stream rings, *Deep Sea Research A*, 39, v-vi, [https://doi.org/10.1016/S0198-0149\(11\)80001-4](https://doi.org/10.1016/S0198-0149(11)80001-4), 1992.
- Wood, S. N.: *Generalized additive models: an introduction with R*, CRC press, 2017.
- 770 Wood, S. N., Pya, N., and Särfken, B.: Smoothing parameter and model selection for general smooth models, *J Am Stat Assoc*, 111, 1548-1563, <https://doi.org/10.1080/01621459.2016.1180986>, 2016.
- Wu, C., Kan, J., Liu, H., Pujari, L., Guo, C., Wang, X., and Sun, J.: Heterotrophic bacteria dominate the diazotrophic community in the Eastern Indian Ocean (EIO) during pre-southwest monsoon, *Microb Ecol*, 78, 804-819, <https://doi.org/10.1007/s00248-019-01355-1>, 2019.
- 775 Zehr, J. P.: Nitrogen fixation by marine cyanobacteria, *Trends Microbiol.*, 19, 162-173, <https://doi.org/10.1016/j.tim.2010.12.004>, 2011.
- Zehr, J. P. and Kudela, R. M.: Nitrogen cycle of the open ocean: from genes to ecosystems, *Ann Rev Mar Sci*, 3, 197-225, <https://doi.org/10.1146/annurev-marine-120709-142819>, 2011.
- Zehr, J. P., Jenkins, B. D., Short, S. M., and Steward, G. F.: Nitrogenase gene diversity and microbial community structure: a cross-system comparison, *Environ. Microbiol.*, 5, 539-554, <https://doi.org/10.1046/j.1462-2920.2003.00451.x>, 2003.
- 780 Zhang, Y., Zhao, Z., Sun, J., and Jiao, N.: Diversity and distribution of diazotrophic communities in the South China Sea deep basin with mesoscale cyclonic eddy perturbations, *FEMS Microbiology Ecology*, 78, 417-427, <https://doi.org/10.1111/j.1574-6941.2011.01174.x>, 2011.