

Compositions of dissolved organic matter in the ice-covered waters above the Aurora hydrothermal vent system, Gakkel Ridge, Arctic Ocean

Muhammed Fatih Sert^{1*}, Helge Niemann^{1,2,3}, Eoghan P. Reeves⁴, Mats A. Granskog⁵, Kevin P. Hand⁶,
5 Timo Kekäläinen⁷, Janne Jänis⁷, Pamela E. Rossel⁸, Bénédicte Ferré¹, Anna Silyakova¹ and Friederike
Gründger⁹

¹Centre for Arctic Gas Hydrate, Environment and Climate (CAGE), Department of Geosciences, UiT The Arctic University
of Norway, Tromsø, Norway

10 ²NIOZ Royal Institute for Sea Research, Department of Marine Microbiology and Biogeochemistry, and Utrecht University,
Texel, the Netherlands

³Department of Earth Sciences, Faculty of Geosciences, Utrecht University, Utrecht, the Netherlands

⁴Department of Earth Science & Centre for Deep Sea Research, University of Bergen, Bergen, Norway

⁵Norwegian Polar Institute, Fram Centre, Tromsø, Norway

15 ⁶Jet Propulsion Laboratory, California Institute of Technology, Pasadena, USA

⁷Department of Chemistry, University of Eastern Finland, Joensuu, Finland

⁸Interface Geochemistry, GFZ German Research Centre for Geoscience, Potsdam, Germany

⁹Arctic Research Centre, Department of Biology, Aarhus University, Aarhus, Denmark

Correspondence to: Muhammed Fatih Sert (muhammed.f.sert@uit.no)

20 **Abstract.** Hydrothermal vents modify and displace subsurface dissolved organic matter (DOM) into the ocean. Once in the
ocean, this DOM is transported together with elements, particles, dissolved gases, and biomass along with the neutrally buoyant
plume layer. Considering the number and extent of actively venting hydrothermal sites in the oceans, their contribution to
oceanic DOM pool may be substantial. Here, we investigate the dynamics of DOM in relation to hydrothermal venting and
related processes at the as-yet unexplored Aurora hydrothermal vent field within the ultraslow spreading Gakkel Ridge in the
25 Arctic Ocean at 82.9°N. We examined the vertical distribution of DOM composition from sea ice to deep waters at six hydrocast
stations distal to the active vent and its neutrally buoyant plume layer. ~~In comparison to background sea water, we found that
the DOM in waters directly affected by the hydrothermal plume was composed of lower numbers of molecular formulas and
5-10 % less abundant compositions associated with the molecular categories related to lipid and protein-like compounds.~~
30 In comparison to background sea water, we found that the DOM in waters directly affected by the hydrothermal plume was
molecularly less diverse and 5-10 % lower in number of molecular formulas associated with the molecular categories related
to lipid and protein-like compounds. Samples that were not directly affected by the plume, on the other hand, were chemically
more diverse and had a higher percentage of chemical formulas associated with the carbohydrate-like category. We suggest,
therefore, that hydrothermal processes at Aurora may influence the DOM distribution in the bathypelagic ocean by spreading
more thermally and/or chemically induced compositions, while DOM compositions in epipelagic and mesopelagic layers are
35 mainly governed by the microbial carbon pump dynamics, and sea ice surface water interactions.

1 Introduction

Dissolved organic matter (DOM) in the vicinity of high temperature hydrothermal vents shows discrepancies in concentration compared to ocean background concentrations, mainly due to thermal decomposition of DOM during hydrothermal circulation (Hawkes et al., 2015; Lang et al., 2006). During the formation of hot hydrothermal fluids in convective circulation systems, cold deep ocean water percolates through permeable pathways (faults/fissures) in the crust and is progressively heated and modified during high temperature alteration of igneous (e.g. mafic, ultramafic) rocks (German and Seyfried, 2014; Simoneit et al., 2004). Continuous exposure to extremes of temperature (up to 400°C) and low pH either completely degrade DOM to volatile species (e.g., CO₂, methane) (Lang et al., 2006) and dissolved hydrocarbon gases (predominantly methane and trace C₂-hydrocarbons) or alters originally stable, high molecular weight, refractory DOM to highly aromatic, unsaturated, oxygen poor, petroleum-like compositions in the hydrothermal vent fluids (Hawkes et al., 2015, 2016; Rossel et al., 2015, 2017). In relatively low temperature hydrothermal systems (<150°C), DOM is not degraded by cracking (Lang et al., 2006), and production of DOM may be more prevalent (Hawkes et al., 2015). DOM may therefore be a significant source of mixed forms of complex reduced carbon in the vent fluids, which can include dissolved free amino acids (Fuchida et al., 2014; Haberstroh and Karl, 1989; Horiuchi et al., 2004), lipids (McCollom et al., 1999, 2015) and bio-labile organic carbon (Hansen et al., 2019; Longnecker et al., 2018; Rossel et al., 2015). Considering that the deep ocean DOM is refractory (e.g. Dittmar and Stubbins, 2014; Hansell, 2013) and/or primarily unavailable-inaccessible for organisms (Dittmar and Stubbins, 2014; Hansell, 2013; e.g., Arrieta et al. 2015), specialized microbial communities may therefore utilize hydrothermally modified DOM as a carbon source and support hydrothermal fauna in the vicinity of hydrothermal vents (Bart et al., 2020; Folkers and Rombouts, 2020; Hestetun et al., 2016; Yahel et al., 2003).

The chemical composition and temperature of venting fluids emanating at the seafloor and contributing to hydrothermal plume formation are mainly controlled by the geological setting and physicochemical constraints including phase separation, water-rock interaction (mineral precipitation/dissolution) and biological processes (German and Seyfried, 2014; Nakamura and Takai, 2014). Hydrothermal fluids emitted from the vents are diluted with ambient water by factors of 10⁴-10⁵ and form a hydrothermal plume that rises hundreds of meters until reaching neutral buoyancy in the water column (Baker et al., 1995). Previous field observations, as well as models, show that rising, neutrally buoyant plumes can disperse horizontally at isopycnal surfaces for tens to even thousands of kilometres (Baker et al., 1995; Lupton and Craig, 1981; Rudnicki and Elderfield, 1993; Speer and Rona, 1989; Tao et al., 2013). Thereby, they form environmental gradients (also called ecotones) conducting particles, nutrients, organic matter, trace elements and biomass (Levin et al., 2016; Ramirez-Llodra et al., 2010). These neutrally buoyant hydrothermal plumes in deep oceans are areas of active chemical cycling that sustain life in the hyper-oligotrophic bathypelagic zones. Plumes fuel chemosynthetic and heterotrophic prokaryotes, archaea, viruses and zooplankton that grow in this ecosystem in a spatio-temporal succession (Burd and Thomson, 1994; Dick, 2019; Levin et al., 2016). Morphological evidence suggest that heterotrophic deep sea communities take over chemosynthetic vent derived communities along the path of the dispersing plume layer. The heterotrophs use DOM in the plume in which subsequent lysis of their cells

Field Code Changed

Field Code Changed

Formatted: Norwegian (Bokmål)

Field Code Changed

further amplify heterotrophic bacteria and DOM release (Ortmann and Suttle, 2005). Controlled incubation experiments
70 showed that the thermal degradation of deep-sea recalcitrant DOM elevates the microbially accessible composition and
stimulate the growth of prokaryotic communities (Hansen et al., 2019).

More than 300 high-temperature venting sites have been identified at mid-ocean ridges (Hannington et al., 2011; Nakamura
and Takai, 2014) and around 800 are estimated to exist, roughly half of which are expected to be found in slow/ultraslow
spreading mid-ocean Ridges (Beaulieu et al., 2015). Because of the limited number of expeditions in the ice-covered Arctic
75 Ocean, surveys of hydrothermal vents at very high latitudes have been extremely rare. However, considering the amount of
fluid venting at hydrothermal vents, and the far-reaching area of impact through plume dispersion, their influence on marine
DOM may be substantial in the oligotrophic Arctic Ocean. There, hydrothermally derived DOM could potentially influence
mesopelagic and epipelagic layers, and surface ocean sea ice interactions.

In this paper, we investigate the influence of hydrothermal activity on the DOM composition in the Arctic Ocean water column
80 at the Aurora hydrothermal vent system. We suggest that the hydrothermal plume distributes thermally altered DOM to greater
areas, where its composition will be altered through admixture with background sea water and through microbial processes.
We characterise the water column DOM composition from the sea floor to the sea ice to assess the vertical extent of
hydrothermal intrusion and its confluence with hydrophysical and chemical parameters.

2 Methods

85 2.1 Study area

The Aurora seamount (82.897 N, 6.255 W) is located at the southern tip of the ultraslow spreading (< 12.0 mm/year) Gakkel
Ridge that extends for 1800 km across the Eurasian basin in the Arctic Ocean (DeMets et al., 2010). The seamount has a height
of 300 m above the seafloor and is elongated in a southeast-northwest direction (Fig.1). The water column at the mount's
summit has a depth of 3800 m and it reaches 4500 m at the southern side of the ridge flank. The sea surface in this region is
90 capped by a perennial ice cover. The Aurora seamount was explored by two research icebreakers in 2001 (Expedition AMORE)
and evidence for high levels of hydrothermal activity were observed (fresh sulfide chimney structures, shimmering water,
abundant biological activity) and anomalies in temperature, dissolved manganese, and light scattering were detected (Edmonds
et al., 2003; Michael et al., 2003). An active 'black smoker' hydrothermal vent at the Aurora site (named after Aurora
seamount) was later located ~100 m southwest of the summit in 2014, and high methane concentrations and temperature
95 anomalies were documented along with a plume layer extending towards northwest of vent field (Boetius et al., 2014,2015;
German and Boetius, 2017). As a continuation of the investigation in the area, and to further understand the role of the Gakkel
Ridge in the scope of the global biogeography of chemosynthetic ecosystems (Vanreusel et al., 2009), the first HACON (Hot
vents in an ice-covered ocean, HACON19) expedition attempted to reach the site again in 2019 (Bünz et al., 2020). During the
cruise, both active and inactive hydrothermal vents at the vent site were visually observed in much greater detail, using [a towed](#)
100 [camera system](#) the (Ocean Floor Observation and Bathymetry System (OFOBS); Purser et al., 2019). The active vent area is

evidently characterized by several vigorously venting black smokers, sulphide mounds, bacterial mats, thinly sedimented outcrops, and benthic communities typical of hydrothermal vent ecosystems (Bünz et al., 2020).

2.2 Water sampling and analyses

2.2.1 Sampling/ sub-sampling

105 We collected water samples from six stations near the Aurora hydrothermal vent field during the HACON19 research cruise
on-board R/V Kronprins Haakon (September - October 2019) (Fig.1). Sea ice and the upper 1000 m of the water column were
sampled at the Upper Layer (UL) station southwest of the vents. Plume (PL) and non-plume (NP1, NP2, NP3) stations were
designated to observe the spreading of the hydrothermal plume and were measured from 2000 m depth down to 10 m above
the seafloor (<4500 m water depth). A Background station (BG) was sampled further away from the seamount (Figure 1e). At
110 this station, measurements were done from 2000 - 4500 m water depth similarly to the PL and NP stations (Table 1).
Seawater samples were collected at selected sampling depths using Niskin bottles attached to a rosette ~~(Error! Reference
source not found.)~~ (Table 1). Sea ice was sampled at the surface of UL, thawed in a high-density polyethylene (HDPE)
container at room temperature and further treated similarly to the seawater samples collected with the rosette sampler.
Duplicate samples for methane concentration and stable C isotope ratios ($\delta^{13}\text{C-CH}_4$) were collected immediately after the
115 recovery of the rosette-sampler into 120 ml airtight syringes and pre-evacuated rubber-stoppered serum vials, respectively.
Care was taken to purge sampling lines of any bubbles, and methane samples were typically the first collected upon opening
of Niskins. Methane concentrations from 2 independent samples (i.e. Niskin bottles closed at the same depth) typically agreed
to within ± 10 -20% (two [standard deviation](#)s; note that we could take duplicate samples only occasionally). For DOM and
nutrient characterization, seawater was first transferred into acid-washed (2% HCl) glass bottles (2 x 1135 ml) and filtered
120 through [pre-combusted \(400 °C, 5 h\)](#) GF/F filters (Whatman) within 2 hours after collection using a low-pressure vacuum.
Nitrate, phosphate, silicate, ammonium, total nitrogen, total phosphorus, and dissolved organic carbon (DOC) samples were
collected in 60 ml HDPE bottles from the filtrate and stored at -20°C . Samples for dissolved inorganic carbon (DIC) and $\delta^{13}\text{C-}$
DIC measurement were taken and poisoned with 10 μl HgCl_2 and stored at 4°C . Seawater aliquots for $\delta^{18}\text{O-H}_2\text{O}$ analysis were
sampled in 20 ml clear glass vials and stored at 4°C . For solid phase extraction of DOM, 1 L of filtrate was acidified to pH 2
125 with HCl (37% v/v, Merck) and extracted with preconditioned (with 6 ml methanol + 12 ml pH 2 water) 500 mg PPL cartridges
(BondElut, Agilent Technologies) as described previously (Dittmar et al., 2008). Loaded cartridges were dried under air
vacuum for ~30 minutes and any organics retained in the cartridges were then eluted into pre-combusted, amber glass vials
with 2 ml methanol and stored at -20°C until analysis in Fourier-transform ion cyclotron resonance mass spectrometer (FT-
ICR MS).




Figure 1: (a) Study site in the Arctic Ocean. (b-c) Close-up of the study site showing the locations of the background station (BG), plume layer station (PL), non-plume stations (NP1 to NP3) and upper layer site (UL) on a bathymetric map (IBCAO; Jakobsson et al., 2008). (d) Direction and the velocity-vectors current velocity obtained from the LADCP (lowered acoustic Doppler current profiler) at PL station where each point represents an average of a depth bin calculated from east and north velocity components. (e) The graphical representation of station locations with respect to the dispersion of the hydrothermal plume, current direction, and the oceanic layers.

130

135

140 **Table 1: Sampling stations, positions, water depth and measured parameters at corresponding sampling depths given in parantheses for methane, $\delta^{13}\text{C}-\text{CH}_4$, dissolved organic matter (DOM) composition and nutrients (see text for details).**

Category	Station name	Latitude N	Longitude W	Water depth (m)	Sampling depths (m) and measured parameters
Background station	BG	82.916	6.255	4407	2000 ^{1,2,3} , 2750 ¹ , 3250 ^{1,2,3} , 3750 ^{1,3} , 4200 ¹ , 4300 ¹ , 4400 ^{1,2,3}
Upper Layer station	UL	82.891	6.324	3982	Sea ice ^{3,4} , 5 ^{3,4} , 25 ^{3,4} , 40 ^{3,4} , 100 ^{3,4} , 250 ^{3,4} , 500 ^{3,4} , 1000 ^{3,4}
Plume Layer station	PL	82.899	6.330	4026	2500 ^{1,3,4} , 3000 ^{1,2,3} , 3250 ^{1,2} , 3300 ^{1,2} , 3350 ^{1,2} , 3400 ^{1,2,3} , 3450 ^{1,2} , 3500 ^{1,2,3} , 3550 ^{1,2} , 3600 ¹ , 4015 ^{1,3,4}
Non-Plume stations	NP1	82.892	6.243	3870	2000 ^{1,4} , 3000 ¹ , 3075 ^{1,2} , 3150 ^{1,2,4} , 3225 ^{1,2} , 3300 ^{1,2} , 3865 ^{1,4}
	NP2	82.894	6.265	4026	2850 ^{1,2} , 2950 ^{1,2} , 3090 ^{1,2} , 3240 ^{1,2,3} , 3530 ^{1,2,3} , 3970 ^{1,2}
	NP3	82.900	6.247	4075	2000 ¹ , 2800 ¹ , 3000 ¹ , 3150 ¹ , 3300 ¹ , 4070 ¹

¹ Methane concentration, ² $\delta^{13}\text{C}-\text{CH}_4$, ³ DOM composition, ⁴ Nutrients.

2.2.2 CTD and ADCP measurements

Sensor-based profiling of the water column was conducted with a Sea-Bird 911 plus CTD (Conductivity Temperature Depth) profiler (accuracies of 0.3 db, 0.001°C, 0.002 for salinity (practical salinity scale, unitless)), SBE 43 coupled with dissolved oxygen sensor (calibrated by Winkler (1888) titration), Wet Labs ECO chlorophyll fluorometer (Excitation/Emission: 470/695 nm; sensitivity and limit of detection 0.01 µg/l) and Wet Labs C-Star beam transmissometer. As a proxy for turbidity, the beam attenuation coefficient (c , [1/m]) at 650 nm was determined by post calibration (see SI1). The sampling rosette was equipped with a HiPAP (High Precision Acoustic Profiler, Kongsberg Maritime) acoustic beacon to obtain real time positioning at depth.

Two Lowered Acoustic Doppler Current Profilers (LADCP) were mounted on the CTD rosette in downward and upward looking configurations. The CTD rosette was stopped at predefined depths for 10 min for LADCP measurements and the raw data were corrected against the continuous drift of the vessel based on HiPAP data (see SI2).

2.2.3 Analysis of sea water constituents and dissolved gases

Nitrate, nitrite, silicate, phosphate, ammonium, total phosphorus, and total nitrogen concentrations were measured with a segmented flow nutrient analyser (ALPKEM Flow Solution IV, OI Analytical), based on colorimetry, with associated detection limits and precision for nitrate (0.5 ± 0.1 µM), nitrite (0.05 ± 0.01 µM), phosphate (0.06 ± 0.01 µM), silicate (0.4 ± 0.1 µM), and ammonium (0.01 ± 0.01 µM). Dissolved organic nitrogen (DON) was determined from total dissolved nitrogen by subtracting the concentrations of dissolved inorganic nitrogen (nitrate + nitrite + ammonium). Similarly, dissolved organic phosphorus (DOP) was calculated from total dissolved phosphorus by subtracting phosphate concentration. Dissolved organic carbon (DOC) concentrations were measured with a TOC analyser (MQ-1001) utilizing a high-temperature combustion technique (Qian and Mopper, 1996). Deep Ocean DOC reference samples (Hansell Laboratory, University of Miami) were used to monitor precision and accuracy. DIC concentrations, $\delta^{13}\text{C}$ -DIC and $\delta^{18}\text{O}$ -H₂O were measured on a Thermo Scientific MAT253 isotope ratio mass spectrometry (IRMS) and presented in the delta notation as $\delta^{13}\text{C}$ relative to Vienna Pee Dee Belemnite (VPDB) and $\delta^{18}\text{O}$ relative to Vienna standard mean ocean water (VSMOW). The IRMS was calibrated with international standards VSMOW2, GISP; Greenland ice sheet precipitation, standard light Antarctic precipitation; SLAP2, VPDB and uncertainty of single measurements is ≤ 0.01 ‰. Dissolved methane concentrations were measured on-board following head-space extraction with nitrogen using a gas chromatograph (GC) equipped with a Flame Ionization Detector following the procedure used in German et al. (2010). For $\delta^{13}\text{C}$ -CH₄ measurements, samples were quantified on-board by Picarro cavity ring down spectrometer instrument (as per McDermott et al., 2017).

2.3 Fourier-Transform Ion Cyclotron Resonance Mass Spectrometry analyses and molecular formula assignments

FT-ICR MS analyses to evaluate the composition of DOM samples were carried out with a 12-T Bruker Solarix FT-ICR mass spectrometer (Bruker Daltonics, Bremen, Germany), equipped with a dynamically harmonized ICR cell (ParaCell) and an Apollo-II electrospray ionization (ESI) source, operated in both positive-ion and negative-ion modes. For the MS analyses,

100 μ l aliquots of DOM samples were diluted 1:10 (v/v) with methanol (HPLC grade). ESI was selected as the ionization technique due to its ability to ionize polar, oxygen-containing compounds present in the DOM samples (Mopper et al., 2007). The samples were directly infused into the ion source by a syringe pump, operating at a flow rate 2 μ L/min. Dry nitrogen was used as the drying and nebulizing gas. The ESI-generated ions were accumulated in the hexapole ion trap and transferred into the ICR cell for trapping, excitation, and detection. For each spectrum, 100 scans were co-added over a mass/charge ratio (m/z) range of 150-2000. The Bruker Compass fmsControl 2.1 software package was used for instrument control and data acquisition.

Initial spectral post-processing was done with the Bruker DataAnalysis 5.0 SR1 software, including an internal mass re-calibration with an in-house calibration list for the DOM samples. The data were then transferred to the PetroOrg IS-18.0.3 software (Omics LLC, Tallahassee, FL, USA) for molecular formula assignments. Only mass peaks with a signal-to-noise ratio (S/N) ≥ 5 were considered. In the assignments of the molecular formulae, monoisotopic compositions were limited to $^{12}\text{C}_{1-100}^{1}\text{H}_{1-200}^{14}\text{N}_{0-4}^{16}\text{O}_{0-30}^{32}\text{S}_{0-2}$ with a double bond equivalent (DBE) of 0-50 and a mass error of ≤ 1.0 ppm. Relative intensities were calculated by normalization with the most abundant ion in each mass spectrum (Kujawinski et al., 2009). Assigned molecular formulas were then generated for each sample at positive and negative ion modes and combined to obtain a single molecular formula list. In case of duplicate formula, relative intensity obtained from the negative ESI was considered. The combination of negative and positive mode ESI provides a better representation of DOM composition for the carboxylic and acidic compounds (negative ESI) and hydrogen saturated aliphatic compounds (positive ESI) (Ohno et al., 2016; Sert et al., 2020).

From the combined formula lists, chemical characterization of DOM compositions was conducted to obtain percentages of (i) three categories of heterogeneous atomic content as CHO, CHON and CHOS, and (ii) H:C and O:C atomic ratios and ranges on van Krevelen diagrams (Kim et al., 2003). Considering the H:C and O:C atomic ratios of major biomolecules, ranges of H:C and O:C were generically associated with four biochemical compound categories, disregarding structural differences and atomic compositions of individual formulas in the group. These categories are (a) lipid- and protein-like (LPD) for $\text{H:C} \geq 1.5$, $\text{O:C} \leq 0.67$, (b) carbohydrates and amino-sugar-like (CAR) for $\text{H:C} \geq 1.5$, $\text{O:C} > 0.67$, (c) unsaturated hydrocarbons and condensed aromatics (UHC) for $\text{H:C} < 1.5$, $\text{O:C} < 0.1$ and $\text{H:C} < 0.7$, $\text{O:C} < 0.67$, and (d) lignin- and tannin -like (LGN) for $1.5 < \text{H:C} < 0.7$, $\text{O:C} > 0.67$. Boundaries were modified from Hockaday et al. (2009) and Hodgkins et al. (2016) as explained previously (Sert et al., 2020). For calculating percentages, the number of formulas that associated with a given category was divided by the total number of formulae in the sample and multiplied by 100. For example, the sum of the percentages of LPD, CAR, LGN and UHC is equal to 100%, and similarly, the sum of the percentages of CHO, CHON and CHOS is equal to 100% for each sample.

2.4 Statistical analyses of DOM samples

Statistical analyses were performed in R (R Core Team, 2018). Diversity indices for DOM samples were calculated by the 'diversity' function analogous to biodiversity in ecology using the Shanon-Weaver formulation (Oksanen, 2019) as explained

previously (Sert et al., 2020). A Bray-Curtis dissimilarity (distance) matrix (Bray and Curtis, 1957) was constructed by using relative abundances-intensities of formulas from 11 DOM samples that were collected >2000 m in stations BG, PL and NP2. Hierarchical cluster analysis was applied for DOM compositions using a Ward clustering algorithm ('hclust' function in Vegan) on a Bray-Curtis dissimilarity matrix. A heat map was constructed to visualize the sample clusters and the relative dissimilarities. Principal coordinate analysis (PCoA) was applied by using 'pcoa' function in the R package APE (Paradis and Schliep, 2019) on the Bray-Curtis dissimilarity matrix. The ordination plot was constructed from the first two main axes that represent the highest variation between samples. Percentages of explained variances on two main axes were calculated by dividing the two highest eigenvalues to the sum of all eigenvalues.


215 3 Results and Discussion

3.1 Water column physical and chemical properties


Water circulation in the Eurasian part of the Arctic Ocean is mainly characterized by Atlantic inflow and Arctic outflow through the Fram Strait, with a monthly mean volume of 9.5 and 11 Sv, respectively (Fahrbach et al., 2001). We categorize the water column in three water layers as epipelagic (0-200 m), mesopelagic (200-1000 m) and bathypelagic (1000-4000 m) considering the depth and the four water masses that are defined by the density layers of $\sigma_t \leq 27.7$ for Polar Surface Waters (PSW), $27.7 < \sigma_t \leq 27.97$ for Arctic Atlantic Waters (AAW), $\sigma_t > 27.97$ and $\sigma_{\theta, S} \leq 30.444$ for Arctic Intermediate Waters (AIW) and $\sigma_{\theta, S} > 30.444$ for Deep Waters (DW) (Marnela et al., 2008; Rudels et al., 2005). The epipelagic layer contains PSW until 165 m depth with temperatures ranging between -1.7 and 0.4°C and salinity from 31.6 to 34.65 (Fig. 2a-e). AAW exists beneath PSW from the bottom of the epipelagic layer down to 450 m and includes a temperature maximum of 1.27°C at 350 m. From 225 450 to 1400 m, the temperature decreases from 0.9 to -0.5°C in a steep thermocline, forming AIW. AAW and AIW together form the East Greenland Current that carries Arctic waters southwest along the Greenland coasts and gradually mixes with the convected surface waters (Rudels et al., 1995). Below the AIW layer, water temperature decreases to -0.73°C at around 2500 m where DW fills the Eurasian basin. Here, the density is largely controlled by salinity rather than temperature and the shape of the density profile is almost identical to the salinity profile (Fig. 2b-c). Despite the dominant sea ice cover at our study area, 230 preventing solar radiation from reaching the water column, a chlorophyll *a* (Chl *a*) fluorescence signal was detected down to 200 m depth (Fig. 2d). At all stations, the first 40 m below the sea surface showed a Chl *a* maximum with an average concentration of 0.17 µg/L, followed by a decrease to 0.10 µg/L at 50 m. From this depth, concentrations gradually decreased further until the detection limit (0.01 µg/L) was reached at the bottom of the epipelagic layer (Fig. 2d).

Nutrient profiles, nutrient ratios, $\delta^{13}\text{C}$ -DIC and $\delta^{18}\text{O}$ all display a typical deep ocean surface-to-bottom gradient and were 235 identical at all stations (Fig. 3). Nitrate, phosphate, and silicate concentrations were lowest at the surface (5 m) with 1, 0.44, and 6.1 µM, respectively, and increased gradually to average bathypelagic concentrations of 14.8, 0.96 and 12.0 µM at 2000 m (Fig. 3a-c). On the contrary, dissolved organic nutrients were highest in the epipelagic layer with the maximum concentrations of 92 µM for DOC at 100 m and 6.2 µM for DON at the surface (Fig. 3e-f). Similarly, ammonium and DOP

240 were detectable only in the epipelagic layer with average concentrations of 0.2 and 0.1 μM , respectively (see Fig. 3d for ammonium, DOP is not plotted). Nitrate, phosphate, and silicate concentrations in sea ice were below detection limit, however detectable concentrations of ammonium (0.6 μM), DOC (4.0 μM), and DON (0.6 μM) were measured in sea ice.



245 **Figure 2:**Depth profiles of (a) in-situ temperature, (b) salinity on practical salinity scale, (c) potential density, (d) Chl *a* fluorescence and (e) potential temperature vs salinity profile along with the characterized water masses, Polar Surface Waters (PSW), Arctic Atlantic Waters and Arctic Intermediate Waters (AAW/AIW), Arctic Deep Waters (DW) in background (BG), upper layer (UL), plume (PL) and non-plume (NP1-NP3) stations.






Figure 3: Depth profiles of (a-d) dissolved inorganic nutrients, (e) dissolved organic carbon (DOC), (f) dissolved organic nitrogen (DON), (g) $\delta^{13}\text{C}$ of dissolved inorganic carbon (DIC) and (h) $\delta^{18}\text{O}$ of water, (i-l) nutrient ratios in sea ice and sampling stations background station (BG), and in sampling stations in the upper layer (UL), plume (PL) and non-plume (NP1, NP2). Axis breaks are used for a better visualization of ammonium concentration and $\delta^{13}\text{C}$ -DIC values. Total dissolved nitrogen (TDN) = nitrate + nitrite + ammonium + DON. Red dashed lines indicate non-buoyant plume layer at PL.

260 $\delta^{18}\text{O}$ values were obtained in three different concentration ranges that changed with depth. Sea ice had a $\delta^{18}\text{O}$ value of -4.53 ‰. At the top 100 m, $\delta^{18}\text{O}$ had an average value of -2.24 ‰. From 250 m to bottom $\delta^{18}\text{O}$ was uniform with an average of 0.36 ‰ (Fig. 3h). This corresponds to water mass properties with contributions from sea ice melt waters and Arctic river runoff with negative $\delta^{18}\text{O}$ values in the upper 100 m within PSW, compared to highly uniform deep-water ratios. Our $\delta^{18}\text{O}$ measurements are comparable to previous measurements in the area that reported ~0.3 ‰ in deep water and -2.6‰ to -1.6 ‰ in the upper 100 m (Bauch et al., 1995; Östlund and Hut, 1984).

265 A $\delta^{13}\text{C}$ -DIC value of -13 ‰ was measured in sea ice. $\delta^{13}\text{C}$ -DIC values decreased from 0.86 ‰ at the surface to 0.44 ‰ at 250 m and gradually increased to 0.62 at 1000 m. $\delta^{13}\text{C}$ -DIC changed in a narrow range from 2000 m to bottom with average composition of 1.03 ‰ (Fig. 3g). Sea ice coverage in the Arctic Ocean has been decreasing by ~9 % per decade, leading to a longer growing period for phytoplankton and associated increase in net primary production (Arrigo and van Dijken, 2015 and references therein). Taken together with the imprint of the anthropogenic CO_2 (with $\delta^{13}\text{C}$ value of ca. -8.5‰), $\delta^{13}\text{C}$ -DIC values
270 obtained at the sea ice and the upper layer of the water column are linked to contributions of several factors, including biological productivity, air sea gas exchange and sea ice or brine formation, which admixes DIC with a lighter $\delta^{13}\text{C}$ signature compared to that of deep waters. E.g., deep water $\delta^{13}\text{C}$ -DIC values of 1.0 ± 0.2 ‰ were found in the Canada Amundsen and Nansen basins (Bauch et al., 2015; de la Vega et al., 2019).

~~No substantial primary production occurred in the~~ At the time of sampling, the net growth of phytoplankton seems to decelerate
275 ~~at the top 50 m layer of the water column, as shown by the~~ considering the nutrient minima depletion and the subsurface
~~maxima of~~ Chl *a* ~~maxima at~~ ~40 m water depth. During the Arctic summer (from March to September), this layer gets depleted in inorganic nutrients and enriched in dissolved organic matter (Thingstad et al., 1997). In the deeper layers, nitrate/phosphate and nitrate/silicate ratios are almost identical to the Redfield ratios of 16:1 and 1:1, respectively (Redfield, 1958) (Fig. 3i-j). Ratios of nitrate/phosphate (2.3:1 – 5.2:1) and nitrate/silicate (1:5 – 1:2.5) in the 0-100 m layer indicate that
280 nitrate limits primary production. With the contribution of the epipelagic DON, however, surface/subsurface TDN/Phosphate (16.3:1) and TDN/Silicate (1.2:1) ratios became comparable with the mesopelagic/bathypelagic ratios (Fig. 3l). This indicates that DON may be used as a nitrogen source to support bacterial growth, considering that such low primary production cannot sustain the bacterial demand in relation to limited light and low water temperature (Fouilland et al., 2018). Besides nitrogen, silicate was also limiting at the surface waters when compared to phosphate (silicate/phosphate ratios of ~13:1). Thus, perhaps
285 due to diatom growth, silicate demand accompanies bacterial phosphate regeneration below the surface, lowering the silicate/phosphate ratios (7.6:1 – 9.6:1) compared to the deep ocean (12.4:1) (Fig. 3k).

3.2 Dispersion of the hydrothermal plume


Turbidity anomalies, elevated methane concentrations and varying $\delta^{13}\text{C}$ - CH_4 values in the vertical profiles imply that the plume rises from the ~3900 m deep vent up to ~3000 m in the water column, and spreads laterally northward in the direction of the
290 prevailing water current (Fig. 1d). Irrespective of the vent setting, tectonic structure, and background currents, previous studies

Formatted: Keep with next

and models imply that the fundamental physics behind the spreading of hydrothermal plumes are commonly characterized by buoyant flow dynamics that have been described in depth for atmospheric systems (Tao et al., 2013). Upon entering to the water column, a multi-phase plume rises and spreads laterally and carries dissolved gases and entrained particles that alter turbidity and methane concentrations in the non-buoyant plume layer (Fig. 4). Fundamental anomalies of venting fluids such as elevated temperature or low salinity can, on the other hand, become rapidly overprinted by ambient seawater background hydrography, making potential anomalies in these parameters undetectable with our instrumentation (Fig. 2). Similarly, parameters that are not strongly influenced by the vent plume intrusion (i.e., nutrients, DOC and DON concentrations) do not indicate any anomaly in relation to plume dispersion distant from the vent due to dilution with seawater (Fig. 3). This seems contradictory to molecular changes in DOM compositions, [\(as detailed further in Section 3.3.3\)](#); however, it must be stated that solid phase extractable DOM represents only a portion of bulk DOM (43 to 62 %; Dittmar et al., 2008), and compositional changes are not necessarily always coupled to large or substantial concentration changes. In essence, DOM molecular compositions may, more subtly, preserve some broad variations in endmember DOM traits than DOC and DON concentrations (Osterholz et al., 2016).

Continuous drifting (often ~0.2 kts or more) of massive (km-scale, ~1 m thick) ice floes prevented the vessel from keeping stable positions on station while taking samples, and, together with time constraints, related to other operational needs of the expedition precluded us from directly sampling the buoyant stem of hydrothermal plume immediately above the active Aurora site. Due to the extreme depths (near 4 km), it furthermore typically took 3–4 hours to complete a single CTD cast, so that the distance between start and end positions of a given cast were often 300–500 m apart. Nonetheless, the exact location of sampling was determined based on HiPAP data. The thickness of the non-buoyant plume also varied depending on the position of the sampling station in relation to the vent location, current direction, and bathymetric features. Based on turbidity anomalies, methane concentrations, and $\delta^{13}\text{C-CH}_4$ contents, the plume layer was most evident at station PL, which was, counterintuitively, the station furthest away from the seamount (Fig. 4a-c), yet consistent with the prevailing current direction (see Fig. 1d). Here, methane concentrations reached a maximum of 5.1 nM at 3363 m, which is >10-fold higher compared to ambient seawater concentrations (0.3 nM) as measured at similar depths at BG and coincide with the sharp maximum in turbidity. A small increase in turbidity was also measured at the other stations from 3000 to 3250 m implying some level of perturbation from the hydrothermal intrusion (Fig. 4c), but these were not as strong as in PL and not accompanied with an increase in $\delta^{13}\text{C-CH}_4$ values or methane concentrations. We found one exception to this at NP1, which was also at the northern side of the vents. Here, methane concentration (1.7 nM) and $\delta^{13}\text{C-CH}_4$ (-43‰) were elevated at 3350 m, i.e., just below the turbidity anomaly.


Formatted: English (United States)



325 **Figure 4: Depth profiles of (a) methane (CH_4) concentration, (b) $\delta^{13}\text{C-CH}_4$ values and (c) beam attenuation coefficient at 650 nm (c) between 2000 and 4500 m depths in background (BG), upper layer (UL), plume (PL) and non-plume (NP1-NP3) stations.**

Deep water methane stable carbon isotope compositions in the Arctic Ocean have previously been reported with values ranging from -63 ‰ to -58‰ in the Beaufort and Chukchi Seas, along with higher concentrations of methane of up to 55 nM (Damm et al., 2008; Kudo et al., 2018; Lorenson et al., 2016). These are attributed to microbial methanogenesis overprinting the atmospheric methane equilibrium signature (typically ~3.5 nM and -47 ‰). We observed a similar range of $\delta^{13}\text{C}\text{-CH}_4$ values with much lower concentrations of methane (e.g., -64.7 ‰ and 0.3 nM) at BG (3250 m) and argue that these values may also be caused by methanogenesis, possibly in sediments or even in the water column (Damm et al., 2010; Grossart et al., 2011; Karl et al., 2008) but we suggest that methanotrophs also consume methane there (Reeburgh, 2007). In contrast, we found relatively heavy methane isotope values together with maximum methane concentrations in the plume layer at PL (e.g., -33 ‰ and 5.1 nM) (Fig. 4). Methane at hydrothermal vents is typically produced non-biologically, either by abiogenic (C reduction) or thermogenic (breakdown) processes and is generally characterised by much higher $\delta^{13}\text{C}$ -values (Whiticar, 1999; Reeves and Fiebig, 2020; Baumberger et al., 2016 and references therein). Abiogenic $\delta^{13}\text{C}\text{-CH}_4$ values of *ca.* -5 to -20 ‰ are typically found in bare-rock (mafic/ultramafic systems), while lower (thermogenic) values of ~ -25 to -56‰ are prevalent in systems influenced by hydrothermal alteration of sedimentary organic matter (Charlou et al. 2010; Baumberger et al. 2016; Wang et al. 2018; and references therein). Hence, the higher $\delta^{13}\text{C}\text{-CH}_4$ values detected at PL provide strong evidence for a similar non-biogenic origin of plume methane venting from the Aurora site. While $\delta^{13}\text{C}\text{-CH}_4$ values measured at the Aurora hydrothermal field are relatively depleted compared to many bare-rock hosted hydrothermal systems, to some degree, this can be explained by both the distal nature of PL samples relative to the more concentrated buoyant plume, and admixing of background sea water with its extremely isotopically depleted $\delta^{13}\text{C}$ signature and low methane concentration values (-64.7 ‰ and 0.3 nM). To further probe the dynamics of water column methane dynamics at Aurora, we applied a 'Keeling' mixing model approach (Keeling, 1961; Keir et al., 2006). This end-member isotope mixing model is widely used in the fields of atmospheric chemistry/physics to analyse admixtures of two isotopically distinct gas sources, and to determine the source isotope composition (e.g., Pataki et al., 2003). In a Keeling plot, the reciprocal of concentration is plotted against isotope composition (Fig. 5). In case of a linear mixing, the values scatter along a straight line and the intercept with the y-axis denotes the isotope value of the source signal. Our values of methane concentration and isotope composition show predominantly linear behaviour, indicating that the mixing and dilution dominates the observed concentration and isotope systematics, as has been found in other Arctic settings (Damm and Budéus, 2003; Graves et al., 2015). More complex mixing scenarios involving mixing and methane oxidation were also suggested to explain methane isotope patterns in hydrothermal plumes (Keir et al., 2009). Though we cannot completely rule out some extent of microbial methane oxidation (particularly in the most distal samples), this process is associated with a significant kinetic isotope effect (Whiticar, 1999), and can even result in extremely high $\delta^{13}\text{C}\text{-CH}_4$ values in microbially 'aged' hydrothermal plumes (Keir et al. 2006; Cowen et al. 2002). If such oxidation were pronounced, it would lead to substantial curvature in a Keeling plot by shifting the residual methane towards heavier isotopic values at lower concentrations (higher $1/[\text{CH}_4]$).

360






Figure 5: Keeling plot for $\delta^{13}\text{C}-\text{CH}_4$ versus $1/[\text{CH}_4]$ for the data obtained in background (BG), upper layer (UL), plume (PL) and non-plume (NP1-NP3) stations (Keeling, 1961; Keir et al., 2006).¹McDermott et al., 2017,²Baumberger et al., 2016. Plotted also are 'closest approach' $\delta^{13}\text{C}-\text{CH}_4$ value reported for the 87°N hydrothermal plume on Gakkel Ridge (McDermott et al., 2017), endmember hydrothermal fluid $\delta^{13}\text{C}-\text{CH}_4$ values from Mid-Atlantic Ridge black smoker systems known to be influenced by ultramafic rock, and the sediment-influenced Loki's Castle system (Baumberger et al., 2016 and references therein).

Formatted: Superscript

Formatted: Superscript

370

Considering a ‘mixing without oxidation’ scenario and under the assumption of linear mixing (Keir et al., 2009), we estimate the $\delta^{13}\text{CH}_4$ value of the hydrothermal fluid source to be ~~about~~ -36‰ using a least squares linear regression of the entire plume methane dataset ($R^2 = 0.59$), with an intercept uncertainty of $\pm 3\text{‰}$ at the Aurora hydrothermal field (Fig. 5). The estimate is interesting, since the Aurora hydrothermal system has, thus far, been postulated to be hosted in ultramafic rock (German and Boetius, 2017; Marcon et al., 2017). The $\delta^{13}\text{CH}_4$ estimate presented here is substantially lower than both plume and endmember $\delta^{13}\text{CH}_4$ values of many Mid Atlantic ridge hydrothermal vents known to be hosted in or influenced by ultramafic rock (ca. -5 to -20 , e.g., Keir et al., 2006; Wang et al., 2018). Such ranges are assumed to be characteristic of putatively ‘abiotic’ methane (e.g., Wang et al., 2018), most likely derived from leaching of rock-hosted gas inclusions (e.g. Grozeva et al., 2020). A $\delta^{13}\text{CH}_4$ value -13.1‰ within this range of has also been reported for a ‘closest approach’ in a CH_4 - and H_2 -rich hydrothermal plume further east on Gakkel Ridge at 87°N (McDermott et al., 2017; confirmed by the same cavity ring-down spectrometry approach used here), also postulated to reflect ultramafic-influence. In contrast, $\delta^{13}\text{CH}_4$ values in hydrothermal fluids either heavily influenced by (or hosted in) sediments undergoing substantial thermal alteration of associated sedimentary organic matter tend to be generally more ^{13}C -depleted (ranging from -25 to -56‰ ; Baumberger et al. 2016 and references therein). Indeed, similarly ‘intermediate’ values of $\delta^{13}\text{CH}_4$ (-27 to -29‰) compared to Mid-Atlantic Ridge hydrothermal systems were also obtained in the bare-rock hosted, but sediment-influenced, Loki’s Castle hydrothermal system farther to the south on the ultraslow spreading Mohns-Knipovich ridge junction of the Arctic Mid Ocean Ridge system (Baumberger et al., 2016; Pedersen et al., 2010). There, exceptionally high methane (15.5 mM) and ammonium (6.1 mM) concentrations together with low C_1/C_2 ratios are attributed to subsurface hydrothermal fluid-sediment interactions (linked to sediment accumulation at ridge flanks and ridge valley, Baumberger et al., 2016; Pedersen et al., 2010).

390 A greater extent of plume dispersion and somewhat higher methane concentrations (≤ 32 nM) were previously reported within ~ 200 m of the Aurora hydrothermal vent area (Boetius, 2015; German and Boetius, 2017) but the maximum concentrations reported here (2.8–5.1 nM, in station PL) do at least coincide with the highest $\delta^{13}\text{CH}_4$ values (-31 to -33‰) and maximum water column turbidity. Hence, we tentatively suggest that hydrothermal fluids at Aurora might, on the basis of these closest approach plume $\delta^{13}\text{CH}_4$ values alone, potentially contain a thermogenic organic matter influence. ~~However, as yet, the endmember methane concentration and carbon isotope composition of the hydrothermal fluids venting at Aurora are not known, precluding a proper assessment of the geochemical processes influencing methane there.~~

3.3 Composition of DOM in the sea ice, water column and in the hydrothermal plume layer


Analyses of DOM extracts revealed 8093 unique formulas in the whole dataset, with the molecular mass range between 150 and 1000 Da. From all unique formulas, 2614 and 2391 were uniquely obtained by negative and positive ESI, respectively and 3088 formulas were assigned in both (see Fig. S3). In agreement with previous findings (Ohno et al., 2016; Sert et al., 2020), unique positive ESI formulas were markedly more hydrogen saturated (average $\text{H:C}=1.51$) and oxygen poor (average $\text{O:C}=0.31$) compared to negative ESI formulas (average $\text{H:C}=1.13$ and $\text{O:C}=0.58$). Numbers of assignments for aliphatic and lipid-


like DOM components were more abundant in positive mode and unsaturated oxygen rich molecular formulas, carbohydrates, lignin-like and aromatic compounds were more abundant in negative mode. Considering the features obtained in different modes (Fig. S4), combining positive and negative ESI datasets provides a considerable advantage for differentiating samples.


3.3.1 Sea ice

The DOM composition of sea ice was determined from a single sample and indicates that it was predominantly more hydrogen-saturated relative to surface water composition (i.e., higher H/C ratios), with lower number of formulae, average relative intensities abundances, average molecular weights, and considerably high LPD content (Fig. 6a-k, Fig. S2). Differences in molecular composition between seawater and sea ice are in agreement with ~~the~~ previous observations in sea ice DOM (Longnecker, 2015; Retelletti Brogi et al., 2018) and lake ice DOM (Liu et al., 2020; Santibáñez et al., 2019; Xu et al., 2020), implying a selective transport of DOM with less complex, aliphatic compounds - rather than larger aromatic oligomers, during ice formation. The hydrodynamics of DOM transport are not known between phases, but previous observations show that DOM segregates between water and ice, favouring lower MW formulae and less aromatic fractions in the ice phase (Belzile et al., 2002; Santibáñez et al., 2019). In the Arctic Ocean, this partition process increases the bioavailability of terrestrial DOM from rivers during initial ice formation and contributes to its removal by retaining biologically active components (Jørgensen et al., 2015). In addition to the abiotic fractionation and transport, biofilm mediated microbial degradation and photooxidation were previously shown to be responsible for fast degradation of petroleum biopolymers in sea ice (Vergeynst et al., 2019). High LPD percentageabundance and low average MW in sea ice DOM may also be attributed to a fresh production by ice algae or heterotrophic degradation of DOM (Hill and Zimmerman, 2016; Li et al., 2019; Retelletti Brogi et al., 2018).

Field Code Changed







425 Figure 6: Depth profiles of the dissolved organic matter (DOM) composition, (a-d) molecular indices, (e-h) percentages of number of formulae in biochemical compound groups defined by H:C and O:C ratios and (i-k) heteroatomic content in background (BG), upper layer (UL), plume (PL) and non-plume (NP2) stations. Abbreviations are MW, molecular weight; LPD, lipid- and protein-like; CAR, carbohydrate-like; LGN, lignin- and tannin-like; UHC, unsaturated hydrocarbon- and condensed aromatic-like; O:C, oxygen to carbon ratio; H:C, hydrogen to carbon ratio. Beam attenuation profile of PL is added to the first panel to indicate non-buoyant plume layer and corresponding boundaries of the plume layer were indicated by red dashed lines in all panels.

430

3.3.2 Upper 1000 m of the water column

From the sea surface down to 1000 m, changes in DOM indices such as the number of formulas, molecular diversity, and average relative abundances-intensities are consistent with changes in nutrient and DOC showing their maxima at 5 m (Fig. 6a-d, Fig. S23a-e). This is in agreement with the expected pace and diversity of biological processes in this water layer, i.e., dissolved nutrient uptake by phytoplankton and the contribution to the synthesis of dissolved and particulate organic matter (Benner et al., 1992; Hedges, 1992). Below this layer, DOM-average relative abundances-intensities of formulas decreased and showed a less diverse composition below the photic zone with decreasing light availability and heterotrophic consumption (Fig. 6b-d, Fig. S2).

The uptake of bioavailable DOM alongside nutrients could explain the distribution of formulas of different chemical classes in the 0-1000 m layer. For example, lower abundances of LPD and LGN at the surface against the higher CAR and UNC contents (Fig. 6e-h) indicate the selective use of the more hydrogenated aliphatic compounds compared to oxygenated or aromatic formulas. Similarly, CHON and CHOS heteroatom contents were maximal-higher at the surface compared to subsurface contents up to 500 m depth (Fig. 6i-k), indicating the increase in the molecular variability and abundance of fresh DOM (Coch et al., 2019; Hertkorn et al., 2016; Jaffé et al., 2012). In contrast to bulk dissolved organic nitrogen (DON) concentrations and CHOS heteroatom contents, percentages of CHON formulas systematically increased from subsurface (~33%) to 1000 m to reach ~50 % of deep-water abundances (Fig. 6j). Taking this together with the CAR and LPD percentages at sub surface layers, surface derived carbohydrates are presumably used with increasing depth by heterotrophs to build up cell materials where N containing proteins dominate intracellular biochemicals (Aluwihare and Meador, 2008) .

For a detailed examination of the DOM structure in the upper layer (i.e., 0-1000 m) a more extensive data set would be required. However, changes in the percentages of the molecular formulas seem to indicate the gradual transformation of semi-labile DOM at the surface to semi-refractory DOM in the mesopelagic zone (200-1000m) (see Hansell, 2013). This trend was evident in most of the profiles displaying molecular percentages (Fig. 6e-k). ~~For instance, when microbial growth is limited by the availability of inorganic nutrients (nitrate in our data) in oligotrophic (e.g. Arctic Ocean) and ultra-oligotrophic regions (e.g. Mediterranean), semi-labile DOM (e.g. polysaccharides and large proteins) resist rapid degradation and temporarily accumulate in the surface layer (Dittmar and Stubbins, 2014). Since, the bacterial community in the surface waters has limited capacity to degrade semi-labile DOM contrary to mesopelagic bacterial communities, which have the required metabolic mechanisms to metabolize semi-labile fractions (Repeta, 2015 and the references therein). While semi-labile DOM gradually gets decomposed with depth, semi-refractory DOM may accumulate with same principle of changing bacterial communities and molecular reactivities along with the microbial pump of refractory DOM to the deep ocean (Jiao et al., 2010).~~


3.3.3 Hydrothermal plume and the surroundings


The influence of the plume dispersion on DOM composition was investigated in 11 samples from three stations below 2000 m: BG (background), PL (plume) and NP2 (non-plume). The approximate locations of these samples relative to hydrothermal

vents and the possible route of the plume dispersion are depicted in Fig. 7a. PCoA and hierarchical clustering indicate a similar composition in the samples obtained from the same station (Fig. 7b-c). Exceptions are samples PL-3000 and BG-2000 which display higher molecular diversity and number of formulas and are therefore more similar to NP2 samples (Fig. 7b). Nevertheless, samples from different depths but from the same station prominently group into three separate clusters (Fig. 7a-c).


The detailed DOM formulae compositions as depicted by van Krevelen diagrams reveal compositional differences between BG, PL and NP2 samples (Fig. 8). All van Krevelen diagrams are populated by a large number of molecular formulas with H:C ratios in the range of 1.0 to 1.8 and O:C ratio of 0.1 to 0.9. In general, the compositions in BG and PL samples display a similar distribution of formulas (Fig. 8a-i). However, BG samples have more formulae, especially at regions $H:C \geq 2.0$ and $H:O:C \leq 1.0$ compared to samples at PL. Contrary to BG and PL, NP2 samples have higher number of formulas and higher frequency of oxygen containing formulas (i.e., O:C ratio extended until 1.5) and almost no formulas with $H:C$ ratio ≤ 0.5 (Fig. 8j-k).

Considering that the plume dispersion mainly occurs along a lateral transect, we collected samples at the same depths along this lateral transect to be able to compare the differences in the DOM composition (e.g., BG-3250 vs NP2-3250 m and PL-3500 vs NP2-3500; Fig. 8b, 8h, and 8j-k, respectively). Formulas that were unique (i.e., formulae that were not present in samples from the same layer in other stations) or distinct molecular indices (i.e., percentages in different molecular classes like LPD or CAR) were used to evaluate changes in DOM composition. Variations in these molecular properties might be related to distinct biogeochemical settings associated with the plume dispersion, as found with turbidity, methane concentrations, and $\delta^{13}C-CH_4$ contents. For example, low molecular diversity, average relative abundances-intensities and LPD content in PL-3500 and PL-3400 (Fig. 8a6a-e) indicate the effect of hydrothermal intrusion or the molecular composition carried by the plume, in line with previous experimental and environmental data suggesting preferential removal of higher molecular weight and oxygen rich compounds under hydrothermal conditions (Hawkes et al., 2016; Rossel et al., 2017).





490 **Figure 7: (a) Relative positions of the dissolved organic matter (DOM) samples below 2000 m and the hydrothermal plume. (b) Heat-**
map representing the relation between hierarchical clustering of DOM compositions based on Bray Curtis dissimilarity. (c) Principal
coordinate analysis (PCoA) based on Bray-Curtis dissimilarity of DOM composition samples. (e) The heat-map represents the
495 **relation between hierarchical clustering of DOM compositions based on Bray-Curtis dissimilarity. Sample names are coded as station**
name and the approximate depth of the sample for background (BG), upper layer (UL), plume (PL) and non-plume (NP2) stations



500 **Figure 88:** Dissolved organic matter (DOM) formulas on van Krevelen diagram showing O:C and H:C ratios of the formulas at stations (a-d) BG, (e-i) PL and (j-k) NP2. Sample names are coded as station name and sample depth. Orange color is given to PL-3400 where the plume effect is expected to be the highest. Blue and green are used for the formulas in samples that were taken from same depths at different stations. Yellow color is for the samples there were taken from the bottom of the water column at BG and PL.

Formatted: Font: 9 pt, Bold

Formatted: Font: 9 pt, Bold

Formatted: Font: 9 pt, Bold

505

Given that the DOM composition throughout the region is thermally altered, the nature of thermal alteration of organic matter at the hydrothermal systems has previously formulated as (i) reduction of organic compounds in low temperature regions (~60-150°C) to generate products from weaker bonds (e.g. formation of petroleum, aliphatic hydrocarbons), (Simoneit, 1992) (ii) oxidation at higher temperature conditions (e.g. PAHs and alkanones) (Dittmar and Koch, 2006; Simoneit, 1995) and (iii) abiotic synthesis of organic matter by thermocatalytic (Fischer-Tropsch-type) reactions (McCollom et al., 1999; McCollom and Seewald, 2007). Therefore, it is difficult to categorize thermal processes as either a source or a sink mechanism for marine DOM over a range of temperatures from ~60°C to ~150°C due to for instance, formation of petroleum products (Simoneit, 1992), water soluble vitamins and amino acids (Longnecker et al., 2018), elevated concentrations of formate, acetate concentrations (Lang et al., 2010; McDermott et al. 2015), fractions of labile compositions (Hansen et al., 2019), and formation of iron and sulphur bearing organic ligands (Yücel et al., 2011). During high temperature hydrothermal circulation, however, ~94 % of the solid phase extractable DOC is typically removed (Hawkes et al., 2015). Findings at several Mid-Atlantic Ridge hydrothermal systems, and temperature-controlled hydrothermal experiments, showed that DOM is highly unstable and effectively removed during hydrothermal circulation, with a reduction in the abundance of recalcitrant DOM, and a decrease in number of formulas and average molecular mass (Hawkes et al., 2015, 2016; Rossel et al., 2015, 2017).

520

This agrees with our observations of a decreasing numbers of molecular formulae, molecular diversity, average relative abundance-intensity and LPD composition in the samples where plume-related hydrochemistry was manifested by a higher level of turbidity, five times higher methane concentration, and markedly heavier methane carbon isotope compositions. We can therefore assume that the compositional differences between background seawater in BG samples and DOM in the plume layer (PL-3400) may be due to hydrothermal degradation, or some other chemical combination of reactions that result in a molecular signal similar to those reported for hydrothermal fluids. Due to the difficult sampling situation in the high Arctic marine environment, we were unfortunately unable to take samples directly from the buoyant plume of the vent to properly illustrate the distribution and composition of hydrothermal endmember DOM. Our analysis, therefore, is solely based on compositional comparison between samples that were different in their relative position to the vent and its non-buoyant plume. However, we do suggest that a portion of the observed composition in the PL is associated with the hydrothermal fluids, which are efficient at scavenging hydrothermal organic compounds from the vent area, and transporting them from the heated subsurface (Simoneit, 1992). Accordingly, the temperature profile at PL implies that the hydrothermal fluids must have been cooled down to the ambient temperature before reaching the sampling point. Therefore, any thermal modifications in DOM compositions must have occurred either before entering the lithosphere or in areas adjacent to the vents themselves, being merely carried along with the plume-transported particles, dissolved gasses, or water masses and gradually diluted with the surrounding waters until it reached the sampling location.

535

Alternatively, unique formula compositions in NP2 with higher oxygen saturation (i.e., $O:C \geq 1$) or ~2% higher CAR content (see Fig. 5f*g*) could be a product of microbial enrichment attributed to chemosynthetic activity at the southern part of the seamount where the faunal communities, sponges, amphipods and traces of biological activity were more abundant as logged

by video and photographic transects from OFOBS dives (Bünz et al., 2020). Nevertheless, based on our data, a clear and direct link between activity of faunal communities and DOM molecular compositions could not be made. Because of this, we cannot yet suggest any mechanism that can transfer modifications of benthic communities to pelagic DOM compositions.

One other plausible explanation for the differences in molecular composition in PL and NP2 compared to BG could be a change in the current direction. The current direction could have been different in the past compared to what we observed on our cruise. If this were the case, the observed composition at the NP2 station could be an imprint of a previous composition of the hydrothermal vent or some other form of environmental gradient that modified and diversified the DOM composition at NP2, while the PL samples reflect the current composition at the sampling location. The higher abundance of CHOS heteroatom composition in the NP2 samples could be evidence of that. A very recent study in Kairei and Pelagia hydrothermal vents in the Indian Ocean indicate 17% more CHOS molecular formulas in DOM from hot vent fluids along with the elevated hydrogen sulphide concentrations compared to surrounding seawater (Noowong et al., 2021). Similarly, reproduction of sulphur containing compounds are found to occur under experimental hydrothermal conditions at several selected temperatures (Hawkes et al., 2016) and at less acidic conditions (Rossel et al., 2017).

4 Summary and Conclusion

Our investigations of the water column above the Aurora hydrothermal vent field (Gakkel Ridge, Arctic Ocean) show that hydrothermal venting in this region displaces thermally degraded DOM compositions within the hydrothermal plume, along with ^{13}C enriched methane and other vent-derived properties. We show that DOM compositions in the plume layer samples had lower numbers of molecular formulae, molecular diversity, average relative abundanceintensity, and percentage of formulae assigned to the LPD molecular class. However, at a site with higher vent related biological imprinting and lower influence from plume dispersion, DOM compositions had higher numbers of formulae, molecular diversity, and CAR percentages. We characterized the water column hydrophysical and biochemical properties, observing that nutrient distributions and DOM compositions mainly reflect vertical changes in primary production, atmospheric gas exchange, riverine influence and water mass distributions in the upper 1000 m layer, while the water column below 2000 m was quite uniform. DOM composition in sea ice was more labile, with markedly high LPD and low molecular weight compositions than the ocean surface waters, suggesting selective transport during ice formation and new production by ice algae.

DOM compositions in the Arctic Ocean are prone to changes in the upper ocean layers, in parallel to the changes in global climate and decline in sea ice extent. However, deep ocean DOM is mostly refractory, displaying remnants of ancient biological processes. Hydrothermal vents in the deep ocean could therefore be a dynamic source of organic matter and perhaps influence overlying pelagic ecosystems. There are several experimental studies on how DOM is modified during hydrothermal processes under different physical and chemical conditions, however, there is a gap of knowledge on to what extent hydrothermally and chemosynthetically modified DOM spreads along plume layers, altering water column biogeochemistry. To the best of our knowledge, our study is the first attempt to differentiate hydrothermal plume and non-plume DOM compositions in the deep

Arctic Ocean; clearly more studies should address this in different hydrothermal systems and other plume-generating deep-sea environments.

Data and code availability

The data and the codes for generating figures and statistical analyses are available in UiT Open Research Data repository (Sert et al., 2021).

Author contributions

MFS designed the study and performed data acquisition with supervision from AS, FG and HN. EPR and KPH analyzed shipboard methane concentrations and methane carbon isotope compositions, respectively. TK and JJ analyzed dissolved organic matter samples, generated formula lists and wrote mass spectrometry methodology. MFS wrote the manuscript with contributions from all coauthors.

Competing Interests

The authors have the following competing interests: One of the (co-)authors is a member of the editorial board of Biogeosciences.

Acknowledgements

We would like to acknowledge the crew of R/V Kronprins Haakon, chief scientist Stefan Bünz and co-chief scientist Eva Ramirez-Llodra for their tremendous efforts during this challenging expedition. We would like to thank Autun Purser and Ulrich Hoge for OFOBS observations, Colin A. Stedmon for DOC and TDN measurements, and Linda Fondes Lunde for nutrient analyses. This study is supported by European Union's Horizon 2020 Research and Innovation Programme (EU FT-ICR MS project; grant agreement 731077). The FT-ICR MS facility is also supported by Biocenter Finland, Biocenter Kuopio and the European Regional Development Fund (grant A70135). KPH contributed this research through support from the Exo-Ocean Worlds project within the Astrobiology Program at NASA, and through support from the Jet Propulsion Laboratory, California Institute of Technology, under a contract with the National Aeronautics and Space Administration (80NM0018D0004). [PR is funded from European research Council \(ERC\) Synergy Grant \(DEEP PURPLE, grant agreement No 856416\) awarded under the European Union's Horizon 2020518 research and innovation programme.](#) The publication charges for this article have been funded by a grant from the publication fund of UiT The Arctic University of Norway.

Formatted: English (United Kingdom)

Funding

This study is funded by the Research Council of Norway through CAGE (Centre for Gas Hydrate, Environment and Climate) project number 223259. The HACON project is funded by the Research Council of Norway through FRINATEK grant (274330).

600 References

- Aluwihare, L. and Meador, T.: Chemical Composition of Marine Dissolved Organic Nitrogen, in: Nitrogen in the Marine Environment, 95–140, <https://doi.org/10.1016/B978-0-12-372522-6.00003-7>, 2008.
- Arrigo, K. R. and van Dijken, G. L.: Continued increases in Arctic Ocean primary production, *Progress in Oceanography*, 136, 60–70, <https://doi.org/10.1016/j.pocean.2015.05.002>, 2015.
- 605 Baker, E. T., German, C. R., and Elderfield, H.: Hydrothermal Plumes Over Spreading-Center Axes: Global Distributions and Geological Inferences, in: Seafloor Hydrothermal Systems: Physical, Chemical, Biological, and Geological Interactions, American Geophysical Union (AGU), 47–71, <https://doi.org/10.1029/GM091p0047>, 1995.
- Bart, M. C., de Kluijver, A., Hoetjes, S., Absalah, S., Mueller, B., Kenchington, E., Rapp, H. T., and de Goeij, J. M.: Differential processing of dissolved and particulate organic matter by deep-sea sponges and their microbial symbionts, *Sci Rep*, 10, 17515, <https://doi.org/10.1038/s41598-020-74670-0>, 2020.
- 610 Bauch, D., Schlosser, P., and Fairbanks, R. G.: Freshwater balance and the sources of deep and bottom waters in the Arctic Ocean inferred from the distribution of H₂180, *Progress in Oceanography*, 35, 53–80, 1995.
- Bauch, D., Polyak, L., and Ortiz, J. D.: A baseline for the vertical distribution of the stable carbon isotopes of dissolved inorganic carbon ($\delta^{13}\text{CDIC}$) in the Arctic Ocean, *Arktos*, 1, 15, <https://doi.org/10.1007/s41063-015-0001-0>, 2015.
- 615 Baumberger, T., Früh-Green, G. L., Thorseth, I. H., Lilley, M. D., Hamelin, C., Bernasconi, S. M., Okland, I. E., and Pedersen, R. B.: Fluid composition of the sediment-influenced Loki's Castle vent field at the ultra-slow spreading Arctic Mid-Ocean Ridge, *Geochimica et Cosmochimica Acta*, 187, 156–178, <https://doi.org/10.1016/j.gca.2016.05.017>, 2016.
- Beaulieu, S. E., Baker, E. T., and German, C. R.: Where are the undiscovered hydrothermal vents on oceanic spreading ridges?, *Deep Sea Research Part II: Topical Studies in Oceanography*, 121, 202–212, <https://doi.org/10.1016/j.dsr2.2015.05.001>, 2015.
- 620 Belzile, C., Gibson, J. A. E., and Vincent, W. F.: Colored dissolved organic matter and dissolved organic carbon exclusion from lake ice: Implications for irradiance transmission and carbon cycling, *Limnol. Oceanogr.*, 47, 1283–1293, <https://doi.org/10.4319/lo.2002.47.5.1283>, 2002.
- Benner, R., Pakulski, J. D., Mccarthy, M., Hedges, J. I., and Hatcher, P. G.: Bulk Chemical Characteristics of Dissolved Organic Matter in the Ocean, *Science*, 255, 1561–1564, <https://doi.org/10.1126/science.255.5051.1561>, 1992.
- 625 Boetius, A.: The Expedition PS86 of the Research Vessel POLARSTERN to the Arctic Ocean in 2014, 2015.
- Boetius, A., Bach, W., Borowski, C., Diehl, A., German, C. R., Kaul, N. E., Koehler, J., Marcon, Y., Mertens, C., Molari, M., Schlindwein, V. S. N., Tuerke, A., and Wegener, G.: Exploring the Habitability of Ice-covered Waterworlds: The Deep-Sea

Hydrothermal System of the Aurora Mount at Gakkel Ridge, Arctic Ocean (82°54' N, 6°15W, 3900 m), AGU Fall Meeting Abstracts, 2014, B24A-02, 2014.

- 630 Bray, J. R. and Curtis, J. T.: An Ordination of the Upland Forest Communities of Southern Wisconsin, *Ecological Monographs*, 27, 325–349, <https://doi.org/10.2307/1942268>, 1957.

- Bünz, S., Ramirez-Llodra, E., German, C., Ferre, B., Sert, F., Kalenickenko, D., Reeves, E., Hand, K., Dahle, H., Kutti, T., Purser, A., Hilario, A., Ramalho, S., Rapp, H. T., Ribeiro, P., Victorero, L., Hoge, U., Panieri, G., Bowen, A., Jakuba, M., Suman, S., Gomez-Ibanez, D., Judge, C., Curran, M., Nalicki, V., Vagenes, S., Lamar, L., Klesh, A., Dessandier, P. A., Steen, I., Mall, A., Vulcano, F., Meckel, E. M., and Drake, N.: RV Kronprins Håkon (cruise no. 2019708) Longyearbyen – Longyearbyen 19.09. – 16.10.2019, UIT - The Arctic University of Norway, 100 pp., 2020.

- 635 Burd, B. J. and Thomson, R. E.: Hydrothermal venting at endeavour ridge: effect on zooplankton biomass throughout the water column, *Deep Sea Research Part I: Oceanographic Research Papers*, 41, 1407–1423, [https://doi.org/10.1016/0967-0637\(94\)90105-8](https://doi.org/10.1016/0967-0637(94)90105-8), 1994.

- 640 Coch, C., Juhls, B., Lamoureux, S. F., Lafrenière, M. J., Fritz, M., Heim, B., and Lantuit, H.: Comparisons of dissolved organic matter and its optical characteristics in small low and high Arctic catchments, *Biogeosciences*, 16, 4535–4553, <https://doi.org/10.5194/bg-16-4535-2019>, 2019.

Damm, E. and Budús, G.: Fate of vent-derived methane in seawater above the Håkon Mosby mud volcano (Norwegian Sea), *Marine Chemistry*, 82, 1–11, [https://doi.org/10.1016/S0304-4203\(03\)00031-8](https://doi.org/10.1016/S0304-4203(03)00031-8), 2003.

- 645 Damm, E., Kiene, R. P., Schwarz, J., Falck, E., and Dieckmann, G.: Methane cycling in Arctic shelf water and its relationship with phytoplankton biomass and DMSP, *Marine Chemistry*, 109, 45–59, <https://doi.org/10.1016/j.marchem.2007.12.003>, 2008.

- Damm, E., Helmke, E., Thoms, S., Schauer, U., Nöthig, E., Bakker, K., and Kiene, R. P.: Methane production in aerobic oligotrophic surface water in the central Arctic Ocean, *Biogeosciences*, 7, 1099–1108, <https://doi.org/10.5194/bg-7-1099-2010>, 2010.

- 650 DeMets, C., Gordon, R., and Argus, D.: Geological current plate motions, *GEOPHYS J INT*, 181, 1–80, <https://doi.org/10.1111/j.1365-246X.2009.04491.x>, 2010.

Dick, G. J.: The microbiomes of deep-sea hydrothermal vents: distributed globally, shaped locally, *Nat Rev Microbiol*, 17, 271–283, <https://doi.org/10.1038/s41579-019-0160-2>, 2019.

- 655 Dittmar, T. and Koch, B. P.: Thermogenic organic matter dissolved in the abyssal ocean, *Marine Chemistry*, 102, 208–217, <https://doi.org/10.1016/j.marchem.2006.04.003>, 2006.

Dittmar, T., Koch, B., Hertkorn, N., and Kattner, G.: A simple and efficient method for the solid-phase extraction of dissolved organic matter (SPE-DOM) from seawater, *Limnol. Oceanogr. Methods*, 6, 230–235, 2008.

- 660 Edmonds, H. N., Michael, P. J., Baker, E. T., Connelly, D. P., Snow, J. E., Langmuir, C. H., Dick, H. J. B., Mühe, R., German, C. R., and Graham, D. W.: Discovery of abundant hydrothermal venting on the ultraslow-spreading Gakkel ridge in the Arctic Ocean, *Nature*, 421, 252–256, <https://doi.org/10.1038/nature01351>, 2003.

Fahrbach, E., Meincke, J., Østerhus, S., Rohardt, G., Schauer, U., Tverberg, V., and Verduin, J.: Direct measurements of volume transports through Fram Strait, *Polar Research*, 20, 217–224, <https://doi.org/10.1111/j.1751-8369.2001.tb00059.x>, 2001.

- 665 Folders, M. and Rombouts, T.: Sponges Revealed: A Synthesis of Their Overlooked Ecological Functions Within Aquatic Ecosystems, in: YOUARES 9 - The Oceans: Our Research, Our Future: Proceedings of the 2018 conference for YOUng MARine RESEARCHer in Oldenburg, Germany, edited by: Jungblut, S., Liebich, V., and Bode-Dalby, M., Springer International Publishing, Cham, 181–193, https://doi.org/10.1007/978-3-030-20389-4_9, 2020.
- Fouilland, E., Floc'h, E. L., Brennan, D., Bell, E. M., Lordsmith, S. L., McNeill, S., Mitchell, E., Brand, T. D., García-Martín, E. E., and Leakey, R. J.: Assessment of bacterial dependence on marine primary production along a northern latitudinal gradient, *FEMS Microbiol Ecol*, 94, <https://doi.org/10.1093/femsec/fiy150>, 2018.
- Fuchida, S., Mizuno, Y., Masuda, H., Toki, T., and Makita, H.: Concentrations and distributions of amino acids in black and white smoker fluids at temperatures over 200°C, *Organic Geochemistry*, 66, 98–106, <https://doi.org/10.1016/j.orggeochem.2013.11.008>, 2014.
- 675 German, C. R. and Boetius, A.: Hydrothermal Exploration of the Gakkel Ridge, 2014 and 2016, *Goldschmidt Abstracts*, 1, 2017.
- German, C. R. and Seyfried, W. E.: Hydrothermal Processes, in: *Treatise on Geochemistry*, Elsevier, 191–233, <https://doi.org/10.1016/B978-0-08-095975-7.00607-0>, 2014.
- 680 German, C. R., Bowen, A., Coleman, M. L., Honig, D. L., Huber, J. A., Jakuba, M. V., Kinsey, J. C., Kurz, M. D., Leroy, S., McDermott, J. M., de Lépinay, B. M., Nakamura, K., Seewald, J. S., Smith, J. L., Sylva, S. P., Van Dover, C. L., Whitcomb, L. L., and Yoerger, D. R.: Diverse styles of submarine venting on the ultraslow spreading Mid-Cayman Rise, *Proc Natl Acad Sci USA*, 107, 14020–14025, <https://doi.org/10.1073/pnas.1009205107>, 2010.
- 685 Graves, C. A., Steinle, L., Rehder, G., Niemann, H., Connelly, D. P., Lowry, D., Fisher, R. E., Stott, A. W., Sahling, H., and James, R. H.: Fluxes and fate of dissolved methane released at the seafloor at the landward limit of the gas hydrate stability zone offshore western Svalbard: Dissolved methane off western Svalbard, *Journal of Geophysical Research: Oceans*, 120, 6185–6201, <https://doi.org/10.1002/2015JC011084>, 2015.
- Grossart, H.-P., Frindte, K., Dziallas, C., Eckert, W., and Tang, K. W.: Microbial methane production in oxygenated water column of an oligotrophic lake, *PNAS*, 108, 19657–19661, <https://doi.org/10.1073/pnas.1110716108>, 2011.
- 690 Grozeva, N. G., Klein, F., Seewald, J. S., and Sylva, S. P.: Chemical and isotopic analyses of hydrocarbon-bearing fluid inclusions in olivine-rich rocks, *Phil. Trans. R. Soc. A.*, 378, 20180431, <https://doi.org/10.1098/rsta.2018.0431>, 2020.
- Haberstroh, P. R. and Karl, D. M.: Dissolved free amino acids in hydrothermal vent habitats of the Guaymas Basin, *Geochimica et Cosmochimica Acta*, 53, 2937–2945, [https://doi.org/10.1016/0016-7037\(89\)90170-1](https://doi.org/10.1016/0016-7037(89)90170-1), 1989.
- Hannington, M., Jamieson, J., Monecke, T., Petersen, S., and Beaulieu, S.: The abundance of seafloor massive sulfide deposits, *Geology*, 39, 1155–1158, <https://doi.org/10.1130/G32468.1>, 2011.
- 695 Hansell, D. A.: Recalcitrant Dissolved Organic Carbon Fractions, *Annu. Rev. Mar. Sci.*, 5, 421–445, <https://doi.org/10.1146/annurev-marine-120710-100757>, 2013.
- Hansen, C. T., Niggemann, J., Giebel, H.-A., Simon, M., Bach, W., and Dittmar, T.: Biodegradability of hydrothermally altered deep-sea dissolved organic matter, *Marine Chemistry*, 103706, <https://doi.org/10.1016/j.marchem.2019.103706>, 2019.
- 700 Hawkes, J. A., Rossel, P. E., Stubbins, A., Butterfield, D., Connelly, D. P., Achterberg, E. P., Koschinsky, A., Chavagnac, V., Hansen, C. T., Bach, W., and Dittmar, T.: Efficient removal of recalcitrant deep-ocean dissolved organic matter during hydrothermal circulation, *Nature Geosci*, 8, 856–860, <https://doi.org/10.1038/ngeo2543>, 2015.

- Hawkes, J. A., Hansen, C. T., Goldammer, T., Bach, W., and Dittmar, T.: Molecular alteration of marine dissolved organic matter under experimental hydrothermal conditions, *Geochimica et Cosmochimica Acta*, 175, 68–85, <https://doi.org/10.1016/j.gca.2015.11.025>, 2016.
- 705 Hedges, J. I.: Global biogeochemical cycles: progress and problems, *Marine Chemistry*, 39, 67–93, [https://doi.org/10.1016/0304-4203\(92\)90096-S](https://doi.org/10.1016/0304-4203(92)90096-S), 1992.
- Hertkorn, N., Harir, M., Cawley, K. M., Schmitt-Kopplin, P., and Jaffé, R.: Molecular characterization of dissolved organic matter from subtropical wetlands: a comparative study through the analysis of optical properties, NMR and FTICR/MS, *Biogeosciences*, 13, 2257–2277, <https://doi.org/10.5194/bg-13-2257-2016>, 2016.
- 710 Hestetun, J. T., Dahle, H., Jørgensen, S. L., Olsen, B. R., and Rapp, H. T.: The Microbiome and Occurrence of Methanotrophy in Carnivorous Sponges, *Front. Microbiol.*, 7, <https://doi.org/10.3389/fmicb.2016.01781>, 2016.
- Hill, V. J. and Zimmerman, R. C.: Characteristics of colored dissolved organic material in first year landfast sea ice and the underlying water column in the Canadian Arctic in the early spring, *Marine Chemistry*, 180, 1–13, <https://doi.org/10.1016/j.marchem.2016.01.007>, 2016.
- 715 Hockaday, W. C., Purcell, J. M., Marshall, A. G., Baldock, J. A., and Hatcher, P. G.: Electrospray and photoionization mass spectrometry for the characterization of organic matter in natural waters: a qualitative assessment, *Limnol. Oceanogr. Methods*, 7, 81–95, 2009.
- Hodgkins, S. B., Tfaily, M. M., Podgorski, D. C., McCalley, C. K., Saleska, S. R., Crill, P. M., Rich, V. I., Chanton, J. P., and Cooper, W. T.: Elemental composition and optical properties reveal changes in dissolved organic matter along a permafrost thaw chronosequence in a subarctic peatland, *Geochimica et Cosmochimica Acta*, 187, 123–140, <https://doi.org/10.1016/j.gca.2016.05.015>, 2016.
- 720 Horiuchi, T., Takano, Y., Ishibashi, J., Marumo, K., Urabe, T., and Kobayashi, K.: Amino acids in water samples from deep sea hydrothermal vents at Suiyo Seamount, Izu-Bonin Arc, Pacific Ocean, *Organic Geochemistry*, 35, 1121–1128, <https://doi.org/10.1016/j.orggeochem.2004.06.006>, 2004.
- 725 Jaffé, R., Yamashita, Y., Maie, N., Cooper, W. T., Dittmar, T., Dodds, W. K., Jones, J. B., Myoshi, T., Ortiz-Zayas, J. R., Podgorski, D. C., and Watanabe, A.: Dissolved Organic Matter in Headwater Streams: Compositional Variability across Climatic Regions of North America, *Geochimica et Cosmochimica Acta*, 94, 95–108, <https://doi.org/10.1016/j.gca.2012.06.031>, 2012.
- 730 Jakobsson, M., Macnab, R., Mayer, L., Anderson, R., Edwards, M., Hatzky, J., Schenke, H. W., and Johnson, P.: An improved bathymetric portrayal of the Arctic Ocean: Implications for ocean modeling and geological, geophysical and oceanographic analyses, *Geophys. Res. Lett.*, 35, <https://doi.org/10.1029/2008GL033520>, 2008.
- Jørgensen, L., Stedmon, C. A., Kaartokallio, H., Middelboe, M., and Thomas, D. N.: Changes in the composition and bioavailability of dissolved organic matter during sea ice formation, *Limnol. Oceanogr.*, 60, 817–830, 2015.
- 735 Karl, D., Beversdorf, L., Orkman, K., Church, M., Martinez, A., and Delong, E.: Aerobic production of methane in the sea, *Nature Geoscience*, 1, <https://doi.org/10.1038/ngeo234>, 2008.
- Keeling, C. D.: The concentration and isotopic abundances of carbon dioxide in rural and marine air, *Geochimica et Cosmochimica Acta*, 24, 277–298, [https://doi.org/10.1016/0016-7037\(61\)90023-0](https://doi.org/10.1016/0016-7037(61)90023-0), 1961.

- Keir, R. S., Sültenfuß, J., Rhein, M., Petrick, G., and Greinert, J.: Separation of ^3He and CH_4 signals on the Mid-Atlantic Ridge at 5°N and 51°N , *Geochimica et Cosmochimica Acta*, 70, 5766–5778, <https://doi.org/10.1016/j.gca.2006.06.005>, 2006.
- 740 Keir, R. S., Schmale, O., Seifert, R., and Sültenfuß, J.: Isotope fractionation and mixing in methane plumes from the Logatchev hydrothermal field, *Geochem. Geophys. Geosyst.*, 10, <https://doi.org/10.1029/2009GC002403>, 2009.
- Kim, S., Kramer, R. W., and Hatcher, P. G.: Graphical Method for Analysis of Ultrahigh-Resolution Broadband Mass Spectra of Natural Organic Matter, the Van Krevelen Diagram, *Anal. Chem.*, 75, 5336–5344, <https://doi.org/10.1021/ac034415p>, 2003.
- 745 Kudo, K., Yamada, K., Toyoda, S., Yoshida, N., Sasano, D., Kosugi, N., Ishii, M., Yoshikawa, H., Murata, A., Uchida, H., and Nishino, S.: Spatial distribution of dissolved methane and its source in the western Arctic Ocean, *J Oceanogr*, 74, 305–317, <https://doi.org/10.1007/s10872-017-0460-y>, 2018.
- Kujawinski, E. B., Longnecker, K., Blough, N. V., Vecchio, R. D., Finlay, L., Kitner, J. B., and Giovannoni, S. J.: Identification of possible source markers in marine dissolved organic matter using ultrahigh resolution mass spectrometry, *Geochimica et Cosmochimica Acta*, 73, 4384–4399, <https://doi.org/10.1016/j.gca.2009.04.033>, 2009.
- 750 Lang, S. Q., Butterfield, D. A., Lilley, M. D., Paul Johnson, H., and Hedges, J. I.: Dissolved organic carbon in ridge-axis and ridge-flank hydrothermal systems, *Geochimica et Cosmochimica Acta*, 70, 3830–3842, <https://doi.org/10.1016/j.gca.2006.04.031>, 2006.
- 755 Lang, S. Q., Butterfield, D. A., Schulte, M., Kelley, D. S., and Lilley, M. D.: Elevated concentrations of formate, acetate and dissolved organic carbon found at the Lost City hydrothermal field, *Geochimica et Cosmochimica Acta*, 74, 941–952, <https://doi.org/10.1016/j.gca.2009.10.045>, 2010.
- Levin, L. A., Baco, A. R., Bowden, D. A., Colaco, A., Cordes, E. E., Cunha, M. R., Demopoulos, A. W. J., Gobin, J., Grupe, B. M., Le, J., Metaxas, A., Netburn, A. N., Rouse, G. W., Thurber, A. R., Tunnicliffe, V., Van Dover, C. L., Vanreusel, A., and Watling, L.: Hydrothermal Vents and Methane Seeps: Rethinking the Sphere of Influence, *Front. Mar. Sci.*, 0, <https://doi.org/10.3389/fmars.2016.00072>, 2016.
- 760 Li, G., Xie, H., Song, G., and Gosselin, M.: Production of Chromophoric Dissolved Organic Matter (CDOM) in Laboratory Cultures of Arctic Sea Ice Algae, *Water*, 11, 926, <https://doi.org/10.3390/w11050926>, 2019.
- 765 Liu, S., He, Z., Tang, Z., Liu, L., Hou, J., Li, T., Zhang, Y., Shi, Q., Giesy, J. P., and Wu, F.: Linking the molecular composition of autochthonous dissolved organic matter to source identification for freshwater lake ecosystems by combination of optical spectroscopy and FT-ICR-MS analysis, *Science of The Total Environment*, 703, 134764, <https://doi.org/10.1016/j.scitotenv.2019.134764>, 2020.
- Longnecker, K.: Dissolved organic matter in newly formed sea ice and surface seawater, *Geochimica et Cosmochimica Acta*, 171, 39–49, <https://doi.org/10.1016/j.gca.2015.08.014>, 2015.
- 770 Longnecker, K., Sievert, S. M., Sylva, S. P., Seewald, J. S., and Kujawinski, E. B.: Dissolved organic carbon compounds in deep-sea hydrothermal vent fluids from the East Pacific Rise at $9^\circ 50'\text{N}$, *Organic Geochemistry*, 125, 41–49, <https://doi.org/10.1016/j.orggeochem.2018.08.004>, 2018.
- Lorenson, T. D., Greinert, J., and Coffin, R. B.: Dissolved methane in the Beaufort Sea and the Arctic Ocean, 1992–2009; sources and atmospheric flux, *Limnol. Oceanogr.*, 61, S300–S323, <https://doi.org/10.1002/lno.10457>, 2016.
- Lupton, J. E. and Craig, H.: A Major Helium-3 Source at 15°S on the East Pacific Rise, *Science*, 214, 13–18, <https://doi.org/10.1126/science.214.4516.13>, 1981.

- 775 Marcon, Y., Purser, A., Albers, E., Türke, A., German, C., Hand, K., Schlindwein, V., Dorschel, B., Boetius, A., and Bach, W.: Geological settings of hydrothermal vents at 6°15'W and 55°30'E on the Gakkel Ridge, Arctic Ocean, *Goldschmidt Abstracts*, 1, 2017.
- Marnela, M., Rudels, B., Olsson, K. A., Anderson, L. G., Jeansson, E., Torres, D. J., Messias, M.-J., Swift, J. H., and Watson, A. J.: Transports of Nordic Seas water masses and excess SF₆ through Fram Strait to the Arctic Ocean, *Progress in Oceanography*, 78, 1–11, <https://doi.org/10.1016/j.pocean.2007.06.004>, 2008.
- 780 McCarthy, M. D., Beaupré, S. R., Walker, B. D., Voparil, I., Guilderson, T. P., and Druffel, E. R. M.: Chemosynthetic origin of 14 C-depleted dissolved organic matter in a ridge-flank hydrothermal system, *Nat. Geosci.*, 4, 32–36, <https://doi.org/10.1038/ngeo1015>, 2011.
- McCollom, T. M. and Seewald, J. S.: Abiotic Synthesis of Organic Compounds in Deep-Sea Hydrothermal Environments, *Chem. Rev.*, 107, 382–401, <https://doi.org/10.1021/cr0503660>, 2007.
- 785 McCollom, T. M., Ritter, G., and Simoneit, B. R. T.: Lipid Synthesis Under Hydrothermal Conditions by Fischer-Tropsch-Type Reactions, *Origins Life Evol Biosphere*, 29, 153–166, 1999.
- McCollom, T. M., Seewald, J. S., and German, C. R.: Investigation of extractable organic compounds in deep-sea hydrothermal vent fluids along the Mid-Atlantic Ridge, *Geochimica et Cosmochimica Acta*, 156, 122–144, <https://doi.org/10.1016/j.gca.2015.02.022>, 2015.
- 790 McDermott, J. M., Albers, E., Bach, W., Diehl, A., German, C. R., Hand, K., Koehler, J., Walter, M., Wegener, G., and Boetius, A.: Geochemistry, physics, and dispersion of a Gakkel Ridge hydrothermal plume, 87°N, 55°30'E, *Goldschmidt Abstracts*, 1, 2017.
- Michael, P. J., Langmuir, C. H., Dick, H. J. B., Snow, J. E., Goldstein, S. L., Graham, D. W., Lehnert, K., Kurras, G., Jokat, W., Mühe, R., and Edmonds, H. N.: Magmatic and amagmatic seafloor generation at the ultraslow-spreading Gakkel ridge, Arctic Ocean, *Nature*, 423, 956–961, <https://doi.org/10.1038/nature01704>, 2003.
- Mopper, K., Stubbins, A., Ritchie, J. D., Bialk, H. M., and Hatcher, P. G.: Advanced Instrumental Approaches for Characterization of Marine Dissolved Organic Matter: Extraction Techniques, Mass Spectrometry, and Nuclear Magnetic Resonance Spectroscopy, *Chem. Rev.*, 107, 419–442, <https://doi.org/10.1021/cr050359b>, 2007.
- 800 Nakamura, K. and Takai, K.: Theoretical constraints of physical and chemical properties of hydrothermal fluids on variations in chemolithotrophic microbial communities in seafloor hydrothermal systems, *Progress in Earth and Planetary Science*, 1, 5, <https://doi.org/10.1186/2197-4284-1-5>, 2014.
- Noowong, A., Gomez-Saez, G. V., Hansen, C. T., Schwarz-Schampera, U., Koschinsky, A., and Dittmar, T.: Imprint of Kairei and Pelagia deep-sea hydrothermal systems (Indian Ocean) on marine dissolved organic matter, *Organic Geochemistry*, 152, 104141, <https://doi.org/10.1016/j.orggeochem.2020.104141>, 2021.
- 805 Ohno, T., Sleighter, R. L., and Hatcher, P. G.: Comparative study of organic matter chemical characterization using negative and positive mode electrospray ionization ultrahigh-resolution mass spectrometry, *Anal Bioanal Chem*, 408, 2497–2504, <https://doi.org/10.1007/s00216-016-9346-x>, 2016.
- Oksanen, J.: *Vegan: ecological diversity*, 2019.

- 810 Ortmann, A. and Suttle, C.: High abundance of viruses in a deep-sea hydrothermal vent system indicates viral mediated microbial mortality, *Deep Sea Research Part I: Oceanographic Research Papers*, 52, 1515–1527, <https://doi.org/10.1016/j.dsr.2005.04.002>, 2005.
- Osterholz, H., Kirchman, D. L., Niggemann, J., and Dittmar, T.: Environmental Drivers of Dissolved Organic Matter Molecular Composition in the Delaware Estuary, *Front. Earth Sci.*, 4, <https://doi.org/10.3389/feart.2016.00095>, 2016.
- 815 Östlund, H. G. and Hut, G.: Arctic Ocean water mass balance from isotope data, *J. Geophys. Res.*, 89, 6373, <https://doi.org/10.1029/JC089iC04p06373>, 1984.
- Paradis, E. and Schliep, K.: ape 5.0: an environment for modern phylogenetics and evolutionary analyses in R, *Bioinformatics*, 35, 526–528, <https://doi.org/10.1093/bioinformatics/bty633>, 2019.
- Pataki, D. E., Ehleringer, J. R., Flanagan, L. B., Yakir, D., Bowling, D. R., Still, C. J., Buchmann, N., Kaplan, J. O., and Berry, J. A.: The application and interpretation of Keeling plots in terrestrial carbon cycle research, *Global Biogeochem. Cycles*, 17, <https://doi.org/10.1029/2001GB001850>, 2003.
- Pedersen, R. B., Rapp, H. T., Thorseth, I. H., Lilley, M. D., Barriga, F. J. A. S., Baumberger, T., Flesland, K., Fonseca, R., Früh-Green, G. L., and Jorgensen, S. L.: Discovery of a black smoker vent field and vent fauna at the Arctic Mid-Ocean Ridge, *Nat Commun*, 1, 126, <https://doi.org/10.1038/ncomms1124>, 2010.
- 825 Pohlman, J. W., Bauer, J. E., Waite, W. F., Osburn, C. L., and Chapman, N. R.: Methane hydrate-bearing seeps as a source of aged dissolved organic carbon to the oceans, *Nat. Geosci.*, 4, 37–41, <https://doi.org/10.1038/ngeo1016>, 2011.
- Purser, A., Marcon, Y., Dreutter, S., Hoge, U., Sablotny, B., Hehemann, L., Lemburg, J., Dorschel, B., Biebow, H., and Boetius, A.: Ocean Floor Observation and Bathymetry System (OFOBS): A New Towed Camera/Sonar System for Deep-Sea Habitat Surveys, *IEEE Journal of Oceanic Engineering*, 44, 87–99, <https://doi.org/10.1109/JOE.2018.2794095>, 2019.
- 830 Qian, J. and Mopper, K.: Automated High-Performance, High-Temperature Combustion Total Organic Carbon Analyzer, *Anal. Chem.*, 68, 3090–3097, <https://doi.org/10.1021/ac960370z>, 1996.
- R Core Team: R: A Language and Environment for Statistical Computing, R Foundation for Statistical Computing, Vienna, Austria, 2018.
- Ramirez-Llodra, E., Brandt, A., Danovaro, R., Mol, B. D., Escobar, E., German, C. R., Levin, L. A., Martinez Arbizu, P., Menot, L., Buhl-Mortensen, P., Narayanaswamy, B. E., Smith, C. R., Tittensor, D. P., Tyler, P. A., Vanreusel, A., and Vecchione, M.: Deep, diverse and definitely different: unique attributes of the world’s largest ecosystem, *Biogeosciences*, 7, 2851–2899, <https://doi.org/10.5194/bg-7-2851-2010>, 2010.
- Redfield, A. C.: The biological control of chemical factors in the environment, *American Scientist*, 46, 230A–221, 1958.
- Reeburgh, W. S.: Oceanic Methane Biogeochemistry, *Chem. Rev.*, 107, 486–513, <https://doi.org/10.1021/cr050362v>, 2007.
- 840 Reeves, E. P. and Fiebig, J.: Abiotic Synthesis of Methane and Organic Compounds in Earth’s Lithosphere, *Elements*, 16, 25–31, <https://doi.org/10.2138/gselements.16.1.25>, 2020.
- Retelletti Brogi, S., Ha, S.-Y., Kim, K., Derrien, M., Lee, Y. K., and Hur, J.: Optical and molecular characterization of dissolved organic matter (DOM) in the Arctic ice core and the underlying seawater (Cambridge Bay, Canada): Implication for increased autochthonous DOM during ice melting, *Science of The Total Environment*, 627, 802–811, <https://doi.org/10.1016/j.scitotenv.2018.01.251>, 2018.
- 845

Formatted: Norwegian (Bokmål)

- Rossel, P. E., Stubbins, A., Hach, P. F., and Dittmar, T.: Bioavailability and molecular composition of dissolved organic matter from a diffuse hydrothermal system, *Marine Chemistry*, 177, 257–266, <https://doi.org/10.1016/j.marchem.2015.07.002>, 2015.
- Rossel, P. E., Stubbins, A., Rebling, T., Koschinsky, A., Hawkes, J. A., and Dittmar, T.: Thermally altered marine dissolved organic matter in hydrothermal fluids, *Organic Geochemistry*, 110, 73–86, <https://doi.org/10.1016/j.orggeochem.2017.05.003>, 2017.
- Rudels, B., Wadhams, P., Dowdeswell, J. A., and Schofield, A. N.: The thermohaline circulation of the Arctic Ocean and the Greenland Sea, *Philosophical Transactions of the Royal Society of London. Series A: Physical and Engineering Sciences*, 352, 287–299, <https://doi.org/10.1098/rsta.1995.0071>, 1995.
- Rudels, B., Björk, G., Nilsson, J., Winsor, P., Lake, I., and Nohr, C.: The interaction between waters from the Arctic Ocean and the Nordic Seas north of Fram Strait and along the East Greenland Current: results from the Arctic Ocean-02 Oden expedition, *Journal of Marine Systems*, 55, 1–30, <https://doi.org/10.1016/j.jmarsys.2004.06.008>, 2005.
- Rudnicki, M. D. and Elderfield, H.: A chemical model of the buoyant and neutrally buoyant plume above the TAG vent field, 26 degrees N, Mid-Atlantic Ridge, *Geochimica et Cosmochimica Acta*, 57, 2939–2957, [https://doi.org/10.1016/0016-7037\(93\)90285-5](https://doi.org/10.1016/0016-7037(93)90285-5), 1993.
- Santibáñez, P. A., Michaud, A. B., Vick-Majors, T. J., D’Andrilli, J., Chiuchiolo, A., Hand, K. P., and Priscu, J. C.: Differential Incorporation of Bacteria, Organic Matter, and Inorganic Ions Into Lake Ice During Ice Formation, *Journal of Geophysical Research: Biogeosciences*, 124, 585–600, <https://doi.org/10.1029/2018JG004825>, 2019.
- Sert, M. F., D’Andrilli, J., Gründger, F., Niemann, H., Granskog, M. A., Pavlov, A. K., Ferré, B., and Silyakova, A.: Compositional Differences in Dissolved Organic Matter Between Arctic Cold Seeps Versus Non-Seep Sites at the Svalbard Continental Margin and the Barents Sea, *Front. Earth Sci.*, 8, <https://doi.org/10.3389/feart.2020.552731>, 2020.
- Sert, M. F., Reeves, E. P., Hand, K. P., and Ferré, B.: Replication data for: Compositions of dissolved organic matter in the ice-covered waters above the Aurora hydrothermal vent system, Gakkel Ridge, Arctic Ocean, <https://doi.org/10.18710/QPGDFW>, 2021.
- Simoneit, B. R. T.: Aqueous organic geochemistry at high temperature/high pressure, *Origins Life Evol Biosphere*, 22, 43–65, <https://doi.org/10.1007/BF01808018>, 1992.
- Simoneit, B. R. T.: Evidence for organic synthesis in high temperature aqueous media — Facts and prognosis, *Origins Life Evol Biosphere*, 25, 119–140, <https://doi.org/10.1007/BF01581578>, 1995.
- Simoneit, B. R. T., Lein, A. Yu., Peresykin, V. I., and Osipov, G. A.: Composition and origin of hydrothermal petroleum and associated lipids in the sulfide deposits of the Rainbow field (Mid-Atlantic Ridge at 36°N) 1 Associate editor: G. A. Logan, *Geochimica et Cosmochimica Acta*, 68, 2275–2294, <https://doi.org/10.1016/j.gca.2003.11.025>, 2004.
- Speer, K. G. and Rona, P. A.: A model of an Atlantic and Pacific hydrothermal plume, *Journal of Geophysical Research: Oceans*, 94, 6213–6220, <https://doi.org/10.1029/JC094iC05p06213>, 1989.
- Tao, Y., Rosswog, S., and Brüggem, M.: A simulation modeling approach to hydrothermal plumes and its comparison to analytical models, *Ocean Modelling*, 61, 68–80, <https://doi.org/10.1016/j.ocemod.2012.10.001>, 2013.
- Thingstad, T. F., Hagström, Å., and Rassoulzadegan, F.: Accumulation of degradable DOC in surface waters: Is it caused by a malfunctioning microbial loop?, *Limnol. Oceanogr.*, 42, 398–404, <https://doi.org/10.4319/lo.1997.42.2.0398>, 1997.

- Vanreusel, A., Andersen, A., Boetius, A., Connelly, D., Cunha, M., Decker, C., Heeschen, K., Hilario, A., Kormas, K., Maignien, L., Olu, K., Pachiadaki, M., Ritt, B., Rodrigues, C., Sarrazin, J., Tyler, P., Van Gaever, S., and Vanneste, H.: Biodiversity of Cold Seep Ecosystems Along the European Margins, *Oceanog.*, 22, 110–127, <https://doi.org/10.5670/oceanog.2009.12>, 2009.
- 885 de la Vega, C., Jeffreys, R. M., Tuerena, R., Ganeshram, R., and Mahaffey, C.: Temporal and spatial trends in marine carbon isotopes in the Arctic Ocean and implications for food web studies, *Glob Chang Biol*, 25, 4116–4130, <https://doi.org/10.1111/gcb.14832>, 2019.
- Vergeynst, L., Christensen, J. H., Kjeldsen, K. U., Meire, L., Boone, W., Malmquist, L. M. V., and Rysgaard, S.: In situ biodegradation, photooxidation and dissolution of petroleum compounds in Arctic seawater and sea ice, *Water Research*, 148, 459–468, <https://doi.org/10.1016/j.watres.2018.10.066>, 2019.
- 890 Wang, D. T., Reeves, E. P., McDermott, J. M., Seewald, J. S., and Ono, S.: Clumped isotopologue constraints on the origin of methane at seafloor hot springs, *Geochimica et Cosmochimica Acta*, 223, 141–158, <https://doi.org/10.1016/j.gca.2017.11.030>, 2018.
- 895 Whiticar, M. J.: Carbon and hydrogen isotope systematics of bacterial formation and oxidation of methane, *Chemical Geology*, 161, 291–314, [https://doi.org/10.1016/S0009-2541\(99\)00092-3](https://doi.org/10.1016/S0009-2541(99)00092-3), 1999.
- Winkler, L. W.: Die Bestimmung des im Wasser gelösten Sauerstoffes, *Berichte der deutschen chemischen Gesellschaft*, 21, 2843–2854, <https://doi.org/10.1002/cber.188802102122>, 1888.
- 900 Xu, W., Gao, Q., He, C., Shi, Q., Hou, Z.-Q., and Zhao, H.-Z.: Using ESI FT-ICR MS to Characterize Dissolved Organic Matter in Salt Lakes with Different Salinity, *Environ. Sci. Technol.*, 54, 12929–12937, <https://doi.org/10.1021/acs.est.0c01681>, 2020.
- Yahel, G., Sharp, J. H., Marie, D., Häse, C., and Genin, A.: In situ feeding and element removal in the symbiont-bearing sponge *Theonella swinhoei*: Bulk DOC is the major source for carbon, *Limnol. Oceanogr.*, 48, 141–149, <https://doi.org/10.4319/lo.2003.48.1.0141>, 2003.
- 905 Yücel, M., Gartman, A., Chan, C. S., and Luther, G. W.: Hydrothermal vents as a kinetically stable source of iron-sulphide-bearing nanoparticles to the ocean, *Nature Geosci*, 4, 367–371, <https://doi.org/10.1038/ngeo1148>, 2011.