

Sediment and carbon accumulation in a glacial lake in Chukotka (Arctic Siberia) during the late Pleistocene and Holocene: Combining hydroacoustic profiling and down-core analyses

Stuart A. Vyse^{1,2}, Ulrike Herzschuh^{1,2,4}, Gregor Pfalz^{1,2,3}, Lyudmila A. Pestryakova⁵, Bernhard Diekmann^{1,2,3}, Norbert Nowaczyk⁶, Boris K. Biskaborn¹

¹Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research, Research Unit Potsdam, Telegrafenberg A45, 14471 Potsdam, Germany

²Institute of Environmental Science and Geography, University of Potsdam, Potsdam, Germany

³Institute of Geosciences, University of Potsdam, Potsdam, Germany

⁴Institute of Biochemistry and Biology, University of Potsdam, Potsdam, Germany

⁵Northeastern Federal University of Yakutsk, Yakutsk, Russia

⁶Helmholtz-Centre Potsdam GFZ, Climate Dynamics and Landscape Evolution, Telegrafenberg, 14473 Potsdam, Germany

Correspondence to: **Stuart Andrew Vyse** (stuart.vyse@awi.de); **Boris K. Biskaborn** (boris.biskaborn@awi.de)

Abstract

Lakes act as important sinks for inorganic and organic sediment components. However, investigations of sedimentary carbon budgets within glacial lakes are currently absent from Arctic Siberia. The aim of this paper is to provide the first reconstruction of accumulation rates, sediment and carbon budgets from a lacustrine sediment core from Lake Rauchaugytyn, Chukotka (Arctic Siberia). We combined multiple sediment-biogeochemical and sedimentological parameters from a radiocarbon-dated 6.5 m sediment core with lake basin hydroacoustic data to derive sediment stratigraphy, sediment volumes, and infill budgets. Our results distinguished three principal sediment and carbon accumulation regimes that could be identified across all measured environmental proxies including Early MIS2 (ca. 29–23.4 cal. ka BP), Mid ~~MIS2-to-late-early~~ MIS12 (ca. 23.4–11.5-69 cal. ka BP), and Holocene (ca. 11.569– present). Estimated organic carbon accumulation rates (OCARs) were higher within Holocene sediments (average 3.53 g OC m⁻² a⁻¹) than Pleistocene sediments (average 1.09-08 g OC m⁻² a⁻¹) and are similar to those calculated for boreal lakes from Quebec and Finland and Lake Baikal but significantly lower than Siberian thermokarst lakes and Alberta glacial lakes. Using a bootstrapping approach, we estimated the total organic carbon pool to 0.26 ± 0.02 Mt and a total sediment pool of 25.7 ± 1.71 Mt within a hydroacoustically derived sediment volume of ca. 32990557 m³. The total organic carbon pool is substantially smaller than Alaskan ~~Yedomeyedoma~~, thermokarst lake sediments, and Alberta glacial lakes but shares similarities with Finnish boreal lakes. Temporal variability in sediment and carbon accumulation dynamics at Lake Rauchaugytyn is controlled predominantly by palaeoclimate variation that regulates lake ice-cover dynamics and catchment glacial, fluvial and permafrost processes through time. These processes, in turn, affect catchment and within-lake primary productivity as well as catchment soil development. Spatial differences to other lake systems at a trans-regional scale likely relates to the high-latitude, mountainous location of Lake Rauchaugytyn.

Kommentar [SV1]:

Dear editor and reviewers,

Thank you very much for your helpful comments, suggestions and proposed changes to our manuscript. In this mark-up version, you will find all edits made to the manuscript that integrate all relevant comments.

Once again, we really appreciate the time you have invested in reviewing our manuscript.

With kind regards and on behalf of the authors,

Stuart Vyse

Feldfunktion geändert

Feldfunktion geändert

Kommentar [SV2]: A small typing mistake was witnessed during the calculation of the accumulation rates. Only a couple of values have changed very slightly and have been updated within the manuscript text and relevant figures where necessary.

Kommentar [SV3]: We stick to the British spelling now throughout the manuscript. Where necessary, American English has been corrected to British English following one major comment from reviewer #2.

1. Introduction

Lakes represent key sentinels of environmental change and can respond rapidly to changes in environmental conditions (Adrian et al., 2009). ~~Lakes act as sinks of clastic sediment derived from local catchment weathering processes as well as from atmospheric deposition and as such gradually accumulate sediment mass over time~~Lakes act as sinks of clastic sediment derived from weathering of their catchments and as such gradually accumulate sediment mass over time (Dietze et al., 2014; Hinderer and Einsele, 2001). They also constitute a significant net sink of carbon, as they can accumulate organic and inorganic carbon within their sediments derived from allochthonous (lake external) and autochthonous (lake internal) pathways (Ferland et al., 2012; Dean and Gorham 1998; Kortelainen et al., 2004; Sobek et al., 2014). Recent syntheses suggest that global lakes represent a carbon sink of around 0.2–0.6 Pg C y⁻¹ (Cole et al., 2007; Battin et al., 2009). In turn, inland waters can also act as major sources of the greenhouse gases CO₂ and CH₄ and thereby contribute further to global climate change (Anthony et al., 2014).

Lake sediment cores contain sedimentological and biogeochemical proxies of environmentally driven sedimentation and carbon dynamics (Birks and Birks, 2006; Biskaborn et al., 2019; Smol et al., 2002). When sedimentation rate data is available via dating of sediment cores, estimations of rates of sediment mass and carbon accumulation can be reconstructed (Ferland et al., 2012). Moreover, if the sediment volume stored within a lake basin can be estimated, sediment and carbon pools can be obtained that allows the lake function as a sediment and carbon sink to be assessed (Campbell et al., 2000; Lehman., 1975; Munroe & Brencher., 2019; Pajunen., 2000).

The region of Chukotka (Arctic Siberia) represents an ~~important environmentally sensitive~~ area with limited lacustrine environmental reconstructions (Lozhkin and Anderson, 2013). Though a number of regional records across Arctic Siberia are becoming increasingly prevalent (Biskaborn et al., 2012; Diekmann et al., 2016; Melles et al., 2012; Pfalz et al., 2021; Subetto et al., 2017), studies have often neglected reconstructions of accumulation rates. This is largely due to limited age-controls related to problematic radiocarbon dating of organic poor systems that often lack dateable macrofossil remains and possess low dating resolutions (Lozhkin et al. 2016, Strunk et al., 2020). Current eastern Siberian research has primarily focussed on the reconstruction of Holocene accumulation rates of carbon using sediment cores derived from thermokarst lake systems (Anthony et al., 2014). Such studies have shown that lake systems may operate in transitional modes between carbon sink and source stages depending on the prevailing climatic background conditions (Anthony, et al., 2014). Recent syntheses have ~~characterized~~characterised the sedimentological characteristics and carbon budgets of Bykovsky peninsula lagoons of northern Yakutia, eastern Siberia and carbon inventories within ~~a~~Alas and ~~Yedoma yedoma~~ deposits from Central Yakutia and are interpreted in terms of palaeoenvironmental variability since the Pleistocene (Jenrich et al., ~~in review~~accepted; Windirsch et al., 2020). Such works have also been extended to drained thermokarst lake basins (DTLB) and Pleistocene ~~y~~Yedoma deposits in north-western Alaska (Jongejeans et al., 2018). ~~The reconstruction of accumulation rates in these syntheses has however been avoided due to significant reworking of carbon material within permafrost landscapes (Strunk et al., 2020; Windirsch et al., 2020). The role of Arctic Siberian glacial lakes as sediment and carbon sinks has not yet been accounted for.~~The reconstruction of accumulation rates in these syntheses has however been avoided due to significant reworking of carbon material within permafrost landscapes (Strunk et al., 2020; Windirsch et al., 2020). The role however, of Arctic Siberian glacial lakes as sediment and carbon sinks has not yet been accounted for.

Several, broad regional studies of dated sediment cores have been applied in an attempt to calculate accumulation rates and carbon pools within small glacial lakes in the Uinta mountains, north America (Munroe and Brencher, 2019) as well as in [proglacial and bedrock-catchment](#) lakes in southern Greenland (Anderson et al., 2009) and from large glacial lakes in Alberta, Canada (Campbell et al., 2000). A major drawback to these studies however is the absence or oversimplification of sediment volume estimation which does not account for lateral variations in underlying sediment stratigraphy resulting from sediment focussing and winnowing (Ferland et al., 2012). Campbell et al. (2000) suggested an empirical equation for the estimation of sediment volume by simplifying lake sediments to an oblate semi-spheroid to account for sediment thinning at the lake margins. However, such approaches have been shown to lead to [significant](#) overestimations of sediment volumes and carbon pools up to four times the true value (Ferland et al., 2012; Munroe and Brencher, 2019). Hence the lack of efficient derivation of sediment volumes represents a major disadvantage of current works. Recent studies that include seismic or hydroacoustic based appraisals of sediment volume account for these disadvantages and show a [significant major](#) improvement of pool calculations (Einola et al., 2011; Ferland et al., 2012; Pajunen et al., 2000). Such methods are simple, rapid and give significant additional insight into lake bathymetry and basin-wide sediment stratigraphy that permits the retrieval of sediment thicknesses and volumes (Ferland et al., 2012). Recent hydroacoustic based studies from central and western Siberia have concentrated on determining Holocene and late Pleistocene sediment distributions and volumes within glacial lakes and have proven highly useful for understanding glacial history in these regions (Hafliðason et al., 2019; Lebas et al., 2019). The inclusion of sediment budgets within investigations of carbon pools is important as it allows the future storage capacity of lakes to be estimated (Hinderer and Einsele, 2001).

This paper aims to provide a first estimation of late Pleistocene-Holocene sediment and carbon accumulation within an Arctic glacial lake basin within the remote Chukotka region (Arctic Siberia). This includes the first reconstruction of Pleistocene-Holocene sediment and organic carbon accumulation rates and pools provided for Chukotka derived by integrating proxy analyses from a high-resolution dated sediment core with high-resolution hydroacoustic data for sediment volumes. Our results are interpreted in the context of regional palaeoenvironmental variability during the late Pleistocene and are compared to trans-regional studies from Yakutia (east Siberia), North America and Europe. As such, our research questions are:

1. How has the palaeoenvironmental variability in Chukotka and resulting sedimentological processes since Marine Isotope Stage 2 (MIS2) influenced the sediment and carbon accumulation dynamics within an Arctic glacial lake?
2. How much sediment and carbon is stored within the Lake Rauchuagytygn basin and how is this proportioned between the glacial and interglacial periods?
3. How do accumulation rates compare with other systems at a trans-regional scale?


2. Study area

Lake Rauchuagytgyn (67.7922° N, 168.7312° E) is situated within the glacially eroded ∇ -shaped, Rauchua mountain valley within the north-western Anadyr Mountains of Chukotka (Arctic Siberia) where elevations reach up to 1600 m a.s.l (Fig. 1). The lake lies at an elevation of ca. 625 m a.s.l with a surface area of 6.1 km² and maximum water depth of 36 m. The lake is supplied by fluvial inflows at the lakes southern margin (Rauchua River) and via an alluvial fan at the lakes south-eastern margin. One major outflow drains the lake to the north and several minor streams drain the basin sides. The basin is divided into several sub-regions (southern and south-eastern Inflows, southern sub-basin, northern sub-basin and northern shelf) based on water depth characteristics. The catchment area comprises 214.5 km². The bedrock surrounding the lake and within the catchment is predominantly composed of cretaceous extrusive and intrusive igneous facies-rocks consisting of silicic-intermediate lithologies dominated by Andesite (Zhuravlev and Kazymir, 1999). Catchment evidence for glaciation includes moraine structures to the north of the lake that denote the maximum extent of glaciation (Glushkova, 2011). Several small glacial cirques are found within the catchment (Glushkova, 2011) (Fig. 1a). The

Kommentar [SV4]: Figure 1 has been updated according to reviewer comments/suggestions from reviewers #1 and #2. We have increased the font size within the figure inset of (a) as well as in the map keys and labels. We have included an additional small inset that marks the position of some glacial cirques within the higher catchment and marked the location on the catchment map. In addition, we have removed the old bathymetry previously designated as figure 1c and replaced this with an overview of the hydroacoustic profiles obtained during 2018 according to reviewer suggestions. The figure caption has been updated to account for these changes. We hope that the figure has improved due to these changes.

Figure 1. (a) Overview map of the Lake Rauchuagytgyn catchment plotted with a digital elevation map of the Anadyr mountains. In #inset is shown the situation of Lake Rauchuagytgyn (1) compared to other studied regional lakes: (2) Lake Hirney and (3) Lake El'gygytyn (ESRI 2020). A second inset shows the outline of a glacial cirque associated with a small proglacial lake within the eastern catchment. Also shown is the situation of Lake Rauchuagytgyn (1) compared to other studied regional lakes: (2) Lake Hirney and (3) Lake El'gygytyn (ESRI 2020). (b) Orthophoto map of the lake and surrounding features. Shown are the locations of surface sampling points and the location of sediment core EN18218 (black dot) as well as an alluvial fan at the lakes south-eastern margin (ESRI 2020). Flow directions of major inflow and outflow river systems are shown (blue arrows). (c) Simplified water depth map adapted from Huang et al., 2020. Lake polygon with a hydroacoustic profile paths retrieved during 2018.

Formatiert: Englisch (Großbritannien)



Arctic continental climate of the area is characterised by mean annual temperatures of -11.8 °C and average July and January temperatures of 13 °C and -30 °C, respectively with low annual precipitation of ca. 200 mm (Menne et al., 2012). A surface ice-layer is present on lakes in this area from October to early July and likely reaches a winter maximum thickness of ca. 1.8 m (Nolan et al., 2002). Catchment vegetation is dominated by open herb- and graminoid tundra with no vegetation at higher elevations. Some forest tundra is found at lower elevations and adjacent to river valleys (Shevtsova et al., 2020).

3. Materials and methods:

3.1 Fieldwork

Fieldwork took place at Lake Rauchuagytyn in July 2018. An SES-2000 compact parametric sub-bottom profiler was used to hydroacoustically survey the basin to identify major acoustic units (AUs) and sediment boundaries and to locate an optimal coring location (Innomar technologies GmbH). A low frequency (8–10 kHz) and low pulse interval (2 seconds) was used for high sediment penetration to retrieve sediment architecture. A high frequency (100 kHz) and high pulse interval (253 microseconds) allowed derivation of lake bathymetry. Depth conversion was conducted assuming a constant sound velocity of 1500 m s⁻¹. In total, 25 hydroacoustic profiles containing > 23000 data points were retrieved.

Sediment core EN18218 (ca. 653 cm) (Fig. 1) was collected from the southern sub-basin using an UWITEC piston coring system. The core was retrieved in parallel overlapping ca. 3 m segments, cut into 1 m sections and transported in transparent 60 mm PVC tubes for further processing. Further core penetration and retrieval was prevented by coarse material comprising of pebble-sized clasts at the core base that most likely represents the low ermost deposited sediments within the basin. ~~Further core penetration and retrieval was prevented by sand-pebble material at the core base most likely representing the basal sediment within the basin.~~

Epilimnetic water samples were collected in 2016 (Huang et al., 2020) and in 2018 (this study) from multiple surface locations and were immediately measured for pH, conductivity and oxygen content (WTW Multilab 540, Germany). Water sub-samples were then filtered (0.45 µm) and transported for analyses of dissolved organic carbon (DOC), anions and cations (Table S1).

3.2 Laboratory analyses

3.2.1 Core processing

Sediment cores were split into halves at the Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research (AWI) with one half being immediately subsampled for radiocarbon dating and subsequently at ca. 10 cm resolution for proxy analyses. The other half was cleaned and logged for macroscopic lithological changes that permitted the derivation of three lithological units (LU-I to LU-III) and prepared for non-destructive scanning approaches.

3.2.2 Radiocarbon dating and age-depth modelling

Due to the lack of suitable plant remains and low organic content of the retrieved sediment core, 25 bulk sediment samples (0.5 cm thickness) and one surface sample (0–0.5 cm sediment depth) were dated for radiocarbon using accelerator mass spectrometry (AMS) with the Mini Carbon Dating System (MICA DAS) at AWI

Formatiert: Englisch (Großbritannien)

Bremerhaven, Germany (Table 1). Samples were prepared following the standard MICADAS chemical pretreatment procedure (see Gentz et al., 2017 for details).

The age-depth relationship was established by using the open-source MATLAB software package 'Undatable' (Lougheed and Obrochta, 2019). For ~~modeling-modelling~~ we used 23 bulk sediment samples. Two samples (Lab-ID: AWI - 3001.1.1; AWI - 3002.1.1) were slightly older than their successive dates further down, which suggest reworking in these depths (81.25 cm and 114.75 cm, respectively) and would lead to unrealistically high sedimentation rates when included within age-depth modelling that is not mirrored by sedimentological proxies.

We treated these two dates as outliers and excluded them from the ~~modeling-modelling~~ process. All dates were calibrated to 'calibrated years Before Present' (years before 1950 C.E., 'cal. a BP') using the IntCal20 calibration curve (Reimer et al., 2020). In order to account for an existing age offset, we added the results from the surface sample (785 ± 31 ^{14}C years BP) to the software as reservoir age (Undatable settings: $\text{nsim} = 10^6$, $\text{bootpc} = 70$, $\text{xfactor} = 0.1$). Median sedimentation rates (SR) (mm a^{-1}) were extracted from the final age-depth model. In order to account for age-depth model uncertainty in subsequent calculations, age-depth model derived 1σ and 2σ SR uncertainty ranges were also extracted.

Table 1. Radiocarbon dates from sediment core EN18218. Calibrated median ages and 2 σ confidence intervals were calculated with CALIB 8.2 (Stuiver et al., 2020) using the IntCal20 (Reimer et al., 2020) calibration curve. Calibrated ages are given as calibrated years Before Present (years before 1950 C.E., 'cal yr BP'). The surface sample used for down-core correction of the reservoir effect was not calibrated. Samples in italic and marked with an asterisk (*) were excluded from the age-depth ~~modeling~~modelling.

Lab code	Sample ID	Composite depth (cm)	Radiocarbon age with error (^{14}C years BP)	Calibrated median age (cal. ^{14}C years BP)	Calibrated 2 σ age range (cal. ^{14}C years BP)	Sample type
AWM - 5627.1.1	EN18218-1 Surface 0-0.5 cm	0.25	785 \pm 31	-	-	Bulk, TOC
AWM - 2998.1.1	EN18218-2_0-100_20-20.5	18.75	2787 \pm 33	2887	2783 – 2961	Bulk, TOC
AWM - 2999.1.1	EN18218-2_0-100_36.5-37	35.25	3629 \pm 33	3942	3842 – 4080	Bulk, TOC
AWM - 3000.1.1	EN18218-2_0-100_61-61.5	59.75	3832 \pm 33	4233	4098 – 4404	Bulk, TOC
AWI - 3001.1.1*	<i>EN18218-2_0-100_82.5-83</i>	81.25	4985 \pm 33	5702	5602 – 5882	<i>Bulk, TOC</i>
AWI - 3002.1.1*	<i>EN18218-2_100-200_116-116.5</i>	114.75	5417 \pm 34	6234	6119 – 6295	<i>Bulk, TOC</i>
AWM - 3003.1.1	EN18218-2_100-200_140-140.5	138.75	5074 \pm 34	5816	5739 – 5908	Bulk, TOC
AWM - 3004.1.1	EN18218-2_100-200_164-164.5	162.75	5382 \pm 34	6201	6008 – 6284	Bulk, TOC
AWM - 3005.1.1	EN18218-2_100-200_189-189.5	187.75	5852 \pm 34	6672	6559 – 6746	Bulk, TOC
AWM - 3006.1.1	EN18218-2_200-240_222.5-223	221.75	6472 \pm 35	7370	7310 – 7460	Bulk, TOC
AWM - 3007.1.2	EN18218-3_0-100_15-15.5	248.75	8872 \pm 37	10011	9779 – 10177	Bulk, TOC
AWM - 3008.1.1	EN18218-3_0-100_37-37.5	270.75	9085 \pm 37	10236	10185 – 10362	Bulk, TOC
AWM - 3009.1.1	EN18218-3_0-100_59.5-60	293.25	9516 \pm 38	10827	10604 – 11074	Bulk, TOC
AWM - 3010.1.1	EN18218-3_0-100_83-83.5	316.75	9901 \pm 39	11298	11214 – 11595	Bulk, TOC
AWM - 3011.1.1	EN18218-3_100-200_105-105.5	338.75	10197 \pm 39	11866	11655 – 11994	Bulk, TOC
AWM - 3012.1.1	EN18218-3_100-200_129-129.5	362.75	11687 \pm 30	13550	13481 – 13598	Bulk, TOC
AWM - 3013.1.1	EN18218-3_100-200_150-150.5	383.75	12205 \pm 46	14112	14026 – 14310	Bulk, TOC
AWM - 3014.1.1	EN18218-3_100-200_171-171.5	404.75	13017 \pm 48	15596	15371 – 15760	Bulk, TOC
AWM - 3015.1.1	EN18218-3_200-292_210-210.5	443.75	14330 \pm 52	17427	17197 – 17803	Bulk, TOC
AWM - 3016.1.1	EN18218-3_200-292_239-239.5	474.75	15686 \pm 48	18941	18847 – 19068	Bulk, TOC
AWM -	EN18218-3_200-	503.75	17708 \pm 56	21473	21191 – 21783	Bulk,

3017.1.1	292_270-270.5					TOC
AWM - 3018.1.1	EN18218-4_0- 100_35-35.5	536.25	18000 ± 55	21945	21740 – 22117	Bulk, TOC
AWM - 3019.1.1	EN18218-4_0- 100_64.5-65	565.75	22649 ± 66	27059	26479 – 27226	Bulk, TOC
AWM - 3020.1.1	EN18218-4_0- 100_95-95.5	596.25	21786 ± 204	26077	25725 – 26442	Bulk, TOC
AWM - 3021.1.1	EN18218-4_100- 163_123-123.5	624.25	25689 ± 325	29941	29205 – 30735	Bulk, TOC
AWM - 3022.1.1	EN18218-4_100- 163_145-145.5	646.25	25081 ± 300	29393	28780 – 29978	Bulk, TOC

3.2.3 Sediment-geochemistry, Magnetic Susceptibility and Grain-Size

Semi-quantitative sediment geochemical data were obtained by X-ray fluorescence line-scanning (XRF) of EN18218 halves. XRF was carried out using an Avaatech core-scanner with a Rh X-ray tube at 0.75 mA and 1.5 mA for 10 and 15 seconds, at 10 kV (no-filter) and 30 kV (Pd-thick filter) at the Federal Institute for Geosciences and Natural Resources (BGR), Germany. A scanning resolution of 5 mm was chosen. The main rock-forming (Aluminium (Al), Silicon (Si), Calcium (Ca), Potassium (K), Titanium (Ti), ~~Manganese (Mn), Iron (Fe)~~, Rubidium (Rb), Strontium (Sr), Zircon (Zr)) and ~~redox~~/productivity (~~Bromine (Br)~~) linked elements (~~Manganese (Mn), Iron (Fe), Bromine (Br)~~) were selected for further processing (Fig. S1). Of these single elements we focus primarily on Br, Ca, K and Ti within this study. The element ratio of K/Ti is used as proxy for clay input and chemical vs physical weathering (Arnaud et al., 2012; Cuvén et al., 2010; Kilian et al., 2013; Marshall et al., 2011). Zr/K and Si/Al ~~is are included as used as a grain-size proxies for coarser grain-sizes~~ and Mn/Fe for redox conditions (Baumer et al., 2020; Cuvén et al., 2010; Pišková et al., 2019). Br/Al is ~~utilized~~ utilised alongside Br as a proxy for productivity (Lenz et al., 2020). ~~Ratios of element intensities were log transformed using the additive log ratio (ALR) transformation within the package "compositions" (version 1.40) in R (Aitchison, 1984; van den Boogaart et al., 2020; Weltje and Tjallingii, 2008)~~. Magnetic susceptibility measurements (MS) were carried out at 1 mm intervals using a Bartington MS2E spot reading sensor integrated in a fully automatic split-core logging device, developed at the GFZ Potsdam, Germany (Bartington Instruments Ltd). Core images and commission on illumination (CIE) l^* (lightness) ~~and~~ b^* (blue-yellow) colour data were retrieved using an Avaatech line scan camera. XRF and MS data were smoothed using 5- and 15-point running means respectively.

Kommentar [SV5]: We have updated the order of supplementary figures as to allow for stratigraphic plots of XRF data within the supplement as suggested by reviewer #1. References to supplementary figures are hence updated throughout the text and supplement.

Kommentar [SV6]: We also decided to add in additionally the Si/Al ratio that was mentioned by reviewer #2 in respect to proxies for coarser sediment grain-size. This is not plotted stratigraphically as to prevent reductions in manuscript readability overall, but is now included within the text and multivariate statistics with the results of the revised manuscript. We also include additional references to support our usage of Si/Al within this manuscript version.

Sediment grain-size analysis was performed on 63 core samples and four surface samples following a 3-week hydrogen peroxide (H₂O₂) treatment to remove organic matter. Samples were homogenised using an elution shaker for 24 hours and split into eight subsamples. At least three subsamples were analysed thrice providing overall nine individual measurements using a Malvern Mastersizer 3000 laser diffraction particle analyser (Malvern Panalytical Ltd). The nine measurements were then averaged to produce a grain-size distribution for each sample. Data was processed using „GRADISTA“ 8.0 software (Blott and Pye, 2001). Intervals of 2 mm–63 µm, 63–2 µm and <2 µm, were used to define percentages of sand, silt and clay respectively (Friedman and Sanders, 1978). The ~~F~~olk and ~~W~~ward method was used for mean grain-size calculation (Folk and Ward, 1957).

3.2.4 Dry bulk density, elemental analyses and accumulation rate calculations

Water contents (wt%) and dry bulk densities (DBD in g cm⁻³) were determined using the volumetric approach of Avnimelech et al. (2001) on 63 samples of known volume obtained with a one cm³ cube during subsampling and calculated from weight-loss following freeze-drying (See also Pajunen, 2000). The total sediment mass accumulation rate (MAR) in g cm⁻² a⁻¹ was calculated according to the ocean drilling project (ODP) methodology (<http://www-odp.tamu.edu/>) (Eq. 1).

Feldfunktion geändert

$$\text{MAR} = \text{DBD} \times \text{SR}$$

(1)

Where DBD is dry bulk density and SR is age-depth model derived sedimentation rate (cm a⁻¹). MAR uncertainty ranges were determined by applying Eq. 1 to sedimentation rate 1σ and 2σ uncertainty ranges extracted from the age-depth model.

225 | Subsequently, 65 dried and milled sub-samples were ~~subsequently~~ analysed for total organic carbon (TOC) and
 total inorganic carbon (TIC) using a Vario SoliTOC cube elemental analyser following combustion at 400 °C
 (TOC), and 900 °C (TIC) (Elementar Corp., Germany). Total carbon (TC) was calculated as the total sum of TOC
 and TIC. Total sulphur (TS) was measured using an Eltra Carbon and Sulfur determinator (Eltra GmbH,
 Germany). Device results are given as weight percent (wt%) in relation to sample mass. Organic carbon
 230 accumulation rates (OCARs) were then calculated by combining TOC (wt%) & DBD to produce organic carbon
 content per volume (g OC cm⁻³) and then with SR for each depth (Eq. 2).

$$OCAR = (DBD \times (\frac{TOC}{100})) \times SR \tag{2}$$

OCARs are reported as grams of organic carbon per square ~~meter-metre~~ per year (g OC m² a⁻¹) for comparison
 with published organic carbon accumulation rates. OCAR uncertainty ranges were calculated identically to MAR
 235 uncertainty ranges.

Average SR, MAR and OCARs were determined for each lithological unit (LU) and the composite sediment
 succession.

3.3 Data analyses

3.3.1 Estimation of lake basin sediment volumes

240 Hydroacoustic profiles were processed and major hydroacoustic boundaries delineated using ISE2 2.9.5
 software ~~software~~ (Innomar technologies GmbH). Boundary depths were extracted using the ISE2 “Depth
 extraction” tool and were imported into ArcMap 10.5.1 software. Data gaps between measurement points and
 profiles were interpolated using the “Topo to Raster” tool to produce interpolated depth surfaces to the sediment-
 water interface and to basin-wide boundary surfaces (Supplement Fig. ~~S2S3~~). The interpolated sediment-water
 245 interface depths were subtracted using the raster calculator tool from interpolated boundary depths to estimate
 total basin and acoustic unit sediment thicknesses (T_{sediment}). Sediment volumes (V_{sediment}) were subsequently
 derived using the “Surface Volume” tool following subtraction of the lake water volume (V_{water}) (Table 2).

250 **Table 2.** Overview of general acoustic (AU) and lithological unit (LU) divisions with associated depth and age ranges and age-
 depth model calculated average sedimentation rates (SR). Additionally shown are basin properties calculated using ArcMap
 10.5.1 software including Interpolated water and sediment thicknesses (T_{water}, T_{sediment}) and water and sediment package
 volumes (V_{water}, V_{sediment}).

Acoustic Unit (AU)	Lithological Unit (LU)	Depth range at core location (cm)	Age range (cal. ka BP)	Average Sedimentation rate (SR) (mm a ⁻¹)	Average Interpolated thickness (m) (T _{water} , T _{sediment})	Estimated volume (m ³) (V _{water} , V _{sediment})
Water	Water	-	-	-	14.5	90103775
1	I	0 - 344 346	0 - 11.569	0.36	2.44	14935205

Kommentar [SV7]: Table 2 has been updated according to reviewer #2 suggestions of moving the LUH/LUH boundary slightly lower. We have thus updated depth-ranges and relevant age-ranges.

2	II	LU-II = 342 346 - 560	11.5-69 - 23.4	0.20	2.96	18055352
	III	LU-III = 561 - 653	23.4 - ca. 29	0.17		

3.3.2 Estimation of carbon and sediment pools using bootstrapping

255 In order to estimate sediment (Sed_{pool} , Mt) and carbon pools (TOC_{pool} , Mt) at Lake Rauchuagytgyn, we used an established bootstrapping approach modified from Strauss et al. (2013), Jongejans et al. (2018) and Windirsch et al. (2020) that excluded ice-wedges not present within the lake sediments and was enhanced to include sediment volumes estimated from GIS-based methods (Eq. 3 & 4).

$$Sed_{pool} = \frac{V_{sediment} \times DBD}{10^6} \quad (3)$$

$$TOC_{pool} = \frac{V_{sediment} \times DBD \times \left(\frac{TOC}{100}\right)}{10^6} \quad (4)$$

260

The bootstrapping approach was carried out using the “boot” (version 1.3-25) and “Bootstrapping-permafrost-OC” packages (Jongejans and Strauss, 2020) in the R environment and included 10,000 iterations of random sampling and replacement of values (Davison and Hinkley, 1997). Combined DBD and TOC values were used for carbon pool estimation and simply DBD for the sediment pool. The approach was applied firstly to the entire sediment core and basin sediment volume and then individually for the LU-II/LU-III (AU2) and LU-I (AU1) sediment packages. The mean sediment and carbon pools (in megatons, Mt), densities ($Kg\ m^{-3}$) and associated standard deviations and confidence intervals of all iterations were computed (See Strauss et al., 2013 for details) (Supplement S4 Code).

265

3.3.3 Multivariate statistics

270 In order to reduce data dimensionality and increase interpretability of presented physicochemical sediment stratigraphic and accumulation rate data obtained from core EN18218, principal component analysis (PCA) was carried out following data centering and ~~standardization~~ standardisation by dividing by standard deviation using the „prcomp“ function in R (R Core team, 2013). Probability ellipses were drawn in the resulting PCA biplot for samples included within each lithological unit.

275

4. Results

4.1 Basin hydroacoustic stratigraphy and sediment distributions

280 Hydroacoustic and bathymetric data (Fig. 2) permit the division of the lake basin into four morphological regions:
 northern shelf, northern sub-basin, southern sub-basin and southern inflow (Fig. 3a). The basin is characterised
 by steep east and west margins and a central basin bathymetric high. A pronounced river delta is present at the
 southern lake margin at the major lake inflow (Fig. 1b). An alluvial fan at the lakes south-eastern margin
 represents drainage from a steep, lake-proximal, west-east oriented valley (Fig. 1b). Hydroacoustic data allow ed
 285 effective imaging of much of the sediment infill except within inflow proximal profiles where acoustic blanking
 limited data retrieval (Fig. 2b).

Kommentar [SV8]: Figure 3 has been updated according to reviewer comments/suggestions chiefly from reviewer #2. This includes increasing font sizes to improve readability. We have also added interpolation artifacts in grey so that they are more clearly distinguished. Moreover, we followed reviewer #2 suggestions and opted to keep a consistent thickness interval (now 0.75 m) spacing in figures 3b-d.




Figure 2. Interpreted hydro-acoustic profiles from Lake Rauchaugytyn (Profile 8 and Profile 18). Both profiles pass close to the drilling location of EN18218. Shown are inferred major hydroacoustic boundaries defined that are connected with major lithological changes recorded by EN18218 and interpreted across the basin.

Average sediment thickness of the entire lake basin infill is estimated as ca. 5.33 m with the thickest infill predominantly within the southern sub-basin and generally correlates with greater water depth. Average total sediment thickness equates to a total sediment volume of ca. 32990557 m³ (0.033 km³) (Fig. 3d). Sediment thicknesses decrease towards the basin margins, particularly within the northern-shelf where water depths < 7.5 m show reduced sedimentation and evidence of ponded sediment lenses and erosion (Figs. 2b, 3b-d). Comparison of hydroacoustic with lithological data and boundaries from core EN18218 allow the subdivision of

basin infill into two acoustic units (AU1 & AU2) corresponding with sediments deposited within lithological units I (AU1) and II & III (AU2) (Table 2). The boundary between AU1 and AU2 represents a major lithological boundary.

295 The base of AU2 likely constitutes the base of sediment infill and is marked by progressively weakening reflectors with depth. Both boundaries trace closely the modern-day lake bathymetry (Fig. S2S3). Acoustic reflectors within AU1 demonstrate generally well-stratified, high-amplitude, continuous, horizontal and sub-parallel reflectors bounded at the top by the sediment-water interface. The thickness of AU1 shows a basin-wide average of 2.44 m and is thickest within the southern sub-basin proximal to the fluvial inflow and alluvial fan (6.5–7 m) and thins towards the northern shelf (Fig. 3b). A volume of ca. 14935205 m³ (0.015 km³) was estimated for AU1.

300 Reflectors within the uppermost portion of AU2 show generally lower amplitude, sub-horizontal and sub-parallel reflectors and show some continuity with AU1 reflectors (Fig. 2a, b). The lowermost part of AU2 consists of highly discontinuous reflectors with little structure and rare sub-horizontal reflectors and likely includes coarse sediments consisting of pebble-sized clasts deposited ~~within the lowermost~~ at the bottom of LU-III directly at the core base and below, including sandy-gravelly material. The upwards transition to sub-parallel reflectors likely represents sediments deposited within the majority of LU-III and throughout LU-II. AU2 is more variable in thickness (average ca. 2.96 m) than AU1 but is generally thickest within the southern sub-basin and to a lesser extent within the northern sub-basin and at localities close to the northern shelf/northern sub-basin boundary observed within hydroacoustic profiles (Fig. 3c). AU2 possesses a volume of ca. 18055352 m³ (0.018 km³) with complex internal architecture with ~~elevation and depression-like hummock-like~~ structures and ridges seen within some areas.

310

315




Figure 3. (a) Interpolated high-resolution lake bathymetry with morphological regions denoted by black boxes. (b) Interpolated AU1 thicknesses. Blue and pink colors denote the thickest sediment within the southern sub-basin. (c) Interpolated sediment thicknesses within AU2. Similarly to AU1, the greatest thicknesses are predominantly located in the southern sub-basin but with other locations in the northern-sub basin showing thick sediments too. (d) Whole sediment package interpolated thickness. Thin sediment is found proximal to the basin margins and at the north-western shelf. The sediment distribution generally reflects sediment focussing into the deeper lake at greater water depths. Interpolation artefacts are attributed to data sparsity and difficulties in tracing reflectors in some regions due to acoustic blanking that led to erroneous interpolation, with larger artefacts being hence excluded.

4.2 Core chronology and sedimentation rates

The age-depth model (Fig. 4a) shows maximum modelled sediment ages of ca. 29 cal. ka BP at the core base and thus the core provides an entire record of the Holocene and MIS2 (Elias and Brigham-Grette, 2013). The occurrence of a surface sample that shows an age older than zero, and two samples (Lab ID: AWI-3001.1.1; AWI-3002.1.1) older than the surrounding samples may reflect sediment mixing or redeposition of organic carbon from the catchment through permafrost processes (Abbott & Stafford, 1996; Björck & Wohlfarth 2002). A similar process may explain the scatter of ages within LU-III samples (Fig 4a). Median sedimentation rates (SR) are plotted alongside 1σ and 2σ uncertainty ranges and vary from a maximum of 0.54 mm a⁻¹ to a minimum of 0.12 mm a⁻¹ (Fig. 4b). Sedimentation rates are highest within LU-I (average 0.36 mm a⁻¹, average 1σ uncertainty

Kommentar [SV9]: Updated the boundary of figure 4a between LU-II/LU-I according to suggestions from reviewer #2.

0.25–0.45 mm a⁻¹). Low -intermediate but progressively increasing rates are seen within LU-II (average 0.20 mm a⁻¹, average 1σ uncertainty 0.16–0.23 mm a⁻¹) with pronounced peaks at 510.5 and 371 cm. LU-III demonstrates low rates, but with larger uncertainty (average 0.17 mm a⁻¹, average 1σ uncertainty 0.15–0.27 mm a⁻¹) due to age model scatter. Changes in sedimentation rates generally correspond with major lithological variations. Sedimentation rate reductions occur at the LU-III/LU-II (ca. 560 cm) boundary and increases are observed across the LU-II/LU-I (ca. 344–346 cm) boundary (Fig 4a, b).




Figure 4. (a) High-resolution age-depth model for sediment core EN 18218 created from 23 radiocarbon dates with sigma uncertainty range. (b) Sedimentation and mass accumulation rates (dark red lines) and 1 and 2σ uncertainty ranges (dark and light grey ribbons).

335

340 **4.3 Core sedimentological and biogeochemical characteristics**

LU-III (Table 2) represents the low ermost unit of core EN18218 ~~comprising sand-gravel and comprises very poorly sorted, coarse material consisting of pebble-sized clasts -directly~~ at the core base of LU-III (ca. 653 cm) that blocked further core retrieval. Coarse basal sediment is then succeeded by ~~overlying, yellow-grey, light-coloured (low b*), and layered silty-clay sediments from ca. 653 cm~~ to 560 cm. Sediments within LU-III show low water contents (average 39.5 %), high DBD (average 1.22 g cm⁻³), very high magnetic susceptibility values (average 681 [10⁻⁶ SI]) and low mean grain-sizes (average 3.9 μm) traced by Zr/K, Si/Al and K/Ti ratios (Figs. 5, 6). The grain-size distribution shows a predominance of silt-clay with minor sand concentrated just above the core base (ca. 650.5 cm). MARs vary between 0.029 g cm⁻² a⁻¹ at the core base and 0.019 g cm⁻² a⁻¹ at the top of LU-III (average 0.022 g cm⁻² a⁻¹, average 1σ uncertainty 0.019–0.033–0.034 g cm⁻² a⁻¹) but are associated with high uncertainty in the positive sigma direction (Fig. 4). TOC contents (wt%) (average 0.27 %), TOC per volume (average 0.0033 g OC cm⁻³), and Br/Al ratio values demonstrate their lowest values at any depth within the core within LU-III, alongside TC (average 0.28 %) and TS (average 0.018 %) (Fig. 6). OCARs are extremely low and stable with low uncertainty (average 0.58 g OC m⁻² a⁻¹, average 1σ uncertainty 0.50–0.92 g OC m⁻² a⁻¹). Equally Mn/Fe shows stable, low values. The LU-III/LU-II boundary is ~~characterized~~ characterized by a decrease in magnetic susceptibility and increase in the K/Ti ratio corresponding to finer grain-sizes at around 560 cm with a grain-size minimum (2.3 μm) at 550 cm. This corroborates a change in sediment lithology to non-layered, very fine-fine silty grey-coloured sediments in LU-II. Sediment grain-size increases around 506 cm and again around 483 cm clearly viewed by an increase in the Zr/K ratio and a peak in mean grain-size marking an increasing contribution of sand and coarser silts to the particle size distribution. Following a unit maximum in grain-size at 431 cm (5.1 μm) a decrease is observed with a value at 391 cm similar to the unit average of 3.82 μm. Grain-size then rises towards the LU-III/LU-I transition at 346 cm. Water content is significantly higher (average 51.8 %) and

Kommentar [SV10]: Figure 5 has now been updated according to reviewer comments. We have corrected the font size issues noted by reviewer #2 as well as reordering the structure of the grain-size diagram to read clay, silt and sand from left to right. We have also moved the LU-II/LU-I boundary to 346 cm as suggested by reviewer #2. Plotted now are also the non-logarithmically transformed element ratios as per reviewer #1 comments to improve comparability to the vast majority of published literature.

Kommentar [SV11]: We have also implemented suggested changes to figure 6. This included removal of one of the TOC curves and also correction of some capitalization errors. Non-log transformed curves are now included. The boundary between LU-II/LU-I has now been shifted to 346 cm as per reviewer #2 suggestions.




Figure 5. Sedimentological and sediment-geochemical proxies obtained from core EN18218 including measured grain-sizes, XRF grain-size proxies, magnetic susceptibility water content and dry bulk density alongside the hydroacoustic profile from the drilling location and lithological unit derivation (Grey-boxes). * in mean grain-size plot refers to three excluded data points (650.5, 341, 321 cm) of very-highcoarser mean grain-size (up to 25 μm).

dry bulk density lower (average 0.87 g cm^{-3}) than in LU-III. MARs are generally lower than in LU-III, associated with lower uncertainty (average $0.017 \text{ g cm}^{-2} \text{ a}^{-1}$, average 1σ uncertainty $0.013\text{--}0.019 \text{ g cm}^{-2} \text{ a}^{-1}$) and follow variations in sedimentation rate with maxima at ca. 510.5 cm and 380–371 cm. TOC & TC values are significantly higher in LU-II (average TOC 0.77 %, average TC 0.78 %) and increase concomitantly to a maximum ($> 2\%$ C) at the LU-II/LU-I transition in line with gradually darkening sediment colour and increasing Br/Al to brown coloured fine-silts at the top of the unit. ~~TOC per volume tracks TOC wt% with a value range of $0.005\text{--}0.008 \text{ g OC cm}^{-3}$ (average $0.0065 \text{ g OC cm}^{-3}$).~~ The average OCAR in LU-II is more than double LU-III (average ~~$1.34\text{--}32 \text{ g OC m}^{-2} \text{ a}^{-1}$~~ , average 1σ uncertainty $1.03\text{--}1.48 \text{ g OC m}^{-2} \text{ a}^{-1}$) and rises from a minimum at 550 cm ($0.72 \text{ g OC m}^{-2} \text{ a}^{-1}$) to a unit maximum at 371 cm ($2.67 \text{ g OC m}^{-2} \text{ a}^{-1}$). TS shows higher values within LU-II (average TS 0.04 %) and increases in the upper part towards the LU-II/LU-I boundary at 346 cm. Nonetheless TS contributes very little to the total core biogeochemistry. The Mn/Fe ratio is more variable within LU-II and gradually increases from the LU-III/LU-II boundary upwards forming a plateau 472–363 cm before decreasing to wards the LU-II/LU-I transition.

LU-I is typified by dark-coloured, fine-silt sediments of coarser grain-size than observed for all other units (average $6.5 \mu\text{m}$) mirrored ~~in~~ by Zr/K, Si/Al and K/Ti ratios. MARs achieve high values particularly between 311 and 251 cm and between 181.5 and 61.5 cm consistent with sedimentation rates (average $0.022 \text{ g cm}^{-2} \text{ a}^{-1}$, 1σ uncertainty $0.015\text{--}0.028 \text{ g cm}^{-2} \text{ a}^{-1}$). TOC & TC contents attain their highest values (average TOC 1.62 %, average TC 1.63 %) with a prominent maximum just after the LU-II/LU-I boundary (ca. 341 cm) and unit minimum (1.2 %) between 261 and 251 cm before increasing to values $> 2\%$ at the modern sediment surface. Br/Al follows TOC and TC and increases slightly throughout the unit towards the sediment surface. ~~TOC per volume values are higher than LU-II and LU-III (average $0.0097 \text{ g OC cm}^{-3}$).~~ TS demonstrates a broad peak between 371 and 271 cm commensurate with extremely low Mn/Fe ratio values. TS values then decrease and remain low with the exception of a maximum at the sediment-water interface. OCARs are significantly higher than in LU-II and LU-III, associated with higher uncertainty and vary between 1.56 to $6.3 \text{ g OC m}^{-2} \text{ a}^{-1}$ (average $3.53 \text{ g OC m}^{-2} \text{ a}^{-1}$)




Figure 6. Biogeochemical proxies measured from samples obtained from core EN18218 (TOC, TC, TS) as well as from XRF (Br/Al, Mn/Fe) and line-scanning (b^*). Average OCAR (Red line) is plotted alongside sigma uncertainty ranges (grey ribbons) synonymous with descriptions in Figure 4.

OC $\text{m}^2 \text{a}^{-1}$, average 1σ uncertainty $2.4650-4.4650 \text{ g OC m}^2 \text{a}^{-1}$) with values increasing across the LU-II/LU-I transition to high values between 311 and 271 cm generally consistent with higher TS and the Mn/Fe minimum. High OCARs are also observed between around 181.5 and 41.5 cm alongside high SRs and MARs in line with increasing TOC contents (maximum $6.3 \text{ g OC m}^2 \text{a}^{-1}$).

390 4.4 Lake carbon and sediment pools

Average sediment thicknesses and volumes are similar between both sediment packages but slightly greater within the LU-II/LU-III (AU2) package (Table 2). The total carbon pool (TOC_{pool}) calculated amounts to $0.26 \pm 0.02 \text{ Mt}$ with a respective average carbon density (TOC-density) of $7.85 \pm 0.60 \text{ kg m}^{-3}$. LU-II & LU-III sediments at Lake Rauchuagygn displayed the lowest estimates with around $0.1 \pm 0.007 \text{ Mt}$ of carbon stored and a TOC-density of $5.65 \pm 0.40 \text{ kg m}^{-3}$ when compared with the more carbon rich LU-I sediments ($\text{TOC}_{\text{pool}} = 0.15 \pm 0.005 \text{ Mt}$, TOC-density = $9.87 \pm 0.34 \text{ kg m}^{-3}$). The total calculated sediment pool for total sediment mass deposited was estimated via bootstrapping as $25.7 \pm 1.71 \text{ Mt}$ with an average sediment density of $780 \pm 52 \text{ kg m}^{-3}$. Individual calculation of sediment pools in LU-I and LU-II/LU-III yielded values of $9.33 \pm 0.32 \text{ Mt}$ and $17.1 \pm 0.89 \text{ Mt}$ respectively.

400 4.5 Results of principal component analysis

The PCA biplot (Fig. 7) of sedimentological, biogeochemical and accumulation rate data shows that PC1 (60.59.72 %) and PC2 (12.40 %) together explain 72.671.7 % of the total variance within the dataset. A clear division and clustering of data is observed that confirms the lithological unit definition and the main data trends described in sect. 4.2 & 4.3. The variance in the positive PC1 direction is controlled chiefly by productivity variables (TOC, TC, Br/Al, Br, TS, b^*), grain-size (Sandsand, Siltsilt, Mean-mean GS, Zr/K, Si/Al) and to a lesser degree sedimentation (SR) and accumulation rates of carbon (OCAR). Negative values of PC1 are explained by reduced organic content, increasing fine-grained contribution of clay, increased DBD, light colouration (l^*) and

Kommentar [SV12]: Updated figure 7 according to reviewer comments. This includes correction for erroneous capitalizations and displaying the lithological units rather than the previous age zones according to reviewer #2 comments. We have also rerun the PCA including Si/Al that was mentioned in a comment from reviewer #2 within the discussion. This has then also been mentioned in the text.




Figure 7. Principal component analysis biplot of sedimentological, biogeochemical and accumulation rate data from core EN18218. Samples are coloured and clustered according to their lithological unit definitions (LU-III, LU-II, LU-I). Clustering according to lithological unit definitions. Also shown are the PC1 scores plotted stratigraphically in relation to depth and modelled age.

MS, Ti, Ca, K and K/Ti. The mass accumulation rate controls chiefly variance within the PC2 direction (Fig. 7, Fig. S3S4).

5. Discussion:

5.1 Accumulation history of Lake Rauchuagytgyn in a palaeoenvironmental context

5.1.1 Glacially influenced sediment and carbon accumulation during the early Marine Isotope Stage 2 (ca. 29–23.4 cal. ka BP)

~~Coarse sediments found directly at the base of core EN18218, that prevented further core retrieval, alongside discontinuous and occasionally chaotic hydroacoustic reflectors within the lowermost portion of AU2, may suggest that the lake bottom substrate represents a diamicton that shares some similarities with basal sediments reported from other Arctic lake sites (cf. Finkenbinder et al., 2014). The presence of pebble-sized clasts, may imply that a lacustrine environment was either totally absent during deposition of the basal sediment, or was dry for long periods of time. Basal structures observed within hydroacoustic data within the lower portion of AU2 may suggest a mixed origin of the coarse sediment that may be related to fluvial or possibly glacio-fluvial activity within the basin that subsequently influenced basin sediment distributions (cf. Lebas et al., 2019; Lebas et al., 2021). A lack of detailed sedimentological data of core base sediment, due to poor retrieval and limited penetration however limits the further comprehensive interpretation of the lake bottom substrate within this study. Although radiocarbon ages show scatter throughout LU-III, a modelled core base age of around 29 cal. ka BP generally overlaps in timing with MIS2 recorded by other regional records and prominent Alaskan lake records (Asikainen et al., 2007; Finkenbinder et al., 2014; Vyse et al., 2020). Moreover, basal ages of several large northern latitude lakes, including dated basal diamicton from Harding Lake, Alaska show generally agreeable age overlap with coarse basal sediment preserved at Lake Rauchuagytgyn (Anderson and Lozhkin, 2015; Brosius et al., 2021; Finkenbinder et al., 2014).~~

~~Overlying layered sediments pervasive throughout LU-III that are characterised by high magnetic susceptibility, dry bulk density and silty-clay composition may suggest the presence of a lake-proximal glacier and the concurrent deposition of minerogenic, predominantly fine-grained, glacially-derived suspended load to the lake basin within a glaciolacustrine setting (cf. Bakke et al., 2005; Gromig et al., 2019; Karlen and Matthews, 1992; Leemann and Niessen, 1994b; Van der Bilt et al., 2015). This may be further supported by an abundance of the minerogenic elements Ca, Ti and K throughout LU-III, and a light sediment colouration and very low organic content that suggest deposition of organic-poor, minerogenic material derived from a catchment glacier (Fig. 7). The fingerprint of minerogenic, glacially-derived suspended load sediment may plausibly be represented by strongly negative PC1 scores from principal component analysis clustered within LU-III of the PCA biplot (Fig. 7) (cf. Van der Bilt et al., 2015). Catchment conditions during deposition of LU-III were likely cold with minimal within-lake and catchment vegetation productivity indicated across all biogeochemical proxies. Extremely low OCARs during the early MIS2 of ca. 0.58 g OC m⁻² a⁻¹ suggest minimal organic carbon input to the basin sediment that was predominantly autochthonous in origin with very limited contribution from catchment vegetation (Kokorowski et al., 2008a). Lacustrine palynological records and vegetation reconstructions of the catchment surrounding Lake Iriney (ca. 50 km south) during MIS2 agrees well with cold conditions with low pollen productivity of catchment flora represented by low pollen concentrations (Andreev et al., 2021). In addition, pollen reconstructed July temperatures were 4–5 °C lower than modern in agreement with unfavourable climate~~

Kommentar [SV13]: We have fully revised section 5.1.1 and have made restructuration's and additions within it to account for comments and suggestions made by reviewer #2. We aimed to make our interpretation of this unit clearer that was not so clear within the first manuscript version. We also provide some extra consideration of the age-scatter within this unit as per reviewer #1 comments. We are however cautious of over-extending the length of the section as to prevent increases in manuscript length and to detract from the carbon story that is a key outcome of our study.

450 conditions for organic carbon productivity (Andreev et al., 2021). Sedimentological records from Lake Iriney during MIS2 also agree with extremely low within-lake productivity represented by very low TC and absent diatoms (Vyse et al., 2020). Preliminary diatom analyses also show a complete absence of diatoms within LU-III in line with cold and possibly turbid water conditions (Biskaborn, unpublished data; Finkenbinder et al., 2014). Organic material arriving at the sediment-water interface was likely preserved by low oxygen conditions. This is recorded by low and stable values of the XRF-derived Mn/Fe ratio that well represents the redox conditions at the sediment-water interface (cf. Baumer et al., 2020; Biskaborn et al., 2019; Fritz et al., 2018; Heinecke et al., 2017; Naeher et al., 2013; Pšková et al., 2019).

455 Finer grain-sizes directly measured by laser diffraction are supported by indirect, XRF-derived grain-size proxies for finer, clay-dominated sediment (K/Ti) and of coarser sediment (Zr/K, Si/Al). Fine-grained composition likely reflects deposition beneath a quasi-permanent lake ice layer that further reduced within-lake productivity through limiting light penetration with insufficient irradiance required by algal communities and hence diminishing organic matter export to the hypolimnion (Andersen et al. 1993; Bouchard et al., 2011; Croudace & Rothwell, 2015; Cuven et al., 2010; McLaren & Bowles, 1985). Thus, during deposition of minerogenic sediment within LU-III the lake may have been of cold-monomict type (Cremer and Wagner, 2003). Sediment input at Lake Rauchaugytgyn was generally low during LU-III deposition as represented by comparatively low SRs and MARs and was likely temporally continuous, with no evidence of unconformities found within sedimentological data. The uncertainty in the positive sigma direction may suggest that rates could be higher when integrating uncertainty associated with age inversions within LU-III. Sedimentation rates reported for Lake El'gygytgyn crater lake (Fig. 1) were low during MIS2 (4.8 cm/ka), compared to higher rates (7.6 cm/ka) during the Holocene (Nowaczyk et al., 2007). A similar finding was made for Harding Lake in Alaska whereby sedimentation rates during the Last Glacial Maximum (30.7 to 15.7 ka) were generally < 0.1 mm/yr but temporally continuous throughout deposition (Finkenbinder et al., 2014). In both of these settings, low sedimentation rates were interpreted as a consequence of thick, quasi-permanent lake surface ice cover and an extremely thin or absent catchment tundra active layer during cold episodes that reduced catchment sediment delivery and wind-driven sediment redistribution (Asikainen et al., 2007; Francke et al., 2013; Melles et al., 2007). Moreover, a similar observation at lake Karakul, Tajikistan of lower sedimentation rates during MIS2 since ca. 29 cal. ka BP (0.15 mm a⁻¹) when compared with Holocene sedimentation rates (0.84 mm a⁻¹) was also explained by reduced sediment input during MIS2 compared to the Holocene alongside reduced organic matter accumulation (Heinecke et al., 2017). The low sedimentation and mass accumulation rates at Lake Rauchaugytgyn during the early MIS2 and the Late Glacial may be supported by the generally equitable thicknesses and sediment volumes between the AU2 (LU-II/LU-III) and AU1 (LU-I) sediment packages derived hydroacoustically (Table 2). Differences are however observed when MIS2 accumulation rates are compared with other Siberian glacial lake sites including Lake Bolshoye Shchuchye in the Ural Mountains (average Last Glacial Maximum (LGM) 1.68 mm a⁻¹). Lake Bolshoye Shchuchye demonstrates a voluminous LGM sediment package (0.325 km³) six times the volume of the Holocene sediment package (0.05 km³) which shows markedly lower sedimentation rates (0.36 mm a⁻¹) and is interpreted to result from intense glacial erosion of the catchment during the LGM (Hafliðason et al., 2019).

485 The difference in glacial sediment accumulation between sites may relate to multiple, interacting factors including the duration of lake surface ice cover, the thickness of the catchment active layer, the existence/absence of deep-water conditions and also differences in glacier dynamics between catchments (Asikainen et al., 2007; Francke et al., 2013). Remote sensing based studies of Chukotkan glacial geomorphology and structures within the Raucha valley have suggested the presence of a passive catchment glacier, ca. 25 km in length that

490 extended along the length of the Rauchua river valley and discharged into the Rauchuaqytgyn basin during
marine isotope stage 2 (Glushkova, 2011). Thus, the catchment glacier may have been predominantly non-
erosive during the early MIS2 and hence not contributed significant sediment volume to the lake basin supporting
the low rates of sediment and mass accumulation discussed here (Gurnell et al., 1996). It must be however noted
that the radiocarbon age scatter in LU-III contributes uncertainty regarding sediment age and hence the SRs and
495 MARs derived for this interval as marked by the wider uncertainty band within the presented age model (Fig. 4b).
Such age scatter may be in part associated with catchment glacial activity that could have led to the reworking of
older catchment organic containing sediments with subsequent deposition within the glaciolacustrine environment
(cf. Lunkka et al., 2001). The usage of bulk sediment radiocarbon dating may additionally exacerbate this
problem and add to radiocarbon uncertainty within LU-III as noted by Oswald et al. 2005 and others regarding
500 dating of organic poor, Last Glacial Maximum lacustrine sediments in Alaska and Siberia. Furthermore, the
presence of layering within LU-III, may also indicate potentially higher sedimentation rates across the unit (cf.
Leemann and Niessen, 1994b). The lack of absolute dating of glacial structures within the Rauchua valley by
Glushkova, 2011, adds additional uncertainty regarding the temporal dynamics of the catchment glacier.

The temporally continuous, fine-grained, minerogenic sediment accumulation preserved within LU-III can likely be
attributed regionally to the MIS2 "Sartan" glaciation recorded within Chukotka (Anderson & Lozhkin 2015;
505 Brigham-Grette et al., 2003; Melles et al., 2012; Vyse et al., 2020) and the Itkillik II glaciation in Alaska, eastern
Beringia (Hamilton and Ashley, 1993). Glaciation during this time was associated with asynchronous but early
glacial advances in Alaskan mountain ranges between 31 and 28 cal. ka BP and advances in western Beringian
glaciers to maximum extents between 27 and 20 cal. ka BP (Brigham-Grette et al., 2003; Elias and Brigham-
Grette, 2013 and references therein). The presence of a valley glacier at Lake Rauchuaqytgyn is in contrast to
510 more westerly, eastern Russia sites from the Verkhoyansk mountains where no glacial activity was witnessed
during MIS2 (Diekmann et al., 2017; Stauch and Lehmkuhl, 2010). This supports previous suggestions that
despite cold conditions, higher moisture derived from the Pacific Ocean permitted the existence of glaciers within
Chukotka (Stauch and Gualtieri, 2008).

515 **5.1.2 Mid MIS2-early MIS1/Mid to late MIS2 accumulation during progressive climate amelioration (ca. 23.4–11.5–69 cal. ka BP)**

Sediment accumulation recorded by LU-II sediments represents a transitional mode between the early MIS2 and
Holocene clearly viewed within PCA results between ca. 23.4 and 11.5–69 cal. ka BP and likely reflects a time-
progressive shift towards the early stages of paraglacial deposition (Fig. 7). Hydroacoustic data show an upward
transition to stable, deep water, lacustrine deposition represented by the increasing prevalence of stronger
520 amplitudes from well-stratified reflectors (cf. Lebas et al., 2021; Hafliðason et al., 2019). A clay maximum and
grain size minimum at ca. 550 cm (ca. 22.7 cal. ka BP) may suggest initial increases in lake water-depth through
glacial melt additions ~~of rock flour rich meltwater~~ that could have led to the observed reduced SRs and MARs at
this time. This ~~is~~ may be supported by the high values of K/Ti (clay contribution) and low values of Zr/K and Si/Al
~~(proxy proxies for coarser grain-sizes) (Kilian et al., 2013; Křibek et al., 2017; Cúven et al., 2010)~~. It is also likely
525 that the persistence of the quasi-permanent ice layer continued to contribute to predominantly fine-grained, low
energy deposition (Asikainen et al., 2007). The strong increase in Zr/K accompanied by increased sand and silt
contribution firstly at ca. 20.5 cal. ka BP consistent with a dramatic peak in SR and MAR values and again ca.
18.5 cal. ka BP likely reflects increasingly rapid glacial retreat (Gromig et al., 2019). The timing of these system
responses is generally consistent with depositional changes recorded at Lake Ilirney in response to deglaciation

Formatiert: Block

530 ca. 20 ka (Vyse et al., 2020) and those proposed for Alaskan glacial retreat interpreted from dated moraine
deposits and lacustrine sediments (20–19 ka BP) (Elias and Brigham-Grette, 2013; Finkenbinder et al., 2015 and
references therein). Progressive deglaciation facilitated the gradual opening of the lake catchment area and
development of fluvial systems that plausibly enhanced sediment load and input to the lake basin through the
initiation of paraglacial processes (Ballantyne, 2002; Dedkov, 2004). Sediment dynamics were likely controlled by
535 rivers controlled by inflowing rivers at the lakes southern margin as indicated by the greatest thicknesses of
sediment within the southern sub-basin deposited within AU2 (Fig. 3c). Thick sediments within AU2 in the
northern sub-basin may indicate additionally enhanced sedimentation derived from the north. The occurrence of
a rhomboidal-shaped deposit similar to preserved lake sediments within other lacustrine and continental shelf
environments at the northern shelf-northern sub-basin transition may represent a mass transport deposit during
540 some part of AU2 deposition (Fig. 2b, Fig. ~~S4S5~~) (Baster et al., 2003). Catchment sediment availability likely also
began to increase during this time due to gradual sub-aerial exposure of subglacial sediments that subsequently
became entrained within paraglacial river systems or transported to the lake via aeolian processes contributing to
progressively increasing sedimentation rates (Tripathi & Rajamani, 1999; Wang et al., 2015).

Organic carbon accumulation rates at the base of LU-II were slightly higher than LU-III signifying increasing
545 organic carbon production and accumulation across the LU-III/LU-II transition. Subsequently increasing rates
alongside up-core sediment darkening and increases in all organic proxies across LU-II points towards
continuous growth in carbon accumulation related to ameliorating conditions during the Mid MIS2-early MIS1
mid-to-late MIS2 onwards. Organic productivity was most likely dominated by autochthonous productivity from within-
lake algae with only minor contributions from catchment vegetation. Though nitrogen was not measured as part
550 of this study, extremely low TOC/TN ratios measured for sediments at the nearby glacial Lake Ilirney during the
late-glacial Late Glacial implied that most organic matter was derived from protein rich algal sources and not from
cellulose-rich and protein-poor vascular land plants from the lake catchment and from macrophytes within the
lake (Baumer et al., 2020; Meyers & Teranes., 2005; Vyse et al., 2020). This is supported by low pollen
concentrations recorded within the Lake Ilirney pollen record and the persistence of *Poaceae* and *Artemisia*
555 during the late-glacial Late Glacial that signify persistently cold conditions and low catchment pollen productivity
(Andreev et al., ~~accepted~~2021). Based on the similarities between the lakes and their catchments, their proximity,
and a lack of macrophytes at both sites, a similar interpretation can be postulated for Lake Rauchagytygn
(Sifeddine et al., 2011). Higher and more variable Mn/Fe ratio values throughout much of LU-II point towards
increased oxygen at the sediment-water interface that could have enhanced degradation of organic matter
560 (Baumer et al., 2020; Fritz et al., 2018). This could be indicative of a gradually increasing summer open water
season with more intensive mixing of the water column through wind and also through enhanced fluvial inflow
during progressive late-glacial Late Glacial climate amelioration (Baumer et al., 2020).

A reduction in accumulation rates ca. 12.6 to ca. 11.695 cal. ka BP, broadly associated with a small-grain-size
findings small reduction in sediment mean grain-size may represent some evidence of reduced accumulation due to
565 cooling associated with the younger Dryas Younger Dryas (YD). The magnitude of this is however not substantial
when all proxies are considered together within principal component analysis (Fig. 7). Moreover, missing
evidence of younger moraines from the catchment implies that there was no glacial re-advance (Glushkova,
2011). These findings may be consistent with recent regional and trans-regional records that suggest a spatially
variable, and possibly more limited younger-dryas Younger Dryas climatic event in east-East & far-east Far East
570 Russia and parts of Eastern Beringia (Anderson & Lozhkin, 2015; Finkenbinder et al., 2015; Kokorowski et al.,
2008b; Lozhkin & Anderson, 2013; Lozhkin et al., 2018). The lack of significant evidence for a strong Younger

Formatiert: Schriftart: Kursiv

Dryas climatic event may also partially reflect seasonal differences in the sensitivity of physical and geochemical proxies to climatic or environmental change, as has been discussed in studies of sediment cores from Burial Lake, Alaska (Finkenbinder et al., 2015).

5.1.3 Holocene controls on basin sediment and carbon dynamics (ca. 11.5–69 cal. ka BP–present)

The Holocene start is marked across all proxies by increasing organic proxy values and sediment grain-size across the LU-II/LU-I boundary (11.69 cal. ka BP) alongside increasing accumulation rate values (Figs 5, 6, 7). High TOC and TS and a dark sediment colour form a prominent peak during the early Holocene between ca. 11.5 and 9.7 cal. ka BP synchronous with the lowest values of Mn/Fe that could reflect increased preservation of organic material under low-oxygen conditions at the sediment-water interface alongside enhanced organic productivity (Fig. 6) (Fritz et al., 2018). Increasing carbon accumulation is supported by rising organic carbon accumulation rates that demonstrate high early Holocene values between ca. 10.2 and 9.2 cal. ka BP. Low K/Ti values during the early Holocene that subsequently persist throughout the mid-late Holocene may reflect increased chemical relative to physical weathering due to the leaching of K from clastic material entering the lake by enhanced catchment pedogenic processes that influenced OCAR values as well as responding to coarser Holocene grain-sizes (Baumer et al., 2020; D'Arcy & Carignan, 1997). The timing of this phase agrees well with the timing of warm and humid environments interpreted during the early Holocene recorded at Lake Emada, central Yakutia (11.5–9.0 cal. ka BP) and coincides with the maximum in local solar insolation (Baumer et al., 2020; Berger & Loutre 1991). Moreover, this finding is consistent with palynological records from Lake Ilirney and from a Holocene short core from Lake Rauchuagytyn which show evidence for a local Holocene thermal maximum ca. 10.6–7 cal. ka BP (Andreev et al., accepted2021).

Increasing early Holocene sediment and mass accumulation rates alongside greater sand contribution may relate to the input of coarser grained fluvial detrital material.~~Increasing early Holocene sediment and mass accumulation rates alongside greater sand contribution to the grain-size distribution may relate to the input of coarser-grained fluvial detrital input~~ from a paraglacial, deglaciated catchment and could result from a water budget change associated with warmer and more humid conditions (Ballantyne, 2002). This is supported by increasing Zr/K and Si/Al ratio values during the early Holocene that support coarser grain sizes that may relate to enhanced input of coarser silts and sands (cf. Pšková et al., 2019). It must however be noted, that higher Si/Al ratio values may also additionally reflect abundant diatoms within Holocene sediments (Procházka et al., 2019).—Annual precipitation (P_{ann}) reconstructed from palynological records at Lake Ilirney suggested an increase from ca. 225 mm during the ~~late-glacial~~Late Glacial to ca. 300 mm in the early Holocene that could have enhanced catchment surface runoff and hence fluvially driven reworking of unstable-metastable glacially deposited sediment by paraglacial processes (Andreev et al., accepted2021; Ballantyne, 2002). Heinecke et al. (2017) proposed a similar scenario to explain increased Holocene sedimentation rates and increased grain-sizes during the end of the ~~late-glacial~~Late Glacial and early Holocene (13.3–6.6 cal. ka BP) at glacial Lake Karakul. Moisture in north-eastern Siberia during interglacials, is derived mostly from warm North Atlantic surface waters and brought via the westerlies to the region (Melles et al., 2007). The waning of the Scandinavian and Barents-Kara ice-sheets likely allowed the passage of moister ~~Westerlies~~westerlies to the region that were blocked during the ~~last-glacial~~maximumLast Glacial Maximum (Elias & Brigham-Grette 2013; Meyer et al., 2002). Increased sedimentation rates also additionally enhanced carbon burial and preservation (Einsele et al., 2001). Enhanced precipitation and fluvial input to Lake Rauchuagytyn during the early Holocene could have led to lake level increase and deep lacustrine conditions at the core location that restricted mixing of the water column, leading to stratification

and may provide a possible explanation for low oxygen conditions (cf. Baumer et al., 2020). An increase in precipitation and rising lake levels was also interpreted for early Holocene sediments preserved at Harding Lake, Alaska (Finkenbinder et al., 2014). Deep lacustrine conditions alongside enhanced early Holocene within-lake and catchment productivity thus likely led to enhanced carbon deposition and preservation at Lake Rauchuagytygn.

A decrease in TOC content (wt%) to a minimum at ca. 8.6 cal. ka BP along with decreasing organic carbon and sediment and mass accumulation rates from 8.9 cal. ka BP to a minimum at ca. 7.7 cal. ka BP alongside ~~low-er~~ grain-size-values finer grain-sizes may reflect some local environmental change and is generally consistent with a Holocene T_{July} minimum at 7.8 cal. ka BP and P_{ann} minimum at 7.6 cal. ka BP recorded within a Lake Rauchuagytygn short core (Andreev et al., ~~accepted~~2021). This may possibly be associated with cooling associated generally with the 8.2 cal. ka BP event recorded at other locations within Siberia (Biskaborn et al., 2016). A degree of similarity between the TOC trend following the early Holocene at both Lake Rauchuagytygn and Lake Eranda may also suggest that bacterial sulphate reduction contributed to the reduced sedimentary organic carbon content during this interval (Fig. 6) (Baumer et al., 2020). A subsequent increase in OCARs from ca. 6.1 cal. ka BP consistent with progressively increasing TOC values towards the modern lake sediment surface represents increased accumulation of organic carbon consistent with increasing mean July temperatures (T_{July}) to a maximum ca. 13.3 °C during the mid-late Holocene by ca. 4.6 cal. ka BP (Andreev et al., ~~accepted~~2021). Synchronous increases in SR and MAR along with coarser grain-sizes may reflect extended periods of ice-free lake surface conditions during summer associated with high T_{July} (Fig. 4,5) (Finkenbinder et al., 2014). Extended open water conditions during this time and throughout the Holocene likely led to longer phases of summer catchment aeolian input, as well as increased wind-driven shoreline erosion and sediment redistribution due to the impact of summer storms (Asikainen et al., 2007; Francke et al., 2013; Vologina et al., 2003). This hypothesis is supported by the highest Mn/Fe ratio values for the core, supportive of enhanced wind and fluvial driven lake mixing processes and weaker stratification (Doran, 1993; Baumer et al., 2020; Regnéll et al., 2019). Francke et al., 2013 argued that coarser grain-sizes and enhanced sedimentation at Lake El'gygytygn during Interglacials was controlled by summer temperatures affecting lake ice cover and hence aeolian input, wind-driven sediment redistribution as well as catchment active layer thicknesses and water availability and was not affected by lake-level variations (~~Francke et al., 2013~~).

Decreasing accumulation rates from ca. 3.6 cal. ka BP to present could reflect reduced late-Holocene sediment input due to neoglacial cooling and is in line with variable T_{July} and low P_{ann} recorded within the Ilirney and Rauchuagytygn short core records and generally lower late Holocene Mn/Fe ratio values (Fig. 6) (Andreev et al., ~~accepted~~2021). This could tentatively be interpreted to represent a reduction in shortening of the summer open water season.

Enhanced fluvial input to the lake basin throughout the Holocene from the Rauchua river valley and from the south-east lake margin are reflected in hydroacoustic and interpolated sediment thickness data (Fig. 3b) from AU1 which show greater thicknesses within the southern sub-basin. A similar distribution has been observed at glacial lakes Bolshoye Shchuchye & Levinson-Lessing (Hafidason et al., 2019; Lebas et al., 2019). The presence of an alluvial fan ca. 600 m in diameter at the south-eastern margin is supportive of intensified Holocene paraglacial processes that resulted in an input source of ~~coarser fluvial detrital material~~ coarser fluvial and alluvial detrital material into the southern sub-basin contributing to the observed increased sedimentation and mass accumulation rates (cf. Doran, 1993; Smith and Jol, 1997). This is further supported by coarse, sand-dominated

655 | surface sediments close to the alluvial fan front (site EN18220) (Fig. 1 & Figs- ~~S5S6~~, ~~S6S7~~). Modern observations at Lake El'gygytgyn have shown that summer fluvial activation associated with snow melt can lead to significant deposition of coarse material of cobble-grade material to the lake basin (Nolan and Brigham-Grette, 2007). Heightened availability of catchment sediment material due to thicker catchment active layer thicknesses during the Holocene may also provide an additional sediment source leading to increased sedimentation. A thick tundra active layer present during the Holocene at Lake El'gygytgyn was suggested to have contributed to greater detrital input to the basin as well as promoting the weathering of clay minerals (Asikainen et al., 2007; Francke et al., 2013). This further supports the low K/Ti ratios during the Holocene that evidence intensified chemical weathering (Regnéll et al., 2019). Acoustic reflectors preserved within AU1 generally reflect well-developed lacustrine deposition under deep-water conditions prevalent across the basin (Fig. 2)(Lebas et al., 2019). Some variability is observed within hydroacoustic data at the northern shelf where sediments are thin and evidence of erosion of sediments are seen (Fig. ~~S4S2~~). This likely represents low deposition due to sediment focussing-transport into the deeper basin and feasibly some Holocene lake-level variability of several meteres permitting localised erosion into shallow areas (Moernaut et al., 2010).

5.2 Comparisons of carbon and sediment dynamics on a trans-regional scale

5.2.1 Organic Carbon Accumulation Rates (OCARs)

670 | The calculated organic carbon accumulation rates for Lake Rauchaugytgyn represent the first calculated values for an Arctic Siberian glacial lake. As such, limited comparable studies exist and are restricted to studies of Siberian thermokarst lake systems that are generally younger and smaller (Anthony et al., 2014) and to the much larger Lake Baikal (Martin et al., 1998). Comparisons must therefore be additionally drawn to boreal and glacial lakes from North America and northern Europe, as well as to proglacial and bedrock-catchment lakes from Greenland. Significant differences however exist between these lake systems relating to contrasting environmental conditions prevailing at different latitudes as well as high variability with regards to lake and catchment spatial extent, lake water depth and catchment environmental and vegetation properties. Comparisons must therefore be drawn to Boreal lakes from North America, Greenland and northern Europe.

680 | The overall Holocene average organic carbon accumulation rate of $3.53 \text{ g OC m}^{-2} \text{ a}^{-1}$ for Lake Rauchaugytgyn is significantly higher than those reported for the Pleistocene units represented by LU-II & LU-III (AU2) of ~~1.09-08~~ $1.09-08 \text{ g OC m}^{-2} \text{ a}^{-1}$ (Figs. 8, 9a). This is in accord to increased organic productivity, carbon accumulation and generally higher SRs and MARs during the Holocene. The calculated Holocene rate is very similar to that obtained for Holocene deposits of boreal lakes from northern Quebec, Canada (mean $3.8 \text{ g OC m}^{-2} \text{ a}^{-1}$) that were also studied using an equivocal hydroacoustic approach to account for sediment focussing and to rates obtained for Lake Baikal (mean $2.8 \text{ g OC m}^{-2} \text{ a}^{-1}$) (Ferland et al., 2012; Ferland et al., 2014; Martin et al., 1998) (Figs. 8, 9a). The range of Holocene rates is on average lower but generally overlaps with Holocene organic carbon accumulation rates of small ($0.033-0.73 \text{ km}^2$) Greenlandic lakes (mean $6 \text{ g OC m}^{-2} \text{ a}^{-1}$) (Anderson et al., 2009), and to small ($0.022-0.145 \text{ km}^2$) Uinta glacial lakes, USA (mean $5.4 \text{ g OC m}^{-2} \text{ a}^{-1}$) (Munroe & Brencher, 2019). A strong resemblance is also observed when comparing to rates of accumulation calculated for Finnish ~~Boreal-boreal~~ lakes that became ~~ice-free~~ ice free at the Holocene start (Fig-s 8, 9a, 9b) (Pajunen, 2000; Kortelainen et al., 2004). The average Holocene and whole core Rauchaugytgyn rates plot well within the range of Finnish lakes and close to the mean of shallow Quebec boreal lakes when considering sediment volumes derived from sub-bottom profiling approaches or estimated from core length and lake surface area (Uinta lakes) (Fig. 9a). Recent

Kommentar [SV14]: A small note here. As mentioned before, a small typo in the calculation of the OCARS led to a very small change in a couple of the values. These have been updated where necessary.

Kommentar [SV15]: Figure 8 has been updated according to comments from reviewer #1. This included increasing the font size within the inset image to improve readability. A slight adjustment to the OCAR values for Lake Rauchaugytgyn has also been carried out associated with the small typo within the calculation of accumulation rates and updated within the figure. We have now made reference to the "Red bar" that represents Alberta lakes reported by Campbell et al. 2000 within the figure caption as pointed out by reviewer #1.

Kommentar [SV16]: Figure 9a and b have been updated and visually improved too with an increase in the font size of the "Holocene, Total and Pleistocene" labels. This has helped to increase figure readability.

Formatiert: Hochgestellt

Formatiert: Hochgestellt

Kommentar [SV17]: Figure 9b was updated with the new carbon storage values of Jenrich et al. accepted.

695 syntheses of average carbon accumulation rates within European lakes also suggest similar mean accumulation rates ca. $5.6 \text{ g OC m}^{-2} \text{ a}^{-1}$ (Kastowski et al., 2011).

Pronounced differences exist when compared with larger lake systems reported from Alberta, Canada (mean 15




Figure 8. Comparison Log-boxplot of average organic carbon accumulation rates (OCARs) for Lake Rauchaugytgn with carbon accumulation rates from other regions. Rauchaugytgn OCARs include a total of 65 downcore points and plot close to Boreal-boreal lake sites and to Lake Baikal (ESRI 2020). The global modern dataset of Mendonça et al., 2017 included OCARs-OCARs from 343 globally distributed lake sites (see Text S1). Only the average OCAR across Alberta lakes was provided in Campbell et al. 2000 and hence the average is displayed as a red bar. Study references are quoted below the study regions.

$\text{g OC m}^{-2} \text{ a}^{-1}$) (Campbell et al., 2000) and global lakes, reservoirs and peatlands (Dean & Gorham, 1998; Mendonça et al., 2017). Furthermore, average Holocene rates calculated for thermokarst lakes from the Cherskii-Kolyma Tundra, far-east Far East Siberia are markedly higher (mean $47 \text{ g OC m}^{-2} \text{ a}^{-1}$) than Rauchaugytgn rates (Anthony et al., 2014) (Fig. 8).

700

5.2.2 Sediment and carbon pools within a Siberian Arctic glacial lake basin

Kommentar [SV18]: We have updated the newly published TOC_{pool} and TOC_{density} values from Jenrich et al. accepted that were updated following revision. This is done in the text and the relevant figures.

705

710

715

720

The total sediment pool of Rauchaugytgn was estimated to 25.7 ± 1.71 Mt within a sediment volume of ca. 0.033 km^3 with an average dry bulk sediment density of 780 kg m^{-3} (0.780 g cm^{-3}). The sediment at Lake Rauchaugytgn is predominantly inorganic as represented clearly by the very low total organic carbon content of sediments Due to the very low organic carbon content of sediments, the majority of the total sediment comprises inorganic detritus. The average calculated sediment density shows high agreement with global syntheses of sediment bulk density measured from lacustrine and fluvial sediments (Avnimelech et al., 2001; Hafidason et al., 2019; Munroe & Brencher, 2019; Sekellick et al., 2013). Sediment densities are however observed to be significantly higher than the average calculated for Holocene Finnish lakes (0.25 g cm^{-3}) mostly due to greater organic carbon and water contents than those remarked for Rauchaugytgn sediments (Pajunen, 2000).

The estimated $\text{TOC}_{\text{pools}}$ (total ca. 0.26 Mt) and TOC -densities (total ca. 7.85 kg m^{-3}) of Lake Rauchaugytgn sediments presented here, represent the first assessment of carbon pools within a Siberian Arctic glacial lake. Our results show that the sediment TOC_{pool} and TOC -density is relatively small, particularly during the deposition of LU-III and LU-II that covers the late Pleistocene and is in agreement with low OCARs discussed in sect. 5.2.1. The small carbon pool is especially evident when compared with carbon budgets and sediment volumes reported for Pleistocene-Holocene age Yedoma-yedoma and thermokarst and drained thermokarst lake basin records (DTLB) from Alaska, as well as Holocene $\text{TOC}_{\text{pools}}$ and sediment volumes of Alberta glacial lakes, Canada (Fig. 9b) (Campbell et al., 2000; Jongejans et al., 2018). Pleistocene-Holocene sediments obtained from thermokarst lagoon and permafrost sediments from the Bykovsky Peninsula, northern Yakutia also show significantly higher




Figure 9. (a) 3D log scatter plot of organic carbon accumulation rate (OCAR) vs lake sediment volumes and surface areas. The average carbon accumulation rate for the Holocene at Lake Rauchaugytgn closely resembles that of Holocene-age Finnish and Quebec lakes and generally lies on a trajectory of increasing OCAR with decreasing sediment volume and area. (b) 3D log scatter plot of Carbon pool vs sediment volume and area. The Rauchaugytgn data-points are highlighted with ellipses in both plots. Rauchaugytgn displays similar carbon pools to Finnish lakes, Bykovsky thermokarst lagoons and Central Yakutian Yedoma-yedoma deposits. Rauchaugytgn Sediments-sediments possess however significantly lower carbon pools relative to volume and area than Alaskan (Baldwin Peninsula thermokarst lake and yedoma deposits and Alberta lakes.

725 | estimated TOC_{pools} (Bykovsky lagoons ~~5-65.72~~ Mt) and TOC-densities (mean ~~14-2415.29~~ kg m⁻³) relative to their
small spatial extents (Jenrich et al., ~~in-review-accepted~~; Schirrmeister et al., 2011). Resemblance is how ever
observed between the TOC_{pools} (~~0-200.23~~ Mt) and TOC-densities (~~40-4512.54~~ kg m⁻³) of Polar Fox thermokarst
lagoon (Bykovsky) and Rauchaugytgyn particularly w ithin Holocene sediment of LU-I (TOC_{pool} 0.15 Mt, TOC-
density 9.87 kg m⁻³) (Jenrich et al., ~~in-review-accepted~~). Moreover, similar values of TOC_{pools} and TOC-density
730 | relative to sediment volume have been estimated for Pleistocene–Holocene ~~Yedoma-yedoma~~ and ~~aA~~las
sediments from central Yakutia, eastern Russia (~~Yedomayedoma~~: 0.057 Mt, 5.27 ~~Kg-kg~~ m⁻³, ~~aA~~las: 0.032 Mt,
6.07 ~~Kg-kg~~ m⁻³) that are closer to the range of the Pleistocene values calculated at Rauchaugytgyn (0.1 Mt, 5.65
kg m⁻³) (Windirsch et al., 2020). The highest degree of similarity is achieved w ith comparisons of TOC_{pools},
730 | surface areas and sediment volumes to Finnish lakes w hereby Lake Rauchaugytgyn pools lie on a distinct trend
in 3D logarithmic space w ithin the range and around the mean of Finnish lakes (Fig. 9b) (Pajunen et al., 2000;
Kortelainen et al. 2004). Estimated carbon pools from Uinta glacial lakes show ~~markedly lower values that~~
~~showsmarkedly lower values that show~~ they store significantly less carbon w ithin their smaller sediment volume
(Munroe and Brencher, 2019).

735 | 5.2.3 Regional and local controls on carbon accumulation and methodological limitations

Lake Rauchaugytgyn stores predominantly inorganic sediment detritus and demonstrates carbon dynamics not
strongly dissimilar from other high-latitude lake systems. Differences are however revealed when comparisons of
carbon dynamics are made to thermokarst and drained thermokarst lakes (DTLB), large Alberta glacial lakes and
global lakes in general. The interplay betw een many, varied factors likely contribute to major differences in
740 | carbon accumulation dynamics between different deposits and regions that are beyond a major discussion here
~~and have been broadly considered elsewhere~~ (Ferland et al., 2012; Kastowski et al., 2011; Sobek et al., 2009).
Carbon accumulation at Lake Rauchaugytgyn is how ever likely affected by several key factors that can be
identified.

745 | Its mountainous, high-latitude, tundra location represents cold and dry environmental conditions that provide a
first order climatic limit on catchment and lake internal primary productivity and hence carbon production. The
expression of a cold climate is marked by open herb- and graminoid tundra vegetation dominated by ~~Poaceae~~
and *Dryas octopetala* L. with the absence of vegetation at higher altitudes on lake proximal slopes and w ithin the
mountainous lake catchment (Shevtsova et al., 2020). The occurrence of a well-developed lake surface ice-layer
up to ca. 1.8 m in thickness that is present for ca. 9 months additionally contributes to reduced rates of algal
750 | primary productivity by limiting the availability of light as w ell as the allochthonous input of particulate (POC) and
dissolved organic carbon (DOC) via fluvial and/or ~~Aeolian-aeolian~~ pathw ays (Baron et al., 1991; Nolan et al.,
2002; Woolway & Merchant, 2019). An additional contributing factor to reduced catchment carbon export to Lake
Rauchaugytgyn is likely connected to the steep, mountainous topography and poor soil development w ithin the
lake catchment. Catchment soil development is limited primarily to thin cryosols but large areas of barren land
755 | exist at higher altitudes (Bünseler, 2019; Shevtsova et al., 2020). D´Arcy and Carignan (1997) suggested that
lakes situated in steep catchments with thin soils demonstrate low DOC exports to lake systems that may lead to
reduced carbon loading and hence reduced carbon accumulation. Low modern carbon export and production is
supported by the extremely low DOC contents of lake surface water samples at Lake Rauchaugytgyn (Table S1)
and at other regional lake sites (Huang et al., 2020) that support low organic matter export from the catchment
760 | (Bouchard et al., 2016). These factors were plausibly enhanced under more strongly continental, colder and drier
Pleistocene climatic conditions that led to extremely low OCARs and a small carbon pool as evidenced by the

Formatiert: Schriftart: Kursiv

multiple environmental proxies discussed throughout sect. 5.1 (Fig. 9) (Anderson & Lozhkin, 2015; Einsele et al., 2001). Changes in sedimentation rates due to temporal variability in available moisture may also regulate the preservation of organic carbon within lacustrine sediments by affecting the time available for oxygen exposure and hence microbial degradation (Einsele et al., 2001). Higher Holocene sedimentation rates at Lake Rauchaugytgn resulting from increased precipitation, active layer thickness and aeolian input may thus have further contributed to enhanced Holocene carbon accumulation. Lake area likely also plays a role in modulating differences in accumulation between lake sites and regions (Kastowski et al., 2011). Generally lakes of greater surface area possess lower rates of carbon accumulation due to reduced carbon loading (Ferland et al., 2012) (Fig. 9a). As geomorphic systems are hierarchical and operate at multiple spatio-temporal scales according to the concept of geomorphic connectivity, the diverse linkages and interrelationships between different catchment components and processes likely plays a large and complex role in regulating sediment and carbon dynamics at Lake Rauchaugytgn (Singh et al., 2021). Anthropogenic factors can also impact upon organic matter dynamics by changing the land use dynamics within lake catchments (Biskaborn et al., 2021). However, the current pristine nature of the Lake Rauchaugytgn catchment and the low occurrence of human activity, limits the anthropogenic influence when compared with lake catchments from more populous, lower latitude regions (Kastowski et al., 2011). Nonetheless, the future impact of exploration and mining operations within the catchment and of climate change within the permafrost landscape could lead to future changes ~~to~~in sediment and carbon dynamics that are yet to be seen (Tranvik et al., 2009).

Kommentar [SV19]: Introducing the full concept of geomorphic connectivity would be beyond the scope of our manuscript and would increase the length. As such, we briefly mention the concept here and the complexity of interacting processes according to reviewer #2 comments.

Differences in accumulation calculation and hence rates can also arise due to diverging methodological approaches that can lead to significant bias. The necessary usage of empirical equations to estimate sediment volumes (Campbell et al., 2000) owed to missing data or the simplification of sediment volumes to a “box” form by combining deposit thickness with area (Jongejans et al., 2018; Windirsch et al., 2020) can lead to the introduction of very large uncertainty regarding pool calculations. Volume uncertainty within our study has been ~~reduced~~minimised through the usage of hydroacoustic methods. Despite this, hydroacoustic methods can be limited to a lesser degree by acoustic blanking and hence data absence in some areas and difficulty in tracing sometimes poorly resolved reflectors. Further methodological uncertainty can arise through oversimplified estimation of dry bulk density (DBD) from empirical equations of DBD and carbon content where discrete, volumetric measurements do not exist (Avnimelech et al., 2001; Kastowski et al., 2011) as well as varied approaches used for the measurement of sample carbon contents (Elemental analyser vs LOI) (Munroe and Brencher, 2019). Limitations associated with radiocarbon dating of Arctic glacial lakes due to the absence of appreciable amounts of datable organic material, particularly during glacial phases as well as the reworking of old organic carbon present within catchment sediments that can be brought to the lake basin through permafrost and glacial processes during warm and cold phases respectively, can additionally lead to uncertainty with regards to sediment ages and hence reconstructed accumulation rates (Abbott & Stafford, 1996; Björck & Wohlfarth 2002; Oswald et al., 2005). ~~from empirical equations of DBD and carbon content where discrete, volumetric measurements do not exist (Avnimelech et al., 2001; Kastowski et al., 2011), varied approaches used for the measurement of sample carbon contents (Elemental analyser vs LOI) and radiocarbon dating limitations that limit effective sedimentation rate derivations (Munroe and Brencher, 2019).~~ Despite this, the calculated OCARs and pools for Lake Rauchaugytgn are largely agreeable with multiple studies from northern ~~Regions~~regions and reinforce the reliability, ~~and~~applicability and novelty of the presented results.

Kommentar [SV20]: We have added more information regarding dating uncertainty here in response to comments and suggestions from reviewer #1.

6. Conclusions:

This study aimed to improve the understanding of accumulation rates and pools within a palaeoenvironmental context ~~within aof a~~ Chukotkan Arctic glacial lake. The major outcomes are as follows:

- Sediment and carbon accumulation at Lake Rauchuagytgyn since MIS2 can be partitioned into three principal regimes (early MIS2, ~~Mid MIS2-early MIS1~~ ~~Mid-to-late MIS2~~ and Holocene accumulation) distinguished clearly from multiple environmental proxy records and calculated accumulation rates and pools.
- Early MIS2 accumulation (ca. 29–23.4 cal. ka BP) was controlled by cold glacial-periglacial conditions, quasi-permanent lake surface ice cover, the presence of a catchment glacier, and low sediment availability that led generally to low mass (MARs) and organic carbon accumulation rates (OCARs) and ~~glæigenic-glacially-derived minerogenic~~ sediment deposition.
- ~~Mid MIS2-early MIS1~~ ~~Mid-to-late MIS2~~ accumulation (ca. 23.4–11.5–69 cal. ka BP) likely reflects the increasing influence of paraglacial processes, longer surface ice-free summers, a thickening catchment active layer and increasing moisture availability. Carbon accumulation increased throughout and accompanied progressive climate amelioration.
- Holocene (ca. 11.5–69 cal. ka BP – present) accumulation reflects a warmer and wetter climate with enhanced runoff within a deglaciated catchment that led to alluvial fan activation and enhanced catchment fluvial dynamics. Ice-free summers likely led to increased wind-driven processes leading to more oxygenated lacustrine conditions during the mid-late Holocene and sediment redistribution. Higher Holocene OCARs reflect the prevailing climatic conditions and possibly the increased preservation under enhanced sedimentation rates.
- Estimated organic carbon pools reflect the presented accumulation regimes and demonstrates the larger carbon pool preserved within Holocene sediments (0.15 ± 0.005 Mt) when compared with Pleistocene sediments (0.1 ± 0.007 Mt). The carbon pool is however significantly small in comparison to the total sediment pool (25.7 ± 1.71 Mt).
- Carbon pools estimated for Finnish lake and Yakutian yedoma sediments show the greatest similarity to pools calculated for Lake Rauchuagytgyn sediments. Baldwin Peninsula (Alaska) permafrost landscapes and Alberta glacial lakes (Canada) represent larger organic carbon pools.
- Lake Rauchuagytgyn OCARs resemble estimations for Lake Baikal, Finnish and Canadian boreal and ~~proglacial and bedrock-catchment Greenlandic lakes~~ ~~Greenlandic lakes~~. Estimated rates are however significantly lower than the global lake mean as well as Cherskii-Kolyma thermokarst lakes in Far East Russia.
- The main drivers of OCAR and carbon pools on a temporal scale are likely related to palaeoclimate variations that control catchment and within-lake processes. Spatial differences with other lake systems ~~is-are~~ related to the high-latitude, mountainous setting of Lake Rauchuagytgyn.

Data availability

Data used in this study will be accessible from the PANGAEA data repository (dataset is currently in review and a DOI will be provided upon dataset publication).

Supplement

See separate supplement file

Author contributions

BB and UH designed the study and together with LP organised field-work. SAV obtained hydroacoustic profiling data and was responsible for opening, splicing and subsampling the core as well as all biogeochemical and sedimentological analyses. He also wrote the first version of the manuscript. GP was responsible for the age-depth model and sedimentation rates. NN provided magnetic susceptibility data. BB supervised the works of SAV. BD assisted in the interpretation of hydroacoustic data. All co-authors contributed to interpretation of the results and commented on the text.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgments

We acknowledge funding from the Past Permafrost Project under the umbrella of the Earth System Knowledge Platform (ESKP) and the initiative and networking fund of the German Helmholtz Association. This research has been supported by the ERC European Union's Horizon 2020 research and innovation programme (grant agreement no. 772852, Project. *GlacialLegacy*) and the BMBF PALMOD (grant no. 01LP1510D) as well as the Russian Foundation for Basic Research (grant no. 18-45-140053) and Ministry of Science and Higher Education of the Russian Federation (grant no. FSRG-2020-0019). The authors would like to thank all members that participated in the Chukotka 2018 expedition. We additionally thank Rebecca Morawietz, Jasmin Weise and Sebastian Golinski (AWI-Potsdam) for their assistance during laboratory work. The authors are grateful for the many helpful comments and suggestions made by Daniel Nývlt and one other anonymous reviewer that helped to improve this manuscript.

10. References

- 865 Abbott, M. B. & Stafford Jr, T. W.: Radiocarbon Geochemistry of Modern and Ancient Arctic Lake Systems, Baffin Island, Canada, *Quat. Res.*, 45, 300–311, <https://doi.org/10.1006/qres.1996.0031>, 1996.
- 870 Adrian, R., O'Reilly, C. M., Zagarese, H., Baines, S. B., Hessen, D. O., Keller, W., Livingstone, D. M., Sommaruga, R., Straile, D., Van Donk, E., Weyhenmeyer, G. A., & Winder, M.: Lakes as sentinels of climate change, *Limnol. Oceanogr.*, 54, 2283–2297, https://doi.org/10.4319/lo.2009.54.6_part_2.2283, 2009.
- ~~Aitchison, J.: The statistical analysis of geochemical compositions, *J. Int. Assoc. Math. Geol.*, 16, 531–564, <https://doi.org/10.1007/BF01029316>, 1984.~~
- 875 Alin, S. R. & Johnson, T. C.: Carbon cycling in large lakes of the world: A synthesis of production, *Global Biogeochem. Cycles*, 21, GB3002, <https://doi.org/10.1029/2006GB002881>, 2007.
- 880 Anthony, K., Zimov, S., Grosse, G. et al.: A shift of thermokarst lakes from carbon sources to sinks during the Holocene epoch, *Nature*, 511, 452–456, <https://doi.org/10.1038/nature13560>, 2014.
- Andersen, D. W., Wharton, R. A. Jr., Squires S. W.: Terrigenous clastic sedimentation in Antarctic Dry Valley Lakes, *Antarct Res Ser.*, 59, 71–81, 1993.
- 885 Anderson, N.J., D'Andrea, W. and Fritz, S.C.: Holocene carbon burial by lakes in SW Greenland, *Glob. Change Biol.*, 15, 2590–2598. <https://doi.org/10.1111/j.1365-2486.2009.01942.x>, 2009.
- Anderson, P.M., Lozhkin, A.V.: Late quaternary vegetation of Chukotka (northeast Russia) implications for glacial and Holocene environments of Beringia, *Quat. Sci. Rev.*, 107, 112–128. <https://doi.org/10.1016/j.quascirev.2014.10.016>, 2015.
- 890 Andreev, A.A., Raschke, E., Biskaborn, B.K., Vyse, S.V., Courtin, J., Böhmer, T., Stoof-Leichsenring, K., Kruse, S., Pestryakova, L.A., Herzschuh, U.H.: Late Pleistocene to Holocene vegetation and climate changes in northwestern Chukotka (Far East Russia) deduced from lakes Ilirney and Rauchuagytyn pollen records, *Boreas*, ~~accepted~~2021.
- 895
- Arnaud, F., Révillon, S., Debret, M., Revel, M., Chapron, E., Jacob, J., Giguët-Covex, C., Poulénard, J., Magny, M.: Lake Bourget regional erosion patterns reconstruction reveals Holocene NW European Alps soil evolution and paleohydrology, *Quat. Sci. Rev.*, 51, 81–92. <https://doi.org/10.1016/j.quascirev.2012.07.025>, 2012.
- 900
- Asikainen, C.A., Francus, P. & Brigham-Grette, J.: Sedimentology, clay mineralogy and grain-size as indicators of 65 ka of climate change from El'gygytyn Crater Lake, Northeastern Siberia. *J. Paleolimnol.*, 37, 105–122. <https://doi.org/10.1007/s10933-006-9026-5>, 2007.
- 905
- Avnimelech, Y., Ritvo, G., Meijer, L., Kochba, M.: Water content, organic carbon and dry bulk density in flooded sediments, *Aquac. Eng.*, 25, 25–33, 2001.
- 910 Bakke, J., Lie, Øyvind, Nesje, A., Dahl, S.O., Paasche, Ø.: Utilizing physical sediment variability in glacier-fed lakes for continuous glacier reconstructions during the Holocene, northern Følgfonna, western Norway, *Holocene*, 15, 161–176. <https://doi.org/10.1191/0959683605hl797rp>, 2005.

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

- Ballantyne, C. K.: Paraglacial geomorphology, *Quat. Sci. Rev.*, 21, 18–19, 1935–2017, [https://doi.org/10.1016/S0277-3791\(02\)00005-7](https://doi.org/10.1016/S0277-3791(02)00005-7), 2002.
- 915
- Battin, T. J., S. Luysaert, L. A. Kaplan, A. K. Aufdenkempe, A. Richter, and L. J. Tranvik, L. J.: The boundless carbon cycle, *Nat. Geosci.*, 2, 598–600, <https://doi.org/10.1038/ngeo618>, 2009.
- Battarbee, R.W., Jones, V.J., Flower, R.J., Cameron, N.G., Bennion, H., Carvalho, L., Juggins, S.: Diatoms, In: *Tracking Environmental Change Using Lake Sediments*, edited by Birks, H.J.B., Last, W.M., Smol, J.P., Kluwer Academic Publishers, Dordrecht, Netherlands, 155–202, 2001.
- 920
- Baron, J., McKnight, D. & Denning, A.S.: Sources of dissolved and particulate organic material in Loch Vale Watershed, Rocky Mountain National Park, Colorado, USA, *Biogeochemistry*, 15, 89–110, <https://doi.org/10.1007/BF00003219>, 1991.
- 925
- Baster I., Girardclos S., Pugin A., Wildi W.: High-resolution seismic stratigraphy of an Holocene lacustrine delta in western Lake Geneva (Switzerland), In: *Lake Systems from the Ice Age to Industrial Time*, edited by Ariztegui, D., Wildi, W., *Eclogae Geologicae Helvetiae / Swiss Journal of Geosciences*, vol 1, Birkhäuser, Basel. https://doi.org/10.1007/978-3-0348-7992-7_3, 2003.
- 930
- Baumer, M. M., Wagner, B., Meyer, H., Leicher, N., Lenz, M., Fedorov, G., Pestryakova, L. A. & Melles, M.: Climatic and environmental changes in the Yana Highlands of north-eastern Siberia over the last c. 57 000 years, derived from a sediment core from Lake Emanda, *Boreas*, 50, 114–133. <https://doi.org/10.1111/bor.12476>, 2020.
- 935
- Berger, A. & Loutre, M.F.: Insolation values for the climate of the last 10 million years, *Quat. Sci. Rev.*, 10, 297–317, 1991.
- 940
- Birks, H.H., Birks, H.J.B.: Multi-proxy studies in palaeolimnology, *Veget Hist Archaeobot*, 15, 235–251 <https://doi.org/10.1007/s00334-006-0066-6>, 2006.
- Biskaborn, B.K., Herzschuh, U., Bolshiyarov, D., Savelieva, L., Diekmann, B.: Environmental variability in northeastern Siberia during the last ~13,300 yr inferred from lake diatoms and sediment-geochemical parameters, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 329–330, 22–36, <https://doi.org/10.1016/j.palaeo.2012.02.003>, 2012.
- 945
- Biskaborn, B.K., Herzschuh, U., Bolshiyarov, D.Y., Schwamborn, G., Diekmann, B.: Thermokarst processes and depositional events in a tundra lake, northeastern Siberia. *Permafrost. Periglac. Process.*, 24, 160–174. <https://doi.org/10.1002/ppp.1769>, 2013.
- 950
- Biskaborn, B.K., Subetto, D., Savelieva, L., Vakhrameeva, P., Hansche, A., Herzschuh, U., Klemm, J., Heinecke, L., Pestryakova, L., Meyer, H., Kuhn, G., Diekmann, B.: Late Quaternary vegetation and lake system dynamics in north-eastern Siberia: Implications for seasonal climate variability, *Quat. Sci. Rev.*, 147, 406–421. <https://doi.org/10.1016/j.quascirev.2015.08.014>, 2016.
- 955
- Biskaborn, B.K., Nazarova, L., Pestryakova, L.A., Syrykh, L., Funck, K., Meyer, H., Chaplignin, B., Vyse, S., Gorodnichev, R., Zakharov, E., Wang, R., Schwamborn, G., Bailey, H.L., Diekmann, B.: Spatial distribution of environmental indicators in surface sediments of Lake Bolshoe Toko, Yakutia, Russia. *Biogeosciences* 16, 4023–4049. <https://doi.org/10.5194/bg-16-4023-2019>, 2019.
- 960
- Biskaborn, B.K., Narancic, B., Stoof-Leichsenring, K.R., Pestryakova, L.A., Appleby, P.G., Pliposian, G.T., Diekmann, B.: Effects of climate change and industrialization on Lake Bolshoe Toko, eastern Siberia. *J. Paleolimnol.*, 65, 335–352, <https://doi.org/10.1007/s10933-021-00175-z>, 2021.
- 965

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

- 970 Björck, S. & Wohlfarth, B.: 14C chronostratigraphic techniques in paleolimnology, In: Tracking environmental change using lake sediments. Basin Analysis, Coring, and Chronological Techniques, edited by Last, W. M. & Smol, J. P., Springer, Dordrecht, Netherlands, 205–245, 2002.
- 975 Blott, S.J. & Pye, K.: GRADISTAT: A grain size distribution and statistics package for the analysis of unconsolidated sediments, *Earth Surf. Proc. Land.*, 26, 1237–1248. <https://doi.org/10.1002/esp.261>, 2001. **Feldfunktion geändert**
- 980 Bouchard, F., Francus, P., Pienitz, R., Laurion, I.: Sedimentology and geochemistry of thermokarst ponds in discontinuous permafrost, subarctic Quebec, Canada. *J. Geophys. Res.*, 116, G00M04, <https://doi.org/10.1029/2011JG001675>, 2011. **Feldfunktion geändert**
- 985 Bouchard, F., MacDonald, L.A., Turner, K.W., Thienpont, J.R., Medeiros, A.S., Biskaborn, B.K., Korosi, J., Hall, R.I., Pienitz, R., and Wolfe, B.B.: Paleolimnology of thermokarst lakes: a window into permafrost landscape evolution, *Arctic Science*, 3(2), 91–117, <https://doi.org/10.1139/as-2016-0022>, 2016. **Feldfunktion geändert**
- 990 Brigham-Grette, J., Gualtieri, L.M., Glushkova, O.Y., Hamilton, T.D., Mostoller, D., Kotov, A.: Chlorine-36 and 14C chronology support a limited last glacial maximum across central Chukotka, northeastern Siberia, and no Beringian ice sheet, *Quat. Res.*, 59, 386–398. [https://doi.org/10.1016/S0033-5894\(03\)00058-9](https://doi.org/10.1016/S0033-5894(03)00058-9), 2003. **Feldfunktion geändert**
- 995 Brosius, L.S., Walter Anthony, K.M., Treat, C.C., Lenz, J., Jones, M.C., Bret-Harte, M.S., Grosse, G.: Spatiotemporal patterns of northern lake formation since the Last Glacial Maximum, *Quat. Sci. Rev.*, 253, <https://doi.org/10.1016/j.quascirev.2020.106773>, 2021. **Feldfunktion geändert**
- 1000 Bunbury, J., Finkelstein, S. A., Bollman, J.: Holocene hydro-climatic change and effects on carbon accumulation inferred from a peat bog in the Attawapiskat River watershed, Hudson Bay Lowlands, Canada. *Quat. Res.*, 78, 275–284, 2012.
- 1005 Bünseler, L.: Interaction of active layer carbon and nitrogen content with vegetation types on a tundra to taiga transect in eastern Siberia, MSc thesis, Potsdam University, Potsdam, Germany, 2019.
- 1010 Campbell, I.D., Campbell, C., Vitt, D.H., Kelker, D., Laird, L.D., Trew, D., Kotak, B., LeClair, D., Bayley, S.: A first estimate of organic carbon storage in Holocene lake sediments in Alberta, Canada, *J. Paleolimnol.*, 24(4), 395–400, <https://doi.org/10.1023/A:1008103605817>, 2000. **Feldfunktion geändert**
- 1015 Canty, A., Ripley, B. D.: boot: Bootstrap R (S-Plus) Functions, R package version 1.3–26, 2020.
- 1020 Cole, J. J., Prairie, Y. T., Caraco, N. F., McDowell, W. H., Tranvik, L. J., Striegl, R. G., Duarte, C. M., Kortelainen, P., Downing, J. A., Middelburg J. J. and Melack, J.: Plumbing the global carbon cycle: Integrating inland waters into the terrestrial carbon pool, *Ecosystems*, 10(1), 172–185, <https://doi.org/10.1007/s10021-006-9013-8>, 2007. **Feldfunktion geändert**
- 1025 Cremer, H., Wagner, B.: The diatom flora in the ultra-oligotrophic Lake El'gygytgyn, Chukotka. *Polar Biol.*, 26, 105–114. <https://doi.org/10.1007/s00300-002-0445-0>, 2003. **Feldfunktion geändert**
- 1030 Croudace, I.W., Rothwell, R.G. (Eds.): *Micro-XRF Studies of Sediment Cores: Applications of a non-destructive tool for the environmental sciences*, Springer, Netherlands, <https://doi.org/10.1007/978-94-017-9849-5>, 2015. **Feldfunktion geändert**

1020 Cuven, S., Francus, P. & Lamoureux, S.F.: Estimation of grain size variability with micro X-ray fluorescence in laminated lacustrine sediments, Cape Bounty, Canadian High Arctic. *J Paleolimnol.*, 44, 803–817. <https://doi.org/10.1007/s10933-010-9453-1>, 2010.

Feldfunktion geändert

D'Arcy, P., and Carignan, R.: Influence of catchment topography on water chemistry in southeastern Québec shield lakes, *Can. J. Fish. Aquat. Sci.*, 54(10), 2215–2227. <https://doi.org/10.1139/f97-129>, 1997.

Feldfunktion geändert

1025 Davison, A.C., Hinkley, D.V. (Eds.): *Bootstrap Methods and Their Applications*, Cambridge University Press, Cambridge. <http://statwww.epfl.ch/davison/BMA/>, 1997.

Feldfunktion geändert

1030 Dean, W. E., and Gorham, E.: Magnitude and significance of carbon burial in lakes, reservoirs, and peatlands, *Geology*, 26(6), 535–538, [https://doi.org/10.1130/00917613\(1998\)026<0535:MASOCB>2.3.CO;2](https://doi.org/10.1130/00917613(1998)026<0535:MASOCB>2.3.CO;2), 1998.

Feldfunktion geändert

Dedkov, A.: The relationship between sediment yield and drainage basin area. *Sediment Transfer through the Fluvial System (Proceedings of a symposium held in Moscow . August 2004)*, IAHS Publ. 288, 2004.

1035 Diekmann, B., Pestryakova, L., Nazarova, L., Subetto, D., Tarasov, P., Stauch, G., Thiemann, A., Lehmkuhl, F., Biskaborn, B., Kuhn, G., Henning, D., Müller, S.: Late Quaternary lake dynamics in the Verkhoyansk Mountains of eastern Siberia: Implications for climate and glaciation history, *Polarforschung*, 86, 97–110. <https://doi.org/10.2312/polarforschung.86.2.97>, 2016.

Feldfunktion geändert

1040 ~~Dietze, E., Maussion, F., Ahlborn, M., Diekmann, B., Hartmann, K., Henkel, K., Kasper, T., Lockett, G., Opitz, S., Haberzettl, T.: Sediment transport processes across the Tibetan Plateau inferred from robust grain-size end members in lake sediments. *Clim. Past.* 10, 91–106. <https://doi.org/10.5194/cp-10-91-2014>, 2014.~~

Formatiert: Englisch (Großbritannien)

Formatiert: Englisch (Großbritannien)

Formatiert: Englisch (Großbritannien)

Formatiert: Englisch (Großbritannien)

Formatiert: Englisch (Großbritannien)

Formatiert: Englisch (Großbritannien)

Formatiert: Englisch (Großbritannien)

Formatiert: Englisch (Großbritannien)

Formatiert: Englisch (Großbritannien)

Formatiert: Englisch (Großbritannien)

1045 Doran, P.T.: Sedimentology of Colour Lake, a Nonglacial High Arctic Lake, Axel Heiberg Island, N.W.T., Canada, *Arctic and Alpine Research*, 25:4, 353–367. <https://doi.org/10.1080/00040851.1993.12003021>, 1993.

Feldfunktion geändert

Feldfunktion geändert

1050 Einola, E., Rantakari, M., Kankaala, P., Kortelainen, P., Ojala, A., Pajunen, H., Mäkelä, S., and Arvola, L.: Carbon pools and fluxes in a chain of five boreal lakes: A dry and wet year comparison, *J. Geophys. Res.*, 116, G03009. <https://doi.org/10.1029/2010JG001636>, 2011.

1055 Einsele, G., Yan, J.P., and Hinderer, M.: Atmospheric carbon burial in modern lake basins and its significance for the global carbon budget, *Glob. Planet. Change*, 30(3–4), 167–195. [https://doi.org/10.1016/S0921-8181\(01\)00105-9](https://doi.org/10.1016/S0921-8181(01)00105-9), 2001.

Feldfunktion geändert

Elias, S. A. & Brigham-Grette, J.: Glaciations: Late Pleistocene glacial events in Beringia, In: *Encyclopedia of Quaternary Science*, edited by: Elias, S. A., Elsevier, Amsterdam, 191–201, 2013.

1060 Ferland, M.E., del Giorgio, P. A., Teodoru, C. R., and Prairie, Y. T.: Long-term C accumulation and total C stocks in boreal lakes in northern Québec, *Global Biogeochem. Cycles*, 26, GB0E04, doi: [10.1029/2011GB004241](https://doi.org/10.1029/2011GB004241), 2012.

Feldfunktion geändert

1065 Ferland, M.E., Prairie, Y. T., Teodoru, C., and del Giorgio, P. A.: Linking organic carbon sedimentation, burial efficiency, and long-term accumulation in boreal lakes, *J. Geophys. Res. Biogeosci.*, 119, 836–847, doi: [10.1002/2013JG002345](https://doi.org/10.1002/2013JG002345), 2014.

Feldfunktion geändert

1070 Finkenbinder, M.S., Abbott, M.B., Edwards, M.E., Langdon, C.T., Steinman, B.A., Finney, B.P.: A 31,000 year record of paleoenvironmental and lake-level change from Harding Lake, Alaska, USA, *Quat. Sci. Rev.*, 87, 98–113, <https://doi.org/10.1016/j.quascirev.2014.01.005>, 2014.

Feldfunktion geändert

1075 [Finkenbinder, M.S., Abbott, M.B., Finney, B.P., Stoner, J.S., Dorfman, J.M.: A multi-proxy reconstruction of environmental change spanning the last 37,000 years from Burial Lake, Arctic Alaska, *Quat. Sci. Rev.*, 126, 227–241, <https://doi.org/10.1016/j.quascirev.2015.08.031>, 2015.](#)

1080 [Folk, R.L. and Ward, W.C.: Brazos River bar \[Texas\]; a study in the significance of grain size parameters. *J. Sediment. Res.*, 27 \(1\): 3–26, doi: <https://doi.org/10.1306/74D70646-2B21-11D7-8648000102C1865D>, 1957.](#)

Formatiert: Einzug:Links: 0 cm

1080 Francke, A., Wennrich, V., Sauerbrey, M., Juschus, O., Melles, M. & Brigham-Grette, J.: Multivariate statistic and time series analyses of grain-size data in quaternary sediments of Lake El'gygytyn, NE Russia, *Clim. Past*, 9, 2459–2470, <https://doi.org/10.5194/cp-9-2459-2013>, 2013.

1085 [Friedman, G.M. and Sanders, J.E. \(Eds\): Principles of Sedimentology, Wiley, New York, 1978.](#)

1090 Frikhat, Y., Hebbeln, D., Jaaidi, El., Mhammdi, N.: Reconstruction of productivity signal and deep-water conditions in Moroccan Atlantic margin (~35°N) from the last glacial to the Holocene, *Springerplus* 4, 69, <https://doi.org/10.1186/s40064-015-0853-6>, 2015.

Feldfunktion geändert

1095 Fritz, M., Unkel, I., Lenz, J., Gajewski, K., Frenzel, P., Paquette, N., Lantuit, H., Körte, L., Wetterich, S.: Regional environmental change versus local signal preservation in Holocene thermokarst lake sediments: A case study from Herschel Island, Yukon (Canada), *J Paleolimnol.*, 60, 77–96 <https://doi.org/10.1007/s10933-018-0025-0>, 2018.

Feldfunktion geändert

1100 Gentz, T., Bonk, E., Hefter, J., Grotheer, H., Meyer, V. and Mollenhauer, G.: Establishment of routine sample preparation protocols at the newly installed MICADAS 14C dating facility at AWI, AMS 14 Conference, Ottawa, 14 August 2017 - 18 August 2017.

1105 Glushkova, O. Y.: Chapter 63 - Late Pleistocene Glaciations in North-East Asia, In: *Developments in Quaternary Sciences*, edited by: Jürgen Ehlers, Philip L. Gibbard, Philip D. Hughes, Elsevier, Amsterdam, 865–875 <https://doi.org/10.1016/B978-0-444-53447-7.00063-5>, 2011.

Feldfunktion geändert

1105 Glückler, R., Herzschuh, U., Kruse, S., Andreev, A., Vyse, S. A., Winkler, B., Biskaborn, B. K., Pestryakova, L., Dietze, E.: Wildfire history of the boreal forest of southwestern Yakutia (Siberia) over the last two millennia documented by a lake-sedimentary charcoal record, *Biogeosciences Discuss.*, <https://doi.org/10.5194/bg-2020-415>, 2020.

Feldfunktion geändert

1110 Gromig, R., Wagner, B., Wennrich, V., Fedorov, G., Savelieva, L., Lebas, E., Krastel, S., Brill, D., Andreev, A., Subetto, D. & Melles, M.: Deglaciation history of Lake Ladoga (northwestern Russia) based on varved sediments, *Boreas*, 48, 330–348, <https://doi.org/10.1111/bor.12379>, 2019.

Feldfunktion geändert

1115 Gurnell, A., Hannah, D., Lawler, D.: Suspended sediment yield from glacier basins, Erosion and Sediment Yield: Global and Regional Perspectives (Proceedings of the Exeter Symposium, July 1996).

1120 Hafidason, H., Zweidorff, J. L., Baumer, M., Gyllencreutz, R., Svendsen, J. I., Gladyshev, V. & Logvina, E.: The Lastglacial and Holocene seismostratigraphy and sediment distribution of Lake Bolshoye Shchuchye, Polar Ural Mountains, Arctic Russia, *Boreas*, 48, 452–469, <https://doi.org/10.1111/bor.12387>, 2019.

Feldfunktion geändert

- Hamilton, T., Ashley, G.: Epiguruk: a late Quaternary environmental record from northwestern Alaska, *GSA Bull.*, 105, 583–602, [https://doi.org/10.1130/0016-7606\(1993\)105<0583:ealqer>2.3.co;2](https://doi.org/10.1130/0016-7606(1993)105<0583:ealqer>2.3.co;2), 1993. **Feldfunktion geändert**
- 1125 Heinecke, L., Mischke, S., Adler, K. et al.: Climatic and limnological changes at Lake Karakul (Tajikistan) during the last ~29 cal ka, *J. Paleolimnol.*, 58, 317–334, <https://doi.org/10.1007/s10933-017-9980-0>, 2017. **Feldfunktion geändert**
- 1130 Hinderer, M., Einsele, G.: The world's large lake basins as denudation-accumulation systems and implications for their lifetimes, *J Paleolimnol.*, 26, 355–372, <https://doi.org/10.1023/A:1012651232541>, 2001. **Feldfunktion geändert**
- Horiuchi K., Kobayashi K., Oda T., Nakamura T. and Fujimura C.: Climate-induced fluctuations of ¹⁰Be concentration in Lake Baikal sediments. *Nucl. Instrum. Meth. Phys. Res.*, 562–567, 2000.
- 1135 Huang, S., Herzsuh, U., Pestryakova, L. A., Zimmermann, H. H., Davydova, P., Biskaborn, B. K., Shevtsova, I., & Stoof-Leichsenring, K. R.: Genetic and morphologic determination of diatom community composition in surface sediments from glacial and thermokarst lakes in the Siberian Arctic. *J Paleolimnol.*, 64, 225–242, <https://doi.org/10.1007/s10933-020-00133-1>, 2020. **Feldfunktion geändert**
- 1140 Jenrich, M., Angelopoulos, M., Grosse, G., Overduin, P.P., Schirrmeister, L., Nitze, I., Biskaborn, B.K., Liebner, S., Grigoriev, M., Murray, A., Strauss, J.: Thermokarst Lagoons on Bykovsky Peninsula: A Core-Based Assessment of Depositional Characteristics and Estimate of Carbon Pools, *Frontiers in Earth Science*, [in review-accepted](#).
- 1145 Jongejans, L., Strauss, J., Lenz, J., Peterse, F., Mangelsdorf, K., Fuchs, M., Grosse, G.: Organic matter characteristics in Yedoma and thermokarst deposits on Baldwin Peninsula, west Alaska. *Biogeosciences*, 15, 6033–6048, <https://doi.org/10.5194/bg-15-6033-2018>, 2018. **Feldfunktion geändert**
- 1150 Jongejans, L., & Strauss, J.: Bootstrapping approach for permafrost organic carbon pool estimation, *Zenodo*, <http://doi.org/10.5281/zenodo.3734247>, 2020. **Feldfunktion geändert**
- 1155 [Karlen, W and Matthews, J.A.: Reconstructing Holocene Glacier Variations from Glacial Lake Sediments: Studies from Nordvestlandet and Jostedalbreen-Jotunheimen, Southern Norway, *Geografiska Annaler. Series A, Physical Geography*, 74, 327–348, 1992.](#) **Formatiert: Einzug:Links: 0 cm**
- Kastowski, M., Hinderer, M. & Vecsei, A.: Long-term carbon burial in European lakes: Analysis and estimate, *Glob. Biogeochem. Cycles*, 25, GB3019, <http://doi:10.1029/2010GB003874>, 2011. **Feldfunktion geändert**
- 1160 Kilian, R., Baeza, O., Breuer, S., Ríos, F., Arz, H., Lamy, F., Wirtz, J., Baque, D., Korf, P., Kremer, K., Ríos, C., Mutschke, E., Simon, M., De Pol-Holz, R., Arevalo, M., Wörner, G., Schneider, C & Casassa, G: Late Glacial and Holocene Paleogeographical and Paleoecological Evolution of the Seno Skyring and Otway Fjord Systems in the Magellan Region, *Anales Instituto Patagonia (Chile)*, 41, 5–26, <http://dx.doi.org/10.4067/S0718-686X2013000200001>, 2013. **Feldfunktion geändert**
- 1165 Kokorowski, H.D., Anderson, P.M., Mock, C.J., Lozhkin, A.V.: A re-evaluation and spatial analysis of evidence for a Younger Dryas climatic reversal in Beringia. *Quat. Sci. Rev.*, 27, 1710–1722, <https://doi.org/10.1016/j.quascirev.2008.06.010>, 2008b. **Feldfunktion geändert**
- 1170 Kokorowski, H.D., Anderson, P.M., Sletten, R.S., Lozhkin, A.V., & Brown T.A.: Late Glacial and Early Holocene Climatic Changes Based on a Multiproxy Lacustrine Sediment Record from Northeast Siberia, *Arct. Antarct. Alp. Res.*, 40:3, 497–505, [https://doi.org/10.1657/1523-0430\(07-036\)](https://doi.org/10.1657/1523-0430(07-036)), 2008a. **Feldfunktion geändert**
- 1175 Kortelainen, P., Pajunen, H., Rantakari, M and Saarnisto M.: A large carbon pool and small sink in boreal Holocene lake sediments, *Global Change Biol.*, 10(10), 1648–1653, <https://doi:10.1111/j.1365-2486.2004.00848.x>, 2004. **Feldfunktion geändert**

- 1180 Kříbek, B., Knésl, I., Rojik, P., Sýkorová, I., Martínek, K.: Geochemical history of a Lower Miocene lake, the Cypris Formation, Sokolov Basin, Czech Republic, *J Paleolimnol.*, 58, 169–190, <https://doi.org/10.1007/s10933-017-9970-2>, 2017.
- 1185 Lebas, E., Krastel, S., Wagner, B., Gromig, R., Fedorov, G., Baumer, M., Kostromina, N., Hafidason, H.: Seismic stratigraphical record of Lake Levinson-Lessing, Tayma Peninsula: evidence for ice-sheet dynamics and lake-level fluctuations since the Early Weichselian. *Boreas*, 48, 470–487, <https://doi.org/10.1111/bor.12381>, 2019.
- 1190 Lebas, E., Gromig, R., Krastel, S., Wagner, B., Fedorov, G., Görtz, C., Averages, T., Subetto, D., Naumenko, M., Melles, M.: Pre-glacial and post-glacial history of the Scandinavian Ice Sheet in NW Russia – Evidence from Lake Ladoga, *Quat. Sci. Rev.*, 251, <https://doi.org/10.1016/j.quascirev.2020.106637>, 2021.
- 1195 ~~Leemann A. Niessen F.: Holocene glacial activity and climatic variations in the Swiss Alps: reconstructing a continuous record from proglacial lake sediments, *The Holocene*, 4(3): 259–268, doi:10.1177/095968369400400305, 1994b.~~
- 1200 Lehman, J. T.: Reconstructing the rate of accumulation of lake sediment: Their effect of sediment focusing, *Quat. Res.*, 5(4), 541–550, [https://doi:10.1016/0033-5894\(75\)90015-0](https://doi.org/10.1016/0033-5894(75)90015-0), 1975.
- 1205 Lenz, M., Savelieva, L., Frolova, L., Cherezova, A., Moros, M., Baumer, M. M., Gromig, R., Kostromina, N., Nigmatullin, N., Kolka, V., Wagner, B., Fedorov, G. & Melles, M.: Lateglacial and Holocene environmental history of the central Kola region, northwestern Russia revealed by a sediment succession from Lake Imandra, *Boreas*, 50, 76–100, <https://doi.org/10.1111/bor.12465>. ISSN 0300-9483, 2020.
- 1210 Lougheed, B.C., Obrochta, S.P.: A Rapid, Deterministic Age-Depth Modeling Routine for Geological Sequences With Inherent Depth Uncertainty. *Paleoceanogr. Paleoclimatol.*, 34, 122–133, <https://doi.org/10.1029/2018PA003457>, 2019.
- 1215 Lozhkin, A.V., Anderson, P.M.: Late quaternary lake records from the Anadyr lowland, central Chukotka (Russia). *Quat. Sci. Rev.*, 68, 1–16. <https://doi.org/10.1016/j.quascirev.2013.02.007>, 2013.
- 1220 Lozhkin, A.V., Brown, T.A., Anderson, P.M., Glushkova, O.Y., Melekestsev, I. V.: The importance of radiocarbon dates and tephra for developing chronologies of Holocene environmental changes from lake sediments, North Far East, *Russ. J. Pac. Geol.*, 10, 249–262. <https://doi.org/10.1134/S1819714016040047>, 2016.
- 1225 Lozhkin, A.V., Anderson, P.M., Minyuk, P., Korzun, J., Brown, T., Pakhomov, A., Tsygankova, V., Burnatny, S., Naumov, A.: Implications for conifer glacial refugia and postglacial climatic variation in western Beringia from lake sediments of the Upper Indigirka basin, *Boreas*, 47, 938–953. <https://doi.org/10.1111/bor.12316>, 2018.
- 1230 ~~Lunkka, J.P., Saarnisto, M., Gey, V., Demidov, I., Kiseleva, V.: Extent and age of the Last Glacial Maximum in the southeastern sector of the Scandinavian Ice Sheet, *Glob. Planet. Change*, 31, 407–425. https://doi.org/10.1016/S0921-8181(01)00132-1, 2001.~~
- 1235 ~~Mangerud, J. and Svendsen, J.I.: Deglaciation chronology inferred from marine sediments in a proglacial lake basin, western Spitsbergen, Svalbard, *Boreas*, 19: 249–272. https://doi.org/10.1111/j.1502-3885.1990.tb00450.x, 1990.~~

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

- 1230 Martin, P., L. Granina, K. Martens, and B. Goddeeris.: Oxygen concentration profiles in sediments of two ancient lakes: Lake Baikal (Siberia, Russia) and Lake Malawi (East Africa), *Hydrobiologia*, 367, 163–174, <https://doi.org/10.1023/A:1003280101128>, 1998.
- 1235 Marshall, M.H., Lamb, H.F., Huws, D., Davies, S.J., Bates, R., Bloemendal, J., Boyle, J., Leng, M.J., Umer, M., Bryant, C.: Late Pleistocene and Holocene drought events at Lake Tana, the source of the Blue Nile. *Glob. Planet. Change*, 78, 147–161, <https://doi.org/10.1016/j.gloplacha.2011.06.004>, 2011.
- Mattsson, T., P. Kortelainen, and Räike, A.: Export of DOM from boreal catchments: Impacts of land use cover and climate, *Biogeochemistry*, 76, 373–394, <https://doi.org/10.1007/s10533-005-6897-x>, 2005.
- 1240 McLaren, P., Bowles, D.: The effects of sediment transport on grain-size distributions, *Int. J. Sediment Res.*, 55 (4), 457–470, <https://doi.org/10.1306/212F86FC-2B24-11D7-8648000102C1865D>, 1985.
- 1245 Melles, M., Brigham-Grette, J., Glushkova, O.Y., Minyuk, P.S., Nowaczyk, N.R., Hubberten, H.W.: Sedimentary geochemistry of core PG1351 from Lake El'gygytgyn—a sensitive record of climate variability in the East Siberian Arctic during the past three glacial-interglacial cycles. *J. Paleolimnol.*, 37, 89–104, <https://doi.org/10.1007/s10933-006-9025-6>, 2007.
- 1250 Melles, M., Brigham-Grette, J., Minyuk, P.S., Nowaczyk, N.R., Wennrich, V., DeConto, R.M., Anderson, P.M., Andreev, A.A., Coletti, A., Cook, T.L., Haltia-Hovi, E., Kukkonen, M., Lozhkin, A. V., Rosén, P., Tarasov, P., Vogel, H., Wagner, B.: 2.8 Million years of arctic climate change from Lake El'gygytgyn, NE Russia, *Science*, 337, 315–320, <https://doi.org/10.1126/science.1222135>, 2012.
- 1255 Mendonça, R., Müller, R.A., Clow, D., Verpoorter, C., Raymond, P., Tranvik, L.J & Sobek, S.: Organic carbon burial in global lakes and reservoirs, *Nat. Commun.*, 8, 1694, <https://doi.org/10.1038/s41467-017-01789-6>
- 1260 Meyer, H., Derevyagin, A. Y., Siegert, C. & Hubberten, H.W.: Paleoclimate studies on Bykovsky Peninsula, North Siberia—hydrogen and oxygen isotopes in ground ice. *Polarforschung*, 70, 37–51, 2002.
- Meyers, P.A., Teranes, J.L.: Sediment organic matter, In: *Tracking Environmental Change Using Lake Sediments*, edited by Smol, J., Kluwer Academic Publishers, Dordrecht, 239–269. https://doi.org/10.1007/0-306-47670-3_9, 2005.
- 1265 Meyers, Philip A.: Preservation of Elemental and Isotopic Source Identification of Sedimentary Organic Matter. *Chem. Geol.*, 114 (3–4), 289–302, [https://doi.org/10.1016/0009-2541\(94\)90059-0](https://doi.org/10.1016/0009-2541(94)90059-0), 1994.
- 1270 Menne, M.J., Durre, I., Korzeniewski, B., McNeal, S., Thomas, K., Yin, X., Anthony, S., Ray, R., Vose, R.S., Gleason, B.E. & Houston, T.G.: Global Historical Climatology Network - Daily (GHCN-Daily), Version 3 NOAA National Climatic Data Center, <https://doi.org/10.7289/V5D21VHZ>, 2012.
- 1275 Moernaut, J., Verschuren, D., Charlet, F., Kristen, I., Fagot, M., De Batist, M.: The seismic-stratigraphic record of lake-level fluctuations in Lake Challa: Hydrological stability and change in equatorial East Africa over the last 140ka, *Earth Planet. Sci. Lett.*, 290, 214–223, <https://doi.org/10.1016/j.epsl.2009.12.023>, 2010.
- Munroe, J., Brencher, Q.: Holocene Carbon Burial in Lakes of the Uinta Mountains, Utah, USA, *Quaternary* 2019, 2, 1–13, <https://doi.org/10.3390/quat2010013>, 2019.

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Formatiert: Deutsch(Deutschland)

Formatiert: Deutsch(Deutschland)

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

- 1280 Naeher, S., Gilli, A., North, R.P., Hamann, Y., Schubert, C.J.: Tracing bottom water oxygenation with sedimentary Mn/Fe ratios in Lake Zurich, Switzerland. *Chem. Geol.*, 352, 125–133. <https://doi.org/10.1016/j.chemgeo.2013.06.006>, 2013.
- 1285 Nolan, M., Liston, G., Prokein, P., Brigham-Grette, J., Sharpton, V. L., and Huntzinger, R.: Analysis of lake ice dynamics and morphology on Lake El'gygytgyn, NE Siberia, using synthetic aperture radar (SAR) and Landsat, *J. Geophys. Res.*, 107, 8162, <https://doi.org/10.1029/2001JD000934>, 2002.
- 1290 Nolan, M., Brigham-Grette, J.: Basic hydrology, limnology, and meteorology of modern Lake El'gygytgyn, Siberia, *J. Paleolimnol.*, 37, 17–35. <https://doi.org/10.1007/s10933-006-9020-y>, 2007.
- Nowaczyk, N.R., Melles, M. & Minyuk, P.: A revised age model for core PG1351 from Lake El'gygytgyn, Chukotka, based on magnetic susceptibility variations tuned to northern hemisphere insolation variations, *J. Paleolimnol.*, 37, 65–76. <https://doi.org/10.1007/s10933-006-9023-8>, 2007.
- 1295 [Oswald, W.W., Anderson, P.M., Brown, T.A., Brubaker, L.B., Hu, F.S., Lozhkin, A.V., Tinner, W and Kaltenrieder, P.: Effects of sample mass and macrofossil type on radiocarbon dating of arctic and boreal lake sediments, *The Holocene*, 15, 758–767, 2005.](#)
- 1300 Pajunen, H.: lake sediments: their carbon store and related accumulations rates, *spec. pap. geol. surv. finl.*, 29, 39–69, 2000.
- Pace, M. L., and Prairie Y. T.: Respiration in lakes, in *Respiration in Aquatic Ecosystems*, edited by P. A. del Giorgio and P. J. L. Williams, 103–121, Oxford Univ. Press, New York, <https://doi.org/10.1093/acprof:oso/9780198527084.003.0007>, 2005.
- 1305 [Pfalz, P., Diekmann, B., Freytag, J.-C., Biskaborn, B.K.: Harmonizing heterogeneous multi-proxy data from lake systems, *Comput Geosci*, 153, <https://doi.org/10.1016/j.cageo.2021.104791>, 2021.](#)
- 1310 [Pšková, A., Roman, M., Bulínová, M., et al.: Late-Holocene palaeoenvironmental changes at Lake Esmeralda \(Vega Island, Antarctic Peninsula\) based on a multi-proxy analysis of laminated lake sediment, *The Holocene*, 2019;29\(7\), 1155–1175. doi:10.1177/0959683619838033, 2019.](#)
- 1315 [Procházka, V., Mizera, J., Kletetschka, G. et al.: Late Glacial sediments of the Stará Jímka paleolake and the first finding of Laacher See Tephra in the Czech Republic, *Int J Earth Sci \(Geol Rundsch\)* 108, 357–378, <https://doi.org/10.1007/s00531-018-1658-y>, 2019.](#)
- R Core Team.: R: A language and environment for statistical computing, R Foundation for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/>, 2013.
- 1320 Reimer, P.J., Austin, W.E.N., Bard, E., Bayliss, A., Blackwell, P.G., Bronk Ramsey, C., Butzin, M., Cheng, H., Edwards, R.L., Friedrich, M., Grootes, P.M., Guilderson, T.P., Hajdas, I., Heaton, T.J., Hogg, A.G., Hughen, K.A., Kromer, B., Manning, S.W., Muscheler, R., Palmer, J.G., Pearson, C., van der Plicht, J., Reimer, R.W., Richards, D.A., Scott, E.M., Southon, J.R., Turney, C.S.M., Wacker, L., Adolphi, F., Büntgen, U., Capano, M., Fahrni, S.M., Fogtmann-Schulz, A., Friedrich, R., Köhler, P., Kudsk, S., Miyake, F., Olsen, J., Reinig, F., Sakamoto, M., Sookdeo, A., Talamo, S.: The IntCal20 Northern Hemisphere Radiocarbon Age Calibration Curve (0–55 Cal ka BP), *Radiocarbon*, 1–33, <https://doi.org/10.1017/RDC.2020.41>, 2020.
- 1325 Regnéll, C., Hafliðason, H., Mangerud, J. & Svendsen, J. I.: Glacial and climate history of the last 24 000 years in the Polar Ural Mountains, Arctic Russia, inferred from partly varved lake sediments, *Boreas*, 48, 432–443. <https://doi.org/10.1111/bor.12369>. ISSN 0300-9483, 2019.
- 1330 Schirrmeister, L., Grosse, G., Wetterich, S., Overduin, P.P, Strauss, J., Edward A. G. Schuur, Hubberten, H. W.: Fossil Organic Matter Characteristics in Permafrost Deposits of the Northeast Siberian Arctic, *J. Geophys. Res.*, 116, G00M02. <https://doi.org/10.1029/2011JG001647>, 2011.
- 1335

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Formatiert: Englisch (Großbritannien)

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

- 1340 Sekellick, A., Banks, W., Myers, M.: Water Volume and Sediment Volume and Density in Lake Linganore between Boyers Mill Road Bridge and Bens Branch, Frederick County, Maryland. Scientific Investigations Report, 2013–5082 U.S. Department of the Interior, U.S. Geological Survey, 2013.
- 1345 Sifeddine, A., Meyers, P., Cordeiro, R., Albuquerque, A., Bernardes, M., Turcq, B., Abraão, J.: Delivery and deposition of organic matter in surface sediments of Lagoa do Caçó (Brazil), *J. Paleolimnol.*, 45, 385–396, <https://doi.org/10.1007/s10933-011-9506-0>, 2011.
- 1350 Singh, M., Sinha, R., Tandon, S.K.: *Geomorphic connectivity and its application for understanding landscape complexities: a focus on the hydro-geomorphic systems of India*, *Earth Surf. Process. Landf.*, 46, 110–130, 2020.
- 1355 Shevtsova, I., Heim, B., Kruse, S., Schröder, J., Troeva, E.I., Pestryakova, L.A., Zakharov E.S. & Herzsuh, U.: Strong shrub expansion in tundra-taiga, tree infilling in taiga and stable tundra in central Chukotka (north-eastern Siberia) between 2000 and 2017, *Environ. Res. Lett.*, <https://doi.org/10.1088/1748-9326/ab9059>, 2020.
- 1360 Smith, D., Jol, H.: Radar structure of a Gilbert-type delta, Peyto Lake, Banff National Park, Canada, *Sediment. Geol.*, 113, 195–209, [https://doi.org/10.1016/S0037-0738\(97\)00061-4](https://doi.org/10.1016/S0037-0738(97)00061-4), 1997.
- 1365 Smol, J.P., Birks, H.J.B., Last, W.M., (Eds.): *Tracking environmental change using lake sediments*, Kluwer academic publishers, New York, USA, 2002.
- 1370 Sobek, S., Tranvik, L., Prairie, Y., Kortelainen, P., Cole, Jonathan J.: Patterns and regulation of dissolved organic carbon: An analysis of 7,500 widely distributed lakes, *Limnol. Oceanogr.*, 52, <https://doi.org/10.4319/lo.2007.52.3.1208>, 2007.
- 1375 Sobek, S., Durisch-Kaiser, E., Zurbrugg, R., Wongfun, N., Wessels, M., Pasche, N., Wehrli, B. Organic carbon burial efficiency in lake sediments controlled by oxygen exposure time and sediment source, *Limnol. Oceanogr.*, 54, <https://doi.org/10.4319/lo.2009.54.6.2243>, 2009.
- 1380 Sobek, S., Anderson, N. J., Bernasconi, S. M., Sontro T. D.: Low organic carbon burial efficiency in arctic lake sediments, *J. Geophys. Res. Biogeosci.*, 119, <https://doi.org/10.1002/2014JG002612>, 2014.
- 1385 Stauch, G., Gualtieri, L.: Late Quaternary glaciations in northeastern Russia, *J Quat Sci.*, 23, 545–558, <https://doi.org/10.1002/jqs.1211>, 2008.
- 1390 Stauch, G., Lehmkuhl, F.: Quaternary glaciations in the Verkhoyansk Mountains, Northeast Siberia, *Quat. Res.*, 74, 145-155, <https://doi.org/10.1016/j.ygres.2010.04.003>, 2010.
- 1395 Strauss, J., Schirrmeister, L., Grosse, G., Wetterich, S., Ulrich, M., Herzsuh, U., and Hubberten, H.-W.: The deep permafrost carbon pool of the Yedoma region in Siberia and Alaska, *Geophys. Res. Lett.*, 40, 6165–6170, doi: [10.1002/2013GL058088](https://doi.org/10.1002/2013GL058088), 2013.
- 1400 Strunk, A., Olsen, J., Sanei, H., Rudra, A., Larsen, N.: Improving the reliability of bulk sediment radiocarbon dating, *Quat. Sci. Rev.*, 242, 106442, <https://doi.org/10.1016/j.quascirev.2020.106442>, 2020.
- 1405 Stuiver, M., Reimer, P.J., Reimer, R.W.: CALIB 8.2, available at: <http://calib.org>, 2020.

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

- 1390 Subetto, D.A., Nazarova, L.B., Pestryakova, L.A., Syrykh, L.S., Andronikov, A.V., Biskaborn, B., Diekmann, B., Kuznetsov, D.D., Sapelko, T.V., Grekov, I.M.: Paleolimnological studies in Russian northern Eurasia: A review. *Contemporary Problems of Ecology*, 10, 327-335, 2017.
- 1395 Tranvik, Lars J., Downing, John A., Cotner, James B., Loiselle, Steven A., Striegl, Robert G., Ballatore, Thomas J., Dillon, Peter, Finlay, Kerri, Fortino, Kenneth, Knoll, Lesley B., Kortelainen, Pirkko L., Kutser, Tiit, Larsen, Soren., Laurion, Isabelle, Leech, Dina M., McCallister, S. Leigh, McKnight, Diane M., Melack, John M., Overholt, Erin, Porter, Jason A., Prairie, Yves, Renwick, William H., Roland, Fabio, Sherman, Bradford S., Schindler, David W., Sobek, Sebastian, Tremblay, Alain, Vanni, Michael J., Verschoor, Antonie M., von Wachenfeldt, Eddie, Weyhenmeyer, Gesa A.: Lakes and reservoirs as regulators of carbon cycling and climate, *Limnol. Oceanogr.* 54, 2298–2314, <https://doi.org/10.4319/lo.2009.54.6.part.2.2298>, 2009.
- 1400 Tripathi, J., Rajamani, V.: Geochemistry of the loessic sediments on Delhi ridge, eastern Thar Desert, Rajasthan: implications for exogenic processes, *Chem. Geol.*, 155, 265–278. [https://doi.org/10.1016/S0009-2541\(98\)00168-5](https://doi.org/10.1016/S0009-2541(98)00168-5), 1999.
- 1405 Van der Bilt, W., Bakke, J., Vasskog, K., D'Andrea, W.J., Bradley, R.S., Ólafsdóttir, S.: Reconstruction of glacier variability from lake sediments reveals dynamic Holocene climate in Svalbard. *Quat. Sci. Rev.*, 126, 15, 201–218. <https://doi.org/10.1016/j.quascirev.2015.09.003>, 2015.
- 1410 ~~Van den Boogaart, K. G., Tolosana-Delgado, R. and Bren, M.: compositions: Compositional Data Analysis. R package version 2.0-0, available at <https://CRAN.R-project.org/package=compositions> 2020.~~
- 1415 Vologina, E.G., Granin, N.G., Lomonosova, T.K., Vorobyeva, SS., Kulikova, N.A., Kalashnikova, I.A., Granina, L.Z.: Input of silt-sand material to the central part of southern lake Baikal by ice transportation, BAIK-SED-2, Gent University, Belgium, 17–18, 2003.
- 1420 Vyse, S.A., Herzsuh, U., Andreev, A.A., Pestryakova, L.A., Diekmann, B., Armitage, S.J., Biskaborn, B.K.: Geochemical and sedimentological responses of arctic glacial Lake Ilirney, chukotka (far east Russia) to palaeoenvironmental change since ~51.8 ka BP, *Quat. Sci. Rev.*, 247, 106607, <https://doi.org/10.1016/j.quascirev.2020.106607>, 2020.
- 1425 Wang, R., Zhang, Y., Wuennemann, B., Biskaborn, B.K., Yin, H., Xia, F., Zhou, L., Diekmann, B.: Linkages between Quaternary climate change and sedimentary processes in Hala Lake, northern Tibetan Plateau, China. *J. Asian Earth Sci.*, 107, 140-150, 2015.
- 1430 ~~Weltje, G.J., Tjallingii, R.: Calibration of XRF core scanners for quantitative geochemical logging of sediment cores: Theory and application, *Earth Planet. Sci. Lett.*, 274, 423–438, <https://doi.org/10.1016/j.epsl.2008.07.054>, 2008.~~
- 1435 Windirsch, T., Grosse, G., Ulrich, M., Schirrmeister, L., Fedorov, A., Konstantinov, P., Fuchs, M., Jongejans, L., Wolter, J., Opel, T., Strauss, J.: Organic carbon characteristics in ice-rich permafrost in alas and Yedoma deposits, central Yakutia, Siberia, *Biogeosciences*, 17, 3797–3814 <https://doi.org/10.5194/bg-17-3797-2020>, 2020.
- 1440 Woolway, R.I., Merchant, C.J.: Worldw ide alteration of lake mixing regimes in response to climate change, *Nat. Geosci.*, 12, 271–276. <https://doi.org/10.1038/s41561-019-0322-x>, 2019.
- 1440 Yu, Z., Loisel, J., Brosseau, D. P., Beilman, D. W., and Hunt, S. J.: Global peatland dynamics since the Last Glacial Maximum, *Geophys. Res. Lett.*, 37, L13402, <https://doi.org/10.1029/2010GL043584>, 2010.
- Zhuravlev, G.F., Kazymín, S.S., Pukalo, P.V.: State geological map of the Russian Federation, scale 1:200 000, Anjuyjsker-Chaunsker Series. Moscow (St. Petersburg)., 1999.

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert